

# Chlorine Radical-Initiated Atmospheric Oxidation of Imines: Implications for Structural Influence on the Nitrosamine Formation

Qian Xu<sup>1</sup>, Fangfang Ma<sup>1,2\*</sup>, Chang Liu<sup>1</sup>, Qiaojing Zhao<sup>1</sup>, Jingwen Chen<sup>1</sup>, Hong-Bin Xie<sup>1\*</sup>

<sup>1</sup>Key Laboratory of Industrial Ecology and Environmental Engineering (Ministry of Education), School of Environmental Science and Technology, Dalian University of Technology, Dalian 116024, China

<sup>2</sup>College of Resources and Environmental Engineering, Guizhou Provincial Key Laboratory for Prevention and Control of Emerging Contaminants, Guizhou University, Guiyang, 550025, PR China

Correspondence to: [ffma@gzu.edu.cn](mailto:ffma@gzu.edu.cn) (Fangfang Ma) and [hbxie@dlut.edu.cn](mailto:hbxie@dlut.edu.cn) (Hong-Bin Xie)

**Abstract.** Chlorine radical ( $\cdot\text{Cl}$ ) initiated oxidation of organic nitrogen compounds (ONCs) plays an important role in carcinogenic nitrosamines formation. Imines are important constituents of ONCs, primarily formed from the atmospheric oxidation of amines. However,  $\cdot\text{Cl}$ -initiated atmospheric oxidation of imines remains poorly understood. Here, we studied the reaction mechanisms and kinetics of  $\cdot\text{Cl}$ -initiated oxidation for five representative imines ( $\text{CH}_2=\text{NH}$ ,  $\text{CH}_3\text{CH}=\text{NH}$ ,  $\text{CH}_3\text{N}=\text{CH}_2$ ,  $(\text{CH}_3)_2\text{C}=\text{NH}$ ,  $\text{HN}=\text{CHCH}_2\text{OH}$ ) to elucidate their atmospheric fate and extend the limited available data of ONCs, thereby establishing a structure-activity relationship for the reactions. The calculated overall reaction rate constants ( $\times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ ) of  $\cdot\text{Cl} + \text{CH}_2=\text{NH}$ ,  $\cdot\text{Cl} + \text{CH}_3\text{CH}=\text{NH}$ ,  $\cdot\text{Cl} + \text{CH}_3\text{N}=\text{CH}_2$ ,  $\cdot\text{Cl} + (\text{CH}_3)_2\text{C}=\text{NH}$ , and  $\cdot\text{Cl} + \text{HN}=\text{CHCH}_2\text{OH}$  are 4.5, 27.2, 7.32, 44.8 and 12.6, respectively, which are consistent with the available experimental values. Importantly, our results show that  $\cdot\text{Cl}$ -initiated reactions of the NH-containing imines mainly produce N-centered radicals. These N-centered radicals exhibit various fates under tropospheric conditions: mainly reacting with NO to form nitrosamines or with  $\text{O}_2$  to form cyanide compounds, which differs substantially from the behavior of previously reported amines. The various fates of the N-centered radicals formed from imines originates from the difference in direct hydrogen abstraction reaction rate constants ( $k_{\text{O}_2}$ ) with  $\text{O}_2$  and the reaction rate ( $k_{\text{NO}}$ ) with NO, both of which are principally governed by the distinct molecular structure of N-centered radicals. The revealed reaction mechanism provides new insights into the atmospheric transformation and risks of imines, and enrich our understanding of  $\cdot\text{Cl}$ /ONCs chemistry.

## 1 Introduction

Chlorine radicals ( $\cdot\text{Cl}$ ) are key atmospheric oxidants that significantly impact air quality and radical budgets by oxidizing volatile organic compounds (VOCs), thereby contributing to the formation of  $\text{O}_3$ ,  $\cdot\text{OH}$  and secondary organic aerosol (SOA), and consequent impacts on climate (Gunthe et al., 2021; Liu et al., 2024; Wang et al., 2019; Yi et al., 2023; Cao et al., 2024; Ma et al., 2025). Historically,  $\cdot\text{Cl}$  was considered to originate mainly from heterogeneous reactions on sea salt particles (Faxon and Allen, 2013). In recent years, diverse sources of  $\cdot\text{Cl}$  precursors have been identified in urban and suburban

environments, e.g., coal combustion, biomass burning, road salt application, chlorine-based fertilizers, and automotive braking processes (Li et al., 2020; Fu et al., 2018; Li et al., 2024; Yin et al., 2022; Thornton et al., 2010; Cooke et al., 2025; Chen et al., 2025). Daytime  $\cdot\text{Cl}$  concentrations have been reported to reach up to  $10^6$  molecule  $\text{cm}^{-3}$ , comparable to typical  $\cdot\text{OH}$  concentrations (Young et al., 2014; Li et al., 2025; Wang et al., 2023). Laboratory and theoretical studies have shown that  $\cdot\text{Cl}$ -initiated oxidation of VOCs proceeds 10-100 times faster than those initiated by  $\cdot\text{OH}$  (Edwards and Young, 2024). Moreover,  $\cdot\text{Cl}$ -initiated oxidation of VOCs often follows distinct pathways and produce products that differ significantly from those of  $\cdot\text{OH}$ -initiated reactions (Chen et al., 2025; Guo et al., 2020; Wang et al., 2022). For example,  $\cdot\text{Cl}$ -initiated oxidation of VOCs could yield chlorine-containing products, potentially altering their toxicity (Fu et al., 2024; Ding et al., 2021). Thus,  $\cdot\text{Cl}$ -initiated reactions represent a crucial role in governing the fate of VOCs and warrant careful consideration.

Organic nitrogen compounds (ONCs), a major subclass of VOCs, account for approximately 65% of VOCs and are commonly detected in the atmosphere (Abudumutailifu et al., 2024). ONCs play significant roles in the formation of particles and hazardous substances, such as hydrogen cyanide (HCN), nitrous oxide ( $\text{N}_2\text{O}$ ), nitrosamines, and nitramines (Ning et al., 2022; Wang et al., 2023; Liu et al., 2021; Abudumutailifu et al., 2024; Sun et al., 2024; Elm et al., 2016). Imines ( $\text{R}_1\text{R}_2\text{C}=\text{NR}_3$ ) comprise about 16% of atmospheric ONCs (Ditto et al., 2022), which originate from multiple sources, including combustion, motor vehicles, cleaning sports equipment and the atmospheric oxidation of amines, the latter being the primary one (Waterman and Hillhouse, 2008; Zhu et al., 2022; You et al., 2022; Onel et al., 2014). In urban areas, imines have been detected at ppt levels, comparable to typical concentrations of amines (Zhu et al., 2022). Given their abundance and their role as first-generation oxidation products of amines, elucidating the atmospheric reaction mechanisms of imines is essential for accurately assessing their environmental impacts.

To date, only limited studies have investigated the atmospheric chemistry of imines (Ali, 2020; Xu et al., 2024; Ali et al., 2016; Bunkan et al., 2014; Akbar Ali et al., 2018; Bunkan et al., 2022; Almeida and Kurtén, 2022; Ditto et al., 2020; Yao et al., 2016). Atmospheric oxidation by  $\cdot\text{OH}$  and  $\cdot\text{Cl}$  are important removal pathways for imines. Previous experimental and theoretical studies have primarily examined the reaction mechanism and kinetics of  $\cdot\text{OH}$ -initiated reactions with simple imines, such as methylimine ( $\text{CH}_2\text{NH}$ ) (Bunkan et al., 2014; Akbar Ali et al., 2018), N-methylmethylimine ( $\text{CH}_3\text{NCH}_2$ ) (Bunkan et al., 2022), and cyclic imine 1,2,3,6-tetrahydropyrazine (THPyz) (Almeida and Kurtén, 2022). These studies demonstrated that the substituents on the  $\text{C}=\text{N}$  bond profoundly influence the reactivity of imines, ultimately leading to varying atmospheric impacts. Nevertheless, no previous studies have addressed  $\cdot\text{Cl}$ -initiated reactions of imines. Since  $\cdot\text{OH}$  and  $\cdot\text{Cl}$ -initiated reactions of ONCs follow distinct mechanisms and yielding different products (Xie et al., 2015; Xue et al., 2022; Xie et al., 2014), the reactivity of imines toward  $\cdot\text{Cl}$  cannot be directly inferred from their  $\cdot\text{OH}$  chemistry.

Our previous studies demonstrated that  $\cdot\text{Cl}$  exhibits a particularly strong interaction with the  $-\text{NH}_x$  group of ONCs, leading to the formation of N-centered radicals, which are recognized precursors of carcinogenic nitrosamines (Xie et al., 2017; Liu et al., 2019). Additionally, the yield of N-centered radicals is strongly dependent on the structure of ONCs. Given the electronic structures of imines,  $\cdot\text{Cl}$ -initiated atmospheric oxidation of imines has the potential to form N-centered radicals. However, the

nitrogen lone pair electrons of imines can be delocalized to some extent to their adjacent C=N bond, which tends to decrease its electron-donor ability compared with previously well-studied ONCs (Carey and Sundberg, 2007), which may alter the reaction mechanisms of  $\cdot\text{Cl} + \text{imines}$  reactions. Moreover, the subsequent reactions of N-centered radicals depended on their specific structures. Therefore, to broaden our comprehension of the atmospheric chemistry of imines and fully assess their environmental risks, it is essential to elucidate the reaction mechanism and kinetics of  $\cdot\text{Cl} + \text{imines}$  reactions.

In this study, the reaction mechanisms and kinetics of imines with  $\cdot\text{Cl}$  were investigated by selecting five simple imines {i.e., methanimine ( $\text{CH}_2=\text{NH}$ ), ethanimine ( $\text{CH}_3\text{CH}=\text{NH}$ ), N-methylmethanimine ( $\text{CH}_3\text{N}=\text{CH}_2$ ), 2-propanimine ( $(\text{CH}_3)_2\text{C}=\text{NH}$ ), and 2-iminoethanol ( $\text{HN}=\text{CHCH}_2\text{OH}$ )} as model compounds. A combination of quantum chemical calculations and kinetic modeling was employed to elucidate these reactions. The parent amines of all selected imines have been detected in the ambient atmosphere (Qiu and Zhang, 2013; Ge et al., 2011; Shen et al., 2023). For the imine radicals yielded in the initial reaction step, their subsequent reactions including isomerization/dissociation, and bimolecular reactions with key atmospheric oxidants ( $\text{O}_2$  and  $\text{NO}$ ) were investigated. These results provide valuable insights into the reaction mechanisms and kinetics of imines, thereby advancing our understanding of their atmospheric chemistry and  $\cdot\text{Cl}$  chemistry.

## 2 Computational Details

**2.1 Global Minimum Search.** The selected imines can exist in multiple distinct gas-phase configurations. To identify their global minima, a multi-step conformational sampling scheme was employed, following our earlier studies (Lu et al., 2024; Ma et al., 2019; Ma et al., 2023) To explore their conformational space, the genitor module within the Molclus program was initially used to form the range of conformations (Lu, 2025). The generated conformers were further optimized at the PM6 and MP2/6-31+G(3df,2p) level (Vereecken and Francisco, 2012). Single-point energy calculations were performed at the CCSD(T)/aug-cc-pVTZ level (Vereecken and Francisco, 2012). The structure with the lowest Gibbs free energy were determined to be the global minima and served as the initial structure for investigating the reaction mechanisms and kinetics. The corresponding global minima of five imines are presented in Figure S1.

**2.2 Electronic Structure Calculations.** All electronic structure calculations were conducted with the Gaussian 09 program package (Frisch et al., 2009). For the Cl-initiated reactions of five imines, geometry optimizations and harmonic vibrational frequency calculations were performed at the MP2/6-31+G(3df,2p) level, followed by single-point energy evaluations at the CCSD(T)/aug-cc-pVTZ level. This is consistent with our previous work on  $\cdot\text{Cl}$ -initiated reactions of ONCs systems, where the combination of MP2 and CCSD(T) methods has proven to yield reliable energies. (Ma et al., 2018). Considering the substantially increased computational cost for the subsequent reactions of the resulting imine radicals, we employed the M06-2X/6-31+G(d,p) level for geometry optimizations and frequency calculations (Zhao and Truhlar, 2008), followed by CCSD(T)/6-311+G(2df,2p) single-point energy evaluations (Pople et al., 1989). The M06-2X functional combined with CCSD(T) method has successfully been applied to predict radical +  $\text{O}_2/\text{NO}$  reactions (Fu et al., 2024). Intrinsic reaction

coordinate (IRC) calculations were used to confirm that each transition state correctly connects the relevant reactants and products at the respective optimization levels (Fukui, 1981). The isolated  $\cdot\text{Cl}$  was treated with a correction value of  $0.8 \text{ kcal mol}^{-1}$  to account for spin-orbit coupling effects (Ma et al., 2018). Natural bond orbital (NBO) analysis was conducted to characterize the atomic charges in the initial reaction transition states (Reed et al., 1985). Unless otherwise specified, the following labels are used throughout the manuscript: ‘reactants’ (R), ‘pre-reaction complex’ (RC), ‘post-reaction complex’ (PC), ‘transition states’ (TS), ‘intermediates’ (IM) and ‘products’ (P).

**2.3 Kinetics Calculations.** MultiWell 2014.1 software was employed to simulate the reaction kinetics (Barker et al., 2014; Barker, 2001). Rate constants for the multi-channel and multi-well chemical reactions involving tight transition states were calculated using Rice-Ramsberger-Kassel-Marcus (RRKM) theory (Barker, 2001), while those for barrierless reactions were obtained with long-range transition-state theory with a dispersion force potential or Inverse Laplace Transformation (ILT) model. Full computational settings are provided in our previous studies (Ma et al., 2018; Ma et al., 2021). For single-step processes, canonical transition state theory (TST) in the Thermo module of MultiWell-2014.1 program suite was applied to calculate the rate constants. Tunneling corrections for H-shift or H-abstraction reactions were included via a one-dimensional asymmetric Eckart potential (Eckart, 1930). Reaction rate constants for  $\cdot\text{Cl}$  + imines reactions were performed across the temperature range of 260 – 330 K. The parameters used in the long-range transition-state theory (LRTST) calculations and Lennard-Jones parameters for various species used in the MultiWell are shown in Table S1 and S2, respectively.

### 3 Results and Discussion

**3.1  $\cdot\text{Cl}$ -Initiated Reactions.**  $\cdot\text{Cl}$  can either abstract H atoms from the  $=\text{CH}_x-$  ( $x = 1\sim 2$ ) and  $=\text{NH}$  groups or add to the  $\text{C}=\text{N}$  bond of the target imines. The calculated energetic data are summarized in Table 1, with the corresponding zero point energy (ZPE) corrected potential energy surfaces (PES) shown in Figure S2. Based on the reaction activation energies ( $E_a$ ) in Table 1, it can be concluded that H-abstractions at the N-sites is the most energetically favorable pathway for  $\cdot\text{Cl}$ -initiated reactions of  $\text{CH}_2=\text{NH}$ ,  $\text{CH}_3\text{CH}=\text{NH}$ ,  $(\text{CH}_3)_2\text{C}=\text{NH}$ , and  $\text{HN}=\text{CHCH}_2\text{OH}$ , which is similar to amines +  $\cdot\text{Cl}$  systems (Ma et al., 2018; Xue et al., 2022; Xie et al., 2015; Xie et al., 2017). Notably, the  $E_a$  values for the formation of N-centered radicals from imines are significantly higher than those from amines. Interestingly,  $\cdot\text{OH}$ -initiated reactions of  $\text{CH}_2=\text{NH}$  can also result in the formation of N-centered radicals, unlike  $\cdot\text{OH}$ -initiated reactions of amines, although the  $E_a$  value for the  $\cdot\text{OH} + \text{CH}_2\text{NH}$  reaction being approximately  $4 \text{ kcal mol}^{-1}$  higher than that of  $\cdot\text{Cl} + \text{CH}_2=\text{NH}$  reaction (Bunkan et al., 2014). For  $\text{CH}_3\text{N}=\text{CH}_2$ , H-abstraction from the  $-\text{CH}_2$  site forming C-centered radicals is the most favorable pathway, consistent with  $\cdot\text{OH} + \text{CH}_3\text{N}=\text{CH}_2$  system (Bunkan et al., 2022).

It should be noted that, despite numerous attempts, the TSs for H-abstraction from the  $\text{CH}_2$  site of  $\text{CH}_3\text{N}=\text{CH}_2$  and for  $\cdot\text{Cl}$  addition to the N site of  $(\text{CH}_3)_2\text{C}=\text{NH}$  could not be located at the MP2/6-31+G(3df,2p) level. By analyzing the thermodynamic data, we found that  $\cdot\text{Cl}$  addition to the N-site is

thermodynamically unfavorable for the  $\cdot\text{Cl} + (\text{CH}_3)_2\text{C}=\text{NH}$  reactions, suggesting that this pathway plays a negligible role. For  $(\text{CH}_3)_2\text{C}=\text{NH}$ , the  $\text{TS}_{3,2}$  of  $\cdot\text{Cl}$  abstracting a H-atom from  $\text{CH}_2$  site was located at the MP2/6-31+G(d,p) level. To evaluate the performance of the MP2/6-31+G(d,p) method for geometry optimization and CCSD(T)/6-311+G(2df,2p) for single point energy calculation in calculating the reaction energies ( $E_a$  and  $\Delta E$ ) of H-abstraction at the  $\text{CH}_2$  site of  $\text{CH}_3\text{N}=\text{CH}_2$ , we randomly selected a pathway involved  $\text{TS}_{3,1}$  to calculate the  $E_a$  and  $\Delta E$  values using the CCSD(T)/6-311+G(2df,2p)//MP2/6-31+G(d,p) and CCSD(T)/aug-cc-pVTZ//MP2/6-31+G(3df,2p) methods. The results show that the difference in  $E_a$  and  $\Delta E$  values between the CCSD(T)/aug-cc-pVTZ//MP2/6-31+G(3df,2p) and CCSD(T)/6-311+G(2df,2p)//MP2/6-31+G(d,p) were within the quantum chemistry method ( $1.0 \text{ mol}^{-1}$ ) (Table S2). These indicate that the more efficient CCSD(T)/6-311+G(2df,2p)//MP2/6-31+G(d,p) method is reliable, and the  $E_a$  values of  $\text{TS}_{3,2}$  obtained with this method have negligible effects on the conclusions.

**Table 1. Calculated reaction activation energies ( $E_a$ , in  $\text{kcal mol}^{-1}$ ), thermodynamic energies ( $\Delta E$ ,  $\text{kcal mol}^{-1}$ ), branch ratios ( $\Gamma$ ) and reaction rate constants ( $k_{\text{Cl}}$ , in  $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ ) for the reactions of  $\cdot\text{Cl}$  with (a)  $\text{CH}_2=\text{NH}$ , (b)  $\text{CH}_3\text{CH}=\text{NH}$ , (c)  $\text{CH}_3\text{N}=\text{CH}_2$ , (d)  $(\text{CH}_3)_2\text{C}=\text{NH}$ , and (e)  $\text{HN}=\text{CHCH}_2\text{OH}$  at the CCSD(T)/aug-cc-pVTZ//MP2/6-31+G(3df,2p) level of theory.**


	Species	TS	$E_a$	$\Delta E$	$\Gamma$	$k_{\text{Cl}} \times 10^{-11}$
(a)	P <sub>1-1</sub>	TS <sub>1-1</sub>	2.35	0.08	0.26%	4.5
	P <sub>1-2</sub>	TS <sub>1-2</sub>	-0.60	-4.30	8.25%	
	P <sub>1-3</sub>	TS <sub>1-3</sub>	-4.39	-12.22	90.85%	
	P <sub>1-4</sub>	TS <sub>1-4</sub>	0.89	-13.13	0.64%	
	P <sub>1-5</sub>	TS <sub>1-5</sub>	5.14	8.97	0.00%	
(b)	P <sub>2-1</sub>	TS <sub>2-1</sub>	7.95	5.68	0.00%	27.2
	P <sub>2-2/3</sub>	TS <sub>2-2</sub>	2.26	-8.59	0.06%	
	P <sub>2-2/3</sub>	TS <sub>2-3</sub>	2.29	-8.59	0.05%	
	P <sub>2-4</sub>	TS <sub>2-4</sub>	-2.22	-5.15	11.96%	
	P <sub>2-5</sub>	TS <sub>2-5</sub>	-5.99	-10.55	87.58%	
	P <sub>2-6</sub>	TS <sub>2-6</sub>	-0.85	-11.12	0.35%	
	P <sub>2-7</sub>	TS <sub>2-7</sub>	6.57	8.46	0.00%	
(c)	P <sub>3-1</sub>	TS <sub>3-1</sub>	1.91	1.87	0.16%	7.32
	P <sub>3-2</sub>	TS <sub>3-2</sub>	-2.57 <sup>a</sup>	-1.66 <sup>a</sup>	59.37%	
	P <sub>3-3</sub>	TS <sub>3-3</sub>	-1.67	-2.01	32.98%	
	P <sub>3-4/5</sub>	TS <sub>3-4/5</sub>	1.39	-12.00	0.98%/0.98%	
	P <sub>3-6</sub>	TS <sub>3-6</sub>	-1.22	-15.02	5.53%	
	P <sub>3-7</sub>	TS <sub>3-7</sub>	4.26	8.19	0.00%	
(d)	P <sub>4-1</sub>	TS <sub>4-1</sub>	6.31	4.35	0.00%	44.8
	P <sub>4-2/3</sub>	TS <sub>4-2/3</sub>	0.89	-6.82	0.01%/0.01%	
	P <sub>4-4</sub>	TS <sub>4-4</sub>	4.19	2.94	0.00%	
	P <sub>4-5/6</sub>	TS <sub>4-5/6</sub>	1.21	-7.13	0.01%/0.01%	
	P <sub>4-7</sub>	TS <sub>4-7</sub>	-6.87	-10.35	99.96%	

	P <sub>4-8</sub>	TS <sub>4-8</sub>	-1.40	-10.53	0.00%	
	P <sub>4-9</sub>	TS <sub>4-9</sub>	—	12.43	—	
	P <sub>5-1/2</sub>	TS <sub>5-1/2</sub>	-1.13	-20.52	6.08%/6.08%	
	P <sub>5-3</sub>	TS <sub>5-3</sub>	-1.53	-2.98	39.46%	
(e)	P <sub>5-4</sub>	TS <sub>5-4</sub>	-2.59	-6.96	48.33%	12.6
	P <sub>5-5</sub>	TS <sub>5-5</sub>	15.09	9.69	0.00%	
	P <sub>5-6</sub>	TS <sub>5-6</sub>	1.16	-8.83	0.05%	
	P <sub>5-7</sub>	TS <sub>5-7</sub>	9.42	12.23	0.00%	

<sup>a</sup>Energies were calculated at the CCSD(T)/6-311+G(2df,2p)//MP2/6-31+G(d,p) level.

Since ·Cl abstracts a H-atom from sp<sup>2</sup>-N site is more favorable than from other sites, it merits further discussion why the  $E_a$  values for generating N-centered radicals are substantially lower than those for C-centered radicals. By analyzing NBO charges for all TSs, we found that larger charge transfers occurred at the most favorable transition states TS<sub>1-3</sub> (-0.428  $e$ ), TS<sub>2-5</sub> (-0.447  $e$ ), TS<sub>3-2</sub> (-0.318  $e$ ) and TS<sub>4-7</sub> (-0.456  $e$ ) than other TSs (see Table S3). This indicates that charge transfer contributes critically to the stabilization of these TSs, thereby facilitating the formation of N-centered radicals in ·Cl-initiated reactions of NH-containing imines. A similar phenomenon was observed in the piperazine (PZ) + ·Cl system (Ma et al., 2018). However, in the case of ·Cl + HN=CHCH<sub>2</sub>OH system, the charge transfer at the TS<sub>5-5</sub> (-0.430  $e$ ) is slightly larger than that at the TS<sub>5-4</sub> (the most favorable one, -0.423  $e$ ) (see Table S3). The presence of intermolecular hydrogen bonds in TS<sub>5-4</sub> may account for its lower  $E_a$  value (see Figure S3).

Using the master equation approach, the overall rate constants ( $k_{Cl}$ ) were determined to be  $4.50 \times 10^{-11}$ 、 $2.72 \times 10^{-10}$ 、 $7.32 \times 10^{-11}$ 、 $4.48 \times 10^{-10}$  and  $1.26 \times 10^{-10}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup> for ·Cl + CH<sub>2</sub>=NH, ·Cl + CH<sub>3</sub>CH=NH, ·Cl + CH<sub>3</sub>N=CH<sub>2</sub>, ·Cl + (CH<sub>3</sub>)<sub>2</sub>C=NH, and ·Cl + HN=CHCH<sub>2</sub>OH reactions at 298 K and 1 atm, respectively. The experimental  $k_{Cl}$  value available for the CH<sub>3</sub>N=CH<sub>2</sub> + ·Cl reaction is  $(1.9 \pm 0.15) \times 10^{-11}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup> (Bunkan et al., 2022), which in good consistency with the corresponding computational result ( $7.32 \times 10^{-11}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup>). This could further support the reliability of our computational approach. Over the temperature range of 260-330 K, CH<sub>2</sub>=NH and (CH<sub>3</sub>)<sub>2</sub>C=NH show positive temperature dependence for  $k_{Cl}$ , whereas CH<sub>3</sub>CH=NH, CH<sub>3</sub>N=CH<sub>2</sub> and HN=CHCH<sub>2</sub>OH exhibit a negative dependence (Figure 1a). By analyzing the substitutions effects on the  $k_{Cl}$  values, it can be found that -CH<sub>3</sub> and -(CH<sub>3</sub>)<sub>2</sub> substitutions at the >CR1= position, as well as the -CH<sub>3</sub> substitution at the =NR2 position, increase the  $k_{Cl}$  values, while substitutions at the R3CCH= position have little effect on the  $k_{Cl}$  values.



190

**Figure 1.** Reaction rate constants ( $k_{Cl}$ ) for the reactions of imines (A) and branching ratios ( $\Gamma$ ) for products for the reactions of (B)  $\text{CH}_2=\text{NH}$ , (C)  $\text{CH}_3\text{CH}=\text{NH}$ , (D)  $\text{CH}_3\text{N}=\text{CH}_2$ , (E)  $(\text{CH}_3)_2\text{C}=\text{NH}$ , and (F)  $\text{HN}=\text{CHCH}_2\text{OH}$  initiated by  $\cdot\text{Cl}$  in the temperature range of 260 – 300 K at 1 atm.

The calculated branch ratios ( $\Gamma$  values) for the N-centered radicals in the reactions of  $\cdot\text{Cl} + \text{CH}_2=\text{NH}$ ,  $\cdot\text{Cl} + \text{CH}_3\text{CH}=\text{NH}$ ,  $\cdot\text{Cl} + (\text{CH}_3)_2\text{C}=\text{NH}$  and  $\cdot\text{Cl} + \text{HN}=\text{CHCH}_2\text{OH}$  are 90.85%, 87.58%, 99.96% and 48.33% at 1 atm and 298 K, respectively. The  $\Gamma$  values of N-centered radicals show a slight decrease with increasing temperature, whereas those of other product species remain very small and negligible across the studied temperature range (Figure 1b, 1c, 1e and 1f). Therefore, N-centered radicals are the main products in these four reactions under atmospheric conditions, similar to  $\cdot\text{Cl} + \text{amines}$  systems (Ma et al., 2018; Xie et al., 2015). For the  $\cdot\text{Cl} + \text{CH}_3\text{N}=\text{CH}_2$  reaction, the calculated  $\Gamma$  values for  $P_{3-1}$ ,  $P_{3-2}$ ,  $P_{3-3}$ ,


200

$P_{3-4/5}$ ,  $P_{3-6}$ , and  $P_{3-7}$  are 0.16%, 59.36%, 32.98%, 0.98%, 5.53%, 0.00%, respectively, indicating a strong preference for the formation of C-centered radicals (Figure 1d). Since previous study have investigated the transformation of  $\text{CH}_3\text{N}=\text{CH}\cdot$  ( $P_{3-2}$ ) (Bunkan et al., 2022), we mainly considered the further transformation of the four N-centered radicals formed in the subsequent sections.

### 205 3.2 Subsequent Reactions of the formed N-centered radicals.

Consistent with previously studied N-centered radicals (Ma et al., 2018; Xie et al., 2015; Da Silva, 2013; Tang and Nielsen, 2012), the four N-centered radicals ( $\text{CH}_2=\text{N}\cdot$ ,  $\text{CH}_3\text{CH}=\text{N}\cdot$ ,  $(\text{CH}_3)_2\text{C}=\text{N}\cdot$ , and  $\cdot\text{N}=\text{CHCH}_2\text{OH}$ ) will subsequently undergo self-isomerization/dissociation or react with key atmospheric oxidants, such as  $\text{O}_2$  and  $\text{NO}$ . From the calculated ZPE-corrected PES of self-isomerization/dissociation  
210 for these four N-centered radicals (Figure S4), it can be found that the  $E_a$  values are in the range of 27.85 ~ 79.58 kcal mol<sup>-1</sup>, suggesting that both the self-isomerization and dissociation processes proceed very slowly. Therefore, in the atmosphere, these four N-centered radicals are more likely to react with  $\text{O}_2$  and  $\text{NO}$ .


Regarding the reactions of the four N-centered radicals with  $\text{O}_2$ , two distinct routes were identified. The  
215 first pathway involves  $\text{O}_2$  directly abstracting aH atom from  $=\text{CH}_x-$  ( $x = 1\sim 2$ ) and  $-\text{CH}_3$  sites, forming cyanide compounds, cyclic imines and  $\text{HO}_2\cdot$ . The second pathway proceeds by  $\text{O}_2$  addition to the  $=\text{N}$ -site, with  $E_a$  values in the range of 5.60 ~ 14.18 kcal mol<sup>-1</sup>, forming adducts with various conformations depending on the direction of  $\text{O}_2$  attack. The calculated ZPE-corrected PES for the four N-centered radicals +  $\text{O}_2$  reactions are shown in Figure 2. With the exception of  $(\text{CH}_3)_2\text{C}=\text{N}\cdot$ , the  $E_a$  values for the  
220 direct H-abstraction pathways in the  $\text{CH}_2=\text{N}\cdot + \text{O}_2$ ,  $\text{CH}_3\text{CH}=\text{N}\cdot + \text{O}_2$ , and  $\cdot\text{N}=\text{CHCH}_2\text{OH} + \text{O}_2$  reactions are 12.25, 10.86 and 10.89 kcal mol<sup>-1</sup>, respectively, significantly lower than those of addition pathways. Therefore, the H-abstraction pathway represents the primary pathway for these three reactions, producing cyanide compounds and  $\text{HO}_2\cdot$ . A similar direct H-abstraction mechanism has been observed in the  
225 reactions of N-centered radicals formed from amines with  $\text{O}_2$  (Ma et al., 2018; Xue et al., 2022; Xie et al., 2015; Xie et al., 2017). For  $(\text{CH}_3)_2\text{C}=\text{N}\cdot$ , the  $E_a$  value of  $\text{O}_2$  addition forming  $\text{IM}_{12-1}$  is lower than that for the direct H-abstraction pathways. Therefore, the formation of adduct  $\text{IM}_{12-1}$  is most favorable in the initial attack of  $\text{O}_2$ . However, the  $\text{IM}_{12-1}$  is unstable and will readily reverts to  $(\text{CH}_3)_2\text{C}=\text{N}\cdot$  and  $\text{O}_2$ , making it difficult to react with  $\text{O}_2$ .



230 **Figure 2. Schematic ZPE-corrected PES for the reactions of O<sub>2</sub> with (a) CH<sub>2</sub>=N·, (b) CH<sub>3</sub>CH=N·, (c) (CH<sub>3</sub>)<sub>2</sub>C=N·, and (d) ·N=CHCH<sub>2</sub>OH at the CCSD(T)/6-311+G(2df,2p)//M06-2X/6-31+G(d,p) level of theory.**

Our previous studies revealed that  $E_a$  values for O<sub>2</sub> directly abstracting H-atoms of N-centered radicals are highly sensitive to the employed theoretical methods, which ultimately affects the reaction rate constant ( $k_{O_2}$ ) for the reactions with O<sub>2</sub> (Liu et al., 2019; Xie et al., 2017). Since  $k_{O_2}$  is a key parameter in determining nitrosamines yields in subsequent reactions (Liu et al., 2019; Wang, 2015), we further evaluated the impact of different computational approaches on the  $E_a$  values for direct H-abstraction pathways. The  $E_a$  values obtained at CCSD(T)/aug-cc-pVTZ//MP2/6-31+G(3df,2p), CCSD(T)/aug-cc-pVTZ//MP2/aug-cc-pVTZ and CCSD(T)/6-311+G(2df,2p)//M06-2X/6-31+G(d,p) are summarized in Table S4. When the geometry optimization method was changed from M06-2X/6-31+G(d,p) to MP2/6-31+G(3df,2p) or MP2/aug-cc-pVTZ, and the single point energy calculation was changed from CCSD(T)/6-311+G(2df,2p) to CCSD(T)/aug-cc-pVTZ, the deviations in  $E_a$  values reach up to 6.2 kcal mol<sup>-1</sup> [between CCSD(T)/aug-cc-pVTZ//MP2/6-31+G(3df,2p) and CCSD(T)/6-311+G(2df,2p)//M06-2X/6-31+G(d,p)] and 6.4 kcal mol<sup>-1</sup> [between CCSD(T)/aug-cc-pVTZ//MP2/aug-cc-pVTZ and CCSD(T)/6-311+G(2df,2p)//M06-2X/6-31+G(d,p)]. By contrast, changing the optimization method from MP2/6-31+G(3df,2p) to MP2/aug-cc-pVTZ lead to only 0.3 kcal mol<sup>-1</sup> deviation of  $E_a$  values between CCSD(T)/aug-cc-pVTZ//MP2/6-31+G(3df,2p) and CCSD(T)/aug-cc-pVTZ//MP2/aug-cc-pVTZ. These results indicate that the combination of the computationally cheaper M06-2X and CCSD(T) does not provide accurate results for these systems. Therefore, the  $E_a$  values calculated at the CCSD(T)/aug-cc-pVTZ//MP2/6-31+G(3df,2p) level were adopted for subsequent kinetics calculations. Notably, H-abstraction in the CH<sub>2</sub>=N· + O<sub>2</sub>, CH<sub>3</sub>CH=N· + O<sub>2</sub>, and ·N=CHCH<sub>2</sub>OH + O<sub>2</sub> systems remain the most favorable reaction pathways even at higher CCSD(T)/aug-cc-pVTZ//MP2/6-31+G(3df,2p) methods.

Since  $E_a$  values of the direct H-abstraction pathways are notably lower than those of the addition pathways for  $\text{CH}_2=\text{N}\cdot$ ,  $\text{CH}_3\text{CHN}\cdot$  and  $\text{HOCH}_2\text{CHN}\cdot$ , the overall  $k_{\text{O}_2}$  are approximated to be the rate constants of the direct H-abstraction pathways. The TST was applied to examine the kinetics of the direct H-abstraction pathways. The calculated  $k_{\text{O}_2}$  of the direct H-abstraction pathway for  $\text{CH}_2=\text{N}\cdot + \text{O}_2$ ,  $\text{CH}_3\text{CHN}\cdot + \text{O}_2$  and  $\text{HOCH}_2\text{CHN}\cdot + \text{O}_2$  reactions based on the energies calculated at the CCSD(T)/aug-cc-pVTZ//MP2/6-31+G(3df,2p) level are  $8.94 \times 10^{-18}$ ,  $1.15 \times 10^{-17}$  and  $1.38 \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at 298 K, respectively. The  $k_{\text{O}_2}$  values of these three N-centered radicals formed from imines oxidation are similar to or higher than those of chain- and cyclic-like N-centered radicals from amines oxidation. Therefore, the  $k_{\text{O}_2}$  values for N-centered radicals with  $\text{O}_2$  varies greatly with their specific molecular structures. Previous studies have shown that the reactions of amines-derived N-centered radicals with  $\text{O}_2$  can compete with their reactions with  $\text{NO}$  under typical tropospheric  $\text{NO}$  concentrations (Ma et al., 2018). Therefore, we further investigated the reactions of these four N-centered radicals ( $\text{CH}_2=\text{N}\cdot$ ,  $\text{CH}_3\text{CHN}\cdot$ ,  $(\text{CH}_3)_2\text{C}=\text{N}\cdot$  and  $\text{HOCH}_2\text{CHN}\cdot$ ) with  $\text{NO}$ .



**Figure 3. Schematic ZPE-corrected PES for the reactions of NO with (a)  $\text{CH}_2=\text{N}\cdot$ , (b)  $\text{CH}_3\text{CH}=\text{N}\cdot$ , (c)  $(\text{CH}_3)_2\text{C}=\text{N}\cdot$ , and (d)  $\cdot\text{N}=\text{CHCH}_2\text{OH}$  at the CCSD(T)/6-311+G(2df,2p)//M06-2X/6-31+G(d,p) level of theory.**

The calculated ZPE-corrected PES for the reactions of these four N-centered radicals with  $\text{NO}$  are presented in Figure 3. As shown in Figure 3, two types of reaction pathways are observed during the initial interaction of  $\text{NO}$  with the N-centered radicals. The first is the direct H-abstraction pathway, where  $\text{NO}$  abstracts a H atom from the  $=\text{CH}_x-$  ( $x = 1\sim 2$ ) sites adjacent to the  $=\text{N}-$  and  $-\text{CH}_3$  groups. These H-abstraction pathways need to overcome at least  $13.35 \text{ kcal mol}^{-1}$   $E_a$  values to form cyanide compounds and  $\text{HNO}$ . The second is the  $\text{NO}$  addition pathway, where  $\text{NO}$  barrierlessly addition to the  $=\text{N}-$  site of the four N-centered radicals, forming nitrosamines adducts with different conformations depending on the approach direction of  $\text{NO}$ . Interconversion between these adducts, such as  $\text{IM}_{14-1}$  and  $\text{IM}_{14-2}$ ,  $\text{IM}_{15-1}$

and IM<sub>15-2</sub>, IM<sub>17-1</sub> and IM<sub>17-2</sub> proceeds with very low energy barriers of 0.53, 1.42 and 1.68 kcal mol<sup>-1</sup>, respectively.

The formed four adducts (nitrosamines) can further undergo isomerization or dissociation reactions. For IM<sub>14-1</sub>/IM<sub>14-2</sub>, IM<sub>15-1</sub>/IM<sub>15-2</sub>, IM<sub>16-1</sub>, IM<sub>17-1</sub>/IM<sub>17-2</sub>, three, five, three and four H-shift pathways are identified, respectively. It is observed that H-shifts from the =CH<sub>x</sub>- (x = 1~2) sites adjacent to the =N- and -CH<sub>3</sub> groups to the O-atom of the -NNO group are the most favorable. However, the formed four adducts need to overcome high barriers to isomerize or dissociate into fragmentation products. This mechanism is analogous to amines-derived N-centered radicals with NO. In addition, the main reaction pathway remains even at high computational methods (see Table S5 and S6).

To maintain consistency with the N-centered + O<sub>2</sub> reactions, the reaction energies calculated at the CCSD(T)/aug-cc-pVTZ//MP2/6-31+G(3df,2p) level are used to calculate the reaction rate constant ( $k_{NO}$ ) for the reactions of four N-centered radicals with NO. The calculated reaction rate constants for NO addition pathways for the reactions of CH<sub>2</sub>=N· + NO, CH<sub>3</sub>CHN· + NO, (CH<sub>3</sub>)<sub>2</sub>C=N· + NO and HOCH<sub>2</sub>CHN· + NO are  $2.20 \times 10^{-16}$ ,  $1.42 \times 10^{-12}$ ,  $1.09 \times 10^{-10}$  and  $4.30 \times 10^{-11}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup> at 298 K, respectively. These are much higher than those for the corresponding direct H-abstraction pathways ( $1.85 \times 10^{-21}$ ,  $2.87 \times 10^{-21}$ ,  $5.57 \times 10^{-27}$  and  $1.04 \times 10^{-19}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup>). Therefore, the  $k_{NO}$  can be assumed to be equal to the reaction rate constant for the addition pathways. The calculated  $\Gamma$  values for the nitrosamine (IM<sub>14-1</sub>, IM<sub>15-1</sub>, IM<sub>16-1</sub> and IM<sub>17-1</sub>) for the reactions of CH<sub>2</sub>=N· + NO, CH<sub>3</sub>CH=N· + NO, (CH<sub>3</sub>)<sub>2</sub>C=N· + NO and ·N=CHCH<sub>2</sub>OH + NO are 0.08%, 6.22%, 52.56% and 35.32% at 1 atm and 298 K, respectively. Consequently, except for CH<sub>2</sub>=N· and CH<sub>3</sub>CH=N·, the reactions of (CH<sub>3</sub>)<sub>2</sub>C=N· and ·N=CHCH<sub>2</sub>OH with NO can lead to carcinogenic nitrosamines with high yields. It is noteworthy that the  $\Gamma$  values for nitrosamine formation in these two reactions are significantly lower than those of PZ-N (99.97%), Monoethanolamine (MEA)-N (86%) and CH<sub>3</sub>NH· (~60%) with NO.

Since (CH<sub>3</sub>)<sub>2</sub>C=N· exclusively reaction with NO, we only assessed the competition of O<sub>2</sub> and NO for the three N-centered radicals using the calculated  $k_{O_2}$  and  $k_{NO}$  values. The required concentrations of NO ([NO]) to equalize the pseudo-first-order rate constants for N-centered radicals with O<sub>2</sub> are  $8.1 \times 10^6$ ,  $1.7 \times 10^3$  and 64 ppb, respectively. For CH<sub>2</sub>=N· and CH<sub>3</sub>CH=N·, the required [NO] are very high, far exceeding the typical [NO] encountered in the atmosphere (Ren, 2003). Therefore, CH<sub>2</sub>=N· and CH<sub>3</sub>CHN· are expected to primarily react with O<sub>2</sub> to form HC≡N and CH<sub>3</sub>C≡N, respectively, consistent with previous studies of CH<sub>2</sub>=NH (Bunkan et al., 2014). In contrast, for ·N=CHCH<sub>2</sub>OH, the required [NO] are achievable under polluted atmospheric conditions. Therefore, both (CH<sub>3</sub>)<sub>2</sub>C=N· and ·N=CHCH<sub>2</sub>OH can react with NO to form nitrosamines. To the best of knowledge, this study represents the first report demonstrating that N-centered radicals formed from OH-initiated reactions of imines can lead to carcinogenic nitrosamine formation. However, the yields of nitrosamine formation are highly dependent on the specific structures of N-centered radicals.

#### 4 Conclusions

Similar to previous studies on amines + ·Cl reaction systems, the reaction of imines containing =NH (CH<sub>2</sub>=NH, CH<sub>3</sub>CH=NH, (CH<sub>3</sub>)<sub>2</sub>C=NH, and HN=CHCH<sub>2</sub>OH) with ·Cl predominantly yield N-centered

radicals. The calculated  $k_{\text{Cl}}$  values for the reactions of  $\text{CH}_2=\text{NH}$ ,  $\text{CH}_3\text{CH}=\text{NH}$ ,  $\text{CH}_3\text{N}=\text{CH}_2$ ,  $(\text{CH}_3)_2\text{C}=\text{NH}$ , and  $\text{HN}=\text{CHCH}_2\text{OH}$  are  $4.50 \times 10^{-11}$ ,  $2.72 \times 10^{-10}$ ,  $7.32 \times 10^{-11}$ ,  $4.48 \times 10^{-10}$  and  $1.26 \times 10^{-10}$   $\text{cm}^3$  molecule $^{-1}$  s $^{-1}$  at 298 K and 1 atm, respectively. Among the five imines studied, only the reaction kinetics of  $\text{CH}_2=\text{NH}$  and  $\text{CH}_3\text{N}=\text{CH}_2$  with  $\cdot\text{OH}$  are available ( $\text{CH}_2=\text{NH}$ :  $3.00 \times 10^{-12}$   $\text{cm}^3$  molecule $^{-1}$  s $^{-1}$ ,  $\text{CH}_3\text{N}=\text{CH}_2$ :  $3.70 \times 10^{-12}$   $\text{cm}^3$  molecule $^{-1}$  s $^{-1}$ ). With these data, we can estimate the contribution of  $\cdot\text{Cl}$  to the degradation of  $\text{CH}_2=\text{NH}$  and  $\text{CH}_3\text{N}=\text{CH}_2$  based on the  $\cdot\text{Cl}$  concentration ( $[\cdot\text{Cl}]$ ). In the marine boundary layer, the  $[\cdot\text{Cl}]$  is typically 1–10% of the  $[\cdot\text{OH}]$ . The contribution of  $\cdot\text{Cl}$  relative to  $\cdot\text{OH}$  ( $k_{\text{Cl}}[\cdot\text{Cl}]/k_{\text{OH}}[\cdot\text{OH}]$ ) to the transformation of  $\text{CH}_2=\text{NH}$  and  $\text{CH}_3\text{N}=\text{CH}_2$  are estimated to be 15% – 150% and 20% – 200%, respectively. Furthermore, the contribution of  $\cdot\text{Cl}$  relative to  $\cdot\text{OH}$  to the formation of  $\text{CH}_2=\text{N}\cdot$  is estimated at 34% – 340% (estimated by  $k_{\text{Cl}}[\cdot\text{Cl}] \times \Gamma_{\text{N,Cl}} / k_{\text{OH}}[\cdot\text{OH}] \times \Gamma_{\text{N,OH}}$ , where  $\Gamma_{\text{N,Cl}}$  and  $\Gamma_{\text{N,OH}}$  represent the yields of N-centered radicals from the reactions initiated by  $\cdot\text{Cl}$  and  $\cdot\text{OH}$ , respectively) This clearly demonstrates the significant role of  $\cdot\text{Cl}$  in the transformation of  $\text{CH}_2=\text{NH}$  and  $\text{CH}_3\text{N}=\text{CH}_2$ . Although complete data on the reactions of other imines ( $\text{CH}_3\text{CH}=\text{NH}$ ,  $(\text{CH}_3)_2\text{C}=\text{NH}$ , and  $\text{HN}=\text{CHCH}_2\text{OH}$ ) with  $\cdot\text{OH}$  are lacking, it is reasonable to believe that  $\cdot\text{Cl}$  also plays a crucial role in their transformation based on their high  $k_{\text{Cl}}$  values .

Unlike N-centered radicals generated from amines oxidation, this study reveals that both  $k_{\text{O}_2}$  and  $k_{\text{NO}}$  values for N-centered radicals generated from imines oxidation are strongly dependent on their specific structures, ultimately affecting nitrosamine formation. This study provides the first evidence that N-centered radical formed from imines oxidation can yield nitrosamines under polluted atmospheric conditions. Therefore, to comprehensively evaluate the formation of nitrosamines from imines, further investigations into the reactions of imine-derived N-centered radicals with  $\text{O}_2$  and  $\text{NO}$  are warranted.

**Data availability.** All data were available in the main text or supplementary materials. The other relevant data are available upon request from the corresponding authors.

**Author contribution.** HBX and MFF designed research; XQ, MFF and HBX performed research; XQ, MFF, LC, ZQJ and HBX analyzed data; XQ and MFF wrote the paper; XQ, MFF, CJW and HBX reviewed and revised the paper.

**Competing interests.** The authors declare that they have no conflict of interest.

**Acknowledgements.** The study was supported by the National Natural Science Foundation of China (22206020, 22236004, 22176022), the National Key Research and Development Program of China (2022YFC3701000), and the Sugon Supercomputing Center.

## References

- Abudumutailifu, M., Shang, X., Wang, L., Zhang, M., Kang, H., Chen, Y., Li, L., Ju, R., Li, B., Ouyang, H., Tang, X., Li, C., Wang, L., Wang, X., George, C., Rudich, Y., Zhang, R., and Chen, J.: Unveiling the Molecular Characteristics, Origins, and Formation Mechanism of Reduced Nitrogen Organic Compounds in the Urban Atmosphere of Shanghai Using a Versatile Aerosol Concentration Enrichment System, *Environ. Sci. Technol.*, 58, 7099-7112, 10.1021/acs.est.3c04071, 2024.
- Akbar Ali, M., M., B., and Lin, K. C.: Catalytic effect of a single water molecule on the  $\text{OH} + \text{CH}_2\text{NH}$  reaction, *Phys. Chem. Chem. Phys.*, 20, 4297-4307, 10.1039/C7CP07091H, 2018.

- Ali, M. A.: Computational studies on the gas phase reaction of methylenimine (CH<sub>2</sub>NH) with water molecules, *Sci. Rep.*, 10, 10995, 10.1038/s41598-020-67515-3, 2020.
- 360 Ali, M. A., Sonk, J. A., and Barker, J. R.: Predicted Chemical Activation Rate Constants for HO<sub>2</sub> + CH<sub>2</sub>NH: The Dominant Role of a Hydrogen-Bonded Pre-reactive Complex, *The Journal of Physical Chemistry A*, 120, 7060-7070, 10.1021/acs.jpca.6b06531, 2016.
- Almeida, T. G., and Kurtén, T.: Atmospheric Oxidation of Imine Derivative of Piperazine Initiated by OH Radical, *ACS Earth Space Chem.*, 6, 2453-2464, 10.1021/acsearthspacechem.2c00170, 2022.
- 365 Barker, J. R.: Multiple-Well, multiple-path unimolecular reaction systems. I. MultiWell computer program suite, *Int. J. Chem. Kinet.*, 33, 232-245, 10.1002/kin.1017, 2001.
- Barker, J. R., Ortiz, N. F., Preses, J. M., Lohr, L. L., Maranzana, A., Stimac, P. J., Nguyen, T. L., and Kumar, T. J. D.: MultiWell-2014, in, edited, 1, New York, 232-245, 2014.
- 370 Bunkan, A. J. C., Reijrink, N. G., Mikoviny, T., Müller, M., Nielsen, C. J., Zhu, L., and Wisthaler, A.: Atmospheric Chemistry of *N*-Methylmethanimine (CH<sub>3</sub>N=CH<sub>2</sub>): A Theoretical and Experimental Study, *The Journal of Physical Chemistry A*, 126, 3247-3264, 10.1021/acs.jpca.2c01925, 2022.
- Bunkan, A. J. C., Tang, Y., Sellevåg, S. R., and Nielsen, C. J.: Atmospheric Gas Phase Chemistry of CH<sub>2</sub>=NH and HNC. A First-Principles Approach, *The Journal of Physical Chemistry A*, 118, 5279-5288, 10.1021/jp5049088, 2014.
- 375 Cao, Y., Liu, J., Ma, Q., Zhang, C., Zhang, P., Chen, T., Wang, Y., Chu, B., Zhang, X., Francisco, J. S., and He, H.: Photoactivation of Chlorine and Its Catalytic Role in the Formation of Sulfate Aerosols, *J. Am. Chem. Soc.*, 146, 1467-1475, 10.1021/jacs.3c10840, 2024.
- Carey, F. A., and Sundberg, R. J.: *Advanced Organic Chemistry FIFTH*, in, edited, Springer Science & Business Media, 2007.
- 380 Chen, G., Fan, X., Lin, Z., Ji, X., Chen, Z., Xu, L., and Chen, J.: Driving factors and photochemical impacts of Cl<sub>2</sub> in coastal atmosphere of Southeast China, *npj Clim. Atmos. Sci.*, 8, 135, 10.1038/s41612-025-01022-y, 2025.
- Chen, X., Jiang, Y., Zong, Z., Wang, Y., Sun, W., Wang, Y., Xia, M., Guan, L., Liu, P., Zhang, C., Chen, J., Mu, Y., and Wang, T.: Atmospheric Reactive Halogens Reshaped by the Clean Energy Policy and
- 385 Agricultural Activity in a Rural Area of the North China Plain, *Environ. Sci. Technol.*, 59, 12775-12785, 10.1021/acs.est.4c13986, 2025.
- Cooke, M. E., Dam, M., Wingen, L. M., Perraud, V., Thomas, A. E., Rojas, B., Nagalingam, S., Ezell, M. J., La Salle, S., Bauer, P. S., Finlayson-Pitts, B. J., and Smith, J. N.: Emissions of Nitrous Acid, Nitryl Chloride, and Dinitrogen Pentoxide Associated with Automotive Braking, *Environ. Sci.*
- 390 *Technol.*, 59, 9167-9177, 10.1021/acs.est.4c13202, 2025.
- Da Silva, G.: Formation of Nitrosamines and Alkyldiazohydroxides in the Gas Phase: The CH<sub>3</sub>NH + NO Reaction Revisited, *Environ. Sci. Technol.*, 47, 7766-7772, 10.1021/es401591n, 2013.
- Ding, Z., Tian, S., Dang, J., and Zhang, Q.: New mechanistic understanding for atmospheric oxidation of isoprene initiated by atomic chlorine, *Sci. Total Environ.*, 801, 149768, 10.1016/j.scitotenv.2021.149768, 2021.
- 395 Ditto, J. C., Joo, T., Slade, J. H., Shepson, P. B., Ng, N. L., and Gentner, D. R.: Nontargeted Tandem Mass Spectrometry Analysis Reveals Diversity and Variability in Aerosol Functional Groups across Multiple Sites, Seasons, and Times of Day, *Environ. Sci. Technol. Lett.*, 7, 60-69, 10.1021/acs.estlett.9b00702, 2020.

- 400 Ditto, J. C., Machesky, J., and Gentner, D. R.: Analysis of reduced and oxidized nitrogen-containing organic compounds at a coastal site in summer and winter, *Atmos. Chem. Phys.*, 22, 3045-3065, 10.5194/acp-22-3045-2022, 2022.
- Eckart, C.: The Penetration of a Potential Barrier by Electrons, *Physical Review*, 35, 1303-1309, 10.1103/PhysRev.35.1303, 1930.
- 405 Edwards, P. M., and Young, C. J.: Primary Radical Effectiveness: Do the Different Chemical Reactivities of Hydroxyl and Chlorine Radicals Matter for Tropospheric Oxidation? *ACS ES&T Air*, 1, 780-788, 10.1021/acsestair.3c00108, 2024.
- Elm, J., Jen, C. N., Kurtén, T., and Vehkamäki, H.: Strong Hydrogen Bonded Molecular Interactions between Atmospheric Diamines and Sulfuric Acid, *The Journal of Physical Chemistry A*, 120, 3693-410 3700, 10.1021/acs.jpca.6b03192, 2016.
- Faxon, C. B., and Allen, D. T.: Chlorine chemistry in urban atmospheres: a review, *Environ. Chem.*, 10, 221, 10.1071/EN13026, 2013.
- Frisch, M. J., Trucks, G. W., Schlegel, H. B., Scuseria, G. E., Robb, M. A., Cheeseman, J. R., Scalmani, G., Barone, V., Petersson, G. A., Nakatsuji, H., Li, X., Caricato, M., Marenich, A. V., Bloino, J., 415 Janesko, B. G., Gomperts, R., Mennucci, B., Hratchian, H. P., Ortiz, J. V., Izmaylov, A. F., Sonnenberg, J. L., Williams-Young, D., Ding, F., Lipparini, F., Egidi, F., Goings, J., Peng, B., Petrone, A., Henderson, T., Ranasinghe, D., Zakrzewski, V. G., Gao, J., Rega, N., Zheng, G., Liang, W., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, J., Ishida, M., Nakajima, T., Honda, Y., Kitao, O., Nakai, H., Vreven, T., Throssell, K., Jr. Montgomery, J. A., Peralta, J. E., Ogliaro, F., Bearpark, M. J., Heyd, J. J., Brothers, E. N., Kudin, K. N., Staroverov, V. N., Keith, T. A., Kobayashi, R., Normand, J., Raghavachari, K., Rendell, A. P., Burant, J. C., Iyengar, S. S., Tomasi, J., Cossi, M., Millam, J. M., Klene, M., Adamo, C., Cammi, R., Ochterski, J. W., Martin, R. L., Morokuma, K., Farkas, O., Foresman, J. B., and Fox, D. J. G. I.: GAUSSIAN 09, in, edited, *Theor Chem Account*, 215-241, 2009.
- 420 Fu, X., Wang, T., Wang, S., Zhang, L., Cai, S., Xing, J., and Hao, J.: Anthropogenic Emissions of Hydrogen Chloride and Fine Particulate Chloride in China, *Environ. Sci. Technol.*, 52, 1644-1654, 425 10.1021/acs.est.7b05030, 2018.
- Fu, Z., Guo, S., Xie, H., Zhou, P., Boy, M., Yao, M., and Hu, M.: A Near-Explicit Reaction Mechanism of Chlorine-Initiated Limonene: Implications for Health Risks Associated with the Concurrent Use of Cleaning Agents and Disinfectants, *Environ. Sci. Technol.*, 58, 19762-19773, 430 10.1021/acs.est.4c04388, 2024.
- Fukui, K.: The path of chemical reactions - the IRC approach, *Acc. Chem. Res.*, 14, 363-368, 10.1021/ar00072a001, 1981.
- Ge, X., Wexler, A. S., and Clegg, S. L.: Atmospheric amines – Part I. A review, *Atmos. Environ.*, 45, 524-546, 10.1016/j.atmosenv.2010.10.012, 2011.
- 435 Gunthe, S. S., Liu, P., Panda, U., Raj, S. S., Sharma, A., Darbyshire, E., Reyes-Villegas, E., Allan, J., Chen, Y., Wang, X., Song, S., Pöhlker, M. L., Shi, L., Wang, Y., Kommula, S. M., Liu, T., Ravikrishna, R., McFiggans, G., Mickleby, L. J., Martin, S. T., Pöschl, U., Andreae, M. O., and Coe, H.: Enhanced aerosol particle growth sustained by high continental chlorine emission in India, *Nat. Geosci.*, 14, 77-84, 10.1038/s41561-020-00677-x, 2021.
- 440 Guo, X., Ma, F., Liu, C., Niu, J., He, N., Chen, J., and Xie, H.: Atmospheric oxidation mechanism and kinetics of isoprene initiated by chlorine radicals: A computational study, *Sci. Total Environ.*, 712,

- 136330, 10.1016/j.scitotenv.2019.136330, 2020.
- Li, C., Yao, L., Wang, Y., Fang, M., Chen, X., Wang, L., Li, Y., Yang, G., and Wang, L.: Formation of chlorinated organic compounds from Cl atom-initiated reactions of aromatics and their detection in suburban Shanghai, *Atmos. Chem. Phys.*, 25, 11247-11260, 10.5194/acp-25-11247-2025, 2025.
- 445 Li, L., Yin, S., Huang, L., Yi, X., Wang, Y., Zhang, K., Ooi, C. G., and Allen, D. T.: An emission inventory for Cl<sub>2</sub> and HOCl in Shanghai, 2017, *Atmos. Environ.*, 223, 117220, 10.1016/j.atmosenv.2019.117220, 2020.
- Li, S., Liu, Y., Zhu, Y., Jin, Y., Hong, Y., Shen, A., Xu, Y., Wang, H., Wang, H., Lu, X., Fan, S., and Fan, Q.: ACEIC: a comprehensive anthropogenic chlorine emission inventory for China, *Atmos. Chem. Phys.*, 24, 11521-11544, 10.5194/acp-24-11521-2024, 2024.
- 450 Liu, C., Ma, F., Elm, J., Fu, Z., Tang, W., Chen, J., and Xie, H.: Mechanism and predictive model development of reaction rate constants for N-center radicals with O<sub>2</sub>, *Chemosphere*, 237, 124411, 10.1016/j.chemosphere.2019.124411, 2019.
- 455 Liu, L., Yu, F., Du, L., Yang, Z., Francisco, J. S., and Zhang, X.: Rapid sulfuric acid–dimethylamine nucleation enhanced by nitric acid in polluted regions, *Proceedings of the National Academy of Sciences*, 118, e2108384118, 10.1073/pnas.2108384118, 2021.
- Liu, X., Liu, L., Zhang, B., Liu, P., Huang, R., Hildebrandt Ruiz, L., Miao, R., Chen, Q., and Wang, X.: Modeling the Global Impact of Chlorine Chemistry on Secondary Organic Aerosols, *Environ. Sci. Technol.*, 58, 23064-23074, 10.1021/acs.est.4c05037, 2024.
- 460 Lu, R., Zhou, P., Ma, F., Zhao, Q., Peng, X., Chen, J., and Xie, H.: Multi-generation oxidation mechanism of m-xylene: Unexpected implications for secondary organic aerosol formation, *Atmos. Environ.*, 327, 120511, 10.1016/j.atmosenv.2024.120511, 2024.
- Lu, T. Molclus program, Version 1.9.3.: <http://www.keinsci.com/research/molclus.html>, (last accessed: 2025-12-12), 2025.
- 465 Ma, F., Ding, Z., Elm, J., Xie, H., Yu, Q., Liu, C., Li, C., Fu, Z., Zhang, L., and Chen, J.: Atmospheric Oxidation of Piperazine Initiated by ·Cl: Unexpected High Nitrosamine Yield, *Environ. Sci. Technol.*, 52, 9801-9809, 10.1021/acs.est.8b02510, 2018.
- Ma, F., Guo, X., Xia, D., Xie, H., Wang, Y., Elm, J., Chen, J., and Niu, J.: Atmospheric Chemistry of Allylic Radicals from Isoprene: A Successive Cyclization-Driven Autoxidation Mechanism, *Environ. Sci. Technol.*, 55, 4399-4409, 10.1021/acs.est.0c07925, 2021.
- 470 Ma, F., Xie, H., Elm, J., Shen, J., Chen, J., and Vehkamäki, H.: Piperazine Enhancing Sulfuric Acid-Based New Particle Formation: Implications for the Atmospheric Fate of Piperazine, *Environ. Sci. Technol.*, 53, 8785-8795, 10.1021/acs.est.9b02117, 2019.
- 475 Ma, F., Xie, H., Zhang, R., Su, L., Jiang, Q., Tang, W., Chen, J., Engsvang, M., Elm, J., and He, X.: Enhancement of Atmospheric Nucleation Precursors on Iodic Acid-Induced Nucleation: Predictive Model and Mechanism, *Environ. Sci. Technol.*, 57, 6944-6954, 10.1021/acs.est.3c01034, 2023.
- Ma, W., Feng, Z., Chen, X., Xia, M., Liu, Y., Zhang, Y., Liu, Y., Wang, Y., Zheng, F., Hua, C., Li, J., Zhao, Z., Yang, H., Kulmala, M., Worsnop, D. R., He, H., and Liu, Y.: Overlooked Significance of Reactive Chlorines in the Reacted Loss of VOCs and the Formation of O<sub>3</sub> and SOA, *Environ. Sci. Technol.*, 59, 6155-6166, 10.1021/acs.est.4c10618, 2025.
- 480 Ning, A., Liu, L., Zhang, S., Yu, F., Du, L., Ge, M., and Zhang, X.: The critical role of dimethylamine in the rapid formation of iodic acid particles in marine areas, *npj Clim. Atmos. Sci.*, 5, 1-9,

- 10.1038/s41612-022-00316-9, 2022.
- 485 Onel, L., Blitz, M., Dryden, M., Thonger, L., and Seakins, P.: Branching Ratios in Reactions of OH Radicals with Methylamine, Dimethylamine, and Ethylamine, *Environ. Sci. Technol.*, 48, 9935-9942, 10.1021/es502398r, 2014.
- Pople, J. A., Head-Gordon, M., Fox, D. J., Raghavachari, K., and Curtiss, L. A.: Gaussian-1 theory: A general procedure for prediction of molecular energies, *The Journal of Chemical Physics*, 90, 5622-5629, 10.1063/1.456415, 1989.
- 490 Qiu, C., and Zhang, R.: Multiphase chemistry of atmospheric amines, *Physical chemistry chemical physics : PCCP*, 15, 5738-5752, 10.1039/c3cp43446j, 2013.
- Reed, A. E., Weinstock, R. B., and Weinhold, F.: Natural population analysis, *The Journal of Chemical Physics*, 83, 735-746, 10.1063/1.449486, 1985.
- 495 Ren, X.: HO<sub>x</sub> concentrations and OH reactivity observations in New York City during PMTACS-NY2001, *Atmos. Environ.*, 37, 3627-3637, 10.1016/S1352-2310(03)00460-6, 2003.
- Shen, X., Chen, J., Li, G., and An, T.: A new advance in the pollution profile, transformation process, and contribution to aerosol formation and aging of atmospheric amines, *Environmental Science: Atmospheres*, 3, 444-473, 10.1039/D2EA00167E, 2023.
- 500 Sun, W., Hu, X., Fu, Y., Zhang, G., Zhu, Y., Wang, X., Yan, C., Xue, L., Meng, H., Jiang, B., Liao, Y., Wang, X., Peng, P., and Bi, X.: Different formation pathways of nitrogen-containing organic compounds in aerosols and fog water in northern China, *Atmos. Chem. Phys.*, 24, 6987-6999, 10.5194/acp-24-6987-2024, 2024.
- Tang, Y., and Nielsen, C. J.: A systematic theoretical study of imines formation from the atmospheric reactions of R<sub>n</sub>NH<sub>2</sub>-<sub>n</sub> with O<sub>2</sub> and NO<sub>2</sub> (R = CH<sub>3</sub> and CH<sub>3</sub>CH<sub>2</sub>; n = 1 and 2), *Atmos. Environ.*, 55, 185-189, <https://doi.org/10.1016/j.atmosenv.2012.02.094>, 2012.
- 505 Thornton, J. A., Kercher, J. P., Riedel, T. P., Wagner, N. L., Cozic, J., Holloway, J. S., Dubé, W. P., Wolfe, G. M., Quinn, P. K., Middlebrook, A. M., Alexander, B., and Brown, S. S.: A large atomic chlorine source inferred from mid-continental reactive nitrogen chemistry, *Nature*, 464, 271-274, 10.1038/nature08905, 2010.
- 510 Vereecken, L., and Francisco, J. S.: Theoretical studies of atmospheric reaction mechanisms in the troposphere, *Chem. Soc. Rev.*, 41, 6259-6293, [doi.org/10.1039/C2CS35070J](https://doi.org/10.1039/C2CS35070J), 2012.
- Wang, C., Liggio, J., Wentzell, J. J. B., Jorga, S., Folkerson, A., and Abbatt, J. P. D.: Chloramines as an important photochemical source of chlorine atoms in the urban atmosphere, *Proceedings of the National Academy of Sciences*, 120, e2074078176, 10.1073/pnas.2220889120, 2023.
- 515 Wang, D. S., Masoud, C. G., Modi, M., and Hildebrandt Ruiz, L.: Isoprene-Chlorine Oxidation in the Presence of NO<sub>x</sub> and Implications for Urban Atmospheric Chemistry, *Environ. Sci. Technol.*, 56, 9251-9264, 10.1021/acs.est.1c07048, 2022.
- Wang, L.: The Atmospheric Oxidation Mechanism of Benzyl Alcohol Initiated by OH Radicals: The Addition Channels, *ChemPhysChem*, 16, 1542-1550, <https://doi.org/10.1002/cphc.201500012>, 2015.
- 520 Wang, S., Zhang, Q., Wang, W., and Wang, Q.: Unexpected enhancement of sulfuric acid-driven new particle formation by alcoholic amines: The role of ion-induced nucleation, *J. Environ. Manage.*, 347, 119079, 10.1016/j.jenvman.2023.119079, 2023.
- 525 Wang, X., Jacob, D. J., Eastham, S. D., Sulprizio, M. P., Zhu, L., Chen, Q., Alexander, B., Sherwen, T., Evans, M. J., Lee, B. H., Haskins, J. D., Lopez-Hilfiker, F. D., Thornton, J. A., Huey, G. L., and Liao,

- H.: The role of chlorine in global tropospheric chemistry, *Atmos. Chem. Phys.*, 19, 3981-4003, 10.5194/acp-19-3981-2019, 2019.
- Waterman, R., and Hillhouse, G. L.:  $\eta^2$ -Organoazide Complexes of Nickel and Their Conversion to Terminal Imido Complexes via Dinitrogen Extrusion, *J. Am. Chem. Soc.*, 130, 12628-12629, 10.1021/ja805530z, 2008.
- 530 Xie, H., Li, C., He, N., Wang, C., Zhang, S., and Chen, J.: Atmospheric Chemical Reactions of Monoethanolamine Initiated by OH Radical: Mechanistic and Kinetic Study, *Environ. Sci. Technol.*, 48, 1700-1706, 10.1021/es405110t, 2014.
- Xie, H., Ma, F., Wang, Y., He, N., Yu, Q., and Chen, J.: Quantum Chemical Study on  $\cdot\text{Cl}$ -Initiated Atmospheric Degradation of Monoethanolamine, *Environ. Sci. Technol.*, 49, 13246-13255, 10.1021/acs.est.5b03324, 2015.
- 535 Xie, H., Ma, F., Yu, Q., He, N., and Chen, J.: Computational Study of the Reactions of Chlorine Radicals with Atmospheric Organic Compounds Featuring  $\text{NH}_x-\pi$ -Bond ( $x = 1, 2$ ) Structures, *The Journal of Physical Chemistry A*, 121, 1657-1665, 10.1021/acs.jpca.6b11418, 2017.
- 540 Xu, Q., Ma, F., Xia, D., Li, X., Chen, J., Xie, H., and Francisco, J. S.: Two-Step Nuncatalyzed Hydrolysis Mechanism of Imines at the Air-Water Interface, *J. Am. Chem. Soc.*, 146, 28866-28873, 10.1021/jacs.4c09080, 2024.
- Xue, J., Ma, F., Elm, J., Chen, J., and Xie, H.: Atmospheric oxidation mechanism and kinetics of indole initiated by  $\bullet\text{OH}$  and  $\bullet\text{Cl}$ : a computational study, *Atmos. Chem. Phys.*, 22, 11543-11555, 10.5194/acp-22-11543-2022, 2022.
- 545 Yao, L., Wang, M., Wang, X., Liu, Y., Chen, H., Zheng, J., Nie, W., Ding, A., Geng, F., Wang, D., Chen, J., Worsnop, D. R., and Wang, L.: Detection of atmospheric gaseous amines and amides by a high-resolution time-of-flight chemical ionization mass spectrometer with protonated ethanol reagent ions, *Atmos. Chem. Phys.*, 16, 14527-14543, 10.5194/acp-16-14527-2016, 2016.
- 550 Yi, X., Sarwar, G., Bian, J., Huang, L., Li, Q., Jiang, S., Liu, H., Wang, Y., Chen, H., Wang, T., Chen, J., Saiz Lopez, A., Wong, D. C., and Li, L.: Significant Impact of Reactive Chlorine on Complex Air Pollution Over the Yangtze River Delta Region, China, *Journal of Geophysical Research: Atmospheres*, 128, e2023JD038898, 10.1029/2023JD038898, 2023.
- Yin, S., Yi, X., Li, L., Huang, L., Ooi, M. C. G., Wang, Y., Allen, D. T., and Streets, D. G.: An Updated Anthropogenic Emission Inventory of Reactive Chlorine Precursors in China, *ACS Earth Space Chem.*, 6, 1846-1857, 10.1021/acsearthspacechem.2c00096, 2022.
- 555 You, B., Zhou, W., Li, J., Li, Z., and Sun, Y.: A review of indoor Gaseous organic compounds and human chemical Exposure: Insights from Real-time measurements, *Environ. Int.*, 170, 107611, 10.1016/j.envint.2022.107611, 2022.
- 560 Young, C. J., Washenfelder, R. A., Edwards, P. M., Parrish, D. D., Gilman, J. B., Kuster, W. C., Mielke, L. H., Osthoff, H. D., Tsai, C., Pikelnaya, O., Stutz, J., Veres, P. R., Roberts, J. M., Griffith, S., Dusanter, S., Stevens, P. S., Flynn, J., Grossberg, N., Lefer, B., Holloway, J. S., Peischl, J., Ryerson, T. B., Atlas, E. L., Blake, D. R., and Brown, S. S.: Chlorine as a primary radical: evaluation of methods to understand its role in initiation of oxidative cycles, *Atmos. Chem. Phys.*, 14, 3427-3440, 10.5194/acp-14-3427-2014, 2014.
- 565 Zhao, Y., and Truhlar, D. G.: The M06 suite of density functionals for main group thermochemistry, thermochemical kinetics, noncovalent interactions, excited states, and transition elements: two new

functionals and systematic testing of four M06-class functionals and 12 other functionals, *Theor. Chem. Acc.*, 120, 215-241, 10.1007/s00214-007-0310-x, 2008.

- 570 Zhu, S., Yan, C., Zheng, J., Chen, C., Ning, H., Yang, D., Wang, M., Ma, Y., Zhan, J., Hua, C., Yin, R., Li, Y., Liu, Y., Jiang, J., Yao, L., Wang, L., Kulmala, M., and Worsnop, D. R.: Observation and Source Apportionment of Atmospheric Alkaline Gases in Urban Beijing, *Environ. Sci. Technol.*, 56, 17545-17555, 10.1021/acs.est.2c03584, 2022.