

**Improving Simulation of Earth System Variability through Weakly
Coupled Ocean Data Assimilation in E3SM**

Pengfei Shi^{1,*}, L. Ruby Leung^{1,*}, Zhaoxia Pu², Samson Hagos¹, and Karthik Balaguru¹

¹ *Atmospheric, Climate, and Earth Sciences Division, Pacific Northwest National Laboratory,
Richland, Washington, USA*

² *Department of Atmospheric Sciences, University of Utah, Salt Lake City, Utah, USA*

* To whom correspondence may be addressed. Pengfei Shi (pengfei.shi@pnnl.gov) and L.
Ruby Leung (Ruby.Leung@pnnl.gov)

Abstract. Accurate initialization of ocean states is essential for skillful prediction of Earth system variability across seasonal-to-decadal timescales. In this study, we evaluate the impact of a newly developed four-dimensional ensemble variational (4DEnVar)-based weakly coupled ocean data assimilation (WCODA) system within the DOE Energy Exascale Earth System Model version 2 (E3SMv2) on global and regional climate variability. By assimilating monthly ocean temperature and salinity from the EN4.2.1 reanalysis into the fully coupled model, we demonstrate substantial improvements in simulating both interannual and decadal climate variability. Compared to the free-running coupled simulation, the assimilation experiment exhibits markedly enhanced interannual correlations with observations for global mean surface air temperature and precipitation anomalies. The temporal variability of key climate modes, including ENSO, the Indian Ocean Dipole, and multidecadal variability in the Pacific and Atlantic Oceans, also shows markedly improved phase agreement with observations. Regional evaluation over the contiguous United States further shows enhanced skill in simulating winter surface air temperature and precipitation, particularly in the northern and southern regions, respectively, linked to improved ENSO simulation. These findings underscore the critical role of coupled forecasts in the data assimilation cycle for propagating observational information across Earth system components. By assimilating ocean reanalysis within the fully coupled framework, the WCODA system enables cross-component information exchange among the ocean, atmosphere, and land, thereby generating physically consistent initial conditions. These improvements contribute to more accurate simulations of Earth system variability across multiple timescales and lay the foundation for providing realistic initial conditions for seasonal-to-decadal prediction applications.

Deleted: representation

Deleted: improves significantly

Deleted: representation

Deleted: integrating ocean observations

Deleted: advance the development of more reliable prediction systems in support of societal resilience

1. Introduction

Accurately representing climate variability across multiple timescales remains a major challenge in climate modeling and prediction. The initial conditions of the climate system, including the ocean, atmosphere, and land surface, play a crucial role in determining the accuracy and reliability of climate predictions (Boer et al., 2016; Taylor et al., 2012). Among these components, the ocean exhibits the longest memory and exerts a dominant influence on climate variability through its interactions with the atmosphere and other components (Wang, 2019). Therefore, improving the initialization of ocean states is widely recognized as a key approach for enhancing the skill of climate predictions (Meehl et al., 2021; Zhu et al., 2017).

Data assimilation methods have been developed to integrate observational data into coupled models, thereby generating realistic initial conditions and enhancing predictive skill. Traditionally, ocean data assimilation has been conducted in an uncoupled framework, where ocean observations are assimilated into a standalone ocean model forced by prescribed atmospheric forcing (Carton and Giese, 2008). The resulting ocean analyses are then combined with independently derived analyses from other uncoupled components (e.g., atmosphere and land) to construct the initial conditions for coupled models. However, because each component is initialized independently, this uncoupled approach often results in dynamical inconsistencies across multiple components, which can trigger initialization shocks that degrade coupled model forecasts (Mulholland et al., 2015; Zhang, 2011). To address these limitations, recent studies have increasingly used coupled data assimilation (CDA) (Wu et al., 2018; Zhang et al., 2012).

CDA has emerged as an advanced framework for performing ocean data assimilation directly within the coupled model. CDA methods are generally categorized into weakly coupled data assimilation (WCDA) and strongly coupled data assimilation (SCDA). In WCDA, ocean observations or reanalysis products are assimilated separately into the ocean component during the analysis step, but the coupled model is utilized in the forecast step to transfer ocean observational information to other components through multi-component interactions (Browne et al., 2019). The fundamental distinction between WCDA and uncoupled data assimilation lies in whether a coupled model is employed in the forecast step to generate the background forecast. In WCDA, the background forecast is produced through coupled model integration, while

69 observations [or reanalysis products](#) are assimilated independently within each individual
70 component. In contrast, uncoupled data assimilation utilizes standalone component models for
71 both the forecast and analysis stages (Shi et al., 2024a; Zhang et al., 2020). Unlike uncoupled
72 approaches, CDA produces more dynamically balanced and self-consistent initial conditions,
73 thereby enhancing forecast skill (Feng et al., 2018; Shi et al., 2022). However, WCDA neglects
74 cross-component background error covariances during the analysis step, which limits its ability
75 to explicitly correct coupled state errors (Tang et al., 2021). In contrast, SCDA leverages these
76 covariances to facilitate cross-domain information propagation, allowing observational data in
77 one component to instantaneously update the state variables of other components (Penny et al.,
78 2019; Lin and Pu, 2020). By treating the coupled system as a unified entity, SCDA typically
79 offers superior assimilation performance over WCDA (Han et al., 2013; Sluka et al., 2016; Lin
80 and Pu, 2019). However, the practical implementation of SCDA faces considerable challenges,
81 primarily owing to the difficulty of estimating cross-component background error covariances.
82 Therefore, WCDA remains more widely adopted in current systems (Zhou et al., 2024).

83 The Energy Exascale Earth System Model (E3SM) is a state-of-the-art Earth system model
84 developed by the U.S. Department of Energy to advance the understanding of Earth system
85 variability and change (Leung et al., 2020). Recently, Shi et al. (2025) developed a new weakly
86 coupled ocean data assimilation (WCODA) system utilizing the four-dimensional ensemble
87 variational (4DEnVar) method. [This 4DEnVar-based WCODA system is intended to provide](#)
88 [realistic initial conditions for developing seasonal-to-decadal \(S2D\) hindcast experiments](#)
89 [using the E3SM model. The primary objective of this study is to evaluate the strengths and](#)
90 [limitations of this newly developed WCODA system and to document its capability in](#)
91 [capturing global and regional climate variability for the broader modeling community, with](#)
92 [particular emphasis on large-scale climate modes and their remote impacts that are essential](#)
93 [for generating realistic initial conditions for S2D hindcast experiments.](#) Specifically, this
94 evaluation focuses on four key aspects: (1) surface air temperature and precipitation anomalies
95 over ocean, land, and global domains; (2) major tropical variability, including the Indian Ocean
96 Dipole (IOD) and El Niño-Southern Oscillation (ENSO); (3) decadal-to-multidecadal
97 variability, such as the Pacific Decadal Oscillation (PDO), Interdecadal Pacific Oscillation

Deleted: Despite this advancement, the impacts of this 4DEnVar-based WCODA system on simulating climate variability have not yet been systematically assessed. The objective of this study is to evaluate the influence of the E3SM coupled ocean data assimilation on the simulation of climate variability across global and regional scales.

104 (IPO), and Atlantic Multidecadal Oscillation (AMO); and (4) regional climate variability over
105 the contiguous United States.

106 The paper is structured as follows. Section 2 describes the E3SMv2 model, datasets, and
107 experimental design. Section 3 presents the results from analyzing major modes of climate
108 variability. Finally, the conclusions and discussion are provided in Section 4.

109

110 **2. Model, data and experimental design**

111 **2.1 Model description**

112 E3SMv2 is a fully coupled Earth system model that includes the atmospheric, ocean, sea
113 ice, land and river transport components. The atmospheric component (EAMv2) is based on a
114 spectral-element dynamical core with 72 vertical levels and is configured on a cubed-sphere
115 grid with approximately 110 km horizontal resolution (Golaz et al., 2022). The ocean
116 component (MPAS-O) employs an unstructured Voronoi mesh with horizontal spacing of ~60
117 km in the midlatitudes and ~30 km near the equator and poles, and includes 60 vertical layers
118 using a z-star coordinate (Reckinger et al., 2015). The sea ice component (MPAS-SI) shares
119 the same horizontal mesh with MPAS-O and provides detailed representations of sea ice
120 thermodynamics and dynamics (Turner et al., 2022). The land component (ELMv2) simulates
121 land surface energy and water fluxes, biophysical processes, and soil hydrological processes
122 (Golaz et al., 2019). The river transport component (MOSARTv2) represents streamflow
123 routing across river basins (Li et al., 2013). These five components are dynamically coupled
124 through the CPL7 coupler, which facilitates the exchange of momentum, heat, and mass fluxes
125 among model components (Craig et al., 2012). The detailed description and assessment of
126 E3SMv2 can be found in Golaz et al. (2022).

127

128 **2.2 Data assimilation scheme**

129 The 4DEnVar method employed in the WCODA system is based on the dimension-reduced
130 projection four-dimensional variational (DRP-4DVar) algorithm (Wang et al., 2010). In DRP-
131 4DVar, the ensemble-based approach is used to replace the adjoint model traditionally required
132 in standard 4DVar, thereby significantly reducing computational cost. The DRP-4DVar method

133 determines the optimal state in the ensemble subspace by fitting observational data to model-
134 generated ensemble samples within a four-dimensional assimilation window (Liu et al., 2011;
135 Shi et al., 2021). Unlike nudging methods, which typically apply empirical relaxation terms to
136 the model's prognostic equations, the DRP-4DVar scheme obtains the optimal analysis by
137 minimizing the cost function of 4DVar. Specifically, observational information is assimilated
138 along the trajectory of the model solution within the assimilation window, ensuring that the
139 optimal analysis is dynamically consistent with the model physics (He et al., 2017). To mitigate
140 ensemble collapse, the DRP-4DVar algorithm employs an inflation technique that introduces a
141 zero initial condition perturbation sample to reconstruct the ensemble projection space as a
142 full-rank matrix (Wang et al., 2010). Due to its flexibility and efficiency, the DRP-4DVar
143 method has been widely applied in diverse numerical models to improve the initialization of
144 ocean, land, and atmospheric states (He et al., 2020; Shi et al., 2024b; Zhu et al., 2022).

145 In this study, monthly mean ocean temperature and salinity from the EN4.2.1 reanalysis
146 (Good et al., 2013) are assimilated into the ocean component of the fully coupled E3SMv2
147 model within a one-month assimilation window. The observational uncertainty of the EN4.2.1
148 reanalysis is represented through the observation error covariance matrix, which is statistically
149 derived from the variance of ocean temperature and salinity data. The spatiotemporal density
150 of observations within the EN4.2.1 product modulates the assimilation impact. Regions with
151 dense observational coverage are expected to exhibit more pronounced improvements, whereas
152 in data-sparse areas such as the Indian Ocean, the assimilation constraint is relatively weaker.
153 Each monthly assimilation cycle consists of the analysis and forecast stages. First, the fully
154 coupled E3SMv2 model is run forward for one month using the background initial conditions
155 to generate model-derived monthly means of ocean temperature and salinity. In the analysis
156 stage, the differences between these model outputs and the EN4.2.1 reanalysis are calculated
157 to form the observational innovation. The DRP-4DVar scheme then applies this innovation to
158 compute the optimal initial conditions for the ocean component at the beginning of the
159 assimilation cycle. During the following forecast stage, the fully coupled model is rewound to
160 the start of the month and re-integrated for the same month using the optimized ocean state and
161 the background fields from other components. Through this coupled forecast, ocean

162 observational information is dynamically propagated to influence the state variables of other
163 components (e.g., atmosphere and land) through multi-component interactions. Thus, while
164 assimilation is confined to the ocean, employing the fully coupled E3SMv2 model in the
165 forecast stage enables the transfer of assimilated ocean information to other Earth system
166 components, classifying this system as a WCDA framework (Zhang et al., 2020). Further
167 technical details of the assimilation system are described in Shi et al. (2025).

168

169 2.3 Numerical experiments

170 Two numerical experiments are conducted to assess the impact of the WCDA system on
171 climate variability: a control experiment (CTRL) and an assimilation experiment (ASSIM).
172 Both experiments are initialized from the same initial condition on January 1, 1950, which is
173 derived from a fully spun-up long-term integration of E3SMv2 (Golaz et al., 2022). Following
174 initialization, the CTRL experiment is integrated freely without re-initialization or further
175 modifications to the model state from 1950 to 2021 under observed historical external forcings.
176 This experiment serves as the reference simulation for evaluating the influence of the
177 assimilation system. Both CTRL and ASSIM experiments are forced with the same historical
178 external forcings prescribed by the CMIP6 protocol (Eyring et al., 2016). The E3SMv2 model
179 had already reached a stable quasi-equilibrium state through a long spin-up and historical
180 integration (Golaz et al., 2022).

181 The ASSIM experiment assimilates monthly mean ocean temperature and salinity from the
182 EN4.2.1 reanalysis into all 60 layers of the ocean component in the fully coupled E3SMv2
183 model from 1950 to 2021 using a one-month assimilation window. At the beginning of each
184 month, monthly mean ocean temperature and salinity from the EN4.2.1 reanalysis are
185 assimilated to update the ocean initial conditions, after which the fully coupled model is
186 integrated freely until the end of the month. During this forecast period, the assimilated ocean
187 information is dynamically propagated to other components through multi-component
188 interactions. Unlike the sea surface temperature (SST) relaxation approach used in pacemaker
189 experiments to isolate the ocean influence on climate variability (e.g., Kosaka and Xie, 2016),
190 our assimilation system is implemented to generate initial conditions for hindcast experiments

Deleted: In the CTRL experiment, the fully coupled E3SMv2 model is integrated freely from 1950 to 2021 under observed historical external forcings.

Deleted: reference

Deleted: In contrast,

Deleted: t

Deleted: Considering sparse observational coverage and large uncertainties, the Arctic Ocean is excluded from the assimilation process.

by incorporating full-depth ocean temperature and salinity reanalysis to constrain the complete three-dimensional ocean state. This approach preserves vertical dynamical consistency and retains the subsurface ocean memory necessary for capturing decadal-to-multidecadal variability. For example, Morioka et al. (2018) demonstrated that skillful prediction of decadal variability in the South Atlantic requires the assimilation of subsurface ocean temperature and salinity, as SST-only assimilation fails to accurately represent zonal heat transport. However, the Arctic Ocean is excluded from the assimilation domain due to sparse observational coverage and the current absence of a coupled ice-ocean assimilation scheme. Much of the Arctic Ocean is characterized by pervasive year-round sea ice cover. Without simultaneously updating sea ice variables, assimilating ocean temperature and salinity alone could introduce physical inconsistencies at the ice-ocean interface. Addressing this technical limitation remains a high priority for future system development.

2.4 Evaluation datasets

Three types of monthly reanalysis datasets are used for evaluation: 1) ERA5 reanalysis data, including surface air temperature, 850 hPa winds, and 500 hPa geopotential height. ERA5 incorporates a wide range of satellite and in situ observations to produce high-quality global atmospheric fields (Hersbach et al., 2020). 2) Global Precipitation Climatology Project (GPCP) monthly precipitation data. This dataset provides reliable global precipitation estimates based on satellite microwave, infrared measurements, and surface rain gauge data (Adler et al., 2003). 3) Hadley Centre Sea Ice and Sea Surface Temperature (HadISST) data. HadISST provides monthly SST analyses derived from a combination of in situ measurements and satellite-based retrievals (Rayner et al., 2003). Since the HadISST dataset and the assimilated EN4.2.1 reanalysis partially share a common pool of underlying in situ ocean observations, HadISST does not serve as a fully independent validation dataset. Nevertheless, the improved agreement with HadISST still demonstrates the effectiveness of the assimilation system in constraining the ocean states.

3. Results

3.1 Surface air temperature and precipitation

The temporal evolution of both non-detrended and linearly detrended annual mean surface air temperature anomalies over the ocean, land, and global domains is shown in Fig. 1. The non-detrended time series (Figs. 1a-c) highlight the long-term warming trend, while the detrended series (Figs. 1d-f) isolate interannual variability by removing the linear trend from each domain-averaged time series. In both cases, ASSIM better captures the observed interannual surface air temperature variability compared to CTRL across all domains. For the non-detrended global annual mean surface air temperature, the correlation of ASSIM with observations is 0.92, slightly higher than the value of 0.85 reported by Douville et al. (2015) in their tropical Pacific pacemaker experiment. After linear detrending, the correlation coefficient between ASSIM and observations reaches 0.47 over the ocean, much higher than that of CTRL (0.09). Over land and global domains, the correlation coefficients for ASSIM are 0.44 and 0.49, respectively, again exceeding those of CTRL (land/global: 0.26/0.18). All three detrended correlations from ASSIM pass the 95% confidence level, indicating statistically significant improvements in simulating interannual surface temperature variability. The improvement is most pronounced over the ocean, where the assimilated ocean state exerts a direct influence on near-surface atmospheric conditions. Notably, the improved correlations over land suggest that the benefits of ocean data assimilation extend beyond oceanic regions, reflecting its broader influence on global surface temperature variability. However, compared with pacemaker experiments in which SST is directly nudged within the tropical Pacific, the relatively weaker atmospheric response to certain ENSO events in ASSIM may be related to the competing influence of global ocean constraints beyond the tropical Pacific, as well as the smoothing effect introduced by the monthly assimilation window.

Figure 2 displays the spatial distribution of climatological mean differences in surface air temperature. Compared to observations, CTRL exhibits widespread cold biases over the tropical and subtropical oceans, as well as notable warm biases at high latitudes. In contrast, these biases are significantly reduced in ASSIM, particularly over the North Pacific, North Atlantic, and the midlatitude Southern Ocean. Specifically, the cold bias in the North Pacific decreases from -1.81 °C in CTRL to -0.76 °C in ASSIM, and from -3.23 °C to -1.09 °C in the

Deleted: These results highlight the effectiveness of ocean data assimilation in enhancing the simulation of interannual temperature variability.

Deleted: enhancing simulations of interannual temperature variability across all domains.

263 North Atlantic. The warm bias over the midlatitude Southern Ocean is notably reduced from
264 1.07 °C in CTRL to 0.13 °C in ASSIM. Improvements are also evident over land regions, such
265 as eastern China, northern Eurasia, northern Africa, and parts of North America. The spatial
266 pattern of reduced surface air temperature bias closely aligns with the reduction in SST bias
267 reported by Shi et al. (2025), highlighting the strong coupling between the ocean surface and
268 the atmosphere. However, the current assimilation experiment does not explicitly isolate the
269 respective contributions of SST and subsurface ocean states. Future SST-only assimilation
270 experiments would be needed to disentangle their individual effects on surface air temperature.
271 The climatological warm biases over the Arctic Ocean appear slightly enhanced in ASSIM
272 compared with CTRL. Since the Arctic Ocean state is not directly constrained by assimilation,
273 it may respond indirectly to remote adjustments introduced elsewhere, potentially leading to a
274 redistribution of temperature biases. Despite this localized degradation, the assimilation
275 maintains the overall stability of the coupled system. As shown in Fig. A1, key global mean
276 diagnostics, including top-of-atmosphere net energy flux, surface air temperature, ocean
277 temperature and salinity, remain stable throughout the ASSIM integration.

278 The interannual variability of linearly detrended precipitation anomalies over the ocean,
279 land, and global domains is illustrated in Fig. 3. Compared to CTRL, ASSIM shows markedly
280 improved agreement with observed interannual precipitation variability across all three
281 domains. Over the ocean, the correlation with observations increases from 0.11 in CTRL to
282 0.52 in ASSIM, while over land it improves from 0.27 to 0.60. At the global scale, the
283 correlation rises from 0.04 in CTRL to 0.55 in ASSIM. All three correlations from ASSIM are
284 statistically significant at the 95% confidence level. These results suggest that assimilating
285 observed ocean states enhances the simulation of interannual precipitation variability over both
286 ocean and land, with the latter improvement possibly related to atmospheric circulation and
287 moisture transport.

288 Figure 4 shows the spatial distribution of climatological mean precipitation and associated
289 model biases. In the observation (Fig. 4a), major precipitation maxima are located along the
290 Intertropical Convergence Zone (ITCZ) in the equatorial Pacific, the South Pacific
291 Convergence Zone (SPCZ) extending northwest–southeast from the western Pacific, and the

Deleted: underscoring the critical role of realistic ocean states in shaping near-surface temperature patterns.

Deleted: Notably, the relatively high correlations for both surface air temperature (0.26) and precipitation (0.27) over land in CTRL suggest some influence of external forcing on land surface climate variability.

298 tropical Indian Ocean. A dry zone is also evident along the west coast of South America.
 299 Compared to observations, CTRL overestimates precipitation over the central Pacific and the
 300 southern equatorial Atlantic, while significantly underestimating rainfall over the tropical
 301 eastern Indian Ocean and the northern equatorial Atlantic (Figs. 4b & d). In contrast, these
 302 biases are substantially reduced in ASSIM (Figs. 4c & e). Precipitation in the tropical eastern
 303 Indian Ocean is enhanced, resulting in better agreement with observations. Moreover, in the
 304 tropical Atlantic, the dry bias over the northern equatorial Atlantic is reduced from -1.27
 305 mm/day in CTRL to -0.34 mm/day in ASSIM, while the wet bias over the southern equatorial
 306 Atlantic decreases from 1.81 mm/day to 0.62 mm/day. These reductions in systematic biases
 307 bring the climatological precipitation in ASSIM into better agreement with observations, with
 308 residual differences over some regions, particularly the tropical Pacific, no longer reaching
 309 statistical significance. The improvement in tropical precipitation is closely linked to changes
 310 in the large-scale meridional circulation. As illustrated in Fig. A2, CTRL exhibits an
 311 anomalously strong ascending branch of the Hadley circulation in the northern tropics, which
 312 sustains a pronounced wet bias over the tropical Pacific. In contrast, this anomalous ascent is
 313 effectively suppressed in ASSIM, contributing to a more realistic precipitation distribution
 314 across the tropical and subtropical regions.

316 3.2 ENSO

317 ENSO is the dominant mode of interannual climate variability in the tropics and exerts
 318 profound impacts on global atmospheric circulation (Chiang and Sobel, 2002). ENSO
 319 variability is commonly monitored by the Niño 3.4 index, defined as the SST anomalies
 320 averaged over the equatorial central Pacific (5°S–5°N, 120°W–170°W) (Trenberth and
 321 Stepaniak, 2001). To evaluate the representation of ENSO variability, we analyze the time
 322 evolutions of the monthly and winter Niño 3.4 index (Fig. 5). As expected, CTRL exhibits very
 323 weak correlations with observations, with correlation coefficients of only 0.02 for the monthly
 324 and 0.07 for the winter Niño 3.4 index, indicating poor phase agreement with the observed
 325 Niño 3.4 index. Compared with CTRL, ASSIM exhibits markedly improved phase agreement
 326 with the observed ENSO variability. The temporal evolution of the Niño 3.4 index in ASSIM

Deleted: the model's limited ability to reproduce the evolution of tropical SST variability in the absence of observational constraints

Deleted: substantial improvements in simulating ENSO variability.

332 closely matches the observation, with correlations increasing to 0.87 for the monthly and 0.95
333 for the winter index. These results demonstrate that ocean data assimilation provides an
334 effective constraint on tropical Pacific SST variability.

335 Figure 6 further evaluates ENSO characteristics through the power spectra and seasonal
336 cycle of the monthly Niño 3.4 index. The observed spectrum exhibits a pronounced peak in the
337 3–5 year band, consistent with typical ENSO periodicities (Fig. 6a). This spectral feature is
338 reasonably well reproduced in ASSIM (Fig. 6b). By comparison, CTRL underestimates the
339 variance in this band and produces an unrealistically amplified peak at 2–3 years (Fig. 6c).
340 Beyond its frequency characteristics, ENSO also exhibits a strong seasonal cycle. The observed
341 Niño 3.4 index shows minimum variance in April and peak amplitude in December, reflecting
342 the well-known phase-locking behavior of ENSO (Fig. 6d). ASSIM captures this seasonal
343 cycle well, although the minimum occurs one month later (Fig. 6e). In contrast, CTRL exhibits
344 a flatter seasonal cycle with markedly weaker variance during the winter peak, and its seasonal
345 minimum is delayed by two months (Fig. 6f).

346 To further assess the representation of ENSO-related atmospheric variability, we examine
347 the Southern Oscillation Index (SOI), which reflects the east–west seesaw pattern in sea level
348 pressure (SLP) between the western and eastern tropical Pacific. The SOI is commonly defined
349 as the normalized difference in SLP anomalies between Tahiti and Darwin, and serves as the
350 atmospheric counterpart to the Niño 3.4 index (Hanley et al., 2003). Figure 7 presents the
351 temporal evolution of the monthly and winter SOI from the observation, ASSIM, and CTRL.
352 The observed SOI exhibits pronounced interannual fluctuations associated with ENSO phases.
353 Consistent with its poor simulation of the Niño 3.4 index, CTRL fails to capture the observed
354 SOI variability, with correlations of -0.02 for the monthly and -0.04 for the winter index. In
355 contrast, ASSIM reproduces the SOI variability more realistically, with correlations increasing
356 to 0.40 for the monthly and 0.62 for the winter index. These results indicate that assimilating
357 ocean reanalysis, not only constrains SST but also enhances the phase agreement of ENSO-
358 related atmospheric variability with observations, in coupled models.

359
360 **3.3 IOD**

Deleted: underscore the critical role of ocean data assimilation in constraining tropical Pacific SST variability and enhancing ENSO representation in coupled models.

Deleted: observed ocean states

Deleted: improves

Deleted: representation

Deleted: circulation anomalies

368 The Indian Ocean Dipole (IOD) is a prominent mode of interannual variability in the
 369 tropical Indian Ocean that strongly influences surrounding climate systems (Saji et al., 2006).
 370 It typically begins to develop in boreal summer and reaches its peak during autumn. A widely
 371 used metric for quantifying IOD variability is the Dipole Mode Index (DMI), defined as the
 372 difference in SST anomalies between the western (50°E–70°E, 10°S–10°N) and eastern (90°E–
 373 110°E, 0°–10°S) equatorial Indian Ocean (Saji et al., 1999). The temporal evolution of the
 374 autumn DMI is further shown in Fig. 8. The observed DMI exhibits pronounced interannual
 375 variations associated with alternating positive and negative IOD phases. The CTRL simulation
 376 substantially overestimates the IOD amplitude and shows a weak correlation of -0.11 with
 377 observations. Such overestimation is the common bias in coupled climate models, as noted in
 378 previous studies (Ju et al., 2025). In contrast, ASSIM shows notable improvements in both the
 379 amplitude and temporal variability of the DMI, with the correlation increasing to 0.56 and
 380 exceeding the 95% confidence level. These results highlight the ~~effectiveness~~ of ocean data
 381 assimilation ~~in~~ improving the ~~temporal agreement of IOD variability with observations.~~
 382 ~~Nevertheless, the lower DMI correlation in ASSIM relative to Niño3.4 is partly attributable to~~
 383 ~~known limitations of the assimilated EN4.2.1 dataset in the Indian Ocean. As noted by Good~~
 384 ~~et al. (2013), the EN4.2.1 reanalysis exhibits regional deficiencies in this basin due to the~~
 385 ~~limited availability of near-surface observations that meet quality-control standards, thereby~~
 386 ~~limiting the assimilation effectiveness in the Indian Ocean.~~

Deleted: benefits

Deleted: for

Deleted: simulation of IOD variability.

388 3.4 PDO, IPO, AMO

389 The Pacific Decadal Oscillation (PDO) is the leading mode of decadal SST variability in
 390 the North Pacific, characterized by basin-scale SST anomalies that typically persist for one to
 391 two decades. The PDO index is derived as the leading empirical orthogonal function (EOF)
 392 mode of detrended SST anomalies over the North Pacific (20°–70°N, 110°E–100°W) (Mantua
 393 et al., 1997). ~~The EOF analysis is performed separately for the observations and each model~~
 394 ~~experiment. Prior to the EOF calculation, all observational datasets and model outputs are~~
 395 ~~interpolated onto a common horizontal grid and linear trends are removed at each grid point.~~
 396 Figure 9 illustrates both the temporal evolution and spatial pattern of the annual PDO index

400 from 1965 to 2016. The observed PDO index displays a canonical sequence of phase transitions:
401 a cool phase before the late 1970s, a persistent warm phase extending into the late 1990s, and
402 a return to negative values thereafter (Fig. 9a). These decadal transitions are reasonably
403 captured by ASSIM, yielding a temporal correlation of 0.67 with observations (Fig. 9b). In
404 contrast, CTRL fails to reproduce the observed phase transitions, with a much lower correlation
405 of 0.19 (Fig. 9c). Besides the temporal evolution, the spatial structure of the PDO index is also
406 analyzed. The observed pattern exhibits a horseshoe-shaped structure with cold anomalies in
407 the central North Pacific and warm anomalies along the western coast of North America (Fig.
408 9d). This spatial pattern is well reproduced in ASSIM, with a spatial correlation of 0.90 (Fig.
409 9e). However, the PDO pattern from CTRL exhibits a distorted structure, and its spatial
410 correlation decreases to 0.68 (Fig. 9f). These improvements in both the temporal evolution and
411 spatial structure of the PDO underscore the potential of the assimilation system to enable more
412 skillful decadal climate predictions by providing improved initial states.

413 Beyond the North Pacific, another major mode of decadal variability affecting the entire
414 Pacific basin is the Interdecadal Pacific Oscillation (IPO). Similar to the PDO, the IPO index
415 is obtained from the leading EOF mode of detrended SST anomalies, but computed over the
416 whole Pacific (50°S–50°N, 100°E–70°W) (Doblas-Reyes et al., 2013). The annual evolution
417 of the IPO index is shown in Fig. 10, indicating the basin-wide coherence of Pacific decadal
418 variability. The observed IPO index undergoes distinct decadal shifts, including a negative
419 phase prior to the late 1970s, a prolonged positive phase extending through the 1980s and 1990s,
420 and a return to negative values in the early 2000s. These phase transitions are closely
421 reproduced by ASSIM, which achieves a high temporal correlation of 0.80 with observations.
422 On the other hand, CTRL fails to track the decadal evolution of the observed IPO and shows a
423 much weaker correlation of 0.20. These results indicate that ocean data assimilation serves as
424 an effective pathway for ~~constraining Pacific-basin SST variability and enhancing the phase~~
425 alignment of the observed Pacific SST variability on decadal timescales.

426 In addition to the Pacific, the North Atlantic also exhibits notable low-frequency SST
427 variability characterized by the Atlantic Multidecadal Oscillation (AMO). The AMO index is
428 computed as the linearly detrended SST anomalies averaged over the North Atlantic (0–60°N,

Deleted: highlight the importance of ocean data assimilation in ...

Deleted: on decadal timescales

Deleted: representation of long-term

Deleted: climate

0–80°W) (Trenberth and Shea, 2006). Figure 11 presents the temporal evolution of the annual AMO index from 1965 to 2016. The observed AMO index shows a pronounced negative phase in the 1970s and 1980s, followed by a persistent positive phase beginning in the mid-1990s. These phase transitions are more accurately reproduced in ASSIM than in CTRL, particularly the timing and amplitude of the warm shift after the mid-1990s. The temporal correlation with observations is 0.71 in ASSIM, much higher than 0.17 in CTRL. This improved phase agreement of AMO variability in ASSIM reinforces its potential for better predicting related climate impacts over the North Atlantic and adjacent regions.

Deleted: e

Deleted: enhanced representation of AMO phases

3.5 U.S. T2m and Precipitation

Building on the improved representation of tropical SST variability through ocean data assimilation, we extend our analysis to assess its impact on regional climate over land. Given the well-established influence of ENSO on U.S. winter climate variability (Ropelewski and Halpert, 1986; Higgins et al., 2000), we evaluate model performance in simulating interannual temperature and precipitation variability over the contiguous U.S. during boreal winter (Figs. 12 & 13). For winter surface air temperature (Fig. 12), ASSIM exhibits widespread increases in temporal correlation with observations across large portions of the northern and western U.S., particularly over the Intermountain West and Great Lakes regions. In these areas, correlation improvements commonly exceed 0.3 and locally surpass 0.4. Consistent with the correlation patterns, RMSE reductions in ASSIM are also evident across the northern and central U.S., with maximum decreases up to 0.6 °C over the central Great Plains.

In contrast to temperature, improvements in winter precipitation simulations are more localized but remain pronounced across the southern U.S. (Fig. 13). Compared with CTRL, ASSIM shows higher correlations with observed precipitation over the Southwest, southern Great Plains, and Southeast. In particular, significant enhancements are evident in parts of California, Texas, and Alabama, where correlation increases exceed 0.4 and locally approach 0.5. These correlation improvements are accompanied by consistent reductions in RMSE across similar regions. Notably, the largest RMSE reductions, exceeding 0.45 mm/day, are found over coastal California. The spatial coherence of these improvements suggests that

465 incorporating realistic ocean states can better capture U.S. winter precipitation variability.

Deleted: highlights the critical role of realistic ocean states in better capturing land

466 To further investigate the mechanisms underlying the improved simulation of U.S. winter
467 climate, we examine the regression of surface air temperature and precipitation onto the Niño
468 3.4 index during boreal winter (Fig. 14). Boreal winter is the focus of this analysis because
469 ENSO teleconnections to North America are strongest and most coherent during winter, while
470 ENSO-related impacts in other seasons are generally weaker and less spatially coherent over
471 the U.S. (Ropelewski and Halpert, 1986; Higgins et al., 2000). The Niño 3.4 index is calculated
472 separately from its corresponding dataset, and all regressions are performed using simultaneous
473 winter averages. The observed patterns reveal that ENSO exerts a pronounced influence on
474 U.S. winter climate through large-scale atmospheric responses. During El Niño events, positive
475 geopotential height anomalies dominate the northern part of the continent, which may suppress
476 the southward intrusion of cold air masses. This circulation pattern could lead to anomalous
477 warming across the northern U.S., producing significant positive correlations with ENSO in
478 these regions (Fig. 14a). Meanwhile, strengthened subtropical westerlies enhance moisture
479 transport into the southern U.S., favoring increased precipitation and yielding strong positive
480 correlations with ENSO across the southern tier of the country (Fig. 14d).

481 These ENSO-related spatial structures are well reproduced in ASSIM (Figs. 14b & e). In
482 particular, the regression patterns of temperature and precipitation in ASSIM closely resemble
483 those in the observations, especially across the northern and southern U.S., respectively. The
484 spatial pattern correlation between ASSIM and observations is 0.86 for surface air temperature,
485 much higher than that of CTRL (0.35). For precipitation, the spatial correlation increases from
486 0.67 in CTRL to 0.72 in ASSIM. Furthermore, the regions with robust ENSO-related signals
487 in ASSIM largely overlap with areas showing improved correlation and reduced RMSE in Figs.
488 12 and 13. The stippled regions in these figures denote statistically significant differences at
489 the 95% confidence level. Specifically, for winter surface air temperature, statistically
490 significant increases in correlation and concurrent reductions in RMSE are primarily found in
491 parts of Arizona and Iowa (Fig. 12), while for winter precipitation, such improvements are
492 mainly located over parts of California and Alabama (Fig. 13). To account for the potential
493 delayed impact of ENSO, a one-month lagged regression analysis is also presented in Fig. A3.

496 ~~The resulting lagged patterns are highly consistent with the simultaneous winter regression~~
497 ~~results (Fig. 14), reinforcing the linkage between ENSO variability and U.S. winter climate~~
498 ~~anomalies. These results demonstrate that the improved phase alignment of ENSO variability~~
499 ~~enhances the simulation of U.S. winter surface air temperature and precipitation.~~

Deleted: , suggesting that enhanced representation of ENSO variability may underpin its superior performance in simulating U.S. winter climate. In contrast, CTRL exhibits notable deviations from the observed ENSO-related patterns, especially in the spatial distribution of temperature responses (Figs. 14c & f).

Deleted: underscore the importance of accurate ENSO representation for improving simulations

Deleted: climate variability

501 4 Conclusions

502 Accurate representation of climate variability across multiple timescales is essential for
503 enhancing the reliability of Earth system predictions. One key factor influencing climate
504 variability is the ocean state due to its long memory and strong coupling with the atmosphere.
505 While ocean data assimilation has proven effective in improving ocean state estimates, many
506 existing systems remain uncoupled and thus have limited ability to support coupled forecasts
507 due to imbalances across multiple components (Mulholland et al., 2015; Zhang et al., 2011).
508 To address this limitation, a new 4DEnVar-based WCODA system ~~has been implemented~~
509 ~~within the E3SM model to provide realistic initial conditions for S2D hindcast experiments.~~
510 ~~Despite this progress, the specific impact of this WCODA system on the simulation of multi-~~
511 ~~scale climate variability requires further evaluation. This study aims to thoroughly document~~
512 ~~the capabilities and limitations of this WCODA system in simulating global and regional~~
513 ~~climate variability, and to demonstrate its potential for improving initialization quality and~~
514 ~~predictive skill in coupled Earth system models.~~

Deleted: Shi et al. (2025) recently developed

Deleted: that enables direct assimilation of ocean observations within a coupled model framework.

Deleted: the extent to which this WCODA system improves simulations of global and regional climate variability has not been fully assessed

Deleted: systematically evaluate its impacts on simulating major modes of climate variability and regional climate over the U.S.,

Deleted: representation

515 Our results show that the WCODA system significantly improves the ~~phase agreement of~~
516 ~~both interannual and decadal climate variability with observations. Beyond phase alignment,~~
517 ~~minor but noteworthy improvements are also found in representing the modes of variability, as~~
518 ~~indicated by the improved Niño3.4 spectrum, PDO EOF pattern, and ENSO teleconnection~~
519 ~~response over the contiguous U.S.~~ Significant enhancements are achieved in capturing
520 interannual variability and reducing model biases of surface air temperature and precipitation
521 over land and ocean domains. Notably, the temporal evolution of major climate modes,
522 including ENSO, IOD, PDO, IPO, and AMO, is markedly improved. Besides better capturing
523 the temporal evolution of various modes, assimilating ocean temperature and salinity also
524 improves the power spectra and seasonal cycle of ENSO and the spatial pattern of PDO.

Furthermore, regional evaluations over the contiguous U.S. reveal increased skill in reproducing the observed interannual variability of winter surface air temperature in the northern U.S. and precipitation in the southern regions, respectively. The improved simulation of U.S. winter temperature and precipitation is closely associated with better-simulated ENSO variability, as supported by spatially coherent regression patterns between Niño 3.4 and U.S. winter climate. These findings demonstrate the capability of the WCODA system to deliver more skillful simulations of Earth system variability through improved ocean initial states and cross-component information exchange. It is worth noting that these improvements primarily reflect enhanced phase alignment of internal variability with observations, and the use of monthly assimilation increments may introduce localized nonconservative effects.

Despite these promising results, several limitations remain and offer opportunities for future development. The current assimilation system does not incorporate atmospheric or sea ice observations, which may limit its ability to fully constrain the coupled model states. While our results demonstrate that assimilating ocean reanalysis provides a sufficient constraint to improve the phase alignment of key climate modes, incorporating additional observational constraints from the atmosphere and sea ice may further enhance the representation of coupled climate variability and associated regional impacts. It should be noted that the improved temporal correlations primarily reflect enhanced phase alignment of the model's internal variability with observations, rather than intrinsic changes in modes of variability, and the assimilation increments may locally affect energy and mass budgets. Furthermore, as a weakly coupled system, the current WCODA framework does not explicitly account for cross-component background error covariances during the analysis step. Transitioning toward a SCDA framework would further improve the physical consistency and accuracy of the generated initial conditions.

Looking ahead, the WCODA system provides a promising foundation for improving the initialization of large-ensemble S2D hindcast experiments. Given that the assimilation of subsurface ocean temperature and salinity is essential for skillful decadal prediction (Morioka et al., 2018; Chikamoto et al., 2019), future efforts could explore systematic comparisons with SST-based pacemaker experiments to elucidate the distinct roles of surface and subsurface

ocean constraints. Such comparisons would provide valuable insights into the mechanisms governing decadal predictability and inform the design of more effective initialization strategies.

Ultimately, these advances will contribute to the development of more reliable seamless climate prediction systems with broad implications for climate services and society.

Code and data availability. The E3SMv2 source code is freely available under an open-source license and can be accessed via Zenodo at <https://zenodo.org/records/16652680>. The ERA5 reanalysis datasets are accessible from the Copernicus Climate Change Service through the Climate Data Store portal: <https://cds.climate.copernicus.eu/datasets/reanalysis-era5-single-levels-monthly-means?tab=download> (Hersbach et al., 2020). Monthly GPCP precipitation data can be obtained from <https://psl.noaa.gov/data/gridded/data.gpcp.html> (Adler et al., 2003). HadISST sea surface temperature data are available at <https://www.metoffice.gov.uk/hadobs/hadisst> (Rayner et al., 2003) and EN4.2.1 ocean temperature and salinity data can be accessed from <https://www.metoffice.gov.uk/hadobs/en4> (Good et al., 2013). Model outputs are available on Zenodo at <https://zenodo.org/records/16740831>.

Author contributions. PS and LRL designed the experiments. PS developed the ocean data assimilation system and performed the model simulations. PS and LRL analyzed the results. PS and LRL drafted the initial manuscript. ZP, SH, and KB contributed to manuscript revisions.

Competing interests. The authors declare no competing interests.

Acknowledgements. This research was supported by the Office of Science, U.S. Department of Energy Biological and Environmental Research (BER) as part of the Water Cycle: Modeling of Circulation, Convection, and Earth System Mechanisms (WACCEM) Scientific Focus Area funded by the Regional and Global Model Analysis program area. Pacific Northwest National Laboratory is operated by Battelle Memorial Institute for the U.S. Department of Energy under contract DE-AC05-76RL01830. This research used computing resources of the National

602 Energy Research Scientific Computing Center, which is supported by the Office of Science of
603 the U.S. Department of Energy under Contract No. DE-AC02-05CH11231, and the BER Earth
604 and Environmental System Modeling program's Compy computing cluster located at Pacific
605 Northwest National Laboratory.

REFERENCES

- Adler, R. F., Huffman, G. J., Chang, A., Ferraro, R., Xie, P. P., Janowiak, J., Rudolf, B., Schneider, U., Curtis, S., Bolvin, D., Gruber, A., Susskind, J., Arkin, P., and Nelkin, E.: The version-2 Global Precipitation Climatology Project (GPCP) monthly precipitation analysis (1979–present), *Journal of Hydrometeorology*, 4, 1147–1167, [https://doi.org/10.1175/1525-7541\(2003\)004<1147:TVGPCP>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<1147:TVGPCP>2.0.CO;2), 2003.
- Boer, G. J., Smith, D. M., Cassou, C., Doblas-Reyes, F., Danabasoglu, G., Kirtman, B., Kushnir, Y., Kimoto, M., Meehl, G. A., Msadek, R., Mueller, W. A., Taylor, K. E., Zwiers, F., Rixen, M., Ruprich-Robert, Y., and Eade, R.: The Decadal Climate Prediction Project (DCPP) contribution to CMIP6, *Geoscientific Model Development*, 9, 3751–3777, <https://doi.org/10.5194/gmd-9-3751-2016>, 2016.
- Browne, P. A., De Rosnay, P., Zuo, H., Bennett, A., and Dawson, A.: Weakly coupled ocean–atmosphere data assimilation in the ECMWF NWP system, *Remote Sensing*, 11, 234, <https://doi.org/10.3390/rs11030234>, 2019.
- Carton, J. A. and Giese, B. S.: A reanalysis of ocean climate using Simple Ocean Data Assimilation (SODA), *Monthly Weather Review*, 136, 2999–3017, <https://doi.org/10.1175/2007MWR1978.1>, 2008.
- Chiang, J. C. H. and Sobel, A. H.: Tropical tropospheric temperature variations caused by ENSO and their influence on the remote tropical climate, *Journal of Climate*, 15, 2616–2631, [https://doi.org/10.1175/1520-0442\(2002\)015<2616:TTTVCB>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<2616:TTTVCB>2.0.CO;2), 2002.
- [Chikamoto, Y., Timmermann, A., Widlansky, M. J., Zhang, S., and Balmaseda, M. A.: A drift-free decadal climate prediction system for the Community Earth System Model, *Journal of Climate*, 32, 5967–5995, <https://doi.org/10.1175/JCLI-D-18-0788.1>, 2019.](#)
- Craig, A. P., Vertenstein, M., and Jacob, R.: A new flexible coupler for Earth system modeling developed for CCSM4 and CESM1, *The International Journal of High Performance Computing Applications*, 26, 31–42, <https://doi.org/10.1177/1094342011428141>, 2012.
- Doblas-Reyes, F. J., Andreu-Burillo, I., Chikamoto, Y., García-Serrano, J., Guemas, V., Kimoto, M., Mochizuki, T., Rodrigues, L. R. L., and van Oldenborgh, G. J.: Initialized near-term regional climate change prediction, *Nature Communications*, 4, 1715,

<https://doi.org/10.1038/ncomms2704>, 2013.

[Douvile, H., Voldoire, A., and Geoffroy, O.: The recent global warming hiatus: What is the role of Pacific variability? *Geophysical Research Letters*, 42, 880–888, <https://doi.org/10.1002/2014GL062775>, 2015.](#)

Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.

Feng, X., Haines, K., and de Boissésou, E.: Coupling of surface air and sea surface temperatures in the CERA-20C reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 144, 195–207, <https://doi.org/10.1002/qj.3194>, 2018.

Golaz, J. C., Caldwell, P. M., Van Roekel, L. P., Petersen, M. R., Tang, Q., Wolfe, J. D., Abeshu, G., Anantharaj, V., Asay-Davis, X. S., Bader, D. C., Baldwin, S. A., Bisht, G., Bogenschutz, P. A., Branstetter, M., Brunke, M. A., Brus, S. R., Burrows, S. M., Cameron-Smith, P. J., Donahue, A. S., Deakin, M., Easter, R. C., Evans, K. J., Feng, Y., Flanner, M., Foucar, J. G., Fyke, J. G., Griffin, B. M., Hannay, C., Harrop, B. E., Hoffman, M. J., Hunke, E. C., Jacob, R. L., Jacobsen, D. W., Jeffery, N., Jones, P. W., Keen, N. D., Klein, S. A., Larson, V. E., Leung, L. R., Li, H. Y., Lin, W., Lipscomb, W. H., Ma, P. L., Mahajan, S., Maltrud, M. E., Mametjanov, A., McClean, J. L., McCoy, R. B., Neale, R. B., Price, S. F., Qian, Y., Rasch, P. J., Reeves Eyre, J. E. J., Riley, W. J., Ringler, T. D., Roberts, A. F., Roesler, E. L., Salinger, A. G., Shaheen, Z., Shi, X., Singh, B., Tang, J., Taylor, M. A., Thornton, P. E., Turner, A. K., Veneziani, M., Wan, H., Wang, H., Wang, S., Williams, D. N., Wolfram, P. J., Worley, P. H., Xie, S., Yang, Y., Yoon, J.-H., Zelinka, M. D., Zender, C. S., Zeng, X., Zhang, C., Zhang, K., Zhang, Y., Zheng, X., Zhou, T., and Zhu, Q.: The DOE E3SM Coupled Model Version 1: Overview and Evaluation at Standard Resolution, *Journal of Advances in Modeling Earth Systems*, 11, 2089–2129, <https://doi.org/10.1029/2018MS001603>, 2019.

Golaz, J. C., Van Roekel, L. P., Zheng, X., Roberts, A. F., Wolfe, J. D., Lin, W. Y., Bradley, A. M., Tang, Q., Maltrud, M. E., Forsyth, R. M., Zhang, C. Z., Zhou, T., Zhang, K., Zender,

664 C. S., Wu, M. X., Wang, H. L., Turner, A. K., Singh, B., Richter, J. H., Qin, Y., Petersen,
 665 M. R., Mametjanov, A., Ma, P., Larson, V. E., Krishna, J., Keen, N. D., Jeffery, N., Hunke,
 666 E. C., Hannah, W. M., Guba, O., Griffin, B. M., Feng, Y., Engwirda, D., Vittorio, A. V.,
 667 Cheng, D., Conlon, L. M., Chen, C., Brunke, M. A., Bisht, G., Benedict, J. J., Asay-Davis,
 668 X. S., Zhang, Y. Y., Zhang, M., Zeng, X. B., Xie, S. C., Wolfram, P. J., Vo, T., Veneziani,
 669 M., Tesfa, T. K., Sreepathi, S., Salinger, A. G., Jack Reeves Eyre, J. E., Prather, M. J.,
 670 Mahajan, S., Li, Q., Jones, P. W., Jacob, R. L., Huebler, G. W., Huang, X. L., Hillman, B.
 671 R., Harrop, B. E., Foucar, J. G., Fang, Y. L., Comeau, D. S., Caldwell, P. M., Bartoletti,
 672 T., Balaguru, K., Taylor, M. A., McCoy, R. B., Leung, L. R., and Bader, D. C.: The DOE
 673 E3SM Model version 2: Overview of the physical model and initial model evaluation,
 674 Journal of Advances in Modeling Earth Systems, 14, e2022MS003156, [https://doi.](https://doi.org/10.1029/2022MS003156)
 675 [org/10.1029/2022MS003156](https://doi.org/10.1029/2022MS003156), 2022.
 676 Good, S. A., Martin, M. J., and Rayner, N. A.: EN4: Quality controlled ocean temperature and
 677 salinity profiles and monthly objective analyses with uncertainty estimates, Journal of
 678 Geophysical Research: Oceans, 118(12), 6704–6716,
 679 <https://doi.org/10.1002/2013JC009067>, 2013.
 680 Han, G., Wu, X., Zhang, S., Liu, Z., and Li, W.: Error covariance estimation for coupled data
 681 assimilation using a Lorenz atmosphere and a simple pycnocline ocean model, Journal of
 682 Climate, 26, 10218–10231, <https://doi.org/10.1175/JCLI-D-13-00236.1>, 2013.
 683 Hanley, D. E., Bourassa, M. A., O'Brien, J. J., Smith, S. R., and Spade, E. R.: A quantitative
 684 evaluation of ENSO indices, Journal of Climate, 16, 1249–1258,
 685 [https://doi.org/10.1175/1520-0442\(2003\)16<1249:AQEOEI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)16<1249:AQEOEI>2.0.CO;2), 2003.
 686 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J.,
 687 Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X.,
 688 Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren,
 689 P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer,
 690 A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux,
 691 P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S.,
 692 and Thépaut, J.-N.: The ERA5 global reanalysis, Quarterly Journal of the Royal

Meteorological Society, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

[He, Y., Wang, B., Liu, M., Liu, L., Yu, Y., Liu, J., Li, R., Zhang, C., Xu, S., Huang, W., Liu, Q., Wang, Y., and Li, F.: Reduction of initial shock in decadal predictions using a new initialization strategy, *Geophysical Research Letters*, 44\(16\), 8538–8547, <https://doi.org/10.1002/2017GL074028>, 2017.](#)

He, Y., Wang, B., Huang, W., Xu, S., Wang, Y., Liu, L., Li, L., Liu, J., Yu, Y., Lin, Y., Huang, X., and Peng, Y.: A new DRP-4DVar-based coupled data assimilation system for decadal predictions using a fast online localization technique, *Climate Dynamics*, 54, 3541–3559, <https://doi.org/10.1007/s00382-020-05190-w>, 2020.

Higgins, R. W., Leetmaa, A., Xue, Y., and Barnston, A.: Dominant factors influencing the seasonal predictability of U.S. precipitation and surface air temperature, *Journal of Climate*, 13, 3994–4017, [https://doi.org/10.1175/1520-0442\(2000\)013<3994:DFITSP>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<3994:DFITSP>2.0.CO;2), 2000.

Ju, X., Chen, S., Chen, W., Wu, R., Chen, L., and Cheng, X.: Origins of the overestimated Indian Ocean Dipole amplitude in current coupled climate models, *Climate Dynamics*, 63, 180, <https://doi.org/10.1007/s00382-025-07673-0>, 2025.

[Kosaka, Y. and Xie, S. P.: The tropical Pacific as a key pacemaker of the variable rates of global warming, *Nature Geoscience*, 9\(9\), 669–673, <https://doi.org/10.1038/ngeo2770>, 2016.](#)

Leung, L. R., Bader, D. C., Taylor, M. A., and McCoy, R. B.: An introduction to the E3SM special collection: Goals, science drivers, development, and analysis, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001821, <https://doi.org/10.1029/2019MS001821>, 2020.

Li, H., Wigmosta, M. S., Wu, H., Huang, M., Ke, Y., Coleman, A. M., and Leung, L. R.: A physically based runoff routing model for land surface and Earth system models, *Journal of Hydrometeorology*, 14, 808–828, <https://doi.org/10.1175/JHM-D-12-015.1>, 2013.

Lin, L. F. and Pu, Z.: Examining the impact of SMAP soil moisture retrievals on short-range weather prediction under weakly and strongly coupled data assimilation with WRF-Noah, *Monthly Weather Review*, 147, 4345–4366, <https://doi.org/10.1175/MWR-D-19-0017.1>, 2019.

722 Lin, L. F. and Pu, Z.: Improving near-surface short-range weather forecasts using strongly
 723 coupled land–atmosphere data assimilation with GSI-EnKF, *Monthly Weather Review*,
 724 148, 2863–2888, <https://doi.org/10.1175/MWR-D-19-0370.1>, 2020.
 725 Liu, J., Wang, B., and Xiao, Q.: An evaluation study of the DRP-4DVar approach with the
 726 Lorenz-96 model, *Tellus A: Dynamic Meteorology and Oceanography*, 63, 256–262,
 727 <https://doi.org/10.1111/j.1600-0870.2010.00487.x>, 2011.
 728 Mantua, N. J., Hare, S. R., Zhang, Y., Wallace, J. M., and Francis, R. C.: A Pacific interdecadal
 729 climate oscillation with impacts on salmon production, *Bulletin of the American*
 730 *Meteorological Society*, 78, 1069–1080, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0477(1997)078<1069:APICOW>2.0.CO;2)
 731 [0477\(1997\)078<1069:APICOW>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<1069:APICOW>2.0.CO;2), 1997.
 732 Meehl, G. A., Richter, J. H., Teng, H., Capotondi, A., Cobb, K., Doblas-Reyes, F., Donat, M.
 733 G., England, M. H., Fyfe, J. C., Han, W., Kim, H., Kirtman, B. P., Kushnir, Y., Lovenduski,
 734 N. S., Mann, M. E., Merryfield, W. J., Nieves, V., Pegion, K., Rosenbloom, N., Sanchez,
 735 S. C., Scaife, A. A., Smith, D., Subramanian, A. C., Sun, L., Thompson, D., Ummenhofer,
 736 C. C., and Xie, S.-P.: Initialized Earth system prediction from subseasonal to decadal
 737 timescales, *Nature Reviews Earth & Environment*, 2, 340–357,
 738 <https://doi.org/10.1038/s43017-021-00155-x>, 2021.
 739 [Morioka, Y., Doi, T., Storto, A., Masina, S., and Behera, S. K.: Role of subsurface ocean in](#)
 740 [decadal climate predictability over the South Atlantic, *Scientific Reports*, 8, 8523,](#)
 741 <https://doi.org/10.1038/s41598-018-26899-z>, 2018.
 742 Mulholland, D. P., Laloyaux, P., Haines, K., and Balmaseda, M. A.: Origin and impact of
 743 initialization shocks in coupled atmosphere–ocean forecasts, *Monthly Weather Review*,
 744 143, 4631–4644, <https://doi.org/10.1175/MWR-D-15-0076.1>, 2015.
 745 Penny, S. G., Bach, E., Bhargava, K., Chang, C.-C., Da, C., Sun, L., and Yoshida, T.: Strongly
 746 coupled data assimilation in multiscale media: Experiments using a quasi-geostrophic
 747 coupled model, *Journal of Advances in Modeling Earth Systems*, 11, 1803–1829,
 748 <https://doi.org/10.1029/2019MS001652>, 2019.
 749 Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P.,
 750 Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night

marine air temperature since the late nineteenth century, *Journal of Geophysical Research: Atmospheres*, 108, 4407, <https://doi.org/10.1029/2002JD002670>, 2003.

Reckinger, S. M., Petersen, M. R., and Reckinger, S. J.: A study of overflow simulations using MPAS-Ocean: Vertical grids, resolution, and viscosity, *Ocean Modelling*, 96, 291–313, <https://doi.org/10.1016/j.ocemod.2015.09.006>, 2015.

Ropelewski, C. F. and Halpert, M. S.: North American precipitation and temperature patterns associated with the El Niño/Southern Oscillation (ENSO), *Monthly Weather Review*, 114, 2352–2362, [https://doi.org/10.1175/1520-0493\(1986\)114<2352:NAPATP>2.0.CO;2](https://doi.org/10.1175/1520-0493(1986)114<2352:NAPATP>2.0.CO;2), 1986.

Saji, N. H., Goswami, B. N., Vinayachandran, P. N., and Yamagata, T.: A dipole mode in the tropical Indian Ocean, *Nature*, 401, 360–363, <https://doi.org/10.1038/43854>, 1999.

Saji, N. H., Xie, S.-P., and Yamagata, T.: Tropical Indian Ocean variability in the IPCC twentieth-century climate simulations, *Journal of Climate*, 19, 4397–4417, <https://doi.org/10.1175/JCLI3847.1>, 2006.

Shi, P., Lu, H., Leung, L.R., He, Y., Wang, B., Yang, K., Yu, L., Liu, L., Huang, W., Xu, S., Liu, J., Huang, X., Li, L., and Lin, Y.: Significant land contributions to interannual predictability of East Asian summer monsoon rainfall, *Earth's Future*, 9(2), e2020EF001762, <https://doi.org/10.1029/2020EF001762>, 2021.

Shi, P., Wang, B., He, Y., Lu, H., Yang, K., Xu, S. M., Huang, W. Y., Liu, L., Liu, J. J., Li, L. J., and Wang, Y.: Contributions of weakly coupled data assimilation-based land initialization to interannual predictability of summer climate over Europe, *Journal of Climate*, 35, 517–535, <https://doi.org/10.1175/JCLI-D-20-0506.1>, 2022.

Shi, P., Leung, L. R., Wang, B., Zhang, K., Hagos, S. M., and Zhang, S.: The 4DEnVar-based weakly coupled land data assimilation system for E3SM version 2, *Geoscientific Model Development*, 17, 3025–3040, <https://doi.org/10.5194/gmd-17-3025-2024>, 2024a.

Shi, P., Leung, L. R., Lu, H., Wang, B., Yang, K., and Chen, H.: Uncovering the interannual predictability of the 2003 European summer heatwave linked to the Tibetan Plateau, *npj Climate and Atmospheric Science*, 7, 242, <https://doi.org/10.1038/s41612-024-00782-3>, 2024b.

Shi, P., Leung, L. R.₂ and Wang, B.: Development and evaluation of a new 4D_{En}Var-based weakly coupled ocean data assimilation system in E3SMv2, *Geoscientific Model Development*, 18, 2443–2460, <https://doi.org/10.5194/gmd-18-2443-2025>, 2025.

Sluka, T. C., Penny, S. G., Kalnay, E.₂ and Miyoshi, T.: Assimilating atmospheric observations into the ocean using strongly coupled ensemble data assimilation, *Geophysical Research Letters*, 43, 752–759, <https://doi.org/10.1002/2015GL067238>, 2016.

Tang, Q., Mu, L., Goessling, H. F., Semmler, T.₂ and Nerger, L.: Strongly coupled data assimilation of ocean observations into an ocean-atmosphere model, *Geophysical Research Letters*, 48, e2021GL094941, <https://doi.org/10.1029/2021GL094941>, 2021.

Taylor, K. E., Stouffer, R. J.₂ and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bulletin of the American Meteorological Society*, 93, 485–498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.

Trenberth, K. E. and Stepaniak, D. P.: Indices of El Niño evolution, *Journal of Climate*, 14, 1697–1701, [https://doi.org/10.1175/1520-0442\(2001\)014<1697:LIOENO>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<1697:LIOENO>2.0.CO;2), 2001.

Trenberth, K. E. and Shea, D. J.: Atlantic hurricanes and natural variability in 2005, *Geophysical Research Letters*, 33, L12704, <https://doi.org/10.1029/2006GL026894>, 2006.

Turner, A. K., Lipscomb, W. H., Hunke, E. C., Jacobsen, D. W., Jeffery, N., Engwirda, D., Ringler, T. D.₂ and Wolfe, J. D.: MPAS-Seaice (v1.0.0): Sea-ice dynamics on unstructured Voronoi meshes, *Geoscientific Model Development*, 15, 3721–3751, <https://doi.org/10.5194/gmd-15-3721-2022>, 2022.

Wang, B., Liu, J., Wang, S., Cheng, W., Liu, J., Liu, C., Xiao, Q., and Kuo, Y. H.: An economical approach to four-dimensional variational data assimilation, *Advances in Atmospheric Sciences*, 27, 715–727, <https://doi.org/10.1007/s00376-009-9122-3>, 2010.

Wang, C.: Three-ocean interactions and climate variability: A review and perspective, *Climate Dynamics*, 53, 5119–5136, <https://doi.org/10.1007/s00382-019-04930-x>, 2019.

Wu, B., Zhou, T.₂ and Zheng, F.: EnOI-IAU initialization scheme designed for decadal climate prediction system IAP-DecPreS, *Journal of Advances in Modeling Earth Systems*, 10, 342–356, <https://doi.org/10.1002/2017MS001132>, 2018.

809 Zhang, S.: A study of impacts of coupled model initial shocks and state-parameter optimization
810 on climate predictions using a simple pycnocline prediction model, *Journal of Climate*,
811 24, 6210–6226, <https://doi.org/10.1175/JCLI-D-10-05003.1>, 2011.

812 Zhang, S., Liu, Z., Rosati, A., and Delworth, T.: A study of enhanceive parameter correction
813 with coupled data assimilation for climate estimation and prediction using a simple
814 coupled model, *Tellus A: Dynamic Meteorology and Oceanography*, 64, 10963,
815 <https://doi.org/10.3402/tellusa.v64i0.10963>, 2012.

816 Zhang, S., Liu, Z., Zhang, X., Wu, X., Han, G., Zhao, Y., Yu, X., Liu, C., Liu, Y., Wu, S., Lu,
817 F., Li, M., and Deng, X.: Coupled data assimilation and parameter estimation in coupled
818 ocean-atmosphere models: A review, *Climate Dynamics*, 54, 5127–5144,
819 <https://doi.org/10.1007/s00382-020-05275-6>, 2020.

820 Zhou, W., Li, J., Yan, Z., Shen, Z., Wu, B., Wang, B., Zhang, R., and Li, Z.: Progress and future
821 prospects of decadal prediction and data assimilation: A review, *Atmospheric and Oceanic*
822 *Science Letters*, 17, 100441, <https://doi.org/10.1016/j.aosl.2023.100441>, 2024.

823 Zhu, J., Kumar, A., Lee, H.-C., and Wang, H.: Seasonal predictions using a simple ocean
824 initialization scheme, *Climate Dynamics*, 49, 3989–4007, [https://doi.org/10.1007/s00382-](https://doi.org/10.1007/s00382-017-3556-6)
825 [017-3556-6](https://doi.org/10.1007/s00382-017-3556-6), 2017.

826 Zhu, S., Wang, B., Zhang, L., Liu, J., Liu, Y., Gong, J., Xu, S., Wang, Y., Huang, W., Liu, L.,
827 He, Y., and Wu, X.: A Four-Dimensional Ensemble-Variational (4DEnVar) Data
828 Assimilation System Based on GRAPES-GFS: System Description and Primary Tests,
829 *Journal of Advances in Modeling Earth Systems*, 14(7), e2021MS002737,
830 <https://doi.org/10.1029/2021MS002737>, 2022.




Figure 1. Time series of non-detrended (left column) and linearly detrended (right column) annual mean surface air temperature anomalies (units: °C) over (a, d) ocean, (b, e) land, and (c, f) global domains from 1965 to 2016. Black line: observation; blue line: CTRL; red line: ASSIM. Temperature anomalies are computed by removing both the climatological mean and long-term trend. For the detrended series, a linear trend is removed from the domain-averaged time series. The correlation coefficients of CTRL and ASSIM with the observation are also shown. The asterisk denotes statistically significant correlation at the 95% confidence level.

Deleted: detrended




Figure 2. Spatial patterns of climatological mean differences in surface air temperature (units: °C) between model simulations and observations for the period 1965–2016. Panel (a) shows the difference between CTRL and observation, and panel (b) shows the difference between ASSIM and observation. Dotted areas denote regions where the differences are statistically significant at the 95% confidence level.




Figure 3. Time series of linearly detrended annual mean precipitation anomalies (units: mm day^{-1}) over (a) ocean, (b) land, and (c) global domains from 1980 to 2016. Black line: observation; blue line: CTRL; red line: ASSIM. Precipitation anomalies are computed by removing both the climatological mean and long-term trend. For the detrended series, a linear trend is removed from the domain-averaged time series. The correlation coefficients of CTRL and ASSIM with the observation are also shown. The asterisk denotes statistically significant correlation at the 95% confidence level.




Figure 4. Spatial patterns of (a–c) climatological mean precipitation (units: mm day^{-1}) and (d, e) precipitation differences for the period 1980–2016. Panels (a–c) show precipitation climatology from observation, CTRL, and ASSIM, respectively. Panels (d, e) represent precipitation differences between model simulations and observations: (d) CTRL minus observation and (e) ASSIM minus observation. Dotted areas indicate regions where the differences are statistically significant at the 95% confidence level.





Figure 5. Time series of the (a) monthly and (b) winter Niño 3.4 index from 1965 to 2016 for the observation (black line), ASSIM (red line), and CTRL (blue line). The correlation coefficients with the observation for ASSIM and CTRL are also shown in parentheses. The asterisk indicates that the correlation is statistically significant at the 95% confidence level.



866
867 **Figure 6.** Power spectra (a–c) and seasonal cycle of the standard deviation (d–f) of the monthly
868 Niño 3.4 index from the observation (left column), ASSIM (middle column), and CTRL (right
869 column) for the period 1965–2016. The theoretical Markov "red noise" spectrum is shown as
870 a solid green line, with the 5% and 95% confidence bounds indicated by dashed blue and red
871 lines, respectively.





Figure 7. Time series of the (a–c) monthly and (d) winter Southern Oscillation Index (SOI) from 1965 to 2016 for the observation (black line), ASSIM (red line), and CTRL (blue line). Panels (a–c) show individual monthly time series from OBS, ASSIM, and CTRL, respectively, while panel (d) shows the winter SOI averaged over December–February (DJF). The correlation coefficients with the observation in ASSIM and CTRL are also indicated in parentheses. The asterisk indicates that the correlation is statistically significant at the 95% confidence level.



880
881 **Figure 8.** Time series of the autumn (September–November) Dipole Mode Index (DMI) from
882 1965 to 2016 for (a) the observation, (b) ASSIM, and (c) CTRL. The correlation coefficients
883 of ASSIM and CTRL with the observed DMI are also shown in the upper-right corner. The
884 asterisk indicates that the correlation is statistically significant at the 95% confidence level.




Figure 9. Time series and spatial patterns of the Pacific Decadal Oscillation (PDO) index from 1965 to 2016 for (a, d) the observation, (b, e) ASSIM, and (c, f) CTRL. The black lines in panels (a–c) denote the 10-year running mean of the annual PDO index. The numbers at the top right of (b) and (c) indicate the temporal correlations of ASSIM and CTRL with observations. The asterisk indicates that the temporal correlation is statistically significant at the 95% confidence level. The percentage of variance explained by EOF1 is shown at the top right of (d–f), and the numbers at the top center of (e) and (f) indicate the spatial correlations with the observed pattern. The EOF analysis is performed separately for the observations and model outputs after interpolating both onto a common horizontal grid and removing linear trends at each grid point.




Figure 10. Time series of the Interdecadal Pacific Oscillation (IPO) index from 1965 to 2016 for (a) the observation, (b) ASSIM, and (c) CTRL. The black line in each panel denotes the 10-year running mean of the annual IPO index. The numbers at the top right of (b) and (c) indicate the temporal correlations of ASSIM and CTRL with observations. The asterisk indicates that the temporal correlation is statistically significant at the 95% confidence level. The IPO index is derived from EOF analysis after interpolating both the observations and model outputs onto a common horizontal grid and removing linear trends at each grid point.




Figure 11. Time series of the Atlantic Multidecadal Oscillation (AMO) index from 1965 to 2016 for (a) the observation, (b) ASSIM, and (c) CTRL. The black line denotes the 10-year running mean. The numbers at the top right of (b) and (c) indicate the temporal correlations of ASSIM and CTRL with observations. The asterisk indicates that the correlation is statistically significant at the 95% confidence level. The AMO index is calculated by subtracting the global mean SST anomalies (60°S–60°N) from the North Atlantic SST anomalies (0–60°N, 0–80°W).




Figure 12. Spatial patterns of the differences in (a) correlation coefficients and (b) root-mean-square errors (RMSE; °C) of detrended winter surface air temperature anomalies between ASSIM and CTRL with observations from 1980 to 2016 over the contiguous U.S. Dotted areas indicate regions where the differences are statistically significant at the 95% confidence level.

Deleted: 90




Figure 13. Similar to Fig. 12 but for detrended winter precipitation. Dotted areas denote regions where the differences pass the 95% confidence level.





Figure 14. Regression patterns of winter surface air temperature (left column; shaded) and precipitation (right column; shaded), 500hPa geopotential height (contours), and 850hPa winds (vectors) on the standardized winter Niño 3.4 index for (a, d) the observation, (b, e) ASSIM, and (c, f) CTRL. In each panel, the Niño 3.4 index is calculated separately from its corresponding dataset and all regressions are based on simultaneous winter (DJF) averages. Dotted areas indicate regions where the regressions are statistically significant at the 95% confidence level.

928 **Appendix A: Supporting Information**



929
930 **Figure A1.** Time series of (a) global mean top-of-atmosphere (TOA) net energy flux (W m^{-2}),
931 (b) global mean surface air temperature ($^{\circ}\text{C}$), (c) global mean ocean temperature averaged over
932 the upper 1000 m ($^{\circ}\text{C}$), and (d) global mean ocean salinity averaged over the upper 1000 m
933 (psu). The blue and red lines denote the CTRL and ASSIM experiments, respectively.




Figure A2. Zonal-mean meridional streamfunction differences (units: $10^{10} \text{ kg s}^{-1}$) between model simulations and observations averaged over the period 1980–2016. Panel (a) shows the difference between CTRL and observations, and panel (b) shows the difference between ASSIM and observations. The vertical axis represents pressure levels (hPa), and the horizontal axis denotes latitude.




Figure A3. Lagged regression patterns of monthly winter (DJF) surface air temperature (left; shaded), precipitation (right; shaded), 500hPa geopotential height (contours), and 850hPa winds (vectors) onto the preceding month's (NDJ) standardized Niño 3.4 index for (a, d) the observation, (b, e) ASSIM, and (c, f) CTRL. Specifically, monthly atmospheric variables from December to February (DJF) are regressed onto the standardized monthly Niño 3.4 index from November to January (NDJ), corresponding to a one-month lead time. The Niño 3.4 index is calculated separately from each corresponding dataset. Dotted areas denote statistical significance at the 95% confidence level.