

Magnetotail response to corotating interaction region driven geomagnetic storms: Cluster observations

Adriane Marques de Souza Franco^{1,3}, Rashmi Rawat², Mauricio José Alves Bolzan³, Ezequiel Echer⁴

¹Federal University of Southern and Southeastern Pará (UNIFESSPA), Marabá, Brazil

²National Centre for Polar and Ocean Research (NCPOR), Goa, India

³Federal University of Jataí (UFJ), Jataí, Brazil

⁴Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, Brazil

Key Points:

- The HILDCAA events that occur during the recovery phase could cause a spread of energy to periodicities of 2 to 12 h in the auroral region.
- During both phases of geomagnetic storms driven by CIRS, cyclic substorm periods (≤ 4 hours) were found as dominant in the coupling between solar wind- magnetotail ($\text{IMF } B_z \times \text{magnetotail } B_x$), solar wind-auroral region ($\text{IMF } B_z \times \text{AE index}$) and magnetotail- auroral region ($\text{magnetotail } B_x \times \text{AE index}$).
- Spectral indices results suggest a strong turbulence in the magnetotail and auroral regions during recovery phases.

Corresponding author: Adriane Marques de Souza Franco, adrianemarquesds@gmail.com

Abstract

In this work we have selected 40 corotating/stream interaction (CIR) driven geomagnetic storms that occurred between 2001 and 2016, and statistically studied their impacts on the magnetotail. The wavelet transform was applied to the interplanetary magnetic field (IMF) B_z component, magnetotail B_x component, and the auroral electrojet (AE) index during geomagnetic storms. The cross-wavelet technique was applied to determine the periods of higher correlation between the IMF $B_z \times$ Tail B_x , IMF $B_z \times$ AE index and Tail $B_x \times$ AE index. More than 80% of the most energetic periods in the IMF B_z and magnetotail B_x are found to be shorter than 4 hours, independently of the storm phase. The AE index presented the range between 2-4 hours as the most common with energetic periods for both storm main and recovery phases. In the recovery phase, periodicities in the AE index are more spread (≤ 12 hours) than for the main phase (≤ 8 hours) probably due to the presence of high-intensity long-duration continuous AE activities (HILDCAAs). From the cross-wavelet analysis (IMF $B_z \times$ magnetotail B_x , IMF $B_z \times$ AE index and magnetotail $B_x \times$ AE index), periods ≤ 4 hours are found to be dominant in both storm phases, which coincide with the cyclic substorm periods. The power spectral analysis showed that the IMF B_z and magnetotail B_x time series follow the Kolmogorov (-5/3) power law. Additionally, the mean values of the spectral indices for the magnetotail B_x and AE index are higher during the recovery phase than the main phase. This suggests that turbulence is more pronounced during the recovery phase of geomagnetic storms driven by CIRs.

1 Introduction

A magnetized plasma is continuously emitted from the solar corona, the solar wind (Parker, 1958). This plasma propagates away from the Sun filling the interplanetary medium, with supermagnetosonic speed and carrying with it the solar magnetic field, called the interplanetary magnetic field (IMF) (Hundhausen, 1995). The IMF has a typical magnitude of the order of 5 to 10 nT near Earth's orbit (Baumjohann & Nakamura, 2007). As of today, the solar wind is broadly classified into two groups: high-speed streams (HSS) and slow streams (Schwenn, 2006). The slow speed streams have typical speed V_{sw} of $\sim 350 - 450 \text{ km s}^{-1}$ and originated mainly from the helmet streamers of the Sun (Gosling et al., 1981; Suess et al., 2009; Abbo et al., 2016). HSSs, on the other hand, with V_{sw} in the range of $\sim 750-850 \text{ km s}^{-1}$ (Phillips et al., 1995; Tsurutani et al., 2006) emanate from solar coronal holes (Krieger et al., 1973). When an HSS overtakes the ambient slower solar wind, a region of compressed plasma with an intense and fluctuating IMF is formed, called stream interaction regions or, if they persist for several solar rotations, corotating interaction region (CIR) (Smith & Wolfe, 1976; Tsurutani et al., 1995; Sanchez-Garcia, 2023, Chian et al., 2024). For brevity and consistency with earlier works we will call all events studied here as CIRs.

In the CIRs there is compression and acceleration of plasma in the slow solar wind streams in the opposite direction to the Sun, and compression and deceleration of the fast solar wind streams towards the Sun. At 1 AU (Astronomical Unit = distance from Earth to the Sun ≈ 150 million kilometers), CIRs, in general, do not present well-developed shocks (Smith and Wolfe, 1976; Tsurutani et al., 1995). However they can lead to the occurrence of geomagnetic storms and other significant geomagnetic activity occurrence (Tsurutani et al., 1995, 2006; Alves et al., 2006). The geoeffectiveness of CIRs is related to fluctuations caused by Alfvén waves that can result in intervals of southward IMF (Richardson, 2006). They are usually followed by moderate storms with smaller percentages of CIRs

97 leading to intense storms (Alves et al., 2006; Echer et al., 2008, 2013).
 98 Another type of geomagnetic activity, high-intensity long-duration continuous
 99 AE activities (HILDCAA) events, have their main cause associated to
 100 alfvénic fluctuations in HSS originating from coronal holes (Tsurutani and
 101 Gonzalez, 1987), since most of the events occurred during CIRs (Hajra et
 102 al., 2013, Souza et al. 2016).

103 Interplanetary Coronal Mass Ejections (ICMEs) are the main
 104 geomagnetic storm drivers during solar maximum and rising phases and
 105 tend to cause more intense events (Gonzalez et al., 2007; Echer et al.,
 106 2008; Rawat et al. 2018). On the other hand, in the declining phase, CIRs
 107 occur more often and become more important in the interaction between
 108 solar wind and the magnetosphere (Gonzalez et al., 1999, 2007). CIR
 109 driven geomagnetic storms are mostly moderate ($-50 \text{ nT} \geq Dst > -100 \text{ nT}$),
 110 but they show to have a higher integrated energy and therefore a long term
 111 higher effectiveness in the solar wind- magnetosphere coupling than
 112 geomagnetic storms driven by ICMEs (Turner et al., 2006). Thus they also
 113 present risks for space technology, communication power and transportation
 114 systems (Strugarek et al., 2014).

115 Koller et al. (2022) studied the relation between jets in the
 116 magnetosheath and large-scale structures from the solar wind, ICMEs and
 117 SIRs (stream interaction regions)/CIRs using THEMIS observations. Their
 118 results suggest that SIR/CIRs and HSS can enhance the number of
 119 geoeffective jets in the magnetosheath.

120 Denton et al. (2006) have found that CIR-driven geomagnetic
 121 storms produce a more significant modulation in the plasma sheet
 122 temperature than events caused by ICMEs.

123 The response of the ionosphere to 79 CIR events during
 124 2004-2009 using GPS-derived ionospheric total electron content (TEC)
 125 values was investigated by Chen et al. (2015). Those authors considered
 126 latitude, longitude, local time and storm onset time dependence. The
 127 daytime TEC response to CIR storms appeared to be different from that of
 128 ionospheric F2 peak densities at high and middle latitudes. The ionospheric

peak density seems to present long periods of negative response of about 2-3 days while the TEC response seems to be mostly positive. Further, a positive response from the electron density in the daytime can be observed, whereas the subauroral region showed a negative response (Chen et al., 2015).

Korth et al. (2006; 2011) studied the response of the magnetotail and polar cusps to CIRs using wavelet and cross-wavelet analysis. Their results showed periods of 2-3 hours of higher correlation between solar wind and magnetotail plasma parameters (Korth et al. 2006), while for the cusps, the periods found were around 15–25 minutes (Korth et al., 2011). The effects in the interior of the Earth's cusp and the recurrent substorm activity are due to the Alfvén waves present in the HSS (Korth et al. 2011).

Echer et al. (2017) studied the magnetosphere response, including magnetotail, to CIR and their HSS occurred in September to October 2003 during HSS-driven storms and substorms. It was found in that work that the magnetosphere exhibits a periodic response in the range 1.8 to 3 hours to solar wind HSS fluctuations. These responses were found to be globally consistent in the magnetotail, geosynchronous orbit, auroral region and ground-based magnetometers.

Typical ionospheric disturbances and magnetotail signatures of a magnetospheric substorm are observed during the passage of CIRs and HSS by Earth. Further, the aurora related to CIRs also showed similar signatures of substorms, with localized onset and expansion toward the pole and both longitudinal directions (Despirak et al., 2013; Despirak et al., 2014). Nevertheless, while auroras formed by substorms are confined to a small region, usually located in the midnight sector, auroras with large sizes were observed during CIR passages (Despirak et al., 2014).

The higher effectiveness of CIR-driven geomagnetic storms in the magnetosphere-ionosphere system is due to the fact that, although the CIR driven geomagnetic storms tend to be weaker than the storms caused by ICMEs, they usually have a longer recovery phase duration. Then CIR-driven storms show a larger value of integrated energy dissipated in the

161 magnetosphere during its duration (Turner et al., 2009).

162 The main aim of this work is to investigate the response of the
 163 magnetotail to geomagnetic storms caused by CIRs using statistical analysis.
 164 We intend to identify the dominant periods of IMF and magnetosphere
 165 fluctuations in the solar wind, tail and auroral region for Cluster tail's intervals
 166 and also find the periods of higher correlation between them. The
 167 characteristic energy distribution during the tail interval of these events, as
 168 well as correlation distributions were also investigated. In our study we have
 169 also divided the Cluster tail's crossings in intervals of main and recovery
 170 phases, with the aim to search for differences and similarities between them.

171

172

173 **2 Data and Methodology of analysis**

174 *2.1 Data sources*

175 In this study we have used data from the IMF B_z component,
 176 magnetotail B_x component and from the auroral electrojet index (AE). The
 177 IMF data were obtained from NASA's OMNI database
 178 (<https://omniweb.gsfc.nasa.gov/>) with a time resolution of 1 minute. The
 179 geomagnetic field B_x component data were obtained from the magnetometer
 180 onboard the SC1 spacecraft of the Cluster constellation from the European
 181 Space Agency (ESA) website (<https://www.cosmos.esa.int/web/csa/access>).
 182 We used 1 minute averaged Cluster magnetic field data in this work. The B_x
 183 component was used here because it can be an indicator of the geometry
 184 and large-scale dynamics of the magnetotail current sheet, providing
 185 signatures of processes like thinning, reconnection, and flapping, which can
 186 be useful in understanding the tail's response to CIR-driven storms
 187 (McPherron et al., 1973, Korth et al., 2006). In order to remove long-term
 188 effects observed in the geomagnetic field B_x , a technique of long-term trend
 189 extraction using the Haar wavelet function was applied in the data (Bolzan et
 190 al., 2020).

191

The auroral electrojet (AE) index was proposed by Davis and Sugiura in 1966 (Davis & Sugiura, 1966) and is used to measure geomagnetic activity in the auroral zone (Rostoker, 1972). The 1 minute AE index data are provided by the World Data Center for Geomagnetism (WDC), Kyoto, Japan (<https://wdc.kugi.kyoto-u.ac.jp/>). For the storm phase classification into main and recovery phases, we have used storm time disturbance indices Dst and $SYM-H$ with temporal resolutions of 1-hour and 1-minute respectively from WDC Kyoto (Sugiura, 1964; Iyemori 1990; Gonzalez et al., 1999). The storm main phase was marked between the times when sharp depression in $SYM-H$ was noted until the minimum value was attained while the recovery phase was defined as the period between time of minimum $SYM-H$ and recovery to pre-storm levels. The selection criterion followed the classification proposed by Gonzalez et al. (1994), where a moderate geomagnetic storm is defined by a minimum Dst peak value in the range $-100\text{nT} \leq Dst \leq -50\text{ nT}$.

In Table 1 the 40 CIR-driven geomagnetic storms that were found between 2001 and 2016 are presented. In addition to the date of occurrence, the peak Dst , V_{sw} , B_s (southward IMF B_z) and E_y (y-component of the electric field) values are also shown. From Table 1 it is noted that the event number 17 exhibited double peaks in Dst . The event occurred on 28 July, 2005, and the first Dst peak (50 nT) could be attributed to the CIR whereas HSS resulted in the second, smaller depression in Dst (peak Dst = - 41 nT). Alfvénic fluctuations in IMF B_z associated with HSS are distinctly visible. The density enhancements indicate the compression region of CIRs.

Table 1. CIR-driven geomagnetic storms during 2001–2016.

No.	Date	Peak Dst (nT)	Peak V_{sw} (km s^{-1})	Peak B_s (nT)	Peak E_y (mV m^{-1})
01	2001-09-23	73	568	9	4.7
02	2001-10-09	64	468	5.7	2.6
03	2002-09-04	109	415	18.3	6.8
04	2002-10-07	115	373	8.4	3
05	2002-10-14	100	356	13.6	4.6
06	2002-10-24	98	733	7.4	4.5
07	2002-10-31	52	513	5.7	2.9
08	2002-11-03	75	484	6.3	3
09	2003-07-12	105	595	12.5	7.4
10	2003-07-16	90	609	10.2	6.1
11	2003-07-27	57	773	14.3	6.3

12	2003-08-06	60	481	11.3	5.0
13	2003-08-07	61	641	8.6	5.4
14	2003-10-14	85	724	8.6	5.3
15	2004-10-13	63	431	8.6	3.5
16	2005-07-10	92	484	19.6	9.1
17	2005-07-28	50 (41)	419	5.2	2.2
18	2005-08-31	131	412	16	6.3
19	2005-09-10	73	748	5.1	3.6
20	2006-09-24	56	667	7.3	2.9
21	2007-10-25	51	694	4	2.3
22	2008-09-04	51	488	8.7	4.3
23	2008-10-11	60	421	11	3.9
24	2009-07-22	83	359	14.5	4.9
25	2010-10-11	75	360	11.6	4.2
26	2011-09-10	55	554	17.1	8.2
27	2013-08-27	54	411	10	4.1
28	2015-07-05	68	575	14	6.7
29	2015-07-23	64	459	11	4.8
30	2015-09-07	71	602	10.1	5.9
31	2015-09-11	82	641	10.8	5.3
32	2015-10-04	56	456	7.7	3.4
33	2015-10-07	93	462	8.7	4.0
34	2015-10-08	58	758	3.2	2.3
35	2015-10-10	50	611	3.1	1.7
36	2016-08-23	74	451	11.1	4.7
37	2016-09-01	59	433	7.5	3.2
38	2016-09-03	50	666	2.6	1.7
39	2016-09-29	66	705	5.2	3.7
40	2016-10-25	60	721	6.6	4.1

2.2 Methodology of analysis

In this section the methodologies used for developing this work are presented: wavelet and Fourier transforms.

2.2.1 Wavelet and cross-wavelet transform

In order to determine the main periodicities occurring in HSS and in the magnetotail and auroral regions, the wavelet transform (WT) was employed to the IMF, Tail and auroral data.

The WT allows us to investigate how the energy is distributed in time and period or frequency ranges in non-stationary time series (Torrence and Compo, 1998). The wavelet functions (known as "daughter wavelets") are generated from a simple generating function (called wavelet-mother):

$$\psi_{a,b} = \frac{1}{\sqrt{a}} \psi\left(\frac{t-b}{a}\right)$$

for $a, b \in \mathbb{Z}$ and $a \neq 0$, which undergoes expansions/contractions and translations in time. The WT transforms the signal $f(t)$ into a function with two variables (scale and time), represented by $WT(a, b)$ from Eq. (1):

$$WT(a, b) = \int f(t) \psi_{a,b}^*(t) dt, \quad (1)$$

where a is the dilation coefficient, b is the coefficient of translation in time, and $\psi_{a,b}^*(t)$ represents the complex conjugate of the mother-wavelet function.

The Cross-Wavelet Transform (XWT) (2) is constructed from two continuous WT of the variable time series (Grinsted et al., 2004; Bolzan, 2005) and can be obtained by Eq. (2).

,

$$XWT^{x,y}(a, b) = W^y(a, b) W^x(a, b)^* \quad (2)$$

In Eq. (2), W^x and W^y represent the WT applied to the time series $x(t)$ and $y(t)$, and $(*)$ represents the complex conjugate of the WT.

The periods with high energy or with higher correlation can be detected using the Global Wavelet Spectrum (GWS), which can be computed by

$GWS = \int |WT(a, b)|^2 db$ for the main periodicities. For the identification of the periods

with higher correlation, the GWS is obtained by $GWS = \frac{\sigma^x \sigma^y}{\sigma^x + \sigma^y} \int |XWT^{xy}(a, b)|^2 db$,

where σ^x and σ^y represent the variances of the time series, $x(t)$ and $y(t)$, respectively.

2.2.1.1 Energy distribution Classification

The wavelet power persistence can be classified into four types: Local, Intermittent, quasi-continuous, and continuous, based on their temporal duration and physical significance. Local distribution indicates power restricted to short intervals, representing transient phenomena like bursty bulk flows. The Intermittent characteristic describes

features appearing and disappearing at intervals, reflecting recurrent or stochastic processes. The Quasi-continuous distribution denotes signals present for most of the duration, suggesting a sustained but modulated response to solar wind drivers. Continuous classification refers to power persisting throughout the entire interval, signifying a steady-state, stable coupling between the solar wind and the magnetosphere (Souza et al., 2016).

269

270

2.2.2 Fourier Transform

271

272

273

274

275

276

277

278

279

280

281

282

In order to study the turbulence of space plasmas, one powerful technique is to apply Fourier transform to the data and obtain a power law spectrum for different frequency ranges. The Fourier spectrum of a time series is fitted by a power-law function as $P(f) = cf^\alpha$, where $P(f)$ is the Fourier spectral power, c is a constant, f is the frequency, and the α exponent is the power law index (Bolzan & Echer, 2014). While the spectral power gives an indication of the amplitude of turbulence fluctuations in different regions and frequency ranges, the power law index indicates how steep is the spectrum, i.e., how fast it changes with frequency, and how the rate of energy is transferred from large to small scales (Bruno et al., 2005). In this study, the Fourier analysis was used in order to obtain the spectral index from the solar wind, tail and auroral index data.

283

284

3 Results and Discussions

285

286

287

In this section, the wavelet, cross-wavelet and Fourier power law analysis results are shown for the main and recovery phases of CIR-driven geomagnetic storms during Cluster tail intervals.

288

289

290

291

292

As expected, for CIR-driven geomagnetic storms, the majority of the events are moderate storms. From the 40 events (Table 1), 87.5 % corresponds to moderate storm classification ($-50 \text{ nT} \geq Dst > -100 \text{ nT}$) while only 12.5 % (5 events) were intense storms ($-100 \text{ nT} \geq Dst > -250 \text{ nT}$). Weak and super intense storms were not identified in this dataset.

293

Figure 1 shows the time series of the IMF B_z (left upper panel),

294 solar wind velocity (left second panel), solar wind pressure (left third panel),
 295 magnetotail B_x (left fourth panel), AE index (left fifth panel), Dst index (left
 296 sixth panel), and $SYM-H$ index (left bottom panel) for the interval when the
 297 Cluster remained in the magnetotail during part of the geomagnetic storm
 298 caused by a CIR that is identified as event 6 from Table 1. The main phase
 299 is indicated by the gray area and corresponds to the interval between 00:00
 300 UT and 19:09 UT, 24 October 2002. From Fig. 1 one can notice that
 301 oscillations in the IMF B_z increase after 08:00 UT, while in the magnetotail B_x
 302 data, high amplitude oscillations are observed from 04:00 UT to 06:00 UT.
 303 These oscillations are related to the southward IMF B_z extended interval
 304 seen from about 01:00 to 06:00 UT. Several quasi-periodic oscillations in the
 305 B_x data can be seen from 06:00 to 18:00 UT. This seems to be related to
 306 the high amplitude IMF B_z fluctuations noted after about 10:00 UT. In the AE
 307 index we can observe peaks higher than 1000 nT in intervals close to both
 308 time ranges, as well as a low (0.31) anti-correlation with the oscillations of
 309 IMF B_z with a lag of 22 minutes. Lower values are observed around 12:00
 310 UT in the IMF B_z , followed by peaks around 13:00 UT, which decreases
 311 after 14:00 UT. While the opposite behavior is observed in the AE index,
 312 where we have peaks around 12:00 UT, lower values observed around
 313 13:00 UT and other peaks are seen after 14:00 UT. There are several
 314 substorm events noted in the AE index. One group seems to be associated
 315 with the first interval of negative B_z , while the second group of substorms is
 316 related to the fluctuating IMF B_z interval. In the Dst index, the characteristic
 317 sharp decrease that occurs during the geomagnetic storm main phase due
 318 to the increase of ionized particles being “trapped” in the magnetic field lines
 319 can be observed. The peak negative Dst (−98 nT) is reached at the end of
 320 the time series (19:09 UT on 24 October 2002). The typical signature of the
 321 main phase of the geomagnetic storm can be also seen in the $SYM-H$ index
 322 time series (bottom-left panel). The right panel shows the Cluster orbit,
 323 where the interval that covers the gray region from the time series is marked
 324 in pink.






Figure 1. (left) Temporal variations of IMF B_z , solar wind velocity (V_{sw}), solar wind pressure (P_{dyn}), magnetotail B_x , AE index, Dst , and $SYM-H$ indices during the main phase of a geomagnetic storm caused by a CIR between on 24 October 2002. The interval when Cluster remained in the magnetotail (00:00 UT and 19:09 UT on 24 October 2002) is shaded in gray (left panel). Right panel shows the Cluster orbit for the same interval marked in pink.

Figure 2 illustrates another example of a Cluster magnetotail during a CIR-driven geomagnetic storm (Event 18, Table 1), focusing on the recovery phase interval (shaded gray area). Following the format of **Fig. 1**, the panels on the left present the time series for IMF B_z (top), solar wind velocity (V_{sw}), and dynamic pressure (P_{dyn}), followed by the magnetotail B_x , the AE index, and the Dst and $SYM-H$ indices (bottom). In the last two panels, the recovery phase of the Dst and $SYM-H$ indices is clearly observed. The right panel displays the Cluster orbit, with the analyzed interval highlighted in pink.

From the top-left panel, as well as in the B_x and AE index plots (fourth and fifth panels, respectively), a prominent positive peak is observed across all three parameters, followed by a subsequent decrease. The IMF B_z exhibits oscillating patterns with several intervals of southward (negative) orientation. Furthermore, both the magnetotail and auroral regions show multiple substorm-like events, characterized by the typical signature of an AE index increase accompanied by a decrease in the tail B_x magnitude (Gonzalez et al., 1994).



Cluster orbit 2005 242 (08/30) 11:00 UT to 2005 244 (09/01) 20:05 UT



349

350 Figure 2. Time series of IMF B_z (left upper panel), solar wind velocity (V_{sw}), solar
 351 wind pressure (P_{dyn}), magnetotail B_x , AE index, Dst , and $SYM-H$ indices during the
 352 geomagnetic storm caused by a CIR between that occurred between 31 August
 353 and 01 September, 2005. The interval when Cluster remained in the magnetotail,
 354 from 19:00 UT, 31 August, to 08:40 UT, 01 September, 2005 is shaded in gray (left
 355 panel). The right panel shows the Cluster orbit. The time series interval is marked in
 356 pink.

357

358

359

360

3.1 Wavelet and Cross-Wavelet Analyses

361

362 The Wavelet transform was applied to the IMF B_z component, to
 363 the magnetotail B_x component and to the AE index for intervals when the
 364 Cluster spacecraft was in the magnetotail during CIR-driven geomagnetic
 365 storms. The analysis is subdivided for intervals of main and recovery
 366 phases. The cross-wavelet between these variables was also computed for
 367 both geomagnetic storm phase intervals in order to investigate at which
 periods the correlation between them is higher.

368

369

370

3.2 Wavelet Analysis

371

3.2.1 Main Phase

372

373 Figure 3 shows the wavelet spectrogram (map) applied to the IMF
 374 B_z component (top panel) and magnetotail B_x component (middle panel) and
 375 AE index (bottom panel) during a moderate geomagnetic storm main phase
 376 interval between 00:00 UT and 19:09 UT, 24 October 2002 (Event 06 from
 377 Table 1). In the upper panel, the left figure shows the wavelet spectrum of

the IMF B_z and the right figure presents the global wavelet spectrum (GWS). In the GWS, three periods with higher energy can be identified. The first, around 0.5 h, can be characterized as having local distribution, since higher energy (integrated wavelet power) is observed in the wavelet spectrum only in a small interval between $\sim 10:00$ UT and $\sim 15:00$ UT. The second period is of 1.5 hours and also presents local distribution, located in a similar region of the first period (from $\sim 10:00$ UT to $\sim 16:00$ UT). The last observed period is around 3.3 hours, and in the wavelet spectrum it can be observed continuously for a long time interval, from around $\sim 05:00$ UT to more than 16:00 UT, characterizing a quasi-continuous behavior. The quasi-continuous nature of the 3.3-hour period can indicate large-scale fluctuations in the solar wind's driving force. This is related to the characteristics of the high-speed stream (HSS) following the CIR, which can contain continuous fluctuations that drive the magnetosphere over a long duration (Borovsky and Denton, 2006).




Figure 3. Wavelet transform analysis for the interval when Cluster remained in the magnetotail during the main phase of a geomagnetic storm caused by a CIR between 00:00 UT and 19:09 UT, 24 October, 2002 for the IMF B_z component (top panel) and magnetotail B_x geomagnetic field component (middle panel) and AE index (bottom panel). Left panels show the wavelet power spectrum and right panels display the GWS.

402

403 In the middle panel of Fig. 3, the wavelet transform periodogram
 404 for the magnetotail B_x is shown. As for the IMF B_z wavelet spectrum, upper
 405 panel, the left figure shows the wavelet spectrum and the right panel shows
 406 the GWS. Four significant periods were identified in the GWS: ~ 0.4 hour, $\sim$
 407 1.3 hour, 2.4 hours and 4.2 hours. From the wavelet spectrum one can see
 408 that local distribution is observed in the two first periods, which are found
 409 from about 03:00 UT to 05:20 UT and from about 01:00 to 08:30 UT,
 410 respectively. The other two periods show quasi-continuous distribution,
 411 where the period of 2.4 hours is the most energetic. The strong energy at
 412 these periods suggests that the magnetotail's internal dynamics, possibly
 413 related to current sheet flapping or other global oscillations, are being
 414 excited and maintained by the external solar wind driver (Runov et al, 2005).
 415 The wavelet analysis of the AE index (bottom panel) shows a single,
 416 significant quasi-continuous period around 3 hours. This timescale is
 417 consistent with the substorm cycle (Borovsky et al., 1993; Korth et al.,
 418 2011).

419 The coincidence of this dominant period with the quasi-continuous
 420 periodicities found in both the solar wind (IMF B_z) and magnetotail (B_x)
 421 provides evidence of a direct and efficient coupling mechanism between the
 422 external driver and the magnetospheric response. It indicates that
 423 fluctuations in the solar wind (IMF B_z) are directly linked to magnetotail
 424 dynamics (B_x) and subsequently drive auroral substorm activity (AE).

425 We also applied the wavelet transform to the solar wind and
 426 magnetospheric data during the main phase intervals of the events from
 427 Table 1 when the Cluster spacecraft remained at least 10 hours in the
 428 magnetotail. A total of 6 events were analyzed (events: 6, 7, 9, 14, 26 and
 429 30). The number of events analysed for each variable varies due to gaps in
 430 the available data of IMF B_z and magnetotail B_x component. For the IMF B_z
 431 and magnetotail B_x , the WT was applied to data in 5 events: 6, 7, 9, 14 and
 432 30 for IMF B_z and 6, 7, 9, 14 and 26 for magnetotail B_x . The statistical result
 433 of this analysis is shown in the subsection below along with the results of the
 434 recovery phase.

3.2.2 Recovery phase

As mentioned before, the WT was also applied to the CIR-driven geomagnetic storm recovery phases during intervals when Cluster was in the magnetotail. Figure 4 shows one example, where the WT was performed for event 18 of Table 1, which occurred from 19:00 UT on 31 August to 08:40 UT on 1 September 2005 (intense storm).

Similarly as it was shown in Fig. 3, the upper panel shows WT for the IMF B_z component, the middle panel for the magnetotail B_x and the bottom panel shows the WT applied to the AE index. Again, the left panels correspond to the wavelet spectrum and the right panels to the global wavelet spectrum.

The GWS, shown in the top panel in Fig. 4, for IMF B_z , shows that two significant spectral peaks are observed, both with local distributions (left panel). The first period, near 1.0 h, is localized between 19:00 and 23:00 UT. The second one is observed at 2.0 hours, with higher energy (integrated wavelet power) concentrated between 23:00 UT on 31 August and ~04:20 UT on 1 September 2005..




Figure 4. Wavelet transform analysis for the interval when Cluster remained in the magnetotail during the recovery phase of a geomagnetic storm caused by a CIR between 19:00 UT on 31 August and 08:40 UT on 1 September 2005 for IMF B_z component (top panel) and magnetotail B_x geomagnetic field component (middle panel) and AE index (bottom panel). Left panels show the wavelet power spectrum and right panels display the GWS.

For the magnetotail B_x , shown in the right middle panel, the GWS

also shows only one significant period, very close to the periodicity observed for IMF B_z , 2.1 hours. However, for the magnetotail B_x component, the energy distribution has a quasi-continuous variation (left middle panel), and is observed from the beginning of Cluster interval until about 05:00 UT of 1 September. In the AE index, (bottom panels), the wavelet analysis shows a period around 3.6 hours in the GWS, which also has a quasi-continuous distribution, from the beginning of the interval to the period at about 06:00 UT on 1 September 2002. Other short periods can be seen in the GWS, but in the wavelet distribution (left bottom panel), the higher energy is observed outside the cone of influence (COI), which delimits the region that is not affected by edge effects. Therefore they were discarded.

While the IMF B_z shows localized, high-energy periods at 1.0 and 2.0 hours, the magnetotail B_x and AE index display quasi-continuous dominant periods at 2.1 and 3.6 hours, respectively. This continuous nature suggests that even in the recovery phase, the magnetosphere is being driven by sustained solar wind fluctuations, a typical characteristic of high-speed streams (Tsurutani and Gonzalez, 1987; Souza et al., 2018).

We followed the same criteria used to compute WT for both the main and recovery phases. For the Cluster tail's periods this interval should be ≥ 10 hours. For the recovery phase, a much higher number of events followed this criterion (29 events). It is worth mentioning that for some cases only one or two variables meet the criterion due to the absence of available data. Figure 5 shows the histograms for period ranges, with bins of 2 hours, for the three variables for the main phase (left panels) and recovery phase (right panels). This figure shows that, except for the magnetotail B_x , longer periods begin to be observed in the recovery phases, right top and bottom panels. For the IMF B_z component histograms, Fig. 5-a (main phase) and Fig. 5-b (recovery phase), we can see the same dominant ranges of periods for both geomagnetic storm phases. Periods ≤ 2 hours represent 50.0 % of the main periods during the main phase, and 54.2 % of the periods during the recovery phase. The range between 2 and 4 hours corresponds to more than 35 % of the periods for the main phase, and 31.3 % for the recovery phase. These two ranges are also predominant for the magnetotail B_x

periodicities. During HILDCAA events (Franco et al., 2019), the stronger magnetotail response to the IMF fluctuations are typically found in the substorm periodicity range (2 - 4 hours) (Borovsky et al., 1993; Franco et al., 2019).




Figure 5. Histograms with period ranges for the main phase (left panels) and recovery phase (right panels). From top to bottom the panels show: IMF B_z , magnetotail B_x and the AE index distributions.

For the magnetotail B_x , although the range between 2 and 4 hours corresponds to 25 % of the periods for the main phase, and 29.2 % for the recovery phase, periods ≤ 2 hours represent 68.7 % of the periods during the main phase and 61.1 % for the recovery phase. From our present results, it seems that during CIR-driven geomagnetic storms the magnetotail

has a stronger response to lower periods than during HILDCAA events. This could occur because the prolonged duration and wave-driven nature of HILDCAAs favor longer characteristic periods, while the shorter, more intense driving of the CIR main phase emphasizes the shorter, internal magnetotail response periods. Further, these results show us that more than 80 % of the most energetic periods in the IMF B_z and magnetotail B_x during geomagnetic storms driven by CIRs are found for periods shorter than 4 hours, independently of the storm phase. Similar periods (1.8 hours to 3.1 hours) were found by Echer et al. (2017) in a study of the solar wind - magnetotail coupling during HSS and CIR intervals.

The AE index analysis revealed that periods between 2 and 4 hours are the most prominent during both the main (42.9 %) and recovery (53.7 %) phases of geomagnetic storms. However, there are notable differences between these phases. During the main phase, periods between 4 and 6 hours are the second range most often observed, followed by periods shorter than 2 hours. In contrast, during the recovery phase, no significant periods shorter than 2 hours were observed. Instead, the distribution of higher-energy periods seems to extend to longer durations, with the 4-6 hour and 6-8 hour ranges sharing the same percentage of periods. This difference may be attributed to the occurrence of HILDCAA events during the recovery phase (Kamide et al., 1998). These events are often associated with Alfvénic trains in the IMF, and typically have periods between 4 and 12 hours (Souza et al., 2016). While the increased number of analyzed recovery phase events could potentially influence the period distribution, we believe this is not the primary cause of the observed distribution. This is supported by the fact that the spread in periods was not observed for the magnetotail B_x component but it was noted only in the IMF B_z component distribution, which oscillations are usually due to Alfvén waves during CIR intervals. Furthermore, the absence of shorter periods during the recovery phase cannot be solely explained by the higher number of analyzed cases.

Figure 6 represents the energy distribution across the main identified periods for each variable and period range during the main and

recovery phases. Considering all ranges, the continuous behavior dominates the IMF B_z energy distribution during the main phase with 35.7 % of the cases, which is also observed for individual ranges, except for the range ≤ 2 hours, at which local distribution dominates. During the recovery phase, the local and intermittent distributions are the most common observed energy distribution types, while the continuous distributions represent only 3.6 % of the most energetic periods identified. Further, note that during the main phase, the local characteristic dominates the range with the larger occurrence of energetic periods (≤ 2 h), while during the recovery phase, the intermittent behavior is dominant for this period range. This distinction in energy distribution characteristics is significant, as the main phase of geomagnetic storms is frequently characterized by enhanced substorm activity and the most intense occurrence of tail dipolarizations (Lee & Min, 2002).




Figure 6- Color maps of energy distribution classification percentage. Maps for IMF B_z (upper panels), magnetotail B_x (middle panels), and the AE index (bottom panels) display the percentage distribution across period ranges (x-axis) and energy distribution classification (y-axis). Results are separated by storm phase: main phase (left column) and recovery phase (right column).

Near Earth, these events can be associated with localized fast plasma jets during intense substorms, which may justify the local distribution observed in the most energetic periods during the main phase (Nakamura et

al., 2002, 2017). The quasi-continuous behavior observed in the recovery phase was also seen in the IMF B_z component during HILDCAA events. For shorter periods (≤ 2 hours), an intermittent distribution was observed, which might be related to the events that occur during geomagnetic storm recovery phases (A. M. Souza, 2015; A. Souza et al., 2016).

For the magnetotail B_x component, the local type distribution of the periodicities was the most often observed for both main and recovery phases considering all ranges. The same characteristic is observed for the range with the higher occurrence of the main periods (≤ 2 h) with 72.7 % for the main phase, which could also be related to localized fast plasma jets in the tail (Nakamura et al., 2017). The same predominant distribution was also observed for the recovery phases (66.7 %). This result differs from those found in a previous magnetotail study during HILDCAAs (Franco et al., 2019), that showed that more than 50 % of the magnetotail B_x periods presented quasi-continuous and intermittent distributions, which is a characteristic of substorms. In the AE index, quasi-continuous and intermittent distributions dominate for all ranges in the main phase. Intermittent behavior was also observed for periods ≤ 2 hours. In the recovery phases, the quasi-continuous characteristic is more frequent for all ranges, which is also seen for the main range, between 2 and 4 hours. Again, these results differ from that observed from HILDCAAs (Souza et al., 2016), which presented intermittent energy distribution as the most observed type of periodicity in the wavelet spectrum. As it was perceived during supersubstorms studied by (Hajra et al., 2023), during geomagnetic storms, in short periods, the local and intermittent behavior is dominating while for longer periods, quasi-continuous and continuous distributions are more often observed.

600

601

602 3.3 Cross-Wavelet Analysis

603 With the aim to identify the periodicities at which there is higher
604 correlation between the solar wind and the magnetotail (IMF $B_z \times$
605 magnetotail B_x), between solar wind and the auroral region (IMF $B_z \times AE$
606 index) and between the magnetotail and the auroral region (magnetotail $B_x \times$

AE index) during the main and recovery phases of these events, the cross-wavelet transform was employed. This study allows us to identify the main periods at which the energy transfer of the solar wind to the magnetotail, as well as the energy transfer from magnetotail-stored energy to the auroral region, occurs during geomagnetic storms caused by CIRs.

612

613

3.3.1 Main Phase

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

The cross-wavelet results applied for the interval of the magnetotail Cluster crossing during the main phase of the event 6 (24 October 2002) of CIR-driven geomagnetic storm presented in Table 1 are shown in Fig. 7. For this same event the wavelet transform was applied (presented in Fig. 3). The cross wavelet spectrum is presented in the left panels: IMF B_z component and magnetotail B_x (upper panel), IMF B_z component and AE index (middle panel) and magnetotail B_x and AE index (bottom panel). The GWS is shown in the right panel of the respective figure. Then the XWT between IMF B_z and magnetotail B_x (top panel) and between magnetotail B_x and AE index (bottom panel) shows three similar main periods of correlation around: 1.30 h, 2.60 h and 4.20 h. The local distribution dominates these periods, except for the period near 2.60 h of the bottom panel (magnetotail $B_x \times$ AE index), where quasi-continuous behavior is observed. In the middle panel (IMF $B_z \times$ AE index) only one significant period is seen, which shows local distribution and a period of 3.21 h. Note that in the cross-wavelet spectrum, the region with higher correlation occurs between 08:00 UT and 15:00 UT for the top and bottom panels, which is also observed in the middle panel, but with a shift in time (10:00 UT and 16:30 UT). These similar dominant periods (1.30 h, 2.60 h and 4.20 h) of correlation between solar wind-magnetosphere-auroral region oscillations can indicate resonant modes of the coupling of the solar-wind–magnetosphere–ionosphere system, and represent the time scales at which the energy transfer is most efficient. This supports the idea of a linked chain of events: solar wind fluctuations drive magnetotail oscillations, which in turn drive substorms and auroral activity at the same time scales.




Figure 7. Cross-wavelet between IMF B_z component and magnetotail B_x (left upper panel), IMF B_z component and AE index (left middle panel) and between magnetotail B_x and AE index (left bottom panel) for the interval when Cluster remained in the magnetotail during the main phase of a geomagnetic storm caused by a CIR that occurred between 00:00 UT and 19:09 UT on 24 October 2002. The right panels show the GWS.

651 Following the WT analysis approach, we initially planned to apply
 652 the cross-wavelet transform to the 6 events with data intervals of at least 10
 653 hours. However, as not all analyzed variables met the duration criterion
 654 within these events, the number of events suitable for cross-wavelet analysis
 655 varied depending on the availability of IMF B_z and magnetotail B_x data. The
 656 XWT was applied between IMF B_z and magnetotail B_x in only 4 events (6, 7,
 657 9 and 14 from Table 1). For IMF B_z and the AE index, it was applied in 5
 658 events (6, 7, 9, 14 and 30 from Table 1), and for magnetotail B_x and the AE
 659 index, it was applied in 5 events (6, 7, 9, 14 and 26 from Table 1). The
 660 results of this analysis are presented in the following subsection, combined
 661 with the recovery phase analysis results.

662

663

664

3.3.2 Recovery Phase

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

Figure 8 shows the cross-wavelet transform applied to the event
 18 (31 August to 1st September 2005) from Table 1, the same event shown
 in Fig. 4. The left panels show the cross wavelet spectrum between the IMF
 B_z component and magnetotail B_x (upper panel), IMF B_z component and AE
 index (middle panel) and between magnetotail B_x and AE index (bottom
 panel). The right panel shows the GWS, where we can note that the period
 near 2.28 hours is present in the three cases. These intervals show
 quasi-continuous behavior for the top and bottom panels, and local
 correlation distributions for the middle panel (IMF $B_z \times AE$ index). Further,
 periods with higher correlation are observed near 0.80 h and ~ 3.5 h in the
 GWS of the middle (IMF $B_z \times AE$ index) and bottom (magnetotail $B_x \times AE$
 index) panels. For both cases, the periodicity around 0.80 h shows local
 distribution, while the 3.5 h, shows a quasi-continuous distribution. In the
 middle panel, (IMF $B_z \times AE$ index), an intermittent higher correlation
 distribution is found for the period of 1.31 h.




Figure 8. Cross-wavelet between IMF B_z component and magnetotail B_x (left upper panel), IMF B_z component and AE index (left middle panel) and between magnetotail B_x and AE index (left bottom panel) for the interval when Cluster remained in the magnetotail during the recovery phase of the geomagnetic storm caused by a CIR between 19:00 UT on 31 August and 08:40 UT on 1

September 2005. The right panels show the GWS.

In Figure 9, histograms with the range of periods with higher correlation obtained from the XWT are presented for the main and recovery phases. As it was seen in the wavelet analysis, in the main phase the periodicities are restricted for periods shorter than 8 hours, while during the recovery phase, except for the magnetotail $B_x \times$ AE index analysis, longer periods are observed. Despite that, even for the recovery phases, more than 95 % of the periods are lower than 8 hours. Similar periods that were found for the IMF B_z WT analysis were obtained in the correlation between the IMF B_z and magnetotail B_x (upper panel). Most of the higher correlation intervals, ~ 54 % for the main phase and $\sim 44\%$ (recovery phase), occurred for shorter periods (≤ 2 hours). Further, the second range (2-4 hours) also represents about 30 % of the periods with higher correlation for main (~ 31 %) and recovery (~ 33 %) phases. These results are in agreement with those results that were found in the XWT analysis performed by (Franco et al., 2019) for HILDCAAs events.

A significant difference can be observed between the main and the recovery phases in the periods of correlation between the IMF B_z and the AE index. Although the dominant range is the same (2 – 4 hours) in both phases, during the recovery phase, periods ≤ 2 hours represent more than 40 % of periods while in the main phase they represent only 21.4 %. Furthermore, considering ranges with longer periods, the main phase has more than 21 % of the periods of correlation between 4 and 8 hours, while less than 15 % of the periods of correlation during the recovery phase are found for periods > 4 hours. This change reflects a transition from external, sustained driving to the magnetosphere's internal process of unloading stored energy through a series of more frequent, cyclic substorms. Further, the dominance of periods ≤ 4 hours in the recovery phase coincide with cyclic substorm periods (Korth et al., 2011). The periodicity with higher correlation observed in the range between 10 and 12 hours may be related to Alfvén waves present in the IMF B_z during periods of HILDCAAs in the tail (Smith et al., 1995; Souza et al., 2016).

In the analysis of the periods at which the energy stored in the

magnetotail is transferred to the auroral region (XWT between magnetotail $B_x \times$ AE index), the histogram with the periods of correlation distribution is presented in the lower panels of Fig. 9. In the main phase (left panel), the percentage of periods of correlation per range decreased as the periods increased. About 47 % of the periods with higher correlation are ≤ 2 hours. The range between 2-4 hours corresponds to almost 30% of the periods, followed by 17.65 % (4-6 hours) and 5.88 % in the range between 6 and 8 hours. In the recovery phase (right panel) the range 0-2 hours and the range 2-4 hours showed the same percentage of periods (41.93 %). The others ~ 16 % intervals are found in the range between 4-6 hours, indicating that all the periods with higher correlation are shorter than 6 hours. As the periods with higher energy from the previous analysis, the periodicities with higher correlation values occur for substorms.





Figure 9. Histograms for period ranges with significant correlation (cross wavelet power) for the main phase (left panels) and recovery phase (right panels). From top to bottom the panels show: IMF $B_z \times$ magnetotail B_x , IMF $B_z \times$ magnetotail B_x and magnetotail $B_x \times$ AE index distributions.

Figure 10 shows the results with the classification of the type of correlation distribution. Considering all periods, the local distribution is dominant in three investigated cases: IMF $B_z \times$ magnetotail B_x , IMF $B_z \times$ AE index and magnetotail $B_x \times$ AE index (also intermittent) during the main phase. For the recovery phase, only the analysis between IMF B_z and magnetotail B_x data did not show the predominance of local distribution, with the intermittent behavior as the most commonly observed, probably related to substorm events.



753

754

Figure 10 – Color maps correlation distribution classification percentage obtained from the XWT analysis. Maps for IMF $B_z \times$ magnetotail B_x (upper panels), IMF $B_z \times$ AE index (middle panels), and magnetotail $B_x \times$ AE index (bottom panels) display the percentage distribution across period ranges (x-axis) and correlation distribution classification (y-axis). Results are separated by storm phase: main phase (left column) and recovery phase (right column).

760

3.4 Power Law analysis

In order to determine the spectral index for the solar wind, magnetospheric and auroral data during the geomagnetic storms main and recovery phases, we applied the Fourier analysis to the IMF B_z , magnetotail B_x and AE index data. The datasets were separated in two categories: main and recovery phases.

766

767

3.4.1 Main Phase

Results from the main phase are summarized in Table 2, and the spectral indices for event 7 are shown in Fig. 11a). According to Table 2, considering all analyzed events, it is possible to observe that mean values of the spectral indices were (-1.62 ± 0.04) for the IMF B_z , (-1.70 ± 0.22) for the magnetotail B_x and (-1.92 ± 0.38) for the AE index. Recently, Rakhmanovaa et al. (2020) studied the dependence of the spectral indices over plasma turbulence and showed that both the solar wind parameters and bow shock geometry have a great influence on the variation on the spectral index values. This fact shows that these values are sensible to this kind of parameters and the geometry from the space environment. These dependencies from the space geometry for the spectral indices are due to the formation of several kinds of phenomena, such as Alfvén waves, and also the coherent structures inside the plasma turbulence. This dependence of the spectral indices over the space environment was also demonstrated in Neptune day side magnetosheath fluctuations during Voyager-2 flyby (Echer et al., 2023) and during solar-wind high-speed streams (HSSs) at high heliolatitudes during the last two solar-cycle minima (SCM) (Echer et al., 2022).

Table 2. Spectral indices (IMF B_z , magnetotail B_x and AE index) obtained for Main Phase.

Spectral index (Main Phase)			
Events	IMF B_z	magnetotail B_x	AE index
6	-	-	-1.65
7	-1.67	-1.98	-2.14
9	-1.63	-1.67	-2.24
21	-	-	-1.29
26	-1.56	-1.45	-2.20
30	-1.63	-	-2.05
Mean $\pm$ std	-1.62 ± 0.04	-1.70 ± 0.22	-1.92 ± 0.38

Bolzan (2005) showed that the coherent structures present in solar wind time series increase (in absolute value) the spectral indices. The idea is that, into the turbulence flow, these organized structures are responsible to lead the energy from large to small scales in the fast and energetic way like a cascade as proposed by (Kolmogorov, 1941). According to Müller & Biskamp (2000), MHD turbulence is driven by the Alfvén wave effect, which couples small-scale velocity and magnetic fluctuations. This results in an energy spectrum consistent with a $k^{-5/3}$ law (the K41

law), which also aligns with observations of solar wind turbulence. Thus, our results from spectral indices mean values found for the AE index are consistent with previous works, where the difference found between $-1.67 = -5/3$ and value $= -1.92$ are due to the intermittency phenomena present in the MHD turbulence.

An interesting aspect observed in this study is the difference found between spectral index mean values obtained for the AE index time series and the solar wind-magnetotail time series (IMF B_z and magnetotail B_x). Three important points are noted: first, the mean values for the IMF B_z and magnetotail B_x series matches the K41 law which corresponds to $k^{-5/3}$; second, despite the standard deviation, there is a notable difference between these spectral indices for these both kinds of time series; third, the mean values for the AE index are in agreement with results from (Tsurutani et al., 1990). Tsurutani et al. (1990) suggested that high amplitude fluctuations in the southward IMF component will generate flux variations in the magnetotail and may be related to the increase of the ionospheric conductivity, which can be associated with the higher spectral index values found for AE index.




Figure 11. Spectral indices for IMF B_z , magnetotail B_x and AE index for a) Main Phase (event 7) and b) Recovery Phase (event 1).

815

816

817 In a previous work, similar results were obtained for the study of the Earth's
 818 magnetotail variability during supersubstorms (Hajra et al., 2023). In that study it was
 819 found that the IMF B_z exhibits the lowest spectral index (compared to the tail B_x and
 820 SML auroral indices), thus, corroborating our results. It is important to mention that the
 821 solar wind turbulence is significantly amplified after crossing the bow shock, and
 822 plasma flows are generated in the magnetosheath region. In the nightside, eddy
 823 vortices are generated due to those large-scale magnetospheric flows, which can drive
 824 the tail turbulence (Antonova & Stepanova, 2021).

824

825 **Table 3.** Spectral indices (IMF B_z , magnetotail B_x and AE index) obtained for
 826 Recovery Phase.

827

Events	Spectral index	(Recovery Phase)	AE index
	IMF B_z	magnetotail B_x	
1	-1.66	-2.09	-2.4
2	-	-	-2.37
3	-	-	-2.36
4	-	-	-2.31
5	-	-	-2.60
6	-	-	-1.85
9	-	-	-2.28
11	-1.53	-2.20	-2.38
12	-1.41	-1.79	-2.20
13	-1.74	-2.13	-2.08
15	-1.56	-1.69	-2.51
18	-1.57	-3.47	-2.32
20	-1.56	-1.81	-2.12
21_2	-1.83	-	-
22_1	-1.74	-	-
22_2	-1.74	-1.45	-
23_1	-1.74	-	-1.97
23_2	-1.65	-1.83	-
24_1	-	-1.46	-2.27
24_2	-1.40	-1.61	-2.42
25	-1.31	-1.98	-2.30
26	-	-	-2.13
27	-	-1.82	-1.97
29	-	-	-2.32
31	-	-	-2.32
32	-	-	-2.38
35	-	-	-2.29
36	-	-	-2.33
38_1	-	-	-2.01
38_2	-	-	-1.82
39	-	-	-1.92
Mean $\pm$ std	-1.60 ± 0.15	-1.95 ± 0.51	-2.23 ± 0.20

3.4.2 Recovery Phase

Similarly to the analyses conducted for the main phase intervals, we performed the Fourier Transform in all data in order to obtain the spectral indices during the recovery phase. An example is shown in Fig. 11b, which displays the spectral indices obtained for the parameters analyzed during Event 1. Table 3 shows the results where the mean values found for IMF B_z , magnetotail B_x and AE index were -1.60 ± 0.15 , -1.95 ± 0.51 and -2.23 ± 0.20 , respectively. For the IMF B_z index, the spectral index is lower in the recovery phase than during the main phase, while for the tail and auroral data, higher mean absolute values are found, where they were -1.95 ± 0.51 and -2.23 ± 0.20 , respectively. A sharp break is observed in the magnetotail B_x and AE index PSDs during the main phase, which is not seen during the recovery phase (Fig. 11). We may conjecture that during the recovery phase the geomagnetic field promotes a fast transfer of the energy from the large to small scales in order to leave the magnetosphere to return to almost steady state.

We also have separated the spectral indices from AE indices time series and magnetic ones (IMF B_z and B_x) where the results were, respectively, -2.23 ± 0.2 and -1.72 ± 0.38 . Note, again, the magnetic time series, IMF B_z and B_x , following the $K41$ law, corroborating results from (Müller & Biskamp, 2000). Furthermore, these results, similar to those observed in the main phase, suggest a characteristic of AE index time series: a distinct behavior compared to magnetic time series, as previously discussed in (Hajra et al., 2023). Also, despite the large standard deviation, it is possible to recognize that the mean values of the spectral indices for the recovery phases are greater if compared with those noted during the main phase. We can associate this result with the presence of the strong turbulence present during the recovery phase. It is just in this phase that the turbulent phenomena need to act in order to promote the transferring of energy from large to small scales like a Kolmogorov cascade.

4 Conclusions

In this work, the magnetotail response to geomagnetic storms driven by CIRs was investigated using wavelet and power spectrum analyses. By examining a

comprehensive dataset of 40 CIR-driven storms over a 15-year period (2001–2016), this study provides a significantly broader statistical basis than previously reported in the literature. The wavelet analyses applied to the IMF B_z and the Tail B_x time series showed that during CIR-driven geomagnetic storms the magnetotail responds stronger to lower periods than for HILDCAA events. Furthermore, we extended this analysis by distinguishing the magnetotail response between the main and recovery phases, allowing for quantitative comparison of turbulence levels.

The results demonstrate that more than 80% of the most energetic periods in the IMF B_z and magnetotail B_x in the tail during geomagnetic storms driven by CIRs were found at ≤ 4 hours, independently of the storm phase. For the AE index, periodicities between 2 and 4 hours dominate during both phases. A notable difference is that, during the main phase, the second range with higher number of periodicities was found in the interval between 4 and 6 hours, followed by periods ≤ 2 hours, while for the recovery phase, no energetic period was observed below 2 hours. Further, besides the dominant range, the number of periods of higher energy seems to be spread in longer periods, with the 4-6 hours and 6-8 hours ranges showing the same percentage of periods. The explanation of this difference may happen because during the recovery phase of geomagnetic storms, HILDCAA events often occur (Kamide et al., 1998 Tsurutani and Gonzalez 1987). It is well known that the main periodicities of the AE index during HILDCAAs are generally found between 4 and 12 h and that they are associated with Alfvénic trains in the IMF (Souza et al., 2016).

The XWT analyses performed to the magnetotail $B_x \times AE$ index and IMF $B_z \times AE$ index data indicated that the percentage of periods of correlation per range decreased as the periods increased. About 47 % of the periods of higher correlation are found at ≤ 2 hours for the main phase. For the recovery phase we obtained almost the same results. This analysis shows that the energy transfer from the solar wind, IMF B_z , to the magnetotail B_x , and finally to the auroral region, AE, occurs mainly at oscillations with periods ≤ 4 hours. Similar values were also found during HILDCAAs by (Franco et al., 2019). The HSS is associated with the generation of CIRs, and the periods found here correspond to cyclic substorm periods caused by Alfvén waves in the IMF during the HSS (Lee et al., 2006; Echer et al., 2017).

Results from spectral indices obtained for the IMF B_z during both the main and recovery phases showed agreement to K41 (-5/3) law. Besides, we observed a

difference between the spectral indices mean values obtained for *AE* indices time series and IMF *B_z* and Tail *B_x* for the main phase. These results show that the mean values for the interplanetary and magnetospheric time series are close to K41 law which corresponds to $k^{-5/3}$ while *AE* spectral index values are close to -1.93 . For the recovery phase, higher values of spectral indices were found for the magnetotail *B_x* (-1.92) and *AE* index -2.23 , corroborating results from Tsurutani et al. (1990). Two results can be summarized:

1) The IMF *B_z* and Tail *B_x* magnetic time series following the K41 ($-5/3$) law during the main phase, corroborating results from Müller & Biskamp (2000);

2) The mean values of the spectral indices for the recovery phase is greater, in the tail and auroral region, if compared with the main phase. We can associate this result with the presence of the strong turbulence present in these regions during the recovery phase.

Several studies were performed about the effect of geomagnetic activity in the Earth's magnetotail and they allowed us to have a general idea of the energy transfer from solar wind to the magnetotail during different kind of events (Nakamura & Kokubun, 2000; Nakamura et al., 2002; Franco et al., 2019; Hajra et al., 2023). Here we conducted a study on geomagnetic storms caused by CIRs in the Earth's magnetotail. The findings of this study support the conclusion that the magnetosphere responds to CIR-driven solar wind forcing mainly at the 2-4 hour timescales characteristic of substorms. It would be interesting to perform this kind of investigation about solar wind magnetosphere coupling on the magnetotail of other planets (Souza Echer et al., 2021).

917

918 Acknowledgments

The *AE* index and IMF *B_z* data used in this manuscript are public available at <https://wdc.kugi.kyoto-u.ac.jp/dstae/index.html> (WDC, 2018) and https://omniweb.gsfc.nasa.gov/form/sc_merge_min1.html (GSFC, 2018). A. M. S. F. would like to thank the Institute of Geosciences and Engineering- UNIFESSPA (project no. n°.23479.018587/2022-14). R. Rawat's work is supported by the Department of Science and Technology, India, under Women Scientist Scheme (project no. DST/WOS-A/EA-24/2020). E. Echer would like to thank Brazilian agencies for research grants: CNPq (contract no.PQ-301883/2019-0 and PQ-303900-2024-5). MJAB acknowledges to Brazilian agency CNPq to financial support (grant no. 304552 /2023 - 2). We thank the Brazilian Ministry of Science, Technology and Innovation and the Brazilian Space Agency as well.

926

927 References

928

Abbo, L., Ofman, L., Antiochos, S. K., Hansteen, V. H., Harra, L., Ko, Y. K., . . . Wang, Y. M. (2016,

- 930 November). Slow solar wind: Observations and modeling. *Space Science Reviews*, 201(1-4), 55–108.
 931 <https://doi.org/10.1007/s11214-016-0264-1>
- 932 Alves, M., Echer, E., & Gonzalez, W. (2006). Geoeffectiveness of corotating interaction regions as measured
 933 by dst index. *Journal of Geophysical Research*, 111. <https://doi.org/10.1029/2005JA011379>
- 934 Antonova, E. E., & Stepanova, M. V. (2021). The impact of turbulence on physics of the geomagnetic tail.
 935 *Frontiers in Astronomy and Space Sciences*, 8, 622570. <https://doi.org/10.3389/fspas.2021.622570>
- 936 Baumjohann, W., & Nakamura, R. (2007). Magnetospheric contributions to the terrestrial magnetic field.
 937 *Developments in Earth and Environmental Sciences*, 5, 77–92.
 938 <https://doi.org/10.1016/B978-044452748-6.00088-2>
- 939 Bolzan, M. J. A. (2005). Statistical and wavelet analysis of the solar wind data. *Brazilian Journal of Physics*,
 940 35, 592–596. <https://doi.org/10.1590/S0103-97332005000400002>
- 941 Bolzan, M. J. A., & Echer, E. (2014). A multifractal approach applied to the magnetic field turbulence in
 942 Jupiter's magnetosheath. *Planetary and Space Science*, 91, 77–82. <https://doi.org/10.1016/j.pss.2013.12.004>
- 943 Bolzan, M. J. A., Franco, A. M. S., & Echer, E. (2020). A wavelet based method to remove the long term
 944 periodicities of geophysical time series. *Advances in Space Research*, 66(2), 299–306.
 945 <https://doi.org/10.1016/j.asr.2020.04.014>
- 946 Borovsky, J. E., & Denton, M. H. (2006). Differences between CME-driven storms and CIR-driven storms.
 947 *Journal of Geophysical Research*, 111(A07S08). <https://doi.org/10.1029/2005JA011447>
- 948 Borovsky, J. E., Nemzek, R. J., & Belian, R. D. (1993). The occurrence rate of magnetospheric-substorm
 949 onsets: Random and periodic substorms. *Journal of Geophysical Research: Space Physics*, 98(A3),
 950 3807–3813. <https://doi.org/10.1029/92JA02556>
- 951 Bruno, R., Carbone, V., Veltri, P., Politano, H., & Malara, F. (2005). Identification of cascade in the solar
 952 wind: The role of the cross-helicity. *Planetary and Space Science*, 53(1–4), 49–54.
- 953 Chen, Y., Wang, W., Burns, A. G., Liu, S., Gong, J., Yue, X., . . . Coster, A. (2015). Ionospheric response to
 954 cir-induced recurrent geomagnetic activity during the declining phase of solar cycle 23. *Journal of*
 955 *Geophysical Research: Space Physics*, 120(2), 1394–1418. <https://doi.org/10.1002/2014JA020657>
- 956 Chian, A., Miranda, R., Bertucci, C., Blanco-Cano, X., Borovsky, J., Dasso, S., . . . Valdivia, J. (2024).
 957 Terrestrial and Martian space weather: A complex systems approach. *Journal of Atmospheric and*
 958 *Solar-Terrestrial Physics*, 259, 106253. <https://doi.org/10.1016/j.jastp.2024.106253>
- 959 Davis, T. N., & Sugiura, M. (1966). Auroral electrojet activity index AE and its universal time variations.
 960 *Journal of Geophysical Research*, 71(3), 785–801. <https://doi.org/10.1029/JZ071i003p00785>
- 961 Denton, M., Borovsky, J., Skoug, R., Thomsen, M., Lavraud, B., Henderson, M., . . . Liemohn, M. (2006).
 962 Geomagnetic storms driven by icme- and cir-dominated solar wind. *Journal of Geophysical Research*, 111.
 963 <https://doi.org/10.1029/2005JA011436>

- Despirak, I., Lubchich, A., & Koleva, R. (2014). Magnetospheric signatures of auroral disturbances during the passage of the solar wind's CIR and sheath regions. *Cosmic Research*, 52, 37–45. <https://doi.org/10.1134/S001095251401002X>
- Despirak, I. V., Lyubchich, A. A., & Koleva, R. (2013). Magnetotail signatures of auroral disturbances associated with Sheath and CIR regions in the solar wind. In K. Georgieva (Ed.), *Proceedings of the fifth workshop “solar influences on the magnetosphere”* (p. E13).
- Echer, E., González, W. D., Tsurutani, B. T., & Alves, M. V. (2013). Interplanetary conditions leading to intense geomagnetic storms ($Dst \leq -100$ nT) during solar cycle 23 (1996–2008). *Journal of Atmospheric and Solar-Terrestrial Physics*, 102, 236–241. <https://doi.org/10.1016/j.jastp.2013.06.002>
- Echer, E., Franco, A. M. S., da Costa Junior, E., Hajra, R., & Bolzan, M. J. A. (2022). Solar-wind high-speed stream (hss) Alfvén wave fluctuations at high heliospheric latitudes: Ulysses observations during two solar-cycle minima. *Advances in Space Research*, 297(143). <https://doi.org/10.1007/s11207-022-02070-w>
- Echer, E., Franco, A. M. S., Magalhães, F. P., Bolzan, M. J. A., & Hajra, R. (2023). Study of Neptune dayside magnetosheath fluctuations during Voyager-2 flyby. *Advances in Space Research*, 71(8), 3468–3478. <https://doi.org/10.1016/j.asr.2022.12.022>
- Echer, E., Gonzalez, W. D., Tsurutani, B. T., & Gonzalez, A. L. C. (2008). Interplanetary conditions causing intense geomagnetic storms ($Dst \leq -100$ nT) during solar cycle 23 (1996–2006). *Journal of Geophysical Research (Space Physics)*, 113(A5), A05221. <https://doi.org/10.1029/2007JA012744>
- Echer, E., Korth, A., Bolzan, M. J. A., & Friedel, R. H. W. (2017). Global geomagnetic responses to the IMF BZ fluctuations during the September/October 2003 high-speed stream intervals. *Annales Geophysicae*, 35(4), 853–868. <https://doi.org/10.5194/angeo-35-853-2017>
- Franco, A. M. S., Echer, E., & Bolzan, M. J. A. (2019). Wavelet analysis of the magnetotail response to solar wind fluctuations during HILDCAA events. *Annales Geophysicae*, 37(5), 919–929. <https://doi.org/10.5194/angeo-37-919-2019>
- Gonzalez, W. D., Joselyn, J. A., Kamide, Y., Kroehl, H. W., Rostoker, G., Tsurutani, B. T., & Vasyliunas, V. M. (1994). What is a geomagnetic storm? *Journal of Geophysical Research: Space Physics*, 99(A4), 5771–5792. <https://doi.org/10.1029/93JA02867>
- Gonzalez, W., Echer, E., Clua-Gonzalez, A., & Tsurutani, B. (2007). Interplanetary origin of intense geomagnetic storms ($dst < -100$ nT) during solar cycle 23. *Geophysical Research Letters*, 34(0). <https://doi.org/10.1029/2006GL028879>
- Gonzalez, W., Tsurutani, B., & Clúa de Gonzalez, A. (1999). Interplanetary origin of geomagnetic storms. *Space Science Reviews*, 88, 529–562. <https://doi.org/10.1023/A:1005160129098>
- Gosling, J. T., Borrini, G., Asbridge, J. R., Bame, S. J., Feldman, W. C., & Hansen, R. T. (1981). Coronal streamers in the solar wind at 1 AU. *Journal of Geophysical Research*, 86(A7), 5438–5448. <https://doi.org/10.1029/JA086iA07p05438>

- 999 Grinsted, A., Moore, J. C., & Jevrejeva, S. (2004). Application of the cross wavelet transform and wavelet
1000 coherence to geophysical time series. *Nonlinear Processes in Geophysics*, 11, 561–566.
1001 <https://doi.org/10.5194/npg-11-561-2004>
- 1002 Hajra, R., Echer, E., Franco, A. M. d. S., & Bolzan, M. J. A. (2023). Earth's magnetotail variability during
1003 supersubstorms (SSSs): A study on solar wind–magnetosphere–ionosphere coupling. *Advances in Space*
1004 *Research*, 72(4), 1208–1223. <https://doi.org/10.1016/j.asr.2023.04.013>
- 1005 Hajra, R., Echer, E., Tsurutani, B. T., & Gonzalez, W. D. (2013). Solar cycle dependence of high-intensity
1006 long-duration continuous AE activity (HILDCAA) events, relativistic electron predictors? *Journal of*
1007 *Geophysical Research: Space Physics*, 118(9), 5626–5638. <https://doi.org/10.1002/jgra.50530>
- 1008 Hundhausen, A. J. (1995). The Solar Wind. In M. G. Kivelson & C. T. Russell (Eds.), *Introduction to Space*
1009 *Physics* (pp. 91–128). Cambridge University Press.
- 1010 Iyemori, T. (1990). Storm-time magnetospheric currents inferred from mid-latitude geomagnetic field
1011 variations. *Journal of Geomagnetism and Geoelectricity*, 42(11), 1249–1265.
- 1012 Kamide, Y., Baumjohann, W., Daglis, I. A., Gonzalez, W. D., Grande, M., Joselyn, J. A., . . . Vasyliunas, V.
1013 M. (1998). Current understanding of magnetic storms: Storm-substorm relationships. *Journal of Geophysical*
1014 *Research: Space Physics*, 103(A8), 17705–17728. <https://doi.org/10.1029/98JA01426>
- 1015 Koller, F., Temmer, M., Preisser, L., Plaschke, F., Geyer, P., Jian, L. K., . . . Lamoury, A. T. (2022).
1016 Magnetosheath jet occurrence rate in relation to CMEs and SIRs. *Journal of Geophysical Research: Space*
1017 *Physics*, 127. [DOI ausente]
- 1018 Kolmogorov, A. N. (1941). The local structure of turbulence in an incompressible viscous fluid for very
1019 large Reynolds numbers. *Doklady Akademii Nauk SSSR*, 30(4), 299–304.
- 1020 Korth, A., Echer, E., Guarnieri, F. L., Franz, M., Friedel, R., Gonzalez, W. D., Mouikis, C. G., & Reme, H.
1021 (2006). Cluster observations of plasma sheet activity during the 14–28 September 2003 corotating high
1022 speed stream event. In A. Syrjäsuo & E. Donovan (Eds.), *Proceedings of the Eight International Conference*
1023 *on Substorms (ICS-8)* (pp. 133–138). University of Calgary.
- 1024 Korth, A., Echer, E., Zong, Q.-G., Guarnieri, F., Fraenz, M., & Mouikis, C. (2011). The response of the polar
1025 cusp to a high-speed solar wind stream studied by a multispacecraft wavelet analysis. *Journal of*
1026 *Atmospheric and Solar-Terrestrial Physics*, 73(1), 52–60. <https://doi.org/10.1016/j.jastp.2009.10.004>
- 1027 Krieger, A. S., Timothy, A. F., & Roelof, E. C. (1973). A coronal hole and its identification as the source of a
1028 high velocity solar wind stream. *Solar Physics*, 29(2), 505–525. <https://doi.org/10.1007/BF00150828>
- 1029 Lee, D.-Y., Lyons, L., Kim, K.-C., Baek, J.-H., Kim, K.-H., Weygand, J., . . . Han, W. (2006). Repetitive
1030 substorms caused by Alfvénic waves of the interplanetary magnetic field during high-speed solar wind
1031 streams. *Journal of Geophysical Research*, 111. <https://doi.org/10.1029/2006JA011685>
- 1032 Lee, D.-Y., & Min, K. W. (2002). Statistical features of substorm indicators during geomagnetic storms.
1033 *Journal of Geophysical Research: Space Physics*, 107(A11), SMP 16-1–SMP 16-12.

- 1034 <https://doi.org/10.1029/2002JA009243>
- 1035 McPherron, R. L., Russell, C. T., & Aubry, M. (1973). Satellite studies of magnetospheric substorms on
1036 August 15, 1968: 5. Phenomenological model for substorms. *Journal of Geophysical Research*, 78(16),
1037 3131–3143.
- 1038 Müller, W.-C., & Biskamp, D. (2000). Scaling properties of three-dimensional magnetohydrodynamic
1039 turbulence. *Physical Review Letters*, 84(3), 475–478.
- 1040 Nakamura, R., Baumjohann, W., Klecker, B., Bogdanova, Y., Balogh, A., Rème, H., . . . Runov, A. (2002).
1041 Motion of the dipolarization front during a flow burst event observed by cluster. *Geophysical Research*
1042 *Letters*, 29(20), 3-1–3-4. <https://doi.org/10.1029/2002GL015763>
- 1043 Nakamura, R., & Kokubun, S. (2000). Tail configuration during storms. *Advances in Space Research*, 25(7),
1044 1631–1638. [https://doi.org/10.1016/S0273-1177\(99\)00677-8](https://doi.org/10.1016/S0273-1177(99)00677-8)
- 1045 Nakamura, R., Nagai, T., Birn, J., Sergeev, V., Contel, O. L., Varsani, A., . . . Turner, D. L. (2017).
1046 Near-earth plasma sheet boundary dynamics during substorm dipolarization. *Earth, Planets, and Space*, 69.
- 1047 Parker, E. N. (1958). Dynamics of the interplanetary gas and magnetic fields. *Astrophysical Journal*, 128,
1048 664. <https://doi.org/10.1086/146579>
- 1049 Phillips, J., Bame, S., Feldman, W., Gosling, J., Hammond, C., McComas, D., . . . Suess, S. (1995). Ulysses
1050 solar wind plasma observations at high southerly latitudes. *Science (New York, N.Y.)*, 268, 1030–1033.
1051 <https://doi.org/10.1126/science.268.5213.1030>
- 1052 Rakhmanovaa, L. S., Riazantsevaa, M. O., Zastenkera, G. N., Yermolaeva, Y. I., & Lodkinaa, I. G. (2020).
1053 Dependence of the properties of a turbulent cascade behind the bow shock on the dynamics of the solar wind
1054 parameters. *Cosmic Research*, 58(6), 478–486.
- 1055 Rawat, R., Echer, E., & Gonzalez, W. D. (2018). How different are the solar wind-interplanetary conditions
1056 and the consequent geomagnetic activity during the ascending and early descending phases of the solar
1057 cycles 23 and 24? *Journal of Geophysical Research: Space Physics*, 123, 6621–6638.
1058 <https://doi.org/10.1029/2018JA025683>
- 1059 Richardson, I. G. (2006). The formation of CIRs at stream-stream interfaces and resultant geomagnetic
1060 activity. In *Recurrent magnetic storms: Corotating solar wind streams* (pp. 45–58). American Geophysical
1061 Union (AGU). <https://doi.org/10.1029/167GM06>
- 1062 Rostoker, G. (1972). Geomagnetic indices. *Reviews of Geophysics*, 10(4), 935–950.
1063 <https://doi.org/10.1029/RG010i004p00935>
- 1064 Runov, A., et al. (2005), Electric current and magnetic field geometry in flapping magnetotail current sheets,
1065 *Ann. Geophys.*, 23, 1391–1403.
- 1066 Sánchez-García, E., Gonzalez-Esparza, J. A., Aguilar-Rodríguez, E., & Corona-Romero, P. (2024). [Título
1067 do Artigo]. *Journal of Geophysical Research: Space Physics*, 129, e2023JA031685.

- 1068 <https://doi.org/10.1029/2023JA031685>
- 1069 Schwenn, R. Space Weather: The Solar Perspective. *Living Rev. Sol. Phys.* 3, 2 (2006).
1070 <https://doi.org/10.12942/lrsp-2006-2>
- 1071 Smith, E. J., Balogh, A., Neugebauer, M., & McComas, D. (1995). Ulysses observations of Alfvén waves in
1072 the southern and northern solar hemispheres. *Geophysical Research Letters*, 22(23), 3381–3384.
1073 <https://doi.org/10.1029/95GL03268>
- 1074 Smith, E. J., & Wolfe, J. H. (1976). Observations of interaction regions and corotating shocks between one
1075 and five au: Pioneers 10 and 11. *Geophysical Research Letters*, 3(3), 137–140.
1076 <https://doi.org/10.1029/GL003i003p00137>
- 1077 Souza, A., Echer, E., Bolzan, M., & Hajra, R. (2016). A study on the main periodicities in interplanetary
1078 magnetic field bz component and geomagnetic AE index during HILDCAA events using wavelet analysis.
1079 *Journal of Atmospheric and Solar-Terrestrial Physics*, 149, 81–86.
1080 <https://doi.org/10.1016/j.jastp.2016.09.006>
- 1081 Souza, A. M., Echer, E., Bolzan, M. J. A., & Hajra, R. (2018). Cross-correlation and cross-wavelet analyses
1082 of the solar wind IMF Bz and auroral electrojet index AE coupling during HILDCAAs. *Annales*
1083 *Geophysicae*, 36, 205–213. <https://doi.org/10.5194/angeo-36-205-2018>
- 1084 Souza, A. M. (2015). *Estudo do acoplamento vento solar-magnetosfera durante eventos HILDCAAs*
1085 *utilizando análise por wavelets*. [Dissertação de Mestrado/Tese de Doutorado, Instituição]. São José dos
1086 Campos.
- 1087 Souza Echer, M. P. A., Echer, E., Domingues, M., Mendes, O., Seo, R. T., & Gonzalez, W. (2021). Wavelet
1088 analysis of low frequency magnetic field fluctuations in the Jupiter's magnetotail. *Advances in Space*
1089 *Research*, 68(1), 246–258. <https://doi.org/10.1016/j.asr.2021.03.003>
- 1090 Strugarek, A., Janitzek, N., Lee, A., Löschl, P., Seifert, B., Hoilijoki, S., . . . Kargl, G. (2014). A space
1091 weather mission concept: Observatories of the solar corona and active regions (OSCAR).
- 1092 Suess, S. T., Ko, Y.-K., von Steiger, R., & Moore, R. L. (2009). Quiescent current sheets in the solar wind
1093 and origins of slow wind. *Journal of Geophysical Research: Space Physics*, 114(A4).
1094 <https://doi.org/10.1029/2008JA013704>
- 1095 Sugiura, M. (1964). Hourly values of equatorial Dst for the IGY. *Annals of the International Geophysical*
1096 *Year*, 35(1), 9–45.
- 1097 Torrence, C., & Compo, G. P. (1998). A practical guide to wavelet analysis. *Bulletin of the American*
1098 *Meteorological Society*, 79, 61–78. <https://doi.org/10.1175/1520-0477>
- 1099 Tsurutani, B. T., & Gonzalez, W. D. (1987). The cause of high-intensity long-duration continuous AE
1100 activity (HILDCAAs): Interplanetary Alfvén wave trains. *Planetary and Space Science*, 35(4), 405–412.
1101 [https://doi.org/10.1016/0032-0633\(87\)90097-3](https://doi.org/10.1016/0032-0633(87)90097-3)

- 1102 Tsurutani, B. T., Gonzalez, W. D., Gonzalez, A. L. C., Guarnieri, F. L., Gopalswamy, N., Grande, M., . . .
 1103 Vasyliunas, V. (2006). Corotating solar wind streams and recurrent geomagnetic activity: A review. *Journal*
 1104 *of Geophysical Research: Space Physics*, 111(A7). <https://doi.org/10.1029/2005JA011273>
- 1105 Tsurutani, B. T., Gonzalez, W. D., Gonzalez, A. L. C., Tang, F., Arballo, J. K., & Okada, M. (1995).
 1106 Interplanetary origin of geomagnetic activity in the declining phase of the solar cycle. *Journal of*
 1107 *Geophysical Research: Space Physics*, 100(A11), 21717–21733. <https://doi.org/10.1029/95JA01476>
- 1108 Tsurutani, B. T., Sugiura, M., Iyemori, T., Goldstein, B. E., Gonzalez, W. D., Akasofu, S. I., & Smith, E. J.
 1109 (1990). The nonlinear response of AE to the IMF BS driver: A spectral break at 5 hours. *Geophysical*
 1110 *Research Letters*, 17(3), 279–282. <https://doi.org/10.1029/GL017i003p00279>
- 1111 Turner, N. E., Cramer, W. D., Earles, S. K., & Emery, B. A. (2009). Geoefficiency and energy partitioning in
 1112 CIR-driven and CME-driven storms. *Journal of Atmospheric and Solar-Terrestrial Physics*, 71(10),
 1113 1023–1031. <https://doi.org/10.1016/j.jastp.2009.02.005>
- 1114 Turner, N. E., Mitchell, E. J., Knipp, D. J., & Emery, B. A. (2006). Energetics of magnetic storms driven by
 1115 corotating interaction regions: A study of geoeffectiveness. In *Recurrent magnetic storms: Corotating solar*
 1116 *wind streams* (pp. 113–124). American Geophysical Union (AGU). <https://doi.org/10.1029/167GM11>
 1117