

Exploring the Potential of LPJmL-5 to Simulate Vegetation Responses to (Multi-Year) Droughts

Denise Ruijsch¹, Sandra Margrit Hauswirth¹, Hester Biemans², and Niko Wanders¹

¹Department of Physical Geography, Utrecht University, Princetonlaan 8a, Utrecht, the Netherlands

²Water Systems and Global Change, Department of Environmental Sciences, Wageningen University & Research, Droevendaalsesteeg 4, Wageningen, The Netherlands

Correspondence: Denise Ruijsch (d.ruijsch@uu.nl)

Abstract.

Climate change is expected to increase the frequency and severity of Multi-Year Droughts (MYDs), but their impacts on vegetation remain poorly understood. While satellite records offer valuable insights, they cover only recent decades, limiting the number of MYDs available for analysis. Dynamic global vegetation models (DGVMs), such as LPJmL-5, can help overcome this limitation by simulating vegetation dynamics over longer timescales. However, their ability to capture drought impacts has not yet been systematically evaluated. In this study, we benchmark LPJmL-5 against MODIS-derived gross primary production (GPP) to assess how well it captures vegetation responses to drought. We find that LPJmL-5 reproduces GPP reasonably well in some regions, but ~~improvements can still be made in~~ there is a performance decline in parts of the Southern Hemisphere and ~~for in regions with~~ croplands. During MYDs, LPJmL-5 captures the key temporal and spatial GPP drought dynamics ~~observed~~ found in MODIS. However, the model tends to overestimate vegetation response at the onset of MYDs and shows some rapid recovery behaviour, resulting in muted overall drought impacts. Vegetation responses also vary by type: croplands show relatively good agreement, while boreal and temperate vegetation underestimate positive and negative impacts, respectively. These discrepancies appear to be linked to simplified model representations of vegetation stress and mortality, which limit long-term vegetation loss. Our results highlight the need to improve how LPJmL-5 simulates vegetation stress and recovery, especially under prolonged drought conditions, in order to better capture ecosystem vulnerability in a changing climate.

1 Introduction

Multi-Year Droughts (MYDs) can have severe impacts on natural vegetation and croplands, ranging from reduced vegetation health to diminished ecosystem resilience and increased tree mortality (Choat et al., 2018; DeSoto et al., 2020; Gessler et al., 2020; Allen et al., 2010; Cooley et al., 2015; Dong et al., 2019; Moravec et al., 2021; Jiao et al., 2020; Wittwer and Waschik, 2021; Hughes et al., 2019). Although less frequent than shorter droughts, MYDs have already been observed worldwide (van Mourik et al., 2025) and are projected to increase in both frequency and severity (van der Wiel et al., 2023).

However, ecosystem responses to drought are not universally negative (Ruijsch et al., 2025). Recent studies reveal a more nuanced picture, with some regions showing weak or even positive feedbacks between drought and vegetation (Ruijsch et al., 2025; van Hateren et al., 2021; Dong et al., 2019; Yang et al., 2021). These findings indicate that vegetation responses to (multi-year) droughts are not uniform, but rather that they vary with local water and energy availability (Ruijsch et al., 2025; van Hateren et al., 2021). Understanding this variability is crucial for assessing ecosystem vulnerability to MYDs in a changing climate and for developing effective adaptation strategies.

To investigate drought responses and to monitor vegetation status, previous studies have made use of satellite remote sensing datasets such as MODIS (Ruijsch et al., 2025; van Hateren et al., 2021; Dong et al., 2019; Yang et al., 2021). These datasets offer global coverage with high spatial and temporal resolution, making them valuable tools for analysing vegetation dynamics during droughts. However, MODIS records generally only extend back to 2000 (Didan, 2015a, b), limiting the number of MYDs available for analysis and reducing the statistical robustness of drought impact assessments based solely on remotely sensed data.

To address this limitation, process-based modelling approaches such as Dynamic Global Vegetation Models (DGVM) can complement satellite observations by simulating vegetation responses over longer timeframes. Among these models, LPJmL-5 (Schaphoff et al., 2018b, a; von Bloh et al., 2018) is widely used and extensively validated to represent establishment, growth and mortality of both natural ecosystems and managed croplands. By integrating carbon, water, nitrogen, and energy fluxes, LPJmL-5 simulates vegetation dynamics under changing climate and land-use conditions, enabling the extension of vegetation records beyond the satellite era and the analysis of a broader range of drought events.

However, ~~the ability of models like LPJmL-5 before DGVMs can be used to study MYD impacts is largely unknown. Therefore over longer historical periods or for future projections, it is essential to evaluate how well it captures vegetation responses to whether their simulated vegetation responses are consistent with observation-constrained signals. In this study, we assess the ability of LPJmL-5 to reproduce satellite-derived drought response patterns for both MYDs and normal droughts (NDs; droughts lasting less than a year), and if it can differentiate the impact of these different droughts on vegetation. In this study, we assess the spatial and temporal agreement between modelled and satellite-observed vegetation responses to both drought types, and investigate differences and similarities in vegetation drought response. Ultimately, it provides a foundation for using. Our aim is to determine whether LPJmL-5 to study can serve as a reliable tool for studying vegetation responses to droughts over extended historical periods and under future climate scenarios multi-year droughts beyond the period covered by satellite observations or whether potential improvements in process representation are required for future drought studies.~~

50 2 Materials and Methods

This section outlines the definitions, vegetation model, data, and methods used to evaluate vegetation responses to (multi-year) droughts (Figure 1). Section 2.1 defines MYDs and NDs, Section 2.2 describes the LPJmL-5 model, followed by model validation against observational data in Section 2.3. Sections 2.4 and 2.5 explain how vegetation responses to (multi-year) droughts are defined. Finally, Section 2.6 introduces the focus regions analysed in this study.




Figure 1. Schematic overview of the methodological workflow used in this study. The processes include drought classification, LPJmL-5 model simulations, comparison with MODIS-derived GPP observations, and the analysis of vegetation responses to both MYDs and NDs across selected focus regions. Blue elements represent observational data, while pink elements indicate model outputs.

55 2.1 Drought identification

Drought indices are essential for studying drought events, as they allow for consistent comparison of drought characteristics across space and time. In this study, we used the Standardized Precipitation Evapotranspiration Index (SPEI), which accounts for both water supply (precipitation, P) and atmospheric water demand (potential evapotranspiration, PET) (Vicente-Serrano et al., 2010). SPEI is particularly suitable for assessing drought impacts on vegetation as vegetation is not only impacted by lack of water supply via P, but also the increase in atmospheric water demand via PET.

SPEI was computed at multiple timescales (1–24 months) using P and PET derived from W5E5 reanalysis data (see Table 1, (Lange et al., 2022)). Both P and PET were aggregated to monthly values to generate monthly SPEI outputs. A detailed description of the SPEI calculation is provided in [Section Supplements S1](#).

To define MYDs and NDs, we focused on SPEI calculated at a 12-month timescale (SPEI-12), as it captures multi-year drought conditions while minimizing seasonal variability. Following Wiel et al. (2022), MYDs are defined as periods where SPEI-12 remains below -1 for at least 12 consecutive months. The threshold of -1 was chosen to represent moderate to extreme droughts (McKee et al., 1993). While the start date of the drought is only identified as the first month where SPEI-12 dips below -1, this definition already takes into account the accumulated P and PET over the 12 months leading up to the start of the drought. Ruijsch et al. (2025) evaluated the impact of different SPEI aggregation periods and concluded that the SPEI-12 is suitable for capturing multi-year drought dynamics.

To distinguish MYDs from shorter drought events, we define NDs as periods where SPEI-12 remains below -1 for less than 12 consecutive months. This definition ensures that each drought event is consistently categorized as either an MYD or ND, allowing us to systematically compare their respective impacts on vegetation.

Table 1. Overview of datasets and variables used in this study. W5E5 climate variables are grouped by their application in either SPEI calculation or LPJmL-5 simulations.

Dataset / Variable	Temporal range	Temporal resolution	Spatial resolution	Use / Reference
W5E5 Climate Forcing Data				
<i>Used for SPEI calculation</i>				
Precipitation [mm/day]	1901–2019	Daily	0.5° (~55 km)	Lange et al. (2022) SPEI input
Temperature [°C]	1901–2019	Daily	0.5°	PET calc. for SPEI
Relative humidity [%]	1901–2019	Daily	0.5°	PET calc. for SPEI
Wind speed [m s ⁻¹]	1901–2019	Daily	0.5°	PET calc. for SPEI
Surface pressure [Pa]	1901–2019	Daily	0.5°	PET calc. for SPEI
Downward shortwave radiation [W m ⁻²]	1901–2019	Daily	0.5°	PET calc. for SPEI
<i>Used in LPJmL-5 simulations</i>				
Precipitation [mm/day]	1901–2019	Daily	0.5°	Lange et al. (2022) LPJmL-5 input
Temperature [°C]	1901–2019	Daily	0.5°	LPJmL-5 input
Wind speed [m s ⁻¹]	1901–2019	Daily	0.5°	LPJmL-5 input
Downward longwave radiation [W m ⁻²]	1901–2019	Daily	0.5°	LPJmL-5 input
Downward shortwave radiation [W m ⁻²]	1901–2019	Daily	0.5°	LPJmL-5 input
Additional LPJmL-5 Inputs				
CO ₂ concentration [ppm]	1901–2019	Annual	Global	Büchner and Reyer (2022)
NH _x deposition [g/m ² /month]	1901–2019	Monthly	0.5°	Yang and Tian (2023)
NO _y deposition [g/m ² /month]	1901–2019	Monthly	0.5°	Yang and Tian (2023)
Soil pH	-	-	0.5°	Volkholz and Müller (2020)
Soil type	-	-	250 m (0.25 km)	Knoben (2021)
Land use	2020	-	5 min (~9.25 km)	International Food Policy Research Institute (IFPRI) (2024)
Land cover	2019	-	300 m (0.3 km)	Harper et al. (2023)
Remote Sensing Datasets				
MODIS GPP	2000–2019	16-day	1 km	Didan (2015a, b)

2.2 Dynamic global vegetation modelling using LPJmL-5

75 In this study, the DGVM LPJmL-5 (v5.7.9, von Bloh et al. (2018)) was used to simulate vegetation responses to MYDs. LPJmL-5 builds on LPJmL-4 and simulates both natural and agricultural vegetation, linking their growth and productivity through consistent water, carbon, and energy fluxes (Schaphoff et al., 2018b). LPJmL-5 makes use of the big-leaf representation for different Plant Functional Types (PFTs) that can coexist within a grid cell and compete for resources such as light, water, and nitrogen (von Bloh et al., 2018). Land cover can either be simulated dynamically or be prescribed, both of which are used in this study. In contrast, Crop Functional Types (CFTs) are always prescribed, with a distinction between irrigated and rainfed crops. Each grid cell can contain both managed and natural vegetation fractions, allowing for a realistic subgrid representation of land-use. CFTs only appear when crops are sown and grow on soil areas separate from natural vegetation (von Bloh et al., 2018). The model also includes key physiological processes such as photosynthesis, gross primary production, evapotranspiration, and plant responses to soil moisture and drought stress (Schaphoff et al., 2018b). The terrestrial nitrogen cycle is incorporated to account for nutrient limitations to plant growth (von Bloh et al., 2018). This makes that LPJmL-5 contains all necessary processes to study vegetation responses to drought, though its performance may still vary across regions and vegetation types.

2.2.1 Drought in the LPJmL-5 model

90 In LPJmL-5, GPP is first calculated as a potential GPP, based on the light-limited (JE) and Rubisco-limited (JC) rates and on the absorbed photosynthetically active radiation (APAR; Schaphoff et al. (2018b)). This potential GPP is then constrained by soil water availability. When the water supply cannot meet atmospheric demand, water stress develops and the potential GPP is reduced to the actual GPP.

95 Drought can further influence GPP through phenology. LPJmL-5 includes a phenology function that controls how active the canopy is, ranging from 0 (no active canopy) to 1 (fully active canopy). When phenology declines, leaf area and FAPAR decrease, thereby reducing GPP. The phenology is determined by four limiting functions: cold temperature, light availability, water availability, and heat stress (Schaphoff et al., 2018b). Each factor varies between 0 (full limitation) and 1 (no limitation), and their combined effect defines the total phenology. The limiting functions follow logistic curves and include a memory term, meaning that canopy responses are delayed and differ across PFTs (Schaphoff et al., 2018b). The associated parameters were calibrated by fitting simulated FAPAR to satellite-derived FAPAR over 30 years (Schaphoff et al., 2018b).

100 Vegetation in LPJmL-5 is represented using the the big-leaf representation, meaning that all leaves of a PFT within a grid cell are treated as a single canopy layer with an average temperature, photosynthesis, and water use. Under drought, reductions in phenology and water supply therefore shrink this “big leaf” and makes it less productive, rather than simulating individual trees.

105 Finally, LPJmL-5 does not simulate explicit drought-induced mortality. Instead, mortality emerges indirectly from low productivity and heat stress, and is capped by a maximum annual mortality rate (Schaphoff et al., 2018b). As a result, prolonged water deficits only lead to die-off when plants repeatedly fail to maintain sufficient biomass. Moreover, this mortality process is only active when land cover is simulated dynamically and is not applied when land cover is prescribed.

2.2.2 LPJmL-5 input and output

In this study, LPJmL-5 was used to simulate vegetation dynamics at a 0.1° spatial resolution (~ 11 km) and a daily temporal resolution. The model was forced with meteorological forcing, soil properties, land use, and land cover derived from observational datasets (Table 1). Daily meteorological forcings are taken from the W5E5 dataset and include temperature, precipitation, downward longwave and shortwave radiation, and wind speed (Lange et al., 2022). Soil inputs include static variables such as soil pH (Volkholz and Müller, 2020) and soil type (Knoben, 2021), as well as dynamic inputs like monthly nitrogen deposition (NH_4 and NO_3) (Yang and Tian, 2023). Annual atmospheric CO_2 concentrations were taken from Büchner and Reyer (2022). Land use and country code data are provided at a 5-minute (~ 9.25 km) spatial resolution, and soil type data at 250 m. All three were regridded to a 0.1° spatial resolution, whereas soil pH and climate forcing data are downscaled using a nearest neighbour interpolation from their original 0.5° (~ 55 km) resolution.

The land use map was primarily derived from International Food Policy Research Institute (IFPRI) (2024), which provides information on crop areas. However, the crop types in this dataset did not directly match the CFTs used in LPJmL-5, so they were reclassified accordingly. Additionally, pasture areas are a land-use type in LPJmL-5 but were not included in the IFPRI dataset. These were obtained from the ESA land cover maps (Harper et al., 2023) and incorporated into the land-use input. For the land cover input, we used the ESA PFT maps (Harper et al., 2023), which provide fractional cover of tree, shrub, and grass types. Because LPJmL-5 separates PFTs by climate zone (tropical, temperate, and boreal), we reclassified the ESA PFT maps to match the PFTs in LPJmL-5 using the Köppen-Geiger classification (Beck et al., 2023), following the approach of Forkel et al. (2019)(see [Section S5](#) [Supplements S6](#)). Shrubs were merged with the corresponding tree PFTs because LPJmL-5 does not distinguish between these growth forms. This procedure ensured that land-use and land-cover inputs were compatible with the vegetation types represented in LPJmL-5.

LPJmL-5 provides a multitude of vegetation-related outputs on ecological, hydrological, and agricultural components (such as leaf area index, soil moisture, runoff, and crop yields). To study vegetation response to drought, this study used the monthly Gross Primary Production (GPP), which is the amount of carbon captured from the atmosphere through vegetation photosynthesis (Beer et al., 2010). The GPP output used in this study was produced at a 0.1° spatial and monthly temporal resolution.

2.3 Model validation

~~Before using the simulated GPP to analyse the response of vegetation to MYDs, we first~~

2.3.1 MODIS GPP as validation dataset

In this study, we evaluated simulated GPP by LPJmL5 against MODIS satellite derived GPP estimates. Although MODIS GPP is widely used, it is not purely observational, but derived using the MOD17 light-use efficiency algorithm and relies on inputs such as MODIS FPAR, meteorological drivers, and land-cover classification (Running and Zhao, 2021). Therefore, before using it as a validation dataset, we assessed how well the ~~model reproduces observed GPP dynamics in general. To do~~

~~this, we compared modeled LPJmL-5 GPP to satellite-based MODIS GPP (see Table 1). The MODIS observations are MODIS GPP product represents vegetation productivity (Supplements S3).~~

140 ~~We compared MODIS GPP with two satellite products that are more directly observation-based: the Enhanced Vegetation Index (EVI; Didan (2015a, b)) and solar-induced fluorescence (SIF; Wang and Zhang (2023)). Both comparisons showed strong and statistically significant relationships, particularly outside tropical regions, indicating that MODIS GPP captures large-scale variability in vegetation productivity. Other validation studies report similar limitations, with reduced performance mainly in tropical ecosystems. A full description of these analyses is provided in Supplements S3. Together, these results support the use~~
145 ~~of MODIS GPP as a validation product for LPJmL5, while acknowledging that uncertainties are larger in tropical regions.~~

~~To ensure consistency with LPJmL5 output, the MODIS GPP data was~~ regridded to a spatial resolution of 0.1° and a monthly temporal resolution.

~~Since-Because~~ LPJmL-5 provides a single GPP value per grid cell, each cell was assigned to its dominant vegetation type. ~~Cropland fractions (CFTs)-CFT fractions~~ were aggregated into one cropland category, while natural PFTs were grouped into
150 broader classes: tropical trees, temperate trees, boreal trees, tropical C4 grass, temperate C3 grass, and polar C3 grass (Appendix A).

2.3.2 Kling-Gupta Efficiency

To quantify model performance, we used the Kling-Gupta Efficiency (Gupta et al., 2009), which combines correlation, bias, and variability into a single metric:

$$155 \text{ KGE} = 1 - \sqrt{(r - 1)^2 + (\alpha - 1)^2 + (\beta - 1)^2}, \quad (1)$$

where r is the linear correlation between observations and simulations, $\alpha = \frac{\sigma_{sim}}{\sigma_{obs}}$ a measure of the variability error, and $\beta = \frac{\mu_{sim}}{\mu_{obs}}$ a bias term. Here, σ_{obs} and σ_{sim} are the standard deviations of the observations and simulations, respectively, while μ_{sim} and μ_{obs} are the corresponding means. KGE values greater than -0.41 indicate that the model performs better than the mean benchmark, while a value of 1 indicates perfect agreement between simulations and observations (Knoben et al., 2019).

160 This allows us to assess how well LPJmL-5 simulates GPP for different PFTs compared to MODIS, and to identify structural patterns in performance across the globe.

2.4 **Vegetation sensitivity to varying drought timescales**

After evaluating how well LPJmL-5 simulates general patterns of GPP compared to MODIS GPP, we examined how well LPJmL-5 can model vegetation responses to drought. Here, we aim to understand modelled vegetation sensitivity to drought
165 without differentiating between "normal" and "multi-year" droughts. To ensure a fair comparison of vegetation responses across regions and to remove seasonal variation, the GPP data was normalized and standardized using z-score normalization resulting in a standardized GPP anomaly (GPP_{SA}). The reference normalization period was chosen to match the MODIS dataset (2000-2019) to avoid errors resulting from GPP trend differences between modelled and observed data.

Vegetation can respond differently to drought events of different durations. To capture this, we used the extreme-based method developed by Deng et al. (2022). This method examines how different drought durations affect vegetation across various SPEI timescales (ranging from 1 to 24 months). Specifically, we calculated the average GPP_{SA} response for each timescale and identified which one has the strongest negative impact during drought periods. This timescale is referred to as the dominant drought timescale and reflects how quickly vegetation responds to drought stress. Shorter timescales indicate faster, more sensitive responses, while longer timescales suggest a slower response and less drought-sensitive vegetation. For this analysis, a SPEI threshold of -1 was chosen to define atmospheric drought conditions, as suggested by McKee et al. (1993).

2.5 Vegetation response to MYDs

After determining how sensitive vegetation is to drought at different timescales, we further investigated how vegetation in LPJmL-5 responds to MYDs specifically. We compared MODIS observations and LPJmL-5 simulations by analysing global spatial patterns together with temporal dynamics in selected focus regions. Vegetation responses during MYDs were quantified by using the mean GPP_{SA} during MYD events.

2.6 Selection of Focus Regions

While this study primarily examines global drought patterns, it also focuses on selected focus regions to better understand local drought dynamics and how regional differences affect vegetation responses. The focus regions used in this study are: California (CAL), the Rhine-Meuse delta in western Europe (WEU), the Brahmaputra River basin spanning Bangladesh, India, Bhutan, and China (BRA), central Argentina (ARG), the Orange River basin in southern Africa (SA), and the Murray-Darling basin in Australia (AUS) (Appendix B). These regions correspond to the focus regions used by Ruijsch et al. (2025) examining MYD patterns. The focus regions were selected based on the number of MYDs that have occurred between 2000-2019, geographic and climatic diversity, and variation in vegetation types, ensuring a broad representation of both water- and energy-limited regions across different climate zones.

The following sections describe different aspects of the analysis on the potential of LPJmL-5 to simulate vegetation response to (multi-year) droughts. Section 3.1 examines LPJmL's ability to reproduce ~~observed~~ satellite-derived GPP, while Sections 3.2 and 3.3 focus on its ability to reproduce (multi-year) droughts.

3.1 General performance of LPJmL-5

195 To evaluate LPJmL-5's ability to simulate GPP, we compared modelled GPP to MODIS-derived GPP using two simulation setups: dynamic and prescribed land cover. The dynamic land cover setup allows vegetation fractions to shift in response to climate and includes background mortality, capturing potential vegetation loss (Schaphoff et al., 2018b). However, dynamic vegetation does not necessarily match observed vegetation types, which can lead to differences in GPP due to mismatched vegetation types. The prescribed land cover simulation maintains fixed vegetation fractions based on observed maps. This
200 approach ensures that variations in GPP result from the model's ability to simulate vegetation productivity. However, this setup does not include background mortality, potentially causing an underestimation of vegetation loss. By comparing both setups, we can determine which one provides a better representation of ~~observed~~ satellite-derived GPP and choose the setup to be used in later analyses. Model performance is evaluated by comparing LPJmL-5 and MODIS GPP using the KGE metric (Section 3.1.1) and time series analysis (Section 3.1.2).

205 3.1.1 Evaluating Model Performance with the KGE metric

Figure 2 shows the spatial distribution of the KGE and its components (correlation, bias, and variability) for both dynamic and prescribed land cover simulations. Overall, the prescribed land cover demonstrates a clear performance improvement over the dynamic simulation (Figures 2a and b).

The dynamic land cover consistently shows low agreement with MODIS across all KGE components (Figures 2c, e and
210 g). The agreement is particularly poor in the Southern Hemisphere, with KGE values often falling below -0.41, which is the threshold for a skilful performance. In contrast, the Northern Hemisphere generally shows a better performance. Switching to prescribed land cover results in improved KGE scores globally. However, the performance difference between the Northern and Southern hemispheres remains.

When examining the individual KGE components, the correlation between modelled and ~~observed~~ satellite-derived GPP
215 is relatively high for both simulations, except in the tropical and semi-arid regions (Figures 2c and d). This suggests that ~~the~~ LPJmL-5 captures intra-annual response to changing meteorological conditions is similar between ~~the model and observations~~ LPJmL-5 and satellite-derived MODIS GPP.

Both the bias (Figure 2e) and variability (Figure 2g) components are overestimated in the dynamic land cover simulation. The bias frequently exceeds 2, except in the tropical regions, suggesting that the model's mean GPP is approximately double
220 that of the ~~observed~~ satellite-derived value. The prescribed land cover simulation (Figure 2f) significantly reduces this bias, particularly in tropical and temperate regions, bringing modelled GPP closer to ~~observations~~ the satellite-derived reference. A

similar trend is observed for the GPP variability, where the dynamic land cover shows overestimated variability (greater than 2) in Figure 2g, while prescribed land cover reduces this variability (Figure 2h).

225 These differences in model performance can be attributed to discrepancies between the modeled and observed vegetation types when using dynamic land cover. For example, we observe that LPJmL-5 simulates tree cover in regions where the ESA land cover map shows predominantly grasslands, such as Western Europe and parts of South Africa (see Appendix C and Supplements [S4S5](#)). These discrepancies in vegetation types lead to differences in GPP that are not related to the model's capability to simulate GPP responses to changing meteorological conditions, but rather to inconsistencies in vegetation cover. By using prescribed land cover fractions based on the ESA land cover map, we observe an improvement in model performance, 230 with remaining errors linked to LPJmL-5's inability to accurately simulate the GPP response to changes in meteorological conditions.




Figure 2. Spatial comparison of LPJmL-5 model performance using the Kling-Gupta Efficiency (KGE) metric and its components for dynamic (left column) and prescribed (right column) land cover simulations. The rows show (a–b) overall KGE, followed by its components: (c–d) correlation, (e–f) bias, and (g–h) variability.

To further assess model performance in the prescribed land cover simulation, we analyse the KGE distribution for each PFT, as shown in Figure 3. This cumulative distribution function (CDF) displays the proportion of grid cells for each dominant vegetation type (Appendix A) that exceed the KGE threshold of -0.41 , which represents the mean benchmark (Knoben et al., 235 2019).

The performance of LPJmL-5 varies significantly across PFTs. Croplands show the lowest agreement with ~~observations~~satellite-derived GPP, with over 60% of grid cells falling below the $KGE = -0.41$ line. In contrast, temperate trees show the best agreement ~~with observed-GPP~~, with 90% showing skilful GPP simulations. Approximately 80% of areas dominated by boreal trees show good agreement with MODIS data. Polar and temperate C3 grasses are the dominant PFTs and exhibit intermediate performance, 240 with around 75% of their area showing significant skill. Tropical trees show moderate KGE values, indicating a moderate performance in simulating the GPP. The corresponding analysis for the dynamic land-cover simulation (Figure C2) shows consistently lower agreement across most PFTs than for prescribed land cover.

Overall, the KGE map and the CDF analysis (Figures 2, 3, and C2) indicate that simulating with prescribed land cover outperforms dynamic land cover in LPJmL-5. However, significant improvements in modelled GPP response can still be made 245 in the Southern Hemisphere and for croplands. Based on these results, all subsequent simulations will be done using prescribed land cover to ensure a more robust comparison with observational data.

3.1.2 Evaluating Model Performance with the GPP Time Series

Figure 4 shows the monthly GPP for LPJmL-5 (with prescribed land cover) and MODIS ~~satellite-observations~~ across the six focus regions between 2000 and 2019. The background shading represents the SPEI-12, highlighting dry periods in red and 250 wet periods in blue. Additionally, the figure includes the correlation between MODIS and LPJmL-5.

Three general patterns can be identified in the timeseries: (1) A temporal shift in the LPJmL-5 GPP time series, with peaks occurring earlier in the season compared to MODIS ~~observations~~; (2) Overestimation of summer GPP peaks, and in some cases underestimation of winter values; (3) Lower model performance during dry periods.

These patterns are reflected to varying degrees across the focus regions. CAL and BRA show the temporal shift, which may 255 be the cause of the lower correlation with MODIS. BRA and WEU exhibit overestimation of summer GPP peaks, with SA also showing the underestimation of winter values. ARG performs relatively well, although it also shows slightly increased summer peaks. In AUS, the contrast between wet and dry periods is largest, with GPP simulated accurately during wet periods, while the model does not capture dry periods well. Despite these issues, WEU, ARG, SA, and AUS show relatively high correlation values, indicating that LPJmL-5 generally captures the seasonal and interannual patterns of satellite-derived GPP 260 in these regions. Overall, based on the KGE values and the time series analysis, we can conclude that LPJmL-5 simulates GPP reasonably well, although notable regional differences remain.




Figure 3. Cumulative distribution function (CDF) of the KGE of the prescribed land cover GPP for the different vegetation types in LPJmL-5. Percentages show the percentage of grid cells with that dominant land cover. Black dashed line indicates KGE = -0.41. Values greater than -0.41 indicate that the model improved upon the mean benchmark.




Figure 4. Timeseries of the GPP for LPJmL-5 with prescribed land cover (green) and MODIS (black) for the focus regions: California (CAL), western Europe (WEU), Brahmaputra (BRA), middle Argentina (ARG), southern Africa (SA) and the Murray Darling Basin in Australia (AUS). SPEI-12 is shaded in the background, where red areas indicate drier periods and blue areas wetter periods. Additionally, the figure includes the correlation between MODIS and LPJmL-5 per region. Note that WEU uses a different y-axis range from the other regions.

3.2 General drought response

To evaluate the ability of LPJmL-5 to capture vegetation drought response, we applied the extreme-based method (Section 2.4).

265 This method identifies the dominant drought timescale, defined as the SPEI timescale (ranging from 1 to 24 months) that has the most negative impact on vegetation during drought periods.

The spatial distribution of dominant drought timescales, shown in Figure 5, shows a clear difference between MODIS ~~observations~~ (a) and the LPJmL-5 ~~model~~ simulations (b). Compared to MODIS, LPJmL-5 ~~shows shorter drought exhibits shorter dominant~~ timescales, indicating a faster ~~vegetation-GPP~~ response to drought. This difference is most pronounced in 270 the Southern Hemisphere. LPJmL-5 also shows a skewed distribution with a clear peak around 2 months (panel c), ~~while whereas~~ the MODIS response is more evenly distributed. ~~In summary, Overall, these patterns highlight differences in temporal sensitivity between~~ LPJmL-5 ~~tends to underestimate drought resistance by simulating and the satellite-derived reference, with LPJmL-5 showing~~ shorter response times ~~than observed in MODIS, especially in the tropical regions and the Southern Hemisphere~~ during drought periods.




Figure 5. Spatial patterns of the dominant drought timescale of GPP_{SA} for a) [the MODIS observations-reference](#) and b) the LPJmL-5 model. The dominant timescale is defined as the shortest SPEI timescale (between 1 and 24 months) that shows the most negative GPP_{SA} during drought periods (with a SPEI threshold of -1). Bare areas and sparse vegetation are filtered out. c) Violin plot comparing the distribution of dominant drought timescales between MODIS and LPJmL-5.

275 3.3 MYD and ND response

After evaluating LPJmL-5's response to droughts in general, this study now shifts its focus to MYDs and NDs.

3.3.1 Temporal patterns

280 Firstly, we examine how well LPJmL-5 captures the timing and magnitude of GPP_{SA} during MYDs and NDs across the focus regions. Figure 6 shows the timeseries of GPP_{SA} for [MODIS-the satellite-derived reference \(MODIS\)](#) and LPJmL-5, with MYDs and NDs indicated by red and orange shaded areas, respectively.

Overall, LPJmL-5 captures the timing and magnitude of GPP_{SA} anomalies reasonably well, showing relatively high correlations with MODIS in most regions. The model aligns particularly well in CAL, WEU, ARG, and AUS, while performance

is somewhat lower in SA and BRA. Correlations during MYDs are higher than those for the entire time series in all regions except BRA, and generally higher than those during NDs, except in ARG and BRA. This suggests that LPJmL-5 simulates MYD dynamics more accurately than ND dynamics.

Nonetheless, some systematic differences remain during MYDs: (1) the model tends to simulate a too fast and severe decline in GPP_{SA} at the onset of MYDs, and (2) it exhibits ~~unrealistic-strong~~ variability during MYD periods, especially in the form of abrupt recovery spikes ~~not supported by observations that are not seen in the satellite-derived reference~~.

These biases are reflected across the focus regions to varying degrees. In WEU, the model simulates an early GPP_{SA} drop that precedes the ~~observed-reference~~ onset. CAL shows a similar offset, likely linked to the timing mismatch seen in Figure 4. In BRA, the first MYD minimum occurs too early in the model, while the second MYD ends with a sudden increase in GPP_{SA} not seen in ~~observations, again pointing to the reference, again indicating~~ an early response and ~~unrealistic-rapid~~ recovery. ARG captures drought severity relatively well but still shows some ~~unrealistic-abrupt~~ recovery spikes during MYDs. SA also shows the rapid onset and overestimated recovery rate, despite persistently low SPEI-12 values. AUS shows the best overall agreement, with most MYDs being captured well. However, the second MYD still shows the ~~unrealistic-strong~~ variability mid-drought, though less severely.

Overall, while LPJmL-5 tends to ~~overestimate-vegetation response at the onset of MYDs and shows some rapid recovery behaviour~~ simulate an overall more dynamic response with faster declines and rapid recovery during MYDs, it successfully ~~captures-reproduces~~ the key temporal dynamics of MYDs relative to the satellite-derived reference.





Figure 6. Timeseries of GPP_{SA} for LPJmL-5 and MODIS. MYDs and NDs are shaded in red and orange, respectively. Correlation coefficients between **model-LPJmL-5** and **observations-MODIS** are shown for the full time series, MYDs, and NDs. The grey line indicates the SPEI-12. The red and blue dashed lines mark -1 and +1, respectively.

3.3.2 Spatial patterns

In addition to the regional time series analysis, we now examine global spatial patterns of vegetation response to MYDs. Figure 7 shows the spatial distribution of the mean GPP_{SA} during MYDs, as derived from (a) ~~MODIS observations~~ the [MODIS reference](#) and (b) the LPJmL-5 model. Negative values (red) indicate below-average GPP during MYD periods, while positive values (green) indicate GPP being higher than average.

305

a) Reference (MODIS)



b) Model (LPJmL-5)

Figure 7. Spatial pattern of the mean GPP_{SA} during MYDs for a) MODIS and b) LPJmL-5. Positive values indicate higher GPP than normal during MYDs, while negative values indicate reduced GPP values. Bare areas and sparse vegetation are filtered out. c) shows a violin plot comparing the distributions of the mean GPP_{SA} during MYDs from MODIS and LPJmL-5.

Overall, LPJmL-5 simulates less severe GPP_{SA} during MYDs compared to MODIS ~~observations~~. However, the Spearman rank correlation (0.326, $p < 0.001$) still indicates a similarity in the overall spatial patterns between the model and the ~~observations~~ reference. This means that the model captures some important hotspot areas, but there are still noticeable differences. We attribute these differences largely to ~~unrealistic~~ the abrupt recovery events in the model, as seen in Figure 6. The violin plot (Figure 7c) shows that MODIS has a slightly lower median. However, LPJmL-5 displays more extreme values at both ends of the distribution, indicating greater variability in the modelled MYD responses.

To further highlight these differences, Figure 8 shows the spatial difference in mean GPP_{SA} during MYDs between MODIS and LPJmL-5 and two-dimensional Kernel Density Estimate (KDE) plots for the different PFTs. Purple shades indicate underestimation of negative drought impacts, while orange shades indicate underestimation of positive impacts or overestimation of negative impacts. Black, grey and white areas indicate close agreement between ~~model and observations~~ the model and the reference.

LPJmL-5 shows a reasonable agreement with MODIS in most regions, as indicated by the black, grey and white areas. These areas include Central North America, Europe, and parts of East and West Asia. However, there are several regions that emerge as hotspots of LPJmL-5 underestimation of GPP anomalies (visible as dark purple areas), including Australia, Northwest India, Pakistan, Southern Africa, Central Argentina, Eastern Brazil, and Northern Mexico. In these regions, LPJmL-5 shows weaker or absent negative responses, suggesting that the model does not fully capture drought severity in these areas. This finding is consistent with the temporal patterns shown in Figure 4. Additionally, LPJmL-5 tends to underestimate the ~~observed satellite-derived~~ positive GPP responses in the arctic boreal zone (visible as orange areas). Here, the model ~~fails to capture the increased GPP during MYDs found in observations~~ does not reproduce the positive GPP anomalies during MYDs seen in the MODIS reference data (Figure 7 and Ruijsch et al. (2025)).

Beyond spatial patterns, differences between LPJmL-5 and MODIS also vary by PFT (Figure 8). Tropical trees show the widest spread in MYD GPP responses, while temperate trees align more closely with MODIS but still occasionally underestimate negative impacts (lower right quadrant). Grasslands show good agreement, with LPJmL-5 capturing most directions of the MYD response correctly. Croplands, although displaying the weakest agreement in absolute GPP (Figure 3), show the best modelled drought response. This indicates that while simulating total productivity remains challenging, LPJmL-5 can capture how croplands respond to drought events.

Overall, LPJmL-5 ~~successfully captures the key dynamics of MYDs observed in MODIS~~ reproduces several key temporal and spatial MYD patterns present in the MODIS reference, but still shows some model bias in both time and space. Temporally, LPJmL-5 often ~~overestimates the speed and severity of vegetation decline~~ simulates faster and stronger vegetation declines at the onset of the MYD than the satellite-derived reference and shows a fast recovery during the drought period ~~compared to MODIS~~. Spatially, LPJmL-5 shows a reasonable agreement with MODIS in most regions and for most PFTs, but also shows some underestimation of both strong negative and strong positive vegetation responses. These biases may be caused by short-term overreactions and rapid recoveries in the model's time series, which reduces the average drought response.




Figure 8. Spatial differences in mean GPP_{SA} during MYDs between MODIS observations and LPJmL-5 simulations. The map highlights regions where LPJmL-5 under-shows weaker or overestimates stronger vegetation responses compared relative to observations the satellite-derived reference. Purple shades indicate underestimated negative impacts, while orange shades indicate underestimated positive or overestimated negative impacts. White, grey and black shades represent regions where the model and observations reference agree closely on vegetation response during MYDs. Insets display two-dimensional kernel density estimates (KDEs) for seven vegetation types, illustrating the distribution of differences in mean MYD responses. KDE levels are set to [0.05, 0.15, 0.25, 0.5, 0.75, 0.9, 1.0].

4 Discussion

340 Overall, this study assesses LPJmL-5's ability to simulate global vegetation responses to (multi-year) droughts. First, comparing dynamic and prescribed land cover settings in LPJmL-5 revealed that prescribed land cover improves agreement with MODIS GPP. The lowest agreement is generally found in the Southern Hemisphere and in cropland-dominated regions. Second, LPJmL-5 and MODIS ~~show GPP exhibit~~ similar broad seasonal and interannual ~~patterns of GPP. There is however a tendency to overestimate variability, although the model tends to simulate stronger~~ GPP dynamics and ~~to introduce temporal~~
345 ~~shifts compared to observations~~ introduces temporal shifts relative to the satellite-derived reference. Third, using the extreme-based method ~~;~~ ~~which identifies to identify~~ the timescale at which drought has the strongest impact on vegetation, we found that LPJmL-5 ~~underestimates drought resistance by simulating~~ simulates shorter dominant drought timescales than MODIS, particularly in tropical regions and the Southern Hemisphere, indicating a faster modelled GPP response to drought conditions. Fourth, the analysis of MYDs revealed a consistent pattern of model bias across both time and space. Temporally, LPJmL-
350 5 overestimates vegetation decline simulates stronger and more rapid GPP declines at the onset of MYDs and shows ~~a fast recovery during the drought period~~ faster recovery during drought periods compared to the reference. Spatially, the model ~~underestimates~~ shows muted representations of both strong negative and strong positive ~~vegetation responses. In general~~ GPP anomalies compared to MODIS.

Overall, LPJmL-5 tends to simulate vegetation that is too responsive in both the drought onset and the recovery. This points
355 to simulates faster GPP declines and recoveries during droughts than the satellite-derived reference, suggesting limitations in how the model represents vegetation resistance and resilience to water induced stress and mortality.

4.1 Uncertainties in the model choices and setup

Different choices in model parametrisation and setup can lead to significant differences in modelled GPP. For example, both the choice of meteorological forcing dataset (ERA5 (Hersbach et al., 2023) vs. W5E5 (Lange et al., 2022)) and the temporal
360 resolution (monthly vs. daily) have an significant impact on simulated GPP patterns (Supplements S3 S4). This indicates that it is important for future drought oriented studies to make clear choices with regard to the modelling setup. In this study we selected the daily W5E5 dataset (Lange et al., 2022), as it is bias-corrected and offers a high temporal resolution and the total global GPP was closest to that found in observational datasets.

Our analyses also showed that GPP output is highly sensitive to the radiation inputs (Supplements S3 S4). GPP values varied
365 substantially depending on whether the model calculated radiation internally or used observed radiation data. We found that errors in GPP are higher from internal estimation, compared to directly providing longwave and shortwave radiation from observations.

Using dynamic land cover in LPJmL-5 resulted in systematically overestimated GPP values across most regions (Figure 2 and Supplements S4 S5), also noted in Schaphoff et al. (2018a). We argue that this difference is the results of a mismatch
370 between modelled and observed vegetation distributions (Appendix C and Supplements S4 S5), which is to be expected as the land cover is not constrained to the observed land cover. For example, LPJmL-5 simulates tree cover in parts of Western

Europe where forests could exist naturally, but are absent today due to long-term human management. To better match **observed** satellite-derived GPP and reduce these discrepancies, we prescribed land cover fractions, rather than letting the model simulate land cover dynamically.

375 4.2 Evaluation of GPP simulations and vegetation drought response in LPJmL-5

Prescribing land cover in LPJmL-5 improved the agreement with MODIS GPP, as indicated by higher KGE values in Figure 2. This is consistent with studies showing that accurate land cover representation is essential for reproducing observed productivity patterns (Krause et al., 2022), and that constraining DGVM simulations with satellite-derived observations (including land cover) enhances their ability to capture observed spatial patterns of vegetation and carbon dynamics (Forkel et al., 2019).

380 Breaking down the differences between LPJmL-5 and MODIS GPP during MYDs by vegetation type further highlights varying model performance (Figure 8). Tropical trees show the largest variability in drought response, showing both under- and overestimations of impacts. Similar results were found by Powell et al. (2013), who showed that terrestrial biosphere models underestimate drought impacts on tropical forest biomass. Temperate trees generally align more closely with observations but still underestimate negative drought impacts. This is consistent with Kolus et al. (2019), who found that land carbon models
385 underestimate the severity and duration of drought effects on productivity across temperate forests in the US and Europe. Similarly, Schaefer et al. (2012) observed that models frequently overpredict GPP under dry conditions, further pointing to an underestimation of drought stress. Boreal trees and grasses are more prone to underestimated positive responses, while croplands, despite poorer overall productivity agreement, show relatively better modelled drought responses.

A key limitation underlying these issues appears to be how vegetation mortality is represented in LPJmL-5. In the model,
390 mortality is only driven by low productivity and heat stress, and can not be directly induced by prolonged water deficits. Drought-induced mortality occurs only if plants fail to produce enough biomass to sustain themselves. However, this natural mortality is capped by a maximum annual rate (Schaphoff et al., 2018b), which restricts how much vegetation can be lost due to die-off each year. Moreover, this mortality mechanism is only active in the dynamic land cover setting, and is not applied when land cover is prescribed. Since the simulations in this study use the prescribed land cover setting, no vegetation is lost
395 due to this background mortality. This means that while GPP may decline sharply during droughts, vegetation does not die, causing unrealistically rapid recovery periods when precipitation (temporarily) returns. Although LPJmL-5 includes a heat-stress mortality mechanism, it only applies to boreal trees (Schaphoff et al., 2018b). As a result, vegetation decline during MYDs is not directly driven by water shortages, which prevents large-scale die-off and causes the model to underestimate total vegetation loss. These limitations reduce the model's ability to simulate the full severity of vegetation decline, especially
400 during MYDs. This issue has been noted by Schaphoff et al. (2018a), who reported that the lack of realistic mortality processes leads to an overestimation of simulated biomass and contributes to biases in GPP. Similarly, McDowell et al. (2013) found that models which combine carbon starvation with hydraulic failure mechanisms are better able to reproduce observed drought-induced tree mortality, and Zhou et al. (2013) highlighted the need to account for both stomatal and non-stomatal limitations to realistically capture plant responses to water stress. Meyer et al. (2025) further showed that including hydraulic processes in

405 LPJ-GUESS improves simulations of drought responses. Together, these findings suggest that LPJmL-5 likely underestimates the true impact of drought on vegetation.

Despite underestimating long-term vegetation loss, LPJmL-5 tends to overestimate vegetation decline at the start of droughts, as shown in Figures 5 and 6. LPJmL-5 does include a water stress factor that influence the growth, which results in a rapid decline in GPP. This stress is determined by soil moisture availability and fixed rooting depths for each PFT, without accounting
410 for dynamic rooting strategies, capillary rise or explicit groundwater processes (Schaphoff et al., 2018b). However, in reality, vegetation often withstands short-term stress periods through drought survival strategies (Santiago et al., 2016; Pivovarov et al., 2016). At the same time, LPJmL-5 also simulates unrealistic recoveries during the drought period. Since there is no mortality, vegetation can survive during a drought period, resulting in no loss of biomass. When conditions improve temporarily, for example due to brief rainfall, the model quickly simulates regrowth. This leads to an unrealistic drought recovery effect. This
415 is in contrast with real vegetation, which often suffers from long-term impacts (Wu et al., 2018). Both the quick decline and recovery are consistent with findings by Kolus et al. (2019), who show that land carbon models underestimate the severity and duration of drought impact on plant productivity.

Together, the combination of overly sensitive short-term GPP decline, limited representation of long-term vegetation loss, and unrealistically fast recovery during drought periods results in a mismatch in LPJmL-5's simulated vegetation drought
420 response. While prescribing land cover improves the overall GPP performance, these structural issues in the mortality representation limit the model's ability to simulate realistic vegetation dynamics during drought events. This is especially limiting when the analysis is focussed on multi-year drought events, which have been shown to cause higher vegetation mortality rates (Češljar et al., 2025) and are expected to become more frequent under climate change (Wiel et al., 2022).

4.3 Uncertainties in methodological choices

425 To improve the overall quality and reliability of our simulations, we used the state-of-the-art dynamic vegetation model LPJmL-5. This model is publicly available with an extensive model description and evaluation (Schaphoff et al., 2018b, a; von Bloh et al., 2018). LPJmL-5 provides a multitude of vegetation-related outputs on ecological, hydrological, and agricultural components, making it suitable for studying vegetation responses at the global scale.

Drought events were identified using the Standardized Precipitation Evapotranspiration Index (SPEI), which incorporates
430 both precipitation (P) and potential evapotranspiration (PET), allowing it to better reflect the impact of temperature on drought severity (Vicente-Serrano et al., 2010). As a standardized index, SPEI also enables consistent spatial comparisons of drought conditions at the global scale. By using a 12 month aggregation period for the SPEI, we also remove seasonal variations and focus on capturing MYDs.

As a standardized metric, SPEI characterizes drought conditions relative to the local climatic baseline, reflecting periods that
435 are characterized by below normal water availability. In energy-limited regions, such as the tropics or boreal forests, negative SPEI values may occur even when absolute water availability remains relatively high (Zang et al., 2019). To ensure these droughts correspond to meaningful reductions in water availability, we compared soil water content (SWC) during droughts

and normal conditions (Supplements S2), finding consistently lower SWC across soil layers. This supports the interpretation that identified MYDs correspond to periods of meaningful vegetation stress, even in energy-limited regions.

440 While there is currently no universally agreed-upon definition of MYDs, we follow the approach of Wiel et al. (2022) (see Section 2.1), which was also adopted in Ruijsch et al. (2025) and van Mourik et al. (2025). SPEI values are sensitive to the method and input data used for calculating PET. We compared results using ERA5 and W5E5 and found differences in the number of MYDs identified for the period 2000–2019 (Supplements S3 S4). Nonetheless, the MYDs identified in our focus regions are broadly consistent with those reported in other studies (van Mourik et al., 2025; Wiel et al., 2022; Luo et al., 2017; 445 Liu et al., 2022; Chikoore and Jury, 2021; Naumann et al., 2023; van Dijk et al., 2013), despite minor differences in timing.

To evaluate vegetation responses, we used ~~the monthly monthly MODIS~~ GPP as a proxy ~~-, which quantifies for ecosystem productivity, representing~~ the amount of carbon captured from the atmosphere through photosynthesis (Beer et al., 2010). ~~Although the Enhanced Vegetation Index (EVI) has been used in previous studies Previous drought studies have often relied on the EVI~~ (Ruijsch et al., 2025; Huang and Xia, 2019; Yang et al., 2024), ~~it is derived from satellite observations but EVI is satellite-derived~~ and not directly available ~~as a model output. GPP has been shown to correlate well with EVI (Rahman et al. (2005); Sims et al. (2006) and Supplements S2), even during MYDs (Ruijsch et al., 2025). from LPJmL-5. Because MODIS GPP and EVI are strongly correlated (Rahman et al., 2005; Sims et al., 2006; see also Supplements S3), including during multi-year droughts (Ruijsch et al., 2025), MODIS GPP can be considered a suitable proxy for vegetation activity.~~ 450

Several studies further indicate that MODIS GPP provides a meaningful representation of productivity across biomes, with most errors linked to input datasets rather than the algorithm itself. Performance is generally good in temperate and mixed forests, while errors are larger in evergreen broadleaf forests and tropical regions (Tang et al., 2015). Wang et al. (2017) showed that much of the error originates from FPAR, land-cover misclassification, and light-use efficiency parameters rather than meteorology. Systematic underestimation has also been reported in grasslands (Zhu et al., 2018). These studies agree that cloud contamination and uncertainties in tropical regions remain key challenges. Taken together, the strong correspondence 455 between MODIS GPP, EVI, and solar-induced fluorescence (Supplement S3) supports the use of MODIS GPP as our validation dataset for LPJmL-5 GPP, while acknowledging that performance is weaker in tropical ecosystems and that results should be interpreted with these limitations in mind.

Although other sources of GPP data exist, such as site-level flux tower measurements, we choose to use satellite derived MODIS GPP because it provides consistent, observation-based estimates at the global scale. To maintain observational consistency and avoid mixing in other models, we did not include GPP estimates derived from other land surface models as they are 460 likely based on similar assumptions and equations as the LPJmL-5 model. Nonetheless, MODIS GPP is subject to uncertainty, particularly in tropical regions (Supplements S3), and this should be kept in mind when interpreting the results.

5 Conclusion

This study evaluated LPJmL-5's ability to simulate global vegetation responses to drought, using MODIS GPP as a benchmark. Prescribing land cover (instead of simulating it dynamically) improves the model's ability to reproduce global GPP dynamics. Overall, LPJmL-5 simulates GPP reasonably well, but improvements can still be made in the Southern Hemisphere and for croplands.

Our analysis shows that LPJmL-5 simulates an accelerated vegetation response to drought onset, especially in tropical regions and the Southern Hemisphere. During MYDs, LPJmL-5 captures the key temporal and spatial dynamics ~~observed in~~ found with MODIS. However, the model tends to overestimate vegetation response at the onset of MYDs and shows some rapid recovery behaviour. This leads to a muted overall vegetation drought response and spatial underrepresentation of both strong negative and strong positive GPP anomalies. Among PFTs, croplands show the best agreement with ~~observed~~ satellite-derived MYD responses. In contrast, boreal vegetation shows underestimated positive drought responses and temperate vegetation underestimated negative ones. Tropical vegetation displays more mixed results, with both over- and underestimation.

These differences between ~~model and observations~~ modelled and satellite-derived GPP may be partly attributed to how LPJmL-5 represents vegetation stress and drought induced mortality. The model responds strongly to short-term productivity losses but limits long-term vegetation decline, resulting in rapid recovery and limited vegetation die-off during (multi-year) drought periods.

In general, LPJmL-5 is able to reproduce the general impacts of MYDs on vegetation but tends to simulate vegetation that is too responsive in both the drought onset and the recovery. This points to limitations in how the model represents vegetation resistance and resilience to water induced stress and mortality. Given the increasing frequency and intensity of MYDs, accurately representing vegetation stress and mortality is critical for reliably assessing MYD impacts on vegetation.

Appendix A: Dominant land cover types

490 Figure A1 shows the dominant land cover types at a 0.1° spatial resolution, based on LPJmL-5's land cover and land use dataset (see Table 1). Cropland fractions were combined into a single "cropland" category, while natural PFTs were grouped into broader vegetation classes: tropical trees (tropical broadleaved evergreen and raingreen trees), temperate trees (needleleaved evergreen, broadleaved evergreen, and broadleaved summergreen trees), boreal trees (needleleaved evergreen, broadleaved summergreen, and needleleaved summergreen trees), tropical C4 grass, temperate C3 grass, and polar C3 grass.




Figure A1. Dominant land cover types at a 0.1° resolution for 2019. Vegetation categories include tropical trees (TTr), temperate trees (TTe), boreal trees (TBo), cropland (Cr), tropical C4 grass (GTr), temperate C3 grass (GTe), and polar C3 grass (GBo).

Appendix B: Number of MYDs and chosen focus regions

495 Figure B1 shows the number of MYDs (2000-2019) at a 0.5° spatial resolution from W5E5 data (Lange et al., 2022)(See Section 2.1 and [Supplements S1](#)). Black outlines indicate the chosen focus regions: California (CAL), the Rhine-Meuse delta in western Europe (WEU), the Brahmaputra River basin in Bangladesh/India/Bhutan/China (BRA), central Argentina (ARG), the Orange River basin in southern Africa (SA), and the Murray-Darling basin in Australia (AUS) (See Section 2.6).




Figure B1. Number of MYDs between 2000–2019 at 0.5° spatial resolution. Bare areas and regions with sparse vegetation are excluded. Black contours indicate the six chosen focus regions: CAL, WEU, BRA, ARG, SA, and AUS.

Appendix C: LPJmL dynamic vs ESA prescribed land cover

500 Figure C1 shows the difference in vegetation cover fractions between LPJmL-5 and ESA. Globally, the ESA dataset shows greater coverage of temperate and polar C3 grasses, as well as boreal needleleaved summergreen forests. In contrast, LPJmL-5's dynamic vegetation simulates more temperate needleleaved evergreen trees, boreal needleleaved evergreen trees, boreal broadleaved summergreen trees, and tropical C4 grasses.




Figure C1. Difference in vegetation cover fractions between LPJmL-5 and ESA. Blue areas indicate where ESA has a higher vegetation fraction; red areas indicate where LPJmL-5 simulates a higher vegetation fraction.

The CDFs in Figure C2 further show the impact of land cover on model performance. Across all vegetation types, the prescribed land cover shows higher KGE values. The difference between prescribed and dynamic land cover is particularly

505

510 pronounced for temperate C3 grasses, which perform poorly with the the dynamic land cover. LPJmL-5 substantially under-represents the spatial extent of temperate C3 grasses, assigning them to only 2.7% of grid cells, compared to 22.2% under the ESA-based classification. This mismatch likely contributes to the poor performance, as misclassified vegetation types lead to unrealistic GPP estimates. Tree-dominated grid cells also show better performance with prescribed land cover, however, the difference is less severe than for grasses.




Figure C2. Cumulative distribution function (CDF) of the KGE of the a) prescribed land cover GPP and b) dynamic land cover for the different vegetation types in LPJmL-5. Percentages show the percentage of grid cells with that land cover. Black dashed line indicates KGE = -0.41. Values greater than -0.41 indicate that the model improved upon the mean benchmark.

Code and data availability. The LPJmL-5 source code is openly available at <https://github.com/PIK-LPJmL/LPJmL>. This study uses version v5.7.9 (released on May 2 2024).

All processed data required to reproduce the analyses are available at: <https://doi.org/10.24416/UU01-U97GDE> (Ruijsch and Wanders, 2026).

515 The raw datasets are publicly available from their respective sources (see Table 1).

All scripts used for postprocessing and analysis in this study are available at <https://doi.org/10.5281/zenodo.17085725> (Ruijsch, 2025).

Author contributions. **DR:** Conceptualisation, methodology, software, validation, formal analysis, data curation, writing–original draft, visualisation. **SH:** Conceptualisation, formal analysis, writing–review and editing. **HB:** Conceptualisation, formal analysis, writing–review and editing. **NW:** Conceptualisation, formal analysis, writing–review and editing, project administration, funding acquisition.

520 *Competing interests.* The authors declare that they have no conflict of interests.

Acknowledgements. Funded/Co-funded by the European Union (ERC, MultiDry, Grant Agreement number: 101075354). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them. DR and NW are supported by the ERC STG MultiDry project. SH is supported by NWO VIDI project Recover VI.Vidi.223.009. HB is supported by the European Union (ERC-CoG,

525 3POLE2SEA, 101126168).

References

- Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., Rigling, A., Breshears, D. D., Hogg, E. T., Gonzalez, P., Fensham, R., Zhang, Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S. W., Semerci, A., and Cobb, N.: A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests, *Forest Ecology and Management*, 259, 660–684, <https://doi.org/https://doi.org/10.1016/j.foreco.2009.09.001>, adaptation of Forests and Forest Management to Changing Climate, 2010.
- 530 Beck, H., McVicar, T., Vergopolan, N., Berg, A., Lutsko, N., Dufour, A., Zeng, Z., Jiang, X., van Dijk, A., and Miralles, D.: High-resolution (1 km) Köppen-Geiger maps for 1901–2099 based on constrained CMIP6 projections, *Scientific Data*, 10, <https://doi.org/10.1038/s41597-023-02549-6>, 2023.
- 535 Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rödenbeck, C., Arain, M. A., Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M., Luysaert, S., Margolis, H., Oleson, K. W., Rouspard, O., Veenendaal, E., Viovy, N., Williams, C., Woodward, F. I., and Papale, D.: Terrestrial Gross Carbon Dioxide Uptake: Global Distribution and Covariation with Climate, *Science*, 329, 834–838, <https://doi.org/10.1126/science.1184984>, 2010.
- Büchner, M. and Reyer, C. P.: ISIMIP3a atmospheric composition input data, <https://doi.org/10.48364/ISIMIP.664235.2>, 2022.
- 540 Chikoore, H. and Jury, M.: South African drought, deconstructed, *Weather and Climate Extremes*, 33, 100334, <https://doi.org/10.1016/j.wace.2021.100334>, 2021.
- Choat, B., Brodribb, T. J., Brodersen, C. R., Duursma, R. A., López, R., and Medlyn, B. E.: Triggers of tree mortality under drought, *Nature*, 558, 531–539, <https://doi.org/10.1038/s41586-018-0240-x>, 2018.
- Cooley, H., Donnelly, K., Phurisamban, R., and Subramanian, M.: Impacts of California’s ongoing drought: agriculture, *Pac. Inst.*, 2015.
- 545 Deng, Y., Wu, D., Wang, X., and Xie, Z.: Responding time scales of vegetation production to extreme droughts over China, *Ecological Indicators*, 136, 108630, <https://doi.org/https://doi.org/10.1016/j.ecolind.2022.108630>, 2022.
- DeSoto, L., Cailleret, M., Sterck, F., Jansen, S., Kramer, K., Robert, E. M. R., Aakala, T., Amoroso, M. M., Bigler, C., Camarero, J. J., Čufar, K., Gea-Izquierdo, G., Gillner, S., Haavik, L. J., Hereş, A.-M., Kane, J. M., Kharuk, V. I., Kitzberger, T., Klein, T., Levanič, T., Linares, J. C., Mäkinen, H., Oberhuber, W., Papadopoulos, A., Rohner, B., Sangüesa-Barreda, G., Stojanovic, D. B., Suárez, M. L., Villalba, R., and Martínez-Vilalta, J.: Low growth resilience to drought is related to future mortality risk in trees, *Nature Communications*, 11, 545, <https://doi.org/10.1038/s41467-020-14300-5>, 2020.
- 550 Didan, K.: MYD13A2 MODIS/Aqua Vegetation Indices 16-Day L3 Global 1km SIN Grid V006, Data set. NASA EOSDIS Land Processes Distributed Active Archive Center, <https://doi.org/10.5067/MODIS/MYD13A2.006>, Accessed 2024-03-19, 2015a.
- Didan, K.: MOD13A2 MODIS/Terra Vegetation Indices 16-Day L3 Global 1km SIN Grid V006, Data set. NASA EOSDIS Land Processes Distributed Active Archive Center, <https://doi.org/10.5067/MODIS/MOD13A2.006>, Accessed 2024-03-19, 2015b.
- 555 Dong, C., MacDonald, G., Willis, K., Gillespie, T., Okin, G., and Williams, A.: Vegetation Responses to 2012–2016 Drought in Northern and Southern California, *Geophysical Research Letters*, <https://doi.org/10.1029/2019GL082137>, 2019.
- Forkel, M., Drüke, M., Thurner, M., Dorigo, W., Schaphoff, S., Thonicke, K., Von Bloh, W., and Carvalhais, N.: Constraining modelled global vegetation dynamics and carbon turnover using multiple satellite observations, *Scientific Reports*, 9, 18757, <https://doi.org/10.1038/s41598-019-55187-7>, 2019.
- 560 Gessler, A., Bottero, A., Marshall, J., and Arend, M.: The way back: recovery of trees from drought and its implication for acclimation, *New Phytologist*, 228, <https://doi.org/10.1111/nph.16703>, 2020.

- Gupta, H., Kling, H., Yilmaz, K., and Martinez, G.: Decomposition of the Mean Squared Error and NSE Performance Criteria: Implications for Improving Hydrological Modelling, *Journal of Hydrology*, 377, 80–91, <https://doi.org/10.1016/j.jhydrol.2009.08.003>, 2009.
- 565 Harper, K. L., Lamarche, C., Hartley, A., Peylin, P., Ottlé, C., Bastrikov, V., San Martín, R., Bohnenstengel, S. I., Kirches, G., Boettcher, M., Shevchuk, R., Brockmann, C., and Defourny, P.: A 29-year time series of annual 300 m resolution plant-functional-type maps for climate models, *Earth System Science Data*, 15, 1465–1499, <https://doi.org/10.5194/essd-15-1465-2023>, 2023.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS), <https://doi.org/10.24381/cds.adbb2d47>, accessed on 11-04-2024, 2023.
- 570 Huang, K. and Xia, J.: High ecosystem stability of evergreen broadleaf forests under severe droughts, *Global Change Biology*, 25, 3494–3503, <https://doi.org/10.1111/gcb.14748>, 2019.
- Hughes, N., Galeano, D., and Hattfield-Dodds, S.: The effects of drought and climate variability on Australian farms, Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra, ISBN 978-1-74323-459-4, <https://doi.org/10.25814/5de84714f6e08>, cC BY 4.0. <http://doi.org/10.25814/5de84714f6e08>, 2019.
- International Food Policy Research Institute (IFPRI): Global Spatially-Disaggregated Crop Production Statistics Data for 2020 Version 1.0, Geospatial Data. Harvard Dataverse, <https://doi.org/10.7910/DVN/SWPENT>, version 1.0, 2024.
- Jiao, T., Williams, C., Rogan, J., De Kauwe, M., and Medlyn, B.: Drought Impacts on Australian Vegetation During the Millennium Drought Measured With Multisource Spaceborne Remote Sensing, *Journal of Geophysical Research: Biogeosciences*, 125, <https://doi.org/10.1029/2019JG005145>, 2020.
- 580 Knoben, W., Freer, J., and Woods, R.: Technical note: Inherent benchmark or not? Comparing Nash-Sutcliffe and Kling-Gupta efficiency scores, *Hydrology and Earth System Sciences Discussions*, pp. 1–7, <https://doi.org/10.5194/hess-2019-327>, 2019.
- Knoben, W. J. M.: Global USDA-NRCS Soil Texture Class Map, HydroShare, <https://doi.org/10.4211/hs.1361509511e44adfba814f6950c6e742>, 2021.
- 585 Kolus, H. R., Huntzinger, D. N., Schwalm, C. R., Fisher, J. B., McKay, N., Fang, Y., Michalak, A. M., Schaefer, K., Wei, Y., Poulter, B., Mao, J., Parazoo, N. C., and Shi, X.: Land carbon models underestimate the severity and duration of drought’s impact on plant productivity, *Scientific Reports*, 9, 2758, <https://doi.org/10.1038/s41598-019-39373-1>, 2019.
- Krause, A., Papastefanou, P., Gregor, K., Layritz, L. S., Zang, C. S., Buras, A., Li, X., Xiao, J., and Rammig, A.: Quantifying the impacts of land cover change on gross primary productivity globally, *Scientific Reports*, 12, 18 398, <https://doi.org/10.1038/s41598-022-23120-0>, 590 2022.
- Lange, S., Mengel, M., Treu, S., and Büchner, M.: ISIMIP3a atmospheric climate input data (v1.0), <https://doi.org/10.48364/ISIMIP.982724>, iSIMIP Repository, 2022.
- Liu, P.-W., Famiglietti, J., Purdy, A., Adams, K. H., McEvoy, A., Reager, J., Bindlish, R., Wiese, D., David, C., and Rodell, M.: Groundwater depletion in California’s Central Valley accelerates during megadrought, *Nature Communications*, 13, <https://doi.org/10.1038/s41467-022-35582-x>, 2022.
- 595 Luo, L., Apps, D., Arcand, S., Xu, H., Pan, M., and Hoerling, M.: Contribution of Temperature and Precipitation Anomalies to the California Drought During 2012–2015: contribution of T and P to CA drought, *Geophysical Research Letters*, 44, <https://doi.org/10.1002/2016GL072027>, 2017.
- McDowell, N., Rosie, F., Xu, C., Domec, J.-C., Hölttä, T., Mackay, D., Sperry, J., Boutz, A., Dickman, L. T., Gehres, N., Limousin, J., Macalady, A., Martinez Vilalta, J., Mencuccini, M., Plaut, J., Ogée, J., Rasse, D., Ryan, M., and Pockman, W.: Evaluating theories of drought- 600

- induced vegetation mortality using a multimodel-experiment framework, *The New phytologist*, 200, <https://doi.org/10.1111/nph.12465>, 2013.
- McKee, T., Doesken, N., and Kleist, J.: The Relationship of Drought Frequency and Duration to Time Scales, paper presented at 8th Conference on Applied Climatology, Am. Meteorol. Soc., Anaheim, Calif., 17, 1993.
- 605 Meyer, B. F., Darela-Filho, J. P., Gregor, K., Buras, A., Gu, Q.-L., Krause, A., Liu, D., Papastefanou, P., Asuk, S., Grams, T. E. E., Zang, C. S., and Rammig, A.: Simulating the drought response of European tree species with the dynamic vegetation model LPJ-GUESS (v4.1, 97c552c5), *Geoscientific Model Development*, 18, 4643–4666, <https://doi.org/10.5194/gmd-18-4643-2025>, 2025.
- Moravec, V., Markonis, Y., Rakovec, O., Svoboda, M., Trnka, M., Kumar, R., and Hanel, M.: Europe under multi-year droughts: How severe was the 2014–2018 drought period?, *Environmental Research Letters*, 16, <https://doi.org/10.1088/1748-9326/abe828>, 2021.
- 610 Naumann, G., Podesta, G., Marengo, J., Luterbacher, J., Bavera, D., Acosta Navarro, J., Arias-Muñoz, C., Barbosa, P., Cammalleri, C., Cuartas, L. A., et al.: Extreme and long-term drought in the La Plata Basin: event evolution and impact assessment until September 2022, 2023.
- Pivovarov, A., Pasquini, S., De Guzman, M., Alstad, K., Stemke, J., and Santiago, L.: Multiple strategies for drought survival among woody plant species, *Functional Ecology*, 30, <https://doi.org/10.1111/1365-2435.12518>, 2016.
- 615 Powell, T., Galbraith, D., Christoffersen, B., Harper, A., Imbuzeiro, H., Rowland, L., Almeida, S., Brando, P., Costa, A. C., Costa, M., Levine, N., Malhi, Y., Saleska, S., Sotta, E., Williams, M., Meir, P., and Moorcroft, P.: Confronting model predictions of carbon fluxes with measurements of Amazon forests subjected to experimental drought, *The New phytologist*, 200, <https://doi.org/10.1111/nph.12390>, 2013.
- Rahman, F., Sims, D., Cordova, V., and El Masri, B.: Potential of MODIS EVI and Surface Temperature for Directly Estimating Per-Pixel
620 Ecosystem C Fluxes, *Geophysical Research Letters*, 32, <https://doi.org/10.1029/2005GL024127>, 2005.
- Ruijsch, D.: deniseruijsch/MultiDry-Chapter2: v1, <https://doi.org/10.5281/zenodo.17085725>, 2025.
- Ruijsch, D. and Wanders, N.: Multi-Dry Chapter 2: Exploring the Potential of LPJmL-5 to Simulate Vegetation Responses to (Multi-Year) Droughts, <https://doi.org/10.24416/UU01-U97GDE>, [Dataset], 2026.
- Ruijsch, D., van Mourik, J., Biemans, H., Hauswirth, S. M., and Wanders, N.: Thrive or Wither: Exploring the Impacts of Multiyear Droughts
625 on Vegetation, *Journal of Geophysical Research: Biogeosciences*, <https://doi.org/10.1029/2025JG008992>, 2025.
- Running, S. W. and Zhao, M.: User's Guide: Daily GPP and Annual NPP (MOD17A2H/A3H) and Year-end Gap-Filled (MOD17A2HGF/A3HGF) Products: NASA Earth Observing System MODIS Land Algorithm (Collection 6.1), Numerical Terradynamic Simulation Group, University of Montana, <https://www.umt.edu/numerical-terradynamic-simulation-group/project/modis/user-guides/mod17c61usersguidev11mar112021.pdf>, 2021.
- 630 Santiago, L., Bonal, D., De Guzman, M., and Ávila-Lovera, E.: Drought Survival Strategies of Tropical Trees, vol. 6, pp. 243–258, ISBN 978-3-319-27422-5, https://doi.org/10.1007/978-3-319-27422-5_11, 2016.
- Schaefer, K., Schwalm, C. R., Williams, C., Arain, M. A., Barr, A., Chen, J. M., Davis, K. J., Dimitrov, D., Hilton, T. W., Hollinger, D. Y., et al.: A model-data comparison of gross primary productivity: Results from the North American Carbon Program site synthesis, *Journal of Geophysical Research: Biogeosciences*, 117, 2012.
- 635 Schaphoff, S., Forkel, M., Müller, C., Knauer, J., von Bloh, W., Gerten, D., Jägermeyr, J., Lucht, W., Rammig, A., Thonicke, K., and Waha, K.: LPJmL4 – a dynamic global vegetation model with managed land – Part 2: Model evaluation, *Geoscientific Model Development*, 11, 1377–1403, <https://doi.org/10.5194/gmd-11-1377-2018>, 2018a.

- Schaphoff, S., von Bloh, W., Rammig, A., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke, J., Jägermeyr, J., Knauer, J., Langerwisch, F., Lucht, W., Müller, C., Rolinski, S., and Waha, K.: LPJmL4 – a dynamic global vegetation model with managed land – Part 1: Model description, *Geoscientific Model Development*, 11, 1343–1375, <https://doi.org/10.5194/gmd-11-1343-2018>, 2018b.
- 640 Sims, D., Rahman, F., Cordova, V., El Masri, B., Baldocchi, D., Flanagan, L., Goldstein, A., Hollinger, D., Misson, L., Monson, R., Oechel, W., Schmid, H., Wofsy, S., and Xu, L.: On the use of MODIS EVI to assess gross primary productivity of North American ecosystems, *Journal of Geophysical Research*, 111, <https://doi.org/10.1029/2006JG000162>, 2006.
- Tang, X., Li, H., Huang, N., Li, X., Xu, X., Ding, Z., and Xie, J.: A comprehensive assessment of MODIS-derived GPP for forest ecosystems using the site-level FLUXNET database, *Environmental Earth Sciences*, 74, 5907–5918, <https://doi.org/10.1007/s12665-015-4615-0>, 2015.
- 645 van der Wiel, K., Batelaan, T. J., and Wanders, N.: Large increases of multi-year droughts in north-western Europe in a warmer climate, *Climate Dynamics*, 60, 1781–1800, <https://doi.org/10.1007/s00382-022-06373-3>, 2023.
- van Dijk, A., Beck, H., Crosbie, R., Jeu, R., Liu, Y., Podger, G., Timbal, B., and Viney, N.: The Millennium Drought in southeast Australia (2001–2009): Natural and human causes and implications for water resources, ecosystems, economy, and society, *Water Resources Research*, 49, <https://doi.org/10.1002/wrcr.20123>, 2013.
- 650 van Hateren, T. C., Chini, M., Matgen, P., and Teuling, A. J.: Ambiguous Agricultural Drought: Characterising Soil Moisture and Vegetation Droughts in Europe from Earth Observation, *Remote Sensing*, 13, <https://doi.org/10.3390/rs13101990>, 2021.
- van Mourik, J., Ruijsch, D., van der Wiel, K., Hazeleger, W., and Wanders, N.: Regional drivers and characteristics of multi-year droughts, *Weather and Climate Extremes*, 48, 100748, <https://doi.org/https://doi.org/10.1016/j.wace.2025.100748>, 2025.
- 655 Vicente-Serrano, S., Beguería, S., and López-Moreno, J.: A Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index, *Journal of Climate*, 23, 1696–1718, <https://doi.org/10.1175/2009JCLI2909.1>, 2010.
- Volkholz, J. and Müller, C.: ISIMIP3 soil input data, <https://doi.org/10.48364/ISIMIP.942125>, 2020.
- von Bloh, W., Schaphoff, S., Müller, C., Rolinski, S., Waha, K., and Zaehle, S.: Implementing the nitrogen cycle into the dynamic global vegetation, hydrology, and crop growth model LPJmL (version 5.0), *Geoscientific Model Development*, 11, 2789–2812, <https://doi.org/10.5194/gmd-11-2789-2018>, 2018.
- 660 Wang, L., Zhu, H., Lin, A., Zou, L., Qin, W., and Du, Q.: Evaluation of the Latest MODIS GPP Products across Multiple Biomes Using Global Eddy Covariance Flux Data, *Remote Sensing*, 9, <https://doi.org/10.3390/rs9050418>, 2017.
- Wang, S. and Zhang, Y.: Temporally corrected long-term satellite solar-induced chlorophyll fluorescence (SIF) dataset (1995–2018), <https://doi.org/10.6084/m9.figshare.21546066.v1>, 2023.
- 665 Wiel, K., Batelaan, T., and Wanders, N.: Large increases of multi-year droughts in north-western Europe in a warmer climate, *Climate Dynamics*, 60, 1–20, <https://doi.org/10.1007/s00382-022-06373-3>, 2022.
- Wittwer, G. and Waschik, R.: Estimating the economic impacts of the 2017–2019 drought and 2019–2020 bushfires on regional NSW and the rest of Australia, *Australian Journal of Agricultural and Resource Economics*, 65, <https://doi.org/10.1111/1467-8489.12441>, 2021.
- 670 Wu, X., Liu, H., Li, X., Ciais, P., Babst, F., Guo, W., Zhang, C., Magliulo, V., Pavelka, M., Liu, S., Huang, Y., Wang, P., Shi, C., and Ma, Y.: Differentiating drought legacy effects on vegetation growth over the temperate Northern Hemisphere, *Global Change Biology*, 24, 504–516, <https://doi.org/10.1111/gcb.13920>, epub 2017 Oct 23, 2018.
- Yang, J. and Tian, H.: ISIMIP3a N-deposition input data (v1.3), <https://doi.org/10.48364/ISIMIP.759077.3>, iSIMIP Repository, 2023.
- Yang, M., He, Z., Pi, G., and You, M.: Spatiotemporal Variations in MODIS EVI and MODIS LAI and the Responses to Meteorological Drought across Different Slope Conditions in Karst Mountain Regions, *Sustainability*, 16, <https://doi.org/10.3390/su16177870>, 2024.
- 675

- Yang, X., Xu, X., Stovall, A., Chen, M., and Lee, J.: Recovery: Fast and Slow—Vegetation Response During the 2012–2016 California Drought, *Journal of Geophysical Research: Biogeosciences*, 126, <https://doi.org/10.1029/2020JG005976>, 2021.
- Zang, C., Buras, A., Esquivel Muelbert, A., Jump, A., Rigling, A., and Rammig, A.: Standardized drought indices in ecological research: Why one size does not fit all, *Global Change Biology*, 26, 322–324, <https://doi.org/10.1111/gcb.14809>, 2019.
- 680 Zhou, S., Duursma, R., Medlyn, B., Kelly, J., and Prentice, I.: How should we model plant responses to drought? An analysis of stomatal and non-stomatal responses to water stress, *Agricultural and Forest Meteorology*, s 182–183, 204–214, <https://doi.org/10.1016/j.agrformet.2013.05.009>, 2013.
- Zhu, X., Pei, Y., Zheng, Z., Dong, J., Zhang, Y., Wang, J., Chen, L., Doughty, R. B., Zhang, G., and Xiao, X.: Underestimates of Grassland Gross Primary Production in MODIS Standard Products, *Remote Sensing*, 10, <https://doi.org/10.3390/rs10111771>, 2018.
- 685 Češljar, G., Baković, Z., Đorđević, I., Eremija, S., Lučić, A., Živanović, I., and Konatar, B.: Impact of Short-Term and Prolonged (Multi-Year) Droughts on Tree Mortality at the Individual Tree and Stand Levels, *Plants*, 14, 1904, <https://doi.org/10.3390/plants14131904>, 2025.