

The impact of essential climate variables on respiration rates in subpolar and polar planktonic foraminifera

¹Diane [V. Armitage*](#), ²Nicolaas Glock, ^{1,3,4}Thomas Weiss, ⁵Mohamed [M. Ezat](#), ⁵Adele Westgård, ⁵Freya E. Sykes, ⁶Julie Meilland, ¹Elwyn de la Vega, ¹Alessio Fabbrini, ⁷Tali Babila, and ^{1,3}Audrey Morley*

¹School of Geography, Archaeology, and Irish Studies, and Ryan Institute, University of Galway, Ireland

²Institute for Geology, University of Hamburg, Germany

³iCRAG – Irish Centre for Research in Applied Geosciences, Belfield, Ireland.

⁴Georgia Institute of Technology, Atlanta, Georgia

⁵iC3: Centre for ice, Cryosphere, Carbon and Climate, Department of Geosciences, UiT, The Arctic University of Norway, Norway

^{6,5}CEREGE, Aix-Marseille Université, CNRS, IRD, INRAE, CEREGE, France

^{7,6}Department of Earth, Environmental and Planetary Sciences, Case Western Reserve University, USA

Correspondence to: Diane V. Armitage (D.Armitage1@universityofgalway.ie) and Audrey Morley (audrey.morley@universityofgalway.ie)

Abstract: This study investigates the impact of Essential Climate Variables (ECVs) on the respiration rate of polar planktonic foraminifera *Neogloboquadrina pachyderma* and subpolar *Turborotalita quinqueloba* and *Neogloboquadrina incompta* to advance our understanding of foraminifera physiology and geochemical proxy interpretation for species living in understudied subpolar and polar environments. Respiration rates were measured on a total of 163 specimens collected during two field campaigns to the Nordic Seas. To size-normalise respiration rates we measured cavity volume and maximum diameter using x-ray microcomputed tomography (micro-CT) ($\sqrt[3]{\text{cavity volume}} = (0.56 (\text{max } \varnothing) - 0.38)$). Our results show that the physiological response of foraminifera sharing overlapping environments is diverse, with *N. pachyderma* exhibiting remarkable stability over large gradients in temperature, salinity, carbonate chemistry, dissolved oxygen and nutrients. Conversely, *N. incompta* and *T. quinqueloba* have a much stronger thermal response. The difference between species is best described by their respective Q_{10} (the factor by which the rate of respiration changes with a 10°C increase in temperature) values of 1.41 for *N. pachyderma* and 3.58 and 4.53 for *N. incompta* and *T. quinqueloba*, respectively. We also find a significant relationship between cavity volume and respiration rate ($\text{Log}_{10} \text{respiration rate} = 0.399 (\text{Log}_{10} \text{cavity volume}) - 5.785$) for all three species analysed here, which is consistent with marine protists globally. We conclude that respiration is unlikely to influence geochemical proxies and therefore past climate reconstructions derived from *N. pachyderma*, however, this may not apply to *N. incompta* and *T. quinqueloba*.

1. Introduction

Planktonic foraminifera are relatively eurythermal and inhabit a wide range of temperatures and environments while maintaining species-specific temperature preferences (Chaabane et al., 2024). Accordingly, many species can be found across large temperature gradients exceeding 10 °C, and yet, studies focussing on temperate and tropical species have found a strong influence of temperature and cell volume on respiration and growth rate

40 indicating a strong influence of changing environments on foraminifera physiology (Rink et al., 1998; Lombard
et al., 2009; Burke et al., 2025). However, little is known about how subpolar and polar planktonic foraminifera
respiration responds to temperatures below 10 °C and other [Essential Climate Variables \(ECVs\)](#) such as salinity,
carbonate chemistry, and nutrients. Specifically, there are no observations of the physiological processes
underlying the growth and response of these species to environmental stressors such as modern [Polar Arctic](#)
45 Amplification (the [faster](#) warming of the [high latitudes Arctic relative to four times](#) the global mean) (Serreze
and Barry, 2011; Rantanen et al., 2022). Critically, this gap in our knowledge limits our understanding of how
foraminifera will adapt to rapid environmental change and our ability to predict the future resilience of these
ecosystems.

Respiration also significantly influences foraminiferal shell chemistry by modifying the seawater chemistry in the
50 microenvironment surrounding the test (Schiebel and Hemleben, 2017). For example, increased respiration has
been shown to alter carbon isotope ($\delta^{13}\text{C}$) values (used in reconstructing past changes in water mass properties or
productivity) (Spero and Lea, 1993, 1996) and Mg/Ca ratios (for inferring past sea surface or bottom water
temperatures) (Rink et al., 1998; Köhler-Rink and Kühl, 2001; Eggins et al., 2004). Respiration influences these
proxies because O_2 consumption lowers the pH in the diffusive boundary layer, which in turn alters the carbonate
55 chemistry transported to the site of calcification (Wolf-Gladrow et al., 1999; de Nooijer et al., 2014). This raises
concerns about non-thermal factors that may alter foraminiferal geochemistry and affect the reliability of
temperature reconstructions. Investigating how respiration in polar and subpolar species responds to temperature
and other ECVs is thus key to evaluating whether such physiological processes may introduce uncertainties into
geochemical proxies recorded in their shells.

60 Within the polar oceans, *Neogloboquadrina pachyderma* is the dominant planktonic foraminifera species (Al-
Sabouni et al., 2007; Husum and Hald, 2012; Chaabane et al., 2024), making up more than 90 % of the total
assemblages in waters < 4 °C (Spindler, 1996; Greco et al., 2019; Bertlich et al., 2021) and up to 23 % of the
calcium carbonate CaCO_3 flux to the sediments north of 50° (Tell et al., 2022). *Neogloboquadrina pachyderma*
maintains a unique adaptation to these extremely cold and diverse environments including low temperatures (-2
65 to +12 °C) and large gradients in salinity (~30 – 35 psu) including brine channels with salinities of up to 80 psu,
and pH (7.8 - 8.8) (e.g., Spindler and Dieckmann, 1986; Manno et al., 2012; Bertlich et al., 2021; Zamelczyk
et al., 2021; Westgård et al., 2023). The closely related foraminiferal species *Neogloboquadrina incompta* typically
inhabits subpolar surface waters (relative abundance >50 %) where sea surface temperatures (SSTs) range
between 10 and 18 °C. The largest abundances of *N. incompta* occur at salinities ranging between 31 and 35 psu
70 (Greco et al., 2020). *Turborotalita quinqueloba* is most abundant in subpolar waters of the West Spitzbergen
Current (Volkman, 2000) and the Barents Sea with recent studies in that area reporting maximum relative
abundances of 26 % (Meilland et al., 2019; Anglada-Ortiz et al., 2025). Its maximum abundance occurs at a
salinity of 35 psu (Natland, 1938). [T. quinqueloba is typically associated with mid to higher latitude ecosystems
at temperatures between 1 °C and 21 °C but predominates at temperatures colder than 12 °C \(Bé and Tolderlund,
1971\), however, it has been found on the Great Barrier Reef and in the Arabian Sea at temperatures as high as
75 29.5 °C \(Darling et al., 2000; Secars et al., 2012\).](#)

The objective of this study is to measure the respiration rates of polar (*N. pachyderma*) and sub-polar (*N. incompta*
and *T. quinqueloba*) foraminifera to assess the relationship between respiration and ECVs (e.g., temperature,

salinity, pH, dissolved oxygen, alkalinity, and the saturation state of calcite in seawater (Ω_{Ca}), including nutrients
80 (Silicate (SiO_2), Dissolved Inorganic Carbon (DIC), Phosphate (PO_4^{3-}), and Total Organic Nitrogen (TON)) for
these species. This will allow us to test the hypothesis that temperature has a direct effect on respiration rates as
observed in temperate and tropical species (Lombard et al., 2009). Similarly, we will test the physiological
dependence of external pH and carbonate chemistry on respiration rates in non-spinose planktonic foraminifera
species, which is hypothesized to be reduced due to higher energy requirements of calcification under low pH
85 conditions (Davis et al., 2017). Although earlier research on nutrients such as PO_4^{3-} and SiO_2 found they are not
directly utilised by foraminifera, with their primary effects instead arising through impacts on primary
productivity (Schiebel et al., 2001), more recent research on benthic foraminifera has found that PO_4^{3-} can be
accumulated for energy metabolism, pH regulation, and osmoregulation (Glock et al., 2020, 2025), a capacity that
may likewise occur in planktonic foraminifera. Including these variables therefore allows us to explore potential
90 indirect effects and assess whether environmental variability contributes to physiological stress or metabolic shifts
which could be demonstrated through changes in respiration. The results will allow us to assess the importance of
respiration for polar and subpolar planktonic foraminifera and thereby further our understanding on their ability
to adapt to rapid environmental change. In addition, our results will allow us to review the importance and
implications of respiration for geochemical proxies measured in these three species.

95 Determining the response of respiration to temperature requires accurate measurements of individual foraminiferal
respiration under controlled settings. Recent advancements in micro-sensor periphery technology, particularly the
nano-respiration and rosette methodology developed by Unisense (Lopes et al., 2005; Nielsen et al., 2007), have
significantly improved the reproducibility required to detect extremely low respiration rates encountered in
smaller sized planktonic foraminifera species e.g. $< 200 \mu m$ living in polar and subpolar environments. The
100 automated rosette allows for more repetition due to faster analysis time compared to traditional manual profiling
and enhanced throughput which minimises stress on specimens during experimentation. In addition, the precise
determination of internal foraminiferal biovolume is required for size normalisation, since cell size may correlate
positively with respiration as in both benthic (Hannah et al., 1994; Geslin et al., 2011) and other planktonic
foraminifera (Burke *et al.*, 2025). Methods for accurately estimating biovolume vary in the literature, often relying
105 on geometric approximations (e.g. Hannah et al., 1994; Cesbron et al., 2016; Geslin et al., 2011; Maciute et al.,
2023), however, the use of high-resolution micro-CT scanning (Burke et al., 2020) provides a more accurate
means to assess internal volume and will enable us to refine the metabolic relationships between size and
respiration rate.

2. Methods

110 2.1. Sample collection and processing

Living specimens of *N. pachyderma*, *N. incompta*, and *T. quinqueloba* analysed in this study were obtained during
two oceanographic cruises on the RV Celtic Explorer in July/August 2023 (CE23011) and on the RV Helmer
Hansen (ARCLIM-24-1) in June 2024 (Table 1 and Fig. 1). Plankton samples were collected using 100 μm
vertical-closing HydroBios multineets and 63 μm vertical-closing WP2 nets, respectively. Seawater samples were
115 collected with Niskin bottles attached to a CTD profiler and filtered using a vacuum pump with 0.2 μm Whatmann
filters Type G/C and refrigerated. Live foraminifera were picked directly from the plankton net samples and

transferred into a petri dish for an initial bath to remove debris and algae attached to the foraminifera. For the first set of experiments in 2023, a small brush and a dissecting microscope were used to select and transfer living specimens into culture wells (CE23011), placed in an incubator set to the towed environment of the samples, and allowed to rest for at least 12 h and no longer than 24 h before respiration rate measurements were performed on board (CE23011). Foraminifera were not fed prior to respiration measurements to be consistent with common practice on similar planktonic foraminifera respiration studies (Davis 2017; Rink 1998; Burke 2025; Lombard 2009).

For the second set of experiments performed in 2024, all collected specimens were transferred from the initial bath into 75 ml culture bottles and placed into a cold room set at 8 °C during the cruise. After the initial 24 h resting period, the foraminifera were fed 40 µL of a live diatom solution, a self-replicating food source (Fabbrini et al., in review). Subsequently, culture bottles were transferred into incubators set at experimental temperatures of 2 °C, 5 °C, 8 °C and 13 °C for at least 48 h before flux measurements were performed in the Culturing Laboratory at UiT, The Arctic University of Norway. An additional feeding of algae (*Nannochloropsis sp.*) took place 7 days after sampling. All experiments in 2024 were carried out within 11 days of sampling.

Table 1 Metadata, ecological and carbonate chemistry data for analysed samples. NA indicates no data collected at these stations.

| Station [ID] | Taxon | Start | End | Lat [N] | Long [E] | Depth [m] | SST [°C] | Sal [psu] | DIC [µmol.kg ⁻¹] | Alk | pH |
|-----------------|---|----------|----------|------------|-------------|--------------|-------------|--------------|---------------------------------|------|------|
| S1 | <i>N. incompta</i> | 23/07/23 | 24/07/23 | 55.64 | -14.01 | 40 | 13 | 35.45 | 2138 | 2309 | 7.98 |
| S2 | <i>N. pachyderma</i> | 28/07/23 | 29/07/23 | 71.63 | -8.42 | 50 | 1 | 34.58 | 2156 | 2276 | 7.93 |
| S8.1 | <i>N. pachyderma</i> | 02/08/23 | 04/08/23 | 74.25 | -10.07 | 75 | 3.5 | 34.92 | 2172 | 2300 | 8.12 |
| S10 | <i>N. pachyderma</i> | 04/08/23 | 05/08/23 | 70.5 | -17.09 | 100 | 0.5 | 34.57 | 2158 | 2293 | 8.07 |
| S11 | <i>N. pachyderma</i> | 05/08/23 | 06/08/23 | 67.87 | -21.77 | 125 | 2.5 | 34.76 | 2169 | 2298 | 8.06 |
| S12 | <i>N. pachyderma</i> | 07/08/23 | 08/08/23 | 65.42 | -28.33 | 20 | 10 | 35.09 | 2110 | 2317 | 8.08 |
| S16 | <i>N. incompta</i> ; <i>N. pachyderma</i> | 10/08/23 | 10/08/23 | 62.75 | -37.51 | 20 | 10 | 34.9 | 2112 | 2305 | 8.09 |
| S17 | <i>N. pachyderma</i> | 11/08/23 | 11/08/23 | 60.18 | -39.13 | 50 | 6 | 34.91 | 2168 | 2299 | 8.09 |
| S18 | <i>N. pachyderma</i> | 12/08/23 | 13/08/23 | 58.25 | -45.64 | 45 | 7.5 | 34.66 | 2129 | 2185 | 7.89 |
| S19 | <i>N. pachyderma</i> | 13/08/23 | 14/08/23 | 57.55 | -48.52 | 35 | 6 | 34.7 | 2134 | 2296 | 8.11 |
| S20 | <i>N. incompta</i> | 16/08/23 | 17/08/23 | 56.36 | -27.89 | 10 | 14 | 35.04 | 2080 | 2311 | 8.09 |
| S856 | <i>N. pachyderma</i> | 06/06/24 | 06/06/24 | 72.13 | 5.1 | 50 | 7.5 | 35.1 | NA | NA | NA |
| S866 | <i>T. quinqueloba</i> | 07/06/24 | 07/06/24 | 71.21 | 11.5 | 50 | 8 | 35 | NA | NA | NA |

135




Figure 1 Image showing the station locations for cruise CE23011 on the RV Celtic Explorer (2023). Stations S856 and S866 are the sampling sites for ARCLIM-24-1 cruise on the RV Helmer Hansen (2024). Map created using ODV (Schlitzer, 2022). Site symbol colour represents the ocean temperature at the tow depth of the sample site.

140 2.2 Carbonate Chemistry

Total alkalinity of seawater samples was determined onboard cruise CE23011 using an Apollo SciTech AS-ALK3 Total Alkalinity Titrator. This instrument operates on the principle of Gran titration, whereby approximately 0.1 M hydrochloric acid is incrementally added to 20 mL aliquots of seawater maintained at 20 °C. The resulting titration curve is used to construct a Gran function from which total alkalinity is calculated. The pH measurements were conducted using an Orion 8302BNUMD Ross Ultra pH/ATC Triode probe, which was calibrated daily using Thermo Scientific buffer solutions at pH 4.01, 7.00, and 10.01. The concentration of the titrant (HCl) was standardised by titrating Certified Reference Material (CRM) Batch 208 for Oceanic CO₂ analysis, prepared by Dr. Andrew Dickson (Scripps Institution of Oceanography), a minimum of three times. Calibration was repeated until the relative standard deviation (RSD) of the calculated HCl concentration from at least two titrations was below 0.01 %. CRM Batch 208 was also used as a quality control standard, analysed approximately every fifth sample. When treated as a sample, Batch 208 yielded a standard deviation of 7.69 μmol kg⁻¹ and an RSD of 0.35 %. Replicate analyses were performed on every second sample, with an average difference of 8.43 μmol kg⁻¹ and a relative average difference of 0.37 %. Following titration, alkalinity values were converted from μmol L⁻¹ to μmol kg⁻¹ using the equation of state and subsequently corrected for instrumental drift using Batch 208.

DIC samples of seawater were collected in 500 ml glass bottles and poisoned on board with 0.2ml mercuric chloride. DIC concentrations were measured with an LI-5350A DIC analyser coupled with an LI-850 infrared gas analyser. Briefly, 1.5 mL aliquots of seawater were acidified to convert all DIC to CO₂ gas, which was then quantified by the gas analyser. Calibration of the DIC analyser was performed at the beginning and end of each analytical batch (comprising eight sample triplicates) using Batch 208 in triplicate volumes of 1.2 mL, 1.5 mL, and 1.8 mL. Additional triplicate analyses of Batch 208 were conducted after each batch of eight sample triplicates to monitor instrument performance. Standard deviation of averages for Batch 208 triplicates was 1.35 μmol kg⁻¹

across all sample runs. Carbonate system parameters, including $\Delta[\text{CO}_3]$, were calculated from measured alkalinity and DIC using CO2sys version 25b06 (Lewis and Wallace, 1998). We used the equilibrium constants of Lueker et al. (2000), the KSO_4 constant from Dickson, total boron from Uppström (1974), the KF from Dickson and Riley (1979), and default values for PO_4^{3-} and SiO_2 of 0 mol kg^{-1} .

165


2.3 Oxygen measurements

During CE23011, we performed single-specimen respiration measurements on board the RV Celtic Explorer within 24 hours of collection, using the same (filtered) seawater and temperature as foraminifera were collected in. Given the large latitudinal gradient of the survey (e.g., $55 - 74^\circ\text{N}$), this allowed us to perform measurements over a large gradient for most ECVs analysed here. The following year, we collected foraminifera from a single station. We incubated them at different temperatures in the laboratory at UiT before performing respiration measurements in a temperature-controlled laboratory setting. This dual approach allowed us to evaluate the importance of food, rest and setting (ship vs lab) for the reproducibility of the results.

170

Measurements on single foraminifera were performed using the NanoRespiration system developed by Unisense™. It includes the Unisense MicroProfiling System coupled with a micro-rosette of seven fused glass capillaries ($\text{Ø} = 0.68 \text{ mm}$, length = $3 \pm 0.2 \text{ mm}$), a rosette holder and a metal frame which allows exact positioning and movement of the microsensor tip into the glass capillary (see Fig. 2 for set-up diagram). Oxygen profiles were measured using Clark-type microsensors of $50 \mu\text{m}$ tip diameter with guard cathodes (Revsbech, 1989), a fx-6 UniAMP multi-channel amplifier, and SensorTrace PRO v1.9 software from Unisense™. Each sensor was calibrated prior to each experiment using a 0 % oxygen solution using either the premade ascorbate solution prepared by Unisense™ or dissolving Sodium Sulphite PryoScience™ capsules in 50 ml of distilled water and a 100% oxygenated seawater solution using filtered seawater and an aquarium pump.

180



185

Figure 2 Diagram adapted from Unisense (2025) showing the nano respiration set-up used to measure oxygen gradients on single foraminifera. The image on the bottom right shows a close-up of the Nanorespirometer Unit (e.g., rosette) and microsensor entering one of the glass capillaries.

Prior to analysis each glass capillary of the sample rosette was flushed with DI water and placed into an ultrasonic bath for 30 seconds. Once cleaned and DI water removed, the rosette was placed into a Petri dish filled with

190 filtered seawater set at the precise temperature of incubation for each experiment. All glass capillaries were flushed
with filtered seawater carefully removing any remaining bubbles in each capillary. Then, one living foraminifera
(colourful, pseudopodia present) was placed per capillary, keeping one well empty as a control. The room
temperature was maintained within 5 °C of the experiment temperature to avoid large fluctuations for the
foraminifera during transfer from the incubator into the rosette. Once the transfer was complete, the loaded sample
rosette was placed into the rosette holder and submerged in a jacketed beaker filled with filtered seawater at the
195 exact temperature chosen for the experiment. Temperature was maintained to within 0.1 °C using a Julabo
circulator, circulating antifreeze into the jacket of the beaker. Foraminifera in the submerged rosette were
acclimatised for 45-60 minutes. Seawater in the beaker was constantly agitated with air using an aquarium pump
to maintain fully oxygenated seawater above the capillaries. Once acclimatised the position of the sensor was
calibrated using a micromanipulator, a PC-controlled motor unit (Woelfel et al., 2009) and a mounted dissecting
200 microscope to ensure the sensor enters each capillary opening. Oxygen profiles were set to begin 400 µm above
and end 2000 µm inside each capillary, with step sizes of 200 µm. Measurements at each depth were performed
after allowing the sensor to equilibrate for 5 seconds, and each profile was repeated three times. The oxygen
gradient in each capillary was measured using the average slope between 800 µm and 2000 µm inside the capillary
for each triplicate measurement to determine individual respiration rate (IRR) measurements (Maciute et al., 2023)
205 using Fick's first law of diffusion $J = -D \times dC/dx$ where D is the oxygen diffusion coefficient at a given
temperature (Broecker and Peng 1974), and dC/dZ is the measured oxygen gradient inside the capillary tube.
To account for background respiration, one capillary in each rosette was maintained as a blank, containing only
seawater without any foraminifera. Respiration rates from these blanks were measured in triplicate, and the
average value was subtracted from the respiration rates of the experimental chambers to yield a blank-corrected
210 average respiration rate. Blank respiration values averaged at $14.83 \pm 11.50 \text{ pmol h}^{-1} \text{ ind}^{-1}$. Additional tests were
conducted using dead foraminifera specimens that had been previously dehydrated to assess background
respiration in the absence of metabolic activity. Five dead specimens were tested, yielding an average respiration
rate of $17.00 \pm 5.83 \text{ pmol h}^{-1} \text{ ind}^{-1}$, which is indistinguishable from the average procedural blank.

2.4. Maximum diameter and cell volume reconstructions

215 Determining biovolumes using X-ray microcomputed tomography (micro-CT) scanning for all individuals
included in this study (n = 163) was not feasible. We therefore opted to establish a robust empirical relationship
between maximum diameter and cube root cavity volume for *T. quinqueloba*, *N. pachyderma*, and *N. incompta*,
as previously suggested by Burke et al., (2020). Since the three species investigated exhibit similar low
trochospiral coiling morphologies (Darling et al., 2006; El Bani Altuna et al., 2018; Pearson and Kucera, 2018),
220 they are suited for this joint methodology and analysis. For micro-CT analysis tests were glued to a Kapton tube
using a mix of tragacanth gum and MQ water ([Milli-Q water – ultra pure laboratory water produced by a Milli-Q
purification system](#)), following a modified protocol from Coletti et al., (2018), Siccha et al., (2023) and Fabbrini
et al., (2025). The Kapton tube was placed on a sample holder and scanned with the ZEISS Xradia 620 Versa, at
the University of Galway.

225 The sample holder was placed between the X-ray source with a source-to-detector distance of 58 mm (Source-
Rotation Axis distance: -20 mm; Detector-Rotation Axis distance 38 mm), providing a voxel resolution of 240
nm per pixel using the 20X objective magnification in binning 1 mode. The instrument was operated at 120 kV

and 17.5 W, employing no energy filter to optimise transmission and the contrast-to-noise ratio. A total of 1601 radiographs were acquired over a 360° sample rotation range with an exposure time of 10 seconds per radiograph.

230 The raw transmission images (.txm) were reconstructed for each specimen using a commercial image reconstruction software package (ZEISS XMReconstructor) (ZEISS, 2024), which employs a filtered back-projection algorithm to generate the final reconstructed and corrected three-dimensional file. The final reconstructed files (.txm) were then exported as a stack of .tif image files for further study.

235 In addition, each foraminifer was imaged immediately after oxygen profiling using a Moticam X5 Plus (Motic Instruments Inc., [2024]) Wi-Fi camera mounted on a Zeiss Stemi 395 and the Motic Images Plus 3.1 ML software. After calibration, this software was used to measure the maximum diameter of the foraminifera in µm, completed under 64x magnification. The maximum diameter refers to the longest straight line that passes through the widest part of the foraminifera, typically measured through the final chamber (Fig. 4).

2.5. Biometry

240 The exported micro-CT image stacks were segmented using the software *Amira 3D Pro* (Stalling et al., 2005). Manual thresholding was applied to isolate the tests and background, and the paintbrush tool was used to remove infilling or extraneous particles, such as cytoplasm. The segmented tests were rendered as volumes using the watershed algorithm. The maximum diameter was measured for all specimens using the measurement tool on the 3D volume rendering of the test in the same orientation as the optical microscope. The internal voids of all 245 chambers were then isolated and rendered as a separate volume using the “Ambient Occlusion” function in *Amira 3D Pro*, which filled the negative volume virtually (Baum and Titschack, 2016; Titschack et al., 2018). The internal volume of the combined chambers and of the pores within the test wall (Fig. 4) was then measured using the label analysis function in *Amira* (Kiss et al., 2023).

250 Respiration rates were analysed in two ways. First, we used cell volume to size-normalise respiration rates and evaluate the response of the specimens to changes in temperature and other essential climate variables. A common way to estimate the influence of temperature on a physiological rate is the use of the Q_{10} value, which quantifies the rate increase for a 10 °C increase. Q_{10} was calculated following Eq. (1) (adapted from Schmidt-Nielsen, 1997):

$$Q_{10} = (\log_{10} R_2 - R_1) \cdot \frac{10}{(T_2 - T_1)} \quad (\text{Eq.1})$$

255 Where R_2 is the respiration rate at the higher temperature, R_1 is the respiration rate at the lower temperature, T_1 is the lower temperature, and T_2 is the higher temperature. It is essential to recognise that, although the use of a Q_{10} is a convenient measure, it varies as a function of the temperature range being considered (Lombard et al., 2009). For example, when calculating the Q_{10} value for *N. pachyderma*, we excluded the lab-based experiment because it included temperatures outside of this species' typical habitat range.

260 Secondly, we normalised respiration rates to 5 °C, 15 °C and 24 °C to evaluate the influence of cell volume on respiration rates. 24 °C was chosen to facilitate comparative analysis with previous work, particularly that of Lombard et al., (2009), thereby enabling meaningful contextual interpretation of the current dataset. 15 °C represented a thermal midpoint across all experiments, minimising physiological deviation and ensuring that species were assessed under temperatures approximating their average environmental exposure. 5 °C served as

the low-temperature condition, selected for its relevance for *N. pachyderma*, the focal species of this investigation.

265 Using a Q_{10} value of 3.18 from Lombard et al., (2009), and Q_{10} values derived for *N. pachyderma*, *N. incompta* and *T. quinqueloba* in this study (see Table 4) following Eq. (2):

$$R_x = R_t \cdot Q_{10}^{\frac{24-t}{10}} \quad (\text{Eq.2})$$

Where R_x is the respiration rate at X °C, R_t is the respiration rate at the measured temperature, and t is the measured temperature.


270 3. Results

3.1. Cell volume reconstructions

Cavity volume and maximum diameter using micro-CT scans were assessed for a subset of 39 specimens (23.5 %). These include 19 new specimens from this study (e.g., *N. pachyderma* (n = 7), *N. incompta* (n = 5), and *T. quinqueloba* (n = 7) and the reanalysis of 20 .tiff stacks of scanned *N. pachyderma* (n = 9), *N. incompta* (n = 8) and *T. quinqueloba* (n = 3) previously published in Burke et al., (2020). An independent paired student t-test comparing the maximum diameters reported in Burke et al., (2020) and the measurements of the same scans using Amira in the current study had a p-value of < 0.01, (95 % Confidence Interval (C.I.) [1.65-7.4 4 μm]) meaning there is a 0.61 % to 2.75 % difference between measurements reported in Burke et al., (2020) and reported here for the same foraminifera.

280 **Table 2** Summary of maximum diameter and cube root volume measurements of the 39 specimens that were scanned for volume reconstructions.

| | <i>N. pachyderma</i> [n = 16] | <i>N. incompta</i> [n = 13] | <i>T. quinqueloba</i> [n = 10] |
|--|---|---|---|
| Mean max Ø [μm] | 244.3 | 284.2 | 195.8 |
| Range Ø [μm] | 165.48 to 315.5 | 168.36 to 349.08 | 136.91 to 263.33 |
| Mean biovolume $\sqrt[3]{\mu\text{m}}$ | 136.3 | 160.14 | 106.8 |
| Biovolume range $\sqrt[3]{\mu\text{m}}$ | 86.33 to 171.11 | 119.19 to 180.99 | 72.76 to 144.25 |
| $\sqrt[3]{\text{cavity volume}} = a(\text{max } \text{Ø}) + b$ | a = 0.61; b = -10.31 r ² = 0.90; p < 0.01 | a = 0.50; b = +13.19 r ² = 0.93; p < 0.01 | a = 0.60; b = -10.51 r ² = 0.99; p < 0.01 |



285 **Figure 3** Combined max \varnothing (μm) vs $\sqrt[3]{\text{cavity volume}}$ (μm) for *N. pachyderma*, *N. incompta* and *T. quinqueloba* showing trendline, slope and R^2 value. The source data for this figure is available in Table S8


The relationship between the $\text{max}\varnothing$ (μm) and $\sqrt[3]{\text{cavity volume}}$ for all 39 specimens analysed here (Fig. 3) follows Eq. (3):

290
$$\sqrt[3]{\text{cavity volume}} = (0.56 \pm 0.02)\text{max}\varnothing + (-0.38 \pm 5.47) \quad (\text{Eq.3})$$

$n = 39$, $r^2 = 0.95$, $p < 0.01$. For species-specific equations, please see Table 2. Volume reconstructions performed in this study are significantly smaller than previous estimates that are based on the assumption that the cavity volume corresponds to 75 % of the closest geometric shape (e.g., sphere for species analysed) that can be fitted around the maximum diameter of the foraminifera (e.g., Hannah et al., 1994; Geslin et al., 2011; Cesbron et al., 2016; Maciute et al., 2023). A comparison of both techniques shows that using 75 % of a sphere to estimate cavity volume overestimates actual biovolume by 47 ± 7.37 %. Specifically, for each of the species analysed, the relationship between the volume of a sphere based on the max \varnothing (μm) and micro-CT-based cavity volume reconstructions was 35.36 ± 4.57 % for *N. pachyderma*, 33.21 ± 6.09 % for *N. incompta* and 37.69 ± 5.67 % for *T. quinqueloba*. We also note that the protocol for analysing cavity volumes used in this study led to significantly smaller cavity volumes (28 % difference, paired t-test $p > 0.01$) when compared to Burke et al., (2020).

The main difference between methods is that we used the ambient occlusion function in Amira (Titschack et al., 2018), which allows the segmentation of the shell and measurements of the internal volume of the tests, excluding the shell, while in Burke et al., (2020) external meshes were imported into MeshLab software, where they were resurfaced and replaced with watertight “wrap” mesh, which closes all apertures and pores, allowing the interment of the cavity volume of the entire test by subtracting the CaCO_3 volume from the wrap volume. This method appears to overestimate cavity volumes, since the reported cavity volumes exceed the volume of a sphere fitted to the maximum diameter of each test by a mean of 115 ± 27 % for *N. incompta*, 137 ± 20 % for *N. pachyderma* and 179 ± 6 % for *T. quinqueloba*.

310 In this study, we define the term “test biovolume” as the internal cavity volume bound by the calcite test,
representing the space theoretically occupied by living cytoplasm in planktonic foraminifera. While cytoplasmic
density and space occupancy may vary within chambers, this metric captures the maximum possible total living
volume of the cell inside its shell rather than carbon-equivalent biomass. In our definition, test biovolume is thus
confined to the interior volume of the test and does not capture the “catchment volume” or the rhizopodial network
315 which can increase the effective cell volume by several orders of magnitude compared to the test alone (Gaskell
et al., 2019). We choose not to scale reported biovolumes at 75 % of the test biovolume (as in Burke et al 2025)
because the majority of specimens analysed in this study exhibited full chambers prior to measurements.
Furthermore, we note that the suggested 75% method arose from the limitations of estimating biovolumes from
total test volumes that included the shell (e.g., Hannah et al. 1994). Now, using micro-CT internal volume
reconstructions we have more advanced tools to directly measure the cavity volume. Finally, there is no empirical
320 evidence that supports a 75% cell occupancy of the cavity volume for asymbiotic planktonic foraminifera, nor
does a validated technique exist that would allow us to measure it.



325 **Figure 4:** Example of volume rendering of a specimen of *N. pachyderma* selected for 3D volume analysis in Amira 3D Pro. This figure shows the internal voids of the chambers, porosity, and the white line indicates the maximum diameter (Max Diameter) for this specimen. Scale bar 50 μm.

3.2. Respiration Rates

Here we use the multispecies relationship between $\sqrt[3]{\text{cavity volume}} (\mu\text{m})$ and $\text{max } \varnothing (\mu\text{m})$ defined in section 3.1. to size-normalise the respiration rates. For the ship-based CE23011 cruise dataset from 2023, we recorded mean size-normalised respiration rates of $59.22 \text{ pmol hr}^{-1} \text{ ind}^{-1}$ (95 % C.I. [53.32 - 65.13]) for *N. pachyderma* over a
330 temperature gradient of 9.5 °C measured between 0.5 – 10 °C. For *N. incompta* we record higher mean size-normalised respiration rates of $198.99 \text{ pmol hr}^{-1} \text{ ind}^{-1}$ with a larger variability (95 % C.I. [169.48-228.50]) analysed

over a smaller temperature gradient of 4 °C measured between 10 °C and 14 °C. For the laboratory-based dataset (Tromsø 2024), the mean size-normalised respiration rates for *N. pachyderma* were comparable to the ship-based data (e.g., 62.71 pmol hr⁻¹ ind⁻¹ (95 % CI: [57.45, 67.97])). Mean size-normalised respiration rates for *T. quinqueloba*, are also low at 45.75 pmol hr⁻¹ ind⁻¹ with low variability (95 % CI: [31.99, 59.50]) over a temperature gradient of 11 °C measured between 2-13 °C for both species. While early studies by Hemleben Hemleben et al., (1989) and Stangeew (2001) noted the possible presence of symbionts in *T. quinqueloba*, these studies did not present direct evidence, and subsequent studies have found no indication of active photosymbiosis, their presence remains elusive in other studies (e.g., Takagi et al., 2019; Hoogakker et al., (2022); Kanbur, 2025). Consistent with this, and none of the specimens analysed in this study bore symbionts. Thus, calculated respiration rates were not adjusted for photosynthesis as in Lombard et al., (2009) for *O. universa*, *G. ruber* and *G. siphonifera*.


The sensitivity of size-normalised respiration rates and temperature for tropical and subtropical planktonic foraminifera is best described by an exponential or Arrhenius relationship, rather than a simple linear one (Lombard et al., 2009). We therefore plot the log₁₀ size-normalised respiration rates against temperature for our analysis, as temperatures are well established to scale logarithmically with respiration rates (e.g. Burke et al., 2025; Lombard et al., 2009). Conversely, empirically supported functional relationships are not currently available for the other environmental variables measured in this study analysis. The log₁₀ size-normalised respiration rates (pmol hr⁻¹ ind⁻¹ μm⁻¹) against temperature are shown in Fig. 5a for all species. For all other essential climate variables such as, pH, salinity, alkalinity, Ω_{Ca}, dissolved oxygen (Fig. 5), SiO₂ (μmol kg⁻¹), Dissolved Inorganic Carbon, PO₄³⁻ (μmol kg⁻¹), and Total Organic Nitrogen (μmol kg⁻¹), size-normalised respiration rates are shown in Fig 5 b-f and Fig. 6.

In Table 3 we report correlation statistics for each variable and show that there is no significant correlation between respiration rates of *N. pachyderma* and any of the ECVs measured here. However, strong covariance between ECVs in the modern ocean may obscure biological responses of *N. pachyderma* and *N. incompta* to complex environmental gradients. Consequently, we conducted a suite of multivariate analyses to evaluate whether size-normalised respiration rates respond to combined environmental gradients or individual variables. Principal component analysis (PCA) of the complete ECVs dataset relevant for each species revealed structured environmental axes, with PC1–PC4 capturing 95 % of environmental variance for the *N. pachyderma* dataset and PC1–PC2 capturing 100% of variance for the *N. incompta* dataset (Supplementary Table S2). Evaluation of the PCA loadings (Table S3) indicates that these principal components correspond to coherent environmental gradients rather than isolated variables. For example, in the *N. pachyderma* dataset, PC1 primarily reflects a water-mass and nutrient gradient, characterized by strong positive loadings of nutrients (PO₄³⁻, TON, SiO₂) and DIC and negative contributions from temperature and oxygen, consistent with colder nutrient-rich waters. PC2 is dominated by carbonate chemistry variables, including alkalinity, pH, and calcite saturation state (Ω_{Ca}), reflecting variability in the carbonate system. PC3 and PC4 capture secondary productivity-related signals, largely associated with fluorescence, oxygen, and temperature. Despite four clear multivariate gradients, size-normalised respiration rates for *N. pachyderma* do not show a significant relationship with any of the four principal components (all *p* > 0.42), and scatterplots of size-normalised respiration rates across PC1–PC4 space are not significantly different from zero (Fig S1). Principal Component Regression explained 0 % of respiration variance

(adjusted $r^2 = 0.005$) with all PC slopes non-significant except PC4 ($p = 0.01$) but with negligible explanatory power (Table S4). Partial least squares regression similarly showed minimal predictive skill (Y-variance $\leq 22.7\%$; RMSEP comparable to the raw standard deviation; all 95 % coefficient intervals overlapping zero; Table S5). Redundancy analysis detected a statistically significant constrained fraction ($p = 0.001$), but this accounted for only 37.4 % of variance and did not improve performance relative to null models (Table S7). These results indicate that respiration remained invariant across the full multivariate environmental space sampled, supporting physiological robustness rather than the masking of univariate relationships by collinearity.

For *N. incompta* we have a more limited dataset measured over a narrower range in environmental gradients. Here we ~~and~~ find significant linear correlations notable for temperature, SiO_2 , salinity, fluorescence, dissolved O_2 , pH and Ω_{Ca} (Table 3, Table S1). Furthermore, size-normalised respiration rates for *N. incompta* were significantly correlated to the environmental gradients represented by both PC1 and PC2 using Principal Component Analysis ($p < 0.01$; see also Table S4 and Fig. S2). PLSR results were consistent, with 75.3 % of Y-variance explained and coefficient intervals not overlapping zero (Table S5). Redundancy analysis also identified a strong environmental signal (73.4 % constrained variance; $p = 0.001$; Table S6). This suggests that unlike *N. pachyderma* size-normalised respiration rates in *N. incompta* are sensitive to integrated environmental structure. These contrasting responses suggest fundamental differences in metabolic plasticity between the two species. However, we acknowledge the reduced environmental dataset available for *N. incompta*. A detailed correlation matrix, which explores these linear relationships further, is available in Table S1.

Correlation statistics, between \log_{10} size-normalised respiration and temperature and Q_{10} values are reported in Table 4 for all species. We find that temperature accounts for a greater proportion of the variance in respiration for *N. incompta* ($r^2 = 0.45$) and *T. quinqueloba* ($r^2 = 0.71$) than for *N. pachyderma* ($r^2 = 0.05$). The sensitivity to temperature is explained by the Q_{10} values determined for each species. *N. pachyderma* exhibits a low Q_{10} of 1.41 between 0.5 °C and 10 °C. In contrast, *N. incompta* and *T. quinqueloba* demonstrate a much higher Q_{10} of 3.58 and 4.53 over 10-14 °C and 2-13 °C, respectively. We caveat that Q_{10} values for *N. incompta* are only tentative due to the small temperature gradient (e.g., $\Delta T = 4^\circ\text{C}$) covered in this dataset. Specifically, the assumptions required for calculating Q_{10} are not fully met for this species, and any resulting Q_{10} values should be interpreted with caution as a preliminary result that needs to be confirmed over a larger temperature gradient (e.g. $\Delta T = 10^\circ\text{C}$).




400

Figure 5: Box and whisker plots of the effect of essential climate variables on Size Normalised Respiration based on biovolume ($\text{pmol hr}^{-1} \text{ind}^{-1}$) in *N. pachyderma*, *N. incompta* and *T. quinqueloba* (a) Temperature $^{\circ}\text{C}$ and (b) pH; (c) Salinity (psu) and (d) Alkalinity ($\mu\text{mol kg}^{-1}$); (e) Ω_{Ca} and (f) Dissolved Oxygen (ml L^{-1}). Boxes extend from the data's lower to upper quartile values, with a line at the median. Whiskers indicate 1.5 times the interquartile distance. Red dots mark the mean. Log₁₀ Size normalised respiration and laboratory data are only available for (a). The source data for this figure is available in Tables S9 and S10.

405

410



415

Figure 6: Box and whisker plots of the effect of nutrients on Size Normalised Respiration based on Biovolume ($\text{pmol hr}^{-1} \text{ind}^{-1}$) in *N. pachyderma* and *N. incompta*. (a) SiO_2 ($\mu\text{mol kg}^{-1}$) and (b) Dissolved Inorganic Carbon ($\mu\text{mol kg}^{-1}$); (c) PO_4^{3-} ($\mu\text{mol kg}^{-1}$) and TON ($\mu\text{mol kg}^{-1}$). Boxes extend from the lower to upper quartile values of the data, with a line at the median. Whiskers indicate 1.5 times the inter-quartile distance. The red dots mark the means. The source data for this figure is available in Table S9 and S10

Table 3: Quantification of the correlations between size-normalised respiration and essential climate variables and nutrients in *N. pachyderma* and *N. incompta* measured in situ, during expedition CE23011.

| | <i>N. pachyderma</i> | | <i>N. incompta</i> | |
|--|----------------------|----------|--------------------|-----------|
| | [n=68] | | [n=25] | |
| | r^2 | p-value | r^2 | p-value |
| TON | 0.001 | $p=0.85$ | 0.007 | $p=0.70$ |
| PO_4^{3-} | 0.025 | $p=0.19$ | 0.003 | $p=0.81$ |
| SiO_2 | 0.024 | $p=0.21$ | 0.372 | $p=0.001$ |
| Salinity | 0.032 | $p=0.15$ | 0.732 | $p<0.001$ |
| Turbidity | 0.003 | $p=0.67$ | 0.005 | $p=0.73$ |
| Fluorescence | 0.001 | $p=0.85$ | 0.642 | $p<0.001$ |
| DIC | 0.035 | $p=0.13$ | 0.182 | $p=0.03$ |
| Dissolved O_2 | 0.001 | $p=0.90$ | 0.464 | $P<0.001$ |
| Alkalinity | 0.001 | $p=0.90$ | 0.199 | $p=0.03$ |
| pH | 0.005 | $p=0.55$ | 0.642 | $p<0.001$ |
| Ω_{Ca} | 0.014 | $p=0.33$ | 0.225 | $p=0.01$ |

420 **Table 4:** Quantification of the correlations and Q_{10} between \log_{10} size-normalised respiration and temperature in *N. pachyderma*, *N. incompta* and *T. quinqueloba* for in-situ (2023) and laboratory (2024) measured respiration.

| Species | Source | n | Range (°C) | Q_{10} | SD | r^2 | p-value |
|-----------------------|----------|----|------------|----------|-------|-------|----------|
| <i>N. pachyderma</i> | In-situ | 68 | 0.5-10 | 1.41 | 0.203 | 0.05 | p = 0.08 |
| <i>N. incompta</i> | In-situ | 25 | 10-14 | 3.58 | 0.162 | 0.45 | p < 0.01 |
| <i>T. quinqueloba</i> | Lab data | 23 | 2-13 | 4.53 | 0.29 | 0.71 | p < 0.01 |

3.3. Comparison to previously published respiration rates on planktonic foraminifera


425 To quantify the effect of temperature on size-normalised respiration rates, we combined our dataset with previous studies carried out by Lombard et al., (2009) on *Orbulina universa*, *Globigerinoides ruber* and *Globigerinella siphonifera*, by Rink et al., (1998) on *O. universa* and Burke et al., (2025) on *Globorotalia menardii*, *Pulleniatina obliquiloculata*, *Hastigerina pelagica*, *O. universa* and *G. ruber*. (Fig. 7). ~~Test bioCell volumes for *N. pachyderma*, *N. incompta* and *T. quinqueloba* were computed using the relationship between cavity volume and maximum diameter (Fig. 3) derived in this study. We order to compare our results to previous studies by adjusting biovolumes reported in Burke et al., (2025) (e.g., 75% of cavity volume) to 100% that estimated biovolumes at Following Burke et al. (2025), biovolumes were taken as 75 % of the calculated cavity volumes (e.g., Burke et al., 2025), we adjusted biovolume estimates to % reflecting that biovolume does not always correspond directly to cavity volume. This ensures a like-for-like comparison across studies. Results show that~~

430 the respiration rates of all planktonic foraminifera included here follow Eq. (4):

$$\text{Log}_{10}R_{\text{biovolume}} = 0.11t - 5.54 \quad (\text{Eq.4})$$

435 Where $R_{\text{biovolume}}$ is the respiration rate normalised by biovolume, and t is the temperature at which the respiration rate was measured ($r^2 = 0.62$, $n = 201$ and $p < 0.001$). We found that when normalised to 5 °C, 15 °C and 24 °C (Fig. 8), respiration rates scale positively with biovolume across all datasets. The relationships are best described by linear regression following:

$$Rx = m \cdot BV + c \quad \text{Eq. (5)}$$



440

Figure 7: The effect of temperature (°C) on log₁₀ size-normalised respiration rate (pmol O₂ hr⁻¹ ind⁻¹ μm⁻¹) in this study and previously published data (Rink et al., 1998; Lombard et al., 2009 and Burke et al., 2025). The grey shaded regions represent the 95% confidence bounds of the regression model. The source data for this figure is available in Table S11

445

The relationships described by Eq. (5) across different temperatures are:


$$(a) R_4 = 0.34 \pm 0.02 BV - 7.05 \pm 0.21 (r^2 = 0.48, n = 201, p < 0.01)$$

$$(b) R_{15} = 0.46 \pm 0.03 BV - 7.71 \pm 0.24 (r^2 = 0.57, n = 201, p < 0.01)$$

$$(c) R_{24} = 0.56 \pm 0.04 BV - 8.25 \pm 0.34 (r^2 = 0.48, n = 201, p < 0.01)$$

450

Where $R_x = \text{Log}_{10}$ respiration rate normalised to $x^\circ\text{C}$ and $BV = \text{Log}_{10}$ Biovolume (μm^3). The steepest slope occurs when normalizing to 24 °C ($R_{24} = 0.56 \pm 0.04$), followed by 15 °C ($R_{15} = 0.46 \pm 0.03$), and 4 °C ($R_4 = 0.34 \pm 0.02$). R^2 values range between 0.48 and 0.57, indicating that while biovolume is a significant predictor of respiration, other biological or environmental factors likely contribute to the observed variability. Nevertheless, the consistent significance ($p < 0.01$) across all regressions underscores the robustness of the biovolume-respiration relationship.



455 **Figure 8** Scaling of individual planktonic respiration rates as a function of estimated biovolume with respiration
 normalised either using species-specific Q_{10} s established in this study or a uniform Q_{10} of 3.18 from Lombard et
 al., 2009. Data includes previously published sources (Rink et al., 1998; Lombard et al., 2009 and Burke et al.,
 2025). Panels show normalisation to (a) 4 °C, (b) 15 °C and (c) 24 °C. Grey shaded regions represent the 95 %
 confidence intervals of the regression models. The source data for this figure is available in Table S12

460 **4. Discussion**

4.1 Respiration Rates and Environmental Stability

One of the most striking findings of this study is the metabolic stability of *N. pachyderma* across a wide range of
 ECVs and nutrients. Respiration rates in *N. pachyderma* are stable across pH (7.89 - 8.12), salinity (34.57 - 35.09
 psu), dissolved oxygen (2.31 - 2.66 ml L⁻¹), alkalinity (2185-2317 $\mu\text{mol kg}^{-1}$), and Ω_{Ca} (1.22 - 3.52) as well as
 465 nutrients (Fig. 6). This physiological resilience suggests that the polar genotype of *N. pachyderma* (Type I in
 Darling et al., 2004) evolved to adapt to the extreme and variable environment of the Arctic Ocean. We did not
 genotype the specimens in this study. However, Darling et al., (2004 and 2007) found only one genotype of *N.*
pachyderma (e.g., Type I) in the subpolar North Atlantic/Arctic Ocean, which was also confirmed by Bird et al.,

(2025). Only when exposed to temperatures outside of its habitat range in the laboratory (e.g., 13 °C), we recorded a small but significant increase in respiration rate.

The low thermal sensitivity of *N. pachyderma*, is reflected in a Q_{10} value of 1.41 (Table 4) across the 0.5 °C to 10 °C range (based on ship-based data) and suggests a successful physiological adaptation to cold environments. Low Q_{10} values have been interpreted as characteristic of the optimal temperature range of a species in its natural habitat (Wieser, 1973). This suggests that *N. pachyderma* (Type I) functions at its metabolic optimum in present-day Arctic conditions. Notably, respiration rates do not decline at the low temperatures measured, unlike in other species, which often exhibit constrained metabolic activity under extreme cold conditions.

Molecular evidence presented by Darling et al., (2004, 2007) suggests that the *N. pachyderma* population presently living at high northern latitudes became isolated during the onset of Northern Hemisphere Glaciations, between 1.8 and 1.5 million years ago, a period marked by the expansion of the polar ice sheets and the establishment of persistently cold oceanic conditions. Furthermore, Kucera and Kennett (2002) proposed that the species-specific cold-water affinity of *N. pachyderma* may have evolved in response to the onset of the 100,000-year glacial-interglacial climate cycles during the Middle Pleistocene. Furthermore, an increase in *N. pachyderma* shell sizes over the last 1.1 million years may represent an adaptive response to cold environments of the Quaternary (Huber et al., 2000). The current polar affinity of *N. pachyderma* may thus represent an evolutionary legacy of this climatic transition.

While we did not genotype *T. quinqueloba*, it is likely that our specimens belong to one of the two Arctic-associated genotypes, with type IIa occurring in both the subpolar Arctic and subpolar Antarctic, and type IIb restricted to the subpolar Arctic (Darling et al., 2000). *T. quinqueloba* displayed a high Q_{10} value of 4.54 in our experiments, indicating a strong temperature dependence that may influence its metabolic performance, growth, and survival (Mundim et al., 2020). This contrasts with the lower Q_{10} of 1.41 observed in *N. pachyderma*, highlighting clear interspecific differences in thermal sensitivity. *T. quinqueloba* is a spinose species, however its ecology and physiology differ markedly from symbiont-bearing tropical taxa for which elevated Q_{10} values have been reported previously (Lombard et al., 2009). Although preliminary, the relatively high Q_{10} observed for the non-spinose *N. incompta* suggests that spine presence or absence does not reliably predict thermal sensitivity. Instead, we posit that temperature responses are species-specific, reflecting distinct ecological strategies rather than broad morphological categories. Such species-level differences likely contribute to the latitudinal partitioning of planktonic foraminiferal assemblages, with taxa occupying thermal niches that reflect their individual physiological tolerances (Bé & Tolderlund, 1971; Ying et al., 2023).

Likewise, spinose planktonic foraminifera such as *T. quinqueloba* exhibit distinct evolutionary and physiological traits that reflect their adaptation to surface ocean environments. Their spines enhance prey capture and buoyancy and often support photosymbiotic relationships with algae (though no symbionts were seen in this study), enabling survival in oligotrophic waters (Anderson & Bé, 1976; Hemleben et al., 1989). In this study, *T. quinqueloba* displayed a high Q_{10} value of 4.54, indicating that this subpolar species exhibits a particularly strong temperature dependence, which is likely to impact its metabolism, potentially affecting its growth and overall survival (Mundim et al. 2020). This contrasts with the lower Q_{10} of 1.41 observed in the non-spinose, *N. pachyderma*, but is consistent with Q_{10} values around 3.18 derived for other spinose species such as *G. ruber*, *G. siphonifera*, and

~~*O. universa* (Lombard et al., 2009), suggesting divergent thermal strategies linked to morphology and trophic mode. The metabolic divergence observed here between spinose and non-spinose planktonic foraminifera likely contributes to the latitudinal partitioning of planktonic foraminiferal assemblages, with spinose taxa dominating tropical zones and non-spinose forms prevailing in subpolar and polar regions (Bé & Tolderlund, 1971; Ying et al., 2023). However, even within closely related species such as *N. incompta* and *N. pachyderma*, our observations suggest that the metabolic response to environmental conditions varies across foraminiferal taxa (Lombard et al., 2009).~~

4.2 Volume Scaling and Metabolic Allometry

While a positive log-log relationship between biomass, weight or volume and respiration is usually observed in micro/meiofauna (e.g. Fenchel and Finlay, 1983; Gerlach et al., 1985; Moens et al., 1999; Moodley et al., 2008), mixed results have been reported for this relationship in foraminifera. Some studies have found a positive correlation (Bradshaw, 1961; Geslin et al., 2011; Maciute et al., 2023), while others have not (Hannah et al., 1994; Nomaki et al., 2007). However, neither Hannah et al., (1994) nor Nomaki et al., (2007) use micro-CT scanning to estimate biovolumes. Nomaki et al., (2007) used the relationship between test size and organic content (Altenbach, 1985), while Hannah et al., (1994) used 75 % of the best-fitting geometric shape to estimate total internal test volume, a methodology with potential issues as suggested in section 3.1. above. Our results showed a positive linear correlation between respiration and biovolume (Fig. 8) across the polar and subpolar populations of *N. pachyderma*, *N. incompta*, and *T. quinqueloba*, consistent with pelagic foraminiferal species exclusively growing in warmer subtropical and tropical oceans (Burke et al., 2025; Lombard et al., 2009 and Rink et al., 1998). This supports earlier findings that body size is a significant determinant of metabolic rate in foraminifera (Rink et al., 1998; Geslin et al., 2011).

Furthermore, our analysis (Fig. 8) reveals a consistent and positive scaling relationship between biovolume and respiration rates across all temperature normalisations (4 °C, 15 °C, and 24 °C). This demonstrates that within a species, larger planktonic foraminifera exhibit higher metabolic rates, even when respiration is normalised to account for temperature effects using both species-specific and uniform Q₁₀ values (Table 4). Comparison of respiration-biovolume relationships for *N. pachyderma* and *N. incompta* in Fig. 8a (4 °C) and Fig. 8b (15 °C) indicates that *N. incompta* surpasses *N. pachyderma* at the higher temperature of 15 °C, showing a crossover between 4 °C and 15 °C that reflects the point at which the relative size-normalised respiration of the two species changes under temperature normalisation. This intersection suggests a potential physiological tipping point below 15 °C, where the metabolic efficiency of *N. pachyderma* begins to exceed that of *N. incompta*. This aligns with observed shifts in assemblage dominance across temperature gradients, with *N. pachyderma* prevailing in colder polar waters and *N. incompta* in subpolar to temperate regions (Bé and Hutson, 1977; Al-Sabouni et al., 2007; Husum and Hald, 2012; Chaabane et al., 2024). ~~These differences likely is crossover may~~ reflect species-specific thermal sensitivities ~~and that~~ could play a role in defining their biogeographic boundaries. It also highlights the value of using species-specific Q₁₀ values to resolve fine-scale metabolic differences that may underlie broader ecological patterns.

Our results are broadly consistent with Kleiber's Law, which predicts sublinear scaling of metabolic rate with body size (Kleiber, 1932). We observed positive, sublinear relationships between respiration and biovolume

545 across all temperature-normalised datasets, with slopes increasing from 0.39 at 5 °C to 0.69 at 24 °C. While
Kleiber’s canonical $\frac{3}{4}$ exponent is not fully met, the trend supports the principle that larger foraminifera have
higher, but less-than-proportional, metabolic rates. Interestingly, DeLong et al., (2010) showed that metabolic
scaling varies across evolutionary transitions—being super linear in prokaryotes, linear in protists, and sublinear
in metazoans. ~~Our findings place planktonic foraminifera, unicellular eukaryotes, closer to metazoan-like~~
550 ~~metabolic behaviour, possibly due to their structural complexity, calcification, or ecological specialisation. This~~
~~suggests that metabolic constraints in foraminifera may reflect both their unicellular nature and their functional~~
~~convergence with more complex organisms. The position of foraminiferal scaling within this framework should~~
~~be interpreted cautiously, as cross-study comparisons involve heterogeneous datasets and methodologies.~~
555 ~~However, the pattern raises the possibility that planktonic foraminifera may exhibit metabolic behaviour~~
~~intermediate between protists (which they are) and metazoans, potentially reflecting their large cell size that~~
~~overlaps with the size of small metazoans, structural complexity, calcification, or ecological specialisation. This~~
~~remains a hypothesis that warrants further targeted investigation.~~

4.3 Proxy Reliability

A central concern for proxy-based climate reconstructions is the potential impact of physiological processes on
560 shell geochemistry (Pérez-Huerta and Andrus, 2010). Variations in calcification and respiration rate in non-
spinose species can alter pH and carbonate chemistry in the foraminiferal microenvironment, potentially affecting
Mg/Ca, $\delta^{11}\text{B}$, $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values in (Wolf-Gladrow et al., 1999; Zeebe and Sanyal, 2002). However, our data
show that respiration in *N. pachyderma* remains stable across both temperature and pH gradients typically
565 encountered in its natural range (Fig. 5). Respiration rates are also consistent and stable across a large range of
other ECVs measured, suggesting that respiration is unlikely to introduce significant uncertainties into
geochemical-based palaeotemperature reconstructions for this species. These results also agree well with the
recent $\delta^{11}\text{B}$ - pH calibration for *N. pachyderma* that shows $\delta^{11}\text{B}$ values for this species are consistently offset from
seawater borate (de la Vega et al, 2025). Consequently, our findings alleviate concerns about physiological
confounding due to respiration. However, this may not be the case for *N. incompta* and *T. quinqueloba*. The
570 elevated Q_{10} values for these two species may call for the development of species-specific calibration equations
that consider the influence of respiration for accurate proxy application. ~~Whether genotype could have~~
~~implications for accurate proxy applications is dependent on the species with both *N. pachyderma* and *N. incompta*~~
~~only having one genotype (Type I) found in the Arctic/North Atlantic while *T. quinqueloba* has two genotypes~~
~~(Type IIa and Type IIb) (Darling et al., 2000; Darling et al., 2006).~~

4.4 Conclusions

By avoiding overestimation biases inherent in geometric models, micro-CT-derived biovolumes strengthen the
foundation for future studies on respiration in foraminifera. These methodological improvements enhance the
accuracy of single-specimen respiration rate assessments and facilitate more accurate interspecies comparisons.
Furthermore, our study demonstrates that there is no significant relationship between respiration and temperature
580 or other ECVs (salinity, pH, dissolved oxygen, alkalinity, and Ω_{Ca}) or nutrients (SiO_2 , DIC, PO_4^{3-} , and TON) in
N. pachyderma, suggesting this polar species has a strong thermal resistance and is well adapted to its unique
environment. On the other hand, *T. quinqueloba* and *N. incompta* exhibit a statistically significant relationship

between respiration and temperature, demonstrating large physiological diversity among planktonic foraminifera inhabiting overlapping climate zones. Additionally, we demonstrate a strong relationship between biovolume and respiration rate, underscoring the importance of size grading when using proxies. We conclude that for *N. pachyderma* respiration is unlikely to influence geochemical climate proxies.

Data and materials availability

All data needed to evaluate the conclusions in the paper are presented in the paper and/or the Supplementary Materials.

Author contributions

D.A. carried out respiration measurements at UiT and carried out micro-CT scanning at the University of Galway. In addition, she compiled and analysed all datasets and wrote the original draft of the manuscript. A.M. conceptualized the project, acquired funding, designed and carried out respiration experiments on board the Celtic Explorer (CE23011) as well as in the laboratory at UiT. In addition, she supervised D.A. in all data analysis and writing of the original draft. N.G. supported method development, experimental design, and respiration measurements during CE23011 and contributed to the final version of this manuscript. T.L.W. and E.D.V. performed Alkalinity and DIC measurements on board CE23011 and at the Marine Institute and contributed to the final version of this manuscript. In addition, T.L.W. supported D.A. with micro-CT scanning and analysis of images using AMIRA. M.M.E. was the chief scientist on the ARCLIM-24-1 cruise on the RV Helmer Hansen, supported the collection of live foraminifera, provided culturing facilities, and contributed to the final version of this manuscript. A.W., F.S. and A.F. supported the collection of live foraminifera during both expeditions and contributed to the final version of this manuscript. In addition, A.F. supported D.A. with micro-CT scanning and analysis of images using AMIRA. J.M. provided support in collection and culturing of live foraminifera during both expeditions and contributed to the final version of this manuscript. T.L.B. contributed to discussions on the initial conceptualization of the experimental design and method development of the nano-respiration system.

Competing interests

The contact author has declared that neither they nor their co-authors have any competing interests.

Funding Sources

A.M. acknowledges funding by Research Ireland and the Geological Survey of Ireland under the SFI Frontiers for the Future Programme 21/FFP-P/10261 and Grant in Aid funding from the Marine Institute for research expedition CE23011 on the RV Celtic Explorer. N.G. acknowledges funding from the Deutsche Forschungsgesellschaft (DFG) under grant number GL 999/3-1. M.M.E, F.S and A.W. acknowledge funding

from [the ARCLIM project](#), a Tromsø Research Foundation (TFS) starting grant project, with grant number: A31720.

Acknowledgements

620 We gratefully acknowledge the support of the crew on the RV Celtic Explorer sailing under Master Anthony Hobin [and the crew of the R/V Helmer Hanssen.](#) -

5. References:

625 Al-Sabouni, N., Kucera, M., and Schmidt, D. N.: Vertical niche separation control of diversity and size disparity in planktonic foraminifera, *Marine Micropaleontology*, 63, 75–90, <https://doi.org/10.1016/j.marmicro.2006.11.002>, 2007.

Altenbach, A. V.: Die biomasse der benthischen foraminiferen: auswertungen von "Meteor"-expeditionen im östlichen nordatlantik, PhD Thesis, Christian-Albrechts-Universität, 1985.

630 Anderson, O. R., Spindler, M., Bé, A. W. H., and Hemleben, Ch.: Trophic activity of planktonic foraminifera, *J. Mar. Biol. Ass.*, 59, 791–799, <https://doi.org/10.1017/S002531540004577X>, 1979.

Anglada-Ortiz, G., Rasmussen, T. L., Chierici, M., Fransson, A., Ziveri, P., Thomsen, E., Zamelczyk, K., Meilland, J., Ezat, M. M., and Garcia-Orellana, J.: Changes in Planktic Foraminiferal Distribution, Productivity, and Preservation in the Barents Sea During the Last Three Millennia, *Paleoceanography and Paleoclimatology*, 40, e2024PA004989, <https://doi.org/10.1029/2024PA004989>, 2025.

635 Baum, D. and Titschack, J.: Cavity and Pore Segmentation in 3D Images with Ambient Occlusion, EuroVis '16: Proceedings of the Eurographics / IEEE VGTC Conference on Visualization: Short Papers, 113-117, <https://dl.acm.org/doi/abs/10.5555/3058878.3058902>, 2016

Bé, A. W. and Hutson, W. H.: Ecology of planktonic foraminifera and biogeographic patterns of life and fossil assemblages in the Indian Ocean, *Micropaleontology*, 369–414, <https://doi.org/10.2307/1485406>, 1977.

640 Bé, A.W. and Tolderlund, D.S. 1971.: Distribution and ecology of living planktonic foraminifera in surface waters of the Atlantic and Indian Oceans, in: *The Micropaleontology of the Oceans*, edits by: Funnel, B.M. and Riedel, W.R., Cambridge University Press, London, 105-149, 1971.

645 Bertlich, J., Gussone, N., Berndt, J., Arlinghaus, H. F., and Dieckmann, G. S.: Salinity effects on cultured *Neogloboquadrina pachyderma* (sinistral) from high latitudes: new paleoenvironmental insights, *Geo-Mar Lett.*, 41, 2, <https://doi.org/10.1007/s00367-020-00677-1>, 2021.

[Bird, C., Darling, K., Thiessen, R., & Pieńkowski, A. J.: The microbiome of the Arctic planktonic foraminifera *Neogloboquadrina pachyderma* is composed of fermenting and carbohydrate-degrading bacteria and an intracellular diatom chloroplast store, *Biogeosciences*, 22\(17\), 4545-4577. <https://doi.org/10.5194/bg-22-4545->](#)

[2025](#), 2025

650 Bradshaw, J. S.: Laboratory experiments on the ecology of foraminifera., *Cushman Found Foram. Res., Contr.*, 12, 87–106, 1961.

Broecker, W. S. and Peng, T.-H.: Gas exchange rates between air and sea, *Tellus A: Dynamic Meteorology and Oceanography*, 26, 21-35, <https://doi.org/10.3402/tellusa.v26i1-2.9733>, 1974.

655 Burke, J. E., Renema, W., Schiebel, R., and Hull, P. M.: Three-dimensional analysis of inter-and intraspecific variation in ontogenetic growth trajectories of planktonic foraminifera, *Marine Micropaleontology*, 155, 101794, <https://doi.org/10.1016/j.marmicro.2019.101794>, 2020.

Burke, J. E., Elder, L. E., Maas, A. E., Gaskell, D. E., Clark, E. G., Hsiang, A. Y., Foster, G. L., and Hull, P. M.: Physiological and morphological scaling enables gigantism in pelagic protists, *Limnology and Oceanography*, 660 70, 461-476, <https://doi.org/10.1002/lno.12770>, 2025.

Cesbron, F., Geslin, E., Jorissen, F. J., Delgard, M. L., Charrieau, L., Deflandre, B., Jézéquel, D., Anschutz, P., and Metzger, E.: Vertical distribution and respiration rates of benthic foraminifera: Contribution to aerobic remineralization in intertidal mudflats covered by *Zostera noltei* meadows, *Estuarine, Coastal and Shelf Science*, 179, 23–38, <https://doi.org/10.1016/j.ecss.2015.12.005>, 2016.

Chaabane, S., de Garidel-Thoron, T., Meilland, J., Sulpis, O., Chalk, T. B., Brummer, G.-J. A., Mortyn, P. G., Giraud, X., Howa, H., Casajus, N., Kuroyanagi, A., Beaugrand, G., and Schiebel, R.: Migrating is not enough for modern planktonic foraminifera in a changing ocean, *Nature*, 636, 390–396, <https://doi.org/10.1038/s41586-024-08191-5>, 2024.

670 Coletti, G., Stainbank, S., Fabbrini, A., Spezzaferri, S., Foubert, A., Kroon, D., and Betzler, C.: Biostratigraphy of large benthic foraminifera from Hole U1468A (Maldives): a CT-scan taxonomic approach, *Swiss J Geosci*, 111, 523–536, <https://doi.org/10.1007/s00015-018-0306-7>, 2018.

[Darling, K. F., Wade, C. M., Stewart, I. A., Kroon, D., Dingle, R., and Brown, A. J. L.: Molecular evidence for genetic mixing of Arctic and Antarctic subpolar populations of planktonic foraminifers, *Nature*, 405, 43–47, <https://doi.org/10.1038/35011002>, 2000.](#)

675 Darling, K. F., Kucera, M., Pudsey, C. J., and Wade, C. M.: Molecular evidence links cryptic diversification in polar planktonic protists to Quaternary climate dynamics, *Proceedings of the National Academy of Sciences*, 101, 7657–7662, <https://doi.org/10.1073/pnas.0402401101>, 2004.

Darling, K. F., Kucera, M., Kroon, D., and Wade, C. M.: A resolution for the coiling direction paradox in *Neogloboquadrina pachyderma*, *Paleoceanography*, 21, PA2011, <https://doi.org/10.1029/2005PA001189>, 2006.

- Darling, K. F., Kucera, M., and Wade, C. M.: Global molecular phylogeography reveals persistent Arctic circumpolar isolation in a marine planktonic protist, *Proc. Natl. Acad. Sci. U.S.A.*, 104, 5002–5007, <https://doi.org/10.1073/pnas.0700520104>, 2007.
- 685 Davis, C. V., Rivest, E. B., Hill, T. M., Gaylord, B., Russell, A. D., and Sanford, E.: Ocean acidification compromises a planktic calcifier with implications for global carbon cycling, *Scientific reports*, 7, 2225, <https://doi.org/10.1038/s41598-017-01530-9>, 2017.
- 690 de la Vega, E., Raitzsch, M., Foster, G., Bijma, J., Ninnemann, U.S., Kucera, M., Babila, T.L., Crumpton Banks, J., Ezat, M.M. and Morley, A., 2025. A $\delta^{11}\text{B}$ -pH calibration for the high-latitude foraminifera species *Neogloboquadrina pachyderma* and *Neogloboquadrina incompta*. *EGUsphere*, 2025, pp.1-32, <https://doi.org/10.5194/egusphere-2025-2443>
- 695 de Nooijer, L. J., Spero, H. J., Erez, J., Bijma, J., and Reichart, G. J.: Biomineralization in perforate foraminifera, *Earth-Science Reviews*, 135, 48–58, <https://doi.org/10.1016/j.earscirev.2014.03.013>, 2014.
- DeLong, J. P., Okie, J. G., Moses, M. E., Sibly, R. M., and Brown, J. H.: Shifts in metabolic scaling, production, and efficiency across major evolutionary transitions of life, *Proceedings of the National Academy of Sciences*, 107, 12941–12945, <https://doi.org/10.1073/pnas.1007783107>, 2010.
- 700 Dickson, A. G. and Riley, J. P.: The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water — K_w , *Marine Chemistry*, 7, 89–99, [https://doi.org/10.1016/0304-4203\(79\)90001-X](https://doi.org/10.1016/0304-4203(79)90001-X), 1979.
- Eggins, S. M., Sadekov, A., and De Deckker, P.: Modulation and daily banding of Mg/Ca in *Orbulina universa* tests by symbiont photosynthesis and respiration: a complication for seawater thermometry?, *Earth and Planetary Science Letters*, 225, 411–419, <https://doi.org/10.1016/j.epsl.2004.06.019>, 2004.
- 705 El Bani Altuna, N., Pieńkowski, A. J., Eynaud, F., and Thiessen, R.: The morphotypes of *Neogloboquadrina pachyderma*: Isotopic signature and distribution patterns in the Canadian Arctic Archipelago and adjacent regions, *Marine Micropaleontology*, 142, 13–24, <https://doi.org/10.1016/j.marmicro.2018.05.004>, 2018.
- Fabbrini, A., Pearson, P. N., Brombacher, A., Iacoviello, F., Ezard, T. H., and Wade, B. S.: Morphology of *Pulleniatina* (planktonic foraminifera) from optical microscopy, micro-CT, and SEM investigations. *Journal of Micropalaeontology*, 44, 213-235, <https://doi.org/10.5194/jm-44-213-2025>, 2025.
- 710 Fabbrini, A., Meilland, J., Westgård, A., Sykes, F.E., Ezat, M.M., and Morley, A.: Extensive culturing experiment on *Turborotalita quinqueloba* (spinose planktonic foraminifera). *Journal of Plankton Research*, *in review*.
- Fenchel, T. and Finlay, B. J.: Respiration rates in heterotrophic, free-living protozoa, *Microbial Ecology*, 9, 99–122, <https://doi.org/10.1007/BF02015125>, 1983.

[Gaskell, D. E., Ohman, M. D., & Hull, P. M.: Zooglider-based measurements of planktonic foraminifera in the California Current System. *Journal of Foraminiferal Research*, 49\(4\), 390-404, <https://doi.org/10.2113/gsjfr.49.4.390>, 2019](#)

720 Gerlach, S. A., Hahn, A. E., and Schrage, M.: Size spectra of benthic biomass and metabolism, *Marine Ecology Progress Series*, 26, 161–173, <http://www.jstor.org/stable/24817617>, 1985.

Geslin, E., Risgaard-Petersen, N., Lombard, F., Metzger, E., Langlet, D., and Jorissen, F.: Oxygen respiration rates of benthic foraminifera as measured with oxygen microsensors, *Journal of Experimental Marine Biology and Ecology*, 396, 108–114, <https://doi.org/10.1016/j.jembe.2010.10.011>, 2011.

725 Glock, N., Romero, D., Roy, A. S., Woehle, C., Dale, A. W., Schönfeld, J., Wein, T., Weissenbach, J., and Dagan, T.: A hidden sedimentary phosphate pool inside benthic foraminifera from the Peruvian upwelling region might nucleate phosphogenesis, *Geochimica et Cosmochimica Acta*, 289, 14–32, <https://doi.org/10.1016/j.gca.2020.08.002>, 2020.

730 Glock, N., Richirt, J., Woehle, C., Algar, C., Armstrong, M., Eichner, D., Firrincieli, H., Makabe, A., Govindankutty Menon, A., Ishitani, Y., Hackl, T., Hubert-Huard, R., Kienast, M., Milker, Y., Mutzberg, A., Ni, S., Okada, S., Rakshit, S., Schmiedl, G., Steiner, Z., Tame, A., Zhang, Z., and Nomaki, H.: Widespread occurrence and relevance of phosphate storage in foraminifera, *Nature*, 638, 1000–1006, <https://doi.org/10.1038/s41586-024-08431-8>, 2025.

735 Greco, M., Jonkers, L., Kretschmer, K., Bijma, J., and Kucera, M.: Depth habitat of the planktonic foraminifera *Neogloboquadrina pachyderma* in the northern high latitudes explained by sea-ice and chlorophyll concentrations, *Biogeosciences*, 16, 3425–3437, <https://doi.org/10.5194/bg-16-3425-2019>, 2019.

Greco, M., Meilland, J., Zamelczyk, K., Rasmussen, T. L., and Kučera, M.: The effect of an experimental decrease in salinity on the viability of the Subarctic planktonic foraminifera *Neogloboquadrina incompta*, *Polar Research*, 39, <https://doi.org/10.33265/polar.v39.3842>, 2020.

740 Hannah, F., Rogerson, R., and Laybourn-Parry, J.: Respiration rates and biovolumes of common benthic Foraminifera (Protozoa), *J. Mar. Biol. Ass.*, 74, 301–312, <https://doi.org/10.1017/S0025315400039345>, 1994.

Hemleben, C., Spindler, M., and Anderson, O. R.: Taxonomy and Species Features, in: *Modern Planktonic Foraminifera*, edited by: Hemleben, C., Spindler, M., and Anderson, O. R., Springer, New York, NY, 8–32, https://doi.org/10.1007/978-1-4612-3544-6_2, 1989.

745 Hoogakker, B. A. A., Anderson, C., Paoloni, T., Stott, A., Grant, H., Keenan, P., Mahaffey, C., Blackbird, S., McClymont, E. L., Rickaby, R., Poulton, A., and Peck, V. L.: Planktonic foraminifera organic carbon isotopes as archives of upper ocean carbon cycling, *Nat Commun*, 13, 4841, <https://doi.org/10.1038/s41467-022-32480-0>, 2022.

Huber, R., Meggers, H., Baumann, K.-H., Raymo, M. E., and Henrich, R.: Shell size variation of the planktonic

- 750 foraminifer *Neogloboquadrina pachyderma* sin. in the Norwegian–Greenland Sea during the last 1.3 Myrs: implications for paleoceanographic reconstructions, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 160, 193–212, [https://doi.org/10.1016/S0031-0182\(00\)00066-3](https://doi.org/10.1016/S0031-0182(00)00066-3), 2000.
- Husum, K. and Hald, M.: Arctic planktic foraminiferal assemblages: Implications for subsurface temperature reconstructions, *Marine Micropaleontology*, 96–97, 38–47, <https://doi.org/10.1016/j.marmicro.2012.07.001>,
755 2012.
- Kanbur, S.: Neglected small foraminifers of the Eastern Mediterranean and their paleoecological importance, *Journal of Foraminiferal Research*, 55, 218–232, <https://doi.org/10.61551/gsjfr.55.2.218>, 2025.
- Kiss, P., Hudáčková, N., Titschack, J., Siccha, M. G. R., Heřmanová, Z., Silye, L., Ruman, A., Rybár, S., and
760 Kučera, M.: Convergent evolution of spherical shells in Miocene planktonic foraminifera documents the parallel emergence of a complex character in response to environmental forcing, *Paleobiology*, 49, 454–470, <https://doi.org/10.1017/pab.2022.48>, 2023.
- Kleiber, M.: Body size and metabolism, *Hilgardia*, 6, 315, 1932.
- 765 Köhler-Rink, S. and Kühl, M.: Microsensor studies of photosynthesis and respiration in the larger symbiont bearing foraminifera *Amphistegina lobifera*, and *Amphisorus hemprichii*, *Ophelia*, 55, 111–122, <https://doi.org/10.1080/00785236.2001.10409478>, 2001.
- Kucera, M. and Kennett, J. P.: Causes and consequences of a middle Pleistocene origin of the modern planktonic foraminifer *Neogloboquadrina pachyderma* sinistral, *Geology*, 30, 539–542,
770 [https://doi.org/10.1130/0091-7613\(2002\)030%3C0539:CACOAM%3E2.0.CO;2](https://doi.org/10.1130/0091-7613(2002)030%3C0539:CACOAM%3E2.0.CO;2). 2002.
- Lewis, E. R. and Wallace, D. W. R.: Program developed for CO₂ system calculations, Environmental System Science Data Infrastructure for a Virtual Ecosystem, <https://doi.org/10.15485/1464255>, 1998.
- Lombard, F., Erez, J., Michel, E., and Labeyrie, L.: Temperature effect on respiration and photosynthesis of the symbiont-bearing planktonic foraminifera *Globigerinoides ruber*, *Orbulina universa*, and *Globigerinella*
775 *siphonifera*, *Limnology and Oceanography*, 54, 210–218, <https://doi.org/10.4319/lo.2009.54.1.0210>, 2009.
- Lopes, A. S., Larsen, L. H., Ramsing, N., Løvendahl, P., Raty, M., Peippo, J., Greve, T., and Callesen, H.: Respiration rates of individual bovine in vitro-produced embryos measured with a novel, non-invasive and highly sensitive microsensor system, *Reproduction*, 130, 669–679, <https://doi.org/10.1530/rep.1.00703>, 2005.
- Lueker, T. J., Dickson, A. G., and Keeling, C. D.: Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K₁ and K₂: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium, *Marine Chemistry*, 70, 105–119, [https://doi.org/10.1016/S0304-4203\(00\)00022-0](https://doi.org/10.1016/S0304-4203(00)00022-0),
780 2000.
- Maciute, A., Holovachov, O., Glud, R. N., Broman, E., Berg, P., Nascimento, F. J. A., and Bonaglia, S.:

- 785 Reconciling the importance of meiofauna respiration for oxygen demand in muddy coastal sediments, *Limnology and Oceanography*, 68, 1895–1905, <https://doi.org/10.1002/lno.12393>, 2023.
- Manno, C., Morata, N., and Bellerby, R.: Effect of ocean acidification and temperature increase on the planktonic foraminifer *Neogloboquadrina pachyderma* (sinistral), *Polar Biol*, 35, 1311–1319, <https://doi.org/10.1007/s00300-012-1174-7>, 2012.
- 790 Meilland, J., Howa, H., Hulot, V., Demangel, I., Salaün, J., and Garlan, T.: Population dynamics of modern planktonic foraminifera in the western Barents Sea, *Biogeosciences*, <https://doi.org/10.5194/bg-2019-429>, 2019.
- Moens, T., Verbeeck, L., and Vincx, M.: Feeding biology of a predatory and a facultatively predatory nematode (*Enoploides longispiculosus* and *Adoncholaimus fuscus*), *Marine Biology*, 134, 585–593, <https://doi.org/10.1007/s002270050573>, 1999.
- 795 Moodley, L., Steyaert, M., Epping, E., Middelburg, J.J., Vincx, M., van Avesaath, P., Moens, T. and Soetaert, K.: Biomass-specific respiration rates of benthic meiofauna: Demonstrating a novel oxygen micro-respiration system, *Journal of Experimental Marine Biology and Ecology*, 357, 41–47, <https://doi.org/10.1016/j.jembe.2007.12.025>, 2008.
- Motic Instruments Inc: <https://moticeurope.com/en/moticam-x5-plus.html>, 2024.
- 800 Mundim, K.C., Baraldi, S., Machado, H.G. and Vieira, F.M. Temperature coefficient (Q_{10}) and its applications in biological systems: Beyond the Arrhenius theory. *Ecol Model.*, 431, 109127. <https://doi.org/10.1016/j.ecolmodel.2020.109127>, 2020.
- 805 Natland, M. L.: New species of foraminifera from off the West Coast of North America and from the later Tertiary of the Los Angeles basin, University of California Press, New species of foraminifera from off the West Coast of North America and from the later Tertiary of the Los Angeles basin | CiNii Research, 1938.
- Nielsen, P., Larsen, L. H., Ramløv, H., and Hansen, B. W.: Respiration rates of subitaneous eggs from a marine calanoid copepod: monitored by nanorespirometry, *Journal of Comparative Physiology B*, 177, 287–296, <https://doi.org/10.1007/s00360-006-0128-1>, 2007.
- 810 Nomaki, H., Yamaoka, A., Shirayama, Y., and Kitazato, H.: Deep-sea benthic foraminiferal respiration rates measured under laboratory conditions, *Journal of Foraminiferal Research*, 37, 281–286, <https://doi.org/10.2113/gsjfr.37.4.281>, 2007.
- 815 Pearson, P.N. and Kucera, M. Taxonomy, biostratigraphy, and phylogeny of Oligocene Turborotalita, in Wade, B.S., Olsson, R.K., Pearson, P.N., Huber, B.T. and Berggren, W.A. (eds.), *Atlas of Oligocene Planktonic Foraminifera*, Cushman Foundation of Foraminiferal Research, Special Publication, No. 46: https://www.ucl.ac.uk/earth-sciences/sites/earth_sciences/files/Chapter_12.pdf, 2018.

- Pérez-Huerta, A. and Andrus, C. F. T.: Vital effects in the context of biomineralization, *Seminarios de la Sociedad de Española de Mineralogía*, 7, 35-45, ir-api.ua.edu/api/core/bitstreams/efc0bd9b-7204-464d-84b7-454793f7c34a/content, 2010.
- 820
- Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Commun Earth Environ*, 3, 168, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- 825
- Revsbech, N. P.: An oxygen microsensor with a guard cathode, *Limnology and Oceanography*, 34, 474–478, <https://doi.org/10.4319/lo.1989.34.2.0474>, 1989.
- Rink, S., Kühl, M., Bijma, J., and Spero, H. J.: Microsensor studies of photosynthesis and respiration in the symbiotic foraminifer *Orbulina universa*, *Marine Biology*, 131, 583–595, <https://doi.org/10.1007/s002270050350>, 1998.
- 830
- Schiebel, R., Waniek, J., Bork, M., and Hemleben, C.: Planktic foraminiferal production stimulated by chlorophyll redistribution and entrainment of nutrients, *Deep Sea Research Part I: Oceanographic Research Papers*, 48, 721–740, [https://doi.org/10.1016/S0967-0637\(00\)00065-0](https://doi.org/10.1016/S0967-0637(00)00065-0), 2001.
- Schiebel, R. and Hemleben, C.: *Planktic Foraminifers in the Modern Ocean*, Springer Berlin / Heidelberg, Berlin, Heidelberg, Germany, 2017.
- 835
- Schlitzer, R.: Ocean data view, <https://odv.awi.de>, last access: 1 September 2025, 2022.
- Schmidt-Nielsen, K.: *Animal Physiology: Adaptation and Environment*, Cambridge University Press, 630 pp., 1997.
- 840
- [Sears, H. A., Darling, K. F., and Wade, C. M.: Ecological partitioning and diversity in tropical planktonic foraminifera, *BMC Evolutionary Biology*, 12, 54, <https://doi.org/10.1186/1471-2148-12-54>, 2012.](#)
- Serreze, M. C. and Barry, R. G.: Processes and impacts of Arctic amplification: A research synthesis, *Global and Planetary Change*, 77, 85–96, <https://doi.org/10.1016/j.gloplacha.2011.03.004>, 2011.
- 845
- Siccha, M., Morard, R., Meilland, J., Iwasaki, S., Kucera, M., and Kimoto, K.: Collection of X-ray micro computed tomography images of shells of planktic foraminifera with curated taxonomy, *Sci Data*, 10, 679, <https://doi.org/10.1038/s41597-023-02498-0>, 2023.
- 850
- Spero, H. J. and Lea, D. W.: Intraspecific stable isotope variability in the planktic foraminifera *Globigerinoides sacculifer*: Results from laboratory experiments, *Marine Micropaleontology*, 22, 221–234, [https://doi.org/10.1016/0377-8398\(93\)90045-Y](https://doi.org/10.1016/0377-8398(93)90045-Y), 1993.
- Spero, H. J. and Lea, D. W.: Experimental determination of stable isotope variability in *Globigerina bulloides*:

- implications for paleoceanographic reconstructions, *Marine Micropaleontology*, 28, 231–246,
855 [https://doi.org/10.1016/0377-8398\(96\)00003-5](https://doi.org/10.1016/0377-8398(96)00003-5), 1996.
- Spindler M.: On the salinity tolerance of the planktonic foraminifer *Neogloboquadrina pachyderma* from the Antarctic Sea Ice (17th Symposium on Polar Biology), *Proceedings of the NIPR Symposium on Polar Biology*, 9, 85–91, <https://doi.org/10.15094/00005306>, 1996.
- Spindler, M. and Dieckmann, G. S.: Distribution and abundance of the planktic foraminifer *Neogloboquadrina*
860 *pachyderma* in sea ice of the Weddell Sea (Antarctica), *Polar Biol*, 5, 185–191,
<https://doi.org/10.1007/BF00441699>, 1986.
- Stalling, D., Westerhoff, M., and Hege, H.-C.: Amira: A Highly Interactive System for Visual Data Analysis, in: *Visualization Handbook*, Elsevier, 749–767, <https://doi.org/10.1016/B978-012387582-2/50040-X>, 2005.
- Stangeew, E.: Distribution and Isotopic Composition of Living Planktonic Foraminifera *N. pachyderma*
865 (sinistral) and *T. quinqueloba* in the High Latitude North Atlantic, Kiel, University, Diss., 2001.,
https://macau.uni-kiel.de/servlets/MCRFileNodeServlet/dissertation_derivate_00000464/d464.pdf
- Takagi, H., Kimoto, K., Fujiki, T., Saito, H., Schmidt, C., Kucera, M., and Moriya, K.: Characterizing
870 photosymbiosis in modern planktonic foraminifera, *Biogeosciences*, 16, 3377–3396, <https://doi.org/10.5194/bg-16-3377-2019>, 2019.
- Tell, F., Jonkers, L., Meilland, J., and Kucera, M.: Upper-ocean flux of biogenic calcite produced by the Arctic planktonic foraminifera *Neogloboquadrina pachyderma*, *Biogeosciences*, 19, 4903–4927,
<https://doi.org/10.5194/bg-19-4903-2022>, 2022.
875
- Titschack, J., Baum, D., Matsuyama, K., Boos, K., Färber, C., Kahl, W.-A., Ehrig, K., Meinel, D., Soriano, C., and Stock, S. R.: Ambient occlusion – A powerful algorithm to segment shell and skeletal intrapores in computed tomography data, *Computers & Geosciences*, 115, 75–87,
<https://doi.org/10.1016/j.cageo.2018.03.007>, 2018.
880
- Unisense.: Nanorespiration system manual. <https://unisense.com/products/nanorespiration-system/>, 2025.
- Uppström, Leif R.: The boron/chlorinity ratio of deep-sea water from the Pacific Ocean, *Deep Sea Research A* 21, no. 2 (1974): 161-162. [https://doi.org/10.1016/0011-7471\(74\)90074-6](https://doi.org/10.1016/0011-7471(74)90074-6)
- Volkman, R.: Planktic foraminifera in the outer Laptev Sea and the Fram Strait- Modern distribution and
885 ecology, *Journal of Foraminiferal Research*, 30, 157–176, <https://doi.org/10.2113/0300157>, 2000.
- Westgård, A., Ezat, M. M., Chalk, T. B., Chierici, M., Foster, G. L., and Meilland, J.: Large-scale culturing of *Neogloboquadrina pachyderma*, its growth in, and tolerance of, variable environmental conditions, *Journal of Plankton Research*, 45, 732–745, <https://doi.org/10.1093/plankt/fbad034>, 2023.

- 890 Wieser, W.: Temperature Relations of Ectotherms: A Speculative Review, in: Effects of Temperature on Ectothermic Organisms, edited by: Wieser, W., Springer Berlin Heidelberg, Berlin, Heidelberg, 1–23, https://doi.org/10.1007/978-3-642-65703-0_1, 1973.
- Woelfel, J., Sørensen, K., Warkentin, M., Forster, S., Oren, A., and Schumann, R.: Oxygen evolution in a hypersaline crust: in situ photosynthesis quantification by microelectrode profiling and use of planar optode spots in incubation chambers, *Aquat. Microb. Ecol.*, 56, 263–273, <https://doi.org/10.3354/ame01326>, 2009.
- 895
- Wolf-Gladrow, D. A., Bijma, J., and Zeebe, R. E.: Model simulation of the carbonate chemistry in the microenvironment of symbiont bearing foraminifera, *Marine Chemistry*, 64, 181–198, [https://doi.org/10.1016/S0304-4203\(98\)00074-7](https://doi.org/10.1016/S0304-4203(98)00074-7), 1999.
- Ying, R., Monteiro, F. M., Wilson, J. D., and Schmidt, D. N.: ForamEcoGENIE 2.0: incorporating symbiosis and spine traits into a trait-based global planktic foraminiferal model, *Geosci. Model Dev.*, 16, 813–832, <https://doi.org/10.5194/gmd-16-813-2023>, 2023.
- 900
- Zamelczyk, K., Fransson, A., Chierici, M., Jones, E., Meilland, J., Anglada-Ortiz, G., and Lødemel, H. H.: Distribution and Abundances of Planktic Foraminifera and Shelled Pteropods During the Polar Night in the Sea-Ice Covered Northern Barents Sea, *Front. Mar. Sci.*, 8, <https://doi.org/10.3389/fmars.2021.644094>, 2021.
- 905
- Zeebe, R. E. and Sanyal, A.: Comparison of two potential strategies of planktonic foraminifera for house building: Mg²⁺ or H⁺ removal?, *Geochimica et Cosmochimica Acta*, 66, 1159–1169, [https://doi.org/10.1016/S0016-7037\(01\)00852-3](https://doi.org/10.1016/S0016-7037(01)00852-3), 2002.
- 910
- ZEISS. Xradia VERSA 620 User Manual. www.zeiss.com. 2024.