

Spatial heterogeneity of GHG dynamics across an estuarine ecosystem

Nicolas-Xavier Geilfus¹, Bruno Delille², Anna Villnäs¹, Alf Norkko¹

¹Tvärminne Zoological Station, University of Helsinki, Finland

²Chemical Oceanography Unit, Université de Liège, Belgium

5 *Correspondence to:* Nicolas-Xavier Geilfus (nicolas-xavier.geilfus@helsinki.fi)

Abstract.

Coastal ecosystems are critical components of the global carbon cycle, exerting a disproportionate influence on the carbon budget despite their limited spatial extent. ~~Estuaries remain understudied despite being dynamic~~ Although they have gained more attention in the past decade, detailed studies of GHG dynamics across the land-ocean continuum, including shallow-
10 water estuaries, remain relatively scarce even though they are active sources of the three most potent greenhouse gases (GHGs): carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O). Such shallow coastal ecosystems are highly heterogeneous, shaped by strong physical, biogeochemical, and biological gradients. ~~Combined Together~~ with the spatial variability variation in coastal biodiversity, these gradients ~~strongly shapes~~ significantly influence carbon cycling at both local and global ~~scales~~ levels. However, large uncertainties persist due to limited ~~and spatially patchy~~ measurements, highlighting ~~emphasizing~~ the need for improved constraints on GHG budgets ~~and their sensitivity to biodiversity loss and climate change~~.
15 ~~We measured surface~~ Surface seawater partial pressure of CO₂ (pCO₂), CH₄, and N₂O concentrations, along with seawater physical and biogeochemical properties, and air-sea gas exchange, were measured at 21 sites in southwest Finland (Baltic Sea). Sampling ~~followed a transition~~ progressed from estuarine inner bays to the outer archipelago, covering diverse soft-sediment habitats, from sheltered to exposed areas, across a salinity gradient. ~~Surface water~~ Seawater pCO₂ and N₂O
20 ~~concentration~~ concentrations ranged from undersaturated (160 ppm and 9 nmol L⁻¹, respectively) to supersaturated (2521 ppm and 25 nmol L⁻¹, respectively), compared to the atmosphere, resulting in an uptake of -36 and -0.0021 mmol m⁻² d⁻¹, and a release up to 220 and 0.0383 mmol m⁻² d⁻¹, respectively. CH₄ concentrations were consistently supersaturated (19 to 469 nmol L⁻¹) compared to the atmosphere, resulting in a net source to the atmosphere from 0.014 to 1.39 mmol m⁻² d⁻¹.
Freshwater input from the Karjaanjoki River and its mixing with seawater ~~mainly determined~~ shaped the overall spatial patterns
25 of GHGs. However, deviations from this salinity-driven control were ~~observed. In~~ seen in sheltered sites within the archipelago, where elevated pCO₂, ~~CH₄~~, and ~~N₂O~~ CH₄ concentrations likely reflected ~~benthic~~ biological processes, including enhanced organic matter respiration and methanogenesis in warm, late-summer shallow waters, where limited oxidation favoured CH₄ accumulation. At exposed and semi-sheltered sites, mixing processes ~~had a stronger~~ exerted greater control, resulting in lower GHG concentrations. Our results show that both physical mixing and ~~benthic~~ biological processes influence coastal GHG
30 dynamics, with benthic ecosystems potentially playing a key but still poorly constrained role. ~~Air~~ The overall budget of air- sea GHG exchanges ~~were~~ was dominated by CO₂, while fluxes, with CH₄ and N₂O ~~contributed differently~~ consistently acting

as a source, and a N_2O alternating between source and sink. The High environmental variability in shallow coastal systems leads to strong fluctuations in the balance between GHG production and consumption processes, particularly within benthic habitats, is therefore critical for understanding coastal contributions to, which needs to be considered when evaluating their role in the global carbon budget.

1. Introduction

Coastal ecosystems are increasingly recognized as critical components of the global carbon cycle (Bauer et al., 2013). Although covering only about 7% of the oceanic area, they play a disproportionately large role in biogeochemical cycling due to their location at the land–sea interface (Gattuso et al., 1998; Wollast, 1998). These environments act as hubs of exchange across biomes (Resplandy et al., 2024), contributing to the global oceanic uptake of anthropogenic carbon by absorbing carbon dioxide (CO_2) from the atmosphere and either burying, transforming, or releasing the carbon that land ecosystems deliver to coastal waters (Regnier et al., 2022). Due to intense inputs of nutrients and carbon from land and open ocean at continental margins, combined with high rates of biological production and degradation, coastal waters are among the most biogeochemically active regions of the biosphere (Wollast, 1998). Shallow coastal waters such as mangroves, salt marshes, and seagrass meadows effectively sequester large amounts of atmospheric carbon through vegetation growth and long-term sediment accumulation, making significant contributions to long-term carbon storage (Lovelock and Duarte, 2019; Macreadie et al., 2021; Mcleod et al., 2011). These so-called “blue carbon” ecosystems have been proposed as nature-based solutions for short-term climate change mitigation (Hoegh-Guldberg et al., 2019; Lovelock and Duarte, 2019).

Vegetated blue carbon ecosystems exemplify the value of healthy habitats in long-term carbon storage. Still, they cover a small part of the coastal oceans (thebluecarboninitiative.org), and measurements from other key habitats remain scarce and largely neglected in carbon cycling budgets (James et al., 2024). Estuarine areas are characterized by strong physical and chemical gradients and host transitional biodiversity that bridges terrestrial and marine ecosystems (Elliott and Whitfield, 2011) and are recognized as hot spots for carbon cycling and GHG exchange with the atmosphere (Borges et al., 2005; 2016; Cai et al., 2013; Frankignoulle et al., 1998; Humborg et al., 2019; Rosentreter et al., 2021, 2021a). The global oceans absorb roughly 25% of anthropogenic CO_2 emissions annually (Watson et al., 2020). While coastal waters contribute to this sink by taking up atmospheric CO_2 (Chen and Borges, 2009; Regnier et al., 2022), many nearshore systems like estuaries have been recognized as net heterotrophic ecosystems with negative net ecosystem production (Gattuso et al., 1998; Testa et al., 2012), leading to the production and release of both CO_2 and methane (CH_4) (Abril and Borges, 2005; Bonaglia et al., 2025; Borges et al., 2018). Despite their small size, emissions from near-shore ecosystems such as estuaries could nearly balance the carbon uptake by marginal seas (Borges et al., 2005; Chen and Borges, 2009). Global CO_2 emissions from estuaries range from 0.1 to 0.6 PgC per year (Borges et al., 2018), an amount equivalent to 5-30% of the oceanic CO_2 sink of ~ 2 PgC per year (Le Quéré et al., 2016), while CH_4 emissions range from 1 to 7 Tg per year, mainly driven by organic matter accumulation, anaerobic decomposition in sediments, and production in submerged plants (Borges et al., 2018; Rosentreter et al., 2018). However, CH_4

emissions are probably underestimated due to difficulties in accounting for ebullition and gas flaring (Borges et al., 2016; Humborg et al., 2019). The complexity of these pathways, combined with a lack of systematic, high-resolution, and long-term measurements, hampers robust global assessments of coastal CH₄ emissions (Roth et al., 2022). Besides CO₂ and CH₄, coastal ecosystems are important but poorly constrained sources of nitrous oxide (N₂O), a potent greenhouse gas with a global warming potential 273 times higher than CO₂ over 100 years (IPCC 2023). N₂O ~~in oceanic environments is formed as a byproduct during primarily produced via~~ nitrification (ammonia oxidation) and ~~as an intermediate during~~ denitrification, ~~both of which depend strongly on oxygen concentration and are microbial processes that occur in the water column, (nitrate reduction) in sediments, and water columns and within suspended particles is controlled by the availability of dissolved inorganic nitrogen and oxygen~~ (Bange, 2006). Eutrophication and hypoxia, often driven by excess nutrient and organic matter inputs, have been shown to promote N₂O emissions (Murray et al., 2015; Brase et al., 2017). Coastal ecosystems are recognized as significant sources of N₂O to the atmosphere (Bange, 2006; Cheung et al., 2025; Resplandy et al., 2024), ~~where denitrification, especially in sediments and on particles, often dominates N₂O production, even in well-oxygenated waters (Wan et al., 2023).~~ However, coastal estimates remain uncertain due to sparse measurements and high spatial heterogeneity- (Wan et al., 2023). Given the strong warming potential of N₂O, understanding its fluxes from estuaries is essential for a complete picture of coastal greenhouse gas dynamics and determining how these emissions, along with those of CH₄ (with a warming potential of 27; IPCC 2023), offset the coastal uptake of CO₂ (Resplandy et al., 2024; Rosentreter et al., 2018; Roth et al., 2023). This questions the efficiency of blue carbon coastal ecosystems in mitigating the rise of rising GHG concentrations in the atmosphere and their related radiative effects (Kristensen et al., 2025; William and Gattuso, 2022). ~~This study investigates~~ We investigate the dynamics of CO₂, CH₄, and N₂O (GHGs) along a salinity gradient and across contrasting coastal habitats within an estuary. We combined detailed field measurements of surface seawater physical and biogeochemical properties with both in situ measurements and calculated estimates of air–sea GHG exchange. This approach allowed us to capture spatial variability, ~~identify and discuss the underlying biogeochemical roles of physical drivers, and examine interactions with coastal habitats, and biological processes in the observed changes in GHG concentrations.~~ In doing so, we aim to ~~disentangle the biogeochemical controls of GHG dynamics in estuarine environments and contribute~~ estimate the contribution of coastal ecosystems to GHG emissions and provide additional data that may be useful in improving global estimates of GHG emissions from the coastal ~~ocean's GHG emissions~~ ocean.

90 2. Materials and methods

2.1 Study area

This study was conducted within a 30 km radius of the Tvärminne Zoological Station (TZS), located at the southwestern tip of mainland Finland. The study area encompasses the coastal waters extending from the cities of Hanko, Tammisaari, and Pohja (Fig. 1), where the coastal environment is characterized by strong spatial heterogeneity due to the intricate bathymetry and geomorphology of the archipelago system (Asmala and Scheinin, 2024) and enhanced human impact. The studied area

receives substantial freshwater from the Karjaanjoki river (in Pohja, runoff $0.59 \text{ km}^3 \text{ yr}^{-1}$, Raike et al., 2012), delivering relatively large amounts of allochthonous carbon and nutrients to the system, resulting in eutrophication and elevated concentrations of organic carbon in the water column across the inner archipelago area (Fleming-Lehtinen et al., 2015). Sampling sites ($N = 21$) were selected to encompass a wide range of soft-sediment habitats (e.g., including both vegetated and non-vegetated sediments with grain size ranging from coarse sand to clay, silt, and mud) and to represent a spatial gradient (50 km) from the outer to innermost archipelago, encompassing exposed to sheltered bays. Exposed sites are characterized. We used the classification by a higher Lammerant et al. (2025), who categorize sampling sites based on their salinity and sediment characteristics to indicate the degree of exposure to environmental forces. Exposed sites (sites 1–4 and 17–21) are likely to experience stronger wave, wind, and current energy, and greater as well as higher water exchange (sites 1–4 and 14–21), while, compared to sheltered sites. Sites within Pojo Bay (sites 1–4) are broadly open to the bay and are directly influenced by wind and water input from the Karjaajoki River and the archipelago. Therefore, those are included with the exposed sites. The sheltered sites (sites 5 and 7–11) are more enclosed and experience more likely to have limited water circulation (sites 5–13). Semi-sheltered sites fall between exposed and sheltered environments (sites 6 and 12–16).

Exposed and semi-sheltered sites in the outer archipelago (between Hanko and TZS, sites n°14–21, Fig. 1) are characterized by sandy sediments (96.2–100% sand) with low organic matter content (0.3–1.3%). These sites host both marine and freshwater plant species (Gustafsson and Norkko, 2019; Lammerant et al., 2025), and their faunal biomass is dominated by bivalves and polychaetes (Gammal et al., 2019; Makelin et al., 2024). Sheltered and semi-sheltered sites within the archipelago (sites n°5–13) are characterized by sediments containing high proportions of clay/silt (20.8–64.3%) and organic matter (2.4–13.2%; Lammerant et al., 2025). At these sites, fast-growing fresh- and brackish-water macrophytes are often found, while faunal biomass is dominated by gastropods (Makelin et al., 2024). In Pojoviken Bay, at the 14 km inlet between Tammissaari and Pohja (sites n°1–4), dense vegetation occurs down to ~2 m depth but becomes sparse at greater depths (>3 m) due to light limitation (Lammerant et al., 2025). Deeper sites are characterized by high organic matter content (>13%) in the sediment and low faunal biomass (Gammal et al., 2019). Sampling was conducted from 14 August to 18 September 2023/12 September 2023. Sites were visited once during the study period, with sampling conducted every two to three days, depending on weather conditions and boat availability.

2.1 Sampling procedure

Physical and biogeochemical properties of surface seawater, including the partial pressure of dissolved CO_2 ($p\text{CO}_2$) and CH_4 concentration, were measured *in situ* using a custom-built flow-through system. Seawater, from a depth of 50 cm, was pumped onboard using a submersible pump and directed into two separate flow-through systems.

The first water flow (3 L min^{-1}) was dedicated to gas extraction using a double showerhead equilibrator (Sunburst) equipped with a temperature probe (Hex fitting thermistor, ThermX, 0.1°C accuracy) and a barometric pressure sensor (BARO-A-4V, All Sensors). From the equilibrator, a continuous air flow (2 L min^{-1}) circulated in a closed loop to an infrared gas analyser (IRGA, LI-COR, LI-7810). The analyser measured the dry mole fractions of CO_2 and CH_4 , denoted as $x\text{CO}_2$ and $x\text{CH}_4$,

respectively, as the gas passed through a Peltier cooler to remove the excess water before entering the IRGA. Both gas concentration and temperature were recorded every second until an equilibrium was reached (up to 45 minutes). The measured $x\text{CO}_2$ was later converted into $p\text{CO}_2$. Equilibration between the seawater and gas phases was monitored in real time using a laptop connected to the IRGA. Equilibrium was considered reached when both CO_2 and CH_4 concentrations stabilized at a clear plateau. Depending on the concentration gradient between sites, this equilibration period could vary substantially. While CO_2 typically reached equilibrium within a few minutes, CH_4 required longer times (up to 45 minutes) to reach a stable plateau.

135 The measured $x\text{CO}_2$ (in ppm) was later converted into $p\text{CO}_2$ (in μatm) and corrected for *in situ* temperature using the Matlab CO2SYS v3 (Sharp et al., 2023). Surface seawater CH_4 concentration (in nmol L^{-1}) was computed from $x\text{CH}_4$ (in ppm), *in situ* temperature, salinity, and the solubility coefficients from Wiesenburg and Guinasso (1979). The IRGA was factory-calibrated, and standard gases for CO_2 (150, 420, and 1500 ppm) and CH_4 (1, 20, and 150 ppm) were passed through the IRGA both before and after deployment. The data were corrected for potential drift.

140 The second water flow (2 L min^{-1}) was directed through a thermosalinograph (SeaBird, TSG045), a fluorometer for chlorophyll *a* (chl-*a*), turbidity, and phyco cyanin concentrations (Chelsea, TriLux), a fluorometer for coloured dissolved organic matter (CDOM, Chelsea, UviLux), and an optode for dissolved oxygen concentration and saturation (Aanderaa 4531). Each sensor was factory-calibrated before deployment, and data were logged at 1-second intervals until equilibrium for $x\text{CO}_2$ and $x\text{CH}_4$ was reached (up to 45 min). A 5-second average was then recorded.

145 Discrete surface water samples were collected for total alkalinity (TA), dissolved inorganic carbon (DIC), methane (CH_4), and nitrous oxide (N_2O) concentration using a peristaltic pump (Cole Palmer, Masterflex Environmental Sampler) equipped with Tygon tubing. Seawater samples were collected into 12 mL gas-tight vials (Exetainers, Labco High Wycombe, UK) for DIC, 60 mL borosilicate vials for TA, and 60 mL serum bottles for dissolved CH_4 and N_2O concentrations. Samples were preserved by adding 1% of the sample volume of a saturated mercuric chloride (HgCl_2) solution. Samples were stored in the dark at room

150 temperature until analysis.

Air-sea exchanges of CO_2 and CH_4 were measured using the accumulation chamber technique (Frankignoulle, 1988). The chamber consists of a polyethylene container (internal diameter: 34 cm, height: 14 cm, total volume = 11.4 L) connected in a closed loop to the IRGA; with 1.5m long Tygon tubing, which allows the chamber to move freely on the water surface. The air partial pressure of CO_2 ($p\text{CO}_2$) and CH_4 ($p\text{CH}_4$) within the chamber was recorded every second for 10 minutes. The flux

155 was computed from the slope of the linear regression of $p\text{CO}_2$ and $p\text{CH}_4$ against time ($R^2 > 0.99$) following the method of Frankignoulle (1988), accounting for the air volume enclosed within the chamber. The uncertainty of the flux calculation, based on the standard error on the regression slope, was approximately 3% on average.

Before conducting the chamber measurements, the IRGA baseline was established by flushing atmospheric air through the analyser. Based on these measurements, we estimated average atmospheric concentrations of 406 ppm for CO_2 and 2.05 ppm

160 for CH_4 during the survey. Reported fluxes represent the mean of two to three individual chamber measurements performed at each site. Direct measurements of N_2O air-sea exchanges were not possible during our survey. Therefore, the air-sea exchange

of N₂O, as well as CO₂ and CH₄ (i.e., F_X, where X represents N₂O, CO₂, or CH₄, respectively) was parameterized using the bulk formula:

$$F_X = kK_{0(X)}(pX_{(sw)} - pX_{(atm)}) \quad (1)$$

165 where k (m s⁻¹) is the gas transfer velocity, K_0 (mol m⁻³ atm⁻¹) is the temperature and salinity dependent gas solubility determined from Weiss (1974) for CO₂, Wiesenburg and Guinasso (1979) for CH₄, and Weiss and Price (1980) for N₂O, and $pX_{(sw)}$ and $pX_{(atm)}$ are the measured partial pressure of gas in the surface seawater and the air, respectively.

This approach strongly relies on selecting an appropriate parameterization for the gas transfer velocity, k . Numerous theoretical, laboratory, and field studies established that k depends on a variety of parameters, including capillary and breaking waves, boundary layer stability, air bubbles, surfactant surface films, evaporation and condensation, precipitation, water currents (including tides), as well as turbulence at the air-water interface (Borges et al., 2004). The parameterization of k is most often expressed as a function of wind speed. This relationship is well established for the open ocean, where wind stress is the main source of turbulence (e.g., Wanninkhof, 2014). In estuarine environments, significant regression functions between k and wind speed have also been reported (Borges et al., 2004; Raymond et al., 2000). However, such formulations have been proven to be site-specific and may not fully account for other controlling processes, such as turbulence at the air-water interface (Borges et al., 2004). Since no k -wind relationship has been developed for our study area, we adopted the formulation proposed for the Randers fjord by Borges et al. (2004), as its physical characteristics, particularly tidal amplitude and freshwater discharge, are most comparable to those of our site:

$$k = 1.2 + 2.3u \left(\frac{Sc_{(balt)}}{660} \right)^{-0.5} \quad (2)$$

180 where u is the wind speed (m s⁻¹) and $Sc_{(balt)}$ is the Schmidt number at the sampling site, based on local temperature, salinity, and gas molecules. $Sc_{(balt)}$ is extrapolated from freshwater and seawater ($S = 35$) coefficients from Wanninkhof (2014) for the salinity observed at the sampling site (Roth et al., 2023).

Wind speed (in m s⁻¹, METEK, uSonic-3 Scientific) and air temperature (in °C, Vaisala, HMP155) are measured at the 3.2 and 2 m height, respectively, above sea level at the newly established Integrated Carbon Observation System (ICOS) coastal site at TZS (ICOS code FI-Tvm; Fig. 1). Daily averaged wind speed, extrapolated to a height of 10m using the relationship from Hsu et al., (1994), assuming a near-neutral atmospheric stability conditions, was used to calculate the air-sea exchanges of GHGs. Atmospheric concentrations of CO₂ and CH₄ were measured at each site before the air-sea flux measurements. For N₂O, we used the monthly average atmospheric concentration of 0.3367 ppm, measured at Pallas-Sammaltunturi, GAW Station, Finland (PAL, <https://gml.noaa.gov/dv/iadv/graph.php?code=PAL&program=ccgg&type=ts>).

190 The recorded data on from the IRGA for both seawater and air-sea flux measurements were filtered by removing to remove measurements taken during the transition period between stations and when switching between ambient air sampling and seawater measurements, as the IRGA requires time to respond to sharp concentration changes. Additionally, data affected by improper functioning (i.e., seawater flow < 1.5 L min⁻¹) were discarded.

2.3 Sample analysis

195 TA was determined by Gran titration (Gran, 1952) using an AS-ALK2 titration system (Apollo SciTech), where a 15-ml sample was titrated with a standard 0.1 M HCl solution. DIC was measured on a DIC analyser (Apollo SciTech) by acidification of a 0.75 ml subsample with 1 ml 10% H₃PO₄, and quantification of the released CO₂ with a non-dispersive infrared CO₂ analyser (LI-COR, LI-7000). Results were then converted from μmol L⁻¹ to μmol kg⁻¹ based on sample density, calculated from salinity and temperature at the time of the analysis. Accuracies of ±3 and ±2 μmol kg⁻¹ were determined for
200 TA and DIC, respectively, from routine analysis of certified reference materials (A.G. Dickson, Scripps Institution of Oceanography, San Diego, CA, USA).

Discrete seawater samples for CH₄ and N₂O concentrations were measured via the headspace equilibrium technique (25 ml N₂ headspace in 60 ml serum bottles) and measured with a gas chromatograph (SRI 8610C) with flame ionization detection, and electron capture detection calibrated with CH₄: CO₂:N₂O: N₂ mixtures (Air Liquide) of 1.0, 10, and 30 ppm CH₄ and 0.2, 1.0,
205 and 6.0 ppm N₂O. The measurement precision was 8 % for CH₄ and 4 % for N₂O.

3. Results

3.1 Atmospheric forcing and seawater physical properties

During the survey period, atmospheric temperatures remained relatively stable, averaging 18.0°C (SD = 1.5, n = 1989; Fig. 1a). A short warm event occurred on the 7th of August, when the air temperature peaked at 26.3°C. In contrast, short cold spells
210 were observed on 5 and 7 September, with temperatures dropping to 13.7 and 11.5°C, respectively. With relatively stable atmospheric temperature, surface seawater temperature (SST) showed limited variation, ranging from 17.2 to 19.5°C (mean = 18.4°C, SD = 0.66, n = 21; Fig. 1c). The Karjaanjoki River, being the main source of freshwater input to the area, created a pronounced salinity gradient, from nearly fresh surface seawater salinity (SSS = 0.07) at the river mouth (in Pohja) to a SSS of 6.36 in Hanko (Fig. 1d).

215 3.2 Seawater biogeochemical properties

Surface seawater chlorophyll-*a* (chl-*a*) concentration ranged from 3.7 to 37.7 μg L⁻¹ (mean = 14.4, SD = 8.1, n = 21, Fig. 1e), with the highest concentration observed in the sheltered bay around Tammisaari (S5 and S6). Turbidity ranged from 5.0 to 8.4 FTU (mean = 6.1, SD = 0.8, n = 21, Fig. 1f), with the highest values observed in sheltered bays within the archipelago (S8, S9, and S11). Coloured dissolved organic carbonmatter (CDOM) ranged from 3.0 to 10.6 μg L⁻¹ (mean = 5.9, SD = 2.6, n =
220 21, Fig. 1g) and exhibited a salinity-driven gradient, with the highest concentrations observed towards Pohja and the lowest towards Hanko. However, one site in the archipelago (S8) and near Hanko (S19) exhibited elevated CDOM concentrations compared to other locations in their respective area, reaching up to 10.0 and 10.6, respectively. Dissolved oxygen (DO) saturation levels ranged from 74.3 to 112.7 % (mean = 94.1, SD = 9.6, n = 21, Fig. 1h), with most of the sites undersaturated

in O₂, excepted for two sites closed by Tammisaari (S3 and S4) and four sites in closed vicinity of TZS (S14, S15, S18, and S19).

Surface water pCO₂ ranged from strong undersaturation (160 ppm) to a significant supersaturation relative to the atmosphere (average atmospheric concentration measured during this survey = 406 ppm) at S2 (1702 ppm) and up to 2521 ppm at the river mouth (triangle on Fig. 2a). Surface water CH₄ concentrations were consistently supersaturated (up from 636 to 14609 %, data not shown) compared to the atmosphere (average atmospheric concentration measured during this survey = 2.05 ppm), with concentrations ranging from 19 to 469 nmol L⁻¹ (Fig. 2b). Undersaturated pCO₂ (from 160 to 403 ppm) and low CH₄ concentrations (from 19 to 34 nmol L⁻¹) were mostly observed at open sites between TZS and Hanko (S14 to S21), except for S17, where pCO₂ and CH₄ reached 485 ppm and 48 nmol L⁻¹, respectively. Seawater pCO₂ and CH₄ concentrations increased across the archipelago and Pojoviken Bay, reaching their maximal values at the mouth of the Karjaanjoki River. Local lows were observed in Pojoviken Bay (S3 and S4), where pCO₂ was undersaturated (192 and 238 ppm, respectively), and CH₄ concentrations were relatively low (83 and 67 nmol L⁻¹, respectively). N₂O concentration ranged from 9 to 25 nmol L⁻¹, with higher saturation level ranging from 93 to 255 %. Higher concentrations and maximum saturation were observed at the mouth of the Karjaanjoki River (Fig. 2c). South of Tammisaari, surface waters were either undersaturated in N₂O (S8, S13, and S16) or close to the atmospheric equilibrium (S6 and S12, data not shown). TA and DIC ranged from 668 to 1631 and from 720 to 1550 μmol kg⁻¹, respectively, with minimum TA and DIC concentration measured at the mouth of the Karjaanjoki River. Both TA and DIC exhibited conservative behaviour with changes in salinity (Fig. 3a).

3.3 Air-sea exchanges of greenhouse gases GHG flux densities

The area presented contrasts in terms of sink and source of CO₂ and N₂O, but was clearly a source of CH₄. Measured air-sea exchanges of CO₂ flux densities ranged from -36 to 150 mmol m⁻² d⁻¹, where negative values indicate net uptake of atmospheric CO₂ (Fig. 2d). In contrast, the study area acted as a net source of CH₄ to the atmosphere, with fluxes flux densities ranging from 0.014 to 1.39 mmol m⁻² d⁻¹ (Fig. 2e). Overall, the calculated fluxes showed a similar trend in term of pattern and range compared to the observed values, ranging from -26.4 to 220.3 mmol m⁻² d⁻¹ for CO₂ (Fig. 2f) and from 0.03 to 1.1 mmol m⁻² d⁻¹ for CH₄ (Fig. 2g). Calculated fluxes of N₂O ranged from -0.0021 to 0.0383 mmol m⁻² d⁻¹, with sites in the archipelago and towards TZS acting as a slight sink of N₂O (Fig. 2h).

3.4 Statistical analysis

The Kendall's τ coefficient has been calculated to investigate the correlation between surface water pCO₂, CH₄, and N₂O concentration, as well as between physical and biogeochemical parameters and all three GHGs (Fig. 4). Surface pCO₂ and CH₄ concentrations show a moderate (τ = 0.5636) and statistically significant (p = 0.0165) positive relationship (Fig. 4a), suggesting that processes driving high pCO₂, such as respiration and organic matter degradation, are likely also contributing to elevated CH₄ levels. N₂O is not correlated with pCO₂ and shows a moderate correlation with CH₄ (τ = 0.4182), although this relationship is only marginally significant (p = 0.0866). Among the tested biogeochemical variables, SSS and O₂ saturation show strong,

significant negative correlations with pCO₂ and CH₄ (Fig. 4b). N₂O is only strongly negatively correlated with salinity ($\tau = -0.8182$, $p = 0.00013$) and shows a marginal correlation with CDOM ($\tau = 0.4545$, $p = 0.060$). CDOM is positively correlated with CH₄ ($\tau = 0.4952$, $p = 0.0013$) while chl-*a* is moderately and significantly correlated with CH₄ ($\tau = 0.4095$, $p = 0.009$). SST and turbidity show weak and non-significant relationships with the three gases.

260 4. Discussion

Surface water pCO₂, CH₄, and N₂O concentrations showed significant spatial variation, spanning several orders of magnitude (Fig. 2). Humborg et al. (2019) reported pCO₂ and CH₄ levels following the salinity gradient driven by the Karjaanjoki River, with values reaching up to 1583 μatm and 70 nmol L^{-1} , respectively. However, these measurements were conducted alongin the deeper part of the archipelago's main channel-of the archipelago, and they did not explore Pojoviken Bay, whereas. In
265 contrast, our study focused on shallow (< 4m depth) nearshore ecosystems. Asmala and Scheinin (2024) observed a similar range of pCO₂ and CH₄ concentrations across a 2000 km² coastal region surrounding the Hanko Peninsula, with the large magnitude of both gases associated with excessive organic matter loads, elevated primary production, trapping and accumulation of allochthonous organic matter, and sedimentary conditions favourable to CH₄ production. Focusing on nearshore shallow (< 4m depth) coastal habitats around the island of Askö (Sweden, northern Baltic Sea), Roth et al. (2023)
270 reported strong spatial variability in surface water pCO₂ and CH₄ concentrations and emphasized that their dynamics were stronglyhighly habitat--specific. The N₂O concentrations we observed were in the same range as those observed by Aalto et al. (2021) in the same area, with concentrations ranging from 25 to 50 nmol L^{-1} within Pojoviken Bay and from 10 to 30 nmol L^{-1} within the archipelago. Aalto et al. (2021) reported that higher N₂O concentrations were associated with higher nitrate concentrations and inputs of allochthonous carbon, while lower N₂O concentrations were associated with efficient internal
275 recycling of N.

4.1 Salinity gradient

~~The Kendall's τ coefficient has been calculated to investigate the correlation between surface water pCO₂, CH₄, and N₂O concentration, as well as between physical and biogeochemical parameters and all three GHGs (Fig. 4). Surface pCO₂ and CH₄ concentrations show a moderate ($\tau = 0.5636$) and statistically significant ($p = 0.0165$) positive relationship (Fig. 4a), suggesting
280 that processes driving high pCO₂, such as respiration and organic matter degradation, are likely also contributing to elevated CH₄ levels. N₂O is not correlated with pCO₂ and shows a moderate correlation with CH₄ ($\tau = 0.4182$), although this relationship is only marginally significant ($p = 0.0866$). Among all tested biogeochemical variables, SSS and O₂ saturation show strong and significant negative correlations with pCO₂ and CH₄ (Fig. 4b). N₂O is only strongly negatively correlated with salinity ($\tau = -0.8182$, $p = 0.00013$) and shows a marginal correlation with CDOM ($\tau = 0.4545$, $p = 0.060$). CDOM is positively correlated
285 with CH₄ ($\tau = 0.4952$, $p = 0.0013$) while chl-*a* is moderately and significantly correlated with CH₄ ($\tau = 0.4095$, $p = 0.009$). SST and turbidity show weak and non-significant relationships with the three gases.~~

4.1 Drivers for variability in surface water pCO₂, CH₄, and N₂O concentration

The Karjaanjoki river is the main source of freshwater into the studied area, delivering large amounts of allochthonous carbon and nutrients to the system (Fleming-Lehtinen et al., 2015). Changes in SSS have been suggested to strongly influence the concentration of all three GHGs (Fig. 4b), with terrestrial runoff likely contributing to elevated CH₄ concentrations in the area (Asmala and Scheinin, 2024). At a first glimpse, only N₂O exhibited a strong relationship with salinity ($R^2 = 0.95$), while pCO₂ ($R^2 = 0.58$) and CH₄ ($R^2 = 0.39$) exhibited weaker correlations (Fig. 5, dotted black line). However, when focusing **exclusively** on exposed **and semi-sheltered** sites, the correlation with salinity strengthened considerably, reaching values up to 0.79 for pCO₂, 0.94 for CH₄, and 0.96 for N₂O (Fig. 5, dashed red line). The impact of water mixing on surface-water pCO₂ can be assessed by estimating the variation in pCO₂ resulting only from the physical mixing of Karjaanjoki River water ($S = 0$, $TA = 668 \mu\text{mol kg}^{-1}$, $DIC = 720 \mu\text{mol kg}^{-1}$) with a seawater endmember (black **symbols** in Fig. 3, $S = 6.42$, $TA = 1635 \mu\text{mol kg}^{-1}$, $DIC = 1586 \mu\text{mol kg}^{-1}$). The seawater endmember was collected on 18 September 2023 at 59.77393 °N, 23.2607167 °E, about 7.5 km south of TZS. This calculation excludes the effects of biological activity, gas exchange, and potential precipitation or dissolution of calcium carbonate. Assuming that changes in SSS reflect a mixing ratio between the Karjaanjoki River and the seawater endmembers and based on the observed linear relationships between salinity and both TA and DIC (Fig. 3a), representing conservative mixing between the two water masses, TA and DIC can be estimated based on salinity. The estimated TA and DIC are then used to compute pCO₂ (blue dashed line, Fig. 5a) using CO2SYS v3 (Sharp et al., 2023), applying the carbonic acid dissociation constants (K_1 and K_2) of Millero et al. (2006) and the KHSO₄ formulation of Dickson (1990). While TA and DIC behaved conservatively during mixing, pCO₂ **showed** a pronounced nonlinear response, similar to the one observed for exposed **and semi-sheltered** sites (Fig. 5a). Such behaviour has already been reported in estuaries (Abril et al., 2021; Cai et al., 2013). This suggests that the input of CO₂, CH₄, and N₂O supersaturated water from the Karjaanjoki River is a major source for all three GHGs across the study area, with dilution playing a dominant role in GHG dynamics at exposed **and semi-sheltered** sites. While changes in SSS appear to explain most of the variability observed in pCO₂, CH₄, and N₂O concentrations, deviations from the expected salinity-driven pattern, such as those observed within the archipelago, at sheltered sites, likely reflect additional local processes.

4.2 Biological drivers

Many processes affecting the carbonate system are best described by examining the associated changes in DIC and TA (Zeebe and Wolf-Gladrow, 2001). However, most of the changes in TA and DIC in surface seawater appeared to be driven by salinity changes, with estimated endmember ($S=0$) values of 662 and 712 $\mu\text{mol kg}^{-1}$ for TA and DIC, respectively (Fig. 3a). These estimates closely match the measured TA and DIC at the mouth of the Karjaanjoki River (668 and 720 $\mu\text{mol kg}^{-1}$, respectively). Therefore, to discard the impact of freshwater inputs, TA and DIC were normalized to the average surface seawater salinity of 5 (denoted as nTA and nDIC, respectively) using the normalization of Friis et al. (2003). In Figures 3b, c, and d, the dotted lines illustrate the expected responses of TA and DIC to different biogeochemical processes. Biological activity

(photosynthesis/respiration) affects both TA and DIC in a ratio of -0.16 (Lazar and Loya, 1991), air-sea exchange of CO₂ only affects DIC, and the precipitation-dissolution of calcium carbonate affects TA and DIC in a 2:1 ratio. However, the effect of calcifying primary producers in the carbon pool can be neglected in the Baltic Sea, except in the benthic zone (Tyrrell et al., 2008). The sampling sites are compared to the seawater endmember (S = 6.42, TA = 1635 μmol kg⁻¹, DIC = 1586 μmol kg⁻¹, black symbol in Fig. 3). Figures 3b and c, show how exposed sites, undersaturated in pCO₂ and low CH₄ concentrations, follow the theoretical trend of photosynthesis and CO₂ release. However, an uptake of CO₂ was both measured and estimated for those sites (Fig. 2), suggesting that gas exchange as release towards the atmosphere could not explain the observed changes in the carbonate system. In contrast, sheltered sites, with supersaturated pCO₂ and elevated CH₄ concentrations, were more scattered around the theoretical respiration line and CO₂ uptake. Gas exchange, as CO₂ uptake, can also be ruled out, as both measured and estimated air-sea fluxes pointed to a net release of CO₂ into the atmosphere. Altogether, this suggests that biological processes—primary production at undersaturated pCO₂ sites and respiration at supersaturated sites—mainly influence the carbonate system, while the direct role of air-sea exchange of CO₂ appears to be minor in shaping the inorganic carbonate dynamics across the study area. The similar pattern observed for both pCO₂ and CH₄ concentrations illustrated the positive relationship between the two gases (Fig. 4), where processes responsible for higher pCO₂, such as respiration and/or organic matter degradation, also contribute to higher CH₄ concentrations (Reeburgh, 2007). No clear pattern could be observed between the nTA: nDIC ratio and N₂O concentration, as expected by the poor Kendall's coefficient of correlation (Fig. 4).

Elevated chl-*a* concentrations (up to 37.7 μg L⁻¹, Fig. 1) in surface seawater were reported across the studied area. Chl-*a* measurements are typically considered as a proxy for pelagic primary production (Cloern et al., 2014). Therefore, elevated chl-*a* could indicate active photosynthesis production, resulting in pelagic O₂ production and pelagic CO₂ uptake. To investigate the relationship between photosynthetic activity and surface water pCO₂, CH₄, and N₂O concentrations, apparent oxygen utilization (AOU), calculated as the difference between oxygen saturation and measured oxygen concentration, was examined with chl-*a* concentrations. ~~No~~ However, no clear relationship was observed between chl-*a* and AOU (Fig. 6a) and negative AOU values, indicative of net photosynthesis, were associated with relatively low chl-*a* concentrations within the observed range (Fig. 6a). Negative values of AOU, indicative of net photosynthesis, were associated with relatively low chl-*a* concentrations within the observed range (Fig. 6a). These negative AOU values corresponded to undersaturated pCO₂ and low CH₄ concentrations (Fig. 6b, c). In contrast, positive AOU values, representing net respiration, were associated with supersaturated pCO₂ and elevated CH₄ concentrations, both of which increased with higher AOU. A strong correlation exists between AOU and surface water pCO₂ (Fig. 6b), which improves when focusing on exposed and semi-sheltered sites (red dashed line), suggesting that ~~pelagic primary production is the main process affecting~~ strongly affects the carbonate system, as already indicated by the nDIC: nTA ratio (Fig. 3). ~~This relation is weaker for CH₄ and N₂O, as respiration is not ultimately associated with CH₄ (3). Primary production. Therefore, it could take place either in the pelagic or the benthic realm. However, the lack of correlation between chl-*a* and AOU and the low chl-*a* concentration is considered as a proxy for associated with negative AOU suggests that pelagic primary production (Cloern et al., 2014), this suggests that primary production within the pelagic realm is not may not be solely responsible for the change of pCO₂. The influence of the pelagic realm is likely limited, as phytoplankton biomass is constrained observed changes in pCO₂. Pelagic primary production may be limited by the shallow~~

water column, which restricts phytoplankton biomass and its productivity is low at the time, as this study was conducted at the end of the summer bloom (Uth et al., 2024). ~~Therefore, processes beyond pelagic production, such as benthic community production and~~ The correlation between AOU and CH₄ is weaker than for CO₂ (R² = 0.51, black dotted line) but improves significantly when only exposed and semi-sheltered sites are considered (R² = 0.78, red dashed line). O₂ depletion, associated with supersaturated pCO₂ and high CH₄ concentration, could be related to organic matter degradation, are highly likely to influence the observed pCO₂ and as large amounts of carbon are turned over during the summer (Attard et al., 2019). ~~However, here as well, limited phytoplankton biomass in the shallow water column could restrict the impact of the pelagic realm on CH₄ dynamics. No concentration. N₂O exhibited a clear relationships between AOU and N₂O could be observed, as expected positive correlation with AOU (R² > 0.7). This correlation is known to reflect the coupling between O₂ consumption (from the Kendall's correlation coefficient. organic matter remineralization) and microbial nitrogen cycling processes that produce N₂O (Kock et al., 2016; Carrasco et al., 2017).~~

Therefore, the absence of a correlation between chl-*a* and AOU, despite AOU correlating with all three GHGs, suggests that pelagic processes alone cannot explain the observed changes in GHG concentrations. Instead, processes beyond the pelagic realm, such as benthic community production and organic-matter degradation, are likely contributing to the observed surface-water pCO₂, CH₄, and N₂O concentrations (e.g. Roth et al. 2022, 2023).

4.3 Exposed and semi-sheltered vs sheltered sites

The inputs of freshwater supersaturated in CO₂, CH₄, and N₂O represent a major source of GHGs to the study area. Freshwater mixing with seawater appears to control the concentration of all three GHGs at exposed and semi-sheltered sites, as much of the variability in surface seawater concentrations can be explained by salinity. In sheltered sites of the archipelago, deviations from the salinity-driven pattern indicate the influence of additional processes. Biological activity is likely the key driver, with pCO₂ ~~undersaturated~~ undersaturation associated with primary production, ~~while and~~ pCO₂ ~~supersaturated~~ supersaturation linked to respiration. However, pelagic production alone is unlikely to be the main contributor to the observed changes. Benthic processes such as community production, organic matter degradation, and sediment-water interactions are likely to contribute to both CO₂ and CH₄ dynamics (Attard et al., 2019); Roth et al., 2023). For CH₄, riverine inputs and subsequent dilution largely determine large-scale patterns, ~~but~~ whereas elevated concentrations at sheltered sites within the archipelago reflect localized production and limited oxidation. Large amounts of carbon are turned over in ~~the~~ habitats from the studied area during the summer months (Attard et al., 2019), and macrophyte tissue ~~becomes~~ may become a direct component of local sediment organic matter pools ~~that, which~~ favour local CH₄ production (Roth et al., 2022; Wallenius et al., 2021). Anoxic degradation of organic-rich sediments (methanogenesis), exacerbated by warm late-summer waters (Roth et al., 2022; Yvon-Durocher et al., 2014), ~~together combined~~ with short water residence times that ~~restrict~~ limit oxidation in both sediment and the overlying water column (Reeburgh, 2007), ~~create~~ creates favourable conditions for CH₄ ~~accumulation.~~ production. Due to the sheltered nature of the sites and limited water exchange, the produced CH₄ can accumulate. Similarly, enhanced respiration of organic carbon in shallow ~~sediments~~ ecosystems elevates CO₂ concentrations (Humborg et al., 2019). While macrophyte

growth during summer can draw down pCO₂ through photosynthetic CO₂ uptake, the subsequent decomposition of deposited tissues may instead enhance CO₂ and CH₄ release. Spatial variability of N₂O mirrors the findings of Aalto et al. (2021), ~~with who linked~~ higher N₂O concentrations near the Karjaanjoki River ~~linked to to~~ higher nitrate inputs and allochthonous carbon inputs ~~and elevated nitrate availability. Aalto et al. (2021).~~ They suggested that ~~within~~ the ratio between nitrate and autochthonous organic carbon controls the balance between N-removing denitrification and N-recycling through Dissimilatory Nitrate Reduction to Ammonium (DNRA), as well as the end-product of denitrification (Aalto et al., 2021). Within the archipelago, ~~reduced where~~ riverine influence ~~and enhanced sedimentary processes, such as dissimilatory nitrate reduction, likely sustain high~~ is limited. DNRA can produce significant amounts of bioavailable ammonium ~~availability and promote active, enhancing~~ nitrogen recycling between sediments and surface waters, ~~particularly water, especially~~ in summer, when autochthonous biomass production and sedimentation are highest.

4.24 Air-sea ~~fluxes~~ flux densities

Given the strong spatial variability observed in GHG concentrations across the study area, strong variations were also observed in the corresponding air-sea fluxes. ~~In At~~ exposed sites between Hanko and TZS, where surface seawater was undersaturated in CO₂ and low in CH₄ concentration, an uptake of atmospheric CO₂ associated with small releases of CH₄ was observed. In contrast, within the archipelago and across Pojoviken Bay, where surface waters were supersaturated in CO₂ and exhibited elevated CH₄ concentrations, releases of both CO₂ and CH₄ were observed. Sites 3 and 4 represented exceptions, being the only locations undersaturated in CO₂, thus acting as a CO₂ sink. Daily CH₄ emissions of ≥ 0.1 mmol m⁻² across all habitats are comparable to, or even higher than, CH₄ fluxes reported from similar (Lundevall-Zara et al., 2021; Roth et al., 2023) or other vegetated coastal ecosystems (Al-Haj and Fulweiler, 2020; Rosentreter et al., ~~2021~~2021a). Regarding N₂O, Pojoviken Bay is estimated to act as a source to the atmosphere. In sheltered sites of the archipelago and exposed sites near TZS, N₂O concentrations close to or slightly below atmospheric equilibrium result in small emissions and occasional uptake, respectively. Our estimated air-sea exchanges overall mirrored the spatial trends of chamber-based fluxes and exhibited a similar range of air-sea exchange. Both chamber-based and estimated fluxes may underestimate the air-sea fluxes, as possible ebullition events are not considered. Humborg et al. (2019) suggested that air-sea CH₄ flux is likely dominated by frequent bubbling from the sediment. ~~While this~~ This process could be dominant in shallow coastal environments, as shallow seafloor depth promotes a short residence time for CH₄ in the water column, ~~preventing which prevents~~ its potential ~~for full oxidation—as, which, in contrast, been~~ observed in the deep open ocean (Reeburgh, 2007), ~~it~~). Bubbling was however not observed during our chamber-based measurements.

The overall contribution of the study area to atmospheric GHG budgets can be assessed by combining the air-sea flux of CO₂ with the CO₂-equivalent fluxes of CH₄ and N₂O. These equivalents are calculated using their respective sustained-flux global warming potential (SGWP), as a greenhouse gas metric to describe the relative radiative impact of a standardized amount of gas over a defined time horizon. Specifically, over a 100-year time horizon, the SGWP of CH₄ and N₂O is 27 and 273 times, respectively, greater than that of CO₂ (IPCC 2023), on a mass basis, based on:

$$F_{CO_2-eq(X)} = F_{(X)}SGWP_{(X)} \quad (3)$$

420 where the CO_{2-eq} flux of X (CH₄ or N₂O), $F_{CO_2-eq(X)}$, is the product of the flux of the gas ($F_{(X)}$) and its respective SGWP (i.e., 27 or 273) over the time horizon of 100 years. As direct measurements of air-sea N₂O fluxes were not available at the time, and to ensure consistency across all three gases, we chose to use the estimated fluxes rather than the chamber-based measurements.

Overall, the study area acted as a net source of GHGs to the atmosphere, with an average release of 1.6 g of CO_{2-eq} m⁻² d⁻¹ 425 over the study period, and ranging from -1.0 g of CO_{2-eq} m⁻² d⁻¹ at S14 to 10.6 g of CO_{2-eq} m⁻² d⁻¹ at the mouth of the Karjaanjoki River (Fig. 7d). Exposed sites between Hanko and TZS generally acted as GHGs sink, primarily due to the substantial CO₂ uptake compensating for CH₄ emissions. An exception was S21, where the small uptake of CO₂ (-0.008 g of CO₂ m⁻² d⁻¹) was offset by the release of CH₄ (0.016 g of CO_{2-eq} m⁻² d⁻¹), resulting in a net source. In Pojoviken Bay, the river mouth and S2 were acting as a strong source of GHGs to the atmosphere, with N₂O fluxes in the same order of magnitude as those of CH₄. 430 However, S3 and S4, both sites undersaturated in pCO₂, remained net GHG sinks, despite releasing similar amounts of CH₄ and N₂O. In the archipelago, all sheltered sites remained a net source of GHGs to the atmosphere. If, from a climate mitigation perspective, it may seem sufficient to focus on CO₂ (as Figure 7a and d are pretty similar), as in the blue carbon approach, it has been shown that CO₂ uptake can be offset by CH₄ emissions (Roth et al., 2023). Our findings demonstrate that, if CH₄ matters in coastal, nearshore environments, so does N₂O, since N₂O fluxes are not only of the same order of magnitude as CH₄ 435 fluxes but can either reinforce the warming effect of CH₄ or partially counterbalance it.

4.3.5. Conclusions

Shallow coastal ecosystems are highly heterogeneous, with their spatial structure and temporal dynamics of benthic communities shaping ecosystem functions (Snelgrove et al., 2014). This heterogeneity drives strong spatial variability in coastal biogeochemical processes, which exert an important influence on the carbon cycle at both local and global scales (Ward et al., 2020). Yet, despite their importance, identifying the locations and processes regulating coastal CO₂, CH₄, and N₂O fluxes 440 remains uncommon, and global estimates still fail to capture the heterogeneous and dynamic nature of these environments- (Rosentreter et al., 2021b). In particular, northern temperate coastal habitats are poorly represented in current GHG budgets- (Resplandy et al., 2024), despite their relative importance: shallow waters (<5 m) in the Baltic Sea cover ~30,000 km² (HELCOM, 2013), an area comparable to 20% of the global distribution of mangroves (Bunting et al., 2018) or seagrass 445 meadows (McKenzie et al., 2020).

Across the study area, freshwater inputs from the Karjaanjoki River and subsequent mixing with seawater largely determined the overall spatial patterns of surface water pCO₂, CH₄, and N₂O concentrations. However, important deviations from this salinity-driven control were observed at the local scale. In sheltered sites within the archipelago, elevated CH₄ concentrations and supersaturated pCO₂ likely reflected benthic biological processes, including enhanced organic matter respiration and 450 methanogenesis in warm, late-summer shallow waters and sediments, where limited oxidation favoured CH₄ accumulation.

By contrast, ~~in more~~ at semi-sheltered and exposed sites, mixing processes exerted stronger control, leading to lower GHG concentrations. N₂O patterns followed riverine influence with higher concentrations near the river mouth associated with inputs of allochthonous carbon and elevated nitrate availability, and lower concentrations within the archipelago reflecting reduced riverine input and active nitrogen recycling between sediments and surface waters (Aalto et al., 2021). Together, these results

455 underline the dual importance of large-scale physical mixing and local ~~benthic~~ biological processes in shaping the spatial heterogeneity of GHG dynamics. Our findings suggest that benthic ecosystems likely play a pivotal role in regulating GHG dynamics, especially in shallow coastal environments, not only through production but also through consumption pathways, which remain insufficiently constrained in current budgets (Rodil et al., 2021; Roth et al., 2022; Wallenius et al., 2021).

When translated into CO₂-eq, air–sea GHG fluxes were dominated by CO₂, while CH₄ and N₂O contributed comparably but in

460 different ways. CH₄ consistently acted as a source, whereas N₂O partially offset the CH₄ release through uptake. This ~~interplay~~ emphasizes interaction highlights that the balance between production and consumption processes, ~~particularly~~ especially within ~~benthic~~ different seafloor habitats, is critical ~~to~~ for understanding coastal contributions to the global carbon budget. There is a critical need to quantify CH₄ and N₂O exchanges more accurately and to deepen our understanding of the environmental and management factors that control their production and consumption. This would help make global estimates less sensitive

465 to statistical assumptions and reduce uncertainties in blue carbon potential estimates (Rosentreter et al. 2021a).

While coastal ecosystems are often recognized as CO₂ sinks, recent work shows that they are also significant sources of CH₄ and N₂O that can offset a substantial portion of the climate benefit of taking up CO₂ (Rosentreter et al., 2021a; Roth et al., 2023; Resplandy et al., 2024). Given their extent, northern temperate coastal ecosystems represent a relevant but overlooked source of GHGs, with the potential to amplify the global ocean carbon budget. ~~by increasing net greenhouse gas emissions to~~

470 the atmosphere. Our results highlight the urgent need for research that integrates GHG with biodiversity, benthic–pelagic interactions, and microbial processes, and that resolves temporal variability across seasonal cycles. Such knowledge is essential to improve predictions of how coastal ecosystems will mediate carbon–climate feedback under future environmental change.

Data availability

The complete dataset will be made publicly available upon acceptance of the manuscript.

475 **Author contribution**

NXG, AN, and AV designed the hypothesis field sampling. Field measurements were ~~conducted~~ carried out by NXG and AV, while laboratory and data analyses were performed by NXG, and BD. NXG prepared the manuscript with ~~contributions~~ input from all co-authors. All authors approved the final version of the manuscript for submission.

Competing interests

480 The authors declare no competing interests.

Acknowledgements

This study was supported by the Jane and Aatos Erkkö Foundation (AN, NXG), Walter and Andrée de Nottbeck Foundation (NXG), and the Sophie von Julin's Foundation (AN, AV) and utilized research infrastructure facilities at Tvärminne Zoological Station, University of Helsinki, as part of FINMARI (Finnish Marine Research Infrastructure Consortium). This is a
485 publication from the Centre for Coastal Ecosystem and Climate Change Research (www.coastclim.org). We thank Kurt Spence for his assistance with field work. CH₄ and N₂O measurements were supported by GreenFeedBack RIA (Greenhouse gas fluxes and earth system feedback) funded by the European Union's HORIZON research and innovation program under grant agreement N°101056921. Views and opinions expressed are, however, those of the authors only and do not necessarily reflect those of the European Union or CINEA. Neither the European Union nor the granting authority can be held responsible for
490 them. BD is a research associate at the F.R.S-FNRS.

References

- Aalto, S. L., Asmala, E., Jilbert, T., and Hietanen, S.: Autochthonous organic matter promotes DNRA and suppresses N₂O production in sediments of the coastal Baltic Sea, *Estuar Coast Shelf Sci*, 255, 107369, <https://doi.org/https://doi.org/10.1016/j.ecss.2021.107369>, 2021.
- 495 Abril, G. and Borges, A. V.: Carbon Dioxide and Methane Emissions from Estuaries, in: *Greenhouse Gas Emissions — Fluxes and Processes: Hydroelectric Reservoirs and Natural Environments*, edited by: Tremblay, A., Varfalvy, L., Roehm, C., and Garneau, M., Springer Berlin Heidelberg, Berlin, Heidelberg, 187–207, https://doi.org/10.1007/978-3-540-26643-3_8, 2005.
- Abril, G., Libardoni, B. G., Brandini, N., Cotovicz, L. C., Medeiros, P. R. P., Cavalcante, G. H., and Knoppers, B. A.: Thermodynamic uptake of atmospheric CO₂ in the oligotrophic and semiarid São Francisco estuary (NE Brazil), *Mar Chem*,
500 233, 103983, <https://doi.org/https://doi.org/10.1016/j.marchem.2021.103983>, 2021.
- Al-Haj, A. N. and Fulweiler, R. W.: A synthesis of methane emissions from shallow vegetated coastal ecosystems, *Glob Chang Biol*, 26, 2988–3005, <https://doi.org/https://doi.org/10.1111/gcb.15046>, 2020.
- Asmala, E. and Scheinin, M.: Persistent hot spots of CO₂ and CH₄ in coastal nearshore environments, *Limnol Oceanogr Lett*, 9, 119–127, <https://doi.org/https://doi.org/10.1002/lol2.10370>, 2024.

- 505 Attard, K. M., Rodil, I. F., Glud, R. N., Berg, P., Norkko, J., and Norkko, A.: Seasonal ecosystem metabolism across shallow benthic habitats measured by aquatic eddy covariance, *Limnol Oceanogr Lett*, 4, 79–86, <https://doi.org/https://doi.org/10.1002/lol2.10107>, 2019.
- Bange, H. W.: Nitrous oxide and methane in European coastal waters, *Estuar Coast Shelf Sci*, 70, 361–374, <https://doi.org/https://doi.org/10.1016/j.ecss.2006.05.042>, 2006.
- 510 Bauer, J. E., Cai, W. J., Raymond, P. A., Bianchi, T. S., Hopkinson, C. S., and Regnier, P. A. G.: The changing carbon cycle of the coastal ocean, <https://doi.org/10.1038/nature12857>, 2013.
- Bonaglia, S., Cheung, H. L. S., Politi, T., Vybernaite-Lubiene, I., McKenzie, T., Santos, I. R., and Zilius, M.: Eutrophication and urbanization enhance methane emissions from coastal lagoons, *Limnol Oceanogr Lett*, 10, 140–150, <https://doi.org/https://doi.org/10.1002/lol2.10430>, 2025.
- 515 Borges, A. V., Delille, B., Schiettecatte, L.-S., Gazeau, F., Abril, G., and Frankignoulle, M.: Gas transfer velocities of CO₂ in three European estuaries (Randers Fjord, Scheldt, and Thames), *Limnol Oceanogr*, 49, 1630–1641, <https://doi.org/https://doi.org/10.4319/lo.2004.49.5.1630>, 2004.
- Borges, A. V., Delille, B., and Frankignoulle, M.: Budgeting sinks and sources of CO₂ in the coastal ocean: Diversity of ecosystems counts, *Geophys Res Lett*, 32, <https://doi.org/https://doi.org/10.1029/2005GL023053>, 2005.
- 520 Borges, A. V., Champenois, W., Gypens, N., Delille, B., and Harlay, J.: Massive marine methane emissions from near-shore shallow coastal areas, *Sci Rep*, 6, 27908, <https://doi.org/10.1038/srep27908>, 2016.
- Borges, A. V., Abril, G., and Bouillon, S.: Carbon dynamics and CO₂ and CH₄ outgassing in the Mekong delta, *Biogeosciences*, 15, 1093–1114, <https://doi.org/10.5194/bg-15-1093-2018>, 2018.
- 525 [Brase, L., Bange, H. W., Lendt, R., Sanders, T., and Dähnke, K.: High Resolution Measurements of Nitrous Oxide \(N₂O\) in the Elbe Estuary, *Front Mar Sci*, Volume 4-2017, 2017.](#)
- [Brown, A. M., Bass, A. M., and Pickard, A. E.: Anthropogenic-estuarine interactions cause disproportionate greenhouse gas production: A review of the evidence base, *Mar Pollut Bull*, 174, 113240, <https://doi.org/https://doi.org/10.1016/j.marpolbul.2021.113240>, 2022.](#)
- 530 [Bunting, P., Rosenqvist, A., Lucas, R. M., Rebelo, L.-M., Hilarides, L., Thomas, N., Hardy, A., Itoh, T., Shimada, M., Finlayson, C. M.: The Global Mangrove Watch – A New 2010 Global Baseline of Mangrove Extent. *Remote Sens.* 10, 1669. <https://doi.org/10.3390/rs10101669>, 2018.](#)
- Cai, W.-J., Arthur Chen, C. T., and Borges, A.: Carbon dioxide dynamics and fluxes in coastal waters influenced by river plumes, in: *Biogeochemical Dynamics at Major River-Coastal Interfaces: Linkages with Global Change*, edited by: Bianchi,

- T. S., Allison, M. A., and Cai, W.-J., Cambridge University Press, Cambridge, 155–173, <https://doi.org/DOI:10.1017/CBO9781139136853.010>, 2013.
- 535 [Cardinale, B. J., Srivastava, D. S., Emmett Duffy, J., Wright, J. P., Downing, A. L., Sankaran, M., and Jouseau, C.: Effects of biodiversity on the functioning of trophic groups and ecosystems, Nature, 443, 989–992, https://doi.org/10.1038/nature05202, 2006.](https://doi.org/10.1038/nature05202)
- 540 [Carrasco, C., Karstensen, J., and Farias, L.: On the Nitrous Oxide Accumulation in Intermediate Waters of the Eastern South Pacific Ocean, Front Mar Sci, Volume 4-2017, 2017.](https://doi.org/10.1007/s12237-014-9829-y)
- Chen, C.-T. A. and Borges, A. V: Reconciling opposing views on carbon cycling in the coastal ocean: Continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO₂, Deep Sea Research Part II: Topical Studies in Oceanography, 56, 578–590, <https://doi.org/https://doi.org/10.1016/j.dsr2.2009.01.001>, 2009.
- Cheung, H. L. S., Zilius, M., Politi, T., Lorre, E., Vybernaite-Lubiene, I., Santos, I. R., and Bonaglia, S.: Nitrate-Driven Eutrophication Supports High Nitrous Oxide Production and Emission in Coastal Lagoons, J Geophys Res Biogeosci, 130, e2024JG008510, <https://doi.org/https://doi.org/10.1029/2024JG008510>, 2025.
- 545 Cloern, J. E., Foster, S. Q., and Kleckner, A. E.: Phytoplankton primary production in the world’s estuarine-coastal ecosystems, Biogeosciences, 11, 2477–2501, <https://doi.org/10.5194/bg-11-2477-2014>, 2014.
- Dickson, A. G.: Thermodynamics of the dissociation of boric acid in synthetic seawater from 273.15 to 318.15 K, Deep Sea Research Part A. Oceanographic Research Papers, 37, 755–766, [https://doi.org/https://doi.org/10.1016/0198-0149\(90\)90004-F](https://doi.org/https://doi.org/10.1016/0198-0149(90)90004-F), 1990.
- 550 Elliott, M. and Whitfield, A. K.: Challenging paradigms in estuarine ecology and management, Estuar Coast Shelf Sci, 94, 306–314, <https://doi.org/https://doi.org/10.1016/j.ecss.2011.06.016>, 2011.
- Fleming-Lehtinen, V., Räike, A., Kortelainen, P., Kauppila, P., and Thomas, D. N.: Organic Carbon Concentration in the Northern Coastal Baltic Sea between 1975 and 2011, Estuaries and Coasts, 38, 466–481, <https://doi.org/10.1007/s12237-014-9829-y>, 2015.
- 555 Frankignoulle, M.: Field measurements of air-sea CO₂ exchange, Limnol Oceanogr, 33, 313–322, <https://doi.org/https://doi.org/10.4319/lo.1988.33.3.0313>, 1988.
- Frankignoulle, M., Abril, G., Borges, A., Bourge, I., Canon, C., Delille, B., Libert, E., and Théate, J.-M.: Carbon Dioxide Emission from European Estuaries, Science (1979), 282, 434–436, <https://doi.org/10.1126/science.282.5388.434>, 1998.
- 560 Friis, K., Körtzinger, A., and Wallace, D. W. R.: The salinity normalization of marine inorganic carbon chemistry data, Geophys Res Lett, 30, <https://doi.org/https://doi.org/10.1029/2002GL015898>, 2003.

Gammal, J., Järnström, M., Bernard, G., Norkko, J., and Norkko, A.: Environmental Context Mediates Biodiversity–Ecosystem Functioning Relationships in Coastal Soft-sediment Habitats, *Ecosystems*, 22, 137–151, 565 <https://doi.org/10.1007/s10021-018-0258-9>, 2019.

Gattuso, J.-P., Frankignoulle, M., and Wollast, R.: Carbon and Carbonate Metabolism in Coastal Aquatic Ecosystems, *Annu Rev Ecol Syst*, 29, 405–434, 1998.

Gran, G.: Determination of the equivalence point in potentiometric titrations. Part II, *Analyst*, 77, 661–671, <https://doi.org/10.1039/AN9527700661>, 1952.

570 Gustafsson, C. and Norkko, A.: Quantifying the importance of functional traits for primary production in aquatic plant communities, *Journal of Ecology*, 107, 154–166, <https://doi.org/https://doi.org/10.1111/1365-2745.13011>, 2019.

HELCOM: Red List of Baltic Sea underwater biotopes, habitats and biotope complexes. *Baltic Sea Environmental Proceedings* No. 138, 2013.

575 Hoegh-Guldberg, O., Jacob, D., Taylor, M., Guillén Bolaños, T., Bindi, M., Brown, S., Camilloni, I. A., Diedhiou, A., Djalante, R., Ebi, K., Engelbrecht, F., Guiot, J., Hijioka, Y., Mehrotra, S., Hope, C. W., Payne, A. J., Pörtner, H.-O., Seneviratne, S. I., Thomas, A., Warren, R., and Zhou, G.: The human imperative of stabilizing global climate change at 1.5°C, *Science* (1979), 365, eaaw6974, <https://doi.org/10.1126/science.aaw6974>, 2019.

[Holland, K. T. and Elmore, P. A.: A review of heterogeneous sediments in coastal environments, *Earth Sci Rev*, 89, 116–134, <https://doi.org/https://doi.org/10.1016/j.earscirev.2008.03.003>, 2008.](https://doi.org/https://doi.org/10.1016/j.earscirev.2008.03.003)

580 Hsu, S. A., Meindl, E. A., and Gilhousen, D. B.: Determining the Power-Law Wind-Profile Exponent under Near-Neutral Stability Conditions at Sea, *J Appl Meteorol Climatol*, 33, 757–765, [https://doi.org/https://doi.org/10.1175/1520-0450\(1994\)033<0757:DTPLWP>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0450(1994)033<0757:DTPLWP>2.0.CO;2), 1994.

585 Humborg, C., Geibel, Marc. C., Sun, X., McCrackin, M., Mörth, C.-M., Stranne, C., Jakobsson, M., Gustafsson, B., Sokolov, A., Norkko, A., and Norkko, J.: High Emissions of Carbon Dioxide and Methane From the Coastal Baltic Sea at the End of a Summer Heat Wave, *Front Mar Sci*, Volume 6-2019, <https://doi.org/10.3389/fmars.2019.00493>, 2019.

IPCC, 2023: Sections. In: *Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II, and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland, pp. 35-115, doi: 10.59327/IPCC/AR6-9789291691647, 2023.

590 James, K., Macreadie, P. I., Burdett, H. L., Davies, I., and Kamenos, N. A.: It's time to broaden what we consider a 'blue carbon ecosystem,' *Glob Chang Biol*, 30, e17261, <https://doi.org/https://doi.org/10.1111/gcb.17261>, 2024.

[Kock, A., Arévalo-Martínez, D. L., Löscher, C. R., and Bange, H. W.: Extreme N₂O accumulation in the coastal oxygen minimum zone off Peru, *Biogeosciences*, 13, 827–840, <https://doi.org/10.5194/bg-13-827-2016>, 2016.](https://doi.org/https://doi.org/10.5194/bg-13-827-2016)

- Kristensen, E., Flindt, M. R., and Quintana, C. O.: Predicting Climate Mitigation Through Carbon Burial in Blue Carbon Ecosystems—Challenges and Pitfalls, *Glob Chang Biol*, 31, e70022, <https://doi.org/https://doi.org/10.1111/gcb.70022>, 2025.
- 595 Lammerant, R., Hölttä, J., Pykäri, J., Nishant, N., Villnäs, A., Wikström, S. A., Norkko, A., and Gustafsson, C.: Environmental gradients strongly affect functional composition and biomass C stocks within aquatic plant meadows, *Ecosphere*, 16, e70327, <https://doi.org/https://doi.org/10.1002/ecs2.70327>, 2025.
- Lazar, B. and Loya, Y.: Bioerosion of coral reefs-A chemical approach, *Limnol Oceanogr*, 36, 377–383, <https://doi.org/https://doi.org/10.4319/lo.1991.36.2.0377>, 1991.
- 600 Lovelock, C. E. and Duarte, C. M.: Dimensions of Blue Carbon and emerging perspectives, *Biol Lett*, 15, 20180781, <https://doi.org/10.1098/rsbl.2018.0781>, 2019.
- Lundevall-Zara, M., Lundevall-Zara, E., and Brüchert, V.: Sea-Air Exchange of Methane in Shallow Inshore Areas of the Baltic Sea, *Front Mar Sci*, Volume 8-2021, <https://doi.org/10.3389/fmars.2021.657459>, 2021.
- Macreadie, P. I., Costa, M. D. P., Atwood, T. B., Friess, D. A., Kelleway, J. J., Kennedy, H., Lovelock, C. E., Serrano, O., and
605 Duarte, C. M.: Blue carbon as a natural climate solution, *Nat Rev Earth Environ*, 2, 826–839, <https://doi.org/10.1038/s43017-021-00224-1>, 2021.
- Mäkelin, S., Lewandowska, A. M., Rodil, I. F., Karlson, A. M. L., Humborg, C., and Villnäs, A.: Linking Resource Quality and Biodiversity to Benthic Ecosystem Functions Across a Land-to-Sea Gradient, *Ecosystems*, 27, 329–345, <https://doi.org/10.1007/s10021-023-00891-9>, 2024.
- 610 McKenzie, L. J., Nordlund, L. M., Jones, B. L., Cullen-Unsworth, L. C., Roelfsema, C., and Unsworth, R. K. F.: The global distribution of seagrass meadows, *Environmental Research Letters*, 15, 74041, <https://doi.org/10.1088/1748-9326/ab7d06>, 2020.
- Mcleod, E., Chmura, G. L., Bouillon, S., Salm, R., Björk, M., Duarte, C. M., Lovelock, C. E., Schlesinger, W. H., and Silliman, B. R.: A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering
615 CO₂, *Front Ecol Environ*, 9, 552–560, <https://doi.org/https://doi.org/10.1890/110004>, 2011.
- Millero, F. J., Graham, T. B., Huang, F., Bustos-Serrano, H., and Pierrot, D.: Dissociation constants of carbonic acid in seawater as a function of salinity and temperature, *Mar Chem*, 100, 80–94, <https://doi.org/https://doi.org/10.1016/j.marchem.2005.12.001>, 2006.
- 620 [Murray, R. H., Erler, D. V., and Eyre, B. D.: Nitrous oxide fluxes in estuarine environments: response to global change, *Glob Chang Biol*, 21, 3219–3245, https://doi.org/https://doi.org/10.1111/gcb.12923, 2015.](https://doi.org/https://doi.org/10.1111/gcb.12923)
- Le Quéré, C., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Peters, G. P., Manning, A. C., Boden, T. A., Tans, P. P., Houghton, R. A., Keeling, R. F., Alin, S., Andrews, O. D., Anthoni, P., Barbero, L., Bopp, L., Chevallier, F., Chini, L.

- P., Ciais, P., Currie, K., Delire, C., Doney, S. C., Friedlingstein, P., Gkritzalis, T., Harris, I., Hauck, J., Haverd, V., Hoppema, M., Klein Goldewijk, K., Jain, A. K., Kato, E., Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D., Melton, J. R., Metzl, N., Millero, F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S., O'Brien, K., Olsen, A., Omar, A. M., Ono, T., Pierrot, D., Poulter, B., Rödenbeck, C., Salisbury, J., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Stocker, B. D., Sutton, A. J., Takahashi, T., Tian, H., Tilbrook, B., van der Laan-Luijkx, I. T., van der Werf, G. R., Viogy, N., Walker, A. P., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget 2016, *Earth Syst Sci Data*, 8, 605–649, <https://doi.org/10.5194/essd-8-605-2016>, 2016.
- 625
- 630 Räike, A., Kortelainen, P., Mattsson, T., and Thomas, D. N.: 36-year trends in dissolved organic carbon export from Finnish rivers to the Baltic Sea, *Science of The Total Environment*, 435–436, 188–201, <https://doi.org/https://doi.org/10.1016/j.scitotenv.2012.06.111>, 2012.
- Raymond, P. A., Bauer, J. E., and Cole, J. J.: Atmospheric CO₂ evasion, dissolved inorganic carbon production, and net heterotrophy in the York River estuary, *Limnol Oceanogr*, 45, 1707–1717, <https://doi.org/https://doi.org/10.4319/lo.2000.45.8.1707>, 2000.
- 635
- Reeburgh, W. S.: Oceanic Methane Biogeochemistry, *Chem Rev*, 107, 486–513, <https://doi.org/10.1021/cr050362v>, 2007.
- Regnier, P., Resplandy, L., Najjar, R. G., and Ciais, P.: The land-to-ocean loops of the global carbon cycle, *Nature*, 603, 401–410, <https://doi.org/10.1038/s41586-021-04339-9>, 2022.
- Resplandy, L., Hogikyan, A., Müller, J. D., Najjar, R. G., Bange, H. W., Bianchi, D., Weber, T., Cai, W.-J., Doney, S. C., Fennel, K., Gehlen, M., Hauck, J., Lacroix, F., Landschützer, P., Le Quéré, C., Roobaert, A., Schwinger, J., Berthet, S., Bopp, L., Chau, T. T. T., Dai, M., Gruber, N., Ilyina, T., Kock, A., Manizza, M., Lachkar, Z., Laruelle, G. G., Liao, E., Lima, I. D., Nissen, C., Rödenbeck, C., Séférian, R., Toyama, K., Tsujino, H., and Regnier, P.: A Synthesis of Global Coastal Ocean Greenhouse Gas Fluxes, *Global Biogeochem Cycles*, 38, e2023GB007803, <https://doi.org/https://doi.org/10.1029/2023GB007803>, 2024.
- 640
- 645 Rodil, I. F., Attard, K. M., Gustafsson, C., and Norkko, A.: Variable contributions of seafloor communities to ecosystem metabolism across a gradient of habitat-forming species, *Mar Environ Res*, 167, 105321, <https://doi.org/https://doi.org/10.1016/j.marenvres.2021.105321>, 2021.
- Rosentreter, J. A., Maher, D. T., Erler, D. V., Murray, R., and Eyre, B. D.: Factors controlling seasonal CO₂ and CH₄ emissions in three tropical mangrove-dominated estuaries in Australia, *Estuar Coast Shelf Sci*, 215, 69–82, <https://doi.org/https://doi.org/10.1016/j.ecss.2018.10.003>, 2018.
- 650
- Rosentreter, J. A., Borges, A. V., Deemer, B. R., Holgerson, M. A., Liu, S., Song, C., Melack, J., Raymond, P. A., Duarte, C. M., Allen, G. H., Olefeldt, D., Poulter, B., Battin, T. I., and Eyre, B. D.: Half of global methane emissions come from highly variable aquatic ecosystem sources, *Nat Geosci*, 14, 225–230, <https://doi.org/10.1038/s41561-021-00715-2>, [2021](https://doi.org/10.1038/s41561-021-00715-2)2021a.

- 655 [Rosentreter, J. A., Al-Haj, A. N., Fulweiler, R. W., and Williamson, P.: Methane and Nitrous Oxide Emissions Complicate Coastal Blue Carbon Assessments, *Global Biogeochem Cycles*, 35, e2020GB006858, <https://doi.org/https://doi.org/10.1029/2020GB006858>, 2021b.](https://doi.org/https://doi.org/10.1029/2020GB006858)
- Roth, F., Sun, X., Geibel, M. C., Prytherch, J., Brüchert, V., Bonaglia, S., Broman, E., Nascimento, F., Norkko, A., and Humborg, C.: High spatiotemporal variability of methane concentrations challenges estimates of emissions across vegetated coastal ecosystems, *Glob Chang Biol*, 28, 4308–4322, <https://doi.org/https://doi.org/10.1111/gcb.16177>, 2022.
- 660 Roth, F., Broman, E., Sun, X., Bonaglia, S., Nascimento, F., Prytherch, J., Brüchert, V., Lundevall Zara, M., Brunberg, M., Geibel, M. C., Humborg, C., and Norkko, A.: Methane emissions offset atmospheric carbon dioxide uptake in coastal macroalgae, mixed vegetation and sediment ecosystems, *Nat Commun*, 14, 42, <https://doi.org/10.1038/s41467-022-35673-9>, 2023.
- Sharp, J. D., Pierrot, D., Humphreys, M. P., Epitalon, J.-M., Orr, J. C., Lewis, E. R., Wallace, D. W. R.: CO2SYSv3 for MATLAB (Version v3.2.1). Zenodo. <http://doi.org/10.5281/zenodo.3950562>, 2023.
- 665 Snelgrove, P. V. R., Thrush, S. F., Wall, D. H., and Norkko, A.: Real world biodiversity–ecosystem functioning: a seafloor perspective, *Trends Ecol Evol*, 29, 398–405, <https://doi.org/https://doi.org/10.1016/j.tree.2014.05.002>, 2014.
- [Snelgrove, P. V. R., Soetaert, K., Solan, M., Thrush, S., Wei, C.-L., Danovaro, R., Fulweiler, R. W., Kitazato, H., Ingole, B., Norkko, A., Parkes, R. J., and Volkenborn, N.: Global Carbon Cycling on a Heterogeneous Seafloor, *Trends Ecol Evol*, 33, 96–105, <https://doi.org/https://doi.org/10.1016/j.tree.2017.11.004>, 2018.](https://doi.org/https://doi.org/10.1016/j.tree.2017.11.004)
- 670 [Stein, A., Gerstner, K., and Krefth, H.: Environmental heterogeneity as a universal driver of species richness across taxa, biomes and spatial scales, *Ecol Lett*, 17, 866–880, <https://doi.org/https://doi.org/10.1111/ele.12277>, 2014.](https://doi.org/https://doi.org/10.1111/ele.12277)
- Testa, J. M., Kemp, W. M., Hopkinson Jr., C. S., and Smith, S. V: Ecosystem Metabolism, in: *Estuarine Ecology*, 381–416, <https://doi.org/https://doi.org/10.1002/9781118412787.ch15>, 2012.
- 675 Tyrrell, T., Schneider, B., Charalampopoulou, A., and Riebesell, U.: Coccolithophores and calcite saturation state in the Baltic and Black Seas, *Biogeosciences*, 5, 485–494, <https://doi.org/10.5194/bg-5-485-2008>, 2008.
- Uth, C., Asmala, E., and Lewandowska, A. M.: Phytoplankton community composition as a driver of annual autochthonous organic carbon dynamics in the northern coastal Baltic Sea, *Mar Ecol Prog Ser*, 745, 13–24, 2024.
- Wallenius, A. J., Dalcin Martins, P., Slomp, C. P., and Jetten, M. S. M.: Anthropogenic and Environmental Constraints on the Microbial Methane Cycle in Coastal Sediments, *Front Microbiol*, Volume 12-2021, <https://doi.org/10.3389/fmicb.2021.631621>, 2021.
- 680

Wan, X. S., Sheng, H.-X., Liu, L., Shen, H., Tang, W., Zou, W., Xu, M. N., Zheng, Z., Tan, E., Chen, M., Zhang, Y., Ward, B. B., and Kao, S.-J.: Particle-associated denitrification is the primary source of N₂O in oxic coastal waters, Nat Commun, 14, 8280, <https://doi.org/10.1038/s41467-023-43997-3>, 2023.

685 Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnol Oceanogr Methods*, 12, 351–362, <https://doi.org/https://doi.org/10.4319/lom.2014.12.351>, 2014.

Ward, M., Saura, S., Williams, B., Ramírez-Delgado, J. P., Arafeh-Dalmau, N., Allan, J. R., Venter, O., Dubois, G., and Watson, J. E. M.: Just ten percent of the global terrestrial protected area network is structurally connected via intact land, *Nat Commun*, 11, 4563, <https://doi.org/10.1038/s41467-020-18457-x>, 2020.

690 Watson, A. J., Schuster, U., Shutler, J. D., Holding, T., Ashton, I. G. C., Landschützer, P., Woolf, D. K., and Goddijn-Murphy, L.: Revised estimates of ocean-atmosphere CO₂ flux are consistent with ocean carbon inventory, *Nat Commun*, 11, 4422, <https://doi.org/10.1038/s41467-020-18203-3>, 2020.

Weiss, R. F.: Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar Chem*, 2, 203–215, [https://doi.org/https://doi.org/10.1016/0304-4203\(74\)90015-2](https://doi.org/https://doi.org/10.1016/0304-4203(74)90015-2), 1974.

695 Weiss, R. F. and Price, B. A.: Nitrous oxide solubility in water and seawater, *Mar Chem*, 8, 347–359, [https://doi.org/https://doi.org/10.1016/0304-4203\(80\)90024-9](https://doi.org/https://doi.org/10.1016/0304-4203(80)90024-9), 1980.

Wiesenburg, D. A. and Guinasso, N. L. Jr.: Equilibrium solubilities of methane, carbon monoxide, and hydrogen in water and sea water, *J Chem Eng Data*, 24, 356–360, <https://doi.org/10.1021/jc60083a006>, 1979.

Williamson, P. and Gattuso, J.-P.: Carbon Removal Using Coastal Blue Carbon Ecosystems Is Uncertain and Unreliable, With
700 Questionable Climatic Cost-Effectiveness, *Frontiers in Climate*, Volume 4-2022, 2022.

Wollast, R.: Evaluation and comparison of the global carbon cycle in the coastal zone and in the open ocean. p., *The Sea*, Vol. 10, 213–252, 1998.

Yvon-Durocher, G., Allen, A. P., Bastviken, D., Conrad, R., Gudasz, C., St-Pierre, A., Thanh-Duc, N., and del Giorgio, P. A.: Methane fluxes show consistent temperature dependence across microbial to ecosystem scales, *Nature*, 507, 488–491,
705 <https://doi.org/10.1038/nature13164>, 2014.

Zeebe, R. and Wolf-Gladrow, D.: CO₂ in Seawater: Equilibrium, kinetics, isotopes, 2001.

Figure 1: Atmospheric conditions: **a)** air temperature ($^{\circ}\text{C}$) and **b)** wind speed (m/s) recorded at the ICOS station FI-Tvm established at the Tvärminne Zoological Station. Spatial variation of **c)** surface seawater temperature (SST, $^{\circ}\text{C}$), and **d)** salinity (SSS), **e)** chlorophyll-*a* (Chl-*a*, $\mu\text{g L}^{-1}$) concentration, **f)** turbidity (FTU), **g)** colored dissolved organic carbon (CDOM, $\mu\text{g L}^{-1}$), and **h)** dissolved oxygen saturation (%) across the sampling area. Triangles represent the surface water conditions at the mouth of the Karjaanjoki River.




Figure 2: Left column presents the spatial variation of surface seawater **a)** $p\text{CO}_2$ (μatm), **b)** CH_4 (nmol L^{-1}), and **c)** N_2O concentration (nmol L^{-1}). The middle column presents measured air-sea fluxes of **d)** CO_2 ($\text{mmol m}^{-2} \text{d}^{-1}$) and **e)** CH_4 ($\text{mmol m}^{-2} \text{d}^{-1}$). The right column presents the calculated air-sea fluxes of **f)** CO_2 ($\text{mmol m}^{-2} \text{d}^{-1}$), **g)** CH_4 ($\text{mmol m}^{-2} \text{d}^{-1}$), and **h)** N_2O ($\text{mmol m}^{-2} \text{d}^{-1}$). Positive values indicate a release from the sea to the atmosphere, while negative values represent an uptake by the sea. Triangles represent the surface water conditions at the mouth of the Karjaanjoki River.




Figure 3: **a)** Relationship between salinity, TA ($\mu\text{mol kg}^{-1}$, circles) and DIC ($\mu\text{mol kg}^{-1}$, squares). Relationship between nDIC and nTA, and their association with **b)** pCO_2 (μatm), **c)** CH_4 (nmol L^{-1}), and **d)** N_2O (nmol L^{-1}) concentrations. Black symbols represent the seawater endmember. Exposed and semi-sheltered sites are represented by circles, while sheltered sites are represented by square symbols.




Figure 4: Heatmap of Kendall correlation coefficient **a)** between surface water pCO₂, CH₄, and N₂O concentration, and **b)** between main seawater biogeochemical properties (SST, SSS, chl-*a* concentration, colored dissolved organic carbon, turbidity, and O₂ saturation) and pCO₂, CH₄, and N₂O concentration.




Figure 5: Relationship between surface seawater salinity and **a)** pCO₂ (μatm), **b)** CH₄ (nmol L⁻¹), and **c)** N₂O (nmol L⁻¹) concentrations. In each panel, the dotted line represents the regression across all sites, while the red dashed line shows the regression limited to exposed and semi-sheltered sites. The blue line in panel **a)** represents the estimated pCO₂ resulting only from the mixing of the Karjaanjoki River water (S = 0) with seawater (S = 6.42).




Figure 6: Relationship between the apparent oxygen utilization (AOU) and **a)** surface water Chl-*a* concentration ($\mu\text{g L}^{-1}$), **b)** surface water pCO_2 (in μatm), **c)** CH_4 (nmol L^{-1}), and **d)** N_2O (nmol L^{-1}) associated with the Chl-*a* concentration ($\mu\text{g L}^{-1}$). In each panel, the dotted line represents the regression across all sites, while the red dashed line shows the regression limited to exposed and semi-sheltered sites. Exposed and semi-sheltered sites are represented by circles, while sheltered sites are represented by square symbols.




Figure 7: Air-sea fluxes of **a)** CO_2 ($\text{g m}^{-2} \text{d}^{-1}$), and CO_2 -equivalent fluxes of **b)** CH_4 ($\text{g m}^{-2} \text{d}^{-1}$), and **c)** N_2O ($\text{g m}^{-2} \text{d}^{-1}$) estimated using their respective sustained-flux warming potential (SGWPs) over a 100-year time horizon ($\text{CH}_4 = 27$; $\text{N}_2\text{O} = 273$, relative to $\text{CO}_2 = 1$, IPCC 2023), and **d)** the overall budget for all three GHGs ($\text{g m}^{-2} \text{d}^{-1}$).

