

Revisiting the global budget of atmospheric glyoxal: updates on terrestrial and marine precursor emissions, chemistry, and impacts on atmospheric oxidation capacity

Aoxing Zhang^{1,2}, Tzung-May Fu^{1,2,3}, Yuhang Wang⁴, Enyu Xiong^{1,2,5}, Wenlu Wu^{1,2,6}, Yumin Li⁷, Lei Zhu^{1,2}, Wei Tao^{1,2}, Kelley C. Wells⁸, Dylan B. Millet⁸, Zhe Wang⁹, Bin Yuan¹⁰, Min Shao¹⁰, Christophe Lerot^{11,12}, Thomas Danckaert¹¹, Ruixiong Zhang¹³, and Kelvin H. Bates¹⁴

¹State Key Laboratory of Soil Pollution Control and Safety, Shenzhen Key Laboratory of Precision Measurement and Early Warning Technology for Urban Environmental Health Risks, School of Environmental Science and Engineering, Southern University of Science and Technology, Shenzhen, Guangdong 518055, China

²Guangdong Provincial Field Observation and Research Station for Coastal Atmosphere and Climate of the Greater Bay Area, Southern University of Science and Technology, Shenzhen, Guangdong 518055, China

³National Center for Applied Mathematics Shenzhen, Shenzhen, Guangdong 518055, China

⁴School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA 30342, USA

⁵Department of Civil and Environmental Engineering, The Hong Kong Polytechnic University, Hong Kong SAR 999077, China

⁶Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds LS2 9JT, UK

⁷Institute for Atmospheric and Climate Science, ETH Zurich, Zurich 8092, Switzerland

⁸Department of Soil, Water, and Climate, University of Minnesota, St Paul, MN 55108, USA

⁹Division of Environment and Sustainability, The Hong Kong University of Science and Technology, Hong Kong SAR, China

¹⁰Institute for Environmental and Climate Research, Jinan University, Guangzhou, Guangdong 511443, China

¹¹Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels 1180, Belgium

¹²Now at constellr S.A., Li-e 4031, Belgium

¹³Atlassian US Inc., San Francisco, CA 94104, USA

¹⁴Department of Mechanical Engineering, University of Colorado Boulder, Boulder, CO 80309, USA

Correspondence: Tzung-May Fu (fuzm@sustech.edu.cn) and Yuhang Wang (yuhang.wang@eas.gatech.edu)

Abstract. Glyoxal Atmospheric glyoxal (CHOCHO); ~~the smallest dicarbonyl~~, plays critical yet incompletely understood roles in tropospheric chemistry. Current models substantially underestimate glyoxal abundance over both land and ocean, indicating knowledge gaps in our understanding of its sources and sinks. Here, we present an improved global simulation of atmospheric glyoxal using the GEOS-Chem model, advanced by recent theoretical, experimental, and observational insights on precursor emissions, chemical pathways, and heterogeneous losses. By applying top-down-constrained biogenic ~~emissions~~; isoprene emissions, enhancing biomass burning emissions, and revising glyoxal yields from isoprene, monoterpenes, and glycolaldehyde oxidation, ~~and enhancing biomass burning emissions~~; we estimated a global atmospheric glyoxal source of ~~44~~ 40 Tg yr⁻¹ and a global burden of 15 Gg, substantially reducing ~~the normalized mean bias (NMB) low bias~~ of simulated glyoxal abundance ~~by more than 20% against in situ~~ against in situ and TROPOMI satellite observations over land. The improved representation ~~increases of glyoxal and its precursors increases simulated~~ global mean surface ozone by ~~1.3 ppb~~ (4.84.5 ppb (17%)) and SOA formation by ~~5.0~~ 17.9 Tg yr⁻¹ (~~3.8~~ 13%), indicating stronger atmospheric oxidation capacity. Further inclusion of a hypothetical secondary marine glyoxal ~~source~~ production to match TROPOMI glyoxal observations over the global oceans increased the

global glyoxal source to 110 source of atmospheric glyoxal to 106 Tg yr⁻¹ and the its global burden to 39 Gg, substantially improving agreement with in-situ *in situ* (NMB from -92% to 12%) and satellite observations (NMB from -88% to -6%) over the ocean. This enhanced marine glyoxal source increased surface HO₂ concentrations and OH reactivity over tropical oceans by 6.8% and 2.3%. Our work reconciles 6.5% and 1.9%. However, this hypothetical marine glyoxal source cannot be accounted for by known marine NMVOC emissions; its existence remains highly uncertain and warrants further investigation. Our work helped reconcile major model-measurement discrepancies for atmospheric glyoxal, enhancing its utility as a volatile organic compound (VOC) proxy and underscoring the need for accurate representation of glyoxal sources and chemistry in atmospheric models.

1 Introduction

Glyoxal (CHOCHO), the smallest dicarbonyl, is produced from the photo-oxidation of non-methane volatile organic compounds (NMVOCs), and plays critical and multifaceted roles in tropospheric chemistry (Fu et al., 2008; Myriokefalitakis et al., 2008; Seakins and Blitz, 2011). Glyoxal photolyzes to produce HO₂ radicals (Atkinson et al., 2006), thereby propagating atmospheric oxidation and ozone production (Edwards et al., 2014). Glyoxal can also heterogeneously react on surfaces of aqueous aerosols and cloud droplets to become part of secondary organic aerosol (SOA) (Liggio et al., 2005; Fu et al., 2008, 2009, 2012). Once in the aqueous phase, glyoxal may react with ammonium, amines, and amino acids to form light-absorbing organic nitrogenous molecules, which radiatively affect climate (Shapiro et al., 2009; De Haan et al., 2020; Li et al., 2023, 2025). In addition, satellite observations of tropospheric glyoxal column concentrations serve as a proxy for NMVOC emissions, offering species-differentiating precursor-differentiating information when analyzed in combination with formaldehyde—another widely used proxy (Stavrakou et al., 2009; DiGangi et al., 2012; Kaiser et al., 2015; Cao et al., 2018).

Despite its importance, the global budget of atmospheric glyoxal remains poorly understood, limiting its utility as a quantitative tracer for NMVOC emissions and hindering assessments of its chemical and climatic impacts. Early model studies by Fu et al. (2008) and Myriokefalitakis et al. (2008) estimated global atmospheric glyoxal sources of 45 to 56 Tg yr⁻¹, with 53% to 70% produced from the oxidation of terrestrial biogenic NMVOCs, predominantly isoprene. A more recent work by Müller et al. (2019) estimated a 47 Tg yr⁻¹ global glyoxal source but with smaller contribution from isoprene and larger contributions from monoterpenes and anthropogenic NMVOCs. In contrast, top-down constraints using glyoxal observations from the SCIAMACHY satellite instrument implied a much larger atmospheric glyoxal source of 107 Tg yr⁻¹ from terrestrial NMVOC emissions and photochemical production, with approximately 26% attributed to unidentified terrestrial biogenic precursors (Stavrakou et al., 2009). More recently, Sfindla et al. (2026) used TROPOMI satellite observations to inversely derive an 110 Tg yr⁻¹ global terrestrial source for atmospheric glyoxal. However, over 40% of this inferred terrestrial source could not be accounted for by known precursors and chemical mechanisms, highlighting the persistent gap in our understanding of atmospheric glyoxal sources.

Over the past decade, laboratory experiments and theoretical studies have advanced our understanding of the photochemistry of key glyoxal precursors, such as isoprene (Wennberg et al., 2018), aromatics (Xu et al., 2020), and other hydrocarbons (Teng

et al., 2015; Kwon et al., 2021). However, discrepancies persist between modeled and observed glyoxal abundances and budgets (Silva et al., 2018; Bates and Jacob, 2019; Ha et al., 2024). For example, the global mean glyoxal yield from isoprene oxidation simulated with a recently developed chemical mechanism was 2% (Wennberg et al., 2018; Bates and Jacob, 2019), markedly below previous laboratory measurements and mechanistic calculations (Fu et al., 2008; Galloway et al., 2011; Li et al., 2016; 50 Chan Miller et al., 2017; Müller et al., 2019; Ha et al., 2024). These discrepancies highlight critical gaps in our understanding of glyoxal's formation pathways from NMVOC precursors.

Another major uncertainty in the global budget of glyoxal is the undetermined oceanic contribution. A number of observations detected elevated concentrations of glyoxal in sea water samples (1.4 to 4.8 nmol L⁻¹) and in the marine boundary layer (MBL) air (5.6 to 80 ppt) (Zhou and Mopper, 1990a, b; Seaman et al., 2006; Sinreich et al., 2010; Coburn et al., 2014; 55 Mahajan et al., 2014; Lawson et al., 2015; Mungall et al., 2017; Zhu and Kieber, 2019; Kluge et al., 2023), with tentative correlations with the dissolved organic carbon (DOC) or chlorophyll-*a* contents of sea water (Zhu and Kieber, 2019). These correlations suggested a biogenic source of glyoxal or its precursor in the MBL, although the mechanistic pathways have remained unclear (Chiu et al., 2017). A few glyoxal precursors, such as isoprene, ethylene, and propene, have been observed in the remote MBL, but their reported concentrations or fluxes cannot explain the observed glyoxal abundance in the MBL (Plass- 60 Dülmer et al., 1995; Pound, 2021; Zhang et al., 2025; Ferracci et al., 2024; Li et al., 2021; Tripathi et al., 2020). Early satellite retrievals also indicated glyoxal enhancements in the tropical MBL, but there were concerns that those glyoxal retrievals may be confounded by water absorption artifacts (Wittrock et al., 2006; Alvarado et al., 2014). Recent satellite retrieval studies have mitigated water artifacts by incorporating temperature-dependent cross-sections and optimized spectral fitting windows, yet they still found glyoxal hotspots over biologically active tropical oceans (Alvarado et al., 2014; Lerot et al., 2021a). In 65 addition, aircraft observations found the enhanced glyoxal to be vertically well-mixed throughout the MBL (Zhou et al., 2014; Volkamer et al., 2015; Zhu and Kieber, 2019). Collectively, these observations form a line of evidence for substantial glyoxal production in the MBL air from yet unidentified marine precursors. The photochemical impact of this additional potential glyoxal production in the MBL have not been explored.

In this study, we developed an updated global atmospheric glyoxal simulation with the goal of better supporting atmospheric 70 chemistry research, as well as the use of glyoxal for inverting NMVOC emissions and for diagnosing photochemical regime. We incorporated recent experimental, observational, and theoretical knowledge advancements in four aspects: (1) the emissions of glyoxal and its precursors from terrestrial sources, (2) a potential marine photochemical source of glyoxal, (3) updated production pathways and yields of glyoxal from key precursors, and (4) mechanistic representations of heterogeneous uptake on cloud droplets and aqueous aerosols. We evaluated the simulated atmospheric glyoxal abundances against global surface, 75 aircraft, and satellite observations. Finally, we assessed the impacts of the revised atmospheric glyoxal budget on tropospheric oxidation capacity, ozone, and SOA formation globally.

2 Methods

2.1 Global atmospheric glyoxal simulations using the GEOS-Chem model

We developed our new global atmospheric glyoxal simulations using the GEOS-Chem global 3D chemical transport model v14.2.3 (<https://geoschem.github.io/>, last accessed: 18 November 2024) (Bey et al., 2001), superseding the obsolete glyoxal simulation in GEOS-Chem v7.3.6 by Fu et al. (2008). The model was driven by NASA's GEOS-FP assimilated meteorology (<https://fluid.nccs.nasa.gov/weather/><https://fluid.nccs.nasa.gov/weather>, last accessed: 30 July 2024, native resolution 0.3125° longitude \times 0.25° latitude), regridded to $5^\circ 2.5'$ \times $4^\circ 2'$ for efficient global simulations. GEOS-Chem vertically resolved 72 hybrid sigma pressure levels, with 8 layers in the bottom 1 km to resolve near-surface processes.

We conducted four sensitivity simulations to assess the impacts of precursor emissions and chemistry on the global atmospheric glyoxal budget:

- GC-CTRL : Control simulation using the standard GEOS-Chem v14.2.3, including its default precursor emissions and gas and aerosol chemistry (Section 2.2).
- GC-TM-E : Simulation incorporating updated terrestrial emissions of glyoxal and its precursors (Sections 4.1 and 4.2) and a hypothetical [marine glyoxal source](#) [secondary glyoxal source over the global oceans](#) (Section 4.3), with standard GEOS-Chem chemistry.
- GC-T-EC : Simulation incorporating updated terrestrial precursor emissions and revised glyoxal chemistry (Section 5); no marine glyoxal source.
- GC-TM-EC : Same as the GC-T-EC simulation but with the addition of [a the](#) hypothetical marine glyoxal source (Section 4.3).

All simulations were from January 2019 to June 2020, with the first six months used as model initialization. Results from July 2019 to June 2020 were analyzed.

2.2 Sources and sinks of glyoxal in the standard GEOS-Chem v14.2.3

In the standard GEOS-Chem v14.2.3, glyoxal was [represented as](#) both directly emitted and produced from several primary precursors, including isoprene, acetylene, ethylene, benzene, toluene, lumped xylenes, and glycolaldehyde (HOCH_2CHO , also an oxidation product of NMVOCs). GEOS-Chem [v14.2.3](#) also simulated the photochemistry of monoterpenes (Fisher et al., 2016) but did not represent their production of glyoxal. Global emissions of glyoxal and its precursors from anthropogenic, biomass burning, and biogenic sources are summarized in Table 1. Global monthly mean anthropogenic NMVOC emissions were from the Community Emissions Data System (CEDS version 2) (McDuffie et al., 2020) for the year 2019, [representing](#) [which included](#) emissions from power generation, industries, transportation, residential activities, agriculture, and biofuel use, [with the exception that ; however,](#) glyoxal and glycolaldehyde [are not included in biofuel emissions](#) [emissions from biofuel use](#) [were not included](#). To better capture regional characteristics, we superseded CEDS data with the following regional emission

inventories: the National Emissions Inventory for 2016 (NEI2016) over the continental United States (National Emissions Inventory Collaborative, 2019), the Multi-resolution emission inventory for China (MEIC) for 2019 (Zheng et al., 2018, 2021), and the MIX inventory for the rest of East and South Asia for the year 2010 (Li et al., 2017). Monthly biomass burning emissions of NMVOCs were from the Global Fire Emissions Database version 4 with small fires (GFED4s) (Van Der Werf et al., 2017) for the years 2019 and 2020. Biomass burning emissions of glyoxal and glycolaldehyde were estimated using GFED4s burned dry mass and fuel-specific emission factors from Akagi et al. (2011) and Carter et al. (2022). Biogenic emissions of NMVOCs were computed online using the MEGAN v2.1 algorithm (Guenther et al., 2012).

The standard GEOS-Chem v14.2.3 included a HO_x—O_x—OCs—zone—aerosol—erosol chemical mechanism (Park et al., 2004; Bates and Jacob, 2019). In particular, oxidation of isoprene, the largest known precursor of global glyoxal, was represented by the Reduced Caltech Isoprene Mechanism (RCIM) (Bates and Jacob, 2019). The RCIM is a condensed version of the explicit mechanism developed by Wennberg et al. (2018), which incorporated mechanistic understanding from laboratory and theoretical studies. RCIM described isoprene oxidation by OH, O₃, and NO₃, along with subsequent formation of major peroxy radicals (RO₂), ≥ C₄ carbonyls, and organic nitrates. However, Wennberg et al. (2018) and RCIM did not update the formation of small fragmented products such as glyoxal and glycolaldehyde. Consequently, [GEOS-Chem-simulated GEOS-Chem simulated](#) only 2% global glyoxal yield from isoprene oxidation using RCIM (Bates and Jacob, 2019), lower than previous global model estimations (Fu et al., 2008; Myriokefalitakis et al., 2008; Taraborrelli et al., 2009; Müller et al., 2019) and potentially contributing to the model's underestimation of glyoxal compared to global observations (Section 3.1). Bates and Jacob (2019) attributed this low global glyoxal yield partially to loss of isoprene's intermediate oxidation products to aerosol uptake or deposition. Evaluating this glyoxal yield from isoprene was a central objective of our study (Section 5.1).

In GEOS-Chem, gaseous glyoxal was chemically removed via oxidation by OH (rate constant $k_{\text{OH}} = 9.9 \times 10^{-12} \text{ cm}^3 \text{ molecules}^{-1}$) and by NO₃ ($k_{\text{NO}_3} = 1.5 \times 10^{-15} \text{ cm}^3 \text{ molecules}^{-1}$), photolysis, dry and wet deposition, and heterogeneous uptake. Photolysis rates were calculated using the Fast-JX scheme (Bian and Prather, 2002), which incorporated the latest IUPAC-recommended cross sections and quantum yields (<https://iupac.aeris-data.fr>) (Atkinson et al., 2006; IUPAC Subcommittee for Gas Kinetic Data Evaluation, 2013). Dry and wet depositions of gases and aerosols were as parameterized in Wesely (1989) and Amos et al. (2012). The Henry's law constants for glyoxal ($4.19 \times 10^5 \text{ M atm}^{-1}$) and glycolaldehyde ($4.14 \times 10^4 \text{ M atm}^{-1}$) were from Ip et al. (2009) and Betterton and Hoffmann (1988), respectively. In addition, glyoxal can be heterogeneously and irreversibly uptaken by cloud droplets and wet aerosols (Fu et al., 2008; Curry et al., 2018). The standard GEOS-Chem v14.2.3 used a daytime uptake coefficient $\gamma = 4.4 \times 10^{-3}$ and a nighttime uptake coefficient of 8.0×10^{-6} , both applied only when the ambient relative humidity (RH) was >35% (Marais et al., 2016).

2.3 DSMACC box model

We used the Dynamically Simple Model of Atmospheric Chemical Complexity (DSMACC) box model (Emmerson and Evans, 2009; Stone et al., 2010) to evaluate glyoxal production from isoprene as represented in three mechanisms: (1) the RCIM as implemented in GEOS-Chem v14.2.3, (2) the Master Chemical Mechanism (MCM) version 3.3.1 (Jenkin et al., 2015), and (3) our updates to the RCIM (Section 5.1). Simulations were configured to represent a subtropical site during fall (Zou et al.,

2023). All non-isoprene gas-phase reactions and all photolysis were identical to those in the standard GEOS-Chem v14.2.3. Deposition and aerosol uptake processes were disabled. Additional details regarding the DSMACC simulations are given in Text S1.

145 2.4 Satellite, aircraft, and surface observations of atmospheric glyoxal and isoprene

We used the tropospheric glyoxal ~~column concentrations~~ vertical column densities (VCDs) observed by the TROPospheric Monitoring Instrument (TROPOMI) during July 2019 to June 2020 to evaluate our global atmospheric glyoxal simulations and to infer ~~a potential~~ the magnitude of a hypothetical secondary glyoxal source in the MBL (Section 4.3). TROPOMI is onboard the Copernicus Sentinel-5 Precursor satellite, which flies in a sun-synchronous orbit over-passing the Equator at 13:30 local
150 time (Veeffkind et al., 2012). Details of the TROPOMI glyoxal retrieval can be found in Alvarado et al. (2020) and Lerot et al. (2021a). Briefly, slant column densities (SCDs) of glyoxal were fitted in the 435 nm to 460 nm window using the Differential Optical Absorption Spectroscopy (DOAS) approach (Platt and Stutz, 2008), accounting for other absorbers in that wavelength range. In particular, absorption by water vapor and liquid water were quantified using optimized, temperature-dependent cross-sections (Lerot et al., 2021a). The glyoxal SCDs were then converted to tropospheric ~~column concentrations~~ VCDs using
155 air mass factors (AMFs), accounting for the radiative transfer through the atmosphere (Palmer et al., 2001). Over land, the TROPOMI retrievals used the *a priori* glyoxal profiles simulated by the Model of Atmospheric composition at Global and Regional scales using Inversion Techniques for Trace gas Emissions (MAGRITTE v1.1, Müller et al. (2019)). Over oceans, a fixed parameterized glyoxal profile based on aircraft observations over the tropical Pacific from Volkamer et al. (2015) was used in the retrieval (Lerot et al., 2021a). The total systematic errors of the tropospheric glyoxal ~~column concentration~~ VCD
160 retrievals were estimated to be between 1×10^{14} and 3×10^{14} molecules cm^{-2} , corresponding to 30% to 70% of the signals over source regions (Lerot et al., 2021a).

The TROPOMI Level 3 monthly mean glyoxal product averaged the native pixels to 0.05° resolution after excluding pixels with quality assurance values less than 0.5 (Lerot et al., 2021b). This Level 3 product has shown better consistency with
165 ground-based measurements and more reliable seasonal variation and local enhancements over regions of strong biogenic, anthropogenic, and biomass burning emissions, compared to earlier satellite retrievals (Lerot et al., 2021a). ~~For comparison with our GEOS-Chem simulations, we~~ We regrided the Level 3 product to ~~5° longitude \times 4° latitude~~ To infer the 2.5° longitude \times 2° latitude for comparison with GEOS-Chem results and for inferring the hypothetical marine source of glyoxal ~~;~~ we regrided (Section 4.3). ~~Model results were sampled at the satellite overpass time. Even after the quality assurance filter,~~
170 ~~the Level 3 product over the global oceans to 2.5° longitude \times 2° latitude to improve the signal-to-noise ratio while retaining large-scale spatial features~~ showed small negative values over remote oceans and deserts where simulated glyoxal VCDs were also very small; these values were retained in the regridding.

We also compared simulated glyoxal concentrations against surface, ship-based, and aircraft measurements of glyoxal over land and ocean. Table S1 compiles the measurements reported in the literature. ~~In addition,~~ To compare model results against observations, the monthly mean simulated glyoxal concentrations were sampled at the coordinates of the measurement site and
175 during the month of measurement.

2.5 Observations of other chemical species for model evaluation

We compared model results to additional observational datasets to evaluate the impacts of revised glyoxal simulation on atmospheric chemistry. The one-year (July 2019 to June 2020) mean global surface CO concentrations were averaged from the Measurements of Pollution In The Troposphere (MOPITT) Level 3 monthly gridded CO product (MOP03TM, Version 9) (Ziskin, 2021). The one-year (July 2019 to June 2020) mean global surface daily maximum 8-hour average ozone concentrations at 0.1° resolution were from Wang et al. (2025), a machine learning-based fusion of surface observations, chemical transport model simulations, meteorological reanalysis, satellite products, and emission-related information. Simulated organic aerosols (OA) and SOA were also evaluated against measurements over China synthesized by Miao et al. (2021), consisting of campaign-average OA measurements from 68 urban, 18 suburban, and 8 remote sites across China during 2011–2019. Finally, we used aircraft measurements of HO_x concentrations from the Airborne Tropospheric Hydrogen Oxides Sensor (Brune et al., 2021) during the Atmospheric Tomography Mission (ATom-2) campaign (Wofsy and Team, 2018) in February 2017 to evaluate the simulated impacts of glyoxal on atmospheric oxidation capacity [in the marine atmosphere](#). Figure S1 shows the aircraft tracks during ATom-2, which transected the remote Pacific and the remote Atlantic.

3 Evaluation of the standard GEOS-Chem glyoxal simulation as guidance for model improvements

We first evaluated the atmospheric glyoxal simulation in the standard GEOS-Chem (GC-CTRL simulation) against global observations to identify **potential model deficiencies and to guide targeted [aspects for model](#)** improvements. Figure 1 compares the annual mean (July 2019—une 2020) tropospheric glyoxal **column concentrations** [VCDs](#) from the GC-CTRL simulation against TROPOMI observations. TROPOMI showed prominent glyoxal enhancements over major biogenic source regions (Southeast U.S., East and South Asia, tropical forests of South America and Africa), seasonal burning areas (boreal forests of North America and Asia, and forests, savannas, and farm and peat lands in South America, Africa and Southeast Asia), and major anthropogenic source regions. These spatial patterns reflected the expected glyoxal production from biogenic isoprene, as well as the emissions of glyoxal and its precursors from biomass burning and anthropogenic emissions (Fu et al., 2008; Myriokefalitakis et al., 2008; Cao et al., 2018).

The GC-CTRL simulation broadly reproduced the TROPOMI-observed spatial patterns over land (Figure 1a,b), but the simulated annual mean glyoxal **columns** [VCDs](#) were systematically lower than TROPOMI observations, with normalized mean biases (NMB) of -77% globally and ~~-61~~-60% over land (Figure S2). In particular, the model substantially underestimated glyoxal over the major source regions of biogenic isoprene in the tropics and the mid-latitudes. Figure 2 evaluated the simulated concentrations of glyoxal against individual ground-based and aircraft measurements over land (Table S1). Some measurements were made in urban areas or areas affected by local biomass burning (Table S1) and could not be reproduced by our simulations at coarse resolution. **Excluding these outliers** [However, even with these outliers excluded](#), the model still showed substantial underestimations of glyoxal concentrations compared to measurements at the surface and in the boundary layer (NMB = -90%), suggesting systematic errors in precursor emissions and/or near-surface chemistry.

In addition, TROPOMI observed glyoxal enhancements over the tropical oceans (Figure 1a) that were spatially correlated with the [MODIS Aqua retrieved](#) chlorophyll-*a* contents (NASA Goddard Space Flight Center, Ocean Ecology Laboratory, Ocean Biology Processing Group, 2014) of surface sea water (Figure S3, spatial correlation $r = 0.5$ over the tropical ocean), potentially indicating a marine biogenic source, which was also implied in previous studies (Wittrock et al., 2006; Alvarado et al., 2014; Zhou et al., 2014; Volkamer et al., 2015; Zhu and Kieber, 2019; Lerot et al., 2021a). This feature was completely absent in the GC-CTRL simulation (Figure 1b) due to the model's lack of marine emissions of glyoxal or its precursors. The NMB of simulated glyoxal concentrations against TROPOMI observations over the global oceans was -88% (Figures 1b). Lengthening the simulated glyoxal production timescale from terrestrial precursors would not reproduce the observed marine pattern, indicating that the glyoxal enhancements over the oceans could not be explained by the long-range transport of terrestrial precursors alone (Fu et al., 2008). Figure 2 shows that the GC-CTRL simulation was also unable to reproduce the magnitudes and variations of individual glyoxal measurements in the MBL (Table S1, NMB = -92%). Figure S4 compares the vertical profiles of simulated glyoxal concentrations against aircraft observations over the Northeast Pacific, North Atlantic, Tropical Atlantic, and the South Atlantic (Volkamer et al., 2015; Kluge et al., 2023). These measurements reported glyoxal concentrations exceeding 10 ppt in the MBL, decreasing gradually with altitude. These observed concentrations of glyoxal in the marine atmosphere were not reproduced in the GC-CTRL simulations.

The evaluation of the GC-CTRL simulation above indicated potential model deficiencies in precursor emissions over land and ocean, as well as in the photochemical production of glyoxal from those precursors. We addressed these issues in the following sections [and discussed the uncertainties given the current levels of understanding](#).

4 Updates to terrestrial and marine emissions of glyoxal and its precursors in GEOS-Chem

4.1 Top-down constraints on biogenic isoprene emissions over major source regions

Biogenic isoprene is the largest known terrestrial precursor of atmospheric glyoxal (Fu et al., 2008; Myriokefalitakis et al., 2008). Figure 3 compares the annual mean tropospheric isoprene ~~column concentrations observed by CrIS~~ [VCDs observed by the CrIS instrument](#) and those in the GC-CTRL simulation. Relative to the CrIS observations, the GC-CTRL simulation underestimated annual isoprene abundances over the mid-latitudes but overestimated isoprene over the Amazon, tropical Africa, and Northern Australia. These ~~apparent regional biases in isoprene abundance likely reflected regional biases in discrepancies between CrIS and the~~ [discrepancies between CrIS and the](#) GEOS-Chem ~~'s isoprene emissions, which would contribute to the model's underestimation of glyoxal over land (Figure 1b). Our findings~~ were similar to those ~~of~~ [pointed out by](#) Wells et al. (2020) based on comparisons between CrIS and an earlier version of GEOS-Chem (v11-02e) ~~and generally consistent with previous evaluations using satellite-based formaldehyde measurements and in situ isoprene observations (Palmer et al., 2006; Marais and Wiedinmyer, 2016; DiMaria et al., 2025) . However, the~~ [The](#) causes of GEOS-Chem's potential biases in regional isoprene emission are yet unclear due to differences in MEGAN algorithm versions and input data, and insufficient evaluation against *in situ* isoprene flux measurements over all source regions. [Potential biases may also](#)

240 [arise from underestimation of land-use-specific isoprene emitters such as oil palm plantations, as well as from chemistry-driven modulation of isoprene columns by co-located soil and biomass-burning NO \(Yoon et al., 2026\)](#) .

We addressed the potential biases in GEOS-Chem's isoprene emissions in a "top-down" fashion by scaling the isoprene emissions over nine major source regions (shown in Figure 3e), such that the simulated annual mean isoprene **column concentrations** $\bar{\Omega}_{i, \text{isoprene, GC}}$ over each region i ($= 1$ to 9) aligned with the CrIS observations ($\bar{\Omega}_{i, \text{isoprene, GC}}$):

$$245 \quad \bar{E}_{i, \text{isoprene, GC-TM-E}} = \frac{\bar{\Omega}_{i, \text{isoprene, CrIS}}}{\bar{\Omega}_{i, \text{isoprene, GC}}} \times \bar{E}_{i, \text{isoprene, GC}} \quad (1)$$

where $\bar{E}_{i, \text{isoprene, GC}}$ was the annual mean isoprene emission for region i calculated by the default MEGAN algorithm in GEOS-Chem; $\bar{E}_{i, \text{isoprene, GC-TM-E}}$ was the top-down-constrained isoprene emission. In practice, we iterated the scaling process of Eq. (1) twice to account for the photochemical non-linearity associated with OH reactivity.

Figure 3 shows the regional differences in isoprene emissions between our top-down estimates and those of the default
250 GEOS-Chem. The top-down constrained global isoprene emission was ~~428~~ [438](#) Tg yr⁻¹, similar to the default ~~443~~ [449](#) Tg yr⁻¹ in GC-CTRL (Table 1). However, over the Northern mid-latitudes, the regional annual mean isoprene emissions were scaled by factors of 1.2 to 3.5, with maximum increases over the temperate ecosystems of Europe and Asia. Conversely, over the tropical ecosystems of the Amazon, Africa, and Northern Australia, our use of CrIS constraints decreased the regional isoprene emissions by 10% to 60%. [These CrIS-based constraints on global isoprene emissions were still lower than the global isoprene emission estimates \(490 to 514 Tg yr⁻¹\) independently derived by Sfendla et al. \(2026\) using TROPOMI formaldehyde and glyoxal retrievals.](#)

We used the top-down constrained isoprene emissions to drive the GC-TM-E, GC-T-EC, and GC-TM-EC experiments (Section 6). ~~We~~ [Our scaling of regional isoprene emissions generally improves the observed and simulated surface isoprene concentrations in several representative regions, as shown in Figure S5 \(Seco et al., 2011; Link et al., 2015; Garg et al., 2026; Zhang et al., 2020; Sun et al., 2025\)](#) . Nevertheless, we emphasize
260 that our approach was not a formal inversion of isoprene emissions, which would require more thorough evaluations of the CrIS retrieval and its uncertainty, a more chemically coherent treatment of the OH-nonlinearity, as well as representations of the potential OH-regeneration pathways from isoprene oxidation under low-NO_x conditions (Peeters et al., 2014; Yang et al., 2024). Instead, our goal was to broadly constrain the spatial distribution of global isoprene emissions, such that our
265 mechanistic improvements on glyoxal chemistry can be meaningfully evaluated against global observations.

4.2 Additional updates to terrestrial emissions of glyoxal, glycolaldehyde, and their precursors

The default CEDS did not include emissions of glyoxal and glycolaldehyde from biofuel use. We implemented [these emissions](#) into ~~GEOS-Chem~~ [these emissions](#) ~~GEOS-Chem~~ by scaling biofuel emissions of ethane with emission ratios from Andreae (2019). Where regionally-resolved inventories were available - including Africa (Marais and Wiedinmyer, 2016), the
270 continental U.S. (National Emissions Inventory Collaborative, 2019), and Asia (Li et al., 2017; Zheng et al., 2018, 2021) - we scaled those biofuel emissions of ethane to incorporate local information. The resulting emission estimates for glyoxal and glycolaldehyde from global biofuel use were 1.3 Tg yr⁻¹ and 1.0 Tg yr⁻¹, respectively (Table 1).

Biogenic methylbutenol (2-methyl-3-buten-2-ol, $\text{CH}_3\text{COH}(\text{CH}_3)\text{CHCH}_2\text{CHCH}_2$), emitted almost exclusively by North American pine trees (Guenther et al., 2012), can be oxidized by OH to produce glycolaldehyde. However, methylbutenol was not represented in the standard GEOS-Chem v14.2.3. We added to the model emissions of methylbutenol calculated by the MEGAN v2.1 algorithm (Guenther et al., 2012), resulting in a global emission of 1.7 Tg yr^{-1} (Table 1).

Several recent studies indicated that the GFED4s inventory underestimated the amount of burned dry mass and the associated pollutant emissions by 30% to 120% over various biomass burning areas of the world (Zhong et al., 2022; Castagna et al., 2023; Chen et al., 2023; Qi et al., 2024). We scaled biomass burning emissions of glyoxal, glycolaldehyde, and all other NMVOCs by a factor of 1.8 globally (Table 1), an average of the scale factors from previous regional evaluations (Zhong et al., 2022; Chen et al., 2023; Castagna et al., 2023; Qi et al., 2024). This scale factor was also consistent with the global NMVOC emission estimate ($79.3 \text{ Tg C yr}^{-1}$) in the latest GFED5 inventory (Chen et al., 2023), which was 82% higher than the estimate of GFED4s ($43.6 \text{ Tg C yr}^{-1}$). The updated global biomass burning emissions for glyoxal and glycolaldehyde from were 3.8 Tg yr^{-1} and 4.0 Tg yr^{-1} , respectively (Table 1).

4.3 Estimation of a hypothetical secondary glyoxal source in the global MBL

TROPOMI and other *in situ* and remote sensing measurements tentatively implied a secondary source of glyoxal over the tropical MBL associated with unknown, potentially biogenic, marine precursors. We estimated this In this work, we used TROPOMI observations to estimate how large that hypothetical secondary glyoxal source might be, and we evaluated whether the inclusion of that hypothetical glyoxal source is consistent with independent *in situ* and remote sensing measurements in the global MBL. Previous studies have tentatively linked the glyoxal in the MBL to biologically active waters or to photochemical production from the dissolved organic matter (DOM) content in sea water. However, currently reported concentrations or fluxes of precursors (e.g., isoprene, ethylene, and propene) appear insufficient to explain the magnitude of glyoxal inferred from satellite observations (Broadgate et al., 2004; Zhang et al., 2025; Pound, 2021). In addition, measurements of NMVOCs in the MBL remain too sparse to provide a quantitative global constraint. We therefore interpret the inferred marine glyoxal source as a diagnostic term representing the glyoxal production from potentially missing marine NMVOCs.

We estimated the potential secondary glyoxal source by hypothesizing a short-lived marine precursor, which was emitted at the sea surface, instantaneously mixed throughout the MBL, and oxidized by OH to produce glyoxal at 100% yield. We started by assuming an *a priori* daytime emission for this hypothetical precursor, scaled to match the TROPOMI-observed glyoxal column concentrations VCDs over the global ocean at 2.5° longitude \times 2° latitude resolution:

$$E_{i,m,a priori} = \frac{\Omega_{i,m, \text{glyoxal, TROPOMI}}}{\tau} \quad (2)$$

where $E_{i,m,a priori}$ was the *a priori* monthly mean emission flux of the unknown precursor on the i^{th} model grid for the m^{th} month; $\Omega_{i,m, \text{glyoxal, TROPOMI}}$ was the TROPOMI-observed glyoxal column concentration VCD for that grid and month. We assumed an OH-oxidation rate constant of $k_{\text{OH}} = 1.5 \times 10^{-10} \text{ cm}^3 \text{ molecules}^{-1}$, such that the daytime lifetime (τ) of this hypothetical precursor was approximately one hour (at a typical OH concentration in the MBL of $1.8 \times 10^6 \text{ molecules cm}^{-3}$), consistent with the mean mixing timescale of the global MBL (Maclean et al., 2017). However, we did not allow the precursor

to consume OH. We then used the simulated glyoxal **column concentrations** VCDs over the oceans to optimize the emission for the hypothetical precursor ($E_{i,m, optimized}$):

$$E_{i,m, optimized} = \frac{\Omega_{i,m, glyoxal, TROPOMI}}{\Omega_{i,m, glyoxal, GC}} \cdot E_{i,m, a priori} \quad (3)$$

This optimization was iterated twice until the simulated marine glyoxal **column concentrations** VCDs agreed with satellite
310 observations within 20%, yielding an estimated daytime secondary glyoxal source of 66 Tg yr⁻¹ over the global oceans that
spatially correlated with TROPOMI observations (Figures 4 and S3).

5 Improvements to the chemical processes of glyoxal and its precursors in GEOS-Chem

5.1 Photochemical production of glyoxal from isoprene

A key objective of this study was to incorporate latest knowledge to better simulate the glyoxal production from isoprene
315 oxidation. We evaluated Wennberg et al. (2018) and the RCIM against latest literature and identified two key mechanistic
improvements to RCIM (as implemented in the standard GEOS-Chem v14.2.3) to reconcile with experiments. We also
highlighted several pathways where the reactions and yields of glyoxal are still uncertain and thus require further study. Figure
5 shows our mechanistic revisions to the RCIM and the simulated glyoxal yields under low- and high-~~NO~~ NO_x concentrations.


5.1.1 First-generation production of glyoxal and glycolaldehyde from the OH-oxidation of isoprene under high-NO_x 320 conditions

Chamber experiments by Galloway et al. (2011) demonstrated that, under high-NO_x conditions, the production of glyoxal and
glycolaldehyde from OH-initiated oxidation of isoprene occurred in **two-stages** two stages. The first stage occurred immediately
upon isoprene consumption, with measured molar yields of 2.1 ± 0.6% and 2.69 ± 0.82% for glyoxal and glycolaldehyde,
respectively. Subsequent photooxidation of C₅ and C₄ carbonyls led to further production of glyoxal and glycolaldehyde.
325 The first-generation production aligned with the theoretical pathways proposed by Dibble (2004a, b), wherein specific *Z*-
 δ -hydroxy-peroxy isoprene radicals (ISOPOO) first react with NO, **undergo rapid** then rapidly undergo 1,5 H-shifts and
~~O₂-addition to form~~ -additions to yield dihydroxy peroxy radicals. Some of these dihydroxy peroxy radicals further react
with NO to produce an alkoxy radical that fragments into pairs of glyoxal and hydroxyacetone or pairs of glycolaldehyde and
methylglyoxal. Similar fast production of glyoxal (3.8% yield inferred) and glycolaldehyde (4.2% yield) was also noted by
330 Paulot et al. (2009), prompting the inclusion of a Dibble-type pathway in an earlier version of GEOS-Chem (v11-02). The
MCM v3.3.1 also incorporated a Dibble-type glyoxal production at 2.2% molar yield under high-NO_x.

The RCIM omitted this first-generation high-NO_x pathway. Wennberg et al. (2018) assigned 45% of the *Z*- δ -derived
dihydroxy peroxy radicals to form C₅ hydroxy carbonyls based on experimental constraints (Teng et al., 2017). However,
the fate of the remaining 55% dihydroxy-peroxy radicals was unknown. Wennberg et al. (2018) postulated that the remaining
335 dihydroxy-peroxy radicals undergo rapid enol H-shift, O₂-addition, and decomposition; they assumed this pathway to be much
faster than the reaction with NO, effectively bypassing the first-generation production of glyoxal and glycolaldehyde.

We revised the RCIM to incorporate the Dibble pathway (Figure 5), redirecting the 55% of dihydroxy peroxy radicals of unknown fate (denoted as DIBOO) to react with NO and fragment into glyoxal (GLYX) + hydroxyacetone (HAC) or glycolaldehyde (GLYC) + methylglyoxal (MGLY) pairs. The corresponding RCIM reactions in GEOS-Chem were revised to:

340



IHOO1 and IHOO4 were the ISOPOOs from OH-addition at carbons 1 and 4 of isoprene; HC5A was a lumped C₅ carbonyl. Branches (R1) and (R2) comprised approximately 4.6% of the total ISOPOO + NO reactions under high-NO_x conditions. Yields and rate constants of GLYX (+ HAC) and GLYC (+ MGLY) in (R3) followed Wennberg et al. (2018). Our revised mechanism resulted in 1.2% and 1.1% first-generation molar yields of glyoxal and glycolaldehyde under high-NO_x,
350 respectively, still lower than the yields measured experimentally (Paulot et al., 2009; Galloway et al., 2011; Sprengnether et al., 2002).

5.1.2 Post first-generation production of glyoxal from OH-oxidation of isoprene under high- and low-NO_x conditions

Beyond the initial OH-oxidation, glyoxal production from isoprene occurs as second-generation product involving its major intermediate products and as third-generation product via glycolaldehyde (Wennberg et al., 2018), as depicted in Figure 5.
355 However, the glyoxal yields from these pathways are still incompletely constrained, especially for low-NO_x conditions. In Wennberg et al. (2018), the relevant intermediates include methyl vinyl ketone (MVK), isoprene nitrates, and C₅ carbonyls such as isoprene hydroxy hydroperoxides (ISOPOOHs), isoprene dihydroxy epoxides (IEPOXs), and hydroperoxy aldehydes (HPALDs). The yields of these major products from isoprene, the yield of glycolaldehyde from MVK, and the yields of IEPOXs from ISOPOOHs were experimentally constrained under both high- and low-NO_x conditions in Wennberg et al. (2018) and
360 represented current-best knowledge. However, the yields of glyoxal and glycolaldehyde from HPALDs were theoretically derived without experimental constraints (Peeters et al., 2014). Also, glycolaldehyde yields from isoprene nitrates were constrained experimentally only for (4-OH,3-ONO₂)-isoprene hydroxy nitrate (IHN) (Jacobs et al., 2014; Lee et al., 2014) and generalized across isomers with limited validation. In particular, the production of glyoxal and glycolaldehyde from HPALDs and isoprene nitrates both involved the reactions of C₄-dihydroperoxy-carbonyls, whose atmospheric fates were structurally
365 derived (Peeters et al., 2014; Wennberg et al., 2018). Under low-NO_x, the yields of glycolaldehyde from IEPOX+OH were subject to uncertainties associated with the radical propagating channels of non-acyl RO₂ + HO₂ reactions (Bates and Jacob,

2019; Müller et al., 2019). We retained RCIM's standard glyoxal and glycolaldehyde productions from isoprene's intermediate products but highlight the pathways involving HPALDs, isoprene nitrates, and IEPOXs for future experimental investigations.

370 More critically, the third-generation glyoxal formation from isoprene occurs via the OH-oxidation of glycolaldehyde (Figure 5). The GEOS-Chem v14.2.3 mechanism assumed an 11% yield of glyoxal from the glycolaldehyde+OH reaction based on Paulot et al. (2009), which was markedly lower than the 20% yield recommended by the IUPAC and JPL and even lower than the 20%-33% yields measured by Chan et al. (2009). We updated this glyoxal yield to 20% in our revised chemical mechanism.

5.1.3 Overall yield of glyoxal from OH-initiated oxidation of isoprene and its sensitivity to NO_x

We conducted box simulations to evaluate glyoxal production from the OH-initiated oxidation of isoprene using three chemical mechanisms: the mechanism of GC-CTRL (i.e., the RCIM as implemented in GEOS-Chem v14.2.3), MCM v3.3.1, and our revised mechanism for GC-T-EC. Figure 6 shows the simulated glyoxal yields under varying OH and NO_x levels. Oxidation of isoprene initiated by other oxidants, depositions, and aerosol uptake processes were disabled in these box simulations. Under a typical moderate-NO_x condition (OH = 1.8×10^6 molecules cm⁻³ and NO = 1 ppb), the GC-CTRL mechanism produced glyoxal at only a 4.0% molar yield, approximately half of the yield simulated by MCM (9.0%). In comparison, our revised mechanism increased the glyoxal yield to 8.7%, an enhancement attributed mostly to our added first-generation production and an increased glyoxal yield from glycolaldehyde oxidation.

385 The NO_x-sensitivity of glyoxal yield from isoprene OH-oxidation stems from the competition among reaction pathways and serves as a key diagnostic for evaluating chemical mechanisms. Our box simulations showed disparate NO_x-sensitivities among the three chemical mechanisms. In the GC-CTRL mechanism, the glyoxal molar yield from isoprene was almost invariant (3-4%) across a wide range of NO concentrations (0.1 to 10 ppb) (Figure 6d). In comparison, the MCM exhibited strong enhancements of glyoxal yield with increasing NO levels, rising from 7.5% at NO = 0.1 ppb to 10.9% at NO = 5 ppb (Figure 6d). This positive NO_x-sensitivity mainly reflected MCM's explicit representation of glyoxal formation from isoprene nitrates and its assumption of no glyoxal formation from IEPOX-derived peroxy radicals.

Our revised mechanism exhibited a milder yet non-monotonic sensitivity of glyoxal yield to NO_x concentration, rising from 7.3% at NO = 0.1 ppb to a peak of 8.7% at NO = 2 ppb, then decreased slightly to 8.6% for OH = 1.8×10^6 molecules cm⁻³ (Figure 6d). Figure 5 compares the branching ratios of glyoxal formation pathways in our revised mechanism at NO concentrations of 0.1 ppb and 5 ppb, respectively. Under high-NO conditions, the DIBOO pathway became effective, and the ISOPOOH reactions shifted moderately toward the production of MVK and isoprene nitrates, whose subsequent yields of glycolaldehyde and glyoxal also increased slightly. At lower NO levels, ISOPOOH and its isomers favored the formation of C₅ carbonyls; however, the glycolaldehyde and glyoxal yields from these carbonyls decreased substantially. This complex modulation of competing pathways explained the mild yet non-monotonic NO_x-sensitivity in the simulated glyoxal yields of our revised mechanism. Bates and Jacob (2019) showed that the glyoxal yield in the RCIM increased sharply to 10% at extremely low NO concentration (<10 ppt), driven by the HO₂ reaction with IEPOX peroxy radicals and the H-shifts of ISOPOO. This large glyoxal yield at extremely low NO would also be present in our revised mechanism, since our modifications of RCIM were mostly associated with high-NO_x pathways. However, ~~that an~~ NO level of <10 ppt

is unrealistically low for current source areas of isoprene and likely rarely realized in the atmosphere (Hudman et al., 2007; Schulz et al., 2018).

405 Furthermore, the glyoxal yield from isoprene varied substantially with OH concentrations (Figure 6a,b,c). At a fixed NO_x level of 1 ppb, the glyoxal yield increased from 7-?% at OH = 1 × 10⁶ molecules cm⁻³ to 9-10% at OH = 1 × 10⁷ molecules cm⁻³ both in our revised mechanism and in MCM, a trend not captured by the GC-CTRL mechanism. This difference reflected the strong influence of OH concentration on the branching between glycolaldehyde's photolysis and OH-oxidation, which was subdued in the GC-CTRL mechanism due to its lower glyoxal yield from glycolaldehyde. In our revised mechanism, a 50% increase in OH led to a 20% enhancement in the glyoxal yield from glycolaldehyde. This result highlighted the potential for substantial glyoxal production from isoprene in regions with concurrently high NO_x and OH levels.

410 Experimental constraints on glyoxal yields from OH-initiated isoprene oxidation and their NO_x-sensitivity remained limited, though available evidence was generally consistent with our revised mechanism. For high-NO_x conditions, Galloway et al. (2011) calculated the total glyoxal yield from OH-initiated isoprene oxidation in an older version of MCM (v3.2), which lacked first-generation production pathways, to be 8% (with O₃ = 320 ppb) to 11% (without O₃). These numbers closely bracketed the high-NO_x glyoxal yield (10.9%) for the current MCM v3.3.1, which did include a 2.2% first-generation glyoxal production (Figure 6b). However, Galloway et al. (2011) reported measured total glyoxal yields equivalent to only 64% of
415 the MCM v3.2 predictions under high-NO_x conditions, corresponding to measured yields of 5.1-?.0% for glyoxal. These measured yields aligned more closely with our revised mechanism and suggested that both MCM v3.2 and v3.3.1 overestimated glyoxal production from post-first-generation pathways under high-NO_x conditions. Using active differential optical absorption spectroscopy (DOAS), Guo et al. (2021) measured the surface formaldehyde and glyoxal concentrations at a urban site during
420 summer. They observed that in isoprene-dominated air masses, both glyoxal and formaldehyde concentrations, as well as the glyoxal-to-formaldehyde ratio (R_{GF}), were insensitive to NO_x levels between 5 and 40 ppb. This result was consistent with the modest NO_x-dependence of glyoxal yields in our revised mechanism at high NO concentrations.

Field observations in low-NO_x, isoprene-dominated environments indicated that glyoxal production from isoprene is likely substantial. Li et al. (2016) and Chan Miller et al. (2017) analyzed the Southeast Nexus (SENEX) aircraft campaign data
425 over Southeast U.S. during summer 2013. Both studies demonstrated that the observed glyoxal levels and R_{GF} values in this lower-NO_x (0.05 to 0.1 ppb between 0 to 5 km altitude) environment could be well simulated using mechanisms where glyoxal yields at low NO_x were comparable or even exceeded those at high NO_x. Although the two studies proposed different chemical pathways to explain this phenomenon, their results consistently supported a sustained or potentially enhanced glyoxal yield under low-NO conditions, such as that in our revised mechanism and in the RCIM. In contrast, However, a recent
430 direct measurement reported a glyoxal yield of 0.52 ± 0.06% from OH-initiated isoprene oxidation under extremely low NO_x (<70 ppt) (Warman, 2024), contradicting the inference from field observations and underscoring significant uncertainties in quantifying low-NO_x glyoxal yields from isoprene.

5.1.4 Potential uncertainties in glyoxal production from oxidation of isoprene by O₃, NO₃, and Cl

Ozonolysis accounts for approximately 10% of the global sink of atmospheric isoprene (Nguyen et al., 2016). In the explicit
435 mechanism of Wennberg et al. (2018), isoprene ozonolysis produces glyoxal predominantly via MVK, whose oxidation would
lead to a 2.2% molar yield of glyoxal under high-NO_x conditions but would be at the timescale of several hours. However, in
an isoprene ozonolysis chamber experiment with seed particles, Bikkina et al. (2021) observed glyoxal in the particulate phase
within half an hour of isoprene consumption, potentially indicating fast glyoxal production followed by aerosol uptake. Recent
experiments showed that *anti*-MVK-oxide, a Criegee intermediates produced by isoprene ozonolysis Nguyen et al. (2016), may
440 undergo unimolecular decay and O₂-addition to produce glyoxal and OH at 3% ± 1% yield (Vansco et al., 2020). However,
because the current estimate for the yield of *anti*-MVK-oxide from isoprene ozonolysis was only 6.7% (Wennberg et al.,
2018), the effective glyoxal production from that pathway would be negligible (<0.2%). We retained the isoprene ozonolysis
mechanism in the RCIM but highlighted this glyoxal production pathway for [further studies](#)[future investigations](#).

Isoprene oxidation by NO₃ at night accounts for approximately 2% of the global isoprene loss (Bates and Jacob, 2019).
445 In the RCIM, the isoprene + NO₃ reaction rapidly formed nitrooxy alkylperoxy radicals (INO₂), which reacted with HO₂ to
yield various isoprene hydroproxy nitrate isomers (IPN) (Schwantes et al., 2015). Glyoxal was produced mainly through MVK
derived from β-1,2-INO₂, as constrained by Schwantes et al. (2015). A minor pathway to glyoxal involved the oxidation of 2,1-
IPN, though the branching ratios of IPN isomers remained uncertain, and subsequent reaction pathways of IPNs lacked strong
experimental validation (Wennberg et al., 2018). Given the complexity of isoprene nitrate speciation and limited experimental
450 constraints, we retained the existing RCIM representation for isoprene + NO₃ chemistry.

5.2 Production of glyoxal from other NMVOC precursors

5.2.1 Acetylene

Acetylene (C₂H₂), with an 18-day global mean lifetime, is an important anthropogenic precursor of glyoxal in the free
troposphere (Fu et al., 2008). In the standard GEOS-Chem v14.2.3, OH-initiated oxidation of acetylene produced glyoxal
455 at a fixed molar yield of 63% following MCM v3.3.1, consistent with experimentally reported yields of 70% ± 30% (Bohn
and Zetzsch, 1998).

5.2.2 Aromatics

Aromatics in the standard GEOS-Chem v14.2.3 included benzene, toluene, and lumped xylenes (Bates et al., 2021). Laboratory
studies indicated that glyoxal produced from OH-initiated aromatic oxidation occurred first through [ring cleavage](#)[ring-cleavage](#),
460 followed by oxidation of [the](#) cleaved C₄ and C₅ intermediate products (Volkamer et al., 2007; Bates et al., 2021). Reported
first-generation glyoxal yields from ring cleavage varied considerably. In the standard GEOS-Chem v14.2.3, Bates et al. (2021)
implemented first-generation glyoxal yields from aromatics by averaging literature-reported values across low- to high-NO_x
conditions: 18%, 12%, 10% for benzene, toluene, and lumped xylenes, respectively. Glyoxal production pathways and yields

from subsequent C₄ and C₅ intermediate photochemistry were partially constrained by experiments and theoretical studies
465 (Newland et al., 2018; Wang et al., 2020). Given the current level of understanding, we retained the standard GEOS-Chem
representation of aromatic chemistry and glyoxal formation pathways.

5.2.3 Ethylene

In the standard GEOS-Chem v14.2.3, atmospheric ethylene (C₂H₄) was emitted from anthropogenic and biogenic sources over
land and underwent oxidation by OH and O₃ described by a condensed version of MCM v3.3.1 (Kwon et al., 2021), with the
470 OH-oxidation producing glycolaldehyde at 100% yield under both high- and low-NO_x conditions. We retained the standard
ethylene oxidation scheme in GEOS-Chem, but our increased glyoxal yield from glycolaldehyde effectively raised the global
molar yield of glyoxal yield from ethylene from 7% in the GC-CTRL simulation to 19% in the GC-TM-EC simulation.

5.2.4 Methylbutenol

Atmospheric methylbutenol is oxidized by OH and O₃, but these reactions were not represented in the standard GEOS-Chem
475 v14.2.3. Fu et al. (2008) previously assumed a constant 63% molar yield of glycolaldehyde from methylbutenol OH-oxidation,
based on early high-NO_x experiments and an early version of MCM (v3.1). However, later experiments by Chan et al. (2009)
found that methylbutenol's glycolaldehyde yield decrease from 64-70% under high-NO_x to 26-32% at ~~low NO~~ low-NO_x levels
(<2 ppb), indicating competition from an HO₂ pathway that bypasses glycolaldehyde formation. We adopted a condensed
scheme of the methylbutenol oxidation mechanism from MCM v3.3.1 (Müller et al., 2019), which was experimentally validated
480 for its HO₂ turnover rate and formaldehyde production (a product from the high-NO_x oxidation of methylbutenol) (Novelli
et al., 2018). This scheme predicted glycolaldehyde yields of 63% under high-NO_x and 0% under low-NO_x conditions. Our
simulated global glyoxal yield from methylbutenol was 5.8%.

5.2.5 Monoterpenes

Monoterpenes in the standard GEOS-Chem v14.2.3 were represented by two lumped species: one for monoterpenes with a
485 single double bond (including α -pinene, β -pinene, sabinene, and δ^3 -carene) and another for those with two double bonds
(including limonene, myrcene, and ocimene). Their photochemical oxidation was described by the mechanism of Fisher et al.
(2016), which omitted the glyoxal production from ozonolysis— pathway established in earlier studies ((Yu et al., 1998; Fick
et al., 2003, 2004).

Reported glyoxal yields from monoterpene ozonolysis were sparse and highly variable, ranging from 4-9% for α -pinene
490 ((Fick et al., 2003, 2004) to 6.2-24% for α -phellandrene (a two-double-bond species) (Mackenzie-Rae et al., 2017). Lower
yields of 0.5-?% were also reported for two monoterpenoids, geraniol and citral (Nunes et al., 2005). Fu et al. (2008) previously
applied in their GEOS-Chem simulation a mid-range 5% glyoxal yield based on α -pinene experiments. However, this
assumption resulted in a simulated glyoxal-to-formaldehyde ratio (RGF) more than three times lower than satellite observations
over boreal forests, where monoterpene emissions dominated (Silva et al., 2018). This discrepancy was significantly larger

495 than that over other NMVOC source regions, leading Silva et al. (2018) to conclude that monoterpene-derived glyoxal was substantially underestimated. More recently, Müller et al. (2019) assumed a 20% glyoxal yield from monoterpene ozonolysis in their chemical mechanism, but the basis for ~~this assumption were~~ that assumption was unclear. In the latest v14.5 of GEOS-Chem, Travis et al. (2024) implemented a complex monoterpene oxidation mechanism condensed from the MCM v3.3.1. However, that mechanism produced glyoxal from monoterpene ozonolysis exclusively via MVK at relatively low yields.

500 Given the top-down constraint of Silva et al. (2018) and the higher glyoxal yields reported for α -phellandrene (Mackenzie-Rae et al., 2017), we assumed a fixed 15% yield to the ozonolysis of all monoterpenes in our revised mechanism. Our simulated global glyoxal yield from monoterpenes was 2.7%.

5.3 Heterogeneous uptake of glyoxal on aqueous aerosols and cloud droplets

Glyoxal undergoes irreversible uptake by aqueous aerosols and cloud droplets, constituting a substantial pathway for SOA formation in the atmosphere (Fu et al., 2008; Ling et al., 2020). This heterogeneous uptake process was characterized in 505 GEOS-Chem by a reactive uptake coefficient γ (Jacob, 2000; Fu et al., 2008):

$$\frac{dC_G}{dt} = aD_g + \frac{1}{4}v\gamma_G AC_G \quad (4)$$

where C_G was the gas-phase glyoxal concentration, a was the radius of the wet particle or cloud droplet, D_g was the gas-phase diffusivity of glyoxal, v was its mean molecular speed, and A was the particle surface area.

510 Experiments have shown that glyoxal's irreversible uptake is RH-dependent due to a salting-in effect (Waxman et al., 2015), which was not represented in the standard GEOS-Chem v14.2.3. Curry et al. (2018) parameterized the reactive uptake coefficient (γ_G) by incorporating thermodynamics, aqueous chemistry, and salting-in effects on mass transfer limitations, resulting in an exponential dependence of γ_G on RH with a quadratic polynomial exponent. This parameterization yielded $\gamma_G = 3.6 \times 10^{-3}$ at RH = 55%, decreasing sharply to $\gamma_G = 2.6 \times 10^{-5}$ at RH = 90%. The RH dependency was consistent with 515 measured gas-particle partitioning effects (Healy et al., 2009), particularly under high-RH conditions (Gen et al., 2018). We adopted this parameterization from Curry et al. (2018) in our revised mechanism.

6 Evaluation of simulated glyoxal concentrations against global satellite and surface measurements

We evaluated our revised atmospheric glyoxal simulations against global satellite and surface measurements and found that ~~our incorporating~~ updated precursor emissions and chemistry ~~substantially improved~~ improved the overall model— 520 observation consistency. Figure 1 compares the annual mean tropospheric glyoxal ~~columns~~ VCDs at satellite overpass time (13:30 local time) from sensitivity experiments ~~against with~~ TROPOMI observations and the standard GEOS-Chem simulation (GC-CTRL). The ~~simulation, which improved terrestrial precursor emissions and chemistry, showed better spatial agreement with TROPOMI over land (Figure 1d, NMB = -47% ; r = 0.85 over land); though global bias remained due to the lack of representation of glyoxal over the oceans. In contrast, the GC-TM-E simulation only updated~~ experiment updated the terrestrial 525 precursor emissions and ~~added~~ included a hypothetical marine boundary layer ~~source; this glyoxal source. This~~ simulation

significantly reduced the global bias against TROPOMI observations (Figure 1c, NMB = -23 – 20% , $r = 0.86$ globally, NMB = -0.09 , $r=0.94$ over the ocean) but still exhibited substantial discrepancies in spatial pattern and magnitudes of glyoxal over land relative to TROPOMI.

530 The simulation, incorporating current-best knowledge on terrestrial precursor emission and chemistry, along with a secondary marine glyoxal source, demonstrated best overall agreement with TROPOMI observations globally (Figure 1e). The model successfully captured 0.84 globally, primarily because oceanic glyoxal VCDs were constrained to TROPOMI-observed values. In addition, the GC-TM-E simulation enhanced glyoxal levels over major forested and biomass burning regions across North America, Europe, Asia, South America, and Africa, and Southeast Asia, with improving consistency against TROPOMI glyoxal observations over land (NMB = -0.35 – -0.47 and $r = 0.9$ against TROPOMI observations over land). The model also reproduced the spatial patterns and concentrations of glyoxal (0.80). The GC-T-EC simulation, which refined both terrestrial precursor emissions and chemistry, showed further improved spatial correlation with TROPOMI over tropical oceans, with NMB land (Figure 1d), $r = -0.06$ and $r = 0.86$ and NMB = 0.94 against TROPOMI observations over the global oceans (-0.47). However, a global bias remained due to the lack of representation of glyoxal over the oceans.

540 The GC-TM-EC simulation, which incorporated current-best knowledge on terrestrial precursor emission and chemistry, along with TROPOMI-constrained marine glyoxal source, showed the best overall agreement with global TROPOMI observations (Figure 1e). The global bias of simulated annual mean glyoxal column concentration VCD against TROPOMI observations was -18 – 17% , with a spatial correlation of $r = 0.90$. The remaining bias of 0.87 . Over land, the NMB was -42% , with a spatial correlation of $r = 0.85$. The slight improvement in NMB in the GC-TM-EC simulation was mostly over land, reflecting the relative to the GC-T-EC simulation over land arose from enhanced glyoxal VCDs over the coastal grids due to the added marine glyoxal source. The remaining land bias in the GC-TM-EC simulation reflected persistent uncertainties in precursor emissions from biogenic, anthropogenic (Qiu et al., 2020), and biomass burning sources (Silva et al., 2018), as well as glyoxal yields from these precursors.

550 Figure 2 compares simulated concentrations of glyoxal against surface and boundary layer measurements over land and ocean. Excluding measurements in urban and biomass burning areas from urban and biomass-burning sites, the GC-TM-EC simulation showed substantially improved better agreement with observations over land (Figure 2a, NMB = -35%) compared to 43% than the GC-CTRL simulation (NMB = -78 – 77%). By including a hypothetical marine glyoxal source, the simulation demonstrated markedly better alignment with MBL observations across the tropical Pacific, Southern Ocean, Caribbean Sea, and North Atlantic, with NMB reduced from -92% to 12% (Figure 2b, Table S1). This alignment with observations independent of the TROPOMI satellite retrievals provided strong evidence for a significant, yet unaccounted-for glyoxal source in the MBL.

560 The experiment substantially improved the simulated glyoxal levels over major isoprene source regions, addressing a key deficiency in the simulation. We found that this improvement over land arose not only from refined isoprene emissions and its glyoxal production, but also from a better quantification of glyoxal and its precursors from biomass burning. Figure 7 compares the satellite-observed and simulated column concentrations VCDs of glyoxal and isoprene, along with the respective observed and simulated glyoxal-isoprene ratios (R_{GI}), over key over four high-glyoxal regions. Figure S5 compares the simulated

surface isoprene concentrations against measurements in key biogenic source regions. Over North America, the top-down CrIS constraints on isoprene emissions corrected the low bias in GC-CTRL -simulated isoprene abundances, which alone led to large improvements in the simulated glyoxal abundance in the experiment VCDs, improving the agreement with observed surface isoprene concentrations, while also improving the agreement between the simulated glyoxal VCDs and the TROPOMI observations. Updates in chemical mechanisms further improved the simulated glyoxal abundances and the R_{GI} , indicating a better representation of glyoxal production from isoprene in the experiment. Over . In contrast, over South America, Africa, and Southeast Asia, isoprene emissions were reduced by the application of top-down constraints CrIS constraints either reduced or only slightly increased isoprene emissions, which brought the simulated surface isoprene concentrations to better agree with measurements (Figure S5). However, the enhanced biomass burning biomass-burning emissions of glyoxal and its precursors over these regions increased substantially raised both the simulated glyoxal abundances and the simulated R_{GI} to become more consistent with observations. Chemical mechanism updates in GC-TM-EC then helped bring both simulated glyoxal abundances and R_{GI} closer to observed values, demonstrating improved partitioning between different glyoxal sources in these regions. only brought minor adjustments to simulated glyoxal abundance over these regions. These results underscore the large contribution of biomass burning emissions to atmospheric glyoxal, which must be better constrained to enable accurate NMVOC emission inversions using glyoxal VCDs.

Over the ocean, inclusion of the hypothetical marine glyoxal source markedly improved the agreement of the GC-TM-EC simulation with independent MBL observations. Compared to the GC-CTRL simulation, the normalized mean bias (NMB) against MBL measurements in the tropical Pacific, Southern Ocean, Caribbean Sea, and North Atlantic was reduced from -22% to 12% (Figure 2b; Table S1). This consistency with observations independent of the TROPOMI retrieval supports the existence of a significant, previously unaccounted—or glyoxal source in the MBL. Figure S4 further compares the simulated glyoxal over the remote ocean with the aircraft observations from Volkamer et al. (2015) and Kluge et al. (2023). With the added marine glyoxal source, GC-TM-EC —imulated glyoxal concentrations aligned better with the limited aircraft data. However, the model overestimated glyoxal in surface air over the Northeast Pacific but underestimated it over the Atlantic. Additionally, the aircraft profiles suggested a more gradual decline in glyoxal concentration with altitude than simulated, which may indicate a longer-lived marine-derived precursor. This inference is inconsistent with the current understanding that marine-emitted VOCs are predominantly short-lived alkenes (e.g., ethylene and propene) and isoprene (Broadgate et al., 2004; Zhang et al., 2025; Pound, 2021).

7 Revised global budgets of atmospheric glyoxal

Table 2 summarizes the simulated global burdens and budgets of atmospheric glyoxal from our sensitivity experiments and compares them with the GC-CTRL simulation and Fu et al. (2008). Figure 4 shows the global patterns of sources and sinks of atmospheric glyoxal in the GC-TM-EC simulation. Assuming that glyoxal and its precursors originated solely from terrestrial sources and using revised chemistry (the GC-T-EC simulation), the estimated global atmospheric glyoxal burden was 15 Gg, 3650% higher than the standard GEOS—em value (+10 Gg). This revised burden coincided with the 15 Gg reported by

Fu et al. (2008), despite large differences in model emissions and chemical mechanisms. The global atmospheric glyoxal source in the GC-T-EC simulation was ~~44~~ 40 Tg yr⁻¹, a ~~76~~74% increase over the ~~25~~ 23 Tg yr⁻¹ in the GC-CTRL simulation. This enhancement resulted from both elevated ~~direct glyoxal emissions~~ (primary glyoxal emissions from biomass and biofuel burning, ~~increased by a factor of 1.4~~) and ~~strengthened secondary chemical production (enhanced by 70%)~~ (Figure 4) ~~as well as strengthened secondary chemical productions (67% enhancement), particularly from isoprene.~~

Our revised global budget still pointed to isoprene as the most important terrestrial precursor of atmospheric glyoxal, contributing ~~17.1~~ 14 Tg yr⁻¹ (~~39~~40%) to the total atmospheric glyoxal source, ~~88~~79% stronger than the ~~9.1~~ 7.8 Tg yr⁻¹ in the GC-CTRL simulation. In the GC-T-EC simulation, 86% of the global glyoxal production from isoprene was through glycolaldehyde, whose global effective yield of glyoxal was 10%, twice of that in the GC-CTRL simulation. Our revision of the isoprene chemistry increased the global glyoxal yield from isoprene to ~~4.0~~3.2% in the GC-T-EC simulation, compared to the ~~2.1~~1.7% in the GC-CTRL simulation. ~~As previously pointed out by Bates and Jacob (2019), the production~~ ~~This yield~~ of glyoxal from isoprene in the global atmosphere ~~would be~~ was smaller than that predicted by box models, because the key intermediates (such as MVK, IEPOXs, and HPALDs) ~~are~~ were removed by deposition or aerosol uptake, limiting their gas-phase reactions (Bates and Jacob, 2019). The total glyoxal production from biogenic precursors ~~were~~ 21.5 ~~was~~ 18 Tg yr⁻¹, ~~49~~45% of the total atmospheric glyoxal source. The anthropogenic and biomass burning precursors, including aromatics, acetylene, and ethylene, collectively contributed ~~39~~41% of the total glyoxal source.

Atmospheric glyoxal was removed predominantly by photolysis (~~52~~58%), followed by OH-oxidation (~~23~~16%), irreversible uptake by aqueous particles (~~15~~14%), and deposition (~~11~~14%) (Figure 4). ~~We found that the contribution of the global OH removal of glyoxal in the simulation was 28% of that in the standard GEOS-Chem (18%); likely reflecting the enhanced presence of glyoxal in photochemically active regions.~~ In addition, the heterogeneous removal of glyoxal by aqueous particles was ~~6.4~~ 5.6 Tg yr⁻¹, an order of magnitude larger than that in the GC-CTRL simulation. This increase was due to a combination of increased global glyoxal formation and enhanced irreversible uptake coefficient of glyoxal over land. The simulated global lifetime of glyoxal in the GC-T-EC experiment was 3.4 h, shortened compared to the 3.8 h in the standard GEOS-Chem due to accelerated removal by OH and heterogeneous uptake.

The inclusion of a ~~potential~~ hypothetical marine glyoxal source in the GC-TM-EC simulation substantially altered the global glyoxal budget, elevating the burden to 39 Gg and the total source to ~~110~~ 106 Tg yr⁻¹. Approximately 29 Gg of this enhanced burden was over the ocean, representing a 5.5-fold increase compared to the simulated glyoxal burden over the ocean in the GC-T-EC simulation. Sixty percent (66 Tg yr⁻¹) of the total glyoxal source in the GC-TM-EC simulation was produced in the MBL (Table 2, Figure 4), constituting a large source of OVOCs in the MBL that could influence the oxidative capacity of the marine atmosphere. Terrestrial source contributions remained similar to those in the GC-T-EC simulation.

The relative strengths of glyoxal sinks also shifted significantly in the GC-TM-EC experiment. Photolysis became an even more important sink for global glyoxal due to the high solar radiation and glyoxal concentrations in the tropical MBL. In contrast, glyoxal removal by heterogeneous uptake became relatively less important, as the added glyoxal was concentrated in high-humidity MBL where the reactive uptake coefficient of glyoxal (γ_G) was suppressed (Curry et al., 2018; Gen et al.,

2018). Overall, the simulated lifetime of atmospheric glyoxal in the GC-TM-EC experiment was ~~3.5~~ 3.2 h, similar to that in the GC-T-EC experiment.

630 **8 Impacts of the revised glyoxal simulation on atmospheric chemistry**

Finally, we examined how ~~our~~ the revised atmospheric glyoxal simulations affected global atmospheric chemistry. Figure 8 compared the simulated global surface concentrations of OH, HO₂, and O₃, along with the surface OH reactivity (OHR) associated with NMVOCs, in the ~~;~~ and model experiments. In the GC-T-EC experiment, regional changes to biogenic and biomass burning emissions and modifications to precursor chemistry ~~mostly~~ strongly affected atmospheric chemistry over land;
635 increasing. The increased regional isoprene and biomass burning NMVOC emissions in the GC-T-EC simulation strengthened OH reactivity across North America and Euroasia; ~~reduced~~, while reduced regional isoprene emissions decreased OH reactivity over South America and Australia (Figure 8h). The simulated surface OH decreased or increased in accordance to these regional changes in OH reactivity, averaging a ~~-3.3~~ -4.2% reduction in global surface OH relative to the GC-CTRL experiment (Figure 8h). These changes in OH-initiated oxidation of NMVOCs subsequently altered the photochemistry that produced
640 HO₂, particularly over relatively high-NO_x areas: surface HO₂ concentrations increased by more than ~~8% over Southeast U.S., Europe,~~ 10% over Europe, the Middle East, and East and South Asia, while decreasing slightly over the Amazon and Australia. Globally, the updates of ~~glyoxal chemistry and precursor emissions~~ precursor emissions and glyoxal chemistry increased the global surface O₃ concentration by 4.8~~17~~%.

The ~~hypothetical~~ revised precursor emissions and glyoxal chemistry also affected global SOA simulation. Simulated total
645 SOA formation increased from 115 Tg yr⁻¹ in the GC-CTRL experiment to 133 Tg yr⁻¹ in the GC-T-EC experiment and 136 Tg yr⁻¹ in the GC-TM-EC experiment. Glyoxal-derived SOA rose from 0.6 Tg yr⁻¹ in the GC-CTRL experiment to 6.4 Tg yr⁻¹ in the GC-T-EC experiment and 7.6 Tg yr⁻¹ in the GC-TM-EC experiment, respectively. Including the hypothetical marine glyoxal source contributed an extra 1.2 Tg yr⁻¹ of glyoxal SOA. Notably, the relative increase in SOA after inclusion of the hypothetical marine glyoxal source was smaller than the corresponding relative increase in gas-phase glyoxal. This was because
650 glyoxal SOA formation is limited by heterogeneous uptake, which was suppressed in high-RH regions by the salting-in effect, thereby reducing SOA production efficiency over humid marine environments (Waxman et al., 2015).

The hypothetical MBL source of glyoxal also affected the atmospheric oxidative capacity over the ocean, especially in the tropics (Figure 8c,f,i,l). The additional glyoxal removed OH in the MBL, while its photolysis produced HO₂. Relative to the GC-T-EC experiment, the added marine glyoxal source (simulated in GC-TM-EC) had a negative impact on the surface
655 OH concentration globally (~~-0.5~~ -0.6%), especially over the tropical ocean (-1.0%). The marine glyoxal increased the surface concentration of HO₂ by ~~6.8~~ 6.5%, the surface NMVOC OH reactivity by ~~2.3~~ 1.9%, and the surface ozone concentration by 0.3%, over the tropical ocean.

Figure S6 compares our simulated HO_x vertical profile against the observations in the marine boundary layer (~~<~~ below 2,500 m) during the ATom-2 mission (Brune et al., 2020). The OH concentrations simulated in the GC-CTRL simulation agreed well
660 with observations, and our improved glyoxal simulation did not significantly change the OH profile over the remote ocean.

The HO₂ concentration simulated in GC-TM-EC ~~improved~~ slightly alleviated the underestimation in GC-TM-E by 5% but still underestimated HO₂ compared to the observations, indicating the potential lack or persistent underestimation of HO₂ sources (~~potentially e.g.~~, OVOCs) over the remote MBL.

665 We also evaluated the broader influence of the revised glyoxal and precursors representations on tropospheric chemistry
(Figures S7, S8, S9). Relative to the GC-CTRL simulation, increased precursor emissions in the GC-TM-EC simulation
substantially raised the global mean surface CO concentrations from 99.1 ppb to 113.5 ppb, reducing the model bias against
MOPITT observations from -25% to 13% (Figure S7), especially over Eastern United States, East and South Asia, and the
tropical South America and Africa. Similarly, enhanced terrestrial precursor emissions moderately increased the simulated
global mean surface ozone concentration over land from 22.8 ppb in GC-CTRL to 29.5 ppb in GC-TM-EC, closer to the
670 34.1 ppb reported in the data—ssimulated product of Wang et al. (2025) (Figure S8). The increased precursor emissions
and atmospheric glyoxal abundance in the GC-T-EC experiment also led to a stronger heterogeneous uptake of glyoxal by
aqueous particles, thereby enhancing SOA formation (Table 2). Over China, for example, the GC-T-EC experiment simulated
significantly higher SOA concentrations than the GC-CTRL experiment, especially at suburban and remote sites (Figure
S9). These increases brought simulations closer to—hough still substantially below—bserved concentrations, indicating that
675 SOA formation involves complex mechanisms beyond glyoxal chemistry (Miao et al., 2021). The apparent improvements in
simulated CO, ozone, and SOA should therefore be interpreted as supporting the chemical plausibility of the revised glyoxal
budget and underscoring glyoxal’s role as a proxy for atmospheric NMVOC emissions.

9 Conclusion

We developed an improved global simulation of atmospheric glyoxal using the GEOS-Chem model, advanced by updated
680 emissions and chemical mechanism and consistent with global observations. Assuming glyoxal precursors were entirely
terrestrial, we estimated a global atmospheric glyoxal source of ~~44~~ 40 Tg yr⁻¹ and a global burden of 15 Gg, effectively
~~reproducing observed glyoxal levels over land~~ improving the consistency between simulated glyoxal VCDs and TROPOMI
satellite observations. Isoprene was identified as the dominant terrestrial precursor, while anthropogenic and biomass burning
sources—including acetylene, ethylene, and aromatics—ollectively contributed 39% of the global glyoxal source. These
685 results highlighted the importance of non-biogenic precursors to ambient glyoxal in urban and biomass burning areas, with
implications for the use of glyoxal for NMVOC emission inversion in such environments.

~~Further inclusion of a hypothetical secondary glyoxal source (66 Tg yr⁻¹) within the global MBL increased the global~~
~~atmospheric source of glyoxal to 110 Tg yr⁻¹ and its burden to 39 Gg. While this addition shows tentative consistency with in~~
~~situ MBL observations, the existence and impacts of such a source remain highly uncertain and warrant focused investigation~~
690 ~~in future studies.~~

Even with our improved precursor emission estimates and chemical representation, our simulated glyoxal remained
approximately ~~35~~ 40% lower than satellite glyoxal retrievals and surface observations over land. This discrepancy likely stems
from both satellite retrieval uncertainties and persistent uncertainties in ~~both~~ emissions and chemistry. Anthropogenic emissions

of glyoxal precursors, particularly aromatic compounds, are variable and uncertain among different emission inventories and demonstrated a consistent low-bias compared to observations (Yan et al., 2019). In addition, while we scaled isoprene and biomass burning emissions by region on an annual basis, this approach may introduce biases across different land cover types and seasons (Silva et al., 2018; Wang et al., 2024; DiMaria et al., 2025). Uncertainties in simulated OH and NO_x concentrations slightly may moderately affect glyoxal production (within 10%, Figure 6), but, but they have a potentially large impact on top-down isoprene estimates, particularly over Amazonthe Amazonian rainforest, where underestimated OH and NO_x lead to isoprene overestimation for a given emission rate (Wells et al., 2020). Further uncertainties arise from incomplete representation of key chemical pathways, including glyoxal production from isoprene under low-NO_x conditions (Warman, 2024), ozonolysis via Criegee intermediates (Vansco et al., 2020), detailed isoprene nitrate chemistry (Schwantes et al., 2015), and heterogeneous uptake processes of glyoxal (Li et al., 2016; Kim et al., 2022).

Our study further underscores highlights the need for enhanced measurement of NMVOCs in the marine boundary layer (MBL) and improved measurement of VOCs in the MBL and for a deeper exploration of their photochemistry. Current evidence indicates that glyoxal over remote oceans is likely secondary in origin and photochemical transformations. Our exploratory addition of a hypothetical secondary glyoxal source (66 Tg yr⁻¹) raised the global atmospheric glyoxal source to 106 Tg yr⁻¹ and its burden to 39 Gg. While this addition shows tentative consistency with *in situ* MBL observations, this hypothetical glyoxal source cannot be explained by direct emissions alone. Concurrently, emerging research known marine precursor emissions. Recent work suggests marine biogenic emissions of precursors like isoprenemay be more substantial than previously estimated (Zhang and Gu, 2022). This, such as isoprene, may be larger than previous estimates (Zhang and Gu, 2022). However, our evaluation of model results against limited aircraft measurements points to the potential existence of an unexplored NMVOC photochemical cycle driven by emissions from marine plants or algae. Addressing possible influence of a more long-live precursor, the nature and impacts of which remain highly uncertain. Resolving this gap in marine NMVOC photochemistry is essential for understanding its impacts on atmospheric oxidative capacity, marine aerosol formation, and background ozone levels quantifying the roles of glyoxal and its precursors in marine and global atmospheric chemistry.

. Our revised GEOS-Chem model for global atmospheric glyoxal is permanently archived at <https://doi.org/10.57760/sciencedb.27761>. CrIS isoprene observations are available upon request to Dylan Millet. TROPOMI glyoxal observations are available upon request to Thomas Danckaert. Other glyoxal measurements used for model evaluation are summarized in Table S1. ATom observations are downloaded from <https://espoarchive.nasa.gov/archive/browse/atom/DC8/BackTraj> (last accessed: 9 September 2025).

. AZ developed the simulations and analyses with help from EX, WW, YL, LZ, WT, and RZ. KCW, DBM, ZW, BY, MS, TD, and CL provided surface and satellite observations. KHB developed the RCIM and implemented it in GEOS-Chem. AZ and TMF wrote the manuscript with inputs from all coauthors. AZ, TMF, and YW conceived the project. TMF supervised and secured funding for the project.

. At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.

725 . This work was supported by the National Natural Science Foundation of China (42325504, 42305188, 42461160326), the National Key Research and Development Program of China (2023YFC3706205), the Shenzhen Science and Technology Program (KQTD20210811090048025, JCYJ20220818100611024), and the High-level Special Funds (G03034K006). DBM and KCW acknowledge support from the U.S. National Aeronautics and Space Administration (NASA; Grant #80NSSC24M0037). Computational resources were supported by the Center for Computational Science and Engineering of the Southern University of Science and Technology.

730 References

- Akagi, S., Yokelson, R. J., Wiedinmyer, C., Alvarado, M., Reid, J., Karl, T., Crounse, J., and Wennberg, P.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmospheric Chemistry and Physics*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- Alvarado, L. M. A., Richter, A., Vrekoussis, M., Wittrock, F., Hilboll, A., Schreier, S. F., and Burrows, J. P.: An improved glyoxal retrieval
735 from OMI measurements, *Atmospheric Measurement Techniques*, 7, 4133–4150, <https://doi.org/10.5194/amt-7-4133-2014>, 2014.
- Alvarado, L. M. A., Richter, A., Vrekoussis, M., Hilboll, A., Kalisz Hedegaard, A. B., Schneising, O., and Burrows, J. P.: Unexpected long-range transport of glyoxal and formaldehyde observed from the Copernicus Sentinel-5 Precursor satellite during the 2018 Canadian wildfires, *Atmospheric Chemistry and Physics*, 20, 2057–2072, <https://doi.org/10.5194/acp-20-2057-2020>, 2020.
- Amos, H. M., Jacob, D. J., Holmes, C. D., Fisher, J. A., Wang, Q., Yantosca, R. M., Corbitt, E. S., Galarneau, E., Rutter, A. P., Gustin,
740 M. S., Steffen, A., Schauer, J. J., Graydon, J. A., Louis, V. L. S., Talbot, R. W., Edgerton, E. S., Zhang, Y., and Sunderland, E. M.: Gas-particle partitioning of atmospheric Hg(II) and its effect on global mercury deposition, *Atmospheric Chemistry and Physics*, 12, 591–603, <https://doi.org/10.5194/acp-12-591-2012>, 2012.
- Andreae, M. O.: Emission of trace gases and aerosols from biomass burning—an updated assessment, *Atmospheric Chemistry and Physics*, 19, 8523–8546, <https://doi.org/10.5194/acp-19-8523-2019>, 2019.
- 745 Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II – gas phase reactions of organic species, *Atmospheric Chemistry and Physics*, 6, 3625–4055, <https://doi.org/10.5194/acp-6-3625-2006>, 2006.
- Bates, K. H. and Jacob, D. J.: A new model mechanism for atmospheric oxidation of isoprene: global effects on oxidants, nitrogen oxides, organic products, and secondary organic aerosol, *Atmospheric Chemistry and Physics*, 19, 9613–9640, <https://doi.org/10.5194/acp-19-9613-2019>, 2019.
750
- Bates, K. H., Jacob, D. J., Li, K., Ivatt, P. D., Evans, M. J., Yan, Y., and Lin, J.: Development and evaluation of a new compact mechanism for aromatic oxidation in atmospheric models, *Atmospheric Chemistry and Physics*, 21, 18 351–18 374, <https://doi.org/10.5194/acp-21-18351-2021>, 2021.
- Betterton, E. A. and Hoffmann, M. R.: Henry’s law constants of some environmentally important aldehydes, *Environmental Science & Technology*, 22, 1415–1418, <https://doi.org/10.1021/es00177a004>, 1988.
755
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *Journal of Geophysical Research: Atmospheres*, 106, 23 073–23 095, <https://doi.org/10.1029/2001JD000807>, 2001.
- Bian, H. and Prather, M. J.: Fast-J2: Accurate simulation of stratospheric photolysis in global chemical models, *Journal of Atmospheric Chemistry*, 41, 281–296, <https://doi.org/10.1023/A:1014980619462>, 2002.
760
- Bikkina, S., Kawamura, K., Sakamoto, Y., and Hirokawa, J.: Low molecular weight dicarboxylic acids, oxocarboxylic acids and α -dicarbonyls as ozonolysis products of isoprene: Implication for the gaseous-phase formation of secondary organic aerosols, *Science of The Total Environment*, 769, 144 472, <https://doi.org/10.1016/j.scitotenv.2020.144472>, 2021.
- Bohn, B. and Zetzsch, C.: Formation of HO₂ from OH and C₂H₂ in the presence of O₂, *J. Chem. Soc., Faraday Trans.*, 94, 1203–1210,
765 <https://doi.org/10.1039/A708536B>, 1998.

- Broadgate, W. J., Malin, G., Küpper, F. C., Thompson, A., and Liss, P. S.: Isoprene and other non-methane hydrocarbons from seaweeds: a source of reactive hydrocarbons to the atmosphere, *Marine Chemistry*, 88, 61–73, <https://doi.org/10.1016/j.marchem.2004.03.002>, 2004.
- 770 Brune, W. H., Miller, D. O., Thames, A. B., Allen, H. M., Apel, E. C., Blake, D. R., Bui, T. P., Commane, R., Crounse, J. D., Daube, B. C., Diskin, G. S., DiGangi, J. P., Elkins, J. W., Hall, S. R., Hanisco, T. F., Hannun, R. A., Hints, E. J., Hornbrook, R. S., Kim, M. J., McKain, K., Moore, F. L., Neuman, J. A., Nicely, J. M., Peischl, J., Ryerson, T. B., St. Clair, J. M., Sweeney, C., Teng, A. P., Thompson, C., Ullmann, K., Veres, P. R., Wennberg, P. O., and Wolfe, G. M.: Exploring oxidation in the remote free troposphere: insights from atmospheric tomography (ATom), *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031685, <https://doi.org/10.1029/2019JD031685>, 2020.
- 775 Brune, W. H., Miller, D. O., and Thames, A. B.: ATom: Measurements from Airborne Tropospheric Hydrogen Oxides Sensor (ATHOS), V2 (Version 2), ORNL Distributed Active Archive Center, <https://doi.org/10.3334/ORNLDAAAC/1930>, publisher: ORNL Distributed Active Archive Center. Last accessed: 2025-09-15, 2021.
- Cao, H., Fu, T.-M., Zhang, L., Henze, D. K., Miller, C. C., Lerot, C., Abad, G. G., De Smedt, I., Zhang, Q., van Roozendaal, M., Hendrick, F., Chance, K., Li, J., Zheng, J., and Zhao, Y.: Adjoint inversion of Chinese non-methane volatile organic compound emissions using space-based observations of formaldehyde and glyoxal, *Atmospheric Chemistry and Physics*, 18, 15 017–15 046, <https://doi.org/10.5194/acp-18-15017-2018>, 2018.
- 780 Carter, T. S., Heald, C. L., Kroll, J. H., Apel, E. C., Blake, D., Coggon, M., Edtbauer, A., Gkatzelis, G., Hornbrook, R. S., Peischl, J., Pfannerstill, E. Y., Piel, F., Reijrink, N. G., Ringsdorf, A., Warneke, C., Williams, J., Wisthaler, A., and Xu, L.: An improved representation of fire non-methane organic gases (NMOGs) in models: emissions to reactivity, *Atmospheric Chemistry and Physics*, 22, 12 093–12 111, <https://doi.org/10.5194/acp-22-12093-2022>, 2022.
- 785 Castagna, J., Senatore, A., Pellis, G., Vitullo, M., Bencardino, M., and Mendicino, G.: Uncertainty assessment of remote sensing- and ground-based methods to estimate wildfire emissions: a case study in Calabria region (Italy), *Air Quality, Atmosphere & Health*, 16, 705–717, <https://doi.org/10.1007/s11869-022-01300-1>, 2023.
- Chan, A. W. H., Galloway, M. M., Kwan, A. J., Chhabra, P. S., Keutsch, F. N., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Photooxidation of 2-Methyl-3-Buten-2-ol (MBO) as a potential source of secondary organic aerosol, *Environmental Science & Technology*, 43, 4647–4652, <https://doi.org/10.1021/es802560w>, 2009.
- 790 Chan Miller, C., Jacob, D. J., Marais, E. A., Yu, K., Travis, K. R., Kim, P. S., Fisher, J. A., Zhu, L., Wolfe, G. M., Hanisco, T. F., Keutsch, F. N., Kaiser, J., Min, K.-E., Brown, S. S., Washenfelder, R. A., González Abad, G., and Chance, K.: Glyoxal yield from isoprene oxidation and relation to formaldehyde: chemical mechanism, constraints from SENEX aircraft observations, and interpretation of OMI satellite data, *Atmospheric Chemistry and Physics*, 17, 8725–8738, <https://doi.org/10.5194/acp-17-8725-2017>, 2017.
- 795 Chen, Y., Hall, J., van Wees, D., Andela, N., Hantson, S., Giglio, L., van der Werf, G. R., Morton, D. C., and Randerson, J. T.: Multi-decadal trends and variability in burned area from the fifth version of the Global Fire Emissions Database (GFED5), *Earth System Science Data*, 15, 5227–5259, <https://doi.org/10.5194/essd-15-5227-2023>, 2023.
- 800 Chiu, R., Tinel, L., Gonzalez, L., Ciuraru, R., Bernard, F., George, C., and Volkamer, R.: UV photochemistry of carboxylic acids at the air-sea boundary: A relevant source of glyoxal and other oxygenated VOC in the marine atmosphere, *Geophysical Research Letters*, 44, 1079–1087, <https://doi.org/10.1002/2016GL071240>, 2017.
- Coburn, S., Ortega, I., Thalman, R., Blomquist, B., Fairall, C. W., and Volkamer, R.: Measurements of diurnal variations and eddy covariance (EC) fluxes of glyoxal in the tropical marine boundary layer: description of the Fast LED-CE-DOAS instrument, *Atmospheric Measurement Techniques*, 7, 3579–3595, <https://doi.org/10.5194/amt-7-3579-2014>, 2014.

- Curry, L. A., Tsui, W. G., and McNeill, V. F.: Technical note: Updated parameterization of the reactive uptake of glyoxal and methylglyoxal by atmospheric aerosols and cloud droplets, *Atmospheric Chemistry and Physics*, 18, 9823–9830, <https://doi.org/10.5194/acp-18-9823-2018>, 2018.
- De Haan, D. O., Jansen, K., Rynaski, A. D., Sueme, W. R. P., Torkelson, A. K., Czer, E. T., Kim, A. K., Rafla, M. A., De Haan, A. C., and Tolbert, M. A.: Brown carbon production by aqueous-phase interactions of glyoxal and SO₂, *Environmental Science & Technology*, 54, 4781–4789, <https://doi.org/10.1021/acs.est.9b07852>, 2020.
- Dibble, T. S.: Intramolecular hydrogen bonding and double H-atom transfer in peroxy and alkoxy radicals from isoprene, *The Journal of Physical Chemistry A*, 108, 2199–2207, <https://doi.org/10.1021/jp0306702>, 2004a.
- Dibble, T. S.: Prompt chemistry of alkenoxy radical products of the double H-atom transfer of alkoxy radicals from isoprene, *Journal of Physical Chemistry A*, 108, 2208–2215, <https://doi.org/10.1021/jp0312161>, 2004b.
- DiGangi, J. P., Henry, S. B., Kammrath, A., Boyle, E. S., Kaser, L., Schnitzhofer, R., Graus, M., Turnipseed, A., Park, J.-H., Weber, R. J., Hornbrook, R. S., Cantrell, C. A., Maudlin III, R. L., Kim, S., Nakashima, Y., Wolfe, G. M., Kajii, Y., Apel, E., Goldstein, A. H., Guenther, A., Karl, T., Hansel, A., and Keutsch, F. N.: Observations of glyoxal and formaldehyde as metrics for the anthropogenic impact on rural photochemistry, *Atmospheric Chemistry and Physics*, 12, 9529–9543, <https://doi.org/10.5194/acp-12-9529-2012>, 2012.
- DiMaria, C. A., Jones, D. B. A., Ferracci, V., Bloom, A. A., Worden, H. M., Seco, R., Vettikkat, L., Yáñez-Serrano, A. M., Guenther, A. B., Araujo, A., Goldstein, A. H., Langford, B., Cash, J., Harris, N. R. P., Brown, L., Rinnan, R., Schobesberger, S., Holst, T., and Mak, J. E.: Optimizing the temperature sensitivity of the isoprene emission model MEGAN in different ecosystems using a Metropolis-Hastings Markov Chain Monte Carlo method, *Journal of Geophysical Research: Biogeosciences*, 130, e2025JG008806, <https://doi.org/10.1029/2025JG008806>, 2025.
- Edwards, P. M., Brown, S. S., Roberts, J. M., Ahmadov, R., Banta, R. M., deGouw, J. A., Dubé, W. P., Field, R. A., Flynn, J. H., Gilman, J. B., Graus, M., Helmig, D., Koss, A., Langford, A. O., Lefer, B. L., Lerner, B. M., Li, R., Li, S.-M., McKeen, S. A., Murphy, S. M., Parrish, D. D., Senff, C. J., Soltis, J., Stutz, J., Sweeney, C., Thompson, C. R., Trainer, M. K., Tsai, C., Veres, P. R., Washenfelder, R. A., Warneke, C., Wild, R. J., Young, C. J., Yuan, B., and Zamora, R.: High winter ozone pollution from carbonyl photolysis in an oil and gas basin, *Nature*, 514, 351–354, <https://doi.org/10.1038/nature13767>, 2014.
- Emmerson, K. M. and Evans, M. J.: Comparison of tropospheric gas-phase chemistry schemes for use within global models, *Atmospheric Chemistry and Physics*, 9, 1831–1845, <https://doi.org/10.5194/acp-9-1831-2009>, 2009.
- Ferracci, V., Weber, J., Bolas, C. G., Robinson, A. D., Tummon, F., Rodríguez-Ros, P., Cortés-Greus, P., Baccarini, A., Jones, R. L., Galí, M., Simó, R., Schmale, J., and Harris, N. R. P.: Atmospheric isoprene measurements reveal larger-than-expected Southern Ocean emissions, *Nature Communications*, 15, 2571, <https://doi.org/10.1038/s41467-024-46744-4>, 2024.
- Fick, J., Pommer, L., Nilsson, C., and Andersson, B.: Effect of OH radicals, relative humidity, and time on the composition of the products formed in the ozonolysis of α -pinene, *Atmospheric Environment*, 37, 4087–4096, [https://doi.org/10.1016/S1352-2310\(03\)00522-3](https://doi.org/10.1016/S1352-2310(03)00522-3), 2003.
- Fick, J., Nilsson, C., and Andersson, B.: Formation of oxidation products in a ventilation system, *Atmospheric Environment*, 38, 5895–5899, <https://doi.org/10.1016/j.atmosenv.2004.08.020>, 2004.
- Fisher, J. A., Jacob, D. J., Travis, K. R., Kim, P. S., Marais, E. A., Chan Miller, C., Yu, K., Zhu, L., Yantosca, R. M., Sulprizio, M. P., Mao, J., Wennberg, P. O., Crounse, J. D., Teng, A. P., Nguyen, T. B., St. Clair, J. M., Cohen, R. C., Romer, P., Nault, B. A., Wooldridge, P. J., Jimenez, J. L., Campuzano-Jost, P., Day, D. A., Hu, W., Shepson, P. B., Xiong, F., Blake, D. R., Goldstein, A. H., Misztal, P. K., Hanisco, T. F., Wolfe, G. M., Ryerson, T. B., Wisthaler, A., and Mikoviny, T.: Organic nitrate chemistry and its implications for nitrogen budgets

- in an isoprene- and monoterpene-rich atmosphere: constraints from aircraft (SEAC⁴RS) and ground-based (SOAS) observations in the Southeast US, *Atmospheric Chemistry and Physics*, 16, 5969–5991, <https://doi.org/10.5194/acp-16-5969-2016>, 2016.
- 845 Fu, T., Jacob, D. J., Wittrock, F., Burrows, J. P., Vrekoussis, M., and Henze, D. K.: Global budgets of atmospheric glyoxal and methylglyoxal, and implications for formation of secondary organic aerosols, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD009505>, 2008.
- Fu, T.-M., Jacob, D. J., and Heald, C. L.: Aqueous-phase reactive uptake of dicarbonyls as a source of organic aerosol over eastern North America, *Atmospheric Environment*, 43, 1814–1822, <https://doi.org/10.1016/j.atmosenv.2008.12.029>, 2009.
- 850 Fu, T.-M., Cao, J. J., Zhang, X. Y., Lee, S. C., Zhang, Q., Han, Y. M., Qu, W. J., Han, Z., Zhang, R., Wang, Y. X., Chen, D., and Henze, D. K.: Carbonaceous aerosols in China: top-down constraints on primary sources and estimation of secondary contribution, *Atmospheric Chemistry and Physics*, 12, 2725–2746, <https://doi.org/10.5194/acp-12-2725-2012>, 2012.
- Galloway, M. M., Huisman, A. J., Yee, L. D., Chan, A. W. H., Loza, C. L., Seinfeld, J. H., and Keutsch, F. N.: Yields of oxidized volatile organic compounds during the OH radical initiated oxidation of isoprene, methyl vinyl ketone, and methacrolein under high-NO_x conditions, *Atmospheric Chemistry and Physics*, 11, 10 779–10 790, <https://doi.org/10.5194/acp-11-10779-2011>, 2011.
- 855 Garg, A., Desservettaz, M., Christodoulou, A., Christoudias, T., Kanawade, V. P., Savvides, C., Vrekoussis, M., Naqui, S., Jokinen, T., Byron, J., Williams, J., Mihalopoulos, N., Liakakou, E., Sciare, J., and Bourtsoukidis, E.: Heat and continental transport shape the variability of volatile organic compounds in the Eastern Mediterranean: insights from multi-year observations and regional modeling, *Atmospheric Chemistry and Physics*, 26, 2597–2622, <https://doi.org/10.5194/acp-26-2597-2026>, 2026.
- Gen, M., Huang, D. D., and Chan, C. K.: Reactive uptake of glyoxal by ammonium-containing salt particles as a function of relative humidity, *Environmental Science & Technology*, 52, 6903–6911, <https://doi.org/10.1021/acs.est.8b00606>, 2018.
- 860 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- Guo, Y., Wang, S., Zhu, J., Zhang, R., Gao, S., Saiz-Lopez, A., and Zhou, B.: Atmospheric formaldehyde, glyoxal and their relations to ozone pollution under low- and high-NO_x regimes in summertime Shanghai, China, *Atmospheric Research*, 258, 105 635, <https://doi.org/10.1016/j.atmosres.2021.105635>, 2021.
- 865 Ha, E. S., Park, R. J., Kwon, H.-A., Lee, G. T., Lee, S. D., Shin, S., Lee, D.-W., Hong, H., Lerot, C., De Smedt, I., Danckaert, T., Hendrick, F., and Irie, H.: First evaluation of the GEMS glyoxal products against TROPOMI and ground-based measurements, *Atmospheric Measurement Techniques*, 17, 6369–6384, <https://doi.org/10.5194/amt-17-6369-2024>, 2024.
- Healy, R. M., Temime, B., Kuprovskite, K., and Wenger, J. C.: Effect of relative humidity on gas/particle partitioning and aerosol mass yield in the photooxidation of *p*-xylene, *Environmental Science & Technology*, 43, 1884–1889, <https://doi.org/10.1021/es802404z>, 2009.
- 870 Hudman, R. C., Jacob, D. J., Turquety, S., Leibensperger, E. M., Murray, L. T., Wu, S., Gilliland, A. B., Avery, M., Bertram, T. H., Brune, W., Cohen, R. C., Dibb, J. E., Flocke, F. M., Fried, A., Holloway, J., Neuman, J. A., Orville, R., Perring, A., Ren, X., Sachse, G. W., Singh, H. B., Swanson, A., and Wooldridge, P. J.: Surface and lightning sources of nitrogen oxides over the United States: Magnitudes, chemical evolution, and outflow, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD007912>, 2007.
- 875 Ip, H. S. S., Huang, X. H. H., and Yu, J. Z.: Effective Henry’s law constants of glyoxal, glyoxylic acid, and glycolic acid, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2008GL036212>, 2009.
- IUPAC Subcommittee for Gas Kinetic Data Evaluation: Gas Kinetic Data Evaluation for Atmospheric Chemistry, <http://iupac.pole-ether.fr>, last accessed: 15 September 2025, 2013.

- Jacob, D.: Heterogeneous chemistry and tropospheric ozone, *Atmospheric Environment*, 34, 2131–2159, [https://doi.org/10.1016/S1352-2310\(99\)00462-8](https://doi.org/10.1016/S1352-2310(99)00462-8), 2000.
- Jacobs, M. I., Burke, W. J., and Elrod, M. J.: Kinetics of the reactions of isoprene-derived hydroxynitrates: gas phase epoxide formation and solution phase hydrolysis, *Atmospheric Chemistry and Physics*, 14, 8933–8946, <https://doi.org/10.5194/acp-14-8933-2014>, 2014.
- Jenkin, M. E., Young, J., Young, J., and Rickard, A. R.: The MCM v3.3.1 degradation scheme for isoprene, *Atmospheric Chemistry and Physics*, 15, 11 433–11 459, <https://doi.org/10.5194/acp-15-11433-2015>, 2015.
- 885 Kaiser, J., Wolfe, G. M., Min, K.-E., Brown, S. S., Miller, C. C., Jacob, D. J., de Gouw, J. A., Degouw, J., Graus, M., Hanisco, T. F., Holloway, J. S., Peischl, J., Pollack, I. B., Ryerson, T. B., Ryerson, T. B., Warneke, C., Washenfelder, R. A., and Keutsch, F. N.: Reassessing the ratio of glyoxal to formaldehyde as an indicator of hydrocarbon precursor speciation, *Atmospheric Chemistry and Physics*, 15, 7571–7583, <https://doi.org/10.5194/acp-15-7571-2015>, 2015.
- Kim, D., Cho, C., Cho, C., Seokhan Jeong, Soojin Lee, Nault, B., Nault, B. A., Jost, P. C., Campuzano-Jost, P., Day, D. A., Day, D. A., 890 Schroder, J. C., Jimenez, J.-L., Jimenez, J. L., Volkamer, R., Donald R. Blake, Armin Wisthaler, Alan Fried, DiGangi, J. P., DiGangi, J. P., Diskin, G. S., Diskin, G. S., Sally E. Pusede, Samuel R. Hall, Kirk Ullmann, Huey, G., Huey, L. G., David J. Tanner, Jack Dibb, Knote, C., Knote, C., and Kyung-Eun Min: Field observational constraints on the controllers in glyoxal (CHOCHO) reactive uptake to aerosol, *Atmospheric Chemistry and Physics*, 22, 805–821, <https://doi.org/10.5194/acp-22-805-2022>, 2022.
- Kluge, F., Hüneke, T., Lerot, C., Rosanka, S., Rotermund, M. K., Taraborrelli, D., Weyland, B., and Pfeilsticker, K.: Airborne glyoxal 895 measurements in the marine and continental atmosphere: comparison with TROPOMI observations and EMAC simulations, *Atmospheric Chemistry and Physics*, 23, 1369–1401, <https://doi.org/10.5194/acp-23-1369-2023>, 2023.
- Kwon, H.-A., Park, R. J., Oak, Y. J., Nowlan, C. R., Janz, S. J., Kowalewski, M. G., Fried, A., Walega, J., Bates, K. H., Choi, J., Blake, D. R., Wisthaler, A., and Woo, J.-H.: Top-down estimates of anthropogenic VOC emissions in South Korea using formaldehyde vertical column densities from aircraft during the KORUS-AQ campaign, *Elementa: Science of the Anthropocene*, 9, 00 109, 900 <https://doi.org/10.1525/elementa.2021.00109>, 2021.
- Lawson, S. J., Selleck, P. W., Galbally, I. E., Keywood, M. D., Harvey, M. J., Lerot, C., Helmig, D., and Ristovski, Z.: Seasonal in situ observations of glyoxal and methylglyoxal over the temperate oceans of the Southern Hemisphere, *Atmospheric Chemistry and Physics*, 15, 223–240, <https://doi.org/10.5194/acp-15-223-2015>, 2015.
- Lee, L., Teng, A. P., Wennberg, P. O., Crounse, J. D., and Cohen, R. C.: On rates and mechanisms of OH and O₃ reactions with isoprene- 905 derived hydroxy nitrates, *The Journal of Physical Chemistry A*, 118, 1622–1637, <https://doi.org/10.1021/jp4107603>, 2014.
- Lerot, C., Hendrick, F., Van Roozendaal, M., Alvarado, L. M. A., Richter, A., De Smedt, I., Theys, N., Vlietinck, J., Yu, H., Van Gent, J., Stavrou, T., Müller, J.-F., Valks, P., Loyola, D., Irie, H., Kumar, V., Wagner, T., Schreier, S. F., Sinha, V., Wang, T., Wang, P., and Retscher, C.: Glyoxal tropospheric column retrievals from TROPOMI – multi-satellite intercomparison and ground-based validation, *Atmospheric Measurement Techniques*, 14, 7775–7807, <https://doi.org/10.5194/amt-14-7775-2021>, 2021a.
- 910 Lerot, C., Stavrou, T., Van Roozendaal, M., Alvarado, L. M., and Richter, A.: GLYoxal Retrievals from TROPOMI (GLYRETRO) Product User Manual, Royal Belgian Institute for Space Aeronomy (BIRA-IASB), https://glyretro.aeronomie.be/UploadDir/documents/CHOCHO_PUM_S5PI_BIRA_v1.1.pdf, last accessed: 15 September 2025, 2021b.
- Li, J., Li, J., Mao, J., Min, K.-E., Washenfelder, R. A., Brown, S. S., Kaiser, J., Keutsch, F. N., Volkamer, R., Wolfe, G. M., Hanisco, T. F., Pollack, I. B., Ryerson, T. B., Ryerson, T. B., Graus, M., Gilman, J. B., Lerner, B. M., Warneke, C., de Gouw, J. A., de Gouw, 915 J. A., Joost A. de Gouw, Joost A. de Gouw, Middlebrook, A. M., Liao, J., Welti, A., Henderson, B. H., McNeill, V. F., Hall, S. R., Ullmann, K., Donner, L. J., Crounse, J. D., Paulot, F., and Horowitz, L. W.: Observational constraints on glyoxal production from isoprene

- oxidation and its contribution to organic aerosol over the Southeast United States., *Journal of Geophysical Research*, 121, 9849–9861, <https://doi.org/10.1002/2016jd025331>, 2016.
- 920 Li, J.-L., Zhai, X., Wu, Y.-C., Wang, J., Zhang, H.-H., and Yang, G.-P.: Emissions and potential controls of light alkenes from the marginal seas of China, *Science of The Total Environment*, 758, 143–165, <https://doi.org/10.1016/j.scitotenv.2020.143655>, 2021.
- Li, M., Zhang, Q., Kurokawa, J.-i., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmospheric Chemistry and Physics*, 17, 935–963, <https://doi.org/10.5194/acp-17-935-2017>, 2017.
- 925 Li, Y., Fu, T.-M., Yu, J. Z., Yu, X., Chen, Q., Miao, R., Zhou, Y., Zhang, A., Ye, J., Yang, X., Tao, S., Liu, H., and Yao, W.: Dissecting the contributions of organic nitrogen aerosols to global atmospheric nitrogen deposition and implications for ecosystems, *National Science Review*, 10, nwad244, <https://doi.org/10.1093/nsr/nwad244>, 2023.
- Li, Y., Fu, T.-M., Yu, J. Z., Zhang, A., Yu, X., Ye, J., Zhu, L., Shen, H., Wang, C., Yang, X., Tao, S., Chen, Q., Li, Y., Li, L., Che, H., and Heald, C. L.: Nitrogen dominates global atmospheric organic aerosol absorption, *Science*, 387, 989–995, <https://doi.org/10.1126/science.adr4473>, 930 2025.
- Liggio, J., Li, S.-M., and McLaren, R.: Reactive uptake of glyoxal by particulate matter, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/10.1029/2004JD005113>, 2005.
- Ling, Z., Xie, Q., Shao, M., Wang, Z., Wang, T., Guo, H., and Wang, X.: Formation and sink of glyoxal and methylglyoxal in a polluted subtropical environment: observation-based photochemical analysis and impact evaluation, *Atmospheric Chemistry and Physics*, 935 20, 11 451–11 467, <https://doi.org/10.5194/acp-2020-212>, 2020.
- Link, M., Zhou, Y., Taubman, B., Sherman, J., Morrow, H., Krintz, I., Robertson, L., Cook, R., Stocks, J., West, M., and Sive, B. C.: A characterization of volatile organic compounds and secondary organic aerosol at a mountain site in the Southeastern United States, *Journal of Atmospheric Chemistry*, 72, 81–104, <https://doi.org/10.1007/s10874-015-9305-5>, 2015.
- Mackenzie-Rae, F. A., Liu, T., Deng, W., Saunders, S. M., Fang, Z., Zhang, Y., and Wang, X.: Ozonolysis of α -phellandrene – Part 1: Gas- and 940 particle-phase characterisation, *Atmospheric Chemistry and Physics*, 17, 6583–6609, <https://doi.org/10.5194/acp-17-6583-2017>, 2017.
- Maclean, A. M., Butenhoff, C. L., Grayson, J. W., Barsanti, K., Jimenez, J. L., and Bertram, A. K.: Mixing times of organic molecules within secondary organic aerosol particles: a global planetary boundary layer perspective, *Atmospheric Chemistry and Physics*, 17, 13 037–13 048, <https://doi.org/10.5194/acp-17-13037-2017>, 2017.
- Mahajan, A. S., Prados-Roman, C., Hay, T. D., Lampel, J., Pöhler, D., Großmann, K., Tschirner, J., Frieß, U., Platt, U., and Johnston, 945 P.: Glyoxal observations in the global marine boundary layer, *Journal of Geophysical Research: Atmospheres*, 119, 6160–6169, <https://doi.org/10.1002/2013JD021388>, 2014.
- Marais, E. A. and Wiedinmyer, C.: Air quality impact of diffuse and inefficient combustion emissions in Africa (DICE-Africa), *Environmental Science & Technology*, 50, 10 739–10 745, <https://doi.org/10.1021/acs.est.6b02602>, 2016.
- Marais, E. A., Jacob, D. J., Jimenez, J. L., Campuzano-Jost, P., Day, D. A., Hu, W., Krechmer, J., Zhu, L., Kim, P. S., Miller, C. C., Fisher, 950 J. A., Travis, K., Yu, K., Hanisco, T. F., Wolfe, G. M., Arkinson, H. L., Pye, H. O. T., Froyd, K. D., Liao, J., and McNeill, V. F.: Aqueous-phase mechanism for secondary organic aerosol formation from isoprene: application to the southeast United States and co-benefit of SO₂ emission controls, *Atmospheric Chemistry and Physics*, 16, 1603–1618, <https://doi.org/10.5194/acp-16-1603-2016>, 2016.
- McDuffie, E. E., Smith, S. J., O'Rourke, P., Tibrewal, K., Venkataraman, C., Marais, E. A., Zheng, B., Crippa, M., Brauer, M., and Martin, R. V.: A global anthropogenic emission inventory of atmospheric pollutants from sector- and fuel-specific sources (1970–2017): an

- 955 application of the Community Emissions Data System (CEDS), *Earth System Science Data*, 12, 3413–3442, <https://doi.org/10.5194/essd-12-3413-2020>, 2020.
- Miao, R., Chen, Q., Shrivastava, M., Chen, Y., Zhang, L., Hu, J., Zheng, Y., and Liao, K.: Process-based and observation-constrained SOA simulations in China: the role of semivolatile and intermediate-volatility organic compounds and OH levels, *Atmospheric Chemistry and Physics*, 21, 16 183–16 201, <https://doi.org/10.5194/acp-21-16183-2021>, 2021.
- 960 Mungall, E. L., Abbatt, J. P. D., Wentzell, J. J. B., Lee, A. K. Y., Thomas, J. L., Blais, M., Gosselin, M., Miller, L. A., Papakyriakou, T., Willis, M. D., and Liggitto, J.: Microlayer source of oxygenated volatile organic compounds in the summertime marine Arctic boundary layer, *Proceedings of the National Academy of Sciences*, 114, 6203–6208, <https://doi.org/10.1073/pnas.1620571114>, 2017.
- Myriokefalitakis, S., Vrekoussis, M., Tsigaridis, K., Wittrock, F., Richter, A., Brühl, C., Volkamer, R., Burrows, J. P., and Kanakidou, M.: The influence of natural and anthropogenic secondary sources on the glyoxal global distribution, *Atmospheric Chemistry and Physics*, 8, 4965–4981, <https://doi.org/10.5194/acp-8-4965-2008>, 2008.
- 965 Müller, J.-F., Stavrou, T., and Peeters, J.: Chemistry and deposition in the Model of Atmospheric composition at Global and Regional scales using Inversion Techniques for Trace gas Emissions (MAGRITTE v1.1) – Part 1: Chemical mechanism, *Geoscientific Model Development*, 12, 2307–2356, <https://doi.org/10.5194/gmd-12-2307-2019>, 2019.
- NASA Goddard Space Flight Center, Ocean Ecology Laboratory, Ocean Biology Processing Group: MODIS-Terra Ocean Color Data, https://doi.org/10.5067/TERRA/MODIS_OC.2014.0, accessed July 28, 2015, 2014.
- 970 National Emissions Inventory Collaborative: 2016v1 Emissions Modeling Platform, <http://views.cira.colostate.edu/wiki/wiki/10202>, last accessed: 15 September 2025, 2019.
- Newland, M. J., Rea, G. J., Thüner, L. P., Henderson, A. P., Golding, B. T., Rickard, A. R., Barnes, I., and Wenger, J.: Photochemistry of 2-butenedial and 4-oxo-2-pentenal under atmospheric boundary layer conditions, *Physical Chemistry Chemical Physics*, 21, 1160–1171, <https://doi.org/10.1039/C8CP06437G>, 2018.
- 975 Nguyen, T. B., Tyndall, G. S., Crouse, J. D., Teng, A. P., Bates, K. H., Schwantes, R. H., Coggon, M. M., Zhang, L., Feiner, P., Müller, D. O., Skog, K. M., Rivera-Rios, J. C., Dorris, M., Olson, K. F., Koss, A., Wild, R. J., Brown, S. S., Goldstein, A. H., de Gouw, J. A., Brune, W. H., Keutsch, F. N., Seinfeld, J. H., and Wennberg, P. O.: Atmospheric fates of Criegee intermediates in the ozonolysis of isoprene, *Physical Chemistry Chemical Physics*, 18, 10 241–10 254, <https://doi.org/10.1039/C6CP00053C>, 2016.
- 980 Novelli, A., Kaminski, M., Rolletter, M., Acir, I.-H., Bohn, B., Dorn, H.-P., Li, X., Lutz, A., Nehr, S., Rohrer, F., Tillmann, R., Wegener, R., Holland, F., Hofzumahaus, A., Kiendler-Scharr, A., Wahner, A., and Fuchs, H.: Evaluation of OH and HO₂ concentrations and their budgets during photooxidation of 2-methyl-3-butene-2-ol (MBO) in the atmospheric simulation chamber SAPHIR, *Atmospheric Chemistry and Physics*, 18, 11 409–11 422, <https://doi.org/10.5194/acp-18-11409-2018>, 2018.
- Nunes, F., Veloso, M., Deppreira, P., and Deandrade, J.: Gas-phase ozonolysis of the monoterpenoids (S)-(+)-carvone, (R)-(-)-carvone, (-)-carveol, geraniol and citral, *Atmospheric Environment*, 39, 7715–7730, <https://doi.org/10.1016/j.atmosenv.2005.04.009>, 2005.
- 985 Palmer, P. I., Jacob, D. J., Chance, K., Martin, R. V., Spurr, R. J. D., Kurosu, T. P., Bey, I., Yantosca, R., Fiore, A., and Li, Q.: Air mass factor formulation for spectroscopic measurements from satellites: Application to formaldehyde retrievals from the Global Ozone Monitoring Experiment, *Journal of Geophysical Research: Atmospheres*, 106, 14 539–14 550, <https://doi.org/10.1029/2000JD900772>, 2001.
- Palmer, P. I., Abbot, D. S., Fu, T., Jacob, D. J., Chance, K., Kurosu, T. P., Guenther, A., Wiedinmyer, C., Stanton, J. C., Pilling, M. J., Pressley, S. N., Lamb, B., and Sumner, A. L.: Quantifying the seasonal and interannual variability of North American isoprene emissions using satellite observations of the formaldehyde column, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006689>, 2006.
- 990

- Park, R. J., Jacob, D. J., Field, B. D., Yantosca, R. M., and Chin, M.: Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the United States: Implications for policy, *Journal of Geophysical Research: Atmospheres*, 109, 995 <https://doi.org/10.1029/2003JD004473>, 2004.
- Paulot, F., Crouse, J. D., Kjaergaard, H. G., Kroll, J. H., Seinfeld, J. H., and Wennberg, P. O.: Isoprene photooxidation: new insights into the production of acids and organic nitrates, *Atmospheric Chemistry and Physics*, 9, 1479–1501, <https://doi.org/10.5194/acp-9-1479-2009>, 2009.
- Peeters, J., Müller, J.-F., Stavrou, T., and Nguyen, V. S.: Hydroxyl radical recycling in isoprene oxidation driven by hydrogen bonding and hydrogen tunneling: the upgraded LIM1 mechanism, *The Journal of Physical Chemistry A*, 118, 8625–8643, <https://doi.org/10.1021/jp5033146>, 2014.
- Plass-Dülmer, C., Koppmann, R., Ratte, M., and Rudolph, J.: Light nonmethane hydrocarbons in seawater, *Global Biogeochemical Cycles*, 9, 79–100, <https://doi.org/10.1029/94GB02416>, 1995.
- Platt, U. and Stutz, J.: Differential absorption spectroscopy, in: *Differential Optical Absorption Spectroscopy: Principles and Applications*, pp. 135–174, Springer Berlin Heidelberg, Berlin, Heidelberg, https://link.springer.com/content/pdf/10.1007/978-3-540-75776-4_6.pdf, last accessed: 15 September 2025, 2008.
- Pound, R. J.: Quantifying the importance of ocean-atmosphere exchange for atmospheric chemistry, phd, University of York, <https://etheses.whiterose.ac.uk/id/eprint/29608/>, 2021.
- Qi, B., Zhang, Z., Long, T., He, G., Wang, G., Peng, Y., and Xu, Z.: High resolution (30 m) burned area product improves the ability for carbon emission estimation in Africa, *Earth's Future*, 12, e2024EF005 051, <https://doi.org/10.1029/2024EF005051>, 2024.
- Qiu, X., Wang, S., Ying, Q., Duan, L., Xing, J., Cao, J., Wu, D., Li, X., Chengzhi, X., Yan, X., Liu, C., and Hao, J.: Importance of wintertime anthropogenic glyoxal and methylglyoxal emissions in Beijing and implications for secondary organic aerosol formation in megacities, *Environmental Science & Technology*, 54, 11 809–11 817, <https://doi.org/10.1021/acs.est.0c02822>, 2020.
- Schulz, C., Schneider, J., Amorim Holanda, B., Appel, O., Costa, A., de Sá, S. S., Dreiling, V., Fütterer, D., Jurkat-Witschas, T., Klimach, T., Knote, C., Krämer, M., Martin, S. T., Mertes, S., Pöhlker, M. L., Sauer, D., Voigt, C., Walser, A., Weinzierl, B., Ziereis, H., Zöger, M., Andreae, M. O., Artaxo, P., Machado, L. A. T., Pöschl, U., Wendisch, M., and Borrmann, S.: Aircraft-based observations of isoprene-epoxydiol-derived secondary organic aerosol (IEPOX-SOA) in the tropical upper troposphere over the Amazon region, *Atmospheric Chemistry and Physics*, 18, 14 979–15 001, <https://doi.org/10.5194/acp-18-14979-2018>, 2018.
- Schwantes, R. H., Teng, A. P., Nguyen, T. B., Coggon, M. M., Crouse, J. D., St. Clair, J. M., Zhang, X., Schilling, K. A., Seinfeld, J. H., and Wennberg, P. O.: Isoprene NO₃ oxidation products from the RO₂ + HO₂ pathway, *The Journal of Physical Chemistry A*, 119, 10 158–10 171, <https://doi.org/10.1021/acs.jpca.5b06355>, 2015.
- Seakins, P. W. and Blitz, M. A.: Developments in laboratory studies of gas-phase reactions for atmospheric chemistry with applications to isoprene oxidation and carbonyl chemistry, *Annual Review of Physical Chemistry*, 62, 351–373, <https://doi.org/10.1146/annurev-physchem-032210-102538>, 2011.
- Seaman, V. Y., Charles, M. J., and Cahill, T. M.: A sensitive method for the quantification of acrolein and other volatile carbonyls in ambient air, *Analytical Chemistry*, 78, 2405–2412, <https://doi.org/10.1021/ac051947s>, 2006.
- Seco, R., Peñuelas, J., Filella, I., Llusà, J., Molowny-Horas, R., Schallhart, S., Metzger, A., Müller, M., and Hansel, A.: Contrasting winter and summer VOC mixing ratios at a forest site in the Western Mediterranean Basin: the effect of local biogenic emissions, *Atmospheric Chemistry and Physics*, 11, 13 161–13 179, <https://doi.org/10.5194/acp-11-13161-2011>, 2011.

- 1030 Sfendla, Y., Stavrakou, T., Müller, J.-F., Oomen, G.-M., Opacka, B., Danckaert, T., De Smedt, I., and Lerot, C.: Global VOC emissions quantified from inversion of TROPOMI spaceborne formaldehyde and glyoxal data, *Atmospheric Chemistry and Physics*, 26, 733–767, <https://doi.org/10.5194/acp-26-733-2026>, 2026.
- Shapiro, E. L., Szprengiel, J., Sareen, N., Jen, C. N., Giordano, M. R., and McNeill, V. F.: Light-absorbing secondary organic material formed by glyoxal in aqueous aerosol mimics, *Atmospheric Chemistry and Physics*, 9, 2289–2300, [https://doi.org/10.5194/acp-9-2289-](https://doi.org/10.5194/acp-9-2289-2009)
1035 2009, 2009.
- Silva, S. J., Heald, C. L., and Li, M.: Space-based constraints on terrestrial glyoxal production, *Journal of Geophysical Research: Atmospheres*, 123, 13,583–13,594, <https://doi.org/10.1029/2018JD029311>, 2018.
- Sinreich, R., Coburn, S., Dix, B., and Volkamer, R.: Ship-based detection of glyoxal over the remote tropical Pacific Ocean, *Atmospheric Chemistry and Physics*, 10, 11 359–11 371, <https://doi.org/10.5194/acp-10-11359-2010>, 2010.
- 1040 Sprengnether, M., Demerjian, K. L., Donahue, N. M., and Anderson, J. G.: Product analysis of the OH oxidation of isoprene and 1,3-butadiene in the presence of NO, *Journal of Geophysical Research: Atmospheres*, 107, ACH 8–1–ACH 8–13, <https://doi.org/10.1029/2001JD000716>, 2002.
- Stavrakou, T., Müller, J.-F., De Smedt, I., Van Roozendaal, M., Kanakidou, M., Vrekoussis, M., Wittrock, F., Richter, A., and Burrows, J. P.: The continental source of glyoxal estimated by the synergistic use of spaceborne measurements and inverse modelling, *Atmospheric*
1045 *Chemistry and Physics*, 9, 8431–8446, <https://doi.org/10.5194/acp-9-8431-2009>, 2009.
- Stone, D., Evans, M. J., Commane, R., Ingham, T., Floquet, C. F. A., McQuaid, J. B., Brookes, D. M., Monks, P. S., Purvis, R., Hamilton, J. F., Hopkins, J., Lee, J., Lewis, A. C., Stewart, D., Murphy, J. G., Mills, G., Oram, D., Reeves, C. E., and Heard, D. E.: HO_x observations over West Africa during AMMA: impact of isoprene and NO_x, *Atmospheric Chemistry and Physics*, 10, 9415–9429, <https://doi.org/10.5194/acp-10-9415-2010>, 2010.
- 1050 Sun, S., Palmer, P. I., Siddans, R., Kerridge, B. J., Ventress, L., Edtbauer, A., Ringsdorf, A., Pfannerstill, E. Y., and Williams, J.: Seasonal isoprene emission estimates over tropical South America inferred from satellite observations of isoprene, *Atmospheric Chemistry and Physics*, 25, 15 801–15 818, <https://doi.org/10.5194/acp-25-15801-2025>, 2025.
- Taraborrelli, D., Lawrence, M. G., Butler, T. M., Sander, R., and Lelieveld, J.: Mainz Isoprene Mechanism 2 (MIM2): an isoprene oxidation mechanism for regional and global atmospheric modelling, *Atmospheric Chemistry and Physics*, 9, 2751–2777, <https://doi.org/10.5194/acp-9-2751-2009>, 2009.
- 1055 Teng, A. P., Crounse, J. D., Lee, L., St. Clair, J. M., Cohen, R. C., and Wennberg, P. O.: Hydroxy nitrate production in the OH-initiated oxidation of alkenes, *Atmospheric Chemistry and Physics*, 15, 4297–4316, <https://doi.org/10.5194/acp-15-4297-2015>, 2015.
- Teng, A. P., Crounse, J. D., and Wennberg, P. O.: Isoprene peroxy radical dynamics, *Journal of the American Chemical Society*, 139, 5367–5377, <https://doi.org/10.1021/jacs.6b12838>, 2017.
- 1060 Travis, K. R., Nault, B. A., Crawford, J. H., Bates, K. H., Blake, D. R., Cohen, R. C., Fried, A., Hall, S. R., Huey, L. G., Lee, Y. R., Meinardi, S., Min, K.-E., Simpson, I. J., and Ullman, K.: Impact of improved representation of volatile organic compound emissions and production of NO_x reservoirs on modeled urban ozone production, *Atmospheric Chemistry and Physics*, 24, 9555–9572, <https://doi.org/10.5194/acp-24-9555-2024>, 2024.
- Tripathi, N., Sahu, L. K., Singh, A., Yadav, R., Patel, A., Patel, K., and Meenu, P.: Elevated Levels of Biogenic Nonmethane Hydrocarbons in the Marine Boundary Layer of the Arabian Sea During the Intermonsoon, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD032 869, <https://doi.org/10.1029/2020JD032869>, 2020.
- 1065

- Van Der Werf, G. R., Randerson, J. T., Giglio, L., Van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., Van Marle, M. J., Morton, D. C., and Collatz, G. J.: Global fire emissions estimates during 1997-2016, *Earth System Science Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
- 1070 Vansco, M. F., Caravan, R. L., Zuraski, K., Winiberg, F. A. F., Au, K., Trongsiwat, N., Walsh, P. J., Osborn, D. L., Percival, C. J., Khan, M. A. H., Shallcross, D. E., Taatjes, C. A., and Lester, M. I.: Experimental evidence of dioxole unimolecular decay pathway for isoprene-derived criegee intermediates, *The Journal of Physical Chemistry A*, 124, 3542–3554, <https://doi.org/10.1021/acs.jpca.0c02138>, 2020.
- Veeffkind, J., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H., de Haan, J., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P.: 1075 TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sensing of Environment*, 120, 70–83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.
- Volkamer, R., San Martini, F., Molina, L. T., Salcedo, D., Jimenez, J. L., and Molina, M. J.: A missing sink for gas-phase glyoxal in Mexico City: Formation of secondary organic aerosol, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2007GL030752>, 2007.
- Volkamer, R., Baidar, S., Campos, T. L., Coburn, S., DiGangi, J. P., Dix, B., Eloranta, E. W., Koenig, T. K., Morley, B., Ortega, I., Pierce, 1080 B. R., Reeves, M., Sinreich, R., Wang, S., Zondlo, M. A., and Romashkin, P. A.: Aircraft measurements of BrO, IO, glyoxal, NO₂, H₂O, O₂-O₂ and aerosol extinction profiles in the tropics: comparison with aircraft-/ship-based in situ and lidar measurements, *Atmospheric Measurement Techniques*, 8, 2121–2148, <https://doi.org/10.5194/amt-8-2121-2015>, 2015.
- Wang, H., Welch, A. M., Nagalingam, S., Leong, C., Czimczik, C. I., Tang, J., Seco, R., Rinnan, R., Vettikkat, L., Schobesberger, S., Holst, T., Brijesh, S., Sheesley, R. J., Barsanti, K. C., and Guenther, A. B.: High temperature sensitivity of Arctic isoprene emissions explained 1085 by sedges, *Nature Communications*, 15, 6144, <https://doi.org/10.1038/s41467-024-49960-0>, 2024.
- Wang, R., Shen, H., Zeng, C., Chen, J., Wang, Y., and Li, Y.: A global land daily 10-km-resolution surface ozone dataset from 2013–2022, *Scientific Data*, 12, 1710, <https://doi.org/10.1038/s41597-025-05990-x>, 2025.
- Wang, S., Newland, M. J., Deng, W., Rickard, A. R., Hamilton, J. F., Muñoz, A., Ródenas, M., Vázquez, M. M., Wang, L., and 1090 Wang, X.: Aromatic photo-oxidation, a new source of atmospheric acidity, *Environmental Science & Technology*, 54, 7798–7806, <https://doi.org/10.1021/acs.est.0c00526>, 2020.
- Warman, W. J.: Chamber studies of glyoxal yields from the oxidation of selected biogenic VOCs, Ph.d. thesis, University of Leeds, <https://etheses.whiterose.ac.uk/35058/>, 2024.
- Waxman, E. M., Elm, J., Kurtén, T., Mikkelsen, K. V., Ziemann, P. J., and Volkamer, R.: Glyoxal and methylglyoxal setschenow salting constants in sulfate, nitrate, and chloride solutions: measurements and gibbs energies, *Environmental Science & Technology*, 49, 11 500– 1095 11 508, <https://doi.org/10.1021/acs.est.5b02782>, 2015.
- Wells, K. C., Millet, D. B., Payne, V. H., Deventer, M. J., Bates, K. H., de Gouw, J. A., Graus, M., Warneke, C., Wisthaler, A., and Fuentes, J. D.: Satellite isoprene retrievals constrain emissions and atmospheric oxidation, *Nature*, 585, 225–233, <https://doi.org/10.1038/s41586-020-2664-3>, 2020.
- Wennberg, P. O., Bates, K. H., Crounse, J. D., Dodson, L. G., McVay, R. C., Mertens, L. A., Nguyen, T. B., Praske, E., Schwantes, R. H., 1100 Smarte, M. D., St Clair, J. M., Teng, A. P., Zhang, X., and Seinfeld, J. H.: Gas-phase reactions of isoprene and its major oxidation products, *Chemical Reviews*, 118, 3337–3390, <https://doi.org/10.1021/acs.chemrev.7b00439>, 2018.
- Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmospheric Environment (1967)*, 23, 1293–1304, [https://doi.org/10.1016/0004-6981\(89\)90153-4](https://doi.org/10.1016/0004-6981(89)90153-4), 1989.

- Wittrock, F., Richter, A., Oetjen, H., Burrows, J. P., Kanakidou, M., Myriokefalitakis, S., Volkamer, R., Beirle, S., Platt, U.,
1105 and Wagner, T.: Simultaneous global observations of glyoxal and formaldehyde from space, *Geophysical Research Letters*, 33,
<https://doi.org/10.1029/2006GL026310>, 2006.
- Wofsy, S. and Team, A. S.: ATom: Aircraft flight track and navigational data, Tech. rep., ORNL Distributed Active Archive Center, Tennessee,
USA, <https://doi.org/10.3334/ORNLDAAAC/1613>, 2018.
- Xu, L., Møller, K. H., Crounse, J. D., Kjaergaard, H. G., and Wennberg, P. O.: New insights into the radical chemistry
1110 and product distribution in the OH-initiated oxidation of benzene, *Environmental Science & Technology*, 54, 13467–13477,
<https://doi.org/10.1021/acs.est.0c04780>, 2020.
- Yan, Y., Cabrera-Perez, D., Lin, J., Pozzer, A., Hu, L., Millet, D. B., Porter, W. C., and Lelieveld, J.: Global tropospheric effects of aromatic
chemistry with the SAPRC-11 mechanism implemented in GEOS-Chem version 9-02, *Geoscientific Model Development*, 12, 111–130,
<https://doi.org/10.5194/gmd-12-111-2019>, 2019.
- 1115 Yang, X., Wang, H., Lu, K., Ma, X., Tan, Z., Long, B., Chen, X., Li, C., Zhai, T., Li, Y., Qu, K., Xia, Y., Zhang, Y., Li, X., Chen, S., Dong,
H., Zeng, L., and Zhang, Y.: Reactive aldehyde chemistry explains the missing source of hydroxyl radicals, *Nature Communications*, 15,
1648, <https://doi.org/10.1038/s41467-024-45885-w>, 2024.
- Yoon, J. Y. S., Wells, K. C., Millet, D. B., Frankenberg, C., Sanghavi, S., Swann, A. L. S., Thornton, J. A., and Turner, A. J.: Inferring
drivers of tropical isoprene: competing effects of emissions and chemistry, *Atmospheric Chemistry and Physics*, 26, 4509–4529,
1120 <https://doi.org/10.5194/acp-26-4509-2026>, 2026.
- Yu, J., Flagan, R. C., and Seinfeld, J. H.: Identification of Products Containing -COOH, -OH, and -CO in Atmospheric Oxidation of
Hydrocarbons, *Environmental Science & Technology*, 32, 2357–2370, <https://doi.org/10.1021/es980129x>, 1998.
- Zhang, W. and Gu, D.: Geostationary satellite reveals increasing marine isoprene emissions in the center of the equatorial Pacific Ocean, *Npj
Climate and Atmospheric Science*, 5, 83, <https://doi.org/10.1038/s41612-022-00311-0>, 2022.
- 1125 Zhang, W., Weber, J., Archibald, A. T., Abraham, N. L., Booge, D., Yang, M., and Gu, D.: Global Atmospheric Composition Effects from
Marine Isoprene Emissions, *Environmental Science & Technology*, 59, 2554–2564, <https://doi.org/10.1021/acs.est.4c10657>, 2025.
- Zhang, Y., Zhang, R., Yu, J., Zhang, Z., Yang, W., Zhang, H., Lyu, S., Wang, Y., Dai, W., Wang, Y., and Wang, X.: Isoprene Mixing
Ratios Measured at Twenty Sites in China During 2012–2014: Comparison With Model Simulation, *Journal of Geophysical Research:
Atmospheres*, 125, e2020JD033523, <https://doi.org/10.1029/2020JD033523>, 2020.
- 1130 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and
Zhang, Q.: Trends in China’s anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmospheric Chemistry and
Physics*, 18, 14095–14111, <https://doi.org/10.5194/acp-18-14095-2018>, 2018.
- Zheng, B., Zhang, Q., Geng, G., Chen, C., Shi, Q., Cui, M., Lei, Y., and He, K.: Changes in China’s anthropogenic emissions and air quality
during the COVID-19 pandemic in 2020, *Earth System Science Data*, 13, 2895–2907, <https://doi.org/10.5194/essd-13-2895-2021>, 2021.
- 1135 Zhong, Q., Schutgens, N., van der Werf, G. R., van Noije, T., Bauer, S. E., Tsigaridis, K., Mielonen, T., Checa-Garcia, R., Neubauer, D.,
Kipling, Z., Kirkevåg, A., Olivíć, D. J. L., Kokkola, H., Matsui, H., Ginoux, P., Takemura, T., Le Sager, P., Rémy, S., Bian, H., and Chin,
M.: Using modelled relationships and satellite observations to attribute modelled aerosol biases over biomass burning regions, *Nature
Communications*, 13, 5914, <https://doi.org/10.1038/s41467-022-33680-4>, 2022.
- Zhou, S., Gonzalez, L., Leithead, A., Finewax, Z., Thalman, R., Vlasenko, A., Vagle, S., Miller, L. A., Li, S.-M., Bureekul, S., Furutani,
1140 H., Uematsu, M., Volkamer, R., and Abbatt, J.: Formation of gas-phase carbonyls from heterogeneous oxidation of polyunsaturated

- fatty acids at the air–water interface and of the sea surface microlayer, *Atmospheric Chemistry and Physics*, 14, 1371–1384, <https://doi.org/10.5194/acp-14-1371-2014>, 2014.
- Zhou, X. and Mopper, K.: Apparent partition coefficients of 15 carbonyl compounds between air and seawater and between air and freshwater; implications for air-sea exchange, *Environmental Science & Technology*, 24, 1864–1869, <https://doi.org/10.1021/es00082a013>, 1990a.
- 1145 Zhou, X. and Mopper, K.: Measurement of sub-parts-per-billion levels of carbonyl compounds in marine air by a simple cartridge trapping procedure followed by liquid chromatography, *Environmental Science & Technology*, 24, 1482–1485, <https://doi.org/10.1021/es00080a004>, 1990b.
- Zhu, Y. and Kieber, D. J.: Concentrations and Photochemistry of Acetaldehyde, Glyoxal, and Methylglyoxal in the Northwest Atlantic Ocean, *Environmental Science & Technology*, <https://doi.org/10.1021/acs.est.9b01631>, 2019.
- 1150 Ziskin, D.: MOPITT CO gridded monthly means (Thermal Infrared Radiances) V009, <https://doi.org/10.5067/TERRA/MOPITT/MOP03TM.009>, 2021.
- Zou, Z., Chen, Q., Xia, M., Yuan, Q., Chen, Y., Wang, Y., Xiong, E., Wang, Z., and Wang, T.: OH measurements in the coastal atmosphere of South China: possible missing OH sinks in aged air masses, *Atmospheric Chemistry and Physics*, 23, 7057–7074, <https://doi.org/10.5194/acp-23-7057-2023>, 2023.




Figure 1. Annual mean (July 2019—une 2020) atmospheric glyoxal column concentrations as (a) observed by TROPOMI, and as simulated in the (b) [GC-CTRL](#), (c) [GC-TM-E](#), (d) [GC-T-EC](#), and (e) [GC-TM-EC](#) experiments. Model statistics against the TROPOMI observations over land, ocean, and globally are shown on the side of panels (b) to (e). [The simulated VCDs were sampled at the daily satellite overpass time \(13:30 local time\) and then averaged to annual means.](#)

1155 **Annual mean sources and sinks of simulated atmospheric glyoxal in the simulation unit: $\text{kg km}^{-2} \text{year}^{-1}$. Global annual sums of individual sources and sinks are shown inset.**




Figure 2. Scatter plots of simulated versus observed (Table S1) glyoxal concentrations (unit: ppt) over (a) land and (b) ocean. Blue and red symbols indicate results from the GC-CTRL and GC-TM-EC experiments, respectively. Observations in urban areas or those strongly affected by fire are denoted by squares (noted as 'Outliers'). Horizontal and vertical error bars indicate standard deviations of the observations and model results, respectively. Dashed lines indicate the 1:1, 1:2, and 2:1 lines. Solid lines indicate the reduced major-axis regression; regression parameters are shown inset. [The references of all corresponding observations are listed in Table S1.](#)




Figure 3. Annual mean tropospheric isoprene ~~column concentrations~~ VCDs (a) as observed by CrIS and as simulated in the (b) GC-CTRL and (d) GC-TM-E experiments, respectively. Also shown are the annual mean isoprene emissions (c) from the GC-CTRL simulation and (e) those scaled and used in the GC-TM-E simulation (also used in GC-T-EC and GC-TM-EC). The areas over water or with no data are shaded white in (a) (b) and (d). The numbers in red boxes indicate the regional mean ratios of isoprene emissions in (e) relative to those in (c).




Figure 4. Annual mean sources and sinks of simulated atmospheric glyoxal in the GC-TM-EC simulation [unit: $\text{kg km}^{-2} \text{ year}^{-1}$]. Global annual sums of individual sources and sinks are shown inset.




Figure 5. Gas-phase production of glyoxal and glycolaldehyde from the oxidation of isoprene as represented in the standard RCIM (black arrows) and as revised in this work (red arrows). Branching ratios outside and inside parentheses were calculated under NO concentrations of 0.1 ppb and 5 ppb, respectively. The solid or dashed arrows denote pathways comprehensively or partially constrained by experiments, respectively. We set $[\text{OH}] = 1.8 \times 10^6 \text{ molecules cm}^{-3}$, $[\text{O}_3] = 40 \text{ ppb}$, and $\text{RH} = 55\%$; other environmental conditions were representative of a subtropical site in fall. Pathways with glyoxal yields less than 1% are not shown in this figure.




Figure 6. Glyoxal yields from isoprene-OH oxidation as a function of NO_x and OH concentrations. Molar yields of glyoxal from isoprene+OH oxidation under different daily mean NO_x and OH concentrations in (a) our revised chemical mechanism used in GC-T-EC, (b) the RCIM as implemented in the GEOS-Chem v14.2.3 mechanism, and (c) MCM v3.3.1. (d) Comparison of simulated glyoxal molar yields from isoprene+OH oxidation under different NO concentrations at a constant daily mean [OH] = 1.8×10^6 molecules cm⁻³.




Figure 7. Comparison of the satellite-retrieved and simulated column concentrations-VCDs of glyoxal (from TROPOMI) and isoprene (from CrIS-ANN) and their ratios over regions dominated by seasonal biogenic NMVOC emissions: (a) Southeast U.S. (August to October), (b) South America (annual mean), (c) Africa (annual mean), and (d) Southeast Asia (annual mean). Error bars indicate standard deviation of monthly mean column concentrations-VCDs and ratios, respectively.




Figure 8. Simulated (a) OH (unit: ppt) and (d) HO₂ (unit: ppt) concentrations, (g) NMHC OHR (unit: s⁻¹), and O₃ in ppb among the GC-CTRL, GC-TM-E and GC-TM-EC simulations

Table 1. Global emissions of glyoxal and its precursors in the revised and standard (in parentheses) GEOS-Chem simulations^a

Species	Emitting sources (Tg yr ⁻¹)				Total
	Biogenic	Anthropogenic (non-biofuel)	Anthropogenic (biofuel)	Biomass burning	
Isoprene	428 (443) 438 (449)	-	-	-	428 (443) 438 (449)
Monoterpenes	140 (140) ^b	-	-	-	140 (140) ^b
Methylbutenol	1.7 (0) ^b	-	-	-	1.7 (0) ^b
Acetylene	-	2.1 (2.1)	1.9 (1.9)	1.9 (1.2)	5.9 (5.2)
Ethylene	-	5.3 (5.3)	0.2 (0.2)	8.3 (5.2)	14 (11)
Benzene	-	3.0 (3.0)	3.0 (3.0)	3.0 (1.9)	9.0 (7.9)
Toluene	-	6.2 (6.2)	1.4 (1.4)	1.8 (1.1)	9.4 (8.7)
Xylenes	-	6.2 (6.2)	0.8 (0.8)	0.8 (0.5)	7.8 (7.5)
Glyoxal (primary)	-	-	1.3 (0)	3.8 (2.1)	5.2 (2.1)
Glycolaldehyde (primary)	-	-	1.0 (0)	4.0 (2.2)	5.0 (2.2)
Hypothetical marine precursor ^c	170 (0)	-	-	-	170 (0)

^a Numbers in the parentheses were emissions in the standard GEOS-Chem (GC-CTRL) simulation. Numbers outside the parentheses were used in the GC-TM-E, GC-T-EC, and GC-TM-EC simulations.

^b The standard GEOS-Chem (GC-CTRL) simulation included biogenic emissions of monoterpenes and methylbutenol but did not include their production of glyoxal.

^c The hypothetical marine precursor was assumed to have a molecular weight of 58, the same as glyoxal. This hypothetical emission was applied only in the GC-TM-E and GC-TM-EC experiments and resulted in a secondary source of glyoxal of 66 Tg yr⁻¹ in the global MBL.

Table 2. Global budget of atmospheric glyoxal estimated in this work compared to Fu et al. (2008).

		GC-CTRL	GC-T-EC	GC-TM-EC	Fu et al., 2008
Burden, Gg	Total	11 <u>10</u>	15	39	15
	Over land	6.8 <u>6.4</u>	10	10	ND ^a
	Over ocean	3.8 <u>3.6</u>	5.3	29	ND ^a
Lifetime, h		3.8	3.4 <u>3.3</u>	3.5 <u>3.2</u>	2.9
Sources, Tg yr⁻¹	Total	25 <u>23</u>	44 <u>40</u>	110 <u>106</u>	45
	Biomass burning emission	2.1	3.8	3.8	5.2
	Biofuel emission	0	1.3	1.3	2.4
	Chemical production	23 <u>21</u>	39 <u>35</u>	39 <u>35</u>	37
	Isoprene	9.1 <u>7.8</u>	17.1 <u>14</u>	17.1 <u>14</u>	21
	Monoterpenes	0	3.8 <u>3.9</u>	3.8 <u>3.9</u>	2.5
	Methylbutenol	0	0.1	0.1	0.3
	Acetylene	6.2	7.0	7.0	8.9
	Ethylene	0.8	2.6	2.6	2.5
	Aromatics	6.8 <u>6.1</u>	7.4 <u>6.9</u>	7.4 <u>6.9</u>	2.0
	Glycolaldehyde (primary)	0.1	0.5	0.5	0.5
	Hypothetical marine precursor	0	0	66	0
	Sinks, Tg yr⁻¹	Total	25 <u>23</u>	44 <u>40</u>	110 <u>106</u>
Photolysis		17 <u>15</u>	23	62 <u>61</u>	28
Oxidation by OH		4.5 <u>4.3</u>	10 <u>6.4</u>	21 <u>20</u>	6.5
Oxidation by NO ₃		<0.1	<0.1	<0.1	<0.1
Uptake on wet aerosol and cloud droplets		0.6 <u>0.8</u>	6.4 <u>5.6</u>	7.6 <u>6.9</u>	6.4
Dry deposition		1.5	2.8 <u>2.6</u>	11 <u>10</u>	2.2
Wet deposition		1.3 <u>1.4</u>	2.0 <u>2.1</u>	8.3 <u>8.2</u>	1.9

^a Not documented.