

Hydrological drivers of hydrogen cyanide wildfire emissions from Indonesian peat fires during the 2015, 2019, and 2023 El Niño events

Antonio G. Bruno^{1,2}, David P. Moore^{1,2}, Jeremy J. Harrison^{1,2}, Ailish Graham^{3,4}, Martyn P. Chipperfield^{3,4}, and Corinne Vigouroux⁵

¹School of Physics and Astronomy, University of Leicester, Leicester, UK

²National Centre for Earth Observation, University of Leicester, Leicester, UK

³School of Earth and Environment, University of Leeds, Leeds, UK

⁴National Centre for Earth Observation, University of Leeds, Leeds, UK

⁵Royal Belgian Institute for Space Aeronomy, Brussels, Belgium

Correspondence: Antonio G. Bruno (agb22@leicester.ac.uk)

Abstract. Indonesian peatlands store vast amounts of carbon that are highly vulnerable to fire during El Niño-driven droughts. When ignited, peat releases large quantities of greenhouse gases and other species with significant environmental impacts, including hydrogen cyanide (HCN), a sensitive tracer of smouldering combustion. In this work, we use new satellite retrievals from the Infrared Atmospheric Sounding Interferometer (IASI) spectra using the University of Leicester IASI retrieval scheme (ULIRS), that we validate against ground-based ~~FTIR~~ Fourier transform infrared (FTIR) measurements at Reunion Maïdo, TOMCAT atmospheric model simulations, hydrological information, and fire activity observations to evaluate the factors driving trace gas emissions during the 2015, 2019, and 2023 El Niño events.

The 2015 El Niño produced large burdens of HCN and carbon monoxide (CO), unprecedented in the satellite observational era and driven by exceptionally low soil moisture, depressed groundwater levels, and deep burn depths. In contrast, the 2019 and 2023 events exhibited markedly weaker emissions despite similar Oceanic Niño Index (ONI) anomalies, reflecting more favourable hydrological conditions. Comparisons of the satellite trace gas observations with simulations of the TOMCAT model show that burned-area-based inventories such as the Global Fire Emissions Database (GFED) version 4 substantially overestimate emissions from peat fires in 2015, while a new peat-specific database, FINNpeatSM (Fire Inventory from NCAR (FINNv1.5) New peat with Soil Moisture), better represents fire season timing and burn depth by incorporating soil moisture constraints. From satellite-derived HCN:CO enhancement ratios, we provide new emission factors for HCN that offer benchmarks for new emission inventories.

Our results show that peat fire intensity and emissions are driven not only by El Niño strength but also by local hydrological conditions such as soil water content and precipitation. Integrating hydrological indicators with satellite observations of atmospheric composition is therefore critical for improving fire emission inventories.

20 1 Introduction

During the South-East Asian dry season, El Niño generally has a strong influence on the conditions driving widespread fires. Fires are commonly used across the region for land clearing to manage fields and prepare the soil for the growing season. However, much of the land in Kalimantan and Sumatra, Indonesia, is underlain by tropical peat soils that have been extensively drained using a network of canals. Dry conditions and elevated temperatures, typical of the dry season, are particularly exacerbated in the Indonesian environment during El Niño, making man-made fires more difficult to control (Parker et al., 2016). Under these conditions, vegetation fires can ignite the underlying carbon-rich peat soils, causing wildfires that are extremely difficult to extinguish and can burn for weeks.

Indonesian peatlands occupy about 55% of the tropical peatland carbon reservoir (Dargie et al., 2017; Page and Hooijer, 2016). They store extremely large amounts of partially decayed organic matter, and release substantial amounts of pyrogenic trace gas species and particulate matter into the atmosphere when burned. The carbon stored can be emitted in the form of trace gases such as carbon dioxide (CO_2), CO and HCN, the latter also badly affecting the air quality across South-East Asia (Watson et al., 2019). In particular, during El Niño years strong peaks in global HCN have been observed (Park et al., 2021; Rosanka et al., 2021), suggesting that it is a good atmospheric tracer for peat fire (Pumphrey et al., 2018; Sheese et al., 2017). Monitoring HCN emissions is therefore important in understanding severe peatland fires, and for understanding global carbon accounting.

Peatlands are particularly vulnerable to smouldering fires, which are characterised by slow, low-temperature, flameless combustion. Peat fires typically develop in three stages (Usup et al., 2004). In the first, surface peat is ignited by a surface fire event. In the second stage a smouldering front burns laterally and downward into the surface (up to 20 cm), and in the third the smouldering extends into deeper peat layers below 20 cm in depth. It is the latter smouldering stages in which HCN is primarily released. Ultimately, it is the peat soil moisture, in particular the ground water level (GWL), that determines the ignition and spread of these smouldering fires.

During the period September to November 2015, under the influence of a strong El Niño, Indonesia experienced the most severe wildfire season of the last three decades. The intensity of the 2015-2016 El Niño event (ONI: +2.6 deg C) was comparable to the 1997-1998 El Niño (ONI: +2.4 deg C)(Anthony G. Bamston and Goldenberg, 1997; Huang et al., 2017), one of the strongest recorded (Field et al., 2016; Santoso et al., 2017). The peatland fires lasted for about three months during which more than 2.6 Mha of forest, peat and other land types were burned, and an equivalent of 5% of the global 2015 fossil fuel CO_2 emissions were released (Vetrita and Cochrane, 2020). In the period 1997-2016, fires in equatorial Asia and particularly Indonesia produced about 8% of the global fire carbon emissions (van der Werf et al., 2017).

The 2023-2024 El Niño period is currently the second strongest event of this century (ONI: +2.0 deg C), falling just behind that of 2015-2016. However, as will be shown, there is a marked difference in the behaviour of the fire plumes and the amount of HCN released. While previous studies have documented the role of El Niño in intensifying Indonesian fire activity (Field et al., 2016; Whitburn et al., 2017; Nurdianti et al., 2021), large uncertainties remain in quantifying trace gas emissions from peatland combustion, particularly for HCN, a key tracer of peat fires.

Existing fire inventories, such as GFEDv4, rely primarily on burned-area estimates that are poorly suited to capturing underground peat combustion, leading to biases in both the timing and magnitude of emissions. These discrepancies often stem from uncertainties in burned carbon estimates, particularly in peatlands (Nechita-Banda et al., 2018; Lohberger et al., 2018). Peat extent and depth estimation remain highly uncertain in Indonesia (Hooijer and Vernimmen, 2013), and may also reflect inaccuracies in biome attribution.

In this paper, we systematically compare multiple El Niño events to evaluate the influence of large-scale climate forcing and local hydrological parameters on peat fire dynamics. We provide new satellite-based constraints on HCN emission factors, evaluate the performance of the GFED emission inventory, introduce a new database based on the method developed for FINNpeatSM (Kiely et al., 2019), and assess how variations in soil moisture and groundwater level influence interannual differences in wildfire dynamics and emissions.

The paper is structured as follows. Section 2 describes the main datasets and models used in this study, including IASI CO and HCN from ULIRS. In Section 3.1, the IASI CO and HCN retrievals are compared with ground-based ~~FTIR~~ Fourier transform infrared (FTIR) measurements at the Reunion Maïdo NDACC station. In Section 3.2, we investigate HCN emissions during the 2015 Indonesian fire season using total columns retrieved with the ULIRS, and compare with recent TOMCAT model simulations (Bruno et al., 2022, 2023). In this section, we also propose a new estimate of HCN emission factors by calculating the enhancement ratio between HCN and CO, $ER_{HCN/CO}$. Section 3.3 discusses the differences between the 2015, 2019, and 2023 fire seasons and explores their underlying causes, focusing on soil moisture and precipitation as the principal hydrological drivers of fire activity, using the FINNpeatSM approach (Kiely et al., 2019). Finally, Section 4 synthesizes the main findings of this work and highlights the implications of hydrological controls for peatland fire dynamics and trace gas emissions.

2 Data

This Section presents the suite of complementary satellite observations, ground-based measurements, reanalysis products, and emission inventories used to characterise the variability of ~~carbon monoxide (CO) and hydrogen cyanide (HCN)~~ CO and HCN during three recent Indonesian fire seasons (2015, 2019 and 2023) influenced by El Niño. The atmospheric composition is primarily constrained using spaceborne thermal infrared retrievals from IASI and ground-based ~~Fourier-transform-infrared (FTIR)~~ FTIR observations at the NDACC Maïdo station, whereas chemical transport model simulations from the TOMCAT model provide a framework for interpretation. To investigate the hydrological drivers of fire activity and peat combustion, satellite-derived precipitation (IMERG), surface and subsurface soil moisture products (SMAP and SWI), and ERA5 wind fields have been used. In particular, precipitation and SWI temporal variability over Indonesia is used to show the influence of the moisture content of the soil on fire emissions, and the SMAP data are used to produce an enhanced emission database that well represents peat fire emissions. Fire occurrence and emissions are quantified using global (GFED4) and regionally enhanced (FINNpeatSM) inventories, together with high-resolution active fire detections from VIIRS. The combined use of these datasets enables the evaluation of fire emissions, atmospheric transport, and the role of moisture conditions in modulating trace gas emissions over Indonesia and the tropical Indian Ocean.

2.1 Atmospheric CO and HCN data

2.1.1 IASI

IASI is a hyperspectral sounder onboard the three polar-orbiting MetOp satellites, launched in 2006 (IASI-A, now decommissioned), 2012 (IASI-B), and 2018 (IASI-C) jointly by CNES (Centre National d'Études Spatiales) and EUMETSAT (European Organisation for the Exploitation of Meteorological Satellites). The main objective of IASI is to provide a continuous and long-term collection of measurements to support meteorology, namely temperature and humidity tropospheric profiles with high vertical resolution and precision, but it has also been used to monitor both the environment and climate on a global scale through observations of atmospheric composition (Clerbaux et al., 2009; Hilton et al., 2012).

IASI measures the spectrally resolved thermal infrared radiation emitted by the Earth and the atmosphere system at the top of the atmosphere (TOA) across the spectral range 645 cm^{-1} to 2760 cm^{-1} at a spectral resolution of 0.5 cm^{-1} and spectral sampling of 0.25 cm^{-1} . Due to its high spectral resolution and spectral sampling, IASI is able to measure a large number of trace gases, including some species observed only sporadically in the measured spectra (Clerbaux et al., 2009; Clarisse et al., 2011). These species include 10 gases with clear spectral signatures that are always present in the satellite measurements, CO_2 , N_2O , CFC-11, CFC-12, OCS, H_2O , CH_4 , O_3 , CO and HNO_3 (Clarisse et al., 2011; Clerbaux et al., 2009) and 14 reactive trace gases. Some of these reactive gases are the typical products of biomass burning, such as HCN, which has been assessed in previous IASI studies, and is retrieved using the absorption band centred at 712 cm^{-1} (ν_2 band) (Coheur et al., 2009; De Longueville et al., 2021; Duflot et al., 2015, 2013).

The sensitivity of IASI retrievals near the surface is limited and exhibits a strong dependence on thermal contrast (with large differences between daytime and nighttime), which constrains the amount of vertical information that can be obtained. For CO, the averaging kernels indicate two distinct vertical contributions: one from the lower troposphere, peaking on average at about 5 km, and another from the upper troposphere, near 10 km. The top panels of Figure 1 show the CO averaging kernels (solid lines) and total column averaging kernel (dashed lines) for IASI measurements on 2 November within the wildfire plume (110°E , 14.5°S) and outside the plume (95.4°E , 35°S). In contrast, HCN retrievals are primarily sensitive to the mid to upper troposphere, with peak sensitivity between 9 and 12 km, as shown in the bottom panels of Figure 1 reporting HCN averaging kernels (solid lines) and total column averaging kernel (dashed lines) for in-plume and out-of-plume scenarios.

2.1.2 Retrieval algorithm

The concentrations of HCN and CO during the 2015, 2019 and 2023 fire seasons were retrieved using the University of Leicester IASI Retrieval Scheme (ULIRS). The ULIRS scheme is a retrieval algorithm originally developed at the University of Leicester and subsequently by the UK National Centre for Earth Observation to study tropospheric trace gas concentrations from IASI satellite measurements. Here we examine results for carbon monoxide and hydrogen cyanide under clear sky and partially cloudy conditions from the radiances measured by the IASI instrument. A full description of the ULIRS and a summary of the main features are given in (Illingworth et al., 2011a, b), with relevant updates outlined in this work. In essence, the ULIRS retrieves atmospheric trace gas partial and total column information from IASI top-of-atmosphere radiances. To do




Figure 1. CO (top panels) and HCN (bottom panels) averaging kernels and total column averaging kernel for IASI measurements on the 2nd of November withing the wildfire plume (110°E, 14.5°S) and outside the plume (95.4°E, 35°S).

120 this, the scheme incorporates an optimal estimation method (Rodgers, 2000) to constrain the inversion with a priori information about the variables to be retrieved, in this case CO and HCN, alongside H₂O, CO₂ and temperature (profile and surface).

The CO retrieval setup largely follows the setup of (Illingworth et al., 2011a) but with a single global average a priori profile constructed with ACE v3.6 data (Bernath et al., 2021) covering 2004 to 2018 for the stratospheric profile at one-sigma uncertainty (above 15 km). This was merged with the troposphere (below 15 km) using TOMCAT model data over the 2006 to
125 2015 period. The number of retrieval levels compared to (Illingworth et al., 2011a) was also reduced to 17 equidistant pressure levels from the surface up to 50 hPa. The ULIRS CO utilises optimal estimation in common with other IASI schemes such as FORLI-CO (Fast Optimal Retrievals on Layers for IASI) which produces the operational CO product for the EUMETSAT AC-SAF. The main difference being in their vertical grids, a priori assumptions, and forward modeling methods. Both schemes use look-up-tables of absorbances to speed up the radiative transfer calculations.

130 The HCN retrieval employs a fixed altitude grid with 13 levels, 3 km spacing from 0 km to 18 km with levels then at 7 km spacing up to and including 60 km. Due to the weak HCN absorption, ULIRS retrievals are calculated in a limited spectral region, 710-715 cm⁻¹, containing the ν_2 band of HCN in order to avoid interferences from the absorption of other species in the spectrum, in particular avoiding line mixing effects of a CO₂ Q branch centred near 720 cm⁻¹. Only 11 selected spectral channels are used in the process: 710.25, 710.50, 710.75, 712.50, 712.75, 713.50, 713.75, 714.00, 714.25, 714.75, 715.00 cm⁻¹.
135 The a priori profile was constructed from INTEx-B aircraft information (Singh et al., 2009) in the troposphere and ACE v4 data (Boone et al., 2020) in the stratosphere over polluted scenes. A loose constraint on the HCN a priori uncertainty of 100% was assumed. Here we limit both the CO and HCN retrievals to a maximum of 10 iterations. Other schemes that produce HCN products from IASI include the ANNI (Artificial Neural Network for IASI) (Dufлот et al., 2013), which provides rapid, global, total column mapping. While ULIRS retrievals often have a Degrees of Freedom for Signal (DOFS) near 1 for HCN,
140 suggesting limited strict vertical profiling, the scheme still provides sensitivity to broad vertical distributions rather than a fixed total column. This allows for analysis of vertical partitioning and high-altitude transport (e.g., biomass burning plumes) in the troposphere.

ULIRS makes use of the Reference Forward Model (RFM), a line-by-line (LBL) radiative transfer model developed at the University of Oxford (Dudhia, 2017), which can be used to simulate the TOA spectrum as measured by a spaceborne sensor. Line-by-line methods for numerically solving RT equations, such as the RFM, can be time-consuming and require large
145 computational power. Retrieval schemes generally require much faster output, so to improve RFM speeds, a set of absorption coefficients for each absorbing gas is calculated beforehand and interpolated onto the IASI grid. RFM calculations in the ULIRS CO and HCN retrievals make use of spectroscopic data in the form of derived look-up tables (LUTs) including H₂O and O₃. For CO, LUTs were used from Vincent and Dudhia (2017). For HCN these were derived using a set of Curtis-Godson
150 equivalent pressures and nine temperature levels spanning ± 60 K from the mid-latitude reference atmosphere of Remedios et al. (2007). These climatological reference atmospheres cover tropical, mid-latitude day/night and polar summer/winter atmospheric conditions. Testing the RFM radiance output via the LUTs, using the climatology files of Remedios et al. (2007), yielded results within 1% of the full LBL calculations. [A summary of the main information of the CO and HCN retrieval is present at Table S1 of the Supplementary Material.](#)

The Maito Observatory (21.08°S, 55.38°E; ~2160 m a.s.l.), located on Réunion Island in the Indian Ocean, is a high-altitude site of the Network for the Detection of Atmospheric Composition Change (NDACC), operational since 2012 and within the programme since 2013. This site produces long-term observations of atmospheric composition in the tropical Southern Hemisphere. Being located on a mountain ridge above the marine boundary layer (MBL), Maito provides conditions suitable for atmospheric composition measurements representative of the free troposphere (FT) and the upper troposphere–lower stratosphere (UTLS) (Baray et al., 2013). A high-resolution Bruker IFS 125HR Fourier transform infrared (FTIR) spectrometer acquires direct solar absorption spectra. These observations enable the retrieval of total and partial columns of several trace gases, including CO and HCN (Duflo et al., 2015; Vigouroux et al., 2012).

The Maito Observatory is also appropriately located to observe the outflow of biomass burning pollution from Southern Africa (Vigouroux et al., 2012), from South America for long-lived species such as CO and HCN, and from South-East Asia (Duflo et al., 2010). To do this the Reunion Maito site makes regular, year-round measurements of trace-gas species including CO and HCN, using the optimal estimation method as implemented in SFIT4 (v0.9.4.4). The FTIR vertical profiles of volume mixing ratios are weighted by the airmasses in each retrieval layer and integrated to give the total or partial columns in molecules/cm². More detail on ground-based the FTIR dataset, retrieval method, and error budget can be found in (Vigouroux et al., 2012).

The datasets produced are widely used to investigate long-range transport and the impact of biomass-burning emissions over the tropical Indian Ocean, satellite measurements, and model validation (Wells et al., 2025; Callewaert et al., 2022; Zhou et al., 2018; De Mazière et al., 2018). These CO and HCN FTIR data sets are well suited for the validation of our ULIRS products.

2.1.4 TOMCAT

The TOMCAT 3-D chemical transport model (CTM) is a global offline Eulerian model widely used for both tropospheric and stratospheric chemistry studies. Originally developed as two separate models TOMCAT and SLIMCAT (Chipperfield et al., 1993), the unified model (Chipperfield, 2006; Monks et al., 2017) has since been applied in numerous studies (Chipperfield et al., 2018; Pope et al., 2020; Bruno et al., 2022). TOMCAT has been included in several model intercomparison studies of tropospheric chemistry transport and has shown good agreement with other models (Ma et al., 2023; Remaud et al., 2023; Krol et al., 2018; Thompson et al., 2014). The model’s meteorological forcings — humidity, temperature, and wind fields — are driven by ERA-Interim reanalysis data from the European Centre for Medium-Range Weather Forecasts (ECMWF), provided at a 6-hour temporal resolution (Berrisford et al., 2011; Dee et al., 2011). These meteorological variables are linearly interpolated to match the temporal resolution and spatial grid of the model. Given the model resolution, the higher spatial resolution of ERA5 compared with ERA-Interim is not expected to significantly affect the simulated transport. Large-scale vertical advection is not driven directly by the reanalysis vertical velocity fields; rather, it is diagnosed from the reanalysis horizontal divergence after averaging to the model grid. Sub-grid-scale transport associated with convection and boundary layer mixing is represented through dedicated parameterizations. No explicit treatment of pyroconvection is included. Surface emissions from both natural

and anthropogenic sources are incorporated at their original resolution and subsequently re-gridded to align with the model's spatial configuration. HCN emissions are derived from several key datasets: anthropogenic and oceanic emissions are sourced from the Coupled Model Intercomparison Project Phase 6 (CMIP6) (Eyring et al., 2016); biogenic emissions are taken from a fixed annual dataset provided by the Chemistry-Climate Model Initiative (CCMI) (Morgenstern et al., 2017); and biomass burning emissions are obtained from the GFED Version 4.1 (Randerson et al., 2017). A complete description of the tracer version of TOMCAT adapted to simulate global atmospheric HCN distributions has been extensively described in Bruno et al. (2022) and Bruno et al. (2023), and evaluated against independent measurements from ACE-FTS and NDACC. This study can be considered a follow-up to previous work; therefore, we used the same setup to ensure consistency with the previous analysis. In the present study, TOMCAT outputs are compared with IASI observations.

This version of the model makes use of the best-fitting ocean uptake scheme defined in Bruno et al. (2023), the Li et al. (2000) fluxes reduced by 75% and the reaction rates proposed by Kleinböhl et al. (2006) for HCN oxidation by OH radicals and $O(^1D)$. Outputs for 2015 were simulated on a horizontal $2.8^\circ \times 2.8^\circ$ (T42 Gaussian) grid on 60 terrain-following vertical levels (surface to ~ 60 km) for HCN. In order to compare the model output (T_HCN) with the IASI daily measurements over a short time period, the model was sampled daily and adapted for use with the monthly GFED v4.1 inputs for HCN biomass burning emissions. The TOMCAT model reads the emissions as monthly means, which are then interpolated in time so that the emissions vary smoothly during the model run.

TOMCAT total columns were constructed from the raw output using the approach proposed by Deeter (2002), in which the model profiles are interpolated onto the ULIRS retrieval grid and smoothed by applying the ULIRS averaging kernels as $x_{TOMCAT}^{smooth} = x_a + A(x_{TOMCAT}^{int} - x_a)$, where x_a is the a priori profile, A the retrieval averaging kernel matrix, and x_{TOMCAT}^{int} the model profile interpolated onto the retrieval grid.

2.2 Precipitation data

The Integrated Multi-satellitE Retrievals for GPM (IMERG) is a NASA precipitation product developed under the joint NASA-JAXA Global Precipitation Measurement (GPM) satellite mission. It provides global surface precipitation estimates at 0.1° spatial and 30-minute temporal resolution from June 2000 onward, using data from a constellation of passive microwave sensors, intercalibrated against data from the Tropical Rainfall Measuring Mission (TRMM; 2000–2014) and the GPM Core Observatory (2014–present). IMERG data cover the majority of the Earth's surface and support a wide range of applications (Pradhan et al., 2022).

This study uses the IMERG Final Run daily product (Huffman et al., 2024), derived by averaging valid half-hourly rates and scaling to 24 hours, which reduces the dry biases present in earlier versions. In particular, the data used here cover the period 2014–2023 over Indonesia.

2.3 Soil moisture data

2.3.1 Soil Moisture Active Passive (SMAP)

220 The SMAP mission, launched by NASA in 2015, is designed to provide global measurements of surface soil moisture and
freeze-thaw conditions every 2–3 days (Entekhabi et al., 2014). Although the radar instrument ceased operation shortly after
launch, the radiometer continues to function reliably, allowing the generation of global soil moisture maps. SMAP data sup-
port a better understanding of the Earth’s water, energy, and carbon cycles by quantifying the amount of liquid water in the
topsoil layer and distinguishing between frozen and unfrozen ground. These observations are essential for improving weather
225 forecasting, climate modeling, and environmental monitoring by capturing temporal changes in soil moisture that influence
land-atmosphere interactions and broader ecological processes.

Recent studies have further expanded the scientific utility of SMAP data. Nayak et al. (2025) demonstrated that SMAP-
derived surface soil moisture variability is a reliable predictor of subsurface water dynamics. Fang et al. (2024) validated
downscaled SMAP products at 1-km resolution using long-term in situ data, confirming their accuracy across heterogeneous
230 landscapes. Additionally, Cho et al. (2024) improved SMAP’s performance in dense vegetation regions by calibrating retrieval
algorithms to reduce bias. Ma et al. (2024) also enhanced SMAP’s global utility by integrating the data with ASCAT ob-
servations through machine learning techniques, resulting in higher resolution and more temporally consistent soil moisture
estimates.

2.3.2 Soil Water Index (SWI)

235 The operational SWI is produced by the Copernicus Land Service from the surface soil moisture (SSM) measured by the
ASCAT scatterometer onboard the MetOp satellites and from Sentinel-1 C-band SAR. The SWI quantifies from these obser-
vations the moisture content of the soil with a spatial resolution of 0.1°. It is a dimensionless index ranging from 0 (dry) to
1 (saturated relative to local climatology), at eight different depths as a function of the characteristic timelength T (1, 5, 10,
15, 20, 40, 60 and 100 days), a parameter expressed in units of time but proportional to the depth of the layer, as described in
240 Bauer-Marschallinger et al. (2018).

Differently from other products such as SMAP, the SWI is more an indicator of the percentage of relative wetness than an
absolute soil water content. It is not limited only to surface soil moisture, as it is able to evaluate water availability at depths
typical of plant roots. For this reason, SWI is a particularly suitable parameter for assessing GWL conditions, since it better
reflects the integrated moisture status of the soil profile and the overall water availability within the root zone.

245 2.4 ERA5 wind data

The fifth generation European Centre for Medium-range Weather Forecasting atmospheric reanalysis (ERA5) is a high-
resolution global atmospheric dataset providing consistent, spatially complete, record of wind and other atmospheric variables
such as temperature from 1940 to the present (Hersbach et al., 2020). ERA5 is based on the Integral Forecasting System (IFS)

Cycle model which simulates atmospheric physics and dynamics constrained by the 4D-Var data assimilation process. The vertical velocity (Pa/s) is calculated based on the divergence of horizontal winds within the model's pressure-based coordinate system, with negative values indicating upward motion. This study uses the ERA5 monthly data on 37 pressure levels, at 0.25°x0.25° spatial resolution, to determine the large-scale vertical velocity across the equator (0°N).

2.5 Fire data

2.5.1 GFED emission database

The GFED4 database quantifies biomass burning emissions by coupling satellite-derived burned area (MODIS), active fire detections, and biogeochemical models (van der Werf et al., 2017). The GFED framework estimates dry matter (DM) consumption as a function of fuel load, combustion completeness, and fire activity across different biomes on a monthly 0.25° global grid. The trace gas emissions are derived by applying biome-specific emission factors derived from field and lab measurements to the DM consumption (Akagi et al., 2011). The spatial distribution of the DM burned is then determined by using satellite observations of BA in combination with biogeochemical modelling. Cloud cover and a limited knowledge of the biome distribution and emission factors over Indonesia and other peat-dominated regions can cause possible errors in estimating fire emissions (Akagi et al., 2011). Another limitation of the approach used to define GFED is its dependence on information about peat consumption by fires — such as the rate, extent, and depth of peat burned — which cannot be easily determined from satellite data (Bruno, 2024a).

2.5.2 FINNpeatSM emission database

FINNpeatSM (Kiely et al., 2019) is a bespoke regional dataset for peat fire emissions in Indonesia. FINNpeatSM uses Fire Inventory from NCAR (FINNv1.5) (Wiedinmyer et al., 2011) for fire detections of above ground vegetation fires. FINNv1.5 combines active fire detections from MODIS, biomass burned and emission factors (EFs) to provide daily fire emissions at 1-km resolution (Wiedinmyer et al., 2011). However, FINNv1.5 only includes emissions from the combustion of above-ground vegetation. Therefore, it does not include emissions from combustion of peat. Kiely et al. (2019) added the emissions from the combustion of peat across Indonesia, to create FINNpeatSM. Details on the method can be found in Kiely et al. (2019). In summary, to add emissions from peat combustion they use a 2-step process. First, they use a peatland distribution map to identify where a fire occurred on peatland. For each fire occurring on peatland they add additional emissions from the peat burning using Eq 1:

$$E_s = BA \times BD \times \rho \times EF_s \quad (1)$$

where E_s is the emissions of a species (s) from an individual fire that occurred on peatland, BA is the burned area and BD is the burn depth, ρ is the peat density and EF_s is the emissions factor of a species (s).

The second step of the method was to scale burn depth relative to soil moisture (from the ESA CCI Soil Moisture Product New Version Release (v04.4): ESA CCI SMv04.4). This step accounts for burn depth increasing as peat dries out and the water table decreases. In FINNpeatSM burn depth is assumed to increase linearly between a minimum of 5 cm to a maximum of 37 cm. Kiely et al. (2019) used soil moisture from ESA CCI SMv04.4, which provided soil moisture retrievals up to 2018. To extend the FINNpeatSM dataset to 2023, Graham et al. (2024) updated the method to use the SMAP level 4 product. In the Level 4 dataset, SMAP measurements of soil moisture in the top 5 cm of the soil column are combined with estimates from a land-surface model. This provides soil moisture in the top 1 m of the soil column. The SMAP is both spatially (9 km) and temporally (3-hourly) complete, the 3-hourly data were aggregated to generate daily-mean values. Daily-mean SMAP values are used to linearly scale the burn depth between a minimum soil moisture of $0.5 \text{ m}^3\text{m}^{-3}$ (5 cm burn depth) and $0.1 \text{ m}^3\text{m}^{-3}$ (37 cm burn depth). In this study, we use this version of FINNpeatSM to explore changes in fire burn depth and emissions between 2015, 2019 and 2023.

2.5.3 VIIRS VNP14IMG Fire Product

The Visible Infrared Imaging Radiometer Suite (VIIRS) active fire product (VNP14IMG) provides the latitude, longitude, time, and a confidence flag for a fire detection pixel. The VNP14IMG product is a development of the MODIS thermal anomaly algorithm and several studies have now shown (Schroeder et al., 2014; Zhang et al., 2017) that it is capable of detecting small fires at a 1 km spatial resolution which MODIS does not have the sensitivity to measure. VIIRS fire products are provided at 375 m, a much higher spatial resolution compared to MODIS, and therefore VIIRS is able to detect small wildfires that MODIS is insensitive to. Only nominal and high confidence fires during nighttime were included in this work.

3 Results

3.1 Comparison of IASI CO and HCN with ground-based FTIR measurements

To test the consistency of the daily CO and HCN measurements from IASI, comparisons have been made to the total column amounts retrieved from the FTIR instrument based at the Reunion Maïdo NDACC station. The FTIR instrument requires clear-sky daylight conditions for measurements; both IASI daytime and night-time measurements are used in these match ups with the only requirement being that IASI measurements are made within 12 hours of the FTIR measurement and that the IASI measurement is within 100 km of the Reunion Maïdo site. IASI cloudy spectra were removed from the dataset using a 25% contamination threshold. As the IASI measurements are scattered within 100 km of the Maïdo site, these data may have a different surface altitude compared to the ground-based site. As we need to compare over the same altitude range, we smooth the FTIR profiles using the IASI averaging kernels. We do this by adjusting the FTIR partial column profile such that the total column corresponds to IASI total columns with different ground level altitudes. The methodology employed to account for the difference between the height of the ground-based measurement station and the IASI measurement site follows the approach of Kerzenmacher et al. (2012), in particular their Fig. 2 where the altitude correction is described in detail. To assess the vertical

sensitivity of the NDACC FTIR dataset, Fig. 2 shows the mean total column averaging kernels at Reunion Maïdo, for HCN and CO measurements made in 2015. The ground-based FTIR shows a good sensitivity to HCN abundance in the upper troposphere – lower stratosphere (UTLS), peaking around 15 km. For CO, the ground-based FTIR shows good sensitivity throughout the troposphere and stratosphere and is constant around one.




Figure 2. Average total column averaging kernels for ground-based FTIR at Reunion Maïdo for CO (red line) and HCN (blue line) for measurements made in 2015.

Figure 3 shows the comparison between IASI and the ground-based measurements. ~~One can see that there is an excellent overall agreement between the IASI and~~ Overall, the IASI observations closely agree with the ground-based FTIR CO measurements. For the 2013-2019 period where FTIR data were available to use, the IASI columns are only biased high by $0.7\% \pm 4.4\%$, where the uncertainty represents the 1-sigma standard deviation on the mean difference. The seasonality of CO is captured well in both datasets (correlation coefficient of 0.97 for the entire daily mean data set) with the peak CO in October and November periods each year, related to the Southern Hemisphere biomass burning season (Vigouroux et al., 2012). The maximum CO observed by IASI was $2.64 \pm 0.052 \times 10^{18}$ molecules cm^{-2} in 2015; the peak FTIR value was $2.71 \pm 0.054 \times 10^{18}$ molecules cm^{-2} in the same year, both instances occurring on 25 November 2015. For HCN at Reunion Maïdo, the IASI seasonality and interannual variability matches very well with the ground-based measurements (correlation coefficient 0.80 for the entire daily mean data set), but with IASI columns biased high by $14.3\% \pm 7.0\%$. This bias could be caused by a lack of

consistency between the HCN spectroscopic line parameters in HITRAN used for the two retrieval schemes. Unlike the ULIRS HCN retrieval, the NDACC retrieval uses the stronger HCN band near 3 micron. The maximum IASI HCN observed in 2015 of $1.66 \pm 0.059 \times 10^{16}$ molecules cm^{-2} is 10.5% higher than the highest HCN values observed in other years during the validation period. Compared to CO, the maximum HCN in 2015 occurred on 18 November, 2015. The CO total columns did not show a high discrepancy between different years in the record.




Figure 3. Time series of CO (top panel) and HCN (bottom panel) measurements for Reunion Maïdo. IASI measurements are shown in red with associated standard deviations (red lines). Ground-based FTIR measurements are shown in light blue with associated total error. The vertical blue (orange) dashed line represents where the IASI L2 (L1C) data processor version changed.

3.2 HCN emissions during the 2015 Indonesian fire season

3.2.1 Satellite observations of HCN in 2015

IASI satellite observations are used to estimate the amount of HCN emitted from the Indonesian region during the 2015 wildfire season; the plume of HCN emitted during the wildfire season is clearly visible in the left panel of Figure 4. In Section 3.3,

these observations are compared with those from the more recent 2019 and 2023 burning seasons, both of which also occurred under El Niño conditions.




Figure 4. IASI global distribution of HCN total column (molecules cm^{-2}) as retrieved by ULIRS for (a, left) 2 November 2015 during daytime and (b, right) the November 2015 monthly mean regridded on a $0.25^\circ \times 0.25^\circ$ grid. The red box indicates the region (12°S - 7°N , 90°E - 127°E) over which the total column values are averaged.

According to the new peatland fire stages classification — stage 1 surface fire, stage 2 shallow peatland fire (depth < 20 cm) and stage 3 deep peatland fire (depth > 20 cm) — proposed by Hayasaka et al. (2020), the Indonesian fires of 2015 were classified as stage 2 for the second half of August and stage 3 from early September to the end of October. This is broadly consistent with the averaged total column of the IASI HCN observations seen in Figure 5, where the highest HCN total columns are observed in September and October. The HCN total columns peak in late October / early November, consistent with the work of Nechita-Banda et al. (2018) on CO emissions from the same peat fires. Their results indicate a sudden increase in CO emissions for the latter half of October 2015.

During neutral conditions, Indonesia is situated under one of the rising branches of the Walker circulation, linked to high rainfall. However, during El Niño, the circulation shifts eastwards and the air over Indonesia tends to sink, resulting in below average rainfall, higher surface pressure, and dryness. A plot of vertical winds at 0° latitude (Indonesia spans 95°E - 140°E longitude) derived from ERA5 for October 2015 (Figure 6) demonstrates this; the air mass over Indonesia is primarily descending. At the end of October / beginning of November, the circulation pattern suddenly changes to one in which the air is predominantly being uplifted. This is clearly observed in the top panels of Figure 6, which shows the monthly mean ERA5 vertical wind vectors averaged along the equator for October 2015 and November 2015. The end result is that in late October, HCN from the peat fires is more readily uplifted to higher altitudes, coinciding with the region of the atmosphere where the IASI sensitivity is greatest, in the upper troposphere between 8 and 12 km. In addition to the increased emissions at the end of October, this contributes to the plume appearing most prominent in the first days of November 2015. The end of October 2015 also coincides with the onset of increased precipitation and an increase in soil moisture. Figure 7a shows that Indonesian deep soil moisture (SWI at the lowest layer corresponding to the characteristic timelength $T=100$, described in Section 2.3.2)




Figure 5. Comparison of the IASI measured HCN averaged total column time series (molecules cm^{-2}) (black line), and HCN TOMCAT model run (T_HCN) using (i) monthly emissions from GFED v4.1 (blue line) and (ii) GFED v4.1 emissions with September values scaled to 25% (orange line). The errors are defined as the standard deviation of the averaged total columns over Indonesia within the region 12°S - 7°N and 90°E - 127°E .

for August - October 2015 was at its lowest level over the last decade, with a minimum at the end of October. The choice of the parameter $T=100$, the lowest layer of SWI, allows us to evaluate the water availability at the plant root depth. This strongly suggests that the GWL during the 2015 El Nino season was at a lower level than in any other year over the last decade. Precipitation for the same period was at the lower end of typical values for the given months as shown in Figure 7b using IMERG data.

3.2.2 Evaluation of 2015 Indonesian HCN emissions using TOMCAT model simulations

In order to estimate the HCN emissions from the 2015 Indonesian peat fires, retrieved HCN total columns from IASI are compared with outputs of a tracer version of the TOMCAT 3D CTM.

Figure 5 compares the averaged total column time series of HCN, calculated over Indonesia in the regional box $[12^{\circ}\text{S}$ - 7°N , 90°E - $127^{\circ}\text{E}]$ for both IASI measurements (black line) and the TOMCAT model outputs smoothed using the IASI averaging kernels (blue line). The box size was defined to cover Indonesia and, at the same time, capture the wildfire emission plumes. The IASI time series exhibits two gaps in November 2015 due to missing measurements. These same gaps also appear in the model time series, as the absence of IASI averaging kernels during this period prevents the smoothing of the model outputs. The T_HCN model run does not compare very well with the IASI measurements, showing a large overestimation of the HCN

amount during the period between September and November 2015. In particular, the large peak simulated for T_HCN during the last week of September is completely absent from the IASI measurements. The left panel of Figure 8 shows the HCN total column distribution on 2 November 2015 produced from T_HCN. The HCN amount produced is globally overestimated compared with the IASI measurements reported in Figure 4, in particular over the latitude band between 20° N and 40° S. The discrepancy observed between the HCN TOMCAT model and IASI measurements is seemingly caused by the HCN emissions used to drive the model, shown in Figure 9, which combine GFED v4.1 (the main component), CMIP6 and CCMI as described in Bruno et al. (2023). These emissions indicate an anomalously large peak of more than 4×10^{11} molecules $\text{cm}^{-2} \text{s}^{-1}$ in September 2015. This value is more than double the October emissions, during which the largest HCN concentrations were observed over Indonesia during the 2015 wildfire season.

In order to seek better agreement between model and observations, we performed a number of model sensitivity runs. These suggested that the most sensitive component for improving the model simulations was the September GFED HCN emissions. For this reason, a new model run was performed ($T_HCN_{0.25Emiss}$) in which the September 2015 emissions were reduced by 75%, based on the analysis performed in Bruno (2024b). The orange line in Figure 5 indicates the averaged total column (smoothed by the IASI HCN averaging kernels) over Indonesia for the $T_HCN_{0.25Emiss}$ model run, showing a substantial reduction in HCN concentration during September and October and reaching a very good agreement with the IASI measurements. However, the background HCN values, e.g. in December, are underestimated in the model, possibly due to neglect of additional sources. The global HCN total column distribution modelled on 2 November 2015, as shown in the bottom right panel of Figure 8, shows a significantly improved agreement with the IASI measured HCN total columns in Figure 4.

The study of CO emissions from the 2015 Indonesian wildfire season by Nechita-Banda et al. (2018) found that GFED is not good at reproducing emissions from large peat fires. GFED estimates monthly fire emissions from satellite-observed burned area and temporally disaggregates them within each month using fire radiative power as a proxy for combustion intensity and timing. For peat-dominated fires, burned area is more sensitive to the initial burning stages and less sensitive to peat burning that occurs underground. Our work supports this conclusion with respect to emissions of HCN from peat fires.

3.2.3 Enhancement ratios and emission factors

The Indonesian peatland fires from 2015 present an opportunity to estimate the HCN wildfire emission factors from satellite data. The fires burned for several months over a limited area, with the prevailing wind conditions resulting in the emitted plumes being predominantly transported over the Indian Ocean. The relatively long lifetimes of HCN and CO, of the order of 2-5 months, allow us to take advantage of satellite measurements over the ocean, where surface emissivity is reasonably homogeneous, and where the polluted air has been uplifted to higher altitudes, corresponding to the maximum sensitivity of IASI to HCN and CO perturbations. Changes in thermal heating over land between day and night cause a change in vertical sensitivity of IASI to HCN and CO. The effect over the oceans is much smaller due to the larger heat capacity of water. As such, in this work we use an average from both IASI daytime and night time measurements to maximise coverage and reduce the random uncertainty of the results.

The emission factor (EF_X) is a quantity defined as the mass of a trace gas (X) emitted per kilogram of biomass burned ($g\ kg^{-1}$ dry matter). In order to derive EF_X , it is first necessary to derive a related parameter called the enhancement ratio (ER) which can be determined directly from satellite measurements. The $ER_{HCN/CO}$ is defined as a ratio of the emitted number of molecules of HCN over the emitted number of molecules of CO (Goode et al., 2000; Whitburn et al., 2017),

$$ER_{HCN/CO} = \frac{[HCN]_{plume} - [HCN]_{background}}{[CO]_{plume} - [CO]_{background}} \quad (2)$$

CO is chosen as the reference gas in this work as it has a similar lifetime to HCN and is emitted from similar sources. The excellent daily coverage of IASI allows a daily estimation of $ER_{HCN/CO}$ to be performed. We have chosen the region $12^\circ S$ - $7^\circ N$, $90^\circ E$ - $127^\circ E$, located downwind of the Indonesian emissions from Sumatra and Kalimantan; only data with a correlation coefficient greater than 0.3 (following threshold methodology of Whitburn et al. (2017)) are included in our analysis. There is no clear trend in the ER across the September 2015 to December 2015 period (Figure 10), although there is some variability with lower ER in early September, where the correlation between HCN and CO measurements is less than 0.4. The phase of fire is a key factor in the emission of pyrogenic species. HCN and CO emissions are generally higher during the smouldering phase of fire (lower FRP) with smouldering peat fire emission factors being approximately 10 times those of flaming savanna fires (Hu et al., 2018). Peat fires generate weakly buoyant smoke plumes (Hu et al., 2019) that generally accumulate close to the ground, although the emissions can migrate great distances via the prevailing winds. The lower ER in early September may be due to the fact that the fires were only recently ignited and high HCN (and CO) levels were closer to the ground. IASI is more sensitive to near surface CO than HCN, so HCN is likely underestimated at that time. Throughout October, the HCN:CO relationship shows correlation coefficients greater than 0.4. By early November, the fire activity ended abruptly, but the correlation remains greater than 0.3 throughout most of November, until the HCN levels drop below the sensitivity of the IASI instrument.

It is estimated that 90% of fire emissions from the late 2015 fire events were dominated by emissions from peat soils (Whitburn et al., 2017), with the remaining 10% from tropical forest. Using a range of ERs derived for differing correlation coefficients, we are able to estimate HCN emission factors, $EF_{HCN/CO}$, from the derived $ER_{HCN/CO}$. Since we are investigating a region close to the emissions, and are not influenced by transport from other regions, the two parameters are related by the equation

$$EF_{HCN} = ER_{HCN/CO} \times \frac{MM_{HCN}}{MM_{CO}} \times EF_{CO}. \quad (3)$$

The quantities MM_{HCN} (27.0253 g/mol) and MM_{CO} (28.01 g/mol) are the molar masses of HCN and CO, respectively. There are a number of datasets of EF_{CO} for peat over Indonesia (Table 1). Stockwell et al. (2016) measured EF_{CO} in situ ($291 \pm 49\ g\ kg^{-1}$), noting that their measurements were solely from smouldering combustion. Huijnen et al. (2016) also measured EF_{CO} in situ (255 ± 39), but noted a few smoke measurements involved occasional contributions from small clumps of ignited dry vegetation, which may account for the smaller value of EF_{CO} . More recently, Yokelson et al. (2022) revisited Kalimantan in

2019, during El Nino conditions, to sample fires burning only peat and measured a 50% larger EF_{CO} compared to the Stockwell et al. (2016) work in a similar region. Both Christian et al. (2003) and Stockwell et al. (2016) took samples from Sumatra and Kalimantan respectively and burned the samples in the lab, both deriving significantly lower EF_{CO} values. The final source
 435 is from a meta-analyses study by Rodriguez Vasquez et al. (2021) who use the data from the in situ and laboratory studies previously listed to derive an updated EF_{CO} based on methodological differences in measurements and weighted to account for different sample sizes.

Table 1. Derived emission factors, EF_{HCN} , in $g\ kg^{-1}$ dry Matter. EF_{HCN} has been calculated via Equation 3 by using a number of EF_{CO} values from the literature. EF_{HCN} is then calculated for all of the ER results presented in Fig. 10 over the region [12°S-7°N, 90°E-127°E].

Author	Study area	Measurement type	EF_{CO} ($g\ kg^{-1}\ DM$)	Calculated in this study			
				$r > 0.5$	$r > 0.6$	$r > 0.7$	$r > 0.8$
Yokelson et al. (2022)	Kalimantan	In situ	315±49	1.19±0.53	1.33±0.52	1.60±0.51	1.71±0.31
Rodriguez Vasquez et al. (2021)	N/A	Meta-analysis	258.68±15.34	0.95±0.34	1.07±0.32	1.30±0.29	1.40±0.12
Stockwell et al. (2016)	Kalimantan	In situ	291±49	1.10±0.50	1.23±0.49	1.48±0.49	1.58±0.31
Huijnen et al. (2016)	Kalimantan	In situ	255±39	0.96±0.43	1.08±0.42	1.30±0.41	1.39±0.25
Stockwell et al. (2014)	Kalimantan	Laboratory	233±72	0.92±0.52	1.02±0.53	1.22±0.55	1.27±0.42
Christian et al. (2003)	Sumatra	Laboratory	210.3	0.76±0.24	0.86±0.21	1.04±0.17	1.04±0.03

Examining the relationship between HCN and CO total column amounts in Figure 10, the correlation coefficient between the two species varies between 0.3 and 0.8 over most of the September to November 2015 period. We use the correlation
 440 coefficient as a filtering metric here to understand whether we see a change in the ER depending on how closely related CO and HCN concentrations are in the same air parcel. A summary of the effect on estimating EF_{HCN} by including only measured data exceeding a chosen correlation coefficient threshold is shown in Table 1. Generally, the lower correlation coefficients are found to be associated with lower ER values. The highest EF_{HCN} values are derived where the correlation coefficient exceeds 0.8, ranging between 1.04 and 1.71 $g\ kg^{-1}$ depending on which value is used for EF_{CO} . Where the correlation coefficient
 445 HCN:CO is greater than 0.5, we see that the EF_{HCN} estimates decrease, ranging between 0.76 and 1.19 $g\ kg^{-1}$. Both ends of the estimates are lower than the (surface) in situ and lab-based estimates of EF_{HCN} . We know that the IASI HCN sensitivity is in the upper troposphere between 8 and 12 km, with very low sensitivity to high concentrations below 4 km which may lead to an underestimate of EF_{HCN} . Field et al. (2016) show that the MLS CO at 215 hPa (~12 km) increased steadily through September with a rapid increase to CO exceeding 400 ppbv at the end of October. The highest correlation HCN:CO in our results is during
 450 this late October period, where we see enhanced vertical uplift of both gases and therefore increased sensitivity to the HCN emissions. Before mid-October, we likely underestimate the HCN amounts as plumes are closer to the surface. This alone cannot explain the large discrepancies between the IASI measurements and the TOMCAT simulations observed in Figure 5, since the TOMCAT outputs have already been smoothed using the IASI averaging kernels to ensure comparable sensitivity.

3.3 Comparison of the 2015 Indonesian fire season with 2019 and 2023

455 According to the ONI, the 2015 El Niño event was one of the strongest in the satellite era. ONI is the primary index used by NOAA to monitor and classify El Niño and La Niña events (NOAA Climate.gov, 2009). Defined as a running 3-month mean sea-surface-temperature anomaly for the Niño 3.4 region (5°N - 5°S , 120° - 170°W), El Niño events are defined as 5 consecutive overlapping 3-month periods above $+0.5^{\circ}$. Figure 11 shows a plot of ONI as a function of time since 1990. It indicates that the recent 2023 burning season in Indonesia, also occurring during an El Niño event, peaked at the third strongest ONI value since
460 1990, behind only the 1997 and 2015 events. The 2019 event was comparatively weak.

In order to compare fire activity in Indonesia during the last three El Niño events, in late 2015, 2019 and 2023, we selected the VNP14IMG fire product across the whole 2012 to 2023 period and looked at the total fire activity across all of the Indonesian islands. Cumulative monthly fire radiative power over Indonesia is plotted in Figure 12, and indicates that even during a notably strong El Nino phase, the fire activity in 2023 was almost eight times less than in 2015. Specifically in 2023 the fire activity
465 was below the seasonal 2012-2023 average across the whole July-December period. For 2019, we observe significant fires in mid-September and a comparable increase in fire power to 2015 between early September and late September. From late September 2019, the fire activity abruptly ends, whereas 2015 maintains a steady increase in fire cumulative power until an abrupt end in fire activity in late October 2015.

We can use the ULIRS HCN to further investigate the emissions from the Indonesian peat fires. Figure 13 represents a 3-day
470 simple moving average total burden HCN (in Gg) and CO (in Tg) for the three El Nino years. The total mass was calculated over the region 12°S - 7°N , 90°E - 127°E following Equation 2 of R'Honi et al. (2013), and shows the highest values are towards the end of October 2015 for both gases. Comparing all three years used in this study, we see the highest HCN values across the September to November period always occur in 2015. The HCN concentrations increase in early September 2015 and are shown to be coincident with an increase in fire activity during the same period (Figure 12), rapidly exceeding 500 Gg by 25
475 September. After this period we see the largest increase in HCN between 22 October and 30 October, to approximately 600 Gg of HCN, peaking at 1056 Gg on 29 October. This increase can also be linked to increasing fire power over that period, although we note that the increase in fire power is also conducive to extra lofting of the elevated HCN to altitudes where the IASI instrument is more sensitive to HCN changes (i.e. the upper troposphere). After 1 November, concentrations of HCN decrease for the remainder of 2015. For 2019, the pattern in early September mirrors that in 2015 with a sudden increase of
480 240 Gg of HCN between 3 September and 26 September. After that, HCN concentrations decrease back to the 300 Gg baseline by 9 October and stay at these lower levels for the rest of 2019. In 2023, we observe no significant enhancements in HCN across the August to December period, relative to the 300 Gg baseline. For CO in 2015 there is a gradual increase in total CO from early September with a peak of 168 Tg in late October, similar to HCN. Throughout November the CO burden decreases and is back at background levels by early December. In 2019, the CO shows a similar variability to HCN, increasing through
485 September but decreasing again in October 2019.

Comparing the circulation patterns across the equator in October 2015 and 2023 in Figure 6, it is clear that the regimes are quite different, with the descending air over Indonesia in 2015 replaced by more neutral conditions in October 2023. These

conditions are less conducive to prolonged dry periods, which would allow peat fires to establish if left unchecked. Over the western Pacific Ocean itself, October 2023 showed much less wide-scale descent of air compared to 2015, suggesting that the Walker circulation did not weaken significantly in late 2023. The circulation patterns across the Eastern Indian Ocean (60°E-110°E) in October 2015 and October 2019 are remarkably consistent. Over Indonesia itself (100°E-120°E) the vertical winds exhibit very different behaviours between 2015 and 2019, with 2019 conditions more conducive to convection and increased rainfall, which would likely suppress fire activity.

Figure 7b indicates that the precipitation levels in August to October for the El Nino years 2015, 2019 and 2023 were very similar. However, Figure 7a demonstrates that the SWI in the lowest soil layer towards the end of 2019 and 2023 was quite typical of values over the last decade, unlike 2015 which was significantly drier. This is consistent with previous work showing that peat moisture, in particular the GWL, is a major factor in determining the ignition and spread of peat fires (Hayasaka et al., 2020).

3.3.1 HCN emission database using the FINNpeatSM method

Peatland fire emissions from GFEDv4s include a simple representation of burn depth based on soil moisture. However, it is not possible to use the dataset to explore soil moisture and burn depth interactions across different years. Therefore limiting our ability to explore the drivers of differences in HCN retrievals between 2015, 2019 and 2023. To address this limitation in GFEDv4s, we examine fire detections (from VIIRS fire hotspots) and SMAP (O'Neill et al., 2021) over Indonesia peatlands. Using the FINNpeatSM method (Kiely et al., 2019), we can combine these datasets to estimate the burn depth of fires in 2015, 2019 and 2023.

Daily VIIRS temperature hotspots are only used if there is a medium or high confidence in the retrieval (Wiedinmyer et al., 2023) and if the fire occurred on peatland in Indonesia. For each retained fire detection, the nearest-neighbour daily mean soil moisture pixel (from SMAP) is determined. The associated soil moisture from an individual fire detection can be used to estimate the burn depth, and emissions, of the fire. The emissions in FINNpeatSM are linearly related to burn depth, so that the deeper a fire burns the higher the emissions.

First, we consider daily mean soil moisture from SMAP across all Indonesian, Kalimantan, and Sumatra peatlands for 2015, 2019 and 2023 (Figure 14). We consider Kalimantan and Sumatra peatlands in addition to all Indonesian peatlands since these regions dominate fire emissions. In 2015 62% of total emissions were from Kalimantan and 33% were from Sumatra (Kiely et al., 2019). Soil moisture is shown for 1 June to 31 December, with the dry season (1 August to 31 October) shaded in grey, since the dry season is when fires are most likely to occur. Across Indonesia and both Kalimantan and Sumatra soil moisture in 2015, 2019 and 2023 were similar from June-July ($0.58\text{--}0.61\text{ m}^3\text{ m}^{-3}$). However, from July to August, soil moisture across all regions in 2015 and 2019 begins to decrease rapidly (from $0.52\text{--}0.55\text{ m}^3\text{ m}^{-3}$ to $0.46\text{--}0.48\text{ m}^3\text{ m}^{-3}$), while in 2023 soil moisture remains high ($0.59\text{ to }0.54\text{ m}^3\text{ m}^{-3}$). From August to mid-September soil moisture continues to decrease across Indonesia, driven by reductions in soil moisture in Kalimantan in 2015 and 2019 (Indonesia from $0.46\text{--}0.48$ to $0.36\text{--}0.39\text{ m}^3\text{ m}^{-3}$ and Kalimantan from $0.39\text{--}0.44$ to $0.27\text{--}0.29\text{ m}^3\text{ m}^{-3}$). In contrast, over the same period in 2023, soil moisture remains high across Indonesia (decreasing from 0.54 to $0.50\text{ m}^3\text{ m}^{-3}$), driven by high soil moisture in Sumatra peatlands. From mid-

September to mid-October soil moisture continues to decrease in 2015 (from 0.39 to 0.33 m³ m⁻³), while in 2019 there is an abrupt increase in soil moisture (from 0.36 to 0.47 m³ m⁻³), closer to the 2023 values (from 0.50 to 0.54 m³ m⁻³). Overall, soil moisture is lowest throughout the dry season in 2015 (1 August to 31 October mean of 0.40 m³ m⁻³), while in 2019 soil
525 moisture is very low at the start of the dry season but then increases in the late dry season (1 August to 31 October mean of 0.43 m³ m⁻³). In contrast, soil moisture in 2023 is the highest of the 3 years (1 August to 31 October mean of 0.52 m³ m⁻³), in agreement with what was observed in SWI data (Figure 7a).

The collocated daily mean soil moisture and daily total number of fire hotspots (from VIIRS) for the same time period (1 June to 31 December 2015, 2019 and 2023) is shown in Figure 15. This clearly indicates that 2015 had the highest number
530 of fire hotspots (>40 collocated pixels with 1,000-10,000 fire hotspots) occurring in areas where soil moisture was low (0.15-0.35 m³ m⁻³). However, in 2019, there are fewer fire pixels with a high number of hotspots (24 pixels with 1,000-10,000 fire hotspots) and they generally had higher soil moisture (0.25-0.31 m³ m⁻³). In 2023, there were both fewer pixels with a high number of fire hotspots (peaking at 100-1,000 fire hotspots), and the soil moisture where fire hotspots occurred was higher (0.21-0.4 m³ m⁻³).

The soil moisture at each fire location can be used to calculate peat burn depth of the fire, using the method developed by Kiely et al. (2019). Figure 16 indicates the burn depth of fires, and the date they occurred. As shown in Figure 15, 2015 has the highest number of fires (up to 25,000 fires) that occur in low soil moisture areas (0.3 m). Figure 16 indicates that these fires burn deep into the peat below (0.25-0.3 m) and occur between September and mid-October 2015 (in line with Figure 14). In contrast, in 2019, there are fewer fires (0-15000), the burn depth is shallower (0-0.25 m), and the fires that burn the deepest
540 occur earlier in the dry season (mid-September) when soil moisture is higher than 2015 (Figure 14). In 2023, far fewer fires (0-5000 fires) burn deep into the peat below (0.05-0.25 m) and deep burning occurs in mid-September to early-October, in line with increased soil moisture in 2023 (Figure 14).

The accumulated daily burn depth of all Indonesian peatland fires from 1 June to 31 December is shown in Figure 17. This plot accounts for both the number of fire hotspots and the burn depth of individual fire hotspots. At the start of the dry season
545 (August to mid-September) the accumulated burn depth in 2015 and 2019 is similar (500-1000 m), compared with <100 m in 2023. This is likely because soil moisture at the start of the dry season (August to mid-September) is similar in both years, and both years have a similar number of fire hotspots at this time. However, from mid-September, 2015 and 2019 deviate. Accumulated burn depth remains high in October 2015, whereas it substantially decreases in 2019. This is likely driven by the large decrease in both fire hotspots and peatland soil moisture in 2019, compared with 2015 where the number of fire hotspots
550 remains high and the peatland soil moisture remains low.

As illustrated in Figure 18, the time series of accumulated daily burn depth, along with the averaged total column of HCN and CO, exhibit similar temporal patterns, a progressive increase from August through October, reaching a peak in late October, and subsequently declining sharply in November.

3.4 Temporal relationship between burn depth and HCN and CO total column during 2015 wildfires

555 The temporal relationship between two time series can be investigated using the cross-correlation function, which quantifies the linear association between two variables as a function of temporal lag (Storch and Zwiers, 1999). For two stationary time series x_t and y_t , the cross-correlation at a given lag k is defined as the normalized covariance between x_t and y_{t-k} :

$$\rho_{xy}(k) = \frac{\text{COV}(x_t, y_{t-k})}{\sigma_x \sigma_y}, \quad (4)$$

where σ_x and σ_y denote the standard deviations of the two series. In practice, the covariance is estimated from the sample values as

$$\rho_{xy}(k) = \frac{\sum_{t=1}^{N-k} (x_t - \bar{x})(y_{t-k} - \bar{y})}{(N-k)\sigma_x \sigma_y}, \quad (5)$$

where $\bar{x}$ and $\bar{y}$ are the sample means and N is the length of the time series. The resulting function $\rho_{xy}(k)$ ranges between -1 and 1 and describes the strength of the linear relationship between the two variables at different lags. Positive lags indicate that variations in x lead those in y , whereas negative lags indicate that variations in y precede those in x .

565 A cross-correlation analysis between the accumulated daily burn depth and the total column concentrations of CO and HCN can provide insights into the temporal dynamics of plume transport. As shown in Figure 19, both HCN and CO exhibit maximum positive correlations with the burn depth at non-zero lag times, indicating that the fire activity at the surface anticipates the appearance of enhanced trace gas signals at the respective altitudes of high sensitivity of the IASI instrument. Specifically, the cross-correlation peaks at a lag of approximately -10 days for both species, suggesting that 10 days is the typical timescale required for emissions from peatland fires to be transported and mixed into the mid- to upper-tropospheric layers. The cross-correlation function (XCF) for HCN (black line) and CO (orange line) displays a broadly similar temporal structure, with both curves rising steadily from negative lag values, peaking near -10 days, and subsequently declining into negative correlation at higher lag values. The slightly larger peak for CO may reflect differences in vertical transport dynamics or sensitivity profiles, given that CO exhibits dual sensitivity in both the lower and upper troposphere, whereas HCN is primarily sensitive to the mid-troposphere (~ 10 – 12 km), as described in Section 2.1.1. These results support the interpretation that the observed trace gas enhancements are closely linked to fire activity and that the lag time to maximum correlation can serve as an estimate of the average vertical and horizontal transport time of the emission plume.

4 Conclusions

This study has provided a comprehensive assessment of Indonesian peatland fire emissions during recent El Niño events, with a particular focus on HCN as a tracer of smouldering combustion during peat fires. By integrating IASI satellite retrievals with TOMCAT model simulations, precipitation and soil moisture observations, and fire activity datasets, we have quantified both the magnitude and variability of HCN emissions and identified the key drivers underlying interannual differences. The results

provide strong evidence that the magnitude of trace gas emissions from Indonesian peatlands is not governed solely by the strength of El Niño but instead arises from the interaction between large-scale ocean–atmosphere dynamics and local hydro-
585 logical conditions. The 2015 El Niño event, one of the strongest in recent decades, produced exceptionally large atmospheric concentration of HCN and CO, reflecting a combination of extremely low soil moisture, depressed groundwater levels, and deep burn depths in peatlands. The resulting emissions were unprecedented in the observational record, with IASI retrievals showing marked enhancements that were sustained through October and November.

In contrast, the 2019 and 2023 El Niño events, despite also being associated with positive Oceanic Niño Index (ONI)
590 anomalies, resulted in markedly lower HCN burdens related to higher ground water content and atmospheric dynamics that make intense peat fires less favourable. The 2023 event, in particular, occurred under atmospheric circulation regimes that did not sustain prolonged dryness, resulting in fire activity that was nearly an order of magnitude almost eight times weaker than in 2015.

Using IASI total column measurements we also derive new satellite-based emission factors (EFs) for HCN enhancement
595 ratios with CO during the 2015 dry season. These values provide independent, observation-based constraints that can improve the representation of peat fire emissions in models and inventories.

Comparisons between IASI-derived HCN total columns and TOMCAT model simulations revealed that the standard GFEDv4-
based emission inventory substantially overestimates HCN emissions from peat fires during the 2015 dry season, particularly in September. A revised model run, applying a 75% reduction to September emissions, produced much better agreement with
600 satellite observations. This finding is consistent with the work of Nechita-Banda et al. (2018), who showed that GFED performs poorly in reproducing emissions from large peat fires because it relies on satellite-observed burned area and uses fire radiative power to temporally disaggregate emissions. In peat-dominated environments, the burned area is more sensitive to the initial stages of combustion and fails to capture prolonged subsurface smouldering. Together, these results suggest that current fire emission inventories misrepresent peat fire dynamics and are not well suited to capturing the contribution of underground
605 smouldering processes to trace gas release. The analysis also confirms that the new FINNpeatSM approach, which explicitly links burn depth to soil moisture, offers a more accurate description of the dynamics of peat fire emissions.

Hydrological conditions emerged as a dominant control on fire intensity and emissions across all three El Niño events. Soil moisture and groundwater level determine not only the susceptibility of peatlands to ignition but also the persistence of smouldering fires and the total mass of peat carbon released to the atmosphere. When soils are dried, the peat layer can ignite
610 and burn deeply, releasing large quantities of HCN and CO that are transported into the mid- to upper- troposphere within about 10 days. Conversely, higher soil water content suppresses combustion depth and limits the emissions. The strong cross-correlation between accumulated burn depth and satellite-observed HCN and CO concentrations further supports the close coupling between hydrology, fire behavior, and wildfire emissions.

The comparison of 2015, 2019, and 2023 further highlights that fire intensity and trace gas burdens are not determined
615 solely by the strength of El Niño events, but rather emerge from the importance of the local drought intensity, soil moisture deficits, and atmospheric circulation patterns. This underscores the critical role of groundwater and soil moisture in regulating peatland vulnerability to ignition, smouldering persistence, and total emissions. Our results therefore emphasize the need for

next-generation emission inventories that explicitly account for soil moisture and groundwater dynamics to better reproduce the peatland fire dynamics. Integrating satellite-derived products of atmospheric composition (e.g. IASI) and soil hydrology (e.g., SMAP, ASCAT, SWI) offers a valuable constraint for emissions and predictions of future fire behaviour under changing climate conditions. Such integrated approaches can reduce uncertainties in global fire emission estimates, improve the representation of smouldering combustion in chemical transport models, and enhance the predictive capacity for El Niño–induced fires during the dry seasons across Indonesia peat-dominated ecosystems. These advances are essential for understanding the contribution of peatland fires to regional air quality, global atmospheric composition, and the carbon–climate feedback system.

Data availability. The VNP14IMG product is available from the Level-1 and Atmosphere Archive & Distribution System Distributed Active Archive Center (LAADS DAAC, <https://ladsweb.modaps.eosdis.nasa.gov/archive/allData/5000/VNP14IMG/>) which archives a number of data archives on Earth atmosphere products for NASA, NOAA and European Space Administration missions. The IMERG data are available from Global Precipitation Measurement (GPM) archive (<https://gpm.nasa.gov/data/directory>) which provide the Product Version 07 (V07). SMAP Level-4 data are distributed by the National Snow and Ice Data Center at <http://nsidc.org/data/smap/>. The SWI data are produced by Copernicus - Land Monitoring Service and available on the platform (https://land.copernicus.eu/en/products/soil-moisture?tab=soil_water_index). GFED4 data are available via NASA's Earthdata platform (<https://www.earthdata.nasa.gov/data/catalog/ornl-cloud-fire-emissions-v4-r1-1293-4.1>). IASI CO total columns retrieved using ULIRS are available on the CEDA archive (<https://catalogue.ceda.ac.uk/uuid/4b31d47716604b9f84714fab39ce9>) the HCN and CO gridded data are available at <https://doi.org/10.5281/zenodo.20584609> (Bruno et al., 2026a). The TOMCAT model data are available at <https://doi.org/10.5281/zenodo.18848199> (Bruno et al., 2026b) and at <https://doi.org/10.5281/zenodo.18848214> (Bruno et al., 2026c).

The ERA5 datasets are available at Copernicus Climate Data Store, “ERA5 monthly data on pressure levels from 1940 to present” (<https://cds.climate.copernicus.eu/datasets/reanalysis-era5-pressure-levels?download>) The NDACC CO and HCN FTIR measurements at Mado are publicly available from the NDACC database (<https://www-air.larc.nasa.gov/missions/ndacc/data.html#>).

Author contributions. This work formed part of the PhD studies of AGB, under the supervision of JJH, DM, and MPC. JJH designed the overall study framework during the PhD, while AGB and DM developed the specific implementation presented in this paper, with important contributions from the other co-authors. Most of the data analysis was performed by AGB. DM performed the HCN and CO retrievals, and carried out the work on emission factors. AG carried out the FINNpeatSM work. Model runs were performed by AGB under the supervision of MPC. The TOMCAT model is maintained and updated by the MPC research group at the University of Leeds. AGB led the writing of the manuscript and prepared the first draft, with contributions from all co-authors in subsequent revisions.

Competing interests. The authors declare that they have no conflicts of interest.

Financial support. Antonio G. Bruno was funded by a Natural Environment Research Council (NERC) studentship at the University of Leicester, awarded through the Central England NERC Training Alliance (CENTA) (grant no. NE/S007350/1) and by a CASE award from the National Centre for Earth Observation (NCEO). This study was also funded as part of the UK Research and Innovation Natural Environment
650 Research Council's support of the NCEO, grant number NE/R016518/1.

Acknowledgements. The data analysis was performed at the ALICE HPC facility at the University of Leicester. The retrievals were performed on JASMIN, the UK's collaborative data analysis environment (<https://www.jasmin.ac.uk>) The model simulations were performed on the University of Leeds Advanced Research Computing (ARC) high-performance computing (HPC) machines. We thank Wuhu Feng (NCAS Leeds), Chris Wilson (NCEO Leeds) and Richard Pope (NCEO Leeds) for help with TOMCAT. The Reunion Maïdo FTIR data used in
655 this publication were obtained from Corinne Vigouroux, Bavo Langerock and Martine De Mazière (Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium) as part of the Network for the Detection of Atmospheric Composition Change (NDACC) and are available through the NDACC website <https://www.ndacc.org> (last access: March 2026). The measurements at Maïdo have been also supported by the Université de La Réunion and CNRS (LACy-UMR8105 and UMS3365).

References

- 660 Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmospheric Chemistry and Physics*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- Anthony G. Bamston, M. C. and Goldenberg, S. B.: Documentation of a highly ENSO-related sst region in the equatorial pacific: Research note, *Atmosphere-Ocean*, 35, 367–383, <https://doi.org/10.1080/07055900.1997.9649597>, 1997.
- 665 Baray, J.-L., Courcoux, Y., Keckhut, P., Portafaix, T., Tulet, P., Cammas, J.-P., Hauchecorne, A., Godin Beekmann, S., De Mazière, M., Hermans, C., Desmet, F., Sellegri, K., Colomb, A., Ramonet, M., Sciare, J., Vuillemin, C., Hoareau, C., Dionisi, D., Duflot, V., Vérémes, H., Porteneuve, J., Gabarrot, F., Gaudot, T., Metzger, J.-M., Payen, G., Leclair de Bellevue, J., Barthe, C., Posny, F., Ricaud, P., Abchiche, A., and Delmas, R.: Maïdo observatory: a new high-altitude station facility at Reunion Island (21° S, 55° E) for long-term atmospheric remote sensing and in situ measurements, *Atmospheric Measurement Techniques*, 6, 2865–2877, [https://doi.org/10.5194/amt-6-2865-](https://doi.org/10.5194/amt-6-2865-2013)
- 670 2013, 2013.
- Bauer-Marschallinger, B., Paulik, C., Hochstöger, S., Mistelbauer, T., Modanesi, S., Ciabatta, L., Massari, C., Brocca, L., and Wagner, W.: Soil Moisture from Fusion of Scatterometer and SAR: Closing the Scale Gap with Temporal Filtering, *Remote Sensing*, 10, <https://doi.org/10.3390/rs10071030>, 2018.
- Bernath, P., Crouse, J., Hughes, R., and Boone, C.: The Atmospheric Chemistry Experiment Fourier transform spectrometer (ACE-FTS) version 4.1 retrievals: Trends and seasonal distributions, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 259, 107 409, <https://doi.org/10.1016/j.jqsrt.2020.107409>, 2021.
- 675 Berrisford, P., Dee, D., Poli, P., Brugge, R., Fielding, M., Fuentes, M., Kållberg, P., Kobayashi, S., Uppala, S., and Simmons, A.: The ERA-Interim archive Version 2.0, Shinfield Park, Reading, 2011.
- Boone, C., Bernath, P., Cok, D., Jones, S., and Steffen, J.: Version 4 retrievals for the atmospheric chemistry experiment Fourier transform spectrometer (ACE-FTS) and imagers, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 247, 106 939, <https://doi.org/10.1016/j.jqsrt.2020.106939>, 2020.
- 680 Bruno, A. G.: Investigating The Trace Gas Emissions Of Biomass Burning In The Earth System, Thesis, University of Leicester, <https://doi.org/10.25392/leicester.data.25680108.v1>, 2024a.
- Bruno, A. G.: Investigating The Trace Gas Emissions Of Biomass Burning In The Earth System, Ph.D. thesis, <https://doi.org/10.25392/LEICESTER.DATA.25680108>, artwork Size: 10687680 Bytes, 2024b.
- 685 Bruno, A. G., Harrison, J. J., Moore, D. P., Chipperfield, M. P., and Pope, R. P.: Satellite observations and modelling of hydrogen cyanide in the Earth’s atmosphere, *Il Nuovo Cimento C*, 6, <https://doi.org/10.1393/ncc/i/2022-22184-6>, 2022.
- Bruno, A. G., Harrison, J. J., Chipperfield, M. P., Moore, D. P., Pope, R. J., Wilson, C., Mahieu, E., and Notholt, J.: Atmospheric distribution of HCN from satellite observations and 3-D model simulations, *Atmospheric Chemistry and Physics*, 23, 4849–4861, <https://doi.org/10.5194/acp-23-4849-2023>, 2023.
- 690 Bruno, A. G., Moore, D. P., and Harrison, J. J.: Hydrological drivers of hydrogen cyanide wildfire emissions from Indonesian peat fires during the 2015, 2019, and 2023 El Niño events - IASI gridded data, <https://doi.org/10.5281/zenodo.20584609>, 2026a.
- Bruno, A. G., Moore, D. P., Harrison, J. J., Graham, A., and Chipperfield, M.: Hydrological drivers of hydrogen cyanide wildfire emissions from Indonesian peat fires during the 2015, 2019, and 2023 El Niño events - TOMCAT data part 1, <https://doi.org/10.5281/zenodo.18848199>, 2026b.
- 695

- Bruno, A. G., Moore, D. P., Harrison, J. J., Graham, A., and Chipperfield, M.: Hydrological drivers of hydrogen cyanide wildfire emissions from Indonesian peat fires during the 2015, 2019, and 2023 El Niño events - TOMCAT data part 2, <https://doi.org/10.5281/zenodo.18848214>, 2026c.
- 700 Callewaert, S., Brioude, J., Langerock, B., Dufлот, V., Fonteyn, D., Müller, J.-F., Metzger, J.-M., Hermans, C., Kumps, N., Ramonet, M., Lopez, M., Mahieu, E., and De Mazière, M.: Analysis of CO₂, CH₄, and CO surface and column concentrations observed at Réunion Island by assessing WRF-Chem simulations, *Atmospheric Chemistry and Physics*, 22, 7763–7792, <https://doi.org/10.5194/acp-22-7763-2022>, 2022.
- Chipperfield, M. P.: New version of the TOMCAT/SLIMCAT off-line chemical transport model: Intercomparison of stratospheric tracer experiments, *Quarterly Journal of the Royal Meteorological Society*, 132, 1179–1203, <https://doi.org/10.1256/qj.05.51>, 2006.
- 705 Chipperfield, M. P., Cariolle, D., Simon, P., Ramaroson, R., and Lary, D. J.: A three-dimensional modeling study of trace species in the Arctic lower stratosphere during winter 1989–1990, *Journal of Geophysical Research: Atmospheres*, 98, 7199–7218, <https://doi.org/10.1029/92JD02977>, 1993.
- Chipperfield, M. P., Dhomse, S., Hossaini, R., Feng, W., Santee, M. L., Weber, M., Burrows, J. P., Wild, J. D., Loyola, D., and Coldewey-Egbers, M.: On the Cause of Recent Variations in Lower Stratospheric Ozone, *Geophysical Research Letters*, 45, 5718–5726, <https://doi.org/10.1029/2018GL078071>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2018GL078071>, 2018.
- 710 Cho, K., Negrón-Juárez, R., Colliander, A., Cosio, E. G., Salinas, N., De Araujo, A., Chambers, J. Q., and Wang, J.: Calibration of the SMAP Soil Moisture Retrieval Algorithm to Reduce Bias Over the Amazon Rainforest, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 17, 8724–8736, <https://doi.org/10.1109/JSTARS.2024.3388914>, 2024.
- Christian, T. J., Kleiss, B., Yokelson, R. J., Holzinger, R., Crutzen, P., Hao, W. M., Saharjo, B., and Ward, D. E.: Comprehensive laboratory measurements of biomass-burning emissions: 1. Emissions from Indonesian, African, and other fuels, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- 715 Clarisse, L., R'Honi, Y., Coheur, P.-F., Hurtmans, D., and Clerbaux, C.: Thermal infrared nadir observations of 24 atmospheric gases, *Geophysical Research Letters*, 38, <https://doi.org/10.1029/2011GL047271>, 2011.
- Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., 720 Wespes, C., and Coheur, P.-F.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmospheric Chemistry and Physics*, 9, 6041–6054, <https://doi.org/10.5194/acp-9-6041-2009>, 2009.
- Coheur, P.-F., Clarisse, L., Turquety, S., Hurtmans, D., and Clerbaux, C.: IASI measurements of reactive trace species in biomass burning plumes, *Atmospheric Chemistry and Physics*, 9, 5655–5667, <https://doi.org/10.5194/acp-9-5655-2009>, 2009.
- Dargie, G. C., Lewis, S. L., Lawson, I. T., Mitchard, E. T. A., Page, S. E., Bocko, Y. E., and Ifo, S. A.: Age, extent and carbon storage of the 725 central Congo Basin peatland complex, *Nature*, 542, 86–90, 2017.
- De Longueville, H., Clarisse, L., Whitburn, S., Franco, B., Bauduin, S., Clerbaux, C., Camy-Peyret, C., and Coheur, P.-F.: Identification of Short and Long-Lived Atmospheric Trace Gases From IASI Space Observations, *Geophysical Research Letters*, 48, e2020GL091742, <https://doi.org/10.1029/2020GL091742>, e2020GL091742 2020GL091742, 2021.
- De Mazière, M., Thompson, A. M., Kurylo, M. J., Wild, J. D., Bernhard, G., Blumenstock, T., Braathen, G. O., Hannigan, J. W., Lambert, 730 J.-C., Leblanc, T., McGee, T. J., Nedoluha, G., Petropavlovskikh, I., Seckmeyer, G., Simon, P. C., Steinbrecht, W., and Strahan, S. E.: The Network for the Detection of Atmospheric Composition Change (NDACC): history, status and perspectives, *Atmospheric Chemistry and Physics*, 18, 4935–4964, <https://doi.org/10.5194/acp-18-4935-2018>, 2018.

- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/https://doi.org/10.1002/qj.828>, 2011.
- Deeter, M. N.: Calculation and Application of MOPITT Averaging Kernels, <https://api.semanticscholar.org/CorpusID:111388701>, 2002.
- Dudhia, A.: The Reference Forward Model (RFM), *Journal of Quantitative Spectroscopy and Radiative Transfer*, 186, 243–253, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2016.06.018>, satellite Remote Sensing and Spectroscopy: Joint ACE-Odin Meeting, October 2015, 2017.
- Duflot, V., Dils, B., Baray, J. L., De Mazière, M., Attié, J. L., Vanhaelewyn, G., Senten, C., Vigouroux, C., Clain, G., and Delmas, R.: Analysis of the origin of the distribution of CO in the subtropical southern Indian Ocean in 2007, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/https://doi.org/10.1029/2010JD013994>, 2010.
- Duflot, V., Hurtmans, D., Clarisse, L., R'honi, Y., Vigouroux, C., De Mazière, M., Mahieu, E., Servais, C., Clerbaux, C., and Coheur, P.-F.: Measurements of hydrogen cyanide (HCN) and acetylene (C₂H₂) from the Infrared Atmospheric Sounding Interferometer (IASI), *Atmospheric Measurement Techniques*, 6, 917–925, <https://doi.org/10.5194/amt-6-917-2013>, 2013.
- Duflot, V., Wespes, C., Clarisse, L., Hurtmans, D., Ngadi, Y., Jones, N., Paton-Walsh, C., Hadji-Lazaro, J., Vigouroux, C., De Mazière, M., Metzger, J.-M., Mahieu, E., Servais, C., Hase, F., Schneider, M., Clerbaux, C., and Coheur, P.-F.: Acetylene (C₂H₂) and hydrogen cyanide (HCN) from IASI satellite observations: global distributions, validation, and comparison with model, *Atmospheric Chemistry and Physics*, 15, 10 509–10 527, <https://doi.org/10.5194/acp-15-10509-2015>, 2015.
- Entekhabi, D., Yueh, S., and De Lannoy, G.: SMAP handbook, 2014.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Fang, B., Lakshmi, V., and Zhang, R.: Validation of downscaled 1-km SMOS and SMAP soil moisture data in 2010–2021, *Vadose Zone Journal*, 23, e20 305, <https://doi.org/10.1002/vzj2.20305>, 2024.
- Field, R. D., van der Werf, G. R., Fanin, T., Fetzer, E. J., Fuller, R., Jethva, H., Levy, R., Livesey, N. J., Luo, M., Torres, O., and Worden, H. M.: Indonesian fire activity and smoke pollution in 2015 show persistent nonlinear sensitivity to El Niño-induced drought, *Proceedings of the National Academy of Sciences*, 113, 9204–9209, <https://doi.org/10.1073/pnas.1524888113>, 2016.
- Goode, J. G., Yokelson, R. J., Ward, D. E., Susott, R. A., Babbitt, R. E., Davies, M. A., and Hao, W. M.: Measurements of excess O₃, CO₂, CO, CH₄, C₂H₄, C₂H₂, HCN, NO, NH₃, HCOOH, CH₃COOH, HCHO, and CH₃OH in 1997 Alaskan biomass burning plumes by airborne Fourier transform infrared spectroscopy (AFTIR), *Journal of Geophysical Research: Atmospheres*, 105, 22 147–22 166, <https://doi.org/10.1029/2000JD900287>, 2000.
- Graham, A. M., Spracklen, D. V., McQuaid, J. B., Smith, T. E. L., Nurrahmawati, H., Ayona, D., Mulawarman, H., Adam, C., Papargyropoulou, E., Rigby, R., Padfield, R., and Choiruzzad, S.: Updated Smoke Exposure Estimate for Indonesian Peatland Fires Using a Network of Low-Cost PM_{2.5} Sensors and a Regional Air Quality Model, *GeoHealth*, 8, e2024GH001 125, <https://doi.org/10.1029/2024GH001125>, 2024.

- 770 Hayasaka, H., Usup, A., and Naito, D.: New Approach Evaluating Peatland Fires in Indonesian Factors, *Remote Sensing*, 12, <https://doi.org/10.3390/rs12122055>, 2020.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J.,
775 Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/https://doi.org/10.1002/qj.3803>, 2020.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 monthly averaged data on pressure levels from 1940 to present, <https://doi.org/10.24381/cds.6860a573>, accessed on 12-03-2026, 2023.
- 780 Hilton, F., Armante, R., August, T., Barnet, C., Bouchard, A., Camy-Peyret, C., Capelle, V., Clarisse, L., Clerbaux, C., Coheur, P.-F., Collard, A., Crevoisier, C., Dufour, G., Edwards, D., Faijan, F., Fourrié, N., Gambacorta, A., Goldberg, M., Guidard, V., Hurtmans, D., Illingworth, S., Jacquinet-Husson, N., Kerzenmacher, T., Klaes, D., Lavanant, L., Masiello, G., Matricardi, M., McNally, A., Newman, S., Pavelin, E., Payan, S., Péquignot, E., Peyridieu, S., Phulpin, T., Remedios, J., Schlüssel, P., Serio, C., Strow, L., Stubenrauch, C., Taylor, J., Tobin,
785 D., Wolf, W., and Zhou, D.: Hyperspectral Earth Observation from IASI: Five Years of Accomplishments, *Bulletin of the American Meteorological Society*, 93, 347 – 370, <https://doi.org/https://doi.org/10.1175/BAMS-D-11-00027.1>, 2012.
- Hooijer, A. and Vernimmen, R.: Peatland maps for Indonesia. Including accuracy assessment and recommendations for improvement, elevation mapping and evaluation of future flood risk. Quick Assessment and Nationwide Screening (QANS) of Peat and Lowland Resources and Action Planning for the Implementation of a National Lowland Strategy - PVW3A10002., Agentschap NL 6201068 QANS Lowland
790 Development, for Government of Indonesia and Partners for Water (Netherlands), 2013.
- Hu, Y., Fernandez-Anez, N., Smith, T. E. L., and Rein, G.: Review of emissions from smouldering peat fires and their contribution to regional haze episodes, *International Journal of Wildland Fire*, 27, 293–312, <https://doi.org/10.1071/WF17084>, 2018.
- Hu, Y., Christensen, E., Restuccia, F., and Rein, G.: Transient gas and particle emissions from smouldering combustion of peat, *Proceedings of the Combustion Institute*, 37, 4035–4042, <https://doi.org/10.1016/j.proci.2018.06.008>, 2019.
- 795 Huang, B., Thorne, P. W., Banzon, V. F., Boyer, T., Chepurin, G., Lawrimore, J. H., Menne, M. J., Smith, T. M., Vose, R. S., and Zhang, H.-M.: Extended Reconstructed Sea Surface Temperature, Version 5 (ERSSTv5): Upgrades, Validations, and Intercomparisons, *Journal of Climate*, 30, 8179 – 8205, <https://doi.org/10.1175/JCLI-D-16-0836.1>, 2017.
- Huffman, G., Stocker, E., Bolvin, D., Nelkin, E., and Tan, J.: GPM IMERG Final Precipitation L3 1 day 0.1 degree x 0.1 degree V07, <https://doi.org/10.5065/7DE2-M746>, artwork Size: 0.000 bytes Pages: 0.000 bytes, 2024.
- 800 Huijnen, V., Wooster, M. J., Kaiser, J. W., Gaveau, D. L., Flemming, J., Parrington, M., Inness, A., Murdiyarso, D., Main, B., and van Weele, M.: Fire carbon emissions over maritime southeast Asia in 2015 largest since 1997, *Scientific reports*, 6, 26 886, 2016.
- Illingworth, S. M., Remedios, J. J., Boesch, H., Ho, S.-P., Edwards, D. P., Palmer, P. I., and Gonzi, S.: A comparison of OEM CO retrievals from the IASI and MOPITT instruments, *Atmospheric Measurement Techniques*, 4, 775–793, <https://doi.org/10.5194/amt-4-775-2011>, 2011a.
- 805 Illingworth, S. M., Remedios, J. J., Boesch, H., Moore, D. P., Sembhi, H., Dudhia, A., and Walker, J. C.: ULIRS, an optimal estimation retrieval scheme for carbon monoxide using IASI spectral radiances: sensitivity analysis, error budget and simulations, *Atmospheric Measurement Techniques*, 4, 269–288, <https://doi.org/10.5194/amt-4-269-2011>, 2011b.

- Kerzenmacher, T., Dils, B., Kumps, N., Blumenstock, T., Clerbaux, C., Coheur, P.-F., Demoulin, P., García, O., George, M., Griffith, D. W. T., Hase, F., Hadji-Lazaro, J., Hurtmans, D., Jones, N., Mahieu, E., Notholt, J., Paton-Walsh, C., Raffalski, U., Ridder, T., Schneider, M., Servais, C., and De Mazière, M.: Validation of IASI FORLI carbon monoxide retrievals using FTIR data from NDACC, *Atmospheric Measurement Techniques*, 5, 2751–2761, <https://doi.org/10.5194/amt-5-2751-2012>, 2012.
- Kiely, L., Spracklen, D. V., Wiedinmyer, C., Conibear, L., Reddington, C. L., Archer-Nicholls, S., Lowe, D., Arnold, S. R., Knote, C., Khan, M. F., Latif, M. T., Kuwata, M., Budisulistiorini, S. H., and Syaifina, L.: New estimate of particulate emissions from Indonesian peat fires in 2015, *Atmospheric Chemistry and Physics*, 19, 11 105–11 121, <https://doi.org/10.5194/acp-19-11105-2019>, 2019.
- Kleinböhl, A., Toon, G. C., Sen, B., Blavier, J.-F. L., Weisenstein, D. K., Strekowski, R. S., Nicovich, J. M., Wine, P. H., and Wennberg, P. O.: On the stratospheric chemistry of hydrogen cyanide, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2006GL026015>, 2006.
- Krol, M., de Bruine, M., Killaars, L., Ouwersloot, H., Pozzer, A., Yin, Y., Chevallier, F., Bousquet, P., Patra, P., Belikov, D., Maksyutov, S., Dhomse, S., Feng, W., and Chipperfield, M. P.: Age of air as a diagnostic for transport timescales in global models, *Geoscientific Model Development*, 11, 3109–3130, <https://doi.org/10.5194/gmd-11-3109-2018>, 2018.
- Li, Q., Jacob, D. J., Bey, I., Yantosca, R. M., Zhao, Y., Kondo, Y., and Notholt, J.: Atmospheric hydrogen cyanide (HCN): Biomass burning source, ocean sink?, *Geophysical Research Letters*, 27, 357–360, <https://doi.org/10.1029/1999GL010935>, 2000.
- Lohberger, S., Stängel, M., Atwood, E. C., and Siegert, F.: Spatial evaluation of Indonesia’s 2015 fire-affected area and estimated carbon emissions using Sentinel-1, *Global Change Biology*, 24, 644–654, <https://doi.org/10.1111/gcb.13841>, 2018.
- Ma, H., Zeng, J., Zhang, X., Peng, J., Li, X., Fu, P., Cosh, M. H., Letu, H., Wang, S., Chen, N., and Wigneron, J.-P.: Surface soil moisture from combined active and passive microwave observations: Integrating ASCAT and SMAP observations based on machine learning approaches, *Remote Sensing of Environment*, 308, 114 197, <https://doi.org/10.1016/j.rse.2024.114197>, 2024.
- Ma, J., Remaud, M., Peylin, P., Patra, P., Niwa, Y., Rodenbeck, C., Cartwright, M., Harrison, J. J., Chipperfield, M. P., Pope, R. J., Wilson, C., Belviso, S., Montzka, S. A., Vimont, I., Moore, F., Atlas, E. L., Schwartz, E., and Krol, M. C.: Intercomparison of Atmospheric Carbonyl Sulfide (TransCom-COS): 2. Evaluation of Optimized Fluxes Using Ground-Based and Aircraft Observations, *Journal of Geophysical Research: Atmospheres*, 128, e2023JD039 198, <https://doi.org/https://doi.org/10.1029/2023JD039198>, e2023JD039198 2023JD039198, 2023.
- Monks, S. A., Arnold, S. R., Hollaway, M. J., Pope, R. J., Wilson, C., Feng, W., Emmerson, K. M., Kerridge, B. J., Latter, B. L., Miles, G. M., Siddans, R., and Chipperfield, M. P.: The TOMCAT global chemical transport model v1.6: description of chemical mechanism and model evaluation, *Geoscientific Model Development*, 10, 3025–3057, <https://doi.org/10.5194/gmd-10-3025-2017>, 2017.
- Morgenstern, O., Hegglin, M. I., Rozanov, E., O’Connor, F. M., Abraham, N. L., Akiyoshi, H., Archibald, A. T., Bekki, S., Butchart, N., Chipperfield, M. P., Deushi, M., Dhomse, S. S., Garcia, R. R., Hardiman, S. C., Horowitz, L. W., Jöckel, P., Josse, B., Kinnison, D., Lin, M., Mancini, E., Manyin, M. E., Marchand, M., Marécal, V., Michou, M., Oman, L. D., Pitari, G., Plummer, D. A., Revell, L. E., Saint-Martin, D., Schofield, R., Stenke, A., Stone, K., Sudo, K., Tanaka, T. Y., Tilmes, S., Yamashita, Y., Yoshida, K., and Zeng, G.: Review of the global models used within phase 1 of the Chemistry–Climate Model Initiative (CCMI), *Geoscientific Model Development*, 10, 639–671, <https://doi.org/10.5194/gmd-10-639-2017>, 2017.
- Nayak, A. K., Xu, X., Frey, S. K., Khader, O., Erler, A. R., Lapen, D. R., Russell, H. A. J., and Sudicky, E. A.: Quantifying the potential of using Soil Moisture Active Passive (SMAP) soil moisture variability to predict subsurface water dynamics, *Hydrology and Earth System Sciences*, 29, 215–244, <https://doi.org/10.5194/hess-29-215-2025>, 2025.

- Nechita-Banda, N., Krol, M., van der Werf, G. R., Kaiser, J. W., Pandey, S., Huijnen, V., Clerbaux, C., Coheur, P., Deeter, M. N., and Röckmann, T.: Monitoring emissions from the 2015 Indonesian fires using CO satellite data, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373, 20170307, <https://doi.org/10.1098/rstb.2017.0307>, 2018.
- NOAA Climate.gov: Climate Variability: Oceanic Niño Index, <https://www.climate.gov/news-features/understanding-climate/climate-variability-oceanic-nino-index>, 2009.
- Nurdiati, S., Sopaheluwakan, A., and Septiawan, P.: Spatial and Temporal Analysis of El Niño Impact on Land and Forest Fire in Kalimantan and Sumatra, *Agromet*, 35, 1–10, <https://doi.org/10.29244/j.agromet.35.1.1-10>, 2021.
- ONeill, P., Chan, S., Njoku, E., Jackson, T., Bindlish, R., and Chaubell, J.: SMAP L3 Radiometer Global Daily 36 km EASE-Grid Soil Moisture, Version 8, <https://doi.org/10.5067/OMHVSRGFX38O>, 2021.
- Page, S. E. and Hooijer, A.: In the line of fire: the peatlands of Southeast Asia, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371, 20150176, <https://doi.org/10.1098/rstb.2015.0176>, 2016.
- Park, M., Worden, H. M., Kinnison, D. E., Gaubert, B., Tilmes, S., Emmons, L. K., Santee, M. L., Froidevaux, L., and Boone, C. D.: Fate of Pollution Emitted During the 2015 Indonesian Fire Season, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033474, <https://doi.org/10.1029/2020JD033474>, 2021.
- Parker, R. J., Boesch, H., Wooster, M. J., Moore, D. P., Webb, A. J., Gaveau, D., and Murdiyarso, D.: Atmospheric CH₄ and CO₂ enhancements and biomass burning emission ratios derived from satellite observations of the 2015 Indonesian fire plumes, *Atmospheric Chemistry and Physics*, 16, 10111–10131, <https://doi.org/10.5194/acp-16-10111-2016>, 2016.
- Pope, R. J., Arnold, S. R., Chipperfield, M. P., Reddington, C. L. S., Butt, E. W., Keslake, T. D., Feng, W., Latter, B. G., Kerridge, B. J., Siddans, R., Rizzo, L., Artaxo, P., Sadiq, M., and Tai, A. P. K.: Substantial Increases in Eastern Amazon and Cerrado Biomass Burning-Sourced Tropospheric Ozone, *Geophysical Research Letters*, 47, e2019GL084143, <https://doi.org/10.1029/2019GL084143>, <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2019GL084143>, 2020.
- Pradhan, R. K., Markonis, Y., Vargas Godoy, M. R., Villalba-Pradas, A., Andreadis, K. M., Nikolopoulos, E. I., Papalexiou, S. M., Rahim, A., Tapiador, F. J., and Hanel, M.: Review of GPM IMERG performance: A global perspective, *Remote Sensing of Environment*, 268, 112754, <https://doi.org/https://doi.org/10.1016/j.rse.2021.112754>, 2022.
- Pumphrey, H. C., Glatthor, N., Bernath, P. F., Boone, C. D., Hannigan, J. W., Ortega, I., Livesey, N. J., and Read, W. G.: MLS measurements of stratospheric hydrogen cyanide during the 2015–2016 El Niño event, *Atmospheric Chemistry and Physics*, 18, 691–703, <https://doi.org/10.5194/acp-18-691-2018>, 2018.
- Randerson, J., Van Der Werf, G., Giglio, L., Collatz, G., and Kasibhatla, P.: Global Fire Emissions Database, Version 4.1 (GFEDv4), <https://doi.org/10.3334/ORNLDAAAC/1293>, 2017.
- Remaud, M., Ma, J., Krol, M., Abadie, C., Cartwright, M. P., Patra, P., Niwa, Y., Rodenbeck, C., Belviso, S., Kooijmans, L., Lennartz, S., Maignan, F., Chevallier, F., Chipperfield, M. P., Pope, R. J., Harrison, J. J., Vimont, I., Wilson, C., and Peylin, P.: Intercomparison of Atmospheric Carbonyl Sulfide (TransCom-COS; Part One): Evaluating the Impact of Transport and Emissions on Tropospheric Variability Using Ground-Based and Aircraft Data, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037817, <https://doi.org/https://doi.org/10.1029/2022JD037817>, e2022JD037817 2022JD037817, 2023.
- Remedios, J. J., Leigh, R. J., Waterfall, A. M., Moore, D. P., Sembhi, H., Parkes, I., Greenhough, J., Chipperfield, M. P., and Hauglustaine, D.: MIPAS reference atmospheres and comparisons to V4.61/V4.62 MIPAS level 2 geophysical data sets, *Atmospheric Chemistry and Physics Discussions*, 7, 9973–10017, <https://doi.org/10.5194/acpd-7-9973-2007>, 2007.

- R'Honi, Y., Clarisse, L., Clerbaux, C., Hurtmans, D., Dufлот, V., Turquety, S., Ngadi, Y., and Coheur, P.-F.: Exceptional emissions of NH_3 and HCOOH in the 2010 Russian wildfires, *Atmospheric Chemistry and Physics*, 13, 4171–4181, <https://doi.org/10.5194/acp-13-4171-2013>, 2013.
- Rodgers, G. J.: *Optimal Estimation*, Springer, New York, NY, 2000.
- 885 Rodriguez Vasquez, M. J., Benoist, A., Roda, J.-M., and Fortin, M.: Estimating greenhouse gas emissions from peat combustion in wildfires on Indonesian peatlands, and their uncertainty, *Global Biogeochemical Cycles*, 35, e2019GB006218, 2021.
- Rosanka, S., Franco, B., Clarisse, L., Coheur, P.-F., Pozzer, A., Wahner, A., and Taraborrelli, D.: The impact of organic pollutants from Indonesian peatland fires on the tropospheric and lower stratospheric composition, *Atmospheric Chemistry and Physics*, 21, 11257–11288, <https://doi.org/10.5194/acp-21-11257-2021>, 2021.
- 890 Santoso, A., Mcphaden, M. J., and Cai, W.: The Defining Characteristics of ENSO Extremes and the Strong 2015/2016 El Niño, *Reviews of Geophysics*, 55, 1079–1129, <https://doi.org/https://doi.org/10.1002/2017RG000560>, 2017.
- Schroeder, W., Oliva, P., Giglio, L., and Csizsar, I. A.: The New VIIRS 375m active fire detection data product: Algorithm description and initial assessment, *Remote Sensing of Environment*, 143, 85–96, <https://doi.org/https://doi.org/10.1016/j.rse.2013.12.008>, 2014.
- Sheese, P. E., Walker, K. A., and Boone, C. D.: A global enhancement of hydrogen cyanide in the lower stratosphere throughout 2016, *Geophysical Research Letters*, 44, 5791–5797, <https://doi.org/https://doi.org/10.1002/2017GL073519>, 2017.
- 895 Singh, H. B., Brune, W. H., Crawford, J. H., Flocke, F., and Jacob, D. J.: Chemistry and transport of pollution over the Gulf of Mexico and the Pacific: spring 2006 INTEX-B campaign overview and first results, *Atmospheric Chemistry and Physics*, 9, 2301–2318, <https://doi.org/10.5194/acp-9-2301-2009>, 2009.
- Stockwell, C., Yokelson, R., Kreidenweis, S., Robinson, A., DeMott, P., Sullivan, R., Reardon, J., Ryan, K., Griffith, D., and Stevens, L.: Trace gas emissions from combustion of peat, crop residue, domestic biofuels, grasses, and other fuels: configuration and Fourier transform infrared (FTIR) component of the fourth Fire Lab at Missoula Experiment (FLAME-4), *Atmospheric chemistry and physics*, 14, 9727–9754, 2014.
- 900 Stockwell, C. E., Jayarathne, T., Cochrane, M. A., Ryan, K. C., Putra, E. I., Saharjo, B. H., Nurhayati, A. D., Albar, I., Blake, D. R., Simpson, I. J., et al.: Field measurements of trace gases and aerosols emitted by peat fires in Central Kalimantan, Indonesia, during the 2015 El Niño, *Atmospheric Chemistry and Physics*, 16, 11711–11732, 2016.
- 905 Storch, H. V. and Zwiers, F. W.: *Statistical Analysis in Climate Research*, Cambridge University Press, 1 edn., ISBN 978-0-521-45071-3 978-0-511-61233-6 978-0-521-01230-0, <https://doi.org/10.1017/CBO9780511612336>, 1999.
- Thompson, R. L., Patra, P. K., Ishijima, K., Saikawa, E., Corazza, M., Karstens, U., Wilson, C., Bergamaschi, P., Dlugokencky, E., Sweeney, C., Prinn, R. G., Weiss, R. F., O'Doherty, S., Fraser, P. J., Steele, L. P., Krummel, P. B., Saunois, M., Chipperfield, M., and Bousquet, P.: TransCom N_2O model inter-comparison – Part 1: Assessing the influence of transport and surface fluxes on tropospheric N_2O variability, *Atmospheric Chemistry and Physics*, 14, 4349–4368, <https://doi.org/10.5194/acp-14-4349-2014>, 2014.
- 910 Usup, A., Hashimoto, Y., Takahashi, H., and Hayasaka, H.: Combustion and thermal characteristics of peat fire in tropical peatland in Central Kalimantan, Indonesia, *Tropics*, 14, 1–19, <https://doi.org/10.3759/tropics.14.1>, 2004.
- van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth System Science Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
- Vetrita, Y. and Cochrane, M. A.: Fire Frequency and Related Land-Use and Land-Cover Changes in Indonesia's Peatlands, *Remote Sensing*, 12, <https://doi.org/10.3390/rs12010005>, 2020.

- Vigouroux, C., Stavrakou, T., Whaley, C., Dils, B., Dufлот, V., Hermans, C., Kumps, N., Metzger, J.-M., Scolas, F., Vanhaelewyn, G., Müller, J.-F., Jones, D. B. A., Li, Q., and De Mazière, M.: FTIR time-series of biomass burning products (HCN, C₂H₆, C₂H₂, CH₃OH, and HCOOH) at Reunion Island (21deg; S, 55deg; E) and comparisons with model data, *Atmospheric Chemistry and Physics*, 12, 10 367–10 385, <https://doi.org/10.5194/acp-12-10367-2012>, 2012.
- Vincent, R. A. and Dudhia, A.: Fast radiative transfer using monochromatic look-up tables, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 186, 254–264, <https://doi.org/10.1016/j.jqsrt.2016.04.011>, 2017.
- Watson, J. G., Cao, J., Chen, L.-W. A., Wang, Q., Tian, J., Wang, X., Gronstal, S., Ho, S. S. H., Watts, A. C., and Chow, J. C.: Gaseous, PM_{2.5} mass, and speciated emission factors from laboratory chamber peat combustion, *Atmospheric Chemistry and Physics*, 19, 14 173–14 193, <https://doi.org/10.5194/acp-19-14173-2019>, 2019.
- Wells, K. C., Millet, D. B., Brewer, J. F., Payne, V. H., Cady-Pereira, K. E., Pernak, R., Kulawik, S., Vigouroux, C., Jones, N., Mahieu, E., Makarova, M., Nagahama, T., Ortega, I., Palm, M., Strong, K., Schneider, M., Smale, D., Sussmann, R., and Zhou, M.: Global decadal measurements of methanol, ethene, ethyne, and HCN from the Cross-track Infrared Sounder, *Atmospheric Measurement Techniques*, 18, 695–716, <https://doi.org/10.5194/amt-18-695-2025>, 2025.
- Whitburn, S., Van Damme, M., Clarisse, L., Hurtmans, D., Clerbaux, C., and Coheur, P.-F.: IASI-derived NH₃ enhancement ratios relative to CO for the tropical biomass burning regions, *Atmospheric Chemistry and Physics*, 17, 12 239–12 252, <https://doi.org/10.5194/acp-17-12239-2017>, 2017.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geoscientific Model Development*, 4, 625–641, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.
- Yokelson, R. J., Saharjo, B. H., Stockwell, C. E., Putra, E. I., Jayarathne, T., Akbar, A., Albar, I., Blake, D. R., Graham, L. L., Kurniawan, A., et al.: Tropical peat fire emissions: 2019 field measurements in Sumatra and Borneo and synthesis with previous studies, *Atmospheric chemistry and physics*, 22, 10 173–10 194, 2022.
- Zhang, T., Wooster, M. J., and Xu, W.: Approaches for synergistically exploiting VIIRS I- and M-Band data in regional active fire detection and FRP assessment: A demonstration with respect to agricultural residue burning in Eastern China, *Remote Sensing of Environment*, 198, 407–424, <https://doi.org/https://doi.org/10.1016/j.rse.2017.06.028>, 2017.
- Zhou, M., Langerock, B., Vigouroux, C., Sha, M. K., Ramonet, M., Delmotte, M., Mahieu, E., Bader, W., Hermans, C., Kumps, N., Metzger, J.-M., Dufлот, V., Wang, Z., Palm, M., and De Mazière, M.: Atmospheric CO and CH₄ time series and seasonal variations on Reunion Island from ground-based in situ and FTIR (NDACC and TCCON) measurements, *Atmospheric Chemistry and Physics*, 18, 13 881–13 901, <https://doi.org/10.5194/acp-18-13881-2018>, 2018.




Figure 6. Monthly mean ERA5 vertical wind vectors (m s⁻¹) averaged along the equator for (a) October 2015, (b) November 2015, (c) October 2019, (d) November 2019, (e) October 2023, and (f) November 2023 (Hersbach et al., 2023). Brown (blue) areas indicate where, on average, the air is descending (ascending) for that particular month. The red dashed lines highlight the approximate meridional extent of the Sumatra and Kalimantan regions.




Figure 7. (a, top) Averaged SWI T=100, over Indonesia in the period 2014-2023. (b, bottom) Daily accumulated precipitation (combined microwave-IR) averaged over Indonesia estimate from the Integrated Multi-satellitE Retrievals for GPM (IMERG) in the period 2014-2023.




Figure 8. Global distribution of HCN total column (molecules cm^{-2}) on 2 November 2015 produced from (a, top left) T_HCN and (b, top right) T_HCN_{0.25Emiss}. The bottom panels shows the total column difference between the two runs and the IASI measurements regridded on the TOMCAT grid.




Figure 9. Monthly average HCN 2015 emissions (molecules $\text{cm}^{-2} \text{s}^{-1}$) used in the T_HCN TOMCAT model experiment (blue line) compared with the scaled emissions used in the T_HCN_{0.25Emiss} experiment (red dashed line).




Figure 10. Daily HCN enhancement ratios (ER) relative to CO calculated from the slope of the linear regression of HCN against CO total columns between 1 September and 1 December 2015, over the Indian Ocean (12°S-7°N, 90°E-127°E). For completeness the error bars, although small, are included and correspond to the standard errors on the regression slope. Any data with a correlation coefficient <0.3 for the slope have been excluded.




Figure 11. Time series from 1990-2023 of ONI index, with warm (red shaded area) and cold (blue shaded area) periods based on a threshold of $\pm 0.5^\circ\text{C}$.




Figure 12. Total cumulative monthly fire radiative power (MW) over Indonesia measured by VIIRS. The fire activity in 2023 (red line) is almost eight times smaller than in 2015 (blue line), and falls below the seasonal long-term (2012-2023) mean (black dotted line).




Figure 13. Daily Infrared Atmospheric Sounding Interferometer (IASI) derived a) HCN burden (Gg) and b) CO burden (Tg) between the 1 August and 31 December for three years: 2015 (blue line), 2019 (yellow line), and 2023 (red line). This is calculated for the region 12°S-7°N, 90°E-127°E.




Figure 14. Timeseries of daily mean SMAP soil moisture ($\text{m}^3 \text{m}^{-3}$) for all Indonesian peatlands, Kalimantan peatlands and Sumatra peatlands for 2015 (blue), 2019 (yellow) and 2023 (red) for the period 1 June to 31 December in years 2015, 2019 and 2023.




Figure 15. Daily Indonesian peatland fire count (from VIIRS) and collocated daily soil moisture ($\text{m}^3 \text{m}^{-3}$) for the period 1 June to 31 December in years 2015 (blue triangles), 2019 (yellow stars) and 2023 (red circles).




Figure 16. Weekly mean burn depth (m) of Indonesia peat fires between 1 June and 31 December in years 2015 (triangles), 2019 (stars) and 2023 (circles).




Figure 17. Accumulated daily burn depth of Indonesian peatland fires in 2015 (blue), 2019 (yellow) and 2023 (red) between 1 June and 31 December.




Figure 18. IASI measured HCN (red line) and CO (blue line) averaged total column and accumulated daily burn depth (black line) of all Indonesia peatland fires from 1 August to 31 December 2015.




Figure 19. Cross-correlation between the IASI HCN total column and the accumulated daily burned depth (black line) and between the IASI CO total column and the accumulated daily burned depth (red line).