

Heat and continental transport shape the variability of volatile organic compounds in the Eastern Mediterranean: Insights from multi-year observations and regional modeling

Formatted: Indent: Left: 0 cm

5 Anchal Garg¹, Maximilien Desservettaz^{1,2}, Alikı Christodoulou^{1,3}, Theodoros Christoudias¹, Vijay Punjaji Kanawade¹, Chrysanthos Savvides⁴, Mihalis Vrekoussis^{1,5}, Shahid Naqui¹, Tuija Jokinen¹, Joseph Byron⁶, Jonathan Williams^{1,6}, Nikos Mihelopoulos^{1,7}, Eleni Liakakou⁷, Jean Sciare¹, and Efstratios Bourtsoukidis¹

Formatted: Top: 1.75 cm

¹Climate and Atmosphere Research Centre, The Cyprus Institute, Nicosia 2121, Cyprus

²Centre for Atmospheric Chemistry, University of Wollongong, Wollongong, NSW 2500, Australia

10 ³PSI, Center for Energy and Environmental Sciences, 5232 Villigen, Paul Scherrer Institute, Switzerland

⁴Department of Labour Inspection, Ministry of Labour and Social Insurance, Nicosia, Cyprus

⁵Institute of Environmental Physics, University of Bremen, Bremen, Germany

⁶Atmospheric Chemistry Department, Max Planck Institute for Chemistry, 55128, Mainz, Germany

15 ⁷Institute for Environmental Research and Sustainable Development, National Observatory of Athens, Palaia Penteli, 15236 Athens, Greece

*Correspondence: Anchal Garg (a.garg@cyi.ac.cy), Efstratios Bourtsoukidis (e.bourtsoukidis@cyi.ac.cy)

20 **Abstract.** Volatile organic compounds (VOCs) are key precursors of tropospheric ozone formation and secondary organic aerosol formation, yet multi-year observations in the Eastern Mediterranean and Middle East (EMME) remain limited, thereby influencing regional air quality and climate. This study investigates the seasonal, diurnal, and temperature-driven variability of VOCs at a rural background site in Cyprus, located in the Eastern Mediterranean. VOC measurements (April 2022–June 2024) conducted with a proton transfer reaction time-of-flight Mass spectrometer (PTR-ToF-MS), along with HYSPLIT air-mass back-trajectory analysis and WRF-Chem atmospheric chemistry simulations. This study presents multi-year (April 25 2022–June 2024) high-resolution measurements of 76 VOCs using PTR-ToF-MS at a rural background site in Cyprus, combined with HYSPLIT air-mass analysis to examine the effects of regional transport on VOCs variability. Oxygenated VOCs (OVOCs) dominated the VOCs burden (~79%), followed by aliphatic hydrocarbons, aromatic hydrocarbons, and terpenes. Seventy-six VOCs were quantified and categorized into classes such as oxygenated VOCs (OVOCs), aliphatic hydrocarbons, aromatic hydrocarbons, terpenes, and nitrogen- and sulfur-containing compounds. Most VOCs exhibited clear distinct diurnal patterns, observed highest during 08:00–14:00 UTC, varying by species, due to enhanced photochemical activity and temperature-driven emissions. Terpenes, Biogenic VOCs, particularly isoprene, increased exponentially peaked in summer within temperature up to the 35–38 °C range but decreased beyond this threshold, indicating heat-stress inhibition. Monoterpenes showed elevated levels both day and night, reflecting contributions from both biogenic and anthropogenic sources. OVOCs, including acetone, acetaldehyde, methanol, and acetic acid, showed sharp enhancement above 35 °C, consistent with intensified primary emissions and secondary formation under extreme heat. Dimethyl sulphide (DMS) increased during warmer months, indicating enhanced marine microbial activity. OVOCs displayed strong seasonal and thermal enhancement during hot, dry summers due to both primary emissions and secondary formation. Aromatic Hydrocarbons were mainly higher during winter, linked to combustion processes, but Benzene levels was highest during summer particularly when temperature rose above 35 °C from evaporative and potential stress-related biogenic sources, whereas toluene and xylene were higher in colder months, linked to combustion processes. HYSPLIT air-mass trajectory analysis revealed dominant contributions from Europe and Northwest Asia (~68%), transporting aged OVOCs, while Middle East winter inflows enhanced aromatic hydrocarbons. While WRF-Chem captured seasonal trends, most VOCs were underestimated, highlighting missing-under-representation of emission sources and oxidation pathways in the model. Overall, the study emphasizes temperature and regional transport as key drivers of VOC variability in the Eastern Mediterranean.

50 1 Introduction

55 Volatile Organic Compounds (VOCs) are a diverse group of carbon-based chemicals with high vapor pressures that play a central role in atmospheric chemistry (Atkinson, 2000; Kamal et al., 2016). Upon release into the atmosphere, VOCs participate in photochemical reactions that contribute to the formation of tropospheric ozone (O₃), aerosol precursors, and secondary organic aerosol (SOA), all both of which have far-reaching implications for air quality, human health, and climate (Atkinson, 2000; Ehn et al., 2014;

Koppmann, 2020; Mellouki et al., 2015; Pennington et al., 2021; Wang et al., 2024). In scientific literature, VOCs are often categorized either by their chemical composition or by their source (Debevec et al., 2017; Koppmann, 2020; Yuan et al., 2024). These compounds with lifetimes ranging from minutes to months originate from both anthropogenic sources such as fossil fuel combustion, solvent use, industrial processes, and biogenic sources such as, including vegetation, soils, and marine emissions, and also emissions from biomass burning and wildfires (Atkinson, 2000; Bourtsoukidis et al., 2018; Capes et al., 2009; Civan et al., 2015; Debevec et al., 2017; Desservettaz et al., 2023; Huang et al., 2020; Kaltsonoudis et al., 2016; Pinthong et al., 2022; Pugliese et al., 2023; Ralf Koppmann, 2007; Sindelarova et al., 2014; Wang et al., 2022a; Yuan et al., 2024).

In scientific literature, these compounds are frequently classified into the broader categories of oxygenated VOCs (OVOCs), biogenic VOCs (BVOCs), and anthropogenic VOCs (AVOCs), each of the species with distinct sources, reactivities, and atmospheric lifetimes (Debevec et al., 2017; Koppmann, 2020; Yuan et al., 2024). OVOCs, including alcohols, aldehydes, ketones, organic acids, and others, are emitted directly from both anthropogenic and biogenic sources but are predominantly formed secondarily via the oxidation of primary VOCs by hydroxyl radicals (OH), O₃, and nitrate radicals (NO₃) (Huang et al., 2020; Mellouki et al., 2015; Wang et al., 2022b). These compounds are important intermediates in the photochemical formation of SOA and contribute to the oxidative capacity of the atmosphere over both local and regional scales (Huang et al., 2020; Liu et al., 2009; Mellouki et al., 2015; Wang et al., 2020, 2022b). On the other hand, BVOCs terpenes (such as isoprene, monoterpenes, and sesquiterpenes) are primarily emitted (90-95%) by vegetation through enzyme-driven processes influenced by environmental drivers (Bourtsoukidis et al., 2024, 2025; Brito et al., 2018; Ciccioli et al., 2023; Sindelarova et al., 2014; Wang et al., 2024), but some studies (Bryant et al., 2022; Panopoulou et al., 2020) shows that they also have smaller anthropogenic contributions. In the Mediterranean, pine and oak forests are major sources of terpenes BVOCs (Mecca et al., 2024; Song et al., 2011). Others VOCs such as AVOCs, comprising alkanes, alkenes, and aromatic hydrocarbons, whose major emission sources are anthropogenic arise primarily from fossil fuel combustion, industrial activities operations, solvent use, biomass burning (both through direct emissions and secondary production) and vehicular traffic, with elevated levels observed in urban areas and near industrial and shipping activities (Atkinson, 2000; Civan et al., 2015; Kajos et al., 2015; Kaltsonoudis et al., 2016; Perrone et al., 2014; Sicre et al., 1987). Oxygenated VOCs (OVOCs) including alcohols, aldehydes, ketones, and organic acids constitute a chemical class with both biogenic and anthropogenic origins, but they are also predominantly formed secondarily via the oxidation of primary VOCs by hydroxyl radicals (OH), O₃, and nitrate radicals (NO₃) (Huang et al., 2020; Mellouki et al., 2015; Wang et al., 2022b). These compounds are important intermediates in the photochemical formation of SOA and contribute to the oxidative capacity of the atmosphere over both local and regional scales (Huang et al., 2020; Liu et al., 2009; Mellouki et al., 2015; Wang et al., 2020, 2022b).

The eastern mediterranean and middle east (EMME) region is a climatically sensitive and chemically dynamic region (Lazoglou et al., 2024; Zittis et al., 2022) owing to high solar radiation, elevated temperatures, and complex meteorology. This regione-EMME experiences mixed air masses from Western and Eastern Europe, the Middle East, and North Africa, bringing emissions from industrial, biogenic, marine, and biomass burning sources (Bourtsoukidis et al., 2018; Germain-Piaulenne et al., 2024; Lelieveld et al.,

2002; Traub et al., 2003). This convergence, combined with the increasing frequency of heatwaves (Lazoglou et al., 2024), enhances ~~VOCs emissions biogenic and evaporative AVOCs emissions~~ and hence, photochemical activity, raising concerns over regional ozone and SOA formation. Cyprus, located at the core of this region, offers a unique setting to investigate these interactions.

~~Multi~~Long-term-year measurements of VOCs in the EMME are scarce. At Finokalia, Greece, the studies of Liakakou et al. (2007, 2009) reported extended records of isoprene and ~~C₂-C₈~~ hydrocarbons, while in Athens, the studies of Panopoulou et al. (2018, 2020) presented multi-year observations of isoprene, terpenes, and ~~C₂-C₈~~ non methane hydrocarbons (NMHCs). In contrast to these gas chromatography-based observations, proton-transfer-reaction time-of-flight mass spectrometry (PTR-ToF-MS) studies in the region have thus far been limited to short-term, campaign-based deployments (Derstroff et al., 2017; Kaltsonoudis et al., 2016). Overall, despite extensive research in diverse regions, the understanding of VOCs variability and drivers in the EMME region remains limited, particularly in terms of long-term and multi-year observations that can capture seasonal and climate change-driven dynamical responses to extreme heat.

Addressing this gap, the current study presents multi-year high-resolution VOCs measurements collected from 2022 to 2024 at a rural background site in Cyprus located in EMME region. Using PTR-ToF-MS, we quantified ~~and classified 76 VOCs species across biogenic, anthropogenic, and oxygenated classes~~. To identify regional ~~transporte~~contributions, we apply hybrid single-particle lagrangian integrated trajectory (HYSPLIT) air mass back-trajectory analysis and examine air mass origins across seven distinct geographical sectors. Furthermore, we evaluated selected VOCs ~~species~~ simulated by the weather research forecast model coupled with chemistry (WRF-Chem) model which includes a high-resolution mapping of local emissions (Georgiou et al., 2022). This integrated approach enables us to (i) characterize diurnal and seasonal VOCs patterns, (ii) assess the influence of temperature on VOCs composition ~~and chemical processing~~, (iii) understand key emission sources, (iv) evaluate the impact of long-range transport on VOCs variability, and (v) assess the model performance in the estimation of VOCs.

2 Methodology

2.1 Study area and VOCs measurement timeline

Cyprus, the third-largest island in the Mediterranean, is situated at the air pollution crossroads of Europe, Asia, and Africa, making it a receptor for long-range atmospheric transport from diverse source regions (Lelieveld et al., 2002). Air masses influencing the island originate from Western and Eastern Europe, Turkey, North Africa, Southwest and Northwest Asia, the Middle East, and the surrounding Mediterranean Sea (Fig. 1a). This convergence of continental and marine influences makes Cyprus an ideal location for investigating the chemical complexity of both long-range transported and local air pollutants (Meusel et al., 2016; Derstroff et al., 2017; Debevec et al., 2017; Vrekoussis et al., 2022).

The present study was conducted at the Cyprus Atmospheric Observatory—Agia Marina Xyliatou (CAO-AMX, 35.0387° N, 33.0579° E), located at 532 m above ~~mean~~ sea level at the foothills of the Troodos Mountains (Fig. 1b–c). Established in 1997 and upgraded in 2015 to a “supersite,” CAO-AMX provides high-quality, long-term measurements of aerosols and their properties, greenhouse gases, meteorological parameters, and trace gases. It is designed and operated as a rural background site (CAO-AMX,

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

135 <https://cao.cyi.ac.cy/>) primarily by the climate and atmosphere research center (CARE-C) of the Cyprus
institute (Cyl), in collaboration with national authorities such as the department of labour inspection (DLI)
of the ministry of labour and social insurance, while it is part of major international atmospheric monitoring
networks, including ACTRIS (<https://www.actris.eu/>); EMEP (<https://www.emep.int/>); and AERONET
140 (<https://aeronet.gsfc.nasa.gov/>) (Sciare J, 2016). The station is situated near the small villages of Agia Marina
(~630 inhabitants) and Xyliatos (~150 inhabitants), surrounded by oak-pine forests and Maquis shrubland,
with agricultural fields ~4 km to the north (Baalbaki et al., 2021). The nearest urban area lies >35 km away,
ensuring minimal influence from local pollution sources aside from limited vehicle traffic. Its rural setting,
surrounded by forested and low-population areas and situated far from major urban sources (Nicosia about
145 35 km to the northeast and Larnaca about 50 km to the southeast) ensures that measurements are minimally
influenced by major nearby local anthropogenic emissions. This allows the observation of representative
background concentrations and the signatures of regional and long-range transported pollutants.

The VOCs observations were conducted at the CAO-AMX site from 27 April ~~April-27~~, 2022, to 15 June
15, 2024, with occasional interruptions caused by other field campaigns or instrument downtime.
Meteorological data were recorded at a temporal resolution of 5 minutes from a dedicated station located in
150 the village of Xyliatos, approximately 2.85 km from the primary observational site at the CAO-AMX. DLI
managed the installation and operation of a relatively dense monitoring network at this station. As shown in
Fig. 1d, monthly data coverage varied but provided efficient coverage across all seasons. Fig. 1e summarizes
the seasonal and annual distribution of data points: summer (32%) was the most represented season, followed
by spring (30%) and autumn (23%), while winter accounted for 15% of the data. In terms of annual
155 contribution, 2022 comprised 44% of the total dataset, 2023 contributed 25%, and 2024 contributed 31%.

2.2 VOCs measurements

Online measurements of VOCs were performed using a high-resolution PTR-ToF-MS (Model: 4000; Ionicon
Analytik GmbH, Innsbruck, Austria) following the methodology as described by Desservettaz et al., (2023).

160 PTR-ToF-MS is a high-sensitivity, real-time technique that measures VOCs by transferring a proton from
H₃O⁺ ions to VOC molecules. It provides rapid, accurate, and high-resolution detection of a wide range of
atmospheric VOCs without sample preparation.

Briefly about the methodology for VOCs measurement, the drift tube of the PTR-ToF-MS was operated
under controlled conditions with an electric field to number density ratio (E/N) of 128 Townsend (Td), a
pressure of 2.2 mbar, an applied voltage of 630 V, and a temperature of 85 °C. Ambient air was sampled at a
165 flow rate of 210 mL min⁻¹ through a 2.0 m long, 1/16" PEEK (polyether ether ketone) heated inlet line,
equipped with a 0.45 µm PTFE particulate filter (PALL, USA). The inlet drew air from a common glass
manifold shared with other atmospheric measurement instruments. The use of chemically inert materials
such as PTFE and PEEK minimized potential compound losses within the sampling line.

The instrument was automatically switched to measure VOC-free air for 20 minutes after every 23 hours,
170 followed by 40 minutes of calibration using a certified gas standard (Apel-Riemer Environmental, USA)
containing major VOCs diluted to ~20 ppbv; this 23-h cycle caused the non-measurement period to shift by
approximately 1 h each day, ensuring that no specific time of day was systematically underrepresented and
that diurnal patterns were not biased., to establish baseline levels. VOCs not included in the gas standard

were calibrated using established relative response factors determined from prior laboratory experiments and, where possible, verified with secondary standards. Mass spectra were acquired at 1-minute time resolution and averaged to 1-hour intervals for the analysis in this study. The instrument sensitivity was interpolated between calibrations and adjusted for humidity dependence based on water cluster ion ratios. Calibrated sensitivities showed modest temporal variability (~10-23%) for most compounds.

Post-acquisition analysis was carried out using the Ionicon data analyzer (IDA), with mass axis calibration based on H_3O^+ and internal reference peaks (e.g., $\text{C}_6\text{H}_5\text{I}^+$). The software retrieved exact masses and chemical formulas for ~1200 peaks, of which 400–500 was assigned tentative structures using the GLOVOCS database, which is a master compound assignment guide that can be referred to as PTR-MS practitioners (Yáñez-Serrano et al., 2021). A set of 131 common species across analysis periods was further corrected using transmission curves derived from zero and calibration air sampling. Mixing ratios (ppbv) were calculated from ion counts, incorporating species-specific reaction rate constants (k-values) based on molecular dipole moments and polarizabilities (Desservettaz et al., 2023; Pagonis et al., 2019). The limit of detection (LOD) for each species was defined as three times the standard deviation observed during zero-air measurements and averaged over the entire campaign. VOC species with more than 30% of their data below the LOD were excluded, resulting in a final dataset of 76 species, with sub-LOD concentrations appropriately flagged. Measurements impacted by filter changes, exhibiting short-term interference (notably acids and certain hydrocarbons), were filtered by removing a conservative 2-hour window following each swap. Formaldehyde was not included in this study due to its inconsistent identification by the IDA software during part of the measurement period. Furthermore, quantification of formaldehyde via PTR-based techniques is known to be highly sensitive to humidity variations (Vlasenko et al., 2010), which complicates reliable detection. Accurate quantification would require water-dependent calibrations and a dedicated formaldehyde calibration standard, conditions that were not met during the measurement period.




Figure 1: Geographic context, air mass origins, and VOC measurement coverage at the CAO-AMX site (2022–2024). (a) Major regional air mass source areas influencing Cyprus; (b) Location of the CAO-AMX in the Troodos Mountains; (c) The CAO-AMX monitoring station, a designated regional background site; (d) Monthly distribution of

number of datapoints for VOC measurements from April 2022 to June 2024; (e) Summary of data coverage by season and year, showing overall representativeness of the dataset for multi-seasonal and multi-annual analysis.

2.3 Meteorological and air pollutant data

205 Meteorological and auxiliary air quality parameters were analyzed to provide contextual insight into atmospheric dynamics and pollutant characteristics. ~~Meteorological data were recorded at a temporal resolution of 5 minutes from a dedicated station located in the village of Xyliatos (35.0141° N, 33.0492° E), approximately 2.85 km from the primary observational site at the CAO-AMX.~~ The meteorological station (mentioned in section 2.1), ~~positioned at 10 meters above ground level,~~ continuously measured key ~~meteorological parameters atmospheric variables,~~ including ambient temperature (AT), relative humidity (RH), solar radiation (SR), wind speed (WS), and wind direction (WD). ~~DLI managed the installation and operation of a relatively dense monitoring network at this station.~~ In addition, a range of trace gases and particulate pollutant levels were measured at a co-located air quality monitoring station (~~~20 m from CAO-AMX, main VOCs measurement container~~), operating under the European monitoring and evaluation programme (EMEP) framework. ~~This station, situated approximately 20 meters from the main VOC measurement container,~~ provides hourly-resolved concentrations of major atmospheric pollutants: ozone (O₃), carbon monoxide (CO), nitrogen oxides (NO, NO₂, and NO_x), sulfur dioxide (SO₂), and particulate matter with aerodynamic diameters less than 10 μm (PM₁₀) and 2.5 μm (PM_{2.5}). These measurements serve as critical auxiliary data for assessing the chemical background of the region, evaluating pollutant transport processes, and validating model simulations. Instrument specifications, calibration protocols, and operational details of the meteorological and air quality monitoring systems are extensively documented in previous studies (Kleanthous et al., 2014; Pikridas et al., 2018; Vrekoussis et al., 2022).

2.4 HYSPLIT backward trajectory

225 To assess the influence of long-range atmospheric transport on local air quality, five-day back trajectories were computed using the HYSPLIT model (version 4.4; Draxler, 1997; Stein et al., 2015). These calculations were performed using the PC-based version of HYSPLIT provided by the national oceanic and atmospheric administration's (NOAA) air resources laboratory (ARL), incorporating meteorological data from the global data assimilation system (GDAS) with a spatial resolution of 1° × 1° and a temporal resolution of 1-hour (Draxler, 1997). Trajectories were initialized every hour at an altitude of 500 m above ground level (AGL), and each point along the trajectory was calculated at a one-hour time step. The air mass trajectory cluster analysis, following the methodology previously applied by Debevec et al., (2017) identified a total of seven air mass clusters: (C0) slow-moving or stagnant local air masses, (C1) North Africa, (C2) Marine-influenced, (C3) Europe, (C4) Northwest Asia, (C5) West Turkey, and (C6) Middle east origins. For the cluster analysis, the altitude of the end point was set at 1000 m AGL for each trajectory (Debevec et al., 2017).

2.5 WRF-Chem model configuration for VOC simulation in Cyprus

235 To assess VOCs variability and compare modeled and observed mixing ratios over Cyprus, we employed the WRF-Chem model as described and evaluated in Georgiou et al. (2022). The setup used three nested domains with horizontal resolutions of 50, 10, and 2 km, the later focusing on Cyprus, with meteorological inputs

from the global forecast system (GFS) every 3 hours. Key parameterizations included [Yonsei University \(YSU\) planetary boundary layer \(PBL\) scheme](#) for boundary layer, Noah land surface model (LSM) for land surface processes, the rapid radiative transfer model for global applications (RRTMG) for radiation, the regional atmospheric chemistry mechanism (RACM) for gas-phase chemistry and [Modal Aerosol Dynamics for Europe \(MADE\) for inorganic aerosols and Volatility Basis Set \(VBS\) MADE/VBS](#) for organic aerosols as described in [\(Georgiou et al., 2022\)](#). Photolysis was calculated using Fast-J. Anthropogenic emissions were sourced from emissions database for global atmospheric research-hemispheric transport of air pollution (EDGAR-HTAP_{v2}) version 2.1 emission inventory, while BVOCs were simulated online using the Model of Emissions of Gases and Aerosols from Nature version (MEGAN) version 2.1 based on weather and land use data (Guenther et al., 2012).

Formatted: English (United States)

For the innermost domain, a high-resolution emission inventory developed by [\(Georgiou et al., 2020\)](#) is used. This emission inventory uses the total reported emissions of CO, NO_x, non-methane VOCs (NMVOCs), SO₂, and PM on a 1 km × 1 km resolution (Georgiou et al., 2020) which is upscaled to the resolution of the innermost domain of the simulations (2 km) using a nearest-neighbor grid-point attribution algorithm, while diurnal, weekly, and monthly emission cycles are applied to each species and the predominant emission activity per season according to the (Schaap et al., 2011). The high-resolution emission inventory was found to adequately simulate the daily profiles of NO_x and O₃ at measurement sites over Cyprus, and compared to CAMS, the WRF-Chem model predicts more accurately NO₂ mixing ratios with 7% during winter and -44% during summer-normalized mean bias during winter/summer, compared to -81% and -84% biases for the CAMS ensemble using the CAMS inventory (Georgiou et al., 2022).

Formatted: Subscript

In the outer model domain, the EDGAR emission inventory is used [for anthropogenic emissions, and MEGAN for biogenic emissions](#). In comparison with other emission inventories for VOCs, EDGAR provides a larger estimate than the EMEP emission inventory. The CAMS-REG-AP inventory also gives a smaller estimate, in comparison to EMEP. Comparing the VOC by the EDGAR inventory to the EMEP inventory, the higher emission estimates are greatest over Belgium, Austria, Switzerland, Germany and Finland, with differences mainly encountered in the industrial combustion sector and fugitive emissions. Regarding the modelled O₃ concentrations over Europe, all emission inventories were shown ~~to show~~ very similar patterns despite the noted differences in terms of emissions, particularly VOCs, indicating a low sensitivity to VOCs emissions (Thunis et al., 2021).

Major VOCs ~~species~~ such as isoprene, monoterpenes, and aromatics were explicitly modeled. Outputs at hourly resolution were validated against PTR-ToF-MS measurements from CAO-AMX, enabling robust interpretation of VOC sources and atmospheric transport in the eastern mediterranean. The VOCs included in this study are based on the RACM (Stockwell et al., 1997) and encompass key anthropogenic and biogenic compounds relevant to regional air quality. These include isoprene (ISO), monoterpenes represented by α -pinene and related monocyclic terpenes with a single double bond (API), and d-limonene, along with other cyclic dienes (LIM). Aromatic compounds are grouped as toluene (TOL), which includes toluene and less reactive aromatics, and xylenes (XYL), representing xylenes and more reactive aromatic species. OVOCs

are categorized as aldehydes (ALD), including acetaldehyde and higher molecular weight aldehydes, ketones (KET), and glyoxal (GLY).

3 Results and Discussion

3.1 Meteorological and atmospheric composition variability

Mean meteorological conditions included a WS of $2.6 \pm 1.4 \text{ m s}^{-1}$, RH of $55 \pm 20\%$, AT of $17.7 \pm 7.9 \text{ }^\circ\text{C}$, and SR of $224.75 \pm 305.85 \text{ W m}^{-2}$ (Fig. 2a). Average AT ($\geq 30.5 \text{ }^\circ\text{C}$) and SR ($> 8600 \text{ W m}^{-2}$) peaked during daytime in summer, while winter had the lowest daytime values ($< 1.5 \text{ }^\circ\text{C}$ and $< 400\text{--}200 \text{ W m}^{-2}$ Fig. 2e-f). RH showed the opposite pattern during daytime ($\sim 60.85\%$ in winter, $< 43.0\%$ in summer) (Fig. 2g). Midday peaks in temperature and radiation were strongest in summer, with RH lowest in the afternoon (Fig. 2e-g). RH and SR were inversely correlated, consistent with previous studies (Emekwuru and Ejohwomu, 2023; Matthew, 2022). Daily average WS ranged from 1 to 4 m s^{-1} , with slightly higher values observed in spring and summer, suggesting enhanced mixing during these periods (Fig. 2h). Diurnal patterns were also pronounced as WS showed afternoon maxima in all seasons, with enhanced values in spring and summer, likely driven by thermal convection and sea breeze circulation (Fig. 2h), promoting more effective dispersion of air pollutants (Deot et al., 2025; Pérez et al., 2020; Soukissian and Sotiriou, 2022).




Figure 2: Seasonal and diurnal variability of meteorological parameters and air pollutants in Cyprus. Panels (a–c) show time series (24-hour rolling average) of (a) ambient temperature (AT) and solar radiation (SR), (b) relative humidity (RH) and wind speed (WS), and (c) gaseous pollutants, including ozone (O₃), carbon monoxide (CO), and nitrogen dioxide (NO₂) for study period. Panel (d) displays wind rose diagrams indicating wind speed and direction for the entire study period. Panels (e–h) present the diurnal profiles of (e) ambient temperature (°C), (f) solar radiation (Wm⁻²), (g) relative humidity (%), and (h) wind speed (ms⁻¹).

²), (g) relative humidity (%), and (h) wind speed (ms^{-1}) across the four seasons (spring, summer, autumn, winter). Shaded regions in panels (e-h) represent the standard error, capturing variability in diurnal cycles.

300 Wind analysis throughout the study period revealed predominantly moderate conditions, with over 70% wind speeds falling below 3 ms^{-1} , and 94% below 5 ms^{-1} . The wind rose diagrams (Fig. 2d) highlight the prevalence of moderate-speed winds and dominant directions ranging from northwesterly to west-southwesterly. Fig. S1 shows seasonal and monthly variability in wind direction, and suggests that during the winter, the winds were mainly coming from the southeast, followed by the southwest, with most wind speeds exceeding 4 ms^{-1} ; whereas, for the other seasons, the winds originated predominantly from the northwest.

305 Atmospheric conditions were characterized by the following mean mixing ratios for gaseous pollutants such as NO at $0.3 \pm 0.2 \text{ ppbv}$, NO₂ at $1.4 \pm 0.8 \text{ ppbv}$, NO_x at $1.7 \pm 0.9 \text{ ppbv}$, SO₂ at $0.5 \pm 0.4 \text{ ppbv}$, O₃ at $49.2 \pm 9.0 \text{ ppbv}$, and CO at $145.8 \pm 49.5 \text{ ppbv}$. During daytime O₃ remained consistently elevated throughout the year, with a significant increase of (8–20 ppbv) of its mixing ratio in summer, indicating a strong influence from regional transport and photochemical formation (Kleanthous et al., 2014; Lee et al., 2021; Pochanart et al., 2003). A notable increase in CO levels was observed between 2022 and 2024, [particularly linked to increase in anthropogenic activities](#). A modest rise in CO observed during winter is likely due to increased combustion emissions, lower sink due to low photochemistry, variable residence time of upto 90 days in winter and 30 days in summer, and weaker atmospheric dispersion (Koppmann et al., 2005; Pochanart et al., 2003). NO_x showed a winter maximum, consistent with increased anthropogenic emissions and reduced photochemical degradation (Atkinson, 2000; Civan et al., 2015; Koppmann et al., 2005).

3.2 Overview of atmospheric VOCs observations

320 A total of 76 VOCs were quantified, and their descriptive statistics for key species ([the most studied VOCs in EMME region and European studies](#)) are summarized in Table 1, while the complete list is provided in supplementary Table S1. These VOCs were grouped into ~~five~~ six major chemical classes: (i) OVOCs (~~including~~ alcohols, [aldehydes, ketones, carbonyls, organic acids, and other oxygenates](#)); (ii) aliphatic hydrocarbons; (iii) aromatic hydrocarbons; (iv) terpenes; and (v) nitrogen- and sulfur-containing VOCs (~~N- & S-containing VOCs~~). An overview of the average mixing ratios is shown in Fig. 3, while Fig. S2 illustrates the time series of selected VOCs. [The selected VOCs shown in Table 1 were most studied in the previous Mediterranean and European studies, whereas Fig S3 shows the 20 most abundant VOCs observed in this study.](#)

325 OVOCs were the dominant chemical class as [also](#) observed by Debevec et al.; (2017). [In our study OVOCs](#) contributed approximately 79% to the total measured VOC burden. Among these, alcohol was the most abundant subgroup, comprising 32% of total VOCs. Methanol was [found as](#) the single most dominant species, with a median mixing ratio of 2.86 ppbv, accounting for 87% of the total alcohol fraction. Its high dominance reflects oxidation of biogenic and biomass burning emissions, consistent with previous findings (Debevec et al., 2017; Mukherjee et al., 2024; Yuan et al., 2024). ~~Carbonyls-compounds~~ [\(aldehydes, and ketones\)](#) represented the second largest OVOCs subgroup (28% of total VOCs), with median levels dominated by acetone (2.11 ppbv, 74% of the carbonyl group), followed by acetaldehyde (0.54 ppbv, 19% of the carbonyl group). These compounds are known to originate from both primary sources and secondary

Commented [VK1]: It appears only one other place so no need to abbreviate it

oxidation of VOCs (Capes et al., 2009; Mellouki et al., 2015; Mukherjee et al., 2024; Pennington et al., 2021; Wang et al., 2024, 2022b). Elevated levels during summer (Fig. 2b) suggest increased biogenic activity and enhanced photochemical production.

Organic acids, dominated by acetic acid (median: 0.68 ppbv) and formic acid (median: 0.42 ppbv), contributed 13% of total VOCs. Their contributions likely arise from a combination of direct biogenic and anthropogenic emissions, as well as secondary formation, which could not be quantitatively distinguished. (Derstroff et al., 2017; Huang et al., 2020; Kamal et al., 2016; Koppmann, 2020; Wang et al., 2024, 2022b). Other oxygenates, such as methyl ethyl ketone (MEK) and acrolein, also contributed to the OVOCs group and may originate from industrial sources and atmospheric oxidation (Schieweck et al., 2021; Torres-Vinces et al., 2020; Yáñez-Serrano et al., 2016).

-Aliphatic hydrocarbons accounted for 12% of total VOCs, with species such as propylene, propyne, 1,3-butadiene, and cyclohexadiene likely reflecting emissions from vehicular traffic, fossil fuel combustion and industrial activities (Perrone et al., 2014; Sicre et al., 1987). Aromatic hydrocarbons, though less abundant (3%), were dominated by benzene (0.14 ppbv, 32% of the aromatic fraction), followed by toluene and xylenes. These compounds are well-known tracers of urban and industrial emissions (Abbasi et al., 2020; Boursoukidis et al., 2019; Boursoukidis et al., 2020; Das et al., 2024), though their overall concentrations were low, consistent with the remote and background nature of the site.

~~The primary BVOCs mainly include~~ terpenes, such as monoterpenes (median= 0.27 ppbv) and isoprene (median= 0.09 ppbv), collectively contributing ~3% of the VOC budget. Their mixing ratio peaked in summer, aligning with increased biogenic emissions driven by higher temperatures and SR (Gu et al., 2021; Liakakou et al., 2007; Mukherjee et al., 2024; Sindelarova et al., 2014; Strada et al., 2023; Wang et al., 2024).

~~Nitrogen- and sulfur-containing N- & S-containing~~ VOCs also contributed approximately 3%, with median acetonitrile (0.11 ppbv) and dimethyl sulfide (DMS, 0.04 ppbv) mixing ratios as the major constituents. Acetonitrile, a tracer for biomass burning, accounted for 40% of this group, while DMS accounted for 17% of this group reflects both marine and biological sources (Deschaseaux et al., 2022; Edtbauer et al., 2020; Huangfu et al., 2021; Robles, 2005).

The analysis of the 20 most abundant VOCs (Fig. S3) also reinforced the dominance of OVOCs, particularly methanol, acetone, and acetic acid, which together formed a substantial fraction of the observed VOC mass. The relatively low levels of aromatics and aliphatic further support the interpretation that this rural background site is primarily influenced by transported VOC precursors, secondary atmospheric processes and regional-scale transport, rather than direct anthropogenic emissions. A major reason for the low burden of aliphatic hydrocarbons in the PTR-ToF-MS measurements is their inherently low proton affinities, which result in weak ionization and significant fragmentation. Consequently, their signals are unreliable or very small, preventing a clear aliphatic hydrocarbon profile from being obtained in this study.

Table 1. Descriptive statistics of selected VOCs (in ppbv) measured in Cyprus (April 2022- June 2024)

Chemical Name	Chemical Formula	Chemical Class/Group	M.H ⁺	Min	Max	Mean	Median	SD
Methanol	CH ₄ O	Alcohol	33.03	0.138	27.82	3.08	2.86	1.71

Formatted: Space Before: 0 pt

Compound	Chemical Formula	Class	M.H ⁺	Concentration (ppbv)	Seasonal Variability	Winter	Spring	Summer	Autumn
Acetonitrile	C ₂ H ₃ N	N-containing VOC	42.04	0.041	0.33	0.12	0.11	0.04	0.04
Acetaldehyde	C ₂ H ₄ O	Aldehyde	45.03	0.123	6.76	0.88	0.54	0.89	0.89
Formic Acid	CH ₂ O ₂	Acids	47.01	0.022	7.99	0.78	0.42	0.93	0.93
Ethanol	C ₂ H ₆ O	Alcohol	47.05	0.009	3.68	0.43	0.35	0.35	0.35
Acrolein	C ₃ H ₄ O	Oxygenated	57.03	0.01	1.93	0.2	0.12	0.23	0.23
Glyoxal	C ₂ H ₂ O ₂	Oxygenated	59.01	0.019	1.22	0.07	0.07	0.04	0.04
Acetone	C ₃ H ₆ O	Ketone	59.05	0.429	10.82	2.35	2.11	1.19	1.19
Acetic Acid	C ₂ H ₄ O ₂	Acids	61.03	0.039	12.54	1.25	0.68	1.56	1.56
DMS	C ₂ H ₆ S	S-containing VOC	63.04	0.005	0.55	0.06	0.04	0.06	0.06
Furan	C ₄ H ₄ O	Oxygenated	69.03	0.004	0.16	0.01	0.01	0.01	0.01
Isoprene	C ₅ H ₈	Terpenes/Biogenic	69.07	0.005	2	0.15	0.09	0.18	0.18
MVK/Methacrolein	C ₄ H ₆ O	Oxygenated	71.05	0.003	0.53	0.08	0.06	0.07	0.07
MEK	C ₄ H ₈ O	Oxygenated	73.07	0.007	2.78	0.36	0.29	0.27	0.27
Benzene	C ₆ H ₆	Aromatics	79.05	0.018	1.09	0.16	0.14	0.11	0.11
Toluene	C ₇ H ₈	Aromatics	93.07	0.005	1.3	0.1	0.07	0.1	0.1
Xylenes	C ₈ H ₁₀	Aromatics	107.09	0.003	1.76	0.06	0.03	0.12	0.12
Monoterpenes	C ₁₀ H ₁₆	Terpenes/Biogenic	137.13	0.026	3.01	0.34	0.27	0.27	0.27

Compounds are grouped by chemical class, with molecular hydronated ion mass (M.H⁺) shown for each. These data provide insight into the variability and typical concentrations of VOCs in the region, relevant to air quality and atmospheric chemistry assessments. These are the most studied VOCs in EMME region and European studies. For clarity, nitrogen and sulfur containing VOCs class has been separated into distinct classes, N-containing VOC and S-containing VOC in this table to explicitly highlight the nitrogen contribution in acetonitrile and the sulfur contribution in DMS.




Figure 3: Overview of VOCs in Cyprus: Composition and seasonal variability in its composition. (a) % contribution of different VOC groups in Cyprus, highlighting the dominance of alcohols (32%), carbonyls (28%), and acids (13%). Subcategories within each VOC group are detailed in smaller pie charts. (b) Seasonal variability of the VOCs mixing ratio. Here carbonyls include aldehydes, and ketones. Overall OVOCs include alcohols, carbonyls, acids, and other oxygenated VOCs. Isoprene and Monoterpenes combines together to form terpenes class.

As shown in Table 2, VOC mixing ratios measured in our study are within the range reported from other European background and rural sites. Winter mixing ratios for acetaldehyde, acetone, MVK, MEK, acetonitrile, benzene, isoprene, and monoterpene were also comparable to those measured during the same

period in 2015 at this site (Debevec et al., 2017), suggesting stable background concentrations and low interannual variability in the Eastern Mediterranean. However, summertime isoprene mixing ratios at Ineia, Cyprus, were about three times lower. Derstroff et al. (2017) demonstrates potential contribution of downwind forest emissions or the effect of hotter summers and stress induced emissions. Elevated levels of most VOCs during summer (Fig. 3b) suggest the influence of increased biogenic activity and enhanced photochemical production, as well as contributions from solvent use and fuel evaporation.

Table 2. Seasonal mean mixing ratios (ppbv) of VOCs measured in our study compared with previous regional and European studies.

Location	Measurement	Study Reference	Year, Season	Terpenes		Aromatic Hydrocarbons			Oxygenated VOCs							N-containing VOCs
				Isoprene	Mono-terpenes	Benzene	Toluene	Xylenes	Methanol	Ethanol	Acetic Acid	Acetaldehyde	Acetone	MVK+ Methacrolein	MEK	
CAO, Cyprus (Background)	PTR-ToF-MS	This study	Spring 2022-24	0.12	0.20	0.14	0.06	0.03	3.22	0.47	0.84	0.97	2.03	0.06	0.35	0.10
			Summer 2022-24	0.18	0.39	1.30	0.07	0.03	3.92	0.35	1.00	0.61	2.90	0.09	0.28	0.13
			Autumn 2022-24	0.06	0.23	0.12	0.06	0.02	2.22	0.23	0.52	0.42	1.90	0.04	0.24	0.12
			Winter 2022-24	0.03	0.22	0.15	0.09	0.04	1.14	0.33	0.27	0.39	1.05	0.03	0.29	0.08
CAO, Cyprus (Background)	PTR-MS, GC-MS	Debevec et al., 2017	Winter, 2015	0.05	0.24	0.12	0.05	0.02	2.93			0.41	1.15	0.03	0.22	0.12
Ineia, Cyprus (Background)	PTR-MS	Derstroff et al., 2017	Summer, 2014	0.06	0.11	0.02	0.01	0.02	2.90		0.95	0.32	2.25	0.03	0.11	0.11
Zurich, Switzerland (Urban background)	GC-MS	Legreid et al., 2007	Spring 2005	0.08		0.41	1.46	0.97	2.18	6.87			1.66	0.04	0.24	
			Summer 2005	0.16		0.23	1.43	0.83	3.18	3.94		0.80	2.12	0.12	0.20	
			Autumn 2005	0.08		0.48	1.70	1.10	1.11	7.61		0.45	1.24	0.04	0.17	
			Winter 2005	0.06		0.75	1.25	0.91	1.21	7.53		0.82	1.17	0.05	0.22	
Athens, Greece (urban)	PTR-MS	Kaltsonoudis et al., 2016	Summer 2012	0.73	0.92	0.22	0.81	0.67		1.52	2.17		4.28	0.35	0.50	0.20
			Winter 2013	1.05	0.43	1.00	2.34	1.69		1.80	2.11		2.24	0.41	0.59	0.16
Cape Corsica, France (urban)	PTR-MS	Michoud et al., 2017	Summer 2013	0.19	0.41	0.03	0.08			0.18			3.43	0.06	0.48	
Oak Mediterranean Forest, France	PTR-MS	Kalogridis et al., 2014	Spring 2012	1.19	0.06	0.07	0.05		2.28			0.38	1.28	0.21		
Oak Mediterranean Forest, Spain	PTR-ToF-MS	Seco et al., 2011	Summer	0.15-0.75	0.13-1.42	0.04-0.09	0.08-0.47		4.14-6.05	1.15-2.87	1.32-2.94	0.54-1.26	2.26-3.83	0.17-0.54		0.16-0.22
			Winter	0.02-0.07	0.005-0.067	0.11-0.19	0.06-0.41		1.28-2.70	0.47-2.08	0.39-1.43	0.23-0.66	0.79-1.55	0.01-0.05		0.08-0.10
Barcelona, Spain (urban background)	PTR-ToF-MS	In't Veld et al., 2024	Spring 2022	0.04	0.05	0.14	0.47		3.50			0.92	1.93	0.09	0.22	0.41
Summer 2022			0.18	0.19	0.08	0.80		3.73			0.31	2.50	0.17	0.24	0.29	
Spring 2022			0.05	0.02	0.13	0.22		1.46			0.48	1.02	0.05	0.19	0.10	
Summer 2022			0.30	0.45	0.04	0.09		3.89			0.84	2.76	0.55	0.30	0.12	
Oak Mediterranean Forest, Spain	PTR-MS	Yáñez-Serrano et al., 2021	Summer 2019	0.42	0.15	0.04	0.13		4.60			0.77	2.08	0.30	0.29	0.12
			Autumn 2019	0.09	0.09	0.04	0.21		1.60			0.30	1.37	0.08	0.27	0.06

Formatted Table

As shown in Table 2, Generally, the Mediterranean forested sites in France and Spain are characterised by higher daily mean isoprene levels up to 1 ppbv (Kalogridis et al., 2014; Seco et al., 2011), consistent with strong direct emissions from oak-dominated ecosystems compared to the conifer-dominated forest in Cyprus. The urban site of Zurich, Switzerland shows similar seasonal pattern as our study, where isoprene, methanol, acetone, and MVK all showed a summer maximum and a winter minimum (Legreid et al., 2007). However, wintertime aromatic (BTX) concentrations were much higher in Zurich presumably due to combustion sources related to anthropogenic activities. Nitriles and MEK concentrations in our study fall in the range of those reported for previous European background studies (Table 2) and confirm their regional well-mixing as resulting from both anthropogenic and biomass-burning sources.

3.3. Seasonal and diurnal variability of VOCs functional groups

The diurnal variation of total VOCs (TVOCs) across all seasons and years (Fig. S4) reveals clear patterns influenced by temperature, photochemistry, boundary layer dynamics, and emission sources. TVOCs mixing ratio typically observed high between 09:00-14:00 UTC, corresponding to enhanced daytime emissions and increased atmospheric reactivity. The highest median TVOCs mixing ratios were observed during summer 2024 (25.76 ppbv), followed by summer 2023 (16.27 ppbv), spring 2024 (13.71 ppbv), and summer 2022 (12.35 ppbv), whereas winter exhibited the lowest levels (5.83 ppbv). ~~The less pronounced seasonal enhancement observed in summer 2022 compared to 2023 and 2024 is likely due to comparatively lower ambient temperature of about 1-3 °C resulting in weaker photochemical activity during that period, and reduced temperature-driven biogenic emissions. In contrast, summer 2024 was characterized by exceptional heat, including record-breaking temperatures (>45°C) across Cyprus, which strongly intensified direct VOCs emissions and secondary production. June 2024 was the warmest June on record in Cyprus, with 43 of 52 stations registering record maximum temperatures; during the 14–15 June heatwave, maximum temperature reached 45.3 °C, resulting in two heat-related deaths and widespread wildfire impacts (Department of Meteorology, Cyprus, 2024). These results underscore the influence of photochemical processes and temperature-driven biogenic emissions during warmer months, with the extreme heatwave in June 2024 likely intensifying VOCs release.~~ During summer, there is also northerlies that promote the transport of species from regions with pronounced anthropogenic emissions, including biomass burning, ~~contributing to elevated levels. Additionally, northerlies may bring pollution from north Cyprus (Vrekoussis et al., 2022). Also, June 2024 was the hottest ever recorded in Cyprus, with multiple daily maximum temperature records broken across the island (Cyprus Mail, 2024).~~ The prolonged heat (summer 2024) was attributed to a persistent heatwave influenced by global climate change, with such extreme events forecasted to become more frequent and intense in the future (Yáñez-Serrano et al., 2016; Zittis et al., 2022). ~~Similarly, the higher VOCs mixing ratio measured during spring 2024 relative to earlier spring campaigns can also be attributed to warmer-than-usual conditions, which enhanced biogenic release and photochemical reactivity.~~ Fig. 4 presents the diurnal profiles of VOCs for multiple seasons (2022–2024), which demonstrate

435 pronounced variability on seasonal and diurnal dynamics, particularly during summer 2024, when elevated temperatures significantly enhanced VOCs levels across most chemical classes.

3.3.1. Terpenes (isoprene and monoterpenes)


440 High isoprene levels occurred during 08:00-14:00 UTC (the time when SR and AT are highest) in summer 2023 and 2024 (Fig. 4a), correlating with peak SR and AT most of the time. In summer 2024, the median isoprene mixing ratio reached approximately 1.0 ppbv during the day, with daily average median values of 0.43 ± 0.36 ppbv, while monoterpenes showed broader diurnal presence, with early morning and evening peaks in some seasons, and a summer median of 0.51 ± 0.20 ppbv. Interestingly, the observed isoprene maximum at 08:00 UTC does not coincide with the peaks in SR and AT. This suggests that under the extreme drought conditions of 2023 and 2024, vegetation may have downregulated photosynthesis unusually early in the day. Such an early shutdown indicates that a physiological threshold was crossed, whereby plants shift their carbon allocation strategy from growth toward survival. While drought stress is often associated with an initial increase in isoprene emissions (Byron et al., 2022), prolonged or severe stress can suppress photosynthesis and consequently reduce isoprene release (Potosnak et al., 2014; Byron et al., 2022). On the other hand, monoterpenes showed broader diurnal presence, with early morning and evening peaks in all seasons, and a summer median of 0.51 ± 0.20 ppbv.

445 Isoprene and monoterpenes exhibited distinctly different temporal patterns, reflecting their contrasting emission drivers and atmospheric dynamics.

450 The seasonal pattern in isoprene align with established findings, where approximately 77% of annual emissions occur in spring (31%) and summer (46%), driven by elevated light intensity and temperature, which activate isoprene synthase enzymes (Fortunati et al., 2008; Monson et al., 1992). Autumn and winter emissions are markedly lower due to reduced photosynthetic activity and enzymatic downregulation (Panopoulou et al., 2020).

460 Monoterpenes, while also of biogenic origin mainly, demonstrated a more even seasonal distribution, with 38% of levels occurring in summer (0.39 ppbv) and 19–22% across other seasons (0.20 – 0.23 ppbv). This distribution reflects not only temperature dependence but also anthropogenic and ecological responses to herbivory, wounding, and water stress, particularly relevant under changing climate conditions (Bourtsoukidis et al., 2025; Peñuelas and Staudt, 2010; Strada et al., 2023; Wang et al., 2022a). The winter levels remain relatively stable, as stored monoterpenes in plant tissues continue to be emitted even during periods of low biosynthesis (Bourtsoukidis et al., 2024; Ciccioioli et al., 2023; Wang et al., 2022a, 2024). Consistent with the findings of Debevec et al., 2018, our observations also revealed elevated nighttime concentrations of monoterpenes, likely attributable to reduced oxidative sinks, boundary layer dynamics, and continuous emissions from vegetative storage pools.

Formatted: Indent: Left: 0 cm



470

475

Figure 4: Diurnal profiles (hourly averages) for mixing ratio (ppbv) of major VOCs. It includes terpenes (a-b), oxygenated VOCs (alcohols, carbonyls, acids, other oxygenated (c-k)), nitrogen and sulfur-containing VOCs (l-m), and aromatic hydrocarbons (n-p), measured at a rural background site in Cyprus from 2022 to 2024. Seasonal variations are color-coded, highlighting year-wise spring and summer trends alongside autumn, winter, and overall averages. Shaded areas indicate standard error for median values. X-axis here represents time in hours in UTC. [The VOCs selected in this figure are serve as representative indicators for their respective chemical families.](#)

3.3.2 Oxygenated VOCs

480

Methanol and ethanol exhibited moderate diurnal variation with elevated daytime and evening levels in summer (Fig. 4c-d). Median methanol observed as $\sim 5.52 \pm 1.27$ ppbv and ethanol as $\sim 0.76 \pm 0.26$ ppbv in summer 2024, reflecting both biogenic emissions and anthropogenic activities such as solvent use and burning. Carbonyls including acetaldehyde (2.38 ± 1.21 ppbv) and acetone (3.82 ± 1.02 ppbv), displayed sharp midday peaks in summer, consistent with secondary formation via photochemical oxidation. Their diurnal and seasonal consistency between 2023 and 2024 highlights photochemical production pathways. Organic acids [such as including](#) formic and acetic acid showed significant daytime enhancements in summer, particularly in 2024. Their median mixing ratios reached 1.76 ± 0.70 ppbv and 5.60 ± 2.75 ppbv, respectively, suggesting combined direct emissions and oxidative formation, possibly influenced by biomass burning or regional wildfires.

485

490

Among other OVOCs, [methyl vinyl ketone \(MVK\)](#), a product of isoprene oxidation, peaked at 0.15 ± 0.08 ppbv during summer midday. Furan, a tracer of biomass burning, showed morning maxima in summer 2023, indicating episodic fire activity. Consistent with isoprene emission, [methyl vinyl ketone \(MVK\)](#) also shows almost 42% of its emission during the summer season. A total of 60-70% levels of acetic acid, acetaldehyde, acetone, formic acid, and methanol emissions occur during summer and spring season, collectively showing their biogenic [origin and secondary formation dominance.](#)

495

3.3.3 Nitrogen and sulfur-containing VOCs

Acetonitrile (C_2H_3N), a N-containing VOC and a biomass burning tracer, showed minimal diurnal variation (Fig. 4l) due to its long atmospheric lifetime, which is approximately 620 days (Robles, 2005). Nevertheless, elevated levels were observed throughout the summer of 2023–2024, pointing to regional fire events during summer or long-range transport. The median acetonitrile seasonal mixing ratios were found to be highest during summer 2024 as 0.17 ± 0.02 ppbv, while lowest during winter as 0.08 ± 0.01 ppbv. DMS (C_2H_6S), a S-containing VOC, exhibited its highest average levels were observed highest during summer 2023 (0.12 ± 0.09 ppbv) (Fig. 4m), likely it could probably be due to enhanced marine biogenic emission from the nearby Mediterranean sea. The pronounced seasonal variation of DMS — strong seasonal signal — supports a temperature-driven biogenic sourceological origin (Deschaseaux et al., 2022; Edtbauer et al., 2020).

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

3.3.4 Aromatic hydrocarbons

Benzene showed summer midday enhancements (0.26 ± 0.12 ppbv), while toluene and xylenes exhibited flatter or bimodal patterns (high during morning and evening, probably due to increased vehicular activities) with higher variability. Toluene and Xylenes and toluene-mixing ratios peaked in winter and spring, consistent with local combustion under shallow boundary layers. The large standard error variation (Fig. 4o-p), especially for xylenes and toluene, indicates episodic anthropogenic inputs from traffic or regional pollution. Compared with benzene, these more reactive aromatics exhibited flat or noisy diurnal profiles with limited seasonal variability, aside from the winter-spring xylenes enhancement.

3.4. Temperature-driven enhancement of VOCs emissions in EMME regionthe eastern mediterranean

3.4.1. Temperature sensitivity of VOCs

The variation in VOCs mixing ratios with temperature revealed clear temperature-dependent trends across most VOCs (Fig. 5). Among terpenes, BVOCs, particularly isoprene, showed strong responses to rising temperatures, as their mixing ratio increased markedly beyond 25 °C, with a peak observed between 35–38 °C (0.53 ± 0.22 ppbv). There is an exponential increase in isoprene levels up to 38 °C, however, a slight decline (0.02–0.06 ppbv) in its level above 38 °C suggests crossing a tipping point through inhibition under extreme heat stress conditions (Fortunati et al., 2008). Monoterpenes exhibited a comparatively low increase in mixing ratios with an increase in temperature, and its trend was more gradual. The moderate rise at 35–38 °C (from 0.34 to 0.41 ppbv, ~21% increase) supports a biogenic contribution during summer, while the relatively stable levels across temperature bins may indicate sustained emissions throughout the day.

DMS levels also rose steadily with temperature, with maximum values observed between 35–38 °C (0.18 ± 0.13 ppbv). This trend is likely driven by enhanced microbial activity in the Mediterranean

sea (Edtbauer et al., 2020) and increased volatilization rates in warm, semi-arid, and coastal conditions typical of the region (Edtbauer et al., 2020).

535 -OVOCs also exhibited strong temperature dependence. Acetone and methanol levels increased progressively, reflecting a combination of direct biogenic emissions and secondary formation via oxidation of precursor compounds (Huang et al., 2020; Liu et al., 2009; Mellouki et al., 2015; Wang et al., 2022b). Methanol levels rose sharply above 38 °C (~19% with mean value of 4.85±0.91 ppbv), consistent with known biogenic sources such as plant metabolism and microbial activity (Dorokhov et al., 2018). Elevated methanol levels at high temperature are primarily consistent with biogenic sources, such as plant metabolism and microbial activity, although emissions from forest fires may also contribute under high-temperature conditions. Methanol is also produced during forest fire events, so one more probable cause could be increased fire activity with average temperature beyond 38 °C (Paton-Walsh et al., 2008; Yokelson et al., 1999; Dorokhov et al., 2018). Ethanol also shows temperature dependence, but it is weaker compared to methanol likely reflects its mixed source profile at the study site. While methanol is largely controlled by temperature-driven biogenic emissions and photochemistry, ethanol receives significant contributions from non-biogenic sources (e.g., solvent use, vehicular exhaust, long-range transport, and agriculture), which are less temperature-sensitive. As shown in Fig. 5, ethanol levels increase markedly only above 38 °C, suggesting additional evaporative emissions from anthropogenic sources. Acetic acid and acetaldehyde showed particularly sharp increases above 35 °C, with acetic acid levels ~2.8 times higher beyond 38 °C (5.4±2.9 ppbv) when compared with the levels at 35–38 °C. These patterns indicate enhanced photochemical production and possible evaporation-driven emissions (Debevec et al., 2017; Xu et al., 2023; Yuan et al., 2024). The increase of OVOCs under warmer conditions (>38 °C) highlights their potentially significant role in SOA formation in the EMME region (Azmi and Sharma, 2023; Pennington et al., 2021).


540
545
550
555
560
565 Among aromatic hydrocarbonsanthropogenic VOCs, benzene displayed a notable rise above 35 °C (0.35 ppbv), suggesting contributions from evaporate emissions, or long-range transport (Dimitriou and Kassomenos, 2020; Gjesteland et al., 2019). Another possible reason for the high daytime benzene levels, particularly during hot days (AT>35 °C) could be heat-stress-induced biogenic emissions (Misztal et al., 2015), where vegetation was found to release diverse benzenoid compounds under stress, potentially for communication or protective functions. However, the exact cause remains uncertain and warrants further investigation. In contrast, both toluene and xylenes levels were highest at lower temperatures (<15 °C), likely reflecting solvent use, decreased photochemistry, or indoor emissions during cooler periods (Abbasi et al., 2020; Song et al., 2011). A minor elevation (~0.01 ppbv) in toluene and xylenes was observed at 35–38 °C, potentially due to evaporative processes or reduced atmospheric dispersion.

3.4.2. Intensified VOC loading during the Summer 2024 heatwaves

June 2024 was the hottest month on record in Cyprus and coincided with the highest TVOCs mixing ratios observed during the 2022–2024 period (25.76 ± 8.29 ppbv), exceeding summer maxima

Formatted: English (India)

570 recorded in 2023 and 2022 (Fig. S4). As illustrated in Fig. 4, most VOCs also showed markedly
 higher emissions in 2024 compared to previous years. Considering that our observations span only
 until 15 June 2024, we focused on the comparison of the first half of June across the three years.
 The data revealed that TVOCs mixing ratio in early June 2024 (23.53 ppbv) were more than double
 compared to those measured during the same period in 2022 (9.98 ppbv) and 2023 (8.59 ppbv). This
 575 sharp rise was accompanied by extreme meteorological conditions: record-breaking average
 temperatures (24.95 °C), intense SR (314.81 Wm⁻²), and very low RH (27%). In contrast,
 temperatures in early June 2022 and 2023 averaged 21.91 °C and 20.61 °C, with RH at 59.73% and
 56.39%, respectively.



580 **Figure 5: Temperature Dependence of Ambient VOC mixing ratio in Cyprus.** Box plots showing the
 variation in ambient mixing ratio (in ppbv) of selected VOCs across different temperature ranges (°C) at a rural
 background site in Cyprus. Each box represents the interquartile range (IQR) with the median marked, and
 whiskers indicating variability outside the upper and lower quartiles. *The VOCs selected in this figure are serve
 as representative indicators for their respective chemical families.*

585 These conditions favored both primary **biogenicBVOCs** emissions and secondary atmospheric
 processes. Rapid vegetative growth in spring, followed by early thermal stress, likely stimulated
 isoprene and monoterpene emissions, and maybe also benzene. Elevated temperatures and oxidant
 availability enhanced both the emission (Bourtsoukidis et al., 2014) and formation of OVOCs such
 as acetaldehyde, acetic acid, and methanol. Concurrently, increased levels of CO (144.77 ppbv) and
 590 O₃ (58.77 ppbv) in 2024, compared to 2022 (132.69 and 45.13 ppbv) and 2023 (119.27 and 51.88
 ppbv), indicate stronger photochemical activity and increased ozone formation potential (**OFP**).

595
600
605
610
Midday peaks in isoprene ~~and monoterpenes~~, along with enhanced OVOCs mixing ratios (e.g., acetone, acetic acid, acetaldehyde), reflect intense photochemical production under elevated SR. Also, the elevated nighttime MVK/Isoprene ratio (0.96) indicates substantial isoprene oxidation, consistent with an aged air mass that has undergone prolonged exposure to OH radicals. This further supports enhanced oxidation of biogenic precursors (Guo et al., 2012). In contrast, the lower daytime ratio (0.62) suggests comparatively fresher isoprene emissions, reflecting active daytime production of isoprene and limited conversion to MVK, which requires time to form through photochemical processes (Apel et al., 2002). Elevated MVK/isoprene ratio further confirm enhanced oxidation of biogenic precursors (Guo et al., 2012). The findings suggest that escalating heatwave intensity and frequency in the EMME region could significantly increase VOC burdens, contributing to secondary pollutant formation (O₃ and SOA) and worsening regional air quality.

Wildfire activity may also exhibit pronounced variability and likely influenced the observed VOC composition at the CAO-AMX site (Fig. S6). From 2022-2024, the regional fire activity began in spring; however, its timing and intensity differed substantially. In 2022, the burned area increased rapidly in early summer, exceeding ~1800 ha in June (EEFIS, 2025). In 2023, fire activity was comparatively weaker during spring and early summer, with the largest burned areas occurring later in the season (August-September, 900 and 700 ha respectively). In contrast, 2024 was characterized by an earlier onset of wildfires, with fires occurring from April onwards and peak burned areas of >700 ha in May and ~2250 ha in June. As a result, springtime fire influence was strongest in 2024 compared to 2022 and 2023 (EEFIS, 2025). These spring and summer wildfire episodes, particularly in 2024, are expected to have enhanced biomass-burning-related VOCs at the site. Fire-influenced air masses typically exhibit elevated concentrations of acetonitrile, methanol, and other OVOCs.

3.5. Inter-species relationship

615
620
Based on the comprehensive correlation analysis in Fig. 6, clear patterns emerge among VOCs, meteorological parameters, and pollutant levels. The overall correlation matrix reveals strong positive associations among OVOCs, including -such as acetaldehyde, acetic acid, ethanol, acetone, methanol, and formic acid (-often with correlation coefficients $r > 0.6-0.9$), suggesting common sources and -secondary formation atmospheric processes. Ethanol does not cluster closely ($r < 0.5$) with other OVOCs in the hierarchical analysis (Fig. S7). Its more mixed source profile, including contributions from both local and transported anthropogenic activities, results in distinct temporal behaviour relative to primarily biogenic or photochemical VOCs. Thus, the clustering and weaker temperature dependence are consistent with ethanol's more complex source influences at the site.

625
-Isoprene strongly correlate with -monoterpenes, and MVK also correlate strongly, especially isoprene and MVK ($r = 0.75$), indicating secondary photochemical formation and biogenic influence during high radiation periods. Isoprene is also positively associated with temperature, and solar radiation ($r = 0.5$) indicating that variability in isoprene and MVK is driven by photochemical processing. On the contrary, monoterpenes shows weak correlation at both interspecies level, and

630 with meteorological parameters, showing its mixed sources from both biogenic and anthropogenic origins.

Aromatic hydrocarbons (e.g., toluene and xylenes) form another cluster with moderate-to-strong correlations ($r = 0.94$), consistent with urban combustion sources. Among aromatic hydrocarbons, benzene show comparatively weaker relation with toluene and xylenes.

635 Seasonal correlation matrices (Fig. S6) highlight dynamic VOC-environment interactions. On the contrary, benzene-toluene ($r = 0.6$) and benzene-xylenes ($r = 0.4$) correlation is moderate in winter, maybe due to low photochemical loss due to low temperature, but weakens in summer ($r < 0.3$) due to faster toluene degradation and variable sources (Gelencsér et al., 1997). Toluene is showing weak correlation ($r < 0.3$) with other VOCs and environmental parameters, showing their main source is anthropogenic in origin. Whereas, benzene shows strong correlation with other VOCs specially OVOCs ($r > 0.6$) showing either similar sources or evaporation process. In summer, BVOCs correlate most strongly with temperature and radiation, indicating dominant temperature-driven emissions. Winter shows weaker correlations due to reduced photochemistry and biogenic activity, while spring and autumn display moderate, transitional patterns. NO₂ and CO maintain consistent moderate correlations with aromatic VOCs (BTX) across seasons, underscoring persistent combustion and vehicular influences, either local or transported.

645 At the rural Cyprus background site, CO shows a diurnal cycle with an early-morning minimum and an afternoon-late evening maximum, which may be driven by boundary-layer dilution after sunrise and subsequent accumulation under regionally influenced air masses. The generally weak correlation between CO and aromatic VOCs suggests that CO variability is primarily controlled by regional transport and its longer atmospheric lifetime, rather than local traffic emissions. Seasonally, CO exhibits moderate correlations with benzene in autumn ($r \approx 0.4$) and winter ($r \approx 0.5$), likely reflecting enhanced regional combustion sources, reduced boundary-layer heights, and slower photochemical removal during colder months. Correlations are weak in spring and summer due to stronger photochemical processing and deeper boundary layers. CO shows no correlation with toluene or xylene in any season, consistent with their shorter lifetimes and the absence of fresh local emissions at this background site.

650 RH shows negative correlations with most VOCs, likely due to emission suppression or enhanced wet removal. O₃ is correlated moderate but positive ($r > 0.25$) with methanol, acids, acetone, isoprene and also with AT and SR positively correlated with reactive VOCs like isoprene and monoterpenes, reflecting active photochemical processes specially during summer. Although in correlation matrix, WS and WD show weaker correlations but may still influence variability, as suggested by backward trajectory analysis in the section 3.6 below. Overall hierarchical clustering in Fig. 6 groups of species by chemical and functional similarity. OVOCs cluster tightly, as do aromatics, while temperature, radiation, and ozone form a separate group. These clusters reflect common sources and atmospheric behaviors. Seasonal correlation matrices (Fig. S7) also highlight dynamic VOC-environment interactions, with most strong correlation among most VOCs during summer.

Formatted: Indent: Left: 0 cm




Figure 6: Correlation matrix and hierarchical clustering of VOCs, meteorological parameters, and air pollutants. Pearson correlation matrix illustrates the relationships among VOCs, meteorological variables (e.g., temperature, solar radiation, humidity), and air pollutants (NO₂, O₃, CO, PM_{2.5}, PM₁₀). Here, cluster refers to the air-mass trajectory cluster index obtained from the HYSPLIT back-trajectory analysis (shown in Fig. 7), where each number corresponds to a distinct air-mass transport pathway. Positive correlations are shown in orange to red, while negative correlations are represented in green to blue. Hierarchical clustering (dendrogram) on the right groups the variables with similar correlation patterns. Strong intra-group correlations are observed among oxygenated-VOCs (e.g., acetone, acetaldehyde, acetic acid), biogenic VOCs (e.g., isoprene, MVK, monoterpenes), and aromatic compounds (e.g., benzene, toluene, xylene), reflecting common emission sources and photochemical behavior.

The diurnal variation of selected interspecies mixing ratios (Fig. S87) reveals distinct patterns driven by emission sources, photochemical processing, and atmospheric reactivity. Ratios such as toluene/benzene, and xylenes/benzene decrease during the day (minima: ~0.45, and ~0.15 respectively), indicating stronger photochemical degradation of toluene and xylenes compared to the more stable benzene (Tiwari et al., 2010). The MVK/isoprene ratio increases from ~0.5 in the early morning to ~1.0 by midday, reflecting active daytime oxidation of isoprene to MVK (Guo et al., 2012). Similarly, the isoprene/monoterpenes ratio peaks around 1.2 between 10:00–14:00 UTC, highlighting the temperature- and radiation-driven nature of isoprene emissions, while monoterpenes remain relatively stable (Kiendler-Scharr et al., 2009). The ozone/NO₂ ratio exhibits a strong daytime increase, rising from ~30 to over 60 at noon, reflecting photochemical ozone formation from NO₂ under solar radiation. Overall, the interspecies ratios effectively differentiate between primary emissions and secondary formation, with photochemically active species (isoprene, MVK, ozone) showing strong diurnal variability.

Formatted: Subscript

670

675

680

685

690

3.6. Air mass origins and transport patterns

Using the HYSPLIT model to calculate air mass backward trajectories, we identified seven air mass clusters (Fig. 7) that influences VOCs transport and composition over Cyprus. The dominant clusters originated from Northwest Asia (C4, 34.3%) and Europe (C3, 33.6%), jointly accounting for more than two-thirds of the long-range air mass transport. These clusters represent strong, recurrent long-range synoptic-scale transport from urbanized and industrialized continental regions, which occur more frequently than locally influenced air trajectories, and thus played a key role in shaping seasonal and spatial VOC variability. Marine-influenced air masses (C2, 13.4%) reflected mixed characteristics, carrying both marine biogenic signatures and aged anthropogenic pollution. Less frequent clusters included air masses from North Africa (C1, 8.1%), Western Turkey (C5, 4.3%), Middle East (C6, 4.3%), and stagnant local conditions (C0, 1.7%). Notably, C6, although infrequent overall, was more prevalent during winter (10.7%) and exhibited elevated levels of benzene, toluene, and xylenes (BTX) during winter, suggesting long-range transport of aromatic hydrocarbons of AVOCs from the Middle East. C0, characterized by low wind speeds and residence near the site, showed high concentrations of primary pollutants such as benzene, acetonitrile, ethanol, and NO₂, consistent with local anthropogenic and biomass burning sources. Despite dominant local Northwest winds (as shown in Windrose diagram Fig. 2d), back trajectories show multi-day synoptic transport. The more frequent large-scale flows (C3-C4) therefore contribute over two-thirds of long-range air mass transport, exceeding the influence of local pathways (C1-C2).




Figure 7: Five-day air mass backward trajectories starting at 500 m above the ground level starting at every hour. These were calculated from the NOAA ARL PC-version HYSPLIT transport and dispersion model using gridded wind fields from the GDAS, which has a spatial resolution of 1° × 1° longitude by latitude and a time resolution of 1 hr (Draxler and Hess, 1999) to demonstrate the influence of long-range transported air masses at the measurement site.

Distinct chemical signatures were observed across clusters. European clusters (C3) exhibited elevated OVOCs such as methanol, acetone, and acetic acid, indicating the presence of aged air masses and secondary formation processes. C3 also carried higher levels of isoprene and MVK, possibly from transported biogenic emissions. C4 was associated with elevated O₃, potentially due to photochemical aging during long-range transport. In contrast, Western Turkey (C5) air masses had generally low VOC mixing ratios, though slight O₃ enhancement was observed, possibly due to regional NO_x-rich emissions and transport.

Formatted: Subscript




Figure 8: Mixing ratios of VOCs, and other air pollutants (in ppbv except PM_{2.5}, which is in $\mu\text{g m}^{-3}$) across air mass clusters in the regional site at Eastern Mediterranean. Boxplots showing the distribution of mixing ratios of selected VOCs, and other air pollutants for different air mass clusters (C0–C6) identified by back trajectory analysis. The black line within each box represents the median; boxes indicate the interquartile range, and whiskers denote variability outside the upper (75th) and lower (25th) quartiles. [The VOCs selected in this figure are serve as representative indicators for their respective chemical families.](#)

Seasonal analysis (Fig. S98–S109) revealed pronounced variability in both air mass distribution and VOC loading. Spring and summer were dominated by clusters from Europe and NW Asia, corresponding with enhanced biogenic emissions, active photochemistry, and increased long-range transport. These seasons showed elevated mixing ratios of isoprene, monoterpenes, methanol, acetic acid, and O₃. Summer, and spring exhibited the highest overall VOC and O₃ levels across most clusters, underscoring strong atmospheric photochemical reactivity. Summer maintained high VOC loadings, particularly in C2 and C3 clusters, driven by elevated temperatures and insolation. Autumn displayed a relatively balanced air mass distribution with moderate VOC levels. Winter was characterized by reduced photochemical activity and lower overall VOC concentrations, except for C6, which exhibited persistently elevated anthropogenic markers such as BTX likely associated with heating-related emissions and regional transport under more stagnant meteorological

745 conditions. These findings underscore the interplay between long-range transport, regional meteorology, and local emissions in shaping atmospheric VOCs variability over Cyprus. The combined influence of continental, marine, and regional sources contributes to complex spatiotemporal variability, with implications for air quality, ozone formation, and atmospheric oxidation capacity in the EMME region.

3.7. WRF-Chem model simulations

750 WRF-Chem simulations were evaluated against high-resolution PTR-ToF-MS observations sampled at CAO-AMX across multiple seasons to assess the model's capability to reproduce VOCs variability in the EMME region. BVOCs (isoprene and monoterpenes) emissions were online estimated by MEGAN v2.1 (Guenther et al., 2012), while anthropogenic emissions were derived from based on EDGAR-HTAP for the outer domains and a Cyprus high-resolution inventory for the innermost domain (Georgiou et al., 2020, 2022). Key biogenic species include are isoprene and monoterpenes, while anthropogenic aromatics include benzene, toluene, and xylenes. These species, together with OVOCs, are evaluated using seasonal distributions, diurnal behavior, and annual averages (Fig. 9, Figs. S110–S121).

760 The model successfully captured the seasonal and diurnal variability of several VOCs, particularly the strong summer enhancement associated with elevated temperatures and photochemical activity. Strong seasonal and daytime enhancements were found for terpenes, particularly during summer in both observed and modelled mixing ratios due to temperature and light driven emissions. These patterns are in good agreement with previous findings at other Mediterranean and European background stations where biogenic fluxes are strongly modulated during periods of photochemical activity. Terpenes exhibited pronounced daytime and summertime maximum levels in both observations and simulations, consistent with temperature- and light-driven biogenic emissions reported at other Mediterranean and European background sites (Ge et al., 2024; Genard-Zielinski et al., 2015). Among the evaluated species, isoprene showed the best agreement, with the model reproducing its temporal variability ($r = 0.52$, $p < 0.05$), although absolute mixing ratios were overestimated by approximately a factor of three. While the model captured the general temporal dependence of isoprene ($r = 0.52$, $p < 0.05$), absolute mixing ratios were overestimated by about a factor of 3. This overestimation is consistent with previous European-scale studies showing that MEGAN-based WRF-Chem simulations tend to overpredict isoprene emissions, particularly in southern and semi-arid regions (Morichetti et al., 2022). The study by Morichetti et al., 2022 observed that WRF-Chem simulations using MEGAN v3.1 overpredicted isoprene emissions by approximately 20–40% across Europe, particularly in southern regions. These biases have been
765 was primarily attributed to the model's lack of an effective drought stress and soil-moisture constraints response in the current MEGAN emission parameterization (Ge et al., 2024; Genard-Zielinski et al., 2015; Morichetti et al., 2022).

780 InBy contrast, Strada et al. (2023) reported strong underprediction of isoprene mixing ratios for Cyprus, that was also attributed to uncertainties in water availability dynamics, demonstrating inadequate


Formatted: Italian (Italy)

Formatted: Indent: Left: 0 cm

understanding of the role of soil moisture in regulating isoprene emissions in the EMME region. ~~These contrasting results emphasize the need for improved, region-specific parameterizations of plant stress responses under extreme meteorological conditions.~~

785 For monoterpenes, the model significantly underestimated the measured mixing ratios, by up to a
factor of 6-7 during ~~daytime~~ ~~the day~~ and 3-4 at night (Fig. S12). The underestimation of
monoterpenes in the model could stem from the assumption that their emissions are primarily
790 controlled by light and temperature. If monoterpenes originate predominantly from storage pools,
their fluxes will exhibit only a weak light dependency, which could explain the observed
discrepancies. In contrast, the overestimation of isoprene in our study may be linked to heat- and
drought-induced stress during the summer months. Such conditions can alter plant carbon allocation
strategies, leading to reduced isoprene emissions and enhanced monoterpene release, consistent with
observations reported by (Byron et al., (2022), ~~but this mechanism may not be represented in
current MEGAN parameterizations. Incorporating storage-driven emissions and stress-dependent
795 emission pathways would therefore be critical for improving monoterpene simulations in hot, semi-
arid regions. Biogenic emission (isoprene, and monoterpenes) parameterisations require improved
representation of high-temperature and water-stress responses (Sindelarova et al., 2014). In
particular, heat-stress emission factors and land-atmosphere coupling should be targeted in future
work, consistent with studies showing that drought stress algorithms improve the capability of
800 MEGAN isoprene emission estimates (Wang et al., 2022c). Photochemical reaction pathways,
chemical ageing and deposition processes that show sensitivity to heat wave atmospheric states can
be investigated to more properly quantify atmospheric removal in WRF-CHEM (Knote et al.,
2015a,b). In addition, wildfire and anthropogenic VOC emissions should be produced using high
805 spatiotemporal (hourly, 1x1 km) resolution to capture the impact of applying diurnal variability and
vertical distribution to fire emissions, along with VOC speciation that better reflects regional source
profiles.~~

OVOCs ~~including acids, aldehydes, and ketones~~ showed strong ~~seasonal and diurnal and seasonal~~
810 ~~enhancements during summer, consistent with intense photochemical production and evaporative
processes, yet were underestimated by factors of 3–12 depending on chemical class, in the
observations, especially during hot, dry summer periods with enhanced photochemical activity but
were systematically and consistently underestimated by the model. Mean mixing ratios of observed
acids, aldehydes, and ketones were calculated 12, 9, and 3 times respectively higher than the model
values. These similar discrepancies are aligned with underprediction of OVOCs across between
observed and modeled OVOCs have been documented throughout Europe due to missing secondary
formation chemistry, incomplete representation of oxidation products, and inadequate
815 representation treatment of temperature-dependent evaporative emissions during elevated
temperatures (Ge et al., 2024). Taken together, these results indicate that there are significant
deficiencies in the representation of OVOCs precursors and multi-phase processing by RACM.~~



Formatted: Justified, Line spacing: single

820 **Figure 9: Seasonal variation among measured and simulated VOCs mixing ratios at rural background site in Cyprus. The selection, grouping, and classification of VOCs presented in this figure follow the compound categories available within the chemical mechanism of the model to ensure consistency between observations and simulations.**

825 Aromatic hydrocarbons VOCs like toluene and xylenes showed relatively flat diurnal profiles with slight morning and evening peaks in cooler months (Fig. S12[†]), indicative of local anthropogenic sources under stable atmospheric conditions. However, WRF-Chem significantly underestimated these species, particularly xylenes, implying incomplete representation of urban emissions, traffic sources, and solvent use in anthropogenic inventories. Glyoxal, an important secondary oxidation product, exhibited a clear midday peak in measurements, but modeled values were significantly lower, even though the timing of the peak was captured, suggesting underestimation of precursor VOCs and low yields in photochemical pathways.

830 Seasonal comparisons (Fig. 9) confirmed these discrepancies. Observed levels of isoprene, aldehydes, ketones, and monoterpenes peaked in spring and summer, consistent with enhanced biogenic and photochemical activity. Modeled outputs, however, underrepresented OVOCs and AVOCs across all seasons. Despite some agreement in seasonal patterns, WRF-Chem was unable to reproduce the magnitude and timing of peak VOC levels. The annual comparison (Fig. S11[†]) showed that while isoprene mean simulated values were relatively aligned with observations, most other VOC groups, particularly OVOCs and aromatics, were systematically underestimated.

840 EDGAR generally reports higher VOC emissions compared to EMEP or CAMS-REG-AP due to differences in industrial combustion and fugitive sector representation (Thunis et al., 2021). However, these differences in VOC emissions have a limited impact on European-scale ozone modelling, and chemical regime and meteorology remain dominant drivers of O₃ variability (Thunis et al., 2021). A high-resolution Cyprus inventory may significantly improve the spatial representation of anthropogenic emissions and reduces NO₂ and O₃ biases compared to simulations based on CAMS (Georgiou et al., 2022), but uncertainties in BVOCs and OVOCs sources remain the largest contributors to discrepancies.

Overall, the analysis reveals that the heat extremes of the Eastern Mediterranean evoke a distinctive VOC emission response that enhances differences between observations and models based on traditional parameterizations developed in temperate conditions. The systematic underprediction of monoterpenes and OVOCs, when taken together with the overestimation of isoprene, suggests that incorporation of drought-sensitive plant physiology, storage-driven emission dynamics, and extended secondary oxidation chemistry within regional air quality models is required. Both focused biogenic emission modeling efforts and OVOC reaction pathways are necessary to appropriately quantify atmospheric reactivity and chemical transport in this climate-sensitive region. Also, as the frequency and intensity of heatwaves are projected to increase under future climate scenarios (Zittis et al., 2021; Lelieveld et al., 2016), the non-linear response of VOC emissions to temperature stress must be better parameterized to prevent significant underestimations of SOA and ozone formation.

4 Conclusions

~~This study provides a comprehensive assessment of VOCs variability in the Eastern Mediterranean based on multi-year high-resolution PTR-ToF-MS observations at a rural background site in Cyprus, supported by meteorological data, HYSPLIT back trajectories, and WRF-Chem model simulations. VOCs were classified into biogenic, anthropogenic, and secondary/oxygenated types to enable interpretation under varying environmental conditions. OVOCs were the dominant chemical class, contributing approximately 79% to the total measured VOC burden. This study provides the first multi-year, high-resolution characterization of volatile organic compounds (VOCs) in the Eastern Mediterranean based on continuous PTR-ToF-MS measurements at the Cyprus Atmospheric Observatory. By integrating detailed chemical observations with meteorological analyses, HYSPLIT air-mass trajectories, and WRF-Chem simulations, we offer a comprehensive assessment of the processes governing VOC variability in a rapidly warming and climatically sensitive region. Across the 2022–2024 period, oxygenated VOCs dominated the total VOC burden (~79%), reflecting the strong influence of direct sources, secondary formation, and long-range transport on regional atmospheric composition.~~

~~Among terpenes, BVOCs, particularly isoprene, exhibited pronounced summer peaks and daytime maxima, closely linked to solar radiation and temperature. Isoprene increased with temperature up to 38 °C before declining, suggesting thermal stress emission suppression, while monoterpenes showed both daytime and nocturnal presence may be due to diverse emission drivers from vegetation, boundary layer dynamics, or anthropogenic contribution. OVOCs (acetone, methanol, acetic acid, acetaldehyde) were enhanced during hot, dry periods, indicating amplified primary emissions and secondary photochemical formation. DMS peaks under high temperatures pointed to microbial activity and volatilization in semi-arid and coastal environments. Among aromatic hydrocarbons, benzene increased sharply above 35 °C, likely from biogenic origin, evaporative emissions and long-range transport, whereas toluene and xylenes were more abundant in cooler periods and degraded rapidly at higher temperatures.~~

Formatted: English (India)

885 HYSPLIT trajectory cluster analysis revealed elevated aromatic hydrocarbonsAVOCs, especially
aromatics, under the influence of long-range transported continental air masses from middle east
cluster (C6), Europe, NW Asia, the Middle East, and SW Asia. The variability in organic acid and
sulfur compound was attributed to marine and North African air masses. Slow-moving local air
masses, although infrequent, showed disproportionately high VOCs mixing ratio, indicating strong
890 local emission influence. Seasonal patterns showed summer dominated by biogenic emissions and
active photochemistry, while winter was characterized by anthropogenic influences under stagnant
conditions; spring emerged as a particularly reactive period.

WRF-Chem model simulations captured general seasonal patterns of biogenic isoprene but
systematically underpredicted OVOCs, aromatics, and secondary VOCs, with notable temporal
895 mismatches in diurnal cycles. These shortcomings reflect limitations in emission inventories,
oxidation schemes, and meteorological coupling, particularly for region-specific and stress-induced
emissions.

Overall, VOC variability over Cyprus is driven by the interplay of heat, seasonality, source type,
meteorology, and long range transport. Overall, this work demonstrates that VOC variability in the
Eastern Mediterranean is governed by the combined influence of extreme heat, seasonal variation,
900 regional transport, and complex photochemistry. The findings underscore the need for improved
VOC parameterizations in regional to global models, region-specific source inventories, and
adaptive air quality management strategies that account for both local and long-range transported
emissions. A dedicated source apportionment analysis would be a valuable follow-up to this work,
allowing for quantitative attribution of the observed VOCs to specific sources. With projected
905 increases in climatic extremes in the EMME region, understanding these processes is critical for
improved tropospheric ozone budget, secondary organic aerosol production, and broader air quality
and climate impacts. With heatwaves expected to intensify across the EMME region, understanding
how VOC emissions respond to climatic extremes is critical for predicting future ozone formation,
910 secondary organic aerosol production, and the broader oxidative capacity of the atmosphere. This
multi-year dataset provides a valuable foundation for future source apportionment analysis,
evaluation of regional model performance, and development of improved emission inventories. As
one of the few long-term VOC records in the region, it represents an important step toward
constraining atmospheric processes in a globally significant climate hotspot.

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold, English (United States)

915 **Data availability.** All data used in this study are openly available at
<https://zenodo.org/records/17483541>

Author Contribution.

920 AG: Conceptualization, Data curation, Methodology, Validation, Formal analysis, Investigation,
Writing – original draft, Writing – review & editing, Visualization. MD: Data curation,
Methodology, Software, Validation, Formal analysis, Writing – review & editing. AC: Methodology,

Data curation, Formal analysis, Writing – review & editing. TC: Data curation, Software, Writing – review & editing. VPK: Methodology, Software, Writing – review & editing. CS: Data curation, Resources, Writing – review & editing. MV: Methodology, Writing – review & editing. SN: Formal analysis, Writing – review & editing. TJ: Writing – review & editing. JB: Writing – review & editing. JW: Writing – review & editing. NM: Writing – review & editing. EL: Writing – review & editing. JS: Funding acquisition, Resources, Writing – review & editing. EB: Conceptualization, Project administrator, Methodology, Validation, Investigation, Writing – review & editing, Supervision.

Competing interests.

The contact author has declared that none of the authors has any competing interests.

Acknowledgements.

This work was supported by the European Union's Horizon 2020 research and innovation program (grant No. 856612) and the Cyprus Government (EMME-CARE). The authors thank the CAO technical team for the routine upkeep of the AMX station. EB and SN acknowledge support from the project EVOCPOLIS funded by The Cyprus Research and Innovation Foundation (RIF), implemented under the Cohesion Policy Programme “Thalia 2021-2027” and co-funded by the EU. TJ and VPK are supported by the European Union ERC-2022-STGERC-BAE-Project: 101076311. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them.

References

- Abbasi, F., Pasalari, H., Delgado-Saborit, J. M., Rafiee, A., Abbasi, A., and Hoseini, M.: Characterization and risk assessment of BTEX in ambient air of a Middle Eastern City, Process Safety and Environmental Protection, 139, 98–105, <https://doi.org/10.1016/j.psep.2020.03.019>, 2020.
- [Apel, E. C., Riemer, D. D., Hills, A., Baugh, W., Orlando, J., Faloona, I., et al.: Measurement and interpretation of isoprene fluxes and isoprene, methacrolein, and methyl vinyl ketone mixing ratios at the PROPHET site during the 1998 intensive, J. Geophys. Res.-Atmos., 107, ACH-7, <https://doi.org/10.1029/2000JD000225>, 2002.](https://doi.org/10.1029/2000JD000225)
- Atkinson, R.: Atmospheric chemistry of VOCs and NO_x, Atmos Environ, 34, 2063–2101, [https://doi.org/10.1016/S1352-2310\(99\)00460-4](https://doi.org/10.1016/S1352-2310(99)00460-4), 2000.
- Azmi, S. and Sharma, M.: Global PM_{2.5} and secondary organic aerosols (SOA) levels with sectorial contribution to anthropogenic and biogenic SOA formation, Chemosphere, 336, <https://doi.org/10.1016/j.chemosphere.2023.139195>, 2023.
- Baalbaki, R., Pikridas, M., Jokinen, T., Laurila, T., Dada, L., Bezantakos, S., Ahonen, L., Neitola, K., Maisser, A., Bimenyimana, E., Christodoulou, A., Unga, F., Savvides, C., Lehtipalo, K., Kangasluoma, J., Biskos, G., Petäjä, T., Kerminen, V.-M., Sciare, J., and Kulmala, M.: Towards understanding the characteristics of new particle formation in the Eastern Mediterranean, Atmos Chem Phys, 21, 9223–9251, <https://doi.org/10.5194/acp-21-9223-2021>, 2021.

- 960 Boutsoukidis, E., Williams, J., Kesselmeier, J., Jacobi, S., and Bonn, B.: From emissions to ambient mixing ratios: online seasonal field measurements of volatile organic compounds over a Norway spruce-dominated forest in central Germany, *Atmos Chem Phys*, 14, 6495–6510, <https://doi.org/10.5194/acp-14-6495-2014>, 2014.
- 965 Boutsoukidis, E., Behrendt, T., Yañez-Serrano, A. M., Hellén, H., Diamantopoulos, E., Catão, E., Ashworth, K., Pozzer, A., Quesada, C. A., Martins, D. L., Sá, M., Araujo, A., Brito, J., Artaxo, P., Kesselmeier, J., Lelieveld, J., and Williams, J.: Strong sesquiterpene emissions from Amazonian soils, *Nat Commun*, 9, 2226, <https://doi.org/10.1038/s41467-018-04658-y>, 2018.
- 970 Boutsoukidis, E., Ernle, L., Crowley, J. N., Lelieveld, J., Paris, J.-D., Pozzer, A., Walter, D., and Williams, J.: Non-methane hydrocarbon (C₂–C₈) sources and sinks around the Arabian Peninsula, *Atmos Chem Phys*, 19, 7209–7232, <https://doi.org/10.5194/acp-19-7209-2019>, 2019.
- Boutsoukidis, E., Pozzer, A., Williams, J., Makowski, D., Peñuelas, J., Matthaios, V. N., Lazoglou, G., Yañez-Serrano, A. M., Lelieveld, J., Ciais, P., Vrekoussis, M., Daskalakis, N., and Sciare, J.: High temperature sensitivity of monoterpene emissions from global vegetation, *Commun Earth Environ*, 5, 23, <https://doi.org/10.1038/s43247-023-01175-9>, 2024.
- 975 Boutsoukidis, E., Guenther, A., Wang, H., Economou, T., Lazoglou, G., Christodoulou, A., Christoudias, T., Nölscher, A., Yañez-Serrano, A. M., and Peñuelas, J.: Environmental Change Is Reshaping the Temperature Sensitivity of Sesquiterpene Emissions and Their Atmospheric Impacts, *Glob Chang Biol*, 31, <https://doi.org/10.1111/gcb.70258>, 2025.
- 980 Brito, J., Freney, E., Dominutti, P., Borbon, A., Haslett, S. L., Batenburg, A. M., Colomb, A., Dupuy, R., Denjean, C., Burnet, F., Bourriane, T., Deroubaix, A., Sellegri, K., Borrmann, S., Coe, H., Flamant, C., Knippertz, P., and Schwarzenboeck, A.: Assessing the role of anthropogenic and biogenic sources on PM_{2.5} over southern West Africa using aircraft measurements, *Atmos Chem Phys*, 18, 757–772, <https://doi.org/10.5194/acp-18-757-2018>, 2018.
- 985 [Bryant, D. J., Nelson, B. S., Swift, S. J., Budisulistiorini, S. H., Drysdale, W. S., Vaughan, A. R., Xu, L., Rickard, A. R., Hewitt, C. N., Nemitz, E., Mullinger, N., Acton, W. J. F., Young, D. E., Allan, J., & Hamilton, J. F.: Biogenic and anthropogenic sources of isoprene and monoterpenes and their secondary organic aerosol in Delhi, India. *Atmos. Chem. Phys. Discuss.*, 2022, 1–35. <https://doi.org/10.5194/acp-2022-xxx>, 2022.](#)
- 990 Byron, J., Kreuzwieser, J., Purser, G., van Haren, J., Ladd, S. N., Meredith, L. K., Werner, C., and Williams, J.: Chiral monoterpenes reveal forest emission mechanisms and drought responses, *Nature*, 609, 307–312, <https://doi.org/10.1038/s41586-022-05020-5>, 2022.
- 995 Capes, G., Murphy, J. G., Reeves, C. E., McQuaid, J. B., Hamilton, J. F., Hopkins, J. R., Crosier, J., Williams, P. I., and Coe, H.: Secondary organic aerosol from biogenic VOCs over West Africa during AMMA, *Atmos. Chem. Phys.*, 3841–3850 pp., 2009.
- 1000 Ciccio, P., Silibello, C., Finardi, S., Pepe, N., Ciccio, P., Rapparini, F., Neri, L., Fares, S., Brilli, F., Mircea, M., Magliulo, E., and Baraldi, R.: The potential impact of biogenic volatile organic compounds (BVOCs) from terrestrial vegetation on a Mediterranean area using two different emission models, *Agric For Meteorol*, 328, 109255, <https://doi.org/10.1016/j.agrformet.2022.109255>, 2023.
- 1005 Civan, M. Y., Elbir, T., Seyfioglu, R., Kuntasal, Ö. O., Bayram, A., Doğan, G., Yurdakul, S., Andiç, Ö., Müezzinoğlu, A., Sofuoğlu, S. C., Pekey, H., Pekey, B., Bozlaker, A., Odabasi, M., and Tuncel, G.: Spatial and temporal variations in atmospheric VOCs, NO₂, SO₂, and O₃ concentrations at a heavily industrialized region in Western Turkey, and assessment of the carcinogenic risk levels of benzene, *Atmos Environ*, 103, 102–113, <https://doi.org/10.1016/j.atmosenv.2014.12.031>, 2015.

Cyprus Mail: Last month hottest June ever in Cyprus., Cyprus Mail, 2024.

- 1010 Das, A., Giri, B. S., and Manjunatha, R.: Systematic review on benzene, toluene, ethylbenzene, and xylene (BTEX) emissions; health impact assessment; and detection techniques in oil and natural gas operations, *Environmental Science and Pollution Research*, 32, 1–22, <https://doi.org/10.1007/s11356-024-35698-1>, 2024.
- 1015 Debevec, C., Sauvage, S., Gros, V., Sciare, J., Pikridas, M., Stavroulas, I., Salameh, T., Leonardis, T., Gaudion, V., Depelchin, L., Fronval, I., Sarda-Esteve, R., Baisnée, D., Bonsang, B., Savvides, C., Vrekoussis, M., and Locoge, N.: Origin and variability in volatile organic compounds observed at an Eastern Mediterranean background site (Cyprus), *Atmos Chem Phys*, 17, 11355–11388, <https://doi.org/10.5194/acp-17-11355-2017>, 2017.
- 1020 Debevec, C., Sauvage, S., Gros, V., Sellegri, K., Sciare, J., Pikridas, M., Stavroulas, I., Leonardis, T., Gaudion, V., Depelchin, L., Fronval, I., Sarda-Esteve, R., Baisnée, D., Bonsang, B., Savvides, C., Vrekoussis, M., and Locoge, N.: Driving parameters of biogenic volatile organic compounds and consequences on new particle formation observed at an eastern Mediterranean background site, *Atmos Chem Phys*, 18, 14297–14325, <https://doi.org/10.5194/acp-18-14297-2018>, 2018.
- 1025 [Department of Meteorology \(Cyprus\). Ministry of Agriculture, Rural Development and Environment. \(2024, 15 July\). *Monthly Weather Report: The Weather in June 2024* \[PDF\]. \[https://www.dom.org.cy/CLIMATOLOGY/English/Monthly%20Weather%20Bulletin/2024/06_Monthly%20Weather%20Report_Jun2024_ENG.pdf\]\(https://www.dom.org.cy/CLIMATOLOGY/English/Monthly%20Weather%20Bulletin/2024/06_Monthly%20Weather%20Report_Jun2024_ENG.pdf\)](https://www.dom.org.cy/CLIMATOLOGY/English/Monthly%20Weather%20Bulletin/2024/06_Monthly%20Weather%20Report_Jun2024_ENG.pdf)
- 1030 Derstroff, B., Hüser, I., Boursoukoudis, E., Crowley, J. N., Fischer, H., Gromov, S., Harder, H., Janssen, R. H. H., Kesselmeier, J., Lelieveld, J., Mallik, C., Martinez, M., Novelli, A., Parchatka, U., Phillips, G. J., Sander, R., Sauvage, C., Schuladen, J., Stöner, C., Tomsche, L., and Williams, J.: Volatile organic compounds (VOCs) in photochemically aged air from the eastern and western Mediterranean, *Atmos Chem Phys*, 17, 9547–9566, <https://doi.org/10.5194/acp-17-9547-2017>, 2017.
- 1035 Deschaseaux, E. S. M., Swan, H. B., Maher, D. T., Jones, G. B., Schulz, K. G., Koveke, E. P., Toda, K., and Eyre, B. D.: The Interplay Between Dimethyl Sulfide (DMS) and Methane (CH₄) in a Coral Reef Ecosystem, *Front Mar Sci*, 9, <https://doi.org/10.3389/fmars.2022.910441>, 2022.
- 1040 Desservettaz, M., Pikridas, M., Stavroulas, I., Bougiatioti, A., Liakakou, E., Hatzianastassiou, N., Sciare, J., Mihalopoulos, N., and Boursoukoudis, E.: Emission of volatile organic compounds from residential biomass burning and their rapid chemical transformations, *Science of The Total Environment*, 903, 166592, <https://doi.org/10.1016/j.scitotenv.2023.166592>, 2023.
- 1045 Dimitriou, K. and Kassomenos, P.: Background concentrations of benzene, potential long range transport influences and corresponding cancer risk in four cities of central Europe, in relation to air mass origination, *J Environ Manage*, 262, 110374, <https://doi.org/10.1016/j.jenvman.2020.110374>, 2020.
- 1050 Dorokhov, Y. L., Sheshukova, E. V., and Komarova, T. V.: Methanol in Plant Life, *Front Plant Sci*, 9, <https://doi.org/10.3389/fpls.2018.01623>, 2018.
- Draxler, R. R. and H. G. D.: Description of the HYSPLIT_4 modeling system., Silver Spring, 24 pp., 1997.
- Edtbauer, A., Stöner, C., Pfannerstill, E. Y., Berasategui, M., Walter, D., Crowley, J. N., Lelieveld, J., and Williams, J.: A new marine biogenic emission: methane sulfonamide (MSAM), dimethyl sulfide (DMS), and dimethyl sulfone (DMSO₂) measured in air over the Arabian Sea, *Atmos Chem Phys*, 20, 6081–6094, <https://doi.org/10.5194/acp-20-6081-2020>, 2020.

[European Forest Fire Information System \(EFFIS\), European Commission, Copernicus Emergency Management Service: Annual fire reports, available at: https://forest-fire.emergency.copernicus.eu/reports-and-publications/annual-fire-reports, last access: 27 November 2025.](https://forest-fire.emergency.copernicus.eu/reports-and-publications/annual-fire-reports)

- 1055 Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurtén, T., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., Canagaratna, M., Maso, M. D., Berndt, T., Petäjä, T., Wahner, A., Kerminen, V.-M., Kulmala, M., Worsnop, D. R., Wildt, J., and Mentel, T. F.: A large source of low-volatility secondary organic aerosol, *Nature*, 506, 476–479, <https://doi.org/10.1038/nature13032>, 2014.
- 1060 Emekwuru, N. and Ejorwomu, O.: Temperature, Humidity and Air Pollution Relationships during a Period of Rainy and Dry Seasons in Lagos, West Africa, *Climate*, 11, 113, <https://doi.org/10.3390/cli11050113>, 2023.
- 1065 Fortunati, A., Barta, C., Brilli, F., Centritto, M., Zimmer, I., Schnitzler, J., and Loreto, F.: Isoprene emission is not temperature-dependent during and after severe drought-stress: a physiological and biochemical analysis, *The Plant Journal*, 55, 687–697, <https://doi.org/10.1111/j.1365-313X.2008.03538.x>, 2008.
- 1070 Gelencsér, A., Siszler, K., and Hlavay, J.: Toluene–Benzene Concentration Ratio as a Tool for Characterizing the Distance from Vehicular Emission Sources, *Environ Sci Technol*, 31, 2869–2872, <https://doi.org/10.1021/es970004c>, 1997.
- Georgiou, G. K., Kushta, J., Christoudias, T., Proestos, Y., and Lelieveld, J.: Air quality modelling over the Eastern Mediterranean: Seasonal sensitivity to anthropogenic emissions, *Atmos Environ*, 222, 117119, <https://doi.org/10.1016/j.atmosenv.2019.117119>, 2020.
- 1075 Georgiou, G. K., Christoudias, T., Proestos, Y., Kushta, J., Pikridas, M., Sciare, J., Savvides, C., and Lelieveld, J.: Evaluation of WRF-Chem model (v3.9.1.1) real-time air quality forecasts over the Eastern Mediterranean, *Geosci Model Dev*, 15, 4129–4146, <https://doi.org/10.5194/gmd-15-4129-2022>, 2022.
- 1080 Germain-Piaulenne, E., Paris, J.-D., Gros, V., Quéhé, P.-Y., Pikridas, M., Baisnée, D., Berchet, A., Sciare, J., and Bourtsoukidis, E.: Middle East oil and gas methane emissions signature captured at a remote site using light hydrocarbon tracers, *Atmos Environ X*, 22, 100253, <https://doi.org/10.1016/j.aeaoa.2024.100253>, 2024.
- 1085 Gjesteland, I., Hollund, B. E., Kirkeleit, J., Daling, P. S., Sørheim, K. R., and Bråtveit, M.: Determinants of airborne benzene evaporating from fresh crude oils released into seawater, *Mar Pollut Bull*, 140, 395–402, <https://doi.org/10.1016/j.marpolbul.2018.12.045>, 2019.
- Gu, S., Guenther, A., and Faiola, C.: Effects of Anthropogenic and Biogenic Volatile Organic Compounds on Los Angeles Air Quality, *Environ Sci Technol*, 55, 12191–12201, <https://doi.org/10.1021/acs.est.1c01481>, 2021.
- 1090 Guo, H., Ling, Z. H., Simpson, I. J., Blake, D. R., and Wang, D. W.: Observations of isoprene, methacrolein (MAC) and methyl vinyl ketone (MVK) at a mountain site in Hong Kong, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2012JD017750>, 2012.
- 1095 Holzinger, R., Williams, J., Salisbury, G., Klüpfel, T., de Reus, M., Traub, M., Crutzen, P. J., and Lelieveld, J.: Oxygenated compounds in aged biomass burning plumes over the Eastern Mediterranean: evidence for strong secondary production of methanol and acetone, *Atmos Chem Phys*, 5, 39–46, <https://doi.org/10.5194/acp-5-39-2005>, 2005.

- Huang, X.-F., Zhang, B., Xia, S.-Y., Han, Y., Wang, C., Yu, G.-H., and Feng, N.: Sources of oxygenated volatile organic compounds (OVOCs) in urban atmospheres in North and South China, *Environmental Pollution*, 261, 114152, <https://doi.org/10.1016/j.envpol.2020.114152>, 2020.
- 1100 Huangfu, Y., Yuan, B., Wang, S., Wu, C., He, X., Qi, J., de Gouw, J., Warneke, C., Gilman, J. B., Wisthaler, A., Karl, T., Graus, M., Jobson, B. T., and Shao, M.: Revisiting Acetonitrile as Tracer of Biomass Burning in Anthropogenic-Influenced Environments, *Geophys Res Lett*, 48, <https://doi.org/10.1029/2020GL092322>, 2021.
- 1105 Kajos, M. K., Rantala, P., Hill, M., Hellén, H., Aalto, J., Patokoski, J., Taipale, R., Hoerger, C. C., Reimann, S., Ruuskanen, T. M., Rinne, J., and Petäjä, T.: Ambient measurements of aromatic and oxidized VOCs by PTR-MS and GC-MS: intercomparison between four instruments in a boreal forest in Finland, *Atmos Meas Tech*, 8, 4453–4473, <https://doi.org/10.5194/amt-8-4453-2015>, 2015.
- 1110 Kaltsonoudis, C., Kostenidou, E., Florou, K., Psichoudaki, M., and Pandis, S. N.: Temporal variability and sources of VOCs in urban areas of the eastern Mediterranean, *Atmos Chem Phys*, 16, 14825–14842, <https://doi.org/10.5194/acp-16-14825-2016>, 2016.
- Kamal, M. S., Razzak, S. A., and Hossain, M. M.: Catalytic oxidation of volatile organic compounds (VOCs) – A review, *Atmos Environ*, 140, 117–134, <https://doi.org/10.1016/j.atmosenv.2016.05.031>, 2016.
- 1115 Kiendler-Scharr, A., Wildt, J., Maso, M. D., Hohaus, T., Kleist, E., Mentel, T. F., Tillmann, R., Uerlings, R., Schurr, U., and Wahner, A.: New particle formation in forests inhibited by isoprene emissions, *Nature*, 461, 381–384, <https://doi.org/10.1038/nature08292>, 2009.
- 1120 Kleanthous, S., Vrekoussis, M., Mihalopoulos, N., Kalabokas, P., and Lelieveld, J.: On the temporal and spatial variation of ozone in Cyprus, *Science of The Total Environment*, 476–477, 677–687, <https://doi.org/10.1016/j.scitotenv.2013.12.101>, 2014.
- [Knote, C., Hodzic, A., and Jimenez, J. L.: The effect of dry and wet deposition of condensable vapors on secondary organic aerosol concentrations over the continental United States. *Atmos. Chem. Phys.*, 15, 1–18. <https://doi.org/10.5194/acp-15-1-2015>, 2015a.](https://doi.org/10.5194/acp-15-1-2015)
- 1125 [Knote, C., Tuccella, P., Curci, G., Emmons, L., Orlando, J. J., Madronich, S., et al.: Influence of the choice of gas-phase mechanism on predictions of key gaseous pollutants during the AQMEII phase-2 intercomparison. *Atmos. Environ.*, 115, 553–568. <https://doi.org/10.1016/j.atmosenv.2014.11.066>, 2015b.](https://doi.org/10.1016/j.atmosenv.2014.11.066)
- 1130 Koppmann, R.: Chemistry of Volatile Organic Compounds in the Atmosphere, in: *Hydrocarbons, Oils and Lipids: Diversity, Origin, Chemistry and Fate*, Springer International Publishing, Cham, 811–822, https://doi.org/10.1007/978-3-319-90569-3_24, 2020.
- Koppmann, R., von Czapiewski, K., and Reid, J. S.: A review of biomass burning emissions, part I: gaseous emissions of carbon monoxide, methane, volatile organic compounds, and nitrogen containing compounds, <https://doi.org/10.5194/acpd-5-10455-2005>, 24 October 2005.
- 1135 Lazoglou, G., Hadjinicolaou, P., Sofokleous, I., Bruggeman, A., and Zittis, G.: Climate change and extremes in the Mediterranean island of Cyprus: from historical trends to future projections, *Environ Res Commun*, 6, 095020, <https://doi.org/10.1088/2515-7620/ad7927>, 2024.
- 1140 Lee, J. D., Squires, F. A., Sherwen, T., Wilde, S. E., Cliff, S. J., Carpenter, L. J., Hopkins, J. R., Bauguitte, S. J., Reed, C., Barker, P., Allen, G., Bannan, T. J., Matthews, E., Mehra, A., Percival, C., Heard, D. E., Whalley, L. K., Ronnie, G. V., Seldon, S., Ingham, T., Keller, C. A., Knowland, K. E., Nisbet, E. G., and Andrews, S.: Ozone production and precursor emission from wildfires in

- Africa, *Environmental Science: Atmospheres*, 1, 524–542, <https://doi.org/10.1039/D1EA00041A>, 2021.
- 1145 Lelieveld, J., Berresheim, H., Borrmann, S., Crutzen, P. J., Dentener, F. J., Fischer, H., Feichter, J., Flatau, P. J., Heland, J., Holzinger, R., Kormann, R., Lawrence, M. G., Levin, Z., Markowicz, K. M., Mihalopoulos, N., Minikin, A., Ramanathan, V., de Reus, M., Roelofs, G. J., Scheeren, H. A., Sciare, J., Schlager, H., Schultz, M., Siegmund, P., Steil, B., Stephanou, E. G., Stier, P., Traub, M., Warneke, C., Williams, J., and Ziereis, H.: Global Air Pollution Crossroads over the Mediterranean, *Science* (1979), 298, 794–799, <https://doi.org/10.1126/science.1075457>, 2002.
- 1150 Liakakou, E., Vrekoussis, M., Bonsang, B., Donousis, Ch., Kanakidou, M., and Mihalopoulos, N.: Isoprene above the Eastern Mediterranean: Seasonal variation and contribution to the oxidation capacity of the atmosphere, *Atmos Environ*, 41, 1002–1010, <https://doi.org/10.1016/j.atmosenv.2006.09.034>, 2007.
- 1155 Liakakou, E., Bonsang, B., Williams, J., Kalivitis, N., Kanakidou, M., and Mihalopoulos, N.: C2–C8 NMHCs over the Eastern Mediterranean: Seasonal variation and impact on regional oxidation chemistry, *Atmos Environ*, 43, 5611–5621, <https://doi.org/10.1016/j.atmosenv.2009.07.067>, 2009.
- Liu, Y., Shao, M., Kuster, W. C., Goldan, P. D., Li, X., Lu, S., and Gouw, J. A. de: Source Identification of Reactive Hydrocarbons and Oxygenated VOCs in the Summertime in Beijing, *Environ Sci Technol*, 43, 75–81, <https://doi.org/10.1021/es801716n>, 2009.
- 1160 Matthew, O. J.: Estimation of diurnal patterns of global solar radiation, temperature, relative humidity and wind speed from daily datasets at a humid tropical location, *Agric For Meteorol*, 322, 109003, <https://doi.org/10.1016/j.agrformet.2022.109003>, 2022.
- Mecca, M., Todaro, L., D’Auria, M., Italiano, S. S. P., Sofo, A., and Ripullone, F.: Volatile Organic Compounds (VOCs) in Mediterranean Oak Forests of Hungarian Oak (*Quercus frainetto* Ten) Affected by Dieback Phenomena, *Forests*, 15, 1072, <https://doi.org/10.3390/f15061072>, 2024.
- 1165 Mellouki, A., Wallington, T. J., and Chen, J.: Atmospheric Chemistry of Oxygenated Volatile Organic Compounds: Impacts on Air Quality and Climate, *Chem Rev*, 115, 3984–4014, <https://doi.org/10.1021/cr500549n>, 2015.
- 1170 Misztal, P. K., Hewitt, C. N., Wildt, J., Blande, J. D., Eller, A. S. D., Fares, S., Gentner, D. R., Gilman, J. B., Graus, M., Greenberg, J., Guenther, A. B., Hansel, A., Harley, P., Huang, M., Jardine, K., Karl, T., Kaser, L., Keutsch, F. N., Kiendler-Scharr, A., Kleist, E., Lerner, B. M., Li, T., Mak, J., Nölscher, A. C., Schnitzhofer, R., Sinha, V., Thornton, B., Warneke, C., Wegener, F., Werner, C., Williams, J., Worton, D. R., Yassaa, N., and Goldstein, A. H.: Atmospheric benzenoid emissions from plants rival those from fossil fuels, *Sci Rep*, 5, 12064, <https://doi.org/10.1038/srep12064>, 2015.
- 1175 Monson, R. K., Jaeger, C. H., Adams, W. W., Driggers, E. M., Silver, G. M., and Fall, R.: Relationships among Isoprene Emission Rate, Photosynthesis, and Isoprene Synthase Activity as Influenced by Temperature, *Plant Physiol*, 98, 1175–1180, <https://doi.org/10.1104/pp.98.3.1175>, 1992.
- 1180 Mukherjee, S., Pandithurai, G., Waghmare, V., Mahajan, A. S., Tinel, L., Aslam, M. Y., Meena, G. S., Patil, S., Buchunde, P., and Kumar, A.: Seasonal variability of volatile organic compounds (VOCs) at a high-altitude station in the Western Ghats, India: Influence of biogenic, anthropogenic emissions and long-range transport, *Atmos Environ*, 331, 120598, <https://doi.org/10.1016/j.atmosenv.2024.120598>, 2024.
- 1185 Pagonis, D., Sekimoto, K., and de Gouw, J.: A Library of Proton-Transfer Reactions of H_3O^+ Ions Used for Trace Gas Detection, *J Am Soc Mass Spectrom*, 30, 1330–1335, <https://doi.org/10.1007/s13361-019-02209-3>, 2019.

- 1190 Panopoulou, A., Liakakou, E., Gros, V., Sauvage, S., Locoge, N., Bonsang, B., Psiloglou, B. E., Gerasopoulos, E., and Mihalopoulos, N.: Non-methane hydrocarbon variability in Athens during wintertime: the role of traffic and heating, *Atmos Chem Phys*, 18, 16139–16154, <https://doi.org/10.5194/acp-18-16139-2018>, 2018.
- Panopoulou, A., Liakakou, E., Sauvage, S., Gros, V., Locoge, N., Stavroulas, I., Bonsang, B., Gerasopoulos, E., and Mihalopoulos, N.: Yearlong measurements of monoterpenes and isoprene in a Mediterranean city (Athens): Natural vs anthropogenic origin, *Atmos Environ*, 243, 117803, <https://doi.org/10.1016/j.atmosenv.2020.117803>, 2020.
- 1195 Paton-Walsh, C., Wilson, S. R., Jones, N. B., and Griffith, D. W. T.: Measurement of methanol emissions from Australian wildfires by ground-based solar Fourier transform spectroscopy, *Geophys Res Lett*, 35, <https://doi.org/10.1029/2007GL032951>, 2008.
- 1200 Pennington, E. A., Seltzer, K. M., Murphy, B. N., Qin, M., Seinfeld, J. H., and Pye, H. O. T.: Modeling secondary organic aerosol formation from volatile chemical products, *Atmos Chem Phys*, 21, 18247–18261, <https://doi.org/10.5194/acp-21-18247-2021>, 2021.
- Peñuelas, J. and Staudt, M.: BVOCs and global change, *Trends Plant Sci*, 15, 133–144, <https://doi.org/10.1016/j.tplants.2009.12.005>, 2010.
- 1205 Pérez, I. A., García, M. Á., Sánchez, M. L., Pardo, N., and Fernández-Duque, B.: Key Points in Air Pollution Meteorology, *Int J Environ Res Public Health*, 17, 8349, <https://doi.org/10.3390/ijerph17228349>, 2020.
- Perrone, M. G., Carbone, C., Faedo, D., Ferrero, L., Maggioni, A., Sangiorgi, G., and Bolzacchini, E.: Exhaust emissions of polycyclic aromatic hydrocarbons, n-alkanes and phenols from vehicles coming within different European classes, *Atmos Environ*, 82, 391–400, <https://doi.org/10.1016/j.atmosenv.2013.10.040>, 2014.
- 1210 Pikridas, M., Vrekoussis, M., Sciare, J., Kleanthous, S., Vasiliadou, E., Kizas, C., Savvides, C., and Mihalopoulos, N.: Spatial and temporal (short and long-term) variability of submicron, fine and sub-10 μm particulate matter (PM₁, PM_{2.5}, PM₁₀) in Cyprus, *Atmos Environ*, 191, 79–93, <https://doi.org/10.1016/j.atmosenv.2018.07.048>, 2018.
- 1215 Pinthong, N., Thepanondh, S., and Kondo, A.: Source Identification of VOCs and their Environmental Health Risk in a Petrochemical Industrial Area, *Aerosol Air Qual Res*, 22, 210064, <https://doi.org/10.4209/aaqr.210064>, 2022.
- Pochanart, P., Akimoto, H., Kajii, Y., Potemkin, V. M., and Khodzher, T. V.: Regional background ozone and carbon monoxide variations in remote Siberia/East Asia, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2001JD001412>, 2003.
- 1220 Pugliese, G., Ingrisich, J., Meredith, L. K., Pfannerstill, E. Y., Klüpfel, T., Meeran, K., Byron, J., Pursler, G., Gil-Loaiza, J., van Haren, J., Dontsova, K., Kreuzwieser, J., Ladd, S. N., Werner, C., and Williams, J.: Effects of drought and recovery on soil volatile organic compound fluxes in an experimental rainforest, *Nat Commun*, 14, 5064, <https://doi.org/10.1038/s41467-023-40661-8>, 2023.
- 1225 Ralf Koppmann: *Volatile Organic Compounds in the Atmosphere*, edited by: Koppmann, R., Wiley, <https://doi.org/10.1002/9780470988657>, 2007.
- Robles, H.: Acetonitrile, in: *Encyclopedia of Toxicology*, Elsevier, 28–30, <https://doi.org/10.1016/B0-12-369400-0/00015-6>, 2005.
- 1230 Salisbury, G., Williams, J., Holzinger, R., Gros, V., Mihalopoulos, N., Vrekoussis, M., Sarda-Estève, R., Berresheim, H., von Kuhlmann, R., Lawrence, M., and Lelieveld, J.: Ground-based

- PTR-MS measurements of reactive organic compounds during the MINOS campaign in Crete, July–August 2001, *Atmos Chem Phys*, 3, 925–940, <https://doi.org/10.5194/acp-3-925-2003>, 2003.
- 1235 Schieweck, A., Uhde, E., and Salthammer, T.: Determination of acrolein in ambient air and in the atmosphere of environmental test chambers, *Environ Sci Process Impacts*, 23, 1729–1746, <https://doi.org/10.1039/D1EM00221J>, 2021.
- Sciare J: The Agia Marina Xyliatou Observatory: A remote supersite in Cyprus to monitor changes in the atmospheric composition of the Eastern Mediterranean and the Middle East, in: EGU General Assembly Conference Abstracts, 2016.
- 1240 Sicre, M. A., Marty, J. C., Saliot, A., Aparicio, X., Grimalt, J., and Albaiges, J.: Aliphatic and Aromatic Hydrocarbons in the Mediterranean Aerosol, *Int J Environ Anal Chem*, 29, 73–94, <https://doi.org/10.1080/03067318708078412>, 1987.
- 1245 Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrou, T., Müller, J.-F., Kuhn, U., Stefani, P., and Knorr, W.: Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years, *Atmos Chem Phys*, 14, 9317–9341, <https://doi.org/10.5194/acp-14-9317-2014>, 2014.
- Song, W., Williams, J., Yassaa, N., Martinez, M., Camero, J. A. A., Hidalgo, P. J., Bozem, H., and Lelieveld, J.: Winter and summer characterization of biogenic enantiomeric monoterpenes and anthropogenic BTEX compounds at a Mediterranean Stone Pine forest site, *J Atmos Chem*, 68, 233–250, <https://doi.org/10.1007/s10874-012-9219-4>, 2011.
- 1250 Soukissian, T. and Sotiriou, M.-A.: Long-Term Variability of Wind Speed and Direction in the Mediterranean Basin, *Wind*, 2, 513–534, <https://doi.org/10.3390/wind2030028>, 2022.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bull Am Meteorol Soc*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- 1255 Stockwell, W. R., Kirchner, F., Kuhn, M., and Seefeld, S.: A new mechanism for regional atmospheric chemistry modeling, *Journal of Geophysical Research: Atmospheres*, 102, 25847–25879, <https://doi.org/10.1029/97JD00849>, 1997.
- 1260 Strada, S., Pozzer, A., Giuliani, G., Coppola, E., Solmon, F., Jiang, X., Guenther, A., Bourtsoukidis, E., Serça, D., Williams, J., and Giorgi, F.: Assessment of isoprene and near-surface ozone sensitivities to water stress over the Euro-Mediterranean region, *Atmos Chem Phys*, 23, 13301–13327, <https://doi.org/10.5194/acp-23-13301-2023>, 2023.
- Thunis, P., Crippa, M., Cuvelier, C., Guizzardi, D., de Meij, A., Oreggioni, G., and Pisoni, E.: Sensitivity of air quality modelling to different emission inventories: a case study over Europe, *Atmos. Environ. X*, 10, 100111, 2021.
- 1265 Tiwari, V., Hanai, Y., and Masunaga, S.: Ambient levels of volatile organic compounds in the vicinity of petrochemical industrial area of Yokohama, Japan, *Air Qual Atmos Health*, 3, 65–75, <https://doi.org/10.1007/s11869-009-0052-0>, 2010.
- 1270 Torres-Vinces, L., Contreras-Zarazua, G., Huerta-Rosas, B., Sánchez-Ramírez, E., and Segovia-Hernández, J. G.: Methyl Ethyl Ketone Production through an Intensified Process, *Chem Eng Technol*, 43, 1433–1441, <https://doi.org/10.1002/ceat.201900664>, 2020.
- Traub, M., Fischer, H., de Reus, M., Kormann, R., Heland, H., Ziereis, H., Schlager, H., Holzinger, R., Williams, J., Warneke, C., de Gouw, J., and Lelieveld, J.: Chemical characteristics assigned to trajectory clusters during the MINOS campaign, *Atmos Chem Phys*, 3, 459–468, <https://doi.org/10.5194/acp-3-459-2003>, 2003.

- 1275 Vlasenko, A., Macdonald, A. M., Sjostedt, S. J., and Abbatt, J. P. D.: Formaldehyde measurements by Proton transfer reaction – Mass Spectrometry (PTR-MS): correction for humidity effects, *Atmos Meas Tech*, 3, 1055–1062, <https://doi.org/10.5194/amt-3-1055-2010>, 2010.
- 1280 Vrekoussis, M., Pikridas, M., Rousogenous, C., Christodoulou, A., Desservettaz, M., Sciare, J., Richter, A., Bougoudis, I., Savvides, C., and Papadopoulos, C.: Local and regional air pollution characteristics in Cyprus: A long-term trace gases observations analysis, *Science of The Total Environment*, 845, 157315, <https://doi.org/10.1016/j.scitotenv.2022.157315>, 2022.
- 1285 Wang, H., Ma, X., Tan, Z., Wang, H., Chen, X., Chen, S., Gao, Y., Liu, Y., Liu, Y., Yang, X., Yuan, B., Zeng, L., Huang, C., Lu, K., and Zhang, Y.: Anthropogenic monoterpenes aggravating ozone pollution, *Natl Sci Rev*, 9, <https://doi.org/10.1093/nsr/nwac103>, 2022a.
- Wang, L., Lun, X., Wang, Q., and Wu, J.: Biogenic volatile organic compounds emissions, atmospheric chemistry, and environmental implications: a review, *Environ Chem Lett*, 22, 3033–3058, <https://doi.org/10.1007/s10311-024-01785-5>, 2024.
- 1290 Wang, N., Edtbauer, A., Stöner, C., Pozzer, A., Bourtsoukidis, E., Ernle, L., Dienhart, D., Hottmann, B., Fischer, H., Schuladen, J., Crowley, J. N., Paris, J.-D., Lelieveld, J., and Williams, J.: Measurements of carbonyl compounds around the Arabian Peninsula: overview and model comparison, *Atmos Chem Phys*, 20, 10807–10829, <https://doi.org/10.5194/acp-20-10807-2020>, 2020.
- 1295 Wang, S., Yuan, B., Wu, C., Wang, C., Li, T., He, X., Huangfu, Y., Qi, J., Li, X.-B., Sha, Q., Zhu, M., Lou, S., Wang, H., Karl, T., Graus, M., Yuan, Z., and Shao, M.: Oxygenated volatile organic compounds (VOCs) as significant but varied contributors to VOC emissions from vehicles, *Atmos Chem Phys*, 22, 9703–9720, <https://doi.org/10.5194/acp-22-9703-2022>, 2022b.
- 1800 [Wang, H., Lu, X., Seco, R., Stavrakou, T., Karl, T., Jiang, X., et al.: Modeling isoprene emission response to drought and heatwaves within MEGAN using evapotranspiration data and by coupling with the Community Land Model, *J. Adv. Model. Earth Syst.*, 14, e2022MS003174, <https://doi.org/10.1029/2022MS003174>, 2022c.](https://doi.org/10.1029/2022MS003174)
- 1305 Xu, Z., Zou, Q., Jin, L., Shen, Y., Shen, J., Xu, B., Qu, F., Zhang, F., Xu, J., Pei, X., Xie, G., Kuang, B., Huang, X., Tian, X., and Wang, Z.: Characteristics and sources of ambient Volatile Organic Compounds (VOCs) at a regional background site, YRD region, China: Significant influence of solvent evaporation during hot months, *Science of The Total Environment*, 857, 159674, <https://doi.org/10.1016/j.scitotenv.2022.159674>, 2023.
- 1310 Yáñez-Serrano, A. M., Nölscher, A. C., Bourtsoukidis, E., Derstroff, B., Zannoni, N., Gros, V., Lanza, M., Brito, J., Noe, S. M., House, E., Hewitt, C. N., Langford, B., Nemitz, E., Behrendt, T., Williams, J., Artaxo, P., Andreae, M. O., and Kesselmeier, J.: Atmospheric mixing ratios of methyl ethyl ketone (2-butanone) in tropical, boreal, temperate and marine environments, *Atmos Chem Phys*, 16, 10965–10984, <https://doi.org/10.5194/acp-16-10965-2016>, 2016.
- 1315 Yáñez-Serrano, A. M., Filella, I., LLusià, J., Gargallo-Garriga, A., Granda, V., Bourtsoukidis, E., Williams, J., Seco, R., Cappellin, L., Werner, C., de Gouw, J., and Peñuelas, J.: GLOVOCS - Master compound assignment guide for proton transfer reaction mass spectrometry users, *Atmos Environ*, 244, 117929, <https://doi.org/10.1016/j.atmosenv.2020.117929>, 2021.
- 1320 Yokelson, R. J., Goode, J. G., Ward, D. E., Susott, R. A., Babbitt, R. E., Wade, D. D., Bertschi, I., Griffith, D. W. T., and Hao, W. M.: Emissions of formaldehyde, acetic acid, methanol, and other trace gases from biomass fires in North Carolina measured by airborne Fourier transform infrared spectroscopy, *Journal of Geophysical Research: Atmospheres*, 104, 30109–30125, <https://doi.org/10.1029/1999JD900817>, 1999.

1325 Yuan, Q., Zhang, Z., Chen, Y., Hui, L., Wang, M., Xia, M., Zou, Z., Wei, W., Ho, K. F., Wang, Z.,
Lai, S., Zhang, Y., Wang, T., and Lee, S.: Origin and transformation of volatile organic compounds
at a regional background site in Hong Kong: Varied photochemical processes from different
source regions, *Science of The Total Environment*, 908, 168316,
<https://doi.org/10.1016/j.scitotenv.2023.168316>, 2024.

1330 Zittis, G., Almazroui, M., Alpert, P., Ciais, P., Cramer, W., Dahdal, Y., Fnais, M., Francis, D.,
Hadjinicolaou, P., Howari, F., Jrrar, A., Kaskaoutis, D. G., Kulmala, M., Lazoglou, G.,
Mihalopoulos, N., Lin, X., Rudich, Y., Sciare, J., Stenchikov, G., Xoplaki, E., and Lelieveld, J.:
Climate Change and Weather Extremes in the Eastern Mediterranean and Middle East, *Reviews of
Geophysics*, 60, <https://doi.org/10.1029/2021RG000762>, 2022.