

# The influence of ammonia ~~emission~~emission inventories on the size-resolved global atmospheric aerosol composition and acidity

Xurong Wang<sup>1</sup>, Alexandra P. Tsimpidi<sup>1</sup>, Zhenqi Luo<sup>2</sup>, Benedikt Steil<sup>3</sup>, Andrea Pozzer<sup>3,4</sup>, Jos Lelieveld<sup>3,4</sup>, and Vlassis A. Karydis<sup>1</sup>

<sup>1</sup> Institute of Climate and Energy Systems: Troposphere (ICE-3), Forschungszentrum Jülich GmbH, Jülich, Germany

<sup>2</sup> School of Integrative Plant Science, Soil and Crop Sciences Section, Cornell University, Ithaca, NY 14853, United States of America

<sup>3</sup> Max Planck Institute for Chemistry, Atmospheric Chemistry Dept., Mainz, Germany.

<sup>4</sup> The Cyprus Institute, Climate and Atmosphere Research Center Nicosia, Nicosia, Cyprus.

Correspondence to:

Vlassis A. Karydis ([v.karydis@fz-juelich.de](mailto:v.karydis@fz-juelich.de), [karydis@fz-juelich.de](mailto:karydis@fz-juelich.de)) and Alexandra P. Tsimpidi ([a.tsimpidi@fz-juelich.de](mailto:a.tsimpidi@fz-juelich.de), [tsimpidi@fz-juelich.de](mailto:tsimpidi@fz-juelich.de))

**Abstract.** Ammonia (NH<sub>3</sub>) is an abundant alkaline gas in the atmosphere and a key precursor in the formation of particulate matter. While emissions of other aerosol precursors such as SO<sub>2</sub> and NO<sub>x</sub> have decreased significantly, global NH<sub>3</sub> emissions are stable or increasing, and this trend is projected to continue. This study investigates the impact of NH<sub>3</sub> emission changes on size-resolved aerosol composition and acidity using the atmospheric chemistry-climate model EMAC. ~~Three~~Rather than directly perturbing NH<sub>3</sub> emissions, we employ three distinct emission schemes are analyzed in inventories: two bottom-up inventories and one derived using an updated top-down method. The results reveal that sulphate-nitrate-ammonium aerosols in two fine mode size ranges (0–1 μm and 1–2.5 μm) show the greatest sensitivity to NH<sub>3</sub> emission changes. Regional responses vary depending on the local chemical environment of secondary inorganic aerosols. In 'NH<sub>3</sub>-rich' regions (e.g., East Asia and Europe), the abundance of NH<sub>3</sub> partially offsets the effects of reduced NH<sub>3</sub> emissions when NO<sub>x</sub> and SO<sub>2</sub> are available, especially for aerosols in the 1–2.5 μm range. This highlightsunderscores the importance of coordinated control strategies for NH<sub>3</sub>, NO<sub>x</sub> and SO<sub>2</sub> emissions. Further, we find that NH<sub>3</sub> has a buffering effect in densely populated areas, maintaining aerosol acidity at moderate levels and mitigating drastic pH shifts. ~~The study emphasizes that~~While pH changes are closely related to NH<sub>3</sub> emission variations, correlate strongly with the highest sensitivity observed NH<sub>3</sub> variability, they are also influenced by concurrent changes in the fine mode size ranges SO<sub>2</sub> and NO<sub>x</sub> emissions. These results highlight the critical role of NH<sub>3</sub> in shaping aerosol acidity ~~and argue, arguing~~ for size-specific approaches to managing particulate matter.

## 1. Introduction

As an abundant alkaline gas in the atmosphere, ammonia (NH<sub>3</sub>) acts as a precursor in the formation of particulate matter by neutralizing atmospheric acids (e.g., H<sub>2</sub>SO<sub>4</sub>, HNO<sub>3</sub>) to form sulfate-nitrate-ammonium (SNA) aerosols (Li et al., 2018; Chen et al., 2016; Wang et al., 2013), which are the main secondary inorganic components of PM<sub>2.5</sub> (particulate matter with a diameter of 2.5 μm or less). By condensing onto freshly nucleated particles, NH<sub>3</sub> enhances the growth rate of new particles as well as their hydrophilicity (Wang et al., 2020; Li et al., 2018). This can degrade

Formatted: Superscript

Formatted: Font: Italic

Formatted: Justified

Formatted: Font color: Auto

40 air quality and change the solar radiative balance by interaction with radiation and clouds (Che et al., 2009; Zhao et al., 2011; Yao et al., 2018). In addition, the conjugate base-acid pair  $\text{NH}_3/\text{NH}_4^+$  acts as the major buffer that inhibits changes in aerosol acidity (Chen et al., 2019; Zheng et al., 2020; Karydis et al., 2021).

Anthropogenic emissions are the main source of atmospheric  $\text{NH}_3$ , with an average global contribution of 76%, dominated by agricultural activities, including livestock farming and the fertilization of soils (Schlesinger and Hartley, 45 1992; Dentener and Crutzen, 1994; Bouwman et al., 1997; Olivier et al., 1998; Van Aardenne et al., 2001; Bleeker et al., 2013). Meanwhile, the importance of non-agricultural sources, such as industrial emissions and fossil fuel combustion, has been highlighted by studies focusing on severe haze episodes in East Asia (Chang et al., 2019; Liu et al., 2018a; Pan et al., 2016). The construction of high-resolution regional or global datasets has characterized the spatiotemporal patterns of  $\text{NH}_3$  emissions, with the most common datasets consisting of bottom-up inventories and 50 top-down modeling inversion methods. Bottom-up emission inventories rely on activity data and emission factors, the latter being sensitive to assumptions about fertilizer types, local soil and meteorological properties (Bouwman et al., 2002; Sogaard et al., 2002; Xu et al., 2024). Zhang et al. (2018) evaluated discrepancies between bottom-up  $\text{NH}_3$  emission inventories to exceed a factor of two, due to the uncertainties in emission factors, meteorological properties, and agricultural statistics (Beusen et al., 2008). Crippa et al. (2018) pointed out that the uncertainty in estimated  $\text{NH}_3$  55 emissions is largest among all pollutants in the Emissions Database for Global Atmospheric Research (EDGAR v4.3.2), with a range of variation from 186% to 294% in 2012. In contrast, the ability of satellites to measure  $\text{NH}_3$  abundance combined with numerical simulations allows a better characterization of the spatial distribution and seasonality of  $\text{NH}_3$  emissions, but the low signal-to-noise ratio over low-emission areas limits the accuracy of retrieval products (Morán et al., 2016; Xu et al., 2016; Kong et al., 2019). Using a hybrid inverse modeling approach, Chen et al. (2021) optimized the  $\text{NH}_3$  emission inventory NEI over the United States by combining CMAQ model simulations 60 with constraints from the Infrared Atmospheric Sounding Interferometer (IASI) retrieval product. They found a 26% low bias of  $\text{NH}_3$  emissions in the NEI, and the optimized  $\text{NH}_3$  emission inventory improved the model performance of  $\text{PM}_{2.5}$  mass concentration in the Midwest US, and the normalized mean bias of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  decreased from 27% to 22%, and 64% to 55%, respectively.

65 Recently, the emissions of  $\text{SO}_2$  and  $\text{NO}_x$  have decreased due to the implementation of related clean air policies in East Asia, North America and Europe (Zheng et al., 2018; Hand et al., 2012; Russell et al., 2012; Aas et al., 2019; Gong et al., 2024), while the emissions of  $\text{NH}_3$  remained stable or have slightly increased in major agricultural regions such as China (Xu et al., 2016), the United States (Yu et al., 2018), and Europe (Fortems-Cheiney et al., 2022). The increase in  $\text{NH}_3$  emissions is associated with increasing fertilizer use and local temperatures (Warner et al., 2017; Xu et al., 2016; Skjøth and Geels, 2013; Xu et al., 2019), and such trends are expected to continue on a global scale 70 throughout the century (IPCC, 2013). Driven by such emission ~~trend~~trends of  $\text{SO}_2$ ,  $\text{NO}_x$ , and  $\text{NH}_3$ , the main composition of SNA has changed in large regions with a shift from an ammonium sulfate to an ammonium nitrate formation regime (Tsimplidi et al., 2024; Lei et al., 2021; Jo et al., 2020; Zhou et al., 2019; Shah et al., 2018; Wang et al., 2013; Hauglustaine et al., 2014). Li et al. (2017) evaluated that during the period 1989 to 2013, the increasing 75 trend of sulfate ( $\text{SO}_4^{2-}$ ) and ammonium ( $\text{NH}_4^+$ ) mass concentration in India and China occurred at a rate  $>0.1 \mu\text{g m}^{-3} \text{yr}^{-1}$ , while decreasing trends were found in North America and Europe at a rate of about  $0.1 \mu\text{g m}^{-3} \text{yr}^{-1}$ . The reduction

of  $\text{SO}_2$  and  $\text{NO}_x$  results in excess  $\text{NH}_3$  being released to the atmosphere because less  $\text{NH}_3$  is required to neutralize emissions decreases the atmospheric availability of  $\text{H}_2\text{SO}_4$  and  $\text{HNO}_3$ , which in turn hinders the formation thereby reducing the demand for  $\text{NH}_3$  to neutralize these acids and form particulate  $\text{NH}_4^+$ . As  $\text{NH}_3$  emissions have remained stable or slightly increased, this leads to an accumulation of  $\text{NH}_4^+\text{NH}_3$  in the aerosol/gas phase and increases the atmospheric  $\text{NH}_3$  concentration (Liu et al., 2018b). Warner et al. (2017) assessed that the average increase rate of  $\text{NH}_3$  mass concentration from 2002 to 2016 in the United States, Europe, and China was  $2.6\% \text{ yr}^{-1}$ ,  $1.8\% \text{ yr}^{-1}$ , and  $2.3\% \text{ yr}^{-1}$ , respectively. Some studies further pointed out that the increase in  $\text{NH}_3$  concentration may offset the effectiveness of  $\text{PM}_{2.5}$  control achieved via  $\text{SO}_2$  and  $\text{NO}_x$  emission reduction by promoting the formation of nitrate ( $\text{NO}_3^-$ ) (Huang et al., 2021; Cai et al., 2017; Zhang and Geng, 2019; Fu et al., 2017).

Several studies concluded that reducing  $\text{NH}_3$  emissions would be a cost-effective way to control  $\text{PM}_{2.5}$  concentrations (Gu et al., 2021; Tsimpidi et al., 2007; Erisman and Schaap, 2004). However, the response of SNA to changes in its precursors is not linear (Pozzer et al., 2017; Wang et al., 2011; Wang et al., 2013; Tsimpidi et al., 2007; West et al., 1999), because the gas-particle partitioning of  $\text{NH}_3/\text{NH}_4^+$  and  $\text{HNO}_3/\text{NO}_3^-$  is influenced by several parameters, such as temperature, liquid water content, and aerosol acidity (Xu et al., 2020; Nenes et al., 2020; Guo et al., 2018). Nenes et al. (2020) developed a conceptual framework to describe the sensitivity of particulate matter to  $\text{NO}_x$  and  $\text{NH}_3$  emissions, highlighting the critical influence of aerosol acidity and liquid water content on particulate matter formation. Based on sensitivity tests, Guo et al. (2018) evaluated that the response of  $\text{NO}_3^-$  to  $\text{NH}_3$  reduction shows an apparent decrease only when the aerosol pH falls below the value of 3.

On the other hand, reductions in  $\text{SO}_2$  and  $\text{NO}_x$  emissions are expected to reduce aerosol acidity, but recent studies revealed that aerosol acidity does not decrease as expected (Chen et al., 2019; Guo et al., 2017a; Zheng et al., 2022; Karydis et al., 2021). Aerosol acidity affects many processes involving the atmosphere and various aspects of the Earth system (Pye et al., 2020; Tilgner et al., 2021; Karydis et al., 2021) as well as human health (Dockery et al., 1993; Dockery et al., 1996; Thurston et al., 1994; Spengler et al., 1996). Weber et al. (2016) found that the acidity of  $\text{PM}_{2.5}$  in the southeastern United States remained at a relatively constant level with a pH value of 0 – 2 over the past 15 years, despite a 70% reduction in  $\text{SO}_4^{2-}$  concentration. A lack of aerosol acidity trend was also reported in China (Zhou et al., 2022). This is mainly caused by the buffering effect of  $\text{NH}_3/\text{NH}_4^+$  (Chen et al., 2019; Zheng et al., 2022; Zheng et al., 2020). To investigate this, Song et al. (2019) derived an equation from the partitioning of  $\text{NH}_3$  and estimated that a unit increase in pH requires a tenfold increase in  $\text{NH}_3$  concentration, which is consistent with the findings of Guo et al. (2017a). The regional variation of aerosol acidity is considerable, and the pH of  $\text{PM}_{2.5}$  in northern China is in the range of 4 to 5, which is higher than in Europe and the United States (Shi et al., 2019; Shi et al., 2017; Liu et al., 2017; Guo et al., 2015; Guo et al., 2016; Guo et al., 2017b; Karydis et al., 2021). This is caused by multiple driving factors, including aerosol mass concentration and composition,  $\text{NH}_3$  mass concentration, aerosol water content, and meteorological factors (Ding et al., 2019; Zhang et al., 2021a). However, the main driver for the difference in aerosol acidity remains controversial. Zheng et al. (2020) pointed out that aerosol water content is the most important factor causing the regional variation of aerosol acidity, while Zhang et al. (2021a) emphasized the equal importance of aerosol mass concentration and chemical composition for the aerosol acidity contrasts between China and the United States.

Formatted: English (United States)

Formatted: English (United States), Not Superscript/ Subscript

115 Almost all recent studies that discuss the response of aerosol composition and acidity to changes in NH<sub>3</sub> emission trends focus on the fine mode (e.g. PM<sub>2.5</sub>). The size-resolved composition of SNA is not uniform (Karydis et al., 2016; Fang et al., 2017; Karydis et al., 2011; Karydis et al., 2010; Guo et al., 2017b), and NH<sub>4</sub><sup>+</sup> and SO<sub>4</sub><sup>2-</sup> are mainly concentrated in the fine mode (Wang et al., 2012; Seinfeld and Pandis, 2016), while NO<sub>3</sub><sup>-</sup> aerosol can be formed on the surface of super-micron particles via heterogeneous chemistry (Allen et al., 2015). Furthermore, Milousis et al. (2024) revealed that the acidity of fine-mode aerosol is more sensitive to NH<sub>3</sub> emission than coarse-mode aerosol. 120 Reducing the NH<sub>3</sub> emissions by half, the simulated pH of fine and coarse mode aerosol decreased by up to 3 and 1.5 units, respectively. Aerosol acidity tends to decrease with increasing particle size, with pH varying up to 6 units (Craig et al., 2018; Fang et al., 2017; Bougiatioti et al., 2016). Size-resolved aerosol acidity is associated with different formation pathways (Tilgner et al., 2021; Cheng et al., 2016). Ding et al. (2019) found that the coarse-mode aerosol shifted from neutral to weakly acidic with the increase of NO<sub>3</sub><sup>-</sup> and SO<sub>4</sub><sup>2-</sup> during severe hazy days. Cheng et al. (2016) 125 further pointed out that the dominant oxidant in SO<sub>4</sub><sup>2-</sup> production by SO<sub>2</sub> oxidation changes with the ambient aerosol acidity. Therefore, it is necessary to comprehensively investigate the response of size-resolved chemical composition and acidity to changes in NH<sub>3</sub> emissions.

In this study, ~~three different we investigate the influence of ammonia (NH<sub>3</sub> emission schemes are used as input to the atmospheric chemistry-climate model) emissions on size-resolved aerosol composition and acidity using the EMAC (ECHAM5/MESSy Atmospheric Chemistry). The three emission schemes include two bottom-up model. The EMAC model, coupled with the ISORROPIA-II thermodynamic module, provides a robust framework for simulating global aerosol-chemistry-climate interactions and estimating aerosol pH under varying environmental conditions. Rather than directly perturbing NH<sub>3</sub> emissions, we employ three distinct emission inventories: two bottom-up inventories (CAM5 and CEDS\_GBD) and one derived using an updated emission inventory produced following a top-down method approach (Luo et al., 2022). Satellite retrievals of NH<sub>3</sub> column concentrations and aerosol composition observational datasets from multiple sites around the world are used to evaluate the aerosol simulations derived from these scenarios allow us to assess the three NH<sub>3</sub> emission schemes. We investigate the responsesensitivity of size-resolved aerosol SNA composition and acidity to changes aerosols to realistic variations in NH<sub>3</sub> emissions across three well-characterized, anthropogenically polluted regions in the Northern Hemisphere. These regions, representing a gradient from relatively less to more polluted conditions, include the United States, in conjunction with co-varying SO<sub>2</sub> and NO<sub>x</sub> trends. Model results are evaluated against satellite-derived NH<sub>3</sub> columns and ground-based aerosol composition observations. Our analysis spans four aerosol size ranges (0–1 μm, 1–2.5 μm, 2.5–5 μm, and 5–10 μm) and includes regional assessments over North America, Europe, and the North China Plain. The examined aerosol sizes range (diameter) from sub-micron (0–1 μm) to super-micron (1–2.5 μm, 2.5–5 μm, and 5–10 μm). East Asia to explore chemical regime shifts and buffering capacity.~~ 130 135 140 145

## 2. Modelling description

EMAC (ECHAM5/MESSy) is a global atmospheric chemistry and climate model, which includes a number of submodels describing atmospheric processes and interactions among oceans, land, and anthropogenic influences (Jöckel et al., 2016). These submodels are linked to the base model, the 5th generation European Centre Hamburg

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Font: +Body (Aptos)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

general circulation model (ECHAM5; Roeckner et al., 2006), via the Modular Earth Submodel System (MESSy; Jöckel et al., 2005). In this study, the horizontal resolution of the EMAC model is T63L31, which corresponds to a grid resolution of about  $1.875^\circ \times 1.875^\circ$  (Jöckel et al., 2010) and 31 vertical layers extending up to 25 km altitude. EMAC is applied for 4 years, from 2009 to 2012 with the first year used as a spin-up. The meteorological reanalysis data ERA5 (Hersbach et al., 2020), with a significantly enhanced horizontal resolution of 31 km and hourly output throughout, is used in EMAC to nudge the simulation.

In EMAC, organic aerosol (OA) formation is simulated by the ORACLE module (Tsimpidi et al., 2014; Tsimpidi et al., 2018), where logarithmically spaced saturation concentration bins are used to describe the organic aerosol components based on their volatility. The aerosol microphysics and gas/aerosol partitioning are calculated by the Global Modal-aerosol eXtension (GMXe) module described by Pringle et al. (2010), which has the same microphysical core as the M7 sub-model (Vignati et al., 2004). The aerosol size distribution is treated by 7 log-normal modes, including 4 hydrophilic and 3 hydrophobic modes, covering nucleation (soluble only), Aitken, accumulation, and coarse modes (both soluble and insoluble). To determine size-resolved aerosol composition and pH, we sum the contributions of each aerosol component, water content, and  $H^+$  concentration across all GMXe modes corresponding to a given size range. This is achieved by calculating the volume fraction of the lognormal distribution of each mode that falls within the specified size limits. The atmospheric chemistry module MECCA (Module Efficiently Calculating the Chemistry of the Atmosphere), which contains a comprehensive atmospheric reaction is used to calculate the gas concentrations (Sander et al., 2019). The SEDI module is used to compute aerosol particle sedimentation (Kerkweg et al., 2006). Dry deposition and wet deposition of gas and particle species are calculated by the DRYDEP module (Kerkweg et al., 2006) and the SCAV module (Tost et al., 2006), respectively. The CLOUD submodel (Roeckner et al., 2006) is used to calculate cloud properties and microphysics, utilizing the microphysical scheme of Lohmann and Ferrachat (2010) and a physically based treatment of liquid (Karydis et al., 2017) and ice crystal (Bacer et al., 2018) activation processes.

An advanced parameterization scheme is incorporated into the EMAC model to calculate the dust emission flux online (Astitha et al., 2012). The scheme uses the online meteorological fields from the EMAC model, such as temperature, pressure, relative humidity, soil moisture, and surface friction velocity, to calculate the threshold friction velocity which is the initial step of dust production. Above the threshold friction velocity, dust emission is possible. Following Karydis et al. (2016), the emissions of individual crustal species in this study are estimated as constant fractions of the dust emission (Klingmüller et al., 2018). These fractional factors depend on the geological information, which includes different dust emission sources. Karydis et al. (2021) pointed out that the crustal ions ( $Ca^{2+}$ ,  $Mg^{2+}$ ,  $K^+$ , and  $Na^+$ ), especially  $Ca^{2+}$ , have significantly contributed to maintaining the particle pH value at the level of 4.5 – 5 in East Asia during the last decade. The importance of crustal ions in determining aerosol acidity and factors such as liquid water content, aerosol mass concentration, and chemical composition, has been highlighted in other studies (Zheng et al., 2020; Zhang et al., 2021a; Ding et al., 2019).

## 2.1 $NH_3$ emission scheme

In this study, three  $NH_3$  emission schemes are applied in specific model simulation cases to

Formatted: English (United States)

quantify the impact of changes in  $\text{NH}_3$  emissions on aerosol composition and acidity (Table 1), which are divided into Bottom-up and Top-down cases. These include two bottom-up inventories used in the Base and Top-CEDS cases, and one top-down scheme inventory applied in the Top-Dep case. In addition, two sensitivity simulations are conducted: the *Meta* case, which assumes metastable aerosol thermodynamics (described in detail in Section 2.2), and the *noNH<sub>3</sub>* case, in which all  $\text{NH}_3$  emissions are turned off. The latter is used to provide insights into the role of  $\text{NH}_3$  in shaping the simulated size-resolved aerosol acidity.

**Table 1. Overview of simulation cases.**

Simulation Case	$\text{NH}_3$ Emission Scheme	Equilibrium State
Base	CAMS-GLOB-ANT, CAMS-GLOB-AIR, GEIA, biomass burning	Stable
CEDS	Same as Base case, but $\text{NH}_3$ , $\text{SO}_2$ , and $\text{NO}_x$ are from CEDS-GBD	Stable
Top-Dep	Luo's method with the lifetime derived from deposition, GEIA-water	Stable
Meta	Same as Base case	Metastable
noNH <sub>3</sub>	No $\text{NH}_3$ emission input	Stable

### 2.1.1 Bottom-up scheme

Bottom-up schemes are applied in the base and the CEDS *easescase*, with different anthropogenic emission inventories. The anthropogenic emission inventory utilized in the base case simulation is CAMS-GLOB-ANT (v4.2, <https://eccad3.sedoo.fr>, abbreviated as “CAMS”), which contains 17 sectors with a spatial resolution of  $0.1 \times 0.1$  degree and monthly temporal resolution. With the basic data of 2010 from the Emission Database for Global Atmospheric Research (EDGAR, v4.3.2), CAMS extends the period to recent years based on the trend factors derived from the Community Emissions Data System (Hoesly et al., 2018). Meanwhile, Hoesly et al. (2018) pointed out that there are limitations in the system, especially in the emission trends for specific sectors, and emphasized the need for more detailed data to be incorporated into regional emission inventories. The other one used in the CEDS sensitivity simulation case for aerosol precursor emissions ( $\text{NH}_3$ ,  $\text{SO}_2$ , and  $\text{NO}_x$ ) is CEDS-GBD-MAPS (McDuffie et al., 2020), abbreviated as “CEDS\_GBD”. The CEDS\_GBD is developed using the Community Emissions Data System and is reported as a function of 11 anthropogenic sectors and 4 fuel categories, with a spatial resolution of  $0.5 \times 0.5$  degrees and monthly temporal resolution. Assuming that the specific regional emission inventories are more accurate, McDuffie et al. (2020) updated the activity data and the core scaling procedure, modified the final emission gridding and aggregation procedures, and then utilized several regional emission inventories to improve the previous version of CEDS\_GBD via the scaling procedure, which can not only reduce the discrepancy with other global emission inventories but also help to maintain the timeliness and regional accuracy of the global estimates. However, they also pointed out that the sources of uncertainty in the CEDS\_GBD are similar to those in the CEDS.

Other  $\text{NH}_3$  emissions include aircraft emissions from the CAMS-GLOB-AIR inventory (v1.1, <https://eccad3.sedoo.fr>), land and water biological emissions from the Global Emissions Inventory Activity (GEIA) inventory, and biomass burning emissions calculated by the BIOBURN submodel (Kaiser et al., 2012). BIOBURN determines the flux based on biomass burning emission factors and dry matter combustion rates from the Global Fire Assimilation System (GFAS), which calculates biomass burning emissions by assimilating Fire Radiative Power (FRP) observations from MODIS (Andreae, 2019).

Formatted: No underline, Font color: Auto

### 2.1.2 Top-down scheme

The NH<sub>3</sub> emission inventory over land is updated by a top-down method with the constraint of IASI satellite observations (<https://iasi.acris-data.fr>) developed by Luo et al. (2022). This fast top-down method updates the prior NH<sub>3</sub> emissions ( $E_{NH_3,mod}$ , molecule m<sup>-2</sup> s<sup>-1</sup>) with a correction term positively proportional to the biases between observed ( $C_{NH_3,obs}$ , molecule m<sup>-2</sup>) and simulated ( $C_{NH_3,mod}$ , molecule m<sup>-2</sup>) monthly averaged NH<sub>3</sub> total column densities and inversely proportional to the NH<sub>3</sub> lifetime ( $\tau_{NH_3,mod}$ , s) (Eq. 1). The  $\tau_{NH_3,mod}$  is calculated as the ratio of the simulated NH<sub>3</sub> column and the sum of the simulated loss rate of the NH<sub>x</sub> family (NH<sub>x</sub> ≡ NH<sub>3</sub>+ NH<sub>4</sub><sup>+</sup>) through dry and wet deposition (Eq. 2).

$$E_{NH_3} = E_{NH_3,mod} + \frac{C_{NH_3,obs} - C_{NH_3,mod}}{\tau_{NH_3,mod}} E_{NH_3} = E_{NH_3,mod} + \frac{C_{NH_3,obs} - C_{NH_3,mod}}{\tau_{NH_3,mod}} E_{NH_3} \quad (1)$$

$$\tau_{NH_3,mod} = \frac{C_{NH_3,mod}}{D_{NH_3,mod} + D_{NH_4^+,mod}} \tau_{NH_3,mod} = \frac{C_{NH_3,mod}}{D_{NH_3,mod} + D_{NH_4^+,mod}} \quad (2)$$


The fast top-down method relies on the total column concentrations retrieved by IASI. According to Dammers et al. (2019), the dominant source of uncertainty in the IASI observational product stems from the systematic bias, with the negative bias estimated between 25% to 40%, compared to site observations (Dammers et al., 2017). While this method simplifies the chemical and physical processes governing NH<sub>3</sub>, Luo et al. (2022) identified large uncertainties in regions like Central Asia and tropical Africa due to poorly constrained sources by IASI observations in these areas. Nonetheless, they demonstrated that simulations driven by the updated top-down emission inventory show better consistency with satellite observations compared to those driven by the prior emission inventory.

### 2.1.3 Emission comparison

Overall, the global NH<sub>3</sub> emissions used in this study in all simulation cases range from 74.73 Tg yr<sup>-1</sup> to 85 Tg yr<sup>-1</sup> (Table 2), which is within the reported range of the current literature from 52 Tg yr<sup>-1</sup> to 91 Tg yr<sup>-1</sup> (Schlesinger and Hartley, 1992; Dentener and Crutzen, 1994; Bouwman et al., 1997; Olivier et al., 1998; Van Aardenne et al., 2001; Bleeker et al., 2013). The distribution of the global NH<sub>3</sub> emission flux derived from the base case and the absolute emission flux difference between the sensitivity simulation cases and the base case are shown in Figure 1. Significant regional variations in NH<sub>3</sub> emission flux are found in Figure 1(a), with the maximum emission flux exceeding 7 g m<sup>-2</sup> yr<sup>-1</sup> in northern India, eastern China, and central Europe, all regions with the highest population density. Other emission hotspots include the eastern United States, southeastern Latin America, and central and eastern Africa. The base case is able to capture the global NH<sub>3</sub> emission hotspots reported by Van Damme et al. (2018). Meanwhile, the minimum flux below 0.01 g m<sup>-2</sup> yr<sup>-1</sup> is located in the Antarctic and Arctic regions and the Sahara Desert, as well as in remote oceans, where there is little impact from human activities. Agricultural activities including livestock and fertilization are the main source of NH<sub>3</sub> emissions in China, India, and the United States (Liu et al., 2022; Khan et al., 2020; Van Damme et al., 2018; Sahoo et al., 2024), while soil emissions, biomass burning and domestic fires are the

Formatted: Subscript

main contributors to  $\text{NH}_3$  emissions in central and eastern Africa (Hickman et al., 2021; Delon et al., 2012).



**Figure 1.** Global distribution of the annual average  $\text{NH}_3$  emission flux for (a) the Base case and the absolute differences between (b) the Base and CEDS and (c) the Base and Top-Dep cases during 2010–2012.

Compared with the base case, the global  $\text{NH}_3$  emissions in both the CEDS and Top-Dep cases show an increase exceed those in the global  $\text{NH}_3$  emission amount base case, with increases of approximately 5% and 16%, respectively. The lower  $\text{NH}_3$  emission fluxes of the CEDS case are found in North America, Europe, and China except for the northeastern and southeastern coastal areas, while emission fluxes are higher throughout India (Figure 1b). Slightly higher fluxes are also found in western and eastern Africa, the northern Middle East, and southeastern Asia. According to McDuffie et al. (2020), the  $\text{NH}_3$  emission flux from the Multi-resolution Emission Inventory for China (MEIC, <http://www.meicmodel.org>), European Monitoring and Evaluation Programme (EMEP), and US EPA are used

to scale the previous emission (the basic data of 2010 from EDGAR) over mainland China, Europe, and the USA. The NH<sub>3</sub> emission flux from India and Africa remains the same as that of the original inventory. Constrained by the IASI satellite observation, the NH<sub>3</sub> emission flux of the Top-Dep case is ~~increased~~<sup>elevated</sup> in most regions of the world, but lower emission fluxes are estimated in regions such as western North America, western and southern Europe, India, China except the northeastern and southeastern coastal areas, and western Australia. The comparison of the absolute and relative difference between the base case and the other emission schemes is summarized in Table 2.

**Table 2.** Comparison of annual NH<sub>3</sub> emissions (Tg yr<sup>-1</sup>) across global and regional scales between the two sensitivity cases ~~cases~~ and the base case.

Region	Base (Tg yr <sup>-1</sup> )	diff <sup>a</sup>	CEDS relative diff <sup>b</sup> (%)	diff <sup>a</sup>	Top-Dep relative diff <sup>b</sup> (%)
Globe	73.273	3.869	5.3	11.444	16
Land	61.141	3.050	5.0	11.263	18
North America	5.566	-1.000	-18	3.232	58
South America	8.202	0.697	8.4	4.182	51
Europe	6.444	-1.505	-23	1.010	16
Middle East	1.051	0.212	20	-0.015	-1.4
South Asia	10.111	3.798	38	-1.098	-9.7
East Asia	15.757	-2.030	-13	-2.586	-16

<sup>a</sup>: absolute difference between sensitivity cases and the base case.

<sup>b</sup>: relative difference between sensitivity cases and the base case.

<sup>c</sup>: below the 0.1 precision threshold.

Formatted Table

## 2.2 ISORROPIA II

The thermodynamic equilibrium model ISORROPIA II is used to calculate the multi-phase mass transfer of the K<sup>+</sup>-Ca<sup>2+</sup>-Mg<sup>2+</sup>-NH<sub>4</sub><sup>+</sup>-Na<sup>+</sup>-SO<sub>4</sub><sup>2-</sup>-NO<sub>3</sub><sup>-</sup>-Cl<sup>-</sup>-H<sub>2</sub>O aerosol system (Nenes et al., 1998; Fountoukis and Nenes, 2007). The process of gas/aerosol partitioning is calculated in two steps (Pringle et al., 2010). In the first step, the amount of gas phase species kinetically able to condense on the aerosol within one timestep is calculated (Vignati et al., 2004). ISORROPIA II then re-distributes the mass between the gas and aerosol phase. In this study, ISORROPIA II runs in the forward mode with the input of relative humidity, temperature, and concentration of aerosol and gas phase species. ISORROPIA II determines the subsystem set of equilibrium equations and solves the equilibrium state by the chemical potential method. It outputs the equilibrium concentration of species in gas, solid, and liquid phases by assuming that the particle phase is in the thermodynamically stable-state mode where salts precipitate once the aqueous phase becomes saturated (Fountoukis and Nenes, 2007).

Meanwhile, a sensitivity case assuming the particle phase in the thermodynamically metastable state mode is performed with the same emission scheme as the base case. (Meta case). In the metastable state, the aerosol may be supersaturated with respect to dissolved salts and always consists only of an aqueous phase (Fountoukis and Nenes, 2007). Karydis et al. (2021) pointed out that more acidic particles (up to 2 pH units) are derived from the metastable assumption in regions affected by high concentrations of crustal cations and consistently low relative humidity values.

According to past studies, the treatment of crustal species (e.g. Ca<sup>2+</sup>, K<sup>+</sup>, Mg<sup>2+</sup>) in ISORROPIA II can improve model predictions (Karydis et al., 2010; Karydis et al., 2011), as both the phase partitioning of NO<sub>3</sub><sup>-</sup> and the thermodynamic interaction between NH<sub>4</sub><sup>+</sup> and the remaining ions in the aqueous phase are significantly affected. Karydis et al. (2016) found that when these crustal species are included in the EMAC model, the increase in global

NO<sub>3</sub><sup>-</sup> tropospheric load can be up to 44% while the global NH<sub>4</sub><sup>+</sup> tropospheric load decreases by 41%.

Formatted: English (United Kingdom)

### 2.2.1 pH calculations

The pH is calculated from the negative decimal logarithm of the hydrogen ion activity,

$$pH = -\log_{10} pH = -\log_{10}(\gamma x_H x_{H^+}) \quad (3)$$

where  $x_{H^+}$  and  $\gamma$  represent the molality of hydrogen ions in the solution and the ionic activity coefficient of hydrogen, respectively. With  $\gamma$  assumed to be unity, the pH value is derived using the hydrogen ion concentration in the aqueous particle phase output by ISORROPIA-II (in mol m<sup>-3</sup>) and the aerosol water content output by GMXe (in mol mol<sup>-1</sup>). The effect of water-soluble organic aerosols on aerosol water content is accounted for via the GMXe module, which includes both inorganic and organic contributions. Organic aerosol formation is simulated using the ORACLE module, and the associated water uptake is calculated assuming a  $\kappa$ -hygroscopicity value of 0.14 for all organic components (Tsimpidi et al., 2017). This influences the total aerosol water content used in pH calculations. Both hydrogen ion and aerosol water content are output every 5 hours, following Karydis et al. (2021). In addition, the temperature threshold of 269 K is set to ensure that the calculations are performed only when liquid water is present in the aerosol.

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: Italic

Formatted: English (United States)

Formatted: Font: Italic

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Discrepancies among thermodynamic models tend to increase as RH decreases, primarily due to differing assumptions about activity coefficients (Pye et al., 2020). ISORROPIA-II assumes constant mean activity coefficients and a unity activity coefficient for H<sup>+</sup>, which results in relatively invariant and often lower pH predictions under dry conditions. Additionally, the choice between stable and metastable phase state assumptions introduces further uncertainty. Under low RH (e.g., <35%), the metastable assumption allows for supersaturated solutions, enabling pH calculations even at low aerosol liquid water content, often resulting in low pH values. Conversely, the stable state permits salt crystallization, and in cases where the aerosol becomes fully solid, pH may no longer be defined. To assess the sensitivity of our results to this assumption, we conducted a simulation using the metastable assumption (Meta case; see Table 1).

EMAC model does not account for chemical interactions between ammonia and organic acids (e.g., ammonium oxalate formation) and treats inorganic and organic aerosol phases separately. As a result, while organics can indirectly influence pH through their contribution to aerosol water content, changes in NH<sub>3</sub> emissions do not affect organic aerosol formation or the associated water content in our simulations. Furthermore, although the influence of organics on hydrogen ion activity coefficients is not explicitly represented, previous studies have shown that water-soluble organic aerosols exert only a minor effect on aerosol pH (Guo et al., 2015; Pye et al., 2018; Vasilakos et al., 2018).

### 2.2.2 Two factors affecting pH value change

According to Equation (3), the pH value is determined by the concentrations of H<sup>+</sup> and H<sub>2</sub>O. To evaluate the impact of each factor on the pH value, we independently calculate the changes in pH arising from two pathways: one driven by H<sup>+</sup> and the other by H<sub>2</sub>O. The corresponding results are presented in Figures S16 and S17.

$$\Delta pH_{H_2O} = \log_{10} \frac{H_2O + \Delta H_2O}{H_2O} - \log_{10} \frac{H_2O}{H^+} pH_{H_2O} = \log_{10} \frac{H_2O + \Delta H_2O}{H^+} - \log_{10} \frac{H_2O}{H^+} \quad (4)$$

$$\Delta pH_{H^+} = \log_{10} \frac{H^+}{H^+ + \Delta H^+} - \log_{10} \frac{H^+}{H^+} \Delta pH_{H^+} = \log_{10} \frac{H_2O}{H^+ + \Delta H^+} - \log_{10} \frac{H_2O}{H^+} \quad (5)$$

Here,  $\Delta pH_{H_2O}$  and  $\Delta pH_{H^+}$  represent the changes in pH caused by variations in  $H_2O$  and  $H^+$  concentrations, respectively. The base case concentrations of  $H_2O$  and  $H^+$  are used as references, and the changes in concentration are expressed as  $\Delta H_2O$  and  $\Delta H^+$ , corresponding to the deviations in  $H_2O$  and  $H^+$  from their base case levels.

### 3. Observations

Multiple observational datasets are used in this study to validate the model simulation in different regions of the world as defined by the IPCC (2023/2022). The information of each dataset is summarized in Table 3 and the site distribution is plotted in Figure 2 and Figure S1. These datasets include satellite retrievals from the Infrared Atmospheric Sounding Interferometer (IASI; [https://iasi.aeris-data.fr/NH3\\_IASI\\_A\\_L3\\_data/](https://iasi.aeris-data.fr/NH3_IASI_A_L3_data/)), and observation site networks Nationwide Nitrogen Deposition Monitoring Network (NNDMN; [https://figshare.com/articles/dataset/Data\\_Descriptor\\_Xu\\_et\\_al\\_20181211\\_Scientific\\_data\\_docx/7451357/5](https://figshare.com/articles/dataset/Data_Descriptor_Xu_et_al_20181211_Scientific_data_docx/7451357/5)), the European Monitoring and Evaluation Programme (EMEP; <https://ebas-data.nilu.no/Default.aspx>), the Central Pollution Control Board (CPCB; <https://cpcb.nic.in/>), the Acid Deposition Monitoring Network in East Asia (EANET; <http://www.eanet.asia/product/index.html>), Ammonia Monitoring Network (AMoN; <https://nadp.slh.wisc.edu/networks/ammonia-monitoring-network/>), the U.S. Environmental Protection Agency (EPA; <https://www.epa.gov/data>), and the Clean Air Status and Trends Network (CASTNET; <http://www.epa.gov/castnet>).

**Table 3.** Information for each observation dataset used to validate model simulation during 2010–2012.

Dataset	Parameters	Location
IASI	NH <sub>3</sub> column concentration	Globe <sup>a</sup>
NNDMN	NH <sub>3</sub> , NH <sub>4</sub> <sup>+</sup> , NO <sub>3</sub> <sup>-</sup> mass concentration	China (29 sites)
EMEP	NH <sub>3</sub> , NH <sub>4</sub> <sup>+</sup> , NO <sub>3</sub> <sup>-</sup> , and SO <sub>4</sub> <sup>2-</sup> mass concentration	Europe (25 sites for NH <sub>3</sub> ; 7 sites of PM <sub>25</sub> matrix <sup>b</sup> )
CPCB	NH <sub>3</sub> mass concentration	India (8 sites)
EANET	NH <sub>4</sub> <sup>+</sup> , NO <sub>3</sub> <sup>-</sup> , and SO <sub>4</sub> <sup>2-</sup> mass concentration	eastern and southeastern Asia (50 sites, PM <sub>25</sub> matrix <sup>b</sup> )
AMoN	NH <sub>3</sub> and NH <sub>4</sub> <sup>+</sup> mass concentration	America (21 sites)
CASTNET	NH <sub>3</sub> , NH <sub>4</sub> <sup>+</sup> , NO <sub>3</sub> <sup>-</sup> , and SO <sub>4</sub> <sup>2-</sup> mass concentration	America (79 sites, PM <sub>25</sub> matrix <sup>b</sup> )
EPA	NH <sub>4</sub> <sup>+</sup> , NO <sub>3</sub> <sup>-</sup> , and SO <sub>4</sub> <sup>2-</sup> mass concentration	America (211 sites, PM <sub>25</sub> matrix <sup>b</sup> )

<sup>a</sup>: Only use the data over land.

<sup>b</sup>: Measurements refer to a chemical or physical property of the total aerosol particle phase in the size fraction less than 2.5 micrometer median aerodynamic diameter.

Due to its better precision resulting from favorable thermal contrast conditions (Clarisse et al., 2009), only the

morning (around 9:30 local time) overland IASI data is used in this study. The original temporal resolution of the various datasets includes hourly, three-day, daily, weekly, bi-weekly, and monthly, which we uniformly convert to monthly. The mean bias (MB), mean absolute gross error (MAGE), normalized mean bias (NMB), normalized mean error (NME), and root mean square error (RMSE) are calculated to evaluate the model performance:

$$MAGE = \frac{1}{N} \sum_{i=1}^N |P_i - O_i| \quad (6)$$


$$MB = \frac{1}{N} \sum_{i=1}^N (P_i - O_i) \quad (7)$$

$$NME = \frac{\sum_{i=1}^N |P_i - O_i|}{\sum_{i=1}^N O_i} \quad (8)$$

$$NMB = \frac{\sum_{i=1}^N (P_i - O_i)}{\sum_{i=1}^N O_i} \quad (9)$$

$$RMSE = \left[ \frac{1}{N} \sum_{i=1}^N (P_i - O_i)^2 \right]^{\frac{1}{2}} \quad (10)$$

where  $P_i$  and  $O_i$  represent the monthly value of model simulation and measurement, respectively.  $N$  is the total number of data points used for comparison.



**Figure 2.** Regions and observation sites used in this study. North America, South America, Europe, Middle East, South Asia, and East Asia are marked represented by purple, navy blue, sky blue, orange, light green, and yellow, respectively. In North America AMoN and CASTNET sites are represented by shown as blue and red circles in North America (the, respectively (EPA sites are shown provided in Figure S3-S1). In Europe, EMEP sites in Europe are shown by blue circles for NH<sub>3</sub> and red circles for PM<sub>2.5</sub> components. In South Asia, CPCB sites are marked by green circles in South. In East Asia, while NNDMN and EANET sites are indicated by dark green and dark blue in East Asia circles, respectively.

#### 4. Observation Model evaluation

##### 4.1 Aerosol comparison composition

The comparison between the simulation derived from the base case and the observational datasets is summarized in Tables 4 – 6. Compared with the NNDMN dataset, of which primarily includes sites are mainly located in urban and rural areas of across China, the base case overestimates the NH<sub>3</sub> mass concentration (NMB = 0.19), underestimates the NH<sub>4</sub><sup>+</sup> mass concentration (NMB = -0.41) though reproduces the NO<sub>3</sub><sup>-</sup> mass concentration well (NMB = -0.02).

Xie et al. (2022) summarized the NMB between observed  $\text{NO}_3^-$  and simulated values in China as ranging from -0.97 to 1.90 based on modelling studies in the last decade. The negligible bias of the simulated  $\text{NO}_3^-$  shows the good performance of the EMAC model in this region. However, the biases in the simulation of  $\text{NH}_3$  and  $\text{NH}_4^+$  indicate that the  $\text{NH}_3/\text{NH}_4^+$  partitioning treatment is not efficient enough or over-simplified, as less  $\text{NH}_4^+$  is produced even with sufficient  $\text{NH}_3$ . Similarly, in Europe and North America, we obtain a positive bias of  $\text{NH}_3$  mass concentration (EMEP dataset: NMB = 2.26; AMoN dataset: NMB = 0.58) and a comparably lower  $\text{NH}_4^+$  mass concentration (EMEP dataset: NMB = 0.05; AMoN dataset: NMB = -0.23). On the other hand, the dry deposition of  $\text{NH}_3$  in China is lower than reported from observations (NMB = -0.28; not shown), which contributes to a higher atmospheric  $\text{NH}_3$  concentration.

**Table 4.** Comparison of the Base case predictions with monthly average observations from China and Europe during 2010–2012.

	NNDMN network <i>Metrie</i>			EMEP network <i>Metrie</i>			
	$\text{NH}_3$	$\text{NH}_4^+$	$\text{NO}_3^-$	$\text{NH}_3$	$\text{NH}_4^+$	$\text{NO}_3^-$	$\text{SO}_4^{2-}$
Observed ( $\mu\text{g m}^{-3}$ )	7.68	7.45	11.92	1.16	1.09	1.29	2.11
Simulated ( $\mu\text{g m}^{-3}$ )	9.14	4.40	11.70	3.79	1.15	3.33	1.24
MAGE ( $\mu\text{g m}^{-3}$ )	5.27	4.21	6.00	2.78	0.60	2.38	0.98
MB ( $\mu\text{g m}^{-3}$ )	1.46	-3.05	-0.22	2.63	0.06	2.04	-0.88
NME	0.69	0.56	0.50	2.40	0.55	1.84	0.46
NMB	0.19	-0.41	-0.02	2.26	0.05	1.58	-0.41
RMSE ( $\mu\text{g m}^{-3}$ )	7.43	6.55	9.07	4.03	0.97	2.95	1.63
Number of comparisons	765	765	765	832	249	320	320

**Table 5.** Comparison of the Base case predictions with monthly average observations from India and eastern Asia ( $\text{PM}_{25}$  matrix) during 2010–2012.

	CPCB network <i>Metrie</i>	EANET network <i>Metrie</i>		
	$\text{NH}_3$	$\text{NH}_4^+$	$\text{NO}_3^-$	$\text{SO}_4^{2-}$
Observed ( $\mu\text{g m}^{-3}$ )	30.07	0.84	1.22	2.95
Simulated ( $\mu\text{g m}^{-3}$ )	22.30	0.83	2.09	1.52
MAGE ( $\mu\text{g m}^{-3}$ )	21.25	0.55	1.58	1.74
MB ( $\mu\text{g m}^{-3}$ )	-7.78	-0.02	0.87	-1.43
NME	0.71	0.65	1.29	0.59
NMB	-0.26	-0.02	0.71	-0.49
RMSE ( $\mu\text{g m}^{-3}$ )	30.57	1.11	2.58	2.48
Number of comparisons	137	908	916	886

**Table 6.** Comparison of the Base case predictions with monthly average observations from America ( $\text{PM}_{25}$  matrix) during 2010–2012.

	AMoN network <i>Metrie</i>		CASTNET network <i>Metrie</i>			EPA network <i>Metrie</i>		
	$\text{NH}_3$	$\text{NH}_4^+$	$\text{NH}_4^+$	$\text{NO}_3^-$	$\text{SO}_4^{2-}$	$\text{NH}_4^+$	$\text{NO}_3^-$	$\text{SO}_4^{2-}$
Observed ( $\mu\text{g m}^{-3}$ )	1.20	1.27	0.69	0.74	1.81	0.83	1.24	1.97
Simulated ( $\mu\text{g m}^{-3}$ )	1.89	0.99	0.90	1.96	1.34	1.02	2.22	1.44
MAGE ( $\mu\text{g m}^{-3}$ )	1.16	1.02	0.34	1.33	0.64	0.42	1.30	0.65
MB ( $\mu\text{g m}^{-3}$ )	0.69	-0.29	0.21	1.22	-0.46	0.19	0.98	-0.53
NME	0.97	0.80	0.49	1.80	0.36	0.51	1.05	0.33
NMB	0.58	-0.23	0.30	1.65	-0.26	0.23	0.79	-0.27
RMSE ( $\mu\text{g m}^{-3}$ )	1.64	1.43	0.45	1.71	0.86	0.60	1.76	0.90
Number of comparisons	553	552	2825	2825	2825	5085	5392	5429

In East and Southeast Asia, the mass concentration of  $\text{NH}_4^+$  is well reproduced (NMB = -0.02), while high and low discrepancies are found in the mass concentrations of  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$  (NMB = 0.71 and -0.49, respectively). Similar results were also found in Europe, with agreement for  $\text{NH}_4^+$  (NMB = 0.05) but an overestimation of the  $\text{NO}_3^-$  mass

concentration (NMB = 1.58) and underestimated  $\text{SO}_4^{2-}$  (NMB = -0.41). The positive bias of the simulated  $\text{NO}_3^-$  is reported by many studies (Xie et al., 2022; Heald et al., 2012; Colette et al., 2011; Bian et al., 2017; Pozzer et al., 2022; Milousis et al., 2025). The negative bias of  $\text{SO}_4^{2-}$  is considered a reason for the positive bias of  $\text{NO}_3^-$ , regarding the thermodynamic equilibrium between  $\text{NH}_4^+$ ,  $\text{SO}_4\text{SO}_4^{2-}$ , and  $\text{NO}_3^-$ . The discrepancy in  $\text{SO}_4\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  is also due to the missing heterogeneous oxidation reactions for  $\text{SO}_2$  and  $\text{NO}_x$  in the model. Several studies have concluded that adding multiphase chemistry can significantly improve the model performance (Zheng et al., 2015; Zhang et al., 2021b). Cheng et al. (2016) and Zheng et al. (2024) pointed out that the multiphase reactions act as an important  $\text{SO}_4^{2-}$  source in haze pollution, while Guo et al. (2017a) argued that the multiphase reactions are not likely limited by the required alkaline environment.

In North America, the base case reproduces the mass concentration of  $\text{SO}_4^{2-}$  (NMB = -0.26), but overpredicts the mass concentrations of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  (NMB = 0.23 and 0.79, respectively), which is in line with the findings of Walker et al. (2012). The excess  $\text{NH}_4^+$  promotes the formation of  $\text{NO}_3^-$ , and the uncertain uptake coefficient of  $\text{N}_2\text{O}_5$  used in models may contribute to more  $\text{NO}_3^-$  (Walker et al., 2012). The highest  $\text{NH}_3$  mass concentration is found in India, especially in the Indo-Gangetic Plain. Our model basically captures the “hot spot”, with a slight negative bias (NMB = -0.26). However, the scarce observation sites and the lack of observed  $\text{NH}_4^+$ ,  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$  hinder further evaluation of the model performance.

#### 4.2 pH-value comparison Aerosol acidity

Due to the lack of direct measurement of aerosol acidity, we collect the  $\text{PM}_{2.5}$  pH value from related studies to compare with our model simulation in Table 7. These pH values are calculated using the thermodynamic equilibrium model ISORROPIA or E-AIM with input from observational datasets. Compared with Guo et al. (2017b), our simulated pH value from the base case is higher in the western USA (4.3 vs 2.7), but the value from the Meta case is much closer (2.6). Both the base and Meta cases predict the same aerosol water content, and the high  $\text{Ca}^{2+}$  concentration from the Great Basin Desert leading to the precipitation of  $\text{CaSO}_4$  is the main reason for the lower aerosol acidity in the base case (Karydis et al., 2021). It is worth noting that the effect of  $\text{Ca}^{2+}$  on aerosol acidity was not considered by Guo et al. (2017b). The summertime southeastern USA is characterized as relatively clean with aerosol dominated by  $\text{NH}_4^+$  and  $\text{SO}_4^{2-}$ , but negligible mineral cations, the aerosol is acidic with the observed pH value around ~0.6 (Guo et al., 2015; Pye et al., 2020). However, due to a low bias in simulated  $\text{SO}_4^{2-}$  mass concentration, the model underestimates aerosol acidity, resulting to a higher simulated pH value of approximately 2.0 during the summer period. In Europe, although the good agreement between our simulated pH and the result of Guo et al. (2018) (both are 3.9) supports the model simulations, we note the a significant overestimation of simulated alkaline species (sum of  $\text{NH}_4^+$  and  $\text{NH}_3$ , NMB = 17%) along with an underestimation of acidic species (sum of  $\text{NO}_3^-$  and  $\text{HNO}_3$ , NMB = -57%;  $\text{SO}_4^{2-}$ , NMB = -51%) in December emphasizes the improvement of the aerosol. These discrepancies emphasize the need for improved representation of aerosol composition in the model. In northern China, the pH value calculated by Wang et al. (2016) was more than reported pH values that are about 2 units higher than our simulations (4.4 vs 6.7), and we suggest that the low bias is due to Guo et al. (2017) found moderate acidic aerosol (pH = 5) during wintertime. We attribute this high bias in simulated  $\text{pH}_a$  to the underestimation of the cation mass

Formatted: English (United States)

Formatted: English (United States)

concentration of cations concentrations (e.g.  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ) in  $\text{PM}_{2.5}$ , as the mass concentrations of  $\text{NH}_4^+$ ,  $\text{SO}_4^{2-}$ , and  $\text{NO}_3^-$  are in the observational ranges of reported by Wang et al. (2016).

Formatted: Font color: Text 1

Table 7. Simulated pH value of  $\text{PM}_{2.5}$  at single points certain locations compared with literature observational estimates of particle pH.


Location	Latitude	Longitude	Time Period	Method used	Field derived mean pH	Base	Meta	Reference
Egbert, ON, Canada	44.23	-79.78	Jul-Sep, 2012	E-AIM	2.1	3.77	1.57	Murphy et al. (2017)
Pasadena, CA, USA	34.14	-118.12	Jun, 2010	ISORROPIA	2.7	4.26	2.58	Guo et al. (2017)
Sao Paulo, Brazil	-23.55	-46.63	Aug-Sep, 2012	E-AIM	4.8	3.85	3.34	Vieira-Filho et al. (2016)
Cabauw, Netherland	51.97	4.93	Dec-Feb, 2012	ISORROPIA	3.9	3.91	3.58	Guo et al. (2018)
Xi'an, China	34.23	108.89	Nov-Dec, 2012	ISORROPIA	6.7	4.41	3.20	Wang et al. (2016)

Note: table extracted in part from Karydis et al. (2021).

5. Secondary inorganic aerosol composition


5.1 Size-resolved composition

The regional mass fractions of size-resolved inorganic aerosol components ( $\text{NH}_4^+$ ,  $\text{SO}_4^{2-}$ , and  $\text{NO}_3^-$ ) are presented as bar charts in Figure 3, while global distribution maps of their size-resolved mass concentrations are shown in Figure S2, both the simulation results are derived from base case. To assess ammonia neutralization of sulfuric and nitric acids, we applied the chemical domain framework defined by Ge et al. (2022) based on SNA molar concentrations in  $\text{PM}_{10}$  with a threshold of  $>1 \mu\text{g m}^{-3}$ . The four chemical domains, illustrated in Figure 4, are defined as follows: “ $\text{SO}_4^{2-}$  very rich” ( $\text{totNH}_3/\text{totSO}_4 < 1$ ,  $\text{totNH}_3$ : sum of  $\text{NH}_3$  and  $\text{NH}_4^+$ ,  $\text{totSO}_4$ : sum of  $\text{SO}_4^{2-}$  and  $\text{HSO}_4^-$ ), “ $\text{SO}_4^{2-}$  rich” ( $\text{totNH}_3/\text{totSO}_4$  between 1 and 2), “ $\text{NO}_3^-$  rich” (free  $\text{NH}_3/\text{totNO}_3$  between 0 and 1, free  $\text{NH}_3$ :  $\text{totNH}_3$  minus double  $\text{totSO}_4$ ,  $\text{totNO}_3$ : sum of  $\text{NO}_3^-$  and  $\text{HNO}_3$ ), and “ $\text{NH}_3$  very rich” (free  $\text{NH}_3/\text{totNO}_3 > 1$ ). Figure S3 shows the ratios used to define these domains.



**Figure 3.** (a) – (c) global distribution of  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ , and  $\text{NH}_4^+$  mass concentration in the size range of  $0 - 10 \mu\text{m}$ , (d) – (f) bar plots for mean mass fractions of size-resolved  $\text{NH}_4^+$ ,  $\text{SO}_4^{2-}$ , and  $\text{NO}_3^-$  over globe, land, ocean and regions (marked in Figure 2), the size range of  $5 - 10 \mu\text{m}$ ,  $2.5 - 5 \mu\text{m}$ ,  $1 - 2.5 \mu\text{m}$ , and  $0 - 1 \mu\text{m}$  are marked by green, orange, purple, and blue, respectively (NAM: North America; SAM: South America; Eu: Europe; ME: Middle East; SAs: South Asia; EAs: East Asia).

Regions with low inorganic aerosol concentrations ( $< 1 \mu\text{g m}^{-3}$ ) are found in Southern Hemisphere oceans and remote areas such as the North Pole and South America, including the Amazon Basin. These areas represent relatively pristine baselines for evaluating anthropogenic impacts (Andreae et al., 1990; Andreae, 2007). In low northern latitude oceans, the “ $\text{SO}_4^{2-}$  very rich” and “ $\text{SO}_4^{2-}$  rich” domains dominate, characterized by  $\text{NH}_3$  fully converted to  $\text{NH}_4^+$  but incomplete neutralization of sulfuric acid where  $\text{NH}_3$  is insufficient to fully neutralize sulfuric acid. As a result, sulfuric acid remains only partially neutralized, contributing to aerosol acidity. This pattern reflects low  $\text{NH}_3$  emissions over oceans (Figure 1) and the contribution of biogenic dimethyl sulfide (DMS) to marine  $\text{SO}_4^{2-}$  (Fiddes et al., 2018; Jackson et al., 2020). Additional  $\text{NO}_x$  and  $\text{SO}_2$  from shipping contribute to marine  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  (Wang et al., 2023; Burgard and Briä, 2016). Average  $\text{PM}_{10}$  SNA concentrations are lower over oceans compared to land ( $\text{NH}_4^+$ :  $0.06 \mu\text{g m}^{-3}$  vs.  $0.36 \mu\text{g m}^{-3}$ ;  $\text{NO}_3^-$ :  $0.37 \mu\text{g m}^{-3}$  vs.  $1.24 \mu\text{g m}^{-3}$ ;  $\text{SO}_4^{2-}$ :  $0.60 \mu\text{g m}^{-3}$  vs.  $0.95 \mu\text{g m}^{-3}$ ). Marine  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  are predominantly in the super-micron mode, with mass fractions of 70% and 58%, respectively. Coastal areas exhibit a “ $\text{NO}_3^-$  rich” domain due to continental outflows (Figure S2S3) (Prospero et al., 1985), consistent with prior findings that marine aerosol in the super-micron mode primarily comprises inorganic salts, including sea salt, non-sea salt sulfate, and nitrate, while organic matter is concentrated in the sub-micron range (Russell et al., 2023; Cavalli et al., 2004).



**Figure 4.** Distribution of ammonia neutralization state of sulfuric and nitric acids, based on SNA molar concentrations in  $\text{PM}_{10}$  with a threshold of  $\geq 1 \mu\text{g m}^{-3}$ , where the SNA mass concentration less than the threshold is measured by blank zone. The “ $\text{SO}_4^{2-}$  very rich” represents the ratio of  $\text{totNH}_3$  to  $\text{totSO}_4$  less than 1 ( $\text{totNH}_3$ : sum of  $\text{NH}_3$  and  $\text{NH}_4^+$ ,  $\text{totSO}_4$ : sum of  $\text{SO}_4^{2-}$  and  $\text{HSO}_4^-$ ), the “ $\text{SO}_4^{2-}$  rich” represents the ratio of  $\text{totNH}_3$  to  $\text{totSO}_4^{2-}$  between 1 and 2, the “ $\text{NO}_3^-$  rich” represents the ratio of free  $\text{NH}_3$  to  $\text{totNO}_3$  between 0 and 1 (free  $\text{NH}_3$ :  $\text{totNH}_3$  minus double  $\text{totSO}_4$ ,  $\text{totNO}_3$ : sum of  $\text{NO}_3^-$  and  $\text{HNO}_3$ ), and the “ $\text{NH}_3$  very rich” represents the ratio of the free  $\text{NH}_3$  to  $\text{totNO}_3$  over 1.

On land, except for areas such as northern Russia, central Africa, and the Arabian Peninsula, the aerosol typically falls within the “ $\text{NH}_3$  very rich” domain. In this domain,  $\text{SO}_4^{2-}$  is fully-neutralized by the available  $\text{NH}_3$ , with sufficient  $\text{NH}_3$  available remaining to additionally neutralize  $\text{NO}_3^-$ , making. As a result,  $\text{NO}_3^-$  becomes the limiting factor in  $\text{NH}_4\text{NO}_3$  the formation of  $\text{NH}_4\text{NO}_3$ . More than 60% of SNA mass is in the sub-micron mode on land, while the super-micron modes (i.e.  $1 - 2.5 \mu\text{m}$ ,  $2.5 - 5 \mu\text{m}$  and  $5 - 10 \mu\text{m}$ ) accounts for a smaller fraction (7% for  $\text{NH}_4^+$ , 15% for

SO<sub>4</sub><sup>2-</sup>, and 18% for NO<sub>3</sub><sup>-</sup>). In polluted regions such as East and South Asia, PM<sub>10</sub> SNA concentrations are three times higher than the global land average. For example, East Asia shows NH<sub>4</sub><sup>+</sup>: 1.88 µg m<sup>-3</sup>, NO<sub>3</sub><sup>-</sup>: 5.31 µg m<sup>-3</sup>, and SO<sub>4</sub><sup>2-</sup>: 2.29 µg m<sup>-3</sup>, while South Asia records NH<sub>4</sub><sup>+</sup>: 1.58 µg m<sup>-3</sup>, NO<sub>3</sub><sup>-</sup>: 3.68 µg m<sup>-3</sup>, and SO<sub>4</sub><sup>2-</sup>: 3.18 µg m<sup>-3</sup>. Free NH<sub>3</sub> is abundant in regions like southern North America, Europe, South Asia, and East Asia, with mean free NH<sub>3</sub>/total NO<sub>3</sub><sup>-</sup> ratios of 2.11, 3.77, 5.30, and 3.78, respectively. Over 75% of SNA mass in these regions is concentrated in the 0–1 µm and 1–2.5 µm size ranges. The NH<sub>3</sub> surplus reflects recent trends in precursor emissions, with stable or increasing NH<sub>3</sub> emissions contrasting with declining SO<sub>2</sub> and NO<sub>x</sub> emissions (Zheng et al., 2018; Hand et al., 2012; Russell et al., 2012). In the Middle East, particularly the Arabian Peninsula, aerosol is dominated by desert dust with negligible NH<sub>3</sub> emissions. The “SO<sub>4</sub><sup>2-</sup> rich” and “NO<sub>3</sub><sup>-</sup> rich” domains predominate, where insufficient NH<sub>3</sub> levels are insufficient to neutralize acidic components fully, limiting NH<sub>4</sub>NO<sub>3</sub> formation. Mean concentrations of NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, and SO<sub>4</sub><sup>2-</sup> in this region are 0.72 µg m<sup>-3</sup>, 2.07 µg m<sup>-3</sup>, and 3.19 µg m<sup>-3</sup>, respectively. Over 70% of NO<sub>3</sub><sup>-</sup> resides in the super-micron modes due to interactions with sea salt and crustal dust, which shift NO<sub>3</sub><sup>-</sup> from sub-micron to super-micron modes (Chen et al., 2020; Koçak et al., 2007; Karydis et al., 2016). Sub-micron NH<sub>4</sub><sup>+</sup> and SO<sub>4</sub><sup>2-</sup> remain dominant, accounting for 96% and 79% of their respective fractions, consistent with Osipov et al. (2022), who identified anthropogenic sources as the primary contributors to fine particles in the region.


## 5.2 NH<sub>3</sub> column concentration

Figure 5-(a)5a shows the global distribution of NH<sub>3</sub> column concentration. The global area-weight mean value of NH<sub>3</sub> column concentration is 0.80 mg m<sup>-2</sup>, with the highest value up to 30 mg m<sup>-2</sup> in the Indo-Gangetic Plain of India and the lowest value of less than 0.01 mg m<sup>-2</sup> in the remote oceans of the Southern Hemisphere and the South Pole. Compared with previous studies that investigated NH<sub>3</sub> column concentrations based on satellite retrievals (Van Damme et al., 2014; Van Damme et al., 2015; Zhou et al., 2024), our results can capture the distribution of NH<sub>3</sub> hotspots worldwide, including the Indo-Gangetic Plain, the North China Plain, and West Africa and Amazonia, where biomass burning is dominant (Van Damme et al., 2018). However, Van Damme et al. (2018) pointed out that two-thirds of the NH<sub>3</sub> emission hotspots are underestimated by at least one order of magnitude in the NH<sub>3</sub> emission inventory EDGAR (CAMSGLOBANT used in this study is derived from EDGAR, see 2.1.1). Given such an underestimation in the current NH<sub>3</sub> emission inventory, we further improve the emission by applying a new inventory and updating the current inventory using a top-down method introduced in Section 2.1. The simulation results are discussed in Section 7.

Formatted: Highlight

Formatted: Font: Times New Roman, 10 pt, English (United States)

Formatted: Indent: First line: 0.51 cm, Line spacing: 1.5 lines, Widow/Orphan control, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers



**Figure 5.** Global distribution of (a) the average  $\text{NH}_3$  column concentration ( $\text{mg m}^{-2}$ ) for the Base Case, and (b) the absolute difference between IASI satellite retrieval and Base case, from 2010 to 2012.

Our simulated  $\text{NH}_3$  column concentrations show good agreement with the IASI satellite observations (Figure 5b), with global land mean values at a comparable level of  $1.66 \text{ mg m}^{-2}$ . However, regional biases can be significant, with an NMB of  $-0.32$ . The model overestimates  $\text{NH}_3$  column concentrations over regions such as India, China, western America, and northeastern Africa, while it has a negative bias in high-latitude regions. It is also important to note that biases in the IASI satellite products have been identified in previous studies (Dammers et al., 2017; Dammers et al., 2019), more details are discussed in Section 2.1.2.

### 5.3 Global $\text{NH}_3$ budget

The global budget for  $\text{NH}_3$  and  $\text{NH}_4^+$  is summarized in Table 8. Given a global  $\text{NH}_3$  emission input of  $74.3773.3 \text{ Tg yr}^{-1}$ , the global burden and lifetime derived from the base case simulation are  $0.414 \text{ Tg}$  and  $2.040 \text{ days}$ , respectively; for  $\text{NH}_4^+$ , the global burden and lifetime are  $0.343 \text{ Tg}$  and  $3.461 \text{ days}$ , respectively. Based on the simulations of nine models, Bian et al. (2017) assessed that given the average  $\text{NH}_3$  emission input of  $76.384 \text{ Tg yr}^{-1}$ , the average global burden and lifetime for  $\text{NH}_3$  and  $\text{NH}_4^+$  are  $0.292 \text{ Tg}$  and  $0.727 \text{ days}$ , and  $0.323 \text{ Tg}$  and  $4.3 \text{ days}$ , respectively. Compared to Bian et al. (2017), our study uses the same  $\text{NH}_3$  emission input; however, the global burden and lifetime of  $\text{NH}_3$  derived from the base case are larger. This potential overestimation may be attributed to the wet deposition scheme used in EMAC. The scavenging scheme (SCAV) accounts for pH adjustments in  $\text{NH}_3$  dissolution. More specifically, the EMAC model implicitly determines the effective Henry's law constant by solving a system of coupled ordinary differential equations, explicitly representing liquid-phase processes in clouds and raindrops, including dissociation, acid-base equilibria, redox reactions, and photolysis (Tost et al., 2006). This approach ensures a comprehensive calculation of total wet deposition for  $\text{NH}_4^+$  and  $\text{NH}_3$ . Notably, the  $\text{NH}_3$  burdens simulated in the AeroCom model intercomparison by Bian et al. (2017) exhibit significant variability, spanning a factor of 17. This wide range underscores the sensitivity of atmospheric  $\text{NH}_4^+$  and  $\text{NH}_3$  burdens and lifetimes to model domain definitions, deposition pathways, and  $\text{NH}_3$  chemical processes across different models (Ge et al., 2022).

**Table 8.** Atmospheric budget of  $\text{NH}_3$ ,  $\text{NH}_4^+$ , and  $\text{NH}_x$  ( $\text{NH}_3 + \text{NH}_4^+$ ).

Simulation Case	Specie	Emission ( $\text{Tg yr}^{-1}$ )	Burden ( $\text{Tg}$ )	Dry Deposition ( $\text{Tg yr}^{-1}$ )	Wet Deposition ( $\text{Tg yr}^{-1}$ )	Lifetime <sup>a</sup> (day)
Base	$\text{NH}_3$	73.273	0.414	28.222	-	2.040
CEDS		77.131	0.434	29.909	-	2.041.9
Top-Dep		84.717	0.414	34.364	-	1.777

Base		-	0.343	1.462	34.697	3.461
CEDS	NH <sub>4</sub> <sup>+</sup>	-	0.354	1.253	35.778	3.459
Top-Dep		-	0.353	1.525	36.192	3.392.9
Base		-	0.757	29.384	34.697	3.685
						(4.270) <sup>b</sup>
CEDS	NH <sub>x</sub>	-	0.788	31.452	35.778	3.668
						(4.254)
Top-Dep		-	0.767	35.909	36.212	3.270
						(3.855)

<sup>a</sup>: NH<sub>3</sub> lifetime = Burden/Emission; NH<sub>4</sub><sup>+</sup> lifetime = Burden/(Dry Deposition + Wet Deposition);


<sup>b</sup>: NH<sub>x</sub> lifetime = NH<sub>x</sub> Burden/ NH<sub>3</sub> Emission; in the parentheses, NH<sub>x</sub> lifetime = NH<sub>x</sub> Burden/ (Dry Deposition + Wet Deposition);

Considering the SCAV scavenging scheme, we further calculate the global budget for NH<sub>x</sub> (NH<sub>3</sub> + NH<sub>4</sub><sup>+</sup>) in Table 8. The global burden of NH<sub>x</sub> is 0.757 Tg. Wet and dry deposition contribute almost equally to the sink accounting for 54% and 46%, respectively. The lifetime of NH<sub>x</sub> is 3.685 – 4.270 days, depending on the calculation method. Ge et al. (2022) estimated a global budget for NH<sub>x</sub>, with an input NH<sub>3</sub> emission of 64.485 Tg yr<sup>-1</sup>. They calculated the global burden of NH<sub>x</sub> to be 0.949 Tg, with a lifetime of 4.9–5.2 days and its wet and dry deposition contributing equally to the sink (i.e., 51% and 49% of total deposition, respectively).

## 6. Aerosol acidity

Figure 6 illustrates the global distributions of size-resolved aerosol pH, with regional averages summarized in Table 9. Aerosol pH exhibits marked spatial variations, influenced by land-sea contrasts and regional sources. Over land in the Northern Hemisphere, excluding deserts, aerosols are generally acidic, whereas marine aerosols are alkaline due to sea salt influence. However, high-latitude marine aerosols are generally more acidic than those over remote ocean regions, primarily due to the low liquid water content/long-range transport of anthropogenic pollutants such as H<sub>2</sub>SO<sub>4</sub> and hydrogen ion outflow/HNO<sub>3</sub> from land-continental sources (Myhre et al., 2013; Karydis et al., 2021). The average pH values for land-based aerosols are 4.2 (5–10 μm), 4.2 (2.5–5 μm), 3.9 (1–2.5 μm), and 4.3 (0–1 μm). In contrast, marine aerosol pH values are 6.2 (5–10 μm), 5.8 (2.5–5 μm), 5.1 (1–2.5 μm), and 5.1 (0–1 μm). The deserts of North Africa, the Middle East, and the Gobi maintain the highest pH values (>7) across all size ranges, driven by non-volatile cations (e.g., Ca<sup>2+</sup>, Mg<sup>2+</sup>) that neutralize acidic components and enhance water uptake.

Regionally, higher pH values (5–5.7) in the Middle East are attributed to airborne dust, while coastal areas like the western Arabian Gulf have lower aerosol pH (<3 in 0–1 μm) due to elevated sulfate concentrations. In South Asia, abundant NH<sub>3</sub> emissions keep pH values between 5.4 and 4.9 despite high SO<sub>2</sub> and NO<sub>x</sub> emissions. East Asia shows a distinct pH gradient, with desert regions in the northwest reaching pH >7 across all sizes, and southeast coastal areas exhibiting low pH (2–4), linked to significant SO<sub>2</sub> emissions and sulfate formation. Europe (pH 3.7–4.1) and North America (pH 3.2–3.6) demonstrate moderate acidity, with the western USA exhibiting higher values (4–6) due to desert influences. In South America, pH ranges from 3.9 to 4.5, with coastal regions exceeding 6 due to sea salt.



**Figure 6.** Global distribution of surface average aerosol pH values in the size ranges: (a) 0 – 1  $\mu\text{m}$ , (b) 1 – 2.5  $\mu\text{m}$ , (c) 2.5 – 5  $\mu\text{m}$ , (d) 5 – 10  $\mu\text{m}$ , derived from Base case from 2010 to 2012.

585 An increasing trend in pH is observed from North America to Europe and East Asia, consistent with prior studies (Zhang et al., 2021a; Ding et al., 2019; Guo et al., 2017a). Aerosol alkalinity is driven by  $\text{NH}_4^+$  and non-volatile cations, which neutralize  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$ , while water-soluble ions (WSIs) enhance liquid water uptake. East Asia exhibits the highest pH among regions, facilitated by relatively lower  $\text{SO}_4^{2-}$ , abundant  $\text{NH}_4^+$  and WSIs, and significant  $\text{NO}_3^-$  and non-volatile cations in the coarse modes. These chemical properties outweigh the influence of meteorological effects such as differences in temperature and humidity.

590

**Table 9.** Size-resolved pH values across the globe and regions from simulation cases.

Region	0 – 1 $\mu\text{m}$			1 – 2.5 $\mu\text{m}$			2.5 – 5 $\mu\text{m}$			5 – 10 $\mu\text{m}$		
	Base <sup>a</sup>	Meta <sup>a</sup>	noNH <sub>3</sub> <sup>a</sup>	Base <sup>a</sup>	Meta <sup>a</sup>	noNH <sub>3</sub> <sup>a</sup>	Base <sup>a</sup>	Meta <sup>a</sup>	noNH <sub>3</sub> <sup>a</sup>	Base <sup>a</sup>	Meta <sup>a</sup>	noNH <sub>3</sub> <sup>a</sup>
Globe	4.9	4.19	3.28	4.83	4.3	4.09	5.38	5	4.82	5.75	5.47	5.46
Land	4.29	2.96	1.06	3.94	2.9	1.71	4.19	3.36	2.51	4.23	3.54	3.11
Ocean	5.1	4.58	3.99	5.11	4.75	4.86	5.76	5.52	5.56	6.24	6.08	6.21
North America	3.62	2.63	1.01	3.16	2.38	1.17	3.31	2.63	1.83	3.38	2.81	2.34
South America	4.11	3.25	1.46	3.85	3.26	2.26	4.4	4	3.25	4.52	4.22	3.84
Europe	4.09	2.97	0.84	3.65	2.8	1.27	3.69	2.92	1.91	3.75	3.08	2.67
Middle East	5.02	1.94	-0.17	5.25	2.08	0.51	5.47	3.59	1.86	5.66	4.18	3.85
South Asia	5.42	2.8	-0.2	4.86	2.33	0.43	5	2.73	1.37	5.04	2.88	2.16
East Asia	5.15	3.23	0.5	4.65	3.32	1.16	4.56	3.24	1.76	4.54	3.26	2.3

<sup>a</sup>: average pH value.

595 Contrary to previous findings (e.g. Kakavas et al., 2021; Ding et al., 2019), which suggest decreasing aerosol acidity with increasing particle size, pH in the base case (stable state) for 0–1  $\mu\text{m}$  exceeds that of 1–2.5  $\mu\text{m}$  over many regions, excluding oceans and the Middle East. This is examined through three perspectives. First, [the assumption of phase state. As described in Section 2.2.1, the pH predictions based on the metastable state tend to be lower than those based on the stable state under low RH conditions. Although previous studies concluded that the](#)

**Formatted:** Indent: First line: 0.51 cm, No widow/orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers, Tab stops: Not at 1.27 cm

choice between stable and metastable assumption in ISORROPIA has no significant impact on pH predictions (Song et al., 2018), our sensitivity simulation (Meta case) assuming metastable conditions revealed (Meta case) yields consistently lower pH values across all sizes, with the greatest particle size ranges. The differences in pH relative to the base case are shown in Figure 7 and Table 9. The most substantial reductions are observed in regions with high mineral cations and/or low relative humidity (RH), such as South Asia and the Middle East. In the metastable state, all anions remain aqueous, resulting in, where the pH decreases by more acidic aerosols than 2 units. These results are consistent with the findings of Karydis et al. (2021) and Milousis et al. (2024). For instance, Karydis et al. (2021) reported that the stable-state assumption yields global average pH values approximately 0.5 units higher than those from the metastable assumption, comparable to the magnitude of differences calculated in our study (Table 9). While the Meta case explains some discrepancies, pH in the 0–1  $\mu\text{m}$  size range remains higher relative to the 1–2.5  $\mu\text{m}$  range in several regions (Table 5). Another sensitivity simulation, removing  $\text{NH}_3/\text{NH}_4^+$  emissions (no  $\text{NH}_3/\text{NH}_4^+$  case), significantly reduced pH across all sizes over land, especially in  $\text{NH}_3$ -rich regions like South Asia and the Middle East, where 0–1  $\mu\text{m}$  pH dropped below zero. This indicates increased pH sensitivity in the fine size ranges (i.e., 0–1  $\mu\text{m}$  and 1–2.5  $\mu\text{m}$ ), particularly in regions where  $\text{NH}_3$  availability is abundant. Excluding  $\text{NH}_3$ , results in a consistent pH increase with particle size across all regions. Finally, as discussed in Section 4.1, the comparison with observed size-resolved mass concentrations from the EMEP network revealed and  $\text{NH}_3/\text{NH}_4^+$  partitioning ratio reveals an underestimation of acidic components ( $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$ ) and an overestimation of alkaline components in the 0–1  $\mu\text{m}$  size range. This, along with an inadequate treatment of  $\text{NH}_3/\text{NH}_4^+$  partitioning. This chemical imbalance contributes to the anomalously high simulated pH in the fine mode (0–1  $\mu\text{m}$ ). For instance, the observed  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$  and  $\text{NH}_4^+$  in 0–1  $\mu\text{m}$  accounts for 91%, 59% and 67% of 0–2.5  $\mu\text{m}$  at the Montseny site (41.78° N, 2.35° E) during 2010–2012 (observed  $\text{NH}_4^+$  only available in 2010), respectively, compared to the accounting of 79%, 54% and 77% in simulation, highlighting a need for improved representation of aerosol composition in fine modes. In addition, the observed mean  $\text{NH}_4^+$  partitioning ratios ( $\text{NH}_4^+ / (\text{NH}_3 + \text{NH}_4^+)$ , in  $\mu\text{mole m}^{-3}$ ) during 2010–2012 from EMEP, NNDMN, and AMoN networks are 49%, 49%, and 50%, respectively. Our model shows low biases, with the mean ratios of 32%, 39%, and 41%. These results reinforce the necessity of refining the treatment of  $\text{NH}_3/\text{NH}_4^+$  partitioning to better capture the size-resolved trend of aerosol acidity.

This study calculates Part of the discrepancy in size-resolved aerosol pH based on a log-normal calculations between models may arise from differences in how particle size distributions and gas-particle partitioning are treated. EMAC model uses a lognormal size distribution, differing from other approaches. For instance, and applies ISORROPIA separately to each size mode to calculate gas-aerosol partitioning. In contrast, other studies (e.g., Kakavas et al., 2021) used the PMCAMx model, which employs a employ sectional approach to track approaches, where gas-aerosol mass and composition across 10 size bins ranging from 40 nm to 40  $\mu\text{m}$ . The partitioning is first five bins represent performed on the  $\text{PM}_{10}$  fraction (0.04–1.25  $\mu\text{m}$ ), while three bins cover sizes up bulk aerosol phase, and the resulting condensed mass is then distributed across size bins based on the available surface area. These fundamental differences in modeling assumptions can lead to 40  $\mu\text{m}$ . In addition variations in the predicted distribution of aerosol components across size ranges, which in turn affects the calculated size-resolved pH. Moreover, previous studies on investigating size-resolved aerosol acidity also have certain limitations. For example instance, a laboratory

Formatted: Font color: Auto


Formatted: Font color: Auto

Formatted: Font color: Auto, English (United Kingdom)

Formatted: Font color: Auto, English (United Kingdom)

Formatted: English (United Kingdom)

study observed increasing aerosol acidity with decreasing particle size, but this trend only holds for pH below 2 (Craig et al., 2018). Similarly, a field study [reached by Fang et al. \(2017\)](#) reported the same [conclusion trend](#) but lacked [concurrent](#) measurements of key gas-phase species [\(Fang et al., 2017\)](#), which limits the interpretability of the results.



**Figure 7.** Global distribution maps of pH value absolute difference between Meta case and base case in the size ranges of 0 – 1 μm, 1 – 2.5 μm, 2.5 – 5 μm, and 5 – 10 μm, averaged from 2010 to 2012.

## 7. Emission Sensitivity Analysis

The formation of secondary inorganic aerosols is strongly linked to  $\text{NH}_3$  emissions (Wu et al., 2016; Chen et al., 2019; Liang et al., 2024). Xu et al. (2020) and Wang et al. (2015) have highlighted the critical role of gas-particle partitioning of  $\text{NH}_3/\text{NH}_4^+$  in SNA formation, which is influenced by factors such as temperature, aerosol water content, and aerosol acidity (Nenes et al., 2020). Notably, the  $\text{NH}_3/\text{NH}_4^+$  partitioning process buffers aerosol acidity, maintaining stability even amid shifts in acidic species like  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$  (Karydis et al., 2021). These findings suggest that the effects of  $\text{NH}_3$  emissions on SNA formation and aerosol acidity remain [a subject of debate/understudied](#) (Weber et al., 2016; Zheng et al., 2022; Fu et al., 2017; Zou et al., 2024). However, most studies have focused on fine particles ( $\text{PM}_{2.5}$  or  $\text{PM}_{10}$ ), polluted regions (e.g., northern China), and haze episodes (Ge et al., 2019; Gao et al., 2016). Modeling studies often evaluate the effects of  $\text{NH}_3$  emission changes by uniformly altering emission levels within each grid (Pozzer et al., 2017; Fu et al., 2017). While such approaches provide valuable insights, they may lack feasibility as  $\text{NH}_3$  abundance correlates with population density, making uniform changes less representative of real-world scenarios. To address this, we compare the effects of two  $\text{NH}_3$  emission schemes CEDS and Top-Dep on size-resolved SNA mass concentration and aerosol acidity relative to [the](#) base case.


The  $\text{NH}_3$  mass concentration serves as a proxy for  $\text{NH}_3$  emissions, and differences in  $\text{NH}_3$  mass concentrations between the scenarios and the base case (Figure [S6S4](#)) align with corresponding emission differences (Figure 1). Figures [78](#) and [89](#) show the responses of size-resolved  $\text{NH}_4^+$ ,  $\text{NO}_3^-$ , and  $\text{SO}_4^{2-}$  mass concentrations to  $\text{NH}_3$  emission changes in the CEDS and Top-Dep cases, respectively. Similarly, Figure [910](#) illustrates the size-resolved aerosol acidity responses for the two cases. Additional details are provided in supplementary figures: Figures [S5 – S7–S9](#)

(CEDS case) and Figures S44–S43S8–S10 (Top-Dep case) depict the size-resolved responses of  $\text{NH}_4^+$ ,  $\text{NO}_3^-$ , and  $\text{SO}_4^{2-}$  mass concentrations, while Figures S40S11 and S44S12 highlight changes in size-resolved aerosol acidity.

The regional gas-particle partitioning ratios for  $\text{NH}_3/\text{NH}_4^+$  ( $\epsilon(\text{NH}_4^+) = \text{NH}_4^+ / (\text{NH}_4^+ + \text{NH}_3)$ ) and  $\text{NO}_3^-/\text{HNO}_3$  ( $\epsilon(\text{NO}_3^-) = \text{NO}_3^- / (\text{NO}_3^- + \text{HNO}_3)$ ) are shown in Figure S45S13. Regional emission amounts of  $\text{NO}_x$  and  $\text{SO}_2$  from the CAMS and CEDS\_GBD inventories are detailed in Table S1.

7.1 Size-Resolved SNA Response


Regarding the atmospheric budget of  $\text{NH}_3$  (Table 8), a modest increase in global  $\text{NH}_3$  emissions (CEDS case) slightly raises the deposition of  $\text{NH}_3$  and shortens the lifetime. Meanwhile, both the burden and deposition of  $\text{NH}_3$  and  $\text{NH}_4^+$ , while the are increased, with a longer lifetime remains unchanged. In contrast, the larger emission increase in the Top-Dep cases does not alter the  $\text{NH}_3/\text{NH}_4^+$  burden but leads to higher deposition and a shorter lifetime for both species.



**Figure 78.** Bar plots for regional surface SNA mass concentration ( $\mu\text{g m}^{-3}$ ) absolute difference between CEDS case and base case in the four size ranges (a) – (d); change ratio in the four size ranges (e) – (h). The calculation of change ratio in the size range of 0–1  $\mu\text{m}$  is based on the mask of 0.1  $\mu\text{g m}^{-3}$ , the change ratio in the size range of 1–2.5  $\mu\text{m}$ , 2.5–5  $\mu\text{m}$  and 5–10  $\mu\text{m}$  is based on the mask of 0.05  $\mu\text{g m}^{-3}$  (Land: global land; NAm: North America; SAm: South America; Eu: Europe; ME: Middle East; SAs: South Asia; EAs: East Asia).

Across land regions, a small increase in  $\text{NH}_3$  emissions (CEDS case), along with rising  $\text{NO}_x$  and  $\text{SO}_2$  emissions, slightly raises  $\epsilon(\text{NH}_4^+)$  while marginally lowering  $\epsilon(\text{NO}_3^-)$  (Figure S45S13). The SNA mass concentration increases consistently across size ranges, with the most notable growth in  $\text{NH}_4^+$  and  $\text{NO}_3^-$  in the 1–2.5  $\mu\text{m}$  range (19% and 11%, respectively) and  $\text{SO}_4^{2-}$  in the submicron particles (9%). Under a larger  $\text{NH}_3$  emission increase (Top-Dep case),  $\epsilon(\text{NH}_4^+)$  drops significantly, and  $\epsilon(\text{NO}_3^-)$  decreases slightly. The SNA response becomes more pronounced, with substantial increases in the 1–2.5  $\mu\text{m}$  range ( $\text{NH}_4^+$ : 104%,  $\text{NO}_3^-$ : 41%,  $\text{SO}_4^{2-}$ : 23%), while  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$  decrease

in coarser particles. These shifts are linked to higher  $\text{NH}_3$  emissions in relatively clean, high-latitude regions with low  $\text{NH}_3$  flux (Figure 1c), which elevate  $\text{NH}_3$  concentrations (Figure S6S4) enhancing  $\text{NH}_4^+$  formation (Figure S4+S8).



**Figure 8.** Bar plots for regional surface SNA mass concentration ( $\mu\text{g m}^{-3}$ ) absolute difference between Top-Dep case and base case in the four size ranges (a)–(d); change ratio in the four size ranges (e)–(h). The calculation of change ratio in the size range of 0–1  $\mu\text{m}$  is based on the mask of 0.1  $\mu\text{g m}^{-3}$ , the change ratio in the size range of 1–2.5  $\mu\text{m}$ , 2.5–5  $\mu\text{m}$  and 5–10  $\mu\text{m}$  is based on the mask of 0.05  $\mu\text{g m}^{-3}$  (Land: global land; NAm: North America; SAm: South America; Eu: Europe; ME: Middle East; SAs: South Asia; EAs: East Asia).

Regional analysis of SNA responses in Europe shows that a 23% reduction in  $\text{NH}_3$  emissions, along with increases in  $\text{NO}_x$  and  $\text{SO}_2$  emissions (CEDS case), raises  $\bar{c}(\text{NH}_4^+)$  slightly while causing a minor drop in  $\bar{c}(\text{NO}_3^-)$ . The SNA mass increases mainly in the 1–2.5  $\mu\text{m}$  range, with a slight decrease in  $\text{SO}_4^{2-}$  in coarser particles. Since Europe has abundant  $\text{NH}_3$ , reductions are offset by existing availability and rising  $\text{NO}_x$  and  $\text{SO}_2$  levels, leading to additional  $\text{NH}_4\text{NO}_3$  and  $(\text{NH}_4)_2\text{SO}_4$  formation. With higher  $\text{NH}_3$  emissions (Top-Dep case), the response becomes more complex: SNA decreases in the submicron particles (Figure 8e9e) but increases in larger particles (Figure 8f9f–g), with the largest growth seen in  $\text{NH}_4^+$  (~50%).

Formatted: Font: Italic

Formatted: Font: Italic




Figure 9. The same as Figure 8, but for the difference between Top-Dep case and base case.

In East Asia, reducing  $\text{NH}_3$  emissions while increasing  $\text{NO}_x$  and  $\text{SO}_2$  emissions (CEDS case) raises  $\epsilon(\text{NH}_4^+)$  and slightly increases  $\epsilon(\text{NO}_3^-)$ . SNA grows across all particle sizes, with the most significant changes in 1–2.5  $\mu\text{m}$  range (Figure 748f). A similar  $\text{NH}_3$  reduction in the Top-Dep case produces comparable  $\epsilon(\text{NH}_4^+)$  and  $\epsilon(\text{NO}_3^-)$  changes, with SNA mainly decreasing in the submicron particles. The response in East Asia resembles that of Europe, where abundant  $\text{NH}_3$  buffers SNA changes. In South Asia,  $\text{NH}_3$  and  $\text{NO}_x$  emissions rise in the CEDS case while  $\text{SO}_2$  emissions decline, slightly lowering  $\epsilon(\text{NH}_4^+)$  and increasing  $\epsilon(\text{NO}_3^-)$ .  $\text{NH}_4^+$  and  $\text{NO}_3^-$  concentrations grow across all size ranges (Figure 8), with the largest  $\text{NH}_4^+$  increase found in 1–2.5  $\mu\text{m}$  particles (66%) and  $\text{NO}_3^-$  rising in both submicron and 1–2.5  $\mu\text{m}$  particles (~60%).  $\text{SO}_4^{2-}$  decreases in submicron particles but increases in coarser ones (25%). Conversely, reducing  $\text{NH}_3$  emissions in the Top-Dep case raises  $\epsilon(\text{NH}_4^+)$  and lowers  $\epsilon(\text{NO}_3^-)$ , leading to SNA declines, especially in submicron particles. South Asia, with abundant  $\text{NH}_3$ , shows  $\text{NO}_3^-$  as the limiting factor for  $\text{NH}_4\text{NO}_3$  formation, driving strong  $\epsilon(\text{NO}_3^-)$  and  $\text{NO}_3^-$  responses in finer particles.

In North America,  $\text{NH}_3$  and  $\text{SO}_2$  emissions decrease, while  $\text{NO}_x$  emissions slightly rise (CEDS case).  $\epsilon(\text{NH}_4^+)$  increases marginally, whereas  $\epsilon(\text{NO}_3^-)$  declines slightly.  $\text{NH}_4^+$  and  $\text{SO}_4^{2-}$  show minor reductions in submicron particles, while  $\text{NO}_3^-$  increases. In the Top-Dep case, a sharp rise in  $\text{NH}_3$  emissions, mainly over Canada and Greenland, significantly lowers  $\epsilon(\text{NH}_4^+)$ , stabilizes  $\epsilon(\text{NO}_3^-)$ , and increases  $\text{NH}_4^+$  and  $\text{SO}_4^{2-}$  in 1–2.5  $\mu\text{m}$  particles. In South America, a small  $\text{NH}_3$  emission rise (CEDS case) has little effect on  $\epsilon(\text{NH}_4^+)$  or  $\epsilon(\text{NO}_3^-)$ , resulting in minimal aerosol composition changes. However, a substantial  $\text{NH}_3$  increase (Top-Dep case) significantly boosts  $\text{NH}_3$  concentrations, reduces  $\epsilon(\text{NH}_4^+)$  and  $\epsilon(\text{NO}_3^-)$ , and shifts aerosol partitioning to smaller particles, particularly in central regions. Decreased  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$  in surrounding areas (Figures 842–859–10) suggest  $\text{NH}_3$  is neutralizing transported species, explaining the observed  $\epsilon(\text{NO}_3^-)$  reduction. Changes in other particle size ranges are minimal. In the Middle East,  $\text{NH}_3$  emissions rise moderately (20%), along with slight  $\text{NO}_x$  and  $\text{SO}_2$  increases (CEDS case).  $\epsilon(\text{NH}_4^+)$  and  $\epsilon(\text{NO}_3^-)$  remain stable, with  $\text{NH}_4^+$  and  $\text{NO}_3^-$  increasing mainly in submicron particles (~40%), while  $\text{SO}_4^{2-}$  grows across

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

sizes (~15%). A minor NH<sub>3</sub> emission drop (Top-Dep case) reduces NH<sub>3</sub> concentrations but enhances SNA in 1–2.5 μm particles, highlighting compensatory NH<sub>3</sub> effects in "NH<sub>3</sub> very rich" domains.


Overall, higher NH<sub>3</sub> emissions enhance SNA formation, particularly in the fine size ranges (0–1 μm and 1–2.5 μm). Greater NH<sub>4</sub><sup>+</sup> formation (e.g., Top-Dep case) depletes NO<sub>3</sub><sup>-</sup> and SO<sub>4</sub><sup>2-</sup> from coarse size ranges, leading to decreases in 2.5–5 μm and 5–10 μm. In low-SNA regions (e.g., South America, Greenland), NH<sub>3</sub> increases have limited SNA impacts but substantially elevate NH<sub>3</sub> concentrations. In "NH<sub>3</sub> very rich" regions (e.g., East Asia, Europe), NH<sub>3</sub> reductions alone may still increase  $\delta(\text{NH}_4^+)$ , promoting further SNA formation. These findings align with Zou et al. (2024), emphasizing the greater effectiveness of NO<sub>x</sub> reductions in PM<sub>2.5</sub> mitigation compared to NH<sub>3</sub> or SO<sub>2</sub> control in ammonia-rich environments. Coordinated control of precursor emissions is crucial for mitigating air pollution, especially in heavily polluted regions (Wen et al., 2024).

## 7.2 Aerosol acidity response

### 7.2.1 Size-resolved pH changes

Among atmospheric buffering agents (e.g., conjugate acid-base pairs like NH<sub>3</sub>/NH<sub>4</sub><sup>+</sup> and CO<sub>2</sub>/HCO<sub>3</sub><sup>-</sup>, as well as organic acids), the NH<sub>3</sub>/NH<sub>4</sub><sup>+</sup> acid-base pair exhibits the largest buffering capacity for aerosols, dominating much of the continental urban areas (Zheng et al., 2023; Zheng et al., 2020). When NH<sub>3</sub> emissions are entirely eliminated, changes in oceanic aerosol pH values are negligible compared to the base case (noNH<sub>3</sub> case, Table S9 and Figure S5S14). However, aerosols over land become significantly more acidic, with pH values in the 0–1 μm size range dropping to -0.2 in South Asia and -0.17 in the Middle East (Figure 10). The most substantial pH decreases occur in 0–1 μm aerosols, primarily in South Asia, East Asia, Europe, North America, and South America, consistent with the "NH<sub>3</sub>-buffered regions" identified by Zheng et al. (2020).

Interestingly, while neither the Middle East nor northern Africa are categorized as "NH<sub>3</sub>-buffered regions," noticeable pH decreases are observed in the 0–1 μm and 1–2.5 μm size ranges over the Middle East, including Egypt and Libya. In these regions, the SNA is dominated by "SO<sub>4</sub><sup>2-</sup> rich" and "NO<sub>3</sub><sup>-</sup> rich" chemical domains (Figure 4), where NH<sub>4</sub><sup>+</sup> cannot fully neutralize the acidic components, resulting in available NO<sub>3</sub><sup>-</sup> and SO<sub>4</sub><sup>2-</sup>. This results in excess acidic components, particularly in the 0–1 μm and 1–2.5 μm size ranges (Figure S2S3). Without NH<sub>3</sub> emissions, the abundance of NO<sub>3</sub><sup>-</sup> and SO<sub>4</sub><sup>2-</sup> further increases aerosol acidity.



**Figure 910.** Bar plots for pH value absolute difference between CEDS/Top-Dep case and base case in the four size ranges (a)–(d), with the two dashed lines representing the value of 0.1 and -0.1, respectively (Land: global land; NAm: North America; SAm: South America; Eu: Europe; ME: Middle East; SAs: South Asia; EAs: East Asia).

Compared to the base case, size-resolved pH values in the CEDS case show an average decrease of -0.13 to -0.15 units over land. The most pronounced decreases occur in remote regions of the Northern Hemisphere (Figure S40S11), where  $\text{NH}_3$  emission flux is relatively low. In contrast, the Top-Dep case exhibits an average pH increase of 0.24 units in the 0–1  $\mu\text{m}$  range over land, driven by the higher  $\text{NH}_3$  emission flux. This  $\text{NH}_3$ -driven alkalization effect is less pronounced in the 1–2.5  $\mu\text{m}$  range due to counteracting acidification effects from newly formed acidic components, as suggested by Zheng et al. (2024). Notably, pH decreases of -0.10 units are found in the 2.5–5  $\mu\text{m}$  and 5–10  $\mu\text{m}$  size ranges.

In East Asia, the pH changes in the CEDS case are concentrated in the 2.5–5  $\mu\text{m}$  and 5–10  $\mu\text{m}$  ranges (-0.10 units in both), with minimal changes in 0–1  $\mu\text{m}$  and 1–2.5  $\mu\text{m}$ . In the Top-Dep case, however, pH changes are more drastic across all size ranges: 0–1  $\mu\text{m}$  (-0.17 units), 1–2.5  $\mu\text{m}$  (-0.31 units), 2.5–5  $\mu\text{m}$  (-0.35 units), and 5–10  $\mu\text{m}$  (-0.33 units). These discrepancies between cases highlight that a reduction in  $\text{NH}_3$  emissions enhances aerosol acidity (as in the Top-Dep case), but this effect can be partially offset by shifts in  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  mass concentrations (as in the CEDS case). For instance, in the CEDS case, decreases in  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  concentrations result in pH rises of 0.15–0.25 units across all size ranges compared to the Top-Dep case.

Our simulation results align with other studies. For example, Song et al. (2019) found that a ~0.3 increase in  $\log_{10}[\text{NH}_3]$  contributed to a 0.3–0.4 unit rise in  $\text{PM}_{10}$  pH in Beijing during winter between 2014/2015 and 2018/2019. Concurrent changes in aerosol composition (increased  $\text{NO}_3^-$  and reduced  $\text{SO}_4^{2-}$  and  $\text{Cl}^-$ ) led to a modest 0.1 units pH increase. Similarly, Zhou et al. (2022) observed a minor  $\text{PM}_{2.5}$  pH decrease of -0.24 units in the Yangtze River Delta region from 2011 to 2019, despite significant changes in aerosol composition resulting from the Air Pollution Prevention and Control Action Plan.

### 7.2.2 Drivers of pH Changes

We further quantify the changes in pH values between the CEDS/Top-Dep cases and the base case, isolating the contributions from changes in  $\text{H}^+$  and  $\text{H}_2\text{O}$  concentrations (detailed in Section 2.2.2). The results are presented in Figures S46S15 and S47S16.

In the CEDS case, the decreased pH values across all four aerosol size ranges are primarily driven by changes in  $\text{H}^+$  concentrations, particularly over the remote regions of the Northern Hemisphere. The effects of  $\text{H}_2\text{O}$  concentration changes are particularly relevant in the Middle East and South Asia. Notably, a substantial pH increase is observed in South Asia, with increases of 0.35, 0.43, 0.44, and 0.63 units in the 0–1  $\mu\text{m}$ , 1–2.5  $\mu\text{m}$ , 2.5–5  $\mu\text{m}$ , and 5–10  $\mu\text{m}$  size ranges, respectively. These increases are predominantly caused by changes in  $\text{H}^+$  concentrations, which fully counteract the pH-decreasing effects of changes in  $\text{H}_2\text{O}$  concentrations.

In the Top-Dep case, although changes in  $\text{H}^+$  concentrations significantly enhance pH values in the 0–1  $\mu\text{m}$  size range, the pH-decreasing effects of changes in  $\text{H}_2\text{O}$  concentrations are distributed across all four size ranges over the entire land surface. In East Asia, both  $\text{H}^+$  and  $\text{H}_2\text{O}$  concentration changes contribute substantially to the observed pH decreases, highlighting their combined impact.

In summary,  $\text{NH}_3$  emissions play a crucial role in maintaining terrestrial aerosols at moderately acidic levels,

particularly in the fine-mode size range of 0–1  $\mu\text{m}$ . When  $\text{NH}_3/\text{NH}_3$  emissions are completely removed in the model, the largest pH decreases are found in this size range. Our findings underscore that, in the fine-mode ranges (0–1  $\mu\text{m}$  and 1–2.5  $\mu\text{m}$ ), pH changes closely correspond to variations in  $\text{NH}_3$  emissions, reflecting the distinct sensitivity of size-resolved pH to  $\text{NH}_3$  levels. In the coarser size ranges (2.5–5  $\mu\text{m}$  and 5–10  $\mu\text{m}$ ), pH changes demonstrate a consistent trend with variations in  $\text{NH}_3$  emissions in the fine-mode ranges (0–1  $\mu\text{m}$  and 1–2.5  $\mu\text{m}$ ) based on our findings. Overall, pH changes in the fine-mode aerosols (0–1  $\mu\text{m}$  and 1–2.5  $\mu\text{m}$ ) show a consistent response to variations in  $\text{NH}_3$  emissions. However, increases in  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  associated with elevated  $\text{NH}_3$  emissions can partially offset the expected rise in pH (e.g., over land areas in the 1–2.5  $\mu\text{m}$  range in the CEDS case). Furthermore, in regions with high  $\text{SO}_2$  and  $\text{NO}_x$  emissions, such as East Asia under the CEDS inventory, the enhanced formation of  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  can counteract the pH-lowering effect of reduced  $\text{NH}_3$  emissions. In the coarser size ranges (2.5–5  $\mu\text{m}$  and 5–10  $\mu\text{m}$ ), however, pH changes are governed by the competing influences of  $\text{H}^+$  and  $\text{H}_2\text{O}$  concentration changes.

## 8. Conclusion

This study applied three distinct ammonia ( $\text{NH}_3$ ) emission schemes to the global atmospheric chemistry and climate model (EMAC) to assess the impact of  $\text{NH}_3/\text{NH}_3$  emissions on the size-resolved sulfate-nitrate-ammonium (SNA) aerosol composition and aerosol acidity. It also explores the synergistic effects arising from concurrent changes in  $\text{NH}_3$ ,  $\text{SO}_2$ , and  $\text{NO}_x$  emissions. The emission schemes included two bottom-up inventories, and one inventory updated through a top-down approach. By simulating the aerosol size ranges of 0–1  $\mu\text{m}$ , 1–2.5  $\mu\text{m}$ , 2.5–5  $\mu\text{m}$ , and 5–10  $\mu\text{m}$ , the study provides a comprehensive analysis/assessment of the role of  $\text{NH}_3$  how  $\text{NH}_3$  emissions influence global aerosol composition/chemistry and acidity across different particle size fractions.

The model accurately captures the distribution of global  $\text{NH}_3$  hotspots, but comparisons with observational datasets reveal positive biases in  $\text{NH}_3/\text{NH}_3$  concentrations and underestimations of  $\text{NH}_4^+/\text{NH}_4^+$  in some regions such as China, North America, and Europe. This discrepancy suggests inefficiencies or oversimplifications in the  $\text{NH}_3/\text{NH}_3/\text{NH}_3/\text{NH}_4^+$  partitioning treatment, with insufficient  $\text{NH}_4^+/\text{NH}_4^+$  generated despite  $\text{NH}_3/\text{NH}_3$  availability. In East and Southeast Asia,  $\text{NH}_4^+/\text{NH}_4^+$  concentrations are well-simulated, but significant discrepancies were found for  $\text{NO}_3^-/\text{NO}_3^-$  and  $\text{SO}_4^{2-}/\text{SO}_4^{2-}$ , likely due to the absence of heterogeneous oxidation processes in the model. The simulated global  $\text{NH}_3$  burden and lifetime are higher than reported in related studies, attributed to the wet deposition scheme used in EMAC, which accounts for pH adjustments for  $\text{NH}_3$  dissolution.

Over oceans,  $\text{NH}_3/\text{NH}_3$  is entirely converted to  $\text{NH}_4^+/\text{NH}_4^+$ , while  $\text{SO}_4^{2-}/\text{SO}_4^{2-}$  remains partially neutralized due to low  $\text{NH}_3/\text{NH}_3$  emissions and high  $\text{SO}_4^{2-}/\text{SO}_4^{2-}$  levels from biogenic dimethyl sulfide (DMS) emissions and  $\text{NO}_3^-/\text{NO}_3^-$  and  $\text{SO}_2/\text{SO}_2$  from shipping. Marine  $\text{SO}_4^{2-}/\text{SO}_4^{2-}$  and  $\text{NO}_3^-/\text{NO}_3^-$  are dominant in the super-micron modes, with mass fractions of 70% and 58%, respectively. Over land,  $\text{SO}_4^{2-}$  is total ammonia (the sum of  $\text{NH}_3$  and  $\text{NH}_4^+$ ) generally fully neutralized by  $\text{NH}_3$ , exceeds  $\text{SO}_4^{2-}$ , enabling more complete neutralization, except in regions such as northern Russia, central Africa, and the Arabian Peninsula. Terrestrial  $\text{NO}_3^-/\text{NO}_3^-$  is also largely neutralized by  $\text{NH}_3/\text{NH}_3$ , resulting in over 60% of SNA concentrated in the sub-micron mode. In polluted areas such as East and South Asia,

sub-micron SNA fractions exceed 70%, with South Asia exhibiting nearly 90% sub-micron  $\text{NH}_4^+/\text{NH}_4^+$ . In contrast, the Middle East, dominated by dust and with minor  $\text{NH}_3$  emissions, shows incomplete neutralization of  $\text{SO}_4^{2-}/\text{SO}_4^{2-}$  and  $\text{NO}_3^-/\text{NO}_3^-$ . Here, nearly 70% of  $\text{NO}_3^-/\text{NO}_3^-$  resides in the super-micron modes, while  $\text{NH}_4^+/\text{NH}_4^+$  and  $\text{SO}_4^{2-}/\text{SO}_4^{2-}$  dominate the sub-micron mode with respective mass fractions of 96% and 79%.

In the Northern Hemisphere, terrestrial aerosols are generally more acidic than marine aerosols, except in desert regions. Remote marine and desert aerosols remain neutral due to the alkaline influence of sea salt and non-volatile cations in dust, which enhance water uptake and neutralize  $\text{SO}_4^{2-}/\text{SO}_4^{2-}$  and  $\text{NO}_3^-/\text{NO}_3^-$ . However, at high latitudes, marine aerosols become more acidic due to the long-range transport of anthropogenic pollutants from continental sources. The 0–1  $\mu\text{m}$  size range exhibits higher pH values than the 1–2.5  $\mu\text{m}$  range in many regions, a trend influenced by several factors. These include the assumption of a thermodynamically stable aerosol phase in the ISORROPIA model, the high sensitivity of aerosol acidity to  $\text{NH}_3$  in the 0–1  $\mu\text{m}$  range and observed lower concentrations of acidic components ( $\text{SO}_4^{2-}/\text{SO}_4^{2-}$  and  $\text{NO}_3^-/\text{NO}_3^-$ ) in this size range compared to larger sizes.

An 18% increase in  $\text{NH}_3$  emissions over land leads to a significant increase in SNA concentrations in the 1–2.5  $\mu\text{m}$  size range ( $\text{NH}_4^+/\text{NH}_4^+$ : 104%,  $\text{NO}_3^-/\text{NO}_3^-$ : 41%,  $\text{SO}_4^{2-}/\text{SO}_4^{2-}$ : 23%), while coarse mode SNA (2.5–10  $\mu\text{m}$ ) decreases. However, changes in size-resolved pH remain minimal, with the largest increase of 0.24 pH units occurring in the 0–1  $\mu\text{m}$  range. In regions with low SNA, such as South America and Greenland, increased  $\text{NH}_3$  emissions only lead to higher  $\text{NH}_3/\text{NH}_3$  concentrations due to the limited availability of  $\text{HNO}_3/\text{HNO}_3$  and  $\text{H}_2\text{SO}_4/\text{H}_2\text{SO}_4$  for SNA formation. Conversely, in  $\text{NH}_3/\text{NH}_3$ -rich regions such as East Asia and Europe, reductions in  $\text{NH}_3$  emissions trigger compensatory effects, as excess  $\text{NH}_3/\text{NH}_3$  and available  $\text{SO}_2/\text{SO}_2$  and  $\text{NO}_x/\text{NO}_x$  help to maintain SNA formation despite the reduced  $\text{NH}_3/\text{NH}_3$  supply. This highlights the buffering effect capacity of  $\text{NH}_3$  plays a crucial role in stabilizing aerosol acidity in densely populated areas, mitigating the effects of fluctuations in regions. However, it is important to note that this buffering effect may be partially offset by concurrent increases in precursor gases,  $\text{SO}_2^{2-}$  and  $\text{NO}_3^-$  concentrations.

In summary, this study underscores the critical influence of  $\text{NH}_3/\text{NH}_3$  emissions on global aerosol composition and acidity, with pronounced impacts on the size-resolved composition and properties of SNA aerosols.  $\text{NH}_3/\text{NH}_3$  emissions significantly modulate aerosol acidity, particularly in sub-micron ranges, highlighting the sensitivity of fine-mode aerosols to  $\text{NH}_3/\text{NH}_3$  levels. By buffering changes in aerosol pH,  $\text{NH}_3$  emissions contribute to maintaining a relatively stable aerosol acidity, especially in heavily densely populated and heavily industrialized regions. These findings emphasize the importance of accurately representing  $\text{NH}_3/\text{NH}_3$  dynamics in models for predicting atmospheric chemistry and climate interactions.

## Code and Data Availability

The usage of MESSy (Modular Earth Submodel System) and access to the source code is licensed to all affiliates of institutions which are members of the MESSy Consortium. Institutions can become a member of the MESSy Consortium by signing the “MESSy Memorandum of Understanding”. More information can be found on the MESSy Consortium website: <http://www.messy-interface.org> (last access: 31 January 2025). The code used in this study has been based on MESSy version 2.55 and is archived with a restricted access DOI (<https://doi.org/10.5281/zenodo.8379120>). The MESSy Consortium, (2023). The data produced in the study is available from the authors upon request.

## Acknowledgements

The work described in this paper has received funding from the Initiative and Networking Fund of the Helmholtz Association through the project “Advanced Earth System Modelling Capacity (ESM)”. The authors gratefully acknowledge the Earth System Modelling Project (ESM) for funding this work by providing computing time on the ESM partition of the supercomputer JUWELS (Alvarez, 2021) at the Jülich Supercomputing Centre (JSC). Xurong Wang has been supported by the China Scholarship Council (CSC).

## Competing Interests

At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.

## Author Contributions

XW and VAK planned the research and wrote the manuscript. XW performed the simulations and analyzed the results, assisted by VAK and APT. XW and APT collected the observational data and conducted the model evaluation. XW and ZL designed the Top-Dep case. All the authors discussed the results and contributed to the paper.

## Reference

- Aas, W., Mortier, A., Bowersox, V., Cherian, R., Faluvegi, G., Fagerli, H., Hand, J., Klimont, Z., Galy-Lacaux, C., Lehmann, C. M. B., Myhre, C. L., Myhre, G., Olivie, D., Sato, K., Quaas, J., Rao, P. S. P., Schulz, M., Shindell, D., Skeie, R. B., Stein, A., Takemura, T., Tsyro, S., Vet, R., and Xu, X.: Global and regional trends of atmospheric sulfur, *Scientific Reports*, 9, 953, 10.1038/s41598-018-37304-0, 2019.
- Alexander, L., Allen, S., Bindoff, N., Breon, F.-M., Church, J., Cubasch, U., Emori, S., Forster, P., Friedlingstein, P., Gillett, N., Gregory, J., Hartmann, D., Jansen, E., Kirtman, B., Knutti, R., Kanikicharla, K., Lemke, P., Marotzke, J., Masson-Delmotte, V., and Xie, S.-P.: Climate change 2013: The physical science basis, in contribution of Working Group I (WGI) to the Fifth Assessment Report (AR5) of the Intergovernmental Panel on Climate Change (IPCC), in, 2013.
- Allen, H. M., Draper, D. C., Ayres, B. R., Ault, A., Bondy, A., Takahama, S., Modini, R. L., Baumann, K., Edgerton, E., Knote, C., Laskin, A., Wang, B., and Fry, J. L.: Influence of crustal dust and sea spray supermicron particle concentrations and acidity on inorganic  $\text{NO}_3^-$  aerosol during the 2013 Southern Oxidant and Aerosol Study, *Atmos. Chem. Phys.*, 15, 10669–10685, 10.5194/acp-15-10669-2015, 2015.
- Andreae, M., Berresheim, H., Bingemer, H., Jacob, D. J., Lewis, B., Li, S. M., and Talbot, R. W.: The atmospheric sulfur cycle over the Amazon Basin: 2. Wet season, *Journal of Geophysical Research: Atmospheres*, 95, 16813–16824, 1990.
- Andreae, M. O.: Aerosols Before Pollution, *Science*, 315, 50–51, doi:10.1126/science.1136529, 2007.
- Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmos. Chem. Phys.*, 19, 8523–8546, 10.5194/acp-19-8523-2019, 2019.
- Asthitha, M., Lelieveld, J., Abdel Kader, M., Pozzer, A., and de Meij, A.: Parameterization of dust emissions in the

Formatted: Indent: Hanging: 0.51 cm

- global atmospheric chemistry-climate model EMAC: impact of nudging and soil properties, *Atmos. Chem. Phys.*, 12, 11057–11083, 10.5194/acp-12-11057-2012, 2012.
- 905 Bacer, S., Sullivan, S. C., Karydis, V. A., Barahona, D., Krämer, M., Nenes, A., Tost, H., Tsimpidi, A. P., Lelieveld, J., and Pozzer, A.: Implementation of a comprehensive ice crystal formation parameterization for cirrus and mixed-phase clouds in the EMAC model (based on MESSy 2.53), *Geosci. Model Dev.*, 11, 4021–4041, 10.5194/gmd-11-4021-2018, 2018.
- Beusen, A. H. W., Bouwman, A. F., Heuberger, P. S. C., Van Drecht, G., and Van Der Hoek, K. W.: Bottom-up uncertainty estimates of global ammonia emissions from global agricultural production systems, *Atmospheric Environment*, 42, 6067–6077, <https://doi.org/10.1016/j.atmosenv.2008.03.044>, 2008.
- 910 Bian, H., Chin, M., Hauglustaine, D. A., Schulz, M., Myhre, G., Bauer, S. E., Lund, M. T., Karydis, V. A., Kucsera, T. L., Pan, X., Pozzer, A., Skeie, R. B., Steenrod, S. D., Sudo, K., Tsigaridis, K., Tsimpidi, A. P., and Tsyro, S. G.: Investigation of global particulate nitrate from the AeroCom phase III experiment, *Atmos. Chem. Phys.*, 17, 12911–12940, 10.5194/acp-17-12911-2017, 2017.
- 915 Bleeker, A., Vries, W. d., Dentener, F., Sutton, M. A., Reis, S., Riddick, S. N., Dragosits, U., Nemitz, E., Theobald, M. R., Tang, Y. S., Braban, C. F., Veno, M., Dore, A. J., Mitchell, R. F., Wanless, S., Daunt, F., Fowler, D., Blackall, T. D., Milford, C., Flechard, C. R., Loubet, B., Massad, R., Cellier, P., Coheur, P. F., Clarisse, L., Damme, M. v., Ngadi, Y., Clerbaux, C., Ambelas Skjoth, C., Geels, C., Hertel, O., Wichink Kruit, R. J., Pinder, R. W., Bash, J. O., Walker, J. D., Simpson, D., Horvath, L., and Misselbrook, T. H.: Towards a climate-dependent paradigm of ammonia emission and deposition, urn:NBN:nl:ui:24-uuid:1516f098-ca7e-402b-a05d-83633f8c2612, 2013.
- 920 Bougiatioti, A., Nikolaou, P., Stavroulas, I., Kouvarakis, G., Weber, R., Nenes, A., Kanakidou, M., and Mihalopoulos, N.: Particle water and pH in the eastern Mediterranean: source variability and implications for nutrient availability, *Atmos. Chem. Phys.*, 16, 4579–4591, 10.5194/acp-16-4579-2016, 2016.
- 925 Bouwman, A. F., Boumans, L. J. M., and Batjes, N. H.: Estimation of global NH<sub>3</sub> volatilization loss from synthetic fertilizers and animal manure applied to arable lands and grasslands, *Global Biogeochemical Cycles*, 16, 8-1-8-14, <https://doi.org/10.1029/2000GB001389>, 2002.
- Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., Van Der Hoek, K. W., and Olivier, J. G. J.: A global high-resolution emission inventory for ammonia, *Global Biogeochemical Cycles*, 11, 561–587, <https://doi.org/10.1029/97GB02266>, 1997.
- 930 Burgard, D. A. and Bria, C. R. M.: Bridge-based sensing of NO<sub>x</sub> and SO<sub>2</sub> emissions from ocean-going ships, *Atmospheric Environment*, 136, 54–60, <https://doi.org/10.1016/j.atmosenv.2016.04.014>, 2016.
- Cai, S., Wang, Y., Zhao, B., Wang, S., Chang, X., and Hao, J.: The impact of the “Air Pollution Prevention and Control Action Plan” on PM<sub>2.5</sub> concentrations in Jing-Jin-Ji region during 2012–2020, *Science of The Total Environment*, 580, 197–209, <https://doi.org/10.1016/j.scitotenv.2016.11.188>, 2017.
- 935 Cavalli, F., Facchini, M. C., Decesari, S., Mircea, M., Emblico, L., Fuzzi, S., Ceburnis, D., Yoon, Y. J., O'Dowd, C. D., Putaud, J.-P., and Dell'Acqua, A.: Advances in characterization of size-resolved organic matter in marine aerosol over the North Atlantic, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2004JD005137>, 2004.
- 940 Chang, Y., Zou, Z., Zhang, Y., Deng, C., Hu, J., Shi, Z., Dore, A. J., and Collett, J. L., Jr.: Assessing Contributions of Agricultural and Nonagricultural Emissions to Atmospheric Ammonia in a Chinese Megacity, *Environmental Science & Technology*, 53, 1822–1833, 10.1021/acs.est.8b05984, 2019.
- Che, H., Zhang, X., Li, Y., Zhou, Z., Qu, J. J., and Hao, X.: Haze trends over the capital cities of 31 provinces in China, 1981–2005, *Theoretical and Applied Climatology*, 97, 235–242, 10.1007/s00704-008-0059-8, 2009.
- 945 Chen, D., Liu, Z., Fast, J., and Ban, J.: Simulations of sulfate–nitrate–ammonium (SNA) aerosols during the extreme haze events over northern China in October 2014, *Atmos. Chem. Phys.*, 16, 10707–10724, 10.5194/acp-16-10707-2016, 2016.
- Chen, Y., Shen, H., and Russell, A. G.: Current and Future Responses of Aerosol pH and Composition in the U.S. to Declining SO<sub>2</sub> Emissions and Increasing NH<sub>3</sub> Emissions, *Environmental Science & Technology*, 53, 9646–9655, 10.1021/acs.est.9b02005, 2019.
- 950 Chen, Y., Cheng, Y., Ma, N., Wei, C., Ran, L., Wolke, R., Größ, J., Wang, Q., Pozzer, A., Denier van der Gon, H. A. C., Spindler, G., Lelieveld, J., Tegen, I., Su, H., and Wiedensohler, A.: Natural sea-salt emissions moderate the climate forcing of anthropogenic nitrate, *Atmos. Chem. Phys.*, 20, 771–786, 10.5194/acp-20-771-2020, 2020.
- Chen, Y., Shen, H., Kaiser, J., Hu, Y., Capps, S. L., Zhao, S., Hakami, A., Shih, J. S., Pavur, G. K., Turner, M. D., Henze, D. K., Resler, J., Nenes, A., Napelenok, S. L., Bash, J. O., Fahey, K. M., Carmichael, G. R., Chai, T., Clarisse, L., Coheur, P. F., Van Damme, M., and Russell, A. G.: High-resolution hybrid inversion of IASI ammonia columns to constrain US ammonia emissions using the CMAQ adjoint model, *Atmos. Chem. Phys.*, 21, 2067–2082, 10.5194/acp-21-2067-2021, 2021.

Cheng, Y., Zheng, G., Wei, C., Mu, Q., Zheng, B., Wang, Z., Gao, M., Zhang, Q., He, K., Carmichael, G., Pöschl, U., and Su, H.: Reactive nitrogen chemistry in aerosol water as a source of sulfate during haze events in China, *Science Advances*, 2, e1601530, doi:10.1126/sciadv.1601530, 2016.

Clarisse, L., Clerbaux, C., Dentener, F., Hurtmans, D., and Coheur, P.-F.: Global ammonia distribution derived from infrared satellite observations, *Nature Geoscience*, 2, 479-483, 10.1038/ngeo551, 2009.

Colette, A., Granier, C., Hodnebrog, Ø., Jakobs, H., Maurizi, A., Nyiri, A., Bessagnet, B., D'Angiola, A., D'Isidoro, M., Gauss, M., Meleux, F., Memmesheimer, M., Mieville, A., Rouil, L., Russo, F., Solberg, S., Stordal, F., and Tampieri, F.: Air quality trends in Europe over the past decade: a first multi-model assessment, *Atmos. Chem. Phys.*, 11, 11657-11678, 10.5194/acp-11-11657-2011, 2011.

Craig, R. L., Peterson, P. K., Nandy, L., Lei, Z., Hossain, M. A., Camarena, S., Dodson, R. A., Cook, R. D., Dutcher, C. S., and Ault, A. P.: Direct Determination of Aerosol pH: Size-Resolved Measurements of Submicrometer and Supermicrometer Aqueous Particles, *Analytical Chemistry*, 90, 11232-11239, 10.1021/acs.analchem.8b00586, 2018.

Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., van Aardenne, J. A., Monni, S., Doering, U., Olivier, J. G. J., Pagliari, V., and Janssens-Maenhout, G.: Gridded emissions of air pollutants for the period 1970–2012 within EDGAR v4.3.2, *Earth Syst. Sci. Data*, 10, 1987-2013, 10.5194/essd-10-1987-2018, 2018.

Dammers, E., McLinden, C. A., Griffin, D., Shephard, M. W., Van Der Graaf, S., Lutsch, E., Schaap, M., Gainairu-Matz, Y., Fioletov, V., Van Damme, M., Whitburn, S., Clarisse, L., Cady-Pereira, K., Clerbaux, C., Coheur, P. F., and Erisman, J. W.: NH<sub>3</sub> emissions from large point sources derived from CrIS and IASI satellite observations, *Atmos. Chem. Phys.*, 19, 12261-12293, 10.5194/acp-19-12261-2019, 2019.

Dammers, E., Shephard, M. W., Palm, M., Cady-Pereira, K., Capps, S., Lutsch, E., Strong, K., Hannigan, J. W., Ortega, I., Toon, G. C., Stremme, W., Grutter, M., Jones, N., Smale, D., Siemons, J., Hrpcek, K., Tremblay, D., Schaap, M., Notholt, J., and Erisman, J. W.: Validation of the CrIS fast physical NH<sub>3</sub> retrieval with ground-based FTIR, *Atmos. Meas. Tech.*, 10, 2645-2667, 10.5194/amt-10-2645-2017, 2017.

Delon, C., Galy-Lacaux, C., Adon, M., Lioussé, C., Serça, D., Diop, B., and Akpo, A.: Nitrogen compounds emission and deposition in West African ecosystems: comparison between wet and dry savanna, *Biogeosciences*, 9, 385-402, 10.5194/bg-9-385-2012, 2012.

Dentener, F. J. and Crutzen, P. J.: A three-dimensional model of the global ammonia cycle, *Journal of Atmospheric Chemistry*, 19, 331-369, 10.1007/BF00694492, 1994.

Ding, J., Zhao, P., Su, J., Dong, Q., Du, X., and Zhang, Y.: Aerosol pH and its driving factors in Beijing, *Atmos. Chem. Phys.*, 19, 7939-7954, 10.5194/acp-19-7939-2019, 2019.

Dockery, D. W., Pope, C. A., Xu, X., Spengler, J. D., Ware, J. H., Fay, M. E., Ferris, B. G., and Speizer, F. E.: An Association between Air Pollution and Mortality in Six U.S. Cities, *New England Journal of Medicine*, 329, 1753-1759, 10.1056/nejm199312093292401, 1993.

Dockery, D. W., Cunningham, J., Damokosh, A. I., Neas, L. M., Spengler, J. D., Koutrakis, P., Ware, J. H., Raizenne, M., and Speizer, F. E.: Health effects of acid aerosols on North American children: respiratory symptoms, *Environmental Health Perspectives*, 104, 500-505, doi:10.1289/ehp.96104500, 1996.

Erisman, J. W. and Schaap, M.: The need for ammonia abatement with respect to secondary PM reductions in Europe, *Environ Pollut*, 129, 159-163, 10.1016/j.envpol.2003.08.042, 2004.

Fang, T., Guo, H., Zeng, L., Verma, V., Nenes, A., and Weber, R. J.: Highly Acidic Ambient Particles, Soluble Metals, and Oxidative Potential: A Link between Sulfate and Aerosol Toxicity, *Environmental Science & Technology*, 51, 2611-2620, 10.1021/acs.est.6b06151, 2017.

Fiddes, S. L., Woodhouse, M. T., Nicholls, Z., Lane, T. P., and Schofield, R.: Cloud, precipitation and radiation responses to large perturbations in global dimethyl sulfide, *Atmos. Chem. Phys.*, 18, 10177-10198, 10.5194/acp-18-10177-2018, 2018.

Fortems-Cheiney, A., Dufour, G., Foret, G., Siour, G., Van Damme, M., Coheur, P.-F., Clarisse, L., Clerbaux, C., and Beekmann, M.: Understanding the Simulated Ammonia Increasing Trend from 2008 to 2015 over Europe with CHIMERE and Comparison with IASI Observations, *Atmosphere*, 13, 1101, 2022.

Fountoukis, C. and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for  $K^{+}$ – $Ca^{2+}$ – $Mg^{2+}$ – $NH_4^{+}$ – $Na^{+}$ – $SO_4^{2-}$ – $SO_3^{2-}$ – $NO_3^{-}$ – $Cl^{-}$ – $H_2O$  aerosols, *Atmos. Chem. Phys.*, 7, 4639-4659, 10.5194/acp-7-4639-2007, 2007.

Fu, X., Wang, S., Xing, J., Zhang, X., Wang, T., and Hao, J.: Increasing Ammonia Concentrations Reduce the Effectiveness of Particle Pollution Control Achieved via SO<sub>2</sub> and NO<sub>x</sub> Emissions Reduction in East China, *Environmental Science & Technology Letters*, 4, 221-227, 10.1021/acs.estlett.7b00143, 2017.

- 1015 Ge, Y., Vieno, M., Stevenson, D. S., Wind, P., and Heal, M. R.: A new assessment of global and regional budgets, fluxes, and lifetimes of atmospheric reactive N and S gases and aerosols, *Atmos. Chem. Phys.*, 22, 8343-8368, 10.5194/acp-22-8343-2022, 2022.
- Gong, C., Tian, H., Liao, H., Pan, N., Pan, S., Ito, A., Jain, A. K., Kou-Giesbrecht, S., Joos, F., Sun, Q., Shi, H., Vuichard, N., Zhu, Q., Peng, C., Maggi, F., Tang, F. H. M., and Zechle, S.: Global net climate effects of anthropogenic reactive nitrogen, *Nature*, 632, 557-563, 10.1038/s41586-024-07714-4, 2024.
- 1020 Gu, B., Zhang, L., Van Dingenen, R., Vieno, M., Van Grinsven, H. J., Zhang, X., Zhang, S., Chen, Y., Wang, S., Ren, C., Rao, S., Holland, M., Winiwarter, W., Chen, D., Xu, J., and Sutton, M. A.: Abating ammonia is more cost-effective than nitrogen oxides for mitigating PM<sub>2.5</sub> air pollution, *Science*, 374, 758-762, doi:10.1126/science.abf8623, 2021.
- 1025 Guo, H., Weber, R. J., and Nenes, A.: High levels of ammonia do not raise fine particle pH sufficiently to yield nitrogen oxide-dominated sulfate production, *Scientific Reports*, 7, 12109, 10.1038/s41598-017-11704-0, 2017a.
- Guo, H., Otjes, R., Schlag, P., Kiendler-Scharr, A., Nenes, A., and Weber, R. J.: Effectiveness of ammonia reduction on control of fine particle nitrate, *Atmos. Chem. Phys.*, 18, 12241-12256, 10.5194/acp-18-12241-2018, 2018.
- Guo, H., Liu, J., Froyd, K. D., Roberts, J. M., Veres, P. R., Hayes, P. L., Jimenez, J. L., Nenes, A., and Weber, R. J.: Fine particle pH and gas-particle phase partitioning of inorganic species in Pasadena, California, during the 2010 CalNex campaign, *Atmos. Chem. Phys.*, 17, 5703-5719, 10.5194/acp-17-5703-2017, 2017b.
- 1030 Guo, H., Sullivan, A. P., Campuzano-Jost, P., Schroder, J. C., Lopez-Hilfiker, F. D., Dibb, J. E., Jimenez, J. L., Thornton, J. A., Brown, S. S., Nenes, A., and Weber, R. J.: Fine particle pH and the partitioning of nitric acid during winter in the northeastern United States, *Journal of Geophysical Research: Atmospheres*, 121, 10,355-310,376, <https://doi.org/10.1002/2016JD025311>, 2016.
- 1035 Guo, H., Xu, L., Bougiatioti, A., Cerully, K. M., Capps, S. L., Hite Jr, J. R., Carlton, A. G., Lee, S. H., Bergin, M. H., Ng, N. L., Nenes, A., and Weber, R. J.: Fine-particle water and pH in the southeastern United States, *Atmos. Chem. Phys.*, 15, 5211-5228, 10.5194/acp-15-5211-2015, 2015.
- Hand, J. L., Schichtel, B. A., Malm, W. C., and Pitchford, M. L.: Particulate sulfate ion concentration and SO<sub>2</sub> emission trends in the United States from the early 1990s through 2010, *Atmos. Chem. Phys.*, 12, 10353-10365, 10.5194/acp-12-10353-2012, 2012.
- 1040 Hauglustaine, D. A., Balkanski, Y., and Schulz, M.: A global model simulation of present and future nitrate aerosols and their direct radiative forcing of climate, *Atmos. Chem. Phys.*, 14, 11031-11063, 10.5194/acp-14-11031-2014, 2014.
- 1045 Heald, C. L., Collett Jr, J. L., Lee, T., Benedict, K. B., Schwandner, F. M., Li, Y., Clarisse, L., Hurtmans, D. R., Van Damme, M., Clerbaux, C., Coheur, P. F., Philip, S., Martin, R. V., and Pye, H. O. T.: Atmospheric ammonia and particulate inorganic nitrogen over the United States, *Atmos. Chem. Phys.*, 12, 10295-10312, 10.5194/acp-12-10295-2012, 2012.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999-2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 1055 Hickman, J. E., Andela, N., Tsigaridis, K., Galy-Lacaux, C., Ossohou, M., Dammers, E., Van Damme, M., Clarisse, L., and Bauer, S. E.: Continental and Ecoregion-Specific Drivers of Atmospheric NO<sub>2</sub> and NH<sub>3</sub> Seasonality Over Africa Revealed by Satellite Observations, *Global Biogeochemical Cycles*, 35, e2020GB006916, <https://doi.org/10.1029/2020GB006916>, 2021.
- 1060 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J. I., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geosci. Model Dev.*, 11, 369-408, 10.5194/gmd-11-369-2018, 2018.
- 1065 Huang, X., Zhang, J., Zhang, W., Tang, G., and Wang, Y.: Atmospheric ammonia and its effect on PM<sub>2.5</sub> pollution in urban Chengdu, Sichuan Basin, China, *Environmental Pollution*, 291, 118195, <https://doi.org/10.1016/j.envpol.2021.118195>, 2021.
- Intergovernmental Panel on Climate, C. (Ed.): *Climate Change 2022 - Mitigation of Climate Change: Working Group III Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, DOI: 10.1017/9781009157926, 2023.
- 1070

- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge, United Kingdom and New York, NY, USA, 1535, 2013.
- 1075 [IPCC, 2022: Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change \[P.R. Shukla, J. Skea, R. Slade, A. Al Khouridajie, R. van Diemen, D. McCollum, M. Pathak, S. Some, P. Vyas, R. Fradera, M. Belkacemi, A. Hasija, G. Lisboa, S. Luz, J. Malley. \(eds.\)\]. Cambridge University Press, Cambridge, UK and New York, NY, USA. doi: 10.1017/9781009157926](#)
- 1080 Jackson, R. L., Gabric, A. J., Cropp, R., and Woodhouse, M. T.: Dimethylsulfide (DMS), marine biogenic aerosols and the ecophysiology of coral reefs, *Biogeosciences*, 17, 2181-2204, 10.5194/bg-17-2181-2020, 2020.
- Jo, Y.-J., Lee, H.-J., Jo, H.-Y., Woo, J.-H., Kim, Y., Lee, T., Heo, G., Park, S.-M., Jung, D., Park, J., and Kim, C.-H.: Changes in inorganic aerosol compositions over the Yellow Sea area from impact of Chinese emissions mitigation, *Atmospheric Research*, 240, 104948, <https://doi.org/10.1016/j.atmosres.2020.104948>, 2020.
- 1085 Jöckel, P., Sander, R., Kerkweg, A., Tost, H., and Lelieveld, J.: Technical Note: The Modular Earth Submodel System (MESSy) - a new approach towards Earth System Modeling, *Atmos. Chem. Phys.*, 5, 433-444, 10.5194/acp-5-433-2005, 2005.
- Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geosci. Model Dev.*, 3, 717-752, 10.5194/gmd-3-717-2010, 2010.
- 1090 Jöckel, P., Tost, H., Pozzer, A., Kunze, M., Kirner, O., Brenninkmeijer, C., Brinkop, S., Cai, D. S., Dyroff, C., and Eckstein, J.: Earth System Chemistry Integrated Modelling (ESCiMo) with the Modular Earth Submodel System (MESSy, version 2.51), *Geoscientific Model Development*, 9, 1153-1200, 2016.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527-554, 10.5194/bg-9-527-2012, 2012.
- 1095 Kakavas, S., Patoulas, D., Zakoura, M., Nenes, A., and Pandis, S. N.: Size-resolved aerosol pH over Europe during summer, *Atmos. Chem. Phys.*, 21, 799-811, 10.5194/acp-21-799-2021, 2021.
- Karydis, V. A., Tsimpidi, A. P., Pozzer, A., and Lelieveld, J.: How alkaline compounds control atmospheric aerosol particle acidity, *Atmos. Chem. Phys.*, 21, 14983-15001, 10.5194/acp-21-14983-2021, 2021.
- 1100 Karydis, V. A., Tsimpidi, A. P., Lei, W., Molina, L. T., and Pandis, S. N.: Formation of semivolatile inorganic aerosols in the Mexico City Metropolitan Area during the MILAGRO campaign, *Atmos. Chem. Phys.*, 11, 13305-13323, 10.5194/acp-11-13305-2011, 2011.
- Karydis, V. A., Tsimpidi, A. P., Pozzer, A., Astitha, M., and Lelieveld, J.: Effects of mineral dust on global atmospheric nitrate concentrations, *Atmos. Chem. Phys.*, 16, 1491-1509, 10.5194/acp-16-1491-2016, 2016.
- 1105 Karydis, V. A., Tsimpidi, A. P., Bacer, S., Pozzer, A., Nenes, A., and Lelieveld, J.: Global impact of mineral dust on cloud droplet number concentration, *Atmos. Chem. Phys.*, 17, 5601-5621, 10.5194/acp-17-5601-2017, 2017.
- Karydis, V. A., Tsimpidi, A. P., Fountoukis, C., Nenes, A., Zavala, M., Lei, W., Molina, L. T., and Pandis, S. N.: Simulating the fine and coarse inorganic particulate matter concentrations in a polluted megacity, *Atmospheric Environment*, 44, 608-620, <https://doi.org/10.1016/j.atmosenv.2009.11.023>, 2010.
- 1110 Kerkweg, A., Buchholz, J., Ganzeveld, L., Pozzer, A., Tost, H., and Jöckel, P.: Technical Note: An implementation of the dry removal processes DRY DEposition and SEDimentation in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 4617-4632, 10.5194/acp-6-4617-2006, 2006.
- Khan, M. A. H., Lowe, D., Derwent, R. G., Foulds, A., Chhantyal-Pun, R., McFiggans, G., Orr-Ewing, A. J., Percival, C. J., and Shallcross, D. E.: Global and regional model simulations of atmospheric ammonia, *Atmospheric Research*, 234, 104702, <https://doi.org/10.1016/j.atmosres.2019.104702>, 2020.
- 1115 Klingmüller, K., Metzger, S., Abdelkader, M., Karydis, V. A., Stenchikov, G. L., Pozzer, A., and Lelieveld, J.: Revised mineral dust emissions in the atmospheric chemistry-climate model EMAC (MESSy 2.52 DU\_Astithal KKDU2017 patch), *Geosci. Model Dev.*, 11, 989-1008, 10.5194/gmd-11-989-2018, 2018.
- Koçak, M., Mihalopoulos, N., and Kubilay, N.: Chemical composition of the fine and coarse fraction of aerosols in the northeastern Mediterranean, *Atmospheric Environment*, 41, 7351-7368, <https://doi.org/10.1016/j.atmosenv.2007.05.011>, 2007.
- 1120 Kong, L., Tang, X., Zhu, J., Wang, Z., Pan, Y., Wu, H., Wu, L., Wu, Q., He, Y., Tian, S., Xie, Y., Liu, Z., Sui, W., Han, L., and Carmichael, G.: Improved Inversion of Monthly Ammonia Emissions in China Based on the Chinese Ammonia Monitoring Network and Ensemble Kalman Filter, *Environmental Science & Technology*, 53, 12529-12538, 10.1021/acs.est.9b02701, 2019.
- 1125 Lei, L., Zhou, W., Chen, C., He, Y., Li, Z., Sun, J., Tang, X., Fu, P., Wang, Z., and Sun, Y.: Long-term characterization

of aerosol chemistry in cold season from 2013 to 2020 in Beijing, China, *Environmental Pollution*, 268, 115952, <https://doi.org/10.1016/j.envpol.2020.115952>, 2021.

1130 Li, C., Martin, R. V., van Donkelaar, A., Boys, B. L., Hammer, M. S., Xu, J.-W., Marais, E. A., Reff, A., Strum, M., Ridley, D. A., Crippa, M., Brauer, M., and Zhang, Q.: Trends in Chemical Composition of Global and Regional Population-Weighted Fine Particulate Matter Estimated for 25 Years, *Environmental Science & Technology*, 51, 11185-11195, [10.1021/acs.est.7b02530](https://doi.org/10.1021/acs.est.7b02530), 2017.

1135 Li, K., Chen, L., White, S. J., Yu, H., Wu, X., Gao, X., Azzi, M., and Cen, K.: Smog chamber study of the role of NH<sub>3</sub> in new particle formation from photo-oxidation of aromatic hydrocarbons, *Science of The Total Environment*, 619-620, 927-937, <https://doi.org/10.1016/j.scitotenv.2017.11.180>, 2018.

Liu, J., Ding, P., Zong, Z., Li, J., Tian, C., Chen, W., Chang, M., Salazar, G., Shen, C., Cheng, Z., Chen, Y., Wang, X., Szidat, S., and Zhang, G.: Evidence of Rural and Suburban Sources of Urban Haze Formation in China: A Case Study From the Pearl River Delta Region, *Journal of Geophysical Research: Atmospheres*, 123, 4712-4726, <https://doi.org/10.1029/2017JD027952>, 2018a.

1140 Liu, L., Xu, W., Lu, X., Zhong, B., Guo, Y., Lu, X., Zhao, Y., He, W., Wang, S., Zhang, X., Liu, X., and Vitousek, P.: Exploring global changes in agricultural ammonia emissions and their contribution to nitrogen deposition since 1980, *Proceedings of the National Academy of Sciences*, 119, e2121998119, [doi:10.1073/pnas.2121998119](https://doi.org/10.1073/pnas.2121998119), 2022.

Liu, M., Song, Y., Zhou, T., Xu, Z., Yan, C., Zheng, M., Wu, Z., Hu, M., Wu, Y., and Zhu, T.: Fine particle pH during severe haze episodes in northern China, *Geophysical Research Letters*, 44, 5213-5221, <https://doi.org/10.1002/2017GL073210>, 2017.

1145 Liu, M., Huang, X., Song, Y., Xu, T., Wang, S., Wu, Z., Hu, M., Zhang, L., Zhang, Q., Pan, Y., Liu, X., and Zhu, T.: Rapid SO<sub>2</sub> emission reductions significantly increase tropospheric ammonia concentrations over the North China Plain, *Atmos. Chem. Phys.*, 18, 17933-17943, [10.5194/acp-18-17933-2018](https://doi.org/10.5194/acp-18-17933-2018), 2018b.

1150 Lohmann, U. and Ferrachat, S.: Impact of parametric uncertainties on the present-day climate and on the anthropogenic aerosol effect, *Atmos. Chem. Phys.*, 10, 11373-11383, [10.5194/acp-10-11373-2010](https://doi.org/10.5194/acp-10-11373-2010), 2010.

Luo, Z., Zhang, Y., Chen, W., Van Damme, M., Coheur, P. F., and Clarisse, L.: Estimating global ammonia (NH<sub>3</sub>) emissions based on IASI observations from 2008 to 2018, *Atmos. Chem. Phys.*, 22, 10375-10388, [10.5194/acp-22-10375-2022](https://doi.org/10.5194/acp-22-10375-2022), 2022.

1155 McDuffie, E. E., Smith, S. J., O'Rourke, P., Tibrewal, K., Venkataraman, C., Marais, E. A., Zheng, B., Crippa, M., Brauer, M., and Martin, R. V.: A global anthropogenic emission inventory of atmospheric pollutants from sector- and fuel-specific sources (1970–2017): an application of the Community Emissions Data System (CEDS), *Earth Syst. Sci. Data*, 12, 3413-3442, [10.5194/essd-12-3413-2020](https://doi.org/10.5194/essd-12-3413-2020), 2020.

Milousis, A., Tsimpidi, A. P., Tost, H., Pandis, S. N., Nenes, A., Kiendler-Scharr, A., and Karydis, V. A.: Implementation of the ISORROPIA-lite aerosol thermodynamics model into the EMAC chemistry climate model (based on MESSy v2.55): implications for aerosol composition and acidity, *Geosci. Model Dev.*, 17, 1111-1131, [10.5194/gmd-17-1111-2024](https://doi.org/10.5194/gmd-17-1111-2024), 2024.

1160 [Milousis, A., Scholz, S. M. C., Fuchs, H., Tsimpidi, A. P., and Karydis, V. A.: Global Perspectives on Nitrate Aerosol Dynamics: A Comprehensive Sensitivity Analysis, \*EGU sphere\*, 2025, 1-42, \[10.5194/egusphere-2025-313\]\(https://doi.org/10.5194/egusphere-2025-313\), 2025.](#)

1165 Morán, M., Ferreira, J., Martins, H., Monteiro, A., Borrego, C., and González, J. A.: Ammonia agriculture emissions: From EMEP to a high resolution inventory, *Atmospheric Pollution Research*, 7, 786-798, <https://doi.org/10.1016/j.apr.2016.04.001>, 2016.

1170 [Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J. F., Lin, G., Liu, X., Lund, M. T., Luo, G., Ma, X., van Noije, T., Penner, J. E., Rasch, P. J., Ruiz, A., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J. H., Zhang, K., Zhang, H., and Zhou, C.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations, \*Atmos. Chem. Phys.\*, 13, 1853-1877, \[10.5194/acp-13-1853-2013\]\(https://doi.org/10.5194/acp-13-1853-2013\), 2013.](#)

Nenes, A., Pandis, S. N., and Pilinis, C.: ISORROPIA: A New Thermodynamic Equilibrium Model for Multiphase Multicomponent Inorganic Aerosols, *Aquatic Geochemistry*, 4, 123-152, [10.1023/A:1009604003981](https://doi.org/10.1023/A:1009604003981), 1998.

1175 Nenes, A., Pandis, S. N., Weber, R. J., and Russell, A.: Aerosol pH and liquid water content determine when particulate matter is sensitive to ammonia and nitrate availability, *Atmos. Chem. Phys.*, 20, 3249-3258, [10.5194/acp-20-3249-2020](https://doi.org/10.5194/acp-20-3249-2020), 2020.

Olivier, J. G. J., Bouwman, A. F., Van der Hoek, K. W., and Berdowski, J. J. M.: Global air emission inventories for anthropogenic sources of NO<sub>x</sub>, NH<sub>3</sub> and N<sub>2</sub>O in 1990, *Environmental Pollution*, 102, 135-148, [https://doi.org/10.1016/S0269-7491\(98\)80026-2](https://doi.org/10.1016/S0269-7491(98)80026-2), 1998.

1180 Osipov, S., Chowdhury, S., Crowley, J. N., Tadic, I., Drewnick, F., Borrmann, S., Eger, P., Fachinger, F., Fischer, H., Predybaylo, E., Fnais, M., Harder, H., Pikridas, M., Vouterakos, P., Pozzer, A., Sciare, J., Ukhov, A., Stenchikov,

- G. L., Williams, J., and Lelieveld, J.: Severe atmospheric pollution in the Middle East is attributable to anthropogenic sources, *Communications Earth & Environment*, 3, 203, 10.1038/s43247-022-00514-6, 2022.
- 1185 Pan, Y., Tian, S., Liu, D., Fang, Y., Zhu, X., Zhang, Q., Zheng, B., Michalski, G., and Wang, Y.: Fossil Fuel Combustion-Related Emissions Dominate Atmospheric Ammonia Sources during Severe Haze Episodes: Evidence from (15)N-Stable Isotope in Size-Resolved Aerosol Ammonium, *Environ Sci Technol*, 50, 8049-8056, 10.1021/acs.est.6b00634, 2016.
- 1190 Pozzer, A., Reifenberg, S. F., Kumar, V., Franco, B., Kohl, M., Taraborrelli, D., Gromov, S., Ehrhart, S., Jöckel, P., Sander, R., Fall, V., Rosanka, S., Karydis, V., Akritidis, D., Emmerichs, T., Crippa, M., Guizzardi, D., Kaiser, J. W., Clarisse, L., Kiendler-Scharr, A., Tost, H., and Tsimpidi, A.: Simulation of organics in the atmosphere: evaluation of EMACv2.54 with the Mainz Organic Mechanism (MOM) coupled to the ORACLE (v1.0) submodel, *Geosci. Model Dev.*, 15, 2673-2710, 10.5194/gmd-15-2673-2022, 2022.
- 1195 Pozzer, A., Tsimpidi, A. P., Karydis, V. A., de Meij, A., and Lelieveld, J.: Impact of agricultural emission reductions on fine-particulate matter and public health, *Atmos. Chem. Phys.*, 17, 12813-12826, 10.5194/acp-17-12813-2017, 2017.
- Pringle, K. J., Tost, H., Message, S., Steil, B., Giannadaki, D., Nenes, A., Fountoukis, C., Stier, P., Vignati, E., and Lelieveld, J.: Description and evaluation of GMX: a new aerosol submodel for global simulations (v1), *Geosci. Model Dev.*, 3, 391-412, 10.5194/gmd-3-391-2010, 2010.
- 1200 Prospero, J. M., Savoie, D. L., Nees, R. T., Duce, R. A., and Merrill, J.: Particulate sulfate and nitrate in the boundary layer over the North Pacific Ocean, *Journal of Geophysical Research: Atmospheres*, 90, 10586-10596, <https://doi.org/10.1029/JD090iD06p10586>, 1985.
- Pye, H. O. T., Nenes, A., Alexander, B., Ault, A. P., Barth, M. C., Clegg, S. L., Collett Jr, J. L., Fahey, K. M., Hennigan, C. J., Herrmann, H., Kanakidou, M., Kelly, J. T., Ku, I. T., McNeill, V. F., Riemer, N., Schaefer, T., Shi, G., Tilgner, A., Walker, J. T., Wang, T., Weber, R., Xing, J., Zaveri, R. A., and Zuend, A.: The acidity of atmospheric particles and clouds, *Atmos. Chem. Phys.*, 20, 4809-4888, 10.5194/acp-20-4809-2020, 2020.
- 1205 Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornbluh, L., Manzini, E., Schlese, U., and Schulzweida, U.: Sensitivity of simulated climate to horizontal and vertical resolution in the ECHAM5 atmosphere model, *Journal of Climate*, 19, 3771-3791, 10.1175/jcli3824.1, 2006.
- 1210 Russell, A. R., Valin, L. C., and Cohen, R. C.: Trends in OMI NO<sub>2</sub> observations over the United States: effects of emission control technology and the economic recession, *Atmos. Chem. Phys.*, 12, 12197-12209, 10.5194/acp-12-12197-2012, 2012.
- Russell, L. M., Moore, R. H., Burrows, S. M., and Quinn, P. K.: Ocean flux of salt, sulfate, and organic components to atmospheric aerosol, *Earth-Science Reviews*, 239, 104364, <https://doi.org/10.1016/j.earscirev.2023.104364>, 2023.
- 1215 Sahoo, P., Sahu, S. K., Mangaraj, P., Mishra, A., Beig, G., and Gunthe, S. S.: Reporting of gridded ammonia emission and assessment of hotspots across India: A comprehensive study of 24 anthropogenic sources, *Journal of Hazardous Materials*, 479, 135557, <https://doi.org/10.1016/j.jhazmat.2024.135557>, 2024.
- 1220 Sander, R., Baumgaertner, A., Cabrera-Perez, D., Frank, F., Gromov, S., Groß, J. U., Harder, H., Huijnen, V., Jöckel, P., Karydis, V. A., Niemeyer, K. E., Pozzer, A., Riede, H., Schultz, M. G., Taraborrelli, D., and Tauer, S.: The community atmospheric chemistry box model CAABA/MECCA-4.0, *Geosci. Model Dev.*, 12, 1365-1385, 10.5194/gmd-12-1365-2019, 2019.
- Schlesinger, W. H. and Hartley, A. E.: A global budget for atmospheric NH<sub>3</sub>, *Biogeochemistry*, 15, 191-211, 10.1007/BF00002936, 1992.
- 1225 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics : from air pollution to climate change*, Third edition, John Wiley & Sons, Inc. Hoboken, New Jersey, Hoboken, New Jersey 2016.
- Shah, V., Jaeglé, L., Thornton, J. A., Lopez-Hilfiker, F. D., Lee, B. H., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L., Guo, H., Sullivan, A. P., Weber, R. J., Green, J. R., Fiddler, M. N., Bililign, S., Campos, T. L., Stell, M., Weinheimer, A. J., Montzka, D. D., and Brown, S. S.: Chemical feedbacks weaken the wintertime response of particulate sulfate and nitrate to emissions reductions over the eastern United States, *Proceedings of the National Academy of Sciences*, 115, 8110-8115, doi:10.1073/pnas.1803295115, 2018.
- 1230 Shi, G., Xu, J., Peng, X., Xiao, Z., Chen, K., Tian, Y., Guan, X., Feng, Y., Yu, H., Nenes, A., and Russell, A. G.: pH of Aerosols in a Polluted Atmosphere: Source Contributions to Highly Acidic Aerosol, *Environmental Science & Technology*, 51, 4289-4296, 10.1021/acs.est.6b05736, 2017.
- 1235 Shi, G., Xu, J., Shi, X., Liu, B., Bi, X., Xiao, Z., Chen, K., Wen, J., Dong, S., Tian, Y., Feng, Y., Yu, H., Song, S., Zhao, Q., Gao, J., and Russell, A. G.: Aerosol pH Dynamics During Haze Periods in an Urban Environment in China: Use of Detailed, Hourly, Speciated Observations to Study the Role of Ammonia Availability and Secondary Aerosol Formation and Urban Environment, *Journal of Geophysical Research: Atmospheres*, 124, 9730-9742,

- <https://doi.org/10.1029/2018JD029976>, 2019.
- 1240 Skjoth, C. A. and Geels, C.: The effect of climate and climate change on ammonia emissions in Europe, *Atmos. Chem. Phys.*, 13, 117–128, 10.5194/acp-13-117-2013, 2013.
- Sogaard, H. T., Sommer, S. G., Hutchings, N. J., Huijsmans, J. F. M., Bussink, D. W., and Nicholson, F.: Ammonia volatilization from field-applied animal slurry—the ALFAM model, *Atmospheric Environment*, 36, 3309–3319, [https://doi.org/10.1016/S1352-2310\(02\)00300-X](https://doi.org/10.1016/S1352-2310(02)00300-X), 2002.
- 1245 Song, S., Nenes, A., Gao, M., Zhang, Y., Liu, P., Shao, J., Ye, D., Xu, W., Lei, L., Sun, Y., Liu, B., Wang, S., and McElroy, M. B.: Thermodynamic Modeling Suggests Declines in Water Uptake and Acidity of Inorganic Aerosols in Beijing Winter Haze Events during 2014/2015–2018/2019, *Environmental Science & Technology Letters*, 6, 752–760, 10.1021/acs.estlett.9b00621, 2019.
- 1250 Spengler, J. D., Koutrakis, P., Dockery, D. W., Raizenne, M., and Speizer, F. E.: Health effects of acid aerosols on North American children: air pollution exposures, *Environmental Health Perspectives*, 104, 492–499, doi:10.1289/ehp.96104492, 1996.
- Thurston, G. D., Ito, K., Hayes, C. G., Bates, D. V., and Lippmann, M.: Respiratory Hospital Admissions and Summertime Haze Air Pollution in Toronto, Ontario: Consideration of the Role of Acid Aerosols, *Environmental Research*, 65, 271–290, <https://doi.org/10.1006/enrs.1994.1037>, 1994.
- 1255 Tilgner, A., Schaefer, T., Alexander, B., Barth, M., Collett Jr, J. L., Fahey, K. M., Nenes, A., Pye, H. O. T., Herrmann, H., and McNeill, V. F.: Acidity and the multiphase chemistry of atmospheric aqueous particles and clouds, *Atmos. Chem. Phys.*, 21, 13483–13536, 10.5194/acp-21-13483-2021, 2021.
- Tost, H., Jöckel, P., Kerkweg, A., Sander, R., and Lelieveld, J.: Technical note: A new comprehensive SCAVenging submodel for global atmospheric chemistry modelling, *Atmos. Chem. Phys.*, 6, 565–574, 10.5194/acp-6-565-2006, 2006.
- 1260 Tsimpidi, A. P., Karydis, V. A., and Pandis, S. N.: Response of Inorganic Fine Particulate Matter to Emission Changes of Sulfur Dioxide and Ammonia: The Eastern United States as a Case Study, *Journal of the Air & Waste Management Association*, 57, 1489–1498, 10.3155/1047-3289.57.12.1489, 2007.
- Tsimpidi, A. P., Karydis, V. A., Pozzer, A., Pandis, S. N., and Lelieveld, J.: ~~Global combustion sources of ORACLE (v1.0): module to simulate the organic aerosols: model comparison with 84 AMS factor analysis data sets, Atmos. aerosol composition and evolution in the atmosphere, Geosci. Model Dev., 7, 3153–3172~~ *Chem. Phys.*, 16, 8939–8962, 10.5194/acp-16-8939-2016, 2016 ~~gmd-7-3153-2014, 2014.~~
- 1265 ~~Tsimpidi, A. P., Karydis, V. A., Pandis, S. N., and Lelieveld, J.: Global-scale combustion sources of organic aerosols: sensitivity to formation and removal mechanisms, Atmospheric Chemistry and Physics, 17, 7345–7364, 10.5194/acp-17-7345-2017, 2017.~~
- 1270 Tsimpidi, A. P., Karydis, V. A., Pozzer, A., Pandis, S. N., and Lelieveld, J.: ORACLE (v2-D (v2.0): an efficient module to simulate compute the volatility and oxygen content of organic aerosol composition and evolution in the atmosphere, Geosci. Model Dev., 7, 3153–3172 Development, 11, 3369–3389, 10.5194/gmd-7-3153-2014, 2014 ~~11-3369-2018, 2018.~~
- 1275 ~~Tsimpidi, A. P., Scholz, S. M. C., Milousis, A., Mihalopoulos, N., and Karydis, V. A.: Aerosol Composition Trends during 2000–2020: In depth insights from model predictions and multiple worldwide observation datasets, EGUsphere, 2024, 1–66, 10.5194/egusphere-2024-3590, 2024.~~
- van Aardenne, J. A., Dentener, F. J., Olivier, J. G. J., Goldewijk, C. G. M. K., and Lelieveld, J.: A 1°×1° resolution data set of historical anthropogenic trace gas emissions for the period 1890–1990, *Global Biogeochemical Cycles*, 15, 909–928, <https://doi.org/10.1029/2000GB001265>, 2001.
- 1280 Van Damme, M., Clarisse, L., Whitburn, S., Hadji-Lazaro, J., Hurtmans, D., Clerbaux, C., and Coheur, P.-F.: Industrial and agricultural ammonia point sources exposed, *Nature*, 564, 99–103, 10.1038/s41586-018-0747-1, 2018.
- Van Damme, M., Erisman, J. W., Clarisse, L., Dammers, E., Whitburn, S., Clerbaux, C., Dolman, A. J., and Coheur, P.-F.: Worldwide spatiotemporal atmospheric ammonia (NH<sub>3</sub>) columns variability revealed by satellite, *Geophysical Research Letters*, 42, 8660–8668, <https://doi.org/10.1002/2015GL065496>, 2015.
- 1285 Van Damme, M., Clarisse, L., Heald, C. L., Hurtmans, D., Ngadi, Y., Clerbaux, C., Dolman, A. J., Erisman, J. W., and Coheur, P. F.: Global distributions, time series and error characterization of atmospheric ammonia (NH<sub>3</sub>) from IASI satellite observations, *Atmos. Chem. Phys.*, 14, 2905–2922, 10.5194/acp-14-2905-2014, 2014.
- 1290 Vasilakos, P., Russell, A., Weber, R., and Nenes, A.: Understanding nitrate formation in a world with less sulfate, *Atmos. Chem. Phys.*, 18, 12765–12775, 10.5194/acp-18-12765-2018, 2018.
- Vignati, E., Wilson, J., and Stier, P.: M7: An efficient size-resolved aerosol microphysics module for large-scale aerosol transport models, *Journal of Geophysical Research*, 109, 10.1029/2003JD004485, 2004.
- Walker, J. M., Philip, S., Martin, R. V., and Seinfeld, J. H.: Simulation of nitrate, sulfate, and ammonium aerosols over

Formatted: Indent: Hanging: 0.51 cm

Formatted: Indent: Hanging: 0.51 cm

- the United States, *Atmos. Chem. Phys.*, 12, 11213–11227, 10.5194/acp-12-11213-2012, 2012.
- 1295 Wang, G., Zhang, R., Gomez, M. E., Yang, L., Levy Zamora, M., Hu, M., Lin, Y., Peng, J., Guo, S., Meng, J., Li, J., Cheng, C., Hu, T., Ren, Y., Wang, Y., Gao, J., Cao, J., An, Z., Zhou, W., Li, G., Wang, J., Tian, P., Marrero-Ortiz, W., Secrest, J., Du, Z., Zheng, J., Shang, D., Zeng, L., Shao, M., Wang, W., Huang, Y., Wang, Y., Zhu, Y., Li, Y., Hu, J., Pan, B., Cai, L., Cheng, Y., Ji, Y., Zhang, F., Rosenfeld, D., Liss, P. S., Duce, R. A., Kolb, C. E., and Molina, 1300 M. J.: Persistent sulfate formation from London Fog to Chinese haze, *Proceedings of the National Academy of Sciences*, 113, 13630–13635, doi:10.1073/pnas.1616540113, 2016.
- Wang, M., Kong, W., Marten, R., He, X.-C., Chen, D., Pfeifer, J., Heitto, A., Kontkanen, J., Dada, L., Kürten, A., Yli-  
Juuti, T., Manninen, H. E., Amanatidis, S., Amorim, A., Baalbaki, R., Baccarini, A., Bell, D. M., Bertozzi, B.,  
Bräkling, S., Brilke, S., Murillo, L. C., Chiu, R., Chu, B., De Menezes, L.-P., Duplissy, J., Finkenzeller, H.,  
1305 Carracedo, L. G., Granzin, M., Guida, R., Hansel, A., Hofbauer, V., Krechmer, J., Lehtipalo, K., Lamkaddam, H., Lampimäki, M., Lee, C. P., Makhmutov, V., Marie, G., Mathot, S., Mauldin, R. L., Mentler, B., Müller, T., Onnela, A., Partoll, E., Petäjä, T., Philippov, M., Pospisilova, V., Ranjithkumar, A., Rissanen, M., Rörup, B., Scholz, W., Shen, J., Simon, M., Sipilä, M., Steiner, G., Stolzenburg, D., Tham, Y. J., Tomé, A., Wagner, A. C., Wang, D. S., Wang, Y., Weber, S. K., Winkler, P. M., Wlasits, P. J., Wu, Y., Xiao, M., Ye, Q., Zauner-Wieczorek, M., Zhou, X., Volkamer, R., Riipinen, I., Dommen, J., Curtius, J., Baltensperger, U., Kulmala, M., Worsnop, D. R., Kirkby, J., Seinfeld, J. H., El-Haddad, I., Flagan, R. C., and Donahue, N. M.: Rapid growth of new atmospheric particles by  
1310 nitric acid and ammonia condensation, *Nature*, 581, 184–189, 10.1038/s41586-020-2270-4, 2020.
- Wang, S., Xing, J., Jang, C., Zhu, Y., Fu, J. S., and Hao, J.: Impact Assessment of Ammonia Emissions on Inorganic  
Aerosols in East China Using Response Surface Modeling Technique, *Environmental Science & Technology*, 45,  
1315 9293–9300, 10.1021/es2022347, 2011.
- Wang, X., Zhang, C., Gao, Y., Ji, X., Su, W., and Liu, C.: Satellite unravels recent changes in atmospheric nitrogen  
oxides emissions from global ocean shipping, *Journal of Cleaner Production*, 429, 139591,  
https://doi.org/10.1016/j.jclepro.2023.139591, 2023.
- Wang, X., Wang, W., Yang, L., Gao, X., Nie, W., Yu, Y., Xu, P., Zhou, Y., and Wang, Z.: The secondary formation of  
1320 inorganic aerosols in the droplet mode through heterogeneous aqueous reactions under haze conditions, *Atmospheric Environment*, 63, 68–76, https://doi.org/10.1016/j.atmosenv.2012.09.029, 2012.
- Wang, Y., Zhang, Q. Q., He, K., Zhang, Q., and Chai, L.: Sulfate-nitrate-ammonium aerosols over China: response to  
2000–2015 emission changes of sulfur dioxide, nitrogen oxides, and ammonia, *Atmos. Chem. Phys.*, 13, 2635–  
2652, 10.5194/acp-13-2635-2013, 2013.
- 1325 Warner, J. X., Dickerson, R. R., Wei, Z., Strow, L. L., Wang, Y., and Liang, Q.: Increased atmospheric ammonia over  
the world's major agricultural areas detected from space, *Geophysical Research Letters*, 44, 2875–2884,  
https://doi.org/10.1002/2016GL072305, 2017.
- Weber, R. J., Guo, H., Russell, A. G., and Nenes, A.: High aerosol acidity despite declining atmospheric sulfate  
concentrations over the past 15 years, *Nature Geoscience*, 9, 282–285, 10.1038/ngeo2665, 2016.
- 1330 West, J. J., Ansari, A. S., and Pandis, S. N.: Marginal PM<sub>2.5</sub>: Nonlinear Aerosol Mass Response to Sulfate Reductions  
in the Eastern United States, *Journal of the Air & Waste Management Association*, 49, 1415–1424,  
10.1080/10473289.1999.10463973, 1999.
- Xie, X., Hu, J., Qin, M., Guo, S., Hu, M., Wang, H., Lou, S., Li, J., Sun, J., Li, X., Sheng, L., Zhu, J., Chen, G., Yin,  
J., Fu, W., Huang, C., and Zhang, Y.: Modeling particulate nitrate in China: Current findings and future directions,  
1335 *Environment International*, 166, 107369, https://doi.org/10.1016/j.envint.2022.107369, 2022.
- Xu, J., Chen, J., Zhao, N., Wang, G., Yu, G., Li, H., Huo, J., Lin, Y., Fu, Q., Guo, H., Deng, C., Lee, S. H., Chen, J.,  
and Huang, K.: Importance of gas-particle partitioning of ammonia in haze formation in the rural agricultural  
environment, *Atmos. Chem. Phys.*, 20, 7259–7269, 10.5194/acp-20-7259-2020, 2020.
- 1340 Xu, P., Li, G., Zheng, Y., Fung, J. C. H., Chen, A., Zeng, Z., Shen, H., Hu, M., Mao, J., Zheng, Y., Cui, X., Guo, Z.,  
Chen, Y., Feng, L., He, S., Zhang, X., Lau, A. K. H., Tao, S., and Houlton, B. Z.: Fertilizer management for global  
ammonia emission reduction, *Nature*, 626, 792–798, 10.1038/s41586-024-07020-z, 2024.
- Xu, P., Liao, Y. J., Lin, Y. H., Zhao, C. X., Yan, C. H., Cao, M. N., Wang, G. S., and Luan, S. J.: High-resolution  
inventory of ammonia emissions from agricultural fertilizer in China from 1978 to 2008, *Atmos. Chem. Phys.*, 16,  
1207–1218, 10.5194/acp-16-1207-2016, 2016.
- 1345 Xu, R., Tian, H., Pan, S., Prior, S. A., Feng, Y., Batchelor, W. D., Chen, J., and Yang, J.: Global ammonia emissions  
from synthetic nitrogen fertilizer applications in agricultural systems: Empirical and process-based estimates and  
uncertainty, *Global Change Biology*, 25, 314–326, https://doi.org/10.1111/gcb.14499, 2019.
- Yao, W., Zhang, C., Hao, H., Wang, X., and Li, X.: A support vector machine approach to estimate global solar  
1350 radiation with the influence of fog and haze, *Renewable Energy*, 128, 155–162,  
https://doi.org/10.1016/j.renene.2018.05.069, 2018.

- Yu, F., Nair, A. A., and Luo, G.: Long-Term Trend of Gaseous Ammonia Over the United States: Modeling and Comparison With Observations, *Journal of Geophysical Research: Atmospheres*, 123, 8315-8325, <https://doi.org/10.1029/2018JD028412>, 2018.
- 1355 Zhang, B., Shen, H., Liu, P., Guo, H., Hu, Y., Chen, Y., Xie, S., Xi, Z., Skipper, T. N., and Russell, A. G.: Significant contrasts in aerosol acidity between China and the United States, *Atmos. Chem. Phys.*, 21, 8341-8356, 10.5194/acp-21-8341-2021, 2021a.
- Zhang, L., Chen, Y., Zhao, Y., Henze, D. K., Zhu, L., Song, Y., Paulot, F., Liu, X., Pan, Y., Lin, Y., and Huang, B.: Agricultural ammonia emissions in China: reconciling bottom-up and top-down estimates, *Atmos. Chem. Phys.*, 18, 339-355, 10.5194/acp-18-339-2018, 2018.
- 1360 Zhang, L., Gong, S., Zhao, T., Zhou, C., Wang, Y., Li, J., Ji, D., He, J., Liu, H., Gui, K., Guo, X., Gao, J., Shan, Y., Wang, H., Wang, Y., Che, H., and Zhang, X.: Development of WRF/CUACE v1.0 model and its preliminary application in simulating air quality in China, *Geosci. Model Dev.*, 14, 703-718, 10.5194/gmd-14-703-2021, 2021b.
- 1365 Zhang, Q. and Geng, G.: Impact of clean air action on PM2.5 pollution in China, *Science China Earth Sciences*, 62, 1845-1846, 10.1007/s11430-019-9531-4, 2019.
- Zhao, P., Zhang, X., Xu, X., and Zhao, X.: Long-term visibility trends and characteristics in the region of Beijing, Tianjin, and Hebei, China, *Atmospheric Research*, 101, 711-718, <https://doi.org/10.1016/j.atmosres.2011.04.019>, 2011.
- 1370 Zheng, B., Zhang, Q., Zhang, Y., He, K. B., Wang, K., Zheng, G. J., Duan, F. K., Ma, Y. L., and Kimoto, T.: Heterogeneous chemistry: a mechanism missing in current models to explain secondary inorganic aerosol formation during the January 2013 haze episode in North China, *Atmos. Chem. Phys.*, 15, 2031-2049, 10.5194/acp-15-2031-2015, 2015.
- Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14095-14111, 10.5194/acp-18-14095-2018, 2018.
- 1375 Zheng, G., Su, H., and Cheng, Y.: Revisiting the Key Driving Processes of the Decadal Trend of Aerosol Acidity in the U.S, *ACS Environmental Au*, 2, 346-353, 10.1021/acsenvironau.1c00055, 2022.
- Zheng, G., Su, H., Andreae, M. O., Pöschl, U., and Cheng, Y.: Multiphase Buffering by Ammonia Sustains Sulfate Production in Atmospheric Aerosols, *AGU Advances*, 5, e2024AV001238, <https://doi.org/10.1029/2024AV001238>, 2024.
- 1380 Zheng, G., Su, H., Wang, S., Andreae, M. O., Pöschl, U., and Cheng, Y.: Multiphase buffer theory explains contrasts in atmospheric aerosol acidity, *Science*, 369, 1374-1377, doi:10.1126/science.aba3719, 2020.
- Zhou, M., Deng, Z., Robert, C., Zhang, X., Zhang, L., Wang, Y., Qi, C., Wang, P., and Mazière, M. D.: The First Global Map of Atmospheric Ammonia (NH<sub>3</sub>) as Observed by the HIRAS/FY-3D Satellite, *Advances in Atmospheric Sciences*, 41, 379-390, 10.1007/s00376-023-3059-9, 2024.
- 1385 Zhou, M., Zheng, G., Wang, H., Qiao, L., Zhu, S., Huang, D., An, J., Lou, S., Tao, S., Wang, Q., Yan, R., Ma, Y., Chen, C., Cheng, Y., Su, H., and Huang, C.: Long-term trends and drivers of aerosol pH in eastern China, *Atmos. Chem. Phys.*, 22, 13833-13844, 10.5194/acp-22-13833-2022, 2022.
- 1390 Zhou, W., Gao, M., He, Y., Wang, Q., Xie, C., Xu, W., Zhao, J., Du, W., Qiu, Y., Lei, L., Fu, P., Wang, Z., Worsnop, D. R., Zhang, Q., and Sun, Y.: Response of aerosol chemistry to clean air action in Beijing, China: Insights from two-year ACSM measurements and model simulations, *Environmental Pollution*, 255, 113345, <https://doi.org/10.1016/j.envpol.2019.113345>, 2019.