

Untangling the effects of vertical mixing schemes and convective adjustment in the Mediterranean Sea: insights from a sensitivity study

Lucia Gualtieri¹, Hans Burchard², Federica Borile³, Aimie Moulin¹, Pietro Miraglio¹, Francesco Maicu¹, Emanuela Clementi¹, Paolo Oddo³

¹CMCC Foundation - Euro-Mediterranean Center on Climate Change, Bologna, Italy.

²Leibniz Institute for Baltic Sea Research Warnemünde, Rostock, Germany.

³University of Bologna, Department of Physics and Astronomy, Bologna, Italy.

Correspondence to: Lucia Gualtieri (lucia.gualtieri@cmcc.it)

Abstract. The Mediterranean Sea provides a natural laboratory for investigating ocean circulation processes of global relevance due to its complex dynamics, active deep and intermediate water formation, and sensitivity to climate variability. Regional ocean circulation models' skill strongly depends on the parameterization of subgrid-scale processes, among which turbulent vertical mixing and convection play a major role. To evaluate their impact in the Mediterranean Sea two-year-long simulations were conducted using three vertical closure schemes: the Richardson-number dependent parameterisation, the Turbulent Kinetic Energy (TKE) scheme, and the Generalised Length Scale (GLS) scheme. Each scheme was tested both with and without the convective adjustment approach, resulting in a set of comparative experiments designed to isolate the contribution of the adjustment process and its combined effect with each parameterization. Model results are evaluated against all available Argo floats data, both at the basin scale and in key deep and intermediate water formation regions. The simulations show that adding a convective adjustment is crucial to accurately reproduce observations with the Richardson-number dependent parameterization, where it improves all key variables, while for the TKE scheme it is particularly important for representing the mixed layer depth across the basin and in deep water formation areas. For more physics-based vertical schemes, like the GLS closure, the convective adjustment is mostly redundant and can occasionally degrade results. Overall, the GLS scheme without any convective adjustment provides the most accurate representation of the mixed layer depth as well as the vertical structure and variability both at basin scale and in key regions of deep and intermediate water formation.

1 Introduction

The Mediterranean Sea is a semi-enclosed basin characterized by intense thermohaline variability, strong mesoscale activity, and complex interactions with the atmosphere and surrounding land masses. Its dynamical features, including dense water formation, flux exchanges through narrow straits, and sub-basin circulations, make it a particularly challenging environment for ocean modeling (e.g., Pinardi and Masetti, 2000; Millot and Taupier-Letage, 2005; Testor et al., 2018). The Mediterranean Sea plays a crucial role in shaping the regional climate (e.g. Pinardi and Navarra, 1993; Lionello et al., 2006; Pinardi and Masetti, 2000) and represents a natural laboratory for investigating key ocean processes with implications at the global scale.

Deleted: Paolo Oddo²,

Deleted: ³

Deleted: ²

Deleted: ³

Deleted: ¶

Deleted: ²

Moved up [2]: ³Leibniz Institute for Baltic Sea Research Warnemünde, Rostock, Germany.¶

Deleted: the context of

Deleted: –

Deleted: –

Deleted: combined with

Deleted: a

Deleted: with the aim to study their interaction

Deleted: Despite its relatively small size, t

In this sense, it is often considered a “miniature ocean” (e.g., Bethoux et al., 1999), where processes such as deep and intermediate water formation, strait exchanges, mesoscale variability, and air–sea interactions can be studied in a confined environment as a small-scale analogue of global processes.

50 Over the past two decades, regional ocean models have been widely employed to simulate the Mediterranean circulation at increasing spatial and temporal resolution. Among these, the physical component of the Mediterranean Forecasting System, developed within the framework of the Copernicus Marine Service, represents a state-of-the-art implementation (e.g., Tonani et al., 2009; Clementi et al., 2017; Coppini et al., 2023). It provides regular, open-access analyses, forecasts (Clementi et al., 2023) and reanalyses (Escudier et al., 2021) of the basin’s physical state and serves as a reference framework for scientific and
55 operational applications.

Due to high computational costs and missing physics, the spatial and temporal resolution of current hydrostatic ocean circulation models, even at regional scale, is unable to explicitly resolve the inherently non-hydrostatic and small-scale turbulent vertical mixing and convective processes. To this end, vertical mixing closure schemes – either based on simplified
60 formulations (e.g., Pacanowski and Philander, 1981) or on more physically based models (e.g., Gaspar et al., 1990; Umlauf and Burchard, 2003) – are employed to mimic these turbulent processes. Also, deep convection, which corresponds to the formation of dense deep water masses, has to be represented through parameterizations to reproduce its timing, intensity, and spatial extent realistically (e.g., Marshall et al., 1999; Villarreal et al., 2005; Herrmann et al., 2008; Luneva et al., 2019). Note that convection causes turbulent vertical mixing under unstably stratified conditions and can be parameterized separately
65 (convective adjustment) or within the same turbulence closure scheme (Umlauf and Burchard, 2003; Legay et al., 2025).

Physics-based turbulent closure schemes include buoyancy production terms and are, in principle, able to represent convective mixing when the stratification becomes unstable. However, in practice these schemes may not always remove static instability rapidly enough, particularly at the temporal and vertical resolution typically used in regional ocean models. This can lead to an underestimation of convective mixing. For this reason, ocean circulation models usually combine turbulence closure schemes with a convective adjustment parameterization (e.g. Clementi et al., 2017; Coppini et al., 2023). The convective adjustment acts by stabilizing the water column through a rapid vertical redistribution of tracers, ensuring a prompt restoration of stable stratification. In this sense, it does not represent an independent physical process, but rather a numerical parameterization designed to mimic the fast overturning associated with unresolved convective events.

70

75 This combined approach has been widely adopted in the modelling of the Mediterranean Sea for several decades (Pinardi et al., 2003; Pinardi and Coppini, 2010), mostly because the relatively simple turbulence closure schemes traditionally employed were not sufficient to adequately represent convective mixing. In this context, it is essential to assess how these earlier, less complete schemes compare with more advanced formulations which are specifically designed to provide a more consistent

and physically based representation of turbulent and convective processes. Depending on the turbulence closure scheme adopted, the use of an explicit convective parameterization may or may not be required. Assessing this aspect is one of the main objectives of the present study.

The choice of mixing scheme and convective adjustment parameterisations has been shown to play a crucial role in shaping the circulation, the temperature and salinity distributions and deep-water formation (DWF) processes (e.g., Herrmann et al., 2008; Refray et al., 2015; Luneva et al., 2019; Robertson et al., 2019; Gutjahr et al., 2021). In particular, vertical mixing controls the redistribution of heat and salt throughout the water column, thereby influencing stratification and mixed layer depth (MLD). Convection, on the other hand, strongly affects deep and intermediate water formation, which in turn impacts large-scale circulation patterns and water mass properties. Since these processes are tightly coupled to air–sea exchanges and mesoscale dynamics, model sensitivity to the choice of mixing and convection schemes can propagate across a wide range of spatiotemporal scales, ultimately influencing the realism of simulated ocean variability.

Only a few previous studies provide a context for understanding the performance of different mixing schemes, but none of these explicitly investigate their combined effect with a convective adjustment, and whether a mixing scheme alone is sufficient to reproduce the vertical structure and variability of the upper ocean. Madec et al. (1991) combined Richardson-number dependent closures with nonpenetrative convective adjustment in studies of deep-water formation in the Northwestern Mediterranean Sea. Refray et al. (2015) investigated the sensitivity of turbulent vertical mixing using a 1-D ocean circulation model at the global scale, comparing various mixing schemes but without including convective adjustment. Gutjahr et al. (2021) compared various global-scale mixing schemes – including among the others, Richardson-number dependent closures and Turbulent Kinetic Energy (TKE) – but did not assess the role of convective adjustment. Storto et al. (2023) focused on the Mediterranean Sea, comparing the Generalized Length Scale (GLS) and TKE schemes, but did not examine cases of active convection and therefore did not evaluate the potential impact of including a convective adjustment. By explicitly investigating the interplay between vertical mixing schemes and convective adjustment, the present study addresses a gap left by these prior works and demonstrates how convective adjustment modulates the behaviour of different schemes in the context of the Mediterranean Sea.

The aim of this study is to investigate the sensitivity of a modified configuration of the Mediterranean Forecasting System (Clementi et al., 2023) to the choice of vertical mixing scheme and to the inclusion of an explicit convection adjustment parameterization. Model performance is evaluated across spatial and temporal scales through an intercomparison of temperature, salinity and mixed layer depth, from basin scale over years to deep and intermediate water formation regions ranging from months to hours. Given that the formation of deep and intermediate waters in the Mediterranean occurs on seasonal timescales, with interannual variability, a two-year model integration is sufficient to capture the main differences between experiments.

Deleted: deep water

Deleted: deep water

Deleted: The experiments are performed using a 1/24° regional ocean model based on the Nucleus for European Modelling of the Ocean (NEMO; Madec et al., 2023, doi:10.5281/zenodo.6334656) coupled with the wind wave model WaveWatch III (WW3; Tolman et al., 2021), and forced by realistic atmospheric conditions over the two-year period 2020–2021. Three turbulence closure schemes are tested: the Richardson-number dependent parameterization (Pacanowski and Philander, 1981), the TKE closure scheme (Gaspar et al., 1990), and the GLS closure scheme (Umlauf and Burchard, 2003). These vertical mixing schemes are employed under two different convective setups: with and without the enhanced vertical diffusion parameterization typically used to mimic deep convection events. ...

130

The rest of the paper is structured as follows. Section 2 describes the region and the observational dataset used to assess our model performances (Subsection 2.1). It also describes the model configuration (Subsection 2.2), with a focus on the choice of the vertical mixing schemes and convective adjustment, and the experimental design performed in this study (Subsection 2.3). Section 3 presents the main results of this study. We first present the convective adjustment (Subsection 3.1). Then we show results at the basin scale over the entire two-year period (Subsection 3.2). We focus then on deep and intermediate water formation regions over months to days (Subsection 3.3), and we investigate summer daily convection events at hourly temporal scales (Subsection 3.4). Finally, Section 4 discusses the key conclusions and outlines perspectives for future work.

Deleted: s

2 Data and methods

2.1 Argo float data in the Mediterranean Sea

140

We use Argo float data (Wong et al., 2020; Poulain et al., 2007) from the Copernicus Marine Service Global Ocean near real-time in situ quality controlled observational product (Global Ocean-In-Situ Near-Real-Time Observations, <https://doi.org/10.48670/moi-00036>) as a reference observational dataset for our model validation. The location of Argo floats in the Mediterranean Sea during the two-year period under investigation (2020–2021) is reported as red dots in Fig. 1a. This dataset consists of 11,053 Argo profiles across the Mediterranean Sea, which allows for a statistically robust analysis of the processes under investigation while ensuring homogeneous quality control. Except for a few regions – such as the North Adriatic Sea, the Aegean Sea, the Strait of Sicily and the Gulf of Sirte – the Argo data in this time period provide good spatial coverage across the major part of the basin, offering a statistically meaningful dataset for our model validation.

Moved down [1]: Our model domain encompasses the entire Mediterranean Sea and extends into a portion of the Atlantic Ocean (Fig. 1a). Including an Atlantic box is essential to accurately resolve water exchanges between the Mediterranean Sea and the Atlantic Ocean through the Strait of Gibraltar, as the inflow of fresher Atlantic water and the outflow of saltier Mediterranean water strongly influence basin-wide salinity, temperature, and density structure, as well as deep water formation. Nevertheless, the analysis performed in this study focuses only on the Mediterranean Sea.

Deleted: ¶

Deleted: basin

Deleted: 2

150

Typical temperature and salinity profiles retrieved by Argo floats in the Mediterranean Sea are shown in Fig. 1 as monthly average vertical profiles of temperature and salinity in January 2021 (Fig. 1b) and July 2021 (Fig. 1c). The observations exhibit strong variability around the mean both in temperature and salinity, especially in the upper ~200 m, highlighted by shading representing one standard deviation of the dataset. It is known that, both temperature and salinity show also a pronounced seasonal variability (e.g., Pinardi and Masetti, 2000). In winter (Fig. 1b) stratification is weakened because of surface cooling and increased vertical mixing, resulting in the formation of a thick mixed layer with relatively uniform temperature and salinity. In summer (Fig. 1c), temperature and salinity values close to the surface become higher with respect to winter as the result of strong heating and evaporation. These processes inhibit vertical mixing and give rise to a well-defined thermohaline stratification. As a result, vertical profiles are more homogeneous in winter and strongly stratified in summer.

Deleted: their

Deleted: warmer and saltier

160

The Mediterranean Sea hosts several key regions of deep and intermediate water formation, including the Gulf of Lion, the South Adriatic Sea, the Rhodes Gyre, and the Aegean Sea (see map in Fig. 1a). In these areas, vertical mixing and convection

180 play a central role in preconditioning and driving dense water formation processes by modulating stratification and contributing to surface buoyancy loss. Fig. 1 also shows the seasonal variability of temperature and salinity profiles from Argo floats for two of these regions: the South Adriatic Pit (Fig. 1d) and the Rhodes Gyre region (Fig. 1e). In both regions, temperature profiles exhibit a pronounced seasonal variability in the upper layers, especially near the surface, closely following the seasonality of the surface heat fluxes. On the other hand, salinity shows distinct seasonal changes associated with deep and intermediate water formation processes. In the South Adriatic Pit (Fig. 1d), the winter period (i.e., December-January-February) is influenced by the intrusion of intermediate water formed in the Levantine basin, visible as a marked salinity increase between about 100 m and 200 m. Deep-water formation is not clearly visible, being largely smoothed by seasonal averaging, reflecting its short duration (order of weeks) and spatially localized nature. In the Rhodes Gyre region (Fig. 1e), where intermediate water forms, salinity shows a persistent increase throughout the year. The salinity maximum is located at about 150 m in winter, during the formation phase, and deepens to about 200 m in spring (i.e., March-April-May) and to about 250 m in summer (i.e., June-July-August) and autumn (i.e. September-October-November).

Deleted: ¶





Figure 1: (a) Model domain including the Mediterranean Sea and an Atlantic box, with bathymetry as a background colour and labels for the main subbasins and deep and intermediate water formation regions. Red dots indicate Argo floats during 2020–2021. The black zonal line at 37.5°N marks the location of a vertical section used throughout the paper. **Middle panels** show the basin-averaged observed temperature (blue) and salinity (orange) for (b) January 2021 and (c) July 2021, where shaded areas represent one standard deviation. **Bottom panels** show the season-averaged observed temperature (blue) and salinity (orange) in (d) the South Adriatic Pit and (e) the Rhodes Gyre region.



Deleted:

Deleted: Bottom

Deleted: pannels

Deleted:

2.2 Model setup

Our model setup builds on the Mediterranean Forecasting System model of the Copernicus Marine Service. The model domain encompasses the entire Mediterranean Sea and extends into a portion of the Atlantic Ocean (Fig. 1a), spanning from 18° 7.50' W to 36° 17.50' E in longitude and 30° 11.25' N and 45° 58.75' N in latitude. Including an Atlantic box is essential to accurately resolve water exchanges between the Mediterranean Sea and the Atlantic Ocean through the Strait of Gibraltar, as the inflow of fresher Atlantic water and the outflow of saltier Mediterranean water strongly influence basin-wide salinity, temperature, and density structure, as well as DWF. Nevertheless, the analysis performed in this study focuses only on the Mediterranean Sea.

The model system consists of a two-way coupled hydrodynamic circulation model and a third-generation spectral wind wave model. The ocean hydrodynamic circulation model is based on Nucleus for European Modelling of the Ocean (NEMO; Madec et al., 2023, doi:10.5281/zenodo.6334656), version 4.2 and the wind wave model on WaveWatch III (WW3; Tolman et al., 2021), version 6.07. The coupling happens hourly online with the exchange of air-sea temperature difference and surface currents from NEMO to WW3, and of neutral drag coefficient used to evaluate the surface wind stress from WW3 to NEMO (Clementi et al., 2017).




Figure 2: Vertical layer thickness as a function of depth. A zoom of the upper 50 m highlights the fine vertical discretization within the mixed layer.

Deleted: Our model domain
Formatted: Font: Times New Roman, 10 pt
Formatted: Font: Times New Roman, 10 pt
Formatted: Font: Times New Roman, 10 pt
Formatted: Font: Times New Roman, 10 pt

Deleted: deep water formation

Deleted: NEMO
Deleted: (Madec et al., 2023)
Deleted: WW3
Deleted: (Tolman et al., 2021)
Deleted: d

Formatted: Keep with next

Formatted: Caption

230 The model has a horizontal resolution of $1/24^\circ$ (about 4 km) in both latitude and longitude, and it includes 141 non-uniformly distributed vertical levels. The vertical layer thickness varies with depth (Fig. 2), with particularly fine resolution in the upper ocean (from about 2.1 m near the surface to 4.6 m at 50 m depth). This dense discretization in the mixed layer allows for an accurate representation of key upper-ocean processes, such as mixing and stratification, which are essential for capturing surface-driven dynamics. The time stepping is achieved within a leapfrog differencing structure associated with an Asselin filter and the baroclinic time step is set to 180 s. To capture the effects of intensified vertical mixing induced by internal wave breaking in the Strait of Gibraltar, particularly near Camarinal Sill (Wesson and Gregg, 1994; Hilt et al., 2020), we also prescribe an enhanced vertical diffusivity farther to the east of the sill.

Horizontal tracer advection is computed using the fourth-order Flux-Corrected Transport (FCT) scheme, which provides a good balance between accuracy and numerical stability by combining a high-order transport method with a monotonicity constraint (Levy, 2001). Lateral diffusion of tracers is implemented as a Laplacian operator along isoneutral surfaces, with a spatially constant diffusivity. Horizontal momentum advection is treated in flux form using the third-order Upstream-Biased Scheme (UBS). As for tracers, momentum lateral diffusion is applied using a Laplacian operator along isoneutral surfaces, again with constant viscosity. These parameterisations are widely used in NEMO-based configurations for large-scale and regional ocean modelling, providing physically consistent dissipation while preserving the large-scale circulation features (e.g., Madec et al., 2023; Le Sommer et al., 2009).

2.2.1 Vertical mixing scheme

In this study, we compare three different parameterisations for vertical mixing, representative of the main families of vertical mixing parameterizations commonly used in ocean modelling. Our aim is to compare more simple approaches – such as the Richardson-number dependent scheme, which is still used in the Mediterranean Forecasting system of the Copernicus Marine Service (Clementi et al., 2017; Coppini et al., 2023) – with more physically based formulations. These physics-based parameterizations rely on a prognostic treatment of turbulence and are designed to capture a wider range of oceanic mixing processes with greater physical accuracy. Physically based vertical mixing schemes typically fall into two categories: one-equation and two-equation turbulence closure models. One-equation schemes solve a prognostic equation for the TKE while relying on diagnostic expressions or empirical assumptions for the mixing length or the dissipation rate (e.g., Gaspar et al., 1990). Two-equation models introduce an additional prognostic variable – usually the turbulence length scale or the dissipation rate of turbulent kinetic energy – which allows for a more flexible and accurate representation of turbulence in stratified and sheared flows (e.g., Mellor and Yamada, 1982; Umlauf and Burchard, 2003). Each of these formulations requires calibrating several parameters. In this study, the best configuration for each scheme was selected from sensitivity tests as the one minimising statistical errors at the basin scale when compared with all available Argo data.

Deleted: , and uses a baroclinic time step of 180 s

Deleted: The system is forced at the ocean surface with the atmospheric fields at $1/10^\circ$ horizontal resolution of the European Centre for Medium-Range Weather Forecasts (ECMWF) made available by the Italian Air Force Meteorological Service. At the Atlantic boundary, the hydrodynamic model is nested into the daily analysis and forecast data of the Copernicus Marine Global product (Global Ocean Physics Analysis and Forecast, <https://doi.org/10.48670/moi-00016>) at $1/12^\circ$ horizontal resolution and 50 vertical levels. River runoff in the modelling implementation originates from 39 rivers. Daily mean data for 38 of these rivers are taken from the European Flood Awareness System dataset (EFAS v5.0; Mazzetti et al., 2023) delivered within the Copernicus Emergency Service, while monthly mean data for the Nile River are taken from a climatology computed over the period 1970–2007 (Said et al., 2011).

Deleted: Gerbeau and LeRoux

Deleted: latter

Deleted: TKE

280 The first vertical mixing scheme used in this study is a Richardson number–dependent formulation originally proposed by Pacanowski and Philander (1981). This scheme was originally developed for the equatorial ocean, using a rectangular box model, and was not designed to provide prognostic mixing. Vertical eddy viscosity and diffusivity are parameterised as functions of the Richardson number (see Appendix A). The Richardson number expresses the ratio between stratification (buoyancy) and vertical shear, and it quantifies their relative dominance. Low Richardson number values indicate that shear overcomes stratification, promoting instability, while high values correspond to strongly stratification. Accordingly, this scheme increases mixing when shear effects dominate over buoyancy and reduces it when buoyancy prevails. Specific parameters control the sensitivity of the mixing to these conditions, while background values ensure a minimum level of vertical mixing even under stable conditions. The Richardson-dependent mixing parameterization does not simulate convective overturning, but just reduces density gradients by enhancing mixing in unstable layers when shear dominates over buoyancy.

285

290 See Appendix B (Table B1), for details on the parameters used in the NEMO namelist for the Richardson number–dependent formulation.

The second vertical mixing scheme adopted here is the one-equation TKE scheme as proposed by Gaspar et al. (1990), in which the evolution of the turbulent kinetic energy k is described by a prognostic equation (see Appendix A). The model

295 accounts for the main four physical processes involved in turbulence dynamics: production by vertical shear, destruction by buoyancy effects, turbulent kinetic energy diffusion, and dissipation parameterised following the Kolmogorov formulation (Kolmogorov, 1942). Vertical eddy viscosity A^{vm} is diagnosed from the TKE using a mixing length approach, where the mixing length is parametrized using a diagnostic formulation, as a function of both a prescribed maximum and the local stratification to ensure numerical stability by limiting its vertical gradients. Tracer diffusivity $A^{vT} = A^{vS}$ for temperature and salinity,

300 respectively, is derived from A^{vm} through a constant turbulent Prandtl number ($A^{vT} = A^{vm} / Pr$). At the ocean surface, we impose a fixed minimum surface mixing length. A simplified representation of Langmuir turbulence is also included. The TKE scheme treats convection prognostically through the calculation of enhanced turbulent kinetic energy production in statically unstable layers, which results from the combined contribution of the four terms described above: shear production, stratification or buoyancy destruction, diffusion and dissipation. The resulting enhanced turbulent kinetic energy production

305 increases the eddy diffusivity coefficients and leads to the homogenization of these layers. See Appendix B (Table B2), for details on the parameters used in the NEMO namelist for the TKE closure scheme.

The third vertical mixing scheme used in this work is a two-equation turbulence closure scheme defined within the GLS framework (Umlauf and Burchard, 2003). The GLS schemes are a family of models that differ mainly in the choice of the

310 second turbulence variable to be worked out through a prognostic equation and the stability functions used to parameterise stratification effects on turbulence. In addition to the turbulent kinetic energy k , the second prognostic variable can be defined in different ways, leading to different two-equation turbulence models. In the $k-\varepsilon$ formulation, the second variable is the turbulent dissipation rate ε , which quantifies the rate at which turbulent kinetic energy is dissipated into heat. In the $k-\omega$

Deleted: ,

Deleted: where v

Deleted: balance

Deleted: (see Appendix A)

Deleted: In this way,

Deleted: enhances

Deleted: in regions of

Deleted: intense shear (or weak stratification)

Deleted: under weak shear (or strong

Deleted: stratification

Deleted:)

Deleted: ,

Deleted: ,

formulation, it is the specific dissipation rate $\omega = \epsilon/k$ (where ϵ represents the turbulent kinetic energy dissipation rate) which has the dimension of an inverse time and represents a characteristic turbulence frequency. Alternatively, in the $k-k_t$ formulation, the second variable is the turbulent length scale k_t , describing the typical size of the energy-containing eddies, usually expressed

330 as $k_t \simeq \frac{k^{3/2}}{\epsilon}$.

In this study, we adopt the k - ϵ formulation, where the prognostic variables are the turbulent kinetic energy k and its dissipation rate ϵ (Umlauf and Burchard, 2003), but we also verified that using the k - ω formulation does not change significantly our conclusions. This model captures turbulence dynamics with enhanced detail, particularly under varying stratification, and is known for its numerical stability and robust performance in oceanic applications (Burchard and Bolding, 2001). The model prognoses the temporal evolution of k and ϵ , accounting for the same four physical processes considered in the TKE scheme: shear production, buoyancy destruction, vertical turbulent diffusion, and dissipation. Vertical eddy viscosity and diffusivity coefficients are computed dynamically based on k and ϵ and modulated by stability functions dependent on the gradient of the Richardson number. These stability functions, which are a common feature of turbulence closures, represent the suppression of turbulence by reducing mixing efficiency under stable stratification. In this study, we adopt the stability function formulation version A by Canuto et al., 2001 (*Canuto A*). Specifically, the momentum mixing efficiency decreases as stratification increases, while the turbulent Prandtl number correspondingly increases, implying a stronger reduction of tracer mixing relative to momentum (Canuto et al., 2001; Umlauf and Burchard, 2003). Boundary conditions at the surface and at the bottom incorporate enhancements to turbulence caused by wave breaking and bottom friction, respectively. Wave-induced mixing is not taken into account. The GLS scheme accounts for convection by prognostically calculating turbulent kinetic energy production and simultaneously estimating a turbulent length scale. While the computation of turbulent kinetic energy is conceptually similar to that in the TKE scheme, the key difference lies in the prognostic treatment of the turbulent length scale, which enables not only a dynamical adjustment of mixing intensity, but also a more physically consistent representation of the vertical extent of convective overturning. We also note that, while the mixed-layer depth is correctly estimated under convective conditions, the stratification is falsely prescribed by the GLS closure to be unstable also in the lower half of the convective boundary layer (Burchard and Petersen, 1999). In that region, typically counter-gradient fluxes are present which need to be parameterised by a higher-order closure (Legay et al., 2025). More details on the GLS closure scheme are in Appendix A and the parameters used in the NEMO namelist are reported in Appendix B (Table B3).

355 2.2.2 Convective adjustment

Convective events are characterized by rapid and intense overturning of the water column, an inherent non-hydrostatic process that cannot be explicitly resolved by hydrostatic ocean models. As a result, convection must be represented through a dedicated parameterization. To better capture convective processes, it is still a common practice in large-scale modelling to apply a

Deleted: ϵ

Deleted: ϵ

Deleted: ,

convective adjustment whenever the water column is statically unstable to mimic the rapid vertical homogenization associated with convective overturning and ensures realistic mixed layer deepening. One of the most common ways is to apply the enhanced vertical diffusivity (EVD) parameterisation (e.g., Lazar et al., 1999), widely adopted in ocean modelling (e.g., Rahmstorf, 1993; Lazar et al., 1999; Griffies et al., 2000). In practice, the vertical eddy diffusivity coefficient is increased to a high constant value (typically between 1 and 100 m² s⁻¹) wherever the Brunt-Väisälä frequency squared N^2 becomes negative. In this study, a value of 10 m² s⁻¹ is prescribed for the vertical eddy diffusivity based on previous sensitivity tests. This is also the value currently used in the physical component of the Mediterranean Forecasting System of the Copernicus Marine Service.

Deleted: Large and Deleersnijder, 1993;

Deleted: in these cases

Since convection has the effect of effectively homogenizing the upper ocean boundary layer, it can be parameterized by applying a high eddy diffusivity which has a similar effect. Although this procedure does not accurately represent the physics of convection with its narrow and dense downward plumes compensated by a broader upwelling in between, leading to non-local fluxes, it effectively mimics the net impact of convective processes, including the rapid vertical redistribution of properties associated with convective overturning. The increased eddy diffusivity coefficient stabilizes the water column, allowing the model to maintain realistic stratification while enabling the development of large-scale circulation patterns.

When the convective adjustment is applied, the vertical eddy diffusivity results from the combined effects of the vertical mixing closure schemes (whether the Richardson-number-dependent parameterization, the one-equation TKE scheme, or the two-equation GLS formulation) and the convective adjustment implemented through the EVD parameterisation. Since the TKE and GLS schemes implicitly include a convection parameterisation via the buoyancy term in the turbulent kinetic energy equation being positive for unstable stratification with N^2 (equations (A3) and (A6), respectively), some double counting is included in this approach. It should be noted that the GLS mixing parameterisation includes further sensitivity to convectively unstable stratification due to the dependence of the stability functions and the length-scale related equation on N^2 .

Deleted: Therefore, w

Deleted: TKE

Deleted: (A3) and (A6)

2.3 Experimental design

We perform six, two-year long simulations over the period 2020–2021. These simulations share the same model setup (as described in Section 2.2), and the same input fields and differ only in the choice of the vertical mixing scheme and whether the convective adjustment is applied.

Deleted: (

Deleted:)

Deleted: ,

The system is forced at the ocean surface with the atmospheric fields at 1/10° horizontal resolution of the European Centre for Medium-Range Weather Forecasts (ECMWF) made available by the Italian Air Force Meteorological Service. Following the configuration of the physical component of the Mediterranean Forecasting System of the Copernicus Marine Service, at the Atlantic boundary, the hydrodynamic model is nested into the daily analysis data of the Copernicus Marine Global product (whose Copernicus product name is *Global Ocean Physics Analysis and Forecast*, <https://doi.org/10.48670/moi-00016>) at 1/12° horizontal resolution and 50 vertical levels. River runoff in the modelling implementation originates from 39 rivers.

Formatted: Font: Italic

405 Daily mean data for 38 of these rivers are taken from the European Flood Awareness System dataset (EFAS v5.0; Mazzetti et al., 2023) delivered within the Copernicus Emergency Service, while monthly mean data for the Nile River are taken from a climatology computed over the period 1970–2007 (Said et al., 2011).

410 Three turbulence closure schemes are tested: the Richardson-number dependent parameterization (Pacanowski and Philander, 1981), the TKE closure scheme (Gaspar et al., 1990), and the GLS closure scheme (Umlauf and Burchard, 2003). These vertical mixing schemes are employed under two different convective setups: with and without the enhanced vertical diffusion parameterization typically used to mimic deep convection events. Therefore, the six simulations performed here differ only in the vertical mixing scheme and in whether convective adjustment is included. This experimental design allows us to assess the interplay between the choice of vertical mixing scheme and the effect of EVD, and to disentangle their individual and combined contributions to the representation of vertical mixing processes. It also enables us to evaluate how different turbulence closure schemes respond to the presence or absence of convective adjustment, and how this affects stratification, heat and salt distribution, and the overall vertical structure of the water column at basin and local scale.

420 Table 1 summarises the experiments. Our control run is defined as the simulation that employs the Richardson-number dependent vertical mixing scheme in combination with convective adjustment. Keeping convective adjustment active, we then replace the vertical mixing scheme with either TKE or GLS parameterisations. The remaining three experiments use the same three mixing schemes, but with convective adjustment disabled. In the experiments where convective adjustment is disabled, adaptive implicit vertical advection is activated to prevent local numerical instabilities (Shchepetkin, 2015, Madec et al., 2023). All simulations are initialized from the same 31 December 2019 restart fields provided by the Mediterranean Forecasting System. These fields are analysis products that incorporate data assimilation of available observations, including satellite and in-situ measurements. As a result, they provide an optimal representation of the Mediterranean Sea state at the start of the simulations, ensuring that all experiments begin from a physically realistic initial condition. The results presented hereafter are not affected by these initial conditions but rather reflect the changes in the vertical mixing schemes and convective adjustment listed in Table 1.

Simulation name	Vertical mixing closure scheme	Convective adjustment	Notes
Exp-PP ⁺	Richardson-dependent scheme	Enhanced Vertical Diffusion (EVD)	Control simulation
Exp-TKE ⁺	Turbulent Kinetic Energy (TKE)	Enhanced Vertical Diffusion (EVD)	
Exp-GLS ⁺	Generalized Length Scale (GLS)	Enhanced Vertical Diffusion (EVD)	
Exp-PP	Richardson-dependent scheme	–	
Exp-TKE	Turbulent Kinetic Energy (TKE)	–	
Exp-GLS	Generalized Length Scale (GLS)	–	

Table 1: Summary of the experiments conducted in this study.

Deleted: varying

Deleted: the inclusion

Deleted: or exclusion of

Deleted: ¶

3. Results

3.1 Impact of the convective adjustment

The EVD parameterization exerts a strong control on the vertical structure of mixing. Figure 3 shows the eddy diffusivity coefficient along a zonal section at 37.5° N (as indicated in Fig. 1a) averaged over January 2021 for different mixing configurations. The presence of convective adjustment together with different mixing schemes (Figs 3a, 3c, 3e) tends to mask the differences among the various vertical mixing parameterisations resulting in a very similar eddy diffusivity pattern regardless of the chosen vertical mixing scheme. Given the different eddy diffusivity scales observed in the experiments with and without EVD (Figs 3a, 3c, 3e versus Figs. 3b, 3d, 3f), it is clear that when EVD is active, the eddy diffusivity produced by the vertical mixing closure is largely overwritten by the EVD parameterization. In contrast, when convective adjustment is disabled (Figs. 3b, 3d, 3f), the vertical eddy diffusivity is determined solely by the mixing schemes and reveals more pronounced differences among schemes. Notably, the Richardson-dependent scheme (Figs. 3b) produces a relatively homogeneous increase in eddy diffusivity within the mixed layer, whereas the TKE (Figs. 3d) and GLS (Figs. 3f) schemes result in more heterogeneous structures. The magnitude of the vertical eddy diffusivity increases going from the Richardson-number dependent scheme (with maximum eddy diffusivity values on the order of 10^{-2} m²/s) to the TKE and then to the GLS schemes (with maximum eddy diffusivity values on the order of 10^{-1} m²/s). Moreover, when the EVD scheme is active, the maximum value of the vertical eddy diffusivity consistently occurs near the surface, while, when EVD is disabled, its maxima are typically found within the mid-depth of the mixed layer, which is more realistic due to the suppression of turbulent length at the top and the bottom of the mixed layer. This distinction underscores the role of EVD in favouring a surface-intensified mixing structure as opposed to the more symmetric and realistic profile observed in its absence, potentially impacting the MLD.

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2

Deleted: 2





Figure 3: Vertical eddy diffusivity coefficient along the zonal section at 37.5°N (black line in Fig. 1a) averaged over January 2021 for different model configurations: (a) Richardson-dependent mixing scheme with EVD activated, and (b) without EVD activated; (c) TKE closure scheme with EVD activated, and (d) without EVD activated; (e) GLS scheme with EVD activated, and (f) without EVD activated. The black contours in each panel denote the depth of the mixed layer, while grey shading indicates land. Note the different eddy diffusivity scales for the experiments with and without EVD.



Deleted:

Deleted: out




Figure 4: Map showing the maximum depth reached by the EVD parameterisation when activated in the control experiment (Exp-PP⁺ in Table 1) during a summer day (1 July 2021). White areas indicate regions where EVD is inactive.

The EVD parameterisation is primarily intended to represent winter convection. However, convection-like vertical mixing occurs year-round, even outside the main winter season, and therefore the EVD parameterisation is active even in summer in some regions of the Mediterranean Sea. Figure 4 illustrates this by showing the maximum depth reached by the EVD on a summer day (1 July 2021) in the simulation using the Richardson-number dependent parameterization. White areas indicate the few regions where EVD is not applied on that day. Elsewhere the EVD is applied down to depths ranging from a few meters to several tens of meters reaching its maximum depth in the eastern Mediterranean Sea, and notably in the semi-permanent Ierapetra eddy region, located southeast of Crete within the Rhodes Gyre region (as defined in Fig 1a). This activity is typically weaker and shallower than winter convection, but it still contributes to reshaping the vertical dynamics, as discussed in the following sections.

3.2 Assessment at the basin scale

3.2.1 Impact on tracers

Among the different experiments (Table 1), the simulation using the Richardson-number dependent vertical mixing scheme combined with a convective adjustment (Exp-PP⁺) is taken as the control run. This configuration for the vertical mixing and

convection corresponds to the current setup of the Mediterranean Forecasting System of the Copernicus Marine Service (Clementi et al., 2023).

To evaluate the performance of our control simulation and identify systematic deficiencies of the model, we compare the model outputs in terms of temperature and salinity with all available Argo data in 2020-2021 (Fig. 1a). We verified (not shown) that removing a spin-up period leads only to negligible differences and does not affect the results. Following the extensive literature on error decomposition for model validation (e.g., Willmott, 1981; Murphy, 1988; Murphy, 1992; Oke et al., 2002; Oddo et al., 2022), we define four key statistical metrics, which provide complementary insights into model skills: the root mean squared error (RMSE), the unbiased root mean squared error (uRMSE), the mean bias (MB), and the cross-correlation coefficient (CC). For more details on their definition, see Appendix C. We compute these four metrics averaging on the entire basin and over the entire two-year period.

The temperature error profiles are shown in Fig. 5 as blue profiles. All metrics exhibit two pronounced decreases in model skill, one in the upper 50 m within the mixed layer and another between 200 and 600 m, corresponding to the Levantine Intermediate Waters (LIW) depth range (e.g., Bryden and Stommel:1982; Robinson and Golnaraghi, 1993; Iascaratos et al., 1993; Taillandier et al., 2022). Within the mixed layer, RMSE and uRMSE peak at 25–30 m, corresponding to these two-year average MLD (30.8 m for observations and 28.3 m in the control simulation – dashed horizontal lines in Fig.5). MB reaches its maximum around the middle of the MLD (about 15 m). CC exhibits a monotonic reduced decrease throughout the mixed layer, with a minimum below the yearly average MLD. A similar behaviour occurs within the LIW depth range, with a minimum in CC around 400 m, while RMSE, uRMSE and MB show their largest deviations slightly above, at approximately 350 m. Overall, this evidence indicates that errors are largest in the surface mixed layer but also reflect difficulties in representing intermediate water masses.

The salinity error profiles (orange) in Fig. 5 show a distinct pattern compared to temperature. RMSE and uRMSE exhibit a single peak confined to the mixed layer, with no significant anomalies at the LIW depth range. Nevertheless, the MB highlights systematic fresh biases at both depth ranges, suggesting persistent model errors also in salinity in representing the LIW. We observe that the minimum salinity MB within the LIW depth range is found near the bottom of the LIW layer (about 500 m), while the minimum temperature MB occurs approximately at mid-depth within the layer (about 350 m). As for temperature, the CC decreases monotonically within the mixed layer; however, unlike temperature, it exhibits only a weak secondary minimum at LIW depths. These findings suggest that salinity shows smaller discrepancies than temperature with respect to observations in the representation of intermediate waters, while it also presents a discrepancy with the observation within the mixed layer.

Deleted: 4

Deleted: Theocharis et al., 2002

Deleted: yearly

Deleted: just

Deleted: in

Deleted: Like

Deleted:

Deleted: and

Deleted: shows

Deleted: is overall less affected

As stated above, overall, for both salinity and temperature, the shape of the error at the basin scale suggests two main decreases of the model skills: the representation of the properties within the mixed layer and the intermediate water masses. The control simulation performs reasonably well in the deep region, while the more pronounced biases are within the mixed layer. These biases are likely related to the vertical mixing scheme, highlighting the need for a more realistic treatment of atmospheric energy input and vertical redistribution of momentum and buoyancy.





Figure 5: Basin-averaged temperature (blue) and salinity (orange) error profiles for 2020–2021 in the control simulation (Exp-PP+ in Table 1). Panels from left to right: root mean squared error (RMSE), unbiased root mean squared error (uRMSE), mean bias (MB), and cross-correlation coefficient (CC). The definitions of these metrics are in Appendix C. Dashed lines represent the two-year basin average values of the MLD for observations and control simulation.

To assess the impact of the different closure schemes and the activation of the convective adjustment at the basin scale, we compare the metrics obtained for the other five simulations (simulation other than Exp-PP+ in Tab. 1) with these of the control



run. Figures 6 and 7 show the basin-averaged error difference for temperature and salinity, respectively. For each experiment and each metric (Equations C1–C4), the error difference (err_{diff}) as a function of depth is computed as

$$err_{diff} = err_{mod}(z) - err_{cs}(z) \quad (1)$$

where $err_{cs}(z)$ denotes the error of the control run (Exp-PP⁺, see Figs. 5), and $err_{mod}(z)$ is the error associated with each model configuration (experiments other than Exp-PP⁺ in Tab. 1). Here, err generically denotes one of the four metrics (RMSE, uRMSE, MB, or CC) detailed in Appendix C. For the MB_{diff} metric, the absolute value of the mean bias is used for both the control run and the model experiments when computing MB_{diff}.

It should be noted that negative values of RMSE_{diff}, uRMSE_{diff}, and MB_{diff} indicate improvement, while positive values indicate deterioration. For CC_{diff}, the interpretation is reversed: higher values (closer to 1) denote improvement, whereas lower values (towards 0) denote deterioration. Accordingly, in Figs. 6 and 7 green and red backgrounds highlight regions of improvement and deterioration relative to the control simulation, respectively.

For temperature (Fig. 6), the error profiles indicate that the GLS closure scheme without convection parameterization (solid green line) provides the largest error reduction, especially in terms of MB_{diff}, a result further confirmed by the CC_{diff}. The greatest improvement occurs within the mixed layer, mostly reflecting a reduction in systematic errors (MB_{diff}), while additional improvements at LIW depths are observed in terms of uRMSE_{diff} and CC_{diff}. Adding the convection parameterization to GLS (dashed green line) diminishes these gains at some depths, and mostly for MB_{diff}. The TKE mixing scheme exhibits intermediate skill: with or without EVD (orange lines), improvements are mostly confined to the mixed layer, while performance below this layer often deteriorates. The Richardson-dependent scheme without EVD (solid blue line) shows the smallest improvement within the mixed layer and a worsening in the LIW layer across all metrics. This behaviour is expected because the scheme estimates turbulence solely from the Richardson number (see Subsection 2.2.1 and Appendix A), that is from the balance between vertical shear and stratification (buoyancy) and does not include convective mixing. Consequently, it is expected that an additional parameterization would be required to properly represent convection.

For salinity (Fig. 7) in the absence of convective adjustment (solid lines), GLS is the only scheme showing improvements above 100 m for RMSE_{diff}, uRMSE_{diff}, CC_{diff}, and MB_{diff}. At the LIW depths, changes are minimal and a small degradation occurs for all experiments between about 100 and 200 m. The TKE and Richardson-number dependent schemes without EVD show smaller improvement in all metrics, particularly within the mixed layer, with the Richardson-number dependent scheme (solid blue line) performing worst, especially for CC_{diff}.

Deleted: 4 and

Deleted: err_{diff}

Deleted: RMSE

Deleted: s

Deleted: across all depths

Deleted: the

Deleted: largest

Deleted: degradation

Deleted: ,

Deleted: consistent with the expectation that a closure scheme based solely on vertical shear and stratification requires additional parameterizations to represent

Deleted: properly

Deleted: convection.

Deleted: at some depths in

Deleted: and

Deleted: the smallest degradation in

Deleted: within the mixed layer

Deleted: except for the configuration with GLS and EVD deactivated, which slightly improves RMSE, uRMSE and CC

Deleted: deterioration across

Deleted: MB





Figure 6: Temperature error difference (equation 1) with respect to the control simulation for the four metrics in equations C1–C4. Dashed lines indicate simulations with EVD and continuous lines without EVD. The transparent background color highlights regions of improvement (green) and worsening (red) compared to the control simulation.







Figure 7: Salinity error difference (equation 1) with respect to the control simulation for the four metrics in equations C1–C4. Dashed lines indicate simulations with EVD and continuous lines without EVD. The transparent background colour highlights regions of improvement (green) and worsening (red) compared to the control simulation.



Deleted:

3.2.2 Representation of the Mixed Layer Depth

Given that the largest impact of the interplay between the mixing closure parameterisation and the convective adjustment occurs within the mixed layer, we also investigate how well the different schemes reproduce the MLD across the Mediterranean Sea. Figure 8 shows the time evolution of the MLD for the years 2020–2021, as computed from all available Argo temperature and salinity observations (black dashed line) and from model outputs (coloured lines) at the corresponding locations. Notably, for each day, model temperature and salinity profiles are first sampled at each Argo location. Density and MLD are then computed individually for each profile in both models and observations, and a basin average MLD is finally calculated.

650 The MLD for both observations and model outputs is computed using the density threshold method, which identifies the depth at which the potential density increases by a fixed amount relative to a near-surface value. Specifically, here, the MLD is defined as the depth where the potential density exceeds that at 10 m by 0.01 kg m^{-3} . With this definition, a minimum MLD of 10 m is imposed, as the reference density is taken at that depth. This approach, commonly used in oceanographic studies, provides a robust estimate of the mixed layer under both stratified and convective conditions (e.g., de Boyer Montégut et al., 2004).

655 The temporal evolution of the basin-averaged MLD over the period 2020–2021 oscillates around a temporal-spatial average value of 31 m and exhibits a pronounced seasonal cycle, with deepening during winter and shoaling in summer. On average, the MLD reaches its maximum around 70–80 m in winter (December to March), when surface cooling and intense wind forcing promote vertical convection and deep mixing. In spring, the increasing solar radiation and weakening winds lead to rapid surface warming and stratification, resulting in a marked shoaling of the MLD. By summer (June to September), the MLD stabilises at its shallowest levels, below 20 meters, reflecting strong thermal stratification and limited vertical mixing. In autumn, cooling and more frequent wind events gradually deepen the MLD again, preparing the system for winter convection.

665 All simulations, colour-coded according to model configuration, reproduce this seasonal pattern, with winter maxima exceeding 70 m and summer minima around 15 m. However, significant differences in the MLD emerge while employing different vertical mixing schemes and varies also significantly depending on whether the EVD convection parameterisation is activated or not.

670 Simulations including EVD (blue, orange, and green lines) behave in a similar way and very close to the observations (black line). They occasionally show deeper (e.g. February–March 2020) or shallower (e.g. March 2021) winter MLDs and overall, they exhibit a good agreement in summer. With EVD activated, the two-year average MLD with TKE (30.9 m) and GLS (31.3 m) are very close to the observed one (30.8 m), while the Richardson-number dependent scheme tends to underestimate the MLD especially in summer (average of 28.3 m).

675 Simulations without EVD (red, purple, and brown lines) show distinct behaviours. The Richardson-dependent scheme (red line) produces the shallowest winter MLD and a low average MLD (average of 25.7 m), significantly underestimating observed deep mixing even in summer. The TKE scheme (purple line) tends to underestimate the MLD, both in winter and in summer. Although it performs better than the Richardson-number dependent scheme yet still fails to capture the basin-average maximum depths seen in observations (average MLD of 25.7 m). On the other hand, the GLS scheme (brown line) yields winter and summer MLDs closely aligned with observations, remaining in good agreement to them throughout the year. With the GLS, the average value of MLD without convective adjustment (32.7 m) is very close to the one obtained with convective adjustment included (31.3 m).

Deleted: 31.1

Deleted: 5

Deleted: 1

Deleted: 0

Deleted: 4

Deleted: the

Deleted: est


Deleted: 5

Deleted: 1


Deleted: 1

Deleted: 5

695 The GLS scheme without convective adjustment reproduces the observed deepening of the mixed layer, while the TKE and
 the Richardson-dependent scheme do not. This evidence demonstrates that this scheme does not need an additional external
 convection parameterisation to reproduce realistic mixed layer deepening. This may be attributed to the two-equation structure
 of the GLS (see Section 2.2.1 and Appendix A): the first equation evolves the turbulent kinetic energy over time, indicating
 where mixing is possible, while the second equation dynamically adjusts the turbulence length scale, controlling how deeply
 mixing penetrates. In contrast, the TKE scheme lacks dynamic depth control, and the Richardson-dependent scheme does not
 700 even include prognostic turbulent kinetic energy; in both cases, the penetration of convective mixing is limited. It is worth
 noting that, in the absence of explicit convective adjustment, the additional physics included in the TKE scheme compared to
 the Richardson-dependent formulation appears to have only a minor impact, likely because, at these scales, TKE still provides
 a relatively simplified representation of convective processes.



705 **Figure 8:** Time evolution of basin-averaged MLD for varying vertical mixing schemes, with and without convection parameterisation (configurations listed in Table 1). The black dashed line represents the MLD from Argo observations. A 7-day moving average has been applied to both model outputs and observations.



Deleted:

3.3 Assessment of water mass formation

The Mediterranean Sea hosts several key sites of deep and intermediate water formation, including the Gulf of Lion, the South Adriatic Sea, the Rhodes Gyre area and the Aegean Sea (e.g., Lascaratos et al., 1999; Simoncelli et al., 2018; Pinardi et al., 2019; Pinardi and Masetti, 2000; Pinardi et al., 2023). In these regions, vertical mixing and convection play a fundamental role in preconditioning and driving dense water formation processes by contributing to surface buoyancy loss and erosion of stratification (e.g., MEDOC group, 1970; Schroeder et al., 2012). The intensity and vertical structure of turbulent mixing directly influence the transformation and ventilation of water masses, making its accurate representation essential for modelling the basin-scale thermohaline circulation (e.g., Simoncelli et al., 2018; Pinardi et al., 2019; Somot et al., 2006).

Deleted: Send et al., 1996;

Deleted: Pinardi et al., 2019;

Deleted: Testor and Mortier, 2005

Deleted: ; Simoncelli et al., 2022

As case studies, to further investigate the impact of the interplay between different vertical mixing schemes and convective adjustment, we select one site of DWF (the South Adriatic Pit) and one site of both deep and intermediate water formation (the Rhodes Gyre region) and analyse them across the various model configurations during two selected time windows in 2021. We note that after more than one year of integration, although all six experiments are initialized from the same initial conditions, the six simulations have already had time to substantially diverge by the beginning of 2020. Consequently, while this section focuses on the behaviour of the schemes within two selected time windows in 2021, the results inherently reflect the prior evolution and memory of the system under each parameterization. This approach allows us to assess the resulting quasi-equilibrium state, accounting for how each scheme steers the model toward a distinct numerical and physical regime through its cumulative effects over time and its interplay with the convective adjustment parameterization. Therefore, our objective in this section is not to evaluate turbulence closure schemes in isolation, but rather to investigate their integrated impact in deep and intermediate water formation regions, within the general circulation of the Mediterranean Sea.

Deleted: deep water formation

Deleted: .

3.3.1 Deep water formation and intermediate water intrusion in the south Adriatic Pit

The South Adriatic Pit (defined as in Simoncelli and Pinardi (2018) between longitudes 17°–19°E and latitudes 41°–42.5°N and shown in Fig. 1a) reaches a depth of 1200 m and is connected to the Eastern Mediterranean Basin through the Strait of Otranto. It is one of the principal sites of DWF in the Mediterranean Sea, where intense cooling and evaporation during winter – especially in February–March – induce strong density increases that drive dense water sinking and convective overturning (e.g., Wüst, 1961; Gačić et al., 2001; Manca et al., 2003). During these events, the MLD can deepen substantially, typically reaching between 600 and 800 m (e.g., Manca et al. 2003; Pinardi et al., 2023).

Deleted: deep water formation

Deleted: Lascaratos et al., 1999

Deleted: ; Roether et al., 1996

Deleted: Lascaratos et al., 1999

Deleted: Roether et al., 1996;





Figure 9: (a) Temperature and (b) salinity model profiles compared to observations (black dashed line) from an Argo float on 25 February 2021. Modelled profiles are color-coded by different vertical mixing schemes, with and without convective adjustment as indicated in the legend. The inset in the left panel shows the position of the Argo float.

In Fig. 9 we show the temperature and salinity profiles as recorded by an Argo float (black dashed line) on 25 February 2021 in comparison with the modelled profiles at the same location for varying vertical mixing schemes and activating (dashed line) or deactivating (continuous line) the EVD. The location of the profile is shown in the insert of Fig. 9a. Argo observations show a nearly uniform layer in the upper ~500 m, followed by a sharp gradient below. This inflection point corresponds to the MLD. Among the models in which EVD is active, the simplest Richardson-dependent scheme performs best, although it underestimates the MLD in the salinity profile of about 50 m. In this case, the GLS scheme underestimates the MLD in both temperature and salinity by about 100 m. In the absence of EVD, we observe a similar behaviour across different mixing scheme, with a broad inflection point corresponding to MLD around 500 m. To investigate how the various configurations of mixing and convection lead to the situation shown in Fig. 9, we examine the time evolution of the salinity field at this location up to the day of the profiles in Fig. 9. Salinity was chosen because it largely reflects the imprint of deep and intermediate water (Fig. 1d).



Deleted:

Deleted: (location shown in the insert of Fig. 9a)

Deleted: data

Deleted: , the situation is reversed. While the Richardson-dependent scheme underestimates the MLD and shows a reversed salinity gradient within the mixed layer, the GLS scheme captures the MLD depth well in both temperature and salinity, indicating that the GLS scheme performs better on its own, without convective adjustment. The TKE scheme, both with and without EVD, underestimates the MLD by about 150 m and shows a reversed salinity gradient within the mixed layer.


In Fig. 10, we show the Hovmöller diagrams of salinity evolution at the site described above from 1 to 25 February 2021. The last day of each Hovmöller corresponds to the day of the profile shown in Fig. 9. The three panels on the left (Fig. 10a, 10c, 10e) show the salinity evolution for the models with EVD activated. The three panels on the right (Fig. 10b, 10d, 10f) correspond to the models with EVD deactivated. In the upper 100 m, the models with EVD activated show similar behaviour, with fresh water persisting until 13 February, whereas without EVD the models show large variability due to a different activation of mixing. In particular, the TKE scheme (Fig. 10d) maintains relatively fresh water up to 25 February, while the Richardson-number dependent scheme (Fig. 10b) and the GLS scheme (Fig. 10f) show increased salinity. However, the most notable differences occur at depth, between approximately 100 and 400 m, where a tongue of salty LIW gradually intrudes and a convective event, initially confined to the surface after 13 February, gradually deepens into intermediate layers. The contrast between simulations with and without EVD is substantial.

In the presence of EVD, convection triggered after 13 February erodes the intruding LIW, so that the LIW is completely mixed thereafter. As shown in Sections 2.2.2 and 3.1, when the convective adjustment is applied, the vertical eddy diffusivity resulting from the mixing schemes, is abruptly replaced with a high constant value, disregarding the temporal evolution of the tracers. This is evident after 13 February regardless of the mixing scheme (Fig. 10a, 10c, 10e), with the salinity field homogenized across both time and depth. Consequently, convection parameterized through EVD does not account for prior stratification: the method is applied uniformly, effectively erasing pre-existing or intruding structure in the region.

Conversely, in the cases without EVD, the LIW persists up to 25 February. In particular, in the GLS scheme (Fig. 10f) the LIW gradually sinks to about 400 m due to its higher density (similarly to what shown in Fig. 1d), exhibiting a more realistic behaviour, whereas in the Richardson-number-dependent scheme (Fig. 10d) it drops abruptly, and in the TKE scheme (Fig. 10b) it remains at roughly the same depth. This behaviour is consistent with the formulation of the mixing schemes. With the GLS scheme (Fig. 10f), which includes a prognostic equation for the mixing length (Section 2.2.1 and Appendix A), both the intensity and depth of convective events appear to be more realistically represented. On the other hand, with the TKE scheme (Fig. 10d), the convective intensity is similar, but the depth of the event is underestimated. For example, on 21 February, the salinity intrusion extends below 300 m with GLS, whereas with TKE the intrusion already begins at 200 m. In the Richardson-dependent formulation (Fig. 10b), the convective event is weaker and more abrupt, producing a less gradual penetration into deeper layers.

Formatted: Centred, Keep with next

Formatted: Caption



Deleted:
Figure 10: Daily water mass formation rate in the south Adriatic Pit. Curves are colour-coded by different vertical mixing schemes, with and without convective adjustment as indicated in the legend.

To monitor the deep convection process in the South Adriatic Pit region we compute the water mass formation rate, following the criteria defined in Von Schuckmann et al. (2018), applied in Simoncelli and Pinardi (2018), and operationally computed yearly as Ocean Monitoring Indicator based on the Mediterranean Reanalysis system delivered within the Copernicus Marine Service. The water mass formation rate is defined as the volume of water above the mixed-layer depth with potential density equal to or greater than 29.10 kg m^{-3} , divided by the number of seconds per year. Since this Ocean Monitoring Indicator is computed daily and very few observations per day are available in this small region, we calculate it using only modelled data for intercomparison. In Fig. 10 we show the water mass formation rate as computed for the six simulations with varying vertical mixing schemes and activation/deactivation of ... [2]

Deleted: .

Deleted: is where the

Deleted: circulates

Deleted: Levantine Intermediate Water forms

Deleted: subsequently




Figure 10: Hovmöller diagrams showing the temporal evolution of salinity at the location indicated in Fig 9a (inset) from 1 to 25 February 2021 for different model configurations: (a) Richardson-dependent mixing scheme with EVD activated, and (b) without EVD activated; (c) TKE closure scheme with EVD activated, and (d) without EVD activated; (e) GLS scheme with EVD activated, and (f) without EVD activated. The last day in each panel corresponds to the day of the profile shown in Fig 9.

3.3.2 Deep and intermediate water formation in the Rhodes Gyre region

As an additional case study, we investigate the capability of the model to reproduce deep and intermediate water in the Rhodes Gyre region, defined here between longitudes 26.3° – 31° E and latitudes 33° – 36.1° N and shown in Fig. 1a. This region, located in the eastern Mediterranean Sea between Rhodes and Cyprus and extending southward into the open Levantine Sea, hosts the Rhodes Gyre and is the key site for the formation and initial spreading of the LIW. Relative to the domain defined by Simoncelli and Pinardi (2018), here the region is expanded by 1° to the east and 1° to the south in order to better resolve the early spreading of the LIW. This water mass, characterised by high salinity and intermediate depth (approximately 200–500 m), begins

spreading, within the defined region and subsequently extends further across the Mediterranean Sea, influencing basin-scale thermohaline circulation (e.g., Lascaratos et al. 1993; Kubin et al., 2019; Pinardi et al., 2023).

Deleted: s

In Fig. 11a, salinity observations from all available Argo floats in the Rhodes Gyre region between January and April 2021 are compiled and arranged to form a Hövmoller diagram. Water with salinity exceeding 39.2 psu is observed above 200 m until approximately mid-February, when a convective event forces the high-salinity layer downward, resulting in a deepening of the entire water column. The LIW, with salinities ranging from 38.95 to 39.05 psu (Lascaratos et al., 1993), descends from approximately 300 m to depth below 400 m.

Deleted: 11

Deleted: we show the Hovmöller diagram of

Deleted: data

Deleted: from all available Argo floats in the Rhodes Gyre region (defined as in Simoncelli and Pinardi (2018) between longitudes 26.3°–30°E and latitudes 34°–36.1°N)

Deleted: In Figs. 11b–11g we present the Hovmöller diagrams of MB (equation C3) of each model configuration with respect to the observations.

In Figs. 11b–11g we present Hovmöller diagrams of salinity for each model configuration. Overall, the model reproduces the observed stratification well and generally reach similar maximum salinity values, with some differences for varying mixing scheme and depending on the presence or absence of convective adjustment. In the upper 100 m, prior to the mid-February convective event, all models tend to overestimate salinity relative to observations, whereas after the event they generally produce fresher waters. At greater depths, model layers are generally narrower and more confined than in the observations, where post-convective mixing causes substantial vertical widening of the layer.

Deleted: All models show the largest errors between approximately 200 and 500 m, corresponding to the depth range of the LIW. We note that this error, originating in this region, spreads over time across the entire basin as shown in Section 3.2 and is clearly visible in the basin-averaged profiles in Figs. 4 and 5. Reducing this source of error in this region is therefore important for the entire Mediterranean Sea. In the presence of convective adjustment, among the configurations (Figs. 11b, 11d and 11f), the GLS closure scheme exhibits the largest errors, whereas without convective adjustment (Figs. 11c, 11e and 11g) the situation is reversed. While the Richardson-number dependent and the TKE schemes perform worse with EVD deactivated, the GLS scheme shows a significant improvement (average MB moves from 0.022 psu to 0.004 psu) demonstrating that while employing physics-based vertical mixing scheme the addition of external convective parameterization can deteriorate model results in this key region.

When EVD is applied (Figs. 11b, 11d, 11f), the convective event is weakly reproduced. The downward displacement of the water column is most evident in the TKE scheme (Fig. 11d), producing a deepening of approximately 100 m, whereas in the Richardson-dependent scheme (Fig. 11b) and GLS scheme (Fig. 11f), the column exhibits almost no abrupt shift, showing only gradual deepening over time. Overall, EVD suppresses the differences among the mixing schemes and results in a very similar salinity response to convection.

In the absence of EVD (Fig. 11c, 11e, 11g), the salinity response differs. With Richardson-dependent scheme (Fig. 11c), the behaviour is very similar to the case with EVD (Fig. 11b), with progressive deepening of the column without a sudden convective shift. This evidence reflects the fact that the Richardson-dependent scheme does not explicitly model convection but only increases mixing when shear dominates over buoyancy (Section 2.2.1). Consequently, convective events are weakly represented: the water column deepens gradually, and high-salinity layers remain confined and less vertically spread compared to observations or schemes that explicitly account for convection. By contrast, the TKE (Fig. 11e) and the GLS (Fig. 11g) schemes alone capture the mid-February convective event, producing a rapid downward displacement of the entire water column, with the GLS scheme better representing the intensity and extend of deepening due to its prognostic treatment of turbulent kinetic energy while simultaneously estimating a turbulence length scale.

930





Figure 11: (a) Hovmöller diagram of salinity observations from available Argo floats in the Rhodes Gyre region (Fig. 1a) between January and April 2021. (b–g) Hovmöller diagrams of salinity for different vertical mixing schemes and convective adjustment settings: (b) Richardson-dependent mixing scheme with EVD activated, and (c) without EVD activated; (d) TKE closure scheme with EVD activated, and (e) without EVD activated; (f) GLS scheme with EVD activated, and (g) without EVD activated. All panels share the colour scale in panel (a).



3.4 Assessment of the nighttime summer convection daily cycle

In summer, the upper layers of the ocean exhibit a pronounced diurnal cycle in temperature, primarily controlled by the alternation of solar heating during the day and cooling processes at night. Surface waters warm under strong solar radiation, leading to a shallow, well-defined thermal stratification, while nocturnal cooling can erode this stratification. These processes occur within the mixed layer, which remains relatively shallow, as the rapid diurnal cycle of formation and destructing does not allow it sufficient time to deepen and night cooling is not as strong as in winter. Consequently, these processes are expected to be strongly influenced by the selected mixing scheme and the presence of a convective adjustment, which can significantly modify both the intensity of the heating and the depth of the near-surface warm layer. As shown in Fig. 4, the EVD during a summer day remains typically around an average of a few meters, but in certain areas – such as the Ierapetra semi-permanent gyre area within the Rhodes Gyre region as defined in the previous section – it can reach tens of meters.

Deleted: 3

The simulations in Tab. 1 were performed again starting from 1 July 2021 for three days, producing hourly outputs. The same initial conditions were used for all simulations, with the goal of monitoring how the mixing closure scheme and the presence of convective adjustment influence the daily cycle. As a case study, we focus on a site (27.5°E, 34.4°N) within the Rhodes Gyre region where the EVD reaches its greatest activation depth during summer (Fig. 4).

Deleted: within the Rhodes Gyre region

Deleted: 3

Figure 12 shows Hovmöller diagrams of the eddy diffusivity coefficient at that location in the upper 50 m for the six simulations. The upper panels correspond to the three simulations with EVD activated, whereas the lower panels show the cases without EVD. In general summer convective events tend to occur during nighttime because of surface cooling associated with absence of solar heating and extend along the water column within the upper few tens of meters (~30 m in Fig. 12). As already observed in Fig. 3, when EVD is activated, the eddy diffusivity coefficient produced by the vertical mixing scheme is largely overwritten by the EVD parameterization (note the three-order-of-magnitude difference between the upper and lower panels in Fig. 12), resulting in similar mixing patterns regardless of the turbulence closure scheme used.

In contrast, when EVD is not applied (bottom panels in Fig.12), the representation of convection differs substantially among the three closure schemes. The Richardson-number dependent scheme produces weak but nearly persistent convective activity that remains present even during daytime. The TKE and GLS schemes instead exhibit a more intermittent pattern of convection, with events occurring less frequently and the strongest one during nighttime. Convective events simulated with the TKE scheme are generally weaker than these produced by the GLS scheme and tend to extend less deeply in the water column.

For example, during the night of 1 July 2021, a convective event reaching approximately 25 m depth occurs in all simulations. In three cases with EVD (upper panels in Fig. 12), this event is represented very similarly and homogeneously throughout the upper 25 m, whereas without EVD (lower panel in Fig. 12) its amplitude and vertical structure differ substantially among the

three closure schemes. In the Richardson-number dependent case, it appears as a relatively weak and vertically homogeneous event down to 25 m. In the TKE simulation, it is confined between a bout 10 and 20 m, with a peak around 15 m. In contrast, the GLS scheme produces a much stronger event, extending all the way down to 25 m.




Figure 12: Hovmöller diagrams of the hourly eddy diffusivity coefficient between 1 and 3 July 2021 (upper 50 m) for the six simulations with (a) Richardson-number dependent scheme with EVD, (b) TKE with EVD, (c) GLS with EVD, (d) Richardson-number dependent scheme without EVD, (e) TKE without EVD, (f) GLS without EVD.

These differences in the representation of summer convection, as seen in Fig. 12, both in terms of amplitude and vertical distribution along the water column, have a direct impact on the temperature field. In particular, the variability in convective activity simulated by the different turbulence closure schemes, and the homogenizing effect of EVD, translates into notable differences in the upper-ocean temperature structure. Figure 13 shows the temperature profiles at that site during the same three days and at different times of the day.

The first profiles (lightest colours) show that at the beginning – on 1 July 2021, between midnight and 1 a.m. – the temperature profiles are well mixed and very similar to each other, with differences that are barely distinguishable, and are not yet influenced by the vertical mixing schemes or convective adjustment. Later the same day – between 6 p.m. and 7 p.m. – we observe that, while the simulations with TKE (orange) and GLS (green) behave in a similar way both with and without convective adjustment, the Richardson-number-dependent scheme (blue) is strongly influenced by the presence or the absence of EVD. In fact, in the case of TKE and GLS, the simulations without EVD (solid lines) produce temperature profiles that are

Formatted: Keep with next

Formatted: No underline

Formatted: Caption

Deleted:

Deleted: 2

Deleted: for

Deleted: days and

Deleted: ¶

Deleted: ¶

only slightly different than those with EVD (dashed lines). By contrast, the Richardson-number dependent scheme without EVD (solid blue line) yields a nearly uniform temperature in the first 5 m, followed by a gradual decrease with depth. With EVD (dashed blue line) the profile is reversed, showing a surface and deep gradient separated by an intermediate quasi-homogeneous layer.

In the following two days (2 and 3 July 2021, both at 6–7 p.m.) the difference between the Richardson-number dependent scheme and the other two schemes increases in both profile shape and amplitude, reaching a sea surface temperature (SST) difference larger than 0.5 °C on 3 July. Moreover, the simulations with TKE and GLS exhibit only minor differences – on the order of ~0.1 °C around 15 m depth – arising from slightly different convective patterns in terms of amplitude and vertical extend (Figs. 12e and 12f). In contrast, the Richardson-number dependent scheme shows much larger discrepancies, especially near the surface, due to the substantially different convective patterns (Fig. 12d).





Figure 13: Temperature profiles at a site (27.5°E, 34.4°N) in the Rhodes Gyre region at different times of the day (see legend in the inset) and for different closure schemes and convective adjustment settings (see bottom legend).

Deleted: while
Deleted: both
Deleted: show
Deleted: with and without EVD, those with
Deleted: display



Deleted:
Deleted: 12
Deleted: in the Rhodes Gyre area

4. Discussion and conclusions

Physics-based turbulence closure schemes can represent convective mixing in principle, but at typical model resolution they may underestimate the rate of instability removal. For this reason, convective adjustment is often applied as a numerical parameterization to rapidly stabilize the water column and mimic the fast overturning associated with unresolved convective events. In this work, we investigated how the combined effect of three vertical mixing schemes – Richardson-number dependent, TKE, and GLS closure schemes – and an external convective adjustment (EVD) affects the performance of a regional ocean circulation model of the Mediterranean Sea. We analysed the model outputs in relation to the physics of vertical mixing represented by the different schemes and assessed the advantages and drawbacks of applying an external convective parameterization alongside each of them.

The results first confirm what is known in the literature: the explicit representation of convection through enhanced vertical diffusion is crucial for simple schemes such as Richardson-number dependent closures, which lack an explicit formulation of convection and only reduce density gradient by enhancing vertical mixing in unstable layers. On the other hand, our results indicate that for more physically based schemes, the inclusion of a convective adjustment should be handled with care, as its application can be redundant or even degrade model performance. For the TKE scheme, we found that the inclusion of EVD can be beneficial in some cases. The TKE scheme has a single prognostic equation for turbulent kinetic energy, while the turbulent mixing length is defined diagnostically, and treats convection prognostically through the calculation of the turbulent kinetic energy; as a result, it often represents the intensity of convection well, but underestimates the penetration depth of convective events. In contrast, the GLS scheme, which treats convection prognostically through the calculation of both enhanced turbulent kinetic energy and turbulent mixing length, is generally better able to simulate both the penetration depth and the intensity of convective processes on its own. Therefore, with the GLS scheme, the convective adjustment is redundant and can often degrade the model's skill. These observations are confirmed both at basin scale and in deep and intermediate water formation regions.

At the basin scale, statistical metrics on temperature and salinity show that the GLS scheme performs best, followed by TKE and the Richardson-number dependent scheme. The inclusion of EVD makes little difference overall. In particular, we observed a more consistent improvement with GLS without EVD across all metrics and over a larger range of depths. The largest impact of including or excluding EVD at the basin scale is observed on the MLD. For both the Richardson-number dependent scheme and the TKE schemes, only the addition of the convective adjustment allows the model to reproduce the observed temporal evolution of the average MLD across the basin. This is because EVD compensates for mixing physical processes in these schemes: either the full convective overturning in the Richardson-number dependent scheme or the prognostic evaluation of the mixing length in TKE, which controls the convection penetration depth. In contrast, the GLS

- Deleted: interplay
- Deleted: between
- Deleted: different
- Deleted: scheme
- Deleted: closure scheme

- Deleted: – such as the GLS closure scheme – which account for buoyancy loss in their equations,
- Deleted: may become
- Deleted: or even degrade
- Deleted: .
- Deleted:
- Deleted: our results show that the addition of a convective adjustment plays a different role depending on the mixing scheme
- Deleted: .

scheme, thanks to its prognostic treatment of both TKE and mixing length, is able to reproduce the observed basin-average MLD over time, including the MLD seasonal variability, without requiring an external convective parameterization.

In DWF areas, such as the south Adriatic Pit, where convective processes play a central role during dense water formation events, we observe that both the choice of mixing scheme and the inclusion of the convective adjustment play a major role in the development of deep water and the intrusion of intermediate water from the Levantine basin. Notably, the inclusion of EVD, with its abrupt and strong increase in vertical eddy diffusivity when unstable layers are present, does not account for previous states or the coherent temporal evolution of the water column: when applied, EVD disrupts pre-existing or intruding structures over time. The different mixing schemes without an external convective adjustment reproduce convection in distinct ways, and a similar behaviour is observed in the Rhodes Gyre region. The Richardson-number dependent scheme produces an abrupt but weak homogenization of the layers affected by surface convection, resulting in less gradual penetration into deeper layers and a partial removal of the intruding LIW. The TKE scheme reproduces convection with a smoother intensity transition compared to the Richardson-number dependent scheme, showing less abrupt effects on salinity. However, the penetration depth in the TKE scheme remains limited, without significant sinking of denser waters, due to the lack of a prognostic control on the penetration depth. In contrast, the GLS scheme more accurately represents both the intensity and penetration depth of convective events, allowing for a smooth sinking of the LIW.

Focusing on nighttime convection during summer, analysed at an hourly timescale, we observe how the nighttime convective adjustment, combined with different vertical mixing schemes, shapes the diurnal evolution of temperature. The three vertical mixing schemes represent summer convection differently: the TKE and GLS schemes largely capture the night-and-day differentiation, with GLS producing homogeneous, strong mixing that penetrates deeper. In contrast, the Richardson-number dependent scheme increases mixing weakly but almost continuously, even during daytime showing almost no day-night contrast. Temperature profiles using the Richardson-number dependent scheme differ noticeably from the other closure schemes, and after three days, the SST reaches a value roughly 0.5 °C higher.

In summary, across all temporal and spatial scales, our results highlight the crucial role of two key features in vertical mixing schemes. First, prognostic treatment of turbulent kinetic energy (as in the TKE and GLS schemes) is essential to accurately reproduce the intensity of overturning convection, retain memory of priori processes and ensure smooth transitions at a site, preserve pre-existing structures as well as the day-night alternation during summer convection. Second, a prognostic additional equation for the mixing length (as in the GLS scheme), rather than a fixed value as in TKE, is necessary to correctly represent the penetration depth of convective processes and to achieve the most accurate representation of the mixed layer depth. The inclusion of a convective adjustment to compensate for the lack of these features only partially addresses the deficiencies, as external convective adjustment are not embedded within the general framework and do not evolve prognostically.

Deleted: For the Richardson-number dependent scheme, EVD is crucial to accurately represent temperature, salinity and mixed layer depth. For the TKE closure scheme, it significantly improves the simulation of salinity and mixed layer depth, while for the GLS closure scheme, it provides some improvement for salinity but has little to no impact on temperature and mixed layer depth. Notably, the GLS scheme alone is able to reproduce the basin-scale seasonal variability of the MLD in good agreement with Argo observations, unlike the TKE scheme.

Deleted: deep water formation

Deleted: a

Deleted: role

Deleted: the water mass formation rate is affected by the presence of EVD mainly when the mixing scheme is the simple Richardson-number dependent one. With TKE and GLS, a comparable amount of dense water is formed during the DWF season. Moreover, the MLD captured by Argo during DWF events is well reproduced only by the GLS without EVD, whereas its addition causes the MLD to be too shallow compared to observations. In areas of LIW formation, such as the Rhodes Gyre region, the representation of dense and salty waters improves significantly when using the GLS scheme without any external convective parameterization.

Deleted: interacts

Deleted: to

Deleted: The EVD modifies the daily cycle and, consequently, the daytime temperature profiles mainly when using the Richardson-number dependent scheme; for the TKE and GLS schemes, it produces little change. Furthermore,

Deleted: t

1155 These findings underscore how critical it is, in regional ocean modelling, to align appropriately the mixing closure and convection parameterisations. More broadly, these results highlight that careful choice of a combination of subgrid-scale parameterizations is essential to reliably simulate key oceanographic features and improve model interpretability in regional applications. Future work could extend this analysis to longer-term simulations and the effect of additional subgrid-scale processes – such as internal waves, eddy formation, and horizontal mixing – in order to further improve model realism and predictive skill. More broadly, our results highlight how the representation of vertical mixing can strongly influence deep and intermediate water formation and properties, with potential implications for regional ocean modelling in other basins with characteristics similar to the Mediterranean Sea. Comparable processes occur in other semi-enclosed or marginal basins, where dense water formation and vertical mixing play a key role in shaping water mass properties. Examples include the Black Sea and the Baltic Sea, as well as convection sites in other marginal seas. In these environments, the choice of vertical mixing parameterization may similarly affect the overall water mass structure in regional ocean models.

Deleted: ,

Deleted: interaction

Deleted: with

Deleted: and to other semi-enclosed basins,

Deleted: ¶

Deleted: ,

Deleted: in order to further improve model realism and predictive skill.

Appendix A

This appendix briefly summarizes the main equations of the three vertical mixing schemes used in this study, outlining how the eddy viscosity coefficient A^{vm} and diffusivity coefficient A^{vT} are determined for each scheme. We note that the eddy diffusivity coefficient for temperature, A^{vT} , is assumed to be equal to that for salinity: $A^{vS} \equiv A^{vT}$.

1. Richardson-number dependent mixing scheme

The first closure scheme used here is based on the Richardson number–dependent formulation originally proposed by Pacanowski and Philander (1981), where the vertical eddy viscosity and diffusivity are parametrized as

$$A^{vm} = \frac{A_{ric}^{vm}}{(1 + \alpha R_l)^n} + A_b^{vm} \quad (A1)$$

$$A^{vt} = \frac{A^{vm}}{(1 + \alpha R_l)} + A_b^{vT} \quad (A2)$$

Where $R_l = \frac{N^2}{(\partial u_h)^2}$ denotes the Richardson number, defined as the ratio between stratification (represented by the squared Brunt-Väisälä frequency, N^2) and vertical shear $(\partial u_h)^2$. The parameter A_{ric}^{vm} represents the maximum value

of the eddy viscosity coefficient reached when $R_t < 0$; α and n are constant parameters. The additive terms A_b^{vm} and A_b^{vT} are background values that ensure a minimum level of vertical mixing even under stable conditions. For further details, see Pacanowski and Philander (1981) for the theoretical framework and Madec et al. (2023) for the practical implementation in NEMO.

2. Turbulent Kinetic Energy (TKE) vertical mixing scheme

The TKE vertical mixing scheme solves a prognostic equation for the turbulent kinetic energy k , while assuming a diagnostic closure for the tubulent length scale l_k . The TKE equation is written as

$$\frac{\partial k}{\partial t} = A^{vm} \left(\frac{\partial U_h}{\partial z} \right)^2 - A^{vT} N^2 + \frac{\partial}{\partial z} \left(A^{vm} \frac{\partial k}{\partial z} \right) - C_\epsilon \frac{k^{3/2}}{l_\epsilon} \quad (A3)$$

where U_h is the horizontal velocity, N^2 is squared Brunt-Väisälä frequency, C_ϵ is a constant and l_ϵ is the dissipative length scale. On the right side of equation (A3), the four terms represent the main physical processes associated with vertical mixing. From left to right: shear production, stratification (or buoyancy) destruction, TKE diffusion and Kolmogorov dissipation (Kolmogorov, 1942). Vertical mixing coefficients are then computed as:

$$A^{vm} = C_k l_k \sqrt{k} \quad (A4)$$

$$A^{vT} = \frac{A^{vm}}{P_{rt}} \quad (A5)$$

where l_k is the mixing length scale, C_k is a model constant, and P_{rt} is the Prandtl number. The turbulent mixing length l_k is computed diagnostically and its value is constrained by stratification (typically $l_k = l_\epsilon = \sqrt{2k}/N$, with additional operational caps imposed). For further details, see Gaspar et al. (1990) for the theoretical framework and Madec et al. (2023) for the practical implementation in NEMO.

3. Generalized Length Scale (GLS) vertical mixing scheme

The GLS vertical mixing schemes solves two prognostic equations, one for the turbulent kinetic energy k and one for the generic length scale ψ . The evolution of the turbulent kinetic energy is governed by

$$\frac{\partial k}{\partial t} = \frac{A^{vm}}{\sigma_\epsilon} \left(\frac{\partial U_h}{\partial z} \right)^2 - A^{vT} N^2 + \frac{\partial}{\partial z} \left(A^{vm} \frac{\partial k}{\partial z} \right) - C_{0\mu} \frac{k^{3/2}}{l_\epsilon} \quad (A6)$$

The equation is coupled with a prognostic equation for the generic length scale:

$$\frac{\partial \psi}{\partial t} = \frac{\psi}{e} \left\{ \frac{C_1 A^{vm}}{\sigma_\psi} \left(\frac{\partial U_h}{\partial z} \right)^2 - C_3 A^{vT} N^2 - C_2 C_{0\mu} \frac{k^{3/2}}{l_\epsilon} F_{wall} \right\} + \frac{\partial}{\partial z} \left(A^{vm} \frac{\partial \psi}{\partial z} \right) \quad (\text{A7})$$

Here, F_{wall} denotes the wall function and C_1, C_2, C_3, σ_k and σ_ψ are constants that depends on the choice of the turbulent model. $C_{0\mu}$ is also a constant. The wall function is only needed in the Mellor-Yamada model.

The eddy viscosity and diffusivity coefficients are then computed from the turbulent kinetic energy and mixing length:

$$A^{vm} = C_\mu l_k \sqrt{k} \quad (\text{A8})$$

$$A^{vT} = C_{\mu'} l_k \sqrt{k} \quad (\text{A9})$$

Where C_μ and $C_{\mu'}$ are the stability functions. The constant $C_{0\mu}$ depends on the choice of the stability functions. For further details, see Burchard and Bolding (2001) for the theoretical framework and Madec et al. (2023) for the practical implementation in NEMO.

Appendix B

In this appendix, we present the relevant portion of the namelist to be given as input to NEMO (version 4.2) corresponding to the three vertical mixing schemes used in this study. One table is provided for each scheme: Tabel B1 refers to the Richardson-number dependent schemes (Pacanowski and Philander, 1981), Table B2 to the TKE closure scheme (Gaspar et al., 1990) and Table B3 to the GLS closure scheme (Burchard and Bolding, 2001).

Parameter	Value	Explanation
rn_avm0	1.2e-6	vertical eddy viscosity [m2/s] (background value)
rn_avt0	1.0e-7	vertical eddy diffusivity [m2/s] (background value)
rn_avmri	100.e-4	maximum value of the vertical viscosity
rn_alp	5.0	coefficient of the parameterization
nn_ri	2	coefficient of the parameterization
ln_mldw	.false.	enhanced mixing in the Ekman layer

Table B1: Richardson-number dependent vertical mixing scheme

- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li

Parameter	Value	Explanation
rn_ediff	0.1	coefficient for vertical eddy diffusivity coefficient (avt=rn_ediff*mxl*sqrt(e))
rn_ediss	0.7	coefficient of the Kolmogoroff dissipation
rn_ebb	3.75	coefficient of the surface input of turbulent kinetic energy
rn_emin	1.e-8	minimum value of turbulent kinetic energy [m2/s2]
rn_emin0	5.e-4	surface minimum value of turbulent kinetic energy [m2/s2]
rn_bshear	1.e-20	background shear (>0) currently a numerical threshold (do not change it)
nn_pdl	1	Prandtl number function of Richardson number (=1, avt=pdl(Ri)*avm) or not (=0, avt=avm)
nn_mxl	2	mixing length: = 0 bounded by the distance to surface and bottom = 1 bounded by the local vertical scale factor = 2 first vertical derivative of mixing length bounded by 1 = 3 as =2 with distinct dissipative mixing length scale
ln_mxl0	.false.	mixing length scale surface value as a function of wind stress or not
rn_mxl0	0.04	surface buoyancy length scale minimum value
ln_mxhsw	.false.	mixing length scale surface value as a function of wave height or not
ln_lc	.true.	Langmuir cell parameterisation (Axell 2002)
rn_lc	0.15	coefficient associated to Langmuir cells
nn_etau	0	penetration of turbulent kinetic energy below the mixed layer (ML) = 0 none ; = 1 add a source of turbulent kinetic energy below the ML = 2 add a source of turbulent kinetic energy just at the base of the ML = 3 as = 1 applied on the high frequency part of the stress
nn_bc_surf	1	surface condition (0/1= Dirichlet/Neumann)
nn_bc_bot	1	bottom condition (0/1= Dirichlet/Neumann)

Table B2: Turbulent Kinetic Energy (TKE) vertical mixing schemes

- Deleted: ¶
- Formatted: Font: Bold
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li
- Formatted: Line spacing: Multiple 1,15 li

Parameter	Value	Explanation
rn_emin	1.e-8	minimum value of turbulent kinetic energy [m2/s2]
rn_epsmin	1.e-12	minimum value of dissipation [m2/s3]
ln_length_lim	.true.	limit on the dissipation rate under stable stratification (Galperin et al., 1988)
rn_clim_galp	0.267	Galperin limit
ln_sigpsi	.true.	Activate or not Burchard (2001) modification for k-ε closure and wave breaking mixing
rn_crban	100	Craig and Banner (1994) constant for surface wave breaking mixing
rn_charn	70000	Charnock constant for wave breaking induced roughness length
rn_hsro	0.01	Minimum surface roughness
rn_hsri	0.03	Ice-ocean roughness
rn_frac_hs	0.4	Fraction of wave height as surface roughness
nn_z0_met	2	Method for surface roughness computation (0/1/2/3) = 0 constant roughness = 1 Charnock formula = 2 Roughness formulae according to Rascle et al., 2008 = 3 Roughness given by the wave model
nn_z0_ice	0	attenuation of surface wave breaking under ice = 0 no impact of ice cover = 1 roughness uses rn_hsri and is weighted by 1-TANH(10*fr_i) = 2 roughness uses rn_hsri and is weighted by 1-fr_i = 3 roughness uses rn_hsri and is weighted by 1-MIN(1,4*fr_i)
nn_mxlice	0	mixing under sea ice = 0 No scaling under sea-ice = 1 scaling with constant Ice-ocean roughness (rn_hsri) = 2 scaling with mean sea-ice thickness = 3 scaling with max sea-ice thickness
nn_bc_surf	1	surface condition (0/1=Dirichlet/Neumann)
nn_bc_bot	1	bottom condition (0/1= Dirichlet/Neumann)
nn_stab_func	2	stability function (0=Galp, 1= KC94, 2=CanutoA, 3=CanutoB)
nn_clos	1	predefined closure type (0=MY82, 1=k-eps, 2=k-w, 3=Gen)

Table B3: Generalized Length Scale (GLS) vertical mixing scheme

Deleted: ¶

... [4]

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Line spacing: Multiple 1,15 li

Formatted: Normal

Appendix C

Throughout this paper, we define four statistics metrics (e.g., Willmott, 1981; Murphy, 1988; Murphy, 1992; Oke et al., 2002, Oddo et al., 2022): the root mean squared error (RMSE), the unbiased root mean squared error (uRMSE), the mean bias (MB), and cross-correlation coefficient (CC). We denote σ and $\bar{m}$ as the mean values of observations (o) and model data (m), N is the total number of observation-model pairs, and σ represents the standard deviation. The metrics are computed as follows:

- The root mean squared error (RMSE), which combines both systematic and random errors into a single measure of overall discrepancy:

$$RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (m_i - o_i)^2} \quad (C1)$$

- The unbiased root mean squared error (uRMSE), which isolates the random component of the model error by removing the bias:

$$uRMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N [(m_i - \bar{m}) - (o_i - \sigma)]^2} \quad (C2)$$

- The mean bias (MB), which quantifies the average systematic difference between the model and observations:

$$MB = \bar{m} - \sigma \quad (C3)$$

- The cross-correlation coefficient (CC), which measures the degree of linear association between model outputs and observations:

$$CC = \frac{1}{\sigma_o} \frac{1}{\sigma_m} \frac{1}{N} \sum_{i=1}^N (m_i - \bar{m})(o_i - \sigma) \quad (C4)$$

Code availability

The ocean hydrodynamic circulation model is based on the Nucleus for European Modelling of the Ocean (NEMO) version 4.2.0 (Madec et al., 2023), available on GitLab (<https://forge.nemo-ocean.eu/nemo/nemo/-/releases/4.2.0>) under [doi:10.5281/zenodo.6334656](https://doi.org/10.5281/zenodo.6334656). The wind-wave model is based on WAVEWATCH IIITM (WW3) version 6.07 (Tolman et al., 2021), accessible through GitHub (<https://github.com/NOAA-EMC/WW3/releases>). NEMO namelist configurations for the vertical mixing closure schemes used in this study are provided in Appendix B.

Author contribution

LG and PO conceived and designed the study with input from HB, who had a central role in the discussions and development strategy. LG implemented and tested the model setup through sensitivity experiments, with valuable contributions from FB and AM. LG performed the simulations and data analysis. PM contributed to the validation and visualization of the results. FM and EC participated in the discussions and provided feedback. LG wrote the manuscript with input and contributions from all authors, who reviewed and approved the final version of the manuscript.

Acknowledgements

The authors would like to thank Prof. Nadia Pinardi for insightful discussions throughout the development of this study and Prof. Paola Cessi for helpful comments.

References

Axell, L. B., Wind-driven internal waves and Langmuir circulations in a numerical ocean model of the southern Baltic Sea, *Journal of Geophysical Research: Oceans*, 107.C11, 25-1, 2002.

Bethoux, J. P., Gentili, B., Morin, P., Nicolas, E., Pierre, C., and Ruiz-Pino, D.: The Mediterranean Sea: a miniature ocean for climatic and environmental studies and a key for the climatic functioning of the North Atlantic, *Progress in Oceanography*, 44, 131–146, 1999.

Bryden, H. L. and Stommel, H. M.: Origin of the Mediterranean outflow, *Journal of Marine Research*, 40, 55–71, 1982.

Burchard, H. and Bolding, K.: Comparative analysis of four second-moment turbulence closure models for the oceanic mixed layer, *Journal of Physical Oceanography*, 31, 1943–1968, [https://doi.org/10.1175/1520-0485\(2001\)031<1943:CAOFSM>2.0.CO;2](https://doi.org/10.1175/1520-0485(2001)031<1943:CAOFSM>2.0.CO;2), 2001.

Burchard, H., Simulating the wave-enhanced layer under breaking surface waves with two-equation turbulence models, *Journal of Physical Oceanography*, 31, 11, 3133–3145, 2001.

Burchard, H. and Petersen, O.: Models of turbulence in the marine environment - A comparative study of two-equation turbulence models. *Journal of Marine Systems*, 21, 29-53, 1999.

Canuto, V. M., Howard, A., Cheng, Y., and Dubovikov, M. S.: Ocean turbulence. Part I: One-point closure model–momentum and heat vertical diffusivities, *Journal of Physical Oceanography*, 31, 1413–1426, [https://doi.org/10.1175/1520-0485\(2001\)031<1413:OTPIOP>2.0.CO;2](https://doi.org/10.1175/1520-0485(2001)031<1413:OTPIOP>2.0.CO;2), 2001.

Clementi, E., Oddo, P., Drudi, M., Pinardi, N., Korres, G., and Grandi, A.: Coupling hydrodynamic and wave models: first step and sensitivity experiments in the Mediterranean Sea, *Ocean Dynamics*, 67, 1293–1312, 2017.

Clementi, E., Drudi, M., Aydogdu, A., Moulin, A., Grandi, A., Mariani, A., Goglio, A. C., Pistoia, J., Miraglio, P., Lecci, R., Palermo, F., Coppini, G., Masina, S., and Pinardi, N.: Mediterranean Sea Physical Analysis and Forecast (CMEMS MED-

Deleted: The formation and circulation of deep water in the Mediterranean Sea

Deleted: Nature

Deleted: 296

Deleted: 49

Deleted: 5

Deleted: <https://doi.org/10.1038/296049a0>,

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, Not Italic

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Deleted: -

Physics, EAS8 system)(Version 1)[Data set],
https://doi.org/10.25423/CMCC/MEDSEA_ANALYSISFORECAST_PHY_006_013_EAS8, 2023.

Coppini, G. and Clementi, E. and Cossarini, G. and Salon, S. and Korres, G. and Ravdas, M. and Lecci, R. and Pistoia, J. and Goglio, A. C. and Drudi, M. and Grandi, A. and Aydogdu, A. and Escudier, R. and Cipollone, A. and Lyubartsev, V. and Mariani, A. and Creti, S. and Palermo, F. and Scuro, M. and Masina, S. and Pinardi, N. and Navarra, A. and Delrosso, D. and Teruzzi, A. and Di Biagio, V. and Bolzon, G. and Feudale, L. and Coidessa, G. and Amadio, C. and Brosich, A. and Mir'ò, A. and Alvarez, E. and Lazzari, P. and Solidoro, C. and Oikonomou, C. and Zacharioudaki, A.: The Mediterranean Forecasting System—part 1: evolution and performance, *Ocean Science*, 19, 1483–1516, 2023.

Craig, P. D., and Banner, M.L., Modeling wave-enhanced turbulence in the ocean surface layer, *Journal of Physical Oceanography*, 24, 12, 2546-2559, 1994.

de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology, *Journal of Geophysical Research: Oceans*, 109, C12003, https://doi.org/10.1029/2004JC002378, 2004.

Drévillon, M., Bourdallé-Badie, R., Derval, C., Lellouche, J., Rémy, E., Tranchant, B., Benkiran, M., Greiner, E., Guinehut, S., Verbrugge, N., Garric, G., Testut, C.-E., Laborie, M., Nouel, L., Bahurel, P., Bricaud, C., Crosnier, L., Dombrowsky, E., Durand, E., Ferry, N., Hernandez, F., Le Galloudec, O., Messal, F., and Parent, L.: The GODAE/Mercator-Ocean global ocean forecasting system: results, applications and prospects, *Journal of Operational Oceanography*, 1, 51–57, https://doi.org/10.1080/1755876X.2008.11020095, 2008.

Escudier, R., Clementi, E., Cipollone, A., Pistoia, J., Drudi, M., Grandi, A., Lyubartsev, V., Lecci, Aydogdu, A., Del Rosso, D., Omar, M., Masina, S., Coppini, G., and Pinardi, N.: A High Resolution Reanalysis for the Mediterranean Sea, *Frontiers in Earth Science*, 9, 1060, https://doi.org/10.3389/feart.2021.702285, 2021.

Gačić, M., Lascaratos, A., Manca, B.B. and Mantziafou, A.: *Adriatic deep water and interaction with the Eastern Mediterranean Sea. In Physical oceanography of the Adriatic Sea: Past, present and future* (pp. 111-142). Dordrecht: Springer Netherlands, 2001.

Galperin, B., Kantha, L. H., Hassid, S., and Rosati, A.: A quasi-equilibrium turbulent energy model for geophysical flows, *Journal of the atmospheric sciences*, 45(1), 55-62, https://doi.org/10.1175/1520-0469(1988)045<0055:AQETEM>2.0.CO;2, 1988.

Gaspar, P., Grégoris, Y., and Lefevre, J. M.: A simple eddy kinetic energy model for simulations of the oceanic vertical mixing: Tests at station Papa and long-term upper ocean study site, *Journal of Geophysical Research: Oceans*, 95, 16179–16193, https://doi.org/10.1029/JC095iC09p16179, 1990.

Global Ocean Physics Analysis and Forecast. E.U. Copernicus Marine Service Information (CMEMS). Marine Data Store (MDS). doi: https://doi.org/10.48670/moi-00016 (Accessed on 15-04-2026)

Global Ocean-In-Situ Near-Real-Time Observations. E.U. Copernicus Marine Service Information (CMEMS). Marine Data Store (MDS). doi: https://doi.org/10.48670/moi-00036 (Accessed on 15-05-2026)

Griffies, S. M., Böning, C., Bryan, F.O., Chassignet, F. P., Gerdes, R., Hasumi, H., Hirst, A., Treguier, A.M. and Webb, D.: Developments in ocean climate modelling, *Ocean Modelling*, 2, 123–192, https://doi.org/10.1016/S1463-5003(00)00014-7, 2000.

Deleted: Coppini, G., Clementi, E., Cossarini, G., Salon, S., Korres, G., Ravdas, M., Lecci, R., Pistoia, J., Goglio, A. C., Drudi, M., et al.

Deleted: m

Deleted: f

Deleted: s

Deleted: Omar, M.,

Deleted: Aydogdu, A.,

Formatted: English (UK)

Deleted: R., Creti, S.,

Deleted: :

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman)

Deleted: ,

Deleted: &

Deleted: Hassid, S.,

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Normal

Formatted: Font: (Default) +Body (Times New Roman), Font colour: Custom Colour (RGB(137,68,3))

Deleted: Gerbeau, J.-F. and Le Roux, A.: A flux-based method for convection-diffusion equations in ocean modelling, *Ocean Modelling*, 3, 229–239, https://doi.org/10.1016/S1463-5003(01)00006-2, 2001.

Deleted: 28

Deleted: 10

Deleted: 5

Deleted: 28

Deleted: 10

Deleted: 5

Deleted: Bryan

Deleted: K

Deleted: Gnanadesikan

Deleted: A

Deleted: Hallberg

Deleted: R

Deleted: W

Deleted: Marshall

Deleted: J

Deleted: and Samuels, B. L.

Gutjahr, O., Brüggemann, N., Haak, H., Jungclauss, J. H., Putrasahan, D. A., Lohmann, K., and von Storch, J.-S.: Comparison of ocean vertical mixing schemes in the Max Planck Institute Earth System Model (MPI-ESM1. 2), *Geoscientific Model Development*, 14, 2317–2349, 2021.

1380 Hermann, M., Somot, S., Sevault, F., Estournel, C., and Déqué, M.: Modeling the deep convection in the northwestern Mediterranean Sea using an eddy-permitting and an eddy-resolving model: Case study of winter 1986–1987, *Journal of Geophysical Research: Oceans*, 113, 2008.

Hilt, M., Auclair, F., Benshila, R., Bordis, L., Capet X., Debreu, L., Dumas, F., Julien, S., Lamarié, F., Marchisiello, P., Nguyen, C.: Numerical modeling of hydraulic control, solitary waves and primary instabilities in the Strait of Gibraltar, *Ocean Modelling*, 151, 101642, <https://doi.org/10.1016/j.ocemod.2020.101642>, 2020.

1385 Kolmogorov, A. N.: Equations of turbulent motion in an incompressible fluid, *Izvestiya Akademiyi Nauk SSSR, Seriya Fizicheskaya*, 6, 56–58, 1942.

Kubin, E., Poulain, P.-M., Mauri, E., Menna, M., and Notarstefano, G.: Levantine Intermediate and Levantine Deep Water Formation: An Argo Float Study from 2001 to 2017, *Water*, 11, 1781, <https://doi.org/10.3390/w11091781>, 2019.

1390 Lascaratos, A., Williams, R.G. and Tragou, E.: A mixed-layer study of the formation of Levantine Intermediate Water, *Journal of Geophysical Research: Oceans*, 98, C8, 14739-14749., 1993

Lascaratos, A., Roether, W., Nittis, K. and Klein, B.: Recent changes in deep water formation and spreading in the eastern Mediterranean Sea: a review. *Progress in oceanography*, 44, 1-3, 5-36, 1999

Lazar, A., Madec, G., and Delecluse, P.: The deep interior downwelling, the Veronis effect, and mesoscale tracer transport parameterizations in an OGCM, *Journal of physical oceanography*, 29, 2945–2961, 1999.

1395 Legay, A., B. Deremble, H. Burchard, Derivation and implementation of a non-local term to improve the oceanic convection representation within the k-ε parameterization, *Journal Advances in Modeling Earth Systems*, 17, e2024MS004243, 2025. <https://doi.org/10.1029/2024MS004243>

1400 Le Sommer, J., Penduff, T., Penduff, T., Theetten, S., Madec, G., and Barnier, B.: How momentum advection schemes include current topography interactions at eddy permitting resolution, *Ocean Modeling*, 29, <https://doi.org/10.1016/j.ocemod.2008.11.007>, 2009.

Lévy, M., Estublier, A. and Madec, G.: Choice of an advection scheme for biogeochemical models, *Geophysical Research Letters*, 28(19), 3725-3728, 2001.

Lionello, P., Malanotte-Rizzoli, P., Boscolo, R., (eds). *Mediterranean climate variability*, vol. 4. Elsevier, 2006.

1405 Luneva, M., Wakelin, S., Holt, J., et al.: Challenging Vertical Turbulence Mixing Schemes in a Tidally Energetic Environment: 1. 3-D Shelf-Sea Model Assessment, *Journal of Geophysical Research: Oceans*, 124, 6360–6387, <https://doi.org/10.1029/2018JC014307>, 2019.

Madec, G. and the NEMO System Team: NEMO Ocean Engine Reference Manual, Tech. rep., <https://doi.org/10.5281/zenodo.8167700>, 2023.

Deleted: Roblou...clair, FL..., Nguyen...enshila, RC..., Marchesiello...ordis, LP..., Lemarié,...apet XF..., Jullien...ebreu, LS..., Dumas, F., Debreu...ulien, SL..., Capet...amarié, FX..., Bordoio...archisiello, PL..., Benshila...guyen, CR., and Auclair, F.:... Numerical modeling of hydraulic control, solitary waves and primary instabilities in the Strait of Gibraltar, *Ocean Modelling*, 151, 101

Deleted: Large, W. G. and Deleersnijder, E.: Stability functions for a mixed-layer model, *Journal of Physical Oceanography*, 23, 778–785, [https://doi.org/10.1175/1520-0485\(1993\)023<0778:SFFAML>2.0.CO;2](https://doi.org/10.1175/1520-0485(1993)023<0778:SFFAML>2.0.CO;2), 1993.

Deleted:

Deleted: gradient ...ocal term to improve the oceanic convection representation within the k-ε parameterization, *Journal....Advances....in Modeling....Earth Systems*.

Deleted: Madec...enduff, TG..., Jourdain...heetten, SN. C..., Beckmann...adec, GA..., and Barnier, B.: Modeling the dynamics of the mesoscale eddy field in the North Atlantic ocean: Scale selection and parameter sensitivity...ow momentum advection schemes include current topography interactions at eddy permitting resolution, *Journal of Climate...cean Modeling*, 2922, 3017–3

Deleted: <https://doi.org/10.1175/2008JCLI5714.1>

Formatted: Font: (Default) +Body (Times New Roman), 10 pt

Formatted

Formatted: Font: Not Italic

Madec, G., Delecluse, P., Crepon, M., and Chartier, M.: A three-dimensional numerical study of deep-water formation in the northwestern Mediterranean Sea, *Journal of Physical Oceanography*, 21, 1349–1371, 1991.

1500 Manca, B.B., Budillon, G., Scarazzato, P. and Ursella, L., Evolution of dynamics in the eastern Mediterranean affecting water mass structures and properties in the Ionian and Adriatic Seas. *Journal of Geophysical Research: Oceans*, 108(C9), 2003.

Marshall, J. and Schott, F.: Open-ocean convection: Observations, theory, and models, *Reviews of Geophysics*, 37, 1–64, <https://doi.org/10.1029/98RG02739>, 1999.

1505 Mazzetti, C., Carton De Wiart, C., Gomes, G., Russo, C., Decremier, D., Grimaldi, S., Disperati, J., Ziese, M., Schweim, C., Garcia Sanchez, R., Jacobson, T., Ramos, A., Prudhomme, C., and Salamon, P.: EFAS v5.0 hydrological reanalysis, JRC Technical Report JRC134686, European Commission, Joint Research Centre (JRC), Ispra, Italy, 2023.

MEDOC Group: Observation of formation of deep water in the Mediterranean Sea, 1969, *Nature*, 227, 1037–1040, <https://doi.org/10.1038/2271037a0>, 1970.

1510 Mellor, G. L. and Yamada, T.: Development of a turbulence closure model for geophysical fluid problems, *Reviews of Geophysics*, 20, 851–875, <https://doi.org/10.1029/RG020i004p00851>, 1982.

Millot, C. and Taupier-Letage, I.: Circulation in the Mediterranean Sea, in: *The Mediterranean Sea*, pp. 29–66, Springer, 2005.

Murphy, A. H.: Skill scores based on the mean square error and their relationships to the correlation coefficient, *Monthly Weather Review*, 116, 2417–2424, [https://doi.org/10.1175/1520-0493\(1988\)116<2417:SSBOTM>2.0.CO;2](https://doi.org/10.1175/1520-0493(1988)116<2417:SSBOTM>2.0.CO;2), 1988.

1515 Murphy, A. H.: Climatology, persistence, and their linear combination as standards of reference in skill scores, *Weather and Forecasting*, 7, 692–698, [https://doi.org/10.1175/1520-0434\(1992\)007<0692:CPATLC>2.0.CO;2](https://doi.org/10.1175/1520-0434(1992)007<0692:CPATLC>2.0.CO;2), 1992.

Oddo, P., Falchetti, S., Viola, S., Pennucci, G., Storto, A., Borriore, I., Giorli, G., Cozzani, E., Russo, A., Tollefsen, C.: Evaluation of different Maritime rapid environmental assessment procedures with a focus on acoustic performance, *The Journal of the Acoustical Society of America*, 152, 5, 2962–2981, 2022.

1520 Oddo, P., Adani, M., Pinardi, N., Fratianni, C., Tonani, M., and Pettenuzzo, D.: A nested Atlantic–Mediterranean Sea general circulation model for operational forecasting, *Ocean Science*, 5, 461–473, <https://doi.org/10.5194/os-5-461-2009>, 2009.

Oke, P. R., Allen, J. S., Miller, R. N., Egbert, G. D., Austin, J. A., Barth, J. A., Boyd, T. J., Kosro, P. M., and Levine, M. D.: A modeling study of the three-dimensional continental shelf circulation off Oregon. Part I: Model-data comparison, *Journal of Physical Oceanography*, 32, 1360–1382, [https://doi.org/10.1175/1520-0485\(2002\)032<1360:AMOSOT>2.0.CO;2](https://doi.org/10.1175/1520-0485(2002)032<1360:AMOSOT>2.0.CO;2), 2002.

1525 Pacanowski, R. C. and Philander, S. G. H.: Parameterization of vertical mixing in numerical models of tropical oceans, *Journal of Physical Oceanography*, 11, 1443–1451, 1981.

1530 Pinardi, N. and Navarra, A., 1993, Baroclinic wind adjustment processes in the Mediterranean Sea. *Deep Sea Research Part II: Topical Studies in Oceanography*, 40(6), pp.1299-1326.

Formatted: Font: (Default) +Body (Times New Roman), English (US)

Formatted: Font: (Default) +Body (Times New Roman)

Deleted: -

Formatted: Font: (Default) Times New Roman, Italian

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, Not Italic

Formatted: Font: (Default) Times New Roman

Pinardi, N., Allen, I., Demirov, E., De Mey, P., Korres, G., Lascaratos, A., Le Traon, P.-Y., Maillard, C., Manzella, G., and Tziavos, C.: The Mediterranean ocean forecasting system: first phase of implementation (1998–2001), *Annales Geophysicae*, 21, 3–20, <https://doi.org/10.5194/angeo-21-3-2003>, 2003.

535 Pinardi, N. and Coppini, G.: Preface "Operational oceanography in the Mediterranean Sea: the second stage of development", *Ocean Sciences*, 6, 263–267, <https://doi.org/10.5194/os-6-263-2010>, 2010.

Pinardi, N. and Masetti, E.: Variability of the large scale general circulation of the Mediterranean Sea from observations and modelling: A review, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 158, 153–173, [https://doi.org/10.1016/S0031-0182\(00\)00048-1](https://doi.org/10.1016/S0031-0182(00)00048-1), 2000.

1540 Pinardi, N., Cessi, P., Borile, F. and Wolfe, C.L.: The Mediterranean sea overturning circulation. *Journal of Physical Oceanography*, 49(7), 1699–1721, 2019.

Pinardi, N., Estournel, C., Cessi, P., Escudier, R. and Lyubartsev, V., Dense and deep water formation processes and Mediterranean overturning circulation. In *Oceanography of the mediterranean sea* (pp. 209–261). Elsevier, 2023.

Poulain, P.-M., Barbanti, R., Font, J., Cruzado, A., Millot, C., Gertman, I., Griffa, A., Molcard, A., Rupolo, V., Le Bras, S., Petit de la Villeon L.: MedArgo: a drifting profiler program in the Mediterranean Sea, *Ocean Science*, 3, 379–395, 2007.

545 Rahmstorf, S.: A fast and complete convection scheme for ocean models, *Ocean Modelling*, 101, 9–11, 1993.

Reffray, G., Bourdallé-Badie, R., and Calone, C.: Modelling turbulent vertical mixing sensitivity using a 1-D version of NEMO, *Geoscientific Model Development*, 8, 69–86, <https://doi.org/10.5194/gmd-8-69-2015>, 2015.

1550 Rasche, N., Arduin, F., Queffelec, P., & Croizé-Fillon, D., A global wave parameter database for geophysical applications. Part 1: Wave-current–turbulence interaction parameters for the open ocean based on traditional parameterizations, *Ocean Modelling*, 25, 3–4, 154–171, 2008.

Robertson, R., Dong, C.: An evaluation of the performance of vertical mixing parameterizations for tidal mixing in the Regional Ocean Modeling System (ROMS), *Geoscience Letters*, 6, 15, <https://doi.org/10.1186/s40562-019-0146-y>, 2019.

555 Robinson, A. R. and Golnaraghi, M. F.: Circulation and dynamics of the eastern Mediterranean Sea: quasi-synoptic data-driven simulations, *Deep Sea Research Part II: Topical Studies in Oceanography*, 40, 6, 1207–1246, 1993.

Said, M. A., Gerges, M. A., Maiyya, I. A., Hussein, M. A., and Radwan, A. A.: Changes in Atlantic Water characteristics in the south-eastern Mediterranean Sea as a result of natural and anthropogenic activities, *Oceanologia*, 53, 81–95, <https://doi.org/10.5697/oc.53-1.81>, 2011.

560 Schroeder, K., Chiggiato, J., Bryden, H. L., Borghini, M., and Ben Ismail, S.: Abrupt climate shift in the Western Mediterranean Sea, *Scientific Reports*, 2, <https://doi.org/10.1038/srep23009>, 2012.

Shchepetkin, A. F.: An adaptive, Courant-number-dependent implicit scheme for vertical advection in oceanic modeling, *Ocean Modelling*, 91, 38–69, 2015.

Simoncelli, S., Pinardi, N., Fratianni, C., Dubois, C., and Notarstefano, G.: Water mass formation processes in the Mediterranean Sea over the past 30 years. Copernicus Marine Service Ocean State Report, Issue 2. K. von Schuckmann et al., Eds., s96- s100, *Journal Operational Oceanography*, 11, s1–s142, 2018.

595 Somot, S., Sevault, F., and Déqué, M.: *Transient climate change scenario simulation of the Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation model*, *Climate Dynamics*, 27, 851–879, <https://doi.org/10.1007/s00382-006-0167-z>, 2006.

Storto, A., Hesham Essa, Y., de Toma, V., Anav, A., Sannino, G., Santoleri, R., and Yang, C.: MESMAR v1: a new regional coupled climate model for downscaling, predictability, and data assimilation studies in the Mediterranean region, *Geoscientific Model Development Discussions*, 1–40, 2023.

600 Taillandier, V., D’Ortenzio, F., Prieur, L., Conan, P., Coppola, L., Cornec, M., Dumas, F., Durrieu de Madron, X., Fach, B., Fourier, M. and Gentil, M.: *Sources of the Levantine intermediate water in winter 2019*, *Journal of Geophysical Research: Oceans*, 127(6), p.e2021JC017506, 2022.

Testor, P., Bosse A., Houpert L., Margirier F., Mortier L., Legoff H., Dausse D., Labaste M., Karstensen J., Hayes D., Olita A.: Multiscale observations of deep convection in the northwestern Mediterranean Sea during winter 2012–2013 using multiple platforms, *Journal of Geophysical Research: Oceans*, 123, 1745–1776, <https://doi.org/10.1002/2016JC012671>, 2018.

1605 Tolman, H. L.: User Manual and System Documentation of WAVEWATCH III™ version 6.07, Tech. rep., NOAA/NWS/NCEP/MMAB, <https://polar.ncep.noaa.gov/waves/wavewatch/wavewatch.shtml>, 2021.

Tonani, M., Pinardi, N., Fratianni, C., Pistoia, J., Dobricic, S., Pensieri, S., de Alfonso, M., and Nittis, K.: Mediterranean Forecasting System: forecast and analysis assessment through skill scores, *Ocean Science*, 5, 649–660, <https://doi.org/10.5194/os-5-649-2009>, 2009.

1610 Umlauf, L. and Burchard, H.: A generic length-scale equation for geophysical turbulence models, *Journal of Marine Research*, 61, 235–265, 2003.

Villarreal, M.R., K. Bolding, Burchard, H., and E. Demirov, Coupling of the GOTM turbulence module to some three-dimensional ocean models, pp. 225-237. In: Baumert, H.Z., J.H. Simpson, and J. Sündermann (eds.), *Marine Turbulence: Theories, Observations and Models*, Cambridge University Press, Cambridge, 630 pp. 2005.

1615 Von Schuckmann, K., Le Traon, P.-Y., Smith, N., Pascual, A., Brasseur, P., Fennel, K., Djavidnia, S., Aaboe, S., Fanjul, E. A., Autret, E., Axell L.: Copernicus marine service ocean state report, *Journal of Operational Oceanography*, 11, S1–S142, 2018.

1620 Wesson, J. C. and Gregg, M. C.: Mixing at Camarinal Sill in the Strait of Gibraltar, *Journal of Geophysical Research: Oceans*, 99, 9847–9878, <https://doi.org/10.1029/94JC00256>, 1994.

Willmott, C. J.: On the validation of models, *Physical Geography*, 2, 184–194, <https://doi.org/10.1080/02723646.1981.10642213>, 1981.

625 Wong, A. P., Wijffels, S. E., Riser, S. C., Pouliquen, S., Hosoda, S., Roemmich, D., Gilson, J., Johnson, G. C., Martini, K., Murphy, D. J., Scanderberg M, Bhaskar TVSU, Buck JJH, Merceur F, Carval T, Maze G, Cabanes C, André X, Poffa N, Yashayaev I, Barker PM, Guinehut S, Belbéoch M, Ignaszewski M, Baringer MO, Schmid C, Lyman JM, McTaggart KE, Purkey SG, Zilberman N, Alkire MB, Swift D, Owens WB, Jayne SR, Hersh C, Robbins P, West-Mack D, Bahr F, Yoshida

Deleted: .

Deleted: r

Deleted: .

Deleted: .

Deleted: Simoncelli, S., Sannino, S., Mazloff, M. R., Vetrano, A., and et al.: Mediterranean Ocean State Report 2, Tech. rep., Euro-Mediterranean Center on Climate Change (CMCC), <https://med.oceanrep.org/>, accessed: 2025-07-31, 2022. ¶

Deleted: G

Deleted: Transitional climate change scenarios for the Mediterranean region

Deleted: , <https://doi.org/10.1007/s00382-006-0126-6>

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: (Default) +Body (Times New Roman)

Deleted: Mortier, L., Bosse, A., et al.

Formatted: Font: (Default) +Body (Times New Roman)

Deleted: Theocharis, A., Lascaratos, A., Nittis, K., and Georgopoulos, D.: The intermediate waters of the Mediterranean Sea: a review, *Progress in Oceanography*, 55, 287–316, [https://doi.org/10.1016/S0079-6611\(02\)00114-3](https://doi.org/10.1016/S0079-6611(02)00114-3), 2002. ¶

Deleted: , <https://doi.org/10.1357/002224003322005087>

Deleted: et al.

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Font: 10 pt

650 S. Sutton PJH, Cancouët R, Coatanoan C, Dobbler D, Juan AG, Gourrion J, Kolodziejczyk N, Bernard V, Bourlès B, Claustre H, D'Ortenzio F, Le Reste S, Le Traon P-Y, Rannou J-P, Saout-Grit C, Speich S, Thierry V, Verbrugge N, Angel-Benavides IM, Klein B, Notarstefano G, Poulain P-M, Vélez-Belchí P, Suga T, Ando K, Iwasaka N, Kobayashi T, Masuda S, Oka E, Sato K, Nakamura T, Sato K, Takatsuki Y, Yoshida T, Cowley R, Lovell JL, Oke PR, van Wijk EM, Carse F, Donnelly M, Gould WJ, Gowers K, King BA, Loch SG, Mowat M, Turton J, Rama Rao EP, Ravichandran M, Freeland HJ, Gaboury I, Gilbert D, Greenan BJW, Ouellet M, Ross T, Tran A, Dong M, Liu Z, Xu J, Kang K, Jo H, Kim S-D and Park H-M: Argo data 1999–2019: Two million temperature-salinity profiles and subsurface velocity observations from a global array of profiling floats, *Frontiers in Marine Science*, 7, 700, 2020.

Deleted: et al.:

655 Wüst, G., On the vertical circulation of the Mediterranean Sea. *Journal of Geophysical Research*, 66(10), pp.3261-3271, 1961.

Formatted: Font: (Default) Times New Roman

Page 17: [1] Deleted	Lucia Gualtieri	12/03/2026 10:09:00
----------------------	-----------------	---------------------

▼

Page 26: [2] Deleted	Lucia Gualtieri	11/03/2026 15:35:00
----------------------	-----------------	---------------------

▼

Page 28: [3] Deleted	Lucia Gualtieri	12/03/2026 10:34:00
----------------------	-----------------	---------------------

▼

Page 28: [3] Deleted	Lucia Gualtieri	12/03/2026 10:34:00
----------------------	-----------------	---------------------

▼

Page 28: [3] Deleted	Lucia Gualtieri	12/03/2026 10:34:00
----------------------	-----------------	---------------------

▼

Page 28: [3] Deleted	Lucia Gualtieri	12/03/2026 10:34:00
----------------------	-----------------	---------------------

▼

Page 28: [3] Deleted	Lucia Gualtieri	12/03/2026 10:34:00
----------------------	-----------------	---------------------

▼

Page 28: [3] Deleted	Lucia Gualtieri	12/03/2026 10:34:00
----------------------	-----------------	---------------------

▼

Page 28: [3] Deleted	Lucia Gualtieri	12/03/2026 10:34:00
----------------------	-----------------	---------------------

▼

Page 28: [3] Deleted	Lucia Gualtieri	12/03/2026 10:34:00
----------------------	-----------------	---------------------

▼

Page 38: [4] Deleted	Lucia Gualtieri	16/04/2026 09:38:00
----------------------	-----------------	---------------------

▼

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼.....

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼.....

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼.....

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼.....

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼.....

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼.....

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼.....

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼.....

Page 42: [5] Deleted	Lucia Gualtieri	20/03/2026 14:50:00
----------------------	-----------------	---------------------

▼

Page 42: [5] Deleted

Lucia Gualtieri

20/03/2026 14:50:00

▼

Page 42: [5] Deleted

Lucia Gualtieri

20/03/2026 14:50:00

▼

Page 42: [5] Deleted

Lucia Gualtieri

20/03/2026 14:50:00

▼

Page 42: [5] Deleted

Lucia Gualtieri

20/03/2026 14:50:00

▼

Page 42: [5] Deleted

Lucia Gualtieri

20/03/2026 14:50:00

▼

Page 42: [5] Deleted

Lucia Gualtieri

20/03/2026 14:50:00

▼

Page 42: [5] Deleted

Lucia Gualtieri

20/03/2026 14:50:00

▼

Page 42: [6] Deleted

Lucia Gualtieri

20/03/2026 14:55:00

▼

Page 42: [7] Deleted

Lucia Gualtieri

20/03/2026 15:44:00

▼

Page 42: [7] Deleted	Lucia Gualtieri	20/03/2026 15:44:00
----------------------	-----------------	---------------------

▼

Page 42: [7] Deleted	Lucia Gualtieri	20/03/2026 15:44:00
----------------------	-----------------	---------------------

▼

Page 42: [7] Deleted	Lucia Gualtieri	20/03/2026 15:44:00
----------------------	-----------------	---------------------

▼

Page 42: [7] Deleted	Lucia Gualtieri	20/03/2026 15:44:00
----------------------	-----------------	---------------------

▼

Page 42: [8] Deleted	Lucia Gualtieri	20/03/2026 15:49:00
----------------------	-----------------	---------------------

▼

Page 42: [8] Deleted	Lucia Gualtieri	20/03/2026 15:49:00
----------------------	-----------------	---------------------

▼

Page 42: [8] Deleted	Lucia Gualtieri	20/03/2026 15:49:00
----------------------	-----------------	---------------------

▼

Page 42: [8] Deleted	Lucia Gualtieri	20/03/2026 15:49:00
----------------------	-----------------	---------------------

▼

Page 42: [8] Deleted	Lucia Gualtieri	20/03/2026 15:49:00
----------------------	-----------------	---------------------

▼

Page 42: [8] Deleted	Lucia Gualtieri	20/03/2026 15:49:00
----------------------	-----------------	---------------------

▼.....

Page 42: [8] Deleted	Lucia Gualtieri	20/03/2026 15:49:00
----------------------	-----------------	---------------------

▼.....

Page 42: [8] Deleted	Lucia Gualtieri	20/03/2026 15:49:00
----------------------	-----------------	---------------------

▼.....

Page 42: [8] Deleted	Lucia Gualtieri	20/03/2026 15:49:00
----------------------	-----------------	---------------------

▼.....

Page 42: [9] Formatted	Lucia Gualtieri	20/03/2026 15:43:00
------------------------	-----------------	---------------------

Font: (Default) +Body (Times New Roman)

Page 42: [9] Formatted	Lucia Gualtieri	20/03/2026 15:43:00
------------------------	-----------------	---------------------

Font: (Default) +Body (Times New Roman)

Page 42: [9] Formatted	Lucia Gualtieri	20/03/2026 15:43:00
------------------------	-----------------	---------------------

Font: (Default) +Body (Times New Roman)

Page 42: [9] Formatted	Lucia Gualtieri	20/03/2026 15:43:00
------------------------	-----------------	---------------------

Font: (Default) +Body (Times New Roman)

Page 42: [9] Formatted	Lucia Gualtieri	20/03/2026 15:43:00
------------------------	-----------------	---------------------

Font: (Default) +Body (Times New Roman)

Page 42: [9] Formatted	Lucia Gualtieri	20/03/2026 15:43:00
------------------------	-----------------	---------------------

Font: (Default) +Body (Times New Roman)

Untangling the effects of vertical mixing schemes and convective adjustment in the Mediterranean Sea: insights from a sensitivity study

Lucia Gualtieri¹, Hans Burchard², Federica Borile³, Aimie Moulin¹, Pietro Miraglio¹, Francesco Maicu¹, Emanuela Clementi¹, Paolo Oddo³

5 ¹CMCC Foundation - Euro-Mediterranean Center on Climate Change, Bologna, Italy.

²Leibniz Institute for Baltic Sea Research Warnemünde, Rostock, Germany.

³University of Bologna, Department of Physics and Astronomy, Bologna, Italy.

Correspondence to: Lucia Gualtieri (lucia.gualtieri@cmcc.it)

Abstract. The Mediterranean Sea provides a natural laboratory for investigating ocean circulation processes of global
10 relevance due to its complex dynamics, active deep and intermediate water formation, and sensitivity to climate variability. Regional ocean circulation models' skill strongly depends on the parameterization of subgrid-scale processes, among which turbulent vertical mixing and convection play a major role. To evaluate their impact in the Mediterranean Sea two-year-long simulations were conducted using three vertical closure schemes: the Richardson-number dependent parameterisation, the
15 Turbulent Kinetic Energy (TKE) scheme, and the Generalised Length Scale (GLS) scheme. Each scheme was tested both with and without the convective adjustment approach, resulting in a set of comparative experiments designed to isolate the contribution of the adjustment process and its combined effect with each parameterization. Model results are evaluated against all available Argo floats data, both at the basin scale and in key deep and intermediate water formation regions. The simulations show that adding a convective adjustment is crucial to accurately reproduce observations with the Richardson-number
20 dependent parameterization, where it improves all key variables, while for the TKE scheme it is particularly important for representing the mixed layer depth across the basin and in deep water formation areas. For more physics-based vertical schemes, like the GLS closure, the convective adjustment is mostly redundant and can occasionally degrade results. Overall, the GLS scheme without any convective adjustment provides the most accurate representation of the mixed layer depth as well as the vertical structure and variability both at basin scale and in key regions of deep and intermediate water formation.

1 Introduction

25 The Mediterranean Sea is a semi-enclosed basin characterized by intense thermohaline variability, strong mesoscale activity, and complex interactions with the atmosphere and surrounding land masses. Its dynamical features, including dense water formation, flux exchanges through narrow straits, and sub-basin circulations, make it a particularly challenging environment for ocean modeling (e.g., Pinardi and Masetti, 2000; Millot and Taupier-Letage, 2005; Testor et al., 2018). The Mediterranean Sea plays a crucial role in shaping the regional climate (e.g. Pinardi and Navarra, 1993; Lionello et al., 2006; Pinardi and
30 Masetti, 2000) and represents a natural laboratory for investigating key ocean processes with implications at the global scale.

In this sense, it is often considered a “miniature ocean” (e.g., Bethoux et al., 1999), where processes such as deep and intermediate water formation, strait exchanges, mesoscale variability, and air–sea interactions can be studied in a confined environment as a small-scale analogue of global processes.

35 Over the past two decades, regional ocean models have been widely employed to simulate the Mediterranean circulation at increasing spatial and temporal resolution. Among these, the physical component of the Mediterranean Forecasting System, developed within the framework of the Copernicus Marine Service, represents a state-of-the-art implementation (e.g., Tonani et al, 2009; Clementi et al., 2017; Coppini et al., 2023). It provides regular, open-access analyses, forecasts (Clementi et al., 2023) and reanalyses (Escudier et al., 2021) of the basin’s physical state and serves as a reference framework for scientific and
40 operational applications.

Due to high computational costs and missing physics, the spatial and temporal resolution of current hydrostatic ocean circulation models, even at regional scale, is unable to explicitly resolve the inherently non-hydrostatic and small-scale turbulent vertical mixing and convective processes. To this end, vertical mixing closure schemes – either based on simplified
45 formulations (e.g., Pacanowski and Philander, 1981) or on more physically based models (e.g., Gaspar et al., 1990; Umlauf and Burchard, 2003) – are employed to mimic these turbulent processes. Also, deep convection, which corresponds to the formation of dense deep water masses, has to be represented through parameterizations to reproduce its timing, intensity, and spatial extent realistically (e.g., Marshall et al., 1999; Villarreal et al., 2005; Herrmann et al., 2008; Luneva et al., 2019). Note that convection causes turbulent vertical mixing under unstably stratified conditions and can be parameterized separately
50 (convective adjustment) or within the same turbulence closure scheme (Umlauf and Burchard, 2003; Legay et al., 2025).

Physics-based turbulent closure schemes include buoyancy production terms and are, in principle, able to represent convective mixing when the stratification becomes unstable. However, in practice these schemes may not always remove static instability rapidly enough, particularly at the temporal and vertical resolution typically used in regional ocean models. This can lead to
55 an underestimation of convective mixing. For this reason, ocean circulation models usually combine turbulence closure schemes with a convective adjustment parameterization (e.g. Clementi et al., 2017; Coppini et al., 2023). The convective adjustment acts by stabilizing the water column through a rapid vertical redistribution of tracers, ensuring a prompt restoration of stable stratification. In this sense, it does not represent an independent physical process, but rather a numerical parameterization designed to mimic the fast overturning associated with unresolved convective events.

60

This combined approach has been widely adopted in the modelling of the Mediterranean Sea for several decades (Pinardi et al., 2003; Pinardi and Coppini, 2010), mostly because the relatively simple turbulence closure schemes traditionally employed were not sufficient to adequately represent convective mixing. In this context, it is essential to assess how these earlier, less complete schemes compare with more advanced formulations which are specifically designed to provide a more consistent

65 and physically based representation of turbulent and convective processes. Depending on the turbulence closure scheme adopted, the use of an explicit convective parameterization may or may not be required. Assessing this aspect is one of the main objectives of the present study.

70 The choice of mixing scheme and convective adjustment parameterisations has been shown to play a crucial role in shaping the circulation, the temperature and salinity distributions and deep-water formation (DWF) processes (e.g., Herrmann et al., 2008; Refray et al., 2015; Luneva et al., 2019; Robertson et al., 2019; Gutjahr et al., 2021). In particular, vertical mixing controls the redistribution of heat and salt throughout the water column, thereby influencing stratification and mixed layer depth (MLD). Convection, on the other hand, strongly affects deep and intermediate water formation, which in turn impacts large-scale circulation patterns and water mass properties. Since these processes are tightly coupled to air–sea exchanges and mesoscale dynamics, model sensitivity to the choice of mixing and convection schemes can propagate across a wide range of spatiotemporal scales, ultimately influencing the realism of simulated ocean variability.

Only a few previous studies provide a context for understanding the performance of different mixing schemes, but none of these explicitly investigate their combined effect with a convective adjustment, and whether a mixing scheme alone is sufficient to reproduce the vertical structure and variability of the upper ocean. Madec et al. (1991) combined Richardson-number dependent closures with nonpenetrative convective adjustment in studies of deep-water formation in the Northwestern Mediterranean Sea. Refray et al. (2015) investigated the sensitivity of turbulent vertical mixing using a 1-D ocean circulation model at the global scale, comparing various mixing schemes but without including convective adjustment. Gutjahr et al. (2021) compared various global-scale mixing schemes – including among the others, Richardson-number dependent closures and Turbulent Kinetic Energy (TKE) – but did not assess the role of convective adjustment. Storto et al. (2023) focused on the Mediterranean Sea, comparing the Generalized Length Scale (GLS) and TKE schemes, but did not examine cases of active convection and therefore did not evaluate the potential impact of including a convective adjustment. By explicitly investigating the interplay between vertical mixing schemes and convective adjustment, the present study addresses a gap left by these prior works and demonstrates how convective adjustment modulates the behaviour of different schemes in the context of the Mediterranean Sea.

The aim of this study is to investigate the sensitivity of a modified configuration of the Mediterranean Forecasting System (Clementi et al., 2023) to the choice of vertical mixing scheme and to the inclusion of an explicit convection adjustment parameterization. Model performance is evaluated across spatial and temporal scales through an intercomparison of temperature, salinity and mixed layer depth, from basin scale over years to deep and intermediate water formation regions ranging from months to hours. Given that the formation of deep and intermediate waters in the Mediterranean occurs on seasonal timescales, with interannual variability, a two-year model integration is sufficient to capture the main differences between experiments.

The rest of the paper is structured as follows. Section 2 describes the region and the observational dataset used to assess our model performances (Subsection 2.1). It also describes the model configuration (Subsection 2.2), with a focus on the choice of the vertical mixing schemes and convective adjustment, and the experimental design performed in this study (Subsection 2.3). Section 3 presents the main results of this study. We first present the convective adjustment (Subsection 3.1). Then we show results at the basin scale over the entire two-year period (Subsection 3.2). We focus then on deep and intermediate water formation regions over months to days (Subsection 3.3), and we investigate summer daily convection events at hourly temporal scales (Subsection 3.4). Finally, Section 4 discusses the key conclusions and outlines perspectives for future work.

2 Data and methods

2.1 Argo float data in the Mediterranean Sea

We use Argo float data (Wong et al., 2020; Poulain et al., 2007) from the Copernicus Marine Service Global Ocean near real-time in situ quality controlled observational product (Global Ocean-In-Situ Near-Real-Time Observations, <https://doi.org/10.48670/moi-00036>) as a reference observational dataset for our model validation. The location of Argo floats in the Mediterranean Sea during the two-year period under investigation (2020–2021) is reported as red dots in Fig. 1a. This dataset consists of 11,053 Argo profiles across the Mediterranean Sea, which allows for a statistically robust analysis of the processes under investigation while ensuring homogeneous quality control. Except for a few regions – such as the North Adriatic Sea, the Aegean Sea, the Strait of Sicily and the Gulf of Sirte – the Argo data in this time period provide good spatial coverage across the major part of the basin, offering a statistically meaningful dataset for our model validation.

Typical temperature and salinity profiles retrieved by Argo floats in the Mediterranean Sea are shown in Fig. 1 as monthly average vertical profiles of temperature and salinity in January 2021 (Fig. 1b) and July 2021 (Fig. 1c). The observations exhibit strong variability around the mean both in temperature and salinity, especially in the upper ~200 m, highlighted by shading representing one standard deviation of the dataset. It is known that, both temperature and salinity show also a pronounced seasonal variability (e.g., Pinardi and Masetti, 2000). In winter (Fig. 1b) stratification is weakened because of surface cooling and increased vertical mixing, resulting in the formation of a thick mixed layer with relatively uniform temperature and salinity. In summer (Fig. 1c), temperature and salinity values close to the surface become higher with respect to winter as the result of strong heating and evaporation. These processes inhibit vertical mixing and give rise to a well-defined thermohaline stratification. As a result, vertical profiles are more homogeneous in winter and strongly stratified in summer.

The Mediterranean Sea hosts several key regions of deep and intermediate water formation, including the Gulf of Lion, the South Adriatic Sea, the Rhodes Gyre, and the Aegean Sea (see map in Fig. 1a). In these areas, vertical mixing and convection

play a central role in preconditioning and driving dense water formation processes by modulating stratification and contributing to surface buoyancy loss. Fig. 1 also shows the seasonal variability of temperature and salinity profiles from Argo floats for two of these regions: the South Adriatic Pit (Fig. 1d) and the Rhodes Gyre region (Fig. 1e). In both regions, temperature profiles exhibit a pronounced seasonal variability in the upper layers, especially near the surface, closely following the seasonality of the surface heat fluxes. On the other hand, salinity shows distinct seasonal changes associated with deep and intermediate water formation processes. In the South Adriatic Pit (Fig. 1d), the winter period (i.e., December-January-February) is influenced by the intrusion of intermediate water formed in the Levantine basin, visible as a marked salinity increase between about 100 m and 200 m. Deep-water formation is not clearly visible, being largely smoothed by seasonal averaging, reflecting its short duration (order of weeks) and spatially localized nature. In the Rhodes Gyre region (Fig. 1e), where intermediate water forms, salinity shows a persistent increase throughout the year. The salinity maximum is located at about 150 m in winter, during the formation phase, and deepens to about 200 m in spring (i.e., March-April-May) and to about 250 m in summer (i.e., June-July-August) and autumn (i.e. September-October-November).




Figure 1: (a) Model domain including the Mediterranean Sea and an Atlantic box, with bathymetry as a background colour and labels for the main subbasins and deep and intermediate water formation regions. Red dots indicate Argo floats during 2020–2021. The black zonal line at 37.5°N marks the location of a vertical section used throughout the paper. Middle panels show the basin-averaged observed temperature (blue) and salinity (orange) for (b) January 2021 and (c) July 2021, where shaded areas represent one standard deviation. Bottom panels show the season-averaged observed temperature (blue) and salinity (orange) in (d) the South Adriatic Pit and (e) the Rhodes Gyre region.

2.2 Model setup

Our model setup builds on the Mediterranean Forecasting System model of the Copernicus Marine Service. The model domain encompasses the entire Mediterranean Sea and extends into a portion of the Atlantic Ocean (Fig. 1a), spanning from 18° 7.50' W to 36° 17.50' E in longitude and 30° 11.25' N and 45° 58.75' N in latitude. Including an Atlantic box is essential to accurately resolve water exchanges between the Mediterranean Sea and the Atlantic Ocean through the Strait of Gibraltar, as the inflow of fresher Atlantic water and the outflow of saltier Mediterranean water strongly influence basin-wide salinity, temperature, and density structure, as well as DWF. Nevertheless, the analysis performed in this study focuses only on the Mediterranean Sea.

The model system consists of a two-way coupled hydrodynamic circulation model and a third-generation spectral wind wave model. The ocean hydrodynamic circulation model is based on Nucleus for European Modelling of the Ocean (NEMO; Madec et al., 2023, [doi:10.5281/zenodo.6334656](https://doi.org/10.5281/zenodo.6334656)) version 4.2 and the wind wave model on WaveWatch III (WW3; Tolman et al., 2021) version 6.07. The coupling happens hourly online with the exchange of air-sea temperature difference and surface currents from NEMO to WW3, and of neutral drag coefficient used to evaluate the surface wind stress from WW3 to NEMO (Clementi et al., 2017).




Figure 2: Vertical layer thickness as a function of depth. A zoom of the upper 50 m highlights the fine vertical discretization within the mixed layer.

The model has a horizontal resolution of $1/24^\circ$ (about 4 km) in both latitude and longitude, and it includes 141 non-uniformly distributed vertical levels. The vertical layer thickness varies with depth (Fig. 2), with particularly fine resolution in the upper ocean (from about 2.1 m near the surface to 4.6 m at 50 m depth). This dense discretization in the mixed layer allows for an accurate representation of key upper-ocean processes, such as mixing and stratification, which are essential for capturing surface-driven dynamics. The time stepping is achieved within a leapfrog differencing structure associated with an Asselin filter and the baroclinic time step is set to 180 s. To capture the effects of intensified vertical mixing induced by internal wave breaking in the Strait of Gibraltar, particularly near Camarinal Sill (Wesson and Gregg, 1994; Hilt et al., 2020), we also prescribe an enhanced vertical diffusivity farther to the east of the sill.

180

Horizontal tracer advection is computed using the fourth-order Flux-Corrected Transport (FCT) scheme, which provides a good balance between accuracy and numerical stability by combining a high-order transport method with a monotonicity constraint (Levy, 2001). Lateral diffusion of tracers is implemented as a Laplacian operator along isoneutral surfaces, with a spatially constant diffusivity. Horizontal momentum advection is treated in flux form using the third-order Upstream-Biased Scheme (UBS). As for tracers, momentum lateral diffusion is applied using a Laplacian operator along isoneutral surfaces, again with constant viscosity. These parameterisations are widely used in NEMO-based configurations for large-scale and regional ocean modelling, providing physically consistent dissipation while preserving the large-scale circulation features (e.g., Madec et al., 2023; Le Sommer et al., 2009).

185

2.2.1 Vertical mixing scheme

In this study, we compare three different parameterisations for vertical mixing, representative of the main families of vertical mixing parameterizations commonly used in ocean modelling. Our aim is to compare more simple approaches – such as the Richardson-number dependent scheme, which is still used in the Mediterranean Forecasting system of the Copernicus Marine Service (Clementi et al., 2017; Coppini et al., 2023) – with more physically based formulations. These physics-based parameterizations rely on a prognostic treatment of turbulence and are designed to capture a wider range of oceanic mixing processes with greater physical accuracy. Physically based vertical mixing schemes typically fall into two categories: one-equation and two-equation turbulence closure models. One-equation schemes solve a prognostic equation for the TKE while relying on diagnostic expressions or empirical assumptions for the mixing length or the dissipation rate (e.g., Gaspar et al., 1990). Two-equation models introduce an additional prognostic variable – usually the turbulence length scale or the dissipation rate of turbulent kinetic energy – which allows for a more flexible and accurate representation of turbulence in stratified and sheared flows (e.g., Mellor and Yamada, 1982; Umlauf and Burchard, 2003). Each of these formulations requires calibrating several parameters. In this study, the best configuration for each scheme was selected from sensitivity tests as the one minimising statistical errors at the basin scale when compared with all available Argo data.

195

200

The first vertical mixing scheme used in this study is a Richardson number–dependent formulation originally proposed by Pacanowski and Philander (1981). This scheme was originally developed for the equatorial ocean, using a rectangular box model, and was not designed to provide prognostic mixing. Vertical eddy viscosity and diffusivity are parameterised as functions of the Richardson number (see Appendix A). The Richardson number expresses the ratio between stratification (buoyancy) and vertical shear, and it quantifies their relative dominance. Low Richardson number values indicate that shear overcomes stratification, promoting instability, while high values correspond to strongly stratification. Accordingly, this scheme increases mixing when shear effects dominate over buoyancy and reduces it when buoyancy prevails. Specific parameters control the sensitivity of the mixing to these conditions, while background values ensure a minimum level of vertical mixing even under stable conditions. The Richardson-dependent mixing parameterization does not simulate convective overturning, but just reduces density gradients by enhancing mixing in unstable layers when shear dominates over buoyancy. See Appendix B (Table B1), for details on the parameters used in the NEMO namelist for the Richardson number–dependent formulation.

The second vertical mixing scheme adopted here is the one-equation TKE scheme as proposed by Gaspar et al. (1990), in which the evolution of the turbulent kinetic energy k is described by a prognostic equation (see Appendix A). The model accounts for the main four physical processes involved in turbulence dynamics: production by vertical shear, destruction by buoyancy effects, turbulent kinetic energy diffusion, and dissipation parameterised following the Kolmogorov formulation (Kolmogorov, 1942). Vertical eddy viscosity A^{vm} is diagnosed from the TKE using a mixing length approach, where the mixing length is parametrized using a diagnostic formulation, as a function of both a prescribed maximum and the local stratification to ensure numerical stability by limiting its vertical gradients. Tracer diffusivity $A^{vT} = A^{vS}$ for temperature and salinity, respectively, is derived from A^{vm} through a constant turbulent Prandtl number ($A^{vT} = A^{vm} / P_t$). At the ocean surface, we impose a fixed minimum surface mixing length. A simplified representation of Langmuir turbulence is also included. The TKE scheme treats convection prognostically through the calculation of enhanced turbulent kinetic energy production in statically unstable layers, which results from the combined contribution of the four terms described above: shear production, stratification or buoyancy destruction, diffusion and dissipation. The resulting enhanced turbulent kinetic energy production increases the eddy diffusivity coefficients and leads to the homogenization of these layers. See Appendix B (Table B2), for details on the parameters used in the NEMO namelist for the TKE closure scheme.

The third vertical mixing scheme used in this work is a two-equation turbulence closure scheme defined within the GLS framework (Umlauf and Burchard, 2003). The GLS schemes are a family of models that differ mainly in the choice of the second turbulence variable to be worked out through a prognostic equation and the stability functions used to parameterise stratification effects on turbulence. In addition to the turbulent kinetic energy k , the second prognostic variable can be defined in different ways, leading to different two-equation turbulence models. In the $k-\varepsilon$ formulation, the second variable is the turbulent dissipation rate ε , which quantifies the rate at which turbulent kinetic energy is dissipated into heat. In the $k-\omega$

formulation, it is the specific dissipation rate $\omega = \varepsilon/k$ (where ε represents the turbulent kinetic energy dissipation rate) which has the dimension of an inverse time and represents a characteristic turbulence frequency. Alternatively, in the $k-k_l$ formulation, the second variable is the turbulent length scale k_l , describing the typical size of the energy-containing eddies, usually expressed as $k_l \simeq \frac{k^{3/2}}{\varepsilon}$.

In this study, we adopt the $k-\varepsilon$ formulation, where the prognostic variables are the turbulent kinetic energy k and its dissipation rate ε (Umlauf and Burchard, 2003), but we also verified that using the $k-\omega$ formulation does not change significantly our conclusions. This model captures turbulence dynamics with enhanced detail, particularly under varying stratification, and is known for its numerical stability and robust performance in oceanic applications (Burchard and Bolding, 2001). The model prognoses the temporal evolution of k and ε , accounting for the same four physical processes considered in the TKE scheme: shear production, buoyancy destruction, vertical turbulent diffusion, and dissipation. Vertical eddy viscosity and diffusivity coefficients are computed dynamically based on k and ε and modulated by stability functions dependent on the gradient of the Richardson number. These stability functions, which are a common feature of turbulence closures, represent the suppression of turbulence by reducing mixing efficiency under stable stratification. In this study, we adopt the stability function formulation version A by Canuto et al., 2001 (*Canuto A*). Specifically, the momentum mixing efficiency decreases as stratification increases, while the turbulent Prandtl number correspondingly increases, implying a stronger reduction of tracer mixing relative to momentum (Canuto et al., 2001; Umlauf and Burchard, 2003). Boundary conditions at the surface and at the bottom incorporate enhancements to turbulence caused by wave breaking and bottom friction, respectively. Wave-induced mixing is not taken into account. The GLS scheme accounts for convection by prognostically calculating turbulent kinetic energy production and simultaneously estimating a turbulent length scale. While the computation of turbulent kinetic energy is conceptually similar to that in the TKE scheme, the key difference lies in the prognostic treatment of the turbulent length scale, which enables not only a dynamical adjustment of mixing intensity, but also a more physically consistent representation of the vertical extent of convective overturning. We also note that, while the mixed-layer depth is correctly estimated under convective conditions, the stratification is falsely prescribed by the GLS closure to be unstable also in the lower half of the convective boundary layer (Burchard and Petersen, 1999). In that region, typically counter-gradient fluxes are present which need to be parameterised by a higher-order closure (Legay et al., 2025). More details on the GLS closure scheme are in Appendix A and the parameters used in the NEMO namelist are reported in Appendix B (Table B3).

2.2.2 Convective adjustment

Convective events are characterized by rapid and intense overturning of the water column, an inherent non-hydrostatic process that cannot be explicitly resolved by hydrostatic ocean models. As a result, convection must be represented through a dedicated parameterization. To better capture convective processes, it is still a common practice in large-scale modelling to apply a

convective adjustment whenever the water column is statically unstable to mimic the rapid vertical homogenization associated with convective overturning and ensures realistic mixed layer deepening. One of the most common ways is to apply the enhanced vertical diffusivity (EVD) parameterisation (e.g., Lazar et al., 1999), widely adopted in ocean modelling (e.g., Rahmstorf, 1993; Lazar et al., 1999; Griffies et al., 2000). In practice, the vertical eddy diffusivity coefficient is increased to a high constant value (typically between 1 and 100 m² s⁻¹) wherever the Brunt-Väisälä frequency squared N^2 becomes negative. In this study, a value of 10 m² s⁻¹ is prescribed for the vertical eddy diffusivity based on previous sensitivity tests. This is also the value currently used in the physical component of the Mediterranean Forecasting System of the Copernicus Marine Service.

Since convection has the effect of effectively homogenizing the upper ocean boundary layer, it can be parameterized by applying a high eddy diffusivity which has a similar effect. Although this procedure does not accurately represent the physics of convection with its narrow and dense downward plumes compensated by a broader upwelling in between, leading to non-local fluxes, it effectively mimics the net impact of convective processes, including the rapid vertical redistribution of properties associated with convective overturning. The increased eddy diffusivity coefficient stabilizes the water column, allowing the model to maintain realistic stratification while enabling the development of large-scale circulation patterns.

When the convective adjustment is applied, the vertical eddy diffusivity results from the combined effects of the vertical mixing closure schemes (whether the Richardson-number-dependent parameterization, the one-equation TKE scheme, or the two-equation GLS formulation) and the convective adjustment implemented through the EVD parameterisation. Since the TKE and GLS schemes implicitly include a convection parameterisation via the buoyancy term in the turbulent kinetic energy equation being positive for unstable stratification with N^2 (equations (A3) and (A6), respectively), some double counting is included in this approach. It should be noted that the GLS mixing parameterisation includes further sensitivity to convectively unstable stratification due to the dependence of the stability functions and the length-scale related equation on N^2 .

2.3 Experimental design

We perform six, two-year long, simulations over the period 2020–2021. These simulations share the same model setup (as described in Section 2.2), and the same input fields and differ only in the choice of the vertical mixing scheme and whether the convective adjustment is applied.

The system is forced at the ocean surface with the atmospheric fields at 1/10° horizontal resolution of the European Centre for Medium-Range Weather Forecasts (ECMWF) made available by the Italian Air Force Meteorological Service. Following the configuration of the physical component of the Mediterranean Forecasting System of the Copernicus Marine Service, at the Atlantic boundary, the hydrodynamic model is nested into the daily analysis data of the Copernicus Marine Global product (whose Copernicus product name is *Global Ocean Physics Analysis and Forecast*, <https://doi.org/10.48670/moi-00016>) at 1/12° horizontal resolution and 50 vertical levels. River runoff in the modelling implementation originates from 39 rivers.

Daily mean data for 38 of these rivers are taken from the European Flood Awareness System dataset (EFAS v5.0; Mazzetti et al., 2023) delivered within the Copernicus Emergency Service, while monthly mean data for the Nile River are taken from a climatology computed over the period 1970–2007 (Said et al., 2011).

Three turbulence closure schemes are tested: the Richardson-number dependent parameterization (Pacanowski and Philander, 1981), the TKE closure scheme (Gaspar et al., 1990), and the GLS closure scheme (Umlauf and Burchard, 2003). These vertical mixing schemes are employed under two different convective setups: with and without the enhanced vertical diffusion parameterization typically used to mimic deep convection events. Therefore, the six simulations performed here differ only in the vertical mixing scheme and in whether convective adjustment is included. This experimental design allows us to assess the interplay between the choice of vertical mixing scheme and the effect of EVD, and to disentangle their individual and combined contributions to the representation of vertical mixing processes. It also enables us to evaluate how different turbulence closure schemes respond to the presence or absence of convective adjustment, and how this affects stratification, heat and salt distribution, and the overall vertical structure of the water column at basin and local scale.


Table 1 summarises the experiments. Our control run is defined as the simulation that employs the Richardson-number dependent vertical mixing scheme in combination with convective adjustment. Keeping convective adjustment active, we then replace the vertical mixing scheme with either TKE or GLS parameterisations. The remaining three experiments use the same three mixing schemes, but with convective adjustment disabled. In the experiments where convective adjustment is disabled, adaptive implicit vertical advection is activated to prevent local numerical instabilities (Shchepetkin, 2015, Madec et al., 2023). All simulations are initialized from the same 31 December 2019 restart fields provided by the Mediterranean Forecasting System. These fields are analysis products that incorporate data assimilation of available observations, including satellite and in-situ measurements. As a result, they provide an optimal representation of the Mediterranean Sea state at the start of the simulations, ensuring that all experiments begin from a physically realistic initial condition. The results presented hereafter are not affected by these initial conditions but rather reflect the changes in the vertical mixing schemes and convective adjustment listed in Table 1.

Simulation name	Vertical mixing closure scheme	Convective adjustment	Notes
Exp-PP ⁺	Richardson-dependent scheme	Enhanced Vertical Diffusion (EVD)	Control simulation
Exp-TKE ⁺	Turbulent Kinetic Energy (TKE)	Enhanced Vertical Diffusion (EVD)	
Exp-GLS ⁺	Generalized Length Scale (GLS)	Enhanced Vertical Diffusion (EVD)	
Exp-PP	Richardson-dependent scheme	–	
Exp-TKE	Turbulent Kinetic Energy (TKE)	–	
Exp-GLS	Generalized Length Scale (GLS)	–	


Table 1: Summary of the experiments conducted in this study.

3.1 Impact of the convective adjustment

The EVD parameterization exerts a strong control on the vertical structure of mixing. Figure 3 shows the eddy diffusivity coefficient along a zonal section at 37.5° N (as indicated in Fig. 1a) averaged over January 2021 for different mixing configurations. The presence of convective adjustment together with different mixing schemes (Figs 3a, 3c, 3e) tends to mask the differences among the various vertical mixing parameterisations resulting in a very similar eddy diffusivity pattern regardless of the chosen vertical mixing scheme. Given the different eddy diffusivity scales observed in the experiments with and without EVD (Figs 3a, 3c, 3e *versus* Figs. 3b, 3d, 3f), it is clear that when EVD is active, the eddy diffusivity produced by the vertical mixing closure is largely overwritten by the EVD parameterization. In contrast, when convective adjustment is disabled (Figs. 3b, 3d, 3f), the vertical eddy diffusivity is determined solely by the mixing schemes and reveals more pronounced differences among schemes. Notably, the Richardson-dependent scheme (Figs. 3b) produces a relatively homogeneous increase in eddy diffusivity within the mixed layer, whereas the TKE (Figs. 3d) and GLS (Figs. 3f) schemes result in more heterogeneous structures. The magnitude of the vertical eddy diffusivity increases going from the Richardson-number dependent scheme (with maximum eddy diffusivity values on the order of 10^{-2} m²/s) to the TKE and then to the GLS schemes (with maximum eddy diffusivity values on the order of 10^{-1} m²/s). Moreover, when the EVD scheme is active, the maximum value of the vertical eddy diffusivity consistently occurs near the surface, while, when EVD is disabled, its maxima are typically found within the mid-depth of the mixed layer, which is more realistic due to the suppression of turbulent length at the top and the bottom of the mixed layer. This distinction underscores the role of EVD in favouring a surface-intensified mixing structure as opposed to the more symmetric and realistic profile observed in its absence, potentially impacting the MLD.



355 **Figure 3: Vertical eddy diffusivity coefficient along the zonal section at 37.5°N (black line in Fig. 1a) averaged over January 2021 for different model configurations: (a) Richardson-dependent mixing scheme with EVD activated, and (b) without EVD activated; (c) TKE closure scheme with EVD activated, and (d) without EVD activated; (e) GLS scheme with EVD activated, and (f) without EVD activated. The black contours in each panel denote the depth of the mixed layer, while grey shading indicates land. Note the different eddy diffusivity scales for the experiments with and without EVD.**



360 **Figure 4: Map showing the maximum depth reached by the EVD parameterisation when activated in the control experiment (Exp-PP⁺ in Table 1) during a summer day (1 July 2021). White areas indicate regions where EVD is inactive.**

The EVD parameterisation is primarily intended to represent winter convection. However, convection-like vertical mixing occurs year-round, even outside the main winter season, and therefore the EVD parameterisation is active even in summer in some regions of the Mediterranean Sea. Figure 4 illustrates this by showing the maximum depth reached by the EVD on a summer day (1 July 2021) in the simulation using the Richardson-number dependent parameterization. White areas indicate the few regions where EVD is not applied on that day. Elsewhere the EVD is applied down to depths ranging from a few meters to several tens of meters reaching its maximum depth in the eastern Mediterranean Sea, and notably in the semi-permanent Ierapetra eddy region, located southeast of Crete within the Rhodes Gyre region (as defined in Fig 1a). This activity is typically weaker and shallower than winter convection, but it still contributes to reshaping the vertical dynamics, as discussed in the following sections.

3.2 Assessment at the basin scale

3.2.1 Impact on tracers

Among the different experiments (Table 1), the simulation using the Richardson-number dependent vertical mixing scheme combined with a convective adjustment (Exp-PP⁺) is taken as the control run. This configuration for the vertical mixing and

convection corresponds to the current setup of the Mediterranean Forecasting System of the Copernicus Marine Service (Clementi et al., 2023).

To evaluate the performance of our control simulation and identify systematic deficiencies of the model, we compare the model
380 outputs in terms of temperature and salinity with all available Argo data in 2020-2021 (Fig. 1a). We verified (not shown) that
removing a spin-up period leads only to negligible differences and does not affect the results. Following the extensive literature
on error decomposition for model validation (e.g., Willmott, 1981; Murphy, 1988; Murphy, 1992; Oke et al., 2002; Oddo et
al., 2022), we define four key statistical metrics, which provide complementary insights into model skills: the root mean
squared error (RMSE), the unbiased root mean squared error (uRMSE), the mean bias (MB), and the cross-correlation
385 coefficient (CC). For more details on their definition, see Appendix C. We compute these four metrics averaging on the entire
basin and over the entire two-year period.

The temperature error profiles are shown in Fig. 5 as blue profiles. All metrics exhibit two pronounced decreases in model
skill, one in the upper 50 m within the mixed layer and another between 200 and 600 m, corresponding to the Levantine
390 Intermediate Waters (LIW) depth range (e.g., Bryden and Stommel:1982; Robinson and Golnaraghi, 1993; Lascaratos et al.,
1993; Taillandier et al., 2022). Within the mixed layer, RMSE and uRMSE peak at 25–30 m, corresponding to these two-year
average MLD (30.8 m for observations and 28.3 m in the control simulation – dashed horizontal lines in Fig.5). MB reaches
its maximum around the middle of the MLD (about 15 m). CC exhibits a monotonic reduced decrease throughout the mixed
layer, with a minimum below the yearly average MLD. A similar behaviour occurs within the LIW depth range, with a
395 minimum in CC around 400 m, while RMSE, uRMSE and MB show their largest deviations slightly above, at approximately
350 m. Overall, this evidence indicates that errors are largest in the surface mixed layer but also reflect difficulties in
representing intermediate water masses.

The salinity error profiles (orange) in Fig. 5 show a distinct pattern compared to temperature. RMSE and uRMSE exhibit a
400 single peak confined to the mixed layer, with no significant anomalies at the LIW depth range. Nevertheless, the MB highlights
systematic fresh biases at both depth ranges, suggesting persistent model errors also in salinity in representing the LIW. We
observe that the minimum salinity MB within the LIW depth range is found near the bottom of the LIW layer (about 500 m),
while the minimum temperature MB occurs approximately at mid-depth within the layer (about 350 m). As for temperature,
the CC decreases monotonically within the mixed layer; however, unlike temperature, it exhibits only a weak secondary
405 minimum at LIW depths. These findings suggest that salinity shows smaller discrepancies than temperature with respect to
observations in the representation of intermediate waters, while it also presents a discrepancy with the observation within the
mixed layer.

As stated above, overall, for both salinity and temperature, the shape of the error at the basin scale suggests two main decreases of the model skills: the representation of the properties within the mixed layer and the intermediate water masses. The control simulation performs reasonably well in the deep region, while the more pronounced biases are within the mixed layer. These biases are likely related to the vertical mixing scheme, highlighting the need for a more realistic treatment of atmospheric energy input and vertical redistribution of momentum and buoyancy.




Figure 5: Basin-averaged temperature (blue) and salinity (orange) error profiles for 2020–2021 in the control simulation (Exp-PP⁺ in Table 1). Panels from left to right: root mean squared error (RMSE), unbiased root mean squared error (uRMSE), mean bias (MB), and cross-correlation coefficient (CC). The definitions of these metrics are in Appendix C. Dashed lines represent the two-year basin average values of the MLD for observations and control simulation.

To assess the impact of the different closure schemes and the activation of the convective adjustment at the basin scale, we compare the metrics obtained for the other five simulations (simulation other than Exp-PP⁺ in Tab. 1) with these of the control

run. Figures 6 and 7 show the basin-averaged error difference for temperature and salinity, respectively. For each experiment and each metric (Equations C1–C4), the error difference (err_{diff}) as a function of depth is computed as

$$err_{diff} = err_{mod}(z) - err_{CS}(z) \quad (1)$$

430 where $err_{CS}(z)$ denotes the error of the control run (Exp-PP⁺, see Figs. 5), and $err_{mod}(z)$ is the error associated with each model configuration (experiments other than Exp-PP⁺ in Tab. 1). Here, err generically denotes one of the four metrics (RMSE, uRMSE, MB, or CC) detailed in Appendix C. For the MB_{diff} metric, the absolute value of the mean bias is used for both the control run and the model experiments when computing MB_{diff}.

435 It should be noted that negative values of RMSE_{diff}, uRMSE_{diff}, and MB_{diff} indicate improvement, while positive values indicate deterioration. For CC_{diff}, the interpretation is reversed: higher values (closer to 1) denote improvement, whereas lower values (towards 0) denote deterioration. Accordingly, in Figs. 6 and 7 green and red backgrounds highlight regions of improvement and deterioration relative to the control simulation, respectively.

440 For temperature (Fig. 6), the error profiles indicate that the GLS closure scheme without convection parameterization (solid green line) provides the largest error reduction, especially in terms of MB_{diff}, a result further confirmed by the CC_{diff}. The greatest improvement occurs within the mixed layer, mostly reflecting a reduction in systematic errors (MB_{diff}), while additional improvements at LIW depths are observed in terms of uRMSE_{diff} and CC_{diff}. Adding the convection parameterization to GLS (dashed green line) diminishes these gains at some depths, and mostly for MB_{diff}. The TKE mixing scheme exhibits
 445 intermediate skill: with or without EVD (orange lines), improvements are mostly confined to the mixed layer, while performance below this layer often deteriorates. The Richardson-dependent scheme without EVD (solid blue line) shows the smallest improvement within the mixed layer and a worsening in the LIW layer across all metrics. This behaviour is expected because the scheme estimates turbulence solely from the Richardson number (see Subsection 2.2.1 and Appendix A), that is from the balance between vertical shear and stratification (buoyancy) and does not include convective mixing. Consequently,
 450 it is expected that an additional parameterization would be required to properly represent convection.

For salinity (Fig. 7) in the absence of convective adjustment (solid lines), GLS is the only scheme showing improvements above 100 m for RMSE_{diff}, uRMSE_{diff}, CC_{diff}, and MB_{diff}. At the LIW depths, changes are minimal and a small degradation occurs for all experiments between about 100 and 200 m. The TKE and Richardson-number dependent schemes without EVD
 455 show smaller improvement in all metrics, particularly within the mixed layer, with the Richardson-number dependent scheme (solid blue line) performing worst, especially for CC_{diff}.




Figure 6: Temperature error difference (equation 1) with respect to the control simulation for the four metrics in equations C1–C4. Dashed lines indicate simulations with EVD and continuous lines without EVD. The transparent background color highlights regions of improvement (green) and worsening (red) compared to the control simulation.

460




Figure 7: Salinity error difference (equation 1) with respect to the control simulation for the four metrics in equations C1–C4. Dashed lines indicate simulations with EVD and continuous lines without EVD. The transparent background colour highlights regions of improvement (green) and worsening (red) compared to the control simulation.

3.2.2 Representation of the Mixed Layer Depth

Given that the largest impact of the interplay between the mixing closure parameterisation and the convective adjustment occurs within the mixed layer, we also investigate how well the different schemes reproduce the MLD across the Mediterranean Sea. Figure 8 shows the time evolution of the MLD for the years 2020–2021, as computed from all available Argo temperature and salinity observations (black dashed line) and from model outputs (coloured lines) at the corresponding locations. Notably, for each day, model temperature and salinity profiles are first sampled at each Argo location. Density and MLD are then computed individually for each profile in both models and observations, and a basin average MLD is finally calculated.

475 The MLD for both observations and model outputs is computed using the density threshold method, which identifies the depth
at which the potential density increases by a fixed amount relative to a near-surface value. Specifically, here, the MLD is
defined as the depth where the potential density exceeds that at 10 m by 0.01 kg m^{-3} . With this definition, a minimum MLD of
10 m is imposed, as the reference density is taken at that depth. This approach, commonly used in oceanographic studies,
480 2004).

The temporal evolution of the basin-averaged MLD over the period 2020–2021 oscillates around a temporal-spatial average
value of 31 m and exhibits a pronounced seasonal cycle, with deepening during winter and shoaling in summer. On average,
the MLD reaches its maximum around 70–80 m in winter (December to March), when surface cooling and intense wind forcing
485 promote vertical convection and deep mixing. In spring, the increasing solar radiation and weakening winds lead to rapid
surface warming and stratification, resulting in a marked shoaling of the MLD. By summer (June to September), the MLD
stabilises at its shallowest levels, below 20 meters, reflecting strong thermal stratification and limited vertical mixing. In
autumn, cooling and more frequent wind events gradually deepen the MLD again, preparing the system for winter convection.

490 All simulations, colour-coded according to model configuration, reproduce this seasonal pattern, with winter maxima
exceeding 70 m and summer minima around 15 m. However, significant differences in the MLD emerge while employing
different vertical mixing schemes and varies also significantly depending on whether the EVD convection parameterisation is
activated or not.

495 Simulations including EVD (blue, orange, and green lines) behave in a similar way and very close to the observations (black
line). They occasionally show deeper (e.g. February–March 2020) or shallower (e.g. March 2021) winter MLDs and overall,
they exhibit a good agreement in summer. With EVD activated, the two-year average MLD with TKE (30.9 m) and GLS (31.3
m) are very close to the observed one (30.8 m), while the Richardson-number dependent scheme tends to underestimate the
MLD especially in summer (average of 28.3 m).

500 Simulations without EVD (red, purple, and brown lines) show distinct behaviours. The Richardson-dependent scheme (red
line) produces the shallowest winter MLD and a low average MLD (average of 25.7 m), significantly underestimating observed
deep mixing even in summer. The TKE scheme (purple line) tends to underestimate the MLD, both in winter and in summer.
Although it performs better than the Richardson-number dependent scheme yet still fails to capture the basin-average
505 maximum depths seen in observations (average MLD of 25.7 m). On the other hand, the GLS scheme (brown line) yields
winter and summer MLDs closely aligned with observations, remaining in good agreement to them throughout the year. With
the GLS, the average value of MLD without convective adjustment (32.7 m) is very close to the one obtained with convective
adjustment included (31.3 m).

The GLS scheme without convective adjustment reproduces the observed deepening of the mixed layer, while the TKE and the Richardson-dependent scheme do not. This evidence demonstrates that this scheme does not need an additional external convection parameterisation to reproduce realistic mixed layer deepening. This may be attributed to the two-equation structure of the GLS (see Section 2.2.1 and Appendix A): the first equation evolves the turbulent kinetic energy over time, indicating where mixing is possible, while the second equation dynamically adjusts the turbulence length scale, controlling how deeply mixing penetrates. In contrast, the TKE scheme lacks dynamic depth control, and the Richardson-dependent scheme does not even include prognostic turbulent kinetic energy; in both cases, the penetration of convective mixing is limited. It is worth noting that, in the absence of explicit convective adjustment, the additional physics included in the TKE scheme compared to the Richardson-dependent formulation appears to have only a minor impact, likely because, at these scales, TKE still provides a relatively simplified representation of convective processes.




Figure 8: Time evolution of basin-averaged MLD for varying vertical mixing schemes, with and without convection parameterisation (configurations listed in Table 1). The black dashed line represents the MLD from Argo observations. A 7-day moving average has been applied to both model outputs and observations.

The Mediterranean Sea hosts several key sites of deep and intermediate water formation, including the Gulf of Lion, the South Adriatic Sea, the Rhodes Gyre area and the Aegean Sea (e.g., Lascaratos et al., 1999; Simoncelli et al., 2018; Pinardi et al., 2019; Pinardi and Masetti, 2000; Pinardi et al., 2023). In these regions, vertical mixing and convection play a fundamental role in preconditioning and driving dense water formation processes by contributing to surface buoyancy loss and erosion of stratification (e.g., MEDOC group, 1970; Schroeder et al., 2012). The intensity and vertical structure of turbulent mixing directly influence the transformation and ventilation of water masses, making its accurate representation essential for modelling the basin-scale thermohaline circulation (e.g., Simoncelli et al., 2018; Pinardi et al., 2019; Somot et al., 2006).

As case studies, to further investigate the impact of the interplay between different vertical mixing schemes and convective adjustment, we select one site of DWF (the South Adriatic Pit) and one site of both deep and intermediate water formation (the Rhodes Gyre region) and analyse them across the various model configurations during two selected time windows in 2021. We note that after more than one year of integration, although all six experiments are initialized from the same initial conditions, the six simulations have already had time to substantially diverge by the beginning of 2020. Consequently, while this section focuses on the behaviour of the schemes within two selected time windows in 2021, the results inherently reflect the prior evolution and memory of the system under each parameterization. This approach allows us to assess the resulting quasi-equilibrium state, accounting for how each scheme steers the model toward a distinct numerical and physical regime through its cumulative effects over time and its interplay with the convective adjustment parameterization. Therefore, our objective in this section is not to evaluate turbulence closure schemes in isolation, but rather to investigate their integrated impact in deep and intermediate water formation regions, within the general circulation of the Mediterranean Sea.

545

3.3.1 Deep water formation and intermediate water intrusion in the south Adriatic Pit

The South Adriatic Pit (defined as in Simoncelli and Pinardi (2018) between longitudes 17°–19°E and latitudes 41°–42.5°N and shown in Fig. 1a) reaches a depth of 1200 m and is connected to the Eastern Mediterranean Basin through the Strait of Otranto. It is one of the principal sites of DWF in the Mediterranean Sea, where intense cooling and evaporation during winter – especially in February–March – induce strong density increases that drive dense water sinking and convective overturning (e.g., Wüst, 1961; Gačić et al., 2001; Manca et al., 2003). During these events, the MLD can deepen substantially, typically reaching between 600 and 800 m (e.g., Manca et al. 2003; Pinardi et al., 2023).





Figure 9: (a) Temperature and (b) salinity model profiles compared to observations (black dashed line) from an Argo float on 25 February 2021. Modelled profiles are color-coded by different vertical mixing schemes, with and without convective adjustment as indicated in the legend. The inset in the left panel shows the position of the Argo float.

In Fig. 9 we show the temperature and salinity profiles as recorded by an Argo float (black dashed line) on 25 February 2021 in comparison with the modelled profiles at the same location for varying vertical mixing schemes and activating (dashed line) or deactivating (continuous line) the EVD. The location of the profile is shown in the insert of Fig. 9a. Argo observations show a nearly uniform layer in the upper ~500 m, followed by a sharp gradient below. This inflection point corresponds to the MLD. Among the models in which EVD is active, the simplest Richardson-dependent scheme performs best, although it underestimates the MLD in the salinity profile of about 50 m. In this case, the GLS scheme underestimates the MLD in both temperature and salinity by about 100 m. In the absence of EVD, we observe a similar behaviour across different mixing scheme, with a broad inflection point corresponding to MLD around 500 m. To investigate how the various configurations of mixing and convection lead to the situation shown in Fig.9, we examine the time evolution of the salinity field at this location up to the day of the profiles in Fig. 9. Salinity was chosen because it largely reflects the imprint of deep and intermediate water (Fig. 1d).

570 In Fig. 10, we show the Hovmöller diagrams of salinity evolution at the site described above from 1 to 25 February 2021. The last day of each Hovmöller corresponds to the day of the profile shown in Fig. 9. The three panels on the left (Fig. 10a, 10c, 10e) show the salinity evolution for the models with EVD activated. The three panels on the right (Fig. 10b, 10d, 10f) correspond to the models with EVD deactivated. In the upper 100 m, the models with EVD activated show similar behaviour, with fresh water persisting until 13 February, whereas without EVD the models show large variability due to a different
575 activation of mixing. In particular, the TKE scheme (Fig. 10d) maintains relatively fresh water up to 25 February, while the Richardson-number dependent scheme (Fig. 10b) and the GLS scheme (Fig. 10f) show increased salinity. However, the most notable differences occur at depth, between approximately 100 and 400 m, where a tongue of salty LIW gradually intrudes and a convective event, initially confined to the surface after 13 February, gradually deepens into intermediate layers. The contrast between simulations with and without EVD is substantial.

580
In the presence of EVD, convection triggered after 13 February erodes the intruding LIW, so that the LIW is completely mixed thereafter. As shown in Sections 2.2.2 and 3.1, when the convective adjustment is applied, the vertical eddy diffusivity resulting from the mixing schemes, is abruptly replaced with a high constant value, disregarding the temporal evolution of the tracers. This is evident after 13 February regardless of the mixing scheme (Fig. 10a, 10c, 10e), with the salinity field homogenized
585 across both time and depth. Consequently, convection parameterized through EVD does not account for prior stratification: the method is applied uniformly, effectively erasing pre-existing or intruding structure in the region.

Conversely, in the cases without EVD, the LIW persists up to 25 February. In particular, in the GLS scheme (Fig. 10f) the LIW gradually sinks to about 400 m due to its higher density (similarly to what shown in Fig. 1d), exhibiting a more realistic
590 behaviour, whereas in the Richardson-number-dependent scheme (Fig. 10d) it drops abruptly, and in the TKE scheme (Fig. 10b) it remains at roughly the same depth. This behaviour is consistent with the formulation of the mixing schemes. With the GLS scheme (Fig. 10f), which includes a prognostic equation for the mixing length (Section 2.2.1 and Appendix A), both the intensity and depth of convective events appear to be more realistically represented. On the other hand, with the TKE scheme (Fig. 10d), the convective intensity is similar, but the depth of the event is underestimated. For example, on 21 February, the
595 salinity intrusion extends below 300 m with GLS, whereas with TKE the intrusion already begins at 200 m. In the Richardson-dependent formulation (Fig. 10b), the convective event is weaker and more abrupt, producing a less gradual penetration into deeper layers.



600 **Figure 10: Hovmöller diagrams showing the temporal evolution of salinity at the location indicated in Fig 9a (inset) from 1 to 25 February 2021 for different model configurations: (a) Richardson-dependent mixing scheme with EVD activated, and (b) without EVD activated; (c) TKE closure scheme with EVD activated, and (d) without EVD activated; (e) GLS scheme with EVD activated, and (f) without EVD activated. The last day in each panel corresponds to the day of the profile shown in Fig 9.**

3.3.2 Deep and intermediate water formation in the Rhodes Gyre region

605 As an additional case study, we investigate the capability of the model to reproduce deep and intermediate water in the Rhodes Gyre region, defined here between longitudes 26.3°–31°E and latitudes 33°–36.1°N and shown in Fig.1a. This region, located in the eastern Mediterranean Sea between Rhodes and Cyprus and extending southward into the open Levantine Sea, hosts the Rhodes Gyre and is the key site for the formation and initial spreading of the LIW. Relative to the domain defined by Simoncelli and Pinardi (2018), here the region is expanded by 1° to the east and 1° to the south in order to better resolve the early spreading
 610 of the LIW. This water mass, characterised by high salinity and intermediate depth (approximately 200–500 m), begins

spreading within the defined region and subsequently extends further across the Mediterranean Sea, influencing basin-scale thermohaline circulation (e.g., Lascaratos et al. 1993; Kubin et al., 2019; Pinardi et al., 2023).

615 In Fig. 11a, salinity observations from all available Argo floats in the Rhodes Gyre region between January and April 2021 are compiled and arranged to form a Hövmoller diagram. Water with salinity exceeding 39.2 psu is observed above 200 m until approximately mid-February, when a convective event forces the high-salinity layer downward, resulting in a deepening of the entire water column. The LIW, with salinities ranging from 38.95 to 39.05 psu (Lascaratos et al., 1993), descends from approximately 300 m to depth below 400 m.

620 In Figs. 11b–11g we present Hovmöller diagrams of salinity for each model configuration. Overall, the model reproduces the observed stratification well and generally reach similar maximum salinity values, with some differences for varying mixing scheme and depending on the presence or absence of convective adjustment. In the upper 100 m, prior to the mid-February convective event, all models tend to overestimate salinity relative to observations, whereas after the event they generally produce fresher waters. At greater depths, model layers are generally narrower and more confined than in the observations,
625 where post-convective mixing causes substantial vertical widening of the layer.

When EVD is applied (Figs. 11b, 11d, 11f), the convective event is weakly reproduced. The downward displacement of the water column is most evident in the TKE scheme (Fig. 11d), producing a deepening of approximately 100 m, whereas in the Richardson-dependent scheme (Fig. 11b) and GLS scheme (Fig. 11f), the column exhibits almost no abrupt shift, showing
630 only gradual deepening over time. Overall, EVD suppresses the differences among the mixing schemes and results in a very similar salinity response to convection.

In the absence of EVD (Fig. 11c, 11e, 11g), the salinity response differs. With Richardson-dependent scheme (Fig. 11c), the behaviour is very similar to the case with EVD (Fig. 11b), with progressive deepening of the column without a sudden
635 convective shift. This evidence reflects the fact that the Richardson-dependent scheme does not explicitly model convection but only increases mixing when shear dominates over buoyancy (Section 2.2.1). Consequently, convective events are weakly represented: the water column deepens gradually, and high-salinity layers remain confined and less vertically spread compared to observations or schemes that explicitly account for convection. By contrast, the TKE (Fig. 11e) and the GLS (Fig. 11g) schemes alone capture the mid-February convective event, producing a rapid downward displacement of the entire water
640 column, with the GLS scheme better representing the intensity and extend of deepening due to its prognostic treatment of turbulent kinetic energy while simultaneously estimating a turbulence length scale.




Figure 11: (a) Hovmöller diagram of salinity observations from available Argo floats in the Rhodes Gyre region (Fig. 1a) between January and April 2021. (b–g) Hovmöller diagrams of salinity for different vertical mixing schemes and convective adjustment settings: (b) Richardson-dependent mixing scheme with EVD activated, and (c) without EVD activated; (d) TKE closure scheme with EVD activated, and (e) without EVD activated; (f) GLS scheme with EVD activated, and (g) without EVD activated. All panels share the colour scale in panel (a).

3.4 Assessment of the nighttime summer convection daily cycle

650 In summer, the upper layers of the ocean exhibit a pronounced diurnal cycle in temperature, primarily controlled by the alternation of solar heating during the day and cooling processes at night. Surface waters warm under strong solar radiation, leading to a shallow, well-defined thermal stratification, while nocturnal cooling can erode this stratification. These processes occur within the mixed layer, which remains relatively shallow, as the rapid diurnal cycle of formation and destructing does not allow it sufficient time to deepen and night cooling is not as strong as in winter. Consequently, these processes are expected
655 to be strongly influenced by the selected mixing scheme and the presence of a convective adjustment, which can significantly modify both the intensity of the heating and the depth of the near-surface warm layer. As shown in Fig. 4, the EVD during a summer day remains typically around an average of a few meters, but in certain areas – such as the Ierapetra semi-permanent gyre area within the Rhodes Gyre region as defined in the previous section – it can reach tens of meters.

660 The simulations in Tab. 1 were performed again starting from 1 July 2021 for three days, producing hourly outputs. The same initial conditions were used for all simulations, with the goal of monitoring how the mixing closure scheme and the presence of convective adjustment influence the daily cycle. As a case study, we focus on a site (27.5°E , 34.4°N) within the Rhodes Gyre region where the EVD reaches its greatest activation depth during summer (Fig. 4).

665 Figure 12 shows Hovmöller diagrams of the eddy diffusivity coefficient at that location in the upper 50 m for the six simulations. The upper panels correspond to the three simulations with EVD activated, whereas the lower panels show the cases without EVD. In general summer convective events tend to occur during nighttime because of surface cooling associated with absence of solar heating and extend along the water column within the upper few tens of meters (~ 30 m in Fig. 12). As already observed in Fig. 3, when EVD is activated, the eddy diffusivity coefficient produced by the vertical mixing scheme is
670 largely overwritten by the EVD parameterization (note the three-order-of-magnitude difference between the upper and lower panels in Fig. 12), resulting in similar mixing patterns regardless of the turbulence closure scheme used.

In contrast, when EVD is not applied (bottom panels in Fig.12), the representation of convection differs substantially among the three closure schemes. The Richardson-number dependent scheme produces weak but nearly persistent convective activity
675 that remains present even during daytime. The TKE and GLS schemes instead exhibit a more intermittent pattern of convection, with events occurring less frequently and the strongest one during nighttime. Convective events simulated with the TKE scheme are generally weaker than these produced by the GLS scheme and tend to extend less deeply in the water column.

For example, during the night of 1 July 2021, a convective event reaching approximately 25 m depth occurs in all simulations.
680 In three cases with EVD (upper panels in Fig. 12), this event is represented very similarly and homogeneously throughout the upper 25 m, whereas without EVD (lower panel in Fig. 12) its amplitude and vertical structure differ substantially among the

three closure schemes. In the Richardson-number dependent case, it appears as a relatively weak and vertically homogeneous event down to 25 m. In the TKE simulation, it is confined between a bout 10 and 20 m, with a peak around 15 m. In contrast, the GLS scheme produces a much stronger event, extending all the way down to 25 m.

685




Figure 12: Hovmöller diagrams of the hourly eddy diffusivity coefficient between 1 and 3 July 2021 (upper 50 m) for the six simulations with (a) Richardson-number dependent scheme with EVD, (b) TKE with EVD, (c) GLS with EVD, (d) Richardson-number dependent scheme without EVD, (e) TKE without EVD, (f) GLS without EVD.

690

These differences in the representation of summer convection, as seen in Fig. 12, both in terms of amplitude and vertical distribution along the water column, have a direct impact on the temperature field. In particular, the variability in convective activity simulated by the different turbulence closure schemes, and the homogenizing effect of EVD, translates into notable differences in the upper-ocean temperature structure. Figure 13 shows the temperature profiles at that site during the same

695

The first profiles (lightest colours) show that at the beginning – on 1 July 2021, between midnight and 1 a.m. – the temperature profiles are well mixed and very similar to each other, with differences that are barely distinguishable, and are not yet influenced by the vertical mixing schemes or convective adjustment. Later the same day – between 6 p.m. and 7 p.m. – we observe that, while the simulations with TKE (orange) and GLS (green) behave in a similar way both with and without convective adjustment, the Richardson-number-dependent scheme (blue) is strongly influenced by the presence or the absence of EVD. In fact, in the case of TKE and GLS, the simulations without EVD (solid lines) produce temperature profiles that are

700

only slightly different than those with EVD (dashed lines). By contrast, the Richardson-number dependent scheme without EVD (solid blue line) yields a nearly uniform temperature in the first 5 m, followed by a gradual decrease with depth. With EVD (dashed blue line) the profile is reversed, showing a surface and deep gradient separated by an intermediate quasi-homogeneous layer.

In the following two days (2 and 3 July 2021, both at 6–7 p.m.) the difference between the Richardson-number dependent scheme and the other two schemes increases in both profile shape and amplitude, reaching a sea surface temperature (SST) difference larger than 0.5 °C on 3 July. Moreover, the simulations with TKE and GLS exhibit only minor differences – on the order of ~0.1°C around 15 m depth – arising from slightly different convective patterns in terms of amplitude and vertical extend (Figs. 12e and 12f). In contrast, the Richardson-number dependent scheme shows much larger discrepancies, especially near the surface, due to the substantially different convective patterns (Fig. 12d).




Figure 13: Temperature profiles at a site (27.5°E, 34.4°N) in the Rhodes Gyre region at different times of the day (see legend in the inset) and for different closure schemes and convective adjustment settings (see bottom legend).

4. Discussion and conclusions

Physics-based turbulence closure schemes can represent convective mixing in principle, but at typical model resolution they may underestimate the rate of instability removal. For this reason, convective adjustment is often applied as a numerical parameterization to rapidly stabilize the water column and mimic the fast overturning associated with unresolved convective events. In this work, we investigated how the combined effect of three vertical mixing schemes – Richardson-number dependent, TKE, and GLS closure schemes – and an external convective adjustment (EVD) affects the performance of a regional ocean circulation model of the Mediterranean Sea. We analysed the model outputs in relation to the physics of vertical mixing represented by the different schemes and assessed the advantages and drawbacks of applying an external convective parameterization alongside each of them.

The results first confirm what is known in the literature: the explicit representation of convection through enhanced vertical diffusion is crucial for simple schemes such as Richardson-number dependent closures, which lack an explicit formulation of convection and only reduce density gradient by enhancing vertical mixing in unstable layers. On the other hand, our results indicate that for more physically based schemes, the inclusion of a convective adjustment should be handled with care, as its application can be redundant or even degrade model performance. For the TKE scheme, we found that the inclusion of EVD can be beneficial in some cases. The TKE scheme has a single prognostic equation for turbulent kinetic energy, while the turbulent mixing length is defined diagnostically, and treats convection prognostically through the calculation of the turbulent kinetic energy; as a result, it often represents the intensity of convection well, but underestimates the penetration depth of convective events. In contrast, the GLS scheme, which treats convection prognostically through the calculation of both enhanced turbulent kinetic energy and turbulent mixing length, is generally better able to simulate both the penetration depth and the intensity of convective processes on its own. Therefore, with the GLS scheme, the convective adjustment is redundant and can often degrade the model's skill. These observations are confirmed both at basin scale and in deep and intermediate water formation regions.

At the basin scale, statistical metrics on temperature and salinity show that the GLS scheme performs best, followed by TKE and the Richardson-number dependent scheme. The inclusion of EVD makes little difference overall. In particular, we observed a more consistent improvement with GLS without EVD across all metrics and over a larger range of depths. The largest impact of including or excluding EVD at the basin scale is observed on the MLD. For both the Richardson-number dependent scheme and the TKE schemes, only the addition of the convective adjustment allows the model to reproduce the observed temporal evolution of the average MLD across the basin. This is because EVD compensates for mixing physical processes in these schemes: either the full convective overturning in the Richardson-number dependent scheme or the prognostic evaluation of the mixing length in TKE, which controls the convection penetration depth. In contrast, the GLS

750 scheme, thanks to its prognostic treatment of both TKE and mixing length, is able to reproduce the observed basin-average MLD over time, including the MLD seasonal variability, without requiring an external convective parameterization.

In DWF areas, such as the south Adriatic Pit, where convective processes play a central role during dense water formation events, we observe that both the choice of mixing scheme and the inclusion of the convective adjustment play a major role in
755 the development of deep water and the intrusion of intermediate water from the Levantine basin. Notably, the inclusion of EVD, with its abrupt and strong increase in vertical eddy diffusivity when unstable layers are present, does not account for previous states or the coherent temporal evolution of the water column: when applied, EVD disrupts pre-existing or intruding structures over time. The different mixing schemes without an external convective adjustment reproduce convection in distinct ways, and a similar behaviour is observed in the Rhodes Gyre region. The Richardson-number dependent scheme produces an
760 abrupt but weak homogenization of the layers affected by surface convection, resulting in less gradual penetration into deeper layers and a partial removal of the intruding LIW. The TKE scheme reproduces convection with a smoother intensity transition compared to the Richardson-number dependent scheme, showing less abrupt effects on salinity. However, the penetration depth in the TKE scheme remains limited, without significant sinking of denser waters, due to the lack of a prognostic control on the penetration depth. In contrast, the GLS scheme more accurately represents both the intensity and penetration depth of
765 convective events, allowing for a smooth sinking of the LIW.

Focusing on nighttime convection during summer, analysed at an hourly timescale, we observe how the nighttime convective adjustment, combined with different vertical mixing schemes, shapes the diurnal evolution of temperature. The three vertical mixing schemes represent summer convection differently: the TKE and GLS schemes largely capture the night-and-day
770 differentiation, with GLS producing homogeneous, strong mixing that penetrates deeper. In contrast, the Richardson-number dependent scheme increases mixing weakly but almost continuously, even during daytime showing almost no day-night contrast. Temperature profiles using the Richardson-number dependent scheme differ noticeably from the other closure schemes, and after three days, the SST reaches a value roughly 0.5 °C higher.

775 In summary, across all temporal and spatial scales, our results highlight the crucial role of two key features in vertical mixing schemes. First, prognostic treatment of turbulent kinetic energy (as in the TKE and GLS schemes) is essential to accurately reproduce the intensity of overturning convection, retain memory of priori processes and ensure smooth transitions at a site, preserve pre-existing structures as well as the day-night alternation during summer convection. Second, a prognostic additional equation for the mixing length (as in the GLS scheme), rather than a fixed value as in TKE, is necessary to correctly represent
780 the penetration depth of convective processes and to achieve the most accurate representation of the mixed layer depth. The inclusion of a convective adjustment to compensate for the lack of these features only partially addresses the deficiencies, as external convective adjustment are not embedded within the general framework and do not evolve prognostically.

These findings underscore how critical it is, in regional ocean modelling, to align appropriately the mixing closure and convection parameterisations. More broadly, these results highlight that careful choice of a combination of subgrid-scale parameterizations is essential to reliably simulate key oceanographic features and improve model interpretability in regional applications. Future work could extend this analysis to longer-term simulations and the effect of additional subgrid-scale processes – such as internal waves, eddy formation, and horizontal mixing – in order to further improve model realism and predictive skill. More broadly, our results highlight how the representation of vertical mixing can strongly influence deep and intermediate water formation and properties, with potential implications for regional ocean modelling in other basins with characteristics similar to the Mediterranean Sea. Comparable processes occur in other semi-enclosed or marginal basins, where dense water formation and vertical mixing play a key role in shaping water mass properties. Examples include the Black Sea and the Baltic Sea, as well as convection sites in other marginal seas. In these environments, the choice of vertical mixing parameterization may similarly affect the overall water mass structure in regional ocean models.

Appendix A

This appendix briefly summarizes the main equations of the three vertical mixing schemes used in this study, outlining how the eddy viscosity coefficient A^{vm} and diffusivity coefficient A^{vT} are determined for each scheme. We note that the eddy diffusivity coefficient for temperature, A^{vT} , is assumed to be equal to that for salinity: $A^{vS} \equiv A^{vT}$.

1. Richardson-number dependent mixing scheme

The first closure scheme used here is based on the Richardson number–dependent formulation originally proposed by Pacanowski and Philander (1981), where the vertical eddy viscosity and diffusivity are parametrized as

$$A^{vm} = \frac{A_{ric}^{vm}}{(1 + \alpha R_i)^n} + A_b^{vm} \quad (A1)$$

$$A^{vt} = \frac{A^{vm}}{(1 + \alpha R_i)} + A_b^{vt} \quad (A2)$$

Where $R_i = \frac{N^2}{(\partial U_h)^2}$ denotes the Richardson number, defined as the ratio between stratification (represented by the squared Brunt-Väisälä frequency, N^2) and vertical shear $(\partial U_h)^2$. The parameter A_{ric}^{vm} represents the maximum value

805 of the eddy viscosity coefficient reached when $R_i < 0$; α and n are constant parameters. The additive terms A_b^{vm} and A_b^{vT} are background values that ensure a minimum level of vertical mixing even under stable conditions. For further details, see Pacanowski and Philander (1981) for the theoretical framework and Madec et al. (2023) for the practical implementation in NEMO.

2. Turbulent Kinetic Energy (TKE) vertical mixing scheme

810 The TKE vertical mixing scheme solves a prognostic equation for the turbulent kinetic energy k , while assuming a diagnostic closure for the tubulent length scale l_k . The TKE equation is written as

$$\frac{\partial k}{\partial t} = A^{vm} \left(\frac{\partial U_h}{\partial z} \right)^2 - A^{vT} N^2 + \frac{\partial}{\partial z} \left(A^{vm} \frac{\partial k}{\partial z} \right) - C_\epsilon \frac{k^{3/2}}{l_\epsilon} \quad (\text{A3})$$

815 where U_h is the horizontal velocity, N^2 is squared Brunt-Väisälä frequency, C_ϵ is a constant and l_ϵ is the dissipative length scale. On the right side of equation (A3), the four terms represent the main physical processes associated with vertical mixing. From left to right: shear production, stratification (or buoyancy) destruction, TKE diffusion and Kolmogorov dissipation (Kolmogorov, 1942). Vertical mixing coefficients are then computed as:

$$A^{vm} = C_k l_k \sqrt{k} \quad (\text{A4})$$

$$A^{vT} = \frac{A^{vm}}{P_{rt}} \quad (\text{A5})$$

820

where l_k is the mixing length scale, C_k is a model constant, and P_{rt} is the Prandtl number. The turbulent mixing length l_k is computed diagnostically and its value is constrained by stratification (typically $l_k = l_\epsilon = \sqrt{2k}/N$, with additional operational caps imposed). For further details, see Gaspar et al. (1990) for the theoretical framework and 825 Madec et al. (2023) for the practical implementation in NEMO.

3. Generalized Length Scale (GLS) vertical mixing scheme

The GLS vertical mixing schemes solves two prognostic equations, one for the turbulent kinetic energy k and one for the generic length scale ψ . The evolution of the turbulent kinetic energy is governed by

$$\frac{\partial k}{\partial t} = \frac{A^{vm}}{\sigma_e} \left(\frac{\partial U_h}{\partial z} \right)^2 - A^{vT} N^2 + \frac{\partial}{\partial z} \left(A^{vm} \frac{\partial k}{\partial z} \right) - C_{0\mu} \frac{k^{3/2}}{l_\epsilon} \quad (\text{A6})$$

830

The equation is coupled with a prognostic equation for the generic length scale:

$$\begin{aligned} \frac{\partial \psi}{\partial t} = \frac{\psi}{e} & \left\{ \frac{C_1 A^{vm}}{\sigma_\psi} \left(\frac{\partial U_h}{\partial z} \right)^2 - C_3 A^{vT} N^2 - C_2 C_{0\mu} \frac{k^{3/2}}{l_\epsilon} F_{wall} \right\} \\ & + \frac{\partial}{\partial z} \left(A^{vm} \frac{\partial \psi}{\partial z} \right) \end{aligned} \quad (A7)$$

Here, F_{wall} denotes the wall function and C_1 , C_2 , C_3 , σ_k and σ_ψ are constants that depends on the choice of the turbulent model. $C_{0\mu}$ is also a constant. The wall function is only needed in the Mellor-Yamada model.

835

The eddy viscosity and diffusivity coefficients are then computed from the turbulent kinetic energy and mixing length:

$$A^{vm} = C_\mu l_k \sqrt{k} \quad (A8)$$

$$A^{vT} = C_{\mu'} l_k \sqrt{k} \quad (A9)$$

Where C_μ and $C_{\mu'}$ are the stability functions. The constant $C_{0\mu}$ depends on the choice of the stability functions. For further details, see Burchard and Bolding (2001) for the theoretical framework and Madec et al. (2023) for the practical implementation in NEMO.

840

Appendix B

845

In this appendix, we present the relevant portion of the namelist to be given as input to NEMO (version 4.2) corresponding to the three vertical mixing schemes used in this study. One table is provided for each scheme: Tabel B1 refers to the Richardson-number dependent schemes (Pacanowski and Philander, 1981), Table B2 to the TKE closure scheme (Gaspar et al., 1990) and Table B3 to the GLS closure scheme (Burchard and Bolding, 2001).

Parameter	Value	Explanation
rn_avm0	1.2e-6	vertical eddy viscosity [m2/s] (background value)
rn_avt0	1.0e-7	vertical eddy diffusivity [m2/s] (background value)
rn_avmri	100.e-4	maximum value of the vertical viscosity
rn_alp	5.0	coefficient of the parameterization
nn_ric	2	coefficient of the parameterization
ln_mldw	.false.	enhanced mixing in the Ekman layer

Table B1: Richardson-number dependent vertical mixing scheme

Parameter	Value	Explanation
rn_ediff	0.1	coefficient for vertical eddy diffusivity coefficient (avt=rn_ediff*mxl*sqrt(e))
rn_ediss	0.7	coefficient of the Kolmogoroff dissipation
rn_ebb	3.75	coefficient of the surface input of turbulent kinetic energy
rn_emin	1.e-8	minimum value of turbulent kinetic energy [m2/s2]
rn_emin0	5.e-4	surface minimum value of turbulent kinetic energy [m2/s2]
rn_bshear	1.e-20	background shear (>0) currently a numerical threshold (do not change it)
nn_pdl	1	Prandtl number function of Richarson number (=1, avt=pdl(Ri)*avm) or not (=0, avt=avm)
nn_mxl	2	mixing length: = 0 bounded by the distance to surface and bottom = 1 bounded by the local vertical scale factor = 2 first vertical derivative of mixing length bounded by 1 = 3 as =2 with distinct dissipative mixing length scale
ln_mxl0	.false.	mixing length scale surface value as a function of wind stress or not
rn_mxl0	0.04	surface buoyancy length scale minimum value
ln_mxhsw	.false.	mixing length scale surface value as a function of wave height or not
ln_lc	.true.	Langmuir cell parameterisation (Axell 2002)
rn_lc	0.15	coefficient associated to Langmuir cells
nn_etau	0	penetration of turbulent kinetic energy below the mixed layer (ML) = 0 none ; = 1 add a source of turbulent kinetic energy below the ML = 2 add a source of turbulent kinetic energy just at the base of the ML = 3 as = 1 applied on the high frequency part of the stress
nn_bc_surf	1	surface condition (0/1= Dirichlet/Neumann)
nn_bc_bot	1	bottom condition (0/1= Dirichlet/Neumann)

Table B2: Turbulent Kinetic Energy (TKE) vertical mixing schemes

Parameter	Value	Explanation
rn_emin	1.e-8	minimum value of turbulent kinetic energy [m2/s2]
rn_epsmin	1.e-12	minimum value of dissipation [m2/s3]
ln_length_lim	.true.	limit on the dissipation rate under stable stratification (Galperin et al., 1988)
rn_clim_galp	0.267	Galperin limit
ln_sigpsi	.true.	Activate or not Burchard (2001) modification for k-ε closure and wave breaking mixing
rn_crban	100	Craig and Banner (1994) constant for surface wave breaking mixing
rn_charn	70000	Charnock constant for wave breaking induced roughness length
rn_hsro	0.01	Minimum surface roughness
rn_hsri	0.03	Ice-ocean roughness
rn_frac_hs	0.4	Fraction of wave height as surface roughness
nn_z0_met	2	Method for surface roughness computation (0/1/2/3) = 0 constant roughness = 1 Charnock formula = 2 Roughness formulae according to Raschle et al., 2008 = 3 Roughness given by the wave model
nn_z0_ice	0	attenuation of surface wave breaking under ice = 0 no impact of ice cover = 1 roughness uses rn_hsri and is weighted by 1-TANH(10*fr_i) = 2 roughness uses rn_hsri and is weighted by 1-fr_i = 3 roughness uses rn_hsri and is weighted by 1-MIN(1,4*fr_i)
nn_mxlice	0	mixing under sea ice = 0 No scaling under sea-ice = 1 scaling with constant Ice-ocean roughness (rn_hsri) = 2 scaling with mean sea-ice thickness = 3 scaling with max sea-ice thickness
nn_bc_surf	1	surface condition (0/1=Dirichlet/Neumann)
nn_bc_bot	1	bottom condition (0/1= Dirichlet/Neumann)
nn_stab_func	2	stability function (0=Galp, 1= KC94, 2=CanutoA, 3=CanutoB)
nn_clos	1	predefined closure type (0=MY82, 1=k-eps, 2=k-w, 3=Gen)

Table B3: Generalized Length Scale (GLS) vertical mixing scheme

Appendix C

Throughput this paper, we define four statistics metrics (e.g., Willmott, 1981; Murphy, 1988; Murphy, 1992; Oke et al., 2002, Oddo et al., 2022): the root mean squared error (RMSE), the unbiased root mean squared error (uRMSE), the mean bias (MB), and cross-correlation coefficient (CC). We denote $\bar{o}$ and $\bar{m}$ as the mean values of observations (o) and model data (m), N is the total number of observation-model pairs, and σ represents the standard deviation. The metrics are computed as follows:

- The root mean squared error (RMSE), which combines both systematic and random errors into a single measure of overall discrepancy:

$$RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (m_i - o_i)^2} \quad (C1)$$

- The unbiased root mean squared error (uRMSE), which isolates the random component of the model error by removing the bias:

$$uRMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N [(m_i - \bar{m}) - (o_i - \bar{o})]^2} \quad (C2)$$

- The mean bias (MB), which quantifies the average systematic difference between the model and observations:

$$MB = \bar{m} - \bar{o} \quad (C3)$$

875

- The cross-correlation coefficient (CC), which measures the degree of linear association between model outputs and observations:

$$CC = \frac{1}{\sigma_o} \frac{1}{\sigma_m} \frac{1}{N} \sum_{i=1}^N (m_i - \bar{m})(o_i - \bar{o}) \quad (C4)$$

Code availability

The ocean hydrodynamic circulation model is based on the Nucleus for European Modelling of the Ocean (NEMO) version 4.2.0 (Madec et al., 2023), available on GitLab (<https://forge.nemo-ocean.eu/nemo/nemo/-/releases/4.2.0>) under [doi:10.5281/zenodo.6334656](https://doi.org/10.5281/zenodo.6334656). The wind-wave model is based on WAVEWATCH IIITM (WW3) version 6.07 (Tolman et al., 2021), accessible through GitHub (<https://github.com/NOAA-EMC/WW3/releases>). NEMO namelist configurations for the vertical mixing closure schemes used in this study are provided in Appendix B.

Author contribution

885 LG and PO conceived and designed the study with input from HB, who had a central role in the discussions and development strategy. LG implemented and tested the model setup through sensitivity experiments, with valuable contributions from FB and AM. LG performed the simulations and data analysis. PM contributed to the validation and visualization of the results. FM and EC participated in the discussions and provided feedback. LG wrote the manuscript with input and contributions from all authors, who reviewed and approved the final version of the manuscript.

890 Acknowledgements

The authors would like to thank Prof. Nadia Pinardi for insightful discussions throughout the development of this study and Prof. Paola Cessi for helpful comments.

References

- Axell, L. B., Wind-driven internal waves and Langmuir circulations in a numerical ocean model of the southern Baltic Sea, *Journal of Geophysical Research: Oceans*, 107.C11, 25-1, 2002.
- Bethoux, J. P., Gentili, B., Morin, P., Nicolas, E., Pierre, C., and Ruiz-Pino, D.: The Mediterranean Sea: a miniature ocean for climatic and environmental studies and a key for the climatic functioning of the North Atlantic, *Progress in Oceanography*, 44, 131–146, 1999.
- Bryden, H. L. and Stommel, H. M.: Origin of the Mediterranean outflow, *Journal of Marine Research*, 40, 55–71, 1982.
- 900 Burchard, H. and Bolding, K.: Comparative analysis of four second-moment turbulence closure models for the oceanic mixed layer, *Journal of Physical Oceanography*, 31, 1943–1968, [https://doi.org/10.1175/1520-0485\(2001\)031<1943:CAOFSM>2.0.CO;2](https://doi.org/10.1175/1520-0485(2001)031<1943:CAOFSM>2.0.CO;2), 2001.
- Burchard, H., Simulating the wave-enhanced layer under breaking surface waves with two-equation turbulence models, *Journal of Physical Oceanography*, 31, 11, 3133–3145, 2001.
- 905 Burchard, H. and Petersen, O.: Models of turbulence in the marine environment - A comparative study of two-equation turbulence models. *Journal of Marine Systems*, 21, 29–53, 1999.
- Canuto, V. M., Howard, A., Cheng, Y., and Dubovikov, M. S.: Ocean turbulence. Part I: One-point closure model—momentum and heat vertical diffusivities, *Journal of Physical Oceanography*, 31, 1413–1426, [https://doi.org/10.1175/1520-0485\(2001\)031<1413:OTPIOP>2.0.CO;2](https://doi.org/10.1175/1520-0485(2001)031<1413:OTPIOP>2.0.CO;2), 2001.
- 910 Clementi, E., Oddo, P., Drudi, M., Pinardi, N., Korres, G., and Grandi, A.: Coupling hydrodynamic and wave models: first step and sensitivity experiments in the Mediterranean Sea, *Ocean Dynamics*, 67, 1293–1312, 2017.
- Clementi, E., Drudi, M., Aydogdu, A., Moulin, A., Grandi, A., Mariani, A., Goglio, A. C., Pistoia, J., Miraglio, P., Lecci, R., Palermo, F., Coppini, G., Masina, S., and Pinardi, N.: Mediterranean Sea Physical Analysis and Forecast (CMEMS MED-

- Physics, EAS8 system)(Version 1)[Data set],
 915 https://doi.org/10.25423/CMCC/MEDSEA_ANALYSISFORECAST_PHY_006_013_EAS8, 2023.
- Coppini, G. and Clementi, E. and Cossarini, G. and Salon, S. and Korres, G. and Ravdas, M. and Lecci, R. and Pistoia, J. and Goglio, A. C. and Drudi, M. and Grandi, A. and Aydogdu, A. and Escudier, R. and Cipollone, A. and Lyubartsev, V. and Mariani, A. and Cret\`{i}, S. and Palermo, F. and Scuro, M. and Masina, S. and Pinardi, N. and Navarra, A. and Delrosso, D. and Teruzzi, A. and Di Biagio, V. and Bolzon, G. and Feudale, L. and Coidessa, G. and Amadio, C. and Brosich, A. and
 920 Mir\`{o}, A. and Alvarez, E. and Lazzari, P. and Solidoro, C. and Oikonomou, C. and Zacharioudaki, A.: The Mediterranean Forecasting System—part 1: evolution and performance, *Ocean Science*, 19, 1483–1516, 2023.
- Craig, P. D., and Banner, M.L., Modeling wave-enhanced turbulence in the ocean surface layer, *Journal of Physical Oceanography*, 24, 12, 2546–2559, 1994.
- de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: An
 925 examination of profile data and a profile-based climatology, *Journal of Geophysical Research: Oceans*, 109, C12003, <https://doi.org/10.1029/2004JC002378>, 2004.
- Drévillon, M., Bourdallé-Badie, R., Derval, C., Lellouche, J., Rémy, E., Tranchant, B., Benkiran, M., Greiner, E., Guinehut, S., Verbrugge, N., Garric, G., Testut, C.-E., Laborie, M., Nouel, L., Bahurel, P., Bricaud, C., Crosnier, L., Dombrowsky, E., Durand, E., Ferry, N., Hernandez, F., Le Galloudec, O., Messal, F., and Parent, L.: The GODAE/Mercator-Ocean global
 930 ocean forecasting system: results, applications and prospects, *Journal of Operational Oceanography*, 1, 51–57, <https://doi.org/10.1080/1755876X.2008.11020095>, 2008.
- Escudier, R., Clementi, E., Cipollone, A., Pistoia, J., Drudi, M., Grandi, A., Lyubartsev, V., Lecci, Aydogdu, A., Del Rosso, D., Omar, M., Masina, S., Coppini, G., and Pinardi, N.: A High Resolution Reanalysis for the Mediterranean Sea, *Frontiers in Earth Science*, 9, 1060, <https://doi.org/10.3389/feart.2021.702285>, 2021.
- 935 Gačić, M., Lascaratos, A., Manca, B.B. and Mantziafou, A.: Adriatic deep water and interaction with the Eastern Mediterranean Sea. In *Physical oceanography of the Adriatic Sea: Past, present and future* (pp. 111–142). Dordrecht: Springer Netherlands, 2001.
- Galperin, B., Kantha, L. H., Hassid, S., and Rosati, A.: A quasi-equilibrium turbulent energy model for geophysical flows, *Journal of the atmospheric sciences*, 45(1), 55–62, [https://doi.org/10.1175/1520-0469\(1988\)045<0055:AQETEM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1988)045<0055:AQETEM>2.0.CO;2), 1988.
 940
- Gaspar, P., Grégoris, Y., and Lefevre, J. M.: A simple eddy kinetic energy model for simulations of the oceanic vertical mixing: Tests at station Papa and long-term upper ocean study site, *Journal of Geophysical Research: Oceans*, 95, 16179–16193, <https://doi.org/10.1029/JC095iC09p16179>, 1990.
- Global Ocean Physics Analysis and Forecast. E.U. Copernicus Marine Service Information (CMEMS). Marine Data Store
 945 (MDS). doi: <https://doi.org/10.48670/moi-00016> (Accessed on 15-04-2026)
- Global Ocean-In-Situ Near-Real-Time Observations. E.U. Copernicus Marine Service Information (CMEMS). Marine Data Store (MDS). doi: <https://doi.org/10.48670/moi-00036> (Accessed on 15-05-2026)
- Griffies, S. M., Böning, C., Bryan, F.O, Chassingnet, E. P., Gerdes, R., Hasumi, H., Hirst, A., Treguier, A.M. and Webb, D.: Developments in ocean climate modelling, *Ocean Modelling*, 2, 123–192, [https://doi.org/10.1016/S1463-5003\(00\)00014-7](https://doi.org/10.1016/S1463-5003(00)00014-7),
 950 2000.

- Gutjahr, O., Brüggemann, N., Haak, H., Jungclaus, J. H., Putrasahan, D. A., Lohmann, K., and von Storch, J.-S.: Comparison of ocean vertical mixing schemes in the Max Planck Institute Earth System Model (MPI-ESM1. 2), *Geoscientific Model Development*, 14, 2317–2349, 2021.
- Herrmann, M., Somot, S., Sevault, F., Estournel, C., and Déqué, M.: Modeling the deep convection in the northwestern Mediterranean Sea using an eddy-permitting and an eddy-resolving model: Case study of winter 1986–1987, *Journal of Geophysical Research: Oceans*, 113, 2008.
- Hilt, M., Auclair, F., Benshila, R., Bordis, L., Capet X., Debreu, L., Dumas, F., Julien, S., Lamarié, F., Marchisiello, P., Nguyen, C.: Numerical modeling of hydraulic control, solitary waves and primary instabilities in the Strait of Gibraltar, *Ocean Modelling*, 151, 101642, <https://doi.org/10.1016/j.ocemod.2020.101642>, 2020.
- 960 Kolmogorov, A. N.: Equations of turbulent motion in an incompressible fluid, *Izvestiya Akademii Nauk SSSR, Seriya Fizicheskaya*, 6, 56–58, 1942.
- Kubin, E., Poulain, P.-M., Mauri, E., Menna, M., and Notarstefano, G.: Levantine Intermediate and Levantine Deep Water Formation: An Argo Float Study from 2001 to 2017, *Water*, 11, 1781, <https://doi.org/10.3390/w11091781>, 2019.
- 965 Lascaratos, A., Williams, R.G. and Tragou, E.: A mixed-layer study of the formation of Levantine Intermediate Water, *Journal of Geophysical Research: Oceans*, 98, C8, 14739–14749., 1993
- Lascaratos, A., Roether, W., Nittis, K. and Klein, B.: Recent changes in deep water formation and spreading in the eastern Mediterranean Sea: a review. *Progress in oceanography*, 44, 1–3, 5–36, 1999
- Lazar, A., Madec, G., and Delecluse, P.: The deep interior downwelling, the Veronis effect, and mesoscale tracer transport parameterizations in an OGCM, *Journal of physical oceanography*, 29, 2945–2961, 1999.
- 970 Legay, A., B. Deremble, H. Burchard, Derivation and implementation of a non-local term to improve the oceanic convection representation within the $k-\epsilon$ parameterization, *Journal Advances in Modeling Earth Systems*, 17, e2024MS004243, 2025. <https://doi.org/10.1029/2024MS004243>
- 975 Le Sommer, J., Penduff, T., Penduff, T., Theetten, S., Madec, G., and Barnier, B.: How momentum advection schemes influence current topography interactions at eddy permitting resolution, *Ocean Modeling*, 29, <https://doi.org/10.1016/j.ocemod.2008.11.007>, 2009.
- Lévy, M., Estublier, A. and Madec, G.: Choice of an advection scheme for biogeochemical models. *Geophysical Research Letters*, 28(19), 3725–3728, 2001.
- Lionello, P., Malanotte-Rizzoli, P., Boscolo, R., (eds). *Mediterranean climate variability*, vol. 4. Elsevier, 2006.
- 980 Luneva, M., Wakelin, S., Holt, J., et al.: Challenging Vertical Turbulence Mixing Schemes in a Tidally Energetic Environment: 1. 3-D Shelf-Sea Model Assessment, *Journal of Geophysical Research: Oceans*, 124, 6360–6387, <https://doi.org/10.1029/2018JC014307>, 2019.
- Madec, G. and the NEMO System Team: NEMO Ocean Engine Reference Manual, Tech. rep., <https://doi.org/10.5281/zenodo.8167700>, 2023.

- 985 Madec, G., Delecluse, P., Crepon, M., and Chartier, M.: A three-dimensional numerical study of deep-water formation in the northwestern Mediterranean Sea, *Journal of Physical Oceanography*, 21, 1349–1371, 1991.
- Manca, B.B., Budillon, G., Scarazzato, P. and Ursella, L., Evolution of dynamics in the eastern Mediterranean affecting water mass structures and properties in the Ionian and Adriatic Seas. *Journal of Geophysical Research: Oceans*, 108(C9), 2003.
- 990 Marshall, J. and Schott, F.: Open-ocean convection: Observations, theory, and models, *Reviews of Geophysics*, 37, 1–64, <https://doi.org/10.1029/98RG02739>, 1999.
- Mazzetti, C., Carton De Wiart, C., Gomes, G., Russo, C., Decremer, D., Grimaldi, S., Disperati, J., Ziese, M., Schweim, C., Garcia Sanchez, R., Jacobson, T., Ramos, A., Prudhomme, C., and Salamon, P.: EFAS v5.0 hydrological reanalysis, JRC Technical Report JRC134686, European Commission, Joint Research Centre (JRC), Ispra, Italy, 2023.
- 995 MEDOC Group: Observation of formation of deep water in the Mediterranean Sea, 1969, *Nature*, 227, 1037–1040, <https://doi.org/10.1038/2271037a0>, 1970.
- Mellor, G. L. and Yamada, T.: Development of a turbulence closure model for geophysical fluid problems, *Reviews of Geophysics*, 20, 851–875, <https://doi.org/10.1029/RG020i004p00851>, 1982.
- Millot, C. and Taupier-Letage, I.: Circulation in the Mediterranean Sea, in: *The Mediterranean Sea*, pp. 29–66, Springer, 2005.
- 1000 Murphy, A. H.: Skill scores based on the mean square error and their relationships to the correlation coefficient, *Monthly Weather Review*, 116, 2417–2424, [https://doi.org/10.1175/1520-0493\(1988\)116<2417:SSBOTM>2.0.CO;2](https://doi.org/10.1175/1520-0493(1988)116<2417:SSBOTM>2.0.CO;2), 1988.
- Murphy, A. H.: Climatology, persistence, and their linear combination as standards of reference in skill scores, *Weather and Forecasting*, 7, 692–698, [https://doi.org/10.1175/1520-0434\(1992\)007<0692:CPATLC>2.0.CO;2](https://doi.org/10.1175/1520-0434(1992)007<0692:CPATLC>2.0.CO;2), 1992.
- 1005 Oddo, P., Falchetti, S., Viola, S., Pennucci, G., Storto, A., Borriore, I., Giorli, G., Cozzani, E., Russo, A., Tollefsen, C.: Evaluation of different Maritime rapid environmental assessment procedures with a focus on acoustic performance, *The Journal of the Acoustical Society of America*, 152, 5, 2962–2981, 2022.
- Oddo, P., Adani, M., Pinardi, N., Fratianni, C., Tonani, M., and Pettenuzzo, D.: A nested Atlantic–Mediterranean Sea general circulation model for operational forecasting, *Ocean Science*, 5, 461–473, <https://doi.org/10.5194/os-5-461-2009>, 2009.
- 1010 Oke, P. R., Allen, J. S., Miller, R. N., Egbert, G. D., Austin, J. A., Barth, J. A., Boyd, T. J., Kosro, P. M., and Levine, M. D.: A modeling study of the three-dimensional continental shelf circulation off Oregon. Part I: Model-data comparison, *Journal of Physical Oceanography*, 32, 1360–1382, [https://doi.org/10.1175/1520-0485\(2002\)032<1360:AMOSOT>2.0.CO;2](https://doi.org/10.1175/1520-0485(2002)032<1360:AMOSOT>2.0.CO;2), 2002.
- Pacanowski, R. C. and Philander, S. G. H.: Parameterization of vertical mixing in numerical models of tropical oceans, *Journal of Physical Oceanography*, 11, 1443–1451, 1981.
- 1015 Pinardi, N. and Navarra, A., 1993. Baroclinic wind adjustment processes in the Mediterranean Sea. *Deep Sea Research Part II: Topical Studies in Oceanography*, 40(6), pp.1299–1326.

- Pinardi, N., Allen, I., Demirov, E., De Mey, P., Korres, G., Lascaratos, A., Le Traon, P.-Y., Maillard, C., Manzella, G., and Tziavos, C.: The Mediterranean ocean forecasting system: first phase of implementation (1998–2001), *Annales Geophysicae*, 21, 3–20, <https://doi.org/10.5194/angeo-21-3-2003>, 2003.
- 1020 Pinardi, N. and Coppini, G.: Preface "Operational oceanography in the Mediterranean Sea: the second stage of development", *Ocean Sciences*, 6, 263–267, <https://doi.org/10.5194/os-6-263-2010>, 2010.
- Pinardi, N. and Masetti, E.: Variability of the large scale general circulation of the Mediterranean Sea from observations and modelling: A review, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 158, 153–173, [https://doi.org/10.1016/S0031-0182\(00\)00048-1](https://doi.org/10.1016/S0031-0182(00)00048-1), 2000.
- 1025 Pinardi, N., Cessi, P., Borile, F. and Wolfe, C.L.: The Mediterranean sea overturning circulation. *Journal of Physical Oceanography*, 49(7), 1699–1721, 2019.
- Pinardi, N., Estournel, C., Cessi, P., Escudier, R. and Lyubartsev, V., Dense and deep water formation processes and Mediterranean overturning circulation. In *Oceanography of the mediterranean sea* (pp. 209–261). Elsevier, 2023.
- Poulain, P.-M., Barbanti, R., Font, J., Cruzado, A., Millot, C., Gertman, I., Griffa, A., Molcard, A., Rupolo, V., Le Bras, S.,
1030 Petit de la Villeon L.: MedArgo: a drifting profiler program in the Mediterranean Sea, *Ocean Science*, 3, 379–395, 2007.
- Rahmstorf, S.: A fast and complete convection scheme for ocean models, *Ocean Modelling*, 101, 9–11, 1993.
- Reffray, G., Bourdallé-Badie, R., and Calone, C.: Modelling turbulent vertical mixing sensitivity using a 1-D version of NEMO, *Geoscientific Model Development*, 8, 69–86, <https://doi.org/10.5194/gmd-8-69-2015>, 2015.
- 1035 Raschle, N., Ardhuin, F., Queffelec, P., & Croizé-Fillon, D., A global wave parameter database for geophysical applications. Part 1: Wave-current–turbulence interaction parameters for the open ocean based on traditional parameterizations, *Ocean Modelling*, 25, 3–4, 154–171, 2008.
- Robertson, R., Dong, C.: An evaluation of the performance of vertical mixing parameterizations for tidal mixing in the Regional Ocean Modeling System (ROMS), *Geoscience Letters*, 6, 15, <https://doi.org/10.1186/s40562-019-0146-y>, 2019.
- 1040 Robinson, A. R. and Golnaraghi, M. F.: Circulation and dynamics of the eastern Mediterranean Sea; quasi-synoptic data-driven simulations, *Deep Sea Research Part II: Topical Studies in Oceanography*, 40, 6, 1207–1246, 1993.
- Said, M. A., Gerges, M. A., Maiyza, I. A., Hussein, M. A., and Radwan, A. A.: Changes in Atlantic Water characteristics in the south-eastern Mediterranean Sea as a result of natural and anthropogenic activities, *Oceanologia*, 53, 81–95, <https://doi.org/10.5697/oc.53-1.81>, 2011.
- 1045 Schroeder, K., Chiggiato, J., Bryden, H. L., Borghini, M., and Ben Ismail, S.: Abrupt climate shift in the Western Mediterranean Sea, *Scientific Reports*, 2, <https://doi.org/10.1038/srep23009>, 2012.
- Shchepetkin, A. F.: An adaptive, Courant-number-dependent implicit scheme for vertical advection in oceanic modeling, *Ocean Modelling*, 91, 38–69, 2015.

- Simoncelli, S., Pinardi, N., Fratianni, C., Dubois, C., and Notarstefano, G.: Water mass formation processes in the Mediterranean Sea over the past 30 years. Copernicus Marine Service Ocean State Report, Issue 2. K. von Schuckmann et al., Eds., s96- s100, Journal Operational Oceanography, 11, s1–s142, 2018.
- Somot, S., Sevault, F., and Déqué, M.: Transient climate change scenario simulation of the Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation model, *Climate Dynamics*, 27, 851–879, <https://doi.org/10.1007/s00382-006-0167-z>, 2006.
- Storto, A., Hesham Essa, Y., de Toma, V., Anav, A., Sannino, G., Santoleri, R., and Yang, C.: MESMAR v1: a new regional coupled climate model for downscaling, predictability, and data assimilation studies in the Mediterranean region, *Geoscientific Model Development Discussions*, 1–40, 2023.
- Taillandier, V., D’Ortenzio, F., Prieur, L., Conan, P., Coppola, L., Cornec, M., Dumas, F., Durrieu de Madron, X., Fach, B., Fourier, M. and Gentil, M.: Sources of the Levantine intermediate water in winter 2019. *Journal of Geophysical Research: Oceans*, 127(6), p.e2021JC017506, 2022.
- Testor, P., Bosse A, Houpert L, Margirier F, Mortier L, Legoff H, Dausse D, Labaste M, Karstensen J, Hayes D, Olita A.: Multiscale observations of deep convection in the northwestern Mediterranean Sea during winter 2012–2013 using multiple platforms, *Journal of Geophysical Research: Oceans*, 123, 1745–1776, <https://doi.org/10.1002/2016JC012671>, 2018.
- Tolman, H. L.: User Manual and System Documentation of WAVEWATCH III™ version 6.07, Tech. rep., NOAA/NWS/NCEP/MMAB, <https://polar.ncep.noaa.gov/waves/wavewatch/wavewatch.shtml>, 2021.
- Tonani, M., Pinardi, N., Fratianni, C., Pistoia, J., Dobricic, S., Pensieri, S., de Alfonso, M., and Nittis, K.: Mediterranean Forecasting System: forecast and analysis assessment through skill scores, *Ocean Science*, 5, 649–660, <https://doi.org/10.5194/os-5-649-2009>, 2009.
- Umlauf, L. and Burchard, H.: A generic length-scale equation for geophysical turbulence models, *Journal of Marine Research*, 61, 235–265, 2003.
- Villarreal, M.R., K. Bolding, Burchard, H., and E. Demirov, Coupling of the GOTM turbulence module to some three-dimensional ocean models, pp. 225-237. In: Baumert, H.Z., J.H. Simpson, and J. Sündermann (eds.), *Marine Turbulence: Theories, Observations and Models*, Cambridge University Press, Cambridge, 630 pp. 2005.
- Von Schuckmann, K., Le Traon, P.-Y., Smith, N., Pascual, A., Brasseur, P., Fennel, K., Djavidnia, S., Aaboe, S., Fanjul, E. A., Autret, E., Axell L. : Copernicus marine service ocean state report, *Journal of Operational Oceanography*, 11, S1–S142, 2018.
- Wesson, J. C. and Gregg, M. C.: Mixing at Camarinal Sill in the Strait of Gibraltar, *Journal of Geophysical Research: Oceans*, 99, 9847–9878, <https://doi.org/10.1029/94JC00256>, 1994.
- Willmott, C. J.: On the validation of models, *Physical Geography*, 2, 184–194, <https://doi.org/10.1080/02723646.1981.10642213>, 1981.
- Wong, A. P., Wijffels, S. E., Riser, S. C., Pouliquen, S., Hosoda, S., Roemmich, D., Gilson, J., Johnson, G. C., Martini, K., Murphy, D. J., Scanderbeg M, Bhaskar TVSU, Buck JJH, Merceur F, Carval T, Maze G, Cabanes C, André X, Poffa N, Yashayaev I, Barker PM, Guinehut S, Belbéoch M, Ignaszewski M, Baringer MO, Schmid C, Lyman JM, McTaggart KE, Purkey SG, Zilberman N, Alkire MB, Swift D, Owens WB, Jayne SR, Hersch C, Robbins P, West-Mack D, Bahr F, Yoshida

- 1085 S, Sutton PJH, Cancouët R, Coatanoan C, Dobbler D, Juan AG, Gournion J, Kolodziejczyk N, Bernard V, Bourlès B, Claustre H, D'Ortenzio F, Le Reste S, Le Traon P-Y, Rannou J-P, Saout-Grit C, Speich S, Thierry V, Verbrugge N, Angel-Benavides IM, Klein B, Notarstefano G, Poulain P-M, Vélez-Belchí P, Suga T, Ando K, Iwasaka N, Kobayashi T, Masuda S, Oka E, Sato K, Nakamura T, Sato K, Takatsuki Y, Yoshida T, Cowley R, Lovell JL, Oke PR, van Wijk EM, Carse F, Donnelly M, Gould WJ, Gowers K, King BA, Loch SG, Mowat M, Turton J, Rama Rao EP, Ravichandran M, Freeland HJ, Gaboury I, Gilbert D, Greenan BJW, Ouellet M, Ross T, Tran A, Dong M, Liu Z, Xu J, Kang K, Jo H, Kim S-D and Park H-M: Argo data 1999–2019: Two million temperature-salinity profiles and subsurface velocity observations from a global array of profiling floats, *Frontiers in Marine Science*, 7, 700, 2020.

Wüst, G., On the vertical circulation of the Mediterranean Sea. *Journal of Geophysical Research*, 66(10), pp.3261-3271, 1961.

1095