



On the use of radar isochrones to improve paleo-ice-sheet model simulations away from ice divides

Julien A. Bodart^{1,2}, Vjeran Višnjević^{1,2}, Steven Franke³, Veit Helm⁴, Olaf Eisen^{4,5}, Antoine Hermant^{1,2}, Alexandra M. Zuhr³, Daniel Steinhage⁴, and Johannes C.R. Sutter^{1,2}

¹Climate and Environmental Physics, Physics Institute, University of Bern, Bern, Switzerland

²Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

³Department of Geosciences, Tübingen University, Tübingen, Germany

⁴Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

⁵Faculty of Geosciences, University of Bremen, Bremen, Germany

Correspondence: Julien A. Bodart (julien.bodart@unibe.ch)

Abstract. Radio-echo sounding of polar ice masses have revealed extensive isochronal surfaces that have primarily been used to constrain paleo-accumulation rates, geothermal heat flux, and changes in ice-sheet dynamics in stable regions of the ice sheet. However, isochrones remain under-utilised to calibrate ice-sheet models over large spatial scales, particularly in areas far from the stable ice-sheet divide where englacial layering is more disrupted and models likely perform less accurately. Here, we illustrate the utility of isochrones to constrain paleo-ice-sheet simulations in two off-divide areas of the East Antarctic Ice Sheet; the Wilkes Subglacial Basin (WSB) and the Dronning Maud Land (DML) regions. Using airborne radio-echo sounding data from both legacy and newly acquired surveys, and the three-dimensional, thermo-mechanically coupled Parallel Ice-Sheet Model (PISM), we show that traced and dated isochrones are essential for calibrating ice-sheet model simulations in faster-flowing areas of the ice sheet. We highlight the imprint of paleo-climate forcing and model parameterisation that lead to widely different model representations of isochrones with comparable present day representations of ice sheet geometry. Associated with this paper are two datasets of 9 and 7 newly traced isochronal surfaces spanning the Holocene and Last Interglacial (~4.8 - 128.4 ka) across the WSB and DML sectors, respectively, which may be used in future modelling studies to assess the paleo-evolution of the Antarctic Ice Sheet. This paper sits at the intersection of data-model integration and highlights further opportunities for using isochrones as boundary conditions in paleo-ice-sheet model simulations.

1 Introduction

In recent years, isochronal surfaces (also known as Internal Reflection Horizons, or IRHs) have been increasingly traced and dated across large swaths of the Antarctic Ice Sheet (AIS), mainly in association with the SCAR (Scientific Committee on Antarctic Research) AntArchitecture Action group (Bingham et al., 2025), which aims to build the first three-dimensional age–depth model of the AIS from isochrones. To date, these traced and dated IRHs have primarily been used to constrain paleo-accumulation rates (e.g., Leysinger Vieli et al., 2004; Eisen et al., 2005; Siegert et al., 2004; Neumann et al., 2008; MacGregor et al., 2009; Leysinger Vieli et al., 2011; Steinhage et al., 2013; Karlsson et al., 2014; Koutnik et al., 2016; Cavitte



et al., 2018; Bodart et al., 2023) and ice-dynamical processes (e.g., Nereson et al., 1998; Catania et al., 2006; Leysinger Vieli et al., 2007; Catania et al., 2010; Leysinger Vieli et al., 2018; Chung et al., 2023).

Being often traceable across distances of several hundred kilometres, IRHs represent time-transgressive states of the ice sheet at large spatial scales. This sets them apart from other paleo-records such as offshore sediment cores (e.g., Hillenbrand et al., 2012; Larter et al., 2014), grounding zone wedges (e.g., Bart et al., 2017), radiocarbon-dated erratics on nunataks (e.g., Johnson et al., 2014; Nichols et al., 2019), and shells and bones on beaches (e.g., Braddock et al., 2022), or ice (e.g., Ruth et al., 2007; Buiron et al., 2011) and bedrock (e.g., Balco et al., 2022) cores. These proxies typically provide valuable local to regional snapshots of past ice-sheet dynamics (sediment cores) or extent (grounding zone wedges), elevation changes (radiocarbon dating of rocks and bedrock cores), or climate evolution at a single site (ice cores); however, their one-dimensional nature makes them susceptible to local biases (e.g., Johnson et al., 2021). Complementary to these proxies, isochrones uniquely provide both spatial and temporal calibration targets at the ice-sheet scale (Bingham et al., 2025), bridging a critical gap in available proxy information on past ice-sheet evolution.

Most modelling studies using traced and dated isochrones have focused on regions near stable ice divides (e.g., Leysinger Vieli et al., 2004; Neumann et al., 2008; Koutnik et al., 2016; Cavitte et al., 2018; Bodart et al., 2023; Chung et al., 2023). However, such settings are also where models perform best, largely because ice dynamics at divide locations can be reproduced without the need for more complex physics (e.g., Waddington et al., 2007; MacGregor et al., 2009; Leysinger Vieli et al., 2011; Bodart et al., 2023; Chung et al., 2023). Away from the interior, however, the vertical position of isochrones in the ice column will be the product of different internal processes influencing their elevations and geometries, including (1) ice-flow disturbances which can stretch or disrupt the planar geometry of IRHs, and (2) variations in surface and basal boundary conditions, such as geothermal heat flux (GHF), basal melting, or surface accumulation, which can alter the natural and slow compaction of isochrones over time; all of which require complex, multi-dimensional models to provide an accurate representation of their respective influences (Leysinger Vieli et al., 2007; Waddington et al., 2007; Leysinger Vieli et al., 2011; MacGregor et al., 2009; Sutter et al., 2021).

Continental-scale ice-sheet models have yet to be fully evaluated for their ability to reproduce transient geometries in fast-flowing ice streams or across entire drainage sectors, where ice-dynamical processes and boundary conditions are less well constrained (Sutter et al., 2021). This evaluation is critical because the type of models used in our study are routinely employed to project future sea-level rise, with large uncertainties arising not only from climate scenarios but also to a large part from model initialisation and parametric uncertainties (Golledge et al., 2015; DeConto and Pollard, 2016; Seroussi et al., 2020; Siegert et al., 2020; DeConto et al., 2021; IPCC, 2023; Morlighem et al., 2024). Consequently, it is essential to assess these models' performance not only against present-day observations of ice-sheet geometry and flow, but also on their ability to reproduce past ice-sheet dynamics across a range of climatic states and transitions (Stokes et al., 2015; Kingslake et al., 2018; Bracegirdle et al., 2019; Johnson et al., 2021; Sutter et al., 2021).

Although IRHs have not yet been widely used to constrain ice-sheet model simulations, they offer a powerful benchmark by allowing direct comparisons between observed isochrone elevations and geometries with those generated in large-scale paleo-ice-sheet simulations. Building on this potential, Born and Robinson (2021) and Sutter et al. (2021) presented the first direct



applications of IRHs to constrain continental scale ice-sheet models for Greenland and Antarctica. These studies demonstrated that IRHs can complement more commonly used paleo-proxy reconstructions (e.g., Bracegirdle et al., 2019; Johnson et al., 2021; Lecavalier et al., 2023) to assess how well models reproduce past ice-sheet geometries.

60 This paper sits at the intersection between data and models, and explores the utility of isochrones for ice-sheet model calibration in more dynamical regions of the ice sheet. Together with this study, we also release two new datasets of traced and dated isochrones tailored for ice-sheet model calibration across two regions of East Antarctica. To make use of these datasets within an ice-sheet modelling framework, we begin by describing the airborne radar datasets onto which the isochrones were traced; summarise the process of tracing, dating, and assigning age–depth uncertainties; and then outline the ice-sheet model
65 set-up. We then discuss our results, first describing the new database of isochronal surfaces traced on both legacy and newly acquired radar data across the Wilkes Subglacial Basin (WSB) and Dronning Maud Land (DML), respectively, and then turn our attention to a specific IRH traced and dated at ~ 90 ka (before the present) and found across both WSB and DML regions, with the aim to assess how its elevation and geometry compares with its equivalent modelled isochrone. We conclude this manuscript with a series of lessons learned regarding modelling the paleo-history of the ice sheet in faster (and more dynamical) regions
70 away from stable ice divides.

2 Data sets and methods

2.1 Airborne radar data

We used extensive radio-echo sounding data acquired over both the WSB and DML regions, as well as existing (i.e., published) IRH stratigraphies from the AntArchitecture database (Bingham et al., 2025) (data coverage of these existing stratigraphies is
75 shown in Figure 1) to trace and constrain the age of our isochrones. An important consideration when discussing the airborne radar surveys conducted over both WSB and DML is that whereas data over WSB was acquired mainly to obtain bedrock topography information in a grid-like fashion irrespective of ice flow, the DML survey's main objective was to acquire data that is tailored to ice-sheet modelling applications. This means acquiring data along and across the main trunk of ice streams to (i) constrain mass flux and (ii) quantify the topographical imprint on isochronal stratigraphy which can be used more widely
80 as a benchmark to assess simulated ice flow patterns. Crucially, both sets of radar data intersect either prominent ice core locations or existing traced and dated isochrones previously extracted from coincident radar data (Winter et al., 2019; Cavitte et al., 2021; Franke et al., 2025a), thus enabling the accurate dating of our isochrones.

2.1.1 Legacy radar data over WSB

The primary dataset used over WSB is the 2005-06 airborne radar survey "WISE-ISODYN", acquired in a grid-like pattern
85 (8×44 km spacing on average) by the British Antarctic Survey (BAS) using the 150-MHz Polarimetric Airborne Survey INstrument (PASIN) operating in two interleaved radar modes: a narrow-sounding, $0.1\mu\text{s}$ pulse ideal for tracing isochrones in the upper 2-km of the ice at high resolution, and a deeper-sounding, $4\mu\text{s}$, 10-MHz chirp primarily used to trace deeper



isochrones in the ice column (Frémand et al., 2022) (Table 1). This survey spans most of the WSB catchment with tie-points to the EPICA Dome C (EDC) and TALDICE (TALos Dome Ice CorE) ice cores, covering a total of 730,000 km² and over 61,000 line-km (Fig. 1a) (Ferraccioli et al., 2009; Jordan et al., 2013; Frémand et al., 2022). For the purpose of linking our stratigraphy and increasing the amount of crossovers in crossover-poor areas of the survey, we also used opportunistic flight lines from: (i) CReSIS (Centre for Remote Sensing of Ice Sheets) NASA Operation IceBridge (OIB) MCoRDS (Multichannel Coherent Radar Depth Sounder) radar systems flown between 2013 and 2019 across the WSB to link major deep ice-cores in the region (MacGregor et al., 2021); (ii) Alfred Wegener Institute (AWI)'s EMR (Electromagnetic Reflection System) radar system flown between EDC and TALDICE ice-core sites in 2008–2009 (Winter et al., 2019); and (iii) the University of Texas Institute for Geophysics (UTIG)'s HiCARS2 (High Capability Airborne Radar Sounder 2) radar system flown across WSB during the ICECAP (Investigating the Cryospheric Evolution of the Central Antarctic Plate) survey of 2011–2012 (Frederick et al., 2016; Young et al., 2016), all of which intersect the WISE-ISODYN survey grid in multiple places (Fig. 1a and Table 1). Importantly, existing IRH datasets were also previously extracted for the (ii) and (iii) surveys (i.e., Winter et al., 2019; Cavitte et al., 2021) which we make full use of here (see Section 2.2).

Table 1. Specifications of all the airborne radar systems used in this study over the WSB and DML regions. For PASIN, we provide the transmit signal and range resolution for the pulse and chirp modes, respectively. All values for the range resolution are calculated using an electromagnetic wave speed in ice of 168.5 m s⁻¹. The horizontal sampling distances are typical values after post-processing. The vertical resolution of the chirp systems was calculated following CReSIS using a scaling factor k accounting for resolution degradation. TU HH in the table refers to the Hamburg University of Technology who designed the early EMR system for AWI.

Region	Radar System	Developer	Frequency (MHz)	Transmit Signal	Vert. Sampling Freq. (MHz)	Range Resolution (m)	Horiz. Sampling Distance (m)
WSB	PASIN	BAS	150	0.1 μ s pulse / 4 μ s chirp	22	8.4 / 12.9	45
WSB	MCoRDS3	CReSIS	180–210	1, 3 and 10 μ s chirp	111	4.3	15–30
WSB	MCoRDS5	CReSIS	150–450	1, 3 and 10 μ s chirp	1600	0.35	15
WSB	HiCARS2	UTIG	52.5–67.5	1 μ s chirp	50	8.6	20
WSB & DML	AWI EMR	TU HH	150	60 ns (short-) pulse	75	5.0	35–75
DML	AWI MCoRDS5	CReSIS	180–210	1, 3 and 10 μ s chirp	160	4.3	15
DML	AWI MCoRDS5	CReSIS	150–520	1, 3 and 10 μ s chirp	1600	0.35	15

In order to optimise and standardise the radar data, we removed the data between the air plane and the surface and flattened the ice surface using the previously traced surface pick. To facilitate our tracing, we also applied a gain function and horizontal averaging filter to reduce incoherent noise and enhance the brightness of isochronal stratigraphy, as previously done elsewhere on BAS PASIN data (e.g., Ashmore et al., 2020; Bodart et al., 2021). The vertical scaling of the radargrams were all adjusted to the BAS PASIN system range and sampling interval before being exported as SEG-Y files for isochrone picking.




Figure 1. Map of the airborne radar datasets used in this study for WSB (a) and DML (b) regions. (a) BAS WISE-ISODYN survey (white solid lines) and a combination of individual flight lines (grey dashed lines) from AWI (EMR 60ns pulse data, otherwise referred to in this paper as "short-pulse"; 2008–09), CReSIS (MCoRDS; 2013–19), and UTIG (HiCARS; 2011–12) (Table 1). (b) newly acquired AWI UWB (narrowband) CHARISO survey lines (white solid lines; 2023–25) and a combination of individual flight lines (grey lines) from AWI's EMR (short-pulse; 1999–03; solid lines) and UWB (wideband mode; 2022–23; dashed lines) radars (Table 1). Also shown are the previously published IRH datasets used in this study to tie our IRHs to existing age–depth profiles from Winter et al. (2019) and Cavitte et al. (2021) over WSB (a) and Franke et al. (2025a) over DML (b), with the colorscale representing the number of IRHs traced at a particular position (ranging from 1 to >8 in both (a) and (b)). The background map shows modern ice-flow speeds from Rignot et al. (2017) in m/a. The black lines represent the RINGS/Bedmap3 grounding line from Pritchard et al. (2025) and the main drainage basin boundaries are from IMBIE (2018). The red triangles depict the location of the deep ice cores used in this study to date IRHs. The two yellow rectangles in (a) and (b) represent the location of profiles A-A' and B-B' shown in Figure 3 for the data-model comparison. Abbreviations on (a) and (b) are as follows: EDC (EPICA Dome C), EDML (EPICA Dronning Maud Land), Mertz Glacier (MG), Ninnis Glacier (NG), Belgium's Princess Elizabeth Station (PE), Ross IS (Ross Ice Shelf), SRG (Sør-Rondanebreen Glacier). Note that throughout the paper, grid-west, grid-east, and grid-south are defined relative to the map grid orientation shown here and not to geographic north.



2.1.2 Newly acquired radar data over DML

Obtaining radar data that is tailored to ice-sheet modelling's needs is relatively rare in Antarctica, primarily because the main motivation of radar surveying in Antarctica has historically been about obtaining information on ice thickness and bedrock topography in an efficient manner, which has typically been characterised by surveying an entire catchment on a regular grid-like pattern, thus also facilitating optimized airborne geophysical data acquisition for gravimetry and magnetics (i.e., Fig. 1a). However, ice-sheet models employed away from ice divides also need tailored information of fast-flowing ice streams and main conduits of ice flow to constrain mass flux. This requires acquiring radar data oriented perpendicular or parallel to the direction of flow to capture the imprint of ice streaming often preserved in the geometry of englacial layers. Additionally, ice-sheet models benefit from high-resolution bedrock data (ideally non-interpolated) and, as in this study, englacial layering, which allow for direct evaluation of how well the model reproduces observed features such as the surface, bed, and isochronal surfaces along specific radar transects. Where the WSB and DML flightlines differ is precisely in the motivation of acquisition; WSL was acquired to obtain ice-thickness and bedrock topography over large scales regardless of the surface flow speeds or modelling needs, whereas the DML lines were acquired specifically for ice-sheet modelling applications to encompass fast ice-streaming motion where glacier dynamics are much more challenging than over stable ice-sheet divides.

In the austral summer of 2023–24 and 2024–25, the AWI UWB (UltraWideBand) radar system was flown on the Basler BT-67 aircraft Polar 6 (Alfred Wegener Institute, 2016) across prominent ice-streams draining parts of the DML region as part of the CHARISO (Swiss (CH) Antarctic Research Initiative - ISOchrones in East Antarctic outflow regions) project, providing an opportunity to trace isochrones there and compare their pattern with modelled isochrones from ice-sheet modelling (Fig. 1b and Table 1). The radar system was operated in narrowband mode (180–210 MHz range; 160 MHz sampling frequency; see Table 1) and the data was processed using synthetic-aperture-radar focusing to enhance layering and reduce incoherent noise (for more details on the radar system and processing, see; ?Hale et al., 2016; Franke et al., 2025a). To facilitate tracing, we used the post-processed layer-enhancing AGC (Automatic Gain Control) version of the AWI radar data where possible and, as per WSB, we cut the data between the air plane and the surface and flattened the ice surface using the surface pick prior to exporting the data to SEG-Y file format. In addition to this, further (non-CHARISO) radar data were also used to extend the layering across the DML lines in the upper section near the ice divide, and over the main ice divide to link up specific isochrones with the EDML and Dome Fuji ice cores (Fig. 1b). These additional profiles were acquired using both the AWI UWB system operating in wideband mode (150–520 MHz range) in 2023–24 (Franke et al., 2025a); and the older AWI EMR system (150 MHz) acquired between 1993 and 2003 and operating in short-pulse mode (Fig. 1b, Table 1) to connect the EDML and Dome Fuji ice cores (Nixdorf et al., 1999; Steinhage et al., 2013; Franke et al., 2025a).

2.2 Isochrone tracing, dating, and validation

2.2.1 IRH Tracing

We performed our isochrone tracing using the Schlumberger Petrel® 3-D seismic software's in-built semi-automatic algorithm, which follows the local maxima in reflected power within a previously specified vertical window. The high density of crossovers



between survey flight lines enabled consistent tracing of the same set of isochrones across extensive portions of the radar surveys. This ensured that isochrones traced along the perpendicular grid direction matched their counterparts in the parallel direction, within the limits of system resolution and variations in isochrone appearance—such as brightness or clarity—across different radar systems (Fig. 1). Where necessary, we also made use of the 3-D capability of the software to find intersecting IRHs that were not visible in the 2-D view. Slightly different tracing strategies were employed for each of the WSB and DML regions.

For WSB, we first attempted to extend the existing isochronal stratigraphy datasets from Winter et al. (2019) and Cavitte et al. (2021) at key crossover points, starting at EDC and progressively travelling downstream into the main WSB grid (Fig. 1a). Due to disruptions in IRH geometries away from the dome region around EDC, and with our objective to trace as many IRHs away from the main ice divide, we also traced distinctively bright isochrones at the onset of the TALDICE ice core which were continuous over long enough distances and were not tied to known isochrones from (e.g., Winter et al., 2019) (Fig. 1a). The age of these new isochrones was transferred using the many crossovers in the BAS radar grid to expand the coverage grid-east into the main WSB catchment (Fig. 1a).

For DML, we mainly made use of the existing isochronal stratigraphy datasets from Franke et al. (2025a), which also included traced profiles from other previous studies (Steinhage et al., 2013; Winter et al., 2019), to extend their stratigraphy onto the CHARISO lines and away from the main ice divide using available crossovers in the radar data (Fig. 1b). We found that most of the IRHs from Franke et al. (2025a) were also distinctively bright over the length of the CHARISO profiles, except for the second shallowest and deepest isochrones we traced, which were both relatively bright but did not exactly correspond to the traced IRHs in the Franke et al. (2025a) dataset at our crossovers. In order to assign ages and age uncertainties to these two isochrones, we traced them to EDML and Dome Fuji ice-core sites using the legacy AWI data line along the main ice divide (Fig. 1b). We come back to this point in Section 3.1.

2.2.2 IRH depths and depth uncertainties

Once the isochrones were traced, we converted the value of each data point from the vertical range of the radar systems (t_{IRH}) to depths in metres (z_{IRH}), following

$$z_{IRH} = \frac{t_{IRH} v_i}{2} + z_f, \quad (1)$$

where v_i is the electromagnetic wave speed in ice in m s^{-1} and z_f is the firm correction calculated in Equation 2 in metres.

Different values for v_i and z_f were used to convert to depths in metres, due to the different characteristics of both the radar surveys and the respective regions of East Antarctica in which these surveys and the accompanying ice-cores were situated. For WSB, we used a v_i value of 168.5 m s^{-1} to match the depth of ice-thickness measurements obtained by BAS with the same value, and a spatially invariant z_f value of 14.6 ± 1.4 metres as previously calculated by Cavitte et al. (2016) for EDC. For TALDICE, there are no reported z_f values, so we employed the firm correction equation from Dowdeswell and Evans (2004),

$$z_f = \frac{K}{n_i} \int_0^{z_{\max}} (\rho_i - \rho(z)) dz, \quad (2)$$



where K is the coefficient calculated from Robin et al. (1969) ($0.00085 \text{ m}^3 \text{ kg}^{-1}$) and later refined by Kovacs et al. (1995), n_i is the refractive index of ice (1.78), ρ_i is the density of pure ice (917 kg m^{-3}), and ρ_z is the density at depth z using the TALDICE density profile for the upper 300 metres of the core in kg m^{-3} (Buiron et al., 2011). Following Equation 2, we found a z_f value of 9.5 m, which we applied to all IRHs dated at TALDICE. To this, we added an approximate 2% uncertainty error in density values at TALDICE, as previously suggested by Barnes et al. (2002) and Cavitte et al. (2016) for EDC and Vostok ice cores, amounting to a z_f uncertainty value at TALDICE of $\pm 1.7 \text{ m}$.

For DML, we used a v_i value of 169.5 m s^{-1} to match the depth in metres of the radar-derived products from AWI in the region (e.g., Franke et al., 2025a) and spatially invariant z_f values of $13 \pm 2.5 \text{ m}$ and $15.5 \pm 0.5 \text{ m}$ for IRHs dated at EDML and Dome Fuji, respectively, as previously calculated by Wang et al. (2023) and Franke et al. (2025a) based on shallow firn cores.

We followed the methods of Cavitte et al. (2016), Winter et al. (2019) and Bodart et al. (2021) to calculate the depth uncertainties associated with our isochrones. These uncertainties originate mainly from three components: (i) the range resolution of each radar system (provided in Table 1), (ii) the uncertainty associated with the speed of radar wave in ice, which can be approximated to 1% based on dielectric permittivity measurements in dry ice (Bohleber et al., 2012); and (iii) the uncertainty associated with the firn density corrections used. These three sources of uncertainties are then combined by calculating the root-sum-square error for each isochrone depth. Assuming that the largest uncertainty in IRH depth will be on the deepest (and thus oldest) isochrone, we report here the maximum IRH depth uncertainty on the oldest isochrone in both sectors, which is 13.8 m for the oldest IRH traced on PASIN over WSB and 7.0 m for the oldest IRH traced on the AWI UWB over DML. Given that this represents a worst-case uncertainty across our datasets from the radar uncertainties alone, we also provide depth and age uncertainties for each IRH at the specific ice-core sites (see Section 2.2.3). These numbers represent a simple measure of the maximum IRH depth uncertainty that dataset users can expect from the radar depths alone. It is worth noting that these uncertainties are up to an order of magnitude smaller than what ice-sheet models can currently represent realistically when their own uncertainties are taken into account. In the modelling domain we thus treat traced and dated isochrones such as those presented here as absolute ground truth for model calibration. We do not consider errors associated with our picking as this is difficult to estimate, but accept that this could amount to several more metres in places. To ensure that IRHs are, on average, the same across intersecting transects, we conducted a crossover analysis of IRH depths across both WSB and DML radar lines (see Appendix A), which showed that median crossover errors are between 3 and 18 m over WSB (Fig. A1) and 1 to 8 m over DML (Fig. A2). Lastly, we do not account for snow accumulation or snow re-distribution between survey years, as we expect this to be smaller than the range resolution of the radar systems used here for the area and timespan considered.

2.2.3 IRH ages and age uncertainties

Following the conversion to depth, we proceeded to date the isochrones closest to intersecting ice-core locations using the new AICC2023 ice-core chronology (Bouchet et al., 2023). Over WSB, we used the EDC and TALDICE age–depth profiles to assign ages to the isochrones traced closest to the each ice core by averaging the depths of each isochrone in a 50-m radius around the ice-core sites. Although some of our IRHs matched the depths of previously ice-core dated isochrones in



the region (i.e., Winter et al., 2019; Cavitte et al., 2021), the ages assigned to these were based on an earlier version of the ice-core chronology (i.e., AICC2012; Bazin et al., 2013; Veres et al., 2013), leading to small discrepancies in the age–depth scale between dataset versions. This motivated our decision to update the ages of the newly traced IRHs using the updated ice-core chronology. Over DML, we primarily focused on identifying previously traced and dated isochrones over the region from Franke et al. (2025a), with the exception of the second-shallowest and the deepest isochrones which differed slightly from Franke et al. (2025a). The second-shallowest isochrone we traced, widely visible in the CHARISO profiles, sits slightly deeper in the ice column than the second-shallowest isochrone in Franke et al. (2025a) and was therefore traced along the divide to EDML for accurate dating. In addition, the deepest isochrone traced over the CHARISO lines did not correspond to any isochrone in Franke et al. (2025a) at the intersection between our radar profiles and those from Franke et al. (2025a). Being widely present over our newly acquired survey lines, we proceeded to trace it along the ice divide on the legacy AWI data to both EDML and Dome Fuji ice cores for accurate dating (Fig. 1b). The other five isochrones in our CHARISO lines matched the depth of the other isochrones traced there by Franke et al. (2025a), and so we relied on their age–depth calculations to assign the ages and age uncertainties to our isochrones.

For the IRHs that were dated directly at ice-core sites, we quantified the combined age uncertainty Δa_{comb} of each IRH at each specific ice-core sites by calculating the root-sum-square combination of the age uncertainty associated with the unweighted mean IRH depths found within 50 m of the ice-core site (Δa_{depth}), and the age uncertainty associated with the AICC2023 ice-core age-scales (Bouchet et al., 2023) at the depth of the specific IRH (Δa_{core}) at the core, as per Equation 3,

$$\Delta a_{\text{comb}} = \sqrt{(\Delta a_{\text{depth}})^2 + (\Delta a_{\text{core}})^2}. \quad (3)$$

This combined age uncertainty is calculated for each IRH and each ice-core site located in WSB and DML, except for those already dated by Franke et al. (2025a) over DML.

2.3 Ice-sheet model simulations

We used the three-dimensional, thermo-mechanically coupled Parallel Ice-Sheet Model (PISM, Bueler and Brown, 2009; Winkelmann et al., 2011) to model the evolution of both sectors between the Last Interglacial (LIG) and the present. Because our objective here is to demonstrate the usefulness of isochrones to constrain ice-sheet model simulations, we do not experiment in this study with different sets of boundary conditions, nor do we conduct new large model ensembles testing different model parameters. Instead, we adopted a set of boundary conditions and model parameters from previous simulations of the paleo-state of the continental AIS ran at 16 km² resolution (Sutter et al., 2021), and which accurately reproduce the transient evolution of the ice sheet.

The model configuration employs a combination of shallow-ice/shallow-shelf approximation (SIA+SSA) (Winkelmann et al., 2011) and makes use of the PICO (Potsdam Ice-shelf Cavity mOdel) ocean box model within PISM to simulate ocean-induced melting beneath ice shelves (Reese et al., 2018). Basal sliding follows a pseudo-plastic sliding law (Schoof, 2010), and the visco-elastic response to ice load changes is computed using the Lingle–Clark model (Lingle and Clark, 1985; Bueler et al.,



2007). All simulations are ran on a grid resolution of 8 km^2 (two times higher resolution than in e.g., Sutter et al. (2021)), and 201 vertical levels, which allows for a good representation of vertical velocity gradients that may impact isochrone elevations.

We first run a 2000-year regional initialisation run using present-day bedrock topography and ice thickness from Bedmap3 (Pritchard et al., 2025), and using GHF fields from Shapiro and Ritzwoller (2004) and Haeger et al. (2022) for WSB and DML, respectively. A transient climate forcing is computed by using climate snapshots of the LIG, Last Glacial Maximum (LGM) and pre-industrial from the COSMOS model (Pfeiffer and Lohmann, 2016) and interpolating between these states using a climate index method based on the EDC temperature-accumulation reconstruction and the deuterium record (Jouzel et al., 2007), as previously used in Sutter et al. (2019) and Sutter et al. (2021). Present-day ocean conditions (namely temperature and salinity) are derived from the World Ocean Atlas as used in ISMIP6 (Seroussi et al., 2020) and snapshots of paleo-ocean conditions come from a previous paleo-continental run at 16 km^2 resolution (Wirhs et al., 2025). Modelled isochrone elevations are computed using PISM's own implementation of the Englacial Layer Simulation Architecture (ELSA), an isochronal model for ice-sheet layer tracing (Born, 2017; Rieckh et al., 2024).

3 Results and Discussions

We begin by presenting the observed radar isochrones (Sect. 3.1), traced in WSB (Sect. 3.1.1) and DML (Sect. 3.1.2), with their spatial and depth-normalised coverage illustrated in Figure 2. To showcase the usefulness of IRHs in ice-sheet modelling applications away from the stable ice divides, we then analyse a common isochrone traced in both regions and compare it with a corresponding modelled isochrone (Sect. 3.2), with the results shown in Figure 3.

3.1 Observed isochrones from radar

3.1.1 IRHs over WSB


Over WSB (Fig. 2a-h), we traced 9 IRHs spanning the Holocene and LIG. The depth and age of our IRHs match closely those from Winter et al. (2019) and Cavitte et al. (2021) and include the $38.0 \pm 0.8 \text{ ka}$, $73.5 \pm 2.2 \text{ ka}$, $90.2 \pm 1.9 \text{ ka}$, $117.4 \pm 1.2 \text{ ka}$, $121.1 \pm 1.8 \text{ ka}$, and $128.4 \pm 3.6 \text{ ka}$ (Fig. 2c,e-i) isochrones from their studies. The small age differences between our isochrones and those from Winter et al. (2019) and Cavitte et al. (2021) are attributed to the use of the newer version of the age–depth scale of Antarctic ice cores by Bouchet et al. (2023), as described in Section 2.2.3. Due to disruptions in the continuity of isochrones in the WSB radar data at the transition between the EDC-outbound lines and the main WSB grid covering the central part of the catchment, we were unable to connect many of the traced isochrones from EDC to the main grid with reasonable accuracy. As a result, and to obtain IRH coverage along the central grid-west to grid-east portion of the WSB radar grid (Fig. 1a), IRHs were also traced via their intersection with the age–depth profile at TALDICE ice core where they were dated. This included IRHs dated at $6.5 \pm 0.2 \text{ ka}$, $13.2 \pm 0.4 \text{ ka}$, and $64.4 \pm 3.0 \text{ ka}$. Unfortunately, this disruption in the continuity of internal layering in the upstream part of the main WSB grid and the lack of many crossovers in the radar data in this area hindered our ability to accurately connect isochrones between the EDC and TALDICE ice-core sites (Fig. 2a-h).



This problem has previously been reported elsewhere in East Antarctica (e.g., Cavitte et al., 2016) and points to important lessons learned with regards to survey planning and enhancing crossovers in the radar data when conducting new surveys.

270 When accounting for age uncertainties from this study, the isochrones traced and dated over WSB match closely the age of IRHs previously traced across other sectors of East Antarctica (Siegert, 2003; Leysinger Vieli et al., 2011; Beem et al., 2021; Wang et al., 2023; Franke et al., 2025a; Sanderson et al., 2024; Yan et al., 2025), an important finding for building a 3-D age–depth model of the ice sheet, as motivated by the SCAR AntArchitecture initiative (Bingham et al., 2025). The most extensive isochrone traced across WSB is the 90.2 ± 1.9 ka, which connects the EDC ice core site with the main WSB grid extensively, including as far grid-west as the foothills of the Transantarctic Mountains and in the upstream reaches of the Ninnis Glacier that drains part of the WSB catchment (Figs. 1a and 2f). Another prominent isochrone, dated at 128.4 ± 3.6 ka, connects EDC with the centre of the WSB grid, extending as far downstream (grid south) as the 90.2 ka IRH (Fig. 2i). This isochrone provides an important time marker, ideally dated to around the same time as the onset of the Last Interglacial (LIG; 130–115 ka before the present), and which can be combined with past geological evidence and model simulations that indicate substantial grounding line retreat and ice dynamics occurring in this area of East Antarctica during the LIG (Blackburn et al., 2020; Sutter et al., 2020; Crotti et al., 2022; Iizuka et al., 2023).

It is worth noting that the three deepest isochrones in the WSB catchment, and particularly the oldest IRH dated at 128.4 ka, vary spatially in their reflection strength, forming first a very clear reflector but quickly transitioning to a thick band of diffuse scattering in the chirp data on approach to the main WSB grid from EDC. This diffuse reflector band for the oldest IRH is approximately 15 samples in the vertical range of the BAS PASIN radar system, which approximates to ~ 50 m in the ice. We picked the upper part of this diffuse reflector where it was continuous, but lost it further grid-south where ice-surface speeds and thick ice make these reflectors difficult to interpret. Although uncertainties in both age and depth are greater for this deepest reflector, this IRH is crucial for assessing deep—and therefore old—ice in the catchment and for calibrating the ice-sheet model for this time period. We therefore accept a larger uncertainty for the deepest isochrone over WSB, of approximately ± 50 m, corresponding to ± 3.6 ka at EDC, which is twice the uncertainty associated with the 127.8 ± 1.8 ka IRH in Cavitte et al. (2021). This uncertainty will be more appropriate in such areas where we pick this diffuse reflector band (i.e. in the downstream section of the WSB grid), but we expect a much smaller uncertainty of ~ 1.8 ka in the upper part of the WSB survey grid and towards EDC due to the fact that the reflection is much more distinct and intersects Cavitte et al. (2021)’s IRH in several places with little difference in depth at those crossovers.





295 **Figure 2.** Spatial and vertical depth distribution (as a percentage of the local ice thickness) of each IRH traced over the legacy
 WSB data and connected to EDC and TALDICE ice cores (a–i) and over the newly acquired DML data and connected to
 EMDL and Dome Fuji ice cores (j–p) (shallow depths in the ice column – blue; deeper depths in the ice column – red). The
 thin grey lines represent the full radar survey datasets onto which IRHs were traced, the red triangles represent the deep ice
 cores used to date the IRHs, and the white lines represent the main drainage basin boundaries (IMBIE, 2018). The background
 300 map is the 2014 MODIS mosaic of Antarctica (Haran et al., 2018).

3.1.2 IRHs over DML

Over DML, we traced a total of 7 IRHs across all survey lines, with ages from younger to older: 4.8 ± 0.6 ka, 7.6 ± 0.8 ka,
 25.6 ± 0.1 ka, 38.1 ± 0.1 ka, 52.2 ± 0.1 ka, 73.4 ± 0.9 ka, and 91.0 ± 1.1 ka (Fig. 2j–p). Most IRHs are the exact same
 isochrones as traced previously by Franke et al. (2025a), and so we did not re-calculate the age and age uncertainties and relied
 305 on Franke et al. (2025a) for the dating of our IRHs. The only exceptions to the above is the second-shallowest and the deepest
 IRHs in our DML dataset, which differ slightly from those of Franke et al. (2025a). Our second-shallowest IRH, dated at 7.6
 ± 0.8 ka isochrone, is slightly deeper in the ice than Franke et al. (2025a)'s 7.3 ± 0.8 ka IRH and so was traced it to EDML
 for dating. We note, however, that these two closely spaced IRHs correspond to a couplet seen for several tens of kilometres
 on the CHARISO lines, but with the shallowest one fading away quickly to leave only one distinct isochrone which we traced
 310 here. This characteristic of being a couplet of two shallow isochrones, compounded with the fact that a couplet of similar age
 was also found widely across the WAIS by others (Ashmore et al., 2020; Bodart et al., 2021), leads us to postulate that this set
 of isochrones may be a reliable widespread marker of a powerful volcanic eruption that may have left signatures across large
 swathes of the WAIS and EAIS. Similarly, our deepest IRH, dated at 91.0 ± 1.1 ka is the same IRH as the 91.0 ± 1.1 ka in
 Franke et al. (2025a) but had to be traced further out of the CHARISO lines as no intersections existed in the vicinity of the
 315 new CHARISO lines (Fig. 2p).

Over our new survey lines, we find that the four deepest IRHs (38.1 ± 0.1 ka, 52.2 ± 0.1 ka, 73.4 ± 0.9 ka, and $91.0 \pm$
 1.1 ka) extend much more downstream and across to the SRG ice stream than the shallowest IRHs which tend to fade away
 sooner (Fig. 1j–k vs l–p). Moreover, IRHs are heavily disrupted over faster-flowing ice and where a subglacial mountain range
 constrains ice-stream flow through deep trenches on approach to the bluff onto which the PE station is located (Fig. 1b). As
 320 shown by Franke et al. (2025a), all the isochrones traced over DML correspond to the approximate age (when uncertainties
 are accounted for) of other previously traced and dated IRHs across both West and East Antarctica (e.g., Siegert, 2003; Siegert
 et al., 2004; Siegert and Payne, 2004; Leysinger Vieli et al., 2011; Steinhage et al., 2013; Muldoon et al., 2018; Ashmore et al.,
 2020; Beem et al., 2021; Bodart et al., 2021; Chung et al., 2023; Wang et al., 2023; Sanderson et al., 2024; Yan et al., 2025),
 and importantly, match the age of major volcanic eruptions. Exemplifying this is the fact that we find over both WSB and DML
 325 three common IRHs (~ 38 , ~ 73 , and ~ 91 ka) across both regions when age uncertainties are accounted for.



3.2 Data–model comparison

It is important to note that here, we do not conduct an in-depth analysis of isochronal elevations and patterns for each time slice shown in (Fig. 2), as is common for purely data-focused isochronal studies (e.g., Cavitte et al., 2016; Winter et al., 2019; Ashmore et al., 2020; Bodart et al., 2021; Franke et al., 2025a; Sanderson et al., 2023). This is because our main objective is to (i) provide isochronal data tailored to ice-sheet modelling purposes which aim to constrain the past evolution of WSB and DML regions, and (ii) exemplify the use of these datasets via comparison of individual isochronal transects with model simulations.

For this purpose, we focus here on a common isochrone which was traced and dated independently and which is present across large swathes of both WSB and DML. This isochrone, dated at 90.2 ± 1.9 ka over WSB (Fig. 2f) and 91.0 ± 1.1 ka over DML (Fig. 2p), is also a widely present isochrone across the EAIS (see Section 3.1.1), and likely results from a large explosive volcanic event originating from Mt. Berlin (West Antarctica) that left tephra deposits in EDC and Dome Fuji ice cores at around the same time (Narcisi et al., 2006). We assume that this isochrone is the same reflection across both sectors when uncertainties are considered, especially because it was consistently imaged by a series of radar systems across distant regions of the EAIS and dated at several ice cores, with the absolute age and age uncertainties reflecting the diversity in tracing, dating, and the multitude of datasets used (e.g., Table 1). For the sake of brevity, we will refer to this isochrone as the 90-ka isochrone for the remaining part of this paper.

One way to demonstrate the usefulness of isochrones in assessing 3-D model simulations in off-divide locations — and to disentangle climatic imprints (e.g., surface mass balance variations) from dynamical or physics-based processes that the model does not accurately represent — is through the interrogation of multiple targeted isochrones dated at or near major climate transitions (e.g., Holocene, LGM, LIG) (MacGregor et al., 2015; Born and Robinson, 2021), or alternatively by examining single isochrones across large regions or long transects (Sutter et al., 2021). Here, we focus on the latter, targeting the 90-ka isochrone observed across both WSB and DML, which was deposited during a period when East Antarctica likely experienced broadly similar conditions across these two extensive sectors, in a phase of ice re-growth following the LIG retreat. Notwithstanding, a more focused model-data intercomparison involving large scale regional modelling ensembles constrained by all the isochrones published here will be pursued in upcoming studies (e.g., Višnjević et al., 2024), and we encourage others to make use of these useful datasets in their own future modelling applications.

3.2.1 Comparison between the observed and modelled 90-ka isochrone in WSB and DML regions

We ran three simulations on a regional domain in WSB and DML by slightly modifying the precipitation scaling which governs the relationship between surface temperature change and precipitation change, and the PISM-specific till effective pressure parameter which governs how much of the overburden pressure from the ice load is transferred to the subglacial till to drain excess water from the sediment column (Bueler and van Pelt, 2015), thereby modulating the effective pressure (Fig. 3).

Whilst the precipitation scaling primarily controls Surface Mass Balance (SMB), which in East Antarctica is dominated by net accumulation of snow and minimal surface melt, the till effective pressure primarily affects ice flow via its interaction



with the substrate underneath the ice. The selection of these two parameters was thus made to illustrate the imprint of SMB
 and ice flow on the elevations of isochrones via a small (but commonly used) subset of the large parameter space available to
 ice-sheet models and climate forcing choices (i.e., Albrecht et al., 2020). On Figure 3, we show two long profiles across WSL
 and DML where the 90-ka isochrone is present and crosses two different ice-dynamical regions: (1) the stable ice divide where
 the position of the isochrone in the ice is primarily the result of SMB variations and changes in GHF (Sutter et al., 2021), and
 (2) the faster-flowing areas of the ice sheet away from the ice-divide where ice dynamical processes such as ice deforming over
 subglacial bedrock at relatively fast speeds is common and where the geometry of isochrones can become heavily distorted
 (e.g., Catania et al., 2005).

Often, ice-sheet modelling studies that run long paleo-simulations on glacial to interglacial timescales will be tuned to
 present ice-sheet states from observations (i.e., modern ice-sheet conditions such as ice thickness and ice-flow speeds) and
 past ice-sheet states from paleo-reconstructions (i.e., grounding-line reconstruction from sediment cores). This ensures that the
 ice-sheet model accurately represents known processes well and evolves following realistic paleo-conditions (Albrecht et al.,
 2020). Once the ice-sheet model is initialized to match such present and past ice-sheet states, it is typically run for several
 thousand to hundreds of thousands of years from past to present day, under the assumption that, because it is calibrated to
 modern ice-sheet conditions and sparse paleo-information, its transient evolution provides a reliable representation of the past
 evolution of the ice sheet. Unfortunately, an acceptable match to present day and sparse paleo-benchmark datasets does not
 guarantee a proper representation of internal ice flow as illustrated in Figure 3, which shows close match to modern ice-sheet
 configuration along a radar transect, while at the same time highlighting considerably deviations to the proxy record (in this
 case from the observed isochrone) in both regions. In this context, a mismatch between the modelled and observed isochrones
 potentially indicates overtuning of the ice-sheet model, something which is difficult to identify via simple comparisons with
 current ice geometries or sparse paleo-reconstructions.

A striking observation is that relying on interpolated gridded products such as Bedmap3 (as used here) or BedMachine
 (Morlighem et al., 2020) as appropriate boundary conditions to run the ice-sheet model can be problematic, particularly in
 areas with sparse underlying data used to generate the gridded product. This is exemplified in Figure 3b where the observed
 bed imaged in the newly acquired radar data (and which was not included in the latest iteration of the Bedmap gridding product)
 differs substantially from the modelled beds due to the lack of reliable Bedmap3 data in the area surrounding the profile (see
 also Figure B1). This is particularly striking in the last ~40 km of the profile where a subglacial mountain imaged by the radar
 is not present in the modelled beds, resulting in the modelled isochrones cutting through the observed bed. This section of the
 profile is also where we observed one of the largest difference between observed and modelled isochrones (Fig. 3b; Table C1).
 This exemplifies the utility of using isochrones as a data-benchmark for ice-sheet modelling to investigate both the minimum
 amount of bedrock offset that is required to produce such contrasting isochronal elevations and to rapidly identify areas where
 the modelling results can be least trusted.

Closer to the ice divide, the modelled isochrones are primarily controlled by SMB, which is where the largest difference
 in simulations employing different precipitation scaling is seen, particularly on the DML profiles where the difference in
 isochronal elevations is greater near the divide than near the end of the profile in a compressive flank-flow regime (Fig. 3b).




Figure 3. Comparison of the 90-ka isochrone from observations and model simulations generated in this study, along profiles A-A' (WSB, a) and B-B' (DML, b) crossing the ice divide (start) to fast-flowing ice downstream (end). The solid lines represent the observed (i.e., picked) ice surface, bed and isochrone elevations from the radar data. The non-solid lines (dashed, dashed-dotted, and dotted) represent three simulations that were ran to simulate an accurate paleo-evolution of the two catchments. For each observed surface, bed and isochrone from the radar, we show the three corresponding modelled surfaces, isochrones, and beds which were transiently simulated by the model. Exact location of profiles A–A' and B–B' is shown in Figure 1. The abbreviations "ps" and "teff" in the plot legends refer respectively to the precipitation scaling and till effective pressure parameters mentioned in Section 3.2.1, and the "iso" numbers (ranging from 1 to 3) preceding these refer to each of the three simulations matching the specific parameter values for "ps" and "teff" in the legends. For simplicity, we refer to each of these simulations in the text and in Table C1 using this "iso" numbering abbreviation.



There, the difference in isochronal elevations between the different simulations is reduced, with none of the simulations able
 395 to match the observed isochrone as a result of the unrealistic representation of the modelled bed compared with the observed
 bed. We also note that selecting higher precipitation scaling than previous estimates from deep ice cores (i.e., Frieler et al.,
 2015) appears to improve the agreement between the average elevation of the observed and modelled isochrone close to the
 ice divide, whereas reducing this value means that the modelled isochrone is too low in the ice column. This result was also
 previously highlighted in Sutter et al. (2021), where a higher precipitation scaling provides a better match with the local
 400 isochronal stratigraphy, likely pointing to the fact that a uniform precipitation scaling across the region is unlikely to correctly
 represent the variable conditions away from ice-divide locations.

Similarly over WSB (Fig. 3a), the difference between each simulation is slightly larger closer to the ice divide, owing to the
 strong control of SMB on the isochrone elevations there, whereas this difference reduces further downstream. While the two
 modelled isochrones iso1 and iso3 appear to better match the observed isochrone in the first half of the profile, ice dynamical
 405 processes downstream result in them being situated too low in the ice column compared with the observed isochrone. This
 is a problem that has previously been identified in deep marine basins below sea level, where the basal drag in the model is
 designed to be linearly reduced as bedrock elevation decreases leading to faster sliding (Sutter et al., 2021). Combined with
 a model resolution of 8 km², which is relatively high for transient paleo-modelling but still coarse compared to the 500 m
 Bedmap3 grid, the smoothing of bedrock obstacles and reduction in basal friction at this scale likely contributes to an overall
 410 drawdown of modelled isochrones relative to observations in this sector (Sutter et al., 2021).

For an acceptable level of differences in modern surface elevation and ice thickness between the modelled elevations (last
 time step in the model) and observations (from radar or gridded products), we find that isochrones can differ more substantially
 and at times in contradictory fashion, pointing to the fact that a good match with modern ice surface elevations and ice thickness
 does not guarantee a good match in isochrone elevation. This is illustrated in Fig. 3a over WSB where the mean Root Mean
 Square Error (RMSE) for surface elevations and ice thickness along the entire length of the profile is 98 and 205 m, respectively,
 415 whereas the mean Root Mean Square Difference (RMSD; with reference to the local ice thickness, see Sutter et al. (2021)) for
 all modelled isochrone elevations is 7.5% (equivalent to an RMSE in isochrone elevations of 221 m) across all three isochrones
 for the entire profile (6.4% or 190 m RMSE for iso1, 10.0% or 286 m RMSE for iso2, and 6.0% or 187 m RMSE for iso3;
 Table C1). It is important to note, that these RMSE values further improve when only considering portions of the transect
 420 where the gridded bedrock at model resolution fits the observed bedrock reasonably well. When calculating RMSE in surface
 elevation and ice thickness for each 100-km segment along the profile, we find that the best matches for surface elevation (4 out
 of 7 segments) and ice thickness (6 out of 7) come from both the best (iso3) and worst (iso2) isochrone RMSD values (Table
 C1). This suggests that, for this particular profile, a good agreement with observed surface elevation and ice thickness does
 not necessarily translate to an accurate transient model evolution. In fact, a simple tweak of just two model parameters (in this
 425 case precipitation scaling and till effective pressure between iso3 and iso2) can nearly double the RMSD for this isochrone,
 while still maintaining a good match with surface elevation and ice thickness, illustrating how sensitive the model's temporal
 evolution is to small parameter changes.



A similar pattern is found over DML (Fig. 3b; Table C1) where the best match in surface elevations (4 out of 6 segments) and ice thickness (3 out of 6 segments) come from the simulation where the isochrone has the highest RMSD value (iso1; 20.4% or 359 m RMSE-equivalent) and thus the poorest match to isochrone observations. This contrasts with only 1/6 and 3/6 of all segments for surface elevation and ice thickness, respectively arising from the simulation where the isochrone has the lowest RMSD value (iso3; 15.2% or 285 m RMSE-equivalent) and thus the best match in isochrone observations (Table C1). We note that a precipitation scaling of 11% as employed in the simulation that produced iso3 is likely an overestimate of the average precipitation scaling calculated for the high plateau areas of the East Antarctic Ice Sheet (including EDML), where the mean continental-scale value was estimated to be $\approx 5 \pm 1\%$ K⁻¹ (Frieler et al., 2015). The reason for a high-end precipitation scaling reducing the overall isochronal mismatch over this profile is likely two-fold: (i) low precipitation scaling are potentially more accurate over high-elevation ice divide locations than in off-divide areas, and (ii) the misrepresentation of the true bedrock topography in the boundary conditions used to run the model compared with our newly acquired radar data, which results in unrealistic simulations of isochronal elevations along this profile. A high-precipitation scaling can compensate for this mismatch in bedrock elevations but is not necessarily the most adequate. This illustrates effectively that even though isochrones are a powerful tool for model calibration, they should only be applied where the model can represent the observed bedrock topography to a reasonable degree. What is also notable in the DML transect is the low sensitivity of the surface elevation mismatch at present day compared with correspondingly strong offsets between the modelled and observed isochrones. Ultimately, parameters and forcing choices are key in optimising modelled isochrone geometry while preserving a good match with present day ice-sheet observations. The example shown here highlights that past SMB variations substantially modulate isochrone geometry while affecting the present-day fit of the ice sheet only moderately. Choosing proper sliding laws and associated parameterisation for isochrone geometry is much more challenging, as already small grounding line mismatches can lead to large model deviations from the observed present day ice sheet geometry. As it only makes sense to compare isochrone geometries if the modelled present day ice sheet matches observations, the operating space in ice dynamic parameters is much more limited.

We conclude that whilst the simulations with the highest precipitation scaling produce the best match in isochronal elevation along these two profiles, this is not necessarily associated with the simulations which exhibit the best match in surface elevations or ice thickness. However, since thickness deviations are acceptable for the experiments considered here (i.e., compared to other present-day spin ups in the ISMIP6 intercomparison exercise (Seroussi et al., 2020)), we can extract several key messages from this analysis: (i) ice-sheet models can relatively accurately simulate ice-sheet evolution through long time-scales at 8 km resolution, as illustrated by the relatively good match of data and model along these two profiles and across all three simulations (where mean RMSD in isochrone elevations in the well-constrained WSB profile ranges between 6.0 and 10.0%; equivalent to an RMSE in isochrone elevations of between 187 and 286 m). This is particularly encouraging considering that this isochrone is relatively deep in the ice column and crosses several boundaries where different processes (SMB vs ice dynamics) dominate and where ice-flow transitions from slow to fast flowing; (ii) the underlying boundary conditions used to constrain the ice-sheet model (i.e., here Bedmap3) can significantly affect the transient evolution of the simulations, particularly in areas where strong interpolation in the underlying data product was required due to the lack of nearby bed elevation data from radars. This has the



effect of strongly misrepresenting the isochronal architecture there, as exemplified towards the end of the DML profile where mean RMSD values for the isochrones exceed 35% in the last ~40 km (equivalent an RMSE of 399 m), leading in turn to a misrepresentation of the mean RMSD value in isochrone elevations for the entire profile (i.e., 18.0%; Table C1); and (iii) the use of isochrones as spatially extensive paleo-proxies must form a greater part in constraining ice-sheet modelling results away from stable ice divides to improve simulations and avoid biases, particularly in areas where little other proxy records such as grounding-line reconstructions (Bentley et al., 2014), ice-elevation change reconstructions (Johnson et al., 2021), or other paleo-proxies exists. This is particularly relevant when refining ice-sheet model parameters, as simply tuning an ice-sheet model to modern ice-sheet states does not guarantee a realistic transiently-calibrated ice-sheet model simulation.

4 Conclusions

Using both legacy radar data acquired over the Wilkes Subglacial Basin and newly acquired radar data over Dronning Maud Land, and the 3-D thermo-mechanically coupled PISM ice-sheet model, we showed that the newly traced and dated radar isochrones present at locations away from the stable ice divide of East Antarctica can be extremely useful at assessing the model's ability to transiently simulate the evolution of the ice sheet. A key conclusion from our analysis is that a relatively good match between observed and modelled surface elevation and ice thickness does not necessarily correspond to the best match in isochronal elevations, pointing to the need for increased integration of isochrones in ice-sheet models, particularly in areas far from the stable ice divide where ice flow is faster and ice-sheet models perform less well. Also released alongside this paper are two extensive datasets of isochronal surfaces (16 in total) spanning the Holocene and Last Interglacial (~4.8–128.4 ka) over both investigated East Antarctic sectors, which can be used for further (and more detailed) assessments of ice-sheet models' ability to accurately model the transient evolution of Antarctica. This study exemplifies the need to include further isochronal surfaces into ice-sheet models as an additional proxy to currently used tuning targets, such as modern ice thickness and ice-flow speeds, or key paleo-climate snapshots.

Code and data availability. PISM is an open-source ice-sheet model (Winkelmann et al., 2011; The PISM authors, 2025) that is freely available on GitHub (<https://github.com/pism/pism>). The isochrones produced as part of this study are freely available on Zenodo (<https://doi.org/10.5281/zenodo.17348094>; <https://doi.org/10.5281/zenodo.17348976>) (Bodart and Sutter, 2025a, b) and are provided following guidance from the AntArchitecture data standardisation protocol for IRH datasets (see Supplementary Materials in Bingham et al., 2025), including a specification of the core site where each data point was dated so that users can re-calculate age–depth or firn correction values if these are updated in the future. The WISE-ISODYN radio-echo sounding data is fully available from the BAS Polar Geophysics Data Portal (<https://www.bas.ac.uk/project/nagdp/>) (Ferraccioli et al., 2021; Frémand et al., 2022). The CReSIS and UTIG radar data over WSB are available from (<https://data.cresis.ku.edu/>; Arnold et al., 2020) and (<https://doi.org/10.5067/017PFBVQOGO5>; Blankenship et al., 2017), respectively. The AWI EMR legacy radar data over WSB and DML are available at AWI's new Radar Data over Polar Ice Sheets viewer, accessible within the Marine Data Portal (<https://marine-data.de/viewer/>; Eisen et al., 2024). The newly acquired CHARISO radar data from the 2023–24 and 2024–25 seasons over DML will also be published in AWI's Radar Data viewer prior to publication. The previously existing



495 isochrones used in this study to guide our tracing and dating are openly available with links to the datasets provided in the reference list (Winter et al., 2018; Cavitte et al., 2020; Franke et al., 2025b). The TALDICE density–depth profile used for the firn correction calculations in the paper was downloaded from the TALDICE website (<https://www.taldice.org/data/>; last visited: 17.10.2025). All other datasets used in the methods and to make the figures are freely available, with links provided in the reference list.

Author contributions. J.C.R.S. and J.A.B. planned the new radar flight lines over DML, with O.E., J.C.R.S. and D.S. securing access and funding for the airborne research infrastructure. V.H., S.F., and A.M.Z. acquired and processed the newly flown lines over DML, under the leadership of O.E. and D.S. who also acquired the legacy AWI EMR data over WSB and DML. J.A.B. post-processed the radar data for IRH tracing and traced and dated the IRHs across both regions. J.C.R.S. and V.V. provided the modelling results over WSB and DML, respectively, and J.A.B. analysed them in the context of data-model comparison. A.H. produced the climate-index data used in the modelling. J.A.B. created the figures and wrote the manuscript, with conceptual and scientific inputs from J.C.R.S. All other authors contributed ideas and edits throughout the manuscript and peer-review process.

Competing interests. J.C.R.S. and O.E. are Editor and Advisory Editor of *The Cryosphere* respectively.

Acknowledgements. We thank the the Kenn Borek crew of AWI's polar research aircrafts for the acquisition of the new CHARISO radar lines over DML, and acknowledge the logistical support of the Neumayer III Station (Germany), Troll Station (Norway), and Princess Elisabeth Station (Belgium) during the duration of the flight campaign. We acknowledge the use of PISM, which is supported by NASA grants 20-CRYO2020-0052 and 80NSSC22K0274, and NSF grant OAC-2118285. Finally, we also acknowledge the use of data and/or data products from CReSIS generated with support from the University of Kansas, NASA Operation IceBridge grant NNX16AH54G, NSF grants ACI-1443054, OPP-1739003, and IIS-1838230, Lilly Endowment Incorporated, and Indiana METACyt Initiative. This study was motivated by the SCAR AntArchitecture Action group.

Financial support. J.A.B., V.V., A.H., and J.C.R.S. were all supported by the Swiss National Science Foundation under the CHARIBDIS (Charting Antarctic Ice Sheet evolution via the ice sheet's internal stratigraphy) project (grant number 211542), awarded to J.C.R.S. The new CHARISO lines acquired over DML were funded by AWI, with grant number: TBC. S.F. and A.M.Z. were funded by the Deutsche Forschungsgemeinschaft (grant numbers 506043073 and 541978456 for S.F. and 522419679 for A.M.Z.).

Appendix A: IRHs crossover-points analysis

In order to assess the accuracy of the IRH tracing across the wider WSB and DML regions, we performed a crossover-point analysis on all the IRHs traced in this study (Figs. A1–A2). For this analysis, we extracted all IRH points that fell within 30 m of a crossover point between two survey flight lines and calculated the depth difference of corresponding IRHs of the same



age. Overall, crossover error is lesser over DML than WSB due to the use of newer radar systems in the former and a consistent processing applied to the radar data (Fig. A2), with a median depth difference ranging from 1 to 8 m, which is well within the uncertainty range of the radar system and assumptions made regarding the electromagnetic wave speed in ice. Comparatively, over WSB, where we used data originating from four very different radar systems used between 2005 and 2020 (Table 1), and where data was acquired using different orientations to ice flow, the median depth differences ranged from 3 to 18 m (Fig. A1). Crossover error in WSB is typically greater for older IRHs (Fig. A1d, f–g, and i) where the IRHs transitioned from specular to diffuse and resulted in a thick layer package further downstream from the stable ice divide.




Figure A1. Crossover differences in IRH depths across the WSB radar lines. The median line is shown in red and the 1st and 3rd quartiles are shown as dashed lines.




Figure A2. Crossover differences in IRH depths across the DML radar lines. The median line is shown in red and the 1st and 3rd quartiles are shown as dashed lines.



Appendix B: Bedmap3 bed uncertainty in the vicinity of the newly acquired CHARISO lines in DML




Figure B1. Bed uncertainty from the Bedmap3 dataset over DML, where dark grey lines represent the underlying data used to grid the final data product, and the light gray spacing represents gaps between known survey lines where the interpolation is conducted. Overlaid on top are the new CHARISO radar lines acquired as part of this paper (and which were not integrated in the latest iteration of the Bedmap gridding product), and in yellow the profile B-B' from Figure 3b, with the last ~40 km circled in red.

530 Appendix C: Differences in elevations between modelled and observed surfaces across the WSB and DML profiles



Table C1. Values for RMSD (isochrone elevations; in percentages) and RMSE (surface elevations or ice thicknesses; in metres) between observed and modelled surfaces, calculated for each of the three simulations shown in Figure 3 for each 100 (WSB) or 50 (DML)-km segments, or for the entire length of the profile (last row). "Av" stands for average, "surf" for surface elevation, and "iceT" for ice thickness.

WSB profile – Part 1						
Segment	RMSD_iso_1	RMSD_iso_2	RMSD_iso_3	RMSE_surf_1	RMSE_surf_2	RMSE_surf_3
0	1.7	6.4	2.4	81	128	104
100	8.1	5.4	10.0	86	158	120
200	2.4	7.1	2.2	52	150	103
300	9.2	13.7	8.7	26	102	54
400	6.2	12.3	5.0	72	65	14
500	4.7	9.6	3.2	117	34	33
600	6.7	11.3	4.3	177	9	86
700	8.3	13.2	5.8	200	14	107
0-700	6.4	10.0	6.0	105	105	84

WSB profile – Part 2					
RMSE_iceT_1	RMSE_iceT_2	RMSE_iceT_3	Av_RMSD_iso	Av_RMSE_surf	Av_RMSE_iceT
177	247	206	3.5	104	210
230	217	217	7.8	121	221
124	206	148	3.9	101	159
285	270	263	10.5	60	273
148	159	115	7.8	50	140
258	173	185	5.8	62	206
287	108	171	7.4	91	189
198	173	131	9.1	107	167
225	201	190	7.5	98	205

DML profile – Part 1						
Segment	RMSD_iso_1	RMSD_iso_2	RMSD_iso_3	RMSE_surf_1	RMSE_surf_2	RMSE_surf_3
0	10.6	8.4	4.6	88	100	133
50	7.9	6.6	5.8	122	129	155
100	5.7	4.7	4.7	126	130	150
150	14.7	12.9	9.5	143	143	156
200	9.5	8	5.7	161	156	160
250	37.4	35.7	32.4	151	145	141
0-250	20.4	18.5	15.2	342	339	337

DML profile – Part 2					
RMSE_iceT_1	RMSE_iceT_2	RMSE_iceT_3	Av_RMSD_iso	Av_RMSE_surf	Av_RMSE_iceT
278	266	245	7.9	107	263
326	327	337	6.8	135	330
305	311	334	5.0	135	316
293	292	289	12.4	147	291
219	220	228	7.7	159	222
364	364	360	35.2	146	363
904	898	890	18.0	339	897



References

- Albrecht, T., Winkelmann, R., and Levermann, A.: Glacial-cycle simulations of the Antarctic Ice Sheet with the Parallel Ice Sheet Model (PISM) – Part 2: Parameter ensemble analysis, *The Cryosphere*, 14, 633–656, <https://doi.org/10.5194/tc-14-633-2020>, 2020.
- Alfred Wegener Institute: Polar aircraft Polar5 and Polar6 operated by the Alfred Wegener Institute, Journal of large-scale research facilities, 2, A87–A87, <https://doi.org/10.17815/jlsrf-2-153>, 2016.
- Arnold, E., Leuschen, C., Rodriguez-Morales, F., Li, J., Paden, J., Hale, R., and Keshmiri, S.: CReSIS airborne radars and platforms for ice and snow sounding, *Annals of Glaciology*, 61, 58–67, <https://doi.org/10.1017/aog.2019.37>, 2020.
- Ashmore, D. W., Bingham, R. G., Ross, N., Siegert, M. J., Jordan, T. A., and Mair, D. W.: Englacial architecture and age-depth constraints across the West Antarctic Ice Sheet, *Geophysical Research Letters*, 47, e2019GL086663, <https://doi.org/10.1029/2019GL086663>, 2020.
- Balco, G., Brown, N., Nichols, K., Venturelli, R. A., Adams, J., Braddock, S., Campbell, S., Goehring, B., Johnson, J. S., Rood, D. H., et al.: Reversible ice sheet thinning in the Amundsen Sea Embayment during the Late Holocene, *The Cryosphere Discussions*, 2022, 1–24, <https://doi.org/10.5194/tc-17-1787-2023>, 2022.
- Barnes, P., Wolff, E. W., Mulvaney, R., Udisti, R., Castellano, E., Röthlisberger, R., and Steffensen, J.-P.: Effect of density on electrical conductivity of chemically laden polar ice, *Journal of Geophysical Research: Solid Earth*, 107, ESE–1, <https://doi.org/10.1029/2000JB000080>, 2002.
- Bart, P. J., Krogmeier, B. J., Bart, M. P., and Tulaczyk, S.: The paradox of a long grounding during West Antarctic Ice Sheet retreat in Ross Sea, *Scientific Reports*, 7, 1262, <https://doi.org/10.1038/s41598-017-01329-8>, 2017.
- Bazin, L., Landais, A., Lemieux-Dudon, B., Toyé Mahamadou Kele, H., Veres, D., Parrenin, F., Martinerie, P., Ritz, C., Capron, E., Lipenkov, V., et al.: An optimized multi-proxy, multi-site Antarctic ice and gas orbital chronology (AICC2012): 120–800 ka, *Climate of the Past*, 9, 1715–1731, <https://doi.org/10.5194/cp-9-1715-2013>, 2013.
- Beem, L. H., Young, D. A., Greenbaum, J. S., Blankenship, D. D., Cavitte, M. G. P., Guo, J., and Bo, S.: Aerogeophysical characterization of Titan Dome, East Antarctica, and potential as an ice core target, *The Cryosphere*, 15, 1719–1730, <https://doi.org/10.5194/tc-15-1719-2021>, 2021.
- Bentley, M. J., Cofaigh, C. O., Anderson, J. B., Conway, H., Davies, B., Graham, A. G., Hillenbrand, C.-D., Hodgson, D. A., Jamieson, S. S., Larter, R. D., et al.: A community-based geological reconstruction of Antarctic Ice Sheet deglaciation since the Last Glacial Maximum, *Quaternary Science Reviews*, 100, 1–9, <https://doi.org/10.1016/j.quascirev.2014.06.025>, 2014.
- Bingham, R. G., Bodart, J. A., Cavitte, M. G. P., Chung, A., Sanderson, R. J., Sutter, J. C. R., Eisen, O., Karlsson, N. B., MacGregor, J. A., Ross, N., Young, D. A., Ashmore, D. W., Born, A., Chu, W., Cui, X., Drews, R., Franke, S., Goel, V., Goodge, J. W., Henry, A. C. J., Hermant, A., Hills, B. H., Holschuh, N., Koutnik, M. R., Leysinger Vieli, G., MacKie, E. J., Mantelli, E., Martín, C., Ng, F. S. L., Oraschewski, F. M., Napoleoni, F., Parrenin, F., Popov, S. V., Rieckh, T., Schlegel, R., Schroeder, D. M., Siegert, M. J., Tang, X., Teisberg, T. O., Winter, K., Yan, S., Davis, H., Dow, C. F., Fudge, T. J., Jordan, T. A., Kulesa, B., Matsuoka, K., Nyqvist, C. J., Rahnmooonfar, M., Siegfried, M. R., Singh, S., Višnjević, V., Zamora, R., and Zuhr, A.: Review article: AntArchitecture – building an age–depth model from Antarctica’s radiostratigraphy to explore ice-sheet evolution, *The Cryosphere*, 19, 4611–4655, <https://doi.org/10.5194/tc-19-4611-2025>, 2025.
- Blackburn, T., Edwards, G., Tulaczyk, S., Scudder, M., Piccione, G., Hallet, B., McLean, N., Zachos, J., Cheney, B., and Babbe, J.: Ice retreat in Wilkes Basin of East Antarctica during a warm interglacial, *Nature*, 583, 554–559, <https://doi.org/10.1038/s41586-020-2484-5>, 2020.



- Blankenship, D., Kempf, S., Young, D., Richter, T., Schroeder, D., Ng, G., Greenbaum, J., van Ommen, T., Warner, R., Roberts, J., et al.: IceBridge HiCARS 2 LIB Time-Tagged Echo Strength Profiles, Version 1 [Data set], <https://doi.org/10.5067/0I7PFBVQOGO5>, 2017.
- Bodart, J., Bingham, R. G., Ashmore, D., Karlsson, N., Hein, A., and Vaughan, D.: Age-depth stratigraphy of Pine Island Glacier
 570 inferred from airborne radar and ice-core chronology, *Journal of Geophysical Research: Earth Surface*, 126, e2020JF005927, <https://doi.org/10.1029/2020JF005927>, 2021.
- Bodart, J. A. and Sutter, J. C.: Dated radar Internal Reflection Horizons (IRHs) from the Wilkes Subglacial Basin (East Antarctica) for ice-sheet model calibration [Data set], <https://doi.org/10.5281/zenodo.17348094>, 2025a.
- Bodart, J. A. and Sutter, J. C.: Dated radar Internal Reflection Horizons (IRHs) from Dronning Maud Land (East Antarctica) for ice-sheet
 575 model calibration [Data set], <https://doi.org/10.5281/zenodo.17348976>, 2025b.
- Bodart, J. A., Bingham, R. G., Young, D. A., MacGregor, J. A., Ashmore, D. W., Quartini, E., Hein, A. S., Vaughan, D. G., and Blankenship, D. D.: High mid-Holocene accumulation rates over West Antarctica inferred from a pervasive ice-penetrating radar reflector, *The Cryosphere*, 17, 1497–1512, <https://doi.org/10.5194/tc-17-1497-2023>, 2023.
- Bohleber, P., Wagner, N., and Eisen, O.: Permittivity of ice at radio frequencies: Part I. Coaxial transmission line cell, *Cold Regions Science
 580 and Technology*, 82, 56–67, <https://doi.org/10.1016/j.coldregions.2012.05.011>, 2012.
- Born, A.: Tracer transport in an isochronal ice-sheet model, *Journal of Glaciology*, 63, 22–38, <https://doi.org/10.1017/jog.2016.111>, 2017.
- Born, A. and Robinson, A.: Modeling the Greenland englacial stratigraphy, *The Cryosphere*, 15, 4539–4556, <https://doi.org/10.5194/tc-15-4539-2021>, 2021.
- Bouchet, M., Landais, A., Grisart, A., Parrenin, F., Prié, F., Jacob, R., Fourré, E., Capron, E., Raynaud, D., Lipenkov, V. Y., Loutre, M.-F.,
 585 Extier, T., Svensson, A., Legrain, E., Martinerie, P., Leuenberger, M., Jiang, W., Ritterbusch, F., Lu, Z.-T., and Yang, G.-M.: The Antarctic Ice Core Chronology 2023 (AICC2023) chronological framework and associated timescale for the European Project for Ice Coring in Antarctica (EPICA) Dome C ice core, *Climate of the Past*, 19, 2257–2286, <https://doi.org/10.5194/cp-19-2257-2023>, 2023.
- Bracegirdle, T. J., Colleoni, F., Abram, N. J., Bertler, N. A., Dixon, D. A., England, M., Favier, V., Fogwill, C. J., Fyfe, J. C., Goodwin, I., et al.: Back to the future: using long-term observational and paleo-proxy reconstructions to improve model projections of Antarctic
 590 climate, *Geosciences*, 9, 255, <https://doi.org/10.3390/geosciences9060255>, 2019.
- Braddock, S., Hall, B. L., Johnson, J. S., Balco, G., Spoth, M., Whitehouse, P. L., Campbell, S., Goehring, B. M., Rood, D. H., and Woodward, J.: Relative sea-level data preclude major late Holocene ice-mass change in Pine Island Bay, *Nature Geoscience*, 15, 568–572, <https://doi.org/10.1038/s41561-022-00961-y>, 2022.
- Bueler, E. and Brown, J.: Shallow shelf approximation as a “sliding law” in a thermomechanically coupled ice sheet model, *Journal of
 595 Geophysical Research: Earth Surface*, 114, <https://doi.org/10.1029/2008JF001179>, 2009.
- Bueler, E. and van Pelt, W.: Mass-conserving subglacial hydrology in the Parallel Ice Sheet Model version 0.6, *Geoscientific Model Development*, 8, 1613–1635, <https://doi.org/10.5194/gmd-8-1613-2015>, 2015.
- Bueler, E., Lingle, C. S., and Brown, J.: Fast computation of a viscoelastic deformable Earth model for ice-sheet simulations, *Annals of Glaciology*, 46, 97–105, <https://doi.org/10.3189/172756407782871567>, 2007.
- Buiron, D., Chappellaz, J., Stenni, B., Frezzotti, M., Baumgartner, M., Capron, E., Landais, A., Lemieux-Dudon, B., Masson-Delmotte, V., Montagnat, M., et al.: TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica, *Climate of the Past*, 7, 1–16, <https://doi.org/10.5194/cp-7-1-2011>, 2011.
- Catania, G., Conway, H., Raymond, C., and Scambos, T.: Surface morphology and internal layer stratigraphy in the downstream end of Kamb Ice Stream, West Antarctica, *Journal of Glaciology*, 51, 423–431, <https://doi.org/10.3189/172756505781829142>, 2005.



- 605 Catania, G., Scambos, T., Conway, H., and Raymond, C.: Sequential stagnation of Kamb ice stream, West Antarctica, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2006GL026430>, 2006.
- Catania, G., Hulbe, C., and Conway, H.: Grounding-line basal melt rates determined using radar-derived internal stratigraphy, *Journal of Glaciology*, 56, 545–554, <https://doi.org/10.3189/002214310792447842>, 2010.
- Cavitte, M. G., Blankenship, D. D., Young, D. A., Schroeder, D. M., Parrenin, F., Lemeur, E., Macgregor, J. A., and Siegert, M. J.: Deep
 610 radiostratigraphy of the East Antarctic plateau: connecting the Dome C and Vostok ice core sites, *Journal of Glaciology*, 62, 323–334, <https://doi.org/10.1017/jog.2016.11>, 2016.
- Cavitte, M. G., Parrenin, F., Ritz, C., Young, D. A., Van Liefferinge, B., Blankenship, D. D., Frezzotti, M., and Roberts, J. L.: Accumulation patterns around Dome C, East Antarctica, in the last 73 kyr, *The Cryosphere*, 12, 1401–1414, <https://doi.org/10.5194/tc-12-1401-2018>, 2018.
- 615 Cavitte, M. G., Young, D. A., Mulvaney, R., Ritz, C., Greenbaum, J. S., Ng, G., Kempf, S. D., Quartini, E., Muldoon, G. R., Paden, J., et al.: A detailed radiostratigraphic data set for the central East Antarctic Plateau spanning from the Holocene to the mid-Pleistocene, *Earth System Science Data*, 13, 4759–4777, <https://doi.org/10.5194/essd-13-4759-2021>, 2021.
- Cavitte, M. M., Blankenship, D. D., Frezzotti, M., Kempf, S. D., and Muldoon, G.: Ice-penetrating radar internal stratigraphy over Dome C and the wider East Antarctic Plateau [Data set], <https://doi.org/10.15784/601411>, 2020.
- 620 Chung, A., Parrenin, F., Steinhage, D., Mulvaney, R., Martín, C., Cavitte, M. G., Lilien, D. A., Helm, V., Taylor, D., Gogineni, P., et al.: Stagnant ice and age modelling in the Dome C region, Antarctica, *The Cryosphere*, 17, 3461–3483, <https://doi.org/10.5194/tc-17-3461-2023>, 2023.
- Crotti, I., Quiquet, A., Landais, A., Stenni, B., Wilson, D. J., Severi, M., Mulvaney, R., Wilhelms, F., Barbante, C., and Frezzotti, M.: Wilkes subglacial basin ice sheet response to Southern Ocean warming during late Pleistocene interglacials, *Nature Communications*, 13, 5328,
 625 <https://doi.org/10.1038/s41467-022-32847-3>, 2022.
- DeConto, R. M. and Pollard, D.: Contribution of Antarctica to past and future sea-level rise, *Nature*, 531, 591–597, <https://doi.org/10.1038/nature17145>, 2016.
- DeConto, R. M., Pollard, D., Alley, R. B., Velicogna, I., Gasson, E., Gomez, N., Sadai, S., Condrón, A., Gilford, D. M., Ashe, E. L., et al.:
 The Paris Climate Agreement and future sea-level rise from Antarctica, *Nature*, 593, 83–89, <https://doi.org/10.1038/s41586-021-03427-0>,
 630 2021.
- Dowdeswell, J. A. and Evans, S.: Investigations of the form and flow of ice sheets and glaciers using radio-echo sounding, *Reports on Progress in Physics*, 67, 1821, <https://doi.org/10.1088/0034-4885/67/10/R03>, 2004.
- Eisen, O., Rack, W., Nixdorf, U., and Wilhelms, F.: Characteristics of accumulation around the EPICA deep-drilling site in Dronning Maud Land, Antarctica, *Annals of Glaciology*, 41, 41–46, <https://doi.org/10.3189/172756405781813276>, 2005.
- 635 Eisen, O., Steinhage, D., Franke, S., Helm, V., Binder, T., Drews, R., Eagles, G., Humbert, A., Jansen, D., Jokat, W., Lambrecht, A., Mieth, M., Riedel, S., and Miller, H.: Collection of datasets from AWT's radio-echo sounding systems on ice sheets and glaciers [Data set], <https://doi.org/10.1594/PANGAEA.972094>, 2024.
- Ferraccioli, F., Armadillo, E., Jordan, T., Bozzo, E., and Corr, H.: Aeromagnetic exploration over the East Antarctic Ice Sheet: a new view of the Wilkes Subglacial Basin, *Tectonophysics*, 478, 62–77, <https://doi.org/10.1016/j.tecto.2009.03.013>, 2009.
- 640 Ferraccioli, F., Forsberg, R., Matsuoka, K., Olesen, A., Jordan, T., Corr, H., Robinson, C., and Kohler, J.: Processed airborne radio-echo sounding data from the POLARGAP survey covering the South Pole, and Foundation and Recovery Glaciers, East



- Antarctica (2015/2016)(Version 1.0)[Data set]. NERC EDS UK Polar Data Centre, <https://doi.org/10.5285/70adab3d-3632-400d-9aa1-ddf2d62a11b3>, 2021.
- 645 Franke, S., Steinhage, D., Helm, V., Zuhr, A. M., Bodart, J. A., Eisen, O., and Bons, P.: Age–depth distribution in western Dronning Maud Land, East Antarctica, and Antarctic-wide comparisons of internal reflection horizons, *The Cryosphere*, 19, 1153–1180, <https://doi.org/10.5194/tc-19-1153-2025>, 2025a.
- Franke, S., Steinhage, D., Helm, V., Zuhr, A. M., and Eisen, O.: Internal Reflection Horizons (IRHs) in western Dronning Maud Land, East Antarctica from airborne radar surveys (4.8 - 91.0 ka) [Data set], <https://doi.org/10.1594/PANGAEA.973266>, 2025b.
- 650 Frederick, B. C., Young, D. A., Blankenship, D. D., Richter, T. G., Kempf, S. D., Ferraccioli, F., and Siegert, M. J.: Distribution of subglacial sediments across the Wilkes Subglacial basin, East Antarctica, *Journal of Geophysical Research: Earth Surface*, 121, 790–813, <https://doi.org/10.1002/2015JF003760>, 2016.
- Frieler, K., Clark, P. U., He, F., Buizert, C., Reese, R., Ligtenberg, S. R., Van Den Broeke, M. R., Winkelmann, R., and Levermann, A.: Consistent evidence of increasing Antarctic accumulation with warming, *Nature Climate Change*, 5, 348–352, <https://doi.org/10.1038/nclimate2574>, 2015.
- 655 Frémand, A. C., Bodart, J. A., Jordan, T. A., Ferraccioli, F., Robinson, C., Corr, H. F. J., Peat, H. J., Bingham, R. G., and Vaughan, D. G.: British Antarctic Survey’s aerogeophysical data: releasing 25 years of airborne gravity, magnetic, and radar datasets over Antarctica, *Earth System Science Data*, 14, 3379–3410, <https://doi.org/10.5194/essd-14-3379-2022>, 2022.
- Golledge, N. R., Kowalewski, D. E., Naish, T. R., Levy, R. H., Fogwill, C. J., and Gasson, E. G.: The multi-millennial Antarctic commitment to future sea-level rise, *Nature*, 526, 421–425, <https://doi.org/10.1038/nature15706>, 2015.
- 660 Haeger, C., Petrunin, A., and Kaban, M.: Geothermal heat flow and thermal structure of the Antarctic lithosphere, *Geochemistry, Geophysics, Geosystems*, 23, e2022GC010 501, <https://doi.org/10.1029/2022GC010501>, 2022.
- Hale, R., Miller, H., Gogineni, S., Yan, J. B., Rodriguez-Morales, F., Leuschen, C., Paden, J., Li, J., Binder, T., Steinhage, D., Gehrmann, M., and Braaten, D.: Multi-Channel Ultra-Wideband Radar Sounder and Imager, 2016 IEEE International Geoscience and Remote Sensing Symposium (IGARSS), pp. 2112–2115, <https://doi.org/10.1109/igarss.2016.7729545>, 2016.
- 665 Haran, T., Klinger, M., Bohlander, J., Fahnestock, M., Painter, T., and Scambos, T.: MEaSUREs MODIS Mosaic of Antarctica 2013-2014 (MOA2014) Image Map, Version 1 [Data set], <https://doi.org/10.5067/RNF17BP824UM>, 2018.
- Hillenbrand, C.-D., Melles, M., Kuhn, G., and Larter, R. D.: Marine geological constraints for the grounding-line position of the Antarctic Ice Sheet on the southern Weddell Sea shelf at the Last Glacial Maximum, *Quaternary Science Reviews*, 32, 25–47, <https://doi.org/10.1016/j.quascirev.2011.11.017>, 2012.
- 670 Iizuka, M., Seki, O., Wilson, D. J., Suganuma, Y., Horikawa, K., van de Flierdt, T., Ikehara, M., Itaki, T., Irino, T., Yamamoto, M., et al.: Multiple episodes of ice loss from the Wilkes Subglacial Basin during the Last Interglacial, *Nature Communications*, 14, 2129, <https://doi.org/10.1038/s41467-023-37325-y>, 2023.
- IMBIE: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219–222, <https://doi.org/10.1038/s41586-018-0179-y>, 2018.
- 675 IPCC: Ocean, Cryosphere and Sea Level Change, p. 1211–1362, Cambridge University Press, <https://doi.org/10.1017/9781009157896.011>, 2023.
- Johnson, J. S., Bentley, M. J., Smith, J. A., Finkel, R., Rood, D., Gohl, K., Balco, G., Larter, R. D., and Schaefer, J.: Rapid thinning of Pine Island Glacier in the early Holocene, *Science*, 343, 999–1001, <https://doi.org/10.1126/science.1247385>, 2014.



- Johnson, J. S., Pollard, D., Whitehouse, P. L., Roberts, S. J., Rood, D. H., and Schaefer, J. M.: Comparing glacial-geological evidence and
 680 model simulations of ice sheet change since the last glacial period in the Amundsen Sea sector of Antarctica, *Journal of Geophysical Research: Earth Surface*, 126, e2020JF005 827, <https://doi.org/10.1029/2020JF005827>, 2021.
- Jordan, T., Ferraccioli, F., Armadillo, E., and Bozzo, E.: Crustal architecture of the Wilkes Subglacial Basin in East Antarctica, as revealed
 from airborne gravity data, *Tectonophysics*, 585, 196–206, <https://doi.org/10.1016/j.tecto.2012.06.041>, 2013.
- Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster, B., Nouet, J., Barnola, J.-M.,
 685 Chappellaz, J., et al.: Orbital and millennial Antarctic climate variability over the past 800,000 years, *Science*, 317, 793–796,
<https://doi.org/10.1126/science.1141038>, 2007.
- Karlsson, N. B., Bingham, R. G., Rippin, D. M., Hindmarsh, R. C., Corr, H. F., and Vaughan, D. G.: Constraining past accumu-
 lation in the central Pine Island Glacier basin, West Antarctica, using radio-echo sounding, *Journal of Glaciology*, 60, 553–562,
<https://doi.org/10.3189/2014JoG13J180>, 2014.
- 690 Kingslake, J., Scherer, R., Albrecht, T., Coenen, J., Powell, R., Reese, R., Stansell, N., Tulaczyk, S., Wearing, M., and Whitehouse, P.: Exten-
 sive retreat and re-advance of the West Antarctic Ice Sheet during the Holocene, *Nature*, 558, 430–434, <https://doi.org/10.1038/s41586-018-0208-x>, 2018.
- Koutnik, M. R., Fudge, T., Conway, H., Waddington, E. D., Neumann, T. A., Cuffey, K. M., Buizert, C., and Taylor, K. C.: Holocene
 accumulation and ice flow near the West Antarctic Ice Sheet Divide ice core site, *Journal of Geophysical Research: Earth Surface*, 121,
 695 907–924, <https://doi.org/10.1002/2015JF003668>, 2016.
- Kovacs, A., Gow, A. J., and Morey, R. M.: The in-situ dielectric constant of polar firn revisited, *Cold Regions Science and Technology*, 23,
 245–256, [https://doi.org/10.1016/0165-232X\(94\)00016-Q](https://doi.org/10.1016/0165-232X(94)00016-Q), 1995.
- Larter, R. D., Anderson, J. B., Graham, A. G., Gohl, K., Hillenbrand, C.-D., Jakobsson, M., Johnson, J. S., Kuhn, G., Nitsche, F. O., Smith,
 J. A., et al.: Reconstruction of changes in the Amundsen Sea and Bellingshausen sea sector of the West Antarctic ice sheet since the last
 700 glacial maximum, *Quaternary Science Reviews*, 100, 55–86, <https://doi.org/10.1016/j.quascirev.2013.10.016>, 2014.
- Lecavalier, B. S., Tarasov, L., Balco, G., Spector, P., Hillenbrand, C.-D., Buizert, C., Ritz, C., Leduc-Leballeur, M., Mulvaney, R., White-
 house, P. L., Bentley, M. J., and Bamber, J.: Antarctic Ice Sheet paleo-constraint database, *Earth System Science Data*, 15, 3573–3596,
<https://doi.org/10.5194/essd-15-3573-2023>, 2023.
- Leysinger Vieli, G. J., Siegert, M. J., and Payne, A. J.: Reconstructing ice-sheet accumulation rates at ridge B, East Antarctica, *Annals of*
 705 *Glaciology*, 39, 326–330, <https://doi.org/10.3189/172756404781814519>, 2004.
- Leysinger Vieli, G. J., Hindmarsh, R. C., and Siegert, M. J.: Three-dimensional flow influences on radar layer stratigraphy, *Annals of*
Glaciology, 46, 22–28, <https://doi.org/10.3189/172756407782871729>, 2007.
- Leysinger Vieli, G. J., Hindmarsh, R. C. A., Siegert, M. J., and Bo, S.: Time-dependence of the spatial pattern of accumulation rate in East
 Antarctica deduced from isochronic radar layers using a 3-D numerical ice flow model, *Journal of Geophysical Research: Earth Surface*,
 710 116, <https://doi.org/10.1029/2010jg001785>, 2011.
- Leysinger Vieli, G. J.-M. C., Martín, C., Hindmarsh, R. C. A., and Lüthi, M. P.: Basal freeze-on generates complex ice-sheet stratigraphy,
Nature Communications, 9, 4669, <https://doi.org/10.1038/s41467-018-07083-3>, 2018.
- Lingle, C. S. and Clark, J. A.: A numerical model of interactions between a marine ice sheet and the solid earth: Application to a West
 Antarctic ice stream, *Journal of Geophysical Research: Oceans*, 90, 1100–1114, <https://doi.org/10.1029/JC090iC01p01100>, 1985.



- 715 MacGregor, J. A., Matsuoka, K., Koutnik, M. R., Waddington, E. D., Studinger, M., and Winebrenner, D. P.: Millennially averaged accumulation rates for the Vostok Subglacial Lake region inferred from deep internal layers, *Annals of Glaciology*, 50, 25–34, <https://doi.org/10.3189/172756409789097441>, 2009.
- MacGregor, J. A., Fahnestock, M. A., Catania, G. A., Paden, J. D., Prasad Gogineni, S., Young, S. K., Rybarski, S. C., Mabrey, A. N., Wagman, B. M., and Morlighem, M.: Radiostratigraphy and age structure of the Greenland Ice Sheet, *Journal of Geophysical Research: Earth Surface*, 120, 212–241, <https://doi.org/10.1002/2014JF003215>, 2015.
- 720 MacGregor, J. A., Boisvert, L. N., Medley, B., Petty, A. A., Harbeck, J. P., Bell, R. E., Blair, J. B., Blanchard-Wigglesworth, E., Buckley, E. M., Christoffersen, M. S., et al.: The scientific legacy of NASA’s Operation IceBridge, *Review of Geophysics*, 59, e2020RG000712, <https://doi.org/10.1029/2020RG000712>, 2021.
- Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., Goel, V., Greenbaum, J. S., Gudmundsson, H., Guo, J., Helm, V., Hofstede, C., Howat, I., Humbert, A., Jokat, W., Karlsson, N. B., Lee, W. S., Matsuoka, K., Millan, R., Mouginot, J., Paden, J., Pattyn, F., Roberts, J., Rosier, S., Ruppel, A., Seroussi, H., Smith, E. C., Steinhage, D., Sun, B., Broeke, M. R. v. d., Ommen, T. D. v., Wessem, M. v., and Young, D. A.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, 13, 132–137, <https://doi.org/10.1038/s41561-019-0510-8>, 2020.
- Morlighem, M., Goldberg, D., Barnes, J. M., Bassis, J. N., Benn, D. I., Crawford, A. J., Gudmundsson, G. H., and Seroussi, H.: The West Antarctic Ice Sheet may not be vulnerable to marine ice cliff instability during the 21st century, *Science Advances*, 10, eado7794, <https://doi.org/10.1126/sciadv.ado7794>, 2024.
- 730 Muldoon, G. R., Jackson, C. S., Young, D. A., and Blankenship, D. D.: Bayesian estimation of englacial radar chronology in Central West Antarctica, *Dynamics and Statistics of the Climate System*, 3, dzy004, <https://doi.org/10.1093/climatesystem/dzy004>, 2018.
- Narcisi, B., Petit, J. R., and Tiepolo, M.: A volcanic marker (92 ka) for dating deep east Antarctic ice cores, *Quaternary Science Reviews*, 25, 2682–2687, <https://doi.org/10.1016/j.quascirev.2006.07.009>, 2006.
- 735 Nereson, N., Raymond, C., Waddington, E., and Jacobel, R.: Migration of the Siple Dome ice divide, West Antarctica, *Journal of Glaciology*, 44, 643–652, <https://doi.org/10.3189/S0022143000002148>, 1998.
- Neumann, T., Conway, H., Price, S., Waddington, E., Catania, G., and Morse, D.: Holocene accumulation and ice sheet dynamics in central West Antarctica, *Journal of Geophysical Research: Earth Surface*, 113, <https://doi.org/10.1029/2007JF000764>, 2008.
- 740 Nichols, K. A., Goehring, B. M., Balco, G., Johnson, J. S., Hein, A. S., and Todd, C.: New last glacial maximum ice thickness constraints for the Weddell Sea Embayment, Antarctica, *The Cryosphere*, 13, 2935–2951, <https://doi.org/10.5194/tc-13-2935-2019>, 2019.
- Nixdorf, U., Steinhage, D., Meyer, U., Hempel, L., Jenett, M., Wachs, P., and Miller, H.: The newly developed airborne radio-echo sounding system of the AWI as a glaciological tool, *Annals of Glaciology*, 29, 231–238, <https://doi.org/10.3189/172756499781821346>, 1999.
- Pfeiffer, M. and Lohmann, G.: Greenland Ice Sheet influence on Last Interglacial climate: global sensitivity studies performed with an atmosphere–ocean general circulation model, *Climate of the Past*, 12, 1313–1338, <https://doi.org/10.5194/cp-12-1313-2016>, 2016.
- 745 Pritchard, H. D., Fretwell, P. T., Fremant, A. C., Bodart, J. A., Kirkham, J. D., Aitken, A., Bamber, J., Bell, R., Bianchi, C., Bingham, R. G., et al.: Bedmap3 updated ice bed, surface and thickness gridded datasets for Antarctica, *Scientific data*, 12, 414, <https://doi.org/10.1038/s41597-025-04672-y>, 2025.
- Reese, R., Albrecht, T., Mengel, M., Asay-Davis, X., and Winkelmann, R.: Antarctic sub-shelf melt rates via PICO, *The Cryosphere*, 12, 1969–1985, <https://doi.org/10.5194/tc-12-1969-2018>, 2018.
- 750 Rieckh, T., Born, A., Robinson, A., Law, R., and Gülle, G.: Design and performance of ELSA v2. 0: an isochronal model for ice-sheet layer tracing, *Geoscientific Model Development*, 17, 6987–7000, <https://doi.org/10.5194/gmd-17-6987-2024>, 2024.



- Rignot, E., Mouginot, J., and Scheuchl, B.: MEaSUREs InSAR-Based Antarctica Ice Velocity Map, v 2.0.0. [Data Set],
<https://doi.org/10.5067/D7GK8F5J8M8R>, 2017.
- 755 Robin, G. D. Q., Evans, S., and Bailey, J. T.: Interpretation of radio echo sounding in polar ice sheets, *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 265, 437–505, <https://doi.org/10.1098/rsta.1969.0063>, 1969.
- Ruth, U., Barnola, J.-M., Beer, J., Bigler, M., Blunier, T., Castellano, E., Fischer, H., Fundel, F., Huybrechts, P., Kaufmann, P., Kipfstuhl, S., Lambrecht, A., Morganti, A., Oerter, H., Parrenin, F., Rybak, O., Severi, M., Udisti, R., Wilhelms, F., and Wolff, E.: "EDML1": a chronology for the EPICA deep ice core from Dronning Maud Land, Antarctica, over the last 150 000 years, *Climate of the Past*, 3,
 475–484, <https://doi.org/10.5194/cp-3-475-2007>, 2007.
- 760 Sanderson, R. J., Winter, K., Callard, S. L., Napoleoni, F., Ross, N., Jordan, T. A., and Bingham, R. G.: Englacial architecture of Lambert Glacier, East Antarctica, *The Cryosphere*, 17, 4853–4871, <https://doi.org/10.5194/tc-17-4853-2023>, 2023.
- Sanderson, R. J., Ross, N., Winter, K., Bingham, R. G., Callard, S. L., Jordan, T. A., and Young, D. A.: Dated radar-stratigraphy between Dome A and South Pole, East Antarctica: old ice potential and ice sheet history, *Journal of Glaciology*, p. 1–14,
 765 <https://doi.org/10.1017/jog.2024.60>, 2024.
- Schoof, C.: Coulomb friction and other sliding laws in a higher-order glacier flow model, *Mathematical Models and Methods in Applied Sciences*, 20, 157–189, <https://doi.org/10.1142/S0218202510004180>, 2010.
- Seroussi, H., Nowicki, S., Payne, A. J., Goelzer, H., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Albrecht, T., Asay-Davis, X., Barthel, A., Calov, R., Cullather, R., Dumas, C., Galton-Fenzi, B. K., Gladstone, R., Golledge, N. R., Gregory, J. M., Greve, R., Hattermann, T.,
 770 Hoffman, M. J., Humbert, A., Huybrechts, P., Jourdain, N. C., Kleiner, T., Larour, E., Leguy, G. R., Lowry, D. P., Little, C. M., Morlighem, M., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Reese, R., Schlegel, N.-J., Shepherd, A., Simon, E., Smith, R. S., Straneo, F., Sun, S., Trusel, L. D., Van Breedam, J., van de Wal, R. S. W., Winkelmann, R., Zhao, C., Zhang, T., and Zwinger, T.: ISMIP6 Antarctica: a multi-model ensemble of the Antarctic ice sheet evolution over the 21st century, *The Cryosphere*, 14, 3033–3070, <https://doi.org/10.5194/tc-14-3033-2020>, 2020.
- 775 Shapiro, N. M. and Ritzwoller, M. H.: Inferring surface heat flux distributions guided by a global seismic model: particular application to Antarctica, *Earth and Planetary Science Letters*, 223, 213–224, <https://doi.org/10.1016/j.epsl.2004.04.011>, 2004.
- Siebert, M., Alley, R. B., Rignot, E., Englander, J., and Corell, R.: Twenty-first century sea-level rise could exceed IPCC projections for strong-warming futures, *One Earth*, 3, 691–703, <https://doi.org/10.1016/j.oneear.2020.11.002>, 2020.
- Siebert, M. J.: Glacial–interglacial variations in central East Antarctic ice accumulation rates, *Quaternary Science Reviews*, 22, 741–750,
 780 [https://doi.org/10.1016/S0277-3791\(02\)00191-9](https://doi.org/10.1016/S0277-3791(02)00191-9), 2003.
- Siebert, M. J. and Payne, A. J.: Past rates of accumulation in central West Antarctica, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004gl020290>, 2004.
- Siebert, M. J., Welch, B., Morse, D., Vieli, A., Blankenship, D. D., Joughin, I., King, E. C., Leysinger Vieli, G. J. L., Payne, A. J., and Jacobel, R.: Ice Flow Direction Change in Interior West Antarctica, *Science*, 305, 1948–1951, <https://doi.org/10.1126/science.1101072>,
 785 2004.
- Steinhage, D., Kipfstuhl, S., Nixdorf, U., and Miller, H.: Internal structure of the ice sheet between Kohnen station and Dome Fuji, Antarctica, revealed by airborne radio-echo sounding, *Annals of Glaciology*, 54, 163–167, <https://doi.org/10.3189/2013aog64a113>, 2013.
- Stokes, C. R., Tarasov, L., Blomdin, R., Cronin, T. M., Fisher, T. G., Gyllencreutz, R., Hättestrand, C., Heyman, J., Hindmarsh, R. C., Hughes, A. L., et al.: On the reconstruction of palaeo-ice sheets: Recent advances and future challenges, *Quaternary Science Reviews*, 125, 15–49,
 790 <https://doi.org/10.1016/j.quascirev.2015.07.016>, 2015.



- Sutter, J., Fischer, H., Grosfeld, K., Karlsson, N. B., Kleiner, T., Van Liefferinge, B., and Eisen, O.: Modelling the Antarctic Ice Sheet across the mid-Pleistocene transition—implications for Oldest Ice, *The Cryosphere*, 13, 2023–2041, <https://doi.org/10.5194/tc-13-2023-2019>, 2019.
- Sutter, J., Eisen, O., Werner, M., Grosfeld, K., Kleiner, T., and Fischer, H.: Limited retreat of the Wilkes Basin ice sheet during the Last Interglacial, *Geophysical Research Letters*, 47, e2020GL088131, <https://doi.org/10.1029/2020GL088131>, 2020.
- Sutter, J., Fischer, H., and Eisen, O.: Investigating the internal structure of the Antarctic ice sheet: the utility of isochrones for spatiotemporal ice-sheet model calibration, *The Cryosphere*, 15, 3839–3860, <https://doi.org/10.5194/tc-15-3839-2021>, 2021.
- The PISM authors: Repository for the Parallel Ice Sheet Model (PISM) [Codes], <https://github.com/pism/pism>, 2025.
- Veres, D., Bazin, L., Landais, A., Toyé Mahamadou Kele, H., Lemieux-Dudon, B., Parrenin, F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., et al.: The Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years, *Climate of the Past*, 9, 1733–1748, <https://doi.org/10.5194/cp-9-1733-2013>, 2013.
- Višnjević, V., Bodart, J., Hermant, A., Wirths, C., Spezia, E., and Sutter, J.: Isochronally constrained ice flow evolution of Dronning Maud Land, Antarctica during the Last Glacial Period, *EGU General Assembly Conference Abstracts*, p. 16702, <https://doi.org/10.5194/egusphere-egu24-16702>, 2024.
- Waddington, E. D., Neumann, T. A., Koutnik, M. R., Marshall, H.-P., and Morse, D. L.: Inference of accumulation-rate patterns from deep layers in glaciers and ice sheets, *Journal of Glaciology*, 53, 694–712, <https://doi.org/10.3189/002214307784409351>, 2007.
- Wang, Z., Chung, A., Steinhage, D., Parrenin, F., Freitag, J., and Eisen, O.: Mapping age and basal conditions of ice in the Dome Fuji region, Antarctica, by combining radar internal layer stratigraphy and flow modeling, *The Cryosphere*, 17, 4297–4314, <https://doi.org/10.5194/tc-17-4297-2023>, 2023.
- Winkelmann, R., Martin, M. A., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., and Levermann, A.: The Potsdam parallel ice sheet model (PISM-PIK)—Part 1: Model description, *The Cryosphere*, 5, 715–726, <https://doi.org/10.5194/tc-5-715-2011>, 2011.
- Winter, A., Steinhage, D., Creyts, T. T., and Eisen, O.: Radio-echo sounding isochrone depths in the East Antarctic Ice Sheet [Data set], <https://doi.org/10.1594/PANGAEA.895528>, 2018.
- Winter, A., Steinhage, D., Creyts, T. T., Kleiner, T., and Eisen, O.: Age stratigraphy in the East Antarctic Ice Sheet inferred from radio-echo sounding horizons, *Earth System Science Data*, 11, 1069–1081, <https://doi.org/10.5194/essd-11-1069-2019>, 2019.
- Wirths, C., Hermant, A., Stepanek, C., Stocker, T. F., and Sutter, J. C.: Sequence of abrupt transitions in Antarctic drainage basins before and during the Mid-Pleistocene Transition, *Nature Communications* [in press], <https://doi.org/10.1038/s41467-025-65375-x>, 2025.
- Yan, S., Koutnik, M. R., Blankenship, D. D., Greenbaum, J. S., Young, D. A., Roberts, J. L., van Ommen, T., Sun, B., and Siegert, M. J.: Holocene hydrological evolution of subglacial Lake Snow Eagle, East Antarctica implied by englacial radiostratigraphy, *Journal of Glaciology*, pp. 1–38, <https://doi.org/10.1017/jog.2025.15>, 2025.
- Young, D., Schroeder, D., Blankenship, D., Kempf, S. D., and Quartini, E.: The distribution of basal water between Antarctic subglacial lakes from radar sounding, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 374, 20140297, <https://doi.org/10.1098/rsta.2014.0297>, 2016.