

Quantifying atmospheric and land drivers of hot temperature extremes through explainable Artificial Intelligence

Arnau Garcia Mesa¹, Lluís Palma¹, Markus Donat^{1,2}, Stefano Materia^{1,3}, Bruna Gràvalos Talló¹, and Raül Marcos Matamoros⁴

¹Earth Sciences Department, Barcelona Supercomputing Center (BSC), Barcelona, Spain

²Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

³CMCC Foundation–Euro–Mediterranean Center on Climate Change, Bologna, Italy

⁴Facultat de Física, Universitat de Barcelona, Diagonal 645, 08028 Barcelona, Spain

Correspondence: Arnau Garcia Mesa¹ (arnau.garcia@bsc.es)

Abstract. Different drivers have been shown to play a central role in modulating the occurrence and intensity of summer temperature extremes, yet their individual contributions remain difficult to quantify. In this study, we develop an explainable machine-learning framework to disentangle the respective influences of large-scale atmospheric circulation, soil-moisture anomalies, and rising CO₂ concentrations on boreal-summer temperature extremes at six locations across Europe and North Africa with different characteristics of land–atmosphere coupling (Córdoba, Lyon, Hannover, Stockholm, Belgrade, and Marrakech). Using SHapley Additive exPlanation (SHAP) values, we find that the atmospheric circulation consistently dominates model explainability across all locations, contributing to ~~67–90~~67–82 % of the total mean SHAP, with the geopotential at 500 hPa field contributing the most. Soil-moisture influence exhibits a northward gradient: negligible at Marrakech (~~0.5–1.4~~ %), moderate at Córdoba (~~7.7–12.0~~ %), and substantial at Lyon (~~15–21.4~~ %). Additionally, ~~negative correlations between~~
5 ~~soil-moisture-standardized the dominant shallow soil-moisture signal shows an inverse relationship between moisture~~ anomalies and SHAP values~~across three depth levels corroborate~~, consistent with the amplifying effect of land drying on heat extremes, while attribution for deeper layers is weaker and less robust across locations. We demonstrate the robustness of these findings to a less stringent (80th percentile) extreme definition. Furthermore, the identified driver contributions are consistent when using alternative observational data for temperature extreme definition and for computing SPI/SPEI drought indices as
15 proxies for soil moisture, with SPEI showing a closer alignment to the original ERA5-Land results. We also illustrate the methodology for case studies of two individual events, heatwaves occurring in Córdoba (Spain) 2021 and Hannover (Germany) 2018, which reveal a pronounced spatial pattern in the distribution of SHAP values for the circulation predictors. They also confirm the enhanced role of the land component in regions of Northern Europe and reveal a contribution of the anthropogenic factor through CO₂ concentrations, even for specific events. These ~~insights enhance our understanding of the physical~~
20 ~~mechanisms behind temperature extremes~~results align with the established physical roles of atmospheric circulation and soil moisture modulation for temperature extremes, and demonstrate the methodological potential of explainable artificial intelligence ~~methods~~ to quantify the ~~contributions from different drivers of hot temperature extremes~~relative contributions of these drivers in a data-driven framework.

1 Introduction

25 Understanding weather extremes is of vital importance due to their impacts on society and ecosystems. Events such as extreme precipitation, severe droughts, and intense heatwaves pose growing threats to biodiversity and sectors like agriculture, energy, health or the economy (e.g., Zscheischler et al., 2018). As the climate warms, evidence from observations and model simulations shows that these events are becoming longer, more intense and more frequent (Intergovernmental Panel On Climate Change, 2023). Consequently, predicting, characterizing, and understanding these events is crucial for the development
30 of early-warning systems and adaptation strategies in a climate change context.

Much progress has been made in identifying the physical processes driving temperature extremes, although many mechanisms are still not well understood due to difficulties in quantifying their interactions and responses to climate change (Barriopedro et al., 2023). The changing behaviour linked to the positive trends in global temperatures poses challenges for establishing a unique definition or common metrics describing hot extremes (Perkins and Alexander, 2013).

35 Hot extremes are primarily driven by a combination of atmospheric, ocean, and land surface conditions, including persistent high-pressure systems and soil moisture deficits (Barriopedro et al., 2023). In this study, we discard the ocean drivers because we are focusing on the short term variability drivers of hot extremes. Additionally, we include atmospheric CO₂ concentrations to capture the multiyear trend of hot extreme events. Many studies focus ~~in on~~ understanding the role of these land–atmosphere interactions, such as the positive soil–moisture–temperature feedback, and its amplification role during high-pressure conditions
40 (Materia et al., 2022; Quesada et al., 2012). This amplification becomes especially relevant in a changing climate due to the evidence of regional shifts of wet, dry and transitional evaporative regimes (Hsu and Dirmeyer, 2023). Seneviratne et al. (2006), for instance, highlight the need of including land–atmosphere interactions in regional simulations to correctly modulate the increase in summer temperature variability (and resulting temperature extremes) in central and Eastern Europe due to increasing greenhouse gas concentrations.

45 Regarding the role of the atmospheric circulation, many studies show the direct link of hot extremes to atmospheric circulation patterns, such as persistent double jets over Eurasia (Rousi et al., 2022), mid-tropospheric deep depressions over the North–Atlantic (D’Andrea et al., 2024), or episodes of Saharan warm air intrusion in the West Mediterranean (Cos et al., 2024).

In addition, growing attention is being paid to heatwave-driving mechanisms related to other elements of the Earth system, like particular patterns in preceding sea surface temperatures identified as precursors of summer heatwaves in Europe (Beobide-
50 Arsuaga et al., 2023).

Complementing physical understanding, new tools and methodologies are being actively explored to improve the identification and prediction of temperature extremes. In particular, Artificial Intelligence (AI) and Machine Learning (ML) approaches are emerging as powerful complements to traditional numerical modelling techniques (Camps-Valls et al., 2025). Moreover, ongoing developments in explainable AI (XAI) offer promising pathways to interpret model behaviour, allowing researchers
55 to assess the contribution of individual drivers to extreme events (Materia et al., 2024). This capability is critical not only for attribution purposes, but also for potentially gaining new insights into the underlying physical mechanisms.

For example, Toms et al. (2020) demonstrate how explainable ML methods such as layerwise relevance propagation (LRP) and backward optimization can be used to interpret neural network predictions in climate applications, revealing scientifically meaningful relationships between input patterns and network outputs. Davenport and Diffenbaugh (2021) employs-employ LRP to analyse large-scale circulation patterns associated with extreme precipitation. Other explainable AI (XAI) techniques such as SHAP have also been used to assess the relative importance of different climate drivers; for instance, Uckan et al. (2024) apply SHAP values within a Random Forest framework to quantify the contribution of multiple heatwave drivers at the global scale.

In this study, we aim to disentangle the respective roles of atmospheric circulation, soil moisture and anthropogenic climate change in driving extreme temperature events during the boreal summer (JJA), using an explainable ML modelling framework. With this goal in mind, we aim to develop a framework that is both predictive and diagnostic.

First, we classify daily extreme events using the 90th percentile threshold of the climatological maximum temperature distribution. Then, we build a robust extreme/no-extreme binary classifier using supervised ML, with lagged anomalies of large-scale circulation variables and soil moisture, and atmospheric CO₂ concentrations as inputs. To understand the drivers learned by the ML model during the prediction task, we apply the SHAP explainability technique to quantify the contribution of each input feature to the model's predictions (Lundberg and Lee, 2017a). First, we compute general aggregated SHAP values for each variable to obtain a general characterization of feature relevance. Then, we proceed by analysing specific SHAP explanations for two case studies to investigate region-specific driver behaviour: the August 2021 heatwave in Córdoba and the July 2018 heatwave in Hannover. These events were selected to represent contrasting soil-moisture feedback regimes – arid-versus-temperate-dry/transitional versus transitional/wet – with differing soil moisture variability and land-atmosphere coupling strengths (Hsu and Dirmeyer, 2023). We extend our explainable ML analysis to evaluate the robustness of our driver attribution to different extreme thresholds (80th and 95th percentiles) and to the use of alternative observational data for the extreme event classification and the land component features.

In section 2 we discuss the data and methodology, including the datasets used, the domain for circulation data, the selected locations based on different soil-moisture feedback regimes, the deep learning model used and the explainability method. In Sect. 3 we show and discuss the results: we display the extreme classification results, give a statistical assessment of the model, and provide the SHAP explainability results. Lastly, conclusions of the study can be found in Sect. 4.

2 Data and methodology

2.1 Datasets

We use data from ERA5 (Soci et al., 2024)(Hersbach et al., 2020; Soci et al., 2024), the fifth-generation atmospheric reanalysis from the European Center for Medium-Range Weather Forecasts (ECMWF), which provides a globally complete, gridded data from 1940 to the present. The long-term availability and spatial completeness of reanalyses make them suitable for studying climate trends, variability, or extreme events despite substantial uncertainties across datasets (Donat et al., 2014). We use daily data using the spatial domain discussed in 2.3. We use atmospheric circulation fields that represent the state of the

90 atmosphere and constitute the large-scale variables in the study: Geopotential at 500 hPa ($g500$) [m^2s^{-2}], Geopotential at 200 hPa (~~$g500$~~ $g200$) [m^2s^{-2}], and Sea Level Pressure (psl) [hPa].

For land-surface variables, we use the ERA5-Land dataset, ~~which has a better representation of land-surface variables due to its 9km horizontal resolution~~. Soil moisture variability is more consistent in ERA5-Land because it does not assimilate screen-level observations to adjust soil moisture (unlike the ERA5 reanalysis) (Muñoz-Sabater et al., 2021). The variables we use are daily maximum 2-meter temperature (TX) to define our target events, and soil moisture at three depth levels ($swvl1$, $swvl2$, and $swvl3$) as predictors. TX from ERA5-Land is only used to define hot extremes, as the prediction target of the ML model.

Lastly, we include seasonal atmospheric CO_2 concentrations to account for the trend of climate change. We employ U.S. National Oceanic and Atmospheric Administration (NOAA) measurements collected at the Mauna Loa Observatory in Hawaii. The concentrations used have a constant value for each summer and are a seasonal mean. By explicitly including CO_2 , our framework controls for the long-term warming trend as a potential additional driver of heat extremes, allowing for a cleaner attribution of the physical drivers.

Circulation and land variables are used as lagged standardized anomalies (explained in [sec. 2.4](#)). For each prediction day, the model is trained on features derived from values recorded on preceding days. For the local-scale, soil moisture is averaged over the previous seven days to account for its relatively low daily variability (Miloshevich et al., 2023). The values of circulation variables are taken from the day before the prediction, corresponding to a one-day lag in atmospheric features. The inclusion of up to three lags for the circulation features was tested, but we decided to keep a simpler model because the accuracy did not improve by including more lags (not shown).

Similar variable sets using atmospheric circulation and land features have been used in Koh et al. (2025), where they developed a machine-learning framework combining spatial extreme-value theory and adapted loss functions to study large-scale heat extremes, or in Miloshevich et al. (2023), who predict the probability of occurrence of long-lasting 14-day heatwaves over France.

2.2 Observational data and drought indices

To test the robustness of our results to the data source, we supplement the analysis based on reanalysis with observational data for the daily maximum temperature and the land component. For the atmospheric circulation features, we keep the data from ERA5. For the local-scale daily maximum temperature used in the extreme event definition, we use directly the E-OBS gridded dataset (version 29.0) of TX (Cornes et al., 2018).

Due to the lack of long-term, consistent observational soil moisture records, we employ the Standardized Precipitation Index (SPI) (McKee et al., 1993) and the Standardized Precipitation Evapotranspiration Index (SPEI) (Vicente-Serrano et al., 2010) as proxies for soil moisture anomalies, computed with the E-OBS data. These indices serve to validate the robustness of our ERA5-Land results against observational data, providing a reliable alternative for historical comparison. These indicators have already been assessed as surrogates of in situ and satellite soil moisture observations (Scaini et al., 2015; Wang et al., 2015; Yuan et al., 2020), with generally larger association when potential evapotranspiration is included. We use the Hargreaves

method (Masmoudi-Charfi and Habaieb, 2014) for computing the evapotranspiration, essential for the SPEI calculation, and
125 the gamma distribution to standardize the data. Precipitation and temperature data are sourced from the E-OBS dataset. As for
soil moisture, both SPI and SPEI grid-points are averaged spatially over a 100kmx100km region.

The timescale for the index computation was selected by computing the Spearman correlation of the drought indices with
the three soil moisture levels of the original model (see figures C1 and C2). Soil moisture is a variable with a memory that
extends beyond a few days (Vicente-Serrano et al., 2010), so the drought index must be computed on a scale that reflects this
130 persistence. Based on the correlation analysis, the selected timescale for the SPEI and SPI indices is 30 days (SPEI30d and
SPI30d). This timescale shows the best correlation of the indices with *swvl1* and *swvl2*, which we show to be in general the
most relevant depth levels in our explainability analysis (see [sec. 3](#)). We also computed the correlation using a 7-day rolling
mean of the soil moisture data—the same preprocessing applied before inputting soil moisture to the model—and obtained
very similar results, with correlation values not differing more than 0.07. Additionally, we computed the drought indices using
135 ERA5 data to validate the correlations computed with the E-OBS gridded data, with correlation between datasets not differing
more than 0.1. In the correlation analysis, we did not lower the timescale of the drought indices computation (e.g. 15 days)
to stay within the original monthly definition of the indices (Vicente-Serrano et al., 2010), and to avoid fitting the normalized
distribution to data that may not be sufficiently smooth.

2.3 Atmosphere data domain and target locations

140 With the atmospheric pressure and geopotential height fields, we aim to capture the general circulation influencing the locations
where we predict extreme temperature events. The large-scale domain must be large enough to cover the area of interest, but
not too large to avoid including unrelated atmospheric variability (Mahlstein et al., 2015). Fig. 1 shows the selected domain
marked as a red box. The full North Atlantic Ocean is included, since studies link this region to the development of extreme
temperature events in various European regions (Bischof et al., 2023). A large part of North Africa is included as well to capture
145 warm and dry intrusions that may enhance the likelihood of extremes (Cos et al., 2024). The northern and eastern limits are
extended enough to include synoptic-scale circulation patterns.

We study the roles of circulation, soil moisture and anthropogenic climate change during extreme temperature events in six
regions around Córdoba, Marrakech, Lyon, Belgrade, Hannover, and Stockholm (see [fig. 1](#)). For each site, local-scale data
is averaged over a surrounding 100 km × 100 km region, to smooth out unwanted variations across space. The chosen locations
150 span different evaporative regimes: from water-limited (Marrakech) to energy-limited (Stockholm). Location choice is also
partly based on literature: Córdoba and Marrakech are studied in Lemus-Canovas et al. (2024), Stockholm's 2018 heatwave in
Wilcke et al. (2020), and Iberia and Central Europe heatwaves in Khodayar Pardo and Paredes-Fortuny (2024). Belgrade was
selected to provide a more eastern perspective, potentially influenced by sea surface temperatures and atmospheric anomalies
in other regions discussed in Beobide-Arsuaga et al. (2023).

155 Lastly, the robustness analysis using observational data (discussed in 2.2) could not be carried out for Marrakech, due to
missing data for several years (see Fig. 4).




Figure 1. Large-scale data domain and locations selected for the local-scale data in the prediction target. The red box indicates the domain used in the study for the ERA5 data, with longitudes ranging from 54° W to 70° E and latitudes from 14° N to 71° N. Locations are illustrative for different climate regimes in Europe.

2.4 Climatology, anomalies and the extreme temperature definition

Both the climatology and 90th percentile are computed only for the reference period 1950–2000, within the model’s training period (1950–2014). This avoids an indirect provision of future information to the model, which would leak evaluation data into training. The training period was extended to 2014 to include a period with a clear trend in the yearly count of extremes (see [fig Fig. 4](#)).

First, we calculate the raw climatology by taking the daily average. Then, as in Mahlstein et al. (2015), we apply a local polynomial regression (LOESS) to obtain a smoothed profile. After a few tests with the window timescale, we selected a 30-day smoothing window for both climatology and standard deviation. Anomalies are computed daily by subtracting the smoothed climatology from the original data, and then standardized using the standard deviation. Subtracting the climatology removes the annual cycle, and applying the LOESS smoothing removes the short-term variability. Extended detail on LOESS is given in appendix A1.

Using the extreme definition in Perkins and Alexander (2013) an extreme event is defined when the daily mean maximum temperature exceeds a specified percentile of the climatological distribution. Following Beobide-Arsuaga et al. (2023), the percentile is calculated over a centered 5-day window using the reference period, and then we apply LOESS smoothing with a 30-day window. We use a 5-day rolling window prior to the LOESS smoothing to increase the number of samples available for estimating the 90th percentile for each calendar day. The window is centered in the day for which the percentile is computed. The primary analysis in this study uses the 90th percentile, but to test the robustness of our results to the extreme event definition, we conduct additional analyses using the 80th and 95th percentiles, and all are computed following the same procedure.




Figure 2. Climatology and percentiles for the region of Córdoba. Day of the year (x-axis) and daily maximum 2-meter temperature from ERA5-Land (y-axis). We show the raw climatology and percentiles, together with the 5-day window pull and LOESS smoothing versions. The text box shows the total frequency of heat wave events, and the separated frequencies for the reference period 1950–2000, and the rest of the time series (2001–2024).

175 2.5 The deep learning model

180 Although we use daily data since The model is trained on data from 1950 ,the number of extreme days based on to 2013. Validation years are chosen by randomly selecting contiguous blocks of years (see Table B1), while the testing period spans 2014 to 2024. Extreme days are defined using the 90th percentile remains low (see fig-2threshold (i.e., approximately 10% of all days), which results in a class imbalance between extreme and non-extreme days (see Table B1). This imbalance is one of the central challenges we need to cope with during the represents a central challenge during model training, to prevent a default to the as machine learning algorithms may default to predicting the majority (non-extreme class. On the other hand, the) class if no specific strategy is used to address it. At the same time, the relatively limited number of extreme days cases increases the risk of overfitting in our binary classification task when using many parameters, something that this binary classification problem when using models with many parameters. This constraint needs to be taken into account when defining the considered when

185 designing the model architecture and training procedure. The imbalance ~~problem is even worse if one uses higher percentiles~~
~~in the extreme definition~~ issue becomes even more pronounced when higher percentile thresholds are used to define extremes,
as the number of available training samples decreases further. (see Table ~~??~~ B1). In this study, we treat the class imbalance
in the training of the ML model with weighting of the Loss function. By doing this, we penalize more the wrongly classified
extreme samples.

190 The main contributor to high parameter dimensionality are the large-scale fields, which include spatial dimensions at 0.25°
resolution, unlike local-scale data that is area-averaged and provides time-only vectors. Flattening large-scale data would result
in $n_{var} \cdot n_{lags} \cdot n_{lat} \cdot n_{lon}$ inputs, making the model overly complex. We first reduce dimensionality by re-gridding large-scale
data to 1° , directly reducing computational cost and overfitting risk. We think this coarser resolution still captures atmospheric
circulation features for extreme temperatures.

195 We use a model architecture designed to capture land–atmosphere coupling (see ~~fig~~Fig. 3). It includes a Multi Layer Percep-
tron (MLP) for land and CO₂ data, and a ConvNeXt convolutional neural network (CNN) (Liu et al., 2022) for spatial features
from circulation, which, to our knowledge, ~~it~~ has not been used before in driver quantification studies. Their outputs are con-
catenated and passed to a final MLP for binary classification of extreme or non-extreme on a daily basis. This architecture is
hereafter referred to as the *CombinedModel*.

200 The MLP for the Land State + CO₂ inputs ~~has receives an input vector of dimension d_{in} , followed by~~ two hidden layers
of 15 and ~~eight nodes, inspired by Mayer and Barnes (2021), and the MLP for the final classification has 8 nodes. The final~~
~~classification MLP then takes the concatenated outputs from the Local and Large-scale branches (with dimensions 8 and 16,~~
~~respectively) as its input. These features are processed through~~ three hidden layers ~~with of~~ 64, 32, and 8 nodes ~~before reaching~~
~~the final output layer. For the spatial feature extraction, a custom ConvNeXt architecture is employed. The model accepts input~~
205 ~~tensors with C channels (corresponding to the number of Large Scale features) and utilizes a patch size of 4. The ConvNeXt~~
~~here consists of four stages with depths of [1, 2, 2, 1] and corresponding layer dimensions (channels) of [4, 6, 6, 16]. A dropout~~
~~rate of 0.05 is applied to prevent overfitting during training.~~ The architecture of the ConvNeXt model is explained in detail in
Liu et al. (2022). ReLU is used as activation function in the MLPs, and softmax applied at the output to yield probabilities
(Goodfellow et al., 2016). A 0.5 threshold determines the predicted class, kept constant throughout the study. Additionally, we
210 use a 20-member ensemble to filter out noise and handle part of the epistemic uncertainty of the model. These ensembles are
generated by changing the initial random seed (detailed in 2.7). Similar hybrid model architectures have been used in the past
(Gordon et al., 2023; Mayer et al., 2024).

The optimizer used for the CombinedModel is the AdamW optimizer (Loshchilov and Hutter, 2019). We train the model
with an 80 % and 20 % random split of the samples for the training and validation respectively, and apply a random shuffle.
215 The Cross-Entropy Loss function is used (see A2), with specific weights applied to the non-extreme and extreme classes to
address the class imbalance. The Optuna Python package is used for the tuning of the following hyperparameters of the ML
model: batch size, learning rate, weight decay and class weights of the predicted classes. The model is trained for 75 epochs
unless validation loss increases for a total of five epochs. The score used in the hyperparameter tuning process is the balanced
accuracy minus the final validation loss, which is the best validation loss found before early-stopping.




Figure 3. Diagram of the Combined Model. Includes a simple MLP for the Land-State plus a ConvNeXt model for the Atmosphere state, which are then combined using a final MLP that does the binary classification. An ensemble of 20 members is used during training to add robustness and treat the model’s epistemic uncertainty.

2.6 Explainability of the machine learning model using SHapley Additive exPlanation values

To quantify the contribution of each input feature to the model’s predictions, we use SHapley Additive exPlanations (SHAP), a game-theoretic approach that explains the output of any machine learning model by attributing it to the input features (Lundberg and Lee, 2017a).

Positive SHAP values mean a positive influence on the likelihood of the predicted class by the model, while a negative SHAP value means that the feature decreases the likelihood of predicting the class. The magnitude of the value quantifies the influence of the feature, and SHAP values are zero for irrelevant. SHAP values are an additive feature attribution method, meaning that the contribution of each feature can be computed independently, and all values summed up.

The sum of the SHAP values across all features equals the model output (prediction) minus the expected (baseline) prediction :

$$f(x) = \phi_0 + \sum_i^N \phi_i, \quad (1)$$

where $f(x)$ is the model prediction for the input x , ϕ_0 is the baseline (computed by averaging the model prediction over training data), ϕ_i is the SHAP value for the feature i and N is the total number of features in the model. The additive nature of SHAP values allows us to compute a value per grid-point, highlighting the regions the model finds most important, and to spatially aggregate these values into global importance or combine them across features, as ~~done in~~ we do in sec. 3.2.

We compute SHAP values in the testing period ~~using GradientExplainer~~(years 2014-2024) using GradientExplainer (Lundberg and Lee, 2017), which is designed for differentiable models, such as deep neural networks, and estimates them by using an extension of the

integrated gradients method (Sundararajan et al., 2017). We believe this choice to be appropriate given the evaluation metrics results for the Integrated Gradients method shown in Bommer et al. (2024), though no XAI method evaluation was carried out for our study. Additionally, the method requires selecting a baseline to serve as a reference in the SHAP computation. Following best practices suggested by Mamalakis et al. (2023a), the baseline was constructed by randomly selecting 300 samples from the training period that had non-extreme labels. Using only non-extreme days for the baseline is appropriate in this study, as our focus is on identifying the drivers of predicted extreme events with the aforementioned ML model.

2.7 Creating robust explanations

Since our focus is to study the contribution of the atmospheric circulation, the soil moisture, and atmospheric CO₂ concentrations during extreme temperature events, we want to ensure that our explanations are done for predictions for which our model is confident. To do so, we take an ensemble of trained models with twenty members, adding robustness to our output probabilities and computed SHAP values. The members are created by setting deterministically twenty different seeds, initializing the random processes present in the framework. On top of that, we analyse the general explainability results for a certain percentage of top most confident predictions, following the approach of studies like Mayer and Barnes (2021).

We assess model performance using balanced accuracy (BA) and the Area Under the Receiver Operating Characteristic Curve (AUC), which are well-suited for imbalanced classification tasks.




Figure 4. Extremes yearly count for the six locations for both ERA5-Land and E-OBS data. Yearly extreme count for the full period of used data in ERA5 and (1950–2024) E-OBS (1950–2023 for E-OBS). Years in the x-axis and count (number of extreme days) in the y-axis. The text box shows the mean count for the period 1950–2000 to verify the percentile definition of extreme in both datasets.

3 Results and discussion

Fig. 4 shows the yearly count of summer extremes (true labels) for the six selected locations, resulting from the extreme definition given in 2.4. We show the results of the extreme classification both for the ERA5-Land and E-OBS datasets. Clear warming trends are apparent for Córdoba and Lyon regions. Belgrade region also shows a visible trend after approximately 1960. To quantify these observations, we performed a formal linear trend analysis across different time windows (see Fig. B1). A generalized acceleration in the frequency of extremes is visible when comparing the full record against the period starting in 1980. In the most recent period (since 2000), a slight decrease in the trend magnitude—and a corresponding loss of statistical significance—is observed in Lyon, Marrakech, and Belgrade. A similar slight decrease in magnitude is noted for Córdoba when compared to the 1990–onwards period, though the trend there remains statistically significant. The average count during summer for the period 1950–2000 is approximately 10 for all locations, which is expected considering our 90th percentile definition of extreme. When comparing the classification results for ERA5-Land and E-OBS datasets, we see a general agreement except for some periods and locations like the 50s decade in Córdoba, or the years 1950 and 1952 in Belgrade.

3.1 Prediction and statistical assessment of the models

For the primary analysis using the 90th percentile threshold, ~~table ?? in ??~~, (see Table 1) gives relevant statistics of the trained CombinedModel on the test data using the 50% most confident predictions. Fig. B2 shows how we assess the CombinedModel model robustness by plotting the evolution of accuracies when varying the confidence threshold. For each member individually, the CombinedModel shows a smooth increase in both balanced and extreme accuracy across all locations. The improvement in both accuracy measures with increasing network confidence is an expected behavior. For some locations, low percentages of confidence (see ~~fig~~Fig. B2) show a trade-off between extreme and non-extreme accuracies, with the former increasing and the latter decreasing, until a minimum is reached and non-extreme accuracy starts improving again. We did not explore this behaviour in depth, but it may be related to how we handle class imbalance.

In Fig. B3 we show the ROC curve change when changing the percentage of confidence. When using the ensemble of 20 members, most locations show a displacement towards area under the ROC curve equal to 1.

The model trained on the 80th percentile definition shows a similar statistical robustness for the accuracies and ROC curves B4. The increased number of extreme events mitigates class imbalance, resulting in smooth and consistent improvements in balanced accuracy and AUC values as the confidence threshold increases. This indicates the model is well-trained, and its predictions are reliable for this definition. In contrast, the model performance for the 95th percentile definition is notably weaker for most locations, with the balanced accuracy behaving poorly when increasing the percentage of confidence in the predictions. The drastically reduced number of extreme events (see ~~table ??~~Table 1) presents a significant challenge for training, leading to less stable behavior in accuracy metrics and ROC curves ~~as a function of prediction confidence due to the strong class imbalance~~ (see Fig. B5). The class imbalance and lack of extreme events samples is a known issue in ML application for weather extremes (Miloshevich et al., 2023; Sha et al., 2025), so the results using the 95th percentile are not surprising.

285 Due to this lack of robust predictive skill, we do not consider the explainability results for the 95th percentile definition to be trustworthy.

Finally, the model trained with observational TX and SPI/SPEI indices for the 90th percentile also demonstrates robust predictive performance, providing confidence in the reliability of its explanatory outputs (not shown).

Table 1. Statistical assessment of the CombinedModel for the different locations, considering the 50% most confident predictions.

For the test data, we show the extreme class accuracy (EA), the non-extreme class accuracy (NEA), and the balanced accuracy (BA), which accounts for false positives and false negatives. TP, FP, TN, and FN stand for true positives, false positives, true negatives, and false negatives respectively. The last column shows the area under the ROC curve (AUC).

Site	EA[%]	NEA[%]	BA[%]	TP	FP	TN	FN	AUC
Cór.	100±0	87±2	93±1	216	37	252	1	0.96
Lyon	100±0	87±2	94±1	216	37	253	0	0.96
Hann.	100±0	75±2	88±1	169	84	253	0	0.96
Stock.	100±0	70±2	85±1	145	108	253	0	0.91
Bel.	100±0	77±2	89±1	178	75	253	0	0.93
Marra.	100±0	77±2	88±1	176	77	253	0	0.95

3.2 SHAP values

290 Taking the 50% most confident extreme predictions, Fig. 5 shows the mean SHAP values across samples in the test dataset for each of the features in the model. The results are not sensitive to the confidence threshold. We calculate a global SHAP value for the Land and Atmosphere compounds by summing the contributions of their respective features, which can be done thanks to SHAP values being an additive feature attribution method. To determine the SHAP percentage for a feature, we first calculate the mean of its absolute SHAP values over the entire time series. This time-averaged mean is then expressed as a
295 percentage of the total mean SHAP value contributed by all features.

The atmospheric circulation has the largest importance (~~67-90~~67-82 %) at all locations. Apart from being intuitive, this result is in line with for instance Uckan et al. (2024), who show that geopotential height at 500 hPa is the dominant driver of hot extremes worldwide, particularly in mid-latitudes, and that the relevance of atmospheric drivers increases with the duration of the hot event. The Land SHAP value is more important in northern regions. In Marrakech, the land SHAP percentage is
300 ~~practically zero~~very small (1.4 %), which is reasonable given the region is generally very dry during the summer, therefore the interannual variability of soil moisture is low and consequently coupling with atmosphere very little. Córdoba shows a higher percentage (~~7.7~~12.0 %) explained by the land SHAP compared to Marrakech. The importance of swvl3 in Córdoba is slightly larger than swvl1 and swvl2, consistent with top soil layers being fully dissicated in summer. The land SHAP value increases for higher-latitude locations, being specially important in Lyon. This result is in line with Jach et al. (2022), who assessed how
305 changes in the mean temperature and moisture influence the land–atmosphere coupling strength over Europe.




Figure 5. Extreme class prediction mean SHAP value percentage for all locations using the CombinedModel taking the 50% most confident predictions. The x-axis represents the percentage contribution of each feature relative to the total sum of mean absolute SHAP values. Error bars indicate the Standard Error of the Mean (SEM) derived from the 20-member ensemble, representing the uncertainty across different model initializations. The colored text box provides the aggregated percentage contributions for Land and Atmospheric components, as well as the CO₂ contribution, with associated ensemble uncertainties ($\pm$ SEM). The balanced accuracy of the model is also indicated.

As for the large-scale fields, there is a predominance of the geopotential at 500hPa, with only Marrakech having sea level pressure being more important than the atmospheric fields at higher altitude.

The atmospheric CO₂ feature has a higher importance in regions that exhibit a stronger trend in extreme temperature days (see Fig. 4). Except for ~~Stockholm and~~ Hannover, the importance of the CO₂ feature is generally high ($>10\%$). These differences among locations align with the work of Sippel et al. (2020) in that different regions show different responses to climate change, with the forced fingerprint manifesting through varying signal-to-noise ratios across the globe.

Lastly, ~~in Fig. 6 we can observe the negative relationship between the~~ illustrates how SHAP values vary with standardized anomalies of the ~~soil moisture levels and the contribution to the extreme class prediction. The results are consistent, with negative anomalies across levels contributing positively to predicting soil-moisture predictors. For the upper soil layer (swvl1), negative anomalies are generally associated with positive SHAP contributions to the extreme class and the other way around for positive anomalies. The figure also reflects the larger spread and importance for the first soil moisture level at this timescale,~~




Figure 6. SHAP values swvl[1,2,3] across samples test phase for the 50% most confident predictions and extreme class. The x-axis represents the SHAP value of the sample and the y-axis the soil moisture level feature used in the model. The colour bar shows the value of the standardized anomalies of the soil moisture level.

especially in northern regions. This key negative correlation is also present in the SHAP results for both, consistent with the expected amplifying role of surface drying on hot extremes. In contrast, the relationship for the intermediate layer (swvl2) is less robust across locations, with weak mean contributions and substantial uncertainty. This behaviour likely reflects the strong correlation between neighbouring soil layers, which limits the model's ability to uniquely attribute predictive skill to swvl2 independently of swvl1. The deepest layer (swvl3) shows a more location-dependent behaviour, with a clearer contribution in Córdoba. The inverse SHAP-anomaly relationship identified for the dominant shallow soil-moisture contribution is also found in the 80th percentile model (not shown) and the model using SPEI (see Fig. D3), further confirming the robustness of the identified-land-drying feedback mechanism identified by the model across different methodological choices.

The discussed explainability results are robust to the definition of an extreme event. For the 80th percentile model, the SHAP values (see Fig. D2) show the same main characteristics: the circulation component remains dominant (64.6–88.9/63.9–83.7%), and the land component still exhibits a northward gradient in importance, though less pronounced. Except for Córdoba and Marrakech, the relative contribution of soil moisture is, however, generally smaller than for the 90th percentile model. This would suggest a more marginal role of soil moisture in less severe heat events for which the land-atmosphere feedbacks, a key amplifier of extreme temperatures, may not be fully activated. The fact that the land component contribution is similar for Marrakech and Córdoba in the two extreme definitions might be an indication of the land-atmosphere coupling being rarely activated there in summer due to the scarce coupling in very dry conditions, independently of the extreme definition. Land variables could play an even more important roles for more restrictive percentiles like the 95th percentile. However, as




Figure 7. Same plot as in Fig. 5 but for the model using E-OBS observational data and the SPEI30d index as a proxy for the soil moisture feature. The 90th percentile is still used for the extreme event definition.

mentioned in section 3.1, we do not show the explainability results for this case due to the lack of stability in the accuracy metrics when using this higher percentile as threshold to identify extremes.

To further constrain our analysis with observational records and evaluate the framework's performance beyond reanalysis data, we examine the SHAP results using observational data from E-OBS for TX and the SPEI30d drought index as a proxy for soil moisture (Fig. 7).

The driver contributions are ~~also overall~~ consistent when using observational data from E-OBS instead of ERA5-Land for TX and the SPEI drought index as a proxy for the soil moisture variables. In general, the XAI results reproduce the established north-south gradient in land driver importance and the dominant importance of the atmospheric circulation component (see Fig. ??). Córdoba is the only location that shows a notable discrepancy from the original model when using the swvl features, with the value being approximately four times smaller when calculated with the SPEI index. Nonetheless, the finding that Córdoba's value is smaller than those of more northern locations remains robust, and the small value is physically consistent considering the very dry conditions in summer for this location. In Fig. D3 we see that we also have a negative correlation between SHAP value and SPEI30d feature values.




Figure 8. SHAP values for extreme class predictions for the Córdoba case study. a): SHAP values temporal evolution for the circulation features and the three soil moisture levels. We additionally include temperature standardized anomalies (shaded orange area). The true and predicted labels are shown at the bottom, with orange indicating and extreme (class 1) and gray indicating a non-extreme day (class 0) b): SHAP spatial distribution for the peak-day of temperature anomaly (14th of August). SHAP values with absolute magnitude smaller than 0.008 are masked to improve visual clarity. g500 contours for the day prior to the peak are included. c): Standardized anomalies of geopotential at 500hPa in colour and data in contours for the geopotential at 500hPa for the day prior to the temperature anomaly peak (13th of August). The magenta dot in panel *b* and *c* marks the location of Córdoba.

When using the SPI index, the XAI results for the land component are less similar to the original ERA5-Land soil moisture results (see Fig. D4). Although a correlation analysis between the indices and soil moisture shows that SPEI generally correlated slightly better with soil moisture than SPI (see Fig. C1), the differences are relatively small. Nevertheless, the XAI outcomes reveal that SPI captures less of the original soil moisture signal. This aligns with physical understanding, as SPEI, by incorporating evapotranspiration, provides a more comprehensive representation of surface moisture balance and its coupling with the atmosphere than SPI, which is based solely on precipitation. This suggests that for the future application of this framework to climate models, SPEI is a suitable proxy that avoids inter-model differences in representation of soil layers, and therefore allows more consistent evaluation of the relationships across different models and observations/reanalysis.

is a suitable proxy that avoids inter-model differences in representation of soil layers in therefore allows more consistent evaluation of the relationships across different models and observations/reanalysis

Overall, the feature with the largest importance shown by the XAI results for the *g500* field. As a result, we decided to test the robustness of the model when using only this field to represent the atmospheric circulation features, obtaining similar results to the ones discussed above (see Fig. D5). The predominance of the *g500* field is not surprising, as it is known to have a key role




Figure 9. SHAP values for extreme class predictions for the Córdoba case study. a): SHAP values temporal evolution for the circulation features and the three soil moisture levels. We additionally include temperature standardized anomalies (shaded orange area). The true and predicted labels are shown at the bottom, with orange indicating and extreme (class 1) and gray indicating a non-extreme day (class 0) b): SHAP spatial distribution for the peak-day of temperature anomaly (7th of August). SHAP values with absolute magnitude smaller than 0.008 are masked to improve visual clarity. g500 contours for the day prior to the peak are included. c): Standardized anomalies of geopotential at 500hPa in colour and data in contours for the geopotential at 500hPa for the day prior to the temperature anomaly peak (6th of August). The magenta dot in panel *b* marks the location of Córdoba.

in determining the state and evolution of the troposphere, as well as a key indicator of the climate change response (Christidis and Stott, 2015). Lemus-Canovas et al. (2024) and Quesada et al. (2012) also show the important role of the atmospheric circulation component in triggering the heatwave event, and the land compound acting as an amplifier of extreme conditions. Moreover, Wilcke et al. (2020) highlight the importance of circulation in the 2018 summer heatwave over Stockholm, and its capability to force warm, long-lasting conditions without much contribution of surface feedbacks.

3.3 Case study: Summer heatwave Córdoba 2021

For the first case study, we test the CombinedModel during the period 10 August 2021 to 16-20 August 2021 in the Córdoba region, when a record-shattering heatwave struck the Iberian Peninsula. During this period, temperatures up to 47.6°C were registered in La Rambla (Córdoba) (AEMET, 2021). ~~The model predicts correctly all the daily labels during this period: 0,1,1,1,1,1,1, with 0 corresponding to non-extreme class and 1 to extreme class.~~

370 Fig. 8a shows the temporal evolution of the extreme class SHAP values for the atmospheric circulation features in terms of mean absolute SHAP (spatially aggregated), and the three soil moisture levels. Additionally, it shows the standardized temperature anomalies and the true and predicted labels. By looking at the predicted labels, we can see that the model correctly predicts all the true positives of this event but yields false positives at the developing and decaying stages of the heatwave event. Fig. 8b ~~represents shows~~ the SHAP values for the *g500* feature ~~for the SHAP peak at the temperature anomaly peak identified~~ in fig. 8a ~~, with statistical significance assessed via a bootstrapping procedure using the 95% confidence interval determined with the 2.5th and 97.5th percentiles. The values outside the confidence interval are not plotted~~ (14 August 2026). Grid points with $|SHAP| < 0.008$ are masked out for clarity. Lastly, Fig. 8c displays the standardized anomalies for the *lag1* (13 August 2021) of the *g500* field (colours) and the geopotential height at 500hPa data (black contours) ~~during the peak day~~. During the thirteenth of August, the positive anomalies and anticyclonic pattern are right above the Iberian Peninsula, and our model gives
380 mainly positive SHAP values for that region, showing strong locality around the predicted location. This signal starts to be strong and positive when the positive anomalies reach the predicted location (Fig. 8a).

The soil moisture anomalies during this event are negative but not especially low (not shown), resulting in SHAP values for the soil moisture levels being small and remaining quite constant. A curious result is the CO₂ being generally the second most important feature. This suggests that even for a specific extreme event like this one, we can identify a clear anthropogenic
385 contribution.

3.4 Case study: Summer heatwave Hannover 2018

We select the Hannover region for the second case study to ~~have a contrasting soil moisture regime~~ represent a different land-atmosphere coupling regime, characterized by higher soil water content, compared to the Córdoba region (Seneviratne et al., 2006). We focus on the 2018 European heatwave, which was an exceptional climatic event that brought prolonged periods of extreme
390 heat and drought to much of Northern and Central Europe, including Germany.

The period selected is ~~29-22 July 2018 to 4 August 2018 and the model predicts correctly all classes, which are a complete week of temperatures above the 90th percentile: 1,1,1,1,1,1,1.~~ 12 August 2018. In Fig. 9 we show the same type of plots as in the former case study, with the same ~~significance treatment~~ filtering for the SHAP values applied in 9bas ~~in the previous case~~. Again we see a strong locality in the spatial SHAP values for *g500*, with positive values surrounding the predicted location, where the positive anomalies are located. The spatial distributions are noisy for the rest of circulation variables, but they all explain less than 10% of the total mean absolute SHAP value. In Fig. 9a we can see that in this case the level one of soil moisture plays a more important role compared to the Córdoba event. The soil moisture at level one has positive SHAP values during the whole ~~week, which period, which is~~ comparable to *g500* in magnitude ~~during the first for some of the~~ days. Regarding the CO₂ concentrations, in this case its relevance is much lower compared to the Córdoba case. ~~Different~~
400 ~~drivers have been proven to play a central role in modulating the occurrence and intensity of summer temperature extremes, yet their individual contributions remain difficult to quantify. In this study, we develop an explainable machine-learning framework to disentangle the respective influences of large-scale atmospheric circulation, soil moisture anomalies, and rising CO₂ concentrations on boreal summer temperature extremes at six locations across Europe and North Africa with different~~

characteristics of land-atmosphere coupling (Córdoba, Lyon, Hannover, Stockholm, Belgrade, and Marrakech). Using SHapley Additive exPlanation (SHAP) values, we find that the atmospheric circulation consistently dominates model explainability across all locations, contributing to 67-90 % of the total mean SHAP, with the geopotential at 500 hPa field contributing the most. Soil-moisture influence exhibits a northward gradient: negligible at Marrakech (0.5 %), moderate at Córdoba (7.7 %), and substantial at Lyon (15 %). Additionally, negative correlations between soil-moisture standardized anomalies and SHAP values across three depth levels corroborate the amplifying effect of land drying on heat extremes. We demonstrate the robustness of these findings to a less stringent (80th percentile) extreme definition. Furthermore, the identified driver contributions are consistent when using alternative observational data for temperature extreme definition and for computing SPI/SPEI drought indices as proxies for soil moisture, with SPEI showing a closer alignment to the original ERA5-Land results. We also illustrate the methodology for case studies of two individual events, heatwaves occurring in Córdoba (Spain) 2021 and Hannover (Germany) 2018, which reveal a pronounced spatial pattern in the distribution of SHAP values for the circulation predictors. They also confirm the enhanced role of the land component in regions of northern Europe and reveal a contribution of the anthropogenic factor through CO₂ concentrations, even for specific events. These insights enhance our understanding of the physical mechanisms behind temperature extremes and demonstrate the potential of explainable artificial intelligence methods to improve the representation of land-atmosphere feedbacks in climate models. Different drivers have been proven to play a central role in modulating the occurrence and intensity of summer temperature extremes, yet their individual contributions remain difficult to quantify. In this study, we develop an explainable machine-learning framework to disentangle the respective influences of large-scale atmospheric circulation, soil-moisture anomalies, and rising CO₂ concentrations on boreal-summer temperature extremes at six locations across Europe and North Africa with different characteristics of land-atmosphere coupling (Córdoba, Lyon, Hannover, Stockholm, Belgrade, and Marrakech). Using SHapley Additive exPlanation (SHAP) values, we find that the atmospheric circulation consistently dominates model explainability across all locations, contributing to 67-90 % of the total mean SHAP, with the geopotential at 500 hPa field contributing the most. Soil-moisture influence exhibits a northward gradient: negligible at Marrakech (0.5 %), moderate at Córdoba (7.7 %), and substantial at Lyon (15 %). Additionally, negative correlations between soil-moisture standardized anomalies and SHAP values across three depth levels corroborate the amplifying effect of land drying on heat extremes. We demonstrate the robustness of these findings to a less stringent (80th percentile) extreme definition. Furthermore, the identified driver contributions are consistent when using alternative observational data for temperature extreme definition and for computing SPI/SPEI drought indices as proxies for soil moisture, with SPEI showing a closer alignment to the original ERA5-Land results. We also illustrate the methodology for case studies of two individual events, heatwaves occurring in Córdoba (Spain) 2021 and Hannover (Germany) 2018, which reveal a pronounced spatial pattern in the distribution of SHAP values for the circulation predictors. They also confirm the enhanced role of the land component in regions of northern Europe and reveal a contribution of the anthropogenic factor through CO₂ concentrations, even for specific events. These insights enhance our understanding of the physical mechanisms behind temperature extremes and demonstrate the potential of explainable artificial intelligence methods to improve the representation of land-atmosphere feedbacks in climate models.

4 Conclusions

In this study, we quantify the contribution of soil moisture, atmospheric circulation and CO₂ forcing to extreme temperature events during the boreal summer using an explainable ML approach. We focus on six locations with different ~~land–atmosphere coupling–evaporative~~ regimes. Based on a percentile definition of extremes, we find an increasing trend in the yearly count of extreme days during summer for Córdoba, Lyon, Belgrade, and Marrakech regions.

To ensure the robustness of the model predictions, we train an ensemble of 20 members and analyse the 50% most confident predictions across all observed cases. Balanced accuracy increases consistently with the percentage of confidence using the CombinedModel, indicating stable predictive skill. The area under the ROC curve also improves with confidence. We further tested this robustness by applying our framework to different extreme definitions and data sources. The model trained on the 80th percentile definition showed equally robust predictive performance, while the model trained on the more stringent 95th percentile definition exhibited less stable behaviour due to the high class imbalance, leading us to disregard its explainability outputs. Furthermore, the framework demonstrated reliable performance when using observational data for TX and for computing SPI/SPEI drought indices as proxys for soil moisture, with SPEI giving qualitatively more similar results to the model using ERA5 data for the atmospheric circulation and the soil moisture levels.

To ~~interpret the predictions of the model~~ help interpret the model prediction, we use SHAP values, which allows us to assess the marginal contributions of the atmospheric circulation, land and atmospheric CO₂ features. The reader must keep in mind that the explainability results represent the interpretations done by the model, and we cannot assume that these results represent the reality. ~~With the strategies explained in 2.7 we~~ We are ensuring a treatment of part of the existing epistemic uncertainty when using this model, but different architectures or different models might differ in the results. The explainability results show that circulation features, especially the geopotential at 500 hPa, dominate the prediction of extreme events across all locations (66.9–89.7 %). The land component shows a regional differing behaviour: its importance increases in temperate and northern regions (10.1–15.0 %), while remaining negligible or moderate in ~~arid locations~~ a dry/transitional location like Córdoba and ~~Marrakecha~~ dry one like Marrakech (Hsu and Dirmeyer, 2023), representing 7.7 % and 0.5 % of the total SHAP respectively.

~~The correlation between SHAP values and standardized soil moisture anomalies is negative across all levels, confirming the~~ The dominant soil-moisture contributions—particularly from the upper layer—show an inverse relationship between moisture anomalies and SHAP values, consistent with the well-documented role of dry conditions in enhancing the probability of extreme heat events (Hirschi et al., 2011; Seneviratne et al., 2006). Attribution for the intermediate layer is weaker and less robust, likely because strong inter-layer correlations make its independent contribution difficult to isolate. The land component importance increases from the 80th to the 90th percentile extreme definition. This suggests that land variables could play an even more important role for more extreme events (e.g. using a 95th percentile extreme definition). This could not be analysed with the presented methodology due to class imbalance. CO₂ SHAP contributions are substantial in locations where we detect a trend in the yearly extreme count, suggesting that the model captures the long-term forcing signal.

Furthermore, the core patterns of driver contributions identified by the model are robust to a different definition of the extreme events and to the use of observational data from E-OBS instead of ERA5-Land. The north–south gradient in land

driver importance and the dominance of atmospheric circulation are consistently reproduced in the analysis using the 80th percentile extreme definition, albeit with a reduced relative contribution from soil moisture, likely due to the inclusion of less severe events. Moreover, the regionality of the land component is robustly recovered when using observational temperature data and the SPEI drought index as a proxy for soil moisture. The SPI index, which lacks the evapotranspiration component of SPEI, resulted in a weaker land signal, underscoring the importance of representing more accurately surface water balance for capturing land–atmosphere feedbacks and confirming that SPEI provides a more physically comprehensive proxy for this purpose.

We also analyse two case studies to further study the explainability technique. For the 2021 Córdoba heatwave, the model attributes the dominant contribution to the atmospheric circulation, with a mainly positive SHAP signal over the Iberian Peninsula when positive 500 hPa geopotential are placed in the region. We identify a clear anthropogenic contribution due to the high CO₂ SHAP value in this event. The land variables play a minor role in this case, with anomalies close to zero for the three soil moisture levels. During the 2018 Hannover heatwave, we see an enhanced importance of the land component, especially the first soil moisture level, with positive SHAP values comparable to the *g500* circulation feature during the first days of the event. Again, we see a strong locality in the spatial distribution of SHAP values. In this second case study, CO₂ plays a minor role.

These results ~~reveal~~are consistent with findings about the central role of atmospheric circulation in driving extreme temperature events and highlight the regionally dependent modulation by soil moisture ([Fischer et al., 2007](#); [Barriopedro et al., 2023](#); [Quesada et al.](#)). The explainability technique used allows us to attribute data-driven predictions to physical drivers, offering a diagnostic framework that can assess the contribution of individual components to extreme events. We have demonstrated that this framework and its conclusions are robust to the choice of extreme event threshold (if the class imbalance is not too strong) and to the use of observational data instead of reanalysis data.

Our findings on the role of atmospheric circulation in driving daily temperature extremes support the findings of Ucan et al. (2024), who showed that the geopotential field is the most relevant feature over 59.9 % of global land at a 1-day scale and 67.9 % at a 7-day scale. We confirm the importance of the atmospheric circulation using a different ML architecture, adding robustness to this result. Additionally, using a ConvNeXt model for atmospheric fields allows us to identify the spatial regions to which the model attributes higher importance, further demonstrating the ability of XAI to reveal spatial patterns in weather and climate problems, as shown by Davenport and Diffenbaugh (2021) and Mayer and Barnes (2021) using LRP. ~~SHAP values, however, remain relatively unexplored for such pattern identification~~While recent studies have increasingly employed XAI to identify regional climate drivers and sources of predictability (e.g., Zhang et al. (2024); Van Straaten et al. (2023); Mamalakis et al. (2023b)), our work specifically leverages XAI to disentangle the relative roles of atmospheric circulation and land-surface conditions in driving localized hot temperature extreme events.

The role of soil moisture in modulating the occurrence of extremes has been highlighted by Materia et al. (2022) and Seneviratne et al. (2006), showing that dry soils intensify and prolong heat extremes, while wet soils suppress heatwaves. Here, we support these findings on the modulating effect of ~~this the~~ land–atmosphere coupling by analysing the correlation between soil moisture anomalies and corresponding SHAP values. Our work contributes to disentangling the components of

this coupling and quantifying their roles using XAI. We further quantify the influence of soil moisture at three different depth levels and find that, on weather time scales, the level closest to the surface ~~generally~~ plays the largest role for humid/transitional regions, whereas the bottom layer is more important for a dry/transitional location like Córdoba.

510 Finally, we include a quantification of the role of atmospheric CO₂, which is assessed by Materia et al. (2025) at seasonal scales but is not usually considered in similar studies using explainable ML. Including this feature allows us to account for the trend in the occurrence of hot extremes due to global warming, without encountering the complications of detrending the data when splitting it into training, validation, and testing sets in ML studies.

Future work should include testing the robustness of the results across different model architectures. Analysing additional
515 locations in future work may help gain insights into the regional variability of the results. Including variables like humidity fluxes or information on aerosol composition in each location could also provide complementary information on the physical processes modulating temperature extremes during boreal summer. The presented XAI methodology could also be applied to evaluate the importance of the different drivers in CMIP6 dynamical models or in Subseasonal to Seasonal to Decadal Prediction Systems.

520 Overall, this study provides a physically interpretable, data-driven framework to disentangle the contributions of atmospheric circulation, land surface conditions, and anthropogenic forcing to summer temperature extremes. By combining a robust machine learning method with an ML explainability technique, we ~~offer insights that can inform both~~ provide a framework that can support model development and climate attribution efforts. Our results underscore the dominant influence of atmospheric circulation and the modulating role of soil moisture, revealing consistent regional signals across different locations. These
525 findings highlight the potential of explainable machine learning as a complementary tool for climate diagnostics and pave the way for more comprehensive, process-based assessments of future temperature extremes.

Code and data availability. The repository with the code used in this work is available in <https://github.com/agarcimes8/QuantifyDriversHW>. The repository contains the scripts needed to pre-process data (anomalies, heatwave detection), the deep learning model code together with the scripts for training the model and computing the explainability results, and scripts or notebooks to post-process the results.

530

Additionally, the repository includes a few datasets with: the file with the data for the CO₂ feature, the files with the SPEI and SPI drought indices, and the files with the optimized values for the hyperparameters of the deep learning model.

Appendix A

A1 LOESS smoothing

535 To improve the robustness of daily climatology percentiles estimates, we use a local polynomial regression (LOESS) as proposed in Mahlstein et al. (2015). This method smooths the raw climatology by fitting a regression locally at each point using a subset of neighbouring days weighted by distance. The weights w_i applied to each point in the local window are based on their

normalized distance d_i to the estimation point, using the tricubic function:

$$w_i = \begin{cases} (1 - |d_i|^3)^3, & \text{if } |d_i| \leq 1 \\ 0, & \text{if } |d_i| > 1 \end{cases}$$

540 where $d_i = \frac{x_i - x}{h}$ is the distance between the input point x_i and the target point x , scaled by the distance h to the farthest neighbour in the local window. This ensures that the weights smoothly decay to zero at the edges of the window and are strictly zero beyond it.

In practice, this means we apply LOESS to the raw daily climatology or percentile values to obtain a smoother and more stable annual profile. This is particularly important when dealing with small sample sizes or high daily variability, where simple
545 averaging can lead to noisy or misleading estimates. This smoothing step ensures that the resulting threshold and climatology estimates are not overly influenced by isolated values or short-term fluctuations, which is especially relevant for extreme event detection on a daily basis.

A2 Cross-entropy loss function

The Cross Entropy Loss function is defined as:

$$550 \quad \mathcal{L} = \frac{-1}{N} \sum_i y_i \cdot \log(p_i) + (1 - y_i) \cdot \log(1 - p_i) \quad (\text{A1})$$

where p_i is the probability of class 1, $(1 - p_i)$ the probability of class 0, and y_i is the true label. This loss function is widely used in binary classification tasks in ML.

Appendix B: Statistical and robustness results

Table B1. Summary of extreme event counts by site and period for different extreme definitions. Yearly extreme count sum is shown for the different locations, for the training (80% of 1950–2013), validation (20% of 1950–2013) and test (2014–2024) periods. **E** corresponds to the extreme class and **NE** to the non-extreme class. Additionally, we show the randomly selected years for the training and validation sets.

Site	Period	Total Days	80th Percentile		90th Percentile		95th Percentile	
			E	NE	E	NE	E	NE
Cordoba	Train (1950–2013)	4692	1162	3530	654	4038	363	4329
	Val (1950–2013)	1196	278	918	141	1055	71	1125
	Test (2014–2024)	1011	506	505	350	661	225	786
Lyon	Train (1950–2013)	4692	1110	3582	649	4043	378	4314
	Val (1950–2013)	1196	267	929	134	1062	76	1120
	Test (2014–2024)	1011	471	540	333	678	245	766
Hannover	Train (1950–2013)	4692	1084	3608	582	4110	314	4378
	Val (1950–2013)	1196	232	964	120	1076	63	1133
	Test (2014–2024)	1011	358	653	217	794	148	863
Stockholm	Train (1950–2013)	4692	1038	3654	524	4168	278	4414
	Val (1950–2013)	1196	257	939	123	1073	55	1141
	Test (2014–2024)	1011	312	699	177	834	119	892
Belgrado	Train (1950–2013)	4692	1103	3589	604	4088	329	4363
	Val (1950–2013)	1196	265	931	135	1061	68	1128
	Test (2014–2024)	1011	385	626	247	764	156	855
Marrakech	Train (1950–2013)	4692	1058	3634	571	4121	298	4394
	Val (1950–2013)	1196	281	915	157	1039	88	1108
	Test (2014–2024)	1011	344	667	211	800	126	885

Training years: 1951, 1953, 1954, 1958–1986, 1988–1991, 1993, 1995–1998, 2000, 2002–2005, 2007, 2009, 2011–2013

Validation years: 1950, 1952, 1955, 1956, 1957, 1987, 1992, 1994, 1999, 2001, 2006, 2008, 2010




Figure B1. Trends in JJA Summer Hot Extremes across different time periods. Bars represent the slope coefficient (trend magnitude) of a linear regression applied to the yearly count of extreme days for four periods: the full available record (dark red), since 1980 (red), since 1990 (light red), and since 2000 (lightest red). Asterisks (*) indicate trends that are statistically significant at the 95% confidence level, determined using a t-test on the slope coefficient.




Figure B2. Change in accuracies for the Combined Model for different percentages of confidence using the 90th percentile extreme definition. In the x-axis the percentage of confidence taken, and in the y-axis the accuracies. EA, NEA and BA in the legend stand for extreme accuracy, non-extreme accuracy and balanced accuracy respectively. The text box shows the sample size and the random chance of the naive classifier.




Figure B3. Change in ROC curves for the Combined Model for different percentages of confidence using the 90th percentile extreme definition. In the x-axis the false positive rate and in the y-axis the true positive rate. The legend shows the different percentages of confidence taken.




Figure B4. Same plot as in Fig. B2 but for the 80th percentile definition of extreme.




Figure B5. Same plot as in Fig. B2 but for the 95th percentile definition of extreme.

Appendix C: Drought indices correlations




Figure C1. Spearman correlation between the three soil moisture levels (*swvl1*, *swvl2*, *swvl3*) and the SPEI index computed at three different time scale using the E-OBS gridded data. In the x-axis the scale used to compute the SPEI index and in the y-axis the resulting spearman correlation. The legend indicates the three original soi moisture levels used in the DL model.




Figure C2. Same plot as Fig. C1 but using the SPI drought index instead.




Figure D1. Same plot as in Fig. 6 but for the atmospheric circulation features. We sum up the SHAP values and averaged the standardized anomalies of the circulation variables in a 400km by 400km region surrounding the target location.




Figure D2. Same plot as in Fig. 5 but for the 80th percentile definition of extreme.




Figure D3. Same plot as in Fig. 6 but for the model using E-OBS observational data and the SPEI 30 days index as a proxy for the soil moisture feature.




Figure D4. Same plot as in Fig. 5 but for the model using E-OBS observational data and the SPI 30 days index as a proxy for the soil moisture feature.




Figure D5. Same plot as in Fig. 5 but for the model using only *g500* in the atmospheric circulation features.

Author contributions. AG, LP and MD conceptualised the study; AG and LP worked on the data and developed the model code and methodology; BG worked on code optimization and code structure for publication; AG trained the ML model and analysed the data; AG, LP, MD, SM and RM analysed the results; AG wrote the manuscript draft; LP, MD, SM, and RM reviewed and edited the manuscript; MD, SM and RM supervised the study; MD did the funding acquisition

560 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. This research was ~~conducted within the framework of the EXPECT project. EXPECT has received~~ supported with funding from the ~~European Union's Horizon Europe Framework Programme (HORIZON)~~ Horizon Europe project EXPECT under Grant Agreement 101137656. Stefano Materia also acknowledges also acknowledges AI4S fellowships within the “Generación D” initiative by Red.es, Ministerio para la Transformación Digital y de la Función Pública of Spain, for talent attraction (C005/24-ED CV1), funded by
565 NextGenerationEU through PRTR. The authors also want to thank Vincent Verjans for reviewing the manuscript.

References

- AEMET: La ola de calor del puente de agosto'21 y los récords de temperaturas en España, <https://aemetblog.es/2021/08/18/la-ola-de-calor-del-puente-de-agosto21-y-los-records-de-temperaturas-en-espana/>, aEMET Blog, 2021.
- Barriopedro, D., García-Herrera, R., Ordóñez, C., Miralles, D. G., and Salcedo-Sanz, S.: Heat Waves: Physical Understanding and Scientific Challenges, *Reviews of Geophysics*, 61, e2022RG000780, <https://doi.org/10.1029/2022RG000780>, 2023.
- Beobide-Arsuaga, G., Düsterhus, A., Müller, W. A., Barnes, E. A., and Baehr, J.: Spring Regional Sea Surface Temperatures as a Precursor of European Summer Heatwaves, *Geophysical Research Letters*, 50, e2022GL100727, <https://doi.org/10.1029/2022GL100727>, 2023.
- Bischof, S., Pilch Kedzierski, R., Hänsch, M., Wahl, S., and Matthes, K.: The Role of the North Atlantic for Heat Wave Characteristics in Europe, an ECHAM6 Study, *Geophysical Research Letters*, 50, e2023GL105280, <https://doi.org/10.1029/2023GL105280>, 2023.
- 575 Bommer, P. L., Kretschmer, M., Hedström, A., Bareeva, D., and Höhne, M. M.-C.: Finding the Right XAI Method—A Guide for the Evaluation and Ranking of Explainable AI Methods in Climate Science, *Artificial Intelligence for the Earth Systems*, 3, e230074, <https://doi.org/10.1175/AIES-D-23-0074.1>, 2024.
- Camps-Valls, G., Fernández-Torres, M.-Á., Cohrs, K.-H., Höhl, A., Castelletti, A., Pacal, A., Robin, C., Martinuzzi, F., Papoutsis, I., Prapas, I., Pérez-Aracil, J., Weigel, K., Gonzalez-Calabuig, M., Reichstein, M., Rabel, M., Giuliani, M., Mahecha, M. D., Popescu, O.-I., Pellicer-Valero, O. J., Ouala, S., Salcedo-Sanz, S., Sippel, S., Kondylatos, S., Happé, T., and Williams, T.: Artificial intelligence for modeling and understanding extreme weather and climate events, *Nature Communications*, 16, 1919, <https://doi.org/10.1038/s41467-025-56573-8>, 2025.
- Christidis, N. and Stott, P. A.: Changes in the geopotential height at 500 hPa under the influence of external climatic forcings, *Geophysical Research Letters*, 42, <https://doi.org/10.1002/2015gl066669>, publisher: American Geophysical Union (AGU), 2015.
- 585 Cornes, R. C., Van Der Schrier, G., Van Den Besselaar, E. J. M., and Jones, P. D.: An Ensemble Version of the E-OBS Temperature and Precipitation Data Sets, *Journal of Geophysical Research: Atmospheres*, 123, 9391–9409, <https://doi.org/10.1029/2017JD028200>, 2018.
- Cos, P., Olmo, M., Campos, D., Marcos-Matamoras, R., Palma, L., Muñoz, A. G., and Doblas-Reyes, F. J.: Saharan warm air intrusions in the Western Mediterranean: identification, impacts on temperature extremes and large-scale mechanisms, <https://doi.org/10.5194/egusphere-2024-3331>, 2024.
- 590 D'Andrea, F., Duvel, J.-P., Rivière, G., Vautard, R., Cassou, C., Cattiaux, J., Coumou, D., Faranda, D., Happé, T., Jézéquel, A., Ribes, A., and Yiou, P.: Summer Deep Depressions Increase Over the Eastern North Atlantic, *Geophysical Research Letters*, 51, e2023GL104435, <https://doi.org/10.1029/2023GL104435>, 2024.
- Davenport, F. V. and Diffenbaugh, N. S.: Using Machine Learning to Analyze Physical Causes of Climate Change: A Case Study of U.S. Midwest Extreme Precipitation, *Geophysical Research Letters*, 48, e2021GL093787, <https://doi.org/10.1029/2021GL093787>, 2021.
- 595 Donat, M. G., Sillmann, J., Wild, S., Alexander, L. V., Lippmann, T., and Zwiers, F. W.: Consistency of Temperature and Precipitation Extremes across Various Global Gridded In Situ and Reanalysis Datasets, *Journal of Climate*, 27, 5019–5035, <https://doi.org/10.1175/JCLI-D-13-00405.1>, 2014.
- Fischer, E. M., Seneviratne, S. I., Lüthi, D., and Schär, C.: Contribution of land-atmosphere coupling to recent European summer heat waves, *Geophysical Research Letters*, 34, 2006GL029068, <https://doi.org/10.1029/2006GL029068>, 2007.
- 600 Goodfellow, I., Bengio, Y., and Courville, A.: Deep Learning, MIT Press, <http://www.deeplearningbook.org>, 2016.
- Gordon, E. M., Barnes, E. A., and Davenport, F. V.: Separating internal and forced contributions to near term SST predictability in the CESM2-LE, *Environmental Research Letters*, 18, 104047, <https://doi.org/10.1088/1748-9326/acfdbc>, 2023.

- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hirschi, M., Seneviratne, S. I., Alexandrov, V., Boberg, F., Boroneant, C., Christensen, O. B., Formayer, H., Orlowsky, B., and Stepanek, P.: Observational evidence for soil-moisture impact on hot extremes in southeastern Europe, *Nature Geoscience*, 4, 17–21, <https://doi.org/10.1038/ngeo1032>, 2011.
- Hsu, H. and Dirmeyer, P. A.: Soil moisture-evaporation coupling shifts into new gears under increasing CO₂, *Nature Communications*, 14, 1162, <https://doi.org/10.1038/s41467-023-36794-5>, 2023.
- Intergovernmental Panel On Climate Change: Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, 1 edn., ISBN 978-1-00-915789-6, <https://doi.org/10.1017/9781009157896>, 2023.
- Jach, L., Schwitalla, T., Branch, O., Warrach-Sagi, K., and Wulfmeyer, V.: Sensitivity of land–atmosphere coupling strength to changing atmospheric temperature and moisture over Europe, *Earth System Dynamics*, 13, 109–132, 2022.
- Khodayar Pardo, S. and Paredes-Fortuny, L.: Uneven evolution of regional European summer heatwaves under climate change, *Weather and Climate Extremes*, 43, 100648, <https://doi.org/10.1016/j.wace.2024.100648>, 2024.
- Koh, J., Steinfeld, D., and Martius, O.: Using spatial extreme-value theory with machine learning to model and understand spatially compounding weather extremes, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 481, 20240763, <https://doi.org/10.1098/rspa.2024.0763>, 2025.
- Lemus-Canovas, M., Insua-Costa, D., Trigo, R. M., and Miralles, D. G.: Record-shattering 2023 Spring heatwave in western Mediterranean amplified by long-term drought, *npj Climate and Atmospheric Science*, 7, 25, <https://doi.org/10.1038/s41612-024-00569-6>, 2024.
- Liu, Z., Mao, H., Wu, C.-Y., Feichtenhofer, C., Darrell, T., and Xie, S.: A ConvNet for the 2020s, <https://doi.org/10.48550/arXiv.2201.03545>, 2022.
- Loshchilov, I. and Hutter, F.: Decoupled Weight Decay Regularization, <https://doi.org/10.48550/arXiv.1711.05101>, 2019.
- Lundberg, S. and Lee, S.-I.: A Unified Approach to Interpreting Model Predictions, <https://doi.org/10.48550/arXiv.1705.07874>, 2017a.
- Lundberg, S. M. and Lee, S.-I.: A Unified Approach to Interpreting Model Predictions, in: *Advances in Neural Information Processing Systems*, edited by Guyon, I., Luxburg, U. V., Bengio, S., Wallach, H., Fergus, R., Vishwanathan, S., and Garnett, R., vol. 30, Curran Associates, Inc., https://proceedings.neurips.cc/paper_files/paper/2017/file/8a20a8621978632d76c43dfd28b67767-Paper.pdf, 2017b.
- Mahlstein, I., Spirig, C., Liniger, M. A., and Appenzeller, C.: Estimating daily climatologies for climate indices derived from climate model data and observations, *Journal of Geophysical Research: Atmospheres*, 120, 2808–2818, <https://doi.org/10.1002/2014JD022327>, 2015.
- Mamalakis, A., Barnes, E. A., and Ebert-Uphoff, I.: Carefully Choose the Baseline: Lessons Learned from Applying XAI Attribution Methods for Regression Tasks in Geoscience, *Artificial Intelligence for the Earth Systems*, 2, e220058, <https://doi.org/10.1175/AIES-D-22-0058.1>, 2023a.
- Mamalakis, A., Barnes, E. A., and Hurrell, J. W.: Using Explainable Artificial Intelligence to Quantify "Climate Distinguishability" After Stratospheric Aerosol Injection, *Geophysical Research Letters*, 50, e2023GL106137, <https://doi.org/10.1029/2023GL106137>, 2023b.

- 640 Masmoudi-Charfi, C. and Habaieb, H.: Rainfall Distribution Functions for Irrigation Scheduling: Calculation Procedures Following Site of Olive (*Olea europaea* L.) Cultivation and Growing Periods, American Journal of Plant Sciences, 05, 2094–2133, <https://doi.org/10.4236/ajps.2014.513224>, 2014.
- Materia, S., Ardilouze, C., Prodhomme, C., Donat, M. G., Benassi, M., Doblas-Reyes, F. J., Peano, D., Caron, L.-P., Ruggieri, P., and Gualdi, S.: Summer temperature response to extreme soil water conditions in the Mediterranean transitional climate regime, Climate Dynamics, 58, 1943–1963, 2022.
- 645 Materia, S., García, L. P., van Straaten, C., O, S., Mamalakis, A., Cavicchia, L., Coumou, D., de Luca, P., Kretschmer, M., and Donat, M.: Artificial intelligence for climate prediction of extremes: State of the art, challenges, and future perspectives, Wiley Interdisciplinary Reviews: Climate Change, 15, e914, 2024.
- Materia, S., Jung, M., Palma, L., Duarte, A., Weber, U., Samso-Cabrè, M., Bancells, P., and Donat, M.: Data-driven seasonal forecasts of European heat-waves, Artificial Intelligence for the Earth Systems, submitted, 2025.
- 650 Mayer, K. J. and Barnes, E. A.: Subseasonal Forecasts of Opportunity Identified by an Explainable Neural Network, Geophysical Research Letters, 48, e2020GL092092, <https://doi.org/10.1029/2020GL092092>, 2021.
- Mayer, K. J., Chapman, W. E., and Manriquez, W. A.: Exploring the Relative Importance of the MJO and ENSO to North Pacific Subseasonal Predictability, Geophysical Research Letters, 51, e2024GL108479, <https://doi.org/10.1029/2024GL108479>, 2024.
- 655 McKee, T. B., Doesken, N. J., and Kleist, J.: THE RELATIONSHIP OF DROUGHT FREQUENCY AND DURATION TO TIME SCALES, 1993.
- Miloshevich, G., Cozian, B., Abry, P., Borgnat, P., and Bouchet, F.: Probabilistic forecasts of extreme heatwaves using convolutional neural networks in a regime of lack of data, <https://doi.org/10.48550/arXiv.2208.00971>, 2023.
- Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., et al.: ERA5-Land: A state-of-the-art global reanalysis dataset for land applications, Earth system science data, 13, 4349–4383, 2021.
- 660 Perkins, S. E. and Alexander, L. V.: On the Measurement of Heat Waves, Journal of Climate, 26, 4500–4517, <https://doi.org/10.1175/JCLI-D-12-00383.1>, 2013.
- Quesada, B., Vautard, R., Yiou, P., Hirschi, M., and Seneviratne, S. I.: Asymmetric European summer heat predictability from wet and dry southern winters and springs, Nature Climate Change, 2, 736–741, <https://doi.org/10.1038/nclimate1536>, 2012.
- 665 Rousi, E., Kornhuber, K., Beobide-Arsuaga, G., Luo, F., and Coumou, D.: Accelerated western European heatwave trends linked to more-persistent double jets over Eurasia, Nature Communications, 13, 3851, <https://doi.org/10.1038/s41467-022-31432-y>, 2022.
- Scaini, A., Sánchez, N., Vicente-Serrano, S. M., and Martínez-Fernández, J.: SMOS-derived soil moisture anomalies and drought indices: A comparative analysis using in situ measurements, Hydrological Processes, 29, 373–383, 2015.
- 670 Seneviratne, S. I., Lüthi, D., Litschi, M., and Schär, C.: Land–atmosphere coupling and climate change in Europe, Nature, 443, 205–209, <https://doi.org/10.1038/nature05095>, 2006.
- Sha, Y., Sobash, R. A., and Ii, D. J. G.: Improving Ensemble Extreme Precipitation Forecasts Using Generative Artificial Intelligence, 4, 2025.
- Sippel, S., Meinshausen, N., Fischer, E. M., Székely, E., and Knutti, R.: Climate change now detectable from any single day of weather at global scale, Nature Climate Change, 10, 35–41, <https://doi.org/10.1038/s41558-019-0666-7>, 2020.
- 675 Soci, C., Hersbach, H., Simmons, A., Poli, P., Bell, B., Berrisford, P., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Radu, R., et al.: The ERA5 global reanalysis from 1940 to 2022, Quarterly Journal of the Royal Meteorological Society, 150, 4014–4048, 2024.

- Sundararajan, M., Taly, A., and Yan, Q.: Axiomatic Attribution for Deep Networks, <https://doi.org/10.48550/arXiv.1703.01365>, 2017.
- Toms, B. A., Barnes, E. A., and Ebert-Uphoff, I.: Physically Interpretable Neural Networks for the Geosciences: Applications to Earth System
680 Variability, 12, e2019MS002 002, <https://doi.org/10.1029/2019MS002002>, 2020.
- Uckan, Y., Ruiz-Vásquez, M., De Polt, K., and Orth, R.: Global relevance of atmospheric and land surface drivers for hot temperature extremes, <https://doi.org/10.5194/egusphere-2024-2540>, 2024.
- Van Straaten, C., Whan, K., Coumou, D., Van Den Hurk, B., and Schmeits, M.: Correcting Subseasonal Forecast Errors with an Explainable ANN to Understand Misrepresented Sources of Predictability of European Summer Temperatures, Artificial Intelligence for the Earth
685 Systems, 2, e220 047, <https://doi.org/10.1175/AIES-D-22-0047.1>, 2023.
- Vicente-Serrano, S. M., Beguería, S., and López-Moreno, J. I.: A Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index, *Journal of Climate*, 23, 1696–1718, <https://doi.org/10.1175/2009JCLI2909.1>, 2010.
- Wang, H., Rogers, J. C., and Munroe, D. K.: Commonly used drought indices as indicators of soil moisture in China, *Journal of Hydrometeorology*, 16, 1397–1408, 2015.
- 690 Wilcke, R. A. I., Kjellström, E., Lin, C., Matei, D., Moberg, A., and Tyrlis, E.: The extremely warm summer of 2018 in Sweden – set in a historical context, *Earth System Dynamics*, 11, 1107–1121, <https://doi.org/10.5194/esd-11-1107-2020>, 2020.
- Yuan, S., Quiring, S. M., and Zhao, C.: Evaluating the utility of drought indices as soil moisture proxies for drought monitoring and land–atmosphere interactions, *Journal of Hydrometeorology*, 21, 2157–2175, 2020.
- Zhang, H., Finkel, J., Abbot, D. S., Gerber, E. P., and Weare, J.: Using Explainable AI and Transfer Learning to Understand and Predict the
695 Maintenance of Atlantic Blocking With Limited Observational Data, *Journal of Geophysical Research: Machine Learning and Computation*, 1, e2024JH000 243, <https://doi.org/10.1029/2024JH000243>, 2024.
- Zscheischler, J., Westra, S., Van Den Hurk, B. J., Seneviratne, S. I., Ward, P. J., Pitman, A., AghaKouchak, A., Bresch, D. N., Leonard, M., Wahl, T., et al.: Future climate risk from compound events, *Nature climate change*, 8, 469–477, 2018.