



Simulated and Observed Transport Estimates Across the Overturning in the Subpolar North Atlantic Program (OSNAP) Section

Gokhan Danabasoglu¹, Frederic S. Castruccio¹, Burcu Boza², Alice M. Barthel³, Arne Biastoch^{4,5}, Adam Blaker⁶, Alexandra Bozec⁷, Diego Bruciaferri⁸, Frank O. Bryan¹, Eric P. Chassignet⁷, Yao Fu^{9,10}, Ian Grooms¹¹, Catherine Guiavarc'h⁸, Hakase Hayashida¹², Andrew McC. Hogg¹³, Ryan M. Holmes¹⁴, Doroteaciro Iovino¹⁵, Andrew E. Kiss¹³, M. Susan Lozier⁹, Gustavo Marques¹, Alex Megann⁶, Franziska U. Schwarzkopf⁴, Dave Storkey⁸, Luke van Roekel³, Jon Wolfe³, Xiaobiao Xu⁷, Rong Zhang^{16,17}

¹ US National Science Foundation (NSF) National Center for Atmospheric Research (NCAR), Boulder, CO, USA

² Eurasia Institute of Earth Sciences, Istanbul Technical University, Istanbul, Türkiye

³ US Department of Energy (DOE), Los Alamos National Laboratory (LANL), Los Alamos, NM, USA

⁴ GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

⁵ Kiel University, Kiel, Germany

⁶ National Oceanography Centre, Southampton, UK

⁷ Center for Ocean-Atmospheric Prediction Studies (COAPS), Florida State University, Tallahassee, FL, USA

⁸ UK Met Office, Exeter, UK

⁹ School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

¹⁰ College of Marine Science, University of South Florida, St. Petersburg, FL, USA

¹¹ Applied Mathematics Department, University of Colorado Boulder, Boulder, CO, USA

¹² Application Laboratory, Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan

¹³ Research School of Earth Sciences and ARC Centre of Excellence for Climate Extremes, Australian National University, Canberra, Australia

¹⁴ Australian Bureau of Meteorology, Sydney, Australia

¹⁵ Fondazione Centro Euro-Mediterraneo sui Cambiamenti Climatici (CMCC), Bologna, Italy

¹⁶ Program in Atmospheric and Oceanic Sciences, Princeton University, Princeton, NJ, USA

¹⁷ National Oceanic and Atmospheric Administration (NOAA) Geophysical Fluid Dynamics Laboratory (GFDL), Princeton, NJ, USA

Correspondence: Gokhan Danabasoglu (gokhan@ucar.edu)



40

Abstract

A comparison of simulated and observed overturning transports and related properties across the Overturning in the Subpolar North Atlantic Program (OSNAP) sections for the 2014-2022 period is presented, considering both depth and density space transports. The effort was motivated by the observational transport estimates at both OSNAP-West (OW) and OSNAP-East (OE) sections which show a minor role for the Labrador Sea (LS) in setting the mean and variability of the overturning in the subpolar North Atlantic. There are 9 participating groups from around the world, contributing a total of 18 ocean – sea simulations with 6 different ocean models. The simulations use a common set of interannually-varying atmospheric forcing datasets. The horizontal resolutions of the simulations range from nominal 1° to eddy-resolving resolutions of 0.1° - 0.05° . While there are many differences between the simulations and observations as well as among the individual simulations in terms of transport properties, the simulations show significantly larger transports at OE than at OW in general agreement with the observations. Analyzing overturning circulations in both depth and density space together provides a more complete picture of the overturning properties and features. This analysis also reveals that, in both the simulations and observations, northward and southward flows substantially cancel each other, producing much smaller residual (total) transports. Such cancellations tend to be much more prominent in depth space than in density space. In general, the observed transport features are captured better at OE than OW. The simulations generally show larger (smaller) transports with positive (negative) temperature and salinity biases in the upper ocean near the OSNAP sections, but with no such relationship with density biases. In high-resolution simulations, the transport profiles agree better with the observations in general, but challenges remain in some other metrics considered in our analysis. When transports are calculated using a density referenced to 2000-m depth, rather than the ocean surface, the relative contributions of transports at OW increase modestly.

65



1 Introduction

70 The Atlantic Meridional Overturning Circulation (AMOC) depicts a simplified, zonally averaged
 view of rather complex, three-dimensional circulation patterns of the Atlantic Ocean, connected
 to the global circulation system primarily across the basin's southern boundary. Through its
 associated heat, salt, and tracer (e.g., carbon) transports and their effects on the ocean state,
 particularly on decadal-to-multi-decadal time scales, the AMOC has profound impacts on the
 75 climate of the surrounding areas and on the global climate (see recent reviews by Buckley &
 Marshall, 2016; Sutton et al., 2018; and Zhang et al., 2019 and references therein). These impacts
 mainly come about through spatial and temporal variations of sea surface temperatures (SSTs).
 They include impacts on Atlantic hurricane activity, shifts in the Intertropical Convergence Zone,
 precipitation changes over the Sahel and Amazon, changes in summertime climate over North
 80 America and Europe, and other modes of climate variability – all with significant societal and
 economic impacts. Furthermore, due to the presence of low-frequency (decadal and longer)
 AMOC variability seen in many model simulations (e.g., Msadek et al., 2010; Danabasoglu et
 al., 2012b), the AMOC is thought to represent an important component of dynamical memory of
 the climate system, thus making its proper initialization – along with associated (upper-ocean)
 85 heat content anomalies in the North Atlantic – important for decadal climate prediction
 simulations (see Meehl et al., 2014 and references therein). Given all these important impacts,
 there is also a growing interest in how the AMOC will change in a warming climate with many
 recent studies focusing on weakening as well as the stability and collapse of the AMOC (e.g.,
 Caesar et al., 2018; Caesar et al. 2019; Weijer et al., 2020; van Westen et al., 2024).

90 In the absence of continuous and trans-basin observations prior to 2004, much of our knowledge
 of the mean state, spatial and temporal variability, latitudinal coherency, and variability
 mechanisms of the AMOC during the historical period comes from model simulations.
 Unfortunately, models remain inconsistent in their representations of the mean AMOC and its
 variability. For example, results from about 20 ocean model simulations forced with common,
 95 interannually varying atmospheric datasets for the 1948-2007 period as part of the second phase
 of the Coordinated Ocean-ice Reference Experiments (CORE-II; Large & Yeager, 2009; Griffies
 et al., 2009) show a rather large range of mean AMOC transports in both depth- and density-
 space, e.g., between ~5 and ~20 Sv at both 26.5° and 45°N in depth space (Danabasoglu et al.,



2014). There is a broad agreement among these simulations in their temporal depictions of
 100 interannual-to-decadal variability and trends – despite significant differences in the spatial
 structures of variability patterns – indicating that simulated variability and trends are largely
 dictated by the imposed forcing (Danabasoglu et al. 2016). In both Danabasoglu et al. (2014) and
 Danabasoglu et al. (2016), these AMOC differences among the simulations were attributed to
 biases in the Labrador Sea (LS) region in upper-ocean potential temperature and salinity
 105 distributions, mixed layer depths, and sea-ice cover. Similar inter-model differences in AMOC
 properties are also seen in ocean reanalysis products for the historical period (Karspeck et al.,
 2017; Jackson et al., 2019) even though each reanalysis ingests largely the same observational
 datasets. In these products, differences arise due to insufficient observational data to constrain
 the deep and abyssal oceans as well as the continental shelf regions. As such, the fidelity of the
 110 underlying ocean model becomes as important as those used in free-running forced ocean
 simulations. Large inter-model differences in AMOC properties also exist in fully coupled
 simulations participating in phases of the Coupled Model Intercomparison Project (e.g., Cheng et
 al., 2013; Weijer et al., 2020; Bryden et al., 2024). Moreover, historical AMOC variability and
 trends over recent decades do not necessarily agree with those from forced-ocean simulations;
 115 the coupled simulations show a somewhat stronger weakening that starts earlier compared to the
 forced simulations.

As discussed in Buckley & Marshall (2016), Zhang et al. (2019), Danabasoglu et al. (2019), and
 references therein, inter-annual AMOC variability in model simulations is primarily dictated by
 momentum fluxes, i.e., winds. In contrast, low-frequency AMOC variability is largely driven by
 120 buoyancy fluxes at high latitudes of the North Atlantic. Specifically, a large majority of model
 simulations identify deep water formation regions – usually determined by deep mixed layers –
 as important regions linked to downstream AMOC variability. The associated low-frequency
 variability mechanism involves surface buoyancy flux anomalies over the LS region that arise
 from long episodes of the positive phase of the North Atlantic Oscillation (NAO). These
 125 anomalies contribute to positive upper-ocean density anomalies and subsequent deepening of
 mixed layers, followed by downstream AMOC intensification, after a few years (see Ortega et
 al., 2021 and above references). Evidence for such a prominent role for the LS in low-frequency
 AMOC variability includes the finding that the LS remains the only active deep water formation
 region after low-pass filtering of mixed layer depths (Danabasoglu, 2008), and that low-



130 frequency AMOC variability can be reproduced just by imposing NAO-related buoyancy flux anomalies over the LS region (Yeager & Danabasoglu, 2014). Furthermore, analysis of eddy-permitting / -resolving model simulations continue to show the LS as a key region on decadal time scales. For example, through sensitivity experiments with spatially refined surface forcing, Böning et al. (2023) conclude that the Irminger Sea deep water comes from the LS region and

135 that AMOC decadal variability is governed by the LS convection, especially during exceptionally cold winters. This conclusion is consistent with Yeager et al. (2021) who find a disproportionately large role for the LS in AMOC decadal variability. Specifically, despite a weak mean surface diapycnal transformation, multidecadal AMOC variability can be traced to anomalous production of dense LS Water, with buoyancy forcing in the western subpolar gyre

140 playing a substantial driving role. A caveat here is that models have many biases in the North Atlantic, including potential over-production of the LS Water (Li et al., 2019) and failure to adequately represent the dense water overflows from the Nordic Seas (Danabasoglu et al., 2010).

While there have been intermittent and regionally confined measurements of the AMOC or its contributors at several latitudes as well as inferences from hydrographic sections (e.g., Larsen &

145 Sanford, 1985; Bryden et al., 2005; Toole et al., 2011; Meinen et al., 2010; Send et al., 2011), continuous, trans-basin, and comprehensive observational efforts have gained momentum only during the last couple of decades. The RAPID Meridional Overturning Circulation and Heat-flux Array at 26.5°N is the first such observational program providing AMOC measurements since 2004 (Cunningham et al., 2007). The South Atlantic MOC Basin-wide Array (SAMBA) at

150 34.5°S has been operational since 2009, but it has a gap during roughly the 2011-2013 period (Meinen et al., 2018) and a large uncertainty due to the highly variable Malvinas and Agulhas systems. The Overturning in the Subpolar North Atlantic Program (OSNAP) array at approximately 57°N (Fig. 1) is the third trans-basin observational effort that has been providing data since August 2014 (Lozier et al., 2017). OSNAP fills a crucial missing observational piece

155 focusing on the northern North Atlantic, a region identified as an important source of low-frequency, buoyancy-driven AMOC variability with deep water formation areas (again see reviews by Buckley & Marshall, 2019 and Zhang et al., 2019).




Figure 1. OSNAP sections. The red and orange dots indicate the locations of the moorings at OSNAP-West and OSNAP-East, respectively. The dot-dash and solid black lines show the closest model grid lines through the OSNAP mooring locations based on the NCAR1 and NCAR10 grids, respectively (see Appendix A7). Model bathymetry from NCAR10 is also shown in color (in m). The dark and light purple boxes indicate the Labrador Sea (left) and the Irminger Sea (right) regions, respectively, used in bias analysis.

OSNAP observations, by design, provide estimates of overturning transport magnitudes for the LS and the eastern subpolar gyre side separately. Now available for the 8-year period from August 2014 to July 2022 (Fu et al., 2025), these observations show that conversion of warm and salty upper-ocean Atlantic waters into colder and fresher deep waters primarily occurs in the eastern subpolar gyre side, dominating both the mean and seasonal variability of the overturning circulation (Lozier et al., 2019; Li et al., 2021; Lozier, 2023; Fu et al., 2023). Thus, OSNAP observations reveal a minor role for the LS during this period in stark contrast with numerous modeling studies which connect deep water formation in the LS to downstream low-frequency AMOC variability. While a few earlier studies (Spall, 2004; Straneo, 2006; Pickart & Spall, 2007) also argued for a minor role for the LS in AMOC variability, OSNAP observations provide the most compelling evidence to date with direct comparisons of transport estimates at both OSNAP west and east sections. As alluded to above, a caveat here is that most modeling studies



typically focus on the role of the LS on decadal-to-multi-decadal time scales, whereas OSNAP observational record is still too short to provide guidance on these longer time scales.

This finding regarding differences in the role of the LS between the OSNAP observations and the long-standing view from model simulations has sparked vast interest in the scientific community during the last few years. Published studies make use of observational datasets, both forced ocean-only and coupled model simulations (including at eddy-rich resolutions), and ocean and atmosphere reanalysis products – or their combinations. They primarily focus on the locations of deep-water formation, water mass transformation, surface buoyancy fluxes, and connectivity of these waters with the downstream AMOC, considering the recent OSNAP period. Among these, several studies identify the Irminger Sea or the broader eastern subpolar gyre as having the strongest AMOC-related variability (e.g., Petit et al., 2020; Menary et al., 2020; Megann et al., 2021; Chafik et al., 2022; Fu et al., 2024), consistent with observations. However, the ultimate sources of these deep waters and their variability remain unclear. For example, Petit et al. (2020) argue that local buoyancy fluxes over these basins mostly account for these dense waters of the subpolar North Atlantic. This view is also supported in a recent study by Fu et al. (2024) which shows that the overturning at the eastern part of OSNAP is significantly correlated with water mass transformation forced by surface buoyancy forcing in the Irminger and Iceland basins. Chafik et al. (2022) also indicates the Irminger Sea as the center of action for subpolar AMOC variability, arguing that the NAO is the main driving mechanism for Irminger Sea density variations on multiple time scales. However, this study finds no clear connections between the Irminger Sea density variations and atmospheric heat losses but instead finds that waters dominating the central Irminger Sea are cooled in the first place in the LS and advected into the Irminger Sea. Thus, an alternative interpretation of findings of this study could be that the Irminger Sea AMOC variability has its origins in the LS through direct local water mass transformation via surface buoyancy fluxes in the LS. In another study, Menary et al. (2020) assert that density anomalies generated by surface forcing in the Irminger Sea propagate into the LS, where they dominate the density variability. More recently, Petit et al. (2023) identify the LS as a key pathway for upper North Atlantic Deep Water with its waters sourced from the eastern subpolar gyre, but undergoing further densification within the LS. We finally note that using observational and reanalysis datasets, Zou et al. (2020) and Lozier (2023) show that due to



substantial compensation of thermal and saline anomalies, the volume of newly formed dense waters exported out of the LS is relatively small over the observational period.

As indicated earlier, model fidelity remains a challenge with substantial biases in the northern North Atlantic (e.g., Danabasoglu et al., 2016). The model representations of diapycnal mixing, the Nordic Seas overflows, complex flow pathways, and local recirculations are found to be important contributing factors that impact model biases (e.g., Lozier et al., 2022; Buckley et al., 2023; Lozier, 2023). Some of these processes are represented in closer agreement with observations in high-resolution models compared to their low-resolution counterparts (Hirschi et al., 2020). Using the OSNAP observations, Li et al. (2021) show that while both the Labrador Current and West Greenland Current exhibit large density anomalies, these anomalies are strongly correlated, resulting in only modest variability for overturning in the LS. Accurately representing such correlations; temperature, salinity, and current properties, along with their pathways, with minimal departures from observations are a rather high bar for both low- and high-resolution ocean models to achieve. Indeed, Jackson & Petit (2022) find that models with a more saline LS tend to produce stronger water formation in this region, with correspondingly stronger connections with downstream AMOC. Nevertheless, they conclude that models are generally in good agreement with the OSNAP transport observations.

In the present study, we take a step back and provide a basic, yet much-needed evaluation of simulated transports against OSNAP observations for the 2014-2022 period using a large set of forced ocean – sea-ice (FOSI) simulations. These simulations largely follow the Ocean Model Inter-comparison Project (OMIP) protocol (Griffies et al., 2016) and are forced using the atmospheric datasets based on the Japanese atmospheric reanalysis product, suitably adjusted to run such FOSI experiments (JRA55-do; Tsujino et al., 2018; 2020). This protocol is intended to provide a controlled and coordinated framework to evaluate ocean components of coupled models that participate in CMIPs (Griffies et al., 2009; Danabasoglu et al., 2014). There are 9 participating groups / centers from around the world, contributing a total of 18 simulations with 6 different ocean models in this study. The horizontal resolutions of the simulations range from nominal non-eddy 1° to eddy-resolving resolutions of 0.1°-0.05°. Thus, a goal is to assess how the representation of overturning transports at OSNAP changes with model horizontal resolution. The present effort aims to provide a benchmark for simulated transports and related properties at the OSNAP sections in comparison to observations, evaluated in depth, σ_0 , and σ_2 .



spaces – in the latter two, potential densities are referenced to the surface and 2000-m depth, respectively. As such, the present manuscript is largely descriptive. We mainly provide a catalog of model solutions; detailed analyses of individual model results and specific reasons for their differences from and similarities to the observations are not fully covered here. However, an analysis of water mass formations and transformations will be presented in a separate subsequent study. Finally, we note that because the observational record is short, we cannot yet address the relative roles of the LS vs. the eastern subpolar gyre in AMOC variability on decadal and longer time scales.

The manuscript is organized as follows. In Sect. 2, we summarize the OSNAP observing system, model simulations and the forcing datasets, how the transport lines are determined, and how the AMOC is calculated in depth and density space. We then present results for transport timeseries in Sect. 3; time-mean transports in σ_0 space in Sect. 4; time-mean transports in depth and σ_2 space in Sect. 5; transport variability in Sect. 6; temperature and salinity diagrams in Sect. 7; and temperature, salinity, and density biases in Sect. 8. We provide a summary and our conclusions in Sect. 9. Brief model descriptions are given in Appendix A. Finally, Appendix B presents a brief analysis of cycle-to-cycle differences in transports.

2 Observational Data, Participating Models, Forcing, and Methods

2.1 OSNAP Observing System

The OSNAP observing system is designed to provide a continuous record of the full-water column, trans-basin transports of volume, heat, and freshwater in the subpolar North Atlantic (Lozier et al., 2017). It consists of two sections: OSNAP-West, which runs from the southeastern Labrador shelf to the southwestern tip of Greenland, and OSNAP-East, which runs from the southeastern tip of Greenland to the Scottish Shelf (Fig. 1). We will refer to these sections as OW and OE, respectively. OSNAP uses a combination of ~60 fixed moorings located at the continental boundaries and on both sides of the Reykjanes Ridge to obtain direct measurements of velocities at the boundaries, and geostrophic and Ekman flows in the interior to estimate meridional volume transports. In regions of complex topography, the moored arrays were supplemented by glider surveys in the early part of the OSNAP observational period. The gridded temperature and salinity data represent a combination of data from Argo floats in the



upper 2000 m; moorings; ship-based Conductivity-Temperature-Depth (CTD) sections for regions below 2000 m without mooring observations; gliders for the Hatton-Rockall Basin; and World Ocean Atlas monthly climatology for shallow coastal waters without data from moorings and Argo (Li et al., 2017). The data are objectively mapped to the OSNAP sections with a 20-m
 270 resolution in the vertical and a nominal 0.25° horizontal resolution with the original mooring positions preserved. The OSNAP record has been recently extended to 8 years, covering the August 2014 – July 2022 period (Fu et al., 2025). We refer to the OSNAP observational data simply as OSNAP.

2.2 Models

275 We use 18 simulations from 9 groups run with 6 different models as summarized in Table 1. The ocean models are the HYbrid Coordinate Ocean Model (HYCOM), the Modular Ocean Model version 5 (MOM5) and version 6 (MOM6), the Model for Prediction Across Scales – Ocean (MPAS-O), the Nucleus for European Modelling of the Ocean (NEMO), and the Parallel Ocean Program version 2 (POP2), which are coupled to a variety of sea-ice models (see Appendix A).
 280 The horizontal resolutions range from non-eddy-resolving resolutions of near 1° to eddy-permitting and eddy-resolving resolutions of 0.25° - 0.05° . We denote each simulation using the acronym of the group which produced the simulation followed by the nominal horizontal resolution of a simulation in degrees. For example, NCAR1, NCAR67, and NCAR10 refer to simulations performed by the NCAR group, using nominal 1° , 0.67° , and 0.10° horizontal
 285 resolution, respectively. With respect to their vertical coordinates, while many of the simulations use depth / level coordinates – both z and z^* , a few simulations employ hybrid coordinates. The number of vertical levels / layers varies from 36 in FSU08 to 98 in CMCC06 with most models using between 46-75, with varying placement of levels in the vertical, usually with finer resolution near the surface. To partly help reduce crowding in the plots, we separate the
 290 simulations into a low-resolution (LR) group, which includes all the simulations with a resolution of 0.25° or coarser (an arbitrary cutoff), and a high-resolution (HR) group, which has the remaining simulations with finer resolutions. There are 12 and 6 simulations in each group, respectively. Brief descriptions of the models along with their configurations, parameter settings, and references, if available, presenting evaluations of their simulations are provided in Appendix
 295 A.



2.3 Forcing Datasets

The FOSI simulations largely follow the OMIP protocol (Griffies et al., 2016). They are forced with the atmospheric datasets based on the Japanese atmospheric reanalysis product (JRA55; Kobayashi et al., 2015) which were adjusted extensively to correct for biases in the reanalysis and for obtaining reasonably balanced heat and freshwater budgets (Tsujino et al., 2018). This adjusted dataset, referred to as JRA55-do, has a spatial resolution of 55 km with a temporal resolution of 3 hr. Except for the runoff dataset, all the fields are directly from the reanalysis product, thus representing a rather self-contained forcing dataset for the 1958-2023 period. The heat, freshwater, and momentum fluxes needed to force the simulations are based on the bulk formulae described in Large and Yeager (2004) and Large and Yeager (2009).

The simulations apply restoring of sea surface salinity (SSS) to an observed monthly-mean climatology. This is because FOSI simulations do not have many of the feedbacks that exist in fully coupled systems with an active atmosphere, as discussed in Griffies et al. (2009) and Danabasoglu et al. (2014). Moreover, the lack of any appreciable local feedbacks between surface freshwater fluxes and SSS can lead to unrealistic salinities locally, mostly due to biases in precipitation. Usually, the upper 10-m average monthly salinity climatology from the World Ocean Atlas version 2 (WOA13; Zweng et al., 2013; Boyer et al., 2015) is used as the restoring field. The OMIP protocol does not prescribe a particular restoring procedure, and the modeling groups can choose their optimal restoring strength based on how their models represent several key features in comparison to available observations, such as the transport and time evolution of the AMOC to ensure that it does not get too weak or collapse. Usually SSS restoring is applied globally, including ice covered regions, and its global mean is subtracted to ensure that the restoring term does not impact the global salt content. The SSS restoring approaches and strengths used by the participating groups are presented in Table 2.

The OMIP protocol calls for multi-century-long simulations forced with repeat cycles – usually 5 – of the forcing datasets. While an intent is to achieve quasi-equilibrated solutions to the extent possible, this approach has a few drawbacks. One is that the repeat cycle introduces an unphysical jump in forcing from the last year of a particular cycle (e.g., 2023) back to the first year of the subsequent cycle (e.g., 1958). So, at the beginning of the next cycle, the ocean initial state reflects that of year 2023 instead of year 1958. Consequently, it is not recommended to use



the first 15-20 years of a forcing cycle in any analysis. Second, even with multiple cycles, the degree of repeatability of solutions from one cycle to the next is both model and variable dependent (e.g., Danabasoglu et al., 2014). Third, such long simulations are not computationally affordable for HR simulations.

330 As detailed in Appendix A, the simulations do not strictly follow the OMIP protocol for various reasons that include which end year was available for forcings when the simulations started. Many simulations use a slightly shorter forcing period for the cycles before the very last cycle. For example, ANU1 and ANU25 simulations go for 6 forcing cycles where the first 5 cycles use the 1958-2018 period, and the simulations were extended to the end of 2022 only for the last
 335 cycle. Similarly, NCAR1 and NCAR10 simulations use the 1958-2018 period for the first 4 cycles and have the last (5th) cycle extended to the end of 2023. Due to the computational costs, most of the HR simulations are integrated only for 1 forcing cycle. In summary, the participating simulations have been integrated for a range of 1 to 6 forcing cycles.

These differences in simulation lengths as well as limited availability of datasets only for specific
 340 forcing cycles for some simulations certainly present challenges for our analysis. However, because our analysis concerns the last decade of a forcing cycle, cycle-to-cycle differences in overturning transports are relatively small during the last decade compared to those of the earlier periods as discussed in Appendix B. Therefore, we are confident that use of different cycles from the participating simulations does not impact our general conclusions. Noting that we analyze the
 345 same forcing cycles for a given LR and HR set of simulations from the same group, we use the following cycles for each simulation: cycle 1 for CMCC25, CMCC06, FSU72, FSU08, NCAR67, NOC1, NOC25, NOC12, and UKMO25; cycle 4 for ANU10; cycle 5 for E3SM27, GEOMAR25, GEOMAR05, NCAR1, and NCAR10; and cycle 6 for ANU1, ANU25, and GFDL25.

350 As indicated in Sect. 2.1, the OSNAP observations cover the August 2014 – July 2022 period. For ease of analysis, we use the full 2014 – 2022 period for model simulations. For E3SM27, GFDL25, and NCAR67, we use shorter periods from 2014 to their respective simulation end years, i.e., 2020, 2018, and 2018, respectively. Finally, we employ monthly mean fields for both OSNAP and model simulations.

355



2.4 Transport Lines

The OSNAP array is not aligned with a latitude line or models' grid lines, requiring a careful method to extract the model outputs along the OE and OW lines. For this purpose, we developed a tool to obtain the best or closest model *broken grid line* through the OSNAP mooring locations for each participating model. The tool returns the grid indices along the line. These indices are then used to index a Python Xarray to extract the desired fields along this broken line for each simulation. As such, the tool provides direct access to the simulated fields across the OSNAP sections, without requiring any interpolations. As an example, the solid and dot-dash black lines in Fig. 1 show the best broken grid lines for NCAR10 and NCAR1, respectively. Because these lines follow a model's grid, both zonal and meridional velocity components are used in transport calculations depending on whichever component is normal to the local grid line. We note that we chose to use this approach based on the actual simulation outputs to calculate transports instead of extracting the sections strictly following the OSNAP observation grid for two reasons. First, it is a straightforward approach, providing the actual transport simulated by each model. Second, comparisons of both methods using NCAR simulations produced very similar results (not shown), providing justification for our approach.

2.5 Calculation of AMOC

The overturning streamfunction ψ is commonly calculated in both depth and density space using the following equations:

$$\psi(y, z, t) = - \int_{z_b}^{z'} \int_{x_w}^{x_e} v(x, y, z, t) \, dx \, dz',$$

$$\psi(y, \sigma, t) = - \int_{\sigma_b}^{\sigma'} \int_{x_w}^{x_e} v(x, y, \sigma, t) \, dx \, \frac{\partial z}{\partial \sigma} \, d\sigma',$$

where v is the plane-normal velocity, positive northwards; t is time; z is the vertical coordinate, positive upwards; x is the zonal coordinate, positive eastwards; y is the meridional coordinate, positive northwards; and σ is density, either in σ_0 or σ_2 . The subscripts e and w denote the eastern and western limits, respectively, of the zonal integral; the subscript b represents bottom,



meaning that the integrations are performed from bottom up, e.g., from the highest density in
 385 density space; and $\partial z / \partial \sigma$ is the isopycnal thickness. In these equations, v is in m s^{-1} and the
 lengths are in m. Thus, the transports are in $\text{m}^3 \text{s}^{-1}$, displayed in Sv ($= 10^6 \text{ m}^3 \text{s}^{-1}$). With the sign
 convention used in the above equations, the negative and positive profile slopes indicate
 poleward and equatorward transports, respectively, in the overturning plots presented here. In our
 transport calculations, we only use the resolved-flow velocities, i.e., the relatively small
 390 parameterized contributions are not included. We note that the bottom-up integration method
 used here is usually adopted in the modeling community to compute overturning transports,
 because the boundary condition at the bottom, i.e., a zero transport, is known. This approach,
 however, is in contrast with the top-down approach employed in the OSNAP observational
 estimates (e.g., Lozier et al., 2019). So, we expect some small differences between our transport
 395 values and those published for OSNAP observations, but all our transports here are consistently
 calculated using the same bottom-up approach, including for OSNAP.

Overturning in density space directly connects with diapycnal water mass transformation.
 However, in both depth and density space, there are substantial cancellations of northward and
 southward flows at constant depth and density, respectively. We believe that analyzing the
 400 AMOC in both depth and density space provides complementary information – an approach also
 advocated by Lozier et al. (2019) and Zhang & Thomas (2021). Therefore, in our analysis here
 we apply both methods.

For depth-space analysis, we use each model's native vertical discretization. For density-space
 analysis, we employ common density ranges and bin increments across all simulations.
 405 Specifically, σ_0 calculations are referenced to the ocean surface, and they use a density range of
 24.0–28.5 kg m^{-3} with 105 density bins. The bin increments are 0.1 and 0.02 kg m^{-3} for densities
 less and greater than 27.0 kg m^{-3} , respectively. σ_2 calculations use a reference depth of 2000 m.
 The density range is between 28.0 and 38.0 kg m^{-3} . We use 85 bins with increments of 0.2 kg m^{-3}
 for 28.0–35.0 kg m^{-3} , 0.1 kg m^{-3} for 35.0–36.0 kg m^{-3} , and 0.05 kg m^{-3} for densities $> 36.0 \text{ kg m}^{-3}$.
 410 Also, for consistency across all simulations, we use monthly-mean potential temperature and
 salinity to calculate densities using the TEOS-10 equation of state (McDougall & Barker, 2011).

As discussed in Lynn & Reid (1968), there is no single potential density that can be used for all
 purposes. While σ_0 is useful to connect surface water mass transformations to deeper overturning



transports, it does not necessarily depict an accurate representation of transports at depth because
 415 it ignores the pressure effects on density. Indeed, especially in the northern North Atlantic,
 potential density referenced to 4000-m depth, i.e., σ_4 , may be more appropriate. In this study, we
 adopt use of σ_2 as a compromise, but we also think that it is more suitable for our purposes as we
 focus on transports primarily between 1000 – 3500 m. Thus, effects of pressure on density and
 transports are appropriately included. σ_0 is employed to expedite comparisons with observations.
 420 In OSNAP, a zero-net-mass (actually volume) transport constraint across the entire section is
 enforced (Lozier et al., 2019). Specifically, a mean southward transport of 1.6 Sv across OW –
 intended to represent long-term transport measurements across the Davis Strait – is compensated
 by an equivalent northward transport at OE. This is accomplished through an a posteriori
 application of a spatially uniform, but time dependent compensation velocity. In this approach,
 425 there is no transport into the Arctic Basin associated with the flow into the Arctic Ocean through
 the Bering Strait. Lozier et al. (2019) report that the OSNAP transports change very little when
 this flow is accounted for. Therefore, when computing the simulated transports, we do not
 impose any constraints. As such, any non-zero transports at the surface reflect the flows in and
 out of the Arctic Basin associated with the Bering Strait Throughflow. We note that in most of
 430 our plots we use a lower limit for density and do not show near surface or upper-ocean
 transports.

These transport calculations, along with the tool to obtain the best broken grid line at the OSNAP
 sections, will be incorporated in a new version of a Python-based Meridional overErTurning
 DiagnostIC (METRIC) Package. METRIC enables consistent calculations of observed and
 435 simulated AMOC estimates at various observational sites in the Atlantic Ocean. The package can
 be used with different ocean models at various resolutions, and it includes a few additional,
 alternative approaches to calculate these transports. It is freely available to the community at
<https://github.com/NCAR/metric> (Castruccio, 2021).

3 Transport Timeseries

440 We start with the time evolution of transport profiles in depth space presented in Fig. 2 as
 transports per unit depth, i.e., in Sv m⁻¹. The OSNAP profiles are based on the gridded velocity
 datasets, and profiles only from NCAR10 and NCAR1 are shown as examples of model



simulations for brevity. The OSNAP profiles at both sites exhibit a rather streaky behavior, primarily due to the methodology used to estimate basin interior geostrophic transports, which are derived from full-depth dynamic height moorings bracketing the basin interior. Temperature and salinity measurements on these moorings are typically hundreds of meters apart vertically. In general, transport magnitudes and the depth ranges of northward and southward transports are in better agreement between OSNAP and NCAR10, particularly at OE, than with NCAR1. The latter shows a very prominent, deeper-penetrating seasonal cycle at both sites which is much weaker, especially at OW, in both OSNAP and NCAR10. Nevertheless, there are differences between OSNAP and NCAR10 such as a band of northward transports between about 1700 – 2300 m present in OSNAP at OW but missing in NCAR10.




Figure 2. Transport profile timeseries in depth space from (a, d) OSNAP, (b, e) NCAR10, and (c, f) NCAR1. The top and bottom rows are for OW and OE, respectively. Positive and negative contours indicate northward and southward transports, respectively. The transports are in Sv m^{-1} . Note that different scales are used for the top and bottom panels.

The corresponding timeseries in density (σ_0) space are provided in Fig. 3. These are the integrated transports in Sv , and the streak-like features seen in depth-space OSNAP (Figs. 2a and 2d) disappear after binning. As in Fig. 2, prominent features include better agreement between OSNAP and NCAR10 than with NCAR1 and a strong seasonal cycle in NCAR1. Indeed, a seasonal cycle in OSNAP is not as pronounced as in both simulations at either site. While the magnitudes of the transports are comparable at OE among observations and model simulations,



the simulated transports are larger than in OSNAP at OW, particularly so for NCAR1. The solid
 465 black lines in the panels show the timeseries of the maximum positive transport which occurs at
 different density classes. The details of these timeseries will be discussed below, considering all
 the participating simulations. Here, we note that the density at which the maximum transports
 occur varies considerably in time in both observations and simulations. So, even if OSNAP and
 simulations have similar maximum transports at a given time, the density at which this occurs
 470 can differ substantially. At OW, OSNAP has two interesting characteristics. First, the positive
 transport is rather weak, particularly prior to 2018, with nearly no positive transport in early
 2016. Second, there are nonnegligible negative transports in density classes $< 27.6 \text{ kg m}^{-3}$
 throughout the timeseries. This latter feature is somewhat present in NCAR1 but with larger
 magnitudes and rather strong seasonality.

475




Figure 3. Transport profile timeseries in density (σ_0) space from (a, d) OSNAP, (b, e) NCAR10, and (c, f) NCAR1. The top and bottom rows are for OW and OE, respectively. The solid black lines in all the panels show the maximum positive transport timeseries. The transports are in Sv. Note that different scales are used for the top and bottom panels.

480 To provide a broader context for the observational transport estimates and their variability,
 annual-mean maximum transport timeseries for the full forcing cycle from the simulations are
 presented in Fig. 4. These transports are constructed from the average of the maximum monthly
 mean transports, where monthly mean temperature and salinity are used to compute monthly
 mean densities. The solid black lines represent OSNAP, while the dashed black lines indicate the



485 respective multi-model means (MMM) for each panel. Not surprisingly, the maximum transports
and their variability differ considerably among the simulations, but they show general agreement
with OSNAP with their stronger transports at OE which dominate the total transports. At OW
(Figs. 4a and 4b), the simulations largely show transports of < 10 Sv. A few of the simulations,
e.g., ANU1, ANU25, E3SM27, and FSU27, have relatively weak transports during various time
490 segments. There seems to be a slight weakening trend at OW roughly after the mid-1990s in
simulated transports with the OSNAP observations coinciding with a period of low transports ($<$
 ~ 5 Sv) in all (but one) of the simulations. NCAR1 is a clear outlier with a mean transport of ~ 12
Sv for the 1958-2022 period with a rather large amplitude. This represents an improvement
compared to a mean transport of 18.7 Sv for the 1961-2007 period reported in Li et al. (2019) for
495 a previous version of the model that was forced with the CORE-II atmospheric datasets (see
Danabasoglu et al., 2014). The simulated transports at OE tend to be between 10 – 20 Sv (Figs.
4c and 4d) with a slight weakening trend again after the mid-1990s. In contrast with OW,
OSNAP at OE is at the higher end of the simulated transport magnitudes. The maximum total
transport magnitudes (Figs. 4e and 4f) primarily reflect those of OE, noting that these total
500 transports in density space do not need to reflect the sum of the OW and OE transports as they
are not obtained at a constant density.




Figure 4. Annual-mean maximum transport timeseries in σ_0 space at (a, b) OW, (c, d) OE, and (e, f) total. The left and right panels are from LR and HR, respectively. The solid black lines show the OSNAP transports. The dashed black lines represent the respective multi-model means.

505 The mean seasonal cycles of maximum transports in σ_0 space from each simulation are presented in Fig. 5. The solid black lines represent OSNAP with the gray shading showing ± 1 monthly standard deviation (SD). The respective MMMs for each panel are also provided by the dashed black lines. At OW (Figs. 5a and 5b), the seasonal cycle is relatively weak, with OSNAP showing generally larger transports in spring and smaller transports in winter, but with a minimum in September. Most of the simulations do not capture this observed phasing, also displaying differing maximum and minimum transport months among each other. Except for NCAR1, LR simulations cluster around the OSNAP spread. NCAR1 is a clear outlier with the largest transport and seasonal cycle amplitude. In contrast, ANU25, E3SM27, and FSU72 have the lowest transports, near the lower bound of the OSNAP spread. HR simulations tend to be

510



515 outside the OSNAP spread with higher magnitudes. Here, ANU10 and CMCC06 are the closest
 to the OSNAP spread. At OE (Figs. 5c and 5d), OSNAP has a better-defined seasonal cycle –
 compared to that at OW – with maximum transports in spring and minimum transports in winter.
 Arguably, the simulations capture the observed seasonal cycle phasing relatively well as reflected
 in the MMM. An exception is the last few months, particularly evident in HR simulations, where
 520 all HR models show an increasing transport after September in contrast with a decreasing trend
 in OSNAP. Most LR and HR simulations have transports that are below the OSNAP spread,
 again as depicted in the MMM. Among all the simulations, only ANU25 remains within the
 OSNAP spread throughout the seasonal cycle. Among LR simulations, NCAR1, and E3SM27
 and GEOMAR25 bracket the OSNAP seasonal cycle at the high and low ends, respectively.
 525 ANU10, FSU08, and GEOMAR05 have the lowest transports at HR with the largest departures
 from OSNAP. For the total transports, while most LR simulations show weaker than observed
 transports, all but two (ANU10 and GEOMAR05) of the HR simulations are largely within the
 OSNAP envelope (Figs. 5e and 5f). The large-amplitude seasonal cycles at OW and OE in
 NCAR1 mostly cancel each other, resulting in a much smaller amplitude for the total transport.
 530 Finally, to provide a quantitative comparison of the simulated seasonal cycles and those of
 OSNAP, we calculate the root-mean-square differences between MMM and OSNAP as 1.5 Sv
 (LR) and 2.8 Sv (HR) for OW, 3.4 Sv (LR) and 2.9 Sv (HR) for OE, and 3.4 Sv (LR) and 1.8 Sv
 (HR) for the total transports.

NCAR1 is a clear outlier with the largest transport and seasonal cycle amplitude at OW among
 535 the simulations. The phasing of its seasonal cycle – which is out-of-phase with OSNAP – reflects
 that of Fig. 3c with a minimum transport in April. This results from incursions of the negative
 transport cells into higher density classes, indicating that the associated upper-limb southward
 flow and lower-limb northward return flow occur at higher density classes in NCAR1 compared
 to those of OSNAP and NCAR10. Such negative transport cell incursions are also present in both
 540 OSNAP and NCAR10, but they are less well organized, and their densities and transports remain
 much smaller than in NCAR1. It is also interesting to note that maximum mixed or boundary
 layer depths in the LS region occur in March (not shown), although the OSNAP section is not
 co-located with the regions of maximum mixed layer depths. Instead, the OSNAP section was
 purposely located near the exit of the LS to capture the bulk of the waters exported from the



545 basin. All these aspects, including such seasonal cycle features, will be investigated in detail in a follow up study, considering water mass formation and transformation analysis.




Figure 5. Time-mean maximum transport seasonal cycles in σ_0 space at (a, b) OW, (c, d) OE, and (e, f) total. The left and right panels are from LR and HR, respectively. The solid black lines show the time-mean OSNAP maximum transport timeseries with gray shading indicating ± 1 monthly standard deviation. The dashed black lines represent the respective multi-model means.

4 Time-Mean Transports in σ_0 Space

Figure 6 shows the time-mean transport profiles in σ_0 density space. As in the previous figures, the solid black lines represent OSNAP with the gray shading showing ± 1 monthly SDs, and the respective MMMs for each panel are also provided by the dashed black lines. The figure clearly demonstrates that the simulated transports are much larger at OE than at OW with the total



transports dominated by those of OE, all in agreement with OSNAP. At OW (Figs. 6a and 6d), the simulations largely follow the OSNAP profile. However, at high density classes ($\sigma_0 > \sim 27.7$ kg m⁻³), there are differences from observations as well as among the simulations. While these high-density transports are relatively small, they can play an outsized role in downstream AMOC variability (Yeager et al., 2021). NCAR1 has the largest maximum transport with about 9 Sv. While FSU72 shows the highest negative transport magnitude with a distinct peak around 27.65 kg m⁻³, a feature not present either in OSNAP or other simulations, FSU08 deviates the most from observations and the other simulations at density classes < 27.5 kg m⁻³. At OE (Figs. 6b and 6e), a large majority of the simulations have positive transports with negative slopes for densities $< \sim 27.3$ kg m⁻³. This contrasts with OSNAP that shows a much steeper negative density slope for $\sim 27.2 < \sigma_0 < \sim 27.4$ kg m⁻³ and nearly no net flow for densities < 27.2 kg m⁻³ (see also Fig. 3d). For $\sim 27.3 < \sigma_0 < \sim 27.8$ kg m⁻³, the simulations tend to agree better with OSNAP with most within the observed range. Here, ANU25, FSU72, and NCAR67 show lower transport magnitudes, and NCAR1 has the largest magnitude with a transport of ~ 18 Sv. Observed transports at high density classes (> 27.8 kg m⁻³) appear to be shifted to somewhat lower densities in the simulations. The total transport features are mostly similar to those at OE (Figs. 6c and 6f). FSU72 has the smallest maximum transport with ~ 7 Sv due to its large negative transport at OW. The maximum transport is < 10 Sv for E3SM27 as well. HR profiles, in general, show better agreement with OSNAP than those of LR for $\sigma_0 > \sim 27.4$ kg m⁻³. Finally, we note that all FSU08 profiles are noisy, particularly at high densities, which may be due to a mismatch between σ_0 increments used in the analysis and the resolution of the model σ_2 coordinates. Specifically, with relatively small σ_0 increments at high density classes, transports within a σ_2 model layer can be binned into different σ_0 layers, thus leading to some noise.




Figure 6. Time-mean transport profiles in σ_0 space at (a, d) OW, (b, e) OE, and (c, f) total. The top and bottom panels are from LR and HR, respectively. The solid black lines show the OSNAP transports with gray shading indicating ± 1 monthly standard deviation. The dashed black lines represent the respective multi-model means.

5 Time-Mean Transports in Depth and σ_2 Space

The time-mean overturning transports in both depth and σ_2 density space from OSNAP and all the simulations at OW and OE are provided in Figs. 7-8 and Figs. 9-10, respectively. In each set of plots, the main panels (color shading) show southward (blue) and northward (red) transports as a function of density and depth. These are transports in Sv, calculated using the common σ_2 bins, listed in section 2.5, and each model's vertical grid. We prefer to use Sv for this purpose, rather than transport per unit depth, i.e., Sv m^{-1} , because this way the transports can be summed easily in either depth or density space. As such, the profile panels on the right then show the summed transports in depth space. Specifically, the thin blue, red, and black lines display the



summed transports in depth space for the southward, northward, and total, i.e., the sum of the southward and northward, transports, respectively. When the total transports are aggregated (or
 595 integrated) in the vertical, starting from the bottom, the familiar overturning transport profiles in depth space are obtained (thick black lines). Similarly, when the transports of the main panels are summed across constant density bins, the top plots are obtained, showing the transport profiles in density space. The thick black lines show the overturning transports in density space, aggregated starting from the densest bin class. These are essentially the same profiles shown in Fig. 6, but
 600 they are now calculated in σ_2 space instead of σ_0 . Therefore, calculated maximum transports differ as noted below. We note that this analysis can be easily extended to depth – temperature or depth – salinity space to obtain respective transports (not shown). As Figs. 7-10 contain a lot of information, we only discuss a few notable features for brevity.

In OSNAP, the OE transports are more distinctly organized in comparison to those of OW, with
 605 continuous southward and northward transport bands occurring at higher and lower density classes, respectively, throughout all depths. At OW, the structure is a bit more complex. A continuous southward transport band between about 1000 – 3500-m depth splits two northward bands occurring at mid-depth and abyssal ocean. With rather similar high-density waters, OE exhibits broader density ranges at all depths due to the presence of lighter density waters
 610 compared to OW. At both sections, the transports can exceed 0.5 Sv in both directions. The summed transports (right panels) reveal substantial cancellations of southward and northward flows in depth space, producing relatively small total transports, at both sections. These profiles have some noise in the upper 500 m. The integrated overturning profile in depth space has a rather odd shape at OW without a well-defined maximum or flow structure, essentially reflecting
 615 the noisy velocity bands of Fig. 2a. Nevertheless, its maximum transport is slightly less than 2 Sv. In contrast, the depth-space overturning profile at OE is well-defined, with a maximum transport of about 8.2 Sv. There are cancellations of southward and northward transports in density space as well (top panels). The maximum transports are 5.3 and 14.0 Sv for OW and OE, respectively. We note that these transports in σ_2 space are larger than the corresponding
 620 transports of 3.8 and 12.9 Sv in σ_0 space presented in Fig. 6. These indicate that while the ratio of the transports at OE and OW is 3.4 in σ_0 , it drops to 2.6 in σ_2 space, representing a 25% reduction. We will come back to this point later in the manuscript.



It is interesting to note that some degree of cancellation of northward and southward flows in OSNAP occurs at all depths. In density space, these cancellations are less uniform, especially at
 625 OW where they are pronounced for $\sigma_2 > 36.9 \text{ kg m}^{-3}$. At OE, any cancellation is confined to intermediate density classes with not much cancellation of northward flow of $\sigma_2 < 36.4 \text{ kg m}^{-3}$ in the upper 1000 m and the southward flow of $\sigma_2 > 37.1 \text{ kg m}^{-3}$ below ~2000 m.

Arguably, the simulations capture observed transport features better at OE (Figs. 9 and 10) than at OW (Figs. 7 and 8). In all simulations, a well-defined southward transport band at high density
 630 classes that extends through almost the entire depth and an upper-ocean northward transport region extending from the surface to about 1500-m depth and diagonally in density from roughly 35.5 to 36.75 kg m^{-3} are captured at OE. However, the northward transport band at deeper levels and at higher densities seen in OSNAP is largely missing in all simulations except for FSU08 and UKMO25. Consequently, summed northward transports do not cancel much of southward
 635 transports below about 1500-m depth and in high density classes. In depth space, the maximum aggregated transports at OE are between 4.5 Sv (GFDL25) and 8.9 Sv (ANU25 and ANU1) occurring between 750 – 1250 m, but mostly around 1000-m depth, noting that the OSNAP estimate is ~8.2 Sv. In σ_2 space, the maximum aggregated transports range from about 10 Sv in E3SM27 to about 15 Sv in ANU1, ANU25, CMCC06, NCAR10, and NCAR1, bracketing the
 640 OSNAP transport of 14 Sv. While most simulations show a density range of 35.75 to 37.25 kg m^{-3} for the positive transport segment in σ_2 profiles in agreement with OSNAP, this range is much narrower in ANU25 and FSU72, remaining between about 36.25 and 36.75 kg m^{-3} (top panels for all simulation sets). The secondary peak near 36.9 kg m^{-3} seen in OSNAP is present in FSU08, NCAR1, and NOC1, likely due to larger northward flows near this density in these simulations
 645 compared to the other simulations. Among the simulations, FSU08 and FSU72 exhibit somewhat noisier transports in depth – density binned transports, likely due to the choice of their hybrid vertical levels.

At OW, the simulations appear to struggle to capture observed binned transport features in depth – σ_2 panels with notable differences among the simulations as well. Although these binned
 650 transports show similar transport magnitudes to those at OE, their summed and aggregated transport magnitudes are smaller than at OE, consistent with observations. There are cancelling southward and northward transports in both depth and σ_2 space and most simulations reproduce



the observed structure of the aggregated transports reasonably well. In depth space, the maximum transports are mostly around 2 Sv within 1000-2000-m depth range, with a few
655 exceptions. Specifically, ANU1, E3SM27, and GFDL25 have rather weak transports of about < 0.5 Sv. In E3SM27, there is essentially no significant transport at OW. In ANU1 and GFDL25, there are significant cancellations between southward and northward transports. While some simulations exhibit noise in their summed southward and northward transport profiles which are cancelled in the aggregated transports, such noise is particularly large and remains in aggregated
660 transport profiles in FSU08 and FSU72. In density space, E3SM27 does not show any notable transports for $\sigma_2 > \sim 36 \text{ kg m}^{-3}$. At the opposite end, NCAR1 has the highest transport with 13.1 Sv (off scale in Fig. 8).




Figure 7. Time-mean transports at OW in depth and density (σ_2) space from OSNAP and from ANU10, ANU25, ANU1, CMCC06, CMCC25, E3SM27, FSU08, FSU72, and GEOMAR05. In each set, the bottom left panels show southward (blue) and northward (red) transports in Sv. The



bottom right panels are the summed transports in depth space with the thin blue, red, and black lines displaying the southward, northward, and total, i.e., the sum of the southward and northward, transports, respectively. Similarly, the top panels show the summed transports in density space. In the right and top panels, the thick black lines are the respective overturning transports with aggregation (integration) starting from the bottom and the densest bin for depth and density space, respectively.




Figure 8. Same as in Fig. 7, but for GEOMAR25, GFDL25, NCAR10, NCAR1, NCAR67, NOC08, NOC25, NOC1, and UKMO25. The density-space aggregated maximum transport in NCAR1 is 13.1 Sv.




Figure 9. Same as in Fig. 7, but for transports at OE.




Figure 10. Same as in Fig. 8, but for transports at OE.



To expand upon our earlier discussion, we assess the impacts of using different potential densities on overturning transports considering the scatter plots presented in Fig. 11. Panels a-c show the time-mean maximum transport magnitudes as a function of σ_0 and σ_2 densities for the OW, OE, and total transports, respectively. For OW and the total, the σ_2 transports are consistently higher than those in σ_0 in all simulations, on average, by 2.47 Sv and 1.74 Sv, respectively. At OE, all but one simulations are larger in σ_2 than in σ_0 with an average increase of 1.19 Sv which is smaller than those for OW and the total. To quantify these transport changes, we first compute the ratios of OW and OE transports for σ_0 and σ_2 separately which can be denoted as R_{σ_0} and R_{σ_2} , respectively. We then produce a scatter plot of $R_{\sigma_2}/R_{\sigma_0}$ presented in Fig. 11d. This measure considers the relative changes in the maximum transports with density at both OW and OE. The figure shows that this ratio exceeds unity for all simulations, with most of them clustering roughly between 1.2 – 1.5, indicating larger OW contributions in σ_2 space compared to those of σ_0 . ANU25 and FSU72 have the largest ratios with ~ 1.9 and 2, respectively. We note that the spread of the LR simulations is much larger than that of the HR simulations, which cluster around a ratio of 1.3. These model-based findings are consistent with the OSNAP observations which are included as black dots in all the panels, showing roughly a middle-of-the-pack value of ~ 1.4 in Fig. 11d.




Figure 11. Scatter plots of time-mean maximum transports in σ_0 vs. σ_2 space for (a) OW, (b) OE, and (c) total. (d) Scatter plot of $R_{\sigma_2}/R_{\sigma_0}$ (see text) as a function of the total transport in σ_0 space. The low- and high-resolution simulations are indicated by dots and stars, respectively. The regression lines (excluding OSNAP) along with their coefficients are included in each panel. The gray one-to-one lines are also shown for reference in (a-c).

6 Transport Standard Deviations in σ_2 Space

We examine the variability of maximum transports, considering the scatter plots of time-mean maximum transports and their monthly SDs given in Fig. 12 for OW, OE, and their total. Obtained in σ_2 space, these plots show statistically significant correlations between the strength of the maximum transports and the amplitudes of variability at both OW and OE. While all the simulations are subject to the same external variability, their SDs differ considerably due to differing internal variabilities, primarily in response to buoyancy forcings: especially at high latitudes, convective / deep-water formation events can differ across simulations. The simulated mean SDs are slightly higher at OE than at OW, but the spread of the simulated SDs is larger at OW than at OE. While OSNAP SD (= 1.64) at OW is lower than those of most simulations, OSNAP SD (= 3.18) at OE is larger than in all simulations. Relatedly, there are 10 and 15 simulations within 1 SD of the observational maximum transports at OW and OE, respectively.



At OW, CMCC25, NCAR67, NOC1, and NOC25 are the closest to OSNAP SD. Here, FSU08 and NCAR1 have the largest SDs (> 3.5), and E3SM27 shows the smallest SD with ~ 0.5 . At OE, SDs of ANU1, NCAR10, and NCAR1 are in better agreement with that of OSNAP, and E3SM27 and GEOMAR25 SDs of < 2 are the lowest among the simulations. For the total transports, OSNAP SD is again larger than those of nearly all simulations, with FSU08 and GFDL25 closest to it. E3SM27, GEOMAR05, GEOMAR25, and CMCC25 have the lowest SDs with < 2.0 .




Figure 12. Scatter plots of time-mean maximum transports vs. their monthly standard deviations (SDs) in σ_2 space for (a) OW, (b) OE, and (c) total. The horizontal dashed lines show the respective mean SDs with shading indicating their ± 1 SDs. The vertical shaded areas indicate $\pm$



1 SD ranges for OSNAP maximum transports. The low- and high-resolution simulations are indicated by dots and stars, respectively. The correlation coefficients (excluding OSNAP) are included in each panel with the bold font indicating statistical significance at the 95% confidence level, calculated using a 2-sided Student's t-test. The regression lines are also shown in panels with statistically significant correlations.

7 Temperature – Salinity Diagrams

We next consider (potential) temperature – salinity (T-S) diagrams for depths > 500 m as depicted in Figs. 13-16. These are based on monthly mean data and densities are in σ_2 space. We note that some of the differences in the spread of T and S values among the panels, especially at mid-depths, are due to the use of each model's native vertical grid. We further note that because we are primarily interested in deep and abyssal ocean properties and for clarify of our analysis, we choose to use a 500-m cutoff depth here to exclude even larger spreads in the upper ocean which result from the models' fine vertical resolutions near the surface. For OW (Figs. 13 and 14), T-S diagrams span 1 – 7 °C in temperature and 34.5 – 35.2 psu in salinity to cover the spread of the simulations. As with the previous figures, we only highlight several key features for brevity. In OSNAP, S is rather uniform around 34.9 psu for depths > ~1500 m. As such, waters with $\sigma_2 > 37.0 \text{ kg m}^{-3}$ attain those densities primarily through cooling from roughly 3 °C to about 1.5 °C. All the simulations also show dense waters with $\sigma_2 > 37 \text{ kg m}^{-3}$ at depth, consistent with OSNAP, but there are differences in their T and S properties. Except for FSU08 and UKMO25, the simulations have a rather thin, line-like, diagonal structure at high density classes. Most simulations show densification through high-to-low T and S. However, in ANU25, CMCC25, FSU08, GEOMAR25, NOC25, and UKMO25 densification occurs primarily through cooling at relatively constant S. In contrast with all the other simulations, and indeed OSNAP, NCAR67 has densification due to both lower T and higher S values. The discontinuous nature of T and S properties in FSU08 and FSU72 reflect the HYCOM's discrete isopycnic coordinates. We also note that FSU27 does not appear to show much density distribution at depth. Among the simulations, UKMO25 attains densities greater than 37.0 kg m^{-3} with the highest S and warmest T values among the simulations likely due to too high S associated with the overflow waters (see below).



At OE (Figs. 15 and 16), we employ wider ranges than at OW for both T and S, i.e., 1 – 12 °C and 34.5 – 36.0 psu, respectively. The observations show a diagonal band extending from high T (~10 °C) and S (~35.4 psu) to lower T (~1.2 °C) and S (~34.9 psu) values with a maximum density of ~37.2 kg m⁻³ around 3000-m depth. Like OW, the deep ocean has a rather uniform S, and densification is due to colder T. This diagonal structure is reproduced by all the simulations. However, all but three have maximum densities of about 37.0 kg m⁻³. In FSU08, NCAR1, and UKMO25, the maximum densities are closer to the observed values, but with slightly saltier and warmer waters. FSU08, FSU72, and NCAR67 show a discontinuous (banded) T and S structure, reflecting their density layers. Again, FSU72 does not seem to capture high-density classes.

High density classes at OE are thought to be associated with the properties of the Denmark Strait (DS) and Faroe Bank Channel (FBC) overflows. Observational estimates of the product water T and S are 2.1 °C and 34.84 psu for DS (Legg et al., 2009; Garton & Sanford, 2003) and 3.3 °C and 35.1 psu for FBC (Legg et al., 2009; Mauritzen et al., 2005). OSNAP shows much colder water masses than the above estimates for DS and FBC overflows. This discrepancy may be due to different observational periods that these estimates are based on. As stated above, among all the simulations, only FSU08, NCAR1, and UKMO25 reach maximum densities closest to the observed values at OE. Among these, NCAR1 uses an overflow parameterization to represent the DS and FBC overflows. In NCAR1, the parameterization (Danabasoglu et al., 2010) appears to maintain distinct properties of the two overflow waters. In UKMO25, the overflow waters are warmer and saltier than observed. This is probably due to a synergy between their hybrid vertical coordinate system and the large-scale T and S biases affecting their North Atlantic subtropical and subpolar gyres. While their hybrid vertical grid is able to significantly reduce numerical diapycnal mixing when simulating the Nordic overflows, this inevitably results in transporting the density-compensated T and S biases of the upper water column down to greater depths (see Bruciaferri et al. 2024 for the details). FSU08 is the only simulation that has minimum temperatures colder than 2 °C, closer to those observed at OE.




Figure 13. Monthly mean potential temperature – salinity diagrams at OW for depths > 500 m from OSNAP and from ANU10, ANU25, ANU1, CMCC06, CMCC25, E3SM27, FSU08,



FSU72, and GEOMAR05. The background density contours are based on σ_2 densities. The
780 colors represent depth.




Figure 14. Same as in Fig. 13, but for GEOMAR25, GFDL25, NCAR10, NCAR1, NCAR67, NOC08, NOC25, NOC1, and UKMO25.




Figure 15. Same as in Fig. 13, but for potential temperature – salinity at OE.




Figure 16. Same as in Fig. 14, but for potential temperature – salinity at OE.



785 8 Temperature, Salinity, and Density Biases

Figure 17 presents the scatter plots of the upper-ocean (0 – 700 m) and section-average T, S, and σ_2 biases against the respective maximum transports in σ_2 space at OW and OE to provide a quantitative assessment of these biases and their relationships with the overturning transports. The biases are computed with respect to the gridded T and S data from the OSNAP observations (Sect. 2.1). At OW, the number of simulations with negative and positive T biases is roughly evenly split with all, but two, within a range of -0.5 and $+0.5$ °C (Fig. 17a). The exceptions are FSU72 and NCAR1 with respective biases of about -1.15 and 0.70 °C. There is no statistically significant relationship between these T biases and the transports. In contrast, the relationship between the S biases and the transports is significant, showing generally larger (smaller) transports with positive (negative) S biases (Fig. 17b). Many of the simulations have biases within a range of -0.1 and $+0.1$ psu, but most (11) have a salty bias. While the largest fresh biases occur in ANU25 and FSU72 with -0.21 psu, the largest salty bias is in NCAR1 with $+0.22$ psu. In density, except for ANU25 and E3SM27, the biases are within a range of about -0.03 and $+0.10$ kg m $^{-3}$, with no significant relationship with the transports (Fig. 17c). ANU25 and E3SM27 show negative density biases of about -0.21 and -0.15 kg m $^{-3}$, respectively. In about half of the simulations, T and S biases compensate each other in their contributions to density. At OE, both T and S biases are strongly correlated with the transports, indicating generally larger transports with warmer and saltier biases (Figs. 17d and e). However, a large majority of the simulations has cold and fresh biases of up to about -0.9 °C and -0.12 psu. NCAR1 has the largest warm ($\sim +1.1$ °C) and salty (0.22 psu) biases among all the models. ANU1 is a close second with warm and salty biases of $+0.6$ °C and $+0.2$ psu, respectively. These biases substantially compensate each other in density, producing no significant relationship between the density biases and the transports (Fig. 17f). Most of the simulations show a positive density bias of $< +0.1$ kg m $^{-3}$. We note that the sign and magnitude of these T, S, and σ_2 biases do not reveal any systematic dependencies on model resolution.

Finally, we show the scatter plots of the spatial-averaged T, S, and σ_2 density biases against the maximum total transport in σ_2 space in Fig. 18, where the biases are with respect to the EN4 dataset (Good et al., 2013). A goal is to briefly explore relationships between the overturning transports and the upper-ocean hydrographic properties considering broader regions. So, the



815 biases are calculated in the upper 700 m separately for the Labrador Sea (60°-45°W and 50°-
 65°N) and Irminger Sea (45°-25°W and 50°-65°N) regions (see Fig. 1 for these regions), chosen
 as two representative areas. In the LS, there are generally larger (smaller) overturning transports
 with positive (negative) T and S biases, with most simulations showing warm and salty biases
 (Figs. 18a and b). While FSU72 has the largest cold bias with -0.75 °C, NCAR1 and UKMO25
 820 are the warmest models with biases of slightly larger than +0.5 °C. In salinity, ANU25 and
 E3SM27 are the freshest models – both with biases of around -0.2 psu, while UKMO25 has the
 largest salty bias with 0.4 psu. These T and S biases partially compensate each other in their
 contributions to density in 13 simulations, but the resulting density biases in the LS are all
 positive except for ANU25 and E3SM27 (Fig. 18c). These positive biases primarily reflect those
 825 of S as density changes are largely determined by S at low temperatures. No statistically
 significant relationship exists between the LS density biases and the transports for the LS. In the
 Irminger Sea, the simulations show generally cold biases, with an even split of fresh and salty
 biases among the simulations (Figs. 18d and e). There is a tendency towards larger (smaller)
 overturning transports with positive (negative) T and S biases, but this relationship is statistically
 830 significant only for S. ANU10 and FSU72 have the largest cold and warm biases with about -
 0.85 °C and +0.65 °C, respectively. E3SM27 and UKMO25 bracket the S biases on opposite
 sides with bias magnitudes of ~0.2 psu. As in the LS, T and S biases partially compensate each
 other in their contributions to density in 13 simulations in the Irminger Sea (Fig. 18f). The
 resulting density biases are largely positive, due either to negative T or positive S biases. Again,
 835 there is no statistically significant relationship between the Irminger Sea density biases and the
 transports. As in Fig. 17, the sign and magnitude of these broader region T, S, and σ_2 biases do
 not show any systematic dependencies on model resolution.




Figure 17. Scatter plots of the upper-ocean (0-700 m) and section average, time-mean (a, d) temperature, (b, e) salinity, and (c, f) σ_2 density biases against the time-mean maximum transports in σ_2 space at OW (top panels) and OE (bottom panels). The black dots in each panel show the OSNAP transports plotted against no bias. The low- and high-resolution simulations are indicated by dots and stars, respectively. The correlation coefficients (excluding OSNAP) are included in each panel with the bold font indicating statistical significance at the 95% confidence level, calculated using a 2-sided Student's t-test. The regression lines are also shown in panels with statistically significant correlations.




Figure 18. Scatter plots of the Labrador Sea (LS; upper panels) and the Irminger Sea (IR; lower panels) upper-ocean (0–700 m average), time-mean (a, d) temperature, (b, e) salinity, and (c, f) σ_2 density biases against the time-mean maximum total transport in σ_2 space. The Labrador and Irminger Sea regions are bounded by 60°–45°W and 50°–65°N, and 45°–25°W and 50°–65°N, respectively (the dark and light purple boxes in Fig. 1, respectively). The low- and high-resolution simulations are indicated by dots and stars, respectively. The correlation coefficients (excluding OSNAP) are included in each panel with the bold font indicating statistical significance at the 95% confidence level, calculated using a 2-sided Student’s t-test. The regression lines are also shown in panels with statistically significant correlations.

9 Summary and Conclusions

We have presented a comparison of simulated and observed transports and related properties, including hydrographic characteristics, across the OSNAP sections for the ~2014–2022 observational period. Our effort aims to provide a benchmark for the modeling community for evaluations of their simulations at OSNAP, considering both depth and density space transports. There are 9 participating groups from around the world providing 18 simulations with 6 different ocean models. These FOSI simulations use a common set of atmospheric datasets, largely following the OMIP protocol. The horizontal resolutions of the simulations range from coarse nominal 1° to eddy-resolving resolutions of 0.1°–0.05°. In the vertical, many of the simulations



use depth / level coordinates, but a few simulations employ hybrid coordinates. The number of
 865 vertical levels / layers varies from 36 to 98, with most models using between 46 and 75.

Our analysis clearly demonstrates that the simulated transports are, in general, in broad
 agreement with those of observational estimates. Specifically, the transports are larger at OE than
 at OW with the total transports dominated by those of OE. There are, of course, many differences
 between the simulations and observations as well as among the individual simulations in details
 870 of the transport properties. These include differences (or biases) in the maximum transport
 densities, transport directions and magnitudes in a given density class or depth, phase of the
 seasonal cycle of the transports, and T and S properties. A notable difference from OSNAP
 occurs at OE where a large majority of the simulations have positive transports with negative
 slopes for densities $< \sim 27.3 \text{ kg m}^{-3}$, contrasting with observations which show a much steeper
 875 negative density slope and nearly no net flow for densities $< 27.2 \text{ kg m}^{-3}$. There are also
 seemingly small differences in high density classes at OW. While these high-density transports
 are themselves relatively small, they can play an outsized role in downstream low-frequency
 AMOC variability (Yeager et al., 2021). NCAR1 seems to be an outlier in its representation of
 transports at OW with its rather large maximum transport magnitude.

880 Analyzing overturning circulations in both depth and density space together provides a more
 complete picture of overturning properties and features. Our preferred approach uses binned
 southward and northward transports in Sv as a function of depth and density as a starting point,
 similar to that used in Zhang & Thomas (2021). These depictions provide a clear view of how
 these transports are distributed across both depth and density space. They can then be summed or
 885 aggregated either in depth or density space to produce the velocity and the more familiar
 overturning transport profiles. In both the simulations and observations, northward and
 southward flows substantially cancel each other, producing much smaller total velocities. Such
 cancellations tend to be much more prominent (larger) in depth space than in density space. As
 discussed in Zhang & Thomas (2021), these cancellation differences in depth- and density-space
 890 depend on density differences of horizontally circulating water masses, i.e., isopycnal slopes. In
 general, the simulations seem to capture the observed transport features better at OE than OW.
 Not surprisingly, the total velocity and transport profiles tend to agree better between the
 simulations and observations due to summations involved. However, actual binned transports
 show large differences in their depth and density ranges among the simulations and with the



895 observations, i.e., similar-looking northward or southward transports may occur at differing depths and / or densities.

The simulations show T and S biases of both signs in the upper ocean at both OW and OE. Statistically significant relationships exist between overturning transports and S biases at OW and transports and both T and S biases at OE. Specifically, larger (smaller) transports are
 900 associated with positive (negative) T and S biases. In about half of the simulations, T and S biases compensate each other in their contributions to density, and there is no evident relationship between density biases and the corresponding section transports. When broader regions are considered for such bias calculations, i.e., the LS and the Irminger Sea rather than just the OSNAP sections, we again find larger (smaller) overturning transports with positive
 905 (negative) T and S biases, but no such evident relationship with density biases. The LS typically shows warm and salty biases. In contrast, the simulations show mostly cold biases in the Irminger Sea region, with roughly an even split of fresh and salty biases among the simulations. While the LS density biases are largely determined by those of salinity, both temperature and salinity biases contribute to the density biases in the Irminger Sea.

910 The transport profiles from high-resolution simulations show better agreement with OSNAP than those of the low-resolution simulations, particularly in the high-density classes. There are also significant improvements in the representation of the seasonal cycle of the transports as well as the transports at OW with the high-resolution version of the NCAR model compared to its low-resolution counterpart. However, for some of the other metrics considered in our analysis, the
 915 high-resolution simulations face similar challenges as the low-resolution ones, with no obvious improvements. These issues may be related to different levels of bias compensations or to differences in the representation of certain processes, such as overflows and (sub)meso-scale eddies, in both low- and high-resolution simulations.

An important finding is that when transports are calculated in σ_2 space, rather than the σ_0 space
 920 used in observations, transports at both OW and OE increase. Specifically, in observations, the OE transport increases by 1.1 Sv from 12.9 to 14.0 Sv, while the OW transport increases by 1.5 Sv from 3.8 to 5.3 Sv, representing a reduction of the OE / OW transport ratio from 3.4 to 2.6. On average, model simulations show increases of 2.47 and 1.19 Sv for OW and OE, respectively. A comparative measure of these changes, also taking the changes in the OE transports into



925 consideration, indicates that the relative contribution of OW transports increase by about ~35%
in both observations and simulations. We think that using potential densities referenced to a
depth of 2000 m provides a more accurate depiction of deep transports relevant for our analysis.
Therefore, we recommend employing σ_2 densities going forward.

When transports are reported in density space, it is important to provide details of exactly how
930 they are obtained for apples-to-apples comparisons. Specifically, the density space overturning
timeseries usually use monthly maximum transports which occur at different densities for each
month. Then, these monthly maximum transports are averaged to produce, say, annual-mean
values. Due to the inherent nonlinearity of this process, when annual-mean transports are
calculated based on monthly mean transport profiles, the latter maximum transports differ
935 modestly from those obtained with the former approach. This difference results from substantial
changes of the density at which the maximum transport occurs each month. In contrast, such a
problem with the nonlinearity of the overturning strengths is much more muted in depth space
calculations as the depth of the maximum transport does not change substantially from month to
month (see Fig. S1 of Lozier et al., 2019). Transports computed from averaged profiles will be
940 always smaller than those derived from averaging the monthly maximum transports. Indeed,
these differences can be ~2 Sv for the OE and total transports in observations (~15 Sv vs. ~13
Sv) as reported in Lozier et al. (2019).

The OSNAP observations have been incredibly valuable in providing separate overturning
transport estimates for the eastern subpolar gyre and the LS sides. These datasets have been used
945 in evaluation of model simulations as described in this manuscript. Now available for the 8-year
period from August 2014 to July 2022, the OSNAP observations provide a comprehensive view
of the respective roles of the eastern and western sections on inter-annual (or sub-decadal) time
scales. While the model simulations are in general agreement with the OSNAP observations
regarding the respective transport magnitudes during this observational period, studies that
950 attribute a prominent role for the LS primarily focus on decadal-to-multi-decadal time scales,
which cannot be evaluated against observations yet. Therefore, it is important to continue the
OSNAP observational program so that they can provide future guidance on longer time scales.



Table 1. Summary of the ocean and sea-ice models in alphabetical order according to the participating group name (first column). The table includes, in order, the label used in this study for the listed configuration and resolution; the name of the combined ocean – sea-ice configuration (if any); the ocean model name and its version number; the sea-ice model name and its version number; vertical coordinate and number of layers / levels in parentheses; and the nominal horizontal resolution in degrees.

Group	Label	Configuration	Ocean Model	Sea-ice Model	Vertical Grid (number of levels)	Nominal Horizontal Resolution (°)
ANU	ANU10	ACCESS-OM2-01	MOM 5.1	CICE 5.1.2	z* (75)	0.1
	ANU25	ACCESS-OM2-025	MOM 5.1	CICE 5.1.2	z* (50)	0.25
	ANU1	ACCESS-OM2	MOM 5.1	CICE 5.1.2	z* (50)	1
CMCC	CMCC06	GLOB16	NEMO 3.6	LIM 2	z (98)	0.0625
	CMCC25	ORCA025	NEMO 3.6	CICE 4.1	z (50)	0.25
FSU	FSU08		HYCOM	CICE 4	hybrid (36)	0.08
	FSU72		HYCOM	CICE 4	hybrid (41)	0.72
GEOMAR	GEOMAR05	VIKING20X	NEMO 3.6	LIM 2	z (46)	0.05 / 0.25 ^a
	GEOMAR25	ORCA025	NEMO 3.6	LIM 2	z (46)	0.25
GFDL	GFDL25	OM4	MOM 6	SIS 2	hybrid (75)	0.25
LANL	E3SM27		MPAS-Ocean	MPAS-SeaIce	z* (64)	~0.27 (~30 km) ^b
NCAR	NCAR10		POP 2	CICE 4	z (62)	0.1
	NCAR1		POP 2	CICE 4	z (60)	1
	NCAR67		MOM 6	CICE 6	hybrid (75)	0.67
NOC	NOC08	GO8p7	NEMO 4.0.4	SI3	z* (75)	0.08



	NOC25	GO8p7	NEMO 4.0.4	SI3	z^* (75)	0.25
	NOC1	GO8p7	NEMO 4.0.4	SI3	z^* (75)	1
UKMO	UKMO25	GO8p7	NEMO 4.0.4	SI3	z^* (75) ^c	0.25

960 ^a Uses a 0.05° Atlantic grid (33.5°S to ~65°N), two-way nested within a global 0.25° grid (ORCA025).

^b Uses an unstructured grid with a nominal horizontal resolution of 30 km which is roughly 0.27° at the Equator.

965 ^c Uses embedded hybrid terrain-following layers only in the Nordic overflow regions via localized multi-envelope generalized vertical coordinates.



Table 2. Summary of the surface freshwater / salt fluxes and sea surface salinity (SSS) restoring choices for each model configuration (first column). The salt vs. water column indicates the type of surface fluxes used for hydrological forcing with water and salt denoting real freshwater and virtual salt fluxes, respectively. The SSS restoring time scales are given in days over a 50 m length scale. The NEMO-based models convert salinity restoring to a freshwater flux (denoted as fw in the column). The other groups apply salinity restoring as a salt flux. The region column indicates the region over which the salinity restoring is used. An asterisk denotes that a limit of 0.5 psu is imposed to constrain the magnitude of mismatch between the model and observed SSS. A superscript + indicates that surface restoration is capped to not exceed 4 mm day⁻¹ in magnitude. The under sea ice column shows whether restoring is applied under sea-ice covered areas. The remove global-mean restoring column indicates whether the model subtracts the global mean of restoring fluxes. Finally, the normalize freshwater fluxes column refers to whether some sort of normalization to the sum of precipitation, evaporation, and runoff fluxes is applied to reduce drift.

Label	Freshwater Fluxes via Virtual Salt vs. Water	SSS Restoring Time Scale (day)	SSS Restoring Region	SSS Restoring Under Sea Ice	Remove Global-Mean SSS Restoring	Normalize Freshwater Fluxes
ANU10 ^a	Water	455	Global*	Y	Y	Y
ANU25 ^a	Water	455	Global*	Y	Y	Y
ANU1 ^a	Water	455	Global*	Y	Y	Y
CMCC06	Water	365 (fw)	Global	N	N	Y
CMCC25	Water	182.5 (fw)	Global	N	N	Y
E3SM27	Water	365	Global	Y	Y	N
FSU08 ^b	Salt	60	Global*	Y	Y	N
FSU72 ^b	Salt	60	Global*	Y	Y	N
GEOMAR05	Water	365	Global*, ^c	N	N	Y
GEOMAR25	Water	365	Global*, ^c	N	N	Y
GFDL25	Salt	300	Global	Y	Y	Y
NCAR10	Salt	365	Global	Y	Y	Y
NCAR1	Salt	365	Global	Y	Y	Y
NCAR67	Salt	300	Global	Y	Y	Y
NOC08	Water	1500 (fw)	Global ⁺	Y	N	N
NOC25	Water	1500 (fw)	Global ⁺	Y	N	N
NOC1	Water	1500 (fw)	Global ⁺	Y	N	N
UKMO25	Water	1500 (fw)	Global ⁺	Y	N	N



^a The restoring dataset is based on the 0.25° World Ocean Atlas 2013 v2 decav monthly climatology, filled into land via a conjugate gradient method. The 0-m and 10-m data are then averaged, horizontally smoothed, and smoothly interpolated onto the model grid.

^b The Generalized Digital Environmental Model (GDEM4) monthly climatology is used as the
985 restoring dataset.

^c Salinity restoring is suppressed where runoff is applied as well as within an 80-km wide band around Greenland to not interfere with enhanced freshwater input from melting ice sheets.



Appendix A: Model descriptions

990 A1 Australian National University (ANU)

ACCESS-OM2, ACCESS-OM2-025, and ACCESS-OM2-01 are global coupled ocean – sea-ice model configurations at nominal horizontal grid spacings of 1° , 0.25° , and 0.1° , respectively, developed by the Consortium for Ocean - Sea Ice Modelling in Australia. The configurations are respectively labeled as ANU1, ANU25, and ANU10 in this manuscript. They are based on those
 995 described by Kiss et al. (2020) but updated with several improvements described below and in the Supporting Information of Solodoch et al. (2022). The two coarser resolutions also comprise the ocean and sea-ice components for Australia’s ACCESS-CM2 (Bi et al., 2020) and ACCESS-CM2-025 (Huneke et al., 2025) coupled climate models.

The ocean model is MOM5.1 (Griffies, 2012) and the sea-ice model is CICE5.1.2 (Hunke et al.,
 1000 2015); these are coupled every ocean baroclinic timestep (equal to the sea ice thermodynamics timestep) via OASIS3-MCT v.2.0 (Valcke et al., 2013). At each resolution, the configurations use a common tripolar (Murray, 1996) horizontal B-grid for MOM and CICE. Grid lines are aligned with latitude and longitude south of 65°N , with Mercator spacing between 65°N and 65°S in ANU25 and ANU10, and they have a more complicated latitudinal dependence with a meridional
 1005 refinement to $1/3^\circ$ near the Equator in ANU1 (Bi et al., 2013). The ocean vertical coordinate is z^* , with a smoothly varying spacing chosen to optimize the resolution of baroclinic modes (Stewart et al., 2017). There are 50 levels ranging from 2.3-m thick at the surface to 219.6 m in the abyss in ANU1 and ANU25, and 75 levels ranging from 1.1 to 198.4-m thick in ANU10. Bathymetry is represented with partial cells (Adcroft et al., 1997; Pacanowski & Gnanadesikan,
 1010 1998). The topography at all three resolutions is based on GEBCO 2014 v20150318 30 arc-second data (replacing the legacy topography files used at 1° and 0.25° in Kiss et al., 2020), with hand-edits in ANU1 to improve the flow in some straits, e.g., the Denmark Strait, Iceland-Faroe Strait, the Strait of Gibraltar, the Lombok, Ombai, Timor, Makassar Straits, the Caribbean, and the Drake Passage.

1015 The configurations are designed to be as similar as possible at the three resolutions, but there are some unavoidable differences due to the differing resolution of eddies. The Gent and McWilliams (GM; 1990) scheme is used in ANU25 and ANU1 to parameterize advective transports due to unresolved eddies via a skew diffusive implementation (Griffies, 1998). The



GM diffusivity has a spatial structure determined according to the baroclinic zone setting of
 1020 Griffies et al. (2005) and has a maximum value of $200 \text{ m}^2 \text{ s}^{-1}$ in ANU25 and $600 \text{ m}^2 \text{ s}^{-1}$ in ANU1.
 A neutral diffusivity for tracers (Griffies et al., 1998) is included with a horizontal scaling
 according to Hallberg (2013) with a maximum value of $200 \text{ m}^2 \text{ s}^{-1}$ in ANU25 and a constant
 value of $600 \text{ m}^2 \text{ s}^{-1}$ in ANU1. Neither of these parameterizations are active in ANU10. The K-
 profile parameterization (KPP; Large et al., 1994), Lee et al. (2006) barotropic tidal drag and
 1025 mixing, and the Simmons et al. (2004) bottom intensified mixing schemes are used, with
 background vertical viscosities of $10^{-4} \text{ m}^2 \text{ s}^{-1}$ in all configurations and background vertical
 diffusivities following Jochum (2009) for ANU1, no background vertical diffusivity in ANU25,
 and a constant value of $10^{-6} \text{ m}^2 \text{ s}^{-1}$ in ANU10. ANU1 also uses downslope mixing (Beckmann
 and Doscher, 1997; Doscher and Beckmann, 2000; Campin and Goosse, 1999). The Fox-Kemper
 1030 et al. (2008) parameterization of submesoscale eddy-driven mixed-layer restratification is used in
 all configurations.

In all three configurations, the ocean equations of state and freezing temperature are from Jackett
 et al. (2006). The prognostic salinity and temperature variables are practical salinity and
 conservative temperature (converted from potential temperature initial conditions via TEOS-10).
 1035 Sea surface salinity is restored to World Ocean Atlas 2013 v2 (WOA13) monthly climatology
 with a piston velocity of 50 m over 455 days (including under sea ice) in all 3 simulations. The
 ocean albedo is latitude dependent (Large and Yeager, 2009). The CICE configuration uses 5
 thickness categories, with four ice layers and one snow layer for the thermodynamics, and an ice
 salinity (for the purposes of coupling) of 5 psu. The wind stress calculation uses relative velocity
 1040 over both the ocean and sea ice.

Runs at all three resolutions started from initial temperature and salinity from the WOA13
 January-mean climatology. ANU1, ANU25, and ANU10 were forced by 6, 6, and 4 consecutive
 61-year cycles, respectively, of the JRA55-do v1.4.0 atmospheric datasets for the 1958-2018
 period. The final cycle was then extended through 2019 using JRA55-do v1.5.0, and then
 1045 through 2020-2022 inclusive using JRA55-do v1.5.0.1. This last cycle constitutes the data used
 in this study. Output from ANU1 and ANU25 were contributed to CMIP6 through the OMIP2
 experiments (see Mackallah et al., 2022).



The performance of ACCESS-OM2 at different resolutions is assessed and compared to other models and / or to available observations for the Arctic Ocean in Wang et al. (2024); for mixed layer depths in Treguier et al. (2023); for the deep Atlantic multidecadal variability in Yang et al. (2024); and (for the older Kiss et al. (2020) configurations) for the AMOC in Hirschi et al. (2020).

A2 Centro Euro-Mediterraneo sui Cambiamenti Climatici (CMCC)

The CMCC foundation provided the global ocean – sea-ice model configurations, both built on the NEMO modeling framework, at nominal horizontal grid spacings of 0.25° (ORCA025) and 0.0625° (GLOB16). The configurations are respectively labeled as CMCC25 and CMCC06 in this manuscript. Details of the latter configuration and an evaluation of its simulation are given in Iovino et al. (2023).

CMCC06 is a global, eddying configuration of the ocean and sea-ice system. The model is based on its first implementation documented in Iovino et al. (2016), where the ocean component is upgraded from version 3.4 to version 3.6 stable (Madec and the NEMO Team, 2016). Its high computational cost does not permit long evaluation simulations and, therefore, it is simply set up using our best practices based on a forecasting application (Cipollone et al., 2020; Masina et al., 2021).

This eddy-rich configuration makes use of a non-uniform tripolar grid with a nominal $1/16^\circ$ horizontal resolution (6.9 km at the Equator, reducing poleward). The grid consists of an isotropic Mercator grid between 60°S and 20°N and a non-geographic quasi-isotropic grid north of 20°N . The minimum grid spacing is ~ 2 km around Victoria Island, and the meridional scale factor is fixed at 3 km south of 60°S . Ocean and sea ice are on the same horizontal grid. The vertical coordinate system is based on fixed depth levels and consists of 98 vertical levels with a grid spacing increasing from approximately 1 m near the surface to 160 m in the deep ocean.

The ocean component is a finite-difference, hydrostatic, primitive-equation general circulation model, with a linearized free sea surface, a free-slip lateral friction condition, and an Arakawa C-grid. A biharmonic viscosity scheme with a coefficient of $-0.5 \times 10^{10} \text{ m}^4 \text{ s}^{-1}$ is used in the horizontal directions. Lateral tracer diffusion, set at $80 \text{ m}^2 \text{ s}^{-1}$, is along isoneutral surfaces using a Laplacian mixing. Tracer advection uses a total variance dissipation (TVD) scheme (Zalesak,



1979). Vertical mixing is achieved using a turbulent kinetic energy (TKE) closure scheme (Blanke & Delecluse, 1993). Background coefficients of vertical diffusion and viscosity are 1.2×10^{-5} and $1.2 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$, respectively. Vertical eddy mixing of both momentum and tracers is enhanced in case of static instability. The turbulent closure model does not apply any specific modifications in ice-covered regions. Bottom friction is quadratic, and a diffusive bottom boundary layer scheme is included. All configurations use the EOS80 equation of state (Fofonoff & Millard, 1983), with potential temperature and practical salinity as prognostic state variables. The ocean component is coupled to the Louvain-la-Neuve sea Ice Model version 2 (LIM2; Timmermann et al., 2005) for its smaller computational cost compared to the more complex LIM3 code (Rousset et al., 2015; Uotila et al., 2017) available in NEMOv3.6. LIM2 is integrated as an internal module in NEMO. It uses a three-layer model for the vertical heat conduction within snow and ice, featuring a single sea ice category. The ice dynamics are calculated according to external forcing from wind stress, ocean stress, and sea surface tilt and internal ice stresses using a C-grid elastic–viscous–plastic rheology (Bouillon et al., 2013).

The eddy-permitting ORCA025 configuration used here is the ocean and sea-ice components of the CMCC coupled climate model (CMCC-CM2; Cherchi et al., 2019) and Earth system model (CMCC-ESM2; Lovato et al., 2022). This framework is based on the Community Earth System Model (CESMv1.2), in which we replaced the original ocean component with NEMOv3.6 (Fogli & Iovino, 2014). The ocean component is coupled to the Community Ice Code CICEv4.1 (Hunke & Lipscomb, 2010). CMCC25 has a nominal resolution of 0.25° , with 50 vertical levels, ranging from 1 to 400 m. The ORCA025 physical core as implemented for this OMIP simulation is similar to the lower resolution configuration described in Tsujino et al. (2020), except for resolution-dependent features, such as the eddy-induced tracer advection term which is not used in ORCA025. This configuration uses a biharmonic horizontal viscosity of $1.8 \times 10^{11} \text{ m}^4 \text{ s}^{-1}$, a Laplacian tracer diffusivity of $300 \text{ m}^2 \text{ s}^{-1}$, and a background vertical diffusivity and viscosity of 1.2×10^{-5} and $1.2 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$, respectively. The sea-ice model includes energy-conserving thermodynamics (Bitz & Lipscomb, 1999), multi-category ice thickness (Bitz et al., 2001) with five thickness categories, and elastic–viscous–plastic ice dynamics (Hunke & Dukowicz, 1997). It is solved on the Arakawa B-grid, with the tracer points aligned with the ocean grid. The coupling interface between NEMO and CICE is described in Cherchi et al. (2019) and references therein (Fogli & Iovino, 2014). To be able to attribute the main differences among model



configurations mainly to the increase in ocean resolution in the horizontal and vertical grids, the two configurations employ, as much as possible, the same numerical schemes and
 1110 parameterizations, except grid-spacing-dependent parameters.

Sea surface salinity is restored to the WOA13 v2 monthly climatology. Salinity restoring is applied globally via an equivalent-surface freshwater flux using restoring time scales of 365 and 182.5 days over 50 m in CMCC25 and CMCC06, respectively. There is no salinity restoring under sea-ice-covered areas. The two sea-ice models used in the two systems employ different
 1115 bulk ice salinity values that affect the salt release from the sea ice to the ocean. In CICE, a reference sea-ice salinity value of 4 psu is used for computing the ice–ocean exchanges. In our version of LIM2, the freshwater (salinity) fluxes between the ice and the ocean assume a constant salinity of 6 psu. Over a sea ice formation and melt cycle, this produces stratification differences among runs and might have an impact on the large-scale ocean circulation. In wind
 1120 stress calculations, relative winds are used over both ocean and sea ice.

Liquid runoff is deposited along the coast and distributed in the upper 20 m in CMCC25 and at the ocean surface in CMCC06, with no specific enhancement of the mixing in either case. The runoff interpolation in both configurations makes use of a globally conserving method, which also spreads the runoff along the coast, to compute offline remapping weights.

Following the OMIP protocol, the ocean is initially at rest, with a sea level of 0 m and with temperature and salinity from the World Ocean Atlas 2013 v2 (WOA13; Locarnini et al., 2013; Zweng et al., 2013) decav product (averaged from 1955–2012) interpolated on a 0.25° grid. For CMCC25, the initial sea-ice properties are taken from spin-up experiments, while initial ice concentration and thickness for CMCC06 are fixed to 100% with thicknesses of ~3 m north of
 1130 70°N and ~1 m south of 60°S.

CMCC25 was integrated for six 61-year cycles (1958-2018) of JRA55-do, while CMCC06 was integrated over a single cycle. Only the first JRA55-do cycle is used for both simulations in this paper which were extended to 2022.

A3 Florida State University (FSU)

FSU-HYCOM is a global configuration of the HYbrid Coordinate Ocean Model (HYCOM)
 1135 (Bleck, 2002; Chassignet et al., 2003). The vertical coordinate is isopycnic in the stratified open



ocean and makes a dynamically smooth and time-dependent transition to terrain following in the shallow coastal regions and to fixed pressure levels in the surface mixed layer and unstratified seas. The sea-ice component uses version 4 of the Community Ice CodE (CICE4; Hunke & Lipscomb, 2010). Two simulations, one with a coarse resolution of 0.72° and the other with an eddy resolution of 0.08° , are performed. They are referred to as FSU72 and FSU08, respectively, in the manuscript. The configurations and general features of these simulations are described in Chassignet et al. (2020). An in-depth evaluation of the Antarctic Circumpolar Current (ACC) and the AMOC in the South Atlantic Ocean are provided in Xu et al. (2020; 2022).

In FSU72, the resolution is refined to 0.33° near the Equator, and a vertical grid of 41 hybrid coordinate layers is used. Vertical mixing uses the KPP (Large et al., 1994) with a background diffusivity of $3 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$. For horizontal mixing, the simulation uses a combination of a Laplacian diffusive velocity of 3 cm s^{-1} and a biharmonic diffusive velocity of 5 cm s^{-1} for momentum and a Laplacian diffusive velocity of 3 cm s^{-1} for tracers. The model also includes interface height smoothing corresponding to the GM isopycnal mixing parameterization, using a biharmonic operator with a diffusive velocity of 2 cm s^{-1} everywhere, except in the North Pacific and North Atlantic where a Laplacian operator with a velocity scale of 1 cm s^{-1} is used.

FSU08 has 36 hybrid layers in the vertical. This vertical layer set up was retained to enable comparisons with previous simulations performed with other atmospheric forcing datasets. While the vertical resolution in both FSU72 and FSU08 is lower than recommended by Stewart et al. (2017) for z-coordinate models, the vertical resolution is adequate to represent the first five baroclinic modes in the mid-latitudes and the large-scale circulation of key water masses in the Atlantic Ocean (Xu et al., 2023). As in the coarse resolution counterpart, vertical mixing in the eddy simulation uses the KPP with a background diffusivity of $3 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$. For horizontal mixing, the simulation uses a combination of a Laplacian viscosity of $20 \text{ m}^2 \text{ s}^{-1}$ and a biharmonic diffusive velocity of 1 cm s^{-1} for momentum, a Laplacian diffusive velocity of 0.5 cm s^{-1} for tracers, and a biharmonic diffusive velocity of 1.5 cm s^{-1} for layer interfaces.

Both simulations are initialized using potential temperature and salinity from the Generalized Digital Environmental Model (GDEM4; Teague et al., 1990; Carnes, 2009) and are integrated using the JRA55-do datasets from 1958 to 2022 for one cycle. The Large & Yeager (2009) bulk



formulation is used for turbulent air–sea fluxes. Wind stress is calculated without surface ocean currents (i.e., using absolute winds). No restoration is applied on the sea surface temperature. The surface salinity is restored to the monthly GDEM4 climatology over the global domain with a piston velocity of 50 m over 60 days. In addition, the salinity flux at each time step is adjusted to ensure a zero net global-mean flux.

A4 GEOMAR

The VIKING20X model configuration is based on NEMO version 3.6. It consists of a 0.05° Atlantic nest extending from 33.5°S to about 65°N, which is embedded using grid refinement (AGRIF; Debreu et al., 2008) into a global curvilinear 0.25°-grid ORCA025 configuration with 46 vertical levels. The model discretizes the ocean variables on an Arakawa C-grid and the sea-ice component LIM2 (Fichefet & Morales Maqueda, 1997) on a B-grid. A parallel ORCA025 configuration without grid refinement in the Atlantic Basin but otherwise with identical settings as those of the coarse grid in VIKING20X, accompanies the eddy-rich simulations. In this manuscript, these simulations are referred to as GEOMAR05 and GEOMAR25, respectively.

Numerics include a momentum advection scheme in vector form which conserves both energy and enstrophy (Arakawa and Hsu, 1990), leading to a good representation of the large-scale, horizontal flow field (Barnier et al., 2006). Tracer advection is formulated as a two-step flux corrected transport, total variance dissipation scheme (Zalesak, 1979), ensuring positive-definite values. Momentum diffusion is given along geopotential surfaces in a bi-Laplacian form with a viscosity of $15 \times 10^{10} \text{ m}^4 \text{ s}^{-1}$. Tracer diffusion is along iso-neutral surfaces in Laplacian form with an eddy diffusivity of $300 \text{ m}^2 \text{ s}^{-1}$. Diffusion parameters are adjusted for the nest grid to take advantage of the increased resolution. As such, the Laplacian parameter for tracers is $60 \text{ m}^2 \text{ s}^{-1}$ and the bi-Laplacian parameter for momentum is $6 \times 10^9 \text{ m}^4 \text{ s}^{-1}$. In both simulations, the background vertical eddy diffusivity and viscosity are set to $1 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ and $1 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$, respectively.

Both simulations were run for 6 forcing cycles for the 1958-2019 period. However, at the end of each cycle, the simulations were extended to 2023. This study uses the solutions from the fifth cycles, noting that the first cycle is described in Biastoch et al. (2021) where it is referred to as VIKING20X-JRA-OMIP. A surface salinity restoring with a piston velocity of 50 m over 365 days is applied together with a freshwater budget correction that balances the global surface



freshwater fluxes to zero. Specifically, the freshwater budget is calculated on the parent grid, and it is then applied to both the parent and nest grids individually, in proportion to their respective restoring fluxes. Runoff is provided by JRA55-do and enters the ocean in the upper-most grid cell in a 0.25° band along the coast with a slightly wider distribution for the largest rivers. Salinity restoring is suppressed where runoff is applied as well as within an 80-km wide band around Greenland to not interfere with enhanced freshwater input from melting ice sheets.

Because the grid resolution is eddy-rich through the whole Atlantic, VIKING20X simulates well the large-scale circulation including realistic representations of western boundary currents, the path of the North Atlantic Current into the Northwest Corner, convection in the subpolar North Atlantic, pathways of deepwater, and the AMOC (Biaostoch et al., 2021; Fox et al., 2022; R  hs et al., 2021; Fr  hle et al., 2022; Petit et al., 2023). B  ning et al. (2023) used VIKING20X and corresponding sensitivity experiments to link Labrador Sea convection to the decadal evolution of the AMOC.

A5 Geophysical Fluid Dynamics Laboratory (GFDL)

The GFDL simulation is based on the OM4 configuration using the Modular Ocean Model version 6 (MOM6) and the Sea Ice Simulator version 2 (SIS2), as documented in Adcroft et al. (2019). MOM6 and SIS2 have identical horizontal grids (Arakawa C-grid) and employ a tri-polar grid following Murray (1996). A Mercator grid spans from about 78°S to 65°N and matches the tri-polar grid at $\sim 65^\circ\text{N}$. MOM6 uses a vertical Lagrangian-remap algorithm (Hirt et al., 1997; Bleck, 2002) to generate a hybrid depth-isopycnal vertical coordinate. The OM4 configuration has a nominal 0.25° horizontal resolution and a hybrid vertical grid with 75 degrees of freedom. The upper ocean is in the depth coordinate, from a 2-m vertical grid interval near the surface to ~ 20 m at 200-m depth. The isopycnal coordinate (referenced to 2,000 dbar) is employed in the region from the upper thermocline to the abyssal ocean. The ocean bathymetry is derived from a combination of the GEBCO 30-s gridded topography (Weatherall et al., 2015) and the International Bathymetric Chart of the Arctic Ocean (Jakobsson et al., 2012). In the manuscript, this contribution is referred to as GFDL25.

No diffusive mesoscale eddy closure is included. The model uses the energetically constrained parameterization of the surface boundary layer (ePBL) of Reichl & Hallberg (2018) and the submesoscale mixed-layer restratification parameterization of Fox-Kemper et al. (2011) with a



submesoscale frontal length scale of 500 m. A biharmonic viscosity is employed, which is the maximum of the dynamic biharmonic viscosity (coefficient = 0.06) (Griffies & Hallberg, 2000) and the static biharmonic viscosity, $u_4 \Delta^3$, with $u_4 = 0.01 \text{ m s}^{-1}$. Here, Δ is the local grid spacing.

1230 SIS2 has five thickness categories and uses four sea ice layers and one snow layer for the vertical thermodynamics. A radiative transfer scheme is included to simulate the vertical profile of shortwave absorption. The OM4 configuration is also used as the ocean – sea-ice component of GFDL’s 4th generation fully coupled climate model CM4 (Held et al., 2019).

The sea surface salinity is restored with a piston velocity of 50 m over 300 days. The simulation
 1235 was run for 6 forcing cycles. The first 3 cycles use 60 years each (from 1958 to 2017) and the last 3 cycles are for 61 years each (from 1958 to 2018). The general evaluation of OMIP simulations across multiple ocean – sea-ice models including GFDL OM4 has been documented in Tsujino et al. (2020).

A6 Los Alamos National Laboratory (LANL)

1240 The Energy Exascale Earth System Model (E3SM) version 3 (E3SMv3) builds on previous versions of E3SM (E3SMv2.1, Smith et al., 2024; E3SMv2; Golaz et al., 2022). The ocean component, MPAS-Ocean, solves the hydrostatic, incompressible equations under the Boussinesq approximation on a staggered Arakawa C-grid, using a two-step Adams-Bashford time-stepping. The vertical grid is structured and uses an arbitrary Lagrangian-Eulerian method
 1245 with several choices of vertical coordinates (Petersen et al., 2015). The production simulations use a free surface z^* coordinate and 64 vertical levels with thicknesses from 10 m at the surface to over 240 m in the deep ocean with partial bottom cells. The horizontal mesh is unstructured. The new version 3 MPAS-Ocean horizontal mesh used in this paper is an icosahedral mesh of nominal horizontal resolution of 30 km and includes sub-ice-shelf cavity geometry around
 1250 Antarctica. The ocean topography is based on GEBCO 2023, updated with BedMachine Antarctica v3 (Morlighem, 2022) on the Antarctic shelves. Because the grid is unstructured and the nominal 30-km horizontal resolution is approximately 0.27° at the Equator, this contribution is referred to as E3SM27 in this manuscript.

The ocean parameterizations of mesoscale eddy effects include isoneutral diffusion following
 1255 Redi (1982), tapered near the surface and slope-limited, with a diffusivity of $400 \text{ m}^2 \text{ s}^{-1}$ and a



modified GM scheme following Ferrari et al. (2010) using a thickness diffusivity of $600 \text{ m}^2 \text{ s}^{-1}$. These coefficients are linearly reduced in regions where the resolution lies between 30 and 20 km, and reach zero for resolutions finer than 20 km. Restratification due to mixed layer eddies is represented by a submesoscale scheme following Fox-Kemper et al. (2011), with a minimum
 1260 frontal width of 1 km and a 2-day mixing time scale. The mixed layer is represented using the KPP scheme with a critical bulk Richardson number of 0.25. A constant background viscosity and diffusivity of $1.0 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$ is applied. The prognostic volume equation includes surface fluxes from the coupler; thus, virtual salinity fluxes are not needed.

Sea ice in E3SM is represented by the version 3 of the MPAS-SeaIce dynamic-thermodynamic
 1265 model. MPAS-SeaIce solves the sea-ice momentum equation on the B-grid of the horizontal mesh described above. It uses the CICE Consortium's Icepack sea-ice column physics, including the mushy layer thermodynamics. From v2.1 to v3, the cut-off for dynamical sea ice modeling was reduced from a minimum concentration of 10^{-3} to 10^{-11} , and an ice mass per unit area of 10^{-2} to 10^{-10} . The ocean – sea-ice coupling uses a constant 4 psu salinity, an ice-ocean drag coefficient
 1270 of 5.36×10^{-3} , and a constant ocean heat transfer coefficient 6×10^{-3} .

The simulation included in this manuscript was run for 5 cycles of the 1958-2020 period. The ocean-sea ice simulation starts from rest, where the ocean hydrography is initialized from the January 1991-2020 climatology of the World Ocean Atlas 2023 (Locarnini et al., 2024; Reagan et al., 2024). The sea-ice state is initialized as a uniform 1-m-thick disk of sea ice for latitudes
 1275 north of 70°N and south of 60°S , and where ocean sea surface temperature does not exceed the freezing point by 0.2°C .

A7 National Center for Atmospheric Research (NCAR)

NCAR has three contributions: 2 simulations with POP2 and 1 simulation with MOM6. POP2-based simulations use nominal horizontal grid spacings of 1° and 0.1° , which are referred to as
 1280 NCAR1 and NCAR10, respectively. These configurations employ the ocean and sea-ice components of the Community Earth System Model version 2 (CESM2; Danabasoglu et al., 2020). The MOM6-based configuration represents a new global coupled ocean – sea-ice model configuration with a nominal horizontal grid spacing of $2/3^\circ$ ($\sim 0.67^\circ$), noting that MOM6 is the new ocean component of the upcoming version of CESM, i.e., CESM3. This configuration,



1285 referred to here as NCAR67, is still under rapid development, and the results presented here provide a first assessment of our forced simulations.

POP2 is a level (z) coordinate model with the hydrostatic and Boussinesq approximations (Smith et al., 2010; Danabasoglu et al., 2012a). The version used here includes several physics and numerical advancements as summarized in Danabasoglu et al. (2020). These include improved
 1290 treatment of continental freshwater discharge into unresolved estuaries (Sun et al., 2019) and a new parameterization of Langmuir mixing (Li et al., 2016). The sea-ice model is CICE version 5.1.2 (CICE5; Hunke et al., 2015), which features new mushy-layer thermodynamics (Turner and Hunke, 2015) with prognostic sea ice salinity and an updated melt pond parameterization (Hunke et al., 2013). CICE5 runs on the same horizontal grid (B-grid) as the ocean model.

1295 NCAR1 utilizes a dipole grid with the grid north pole displaced into Greenland. The horizontal resolution (nominal 1°) is uniform in the zonal direction at 1.125° but varies in the meridional direction from 0.27° at the Equator to $\sim 0.5^\circ$ at midlatitudes. There are 60 levels in the vertical, varying from 10 m at the surface to 250 m in the deep ocean with a maximum depth of 5500 m. NCAR1 employs the skew-flux form of GM (Griffies, 1998), with depth-dependent GM and
 1300 isopycnal diffusivities varying from $3000 \text{ m}^2 \text{ s}^{-1}$ near the surface to $300 \text{ m}^2 \text{ s}^{-1}$ ($600 \text{ m}^2 \text{ s}^{-1}$ for isopycnal) in the deep ocean (Ferreira et al., 2005; Danabasoglu and Marshall, 2007). Near-surface diabatic mesoscale fluxes are parameterized following Ferrari et al. (2008) and Danabasoglu et al. (2008) with a diffusivity of $3000 \text{ m}^2 \text{ s}^{-1}$. Fox-Kemper et al. (2008; 2011) parameterization is used to include the near-surface restratification effects of submesoscale
 1305 mixed-layer eddies. The KPP scheme – via the Community ocean Vertical Mixing (CVmix) framework – is employed to prescribe the vertical viscosity and diffusivity coefficients as detailed in Danabasoglu et al. (2012a) with a turbulent Pr of 10. The background values have a latitudinal structure but are constant in the vertical. A tidal mixing parameterization provides enhanced abyssal mixing from tidally generated breaking waves (Jayne, 2009). NCAR1 uses an
 1310 overflow parameterization to represent the density-driven flows of the Denmark Strait, Faroe Bank Channel, the Ross Sea, and the Weddell Sea (Danabasoglu et al., 2010). Further details of this configuration can be found in Danabasoglu et al. (2012a; 2014; 2020).

NCAR10 utilizes a tripole grid with the grid north poles located in North America and Asia. It is based on versions documented in Small et al. (2014) but has been updated to the CESM2 code



base. The nominal 0.1° horizontal grid varies from 11 km at the Equator to 2.5 km at high latitudes. The vertical grid is the same as the one used in NCAR1, but it extends deeper to 6000 m with 2 additional levels, i.e., there are 62 vertical levels. NCAR10 uses a partial bottom cell formulation for more accurate representation of bathymetry, but there is no overflow parameterization. It uses a biharmonic horizontal mixing for tracers and momentum with respective coefficients of $-3.0 \times 10^9 \text{ m}^4 \text{ s}^{-1}$ and $-2.7 \times 10^{10} \text{ m}^4 \text{ s}^{-1}$. There is no additional parameterization of eddy-induced mixing in this configuration. A modified virtual salt flux formulation that uses a local reference salinity and allows redistribution of continental freshwater fluxes over several vertical layers near the surface is used in NCAR10, but the estuary model employed in NCAR1 to better represent the riverine freshwater flux is not available in NCAR10.

No Langmuir mixing parameterization is used, either. Furthermore, despite using the CICE5 code base, sea-ice physics is based on its previous CICE4 version to maintain consistency with other fully coupled high-resolution simulations that were run with earlier versions of CESM. Additional details of this configuration can be found in Chassignet et al. (2020).

In both NCAR1 and NCAR10, sea surface salinity is restored to World Ocean Atlas 2013 (WOA13; Zweng et al., 2013) monthly climatology with a piston velocity of 50 m over 1 year with subtraction of the global mean restoring. Both configurations use of a half-hour coupling frequency. The initial conditions are derived from the WOA13 (Locarnini et al., 2013; Zweng et al., 2013). NCAR1 and NCAR10 were integrated through 5 and 1 repeat cycles of the 1958-2018, respectively. In each, the last cycle was extended through 2023.

MOM6 (Adcroft et al., 2019) is the new ocean component of the forthcoming CESM3. NCAR67 uses a tripolar grid based on the semi-analytical method of Madec & Imbard (1996). The two northern grid poles are positioned over land in Canada and Russia, resulting in a global orthogonal curvilinear ocean mesh that has no singularity point within the computational domain with an aspect ratio approaching unity throughout the ocean, and it exhibits no loss of continuity in either the grid lines or the scale factors. Tropical mesh refinement is used, bringing the meridional resolution to 0.25° at the Equator to better capture the equatorial ocean dynamics. The ocean vertical coordinate is hybrid, using z^* near the surface and potential density in deeper layers, with a total of 75 layers. The depth of transition between z^* and isopycnals is the shallowest in the tropics (~ 50 m) and deepens towards high latitudes (~ 1200 and 2000 m in the Southern and Northern Hemispheres, respectively).



Bottom topography and coastlines are derived from the Global Bathymetry and Topography Dataset (SRTM15+ version 2.4; Tozer et al., 2019), with hand edits to improve the flow in several key regions. The minimum and maximum depths are set to 10 and 6000 m, respectively.

1350 The KPP parameterization for vertical mixing is enabled via the CVMix framework. Following the work described in Large et al. (2021), KPP has been updated to account for 1) a non-local momentum flux in the wind direction when the local shear is not aligned with the wind, and 2) the mixing induced by waves following the Monin-Obukhov similarity theory expanded to include the Stokes drift. Interior mixing components include convection, double-diffusion, and vertical shear of the horizontal velocity. The latitude-dependent diapycnal diffusivity due to
 1355 internal wave mixing described in Danabasoglu et al. (2012a) is included with diffusivity values in the range of $1.0 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$ (in the Banda Sea) to $1.0 \times 10^{-6} \text{ m}^2 \text{ s}^{-1}$ (background) with a turbulent Pr of 10. Energy dissipation from tidally induced breaking internal gravity waves is represented using the scheme developed by Simmons et al. (2004). The restratifying effects of baroclinic eddies in the mixed layer are represented using the parameterization developed by
 1360 Bodner et al. (2023).

Mesoscale eddies are parameterized via GM in conjunction with the diffusive mixing of tracers along neutral directions (Redi, 1982). GM is applied using the streamfunction formulation of Ferrari et al. (2010). Mesoscale eddy diffusion is incorporated using the neutral diffusion algorithm described in Shao et al. (2020) below the surface boundary layer, and the horizontal
 1365 diffusion algorithm described in Marques et al. (2023) within the surface boundary layer and transition zones. Both the GM and eddy diffusion coefficients are informed via the geometric formalism of Marshall et al. (2012), except that a prognostic eddy kinetic energy (Jansen et al., 2015) is employed, instead of the full (kinetic plus potential) eddy energy. The GM and Redi diffusivities have a minimum value of $50 \text{ m}^2 \text{ s}^{-1}$, typically reach $\sim 1500 \text{ m}^2 \text{ s}^{-1}$ in regions of strong
 1370 eddy kinetic energy, and peak at $6500 \text{ m}^2 \text{ s}^{-1}$ in only a few locations. Further details on how mesoscale eddies are represented in this configuration are given in Marques et al. (2023).

The computation of density is corrected to account for the interaction of grid-cell averaging with nonlinearities in the equation of state using the method of Stanley et al. (2020). The stochastic implementation described by Kenigson et al. (2022) is adopted both for hydrostatic pressure and



1375 for the computation of isopycnal slopes in GM, as described by Agarwal et al. (2023). The computation of density is not corrected in any other parts of the model.

The model lateral momentum equations use a biharmonic viscosity. The coefficient of this hyper-viscous closure is modeled using the 2D Leith scheme (Fox-Kemper & Menemenlis, 2008). No backscatter is applied, but the coefficient is smoothed as described by Grooms (2023). The background biharmonic viscosity is set to $1.0 \times 10^{12} \text{ m}^4 \text{ s}^{-1}$, with increased values in the tropical regions, reaching a maximum of $\sim 3.3 \times 10^{13} \text{ m}^4 \text{ s}^{-1}$. Momentum is extracted via a quadratic drag law with a constant bottom friction coefficient of 3×10^{-3} . The non-linear equation of state for seawater defined by Wright (1997) is applied.

1385 NCAR67 configuration uses the CICE Consortium model version 6.6 (CICE6). This includes several new physics options as well as a C-grid capability (not turned on currently). Presently, CICE6 needs POP2-style grid and topography files, but work is in progress to read the MOM6 grid and bathymetry files.

The simulation starts from rest with the initial potential temperature and salinity fields from the January-mean climatology of the World Ocean Atlas 2018 (WOA18). The sea surface salinity is restored to the monthly climatology of the upper 10-m averaged salinity from WOA18 using a piston velocity 50 m over 300 days. The simulation covers one forcing cycle for the 1958-2018 period.

A8 National Oceanography Centre (NOC)

1395 The NOC simulations used a code and configuration denoted as GO8p7 from a late stage of the development cycle of the Joint Marine Modelling Programme (JMMP) Global Ocean Sea Ice version 9 (GOSI9; Guiavarc'h et al., 2025). The JMMP is a partnership between the UK Met Office and UK research centres: the NOC, the British Antarctic Survey, and the Centre for Polar Observation and Modelling. The 0.25° and $1/12^\circ$ simulations were delivered as part of the UK NERC-funded ACSIS Programme, and the 1° simulation was run later specifically for this study.

1400 These simulations are referred to as NOC25, NOC12, and NOC1, respectively, in this manuscript.

GO8p7 is based on NEMO v4.0.4 (Madec & System Team, 2019) and uses NEMO's new native sea-ice model, SI3. The models use the extended eORCA grids, which have 75 vertical levels



with partial steps, with level thicknesses varying from 1 m at the surface to 200 m in the abyss.

1405 The 1° (eORCA1) bathymetry is derived from the ETOPO2 data set (NOAA, 2006) with the bathymetry on the Antarctic shelf based on IBSCO (Arndt et al., 2013). At 0.25° (eORCA025), the bathymetry is derived from the ETOPO1 data set (Amante & Eakins, 2009), with modifications in coastal regions based on GEBCO (IOC, IHO, & BODC, 2003) and the bathymetry on the Antarctic shelf derived from IBSCO. Finally, at 1/12° (eORCA12), the
 1410 bathymetry is derived from GEBCO_2014 (Weatherall et al., 2015).

GO8p7 uses a vector-invariant momentum advection scheme and a fourth-order TVD tracer advection and a weak grid-scale-aware GM eddy parameterization, with a coefficient that is scaled with the ratio of the cell size to the Rossby radius, capped at a maximum value of $70 \text{ m}^2 \text{ s}^{-1}$. A modified version of the Gaspar et al. (1990) TKE scheme (Madec et al., 2019) is used for
 1415 vertical mixing, and the parameterization of Simmons et al. (2004) for tidal mixing is applied. We use adaptive-implicit vertical advection (Shchepetkin & McWilliams, 2005). Lateral mixing of tracers is along neutral surfaces using the scheme of Redi (1982) and Cox (1987) with a coefficient equal to the cell size multiplied by 0.011 m s^{-1} . At 1°, the viscosity is Laplacian with a lateral viscosity coefficient of $20\,000 \text{ m}^2 \text{ s}^{-1}$, while at 0.25° and 1/12°, a bi-Laplacian viscosity is
 1420 used, with the coefficient set at the Equator to -1.5×10^{11} and $-1.25 \times 10^{11} \text{ m}^4 \text{ s}^{-1}$, respectively, and reduces with the inverse cube of the cell size. The equation of state is TEOS-10 (IOC et al., 2010).

Between GO8p7 and the official release GOSI9, some additional tuning of parameters took place to reduce some of the larger biases. These include: a reduction of the TKE mixing depth and an
 1425 increase in the globally uniform chlorophyll concentration to improve mixed layer depth and reduce a subsurface warm bias in the 100-200 m depth range; and increases in snow conductivity, lateral melt, heat in ice leads, and albedo parameters to improve the ice representation.

In all cases, the model started from rest using January-mean for the 1995-2014 period from EN4
 1430 (Good et al., 2013) climatological conditions and completed one pass through the JRA55-do forcing dataset (Tsujino et al., 2018). A sea surface salinity (SSS) restoring towards monthly mean climatology is applied with a piston velocity of 50 m over 1500 days.



A9 UK Met Office (UKMO)

1435 The UKMO 0.25° model simulation differs from the NOC simulations described in Appendix A8
 in the following four aspects. First, it uses a hybrid quasi-Eulerian vertical coordinate system
 where model levels are “nearly” terrain-following in the proximity of the Greenland-Scotland
 ridge area while they follow z^* -coordinates with partial steps in the rest of the domain
 (Bruciaferri et al., 2018; 2024). Second, it adopts a “log-layer” formulation for the bottom drag
 1440 coefficient C_D . In NEMO, for stability reasons, the minimum and maximum values of C_D are
 capped. With z -coordinates, the bottom topography is represented as a series of step-like
 structures, resulting in friction being exerted both at the bottom and at the lateral wall-sides via a
 no-slip boundary condition. Conversely, with terrain-following levels the representation of the
 bathymetry is more realistic, making the usage of lateral boundary conditions unnecessary. For
 1445 this reason, in the area where model levels are terrain-following, the simulation uses a more
 realistic C_D minimum value of 3×10^{-3} (see, e.g., observational estimates by Garton & Sanford
 2003, Mauritzen et al. (2005), or numerical studies by Riemenschneider & Legg (2007) and
 Danabasoglu et al. (2010)), while in the rest of the domain it is set to 10^{-3} , the same value of
 NOC25 simulation. Third, it employs the Griffies et al. (1998) triad formulation for the iso-
 1450 neutral diffusion because it is the only available option for using iso-neutral mixing with hybrid
 vertical coordinates in the current release of NEMO (see Appendix D of Bruciaferri et al. (2024)
 for a comparison between this formulation and the one used in the standard GOSI9). Finally, it
 uses the Density Jacobian with Cubic (DJC) polynomial scheme of Shchepetkin & McWilliams
 (2003) as implemented in NEMO by Bell & Bruciaferri (2022) for a more accurate calculation of
 1455 horizontal pressure forces when using steeply inclined vertical levels. This simulation is referred
 to as UKMO25 in the manuscript.

Appendix B: Cycle-to-cycle transport differences

We provide an assessment of cycle-to-cycle differences in overturning transports, using the first
 and fifth cycle transports in σ_0 space from GEOMAR05 and GEOMAR25 in Fig. B1. The top
 1460 panel presents the annual-mean maximum total transport timeseries, showing that the transports
 decrease from the first to the fifth cycle. The largest differences of > 5 Sv occur during the first
 decade of a forcing cycle in both simulations and during the 1980s to the early 2000s in
 GEOMAR05. By 2010, the transport differences become relatively small at ~ 2 Sv between the



cycles. The time-mean transport profiles are provided in the bottom panels of Fig. B1. As in Fig.
 1465 6, the solid black lines represent OSNAP with the gray shading showing ± 1 SDs, and the colored
 lines show the profiles for the first and fifth cycles from GEOMAR05 and GEOMAR25. In
 GEOMAR05, transport differences emerge for densities $> 27.3 \text{ kg m}^{-3}$, likely reflecting the long
 adjustment timescales of the deep ocean. The maximum transports are smaller in the fifth cycle
 by $\sim 0.6 \text{ Sv}$ at OW, $\sim 1.2 \text{ Sv}$ at OE, and $\sim 1.6 \text{ Sv}$ for the total. In GEOMAR25, the cycle-to-cycle
 1470 profile differences exist for low density classes as well. The maximum transports are smaller in
 the fifth cycle by $\sim 1.3 \text{ Sv}$ at OW, $\sim 1.8 \text{ Sv}$ at OE, and $\sim 2.1 \text{ Sv}$ for the total – these are larger
 differences than those of GEOMAR05. Nevertheless, except for the fifth cycle of GEOMAR25,
 the maximum transports are within the observational SD ranges. Interestingly, while the first
 cycle profile has a better agreement with OSNAP at OE, the fifth cycle profile matches that of
 1475 OSNAP better at OW in GEOMAR05. In contrast, GEOMAR25 has a better agreement with
 OSNAP in the fifth cycle for densities $< 27.4 \text{ kg m}^{-3}$ and in the first cycle for larger density
 classes. These cycle-to-cycle transport differences are certainly smaller than the respective inter-
 model spreads shown in Figs. 4 and 6 for the observational period. We also note as a caveat that
 these differences, i.e., their magnitudes and signs, are likely model dependent as discussed in
 1480 Danabasoglu et al. (2014).




Figure B1. Transport timeseries and profiles in σ_0 space for the first and fifth cycles from GEOMAR05 and GEOMAR25: (top) Annual-mean maximum total transport timeseries and (bottom) time-mean transport profiles for OW (left), OE (middle), and total (right). The solid black lines show the OSNAP transports. In (b), gray shading indicates ± 1 monthly standard deviation.

Code Availability. The METRIC Package is available at <https://github.com/NCAR/metric>. The TEOS-10 equation of state is available at <https://github.com/TEOS-10/GSW-Python>.

Data Availability. The simulation datasets for our analysis region are available at <https://doi.org/10.5281/zenodo.17457323> (Castruccio and Danabasoglu, 2025). For some of the simulations, entire global datasets for the full integration lengths are available as follows. ACCESS-OM2 (ANU1), ACCESS-OM2-025 (ANU25), and GFDL25 data were contributed to CMIP6 and are available on ESGF at <https://aims2.llnl.gov/search/cmip6/>. ACCESS-OM2-01 (ANU10) data are available via the NCI Data Catalogue at <https://dx.doi.org/10.25914/608097cb3433f>. Output from the NOC25 and NOC08 simulations



1495 are available from CEDA at <https://data.ceda.ac.uk/bode/SOC220065>. The OSNAP observations
 are available at Georgia Tech Digital Repository at <https://doi.org/10.35090/gatech/78023>.

Author contributions. GD and FC conceived and proposed the idea for this study. GD led writing
 of the manuscript. FC led the analysis of all model simulations. BB helped with the analysis. YF
 and MSL provided observational contributions. The following authors provided the datasets from
 1500 the respective simulations along with their brief model descriptions, listed here based on their
 center names: AH, RMH, HH, and AEK for ANU; DI for CMCC; ABo, EPC, and XX for FSU;
 ABi and FUS for GEOMAR; RZ for GFDL; AMB, LvR, and JW for LANL; GD, FC, FB, IG,
 and GM for NCAR; ABi and AM for NOC; and DB, CG, and DS for UKMO. All authors
 contributed to writing and editing of the manuscript.

1505 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. This effort at the US National Science Foundation (NSF) National Center for
 Atmospheric Research (NCAR) was supported by the contract 1947282 from the US Department
 of Energy (DOE) for the ImPACTS Project and by the grant NA18OAR4310429 from the US
 National Oceanic and Atmospheric Administration (NOAA), Climate Program Office (CPO),
 1510 Climate Variability and Predictability Program; Modeling Analysis, Predictions, and Projections
 Program; and the NOAA Global Ocean Monitoring and Observing (GOMO) Program, and by the
 DOE, Earth and Environmental System Modeling, Regional and Global Model Analysis
 Program. The US NSF NCAR is a major facility sponsored by the US NSF under Cooperative
 Agreement No. 1852977. The project was also partly funded by the US NSF – UK Natural
 1515 Environment Research Council (NERC) project entitled “Wider Impacts of Subpolar North
 Atlantic Decadal Variability on the Ocean and Atmosphere (WISHBONE)”. OSNAP data were
 collected and made freely available by the Overturning in the Subpolar North Atlantic Program
 and all the national programs that contribute to it (www.o-snap.org). The authors extend gratitude
 to everyone involved in the OSNAP program, including those who spend many hours at sea
 1520 collecting data. Dr. Alistair Adcroft and Raphael Dussin are thanked for conducting the GFDL
 OM4 OMIP-2 simulation. Dr. Alistair Adcroft is also thanked for his review of the manuscript.
 Bill Hurlin is thanked for publishing the GFDL OM4 OMIP-2 data. We acknowledge the
 contributions of Filipe Pereira, who passed away before completion of this manuscript. AEK was



supported by the Australian Research Council (ARC) grant LP200100406. RMH was supported
 1525 by the ARC grant DE21010004. AM was supported by the UK Natural Environment Research
 Council under the Atlantic Climate System Integrated Study (ACSIS; grant number
 NE/N018044/1) and the Climate Linked Atlantic Sector Science (CLASS; grant number
 NE/R015953/1) marine research programmes. ABI was supported by CLASS (NE/R015953/1)
 and the WISHBONE project (NE/T013540/1). BB was partly supported by the Turkish Scientific
 1530 and Technological Research Council (TUBITAK) via an International Research Fellowship
 Programme for PhD Students (2214-A). MSL and YF acknowledge funding from the Physical
 Oceanography Program of the US NSF (OCE-1948335). The ACCESS-OM2 model runs were
 undertaken by the Consortium for Ocean-Sea Ice Modelling in Australia (COSIMA)
 (http://www.cosima.org.au) using computational resources provided by the Australian
 1535 Government through the National Computational Infrastructure (NCI) under the National
 Computational Merit Allocation Scheme and ANU Merit Allocation Scheme. The GEOMAR
 simulations were funded by the Earth System Modelling Project (ESM) by providing computing
 time on the ESM partition of the supercomputer JUWELS at Jülich Supercomputing Center
 (JSC). The MPAS-O simulations were produced as part of the DOE Energy Exascale Earth
 1540 System Model (E3SM) project funded by the DOE Earth System Model Development program
 area of the Earth and Environmental System Modeling program, using a high-performance
 computing cluster operated by the Laboratory Computing Resource Center at the Argonne
 National Laboratory. This effort benefited greatly from the discussions that occurred during the
 joint WISHBONE and SNAP-DRAGON (Subpolar North Atlantic Processes – Dynamics and
 1545 Predictability of Variability in Gyre and Overturning) project meetings.

References

- Adcroft, A., Hill, C., and Marshall, J.: Representation of topography by shaved cells in a height
 coordinate ocean model, *Mon. Wea. Rev.*, 125, 2293–2315, [https://doi.org/10.1175/1520-0493\(1997\)125<2293:ROTBSC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1997)125<2293:ROTBSC>2.0.CO;2), 1997.
- 1550 Adcroft, A., Anderson, W., Balaji, V., Blanton, C., Bushuk, M., Dufour, C. O., Dunne, J. P.,
 Griffies, S. M., Hallberg, R., Harrison, M. J., Held, I. M., Jansen, M. F., John, J. G., Krasting,
 J. P., Langenhorst, A. R., Legg, S., Liang, Z., McHugh, C., Radhakrishnan, A., Reichl, B. G.,
 Rosati, T., Samuels, B. L., Shao, A., Stouffer, R., Winton, M., Wittenberg, A. T., Xiang, B.,
 Zadeh, N., and Zhang, R.: The GFDL global ocean and sea ice model OM4.0: Model
 1555 description and simulation features, *J. Adv. Model. Earth Sy.*, 11(10), 3167–3211,
<https://doi.org/10.1029/2019MS001726>, 2019.



- Agarwal, N., Small, R. J., Bryan, F. O., Grooms, I., and Pegion, P.J.: Impact of stochastic ocean density corrections on air-sea flux variability, *Geophys. Res. Lett.*, 50, e2023GL104248, 2023.
- 1560 Amante, C. and Eakins, B. W.: ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis, NOAA Technical Memorandum NESDIS NGDC-24, 19 pp., 2009.
- Arakawa, A. and Hsu, Y.-J. G.: Energy conserving and potential-entropy dissipating schemes for the shallow water equations, *Mon. Wea. Rev.*, 118, 1960–1969, [https://doi.org/10.1175/1520-0493\(1990\)118<1960:ECAPED>2.0.CO;2](https://doi.org/10.1175/1520-0493(1990)118<1960:ECAPED>2.0.CO;2), 1990.
- 1565 Arndt, J. E., Schenke, H. W., Jakobsson, M., Nitsche, F. O., Buys, G., Goleby, B., Rebesco, M., Bohoyo, F., Hong, J., Black, J., Greku, R., Udintsev, G., Barrios, F., Reynoso-Peralta, W., Taisei, M., and Wigley, R.: The International Bathymetric Chart of the Southern Ocean (IBCSO) Version 1.0 – A new bathymetric compilation covering circum-Antarctic waters, *Geophys. Res. Lett.*, 40, 3111–3117, <https://doi.org/10.1002/grl.50413>, 2013.
- 1570 Bamber, J. L., Tedstone, A. J., King, M. D., Howat, I. M., Enderlin, E. M., van den Broeke, M. R., and Noel, B.: Land ice freshwater budget of the Arctic and North Atlantic Oceans: 1. Data, methods and results, *J. Geophys. Res.*, 123, 1827–1837, <https://doi.org/10.1002/2017JC013605>, 2018.
- 1575 Barnier, B., Madec, G., Penduff, T., Molines, J., Treguir, A.-M., Le Sommer, J., Beckmann, A., Biastoch, A., Böning, C., Dengg, J., Derval, C., Durand, E., Gulev, S., Remy, E., Talandier, C., Theetten, S., Maltrud, M., McClean, J., and de Cuevas, B.: Impact of partial steps and momentum advection schemes in a global ocean circulation model at eddy permitting resolution, *Ocean Dynam.*, 56, 543–567, <https://doi.org/10.1007/s10236-006-0082-1>, 2006.
- 1580 Beckmann, A. and Doscher, R.: A method for improved representation of dense water spreading over topography in geopotential-coordinate models, *J. Phys. Oceanogr.*, 27, 581–591, [http://dx.doi.org/10.1175/1520-0485\(1997\)027<0581:AMFIRO>2.0.CO;2](http://dx.doi.org/10.1175/1520-0485(1997)027<0581:AMFIRO>2.0.CO;2), 1997.
- Bell, M. J. and Bruciaferri, D.: Accurate calculation of pressure forces on cells defined by steeply sloping coordinates, EGU General Assembly 2022, Vienna, Austria, 23–27 May 2022, EGU22-5012, <https://doi.org/10.5194/egusphere-egu22-5012>, 2022.
- 1585 Bi, D., Marsland, S., Uotila, P., O’Farrell, S., Fiedler, R., Sullivan, A., Griffies, S., Zhou, X., and Hirst, A.: ACCESS-OM: the ocean and sea-ice core of the ACCESS coupled model, *Australian Meteorological and Oceanographic Journal*, 63, 213–232, <http://www.bom.gov.au/jshess/papers.php?year=2013>, 2013.
- 1590 Bi, D., Dix, M., Marsland, S., O’Farrell, S., Sullivan, A., Bodman, R., Law, R., Harman, I., Sribinovsky, J., Rashid, H. A., Dobrohotoff, P., Mackallah, C., Yan, H., Hirst, A., Savita, A., Dias, F. B., Woodhouse, M., Fiedler, R., and Heerdegen, A.: Configuration and spin-up of ACCESS-CM2, the new generation Australian Community Climate and Earth System Simulator Coupled Model, *J. Southern Hemisphere Earth Sy. Sci.*, 70, 225–251, <https://doi.org/10.1071%2Fes19040>, 2020.
- 1595 Biastoch, A., Schwarzkopf, F. U., Getzlaff, K., Ruhs, S., Martin, T., Scheinert, M., Schulzki, T., Handmann, P., Hummels, R., and Böning, C.: Regional imprints of changes in the Atlantic meridional overturning circulation in the eddy-rich ocean model VIKING20X, *Ocean Sci.*, 17, 1177–1211, <https://doi.org/10.5194/os-17-1177-2021>, 2021.
- 1600 Bitz, C. M. and Lipscomb, W. H.: An energy-conserving thermodynamic model of sea ice, *J. Geophys. Res.*, 104, 15669–15677, 1999.



- Bitz, C. M., Holland, M., Eby, M., and Weaver, A. J.: Simulating the ice-thickness distribution in a coupled climate model, *J. Geophys. Res. Oceans*, 106, 2441–2463, <https://doi.org/10.1029/1999JC000113>, 2001.
- 1605 Blanke, B. and Delecluse, P.: Variability of the tropical Atlantic Ocean simulated by a general circulation model with two different mixed layer physics, *J. Phys. Oceanogr.*, 23, 1363–1388, 1993.
- Bleck, R.: An oceanic general circulation model framed in hybrid isopycnic-Cartesian coordinates, *Ocean Model.*, 4, 55–88, [https://doi.org/10.1016/S1463-5003\(01\)00012-9](https://doi.org/10.1016/S1463-5003(01)00012-9), 2002.
- 1610 Bodner, A. S., Fox-Kemper, B., Johnson, L., Van Roekel, L. P., McWilliams, J. C., Sullivan, P. P., Hall, P. S., and Dong, J.: Modifying the mixed layer eddy parameterization to include frontogenesis arrest by boundary layer turbulence, *J. Phys. Oceanogr.*, 53, 323–339, <https://doi.org/10.1175/JPO-D-21-0297.1>, 2023.
- Bouillon, S., Fichet, T., Legat, V., and Madec, G.: The elastic–viscous–plastic method revisited, *Ocean Model.*, 71, 2–12, <https://doi.org/10.1016/j.ocemod.2013.05.013>, 2013.
- 1615 Boyer, T., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Reagan, J. R., Antonov, J. I., Garcia, H. E., Baranova, O. K., Johnson, D. R., Seidov, D., Biddle, M. M., and Hamilton, M.: Changes to calculations of the World Ocean Atlas 2013 for version 2. Technical Report. NOAA, https://data.nodc.noaa.gov/woa/woa13/DOC/woa13v2_changes, 2015.
- 1620 Böning, C. W., Wagner, P., Handmann, P., Schwarzkopf, F. U., Getzlaff, K., and Biastoch, A.: Decadal changes in Atlantic overturning due to the excessive 1990s Labrador Sea convection, *Nat. Comm.*, 14, 1–10, <https://doi.org/10.1038/s41467-023-40323-9>, 2023.
- Bruciaferri, D., Shapiro, G. I., and Wobus, F.: A multi-envelope vertical coordinate system for numerical ocean modelling, *Ocean Dynamics*, 68, 1239–1258, <https://doi.org/10.1007/s10236-018-1189-x>, 2018.
- 1625 Bruciaferri, D., Guiavarc’h, C., Hewitt, H. T., Harle, J., Almansi, M., Mathiot, P., and Colombo, P.: Localized general vertical coordinates for quasi-Eulerian ocean models: The Nordic overflows test-case. *J. Adv. Model. Earth Sy.*, 16, e2023MS003893, <https://doi.org/10.1029/2023MS003893>, 2024.
- 1630 Bryden, H. L., Longworth, H. R., and Cunningham, S. A.: Slowing of the Atlantic meridional overturning circulation at 25°N, *Nature*, 438, 655–657, <https://doi.org/10.1038/nature04385>, 2005.
- Bryden, H., Beunk, J., Drijfhout, S., Hazeleger, W., and Mecking, J.: Comparing observed and modelled components of the Atlantic Meridional Overturning Circulation at 26° N, *Ocean Sci.*, 20, 589–599, <https://doi.org/10.5194/os-20-589-2024>, 2024.
- 1635 Buckley, M. W. and Marshall, J.: Observations, inferences, and mechanisms of Atlantic meridional overturning circulation variability: A review, *Rev. Geophys.*, 54, 5–63, <https://doi.org/10.1002/2015RG000493>, 2016.
- Buckley, M. W., Lozier, M. S., Desbruyeres, D., and Evans, D. G.: Buoyancy forcing and the subpolar Atlantic meridional overturning circulation, *Philos. Trans. R. Soc. A*, 381, 20220181, <https://doi.org/10.1098/rsta.2022.0181>, 2023.
- 1640 Caesar, L., Rahmstorf, S., Robinson, A., Feulner, G., and Saba, V.: Observed fingerprint of a weakening Atlantic Ocean overturning circulation, *Nature*, 556, 191–196, <https://doi.org/10.1038/s41586-018-0006-5>, 2018.



- 1645 Caesar, L., McCarthy, G. D., Thornalley, D. J. R., Cahill, N., and Rahmstorf, S.: Current Atlantic meridional overturning circulation weakest in last millennium, *Nature Geoscience*, 14, 118–120, <https://doi.org/10.1038/s41561-021-00699-z>, 2021.
- Campin, J.-M. and Goosse, H.: Parameterization of density-driven downsloping flow for a coarse-resolution ocean model in z-coordinate, *Tellus A: Dynam. Meteor. Oceanogr.*, 51, 412–430, <http://dx.doi.org/10.3402/tellusa.v51i3.13468>, 1999.
- 1650 Carnes, M. R.: Description and evaluation of GDEM-V3.0. Report number: NRL/MR/7330-09-9165, United States: Naval Research Laboratory, 2009.
- Castruccio, F. S.: NCAR/metric: metric v0.1. doi/10.5281/zenodo.4708277, 2021.
- Castruccio, F. and Danabasoglu, G.: Datasets to reproduce the figures in “Simulated and observed transport estimates across the Overturning in the Subpolar North Atlantic Program (OSNAP) section”, available at <https://doi.org/10.5281/zenodo.17457323>, 2025.
- 1655 Chafik, L., Holliday, N. P., Bacon, S., and Rossby, T.: Irminger Sea is the center of action for subpolar AMOC variability, *Geophys. Res. Lett.*, 49, e2022GL099133, <https://doi.org/10.1029/2022GL099133>, 2022.
- Chassignet, E. P., Smith, L. T., Halliwell, G. R., and Bleck, R.: North Atlantic simulations with the Hybrid Coordinate Ocean Model (HYCOM): Impact of the vertical coordinate choice, reference pressure, and thermobaricity, *J. Phys. Oceanogr.*, 33, 2504–2526, [https://doi.org/10.1175/1520-0485\(2003\)0332.0.CO;2](https://doi.org/10.1175/1520-0485(2003)0332.0.CO;2), 2003.
- Chassignet, E. P., Yeager, S. G., Fox-Kemper, B., Bozec, A., Castruccio, F., Danabasoglu, G., Kim, W. M., Koldunov, N., Li, Y., Lin, P., Liu, H., Sein, D., Sidorenko, D., Wang, Q., and Xu, X.: Impact of horizontal resolution on global ocean-sea-ice model simulations based on the experimental protocols of the Ocean Model Intercomparison Project phase 2 (OMIP-2), *Geosci. Model Dev.*, 13, 4595–4637, <https://doi.org/10.5194/gmd-13-4595-2020>, 2020.
- Cheng, W., Chiang, J. C. H., and Zhang, D.: Atlantic meridional overturning circulation (AMOC) in CMIP5 models: RCP and historical simulations, *J. Climate*, 26, 7187–7197, <https://doi.org/10.1175/JCLI-D-12-00496.1>, 2013.
- 1670 Cherchi, A., Fogli, P. G., Lovato, T., Peano, D., Iovino, D., Gualdi, S., Masina, S., Scoccimarro, E., Materia, S., Bellucci, A., and Navarra, A.: Global mean climate and main patterns of variability in the CMCC-CM2 coupled model, *J. Adv. Model. Earth Sy.*, 11, 185–209, <https://doi.org/10.1029/2018MS001369>, 2019.
- 1675 Cipollone, A., Storto, A., and Masina, S.: Implementing a parallel version of a variational scheme in a global assimilation system at eddy-resolving resolution, *J. Atmos. Ocean. Tech.*, 37, 1865–1876, 2020.
- Cox, M. D.: Isopycnal diffusion in a z-coordinate ocean model, *Ocean Modelling*, 74, 1–9, 1987.
- Cunningham, S. A., Kanzow, T., Rayner, D., Baringer, M. O., Johns, W. E., Marotzke, J., Longworth, H. R., Grant, E. M., Hirschi, J. J.-M., Beal, L. M., Meinen, C. S., and Bryden, H. L.: Temporal variability of the Atlantic meridional overturning circulation at 26.5°N, *Science*, 317, 935–938, <https://doi.org/10.1126/science.1141304>, 2007.
- 1680 Danabasoglu, G.: On multi-decadal variability of the Atlantic meridional overturning circulation in the Community Climate System Model version 3 (CCSM3), *J. Climate*, 21, 5524–5544, <https://doi.org/10.1175/2008JCLI2019.1>, 2008.
- 1685



- Danabasoglu, G. and Marshall, J.: Effects of vertical variations of thickness diffusivity in an ocean general circulation model, *Ocean Model.*, 18, 122–141, <https://doi.org/10.1016/j.ocemod.2007.03.006>, 2007.
- 1690 Danabasoglu, G., Ferrari, R., and McWilliams, J. C.: Sensitivity of an ocean general circulation model to a parameterization of near-surface eddy fluxes, *J. Climate*, 21, 1192–1208, <https://doi.org/10.1175/2007JCLI1508.1>, 2008.
- Danabasoglu, G., Large, W. G., and Briegleb B. P.: Climate impacts of parameterized Nordic Sea overflows, *J. Geophys. Res.*, 115, C11005, <http://doi.org/10.1029/2010JC006243>, 2010.
- 1695 Danabasoglu, G., Bates, S. C., Briegleb, B. P., Jayne, S. R., Jochum, M., Large, W. G., Peacock, S., and Yeager, S. G.: The CCSM4 ocean component, *J. Climate*, 25, 1361–1389, <https://doi.org/10.1175/JCLI-D-11-00091.1>, 2012a.
- Danabasoglu, G., Yeager, S. G., Kwon, Y.-O., Tribbia, J. J., Phillips, A. S., and Hurrell, J. W.: Variability of the Atlantic meridional overturning circulation in CCSM4, *J. Climate*, 25, 5153–5172, <https://doi.org/10.1175/JCLI-D-11-00463.1>, 2012b.
- 1700 Danabasoglu, G., Yeager, S. G., Bailey, D., Behrens, E., Bentsen, M., Bi, D., Biastoch, A., Böning, C., Bozec, A., Canuto, V. M., Cassou, C., Chassignet, E., Coward, A. C., Danilov, S., Diansky, N., Drange, H., Farneti, R., Fernandez, E., Fogli, P. G., Forget, G., Fujii, Y., Griffies, S. M., Gusev, A., Heimbach, P., Howard, A., Jung, T., Kelley, M., Large, W. G., Leboissetier, A., Lu, J., Madec, G., Marsland, S. J., Masina, S., Navarra, A., Nurser, A. J. G., Pirani, A., Salas y Mélia, D., Samuels, B. L., Scheinert, M., Sidorenko, D., Treguier, A.-M., Tsujino, H., Uotila, P., Valcke, S., Voldoire, A., and Wang, Q.: North Atlantic simulations in Coordinated Ocean-ice Reference Experiments phase II (CORE-II). Part I: Mean states, *Ocean Model.* 73, 76–107. <https://doi.org/10.1016/j.ocemod.2013.10.005>, 2014.
- 1705 Danabasoglu, G., Yeager, S. G., Kim, W. M., Behrens, E., Bentsen, M., Bi, D., Biastoch, A., Bleck, R., Böning, C., Bozec, A., Canuto, V. M., Cassou, C., Chassignet, E., Coward, A. C., Danilov, S., Diansky, N., Drange, H., Farneti, R., Fernandez, E., Fogli, P. G., Forget, G., Fujii, Y., Griffies, S. M., Gusev, A., Heimbach, P., Howard, A., Ilıcak, M., Jung, T., Karspeck, A., R., Kelley, M., Large, W. G., Leboissetier, A., Lu, J., Madec, M., Marsland, S. J., Masina, S., Navarra, A., Nurser, A. J. G., Pirani, A., Romanou, A., Salas y Melia, D., Samuels, B. L., Scheinert, M., Sidorenko, D., Sun, S., Treguier, A.-M., Tsujino, H., Uotila, P., Valcke, S., Voldoire, A., Wang, Q., and Yashayaev, I.: North Atlantic simulations in Coordinated Ocean-ice Reference Experiments phase II (CORE-II). Part II: Inter-annual to decadal variability, *Ocean Model.*, 97, 65–90, <https://doi.org/10.1016/j.ocemod.2015.11.007>, 2016.
- 1715 Danabasoglu, G., Landrum, L., Yeager, S. G., and Gent, P. R.: Robust and non-robust aspects of Atlantic meridional overturning circulation variability and mechanisms in the Community Earth System Model, *J. Climate*, 32, 7349–7369, <https://doi.org/10.1175/JCLI-D-19-0026.1>, 2019.
- 1720 Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: Community
- 1725



- 1730 Earth System Model version 2 (CESM2). *J. Adv. Model. Earth Sys.*, 12, e2019MS001916, <https://doi.org/10.1029/2019MS001916>, 2020.
- Debreu, L., Vouland, C., and Blayo, E.: AGRIF: Adaptive grid refinement in Fortran. *Comput. Geosci.*, 34, 8–13, 2008.
- 1735 Depoorter, M. A., Bamber, J. L., Griggs, J. A., Lenaerts, J. T. M., Ligtenberg, S. R. M., van den Broeke, M. R., and Moholdt, G.: Calving fluxes and basal melt rates of Antarctic ice shelves, *Nature*, 502, 89–92, <https://doi.org/10.1038/nature12567>, 2013.
- Doscher, R. and Beckmann, A.: Effects of a bottom boundary layer parameterization in a coarse-resolution model of the North Atlantic Ocean, *J. Atmos. Oceanic Tech.*, 17, 698–707, [http://dx.doi.org/10.1175/1520-0426\(2000\)017<0698:EOABBL>2.0.CO;2](http://dx.doi.org/10.1175/1520-0426(2000)017<0698:EOABBL>2.0.CO;2), 2000.
- 1740 Ferrari, R., McWilliams, J. C., Canuto, V. M., and Dubovikov, M.: Parameterization of eddy fluxes near oceanic boundaries, *J. Climate*, 21, 2770–2789, <https://doi.org/10.1175/2007JCLI1510.1>, 2008.
- Ferrari, R., Griffies, S. M., Nurser, A. G., and Vallis, G. K.: A boundary-value problem for the parameterized mesoscale eddy transport, *Ocean Modelling*, 32, 143–156, <https://doi.org/10.1016/j.ocemod.2010.01.004>, 2010.
- 1745 Ferreira, D., Marshall, J., and Heimbach, P.: Estimating eddy stresses by fitting dynamics to observations using a residual-mean ocean circulation model and its adjoint, *J. Phys. Oceanogr.*, 35, 1891–1910, <https://doi.org/10.1175/JPO2785.1>, 2005.
- Fichefet, T. and Morales Maqueda, M. A.: Sensitivity of a global sea ice model to the treatment of ice thermodynamic and dynamics, *J. Geophys. Res.*, 102, 12609–12646, 1997.
- 1750 Fofonoff, N. P. and Millard Jr., R. C.: Algorithms for the computation of fundamental properties of seawater, Paris, France, UNESCO, UNESCO Technical Papers in Marine Sciences, 44, 53 pp., <https://doi.org/10.25607/OBP-1450>, 1983.
- Fogli, P. G. and Iovino, D.: CMCC–CESM–NEMO: Toward the new CMCC Earth System Model, CMCC Research Paper no. 248, <https://doi.org/10.2139/ssrn.2603176>, 2014.
- 1755 Fox-Kemper, B. and Menemenlis, D.: Can large eddy simulation techniques improve mesoscale rich ocean models? *Geophys. Monograph Series*, 177, 319–337, 2008.
- Fox-Kemper, B., Ferrari, R., and Hallberg, R.: Parameterization of mixed layer eddies. Part I: Theory and diagnosis, *J. Phys. Oceanogr.*, 38, 1145–1165, <http://dx.doi.org/10.1175%2F2007JPO3792.1>, 2008.
- 1760 Fox-Kemper, B., Danabasoglu, G., Ferrari, R., Griffies, S. M., Hallberg, R. W., Holland, M. M., Maltrud, M. E., Peacock, S., and Samuels, B. L.: Parameterization of mixed layer eddies. III: Implementation and impact in global ocean climate simulations, *Ocean Model.*, 39, 61–78, <https://doi.org/10.1016/j.ocemod.2010.09.002>, 2011.
- 1765 Fox, A. D., Handmann, P., Schmidt, C., Fraser, N., Ruhs, S., Sanchez-Franks, A., Martin, T., Oltmanns, M., Johnson, C., Rath, W., Holliday, N. P., Biastoch, A., Cunningham, S. A., and Yashayaev, I.: Exceptional freshening and cooling in the eastern subpolar North Atlantic caused by reduced Labrador Sea surface heat loss, *Ocean Sci.*, 18, 1507–1533, <https://doi.org/10.5194/OS-18-1507-2022>, 2022.
- 1770 Fröhle, J., Handmann, P. V. K., and Biastoch, A.: Major sources of North Atlantic Deep Water in the subpolar North Atlantic from Lagrangian analyses in an eddy-rich ocean model, *Ocean Sci.*, 18, 1431–1450, <https://doi.org/10.5194/OS-18-1431-2022>, 2022.



- 1775 Fu, Y., Lozier, M. S., Biló, T. C., Bower, A. S., Cunningham, S. A., Cyr, F., de Jong, M. F., deYoung, B., Drysdale, L., Fraser, N., Fried, N., Furey, H. H., Han, G., Handmann, P., Holliday, N. P., Holte, J., Inall, M. E., Johns, W. E., Jones, S., Karstensen, J., Li, F., Pacini, A., Pickart, R. S., Rayner, D., Straneo, F., and Yashayaev, I.: Seasonality of the meridional overturning circulation in the subpolar North Atlantic, *Comm. Earth & Environment*, 4, 181., <https://doi.org/10.1038/s43247-023-00848-9>, 2023.
- 1780 Fu, Y., Lozier, M. S., Majumder, S., and Petit, T.: Water mass transformation and its relationship with the overturning circulation in the eastern subpolar North Atlantic, *J. Geophys. Res.: Oceans*, 129, <https://doi.org/10.1029/2024JC021222>, 2024.
- Fu, Y., Lozier, M. S., Bower, A. S., et al.: Characterizing the interannual variability of North Atlantic subpolar overturning, *Geophys. Res. Lett.*, in review, 2025.
- 1785 Gaspar, P., Grégoris, Y., and Lefevre, J.-M.: A simple eddy kinetic energy model for simulations of the oceanic vertical mixing: Tests at station Papa and long-term upper ocean study site, *J. Geophys. Res.*, 95, 16179–16193, <https://doi.org/10.1029/jc095ic09p16179>, 1990.
- GEBCO Compilation Group: GEBCO 2023 Grid, doi:10.5285/f98b053b-0cbc-6c23-e053-6c86abc0af7b, 2023.
- 1790 Gent, P. R. and McWilliams, J. C.: Isopycnal mixing in ocean circulation models, *J. Phys. Oceanogr.*, 20, 150–155, [https://doi.org/10.1175/15200485\(1990\)020<0150:imicm>2.0.co;2](https://doi.org/10.1175/15200485(1990)020<0150:imicm>2.0.co;2), 1990.
- Girton, J. B. and Sanford, T. B.: Descent and modification of the overflow plume in the Denmark Strait, *J. Phys. Oceanogr.*, 33, 1351–1363, [https://doi.org/10.1175/1520-0485\(2003\)033<1351:DAMOTO>2.0.CO;2](https://doi.org/10.1175/1520-0485(2003)033<1351:DAMOTO>2.0.CO;2), 2003.
- 1795 Golaz, J. C., Van Roekel, L. P., Zheng, X., Roberts, A. F., Wolfe, J. D., Lin, W., Bradley, A. M., Tang, Q., Maltrud, M. E., Forsyth, R. M., Zhang, C., Zhou, T., Zhang, K., Zender, C. S., Wu, M., Wang, H., Turner, A. K., Singh, B., Richter, J. H., Qin Y., Petersen, M. R., Mametjanov, A., Ma, P. L., Larson, V. E., Krishna, J., Keen, N. D., Jeffery, N., Hunke, E. C., Hannah, W. M., Guba, O., Griffin, B. M., Feng, Y., Engwirda, D., Di Vittorio, A. V., Dang, C., Conlon, L.
- 1800 A. M., Chen, C. C. J., Brunke, M. A., Bisht, G., Benedict, J. J., Asay-Davis, X. S., Zhang, Y., Zhang, M., Zeng, X., Xie, S., Wolfram, P. J., Vo, T., Veneziani, M., Tesfa, T. K., Sreepathi, S., Salinger, A. G., Eyre, J. E. J. R., Prather, M. J., Mahajan, S., Li, Q., Jones, P. W., Jacob, R. L., Huebler, G. W., Huang, X., Hillman, B. R., Harrop, B. E., Foucar, J. G., Fang, Y., Comeau, D. S., Caldwell, P. M., Bartoletti, T., Balaguru, K., Taylor, M. A., McCoy, R. B., Leung, L.
- 1805 R., and Bader, D. C.: The DOE E3SM model version 2: Overview of the physical model and initial model evaluation, *J. Adv. Model. Earth Sy.*, 14, e2022MS003156, <https://doi.org/10.1029/2022MS003156>, 2022.
- Good, S. A., Martin, M. J., and Rayner, N. A.: EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates, *J. Geophys. Res. Oceans*, 118, 6704–6716, <https://doi.org/10.1002/2013JC009067>, 2013.
- 1810 Griffies, S. M.: The Gent-McWilliams skew flux, *J. Phys. Oceanogr.*, 28, 831–841, [https://doi.org/10.1175/1520-0485\(1998\)028%3C0831:TGMSF%3E2.0.CO;2](https://doi.org/10.1175/1520-0485(1998)028%3C0831:TGMSF%3E2.0.CO;2), 1998.
- Griffies, S. M.: Elements of the Modular Ocean Model (MOM) 5 (2012 release with updates), Technical Report 7, NOAA/Geophysical Fluid Dynamics Laboratory Ocean Group, 2012.
- 1815 Griffies, S. M., Gnanadesikan, A., Pacanowski, R. C., Larichev, V., Dukowicz, J. K., and Smith, R. D.: Isoneutral diffusion in a z-coordinate ocean model, *J. Phys. Oceanogr.*, 28, 805–830, [https://doi.org/10.1175/1520-0485\(1998\)028%3C0805:IDIAZC%3E2.0.CO;2](https://doi.org/10.1175/1520-0485(1998)028%3C0805:IDIAZC%3E2.0.CO;2), 1998.



- Griffies, S. M., Gnanadesikan, A., Dixon, K. W., Dunne, J. P., Gerdes, R., Harrison, M. J.,
 Rosati, A., Russell, J. L., Samuels, B. L., Spelman, M. J., Winton, M., and Zhang, R.:
 1820 Formulation of an ocean model for global climate simulations, *Ocean Sci.*, 1, 45–79,
<https://doi.org/10.5194/os-1-45-2005>, 2005.
- Griffies, S. M., Biastoch, A., Böning, C., Bryan, F., Danabasoglu, G., Chassignet, E. P., England,
 M. H., Gerdes, R., Haak, H., Hallberg, R. W., Hazeleger, W., Jungclaus, J., Large, W. G.,
 1825 Madec, G., Pirani, A., Samuels, B. L., Scheinert, M., Sen Gupta, A., Severijns, C. A.,
 Simmons, H. L., Treguier, A. M., Winton, M., Yeager, S., and Yin, J.: Coordinated Ocean-ice
 Reference Experiments (COREs), *Ocean Model.* 26, 1–46,
<https://doi.org/10.1016/j.ocemod.2008.08.007>, 2009.
- Griffies, S. M., Danabasoglu, G., Durack, P. J., Adcroft, A. J., Balaji, V., Böning, C. W.,
 Chassignet, E. P., Curchitser, E., Deshayes, J., Drange, H., Fox-Kemper, B., Gleckler, P. J.,
 1830 Gregory, J. M., Haak, H., Hallberg, R. W., Hewitt, H. T., Holland, D. M., Ilyina, T.,
 Jungclaus, J. H., Komuro, Y., Krasting, J. P., Large, W. G., Marsland, S. J., Masina, S.,
 McDougall, T. J., Nurser, A. J. G., Orr, J. C., Pirani, A., Qiao, F., Stouffer, R. J., Taylor, K. E.,
 Treguier, A. M., Tsujino, H., Uotila, P., Valdivieso, M., Winton, M., and Yeager, S. G.: OMIP
 1835 contribution to CMIP6: Experimental and diagnostic protocol for the physical component of
 the Ocean Model Intercomparison Project, *Geosci. Model Dev.*, 9, 3231–3296,
<https://doi.org/10.5194/gmd-9-3231-2016>, 2016.
- Grooms, I.: Backscatter in energetically-constrained Leith parameterizations, *Ocean Modelling*,
 186, 102265, 2023.
- Guiavarc’h, C., Storkey, D., Blaker, A. T., Blockley, E., Megann, A., Hewitt, H. T., Bell, M. J.,
 1840 Calvert, D., Copsey, D., Sinha, B., Moreton, S., Mathiot, P., and An, B.: GOSI9: UK Global
 Ocean and Sea Ice configurations, *Geosci. Model Dev.*, 18, 377–403,
<https://doi.org/10.5194/gmd-18-377-2025>, 2025.
- Hallberg, R.: Using a resolution function to regulate parameterizations of oceanic mesoscale
 eddy effects, *Ocean Model.*, 72, 92–103, <https://doi.org/10.1016/j.ocemod.2013.08.007>,
 1845 2013.
- Held, I. M., Guo, H., Adcroft, A., Dunne, J. P., Horowitz, L. W., Krasting, J., Shevliakova, E.,
 Winton, M., Zhao, M., Bushuk, M., Wittenberg, A. T., Wyman, B., Xiang, B., Zhang, R.,
 Anderson, W., Balaji, V., Donner, L., Dunne, K., Durachta, J., Gauthier, P. P. G., Ginoux, P.,
 1850 Golaz, J.-C., Griffies, S. M., Hallberg, R., Harris, L., Harrison, M., Hurlin, W., John, J., Lin,
 P., Lin, S.-J., Malyshev, S., Menzel, R., Milly, P. C. D., Ming, Y., Naik, V., Paynter, D.,
 Paulot, F., Rammaswamy, V., Reichl, B., Robinson, T., Rosati, A., Seman, C., Silvers, L. G.,
 Underwood, S., and Zadeh, N.: Structure and performance of GFDL's CM4. 0 climate
 model, *J. Adv. Model. Earth Sy.*, 11, 3691–3727, <https://doi.org/10.1029/2019MS001829>,
 2019.
- Hirschi, J. J.-M., Barnier, B., Böning, C., Biastoch, A., Blaker, A. T., Coward, A., Danilov, S.,
 Drijfhout, S., Getzlaff, K., Griffies, S. M., Hasumi, H., Hewitt, H., Iovino, D., Kawasaki, T.,
 Kiss, A. E., Koldunov, N., Marzocchi, A., Mecking, J. V., Moat, B., Molines, J.-M., Myers, P.
 G., Penduff, T., Roberts, M., Treguier, A.-M., Sein, D. V., Sidorenko, D., Small, J., Spence,
 P., Thompson, L., Weijer, W., and Xu, X. (2020), The Atlantic Meridional Overturning
 1860 Circulation in high resolution models, *J. Geophys. Res. Oceans*, 125, e2019JC015522,
<https://dx.doi.org/10.1029/2019JC015522>, 2020.



- Hirt, C. W., Amsden, A. A., and Cook, J. L.: An arbitrary Lagrangian-Eulerian computing method for all flow speeds, *J. Comput. Phys.*, 135, 203–216, <https://doi.org/10.1006/jcph.1997.5702>, 1997.
- 1865 Huneke, W. G. C., Hogg, A. McC., Dix, M., Bi, D., Sullivan, A., McGregor, S., Holgate, C., O’Farrell, S. P., and Oliveira, M. J. T.: The ACCESS-CM2 climate model with a higher resolution ocean-sea ice component (1/4°), *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2025-1006>, 2025.
- 1870 Hunke, E. C. and Dukowicz, J. K.: An elastic–viscous–plastic model for sea ice dynamics, *J. Phys. Oceanogr.*, 27, 1849–1867, 1997.
- Hunke, E. C. and Lipscomb, W. H.: CICE: the Los Alamos Sea Ice Model documentation and software user’s manual version 4.1, Technical Report, LA-CC-06-012, 76 pp., available at: <https://github.com/CICE-Consortium/CICE-svn-trunk/releases/tag/cice-4.1>, 2010.
- 1875 Hunke, E. C., Hebert, D. A., and Lecomte, O.: Level-ice melt ponds in the Los Alamos sea ice model, *CICE, Ocean Model.*, 71, 26–42, <https://doi.org/10.1016/j.ocemod.2012.11.008>, 2013.
- Hunke, E. C., Lipscomb, W. H., Turner, A. K., Jeffery, N., and Elliott, S.: CICE: the Los Alamos Sea Ice Model documentation and software user’s manual version 5.1, Technical Report LA-CC-06-012, Los Alamos National Laboratory, Los Alamos, NM 87545, 2015.
- 1880 IOC, IHO and BODC: Centenary Edition of the GEBCO Digital Atlas, published on CD-ROM on behalf of the Intergovernmental Oceanographic Commission and the International Hydrographic Organization as part of the General Bathymetric Chart of the Oceans, British Oceanographic Data Centre, Liverpool, UK, 2003.
- 1885 IOC, SCOR, and IAPSO: The international thermodynamic equation of seawater – 2010: Calculation and use of thermodynamic properties, 2010.
- Iovino, D., Masina, S., Storto, A., Cipollone, A., and Stepanov, V. N.: A 1/16° eddying simulation of the global NEMO sea-ice–ocean system, *Geosci. Model Dev.*, 9, 2665–2684, <https://doi.org/10.5194/gmd-9-2665-2016>, 2016.
- 1890 Iovino, D., Fogli, P. G., and Masina, S.: Evaluation of the CMCC global eddying ocean model for the Ocean Model Intercomparison Project (OMIP2), *Geosci. Model Dev.*, 16, 6127–6159, <https://doi.org/10.5194/gmd-16-6127-2023>, 2023.
- Jackett, D. R., McDougall, T. J., Feistel, R., Wright, D. G., and Griffies, S. M.: Algorithms for density, potential temperature, conservative temperature, and the freezing temperature of seawater, *J. Atmos. Oceanic Tech.*, 23, 1709–1728, <http://dx.doi.org/10.1175/JTECH1946.1>, 2006.
- 1895 Jackson, L. C. and Petit, T.: North Atlantic overturning and water mass transformation in CMIP6 models, *Clim. Dyn.*, <https://doi.org/10.1007/s00382-022-06448-1>, 2022.
- Jackson, L. C., Dubois, C., Forget, G., Haines, K., Harrison, M., Iovino, D., Kohl, A., Mignac, D., Masina, S., Peterson, K. A., Picuch, C. G., Roberts, C. D., Robson, J., Storto, A., 1900 Toyoda, T., Valdivieso, M., Wilson, C., Wang, Y., and Zuo, H.: The mean state and variability of the North Atlantic circulation: A perspective from ocean reanalyses, *J. Geophys. Res.: Oceans*, 124, 9141–9170, <https://doi.org/10.1029/2019JC015210>, 2019.
- Jakobsson, M., Mayer, L., Coakley, B., Dowdeswell, J. A., Forbes, S., Fridman, B., Hodnesdal, H., Noormets, R., Pedersen, R., Rebescio, M., Schenke, H. W., Zarayskaya, Y., Accettella, 1905 D., Armstrong, A., Anderson, R. M., Bienhoff, P., Camerlenghi, A., Church, I., Edwards, M.,



- Gardner, J. V., Hall, J. K., Hell, B., Hestvik, O., Kristoffersen, Y., Marcussen, C., Mohammad, R., Mosher, D., Nghiem, S. V., Pedrosa, M. T., Travaglini, P. G., and Weatherall, P.: The international bathymetric chart of the Arctic Ocean (IBCAO) Version 3.0. *Geophys. Res. Lett.*, 39, L1260, <https://doi.org/10.1029/2012GL052219>, 2012.
- 1910 Jansen, M. F., Adcroft, A. J., Hallberg, R., and Held, I. M.: Parameterization of eddy fluxes based on a mesoscale energy budget, *Ocean Modelling*, 92, 28–41, <https://doi.org/10.1016/j.ocemod.2015.05.007>, 2015.
- Jayne, S. R.: The impact of abyssal mixing parameterizations in an ocean general circulation model, *J. Phys. Oceanogr.*, 39, 1756–1775, <https://doi.org/10.1175/2009JPO4085.1>, 2009.
- 1915 Jochum, M.: Impact of latitudinal variations in vertical diffusivity on climate simulations, *J. Geophys. Res.*, 114, C01010, <https://doi.org/10.1029/2008JC005030>, 2009.
- Karspeck, A., Stammer, D., Kohl, A., Danabasoglu, G., Balmaseda, M., Smith, D. M., Fujii, Y., Zhang, S., Giese, B., Tsijino, H., and Rosati, A.: Comparison of the Atlantic meridional overturning circulation between 1960 and 2007 in six ocean reanalysis products, *Clim. Dyn.*, 49, 957–982, <https://doi.org/10.1007/s00382-015-2787-7>, 2017.
- 1920 Kenigson, J. S., Adcroft, A., Bachman, S.D., Castruccio, F., Grooms, I., Pegion, P., and Stanley, Z.: Parameterizing the impact of unresolved temperature variability on the large-scale density field: 2. Modeling, *J. Adv. Model. Earth Syst.*, 14, e2021MS002844, 2022.
- Kiss, A. E., Hogg, A. M., Hannah, N., Boeira Dias, F., Brassington, G. B., Chamberlain, M. A., Chapman, C., Dobrohotoff, P., Domingues, C. M., Duran, E. R., England, M. H., Fiedler, R., Griffies, S. M., Heerdegen, A., Heil, P., Holmes, R. M., Klocker, A., Marsland, S. J., Morrison, A. K., Munroe, J., Nikurashin, M., Oke, P. R., Pilo, G. S., Richet, O., Savita, A., Spence, P., Stewart, K. D., Ward, M. L., Wu, F., and Zhang, X.: ACCESS-OM2 v1.0: A global ocean–sea ice model at three resolutions, *Geosci. Model Dev.*, 13, 401–442, <https://doi.org/10.5194/gmd-13-401-2020>, 2020.
- 1930 Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoka, K., and Takahashi, K.: The JRA-55 reanalysis: General specifications and basic characteristics, *J. Meteor. Soc. Jpn*, 93, 5–48, <https://doi.org/10.2151/jmsj.2015-001>, 2015.
- 1935 Large, W. G. and Yeager, S. G.: Diurnal to decadal global forcing for ocean and sea-ice models: the data sets and flux climatologies, NCAR Technical Note, Boulder, Colorado, 112 pp., <https://doi.org/10.5065/D6KK98Q6>, 2004.
- Large, W. G. and Yeager, S. G.: The global climatology of an interannually varying air – sea flux data set, *Clim. Dynam.*, 33, 341–364, <https://doi.org/10.1007/s00382-008-0441-3>, 2009.
- 1940 Large, W. G., McWilliams, J. C., and Doney, S. C.: Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parameterization, *Rev. Geophys.*, 32, 363–403, <https://doi.org/10.1029/94RG01872>, 1994.
- Large, W. G., Patton, E. G., and Sullivan, P. P.: The diurnal cycle of entrainment and detrainment in LES of the Southern Ocean driven by observed surface fluxes and waves, *J. Phys. Oceanogr.*, 51, 3253–3278, <https://doi.org/10.1175/JPO-D-20-0308.1>, 2021.
- 1945 Larsen, J. C. and Sanford, T. B.: Florida current volume transports from voltage measurements, *Science*, 227, 302–304, <https://doi.org/10.1126/science.227.4684.302>, 1985.



- 1950 Lee, H.-C., Rosati, A., and Spelman, M. J.: Barotropic tidal mixing effects in a coupled climate model: Oceanic conditions in the Northern Atlantic, *Ocean Model.*, 11, 464–477, <http://dx.doi.org/10.1016/j.ocemod.2005.03.003>, 2006.
- 1955 Legg, S., Briegleb, B., Chang, Y., Chassignet, E. P., Danabasoglu, G., Ezer, T., Gordon, A. L., Griffies, S., Hallberg, R., Jackson, L., Large, W., Ozgokmen, T. M., Peters, H., Price, J., Riemenschneider, U., Wu, W., Xu, X., and Yang, J.: Improving oceanic overflow representation in climate models: The gravity current entrainment climate process team, *Bull. Amer. Meteor. Soc.*, 90, 657–670, <https://doi.org/10.1175/2008BAMS2667.1>, 2009.
- 1960 Li, F., Lozier, M. S., and Johns, W. E.: Calculating the meridional volume, heat and freshwater transports from an observing system in the subpolar North Atlantic: Observing system simulation experiment, *J. Atmos. Oceanic Tech.*, 34, 1483–1500, <https://doi.org/10.1175/JTECH-D-16-0247.1>, 2017.
- 1965 Li, F., Lozier, M. S., Danabasoglu, G., Holliday, N. P., Kwon, Y.-O., Romanou, A., Yeager, S. G., and Zhang, R.: Local and downstream relationships between Labrador Sea water volume and North Atlantic meridional overturning circulation variability, *J. Climate*, 32, 3883–3898, <https://doi.org/10.1175/JCLI-D-18-0735.1>, 2019.
- 1970 Li, F., Lozier, M. S., Bacon, S., Bower, A. S., Cunningham, S. A., de Jong, M. F., de Young B., Fraser, N., Fried, N., Han, G., Holliday, N. P., Holte, J., Houptert, L., Inall, M. E., Johns, W. E., Jones, S., Johnson, C., Karstensen, J., Le Bras, I. A., Lherminier, P., Lin, X., Mercier, H., Oltmanns, M., Pacini, A., Petit, T., Pickart, R. S., Rayner, D., Straneo, F., Thierry, V., Visbeck, M., Yashayaev, I., and Zhou, C.: Subpolar North Atlantic western boundary density anomalies and meridional overturning circulation, *Nat. Comm.*, 12:3002, <https://doi.org/10.1038/s41467-021-23350-2>, 2021.
- 1975 Li, Q., Webb, A., Fox-Kemper, B., Craig, A., Danabasoglu, G., Large, W. G., and Vertenstein, M.: Langmuir mixing effects on global climate: WAVEWATCH III in CESM, *Ocean Model.*, 103, 145–160, <https://doi.org/10.1016/j.ocemod.2015.07.020>, 2016.
- 1980 Li, Y., Liu, H., Ding, M., Lin, P., Yu, Z., Yu, Y., Meng, Y., Li, Y., Jian, X., Jiang, J., Chen, K., Yang, Q., Wang, Y., Zhao, B., Wei, J., Ma, J., Zheng, W., and Wang, P.: Eddy-resolving simulation of CAS-LICOM3 for phase 2 of the Ocean Model Intercomparison Project, *Adv. Atmos. Sci.*, 37, 1067–1080, <https://doi.org/10.1007/s00376-020-0057-z>, 2020.
- 1985 Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., Zweng, M. M., Paver, C. R., Reagan, J. R., Johnson, D. R., Hamilton, M., and Seidov, D.: *World Ocean Atlas 2013, Vol. 1: Temperature*, NOAA Atlas NESDIS 73, <https://doi.org/10.7289/V55X26VD>, 2013.
- 1990 Locarnini, R. A., Mishonov, A. V., Baranova, O. K., Reagan, J. R., Boyer, T. P., Seidov, D., Wang, Z., Garcia, H. E., Bouchard, C., Cross, S. L., and Paver, C. R.: *World Ocean Atlas 2023, volume 1: Temperature*, 2024.
- 1995 Lovato, T., Peano, D., Butenschön, M., Materia, S., Iovino, D., Scoccimarro, E., Fogli, P. G., Cherchi, A., Bellucci, A., Gualdi, S., and Masina, S.: CMIP6 simulations with the CMCC Earth System Model (CMCC-ESM2), *J. Adv. Model. Earth Sy.*, 14, e2021MS002814, <https://doi.org/10.1029/2021MS002814>, 2022.
- 1999 Lozier, M. S.: Overturning in the subpolar North Atlantic: A review, *Philos. Trans. R. Soc. A*, 381, <https://doi.org/10.1098/rsta.2022.0191>, 2023.



- 1995 Lozier, M. S., Bacon, S., Bower, A. S., Cunningham, S. A., de Jong, M. F., de Steur, L., DeYoung, B., Fischer, J., Gary, S. F., Greenan, B. J. W., Heimbach, P., Holliday, N. P., Houpert, L., Inall, M. E., Johns, W. E., Johnson, H. L., Karstensen, J., Li, F., Lin, X., Mackay, N., Marshall, D. P., Mercier, H., Myers, P. G., Pickart, R. S., Pillar, H. R., Straneo, F., Thierry, V., Weller, R. A., Williams, R. G., Wilson, C., Yang, J., Zhao, J., and Zika, J. D.: Overturning in the Subpolar North Atlantic Program: A new international ocean observing system, *Bull. Amer. Meteor. Soc.*, 98, 737–752, <https://doi.org/10.1175/BAMS-D-16-0057.1>, 2017.
- 2000 Lozier, M. S., Li, F., Bacon, S., Bahr, F., Bower, A. S., Cunningham, S. A., de Jong, M. F., de Steur, L., DeYoung, B., Fischer, J., Gary, S. F., Greenan, B. J. W., Holliday, N. P., Houk, A., Houpert, L., Inall, M. E., Johns, W. E., Johnson, H. L., Johnson, C., Karstensen, J., Koman, G., Le Bras, I. A., Lin, X., Mackay, N., Marshall, D. P., Mercier, H., Olmanns, M., Pickart, R. S., Ramsey, A. L., Rayner, D., Straneo, F., Thierry, V., Torres, D. J., Williams, R. G., Wilson, C., Yang, J., Yashayaev, I., and Zhao, J.: A sea change in our view of overturning in the subpolar North Atlantic, *Science*, 363, 516–521, doi: 10.1126/science.aau6592, 2019.
- 2005 Lozier, M. S., Bower, A. S., Furey, H. H., Drouin, K. L., Xu, X., and Zhou, S.: Overflow water pathways in the North Atlantic, *Prog. Oceanogr.*, 208, 102874, <https://doi.org/10.1016/j.pcean.2022.102874>, 2022.
- 2010 Lynn, R. J. and Reid, J. L.: Characteristics of circulation of deep and abyssal waters, *Deep-Sea Res.*, 15, 577–598, 1968.
- 2015 Mackallah, C., Chamberlain, M. A., Law, R. M., Dix, M., Ziehn, T., Bi, D., Bodman, R., Brown, J. R., Dobrohotoff, P., Druken, K., Evans, B., Harman, I. N., Hayashida, H., Holmes, R., Kiss, A. E., Lenton, A., Liu, Y., Marsland, S., Meissner, K., Menviel, L., O’Farrell, S., Rashid, H. A., Ridzwan, S., Savita, A., Srbinovsky, J., Sullivan, A., Trenham, C., Vohralik, P. F., Wang, Y.-P., Williams, G., Woodhouse, M. T., and Yeung, N.: ACCESS datasets for CMIP6: Methodology and idealised experiments. *J. Southern Hemisphere Earth Sy. Sci.*, 72, 93–116, <https://doi.org/10.1071/ES21031>, 2022.
- Madec, G. and Imbard, M.: A global ocean mesh to overcome the North Pole singularity, *Clim. Dyn.*, 12, 381–388, 1996.
- 2020 Madec, G. and the NEMO Team: NEMO ocean engine - version 3.6. Note du Pôle de modélisation, Institut Pierre-Simon Laplace, 27, 2016.
- Madec, G. and System Team: Nemo Ocean Engine - version 4.0.1, Notes du Pole de modelisation de l’Institut Pierre-Simon Laplace (IPSL), 27, <https://doi.org/10.5281/zenodo.3878122>, 2019.
- 2025 Marques, G. M., Shao, A. E., Bachman, S. D., Danabasoglu, G., and Bryan, F. O.: Representing eddy diffusion in the surface boundary layer of ocean models with general vertical coordinates, *J. Adv. Model. Earth Syst.*, 15, e2023MS003751, <https://doi.org/10.1029/2023MS003751>, 2023.
- 2030 Marshall, D. P., Maddison, J. R., and Berloff, P. S.: A framework for parameterizing eddy potential vorticity fluxes, *J. Phys. Oceanogr.*, 42, 539–557, <https://doi.org/10.1175/JPO-D-11-048.1>, 2012.
- Masina, S., Cipollone, A., Iovino, D., Ciliberti, S., Coppini, G., Lecci, R., Creti, S., Palermo, F., Viola, F., Lyubartsev, V., Baordo, F., Trotta, F., and Pinardi, N.: A global ocean eddying forecasting system at 1/16°, in *Advances in operational oceanography: expanding Europe's*



- 2035 ocean observing and forecasting capacity. Proceedings of the 9th EuroGOOS International Conference, EuroGOOS, Brussels, Belgium, 3–5 May 2021, 524–531, <https://doi.org/10.13155/83160>, 2021.
- Mauritzen, C., Price, J., Sanford, T., and Torres, D.: Circulation and mixing in the Faroese Channels, *Deep-Sea Res. I*, 52, 883–913, <https://doi.org/10.1016/j.dsr.2004.11.018>, 2005.
- 2040 McDougall, T. J., and Barker, P. M.: Getting started with TEOS-10 and the Gibbs Seawater (GSW) Oceanographic Toolbox, 28pp., SCOR/IAPSO WG127, ISBN 978-0-646-55621-5, 2011.
- Meehl, G. A., Goddard, L., Boer, G., Burgman, R., Branstator, G., Cassou, C., Corti, S., Danabasoglu, G., Doblas-Reyes, F., Hawkins, E., Karspeck, A., Kimoto, M., Kumar, A., 2045 Matei, D., Mignot, J., Msadek, R., Navarra, A., Pohlmann, H., Rienecker, M., Rosati, A., Schneider, E., Smith, D., Sutton, R., Teng, H., van Oldenborgh, G. J., Vecchi, G., and Yeager, S.: Decadal climate prediction: An update from the trenches, *Bull. Amer. Meteor. Soc.*, 95, 243–267, <https://doi.org/10.1175/BAMS-D-12-00241.1>, 2014.
- Megann, A., Blaker, A., Josey, S., New, A., and Sinha, B.: 2021. Mechanisms for late 20th and 2050 early 21st century decadal AMOC variability, *J. Geophys. Res.: Oceans*, 126, e2021JC017865, <https://doi.org/10.1029/2021JC017865>, 2021.
- Meinen, C. S., Baringer, M. O., and Garcia, R. F.: Florida current transport variability: An analysis of annual and longer-period signals, *Deep Sea Research Part I: Oceanographic Res. Papers*, 57, 835–846, <https://doi.org/10.1016/j.dsr.2010.04.001>, 2010.
- 2055 Meinen, C. S., Speich, S., Piola, A. R., Ansorge, I., Campos, E., Kersalé, M., Terre, T., Chidichimo, M. P., Lamont, T., Sato, O. T., Perez, R. C., Valla, D., van den Berg, M., Le Henaff, M., Dong, S., and Garzoli, S.: Meridional overturning circulation transport variability at 34.5°S during 2009–2017: Baroclinic and barotropic flows and the dueling influence of the boundaries, *Geophys. Res. Lett.*, 45, 4180–4188, 2060 <https://doi.org/10.1029/2018GL077408>, 2018.
- Menary, M. B., Jackson, L. C., and Lozier, M. S.: Reconciling the relationship between the AMOC and Labrador Sea in OSNAP observations and climate models, *Geophys. Res. Lett.*, 47, e2020GL089793, <https://doi.org/10.1029/2020GL089793>, 2020.
- Morlighem, M.: MEaSUREs BedMachine Antarctica (NSIDC-0756, Version 3), NASA National 2065 Snow and Ice Data Center Distributed Active Archive Center, <https://doi.org/10.5067/FPSU0V1MWUB6>, 2022.
- Msadek, R., Dixon, K. W., Delworth, T. L., and Hurlin, W.: Assessing the predictability of the Atlantic meridional overturning circulation and associated fingerprints, *Geophys. Res. Lett.*, 37, L19608, <https://doi.org/10.1029/2010GL044517>, 2010.
- 2070 Murray, R. J.: Explicit generation of orthogonal grids for ocean models. *J. Comput. Phys.*, 126(2), 251–273, <http://dx.doi.org/10.1006/jcph.1996.0136>, 1996.
- NOAA: available at: <https://www.ngdc.noaa.gov/mgg/global/etopo2.html> (last access: 6 August 2018), 2006.
- 2075 Ortega, P., Robson, J. I., Menary, M., Sutton, R. T., Blaker, A., Germe, A., Hirschi, J. J.-M., Sinha, B., Hermanson, L., and Yeager, S.: Labrador Sea subsurface density as a precursor of multidecadal variability in the North Atlantic: A multi-model study, *Earth Syst. Dynam.*, 12, 419–438, <https://doi.org/10.5194/esd-12-419-2021>, 2021.



- Pacanowski, R. C., and Gnanadesikan, A.: Transient response in a z-level ocean model that resolves topography with partial cells, *Monthly Weather Rev.*, 126, 3248–3270, [http://dx.doi.org/10.1175/1520-0493\(1998\)126<3248:TRIAZL>2.0.CO;2](http://dx.doi.org/10.1175/1520-0493(1998)126<3248:TRIAZL>2.0.CO;2), 1998.
- Petersen, M. R., Jacobsen, D. W., Ringler, T. D., Hecht, M. W., and Maltrud, M. E.: Evaluation of the arbitrary Lagrangian–Eulerian vertical coordinate method in the MPAS-Ocean model, *Ocean Model.*, 86, 93–113, 2015.
- Petit, T., Lozier, M. S., Josey, S. A., and Cunningham, S. A.: Atlantic deep water formation occurs primarily in the Iceland Basin and Irminger Sea by local buoyancy forcing, *Geophys. Res. Lett.*, 47, e2020GL091028, <https://doi.org/10.1029/2020GL091028>, 2020.
- Petit, T., Lozier, M. S., Rühls, S., Handmann, P., and Biastoch, A.: Propagation and transformation of upper North Atlantic Deep Water from the subpolar gyre to 26.5°N, *J. Geophys. Res. Oceans*, 128, e2023JC019726, <https://doi.org/10.1029/2023JC019726>, 2023.
- Pickart, R. S. and Spall, M. A.: Impact of Labrador Sea convection on the North Atlantic meridional overturning circulation, *J. Phys. Oceanogr.*, 37, 2207–2227, <https://doi.org/10.1175/JPO3178.1>, 2007.
- Reagan, J. R., Seidov, D., Wang, Z., Dukhovskoy, D., Boyer, T. P., Locarnini, R. A., Baranova, O. K., Mishonov, A. V., Garcia, H. E., Bouchard, C., and Cross, S. L.: *World Ocean Atlas 2023, volume 2: Salinity*, 2024.
- Redi, M. H.: Oceanic isopycnal mixing by coordinate rotation, *J. Phys. Oceanogr.*, 12, 1154–1158, [https://doi.org/10.1175/1520-0485\(1982\)012<1154:OIMBCR>2.0.CO;2](https://doi.org/10.1175/1520-0485(1982)012<1154:OIMBCR>2.0.CO;2), 1982.
- Reichl, B. G. and Hallberg, R.: A simplified energetics based planetary boundary layer (ePBL) approach for ocean climate simulations, *Ocean Model.*, 132, 112–129, <https://doi.org/10.1016/j.ocemod.2018.10.004>, 2018.
- Riemenschneider, U. and Legg, S.: Regional simulations of the Faroe Bank Channel overflow in a level model, *Ocean Model.*, 17, 93–122, <https://doi.org/10.1016/j.ocemod.2007.01.003>, 2007.
- Rousset, C., Vancoppenolle, M., Madec, G., Fichefet, T., Flavoni, S., Barthélemy, A., Benshila, R., Chanut, J., Levy, C., Masson, S., and Vivier, F.: The Louvain-La-Neuve sea ice model LIM3.6: Global and regional capabilities, *Geosci. Model Dev.*, 8, 2991–3005, <https://doi.org/10.5194/gmd-8-2991-2015>, 2015.
- Rühls, S., Oliver, E., Biastoch, A., Böning, C. W., Dowd, M., Getzlaff, K., and Myers, P. G.: Changing spatial patterns of deep convection in the subpolar North Atlantic, *J. Geophys. Res.*, 126, e2021JC017245, <https://doi.org/https://doi.org/10.1029/2021JC017245>, 2021.
- Send, U., Lankhorst, M., and Kanzow, T.: Observation of decadal change in the Atlantic meridional overturning circulation using 10 years of continuous transport data, *Geophys. Res. Lett.*, 38, 24606, <https://doi.org/10.1029/2011GL049801>, 2011.
- Shao, A. E., Adcroft, A., Hallberg, R., and Griffies, S. M.: A general-coordinate, nonlocal neutral diffusion operator, *J. Adv. Model. Earth Syst.*, 12, e2019MS001992, <https://doi.org/10.1029/2019MS001992>, 2020.
- Shchepetkin, A. F. and McWilliams, J. C.: A method for computing horizontal pressure-gradient force in an oceanic model with a nonaligned vertical coordinate, *J. Geophys. Res.*, 108(C3), 3090, doi:10.1029/2001JC001047, 2003.



- 2120 Shchepetkin, A. F. and McWilliams, J. C.: The regional oceanic modeling system (ROMS): A split-explicit, free-surface, topography following-coordinate oceanic model, *Ocean Model.*, 9, 347–404, <https://doi.org/10.1016/j.ocemod.2004.08.002>, 2005.
- Simmons, H. L., Jayne, S. R., Laurent, L. C. S., and Weaver, A. J.: Tidally driven mixing in a numerical model of the ocean general circulation, *Ocean Model.*, 6, 245–263, [https://doi.org/10.1016/S1463-5003\(03\)00011-8](https://doi.org/10.1016/S1463-5003(03)00011-8), 2004.
- 2125 Small, R. J., Bacmeister, J., Bailey, D., Baker, A., Bishop, S., Bryan, F., Caron, J., Dennis, J., Gent, P., Hsu, H., Jochum, M., Lawrence, D., Muñoz, E., DiNezio, P., Scheitlin, T., Tomas, R., Tribbia, J., Tseng, Y., and Vertenstein, M.: A new synoptic scale resolving global climate simulation using the Community Earth System Model, *J. Adv. Model. Earth Sys.*, 6, 1065–1094, <https://doi.org/10.1002/2014MS000363>, 2014.
- 2130 Smith, K., Barthel, A. M., Conlon, L. M., Van Roekel, L. P., Bartoletti, A., Golez, J. C., Zhang, C., Begeman, C. B., Benedict, J. J., Bisht, G., and Feng, Y.: The DOE E3SM version 2.1: Overview and assessment of the impacts of parameterized ocean submesoscales, *Geosci. Model Dev. Discussions*, 1–38, 2024.
- 2135 Smith, R., Jones, P., Briegleb, B., Bryan, F., Danabasoglu, G., Dennis, J., Dukowicz, J., Eden, C., Fox-Kemper, B., Gent, P., Hecht, M., Jayne, S., Jochum, M., Large, W., Maltrud, M., Norton, N., Peacock, S., Vertenstein, M., and Yeager, S.: The parallel ocean program (POP) reference manual: Ocean component of the community climate system model (CCSM) and Community Earth System Model (CESM), LANL Tech. Rep. LAUR-01853, 141 pp., 2010.
- 2140 Solodoch, A., Stewart, A. L., Hogg, A. M., Morrison, A. K., Kiss, A. E., Thompson, A. F., Purkey, S. G., and Cimoli, L.: How does Antarctic Bottom Water cross the Southern Ocean? *Geophys. Res. Lett.*, 49, e2021GL097211, <https://doi.org/10.1029/2021GL097211>, 2022.
- Spall, M. A.: Boundary currents and watermass transformation in marginal seas, *J. Phys. Oceanogr.*, 34, 1197–1213, 2004.
- 2145 Stanley, Z., Grooms, I., Kleiber, W., Bachman, S. D., Castruccio, F., and Adcroft, A.: Parameterizing the impact of unresolved temperature variability on the large-scale density field: Part 1. Theory, *J. Adv. Model. Earth Syst.*, 12, e2020MS002185, 2020.
- Stewart, K. D., Hogg, A. M., Griffies, S. M., Heerdegen, A. P., Ward, M. L., Spence, P., and England, M. H.: Vertical resolution of baroclinic modes in global ocean models, *Ocean Model.*, 113, 50–65, 2017.
- 2150 Straneo, F.: On the connection between dense water formation, overturning, and poleward heat transport in a convective basin, *J. Phys. Oceanogr.*, 36, 1822–1840, <https://doi.org/10.1175/JPO2932.1>, 2006.
- Sun, Q., Whitney, M. M., Bryan, F. O., and Tseng, Y.: Assessing the skill of the improved treatment of riverine freshwater in the Community Earth System Model (CESM) relative to a new salinity climatology, *J. Adv. Model. Earth Sys.*, 11, 1189–1206, <https://doi.org/10.1029/2018MS001349>, 2019.
- 2155 Sutton, R. T., McCarthy, G. D., Robson, J., Sinha, B., Archibald, A. T., and Gray, L. J.: Atlantic multidecadal variability and the U.K. ACSIS program, *Bull. Amer. Meteor. Soc.*, 99, 415–425, <https://doi.org/10.1175/BAMS-D-16-0266.1>, 2018.
- 2160 Suzuki, T., Yamazaki, D., Tsujino, H., Komuro, Y., Nakano, H., and Urakawa, S.: A dataset of continental river discharge based on JRA-55 for use in a global ocean circulation model, *J. Oceanogr.*, 74, 421–429, <https://doi.org/10.1007/s10872-017-0458-5>, 2018.



- 2165 Teague, W. J., Carron, M. J., and Hogan, P. J.: A comparison between the Generalized Digital Environmental Model and Levitus climatologies, *J. Geophys. Res.*, 95, 7167–7183, <https://doi.org/10.1029/jc095ic05p07167>, 1990.
- Timmermann, R., Goosse, H., Madec, G., Fichefet, T., Etche, C., and Dulière, V.: On the representation of high latitude processes in the ORCA-LIM global coupled sea ice-ocean model, *Ocean Model.*, 8, 175–201, 2005.
- 2170 Toole, J. M., Curry, R. G., Joyce, T. M., McCartney, M., and Peña-Molino, B.: Transport of the North Atlantic deep western boundary current about 39°N, 70°W: 2004–2008, *Deep Sea Res. Part II: Topical Studies in Oceanography*, 58, 1768–1780, <https://doi.org/10.1016/j.dsr2.2010.10.058>, 2011.
- 2175 Tozer, B., Sandwell, D. T., Smith, W. H. F., Olson, C., Beale, J. R., and Wessel, P.: Global bathymetry and topography at 15 arc sec: SRTM15+, *Earth and Space Science*, 6, <https://doi.org/10.1029/2019EA000658>, 2019.
- Treguier, A. M., de Boyer Montegut, C., Bozec, A., Chassignet, E. P., Fox-Kemper, B., Hogg, A. M., Iovino, D., Kiss, A. E., Sommer, J. L., Li, Y., Lin, P., Lique, C., Liu, H., Serazin, G., Sidorenko, D., Wang, Q., Xu, X., and Yeager, S.: The mixed-layer depth in the ocean model intercomparison project (OMIP): Impact of resolving mesoscale eddies. *Geosci. Model Dev.*, 16, 3849–3872, <https://doi.org/10.5194/gmd-16-3849-2023>, 2023.
- 2180 Tsujino, H., Urakawa, S., Nakano, H., Small, R. J., Kim, W. M., Yeager, S. G., Danabasoglu, G., Suzuki, T., Bamber, J. L., Bentsen, M., Böning, C. W., Bozec, A., Chassignet, E. P., Curchitser, E., Dias, F. B., Durack, P. J., Griffies, S. M., Harada, Y., Ilıcak, M., Josey, S. A., Kobayashi, C., Kobayashi, S., Komuro, Y., Large, W. G., Sommer, J. L., Marsland, S. J., Masina, S., Scheinert, M., Tomita, H., Valdivieso, M., and Yamazaki, D.: JRA-55 based surface dataset for driving ocean - sea-ice models (JRA55-do), *Ocean Model.*, 130, 79–139, <http://dx.doi.org/10.1016/j.ocemod.2018.07.002>, 2018.
- 2185 Tsujino, H., Urakawa, L. S., Griffies, S. M., Danabasoglu, G., Adcroft, A. J., Amaral, A. E., Arsouze, T., Bentsen, M., Bernardello, R., Böning, C. W., Bozec, A., Chassignet, E. P., Danilov, S., Dussin, R., Exarchou, E., Fogli, P. G., Fox-Kemper, B., Guo, C., Ilıcak, M., Iovino, D., Kim, W. M., Koldunov, N., Lapin, V., Li, Y., Lin, P., Lindsay, K., Liu, H., Long, M. C., Komuro, Y., Marsland, S. J., Masina, S., Nummelin, A., Rieck, J. K., Ruprich-Robert, Y., Scheinert, M., Sicardi, V., Sidorenko, D., Suzuki, T., Tatebe, H., Wang, Q., Yeager, S. G., and Yu, Z.: Evaluation of global ocean-sea model simulations based on the experimental protocols of the Ocean Model Intercomparison Project phase 2 (OMIP-2), *Geosci. Model Dev.*, 13, 3643–3708, <https://doi.org/10.5194/gmd-13-3643-2020>, 2020.
- 2195 Turner, A. K. and Hunke, E. C.: Impacts of a mushy-layer thermodynamic approach in global sea-ice simulations using the CICE sea-ice model, *J. Geophys. Res.-Oceans*, 120, 1253–1275, <https://doi.org/10.1002/2014JC010358>, 2015.
- 2200 Uotila, P., Iovino, D., Vancoppenolle, M., Lensu, M., and Rousset, C.: Comparing sea ice, hydrography and circulation between NEMO3.6 LIM3 and LIM2, *Geosci. Model Dev.*, 10, 1009–1031, <https://doi.org/10.5194/gmd-10-1009-2017>, 2017.
- 2205 Valcke, S., Craig, T., and Coquart, L.: OASIS3-MCT User Guide: OASIS3-MCT 2.0. CERFACS/CNRS SUC URA No1875, CERFACS TR/CMGC/13/17, CERFACS/CNRS, 2013.



- van Westen, R. M., Kliphuis, M., and Dijkstra, H. A.: Physics-based early warning signal shows that AMOC is on tipping course, *Sci. Adv.*, 10(6), eadk1189, <https://doi.org/10.1126/sciadv.adk1189>, 2024.
- 2210 Wang, Q., Shu, Q., Bozec, A., Chassignet, E. P., Fogli, P. G., Fox-Kemper, B., Hogg, A. M., Iovino, D., Kiss, A. E., Koldunov, N., Le Sommer, J., Li, Y., Lin, P., Liu, H., Polyakov, I., Scholz, P., Sidorenko, D., Wang, S., and Xu, X.: Impact of increased resolution on Arctic Ocean simulations in Ocean Model Intercomparison Project phase 2 (OMIP-2), *Geosci. Model Dev.*, 17, 347–379. <https://doi.org/10.5194/gmd-17-347-2024>, 2024.
- 2215 Weatherall, P., Marks, K. M., Jakobsson, M., Schmitt, T., Tani, S., Arndt, J. E., Rovere, M., Chayes, D., Ferrini, V., and Wigley, R.: A new digital bathymetric model of the world's oceans. *Earth and Space Science*, 2, 331–345, <https://doi.org/10.1002/2015EA000107>, 2015.
- Weijer, W., Cheng, W., Garuba, O. A., Hu, A., and Nadiga, B. T.: CMIP6 models predict significant 21st century decline of the Atlantic Meridional Overturning Circulation, *Geophys. Res. Lett.*, 47, e2019GL086075. <https://doi.org/10.1029/2019GL086075>, 2020.
- 2220 Wright, D. G.: An equation of state for use in ocean models: Eckart's formula revisited, *J. Atm. Ocean. Tech.*, 14, 735–740, [https://doi.org/10.1175/1520-0426\(1997\)014<0735:AEOSFU>2.0.CO;2](https://doi.org/10.1175/1520-0426(1997)014<0735:AEOSFU>2.0.CO;2), 1997.
- Xu, X., Chassignet, E. P., Firing, Y. L., and Donohue, K.: Antarctic Circumpolar Current transport through Drake Passage: What can we learn from comparing high-resolution model results to observations? *J. Geophys. Res. Oceans*, 125, e2020JC016365, <https://doi.org/10.1029/2020JC016365>, 2020.
- 2225 Xu, X., Chassignet, E. P., Dong, S., and Baringer, M. O.: Transport structure of the South Atlantic Ocean derived from a high-resolution numerical model and observations, *Front. Mar. Sci.*, 9, 811398, <https://doi.org/10.3389/fmars.2022.811398>, 2022.
- 2230 Xu, X., Chassignet, E. P., and Wallcraft, A. J.: Impact of vertical resolution on representing baroclinic modes and water mass distribution in the North Atlantic, *Ocean Model.*, 186, <https://doi.org/10.1016/j.ocemod.2023.102261>, 2023.
- 2235 Yang, J., Li, J., and An, Q.: Deep Atlantic multidecadal variability, *Geophys. Res. Lett.*, 51, e2023GL106367, <http://dx.doi.org/10.1029/2023GL106367>, 2024.
- Yeager, S. and Danabasoglu, G.: The origins of late 20th century variations in the large-scale North Atlantic circulation, *J. Climate*, 27, 3222–3247, <https://doi.org/10.1175/JCLI-D-13-00125.1>, 2014.
- 2240 Yeager, S., Castruccio, F., Chang, P., Danabasoglu, G., Maroon, E., Small, J., Wang, H., Wu, L., and Zhang, S.: An outsized role for the Labrador Sea in the multidecadal variability of the Atlantic overturning circulation, *Sci. Adv.*, 7, eabh3592, <https://doi.org/10.1126/sciadv.abh3592>, 2021.
- Zalesak, S. T.: Fully multidimensional flux corrected transport algorithms for fluids, *J. Comput. Phys.*, 31, 335–362, 1979.
- 2245 Zhang, R. and Thomas, M.: Horizontal circulation across density surfaces contributes substantially to the long-term mean northern Atlantic meridional overturning circulation, *Comm. Earth Environment*, 2:112, <https://doi.org/10.1038/s43247-021-00182-y>, 2021.
- Zhang, R., Sutton, R., Danabasoglu, G., Kwon, Y.-O., Marsh, R., Yeager, S. G., Amrhein, D. E., and Little, C. M.: A review of the role of Atlantic meridional overturning circulation in



- 2250 Atlantic multidecadal variability and associated climate impacts, *Rev. Geophys.*, 57, 316–375, <https://doi.org/10.1029/2019RG000644>, 2019.
- Zou, S., Lozier, M. S., Li, F., Abernathey, R., and Jackson, L.: Density-compensated overturning in the Labrador Sea, *Nat. Geosci.*, 13, 121–126, <https://doi.org/10.1038/s41561-019-0517-1>, 2020.
- 2255 Zweng, M., Reagan, J., Antonov, J., Locarnini, R., Mishonov, A., Boyer, T., Garcia, H., Baranova, O., Johnson, D., Seidov, D., and Biddle, M.: World Ocean Atlas 2013, Vol. 2: Salinity, NOAA Atlas NESDIS 74, 39 pp., <https://doi.org/10.7289/V5251G4D>, 2013.