

Measurement Report: Seasonal trends and chemical speciation of chromium-(III/VI) in different fractions of urban particulate matter – a case study of Radom, Poland

Monika Łożyńska¹, Marzena Trojanowska², Artur Molik², Ryszard Świetlik²

¹Łukasiewicz Research Network – Institute for Sustainable Technologies, Bioeconomy and Ecoinnovation Centre, 26-600 Radom, Pulaskiego 6/10, Poland

²Casimir Pulaski Radom University, Faculty of Applied Chemistry, 26-600 Radom, Chrobrego 27, Poland

Correspondence to: Łożyńska Monika (monika.lozynska@itec.lukasiewicz.gov.pl)

Abstract. The paper assesses chromium occurrence in urban particulate matter: PM₁₀, PM_{2.5}, PM₄, and PM_{0.25} over a calendar year. The seasonality of both pseudo-total chromium content and its valence speciation are studied. Airborne aerosol was sampled in the city of Radom (Poland) using a Hi-Flow MOUDI Model 130 cascade impactor from Copley. The pseudo-total chromium concentration (Cr_{tot}) was assayed with GF-AAS and the concentration of Cr(VI) with CCSV-DTPA techniques. Cr_{tot} concentrations in the particulate matter fractions investigated ranged from 0.08 to 4.09 ng/m³. The results point to a seasonality of Cr_{tot} concentration changes in particulate matter. The concentration Cr_{tot} in PM₁₀ was maximum in winter (2.23±0.53 ng/m³ on average), while-and Cr_{tot} in PM₁₀ averaged 1.71±0.83 ng/m³ in the whole measurement period. The average Cr(VI) concentration did not exceed 0.40 ng/m³ and was maximum in winter, too (max. 1.354 ng/m³). The Cr(VI) share in PM in the particular seasons varied a lot-It was_-minimum in summer_(9.1% of Cr_{tot}) and maximum in winter (40% of Cr_{tot}). The carcinogenic risk for the urban residents based on the Cr(VI) concentration in PM₁₀ was below the US EPA threshold for significant risk (between 1·10⁻⁶ and 1·10⁻⁴) and amounted to between 1.11·10⁻⁶ and 5.78·10⁻⁶ for children and from 3.69·10⁻⁶ to 1.92·10⁻⁵ for adults. The non-carcinogenic health risk caused by the presence of Cr_{tot} was also lower than the safe level of 1 - the HQ values for both adults and children ranged from 1.32·10⁻² to 3.92·10⁻².The carcinogenic risk for the urban residents based on the Cr(VI) concentration in PM₁₀ was within the acceptable range (between 1·10⁻⁶ and 1·10⁻⁴) and amounted to between 1.11·10⁻⁶ and 5.78·10⁻⁶ for children and from 3.69·10⁻⁶ to 1.92·10⁻⁵ for adults. The non-carcinogenic health risk caused by the presence of Cr_{tot} was acceptable as well. The HQ values for both adults and children were lower than the safe level of 1 and ranged from 1.57·10⁻² to 3.92·10⁻².

1 Introduction

Air quality has continued to deteriorate lately, especially in urban and industrial areas. Particulate matter (PM) is a major pollutant whose substantial quantities are emitted into the air, affecting adversely both the climate and all parts of the environment (Fang et al., 2005; Wagner et al., 2008; Arhami et al., 2017). Particulate matter, considered one of the most

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks górny

— sformatowano: Indeks górny

— sformatowano: Indeks górny

— sformatowano: Indeks górny

— sformatowano: Indeks górny

— sformatowano: Indeks górny

— sformatowano: Indeks dolny

— sformatowano: Indeks górny

— sformatowano: Indeks górny

dangerous environment pollutants (Park et al., 2009; Waheed et al., 2011; Izhar et al., 2016), has a complex and heterogeneous composition, dependent on the season, source of emissions, and weather conditions. Knowledge of the composition, concentration, and sources of particulate matter suspended in the air is of great importance ~~to residents of urban-industrial areasto-city-and-agglomeration-residents~~, since breathing these particles ~~in~~ can increase mortality or morbidity due to respiratory and pulmonary conditions. The particulate matter's coarse fraction (2.5-10 µm) is assumed to be of natural origin, while the fine fraction (0.1–2.5 µm, especially the particles of less than 1 µm) comes from anthropogenic processes (Nocoń et al., 2018). The distribution of particulate matter particle sizes and its loading with different chemical compounds determine the aerosol's effect on human health and the environment. ~~The p~~Particulate matter with grain diameters below 2.5 µm is particularly dangerous to living organisms as it penetrates alveoli pulmonis and thence potentially enters the bloodstream (Wagner et al., 2008; Pan et al., 2015). For this reason, this particulate matter fraction is the most common object of environmental studies (Feng et al., 2009; Rogula- Kozłowska et al., 2013a; Zajusz-Zubek and Mainka, 2015; Li et al., 2016; Sah et al., 2019; Xie et al., 2019; Anake et al., 2020; Wu et al., 2021; Wang et al., 2023;) (Table S1, Supplementary Material). Researchers frequently examine ~~PM₁₀PM₁₀~~ too (e.g. Richter et al., 2007; Catrambone et al., 2013; Huang et al., 2014a; Pandey et al., 2017; Rubio et al., 2018; Conca et al., 2020). The finer fractions, e.g. ~~PM₁~~, are investigated far more seldom (Rogula-Kozłowska et al., 2013b; Zajusz-Zubek et al., 2015; Zajusz-Zubek, 2017). Scientists rarely assay two (~~PM₁₀PM₁₀~~ and ~~PM_{2.5}~~; Canepari et al., 2009; Dos Santos et al., 2009; Jan et al., 2018; Gunchin et al., 2021) or more grain fractions (~~PM₁~~, ~~PM_{2.5}~~, and ~~PM₁₀PM₁₀~~; Samara and Voutsas, 2005; Rogula-Kozłowska et al., 2013b; Rogula-Kozłowska et al., 2015) in parallel. ~~Various organic and inorganic compounds are also transported together with the particulate matter, some of which are toxic. A variety of toxic compounds, both organic and inorganic, are also transported along with the particulate matter.~~ Special attention is paid to heavy metals (Wagner et al., 2008; Pan et al., 2015; Samara et al., 2016) that may contribute to oxidative DNA damage and cause carcinogenic lesions in effect (Somers, 2011; IARC, 2012; Arhami et al., 2017). ~~The element Chromium occurs in the air in two valence states: Cr(III) and Cr(VI), greatly varying in their physical and chemical properties and toxicity. Chromium-(III) is a microelement necessary for living organisms, whereas chromium-(VI) is toxic and classified as a carcinogen (Katz, 1991; Barceloux, 1999; Kotaś and Stasicka, 2000).~~ Epidemiological research has shown a close connection between chromium(VI) exposure and lung cancer (IARC, 2012). Chromium air presence has been studied a lot given its ~~harmfulness-toxicity~~ (Nriagu and Nieboer, 1988; Nusko and Heumann, 1997; Świetlik et al., 2011; Tirez et al., 2011; Torkmahalleh et al., 2013; Huang et al., 2014a, 2014b; Kang et al., 2016; Widziewicz et al., 2016; Molik et al., 2018; Nocoń et al., 2018). ~~The element occurs in the air in two valence states: Cr(III) and Cr(VI), greatly varying in their physical and chemical properties and toxicity. Chromium (III) is a microelement necessary for living organisms, whereas chromium (VI) is toxic and classified as a carcinogen (Katz, 1991; Barceloux, 1999; Kotaś and Stasicka, 2000).~~ Regionally, natural sources account for 30-40% of total chromium emissions (Pacyna, 1986; Kotaś and Stasicka, 2000). Authors report– that fuel combustion in stationary sources is the chief source of Cr emissions in Europe and more than a half of all anthropogenic emissions (Pacyna et al., 2007). In Poland, the burning of fuels is responsible for 60% of all the Cr emissions to the environment (Ministry of Climate and Environment, 2020).

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

65 The natural air content of Cr_{tot} is estimated to range from 0.1 ng/m^3 to 1 ng/m^3 (Nriagu et al., 1988). Nriagu et al. (1988) report an average atmospheric concentration of Cr starting from 1 ng/m^3 in rural areas to 10 ng/m^3 in polluted urban areas. Świetlik and Trojanowska (2022) review the average Cr in the urban air of diverse regions globally. Given the commonly encountered pollution levels worldwide, Cr_{tot} in the urban air averaged $13\pm2\text{ ng/m}^3$, whereas Cr_{tot} in heavily polluted areas was many times higher, $350\pm70\text{ ng/m}^3$ (Świetlik and Trojanowska, 2022). Average Cr(VI) concentration is far lower – 0.5 ng/m^3 (Seinfeld and
70 Pandis, 2006). It is normally higher close to industrial sources (Proctor et al., 2021). According the US EPA National Air Toxics Assessment in 2017, the median, mean, and maximum Cr(VI) were 0.03 ng/m^3 , 0.1 ng/m^3 and 3.18 ng/m^3 , respectively (Proctor et al., 2021).
To assure legal protection against atmospheric chromium in Poland, reference values were set as follows: $20\text{ }\mu\text{g/m}^3$ and $2.5\text{ }\mu\text{g/m}^3$ for Cr(III) and Cr(VI) compounds and $4.6\text{ }\mu\text{g/m}^3$ and $0.4\text{ }\mu\text{g/m}^3$ for Cr(VI) over 1 h and 1 year, respectively (The Minister
75 of the Environment Regulation, 2010).
Although chromium occurrence in urban air has been extensively studied and a range of publications have appeared recently (Catrambone et al., 2013; Widziewicz et al., 2016; Nocoń et al., 2018), investigations of total Cr occurrence and its valence speciation in particulate matter of different particle sizes are still unavailable. We have only examined Cr-(III/VI) speciation in total suspended particulate (TSP) before (Świetlik et al., 2011). Our current study is- designed to: (I) assess chromium
80 occurrence and speciation in urban particulate matter fractions of varied particle sizes (PM_{10} , $PM_{2.5}$, PM_1 , $PM_{0.25}$); (II) examine the fluctuations and seasonality of Cr concentrations over one year, and (III) estimate health risk caused by inhalation exposure to airborne Cr in two different exposure schemes: 1) the risk that comes from the Cr_{tot} ambient concentrations; 2) the risk that comes exclusively from Cr(VI) species. Radom is an interesting location for such research as the atmospheric chromium levels
are chromium environment pollution is a result not only of an aged urban structure relying on private hard coal heating,
85 considerable road transit, and the operation of multiple metal working factories, but also tanneries clustered in the region for more than 70 years.
These are the results of the first-ever hexavalent chromium (Cr(VI)) measurements in the particulate matter of different sizes in the air of a medium-sized city in central Poland.

2 Experimental

90 2.1 Sampling area and the collection of PM samples

The airborne particulate matter was sampled in Radom, a medium-sized town (a population of 196 000, an area of 112 km^2) in central Poland, 100 km south of Warsaw. The local sources of chromium emissions are: road traffic, coal burning in homes, coal-fired municipal heating plants, and multiple metalworks. The sampling points of the suspended airborne particulate matter were located on the University of Radom premises. The area is adjacent to a housing estate with a prevalence of private houses
95 heated with traditional coal furnaces, but not under an immediate impact of local sources of industrial emissions or streets with heavy traffic.

— sformatowano: Indeks górny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

100 The airborne particulate matter was sampled in spring (variable weather: cool, sunny, wet; the heating campaign until the middle of April), summer (sunny, warm weather), autumn (variable weather: sunny, warm or cool, wet; heating campaign from the middle of October), and winter (cold, overcast and wet weather; heating campaign). Averaged weather conditions for each weekly sampling cycle are presented in Table S2 (Supplementary Material). A Hi-Flow MOUDI Model 130 cascade impactor from Copley was used, positioned at a height of 1.5 m above the ground level. It enabled the separation of four grain fractions of atmospheric dust: ~~PM₁₀~~₁₀, PM_{2.5}, PM₁ and PM_{0.25}. The grain fractions of atmospheric particulate matter were sampled from February 2020 to April 2021 in 29 weekly cycles. A total of 116 samples were collected. The sampling time averaged 70 h ± 4 h. The air rate was maintained in the range of 0.35–0.5 m³/min. The airborne particulate matter was collected on cellulose filters (POCH). The filters were weighed before and after sampling, the weight was 0.01 mg (Microbalance MX5 105 Mettler Toledo) in a temperature and relative humidity controlled environment (20±3°C and 50±10%, respectively). Then the filters were stored in a plastic CD-case. Before the analysis, the filters with the deposit were sectioned into four equal parts. The samples were only withdrawn when there was no precipitation.

2.2 The determination of the contents of the particular airborne particulate matter fractions

110 The concentrations of the ~~particular-specific~~ particulate matter fractions were calculated by dividing the difference between filter weight prior to and following the exposure by the mean air flow at the time of atmospheric aerosol sampling.

2.3 The determination of pseudo-total chromium

In order to assay Cr_{tot} by means of graphite furnace atomic absorption spectrometry (GF-AAS)~~GF-AAS~~, the particulate matter samples collected on filters were mineralised using microwave energy. A quarter of the deposited filter was placed in a 100 115 mL Teflon vessel with 5 mL 65% HNO₃ and 3 mL 30% H₂O₂. The samples were digested in a microwave oven (Milestone MLS 1200 Mega) according to the following program: 1) 6 min., 250 W; 2) 1 min., 0W; 3) 6 min., 400 W; 4) 6 min., 650 W; 5) 6 min., 250 W; 6) 5 min., ventilation. After the digestion, the solutions were filtered ~~the solutions~~ through a 0.45 µm Millex-HV syringe filter (Millipore), transferred them into a polypropylene volumetric flask, and diluted to 25.0 mL with deionized water.

120 A Perkin-Elmer 3100 AAS graphite furnace system equipped with HGA 600 and an AS-60 autosampler were used for the determination of pseudo-total chromium concentration. The standard chromium solution (J.T. Baker Inc.) at a concentration of 1000 µg/mL served to calibrate the device. Limit of detection (LOD) (instrumental) ~~(instrumental)~~ was found to be 0.2 µg/L of Cr or 0.03 ng/m³ expressed as the concentration of Cr in the air. The quality of the chromium results was characterized with a recovery test of the CRM BCR-701 (river sediment). It was found to be 259±18 mg/kg Cr_{tot} (n = 3), while the certified value 125 is 272±20 mg/kg, hence the recovery for extractable Cr was 95.2%.

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

2.4 The determination of chromium(VI)

The filters with the collected airborne particulate matter were extracted as per a modified US EPA 3060A standard by means of alkaline digestion (US EPA, 1996). The modification was necessary given the low chromium(VI) contents in the suspended particulate matter samples. For this purpose, a quarter of the exposed filter was placed in the Teflon vessel, 5 mL of 0.056 M Na₂CO₃/0.08 M NaOH was added and extracted in a microwave oven MLS 1200 Mega (3 min, 400W). The leachates were then centrifuged (MPW 342) and filtered through a 0.45 µm Millex-HV syringe filter (Millipore) into 10 mL test tubes. Cr(VI) in the leachates was determined using the technique of catalytic cathodic stripping voltammetry with the adsorption of Cr(III)-DTPA complexes (CCSV-DTPA) [Li and Xue, 2001]. The voltammograms were recorded with a Trace Analyzer Model 394 connected to a hanging mercury drop working electrode Model 303A SMDE (EG&G Princeton Applied Research Electrochemical).

The supporting electrolyte (5 mL water, 500 µL saturated solution of KNO₃, 200 µL 0.1 M AcONa, 200 µL 0.1 M DTPA) and a 100 µL sample of Cr(VI) extract were transferred into an electrochemical cell, pH was adjusted to 6.00±0.05, and the solution was purged for 4 min with high purity argon. Deposition on a mercury drop was carried out for 60s at -0.95 V. Then the stirring was discontinued and an equilibration time of 10 s was allowed. The differential pulse scan was carried out from -0.95 V to -1.65 V, the pulse height was 50 mV and the rate 6 mV/s. The peak corresponding to the reduction of adsorbed Cr(III)-DTPA complex appeared at -1.20 V to -1.30 V. The chromium(VI) concentration was assayed using the technique of double spike-standard addition (2-10 µL, depending on the expected analyte concentration) with a concentration of 0.2 mg/L Cr(VI). LOD (instrumental) was found to be 2.9 µg/L or 0.010 ng/m³ expressed as the concentration of Cr(VI) in the air. CMR061-030 (sandy loam) was used as the certified reference material to validate the analytical procedure. It was found to equal 239.3 mg/kg Cr(VI) (n =3), the certified value was 241.00±9.00 mg/kg, hence the recovery —of 99.3%.

3 Results and discussion

3.1 The mass concentration of PM

The concentrations of the following particulate matter fractions were assayed: PM₁₀, PM_{2.5}, PM₁, and PM_{0.25}. The World Health Organization (WHO) claims that there is no evidence of a safe exposure level or threshold below which there are no adverse effects of exposure to particulate matter (WHO, 2013).

In Radom, the PM₁₀ fraction concentration ranged widely from 5.2 to 68.2 µg/m³ (40±17 µg/m³ on average). The PM_{2.5} concentrations were lower, from 3.1 to 59.2 µg/m³ (33±15 µg/m³ on average). The concentrations of finer particulate matters were in the following ranges: PM₁ from 2.4 to 45.4 µg/m³ (26±13 µg/m³ on average-), and PM_{0.25} – from 1.3 to 27.6 µg/m³ (12.6±6.6 µg/m³ on average) (Table 1). Comparable concentrations of urban PM_{2.5} are reported by other authors, e.g., Budapest (Hungary) 23 µg/m³, Istanbul (Turkey) 40 µg/m³ (Szigeti et al., 2013), Tehran (Iran) 33±11 µg/m³ (Arhami et al., 2017), Katowice (Poland) 31 µg/m³ (Rogula-Kozłowska et al., 2013a).

- sformatowano: Indeks dolny
- sformatowano: Indeks dolny
- sformatowano: Indeks dolny
- sformatowano: Indeks dolny
- sformatowano: Indeks dolny
- sformatowano: Indeks dolny
- sformatowano: Indeks dolny
- sformatowano: Indeks dolny

175 The high airborne particulate matter concentrations in the urban area investigated during the heating campaign winter season suggest municipal emissions including coal burning in household furnaces were the chief source of particulate matter pollution. The higher suspended particulate matter concentrations in the heating-winter season have also been found by others, e.g., Rogula-Kozłowska et al. (2013a) reports PM_{2.5} concentrations in Zabrze and Katowice (Poland) usually ranging between 10 and 20 µg/m³ and far greater in winter, from 50 to 110 µg/m³ (Rogula-Kozłowska et al., 2013a). The higher suspended particulate matter concentrations in winter (pointing to a greater share of fuel combustion sources) have been noted by other authors, too (Krzemińska-Flowers et al., 2006; Widziewicz et al., 2016).

— sformatowano: Indeks dolny

3.2 Chromium concentration

Given the environmental aspect of this study, the environmentally available chromium content Cr_{tot} (pseudo-total concentration of Cr) in the PM samples was assayed. All the results for Cr_{tot} concentrations in PM₁₀, PM_{2.5}, PM₄ and PM_{0.25} are listed in Table S2-S3 (Supplementary Material).

185 Cr_{tot} concentrations in the particulate matter fractions studied varied quite widely-ranged from 0.08 to 4.09 ng/m³. Like in the case of particulate matter concentrations, the results were several times lower than the chromium concentrations given in our previous work for the non-industrial zone (15 ng/m³ on average), when we studied TSP (Świetlik et al., 2011). It should be pointed out that particulate matter pollution has been substantially reduced in Poland in recent years, owing to the application of state-of-the-art, efficient, and environment-friendly technological solutions. The continuing modernization of the energy, heating, and industrial sectors - such as the EU Clean Air Program (since 2018) and provincial anti-smog resolutions (since 2017) - along with improved fuel quality regulations (established by the Minister of Industry and the Minister of Climate and Environment regarding the quality of solid fuels since 2018), has led to a consistent reduction in the amount of particulate matter pollution emitted into the air each year (EU Clean Air Program, 2024; Regulation of the Minister of Industry and the Minister of Climate and Environment, 2024). It should be pointed out particulate matter pollution has been substantially reduced in Poland in recent years owing to the application of state-of-the-art, efficient, and environment-friendly technological solutions. The continuing modernisation of the energy, heating, and industrial sectors, as well as an improved quality of fuel, have helped to cut the quantity of particulate matter pollution emitted into the air year by year.

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

The results indicate a marked impact of the heating season on Cr_{tot} concentrations in all the PM fractions (Figure 1).




Figure 1. The concentrations of pseudo-total chromium in PM_{10} , $PM_{2.5}$, PM_1 and $PM_{0.25}$ depending on the seasons.

The maximum Cr_{tot} concentration relating to PM_{10} was found in winter (2.23 ± 0.53 ng/m³ on average), whereas it averaged 1.71 ± 0.83 ng/m³ in the entire measurement period. The mean concentrations relating to PM_{10} in Radom were similar to those determined in other cities in Europe in urban areas: Edinburgh (U.K.) 1.6 ng/m³ (Heal et al., 2005), Katowice (Poland) 2.39 ng/m³ (Rogula-Kozłowska, 2015), Rome (Italy) 2-5 ng/m³ (Catrambone et al., 2013).

The $PM_{2.5}$ fraction contains approximately 80% of the total chromium content. A similar correlation was observed for the PM concentration in the winter season, where 86% of PM_{10} was the $PM_{2.5}$ fraction (Table 1). The $PM_{2.5}$ fraction is assumed to be the result of emissions from anthropogenic sources (Nocoń et al., 2018). The sampling point was located near a busy street and a single-family housing estate (from the west and south-west) and far away from an industrial zone (approx. 8-10 km from the south and south-west). Considering that Radom is a city with prevailing westerly winds, especially in autumn and winter (WeatherSpark, 2025), it can be supposed that municipal emissions, mainly stationary coal combustion sources, and road

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

255 Cr(VI) concentrations were maximum in winter (the heating campaign) and averaged: $PM_{10} - 0.84 \pm 0.43 \text{ ng/m}^3$, $PM_{2.5} - 0.70 \pm 0.39 \text{ ng/m}^3$, $PM_1 - 0.58 \pm 0.31 \text{ ng/m}^3$, $PM_{0.25} - 0.36 \pm 0.21 \text{ ng/m}^3$. Maximum Cr(VI) concentrations in PM_{10} reached 1.354 ng/m^3 . For the sake of legal protection against airborne chromium(VI), a reference value of $0.4 \text{ } \mu\text{g/m}^3$ (for 1 year) was introduced in Poland (The Minister of the Environment Regulation, 2010). The World Health Organization (WHO) recommends the baseline limit of hexavalent chromium with an excess lifetime risk (RR values) of 1:10,000, 1:100,000 and 1:1,000,000 to be 2.5, 0.25, ~~0.0025~~ ng/m^3 , respectively (WHO, 2000).
260 Regardless of the particulate matter fraction, chromium(VI) accounted for approximately 20% of Cr_{tot} content in the particulate matter ($PM_{10} - 19.3\%$, $PM_{2.5} - 19.9\%$, $PM_1 - 21.1\%$, $PM_{0.25} - 22.4\%$) during the whole measurement period (Table 2). Cr(VI) share in the particular seasons varied, however. It was minimum in summer, averaging 9.1%, somewhat greater in spring (11.3%), to rise to 22.7% (from 18.1% to 26.9%) in the autumn. It was maximum in winter, constituting 40% of total chromium on average (from 37.6% to 42.6%). Our results are not different from those reported in the literature, e.g., Bell and Hipfner (1997) and Talebi (2003) claim that circa 20% airborne chromium was Cr(VI), Nusko and Heumann (1997) give approx. 30%. Tirez et al. (2011) reported that, in the case of $PM_{2.5}$ from coal combustion, the share of Cr(VI) ranged from 6 to 43% of the total Cr (Tirez et al., 2011). Torkmahalleh et al. (2013) state the average Cr(VI) to total Cr ratios varied from 1 to 30%, whereas Kang et al. (2016) say the concentration of Cr(VI) measured accounted for 0.7 to 9.4 percent of the total chromium level, 270 which is a low percentage compared to those in other urban areas around the world.

Table 2. The percentage share of chromium speciation.

Season	PM ₁₀		PM _{2.5}		PM ₁		PM _{0.25}	
	Share of Cr(VI) [%]	Share of Cr(III) [%]	Share of Cr(VI) [%]	Share of Cr(III) [%]	Share of Cr(VI) [%]	Share of Cr(III) [%]	Share of Cr(VI) [%]	Share of Cr(III) [%]
Spring	9.8	90.2	9.5	90.5	11.1	88.9	14.8	85.2
Summer	11.6	88.4	10.9	89.1	8.5	91.5	5.3	94.7
Autumn	18.1	81.9	21.1	78.9	24.6	75.4	26.9	73.1
Winter	37.6	62.4	38.1	61.9	40.4	59.6	42.6	57.4
Average value	19.3	80.7	19.9	80.1	21.1	78.9	22.4	77.6

Cr(III) averaged 79.3% of Cr_{tot} (Table 2). It was maximum in summer (90.9% on average) and a little lower in spring (88.7%). It was minimum in winter, 60.3% of the total chromium content (Figure S3, Supplementary Material).
275 Based on the difference of Cr(VI) concentrations assayed in PM_{10} , $PM_{2.5}$, PM_1 and $PM_{0.25}$, its concentrations in: $PM_{2.5-10}$, $PM_{1-2.5}$, $PM_{0.25-1}$ and $PM_{0.25}$ were calculated (Table S3S4, Supplementary Material). Like in the case of Cr_{tot} , regardless of the season, Cr(VI) concentrations were normally maximum in the finest fraction, $PM_{0.25}$ (Figure S4, Supplementary Material). Some authors report that fine particulate matter ~~partieles~~, which can be carried over long distances, contain more Cr(VI) (Wang et al., 2023).

— sformatowano: Indeks dolny
— sformatowano: Indeks dolny
— sformatowano: Indeks dolny
— sformatowano: Indeks dolny
— sformatowano: Indeks dolny

— sformatowano: Indeks dolny
— sformatowano: Indeks dolny
— sformatowano: Indeks dolny
— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny
— sformatowano: Indeks dolny
— sformatowano: Indeks dolny
— sformatowano: Indeks dolny

— sformatowano: Indeks dolny
— sformatowano: Indeks dolny
— sformatowano: Indeks dolny
— sformatowano: Indeks dolny
— sformatowano: Indeks dolny

310 ~~risk indicate no significant cancer risk~~. *HQ* below one suggests no significant risk of non-carcinogenic effects. If *HQ* is equal to or more than 1, the non-carcinogenic effects are possible in the future.

The estimated potential ~~non-carcinogenic (HQ) carcinogenic (CR)~~ and ~~non-carcinogenic (HQ) carcinogenic (CR)~~ risk of inhalatory exposure to Cr ~~and Cr(VI)~~ present in PM_{10} for urban residents is shown in Table 3. Both were maximum in winter, when Cr concentrations become highest. ~~The World Health Organization (WHO) recommends the baseline limit of hexavalent chromium with an excess lifetime risk (RR values) of 1:10,000, 1:100,000, and 1:1,000,000 to be 2.5, 0.25, and 0.025 ng/m³, respectively [WHO, 2000].~~ In the light of the standard interpretation, however, regardless of the season, the carcinogenic risk to residents based on Cr(VI) concentration in PM_{10} ~~was below the US EPA threshold for significant risk was within the~~ ~~acceptable range~~ (between $1 \cdot 10^{-6}$ and $1 \cdot 10^{-4}$) and amounted to between $1.11 \cdot 10^{-6}$ and $5.78 \cdot 10^{-6}$ for children and from $3.69 \cdot 10^{-6}$ to $1.92 \cdot 10^{-5}$ for adults. Even in the case of maximum Cr(VI) concentration in PM_{10} during the winter season, the estimated carcinogenic risk to the population of Radom ~~was lower than the upper limit was acceptable~~: $CR=9.24 \cdot 10^{-6}$ for children and $CR=3.12 \cdot 10^{-5}$ for adults. ~~However, although the risk of cancer in children is generally low, it can be misleading because exposures during development can have disproportionately dramatic effects in adulthood.~~

320 **Table 3. Carcinogenic and non-carcinogenic risk from chromium via inhalation exposure to the airborne particles of PM_{10} .**

Season	Carcinogenic risk - CR			Non-carcinogenic risk - HQ		
	Cr(VI) [ng/m ³]	Children	Adult	Cr _{tot} [ng/m ³]	Children	Adult
Average concentration of chromium in PM_{10} in each season						
Spring	0.1604	$1.11 \cdot 10^{-6}$	$3.69 \cdot 10^{-6}$	1.64	$1.57 \cdot 10^{-2}$	$1.57 \cdot 10^{-2}$
Summer	0.1823	$1.26 \cdot 10^{-6}$	$4.19 \cdot 10^{-6}$	1.57	$1.51 \cdot 10^{-2}$	$1.51 \cdot 10^{-2}$
Autumn	0.2508	$1.73 \cdot 10^{-6}$	$5.77 \cdot 10^{-6}$	1.38	$1.32 \cdot 10^{-2}$	$1.32 \cdot 10^{-2}$
Winter	0.8369	$5.78 \cdot 10^{-6}$	$1.92 \cdot 10^{-5}$	2.23	$2.14 \cdot 10^{-2}$	$2.14 \cdot 10^{-2}$
Av.		$2.47 \cdot 10^{-6}$	$8.21 \cdot 10^{-6}$		$1.64 \cdot 10^{-2}$	$1.64 \cdot 10^{-2}$
Maximum chromium concentration in PM_{10} throughout the measurement period						
	1.3544	$9.24 \cdot 10^{-6}$	$3.12 \cdot 10^{-5}$	4.09	$3.92 \cdot 10^{-2}$	$3.92 \cdot 10^{-2}$

The estimated non-carcinogenic inhalation risks from chromium for the residents of Radom posed ~~little risk no threat~~, either. The *HQ* values calculated ~~on the basis based on of~~ the total Cr concentration in PM_{10} were lower than the safe level (*HQ* = 1) and ranged from $1.32 \cdot 10^{-2}$ to $2.14 \cdot 10^{-2}$, indicating ~~no little~~ non-carcinogenic risks from chromium. The non-carcinogenic health risk based on the maximum Cr_{tot} concentration in PM_{10} was ~~acceptable low~~, too, ~~as it was not more than one as it was not in excess of one~~ (Table 3). These risk values are no different ~~to from~~ those found in the literature, e.g., *HQ* in Italy was in the range $1.5 \cdot 10^{-2} - 5.4 \cdot 10^{-2}$ (Diana et al., 2023).

EU air quality directives do not provide for separate normative values for chromium. Directive 2024/2881 specifies quality objectives for arsenic, cadmium, lead, mercury, nickel, and PAHs, but does not cover chromium (Directive 2024/2881). Within

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

the framework of EU policy, chromium(VI) is regulated mainly in the context of occupational safety – the binding occupational exposure limit has been set at 0.005 mg/m³ by 2025 (Directive 2017/2398). For comparison, the highest obtained concentration of Cr(VI) in PM₁₀ (1.3544 ng/m³) is almost three orders of magnitude lower than this value. Consequently, the estimated health risk of chromium is low and does not raise significant health concerns.

In conclusion, the estimated potential non-carcinogenic (HQ) and carcinogenic (CR) risk of inhalatory exposure to Cr and Cr(VI), respectively, present in PM₁₀ for the urban residents of Radom indicates that risks of adverse health effects from chromium and chromium(VI) exposure in Radom are relatively low and complies with the applicable WHO and EU regulatory framework (WHO, 2021; Directive 2017/2398).

— sformatowano: Indeks górny

— sformatowano: Indeks dolny

— sformatowano: Indeks górny

— sformatowano: Indeks dolny

4 Conclusion

The mean concentrations of PM₁₀ (40±17 µg/m³) and PM_{2.5} (33±15 µg/m³) in the city were above the national (40 µg/m³ and 20 µg/m³, respectively) and European (40 µg/m³ and 25 µg/m³, respectively). The total chromium content in PM₁₀ ranged widely from 0.56 to 4.09 ng/m³ and averaged 1.71±0.83 ng/m³. The mean Cr(VI) concentration in PM₁₀ was 0.38 ng/m³ in the entire measurement period. Chromium(VI) accounted for ca. 20% of Cr_{tot} content in the particulate matter. The seasonality of Cr_{tot} and Cr(VI) concentration changes could be noted. The concentrations were maximum in the winter (heating) season: 2.23 ng/m³ and 0.84 ng/m³, respectively, most likely a result of a greater share of the air polluted by fuel combustion sources in the air emissions of this type of pollution. The share of Cr(VI) in PM in the individual seasons varied as well: it was minimum in summer (9.1% of Cr_{tot}) and maximum in winter (40% of Cr_{tot}).

Out of the airborne particulate matter fractions investigated (PM_{2.5-10}, PM_{1-2.5}, PM_{0.25-1}, PM_{0.25}), both Cr_{tot} and Cr(VI) concentrations were maximum in the finest fraction – PM_{0.25}. Toxicity related to ~~transitive-transition~~ metals is greater for fine than for coarse dust since, once they penetrate the respiratory system, the finest ~~particulate-matter~~ particles reach pulmonary alveoli, where 60-80% of a transmitted element enters the blood (Rogula-Kozłowska, 2013b). The health risk to urban residents, both carcinogenic and non-carcinogenic, is estimated to be ~~gravest-highest~~ in winter as Cr_{tot} and Cr(VI) concentrations reach their top values. Regardless of the season, however, these risk levels to the residents of Radom were within the acceptable risk range.

Research into the forms of atmospheric Cr is limited and atmospheric Cr(VI) needs to be further investigated.

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

Data availability

Data sets are available at <https://doi.org/10.5281/zenodo.14808852> (Łożyńska, M., et al., 2025)

CRedit authorship contribution statement

M.L. - Data curation, Formal analysis, Methodology, Writing – original draft, Writing – review and editing, Conceptualization, Investigation, Validation; **M.T.** – Data curation, Methodology, Writing – original draft, Writing – review and editing, Conceptualization, Investigation, Validation; **A.M.** - Data curation, Methodology, Visualization, Investigation; **R.Ś.** – Conceptualization, Methodology, Project administration, Supervision.

Declaration of Competing Interest

The authors declare that they have no conflict of interest.

References

Arhami, M., Hosseini, V., Shahne, M.Z., Bigdeli, M; Lai, A. and Schauer, J.J.: Seasonal trends, chemical speciation and source apportionment of fine PM in Tehran, Atmos. Environ., 153, 70-82, <http://dx.doi.org/10.1016/j.atmosenv.2016.12.046>, 2017.

Anake, W.U., Benson, N.U., Tenebe, I.T., Emenike, P.Ch., Ana, G.R.E.E. and Zhang S.: Chemical speciation and health risks of airborne heavy metals around an industrial community in Nigeria, Hum. Ecol. Risk Assess., 26(1), 242-254, <https://doi.org/10.1080/10807039.2018.1504672>, 2020.

Barceloux, D.G. and Barceloux, D.: Chromium. J. Toxicol. Clin. Toxicol., 37(2), 173-94. <https://doi.org/10.1081/CLT-100102418>, 1999.

Bell, R.W. and Hipfner, J.C.: Airborne Hexavalent Chromium in Southwestern Ontario, J. Air Waste Manag. Assoc., 47, 905–910, <http://dx.doi.org/10.1080/10473289.1997.10464454>, 1997.

Canepari, S., Pietrodangelo, A., Perrino, C., Astolfi, M.L. and Marzo M.L.: Enhancement of source traceability of atmospheric PM by elemental chemical fractionation, Atmos. Environ., 43, 4754–4765, <https://doi.org/10.1016/j.atmosenv.2008.09.059>, 2009.

Catrambone, M., Canepari, S. and Perrino C.: Determination of Cr(III), Cr(VI) and total chromium in atmospheric aerosol samples, E3S Web of Conferences, 1, 07005, <http://dx.doi.org/10.1051/e3sconf/20130107005>, 2013.

Conca, E., Malandrino, M., Giacomino, A., Costa, E., Ardini, F., Inaudi, P. and Abollino, O.: Optimization of a sequential extraction procedure for trace elements in Arctic PM₁₀, Anal. Bioanal. Chem., 412, 7429–7440, <https://doi.org/10.1007/s00216-020-02874-4>, 2020.

Diana, A., Bertinetti, S., Abollino, O., Giacomino, A., Buoso, S., Favilli, L., Inaudi, P. and Malandrino, M.: PM₁₀ Element Distribution and Environmental-Sanitary Risk Analysis in Two Italian Industrial Cities, Atmosphere, 14, 48, <https://doi.org/10.3390/atmos14010048>, 2023.

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

- 385 [Directive \(EU\) 2017/2398 of the European Parliament and of the Council of 12 December 2017 amending Directive 2004/37/EC on the protection of workers from the risks related to exposure to carcinogens or mutagens at work. Official Journal of the European Union, L 345/87.](#)
- [Directive \(EU\) 2024/2881 Directive 2008/50/EC of the European Parliament and of the Council on ambient air quality and cleaner air for Europe of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe.](#) Official Journal of the European Union, [32024L2881L-152/4](#).
- 390 Dos Santos, M., Gómez, D., Dawidowski, L., Gautiera, E. and Smichowski, P.: Determination of water-soluble and insoluble compounds in size classified airborne particulate matter, *Microchem. J.*, 91, 133–139, <https://doi.org/10.1016/j.microc.2008.09.001>, 2009.
- [EU Clean Air Program, <https://czystepowietrze.gov.pl/>, last access: 23.04.2025.](#)
- 395 Fang, G.C., Wu, Y.S., Wen, C.C., Lin, C.K., Huang, S.H., Rau, J.H. and Lin, Ch.P.: Concentrations of nano and related ambient air pollutants at a traffic sampling site, *Toxicol. Ind. Health*, 21, 259-271, <https://doi.org/10.1191/0748233705th234oa>, 2005.
- Feng, X.D., Dang, Z., Huang, W.L. and Yang, C.: Chemical speciation of fine particle bound trace metals, *Int. J. Environ. Sci. Tech.*, 6(3), 337-346, <https://doi.org/10.1007/BF03326071>, 2009.
- Gunchin, G., Osan, J., Migliori, A., Shagjamba, D. and Strel, C.: Chromium and Zinc Speciation in Airborne Particulate Matter Collected in Ulaanbaatar, Mongolia, by X-Ray Absorption Near-edge Structure Spectroscopy, *Aerosol Air Qual. Res.*, 21, 8, 210018, <https://doi.org/10.4209/aaqr.210018>, 2021.
- 400 [Heal, M.R., Hibbs, L.R., Agius, R.M., Beverland, I.J.: Total and water-soluble trace metal content of urban background PM10, PM2.5 and Black Smoke in Edinburgh, U.K, *Atmos. Environ.*, 39, 1417-1430, doi:10.1016/j.atmosenv.2004.11.026, 2005.](#)
- 405 Huang, L., Yu, C.H., Hopke, P.K., Lioy, P.J., Buckley, B.T., Shin, J.Y. and Fan, Z.: Measurement of Soluble and Total Hexavalent Chromium in the Ambient Airborne Particles in New Jersey, *Aerosol Air Qual. Res.*, 14, 1939-1949, <https://doi.org/10.4209/aaqr.2013.10.0312>, 2014a.
- Huang, L., Yu, C.H., Hopke, P.K., Shin, J.Y. and Fan, Z.: Trivalent chromium solubility and its influence on quantification of hexavalent chromium in ambient particulate matter using EPA method 6800, *J. Air Waste Manage.*, 64, 1439–1445, <https://doi.org/10.1080/10962247.2014.951745>, 2014b.
- 410 IARC 2012, Chromium(VI) compounds. IARC Monographs on the evaluation carcinogenic risk to humans. IARC Press, Lyon, 100C.
- IARC 2023, Agents Classified by the IARC Monographs, <https://monographs.iarc.who.int/agents-classified-by-the-iarc/>
- Izhar, S., Goel, A., Chakraborty, A. and Gupta, T.: Annual trends in occurrence of submicron particles in ambient air and health risk posed by particle bound metals, *Chemosphere*, 146, 582-590, <http://dx.doi.org/10.1016/j.chemosphere.2015.12.039>, 2016.
- 415 Jan, R., Roy, R., Yadav, S. and Satsangi, P.G.: Chemical fractionation and health risk assessment of particulate matter-bound metals in Pune, India, *Environ. Geochem. Health*, 40, 255-270, <https://doi.org/10.1007/s10653-016-9900-7>, 2018.

— sformatowano: Domyślina czcionka akapitu, Czcionka: (Domyślny) Times New Roman

Kang, B.W., Lee, H.S., Kim, J.H., Hong, J.H., Kim, R.H., Seo, Y.K., Han, J.S., Baek, K.M., Kim, M.J. and Baek, S.O.:
420 Distribution of Airborne Hexavalent Chromium Concentrations in Large Industrial Complexes in Korea, *Asian J. Atmos. Environ.*, 10, 208-216, <https://doi.org/10.5572/ajae.2016.10.4.208>, 2016.

Katz, S.A.: The Analytical Biochemistry of Chromium, *Environ. Health Persp.*, 92, 13-16. <https://doi.org/10.1289/ehp.919213>, 1991.

Khlystov, A. and Ma, Y.: An on-line instrument for mobile measurements of the spatial variability of hexavalent and trivalent
425 chromium in urban air, *Atmos. Environ.*, 40, 8088–8093, <https://doi.org/10.1016/j.atmosenv.2006.09.030>, 2006.

Kotaś, J. and Stasicka, Z.: Chromium occurrence in the environment and methods of its speciation, *Environ. Pollut.*, 107, 263–
283, [https://doi.org/10.1016/S0269-7491\(99\)00168-2](https://doi.org/10.1016/S0269-7491(99)00168-2), 2000.

Krzemińska-Flowers, M.H., Bem, H. and Górecka, H.: Trace metals concentration in size-fractioned urban air particulate
matter in Lodz. Poland. I. Seasonal and site fluctuations, *Pol. J. Environ. Stud.*, 15(5), 759-767. 2006.

430 Li, H., Wang, Q., Shao, M., Wang, J., Wang, C., Sun, Y., Qian, X., Wu, H., Yang, M. and Li, F.: Fractionation of airborne
particulate-bound elements in haze-fog episode and associated health risks in a megacity of southeast China, *Environ. Pollut.*,
208, 655-662, <https://doi.org/10.1016/j.envpol.2015.10.042>, 2016.

Li, Y. and Xue, H.: Determination of Cr(III) and Cr(VI) species in natural waters by catalytic cathodic stripping voltammetry,
Anal. Chim. Acta, 448, 121e134, [https://doi.org/10.1016/S0003-2670\(01\)01314-9](https://doi.org/10.1016/S0003-2670(01)01314-9), 2001.

435 Łożyńska, M., Trojanowska, M., Molik, A. and Świątlik, R.: Measurement Report: Seasonal trends and chemical speciation
of chromium(III/VI) in different fractions of urban particulate matter – a case study of Radom, Poland [data set],
<https://doi.org/10.5281/zenodo.14808852>, 2025.

Majewski, G. and Rogula-Kozłowska, W.: The elemental composition and origin of fine ambient particles in the largest Polish
conurbation: first results from the short-term winter campaign, *Theor. Appl. Climatol.*, 125, 79–92,
440 <https://doi.org/10.1007/s00704-015-1494-y>, 2016.

Ministry of Climate and Environment, National emission balance of SO₂, NO_x, CO, NH₃, NMLZO, dust, heavy metals and
POPs for the years 1990-2020. Synthetic Report, Warsaw 2022.

Molik, A., Trojanowska, M., Łożyńska, M. and Świątlik, R.: Seasonal variations in chromium concentration in urban
atmospheric aerosol in the city of Radom, *E3S Web of Conferences – Air Protection in Theory and Practice* 28, 01024.
445 <https://doi.org/10.1051/e3sconf/20182801024>, 2018.

Nocoń, K., Rogula-Kozłowska, W. and Widziewicz, K.: Research on chromium and arsenic speciation in atmospheric
particulate matter: short review, *E3S Web of Conferences – Air Protection in Theory and Practice*, 28, 01026.
<https://doi.org/10.1051/e3sconf/20182801026>, 2018.

Nriagu, J.O., Pacyna, J.M., Milford, J.B. and Davidson C.I.: Distribution and characteristic features of chromium in the
450 atmosphere, in *Chromium in Natural and Human Environments*, in: Nriagu, J.O., Nieboer, E. (ed), *Chromium in the Natural
and Human Environments*, Wiley, New York, 125. 1988.

- Nriagu, J.O. and Pacyna, J.M.: Quantitative assessment of worldwide contamination of air, water and soils by trace metals, *Nature*, 333, 134-139, <https://doi.org/10.1038/333134a0>, 1988.
- Nriagu, J.O. and Nieboer, E.: *Chromium in the Natural and Human Environments*, John Wiley & Sons, New York, 20, 1988.
- 455 Nusko, R. and Heumann, K.G.: Cr(III)/Cr(VI) speciation in aerosol particles by extractive separation and thermal ionization isotope dilution mass spectrometry, *Fresenius J. Anal. Chem.*, 357, 1050–1055, <https://doi.org/10.1007/s002160050303>, 1997.
- Pacyna, E.G., Pacyna, J.M., Fudala, J., Strzelecka-Jastrzab, E., Hlawiczka, S., Panasiuk, D., Nitter, S., Pregger, T., Pfeiffer, H. and Friedrich, R.: Current and future emissions of selected heavy metals to the atmosphere from anthropogenic sources in Europe, *Atmos. Environ.*, 41, 8557-8566, <https://doi.org/10.1016/j.atmosenv.2007.07.040>, 2007.
- 460 Pacyna, J.M.: Atmospheric trace elements from natural and anthropogenic sources, in: Nriagu J.O., Davidson, C.I. (Eds.). *Toxic Metals in the Atmosphere*, John Wiley & Sons, New York, 1986.
- Pan, Y., Tian, S., Li, X., Sun, Y., Li, Y., Wentworth, G.R. and Wang, Y.: Trace elements in particulate matter from metropolitan regions of Northern China: sources, concentrations and size distributions, *Sci. Total Environ.*, 537, 9-21, <http://dx.doi.org/10.1016/j.scitotenv.2015.07.060>, 2015.
- 465 Pandey, M., Pandey, A.K., Mishra, A. and Tripathi, B.D.: Speciation of carcinogenic and non-carcinogenic metals in respirable suspended particulate matter (PM₁₀) in Varanasi, India, *Urban Climate*, 17, 141-154, <http://dx.doi.org/10.1016/j.uclim.2017.01.004>, 2017.
- Park, K., Cho, G. and Kwak, J.: Development of an Aerosol Focusing-Laser Induced Breakdown Spectroscopy (Aerosol Focusing-LIBS) for Determination of Fine and Ultrafine Metal Aerosols, *Aerosol. Sci. Tech.*, 43, 375-386.
- 470 <https://doi.org/10.1080/02786820802662947>, 2009.
- Proctor, D.M., Bhat, V., Suh, M., Reichert, H., Jiang, X. and Thompson, C.M.: Inhalation cancer risk assessment for environmental exposure to hexavalent chromium: Comparison of margin-of-exposure and linear extrapolation approaches, *Regul. Toxicol. Pharm.*, 124, 104969, <https://doi.org/10.1016/j.yrtph.2021.104969>, 2021.
- Richter, P., Grino, P., Ahumada, I. and Giordano, A.: Total element concentration and chemical fractionation in airborne particulate matter from Santiago, Chile, *Atmos. Environ.*, 42, 6729-6738, <https://doi.org/10.1016/j.atmosenv.2007.04.053>, 2007.
- 475 [Regulation of the Minister of Industry and the Minister of Climate and Environment of November 4, 2024 regarding quality requirements for solid fuels, 2024, Poland.](#)
- Regulation of the Minister of the Environment of 26 January 2010 on reference values for certain substances in the air, 2010,
- 480 Poland.
- Rogula-Kozłowska, W., Błaszczak, B., Szopa, S., Klejnowski, K., Sówka, I., Zwoździak, A., Jabłońska, M. and Mathews, B.: PM_{2.5} in the central part of Upper Silesia. Poland: concentrations. elemental composition. and mobility of components, *Environ. Monit. Assess.*, 185, 581–601, <https://doi.org/10.1007/s10661-012-2577-1>, 2013a.
- Rogula-Kozłowska, W., Kozielska, B., Klejnowski, K. and Szopa, S.: Hazardous Compounds in Urban PM in the Central Part of Upper Silesia (Poland) in Winter, *Arch. Environ. Prot.*, 39(1), 53-65, <https://doi.org/10.2478/aep-2013-0002>, 2013b.
- 485

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

Rogula-Kozłowska, W.: Chemical composition and mass closure of ambient particulate matter at a crossroads and a highway in Katowice, Poland, *Environ. Prot. Eng.*, 2(41), 15-29, <https://doi.org/10.5277/epe150202>, 2015.

Regulation of the Minister of the Environment of 24 August 2012 on the levels of certain substances in the air. *Journal of Laws*, item 1031, Poland.

490 Rubio, M.A., Sánchez, K., Richter, P., Pey, J. and Gramsch, E.: Partitioning of the water soluble versus insoluble fraction of trace elements in the city of Santiago, Chile, *Atmosphere*, 31(4), 373-387, <https://doi.org/10.20937/ATM.2018.31.04.05>, 2018.

Sah, D., Verma, P.K., Kandikonda, M.K. and Lakhani, A.: Chemical fractionation, bioavailability, and health risks of heavy metals in fine matter at a site in the Indo-Gangetic Plain, India, *Environ. Sci. Pollut. R.*, 26, 19749-19762, <https://doi.org/10.1007/s11356-019-05144-8>, 2019.

495 Samara, C., Kantiranis, N., Kollias, P., Planou, S., Kouras, A., Besis, A., Manoli, E. and Voutsas, D.: Spatial and seasonal variations of the chemical, mineralogical and morphological features of quasi-ultrafine particles (PM_{0.49}) at urban sites, *Sci. Total Environ.*, 553, 392-403, <http://dx.doi.org/10.1016/j.scitotenv.2016.02.080>, 2016.

Samara, C. and Voutsas, D.: Size distribution of airborne particulate matter and associated heavy metals in the roadside environment, *Chemosphere*, 59, 1197-1206, <https://doi.org/10.1016/j.chemosphere.2004.11.061>, 2005.

500 Seinfeld, J.H. and Pandis, S.N.: *Atmospheric Chemistry and Physics – from Air Pollution to Climate Change*, John Wiley & Sons., New York, 2006.

Somers, C.M.: Ambient air pollution exposure and damage to male gametes: human studies and in situ 'sentinel' animal experiments, *Syst. Biol. Reprod. Med.*, 57, 63-71, <https://doi.org/10.3109/19396368.2010.500440>, 2011.

| Szigeti, T., Mihucz, V.G., Óvári, M., Baysal, A., Atilgan, S., Akman, S. and Záray, G.: Chemical characterization of PM_{2.5} fractions of urban aerosol collected in Budapest and Istanbul, *Microchem. J.*, 107, 86-94, <https://doi.org/10.1016/j.microc.2012.05.029>, 2013.

Świetlik, R., Molik, A., Molenda, M., Trojanowska, M. and Siwiec, J.: Chromium(III/VI) speciation in urban aerosol, *Atmos. Environ.*, 45, 1364-1368, <https://doi.org/10.1016/j.atmosenv.2010.12.001>, 2011.

510 Świetlik, R. and Trojanowska, M.: Chemical Fractionation in Environmental Studies of Potentially Toxic Particulate-Bound Elements in Urban Air: A Critical Review, *Toxics*, 10(3), 124, <https://doi.org/10.3390/toxics10030124>, 2022.

Talebi, S.M.: Determination of total and hexavalent chromium concentrations in the atmosphere of the city of Isfahan, *Environ. Res.*, 92, 54-56, [https://doi.org/10.1016/S0013-9351\(02\)00036-1](https://doi.org/10.1016/S0013-9351(02)00036-1), 2003.

515 Tirez, K., Silversmit, G., Bleux, N., Adriaenssens, E., Roekens, E., Servaes, K., Vanhoof, C., Vincze, L. and Berghmans, P.: Determination of hexavalent chromium in ambient air: A story of method induced Cr(III) oxidation, *Atmos. Environ.*, 45, 5332-5341, <https://doi.org/10.1016/j.atmosenv.2011.06.043>, 2011.

Torkmahalleh, M.A., Yu, C.H., Lin, L., Fan, Z.T., Swift, J.L., Bonanno, L., Rasmussen, D.H., Holsen, T.M. and Hopke, P.K.: Improved Atmospheric Sampling of Hexavalent Chromium, *J. Air Waste Manag. Assoc.*, 63(11), 1313-1323, <https://doi.org/10.1080/10962247.2013.823894>, 2013.

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

US EPA: Risk Assessment Guidance for Superfund, Vol. I, Human Health Evaluation Manual (Part A), EPA/540/1-89/002.

520 Office of Emergency and Remedial Response, Washington, D.C. 20450, 1989.

US EPA: Method 3060A. Alkaline digestion for hexavalent chromium, 1996.

US EPA: Risk Assessment Guidance for Superfund Volume I: Human Health Evaluation Manual (Part F, Supplemental Guidance for Inhalation Risk Assessment). Office of Superfund Remediation and Technology Innovation Environmental Protection Agency Washington, D.C, 2009.

525 US EPA: Recommended Default Exposure Factors. US Environmental Protection Agency, February 2014.

US EPA: Regional Screening Levels (RSLs) – Generic Tables, 2022. (available in November 2022, <https://semspub.epa.gov/work/HQ/403628.pdf>).

Wagner, A., Boman, J. and Gatari, M.J.: Elemental analysis of size-fractionated particulate matter sampled in Göteborg, Sweden, Spectrochim. Acta B, 63, 1426-1431, <https://doi.org/10.1016/j.sab.2008.10.010>, 2008.

530 Waheed, A., Li, X., Tan, M., Bao, L., Liu, J., Zhang, Y., Zhang, G. and Li, Y.: Size Distribution and Sources of Trace Metals in Ultrafine/Fine/Coarse Airborne Particles in the Atmosphere of Shanghai, Aerosol. Sci. Tech., 45. 163-171, <https://doi.org/10.1080/02786826.2010.528079>, 2011.

Wang, K., Wang, W., Li, L., Li, J., Wei, L., Chi, W., Hong, L., Zhao, Q. and Jiang, J.: Seasonal concentration distribution of PM_{1.0} and PM_{2.5} and a risk assessment of bound trace metals in Harbin, China: Effect of the species distribution of heavy metals and heat supply, Sci. Rep., 10, 8160, <https://doi.org/10.1038/s41598-020-65187-7>, 2020.

535 Wang, L., Guo, J., Zhang, W., Chen, B., Wang, H. and Li, H.: Pollution Levels for Airborne Hexavalent Chromium of PM_{2.5} in Typical Cities of China, Atmosphere, 14, 209, <https://doi.org/10.3390/atmos14020209>, 2023.

Weather Spark, <https://pl.weatherspark.com/>, last access: 28.04.2025.

WHO human health risk assessment toolkit: chemical hazards, second edition (IPCS harmonization project document, no. 8) Geneva, Switzerland, 2021.

540 WHO: Air Quality Guidelines for Europe. Geneva, Switzerland, 125–162, 2000.

WHO: Health effects of particulate matter, Policy implications for countries in eastern Europe, Caucasus and Central Asia, 2013.

WHO: Health effects of particulate matter, Policy implications for countries in eastern Europe, Caucasus and Central Asia, 2013.

545 WHO: Air Quality Guidelines for Europe. Geneva, Switzerland, 125–162, 2000.

Widziewicz, K., Rogula-Kozłowska, W. and Loska, K.: Cancer risk from arsenic and chromium species bound to PM_{2.5} and PM₁. Polish case study, Atmos. Pollut. Res., 7, 884-894, <http://dx.doi.org/10.1016/j.apr.2016.05.002>, 2016.

550 Wu, T., Liu, P., He, X., Xu, H. and Shen, Z.: Bioavailability of heavy metals bounded to PM_{2.5} in Xi'an, China: seasonal variation and health risk assessment, Environ. Sci. Pollut. R., 28(27), 35844-35853, <https://doi.org/10.1007/s11356-021-13198-w>, 2021.

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

Xie, J.J., Yuan, C.G., Xie, J., Shen, Y.W., He, K.Q. and Zhang, K.G.: Speciation and bioaccessibility of heavy metals in PM_{2.5} in Baoding city, China, Environ. Poll., 252, 336-343, <https://doi.org/10.1016/j.envpol.2019.04.106>, 2019.

555 Zajusz-Zubek, E. and Mainka, A.: Analysis of Trace metals in the Mobile Form of Respirable Fraction PM_{2.5} Collected in the Surroundings of Power Plant, Engineering and Protection of Environment, 18(2), 245-258, (in Polish), 2015.

Zajusz-Zubek, E., Kaczmarek, K. and Mainka A.: Trace Elements Speciation of Submicron Particulate Matter (PM_{2.5}) Collected in the Surroundings of Power Plants, Int. J. Environ. Res. Public Health, 12, 13085-13103. <https://doi.org/10.3390/ijerph121013085>, 2015.

560 Zajusz-Zubek, E., Radko, T. and Mainka, A.: Fractionation of trace elements and human health risk of submicron particulate matter (PM_{2.5}) collected in the surroundings of coking plants, Environ. Monit. Assess., 189, 389, <https://doi.org/10.1007/s10661-017-6117-x>, 2017.

Zwoździak, A., Sówka, I., Krupińska, B., Zwoździak, J. and Nych, A.: Infiltration or indoor sources as determinants of the elemental composition of particulate matter inside a school in Wrocław, Poland, Build. Environ., 66, 173-180, <http://dx.doi.org/10.1016/j.buildenv.2013.04.023>, 2013.

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny

— sformatowano: Indeks dolny