

Development and Testing of Ensemble-Variational Data Assimilation Capabilities for Radar Data within JEDI coupled with FV3-LAM Model

Jun Park^{1,+,} Chengsi Liu^{1,*}, Ming Xue^{1,2}

1 Center for Analysis and Prediction of Storms, University of Oklahoma, Norman, Oklahoma, 73072, USA
2 School of Meteorology, University of Oklahoma, Norman, Oklahoma, 73072, USA

⁺ Current affiliation: NSF National Center for Atmospheric Research, Boulder, Colorado 80301

* Current affiliation: Cooperative Institute for Severe and High-Impact Weather Research and Operations, University of Oklahoma, Norman, OK.

Correspondence to: Chengsi Liu (cliu@ou.edu)

Abstract. This study presents the first implementation and evaluation of radar reflectivity data assimilation capabilities within the ensemble three-dimensional variational (En3DVar) data assimilation (DA) system of the Joint Effort for Data assimilation Integration (JEDI) framework. Building on our earlier works that assimilated reflectivity in JEDI LETKF and in GSI En3DVar, this study focuses on the JEDI En3DVar algorithm when coupled with the FV3-LAM model using the Thompson microphysics scheme. The radar reflectivity observation operator is refined by modifying the snow and graupel reflectivity formulations to improve consistency with Thompson microphysics. The new operator notably improves reflectivity analyses at the upper levels and reduces root-mean-square innovations for both reflectivity and radial velocity during the DA cycles. A high-impact convective storm event is used to evaluate the new implementation. DA experiments are conducted using both the JEDI and GSI En3DVar systems, employing identical observation operators and similar configurations. The resulting analyses and short-range forecasts from the two systems are comparable, supporting the validity of the new implementation of JEDI En3DVar for reflectivity and radial velocity assimilation. Additional comparisons with real-time High-Resolution Rapid Refresh (HRRR) and experimental Rapid Refresh Forecast System (RRFS) forecasts are made. The JEDI-based experiment captures the storm structure and placement with accuracy similar to or better than the HRRR and RRFS forecasts. Improvements are especially evident in the depiction of convective cores and stratiform rainbands, where reflectivity intensity and coverage are better aligned with radar observations.

1 Introduction

Accurate prediction of high-impact convective-scale weather events such as supercells, tornadoes, and heavy precipitation remains a critical challenge for both research and operational numerical weather

Style Definition: Normal: Indent: First line: 0.25"

Formatted: Centered, Indent: First line: 0"

Formatted: Indent: First line: 0.25"

Formatted: Centered, Indent: First line: 0"

prediction (NWP). High-resolution radar observations, particularly radar reflectivity and radial velocity, offer invaluable information on the internal structure and rapid evolution of convective storms. The assimilation of such radar observations into NWP models has been shown to significantly improve the quality of initial conditions, thereby enhancing short-range forecast skill (e.g., Sun and Crook 1997; Schenkman et al. 2011; Snook et al. 2012; Sun et al. 2014; Kong et al. 2018; Tong et al. 2020; Labriola et al. 2021; Park et al. 2023; Gao et al. 2024).

—Ensemble Kalman filter (EnKF; Evensen 2003) and variational methods are two primary approaches for directly assimilating radar reflectivity. EnKF methods are relatively straightforward to implement and provide [ensembles of initial condition perturbations for ensemble forecasting systems](#) that can be used to initialize ensemble forecasting systems, while variational and ensemble-variational methods remain central to many operational regional forecasting systems. Both approaches typically assume Gaussian error statistics, and EnKF performance can be limited by finite ensemble size, leading to rank deficiency and under-dispersion (Houtekamer and Zhang 2016). For variational and hybrid En3DVar methods, a major challenge in radar reflectivity assimilation arises from the strong nonlinearity of reflectivity observation operators, which requires the development of tangent-linear and adjoint models and can lead to convergence difficulties or instability during minimization (Sun and Crook 1997; Liu et al. 2020). Although double-loop algorithms can partially mitigate linearization errors through iterative re-linearization (Courtier et al. 1994), these complexities have motivated alternative formulations. One such approach treats radar reflectivity as a state variable within pure En3DVar systems (Wang and Wang 2017), thereby avoiding explicit use of nonlinear reflectivity operators, but relying entirely on ensemble-derived cross covariances and complicating the inclusion of static background error covariance.

Several studies have focused on improving direct reflectivity assimilation in variational frameworks. Liu et al. (2019) introduced temperature-dependent background error profiles for hydrometeor variables, while Liu et al. (2020) addressed minimization convergence issues through lower bounds on reflectivity and separate analysis passes for radial velocity and reflectivity. Chen et al. (2021) and Li et al. (2022) further reduced nonlinearity by applying power transforms to hydrometeor control variables. In Liu et al. (2022), the reflectivity observation operator was reformulated to be consistent with the Thompson microphysics scheme and implemented in the [Gridpoint Statistical Interpolation analysis system \(GSI\)](#) En3DVar system, leading to improved analyses and short-range forecasts compared to single-moment-based operators. While Liu et al. (2022a) focused on formulation and implementation within GSI, the impacts of modifying the operator formulation and systematically comparing different operator treatments within [the Joint Effort for Data assimilation Integration \(JEDI, Trémolet and Auligné 2020\) a next-generation DA](#) framework were not examined.

The GSI data assimilation (DA) system is currently used in U.S. operational numerical weather prediction systems. Three methods for assimilating radar reflectivity have been developed and implemented in GSI: indirect assimilation through cloud analysis (e.g., Banos et al. 2022; Dowell et al. 2022), which is used operationally in the High-Resolution Rapid Refresh (HRRR) system, together with additional radar-based latent heating during a one-hour spin-up cycle (Dowell et al. 2022; Weygandt et al. 2022); direct method based on the reflectivity-state-variable approach (Wang and Wang 2017), which has been adopted in the initial version of the Rapid Refresh Forecast System (RRFSv1, Banos et al. 2022; Carley et al. 2023); and a traditional variational approach that includes the reflectivity

Formatted

Formatted: Indent: First line: 0"

observation operator along with its tangent-linear and adjoint components in the cost function, which are formulated to be consistent with model microphysics (Liu et al. 2022a).

The ~~Joint Effort for Data assimilation Integration (JEDI; Trémolet and Auligné 2020)~~ is being developed as the next-generation DA system for NOAA’s Unified Forecast System (UFS), with the goal of replacing the current GSI framework. JEDI is designed to be modular, extensible, and model-agnostic, enabling community-driven development and supporting both research and operational applications across coupled Earth system components. Recent progress has demonstrated the feasibility of assimilating radar observations within JEDI using variants of the EnKF method. Park et al. (2023) implemented radar DA in JEDI coupled with the FV3-LAM model (Lin 2004) using the Local Ensemble Transform Kalman Filter (LETKF, Hunt et al. 2007) and the gain-form localization version of LETKF (LGETKF, Bishop et al. 2017), showing promising results in the analysis and short-range prediction of convective storms. However, Park et al. (2023) was limited to demonstrating reflectivity and radial velocity assimilation using the LETKF and LGETKF ensemble algorithms in JEDI. In contrast, the assimilation with En3DVar requires the implementation and testing of tangent-linear and adjoint components, which are highly desirable within JEDI. In fact, the core of the data assimilation systems in current NWS operational regional forecasting models is the hybrid En3DVar, although it is based on the GSI framework (Hu et al. 2017; Wu et al. 2017; Dowell et al. 2022). Therefore, the development of direct reflectivity assimilation capabilities within the JEDI variational framework is both desirable and timely.

This work addresses this gap by developing and implementing the tangent-linear and adjoint of the reflectivity operator in the JEDI variational system, enabling variational assimilation of radar reflectivity. Specifically, this study refines a radar reflectivity observation operator to improve its consistency with the double-moment Thompson microphysics scheme, which is used in the current operational HRRR and the next-generation RRFS. The refined operator is implemented in both the GSI and JEDI En3DVar systems, enabling direct variational assimilation of reflectivity in JEDI and facilitating a consistent comparison between the two systems thereby enabling direct variational assimilation of reflectivity within JEDI. A systematic comparison between the original and modified operator formulations is conducted, and their impacts on hydrometeor analysis increments and short-range forecasts are evaluated using a high-impact severe weather event~~convective storm case~~. The JEDI 3DVar forecasts are further compared with forecasts from the operational HRRR and RRFS, which use the previously discussed alternative approaches discussed previously for radar reflectivity assimilation~~The forecasts are further compared with those from the operational HRRR, which employs indirect assimilation of radar reflectivity through cloud analysis and radar-based latent heating during a one-hour spin-up cycle (Dowell et al. 2022; Weygandt et al. 2022), and an experimental RRFS configuration that uses the reflectivity state variable approach~~. The capabilities developed in this study have been incorporated into the official JEDI repository, and this paper serves as technical documentation to support further community evaluation and development.

The remainder of this paper is organized as follows. Section 2 describes the refinement of the reflectivity observation operator. Section 3 introduces the test case, and presents the forecast model and DA configurations, and the design of the DA experiments. Section 4 presents the results, including the evaluations of the refined operator, intercomparisons between the JEDI and GSI systems, and baseline forecast comparisons against independent operational HRRR and RRFS prototype runs from the 2022

Hazardous Weather Testbed Spring Forecast Experiment (NOAA 2022). Section 5 summarizes the key findings and discusses their implications, along with potential directions for future work.

120 **2 Reflectivity observation operator for Thompson microphysics**

The radar reflectivity factor expressed in logarithmic units (dBZ) is defined as

$$Z = 10 \log_{10} Z_e, \tag{1}$$

where Z_e is the equivalent reflectivity factor that contains contributions from various hydrometeor species. The Thompson microphysics scheme employs a double-moment formulation for rainwater that predicts the mixing ratios and total number concentrations. Therefore, Z_e is computed from the mixing ratios of precipitating hydrometeors of rainwater (q_r), snow (q_s), and graupel (q_g), along with the total number concentration of rainwater (N_{tr}), according to

$$Z_e = Z_{er}(q_r, N_{tr}) + Z_{es}(q_s) + Z_{eg}(q_g). \tag{2}$$

The rainwater contribution to the equivalent reflectivity, Z_{er} , is computed from the mixing ratio q_r and the total number concentration N_{tr} (in kg^{-1}), which are explicitly predicted. Here, the equivalent reflectivity factor Z_e is expressed in units of $\text{mm}^6 \text{m}^{-3}$, following Tong and Xue (2005). Assuming spherical raindrops and using the Rayleigh scattering approximation, Z_{er} is calculated as

$$Z_{er} = \frac{10^{18} \times 720 (\rho q_r)^2}{\pi^2 \rho_r^2 N_{tr}}, \tag{3}$$

where ρ (in kg m^{-3}) is the air density, and ρ_r (in kg m^{-3}) is the density of rainwater. The factor 10¹⁸ arises from converting drop diameters from meters to millimeters in the Rayleigh-scattering expression.

In the Thompson microphysics, the snow particle size distribution is represented as a sum of an exponential and a gamma distribution, with parameters that depend on temperature and snow mixing ratio (Thompson et al. 2008). For graupel, the intercept parameter varies with graupel mixing ratio and temperature, and distinct densities are assumed above and below 0 °C (dry/wet states). These scheme-specific details yield piecewise and state-dependent formulations in WRF that are not well suited for constructing smooth, differentiable tangent-linear and adjoint operators required by En3DVar. Therefore, where closed-form mappings exist (e.g., rain and dry snow) we apply them directly, and for wet snow and temperature-dependent graupel we use compact least-squares power-law relations fit to reflectivity diagnosed from the Thompson scheme. This approach preserves consistency with microphysics while providing the smooth derivatives needed for stable variational minimization.

In this study, the contributions of snow and graupel to reflectivity are treated differently than in Liu et al. (2022a), which employed power-law approximations to the model-predicted mixing ratios using fitted curves directly derived from Thompson microphysics output. While parameterized relationships are still used here, a key improvement of the updated operator is the explicit incorporation of temperature-dependent melting effects, which distinguish between wet and dry states of hydrometeors by applying separate formulations above and below the freezing level.

For snow, reflectivity calculations are separated into wet and dry snow conditions based on air temperature. For the wet snow state (air temperature $T > 0^\circ\text{C}$), we developed a simplified power-law function based on the snow mixing ratio q_s , expressed as

$$Z_{es} = a_{wet}(\rho q_s)^{b_{wet}}, \quad (4)$$

where ρ is the air density, and the empirically derived coefficients are $a_{wet} = 1.47 \times 10^5$ and $b_{wet} = 2.67$. These values were obtained by curve fitting to simulated reflectivity data from the Thompson microphysics scheme under wet snow conditions.

For the dry snow state (air temperature $T \leq 0^\circ\text{C}$), the radar reflectivity factor is computed following Thompson et al. (2008) and Field et al. (2005). It is given by

$$Z_{es} = C_{dry} a(T) (\rho q_s)^{b(T)}, \quad (5)$$

where $a(T)$ and $b(T)$ are temperature-dependent coefficient and exponent, respectively, in the power-law moment relation for dry snow derived by Field et al. (2005). and The constant C_{dry} is defined as

$$C_{dry} = \left(\frac{0.176}{0.93} \right) \left(\frac{6}{\pi} \right)^2 \left(\frac{0.069}{900} \right)^2 = 4.06 \times 10^{-9}. \quad (6)$$

For graupel, the equivalent reflectivity factor Z_{eg} is also separated into two regimes depending on air temperature. Following the Thompson microphysics scheme, different graupel densities are assumed for temperatures above and below freezing. Accordingly, we write apply two separate expressions for Z_{eg} as:

$$Z_{eg} = \frac{10^A \times 720 \rho^{1.75} q_g^{2.5}}{\pi^{1.75} \rho_g^B} \begin{cases} \frac{10^{15.25} \times 720 \rho^{1.75} q_g^{2.5}}{\pi^{1.75} \rho_g^{1.85}} & T > 0^\circ\text{C}, \\ \frac{10^{16.5} \times 720 \rho^{1.75} q_g^{2.5}}{\pi^{1.75} \rho_g^{2.3}} & T \leq 0^\circ\text{C}. \end{cases} \quad (7)$$

where ρ_g is the graupel density, Here $A = 15.25$ and $B = 1.85$ for $T > 0^\circ\text{C}$, while $A = 16.5$ and $B = 2.3$ for $T \leq 0^\circ\text{C}$.





Figure 14. Scatterplots comparing the fitted relationships between radar reflectivity (dBZ) and mixing ratio (g kg^{-1}) for (a) snow and (b) graupel predicted using the Thompson microphysics scheme. Black dots show reflectivity values diagnosed directly from WRF model output. The red and blue circles indicate reflectivity computed using the updated and original observation operators, respectively. Coefficients of determination (r^2) measure the agreement between each operator and the WRF-diagnosed reflectivity.

To assess the consistency of the reflectivity observation operator with the Thompson microphysics scheme, reflectivity values computed using the updated and original operators are compared against reflectivity diagnosed directly from WRF (v4.0) model output using the native Thompson microphysics reflectivity calculation (Fig. 1). For snow, the diagnosed reflectivity exhibits a wide-spread at a given mixing ratio, reflecting the complex treatment of snow microphysics in the Thompson scheme, including temperature-dependent behavior and transitions between dry and wet snow. The original operator, which does not explicitly account for these effects, shows poor agreement with the diagnosed values ($r^2 = 0.165$). In contrast, the updated operator introduces separate temperature-dependent relations for dry and wet snow, resulting in a much closer fit to the diagnosed reflectivity, particularly for the wet-snow branch, and substantially improves the agreement ($r^2 = 0.996$).

For graupel, the diagnosed reflectivity also displays considerable spread due to state-dependent variations in particle density and intercept parameter. By explicitly incorporating these temperature- and density-dependent effects, the updated graupel formulation better captures the spread of diagnosed reflectivity and improves the agreement relative to the original operator ($r^2 = 0.968$ versus 0.781). Overall, the updated relations provide compact power-law approximations to the Thompson microphysics while retaining its key physical dependencies and ensuring smooth behavior suitable for tangent-linear and adjoint implementations.


3 Test case and experiment setup

3.1 Case overview

Figure 2 shows the observed composite reflectivity from the Multi-Radar Multi-Sensor (MRMS; Smith et al. 2016) at 0000 UTC 13 May 2022. The event is a robust derecho-producing convective system, characterized by a well-organized, north-south-oriented convective line with a pronounced bowing segment over the Upper Midwest. A negatively tilted upper-level trough over the Midwest, combined with a low-level jet under an omega block pattern, facilitated northward transport of moisture and instability (synoptic-scale figures are not shown for brevity). A surface low formed over the western Midwest with an associated southward-extending cold front and a warm front located in Minnesota.

Severe weather events and observations were reported across the midwestern contiguous United States (CONUS) within the verification domain (black box in Fig. 2). For example, at 2125 UTC 12 May, a Road Weather Information System weather station in Hutchinson County, South Dakota, recorded a wind gust of 107 mph. Additionally, seven tornadoes in South Dakota and four in Minnesota were reported between 2300 UTC 12 May and 0000 UTC 13 May. Between 0000 and 0200 UTC 13

215 May 2022, the primary forecast period of interest in this study, 17 tornadoes and 128 severe wind reports were observed across the north-central CONUS and the Great Plains, according to reports from the Storm Prediction Center.



220 Figure 2. MRMS composite reflectivity (dBZ) at 0000 UTC 13 May 2022, over the FV3-LAM model domain. The bold black box marks the verification domain, and WSR-88D radar locations are indicated by black dots.

3.2 Experiment setup

3.2.1 Model configuration

225 The numerical experiments utilize the FV3-LAM configured with the RRFS_v1beta physics suite from the Common Community Physics Package (CCPP, Bernardet et al. 2024). The physics parameterizations include the Thompson microphysics scheme (Thompson et al. 2008) without aerosol coupling, the Mellor–Yamada–Nakanishi–Niino (MYNN) planetary boundary layer (PBL) scheme (Olson et al. 2019), the Rapid Radiative Transfer Model for GCMs (RRTMG) radiation scheme (Iacono et al. 2008), and the Noah-MP land surface model (Niu et al. 2011). The computational domain covers the midwestern United States, spanning 1050×850 horizontal grid points on an extended Schmidt
230 [gnomonic](#) projection grid at approximately 3-km horizontal grid spacing (Fig. 2). In the vertical, a terrain-following hybrid sigma-pressure coordinate with 65 vertical layers is employed, with the model top located near 2 hPa, consistent with RRFS_v1beta.

Formatted: Centered

Formatted: Font: 12 pt

3.2.2 Data assimilation system

The radar DA experiments are conducted using En3DVar algorithms implemented within the GSI and JEDI frameworks. In both systems, ensemble perturbations for constructing flow-dependent background error covariances in En3DVar are generated using a variant of EnKF: the ensemble square root filter (EnSRF; Whitaker and Hamill 2002) in GSI and the LETKF (Hunt et al. 2007) in JEDI. These ~~ensemble~~ DA cycles are run in parallel with the En3DVar systems, and ~~ensemble~~ forecasts initialized from ensemble analyses in each cycle provide ~~ensemble-the~~ perturbations for En3DVar. Details of the EnVar formulation implemented in JEDI can be found in Liu et al. (2022b). Relaxation to prior spread (RTPS; Whitaker and Hamill 2012) is applied with a coefficient of 0.99 to help maintain the ensemble spread. Ensemble localization is performed using a recursive filter in GSI En3DVar (Purser et al. 2003) and Background error on an Unstructured Mesh Package (BUMP; Ménétrier 2020) in JEDI, with horizontal and vertical localization scales of 18 km and 0.7 ln(P), respectively.

As part of this study, the radar reflectivity observation operator previously implemented within GSI (Liu et al. 2022a) is first migrated into the Unified Forward Operator (UFO) module of JEDI. This operator follows the formulations introduced in section 2 and includes both forward operators and their tangent linear and adjoint components. The implementation enables variational assimilation of reflectivity in JEDI, and these capabilities have been publicly released as part of the JEDI community codebase after testing described in this paper as well as additional testing were completed. The analysis variables updated by the DA include zonal wind, meridional wind, air temperature, specific humidity, layer pressure thickness, [mixing ratio of](#) rainwater, snow, graupel, cloud water, cloud ice, and number concentration of rainwater. To address nonlinearity in the reflectivity observation operator, power-transformed control variables are adopted for hydrometeor fields following Chen et al. (2021) and Li et al. (2022).

Formatted: Indent: First line: 0"

3.2.3 Radar observations

Radar reflectivity (Z) and radial velocity (V_r) observations are assimilated within a ± 3 -minute window around each analysis time. Reflectivity data are prepared from the MRMS mosaic, and radial velocity data from 40 operational WSR-88D radars are used, after automated quality control processes that include noise removal, clutter suppression, despeckling, and velocity dealiasing (Brewster et al. 2005). Observation errors are assumed constant at 5 dBZ for reflectivity and 3 m s⁻¹ for radial velocity. Reflectivity in clear-air regions in the MRMS data are set to 0 dBZ during assimilation, and radial velocity observations are used only where observed reflectivity exceeds 5 dBZ.

3.2.4 Experiment design

The DA experiments consist of a 1-hour model spin-up followed by five hourly assimilation cycles between 1900 UTC 12 May and 0000 UTC 13 May 2022 (Fig. 3). The initial ensemble is generated from a 30-member Global Ensemble Forecast System (GEFS, Zhou et al. 2022) analysis ensemble initialized at 1800 UTC 12 May. Each member is integrated forward using FV3-LAM for one hour to

provide the background for the subsequent assimilation cycle. Lateral boundary conditions are taken from 3-hourly GEFS ensemble forecasts initialized 1800 UTC 12 May.

Four experiments are conducted: (1) NODA, a baseline run without data assimilation; (2) GSI-New, which assimilates radar reflectivity and radial velocity only using EnSRF-updated ensembles and a modified reflectivity operator in GSI En3DVar; (3) JEDI-Org, which assimilates radar observations only using LETKF-updated ensembles and the original reflectivity operator in JEDI En3DVar; and (4) JEDI-New, which is similar to JEDI-Org but uses the modified reflectivity operator. No conventional or surface observations are assimilated in any of the experiments. In the NODA experiment, the model is integrated freely over the same 6-h period without any data assimilation cycles, while the other experiments perform hourly radar data assimilation cycles. In all En3DVar experiments, no static background error covariance is included; thus, the algorithms are pure En3DVar rather than hybrid. For baseline comparison, forecasts from HRRR and RRFs prototype 2 during the 2022 HWT Spring Forecast Experiment are also included. All experiments produce 6-hour deterministic forecasts initialized from the final analyses at 0000 UTC 13 May.




Figure 3. Flow diagram of the radar data assimilation experiment. A 1-hour model spin-up from GEFS ensemble initial conditions is followed by five hourly assimilation cycles between 1900 UTC 12 May and 0000 UTC 13 May 2022. Final analyses are used to launch 6-hour ensemble forecasts.

4 Results and discussions

4.1 Impact of radar data assimilation on En3DVar analyses

Figure 4 shows the root-mean-square innovation (RMSI) and mean innovation diagnostics for assimilated radar reflectivity and radial velocity during DA cycles. Compared to the NODA baseline, all radar assimilation experiments substantially reduce RMSIs for both Z and V_r (Figs. 4a, c), with steady

decreases in Z RMSI throughout the assimilation period (Fig. 4a). Notably, JEDI-New consistently shows improved performance over JEDI-Org, resulting in a reduction in RMSI of approximately 1.5 dBZ in the background and 3 dBZ in the analysis at the final cycle (Figs. 4a, b). This reduction highlights the impact of the updated reflectivity observation operator in JEDI. Mean innovations further confirm this improvement. ~~The large positive reflectivity biases seen in the NODA experiment are substantially reduced in all radar assimilation experiments. The positive mean innovations in NODA indicate that the forecast reflectivity is generally too weak and/or too limited in spatial coverage relative to the observations, although displacement and structural errors also contribute to the overall mismatch. These positive biases in NODA are primarily attributed to displacement and structural errors in the convective system, rather than to systematic overprediction of reflectivity intensity.~~ JEDI-New consistently outperforms JEDI-Org, especially in the background forecasts of reflectivity (Figs. 4e, f).

As an additional sensitivity check, a 15-min cycling experiment was conducted for JEDI-New over the same 1900–0000 UTC DA window (not shown). Although the reflectivity RMSIs do not decrease monotonically after the initial reduction, the analysis RMSI remains consistently lower than the corresponding background RMSI throughout the cycling period, indicating that the DA updates continue to improve the fit to observations and that no evidence of filter divergence is found.

For V_r , all DA experiments reduce the background RMSI relative to NODA (Fig. 4c), and their analysis RMSIs remain around 2.0 m s^{-1} (Fig. 4d). However, JEDI-Org exhibits slightly higher RMSIs and mean innovations than JEDI-New and GSI-New (Figs. 4d and 4h), and difference is the largest in the V_r analyses between JEDI-New and JEDI-Org (Fig. 4d). This result suggests that inaccuracies in the reflectivity analysis associated with the use of a less accurate observation operator can lead to degraded wind analyses through multivariate correlations in the variational update. Notably, JEDI-New and GSI-New produce nearly identical diagnostics throughout the assimilation period, despite differences in the ensemble DA methods and assimilation frameworks. These results indicate that JEDI-New can match the performance of GSI En3DVar when the same improved reflectivity observation operator is used. Since the reflectivity assimilation capabilities in GSI had been tested and reported in previous papers (Chen et al. 2021; Li et al. 2022; Liu et al. 2022a), the comparison serves to demonstrate the validity of our implementation within JEDI.

Formatted: Font: (Asian) +Body Asian (SimSun)

Formatted: Indent: First line: 0"




Figure 4. The observation-space diagnostics from the backgrounds and analyses for the assimilated reflectivity and radial wind observations. For the Z diagnostics, they are calculated when observed Z or calculated $H(x)$ is greater than or equal to 15 dBZ.

Figure 5 compares composite reflectivity fields at the final analysis time (00 UTC 13 May 2022) among different DA experiments with the MRMS observations. The MRMS field (Fig. 5a) features a well-organized convective system across the Upper Midwest, including a north-south-oriented convective line with a bowing structure in the north and a localized curved reflectivity feature in the south. Without radar DA, NODA (Fig. 5e) fails to reproduce the observed convective organization. In particular, it depicts a more merged, connected and narrower linear convective system, rather than the two larger linear systems with larger areas of stratiform precipitation as evident in the MRMS observations. NODA also underrepresents the reflectivity coverage in western Minnesota and near the South Dakota-North Dakota border. Without radar DA, the NODA (Fig. 5e) fails to capture the key storm structures. It misses the echo coverage in western Minnesota and along the South and North Dakota border and displays weak, poorly organized convection that significantly deviates from the observed reflectivity pattern.

In contrast, the DA experiments substantially improve the representation of the convective system. The benefits of the updated reflectivity operator are clearly evident in the background fields. The impacts of the modified reflectivity operator on the background fields are illustrated in Fig. 5. Compared to JEDI-Org (Fig. 5c), JEDI-New (Fig. 5d) exhibits locally stronger reflectivity along portions of the leading convective line and a broader area of moderate reflectivity in the post-line region. In contrast, JEDI-Org shows weaker reflectivity coverage in the trailing stratiform or weak-echo region in this case, leading to differences in the representation of the convective system.

In the analysis fields, both GSI-New (Fig. 5f) and JEDI-New (Fig. 5h) produce well-organized convective structures that closely resemble the MRMS observation, successfully capturing the bowing segment in the north and a curved, localized reflectivity enhancement in the southern portion of the convective line. In contrast, the JEDI-Org analysis (Fig. 5g) shows less coherent storm organization, particularly in the southern segment, where ~~the~~ this reflectivity enhancement hook echo is poorly represented.




Figure 5. The composite reflectivity at final analysis time (00 UTC on May 13, 2022) from the observation and background forecasts and the analyses of the NODA, GSI En3DVar and JEDI En3DVar experiments. The dashed lines inside the panels show the locations of vertical cross sections shown in Figs. 6 and 7. (a) MRMS observations, (b) GSI-New background, (c) JEDI-Org background, (d) JEDI-New background, (e) NODA forecast, (f) GSI-New analysis, (g) JEDI-Org analysis, and (h) JEDI-New analysis. The dashed line in each panel indicates the location of the vertical cross sections shown in Figs. 6 and 7. Because no data assimilation is performed in NODA, no analysis field is available for that experiment.

Figure 6 shows vertical cross sections of composite reflectivity at the final analysis time (00 UTC 13 May 2022) along the dashed line in Figure 5. The MRMS observation (Fig. 6a) displays a well-developed deep convective system, featuring two upright convective towers: an dominant eastern one tower near 95°W and a secondary western one near 99.6-5-97°W. Both towers exhibit echo tops

exceeding 14 km and strong reflectivity cores (≥ 50 dBZ) through the mid-levels. In contrast, the NODA analysis (Fig. 6e), which does not assimilate radar observations, shows a convective reflectivity structure that is less well aligned with the fixed cross section relative to MRMS, reflecting differences in the analyzed system position. As a result, the main convective core is located outside the cross-section domain shown in Fig. 5e, leading to the apparent absence of deep echoes in the vertical structure.

In the background fields, all DA experiments reproduce the main convective tower near 95°W but overestimate its midlevel reflectivity intensity. This bias is most evident in JEDI-New (Fig. 6d), where the eastern convective column exhibits stronger reflectivity (≥ 50 dBZ) from about 2 to 8 km, ~~compared with GSI-New (Fig. 6b) and JEDI-Org (Fig. 6e).~~ These overamplified cores are likely related to excessive graupel reflectivity ~~associated with the Thompson microphysics scheme,~~ consistent with findings in Liu et al. (2022a) and Grim et al. (2023). Additionally, all three background fields slightly underpredict the top of the precipitation region, with the 5-dBZ echo top generally reaching only about 11–12 km, compared with about 12–13 km in MRMS. The western ~~secondary~~ tower near 99°W that is evident in the MRMS cross section is not clearly reproduced in any of the background fields.

The analysis fields indicate that radar DA partially recovers the previously missing western convective feature. In JEDI-New (Fig. 6h), the ~~secondary-western~~ tower is more evident than in the background fields and the NODA experiment, although the tower is too short in all experiments compared to MRMS ~~differences remain relative to MRMS.~~ GSI-New (Fig. 6f) and JEDI-Org (Fig. 6g) additionally underestimate its strength. In the ~~primary-eastern~~ tower region near 95°W, GSI-New and JEDI-New also effectively reduces the midlevel overestimation seen in the background, but the resulting structure looks arguably better in JEDI-New compared to GSI-New. By contrast, JEDI-Org retains an overly strong midlevel core.




Figure 6. Vertical cross sections of composite reflectivity at the final analysis time (00 UTC 13 May 2022) along the dashed line shown in Fig. 5, from [\(a\) MRMS observations](#).

Formatted: Font: (Asian) +Body Asian (SimSun), (Asian) Chinese (Simplified, Mainland China)

(b) GSI-New background, (c) JEDI-Org background, (d) JEDI-New background, (e) NODA forecast, (f) GSI-New analysis, (g) JEDI-Org analysis, and (h) JEDI-New analysis. the MRMS observation, and the background and analysis fields of the NODA, GSI-New, JEDI-Org, and JEDI-New experiments.


To further understand the reflectivity differences noted in Fig. 6, we examine vertical cross sections of the analysis increments for rainwater, snow, and graupel mixing ratios shown in Fig. 7. As noted in Fig. 6, the background fields in all three DA experiments slightly underpredict the top of the precipitation region, while the analysis update increases the storm-top height and brings it closer to that shown in MRMS. Since radar reflectivity is a nonlinear function of multiple hydrometeor species, the hydrometeor analysis increments help explain both the changes in analyzed reflectivity structure and the increased vertical extent of the convective system. Since radar reflectivity is a nonlinear function of multiple hydrometeor species, differences in its assimilation impacts can help explain the variations in analyzed reflectivity structures.

Among the three hydrometeor species, snow shows the most prominent differences between JEDI-New and JEDI-Org. In the snow analysis increments (Figs. 7b, e, h), JEDI-Org (Fig. 7b) exhibits weak and spatially fragmented increments, suggesting minimal relatively limited adjustments to the snow field. In contrast, JEDI-New (Fig. 7e) produces broad and coherent positive snow increments in the mid- to upper levels (model levels ~25–45), especially over and between the main eastern and western convective towers. The magnitude and coverage of these increments are comparable to those from GSI-New (which uses the updated Z operator) (Fig. 7h), indicating that the updated reflectivity operator in JEDI-New enables more effective retrieval and correction of snow content, and that these positive upper-level snow increments are consistent with the increased storm-top height seen in the analysis cross sections of Fig. 6. This contributes significantly to the improved vertical structure of reflectivity shown in Fig. 6, which contributes significantly to the improved vertical structure of reflectivity shown in Figure 6.

For graupel (Figs. 7c, f, i), all three DA experiments show negative increments collocated with the eastern convective tower. The negative graupel increments are weaker and more localized in JEDI-Org (Fig. 7c), whereas JEDI-New (Fig. 7f) and GSI-New (Fig. 7i) exhibit stronger and more spatially coherent negative increments centered in the midlevels of the primary eastern tower. This indicates a reduction in excessive background graupel, which likely contributes to the overly strong midlevel reflectivity in the background field. In contrast, the rainwater increments (Figs. 7a, d, g) are relatively weak and similar across all experiments, reflecting the fact that the treatment of rainwater is unchanged between the original and updated operators. However, the positive rainwater increments near the western tower appear to contribute to the restored reflectivity structure and vertical development of that tower in the analyses, which was underrepresented in the background fields of all DA experiments. Thus, while differences among experiments still arise mainly from the handling of snow and graupel rather than from rainwater, the rainwater increments also play a non-negligible role in improving the western tower. Thus, differences among experiments arise mainly from the handling of snow and graupel rather than from rainwater.

Formatted: Indent: First line: 0"

445 ~~These results confirm that the improved reflectivity analysis in JEDI-New is primarily associated~~
~~with the refined representation and updating of the snow and graupel fields. The updated Z~~
~~observation operator ensures greater consistency with the Thompson microphysics scheme,~~
~~particularly in the treatment of frozen hydrometeors, leading to analyses that better align with~~
~~model-computed reflectivity fields and more accurate reconstructions of convective vertical~~
~~structures.~~






Figure 7. The vertical cross sections along the line given in Fig. 5 of analysis increments (kg kg^{-1}) for hydrometeor mixing ratios of rainwater, snow, and graupel at final analysis time (00 UTC on May 13, 2022). The three columns correspond to rainwater, snow, and

Formatted: Font: (Asian) +Body Asian (SimSun), (Asian) Chinese (Simplified, Mainland China)

Formatted: Indent: First line: 0"

[graupel, respectively, and the three rows correspond to JEDI-Org, JEDI-New, and GSI-New, respectively. from GSI En3DVar and two JEDI En3DVar experiments.](#)

460 4.2 Impacts on forecasts

For qualitative assessment of forecast performance, the forecasts initialized from the final En3DVar analyses from GSI-new and JEDI-new are compared with those from the no-DA baseline experiment (NODA), and two reference forecasts representative of the current and potential future operational configurations, i.e., the HRRR version 4 (HRRRv4) and the RRFS prototype 2 control run (RRFSp2 CNTL). As shown in Fig. 8, the NODA experiment fails to reproduce key convective features observed in the MRMS composite reflectivity. In the 1-h forecast, it entirely misses the convective line over Minnesota and instead produces a broad, unrealistically merged system (Fig. 8d). The forecast also exhibits notable displacement of the bow echo structures across the central CONUS throughout the 3 and 5 hours of forecast (Figs. 8e–f). In contrast, both the GSI and JEDI En3DVar DA experiments successfully capture the convective line in Minnesota at 1 h (Figs. 8g, j) and reproduce a trailing stratiform precipitation region behind the leading convective line. At later lead times, the En3DVar experiments maintain a better alignment of high-reflectivity structures over Kansas and Nebraska (Figs. 8h–i, k–l), although some degradation in intensity and coverage is evident compared to the 1-h forecasts.

475 The two reference forecasts, HRRRv4 and RRFS p2 CNTL, also capture the general structure of the observed convective systems, particularly the split into northern and southern convective segments. However, positional and structural differences are apparent upon closer inspection. The HRRRv4 forecast shows a weaker bow echo and a southeastward displacement of the Minnesota line as early as 1 h (Fig. 8m), with this error persisting through 5 h. Compared to HRRRv4, RRFS p2 CNTL better captures the structure of the northern convective line at 1 h, but tends to underpredict reflectivity intensity and spatial extent, especially in the southern segment at longer lead times. Additionally, both reference forecasts display smaller overall high-reflectivity coverage relative to those of En3DVar experiments. This difference is likely attributable to the use of cloud analysis for radar DA in HRRRv4 and the Z-state-variable approach in RRFS p2 CNTL, both of which differ from the direct reflectivity assimilation employed in the En3DVar experiments here. Nonetheless, spurious echo generation is evident in the En3DVar runs at later lead times, particularly over the northwestern portion of the domain (Figs. 8f, i, l), where weak, largely non-convective reflectivity appears that is not present in the MRMS observations. These differences may also reflect intrinsic model characteristics; composite reflectivity forecasts from FV3-LAM tend to be stronger than those from HRRRv4, which is based on the WRF model, consistent with findings in Grim et al. (2023). Overall, the results shows the benefits of directly assimilating reflectivity and radial velocity in capturing key convective structures in the analysis and forecast while also highlighting certain sensitivities to model and assimilation methodologies.

Formatted: Indent: First line: 0"




Figure 8. Composite reflectivity at 1 h (left column), 3 h (middle column) and 5 h (right column) from MRMS observation (first row), and forecasts from NODA, GSI-New, JEDI-New experiments, and from HRRRv4, and RRFSp2 CNTL as labeled. Observed composite reflectivity of 40 dBZ is contoured in black in the forecasts. Composite reflectivity forecasts and observations at 1, 3, and 5 h lead times, valid at 0100, 0300, and 0500 UTC 13 May 2022, respectively. Rows show MRMS observations, NODA, GSI-New, JEDI-New, HRRRv4, and RRFS p2 CNTL, and columns show the 1-, 3-, and 5-h forecasts. In the forecast panels, the observed 40-dBZ composite reflectivity contour is overlaid in black.

Formatted: Centered

Formatted: Font: (Asian) +Body Asian (SimSun), (Asian) Chinese (Simplified, Mainland China), (Other) English (United States)




Figure 8. Continued.

Although radar reflectivity assimilation primarily constrains hydrometeor fields, the maintenance and evolution of convective systems also depend on adjustments to thermodynamic and kinematic variables such as temperature, water vapor, pressure, and wind. Since 2-m equivalent potential temperature θ_e and 10-m wind fields are only indirectly impacted by radar DA (via the cross covariances), the fields are partly the results of model response to other more directly impacted fields, through the DA cycles. We examine these fields in the final analyses to see if the DA is producing physically consistent near-surface analyses, in particular in the presentation of the cold pools.

Figure 9 shows 2-m θ_e and 10-m wind fields from UnRestricted Mesoscale Analysis (URMA, NOAA 2018) and the final analyses from the NODA, GSI-New, and JEDI-New experiments and GSI and JEDI En3DVar DA experiments. Compared to URMA (Fig. 9a), NODA (Fig. 9b) exhibits a less expansive region of low- θ_e air associated with the cold pool, and its leading edge or gust-front zone remains closer to the South Dakota–Minnesota border. In contrast, both DA experiments (Figs. 9c,d) show broader low- θ_e coverage and a farther southward extension of the near-surface cold-pool boundary in better agreement with URMA, produces cold pools that are much smaller in extent than URMA, and the leading edge of the cold pool or the gust front stays close to the South Dakota–Minnesota border while those in URMA and the two DA experiments are much closer in cold-pool extents and gust front positions (Fig. 9a,e,d). In the DA experiments, relatively stronger surface winds are present in the cold pool immediately behind the gust front compared to than in NODA, and this feature is more consistent with the analyzed near-surface pattern and the overall cold-pool structure qualitatively resembles that in URMA (Figs. 9c, and 9d). These results indicate that radar DA can indirectly improve near-surface temperature, moisture, and wind analyses by improving the placement and structure of the convective system through the DA cycles. This behavior is consistent with cold pool dynamics, in which stronger cold pools are associated with enhanced near-surface outflow winds.

Further south and east of the gust front, over Kansas, Missouri, and Iowa, the DA experiments exhibit stronger southerly inflow relative to NODA, with flow patterns that are qualitatively closer to URMA. The improved analyses of near surface fields indicate that the radar DA can indirectly improve surface temperature, moisture, and wind fields by leveraging cross-variable covariances derived from the ensemble. The relatively low θ_e over northwest Nebraska and southern South Dakota in both NODA and the DA experiments is likely associated with the synoptic-scale frontal environment. This interpretation is consistent with the Weather Prediction Center surface analysis at 00 UTC, which depicts a cold front and an associated low-pressure system extending across the Dakotas, accompanied by westerly to northwesterly postfrontal flow. This suggests that the θ_e bias in this region is more closely related to errors in the large-scale air-mass structure, such as the frontal position or intensity, which are not directly constrained by radar DA. Assimilating additional conventional surface observations may help reduce these near-surface thermodynamic biases (e.g., Schenkman et al. 2011).




Figure 9. Equivalent Potential Temperature (shaded; K) at 2 m and wind vectors at 10 m (m s⁻¹) for (a) URMA, (b) NODA, (c) GSI-New, and (d) JEDI-New. Black contours denote areas where observed MRMS composite reflectivity exceeds 30 dBZ and are included in each panel as a common observational reference. ~~in plots denote the areas of observed~~

composite reflectivity exceeding 30 dBZ threshold. Yellow contours show relative vorticity exceeding $100 \times 10^{-5} \text{ s}^{-1}$ from model level 1 (lowest model level).

Figure 10 presents the fractions skill scores (FSS; Roberts and Lean 2008) of forecast composite reflectivity, computed at hourly intervals and averaged over the 6 hours of verification period, for the 25 and 45 dBZ thresholds. Both DA experiments consistently outperform HRRRv4, RRFS p2 CNTL, and NODA across all tested neighborhood radii (up to 72 km). GSI-New achieves the minimum skillful threshold at a radius of 57 km for the 45 dBZ threshold, while JEDI-New reaches it at 66 km. For the 25 dBZ threshold, both DA experiments exceed the minimum skillful threshold at the smallest evaluated neighborhood radius (3 km), suggesting improved representation of convective-scale features in a neighborhood sense when averaged over the 6-h verification period. The NODA forecasts are non-skillful for all tested radii at all radii for the 45 dBZ, and dBZ and are only skillful at radii larger than 64 km for the 25 dBZ threshold.

The two reference forecasts also fail to reach the skillful threshold within the tested range of radii at all radii for the 45 dBZ case, though their FSSs are above those of NODA for radii above 21 km and the differences increase with neighborhood radius (Fig. 10b). The reduced difference in FSS between our DA experiments and the reference forecasts at larger radii may be partly due to reflectivity bias. As discussed in Mittermaier and Roberts (2010), models that underpredict high reflectivity can appear more skillful at large spatial scales because FSS emphasizes spatial agreement over intensity accuracy. This possible bias effect is also consistent with the precipitation performance diagrams discussed below.




Figure 10. Fractions skill scores (FSS) of hourly composite reflectivity averaged over a 6-h verification period, shown as a function of neighborhood radius (km) for reflectivity thresholds of (a) 25 dBZ and (b) 45 dBZ. Verification is performed against MRMS

Formatted: Font: (Asian) +Body Asian (SimSun), (Asian) Chinese (Simplified, Mainland China)

Formatted: Indent: First line: 0"

Formatted: Centered

575 observations. Results are shown for HRRRv4 (orange), RRFS p2 CNTL (gray), JEDI-New
(blue), GSI-New (red), and NODA (black). Thin horizontal line in each panel indicates the
minimum skillful forecast threshold.

580 Figure 11 presents performance diagrams for hourly accumulated precipitation again averaged over
the 6 h forecast period, verified against the NCEP Stage IV quantitative precipitation estimates (Lin and
Mitchell 2005) at thresholds of 0.04 in. (≈ 1.0 mm) and 0.25 in. (≈ 6.4 mm). At the lower threshold (Fig.
11a), the GSI-New and JEDI-New experiments demonstrate the highest critical success index (CSI),
both exceeding 0.6. GSI-New exhibits slightly better skill than JEDI-New, with a CSI above 0.65 and a
585 small high bias of close to about 1.1. JEDI-New has a similar high bias, also close to 1.1, whereas
HRRRv4, RRFS p2 CNTL, and NODA have lower biases of about 0.8. HRRRv4 achieves a CSI of
approximately 0.58, while RRFS p2 CNTL and NODA trail behind with CSI values near 0.55 and 0.52,
respectively. The DA experiment forecasts demonstrate a favorable combination of detection rate and
false alarm control, indicating improved representation of precipitation extent and location compared to
other forecasts.

590 At the higher threshold of 6.35 mm (Fig. 11b), forecast skill decreases for all systems, but the
relative ranking remains the same. Both DA experiments maintain CSI values near 0.5, while the
reference forecasts drop to around 0.4, and NODA falls below 0.3. The DA experiments also display
higher bias (~ 1.4), suggesting more aggressive precipitation production, consistent with the more active
reflectivity fields noted in Fig. 8. In contrast, HRRRv4 exhibits a low bias below 1.0 (of ~ 0.8),
595 indicating a tendency to underpredict intense precipitation amounts, while RRFS p2 CNTL remains
closer to unity. Because these forecast systems differ in both dynamical core and DA method, the
present comparison does not allow us to disentangle their respective contributions to the bias
differences. These differences may reflect variations in model configurations and DA approaches; our
DA experiments and RRFS p2 CNTL use the FV3-LAM model, which assimilate reflectivity using
different approaches while the WRF-based HRRRv4 uses cloud analysis for reflectivity DA and
600 produces weaker precipitation overall. The tendency for FV3-LAM to overpredict precipitation and
convective activity has been noted in both cool-season (Supinie et al. 2022) and warm-season ensemble
forecasts during the NOAA HWT Spring Forecasting Experiment (Johnson et al. 2025).

Formatted: Indent: First line: 0"

Commented [MX1]: bias above 1 is high bias, low bias if below 1.

Formatted: Indent: First line: 0"




Figure 11. Performance diagrams of hourly accumulated precipitation for a) 1.016 mm and b) 6.35 mm thresholds over the 6-hr forecast period. HRRRv4 (orange square), RRFS p2 CNTL (gray square), JEDI-New (blue circle), GSI-New (red circle), and NODA (black circle) are verified against Stage-IV observation. Black dashed lines and solid contours denote bias and CSI, respectively.

As described in section 3, our test case involved multiple tornadoes. For convection-allowing predictions, updraft helicity (UH; Kain et al. 2008) is a widely used proxy for diagnosing meso-cyclonic rotation of strong updrafts associated with severe convection, including tornadoes and large hail (Sobash et al. 2011; Gallo et al. 2016). Figure 12 shows the neighborhood probability (NP) of 2-h maximum 2-5 km UH from 0000 to 0200 UTC 13 May 2022 that exceeds $75 \text{ m}^2 \text{ s}^{-2}$, overlaid with Storm Prediction Center reports of tornados, hail, and strong winds. The NP fields are computed using a 42-km neighborhood and subsequently smoothed with an 81-km Gaussian filter (e.g., Tong et al. 2020; Park et al. 2023). Compared to NODA and the reference forecasts, both GSI-New and JEDI-New produce show better spatial alignment between the forecast probability maxima and higher neighborhood probabilities and improved spatial alignment with the observed severe reports. GSI-New shows a well-defined swath of probabilities exceeding 560% extending from northern Iowa into southern Minnesota, broadly collocated with numerous tornado and wind reports. JEDI-New exhibits a similar probability distribution, but with a slightly narrower core. JEDI-New shows a similar pattern to GSI-New, with a slightly narrower probability core, while both DA experiments capture the main swath of elevated probabilities collocated with the clusters of severe storm reports.

The reference forecasts demonstrate weaker performance. HRRRv4 exhibits a reasonably placed UH signal, but with substantially lower probabilities (peak NP $\sim 50\%$), likely attributable to the use of the WRF model rather than FV3-LAM; the latter is known to produce higher UH values (Potvin et al.

Formatted: Indent: First line: 0"

2019). RRFS p2 CNTL generates a broader, westward-shifted UH swath centered over southeastern North Dakota, missing many of the observed reports across Iowa and southern Minnesota, including a peak-probability region in southeastern North Dakota associated with few severe reports and probabilities below 10% in a cluster of hail and wind reports over eastern Nebraska. The NODA experiment broadly captures the spatial envelope of the observed severe weather reports, but the UH probabilities are too diffuse and exhibit a southwestward displacement relative to the report clusters, particularly over southwestern Minnesota. These results again demonstrate the positive impacts of directly assimilating radar reflectivity and radial velocity data, using the GSI- and JEDI-based En3DVar, in capturing the location and strength of rotating updrafts associated with severe weather events.




Figure 12. Neighborhood probability (%; shaded) of 2-hr maximum updraft helicity exceeding 75 m2 s-2 from 00 to 02 UTC on May 13, 2022 from the forecasts of a) NODA,

b) GSI-New, c) JEDI-New, d) HRRRv4 and e) RRFS p2 CNTL. The red, green, and blue dots denote the SPC reports of tornado, hail, and wind, respectively.

5. Summary and conclusions

This study presents the first implementation and evaluation of direct radar reflectivity and radial velocity assimilation within the JEDI En3DVar system, using a microphysics-consistent observation operator for the Thompson double-moment scheme. A high-impact squall-line event that occurred over the Midwest on 13 May 2022 was selected as a test case to assess the performance of the JEDI En3DVar system, using the GSI counterpart and operational forecasts from HRRR and experimental RRFS as references/benchmarks.

The DA experiments assimilated radar observations over a 5-hour window, starting from 1-hour spin-up ensemble forecasts initialized at 1800 UTC and assimilated data at hourly intervals using ensemble covariances generated by parallel ensemble-based DA systems. An updated radar reflectivity operator, consistent with the Thompson microphysics scheme, was used to directly assimilate reflectivity observations into both JEDI and GSI En3DVar systems. Compared to an original operator carried over from GSI En3DVar, the updated version substantially reduced root-mean-square innovations and biases for reflectivity in both the background forecast and analysis fields. These improvements stemmed from enhanced treatment of snow and graupel contributions, which led to more consistent hydrometeor adjustments during analysis. As demonstrated by Liu et al. (2022a), the physical consistency between the operator and forecast model is crucial for reducing spurious analysis increments and improving assimilation stability.

Forecasts initialized from JEDI and GSI En3DVar analyses using the update reflectivity operator captured two primary mesoscale convective systems with comparable skill. Both systems produced more intense cold pools and better-defined surface wind structures by 0000 UTC, which contributed to improved verification scores for reflectivity and precipitation forecasts over the first 6 hours across multiple thresholds. Additionally, the JEDI-based forecasts showed greater spatial alignment between updraft helicity ($UH \geq 75 \text{ m}^2 \text{ s}^{-2}$) and observed severe weather reports during the first two forecast hours. Despite differences in ensemble generation, localization strategy, and minimization procedures between the JEDI and GSI En3DVar systems, their analyses and short-range forecasts were similarly skillful, indicating that JEDI En3DVar is a viable alternative to the more extensively tested GSI En3DVar for radar DA.

When compared against operational forecasts from HRRR and experimental RRFS, which employed cloud analysis and a reflectivity-state-variable En3DVar approaches, respectively, both JEDI and GSI-based experiments demonstrated slightly improved quantitative forecast skill for reflectivity and precipitation. The probabilistic forecasts of UH also aligned more closely with observed severe weather locations, reinforcing the benefit of direct radar DA in convective-scale systems while demonstrating the implementation correctness of the radar DA capabilities in JEDI En3DVar.

Although the present results are encouraging, they are based on a single convective event and a limited cycling configuration. Broader evaluation using additional cases and extended cycling would be

Formatted: Indent: First line: 0"

valuable to assess robustness across different storm types and to further characterize spin-up behavior of the innovation diagnostics and is left for future work.

While this study uses FV3-LAM, NOAA has initiated a transition toward the Model for Prediction Across Scales (MPAS) dynamical core for RRFS version 2 (Alexander et al., 2023). Similar tests that couple JEDI En3DVar with MPAS-based convective-scale forecasts will be an important next step.

Code and data availability. The source code of JEDI-FV3 version 1.0.0 used in this study is available on Zenodo at <https://doi.org/10.5281/zenodo.18272913> (Park and Liu, 2026). For reproducibility, the processed radar observations in IODA netCDF format and the YAML configuration files used in the JEDI EnVar experiments are available on Zenodo at <https://doi.org/10.5281/zenodo.18272368> (Park and Liu, 2026). Global Ensemble Forecast System (GEFS) ensemble analyses and forecasts are available from the NOAA National Centers for Environmental Information (NCEI) at <https://www.ncei.noaa.gov/products/weather-climate-models/global-ensemble-forecast> (last access: 16 January 2026; NOAA, 2026). Reflectivity data from the NOAA Multi-Radar Multi-Sensor (MRMS) system are available from NOAA NCEI at <https://mrms.nssl.noaa.gov/> (last access: 16 January 2026). NEXRAD Level II radial velocity data are available from the NOAA NCEI radar archive at <https://www.ncei.noaa.gov/products/radar/next-generation-weather-radar> (last access: 16 January 2026). These MRMS and NEXRAD datasets are also accessible via the NOAA Open Data Program, including through Amazon Web Services (AWS), at <https://registry.opendata.aws/noaa-mrms-pds> and <https://registry.opendata.aws/noaa-nexrad/>.

Author contributions. JP developed the data assimilation codes, conducted the experiments, performed the diagnostics and plotting, and led the manuscript writing. CL designed the overall study and experiments, analyzed the results, and contributed to the writing. MX contributed to the interpretation of the results and manuscript revision.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Acknowledgments. The authors acknowledge the Texas Advanced Computing Center (TACC) at the University of Texas at Austin for providing high-performance computing resources on the Stampede2 supercomputer through the NSF XSEDE and ACCESS programs.

Financial support. This research has been supported by the National Oceanic and Atmospheric Administration (NOAA) through the Joint Technology Transfer Initiative (JTTI) program (grant no. NA21OAR4590171) and the Research-to-Operations (R2O) program (grant no. NA21OAR4320204).

Appendix A: Brief formulation of the incremental En3DVar used in this study

In this study, both JEDI and GSI use an incremental En3DVar formulation. Following Liu et al. (2022b), the incremental cost function is written as

Formatted: Font: Bold

$$J(\delta x) = \frac{1}{2}(\delta x - \delta x^g)^T \mathbf{B}^{-1}(\delta x - \delta x^g) + \frac{1}{2}(\mathbf{H}\delta x - d)^T \mathbf{R}^{-1}(\mathbf{H}\delta x - d),$$

where δx is the analysis increment, $\delta x^g = x^b - x^g$, x^b is the background state, x^g is the first guess, $d = y - h(x^g)$ is the innovation vector, $\mathbf{H}$ is the linearized observation operator, and $\mathbf{B}$ and $\mathbf{R}$ are the background and observation error covariance matrices, respectively.

The gradient of the cost function with respect to δx is

$$\nabla_{\delta x} J(\delta x) = \mathbf{B}^{-1}(\delta x - \delta x^g) + \mathbf{H}^T \mathbf{R}^{-1}(\mathbf{H}\delta x - d).$$

Setting the gradient to zero yields the linear system for the analysis increment:

$$(\mathbf{B}^{-1} + \mathbf{H}^T \mathbf{R}^{-1} \mathbf{H}) \delta x^a = \mathbf{B}^{-1} \delta x^g + \mathbf{H}^T \mathbf{R}^{-1} d.$$

In this study, no static background covariance is included, so the En3DVar configuration is a pure En3DVar, with the background error covariance represented by the localized ensemble covariance,

$$\mathbf{B} = \mathbf{L} \circ \mathbf{B}_e,$$

where $\mathbf{B}_e$ is the ensemble covariance estimated from the ensemble perturbations and $\mathbf{L}$ denotes the localization matrix. The two systems differ mainly in the ensemble generation and localization implementation: GSI uses EnSRF perturbations together with recursive-filter localization, whereas JEDI uses LETKF perturbations and BUMP-based localization. More complete implementation details of the JEDI EnVar framework are given in Liu et al. (2022b).

References

- Alexander, C. R., Carley, J. R., and Pyle, M. E.: The Rapid Refresh Forecast System: Looking Beyond the First Operational Version, in: *Unified Forecast System Implementations and Future Capabilities Workshop (UFCW 2023)*, Boulder, CO, USA, 25 July 2023, available at: https://epic.noaa.gov/wp-content/uploads/2023/08/UFCW-2023-Tue-9.-Alexander_UFS_UFCW_2023_Final-2.pdf (last access: 10 November 2025), 2023.
- Banos, I. H., Mayfield, W. D., Ge, G. Q., Sapucci, L. F., Carley, J. R., and Nance, L.: Assessment of the data assimilation framework for the Rapid Refresh Forecast System v0.1 and impacts on forecasts of a convective storm case study, *Geosci. Model Dev.*, 15, 6891–6917, <https://doi.org/10.5194/gmd-15-6891-2022>, 2022.
- Bernardet, L., Bengtsson, L., Reinecke, P. A., Yang, F., Zhang, M., Hall, K., Doyle, J., Martini, M., Firl, G., and Xue, L.: Common Community Physics Package: Fostering collaborative development in physical parameterizations and suites, *Bull. Am. Meteorol. Soc.*, 105, E1490–E1505, <https://doi.org/10.1175/BAMS-D-23-0227.1>, 2024.
- Bishop, C. H., Whitaker, J. S., and Lei, L.: Gain form of the ensemble transform Kalman filter and its relevance to satellite data assimilation with model space ensemble covariance localization, *Mon. Weather Rev.*, 145, 4575–4592, <https://doi.org/10.1175/MWR-D-17-0102.1>, 2017.

Formatted: Font: Bold

- Brewster, K., Hu, M., Xue, M., and Gao, J.: Efficient assimilation of radar data at high resolution for short-range numerical weather prediction, in: *World Weather Research Programme Symposium on Nowcasting and Very Short-Range Forecasting (WSN05)*, Toulouse, France, WMO, 2005.
- Carley, J., Alexander, C., Wicker, L., Jablonowski, C., Clark, A., Nelson, J., Jirak, I., and Viner, K.: Mitigation efforts to address rapid refresh forecast system (RRFS) v1 dynamical core performance issues and recommendations for RRFS v2, ON516, <https://doi.org/10.25923/ccgj-7140>, 2023.
- Chen, L., Liu, C., Xue, M., Zhao, G., Kong, R., and Jung, Y.: Use of power transform mixing ratios as hydrometeor control variables for direct assimilation of radar reflectivity in GSI En3DVar and tests with five convective storm cases, *Mon. Weather Rev.*, 149, 645–659, <https://doi.org/10.1175/MWR-D-20-0149.1>, 2021.
- Courtier, P., Thépaut, J. N., and Hollingsworth, A.: A strategy for operational implementation of 4D-Var, using an incremental approach. *Quarterly Journal of the Royal Meteorological Society*, 120(519), 1367–1387, <https://doi.org/10.1002/qj.49712051912>, 1994.
- Dowell, D. C., Alexander, C. R., James, E. P., Weygandt, S. S., Benjamin, S. G., Manikin, G. S., Blake, B. T., Brown, J. M., Olson, J. B., Hu, M., Smirnova, T. G., Ladwig, T., Kenyon, J. S., Ahmadov, R., Turner, D. D., Duda, J. D., and Alcott, T. I.: The High-Resolution Rapid Refresh (HRRR): An hourly updating convection-allowing forecast model. Part I: Motivation and system description, *Wea. Forecasting*, 37, 1371–1395, <https://doi.org/10.1175/WAF-D-21-0151.1>, 2022.
- Evensen, G.: The ensemble Kalman filter: Theoretical formulation and practical implementation. *Ocean dynamics*, 53(4), 343–367, <https://doi.org/10.1007/s10236-003-0036-9>, 2003.
- Field, P. R., Hogan, R. J., Brown, P. R. A., Illingworth, A. J., Choullarton, T. W., and Cotton, R. J.: Parametrization of ice-particle size distributions for mid-latitude stratiform cloud. *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 131(609), 1997–2017, <https://doi.org/10.1256/qj.04.134>, 2005.
- Gallo, B. T., Clark, A. J., and Dembek, S. R.: Forecasting tornadoes using convection-permitting ensembles, *Wea. Forecasting*, 31, 273–295, <https://doi.org/10.1175/WAF-D-15-0134.1>, 2016.
- Gao, J., Heinselman, P. L., Xue, M., Wicker, L. J., Yussouf, N., Stensrud, D. J., and Droegemeier, K. K.: The numerical prediction of severe convective storms: Advances, challenges, and outlook, in: *Elsevier Reference Collection in Earth Systems and Environmental Sciences*, Elsevier, 1–22, <https://doi.org/10.1016/B978-0-323-96026-7.00127-2>, 2024.
- Grim, J. A., Pinto, J. O., and Dowell, D. C.: Assessing RRFS versus HRRR in predicting widespread convective systems over the eastern CONUS, *Wea. Forecasting*, 39, 121–140, <https://doi.org/10.1175/WAF-D-23-0112.1>, 2024.
- Houtekamer, P. L., and Zhang, F.: Review of the ensemble Kalman filter for atmospheric data assimilation, *Mon. Weather Rev.*, 144, 4489–4532, <https://doi.org/10.1175/MWR-D-15-0440.1>, 2016.
- Hu, M., Benjamin, S. G., Ladwig, T. T., Dowell, D. C., Weygandt, S. S., Alexander, C. R., and Whitaker, J. S.: GSI three-dimensional ensemble–variational hybrid data assimilation using a global ensemble for the regional Rapid Refresh model, *Mon. Weather Rev.*, 145, 4205–4225, <https://doi.org/10.1175/MWR-D-16-0418.1>, 2017.

- 800 Hunt, B. R., Kostelich, E. J., and Szunyogh, I.: Efficient data assimilation for spatiotemporal chaos: A
local ensemble transform Kalman filter, *Physica D*, 230, 112–126,
<https://doi.org/10.1016/j.physd.2006.11.008>, 2007.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.:
Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer
805 models, *J. Geophys. Res. Atmos.*, 113, D13103, <https://doi.org/10.1029/2008JD009944>, 2008.
- Johnson, M., Snook, N., Park, J., Xue, M., Brewster, K. A., Supinie, T., and Hu, X.: Severe weather
verification of an FV3-LAM regional ensemble during the 2022 NOAA Hazardous Weather
Testbed Spring Forecasting Experiment, *Wea. Forecasting*, 40, 577–592,
<https://doi.org/10.1175/WAF-D-24-0034.1>, 2025.
- 810 Kain, J. S., et al.: Some practical considerations regarding horizontal resolution in the first generation of
operational convection-allowing NWP, *Wea. Forecasting*, 23, 931–952,
<https://doi.org/10.1175/WAF2007106.1>, 2008.
- Kong, R., Xue, M., and Liu, C.: Development of a hybrid En3DVar data assimilation system and
comparisons with 3DVar and EnKF for radar data assimilation with observing system simulation
815 experiments, *Mon. Weather Rev.*, 146, 175–198, <https://doi.org/10.1175/MWR-D-17-0164.1>, 2018.
- Labriola, J., Jung, Y., Liu, C., and Xue, M.: Evaluating forecast performance and sensitivity to the GSI
EnKF data assimilation configuration for the 28–29 May 2017 mesoscale convective system case,
Wea. Forecasting, 36, 127–146, <https://doi.org/10.1175/WAF-D-20-0071.1>, 2021.
- 820 Li, H., Liu, C., Xue, M., Park, J., Chen, L., Jung, Y., Kong, R., and Tong, C.-C.: Use of power-
transform total number concentration as control variable for direct assimilation of radar reflectivity
in GSI En3DVar and tests with six convective storms cases, *Mon. Weather Rev.*, 150, 821–842,
<https://doi.org/10.1175/MWR-D-21-0041.1>, 2022.
- Lin, S.-J.: A “vertically Lagrangian” finite-volume dynamical core for global models, *Mon. Weather
Rev.*, 132, 2293–2307, [https://doi.org/10.1175/1520-0493\(2004\)132<2293:AVLFDC>2.0.CO;2](https://doi.org/10.1175/1520-0493(2004)132<2293:AVLFDC>2.0.CO;2),
825 2004.
- Lin, Y., and Mitchell, K. E.: The NCEP Stage II/IV hourly precipitation analyses: Development and
applications, in: *19th Conf. on Hydrology*, San Diego, CA, Amer. Meteor. Soc., 1.2, available at:
https://ams.confex.com/ams/Annual2005/techprogram/paper_83847.htm (last access: 25 September
2025), 2005.
- 830 Liu, C., Xue, M., and Kong, R.: Direct assimilation of radar reflectivity data using 3DVAR: Treatment
of hydrometeor background errors and OSSE tests, *Mon. Weather Rev.*, 147, 17–29,
<https://doi.org/10.1175/MWR-D-18-0033.1>, 2019.
- Liu, C., Xue, M., and Kong, R.: Direct variational assimilation of radar reflectivity and radial velocity
data: Issues with nonlinear reflectivity operator and solutions, *Mon. Weather Rev.*, 148, 1483–1502,
835 <https://doi.org/10.1175/MWR-D-19-0149.1>, 2020.
- Liu, C., Li, H., Xue, M., Jung, Y., Park, J., Chen, L., Kong, R., and Tong, C.: Use of a reflectivity
operator based on double-moment Thompson microphysics for direct assimilation of radar
reflectivity in GSI-based hybrid En3DVar, *Mon. Weather Rev.*, 150, 907–926,
<https://doi.org/10.1175/MWR-D-21-0040.1>, 2022a.
- 840 Liu, Z., Snyder, C., Guerrette, J. J., Jung, B.-J., Ban, J., Vahl, S., Wu, Y., Trémolet, Y., Auligné, T.,
Ménétrier, B., Shlyayeva, A., Herbener, S., Liu, E., Holdaway, D., and Johnson, B. T.: Data

- assimilation for the Model for Prediction Across Scales – Atmosphere with the Joint Effort for Data
assimilation Integration (JEDI-MPAS 1.0.0): EnVar implementation and evaluation, *Geosci. Model*
Dev., 15, 7859–7878, <https://doi.org/10.5194/gmd-15-7859-2022>, 2022b.
- 845 Ménétrier, B.: Normalized Interpolated Convolution from an Adaptive Subgrid (NICAS)
documentation, available at: https://github.com/benjaminmenetrier/nicas_doc (last access: 25
September 2025), 2020.
- Mittermaier, M., and Roberts, N.: Intercomparison of spatial forecast verification methods: Identifying
skillful spatial scales using the Fractions Skill Score, *Wea. Forecasting*, 25, 343–354,
850 <https://doi.org/10.1175/2009WAF2222260.1>, 2010.
- Niu, G.-Y., et al.: The community Noah land surface model with multiparameterization options (Noah-
MP): 1. Model description and evaluation with local-scale measurements, *J. Geophys. Res. Atmos.*,
116, D12109, <https://doi.org/10.1029/2010JD015139>, 2011.
- NOAA: 2022 Spring Forecasting Experiment, Hazardous Weather Testbed, National Severe Storms
855 Laboratory, <https://doi.org/10.25923/r70a-aw71>, 2022.
- NOAA/NCEP Environmental Modeling Center: V2.7 RTMA/URMA/RTMA-RU NCEP OD Science
Brief, available at: https://www.emc.ncep.noaa.gov/emc/docs/OD_Science_Brief_RTMA_v2.7.pdf
(last access: 25 September 2025), 2018.
- 860 Olson, J. B., Kenyon, J. S., Angevine, W., Brown, J. M., Pagowski, M., and Sušelj, K.: A description of
the MYNN-EDMF scheme and the coupling to other components in WRF–ARW, NOAA Office of
Oceanic and Atmospheric Research, available at:
<https://repository.library.noaa.gov/view/noaa/19837> (last access: 25 September 2025), 2019.
- Park, J., Xue, M., and Liu, C.: Implementation and testing of radar data assimilation capabilities within
the Joint Effort for Data assimilation Integration (JEDI) framework with ensemble transformation
865 Kalman filter coupled with FV3-LAM model, *Geophys. Res. Lett.*, 50, e2022GL102709,
<https://doi.org/10.1029/2022GL102709>, 2023.
- Park, J., and Liu, C.: JEDI–FV3 v1.0.0 with radar data assimilation implementation (v1.0.0), Zenodo
[code], <https://doi.org/10.5281/zenodo.18272913>, 2026.
- Park, J., and Liu, C.: Processed radar observations and configuration files for JEDI–FV3 v1.0.0
870 (v1.0.0), Zenodo [dataset], <https://doi.org/10.5281/zenodo.18272368>, 2026.
- Potvin, C. K., et al.: Systematic comparison of convection-allowing models during the 2017 NOAA
HWT Spring Forecasting Experiment, *Wea. Forecasting*, 34, 1395–1416,
<https://doi.org/10.1175/WAF-D-19-0056.1>, 2019.
- 875 Purser, R. J., Wu, W.-S., Parrish, D. F., and Roberts, N. M.: Numerical aspects of the application of
recursive filters to variational statistical analysis. Part I: Spatially homogeneous and isotropic
Gaussian covariances, *Mon. Weather Rev.*, 131, 1524–1535, <https://doi.org/10.1175/2543.1>, 2003.
- Roberts, N. M., and Lean, H. W.: Scale-selective verification of rainfall accumulations from high-
resolution forecasts of convective events, *Mon. Weather Rev.*, 136, 78–97,
<https://doi.org/10.1175/2007MWR2123.1>, 2008.
- 880 Schenkman, A. D., Xue, M., Shapiro, A., Brewster, K., and Gao, J.: Impact of CASA Radar and
Oklahoma Mesonet data assimilation on the analysis and prediction of tornadic mesovortices in an
MCS, *Mon. Weather Rev.*, 139, 3422–3445, <https://doi.org/10.1175/MWR-D-10-05051.1>, 2011.

- Smith, T. M., et al.: Multi-Radar Multi-Sensor (MRMS) severe weather and aviation products: Initial operating capabilities, *Bull. Am. Meteorol. Soc.*, 97, 1617–1630, <https://doi.org/10.1175/BAMS-D-14-00173.1>, 2016.
- Snook, N., Xue, M., and Jung, J.: Ensemble probabilistic forecasts of a tornadic mesoscale convective system from ensemble Kalman filter analyses using WSR-88D and CASA radar data, *Mon. Weather Rev.*, 140, 2126–2146, <https://doi.org/10.1175/MWR-D-11-00117.1>, 2012.
- Sobash, R. A., Kain, J. S., Bright, D. R., Dean, A. R., Coniglio, M. C., and Weiss, S. J.: Probabilistic forecast guidance for severe thunderstorms based on the identification of extreme phenomena in convection-allowing model forecasts, *Wea. Forecasting*, 26, 714–728, <https://doi.org/10.1175/WAF-D-10-05046.1>, 2011.
- Sun, J., and Crook, N. A.: Dynamical and microphysical retrieval from Doppler radar observations using a cloud model and its adjoint. Part I: Model development and simulated data experiments, *J. Atmos. Sci.*, 54, 1642–1661, [https://doi.org/10.1175/1520-0469\(1997\)054<1642:DAMRFD>2.0.CO;2](https://doi.org/10.1175/1520-0469(1997)054<1642:DAMRFD>2.0.CO;2), 1997.
- Sun, J., Xue, M., Wilson, J. W., Zawadzki, I., Onvlee-Hoomeyer, J., Ballard, S. P., Joe, P., Barker, D., Lee, P.-H., Golding, B., Xu, M., and Pinto, J.: Use of NWP for nowcasting precipitation: Recent progress and challenges, *Bull. Am. Meteorol. Soc.*, 95, 409–426, <https://doi.org/10.1175/BAMS-D-11-00263.1>, 2014.
- Supinie, T. A., Park, J., Snook, N., Hu, X., Brewster, K. A., and Xue, M.: Cool-season evaluation of FV3-LAM-based CONUS-scale forecasts with physics configurations of experimental RRFS ensembles, *Mon. Weather Rev.*, 150, 2379–2398, <https://doi.org/10.1175/MWR-D-21-0331.1>, 2022.
- Thompson, G., Field, P. R., Rasmussen, R. M., and Hall, W. D.: Explicit forecasts of winter precipitation using an improved bulk microphysics scheme. Part II: Implementation of a new snow parameterization, *Mon. Weather Rev.*, 136, 5095–5115, <https://doi.org/10.1175/2008MWR2387.1>, 2008.
- Trémolet, Y., and Auligné, T.: The Joint Effort for Data Assimilation Integration (JEDI), *JCSDA Quarterly Newsletter*, 66, 1–5, <https://doi.org/10.25923/RB19-0Q26>, 2020.
- Tong, C.-C., Jung, Y., Xue, M., and Liu, C.: Direct assimilation of radar data within the National Weather Service operational GSI EnKF and hybrid En3DVar systems for the stand-alone regional FV3 model at a convection-allowing resolution, *Geophys. Res. Lett.*, 47, e2020GL090179, <https://doi.org/10.1029/2020GL090179>, 2020.
- Wang, Y. and Wang, X.: Direct assimilation of radar reflectivity without tangent linear and adjoint of the nonlinear observation operator in the GSI-based EnVar system: Methodology and experiment with the 8 May 2003 Oklahoma City tornadic supercell. *Monthly Weather Review*, 145(4), 1447–1471, <https://doi.org/10.1175/MWR-D-16-0231.1>, 2017.
- Weygandt, S. S., Benjamin, S. G., Hu, M., Alexander, C. R., Smirnova, T. G., and James, E. P.: Radar reflectivity-based model initialization using specified latent heating (Radar-LHI) within a diabatic digital filter or pre-forecast integration. *Weather and Forecasting*, 37(8), 1419–1434, <https://doi.org/10.1175/WAF-D-21-0142.1> 2022.
- Whitaker, J. S., and Hamill, T. M.: Ensemble data assimilation without perturbed observations, *Mon. Weather Rev.*, 130, 1913–1924, [https://doi.org/10.1175/1520-0493\(2002\)130<1913:EDAWPO>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<1913:EDAWPO>2.0.CO;2), 2002.

- 925 Whitaker, J. S., and Hamill, T. M.: Evaluating methods to account for system errors in ensemble data
assimilation, *Mon. Weather Rev.*, 140, 3078–3089, <https://doi.org/10.1175/MWR-D-11-00276.1>,
2012.
- Wu, W.-S., Parrish, D. F., Rogers, E., and Lin, Y.: Regional ensemble–variational data assimilation
using global ensemble forecasts, *Wea. Forecasting*, 32, 83–96, [https://doi.org/10.1175/WAF-D-16-](https://doi.org/10.1175/WAF-D-16-0045.1)
930 [0045.1](https://doi.org/10.1175/WAF-D-16-0045.1), 2017.
- Zhou, X., et al.: The development of the NCEP Global Ensemble Forecast System version 12, *Wea.*
Forecasting, 37, 1069–1084, <https://doi.org/10.1175/WAF-D-21-0112.1>, 2022.