

Divergent carbon use efficiency-growth rate tradeoff in popular biological growth models

Jinyun Tang, William J. Riley, Gianna L. Marschmann and Eoin L. Brodie

5 Department of Molecular Ecology and Biogeochemistry, Climate and Ecosystem Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA.

Correspondence to: Jinyun Tang jinyuntang@lbl.gov

Running title: Divergent CUE.

10

Abstract. Carbon use efficiency (CUE) is an important trait emerging from processes regulating biological growth. CUE can be computed either based on the growth of structural biomass or total biomass divided by substrate uptake rate. Nonequilibrium thermodynamics and observations suggest that, for an exponentially growing population, structural biomass CUE should first increase, then peak, and finally decrease with specific growth rate; meanwhile, total biomass CUE increases asymptotically with specific growth rate. We compared predictions from six popular models that are often used for plant and microbial growth in existing ecosystem models. We found that, for an exponentially growing population of biological cells, (1) the [source-driven Pirt and Compromise models](#) predict that structural biomass CUE increase asymptotically with growth rate; (2) the [apparent sink-driven modified Droop model](#) predicts that structural biomass CUE decreases with growth rate; and (3) the [sink-driven variable internal storage model](#) and two dynamic energy budget models predict that structural biomass CUE first increases, then peaks, and finally decreases with growth rate. Moreover, the modified Droop model predicts that total biomass CUE is constant with growth rate, while all other five models predict that total biomass CUE increases with growth rate asymptotically. For non-exponential biological growth, we show that there is no deterministic relationship between total biomass CUE or structural biomass CUE with respect to either growth rate or temperature. Therefore, we contend that biological growth models should explicitly represent interactions between substrate acquisition, substrate transformation, and maintenance respiration to better capture observed CUE dynamics, and [the sink-driven model should be preferred for general ecosystem biogeochemistry modeling](#).

15

20

25

Deleted: model

Deleted: thereby

1 Introduction

For a growing biological organism, carbon use efficiency (CUE) is [usually](#) defined as the fraction of a unit mass of newly captured carbon that is retained as newly synthesized biomass (for simplicity, our discussion here regards all excreted organic polymers, such as exoenzymes, as part of the total biomass), while the remaining uptake is respired to support this growth (e.g., Manzoni et al., 2018). Because the carbon used to support new biomass synthesis also fulfills the metabolic needs of maintaining existing structural biomass, and, due to environmental fluctuations, the cost of such maintenance is generally dynamic, CUE is inferred to be dynamic even for an organism that is of fixed chemical elemental stoichiometry. Nevertheless, biogeochemical (BGC) models differ widely in their representation of CUE, for example, as a constant parameter (Wieder et al., 2015), a deterministic function of various regulating factors (e.g., temperature, macronutrient concentration; Allison et al., 2010), or an emergent variable that changes according to the chosen way of analysis (Tang and Riley, 2015; Manzoni et al., 2018;

30

35

40 Allison, 2025). Moreover, a few studies using variable internal storage models differentiated between structural and reserve biomass (e.g., Tang and Riley, 2015; Kooijman, 2009; Nev and Van Den Berg, 2017), and reasoned that, since only the structural biomass requires maintenance, CUE can be either defined with respect to the growth of structural biomass or total biomass. However, models that do not differentiate between structural biomass and reserve biomass (e.g., all microbe-implicit soil BGC models; Koven et al., 2013; Parton et al., 1988; Jenkinson, 1990), the Pirt model (Pirt, 1982), and the Compromise model (Beefink et al., 1990; Wang and Post, 2012)) implicitly assume an equal CUE for structural and total biomass growth.

45 In the variable internal storage models, structural biomass refers to DNA, cell wall and membrane material, and any biomass that requires maintenance to support an organism's normal function, while reserve biomass includes lipids, glycogen, circulating metabolites, short-lived rRNA and any biomass that stores energy and acts as a precursor of structural biomass (Kooijman, 2009). Since it is the structural biomass that represents the effect of microbial population on substrate uptake, it is important to recognize how the method of CUE computation affects the interpretation of CUE dynamics. Specifically, is it the structural biomass CUE that focuses on population growth, or is it the total biomass that focuses on biomass growth? Among the existing empirical methods measuring microbial CUE, for the respiration induced by a given amount of substrate addition, the Chloroform fumigation-extraction method and the ¹³C or ¹⁴C-labeling method focus on total biomass increase after substrate addition, thus they derive CUE conditioned on total biomass (Geyer et al., 2019; Hagerty et al., 2022). In contrast, the PLFA (phospholipid fatty acids) approach quantifies structural biomass growth by measuring how much new membrane is synthesized upon the substrate addition (Ibekwe and Kennedy, 1998; Liu et al., 2020). The ¹⁸O-H₂O labeling method and ¹³C-DNA stable isotope probing (SIP) method quantify the amount of microbial growth based on new DNA synthesis (Xu et al., 2024; Sayre et al., 2026). However, DNA synthesis is not equivalent to increase in cell population size. Therefore, only the PLFA method measures the structural biomass CUE, while the ¹⁸O-H₂O method and ¹³C-DNA SIP method measure what is more accurately described as "apparent" structural biomass CUE. Despite the difference in those approaches, the measured CUE values are often treated as equivalent in the literature. For plants, the CUE is often defined as the ratio between net primary productivity and gross primary productivity, signifying the total biomass CUE (Sinsabaugh et al., 2017).

60 In models of soil organic matter decomposition dynamics, CUE defined for the representative microbe in a population is regarded as an essential parameter to determine how much carbon can be retained in soils after respiration. Recently, Tao et al. (2023), adopting the model structure from (Allison et al., 2010), highlighted microbial CUE as the most important parameter to be better constrained to improve the estimation of global soil carbon storage. Overall, microbe-explicit soil BGC models are demonstrating very diverse variations of predicted microbial CUE (some of their prototypes are analyzed later in this manuscript). In general, these microbe-explicit models can be organized into four classes: (1) the Herbert model (Dawes and Ribbons, 1964), (2) the Pirt-model (Pirt, 1982, 1965), (3) the Compromise model (Beefink et al., 1990), and (4) variable internal storage models, including the plant growth model by Thornley (1972), dynamic energy budget models (Tang and Riley, 2015, 2023; Kooijman, 2009), and the Droop model (Thingstad, 1987). Due to the missing of reserve biomass, the first three models equate total biomass CUE to structural biomass CUE, while the fourth group of models can differentiate total biomass CUE from structural biomass CUE.

75 As another essential group of biological organisms in the biosphere, plants are structurally more complex than microbes, and, naturally, the CUE dynamics during plant growth is also more complicated. However, since plants are constrained by the same physical rules as nature enforces onto the microbes, plant and microbial CUEs should be treated similarly both in modeling and measurements. (As it is argued in Tang et al. (2024), a consistent treatment of similar processes in different

Formatted: Normal (Web)

Formatted: Font: (Asian) +Body Asian (DengXian), Superscript, Ligatures: Standard + Contextual

Deleted: the

Deleted: ing

Deleted: of

Deleted: using a linear turnover pool-based soil carbon model, ...

Formatted: Font: (Asian) +Body Asian (DengXian), Ligatures: Standard + Contextual

Deleted: (by the microbe-implicit model they used)

Deleted: Meanwhile

Deleted: more

Formatted: Font: (Asian) +Body Asian (DengXian), Ligatures: Standard + Contextual

Formatted: Font: (Asian) +Body Asian (DengXian), Ligatures: Standard + Contextual

Formatted: Font: (Asian) +Body Asian (DengXian), Ligatures: Standard + Contextual

Formatted: Font: (Asian) +Body Asian (DengXian), Ligatures: Standard + Contextual

Formatted: Font: (Asian) +Body Asian (DengXian), Ligatures: Standard + Contextual

Formatted: Font: (Asian) +Body Asian (DengXian), Ligatures: Standard + Contextual

Deleted: We note that because the variable internal storage models have both structural and reserve biomass, their CUE may be computed in two different ways by either focusing on structural biomass or total biomass. Since it is the structural biomass that represents the effect of microbial population on substrate uptake, it is important to recognize how the way of CUE computation affects the interpretation of CUE dynamics. ...

Deleted: Since

contexts will make the resultant ecosystem models more robust by improving the coupling coherency between the models'

different components.) Theoretically, the whole plant CUE should be defined as the fraction of allocated carbon to various organs that becomes newly synthesized biomass (Thornley, 1972). Because the relative carbon allocation to each plant organ varies with environmental conditions and different organs usually have different CUEs, the whole plant CUE is generally dynamic (Manzoni et al., 2018). Even when all plant organs are assumed to have the same CUE, the dynamics of relative carbon allocation for maintenance and growth will still result in dynamic whole plant CUE. In the literature, the mean whole plant CUE has been mapped for some regions using remote sensing imagery (Bloom et al., 2016; Zhang et al., 2009) and field data (Liu et al., 2022).

Conceptually, the growth of any biological organism is thermodynamically controlled and therefore naturally resembles how a thermal or mechanical engine works with its supplied energy (Roach et al., 2018). For a Carnot thermal engine, classical thermodynamics has shown that its energy use efficiency has an upper bound of $1 - T_L/T_H$, with T_L and T_H being the low and high temperatures of two heat reservoirs, respectively (Feynman et al., 2011). However, this classical thermodynamic result assumes that all processes are reversible. That is, the engine is working so slowly that its power is *de facto* zero. This assumption disagrees with the empirical observation (and desire) that a working engine should be able to deliver a finite amount of work during any given time. Thus, nonequilibrium thermodynamics was developed to analyze the efficiency of various engines (Bejan, 1996; Andresen et al., 1977; Andresen et al., 1984; Onsager, 1931b, a). One important inference from nonequilibrium thermodynamics is that there exists a tradeoff between power and energy efficiency of an engine, so that one sub-optimal efficiency may correspond to both a low and high input power (Roach et al., 2018; Yuan et al., 2022).

The real-life experience of driving can help understand the power-yield tradeoff mentioned above. In this example, the yield is fuel use efficiency (FUE), i.e., distance traveled per volume fuel consumed by an internal combustion vehicle (e.g., km per liter gas). When a vehicle's engine is off, its output power is zero, and the vehicle has zero FUE. When the engine is on but the car is not moving, FUE is still zero (and the fuel use just keeps the engine idling). When the vehicle gradually accelerates on a flat road, FUE gradually increases, and then stabilizes at a value that is more or less as advertised by the vehicle manufacturer. When the driver is overtaking another vehicle or climbing a slope, greater power is demanded. This is done by pressing the pedal (and probably shifting the gear into sport mode at the same time), so that the vehicle may pick up further speed (with the greater power) at the cost of lower FUE. However, in the real-world, because conditions where a car operates usually fluctuate widely, one is more likely to find the car's power-yield tradeoff varies continuously. We note that internal combustion engines generally cannot store the energy produced from burning the fuel, thus this energy is either wasted or used for moving the vehicle. A hybrid vehicle may store some of the kinetic energy generated from fuel burning for later use. As we will see later, biological cells work more likely a hybrid vehicle. Consistent with this analogy, for mean values based on 398 observations of foreign and US domestic automobiles from the model years 1970 to 1982, Boland and Schreyer (2024) found a clear decrease of FUE with increasing horsepower.

Although biological organisms and vehicles' mechanical engines use energy in different forms, they must adhere to the same thermodynamic principles. Therefore, it is reasonable to infer that the CUE of biological growth should also be a dynamic variable, and follow a similar power-yield tradeoff. This pattern has been hypothesized for microbial CUE by Lipson (2015) based on a synthesis of empirical studies from the literature, and was predicted by the dynamic energy budget models (Tang and Riley, 2023, 2025). Nonetheless, we do expect the CUE dynamics of biological growth to be more complex than the energy use efficiency dynamics of thermal engines. Thermal engines, like internal combustion engines, generally do not include energy storage. In contrast, biological growth is often compounded with processes like dormancy, adaptation and acclimation to environmental changes, giving rise to hysteretic CUE dynamics. Particularly for trees in winter, their roots can grow vigorously while the shoots are dormant (Marchand et al., 2025). Still, these complex CUE dynamics can analogously be conceptualized as

Deleted: is often quantified as the ratio of net primary productivity to gross primary productivity (Sinsabaugh et al., 2017), and

Deleted: ¶

[the coupling of multiple thermal engines, such as the coupling of an electric motor and an internal combustion engine in hybrid vehicles, which are more resilient in terms of energy expenditure than either electric vehicles or internal combustion engine vehicles \(under mixed disruption conditions of energy supply; Orecchini et al. \(2018\)\).](#)

45 Considering the significance of CUE in predicting ecosystem biogeochemistry, it is [thus](#) important to answer the following questions: are the formulations commonly used by existing ecosystem biogeochemical models representing CUE dynamics consistently with thermodynamics? If not, what important features could have been missed by those formulations? In the following sections we analyze the tradeoff between growth rate and CUE in six mathematical formulations, some of which underlie most existing ecosystem biogeochemical models. We then highlight the importance of resolving the growth rate-CUE tradeoff dynamically and discuss how it can be properly formulated in future models.
150

2 Theory

We first introduce a modified version of the model used by Odum and Pinkerton (1955) for their analysis of diverse systems that involve the coupling between energy supply and energy use. We describe that model to illustrate the generic features of coupled energy supply and use as implied by nonequilibrium thermodynamics. Then we introduce the biological growth models that are used in many ecosystem models, and analyze the relationship between growth rate and CUE implied by those growth models. All mathematical symbols are described in [Table A1](#).

2.1 The modified Odum-Pinkerton model

The model is formulated using Onsager's reciprocal relations (Onsager, 1931b):

$$J_1 = (l + cf^2)X_1 - cfX_2, \quad (1)$$

$$J_2 = -cfX_1 + (c + \epsilon)X_2, \quad (2)$$

$$T \frac{dS_e}{dt} = J_1X_1 + J_2X_2. \quad (3)$$

60 In the above model, l , f , c and ϵ are constant parameters (whose definitions depend on the application), X_1 and X_2 are forces, J_1 is the [supply](#) flux into the system under the influence of force X_1 , and J_2 is the [output](#) flux, with force X_2 . The sum of products of fluxes and forces contributes to the heat dissipation of the system as measured by the entropy production rate dS_e/dt multiplied with temperature T in equation (3). Note that equation (3) follows directly from the first law of thermodynamics (aka energy conservation), so that given the [supply](#) power $P_1 = J_1X_1$, and the heat dissipation rate $T \frac{dS_e}{dt}$, the useful [output](#) power is $P_2 = -J_2X_2$. Consequently, the system's energy use efficiency (η) is

$$\eta = \frac{P_2}{P_1} = \left(1 + \frac{l}{cf^2} - p\right) \left[1 - \frac{l}{pcf^2} - \frac{\epsilon}{c} \left(\frac{1+cf^2}{pcf^2} - 1\right)\right], \quad (4)$$

65 where $p = P_1/c(fX_1)^2$. Further, we note that

$$p = \frac{P_1}{c(fX_1)^2} = \left(1 + \frac{l}{cf^2}\right) - \frac{X_2}{fX_1} < 1 + \frac{l}{cf^2}, \quad (5)$$

so that η may vary from being negative to a positive maximum value, [depending on the value of terms in \[\]](#). Particularly, when η is plotted against p , we obtain the non-monotonic tradeoff curve ([Figure 1](#)).

Deleted: Table A1


Deleted: output

Deleted: (3)

Deleted: input

Deleted: output

Deleted: Figure 1



175 Figure 1. The power-yield tradeoff predicted by the modified Odum-Pinkerton model. Here, we set $l = 0$ and $\epsilon = 0.03$, so that
 the model is describing the 7th example in Odum and Pinkerton (1955), which represents an organism that captures (via J_1) and
 uses food for its maintenance (signified by ϵ) and growth (via J_2), except that we here assume that the maintenance cost is paid
 by X_2 (analogous to the reserve biomass in the variable internal storage models, e.g., the DEB models in this study). While we do
 80 not plot negative values of η , it may occur resembling to a starving organism that keeps respiring at the expense of biomass
 under insufficient food supply.

Odum and Pinkerton (1955) (hereafter OP1955) used the above model to analyze (1) Atwood’s machine (e.g., Monteiro
 et al., 2015), (2) a water wheel turning a grindstone, (3) one battery charging another battery, (4) a thermocouple running an
 electric motor, (5) a thermal diffusion engine, (6) the metabolism of a pseudo-organism (with no self-repair), (7) food capture by
 an organism for its maintenance and growth, (8) a model of photosynthesis, (9) primary production in a self-sustaining climax
 185 community, and (10) growth and maintenance of a civilization. Among them, the 7th example is closest to the biological growth
 we discuss here. (Moreover, the fact that these equations can describe so many types of systems suggests what can be inferred
 from these equations for the energy use efficiency is a universal feature that is guaranteed by nonequilibrium thermodynamics.
 Consequently, the 7th example and the subject of our analysis—biological growth—should have the same characteristics in terms
 of energy use.) Following the interpretation by OP1955, in this example, J_1 describes the rate of food supply, X_1 corresponds to
 90 the energy drop in metabolism of captured food units (thus X_1 is equivalent to Gibbs free energy, e.g. that of glucose,
 representing the energy quality of the substrate taken from the environment), J_2 represents the rate of effective food use for
 biomass synthesis, X_2 is the energy drop inherent in the metabolism of a unit of food (thus X_2 is analogous to the Gibbs free
 energy of reserve biomass in the DEB models), l designates the basal metabolism spent in self-repair (as paid by X_1 in the
 original OP1955 formulation), c is related to the effectiveness of food concentrating method, and f equals to X_2/X_1 , measuring
 95 the metabolic efficiency of the food capture process. However, considering that biological organisms may keep respiring in the
 absence of food uptake (aka when $J_1 = 0$; e.g., Dawes and Ribbons (1964)), it is more reasonable to pay the basal metabolism
 with X_2 , so that $l = 0$ while $\epsilon > 0$. Nonetheless, this change does not alter the non-monotonic tradeoff between efficiency η and
 the input power P_1 (Figure 1). Moreover, in the OP1955 model, all input power P_1 goes to output power P_2 and waste, thus there
 200 is no change in storage energy density (aka X_2). Therefore, the implied non-monotonic power-yield tradeoff corresponds to
 exponential biological growth, as discussed below.

- Deleted: For
- Deleted: specifically,
- Deleted: assimilation
- Deleted: represents
- Deleted: capture
- Deleted: capture
- Deleted: and
- Deleted: is
- Deleted: equivalent
- Deleted: Figure 1

2.2 Biological growth models

We next analyze six biological growth models: (1) Pirt model (Pirt, 1965), (2) Compromise model (Beefink et al., 1990), (3) Modified Droop (mDroop) model (Thingstad, 1987), (4) Variable Internal Storage (VIS) model modified from Thornley's plant model (Thornley, 1972), (5) the standard dynamic energy budget (sDEB) model (Kooijman, 2009), and (6) the modified dynamic energy budget (mDEB) model (Tang and Riley, 2025). Both the Pirt and Compromise models consider a single biomass pool, and treat growth and maintenance as independent processes, having the same priority in using the substrate. As the growth in the Pirt model and Compromise model is directly driven by substrate uptake, they are in the category of source-driven growth models (Fatichi et al., 2014). The other four models assume biomass consists of structural and reserve biomass. Among them, the VIS, sDEB and mDEB models drive structural biomass growth using reserve biomass, thus they are in the category of sink-driven models (Fatichi et al., 2014). Following Thingstad (1987), the mDroop model has the most parameters, and uses the total biomass to compute the growth rate (through the total-biomass-based substrate quota, even though it does have reserve biomass), thus it is a sink-driven model only apparently. For the Pirt and Compromise models [under general conditions](#), their equations stay the same as in [Table 1](#). The general forms of the other four models differ from those (for exponential growth) in [Table 1](#) and are detailed (from section A to D) in the supplemental material.

We compared the six models' predictions of CUE dynamics for (1) an exponentially growing population under constant carbon inputs using equations in [Table 1](#), (2) 50-year simulations using equations from section E of the supplemental material that mimic microbe-driven soil carbon dynamics under monthly [varying but annually constant](#) carbon inputs ([with a rate of of 350 gC m⁻² yr⁻¹](#); [red line in Figure S1a](#)), and (3) 50-year simulations that further consider the influence of [smoothed daily variable](#) temperatures ([red line in Figure S1c](#)) on related temperature-dependent kinetic parameters (whose parameterizations are detailed in section F of the supplemental material).

Deleted: Table 1

Deleted: Table 1

Deleted: Table 1

Deleted: -to-month

Deleted: generated by random perturbations around the base annual ...

Deleted: input

Deleted: 350

Deleted: , with a standard deviation of 50 gC yr⁻¹ and fixed monthly allocation

Deleted: intra-

Formatted: Superscript

Deleted: y

Deleted: variation

Deleted: S1b

Deleted: ¶

Deleted: ¶

... [1]

Table 1. Models for exponential biological growth $\mu(S)$ as a function of substrate uptake $q(S)$. Listed in parentheses by each model's name are key model parameters, whose definitions are detailed in the nomenclature Table in Appendix A. Notably, Y_V and Y_B are CUEs for structural biomass and total biomass, respectively, both of which are normalized with substrate assimilation efficiency Y_S . More detailed model descriptions are in the supplemental material.

Pirt model: $(\mu_{max}, m_V, Y_S), h(S)$	Pirt (1965, 1982)
$\mu(S) = \mu_{max} h(S)$ $q(S) = \frac{\mu}{Y_S} + \frac{m_V}{Y_S} = \frac{\mu}{Y_S} \left(1 + \frac{m_V}{\mu}\right)$ $\frac{Y_V}{Y_S} = \frac{Y_B}{Y_S} = \frac{\mu(S)}{q(S)Y_S} = \frac{\mu}{\mu + m_V}$	
<p>Description: This source-driven model assumes that specific growth rate $\mu(S)$ is down-regulated linearly from the maximum growth rate μ_{max} by the substrate response function $h(S)$. Substrate uptake $q(S)$ is partitioned between that for biomass synthesis μ/Y_S and maintenance m_V/Y_S. As a single biomass compartment model, its structural biomass yield Y_V and total biomass yield Y_B are equal.</p>	
Compromise (Comp) model: $(\mu_{max}, m_V, Y_S), h(S)$	Wang and Post (2012)
$\mu(S) = \mu_{max} h(S) - m_V [1 - h(S)]$ $q(S) = \frac{\mu}{Y_S} + \frac{m_V}{Y_S}$ $\frac{Y_V}{Y_S} = \frac{Y_B}{Y_S} = \frac{\mu(S)}{q(S)Y_S} = \frac{\mu}{\mu + m_V}$	
<p>Description: This source-driven model assumes that specific growth rate $\mu(S)$ consists of two parts, where the first part linearly scales the maximum growth rate μ_{max} using the substrate response function $h(S)$, and the second part signifies negative growth due to maintenance $m_V [1 - h(S)]$ that decreases linearly with the substrate response function $h(S)$. Substrate uptake $q(S)$ is partitioned between that for biomass synthesis μ/Y_S and maintenance m_V/Y_S. As a single-biomass-compartment model, its structural biomass yield Y_V and total biomass yield Y_B are equal.</p>	
Modified Droop (mDroop) model: $(\mu_{max}, c_1, c_2, Q_{min}, Q_{max}, Y_S), j_A(S)$	Thingstad (1987)
$\mu(S) = \frac{\mu_{max} j_A (Q_{max} - Q_{min})}{j_A Q_{max} + [c_2 + (1 + c_1) \mu_{max}] Q_{min} (Q_{max} - Q_{min})}$ $q(S) = \frac{(1 + c_1) \mu_{max} + c_2 Q_{min}}{\mu_{max} - \mu} \frac{\mu}{Y_S}$ $\frac{Y_V}{Y_S} = \frac{\mu_{max} - \mu}{(1 + c_1) \mu_{max} + c_2 Q_{min}} \frac{1}{Y_S}$ $\frac{Y_B}{Y_S} = \frac{\mu_{max}}{(1 + c_1) \mu_{max} + c_2}$	
<p>Description: This apparent sink-driven model assumes that specific growth rate $\mu(S)$ is linearly down-regulated from the maximum growth rate μ_{max} by the carbon quota function $1 - Q_{min}/Q$, where the carbon quota Q is defined as the ratio between total biomass B_X and structural biomass B_V (see eq.(A3) in section A of the supplemental material). The total biomass B_X increases due to substrate uptake $j_A \frac{Q_{max}-Q}{Q_{max}-Q_{min}} B_V$, and decreases due to respiration $R B_V$ (see eq. (A2) in the supplemental material), where the specific respiration R is a linear function of specific growth rate $\mu(S)$, whose coefficients depend linearly on carbon quota (eq.(A5) in the supplemental material). As a two-biomass-compartment model, its total biomass yield Y_B and structural biomass yield Y_V differ, and they can be analytically represented as a function of specific growth rate μ only for an exponentially growing population.</p>	
Variable-internal-stores (VIS) model: $(v_E, m_V, Y_S, Y_{RV}), j_A(S)$	Williams (1967); Thornley (1972)

Deleted: n

Deleted: up

Deleted: with

Deleted: n

Deleted: up

Deleted: (

$$\mu(S) = \frac{v_E}{2} \left(-1 + \sqrt{1 + \frac{4(j_A Y_S - m_V) Y_{RV}}{v_E}} \right)$$

$$q(S) = \frac{\mu}{Y_S Y_{RV}} \left(1 + \frac{\mu}{v_E} \right) + \frac{m_V}{Y_S}$$

$$\frac{Y_V}{Y_S} = \frac{v_E \mu}{(v_E + \mu) \mu + v_E m_V Y_{RV}} Y_{RV}$$

$$\frac{Y_B}{Y_S} = \frac{(v_E Y_{RV} + \mu) \mu}{(v_E + \mu) \mu + v_E m_V Y_{RV}} Y_S$$

Description: This [sink-driven](#) model assumes that reserve biomass B_X is mobilized to parallelly meet the demand from maintenance and growth of structural biomass B_V (see eq. (B1) in section B of the supplemental material). As a two-biomass-compartment model, its total biomass yield Y_B and structural biomass yield Y_V differ, and they can be analytically represented as a function of specific growth rate μ only for an exponentially growing population.

Standard dynamic energy budget (sDEB) model: $(v_E, m_V, Y_S, Y_{RV}), j_A(S)$ Tang and Riley (2023); Kooijman (2009)

$$\mu(S) = \frac{Y_{RV} Y_S j_A - m_V}{1 + Y_{RV} Y_S j_A / v_E}$$

$$q(S) = \frac{v_E (m_V + \mu)}{Y_S Y_{RV} (v_E - \mu)}$$

$$\frac{Y_V}{Y_S} = \frac{\mu}{\mu + m_V} \left(1 - \frac{\mu}{v_E} \right) Y_{RV}$$

$$\frac{Y_B}{Y_S} = \left(\frac{\mu}{\mu + m_V} \right) \left(\frac{Y_{RV} v_E + m_V + (1 - Y_{RV}) \mu}{v_E} \right)$$

Description: This [sink-driven](#) model assumes that, when the reserve biomass B_X is mobilized to meet the demand from maintenance first and then the growth of structural biomass B_V , the reserve density (i.e. B_X/B_V) follows the first order kinetics (eq. (C1) in section C of the supplemental material). As a two-biomass-compartment model, its total biomass yield Y_B and structural biomass yield Y_V differ, and they can be analytically represented as a function of specific growth rate μ only for an exponentially growing population.

Modified dynamic energy budget (mDEB) model: $(v_E, m_V, Y_S, Y_{RV}), j_A(S)$ Tang and Riley (2025)

$$\mu(S) = \frac{m_V + v_E}{2} \left(-1 + \sqrt{1 + \frac{4v_E(j_A Y_S Y_{RV} - m_V)}{(m_V + v_E)^2}} \right)$$

$$q(S) = \left(1 + \frac{\mu}{v_E} \right) \left(\frac{\mu + m_V}{Y_{RV} Y_S} \right)$$

$$\frac{Y_V}{Y_S} = \left(\frac{\mu}{\mu + m_V} \right) \left(\frac{v_E}{v_E + \mu} \right) Y_{RV}$$

$$\frac{Y_B}{Y_S} = \left(\frac{\mu}{\mu + m_V} \right) \left(\frac{Y_{RV} v_E + \mu + m_V}{v_E + \mu} \right)$$

Description: This [sink-driven](#) model assumes that reserve biomass B_X is mobilized kinetically to meet the demand from maintenance first and then the growth of structural biomass B_V . Unlike the sDEB model, it does not restrict the reserve density (i.e. B_X/B_V) to follow the first order kinetics. As a two-biomass-compartment model, its total biomass yield Y_B and structural biomass yield Y_V differ, and they can be analytically represented as a function of specific growth rate μ only for an exponentially growing population.

Deleted: ¶

... [2]

3 Results of model comparisons

3.1 Predicted CUE-growth rate tradeoffs for exponentially growing populations




Figure 2. Comparison of relationships between CUE and specific growth rate predicted by the six models. Shown in panels (c)-(f) are parameter values to construct the plots. Structural biomass CUE Y_V is defined as the fraction of captured carbon substrate that is retained as structural biomass. Total biomass CUE Y_B is defined as the fraction of captured carbon substrate that is retained in structural and reserve biomass. Note that the specific growth rate for each model is normalized by parameters chosen by convenience.

For an exponentially growing population of biological cells, predictions by these six models can be sorted into three groups: (1) the Pirt and Compromise models predict that (both structural and total biomass) CUE increases asymptotically with increasing specific growth rate; (2) the mDroop model predicts that the total biomass CUE is insensitive to specific growth rate, while the structural biomass CUE decreases with increasing specific growth rate; and (3) the VIS, sDEB, and mDEB models predict that structural biomass CUE first increases then decreases with specific growth rate, while the total biomass CUE increases asymptotically with specific growth rate. Particularly, the increasing total biomass CUE as a function of specific growth rate aligns well with empirical measurements (e.g., Zheng et al., 2019; Hu et al., 2025; Collado et al., 2014), and the Pirt and Compromise models were specifically created to capture this feature observed in batch experiments (Pirt, 1965; Beftink et al., 1990). However, the Compromise model is argued to be more realistic than the Pirt model by allowing negative growth resulting from biomass degradation under low to no substrate conditions (Beftink et al., 1990; Wang and Post, 2012), while the Pirt model forces the growth rate to be zero under the no-supply substrate condition.

3.2 Dynamic CUE in transient soil carbon dynamics under constant temperature

Among the six models, predicted relationships between total biomass CUE and specific growth rate can also be organized in three groups (Figure 3): (a) the Pirt and Compromise models predict essentially identical, almost linearly increasing, relationships (Figure 3a); (b) the mDroop model predicts a curve that traces like the silhouette of a mussel (Figure 3b); and (c)

- Deleted: Figure 3
- Deleted: Figure 3
- Deleted: a
- Deleted: pair of crossed-eyes
- Deleted: Figure 3

the VIS, sDEB, and mDEB models predict relationships look like baguettes that are slanting slightly up and to the right, and those by the two DEB models are nearly overlapping each other (Figure 3c). (The predicted soil carbon dynamics are presented in Figure S2. Even though the temporal variability of fast pool C, slow pool C, and total biomass are different but arguably quite consistent among all six models by our choice to share as many parameters as possible among the models. Note that the use of fast and slow pools here simply means the former carbon substrate is kinetically more favorable than the latter to the microbes, and they do not have the same meaning as in microbe-implicit models, which are directly associated with turnover times (Parton et al., 1988).)





Figure 3. Relationship between total biomass CUE and specific growth rate. The evolution of different microbial carbon pools and total microbial biomass is presented in Figure S2 in the supplemental material. Note that results for the Pirt model and the Compromise model overlap each other in panel (a).

Since the Pirt and Compromise models each have only one biomass pool, their predicted microbes have only one CUE for each specific growth rate (Figure 3a). In contrast, the representation of reserve biomass makes the other four models predict

Deleted: Figure 3

Deleted: ,

Deleted: where their predictions again fall into three groups, e...



Deleted:

Deleted: Figure 3

multiple total biomass CUE values for each specific growth rate (Figure 3b, c), which gradually converge to a closed curve due to repeating growth pattern (Figure S2d) driven by the constant year-to-year carbon input (Figure S2a).





Figure 4. Relationship between structural biomass carbon use efficiency (CUE) and specific growth rate. Across all panels, blue arrows indicate the period when structural biomass CUE decreases with specific growth rate, while red arrows indicate the period when structural biomass CUE increases with specific growth rate. Black arrows in the figure panels indicate the temporal progression in the simulations.

When the structural biomass CUE is analyzed with respect to the specific growth rate (Figure 4), the mDroop model predicts this relationship to have a left-leaning egg shape (Figure 4a). Meanwhile, the VIS, sDEB, and mDEB models all predict the relationship to be more like a kiwifruit (Figure 4b, c, d). In all, just like that for the total biomass CUE, these four models predict that there is not a one-to-one relationship between structural biomass CUE and specific growth rate under constant temperature and monthly varying but annually constant carbon input.

Deleted: Figure 3

Deleted: due to the non-

Deleted: different



Deleted:

Deleted: In

Deleted: Figure 4

Deleted: right-downward slanting beehive

Deleted: Figure 4

Deleted: n

Deleted: apple

Deleted: Figure 4

Deleted: -to-month

3.3 Dynamic CUE in transient state soil carbon dynamics under monthly varying but annually constant carbon input and day-to-day varying temperatures





Figure 5. Variation of total biomass CUE with respect to specific growth rate. Data are extracted from the last 40 years of the 50-year simulations driven by monthly varying but annually constant carbon input and day-to-day varying temperatures. The corresponding soil carbon dynamics are shown in Figure S3 in the supplemental material.

When the temperature effect on kinetic parameters is further considered (on top of the 50-year monthly variable but annually constant carbon input), we find much richer variation of the relationship between total biomass CUE and specific growth rate. Still, the predicted relationships can be organized into the same three groups: (a) those by the Pirt and Compromise models (Figure 5a), which look like a fishhook, and no longer overlap, (b) that by the mDroop model (Figure 5b), which looks like a thread with an open knot, and (c) those by the VIS, sDEB, and mDEB models (Figure 5c), which look like entangled threads. Among these three groups, the relationship predicted by the mDroop model has the largest spread. In particular, the mDroop model predicts significant occurrence of negative CUE, where excessive maintenance cost leads to the loss of biomass in the presence of active carbon substrate uptake. Meanwhile, predictions by the VIS, sDEB, and mDEB models show a much tighter spread, and, due to the chosen parameter combinations, do not involve negative CUE (Figure 5c) as also found for the Pirt and Compromise models (Figure 5a). Nonetheless, by considering temporally varying carbon inputs and temperatures, one specific growth rate can correspond to many values of CUE.

When the relationship between total biomass CUE and temperature is analyzed, the patterns obtained are even more complex (Figure 6). (Note that temperature was varied daily, and repeated over the 50-year simulation period.) The Pirt and Compromise models both show a pattern of larger spread at low and high temperatures, and a general pattern that first increases

Deleted: [3]

Formatted: Font: Bold



- Deleted:
- Deleted: -
- Deleted: to-month
- Deleted: -
- Deleted: to-month
- Deleted: Figure 5
- Deleted: toucan's head
- Deleted: Figure 5
- Deleted: a left-looking dog that has unusually strong rear legs and a fluffy tail
- Deleted: Figure 5
- Deleted: heads of right-looking eagles
- Deleted: Figure 5
- Deleted: Figure 5
- Deleted: Figure 5
- Deleted: Figure 6
- Deleted: (looks like an iron bridge and)

and then decreases with temperature (Figure 6a). The relationships predicted by the VIS, sDEB, and mDEB models have a much tighter pattern and smaller variation (Figure 6c), a phenomenon that is consistent with observations that biological organisms evolutionarily develop reserve dynamics to stabilize performance (thereby less variable CUE) in the presence of fluctuating environmental conditions (Kooijman, 2009). Predictions by the mDroop model show the largest variation, although the CUE-temperature curve is well-defined, and the spread are largest under low temperature, followed by intermediate and high temperatures with a cross over at around 22.5°C (Figure 6b). Overall, all models predict that there are multiple CUE values almost at any given temperature.




Figure 6. Variation of total biomass CUE with respect to temperature for different models. Results were plotted using the last 40 years of the 50-year simulations (shown in Figure S3).

Deleted: Figure 6

Deleted: Figure 6

Deleted: with

Deleted: s


Deleted: aggregated into many distorted stripes

Deleted: high and

Deleted: Figure 6

Deleted: many

Formatted: Superscript



Deleted:

Formatted: Font: 12 pt

Formatted: Normal

4 Discussion

4.1 No deterministic relationship of CUE with respect to specific growth rate, or other environmental factors

425 It has long been recognized that CUE has a strong control on carbon and nutrient cycling through biological organisms and their host systems. However, many existing microbial models treat CUE as either a constant or a function depending deterministically on its controlling factors (Allison et al., 2010; Wang et al., 2021; He et al., 2015; Wieder et al., 2015). Our analysis here suggests that due to the indirect connection between maintenance respiration and substrate uptake, under time-varying conditions (like the varying carbon input and temperature considered here), there is not likely a deterministic relationship between CUE and its controlling factors or specific growth rate. Note, in systems that couple multiple groups of microorganisms (Graham et al., 2007), we expect that CUE dynamics could even be chaotic. Therefore, depending on the sampling time and location (e.g. the line segments indicated by blue or red arrows in [Figure 4](#)), as well as the temporal averaging method involved in analysis (Tang and Riley, 2015), we may find CUE to have various trends with respect to the controlling variable being investigated (He et al., 2024). While it is not investigated here how much the simulated soil organic matter dynamics will change if the CUE is otherwise parameterized with a deterministic function, based on results from Tang and Riley (2015), we expect the change will be significant. For test, when we drive the models with monthly and annually variable carbon input and non-smoothed daily variable temperatures, the spread of CUE curves becomes much wilder (Figure S5). In all, we contend that CUE should be resolved dynamically by explicitly representing the acquired carbon use for processes other than assimilation. This representation has been conceptually adopted in plant growth models, and soil biogeochemical models should also adopt it to ensure the mechanistic coherency of the overall ecosystem models. However, as we argue below, current plant growth models are generally formulated inappropriately.

Deleted: the figures

Deleted: averaging

Deleted: generally

4.2 The source-driven vs sink-driven models of biological growth

445 In the literature, due to their simple mathematical formulation and relatively good performance, the source-driven Pirt and Compromise models have been used much more frequently than the other four models in modeling microbes and plants. For instance, the land component (ELM) of the Energy Exascale Earth Model (Zhu et al., 2019) represents plant growth in a form that resembles the Compromise model, where new structural growth is driven by the residual gross primary productivity flux after subtracting the carbon requirement for maintenance and storage-replenishment. This source-driven approach is also adopted for plant growth by the BiomE model (Weng et al., 2019) and the FATES model (Knox et al., 2024). Sierra et al. (2022) criticized that this source-driven approach may lead to overly fast plant carbon turnover as indicated by a modeled too young radiocarbon signal in plant respiration compared to observations. Indeed, these source-driven models assume that carbon storage is negligible, so that biological growth is dictated by carbon supply. For plants, this approach contrasts with the existence of nonstructural carbon, which leads to observed phenomenon like delayed growth under nutrient limitation (Li et al., 2021; Boussadia et al., 2010), or coarse root growth during the dormant winter (Marchand et al., 2025). In the relative demand configuration of ELM, in order to match observed nighttime root growth (when carbon supply from photosynthesis is zero), its source-driven approach forces the model to introduce a carbon storage pool that may often (unrealistically) become negative at night and be replenished by new photosynthates in the following daytime (Burrows et al., 2020). When these source-driven models are extended to include nitrogen and phosphorus regulation of plant growth, they may be forced to adopt the law of the minimum (Yang et al., 2014), making predictions that contradict the often-observed multiple nutrients co-limited plant growth (Fay et al., 2015). Additionally, the source-driven models may also be forced to adopt the carbon overflow mechanism under nutrient limitation and forbid luxury nutrient uptake under carbon limitation (Jarrell and Beverly, 1981), both artificially accelerating the carbon cycling. When microbes are also modeled with the source-driven approach, such as in the traditional

Deleted: ¶

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Line spacing: 1.5 lines

Moved down [1]: 4.2 The mDEB model seems to be superior for representing general biological growth¶

Deleted: source driven

Deleted: We note that, in order to match observed nighttime root growth, the relative demand configuration in ELM has to introduce a carbon storage pool that may often become (unrealistically) negative at night and be replenished by new photosynthates in the following daytime (Burrows et al., 2020). ...

Deleted: source driven

CENTURY-like models (Koven et al., 2013; Parton et al., 1988), or the more recent MIMICS and Millennial models (Wieder et al., 2014; Abramoff et al., 2018), by induction, they will also predict too-fast soil carbon cycling unless being compensated by inappropriate model parameter values.

It has long been advocated that plant growth is better modeled using the sink-driven approach, where photosynthesis product is first stored as nonstructural carbon, which is then mobilized to drive the growth of plant organs. (Faticchi et al., 2014; Fourcaud et al., 2008). Our analysis here suggests that the Pirt and Compromise models, although treating growth as driven directly by substrate uptake, are able to produce very rich variation in microbial CUE when analyzed with respect to either specific growth rates or temperatures (Figure 5a and Figure 6a). Interestingly but not surprisingly, their predicted CUE patterns vary more widely than those predicted by the VIS, sDEB, and mDEB models (Figure 5c and Figure 6c), where structural biomass growth is sink-driven as fueled by reserve biomass, which is evolutionarily developed by biological organisms to stabilize their metabolic performance in the face of environmental fluctuations (Kooijman, 2009). From the greater CUE variation by the Pirt and Compromise models, we deduce that, due to the missing of buffering effect from the reserve biomass, the source-driven models may very likely overestimate the plant and microbial response to environmental change, such as elevated CO₂, nutrient limitation or warming. (Some support is shown in Figures S4 and S5, which show that by increasing the variability of carbon input and temperature, the source-driven models respond stronger.) Among the four sink-driven models, the VIS model considers maintenance respiration and growth as two-parallel processes of equal priority, while the sDEB and mDEB models regard maintenance respiration to have priority over growth. Thus, at the cellular level, the VIS model seems to be mechanistically less reasonable by triggering earlier death through maintenance and growth competition. The mDroop model represents biological growth as sink-driven only apparently (because it computes the carbon quota based on total biomass, rather than reserve biomass, to drive the growth), and predicted transient CUE dynamics that are quite different from the three truly sink-driven models (Figure 5 and Figure 6, and also Figures S4 and S5). We believe that predictions by the mDroop model are likely to be more easily falsified empirically.

4.3 The mDEB model seems to be superior for representing general biological growth.

Our analysis indicates that only the sink-driven models (i.e. the VIS model and two DEB models) are able to capture the feature that structural biomass CUE first increases and decreases with specific growth rate as inferred from the synthesis of empirical observations (Lipson, 2015). Nonetheless, when exponential growth is considered, the VIS, sDEB, and mDEB models are found to be more consistent with the Odum-Pinkerton model (Odum and Pinkerton, 1955) that is formulated using non-equilibrium thermodynamics. Our numerical analysis here shows that the VIS, sDEB and mDEB models are equally good (as evidenced by their very similar performance for all cases analyzed here), because maintenance respiration is usually a small fraction of total respiration. As the VIS model was initially proposed to model plant growth (Thornley, 1972), we hypothesize that the sDEB and mDEB models would be comparably good for modeling plant growth. However, if we consider previous comparisons between the sDEB and mDEB models for microbes (Tang and Riley, 2025), then the mDEB model has better mechanistic coherency with the first law of thermodynamics, and is numerically much easier to incorporate multiple nutrients co-limited growth. Therefore, mDEB model seems to be a better formulation for representing biological growth under general conditions. In particular, the mDEB model will enable the growth of plant shoots and roots to be modeled separately, while the two are modulated with shoot-plant exchange of water, carbon, and nutrients. Although the resultant model may appear to be more sophisticated, counter-intuitively (as we observed in EcoSIM), it will be better constrained by using more directly measurable parameters (e.g. nutrient kinetic parameters, root morphological size and hydraulic resistance) and state and flux variables (e.g. phloem transport rate of carbon and nitrogen, root biomass and morphology profile). For instance, the coarse root

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto

Deleted: Meanwhile, i

Deleted: sink driven

Deleted: carbon storage

Deleted: s

Deleted: and photosynthesis effects growth indirectly through replenishing the carbon storage

Deleted: Figure 5

Deleted: Figure 6

Deleted: Figure 5

Deleted: Figure 6

Deleted: sink driven

Formatted: Subscript

Deleted: Note that

Deleted: However, if one chooses t

Deleted: , which

Deleted: sink driven

Deleted: then

Deleted: the

Deleted: can

Deleted: be

Deleted: Figure 5

Deleted: Figure 6

Deleted: ,

Deleted: which

Deleted: 2

Deleted: ¶

Formatted: Indent: First line: 0", Line spacing: Double

Moved up [2]: However, if one chooses the mDroop model, which represents biological growth as sink driven only apparently (because it computes the carbon quota based on total biomass, rather than reserve biomass, to drive the growth), then the predicted transient CUE dynamics can be quite different (Figure 5 and Figure 6), which are likely to be more easily falsified empirically.

Deleted: Additionally

Deleted: with its

Deleted: higher efficiency in numerical solution

Deleted: ,

Deleted: the

dynamics can then be better represented, while the source-driven models forbid coarse root growth in the absence of photosynthesis, a structural deficiency unlikely to be compensated by parameter tuning.

5 Conclusions

By starting from non-equilibrium thermodynamics, we infer that under no change of internal energy storage density (as usually is the case for internal combustion engines), the power use efficiency of a thermal engine as defined by the ratio of useful power output and power input will first increase, then plateau and finally decrease with increasing input power. Following this, we deduced that (which was also inferred by (Lipson, 2015) based on a synthesis of empirical observations), for an exponentially growing population of cells, the relationship between structural biomass carbon use efficiency and specific growth rate should follow the same pattern. In our analysis of six biological growth models adopted in currently popular ecosystem biogeochemical models, we found that only the VIS, sDEB, and mDEB models correctly capture this relationship. Considering transient carbon inputs and temperatures, all six models suggest that there is not a deterministic relationship of carbon use efficiency with respect to specific growth rate, substrate uptake rate, carbon inputs, or temperature. Therefore, combining with our previous analysis (Tang and Riley, 2015), we contend that, in order to robustly simulate the emergent carbon use efficiency dynamics and its consequent influence on ecosystem biogeochemistry, models should represent biological growth as sink-driven, and the effect of controlling factors like temperature and moisture should not be applied as multiplier functions, rather they should be applied to directly modify the kinetic parameters of different metabolic processes. Our analysis shows that the mDEB model is a good candidate framework to meet all these needs, and will potentially help ecosystem biogeochemistry models to improve their simulated variability of ecosystem dynamics in response to changing driving conditions.

Appendix A: Nomenclature

Table A1: Definition of symbols used in the modified Odum-Pinkerton model and models in [Table 1](#).

Symbol	Unit	Description
c	Application dependent	Parameter for the modified Odum-Pinkerton model
c_1	None	Scaling parameter for growth respiration in mDroop model
c_2	year ⁻¹	Scaling parameter for maintenance respiration in mDroop model
f	Application dependent	Parameter for the modified Odum-Pinkerton model
$h(S)$	None	Substrate response function in Pirt model and Compromise model
$j_A(S)$	year ⁻¹	Specific substrate uptake in the mDroop model, VIS model and DEB models.
l	Application dependent	Parameter for the modified Odum-Pinkerton model
m_V	year ⁻¹	Specific maintenance respiration rate for Pirt model, Compromise model, VIS model, and DEB models
p	None	Normalized input power in the modified Odum-Pinkerton model

Deleted: Table 1

$q(\mathcal{S})$	year ⁻¹	Specific substrate uptake rate in all growth models
v_E	year ⁻¹	Specific reserve mobilization rate in the VIS model and DEB models
J_1	Application dependent	Input flux in the modified Odum-Pinkerton model
J_2	Application dependent	Input flux in the modified Odum-Pinkerton model
P_1	Application dependent	Input power in the modified Odum-Pinkerton model
P_2	Application dependent	Useful output power in the modified Odum-Pinkerton model
Q_{min}	None	Minimum carbon quota for growth in the mDroop model
Q_{max}	None	Maximum carbon quota for growth in the mDroop model
S	gC m ⁻³	Substrate concentration in the growth models
S_e	J K ⁻¹	Entropy
T	K	Temperature
X_1, X_2	Application dependent	Forces in the modified Odum-Pinkerton model
Y_{RV}	None	Conversion efficiency from reserve biomass to structural biomass in VIS model and DEB models
Y_S	None	Substrate assimilation efficiency in all models
Y_V	None	Structural biomass carbon use efficiency in all models
Y_B	None	Total biomass carbon use efficiency in all models
ϵ	Application dependent	Parameter for the modified Odum-Pinkerton model
η	None	Power efficiency in the modified Odum-Pinkerton model
μ_{max}	year ⁻¹	Maximum specific growth rate in the Pirt model, Compromise model and mDroop model
$\mu(\mathcal{S})$	year ⁻¹	Specific growth rate in all growth models

Deleted: ²

Formatted: Superscript

Formatted: Superscript

580 Author contributions.

JYT formulated the idea, conduct the analysis and wrote the paper. WJR, GM, ELB discussed the results and edited the manuscript.

Competing interests.

The contact author has declared that neither of the authors has any competing interests

585 Code/data availability.

The code and data used in this study are available at: https://github.com/jinyun1tang/cue_paper.

Disclaimer.

Financial support does not constitute an endorsement by the Department of Energy and National Science Foundation of the
590 views expressed in this study.

Financial support.

The research was supported by the director of the Office of Science, Office of Biological and Environmental Research, of the US
Department of Energy (contract no. DEAC02-05CH11231) and the National Science Foundation (grant no. 2125069).

Reference

- Abramoff, R., Xu, X. F., Hartman, M., O'Brien, S., Feng, W. T., Davidson, E., Finzi, A. C., Moorhead, D., Schimel, J., Torn, M., and Mayes, M. A.: The Millennial model: in search of measurable pools and transformations for modeling soil carbon in the new century, *Biogeochemistry*, 137, 51-71, 10.1007/s10533-017-0409-7, 2018.
- 600 Allison, S. D.: Rethinking microbial carbon use efficiency in soil models, *Nature Climate Change*, 15, 10-12, 10.1038/s41558-024-02217-6, 2025.
- Allison, S. D., Wallenstein, M. D., and Bradford, M. A.: Soil-carbon response to warming dependent on microbial physiology, *Nat Geosci*, 3, 336-340, 10.1038/Ngeo846, 2010.
- 605 Andresen, B., Salamon, P., and Berry, R. S.: Thermodynamics in finite-time, *Phys Today*, 37, 62-70, 10.1063/1.2916405, 1984.
- Andresen, B., Berry, R. S., Nitzan, A., and Salamon, P.: Thermodynamics in finite time .1. Step-Carnot cycle, *Phys Rev A*, 15, 2086-2093, 10.1103/PhysRevA.15.2086, 1977.
- Beeftink, H. H., Vanderheijden, R. T. J. M., and Heijnen, J. J.: Maintenance requirements - energy supply from simultaneous endogenous respiration and substrate consumption, *Fems Microbiol Ecol*, 73, 203-209, 10.1016/0378-1097(90)90731-5, 1990.
- 610 Bejan, A.: Entropy generation minimization: The new thermodynamics of finite-size devices and finite-time processes, *J Appl Phys*, 79, 1191-1218, 10.1063/1.362674, 1996.
- Bloom, A. A., Exbrayat, J. F., van der Velde, I. R., Feng, L., and Williams, M.: The decadal state of the terrestrial carbon cycle: Global retrievals of terrestrial carbon allocation, pools, and residence times, *P Natl Acad Sci USA*, 113, 1285-1290, 10.1073/pnas.1515160113, 2016.
- 615 Boland, B. and Schreyer, S.: The Relationship Between Horsepower & Fuel Efficiency, 2024.
- Boussadia, O., Steppe, K., Zgallai, H., El Hadj, S. B., Braham, M., Lemeur, R., and Van Labeke, M. C.: Effects of nitrogen deficiency on leaf photosynthesis, carbohydrate status and biomass production in two olive cultivars 'Meski' and 'Koroneiki', *Sci Hortic-Amsterdam*, 123, 336-342, 10.1016/j.scienta.2009.09.023, 2010.
- 620 Burrows, S. M., Maltrud, M., Yang, X., Zhu, Q., Jeffery, N., Shi, X., Ricciuto, D., Wang, S., Bisht, G., Tang, J., Wolfe, J., Harrop, B. E., Singh, B., Brent, L., Baldwin, S., Zhou, T., Cameron-Smith, P., Keen, N., Collier, N., Xu, M., Hunke, E. C., Elliott, S. M., Turner, A. K., Li, H., Wang, H., Golaz, J. C., Bond-Lamberty, B., Hoffman, F. M., Riley, W. J., Thornton, P. E., Calvin, K., and Leung, L. R.: The DOE E3SM v1.1 Biogeochemistry configuration: description and simulated ecosystem-climate responses to historical changes in forcing, *J Adv Model Earth Sy*, 12, ARTN e2019MS001766 10.1029/2019MS001766, 2020.
- 625 Collado, S., Rosas, I., González, E., Gutierrez-Lavin, A., and Diaz, M.: response in membrane bioreactors under salicylic acid-induced stress conditions, *J Hazard Mater*, 267, 9-16, 10.1016/j.jhazmat.2013.12.034, 2014.
- 630 Dawes, E. A. and Ribbons, D. W.: Some aspects of endogenous metabolism of bacteria, *Bacteriol Rev*, 28, 126-149, 10.1128/Mmbr.28.2.126-149.1964, 1964.
- Fatichi, S., Leuzinger, S., and Körner, C.: Moving beyond photosynthesis: from carbon source to sink-driven vegetation modeling, *New Phytol*, 201, 1086-1095, 10.1111/nph.12614, 2014.

635 Fay, P. A., Prober, S. M., Harpole, W. S., Knops, J. M. H., Bakker, J. D., Borer, E. T., Lind, E. M.,
MacDougall, A. S., Seabloom, E. W., Wragg, P. D., Adler, P. B., Blumenthal, D. M., Buckley, Y., Chu, C. J.,
Cleland, E. E., Collins, S. L., Davies, K. F., Du, G. Z., Feng, X. H., Firn, J., Gruner, D. S., Hagenah, N.,
Hautier, Y., Heckman, R. W., Jin, V. L., Kirkman, K. P., Klein, J., Ladwig, L. M., Li, Q., McCulley, R. L.,
Melbourne, B. A., Mitchell, C. E., Moore, J. L., Morgan, J. W., Risch, A. C., Schütz, M., Stevens, C. J.,
Wedin, D. A., and Yang, L. H.: Grassland productivity limited by multiple nutrients, *Nat Plants*, 1, ArtN
640 15080
10.1038/Nplants.2015.80, 2015.
Feynman, R. P., Leighton, R. B., and Sands, M.: *The Feynman Lectures on Physics*, The New Millennium
Edition, Basic Books 2011.
645 Fourcaud, T., Zhang, X., Stokes, A., Lambers, H., and Körner, C.: Plant growth modelling and
applications: The increasing importance of plant architecture in growth models, *Ann Bot-London*, 101,
1053-1063, 10.1093/aob/mcn050, 2008.
Geyer, K. M., Dijkstra, P., Sinsabaugh, R., and Frey, S. D.: Clarifying the interpretation of carbon use
efficiency in soil through methods comparison, *Soil Biol Biochem*, 128, 79-88,
10.1016/j.soilbio.2018.09.036, 2019.
650 Graham, D. W., Knapp, C. W., Van Vleck, E. S., Bloor, K., Lane, T. B., and Graham, C. E.: Experimental
demonstration of chaotic instability in biological nitrification, *Isme J*, 1, 385-393, 10.1038/ismej.2007.45,
2007.
Hagerty, S. B., Allison, S. D., and Schimel, J. P.: Testing microbial models with data from a
C glucose tracer experiment, *Soil Biol Biochem*, 172, ARTN 108781
655 10.1016/j.soilbio.2022.108781, 2022.
He, X. J., Abs, E., Allison, S. D., Tao, F., Huang, Y. Y., Manzoni, S., Abramoff, R., Bruni, E., Bowring, S. P.
K., Chakrawal, A., Ciaia, P., Elsgaard, L., Friedlingstein, P., Georgiou, K., Hugelius, G., Holm, L. B., Li,
W., Luo, Y. Q., Marmasse, G., Nunan, N., Qiu, C. J., Sitch, S., Wang, Y. P., and Goll, D. S.: Emerging
660 multiscale insights on microbial carbon use efficiency in the land carbon cycle, *Nat Commun*, 15, ARTN
8010 10.1038/s41467-024-52160-5, 2024.
He, Y. J., Yang, J. Y., Zhuang, Q. L., Harden, J. W., McGuire, A. D., Liu, Y. L., Wang, G. S., and Gu, L. H.:
Incorporating microbial dormancy dynamics into soil decomposition models to improve quantification of
soil carbon dynamics of northern temperate forests, *J Geophys Res-Biogeophys*, 120, 2596-2611,
10.1002/2015jg003130, 2015.
665 Hu, J. X., Cui, Y. X., Manzoni, S., Zhou, S. X., Cornelissen, J. H. C., Huang, C. D., Schimel, J., and
Kozlov, Y.: Microbial carbon use efficiency and growth rates in soil: global patterns and drivers, *Global
Change Biol*, 31, ARTN e70036 10.1111/gcb.70036, 2025.
Ibekwe, A. M. and Kennedy, A. C.: Phospholipid fatty acid profiles and carbon utilization patterns for
analysis of microbial community structure under field and greenhouse conditions, *Fems Microbiol Ecol*,
670 26, 151-163, DOI 10.1111/j.1574-6941.1998.tb00501.x, 1998.
Jarrell, W. M. and Beverly, R. B.: *The Dilution Effect in Plant Nutrition Studies*, *Adv Agron*, 34, 197-224,
Doi 10.1016/S0065-2113(08)60887-1, 1981.
Jenkinson, D. S.: The turnover of organic-carbon and nitrogen in soil, *Philos T R Soc B*, 329, 361-368,
10.1098/rstb.1990.0177, 1990.
675 Knox, R. G., Koven, C. D., Riley, W. J., Walker, A. P., Wright, S. J., Holm, J. A., Wei, X. Y., Fisher, R. A.,
Zhu, Q., Tang, J. Y., Ricciuto, D. M., Shuman, J. K., Yang, X. J., Kueppers, L. M., and Chambers, J. Q.:
Nutrient dynamics in a coupled terrestrial biosphere and land model (ELM-FATES-CNP), *J Adv Model
Earth Sy*, 16, ARTN e2023MS003689 10.1029/2023MS003689, 2024.
Kooijman, S. A. L. M.: *Dynamic Energy Budget Theory for Metabolic Organisation*, Cambridge University
680 Press, Cambridge, 10.1017/CBO9780511805400, 2009.
Koven, C. D., Riley, W. J., Subin, Z. M., Tang, J. Y., Torn, M. S., Collins, W. D., Bonan, G. B., Lawrence,
D. M., and Swenson, S. C.: The effect of vertically resolved soil biogeochemistry and alternate soil C and
N models on C dynamics of CLM4, *Biogeosciences*, 10, 7109-7131, 10.5194/bg-10-7109-2013, 2013.

685 Li, J. M., Du, A. P., Liu, P. H., Tian, X. P., Jin, Y. L., Yi, Z. L., He, K. Z., Fang, Y., and Zhao, H.: High starch accumulation mechanism and phosphorus utilization efficiency of duckweed () under phosphate starvation, *Ind Crop Prod*, 167, ARTN 113529 10.1016/j.indcrop.2021.113529, 2021.

Lipson, D. A.: The complex relationship between microbial growth rate and yield and its implications for ecosystem processes, *Front Microbiol*, 6, ARTN 615 10.3389/fmicb.2015.00615, 2015.

690 Liu, Z. G., Chen, Z., Yu, G. R., Yang, M., Zhang, W. K., Zhang, T. Y., and Han, L.: Ecosystem carbon use efficiency in ecologically vulnerable areas in China: Variation and influencing factors, *Front Plant Sci*, 13, ARTN 1062055 10.3389/fpls.2022.1062055, 2022.

Liu, Z. W., Wu, X. L., Liu, W., Bian, R. J., Ge, T. D., Zhang, W., Zheng, J. F., Drosos, M., Liu, X. Y., Zhang, X. H., Cheng, K., Li, L. Q., and Pan, G. X.: Greater microbial carbon use efficiency and carbon sequestration in soils: Amendment of biochar versus crop straws, *Gcb Bioenergy*, 12, 1092-1103, 10.1111/gcbb.12763, 2020.

Manzoni, S., Capek, P., Porada, P., Thurner, M., Winterdahl, M., Beer, C., Brüchert, V., Frouz, J., Herrmann, A. M., Lindahl, B. D., Lyon, S. W., Santrucková, H., Vico, G., and Way, D.: Reviews and syntheses: Carbon use efficiency from organisms to ecosystems - definitions, theories, and empirical evidence, *Biogeosciences*, 15, 5929-5949, 10.5194/bg-15-5929-2018, 2018.

700 Marchand, L. J., Gricar, J., Zuccarini, P., Dox, I., Marien, B., Verlinden, M., Heinecke, T., Prislán, P., Marie, G., Lange, H., van den Bulcke, J., Penuelas, J., Fonti, P., and Campioli, M.: No winter halt in below-ground wood growth of four angiosperm deciduous tree species, *Nat Ecol Evol*, 9, 10.1038/s41559-024-02602-6, 2025.

705 Monteiro, M., Stari, C., Cabeza, C., and Marti, A. C.: The Atwood machine revisited using smartphones, *Phys Teach*, 53, 373-374, 10.1119/1.4928357, 2015.

Nev, O. A. and van den Berg, H. A.: Variable-Internal-Stores models of microbial growth and metabolism with dynamic allocation of cellular resources, *J Math Biol*, 74, 409-445, 10.1007/s00285-016-1030-4, 2017.

710 Odum, H. T. and Pinkerton, R. C.: Times speed regulator - the optimum efficiency for maximum power output in physical and biological systems, *Am Sci*, 43, 331-343, 1955.

Onsager, L.: Reciprocal relations in irreversible processes. II, *Phys Rev*, 38, 2265-2279, 10.1103/PhysRev.38.2265, 1931a.

715 Onsager, L.: Reciprocal relations in irreversible processes. I., *Phys Rev*, 37, 405-426, 10.1103/PhysRev.37.405, 1931b.

Orecchini, F., Santiangeli, A., Zuccari, F., Ortenzi, F., Genovese, A., Spazzafumo, G., and Nardone, L.: Energy consumption of a last generation full hybrid vehicle compared with a conventional vehicle in real drive conditions, *Energy Proced*, 148, 289-296, 10.1016/j.egypro.2018.08.080, 2018.

720 Parton, W. J., Stewart, J. W. B., and Cole, C. V.: Dynamics of C, N, P and S in grassland soils - a model, *Biogeochemistry*, 5, 109-131, 10.1007/Bf02180320, 1988.

Pirt, S. J.: Maintenance energy of bacteria in growing cultures, *Proc R Soc Ser B-Bio*, 163, 224-231, 10.1098/rspb.1965.0069, 1965.

Pirt, S. J.: Maintenance energy - a general-model for energy-limited and energy-sufficient growth, *Arch Microbiol*, 133, 300-302, 10.1007/Bf00521294, 1982.

725 Roach, T. N. F., Salamon, P., Nulton, J., Andresen, B., Felts, B., Haas, A., Calhoun, S., Robinett, N., and Rohwer, F.: Application of finite-time and control thermodynamics to biological processes at multiple scales, *J Non-Equil Thermody*, 43, 193-210, 10.1515/jnet-2018-0008, 2018.

Sayre, J. M., Wang, D. Y., Erikson, C., Gontijo, J. B., Scow, K., and Rodrigues, J. L. M.: Increased microbial carbon use efficiency and metabolic capacity in manure amended soils: A 665-day field experiment, *Biol Fert Soils*, 10.1007/s00374-026-01986-w, 2026.

730 Sierra, C. A., Ceballos-Núñez, V., Hartmann, H., Herrera-Ramírez, D., and Metzler, H.: Ideas and perspectives: Allocation of carbon from net primary production in models is inconsistent with observations of the age of respired carbon, *Biogeosciences*, 19, 3727-3738, 10.5194/bg-19-3727-2022, 2022.

735 Sinsabaugh, R. L., Moorhead, D. L., Xu, X. F., and Litvak, M. E.: Plant, microbial and ecosystem carbon use efficiencies interact to stabilize microbial growth as a fraction of gross primary production, *New Phytol*, 214, 1518-1526, 10.1111/nph.14485, 2017.

Tang, J. Y. and Riley, W. J.: Weaker soil carbon-climate feedbacks resulting from microbial and abiotic interactions, *Nature Climate Change*, 5, 56-60, 10.1038/Nclimate2438, 2015.

740 Tang, J. Y. and Riley, W. J.: Revising the dynamic energy budget theory with a new reserve mobilization rule and three example applications to bacterial growth, *Soil Biol Biochem*, 178, ARTN 108954 10.1016/j.soilbio.2023.108954, 2023.

Tang, J. Y. and Riley, W. J.: Technical note: A modified formulation of dynamic energy budget theory for faster computation of biological growth, *Biogeosciences*, 22, 1809-1819, 10.5194/bg-22-1809-2025, 2025.

745 Tang, J. Y., Riley, W. J., Manzoni, S., and Maggi, F.: Feasibility of Formulating Ecosystem Biogeochemical Models From Established Physical Rules, *J Geophys Res-Biogeophys*, 129, ARTN e2023JG007674 10.1029/2023JG007674, 2024.

Tao, F., Huang, Y. Y., Hungate, B. A., Manzoni, S., Frey, S. D., Schmidt, M. W. I., Reichstein, M., Carvalhais, N., Ciais, P., Jiang, L. F., Lehmann, J., Wang, Y. P., Houlton, B. Z., Ahrens, B., Mishra, U., Hugelius, G., Hocking, T. D., Lu, X. J., Shi, Z., Viatkin, K., Vargas, R., Yigini, Y., Omuto, C., Malik, A. A., 750 Peralta, G., Cuevas-Corona, R., Di Paolo, L. E., Luotto, I., Liao, C. J., Liang, Y. S., Saynes, V. S., Huang, X. M., and Luo, Y. Q.: Microbial carbon use efficiency promotes global soil carbon storage, *Nature*, 618, 981-985, 10.1038/s41586-023-06042-3, 2023.

Thingstad, T. F.: Utilization of N, P, and organic C by heterotrophic bacteria .1. Outline of a chemostat theory with a consistent concept of maintenance metabolism, *Mar Ecol Prog Ser*, 35, 99-109, 10.3354/meps035099, 1987.

755 Thornley, J. H. M.: A balanced quantitative model for root: shoot ratios in vegetative plants, *Ann. Bot.*, 36, 431-441, 10.1093/oxfordjournals.aob.a084602, 1972.

Wang, G. S. and Post, W. M.: A theoretical reassessment of microbial maintenance and implications for microbial ecology modeling, *Fems Microbiol Ecol*, 81, 610-617, 10.1111/j.1574-6941.2012.01389.x, 2012.

760 Wang, Y. P., Zhang, H. C., Ciais, P., Goll, D., Huang, Y. Y., Wood, J. D., Ollinger, S. V., Tang, X. L., and Prescher, A. K.: Microbial activity and root carbon inputs are more important than soil carbon diffusion in simulating soil carbon profiles, *J Geophys Res-Biogeophys*, 126, ARTN e2020JG006205 10.1029/2020JG006205, 2021.

765 Weng, E. S., Dybzinski, R., Farris, C. E., and Pacala, S. W.: Competition alters predicted forest carbon cycle responses to nitrogen availability and elevated CO₂: simulations using an explicitly competitive, game-theoretic vegetation demographic model, *Biogeosciences*, 16, 4577-4599, 10.5194/bg-16-4577-2019, 2019.

Wieder, W. R., Grandy, A. S., Kallenbach, C. M., and Bonan, G. B.: Integrating microbial physiology and physio-chemical principles in soils with the Microbial-MIneral Carbon Stabilization (MIMICS) model, *Biogeosciences*, 11, 3899-3917, 10.5194/bg-11-3899-2014, 2014.

770 Wieder, W. R., Grandy, A. S., Kallenbach, C. M., Taylor, P. G., and Bonan, G. B.: Representing life in the Earth system with soil microbial functional traits in the MIMICS model, *Geosci Model Dev*, 8, 1789-1808, 10.5194/gmd-8-1789-2015, 2015.

Williams, F. M.: A model of cell growth dynamics, *J Theor Biol*, 15, 190-207, Doi 10.1016/0022-5193(67)90200-7, 1967.

775 Xu, Q. C., Li, L., Guo, J. J., Guo, H. Y., Liu, M. Q., Guo, S. W., Kuzyakov, Y., Ling, N., and Shen, Q. R.: Active microbial population dynamics and life strategies drive the enhanced carbon use efficiency in high-organic matter soils, *Mbio*, 15, 10.1128/mbio.00177-24, 2024.

780 Yang, X., Thornton, P. E., Ricciuto, D. M., and Post, W. M.: The role of phosphorus dynamics in tropical forests - a modeling study using CLM-CNP, *Biogeosciences*, 11, 1667-1681, 10.5194/bg-11-1667-2014, 2014.

Yuan, H., Ma, Y. H., and Sun, C. P.: Optimizing thermodynamic cycles with two finite-sized reservoirs, *Phys Rev E*, 105, ARTN L022101 10.1103/PhysRevE.105.L022101, 2022.

- 785 Zhang, Y. J., Xu, M., Chen, H., and Adams, J.: Global pattern of NPP to GPP ratio derived from MODIS data: effects of ecosystem type, geographical location and climate, *Global Ecol Biogeogr*, 18, 280-290, 10.1111/j.1466-8238.2008.00442.x, 2009.
- Zheng, Q., Hu, Y. T., Zhang, S. S., Noll, L., Böckle, T., Richter, A., and Wanek, W.: Growth explains microbial carbon use efficiency across soils differing in land use and geology, *Soil Biol Biochem*, 128, 45-55, 10.1016/j.soilbio.2018.10.006, 2019.
- 790 Zhu, Q., Riley, W. J., Tang, J. Y., Collier, N., Hoffman, F. M., Yang, X. J., and Bisht, G.: Representing nitrogen, phosphorus, and carbon interactions in the E3SM land model: development and global benchmarking, *J Adv Model Earth Sy*, 11, 2238-2258, 10.1029/2018ms001571, 2019.

Page 6: [1] Deleted	Jinyun Tang	2/19/26 6:32:00 PM
Page 8: [2] Deleted	Jinyun Tang	2/19/26 6:32:00 PM
Page 12: [3] Deleted	Jinyun Tang	2/21/26 8:38:00 PM