

Degradation of anhydro-saccharides and the driving factors in real atmospheric conditions: A cross-city study in China

Biao Zhou ^a, Kun Zhang ^a, Qiongqiong Wang ^b, Jiqi Zhu ^a, Li Li ^{a,*} and Jian Zhen Yu ^{c,d,*}

^a School of Environmental and Chemical Engineering, Shanghai University, Shanghai, 200444, China

^b Department of Atmospheric Science, School of Environmental Studies, China University of Geosciences, Wuhan, China

^c Department of Chemistry, The Hong Kong University of Science & Technology, Hong Kong, China

^d Division of Environment & Sustainability, The Hong Kong University of Science & Technology, Hong Kong, China

Correspondence to: Li Li (Lily@shu.edu.cn) and Jian Zhen Yu (jian.yu@ust.hk)

Abstract

Anhydro-saccharides (levoglucosan, mannosan, and galactosan), as important components of organic aerosol, have been widely used as molecular markers for biomass burning. Previous studies have shown that levoglucosan degrades in the atmosphere, but most of the results are derived from laboratory experiments, little is known about the decay rates and their driving factors in the real complex ambient environment. In this study, a Thermal Desorption Aerosol Gas Chromatography-Mass Spectrometry (TAG-GC/MS) was utilized to collect PM_{2.5}-bound saccharides in three typical cities across the major city clusters in eastern China (Zibo, North China Plain; Changzhou, Yangtze River Delta; and Hong Kong, Pearl River Delta region) during the autumn and winter seasons, with bihourly time resolution. **A total of 31, 21, and 69 valid decay days were identified for Zibo, Changzhou, and Hong Kong, respectively.** With the relative rate constant method, we found the daytime (8:00–16:00 LST) decay rate of levoglucosan was fastest in Changzhou, reaching $0.13 \pm 0.05 \text{ h}^{-1}$ (with a range of $0.01\text{--}0.55 \text{ h}^{-1}$), and the maximum decay rates of mannosan ($0.14 \pm 0.05 \text{ h}^{-1}$, range: $0.04\text{--}0.29 \text{ h}^{-1}$) and galactosan ($0.15 \pm 0.06 \text{ h}^{-1}$, range: $0.04\text{--}0.33 \text{ h}^{-1}$) were observed in Hong Kong. Results from the generalized additive model indicate that the daytime decay rate of anhydro-

30 saccharides is primarily influenced by aerosol liquid water content, relative humidity, and
31 atmospheric oxidation capacity, while temperature and solar surface radiation also contribute to an
32 increase in the decay rates. This study provides valuable field data on the degradation rates of
33 saccharides in real ambient environments and demonstrates that their degradation results are driven
34 by the combined effects of multiple oxidation pathways.

35 **Keywords:** Anhydro-saccharides; Degradation rates; Real atmosphere; TAG-GC/MS

36 **Highlights**

- 37 • Anhydro-saccharides (levoglucosan, mannosan, and galactosan) exhibit decreasing trends
38 during the daytime in the ambient environment.
- 39 • The decay rate of levoglucosan is fastest in Changzhou, while the decay rates of mannosan and
40 galactosan are fastest in Hong Kong.
- 41 • The daytime decay of anhydro-saccharides is primarily influenced by aerosol liquid water
42 content, relative humidity, and oxidants.

43 **1. Introduction**

44 Organic aerosol (OA) constitutes a significant component of PM_{2.5}, accounting for 20%~80%
45 of PM_{2.5} mass (He et al., 2020; Zhang et al., 2007). Biomass burning (BB) is one of the primary
46 sources of OA in the atmosphere and has significant impacts on air quality, visibility, and climate
47 (Bao et al., 2021; Liu et al., 2019). **BB releases various organic compounds, such as anhydro-**
48 **saccharides, polycyclic aromatic hydrocarbons, nitro phenolic and n-alkanes, with levoglucosan**
49 **generally being the most abundant anhydro-saccharide (Chen et al., 2017; Fang et al., 2024; Yan et**
50 **al., 2019; Zhang et al., 2022).** Levoglucosan and its two isomers, mannosan and galactosan, which
51 are produced during the pyrolysis of cellulose and hemicellulose at temperatures ranging from 150
52 to 350 °C (Fabbri et al., 2009; Hong et al., 2022; Stevens et al., 2024), have been widely used as
53 molecular markers of BB aerosol in PM_{2.5} source apportionment (Alvi et al., 2020; Cheng et al.,
54 2022; Hong et al., 2022; Kang et al., 2018; Li et al., 2021; Liang et al., 2016).

55 For decades, the anhydro-saccharides are considered as stable compounds and the majority of
56 previous studies did not consider the degradation of anhydro-saccharides during the source
57 apportionment of PM_{2.5}. However, researchers found that levoglucosan can undergo oxidation in
58 both the gas phase and liquid phase, or undergo heterogeneous oxidation on the aerosol surface (Bai

59 et al., 2013; Li et al., 2021; Zhao et al., 2014). Recent studies have proved that the contribution of
60 BB to organic carbon (OC) and PM_{2.5} could be underestimated if the degradation of anhydro-
61 saccharides was ignored (Hong et al., 2022; Li et al., 2023a). For example, Li et al. (Li et al., 2023a)
62 found that approximately 87% of levoglucosan had already degraded before reaching the receptor
63 site, causing a 14.9 % underestimation of BB-derived OC. A recent study (Wang et al., 2025) in
64 Hong Kong calculated the degradation rate constants and estimated the atmospheric lifetime of
65 levoglucosan, **but the universal impact of this phenomenon, and their driving factors that influence**
66 **this degradation process have not been thoroughly investigated (Wang et al., 2025)**. Furthermore, it
67 is currently unclear how the variations of atmospheric conditions influence the degradation rate of
68 anhydro-saccharides. Traditional studies on levoglucosan degradation typically rely on offline
69 sampling, laboratory experiments, and model simulations (Arangio et al., 2015; Bai et al., 2013;
70 Hennigan et al., 2010; Liu et al., 2019; Xu et al., 2020). However, low temporal resolution
71 observation limits the in-depth understanding of its degradation process, and discrepancies between
72 laboratory experiments and theoretical studies can introduce large biases in model simulations. With
73 the development of the Thermal Desorption Aerosol Gas Chromatography-Mass Spectrometry
74 (TAG-GC/MS) system, it is now possible to collect organic marker data with high temporal
75 resolution, capturing dynamic changes in emission sources and the aging process of organic aerosols
76 (He et al., 2020; Li et al., 2020; Wang et al., 2020; Zhang et al., 2021a; Zhao et al., 2013; Zhu et al.,
77 2021). Therefore, online observations across multiple cities under different atmospheric
78 environments are essential to better understand the degradation rates of anhydro-saccharides.

79 Zibo, Changzhou, and Hong Kong are three representative cities located in the North China
80 Plain (NCP), the Yangtze River Delta (YRD) region, and the Pearl River Delta (PRD) region,
81 respectively. These cities exhibit significant disparities in meteorological conditions, anthropogenic
82 sources, and ambient air pollution levels. These divergent environmental conditions imply that the
83 degradation process of anhydro-saccharides in the atmosphere may be influenced by different
84 factors. To investigate this, a comparative analysis of anhydro-saccharides degradation rates in these
85 cities was conducted. **The bihourly resolution data of PM_{2.5}-bound levoglucosan, mannosan and**
86 **galactosan were collected at Zibo, Changzhou and Hong Kong during various cold seasons using**
87 **the TAG-GC/MS system**. The daytime decay rates of anhydro-saccharides in three cities were
88 calculated using the relative rate constant method, and the driving factors influencing the daytime

89 decay rates of anhydro-saccharides were analyzed with the generalized additive model (GAM). The
90 findings of this study enhance our understanding of atmospheric degradation mechanisms of
91 anhydro-saccharides and provide a scientific foundation for precise source apportionment,
92 especially for refining our knowledge on the source contributions to PM_{2.5} from biomass burning.

93 **2. Methodology**

94 **2.1 Site description and field observation**

95 **Field observations using TAG-GC/MS were conducted during multiple autumn and winter**
96 **season deployments in three different typical cities in three regions (Zibo, Shandong province, NCP;**
97 **Changzhou, Jiangsu province, YRD; and Hong Kong, PRD region) over eastern China (Fig. S1).**
98 The sampling site in Zibo was situated at the Zibo Ecological Environment Monitoring Station
99 (36°50'N, 118°07'E). This location is bordered by Lutai Avenue approximately 500 m to the east,
100 Lushan Avenue approximately 100 m to the south, and residential neighborhoods approximately 1
101 km to the north. It represents the urban atmospheric environment influenced by multiple pollution
102 sources including both anthropogenic and biogenic emissions. The vegetation in the vicinity mainly
103 consists of temperate deciduous broadleaf forests. Organic compounds emitted from this area may
104 significantly impact ground-level aerosols. **The observation period was from November 2022 to**
105 **February 2023.**The sampling site in Changzhou was situated at the Changzhou Environmental
106 Monitoring Center (119°59.730'E, 31°45.510'N). The surrounding environment includes numerous
107 commercial and residential districts, as well as major roads such as Zhongwu Avenue, Heping
108 Middle Road, and Guanghua Road, representing an urban environment affected by various pollution
109 sources. Data from January to March 2021 was collected, and the detailed information of this field
110 campaign can be found in our previous studies (Li et al., 2023b; Yi et al., 2024). The sampling site
111 in Hong Kong was at the Hong Kong University of Science and Technology Super Station (114°16'E,
112 22°19'N), situated in a suburban area with relatively limited local emissions. Observations were
113 conducted from October 2020 to January 2021. During the observation periods, the anhydro-
114 saccharides data had a temporal resolution of 2 hours; however, some data were lost due to
115 instrument malfunctions or maintenance during certain intervals.

116 During the field campaign, we conducted online measurements at the three stations to collect
117 auxiliary data, including meteorological conditions, air pollutant and PM_{2.5} concentrations. **At the**

118 **Changzhou site**, meteorological parameters including wind speed (WS), wind direction (WD),
119 relative humidity (RH), temperature (T), atmospheric pressure (P), rainfall (RF) were monitored by
120 a WXT520 (VAISALA, FL); PM_{2.5} by a BAM1020 (Met One, US) via the beta-ray method; O₃, and
121 NO_x by a MODEL 49i and MODEL450i (Thermo Fisher Scientific, US), respectively; OC/EC by a
122 RT-4 (Sunset Laboratory, US), water-soluble ions (Cl⁻, NO₃⁻, SO₄²⁻, Na⁺, NH₄⁺, K⁺, Mg²⁺, Ca²⁺) and
123 NH₃ by an ADI2080 (Metrohm, CHN). Solar surface radiation (SSR) data were obtained from the
124 ERA5-Reanalysis (<https://cds.climate.copernicus.eu/datasets>). In Zibo, meteorological data
125 including wind speed (WS), wind direction (WD), relative humidity (RH), temperature (T),
126 atmospheric pressure (P), rainfall (RF) were obtained from the China Meteorological
127 Administration (<https://www.cma.gov.cn/>); PM_{2.5} by a MODEL 5014i (Thermo Fisher Scientific,
128 US); O₃, and NO_x by a MODEL 49i, and MODEL 42i (Thermo Fisher Scientific, US), respectively;
129 OC/EC by a MODEL ECOC-610 (Hangzhou Pengpu Technology Co., Ltd., China), water-soluble
130 ions (Cl⁻, NO₃⁻, SO₄²⁻, Na⁺, NH₄⁺, K⁺, Mg²⁺, Ca²⁺) and NH₃ by a MODEL S611 (Fortelice
131 International Co., Ltd., Taiwan, China); solar surface radiation (SSR) by CMP11 (Kipp & Zonen,
132 Zuid-Holland, Netherlands). At the Hong Kong site, PM_{2.5} was measured by a MODEL 5030i
133 (Thermo Fisher Scientific, US); water-soluble ions (Cl⁻, NO₃⁻, SO₄²⁻, Na⁺, NH₄⁺, K⁺, Mg²⁺, Ca²⁺),
134 NH₃ and OC/EC by an ADI2080 (Metrohm, CHN) and RT-4 (Sunset Laboratory, US), respectively;
135 O₃, and NO_x by a MODEL 49i, and MODEL 42i (Thermo Fisher Scientific, US); meteorological
136 parameters wind speed (WS), wind direction (WD), relative humidity (RH), temperature (T),
137 atmospheric pressure (P), rainfall (RF), and solar surface radiation (SSR) by the 10 m automatic
138 weather station tower. Standard calibrated sensors on the automatic weather station (AWS) tower
139 were used for measurements: a temperature/humidity sensor (HMP155) for air temperature and
140 relative humidity, an ultrasonic anemometer (CSAT3A) for wind speed and direction, a barometric
141 pressure sensor (CS106) for atmospheric pressure, a pyranometer (LI200R) for shortwave solar
142 radiation, and a tipping-bucket rain gauge (TE525MM) for precipitation at the tower base. The
143 elemental species (K, Ca) by an Xact 625i (Cooper Environmental Services) via an X-ray method.
144 Detailed information about online observations can be found in Table S1.

145 The TAG-GC/MS system is applied for the online measurement of levoglucosan, galactosan
146 and mannosan. A detailed description and the schematic diagram of the TAG can be found in our
147 previous studies (He et al., 2020; Li et al., 2020; Wang et al., 2020; Wang et al., 2025; Zhang et al.,

148 2021a). During the observations, a deuterium-labeled internal standard solution was injected into
149 each sample to monitor instrument condition and analyze the contamination levels of key species.
150 The detailed description is provided in Text S1. This study consistently utilizes deuterated
151 levoglucosan (Levoglucosan-d₇) as an internal standard to quantify three anhydro-saccharides.
152 Given that these three target analytes share the same molecular formula and possess highly similar
153 molecular structures, Levoglucosan-d₇ can effectively correct for recovery rates during extraction,
154 separation, and detection processes for all three compounds. Furthermore, this internal standard
155 shows no background interference in actual atmospheric samples, ensuring the accuracy and
156 reliability of the quantitative results. The identification of the target saccharides is achieved by
157 comparing their retention times and mass spectra with those of authentic standards. Subsequently,
158 quantitative analysis is performed using internal standard calibration. We plotted the relationship
159 between the peak area ratios of external standard solutions and the concentrations of target
160 compounds in the standard mixture to generate a calibration curve, and the correlation coefficients
161 (R) ranges from 0.92 to 0.99. The detailed information on the preparation of external standard
162 solutions and internal standard solutions for the saccharide compounds analyzed in the field
163 campaign can be found in Table S2 and Table S3.

164 2.2 ISORROPIA-II Model

165 Aerosol acidity (pH_{is}) and aerosol liquid water content (ALWC) were calculated using the
166 forward mode of the ISORROPIA-II model (<http://isorro피아.eas.gatech.edu>) (Fountoukis and Nenes,
167 2007). The input parameters required for the model primarily included water-soluble inorganic ions
168 (SO₄²⁻, NO₃⁻, NH₄⁺, K⁺, Ca²⁺, Na⁺, Mg²⁺, and Cl⁻), NH₃, temperature (T), and relative humidity
169 (Hennigan et al., 2015). The calculation formula is as follows:

$$170 \quad pH_{is} = -\lg\left(\frac{1000 \times H^+}{ALWC}\right) \quad (1)$$

171 where H⁺ represents the liquid-phase mass concentration of hydrogen ions, expressed in μg m⁻³,
172 ALWC denotes the aerosol liquid water content, expressed in μg m⁻³.

173 2.3 Relative Rate Constant Method

174 Previous studies have indicated that K⁺ can serve as a tracer for biomass burning (Hong et al.,
175 2022; Pio et al., 2008). Moreover, the ratio of levoglucosan to BB-derived K⁺ (K^{+_{BB}}) observed in
176 the environment can be used to distinguish different types of biomass burning, such as crop residue

177 burning and wood combustion (Cheng et al., 2013). Additionally, due to the chemical stability of
 178 K^+ in the atmosphere, this ratio can also serve as an indicator of BB aerosol aging. However, since
 179 potassium ions can also originate from sea salt and dust (Karavoltos et al., 2020; White, 2008). To
 180 estimate the potassium ions derived from biomass burning (K^+_{BB}), this study subtracted the
 181 contributions of sea salt and dust by Equations (2)~(7).

$$182 \quad K^+_{BB} = (K^+_{Nss}) - (K^+_{Dust}) \quad (2)$$

$$183 \quad K^+_{Nss} = K^+_{aerosol} - 0.037 * Na^+_{aerosol} \quad (3)$$

$$184 \quad K^+_{Dust} = 0.04 * [(Ca^{2+}_{Nss}) - Ca^{2+}_{BB}] \quad (4)$$

$$185 \quad Ca^{2+}_{BB} = K^+_{Nss} / z \quad (5)$$

$$186 \quad Ca^{2+}_{Nss} = Ca^{2+}_{aerosol} - 0.038 * Na^+_{aerosol} \quad (6)$$

$$187 \quad z = (K^+_{Nss} / Ca^{2+}_{Nss})_{max} - (K^+_{Nss} / Ca^{2+}_{Nss})_{min} \quad (7)$$

188 In Equations (2), K^+_{Nss} and K^+_{Dust} refer to non-sea-salt potassium and potassium originating
 189 from dust, respectively. In Equation (3), $K^+_{aerosol}$, $Na^+_{aerosol}$, and $Ca^{2+}_{aerosol}$ represent the
 190 concentrations of potassium, sodium, and calcium in the aerosol samples, which are the measured
 191 values. Based on previous literature, the mass ratios of (K^+ / Na^+) and (Ca^{2+} / Na^+) in seawater are
 192 0.037 and 0.038, respectively, and are used for the correction of sea salt aerosols (Kumar et al., 2018;
 193 Pio et al., 2007). The study of Kumar et al. (2018) suggests that the maximum and minimum
 194 differences in the mass ratio of ($K^+_{Nss} / Ca^{2+}_{Nss}$) are considered to represent emissions from biomass
 195 burning (Kumar et al., 2018; Pio et al., 2008; Pio et al., 2007). The Ca^{2+} originating from biomass
 196 burning is calculated by using ($K^+_{Nss} / Ca^{2+}_{Nss}$)_{max} minus ($K^+_{Nss} / Ca^{2+}_{Nss}$)_{min} as the denominator.

197 In this study, the calculation method for the anhydro-saccharides decay rate was adopted from
 198 Wang et al. (2025), which is a variant of the relative rate constant approach utilizing inert K^+_{BB} as
 199 the reference substance (Donahue et al., 2005; Wang et al., 2025). The validity of this method has
 200 been demonstrated in previous studies (Wang and Yu, 2021). The detailed derivation of the formula
 201 can be found in Text S2. The final derived expression is presented as Equation (8). According to
 202 Wang et al. (2025), the C_i represents the particle phase concentration of anhydro-saccharides, $C_{K^+_{BB}}$
 203 represents the concentration of K^+ from biomass combustion, k_2 represents the second-order
 204 reaction rate constant between anhydro-saccharides and oxidants, and C_{OX} represents the average
 205 concentration of oxidants in the aerosol phase. The calculated k corresponds to the effective total
 206 decay rate of anhydro-saccharides, which results from various atmospheric processes (such as

207 heterogeneous oxidation and aqueous-phase oxidation). We assume that the emission of anhydro-
208 saccharides and K^+_{BB} is equivalent, or that no new pollutants are emitted, or that such emissions are
209 negligible within the time frame of the study (i.e., 8 hours). Compared to previous studies applying
210 the relative rate constant method to a pair of target and unknown compounds (Donahue et al., 2005;
211 Huff Hartz et al., 2007), this method can be regarded as a special case where the reference species
212 is inert and zero-corrected.

$$213 \quad \frac{\partial \ln(C_i/C_{K^+_{BB}})}{\partial t} = -k, k = k_2 \times C_{OX} \quad (8)$$

214 2.4 Generalized Additive Models

215 Generalized **additive models** (GAM) are used to construct nonlinear regression relationships
216 between explanatory variables and response variables. Unlike statistical distribution-based models,
217 GAM is primarily data-driven, allowing for flexible adjustment of the functional form of the
218 response variable based on the specific context (Stone, 1985). Compared to other statistical models,
219 GAM offers higher flexibility and degrees of freedom, which does not require a pre-defined
220 parametric model, and can be applied to various distribution types, and can directly handle complex
221 nonlinear relationships between explanatory and response variables (Zhai et al., 2019). The GAM
222 model has been widely applied in studies investigating the influencing factors of nonlinear
223 atmospheric pollutants, such as $PM_{2.5}$, O_3 , and SOA (Hu et al., 2022; Zhang et al., 2021b). The basic
224 form of the GAM model is shown in Equation (9).

$$225 \quad g(\mu) = f_1(X_1) + f_2(X_2) + \dots \dots f_n(X_n) + \beta \quad (9)$$

226 In the equation, $g(\mu)$ is a continuous function representing the relationship between the
227 nonlinear formula and the expected value; μ denotes the response variable, i.e., the mass
228 concentration of the target substance; β is the intercept; f_n ($n=1, 2, \dots, n$) is the smoothing function
229 connecting the explanatory variables; X_n ($n=1, 2, \dots, n$) refers to the different explanatory variables.
230 The significance of the explanatory variables is tested using the **akaike** information criterion, and
231 the most appropriate $g(\mu)$ and X_n are selected through multiple linear tests. The R^2 , deviance
232 explained (%), and p-value calculated from the GAM model are used to assess the significance level,
233 importance of X_n , and the model's goodness of fit. **In this study, the generalized additive model**
234 **(GAM) was implemented using the LinearGAM class from the Python library pygam, with cubic**
235 **regression splines as the default basis function. The model utilized a linear link function (the default**

236 setting for LinearGAM), aligning with the assumption of a linear relationship between the response
237 variable and the smooth terms of the predictor variables. All variables were retained in their original
238 units and ranges, and no scaling was applied before model fitting. To optimize the spline-related
239 parameters, we conducted a two-dimensional grid search over the basis dimension (defined as the
240 number of splines k) and the penalty parameter (λ). The enumerated range for k was set at 8, 10,
241 and 12, while λ was exhaustively tested across a predetermined logarithmic range (from 10^{-3} to 10^3 ,
242 encompassing 10 logarithmic points). This process generated 30 parameter combinations, which
243 were evaluated by minimizing the generalized cross-validation (GCV) loss function. The results
244 indicated that when $k = 10$, the model achieved optimal performance, with the corresponding
245 optimal penalty parameter $\lambda = 46.42$. Moreover, with $k = 10$, all smooth curves of the variables
246 exhibited no anomalous oscillations or overfitting artifacts, clearly reflecting the true nonlinear
247 trends of the variables. This parameter combination achieved the best balance between fitting
248 accuracy and structural complexity, leading to the final selection of $k = 10$ for the basis dimension
249 of all predictor variables, thereby determining the optimal effective degrees of freedom (Edf) for
250 each smooth term of the predictors. This optimization step effectively balanced the accuracy of the
251 model fit with its structural complexity, enhancing the model's generalization performance on
252 unseen data. Furthermore, to optimize the set of predictor variables, a multicollinearity check was
253 performed using the variance inflation factor (VIF) before model fitting, with a conservative
254 threshold of 4 used to identify significant multicollinearity; variables with $VIF > 4$ were carefully
255 evaluated and excluded from the final model. The retained predictor variables demonstrated
256 acceptable levels of collinearity, ensuring the stability of model parameter estimates and improving
257 the reliability and interpretability of subsequent analytical results.

258 **3. Results and Discussions**

259 **3.1 Overview of the field campaign**


260 The overall environmental condition and air pollutant levels during the three field campaigns
261 are summarized in Table S4. During the field observation period, $PM_{2.5}$ pollution was the most
262 severe in Zibo, with an average concentration of $69.4 \pm 58.0 \mu g m^{-3}$. In comparison, the average
263 $PM_{2.5}$ concentrations in Changzhou and Hong Kong were $49.9 \pm 26.4 \mu g m^{-3}$ and $31.5 \pm 20.5 \mu g m^{-3}$,
264 respectively. Zibo is a traditional heavy industrial city with more than 8,200 industrial plants,

265 including power plants, chemical factories, and building materials industries (Pan et al., 2025). Due
266 to its high emission intensity, Zibo has been suffering from poor air quality, consistently ranking
267 last in air quality within Shandong Province in recent years. In contrast, the air quality in Hong
268 Kong was relatively good, which is likely related to relatively low emission intensity and favorable
269 meteorological conditions. The average winter temperature in Zibo (-0.2 ± 6.1 °C) was the lowest
270 among the three cities, with a maximum wind speed of 2.2 ± 1.7 m s⁻¹. As a city in northern China,
271 Zibo's cold and windy climate is closely associated with the influence of northern continental air
272 masses during winter. In contrast, Changzhou (10.9 ± 4.9 °C, 1.3 ± 0.7 m s⁻¹) and Hong Kong (15.6
273 ± 10.5 °C, 1.6 ± 0.5 m s⁻¹), located in southern and coastal regions, are influenced by the subtropical
274 monsoon and oceanic climate, leading to significantly higher winter temperatures and lower wind
275 speeds, resulting in a milder and more stable climate. As shown in Fig. 1, levoglucosan is the
276 dominant **anhydro-saccharides** in all three cities, with concentrations of 45.5 ± 32.3 ng m⁻³ in Zibo,
277 45.1 ± 38.7 ng m⁻³ in Changzhou, and 27.5 ± 15.6 ng m⁻³ in Hong Kong, respectively. The average
278 concentration of mannosan in Changzhou (3.6 ± 3.2 ng m⁻³) is higher than in Zibo (2.4 ± 1.7 ng m⁻³)
279 and Hong Kong (1.9 ± 1.5 ng m⁻³). Conversely, the concentration of galactosan in Zibo (4.5 ± 3.4
280 ng m⁻³) is significantly higher than in Changzhou (2.4 ± 2.0 ng m⁻³) and Hong Kong (0.9 ± 0.7 ng
281 m⁻³). These differences may reflect varying BB source types across the cities. The total
282 concentration of three **anhydro-saccharides** (levoglucosan, mannosan, and galactosan) in these three
283 cities accounts for 0.7% (Zibo), 0.9% (Changzhou), and 0.8% (Hong Kong) of the measured total
284 organic carbon (OC) mass. Although their proportion is relatively low, these **anhydro-saccharides**,
285 as characteristic tracers of biomass burning, can be used to inversely estimate the contribution of
286 BB sources to atmospheric OC. Thus, they are key indicators for quantifying the impact of BB
287 source emissions (Cheng et al., 2022; Fabbri et al., 2009; Li et al., 2023a).

288 By analyzing the ratio of levoglucosan (Lev) to mannosan (Man), different types of biomass
289 combustion sources can be identified. For example, previous studies have shown that the Lev/Man
290 ratio from crop straw combustion can exceed 40, while the Lev/Man ratios from hardwood and
291 softwood combustion range from 15 to 25 and 3 to 10, respectively (Engling et al., 2009; Fu et al.,
292 2012; Sang et al., 2013; Xu et al., 2020). We calculated the Lev/Man and Man/Gal ratios for the
293 three cities, as shown in Fig. 1. The average Lev/Man ratios in Zibo, Changzhou, and Hong Kong
294 were 19.3 ± 5.6 (range: 5.2~43.3), 13.9 ± 6.6 (0.9~56.8), and 17.8 ± 6.5 (5.6~51.4), respectively.

295 Fig. S2 illustrates the parameter ratio space of Lev/Man and Lev/K⁺ to characterize biomass burning
296 characteristics and distinguish different combustion types. The tracer ratio space diagram for
297 coniferous trees, deciduous trees, hardwood, softwood, and crop residues, proposed by Cheng et al.
298 (2013), overcomes the limitation of relying solely on a single feature ratio (such as Lev/K⁺ or
299 Lev/Man) for distinguishing biomass burning types (Cheng et al., 2013). It is important to note that,
300 due to insufficient observational data for K⁺ in Hong Kong, we selected total potassium for the
301 calculation, and after excluding the effects of sea salt and dust, the final calculation yielded
302 potassium produced by biomass combustion (K_{BB}). The calculated K_{BB} accounted for 82% of total
303 K, confirming that biomass burning (BB) was the dominant source of K in PM_{2.5} during the
304 observation period. Furthermore, K_{BB} exhibited a good correlation with levoglucosan (R_p = 0.63),
305 which further reinforces that they share the same primary source. Thus, it is reasonable to use K_{BB}
306 as a tracer for BB at the Hong Kong site. However, daily heterogeneity in the sources of potassium
307 cannot be ruled out, which may introduce systematic biases in the estimation of K_{BB} and the
308 inference of k's. Detailed information can be found in our previous study (Wang et al., 2025). The
309 ratio ranges for Zibo, Changzhou, and Hong Kong all fall within the range typically associated with
310 crop residue burning, which is consistent with previous studies. The study period coincided with the
311 autumn and winter seasons, which correspond to the typical period of crop residue burning (Cheng
312 et al., 2013; Wang et al., 2020). In addition, the ratio of mannosan to galactosan (Man/Gal) has been
313 used as an auxiliary method for distinguishing biomass combustion sources. During the observation
314 period, the average Man/Gal values for Changzhou and Hong Kong were 1.56 ± 0.75 (range:
315 $0.32\sim 13.81$) and 2.30 ± 0.64 (range: $0.84\sim 6.38$), respectively, consistent with previous studies which
316 indicate that in the combustion emissions of crop straw, grass, and coal pellets, the content of
317 mannosan (Man) is usually higher than that of galactosan (Gal) (Fabbri et al., 2009; Vicente et al.,
318 2018; Xu et al., 2020). However, the average Man/Gal value in Zibo was 0.56 ± 0.19 (range:
319 $0.23\sim 2.24$), significantly lower than those in Changzhou and Hong Kong, with a relatively higher
320 concentration of galactosan, which may be related to differences in the type of combustion source
321 or combustion conditions (Haque et al., 2022; Kuo et al., 2011; Yan et al., 2018). For example, the
322 combustion of coal and certain industrial fuels may lead to higher galactosan content due to
323 differences in the organic composition of these fuels compared to biomass fuels (Yan et al., 2018).
324 Furthermore, incomplete combustion or low-temperature combustion may increase galactosan

325 concentration (Haque et al., 2022), which could be a characteristic feature of combustion in Zibo.
 326 As a heavy industrial city, Zibo has more industrial combustion sources and incomplete combustion
 327 phenomena, leading to the relative enrichment of galactosan and exhibiting distinct chemical
 328 characteristics compared to common biomass combustion. In addition, residential coal combustion
 329 for heating was also an important emission source in suburban areas of North China in winter, which
 330 is corresponding to the low-temperature combustion scenario in Zibo.



331
 332 **Fig. 1** Boxplot of the concentrations of levoglucosan (Lev), mannosan (Man), galactosan (Gal),
 333 and their ratios (Lev/Man, Man/Gal) in the cities of Zibo, Changzhou, and Hong Kong.

334 **3.2 Diurnal variations of anhydro-saccharides and ratios of Lev/K⁺_{BB}, Man/K⁺_{BB},**
 335 **Gal/K⁺_{BB}**


336 Fig. 2 shows the detailed diurnal variations of anhydro-saccharides across three cities. The
 337 anhydro-saccharides in all three cities exhibit similar diurnal variation characteristics, indicating the
 338 similar source and atmospheric degradation processes. Specifically, the concentrations of anhydro-
 339 saccharides in the three cities generally decrease during the daytime (8:00–16:00 LST) and increase
 340 at night. Notably, in Fig. 2(b), the decline of levoglucosan in Changzhou is most pronounced during
 341 the daytime. In Changzhou, daytime levoglucosan concentrations peak at 08:00 LST (49.26 ± 5.71
 342 ng m^{-3}) before gradually declining to a minimum of $29.53 \pm 4.54 \text{ ng m}^{-3}$ by 14:00 LST. A slight
 343 rebound to $30.56 \pm 4.56 \text{ ng m}^{-3}$ occurs at 16:00 LST, followed by a significant increase to a
 344 secondary peak of $59.88 \pm 5.71 \text{ ng m}^{-3}$ at 20:00 LST. In addition, Fig. 2(c) show the diurnal variation

345 of levoglucosan in Hong Kong exhibits a pronounced pattern, with peak concentrations occurring
346 in the morning at 8:00 LST, ($32.65 \pm 1.72 \text{ ng m}^{-3}$). Similar to Changzhou, the levoglucosan
347 concentration drops to a minimum at 14:00 LST ($22.40 \pm 1.24 \text{ ng m}^{-3}$), followed by a slight rebound
348 to $23.05 \pm 1.20 \text{ ng m}^{-3}$ at 16:00 LST. The diurnal variation of levoglucosan in Zibo is relatively flat,
349 peaking at 8:00 LST ($48.24 \pm 4.29 \text{ ng m}^{-3}$) and declining to $39.83 \pm 3.30 \text{ ng m}^{-3}$ at 16:00 LST.
350 Similarly, the concentration of mannosan in Changzhou shows the most significant daytime
351 decrease, from a peak at 8:00 ($3.89 \pm 0.40 \text{ ng m}^{-3}$) to a minimum at 16:00 ($2.42 \pm 0.45 \text{ ng m}^{-3}$). A
352 significant decline was also observed in Hong Kong, from $2.42 \pm 0.20 \text{ ng m}^{-3}$ at 8:00 to 1.47 ± 0.10
353 ng m^{-3} at 16:00 LST. In contrast, the diurnal variation of mannosan in Zibo is relatively smooth,
354 with an initial concentration of $2.38 \pm 0.19 \text{ ng m}^{-3}$ at 08:00 LST, and decreases to $2.28 \pm 0.18 \text{ ng m}^{-3}$
355 at 16:00 LST. Galactosan in all three cities exhibited a congruent diurnal trend to the other anhydro-
356 saccharides, characterized by a daytime decrease. For instance, galactosan concentrations in Zibo
357 declined from a peak of $4.60 \pm 0.38 \text{ ng m}^{-3}$ at 8:00 LST to $4.00 \pm 0.33 \text{ ng m}^{-3}$ at 16:00 LST. In
358 Changzhou, the galactosan concentration was $2.65 \pm 2.39 \text{ ng m}^{-3}$ at 8:00 LST, and it dropped to 1.79
359 $\pm 0.30 \text{ ng m}^{-3}$ at 16:00 LST. In Hong Kong, the galactosan concentration decreased from 1.08 ± 0.09
360 ng m^{-3} at 8:00 LST to $0.67 \pm 0.06 \text{ ng m}^{-3}$ by 16:00 LST. The concentration and compositional
361 characteristics of organic aerosols in real atmospheric environments are governed by a combination
362 of factors, including the intensity of air pollutant source emissions, variations in meteorological
363 conditions, atmospheric chemical reactions, atmospheric diffusion capabilities, and deposition
364 processes. These interrelated influences collectively impart high complexity and uncertainty to both
365 the concentration levels and chemical composition of organic aerosols (Chen et al., 2022; Kim et
366 al., 2017; Zhang et al., 2013). However, relative concentration is a more effective indicator for
367 detecting the loss of organic species particles. In addition, previous studies have indicated that the
368 ratio of levoglucosan to potassium ions (K^+) from BB sources ($\text{levoglucosan}/\text{K}^+_{\text{BB}}$) is an effective
369 indicator of the aging degree of BB aerosols (Cheng et al., 2013; Li et al., 2021; Mochida et al.,
370 2010). Therefore, to investigate the diurnal decay pattern of anhydro-saccharides, we selected
371 $\text{C}_i/\text{K}^+_{\text{BB}}$ to examine the decay of anhydro-saccharides. As shown in Fig. 2, the diurnal variations of
372 anhydro-saccharides/ K^+_{BB} in the three cities also exhibit a decreasing trend from 08:00 to 16:00,
373 providing a key basis for the subsequent calculation of the decay rate of anhydro-saccharides.
374 Furthermore, the utilization of this ratio can effectively eliminate the confounding effects of

375 boundary layer dynamics (e.g., diurnal variations in mixing height and vertical diffusion), since both
376 anhydro-saccharides and K^+_{BB} are derived from biomass burning and thus subject to the same
377 boundary layer-driven dilution processes. In Zibo and Changzhou, the K^+_{BB} concentrations were
378 calculated using Equations (2)–(7), and the results showed that K^+_{BB} accounted for 93.4% and 91.4%
379 of the total K^+ , respectively, thus biomass burning as the major source of K^+ in $PM_{2.5}$ during the
380 sampling campaign. Correlation analysis results showed a significant positive correlation between
381 levoglucosan and K^+_{BB} across different sites (as shown in Fig. S3). In Zibo, the Pearson correlation
382 coefficients between levoglucosan, mannosan, and galactosan with K^+_{BB} were 0.65, 0.52, and 0.52,
383 respectively. In Changzhou, the Pearson correlation coefficients between levoglucosan, mannosan,
384 and galactosan with K^+_{BB} were 0.76, 0.58, and 0.54, respectively, further confirming the validity
385 and reliability of the K^+_{BB} calculation formulas for Zibo and Changzhou.

386 The detailed time series of anhydro-saccharides and K^+_{BB} for the three cities is presented in
387 Fig. S4. Overall, the concentrations of anhydro-saccharides and K^+_{BB} in these three locations exhibit
388 a synchronous increase and decrease trend, verifying the similarity in their sources. From Fig. S4
389 (a) and (b), it is evident that the concentrations of anhydro-saccharides and K^+_{BB} in Zibo and
390 Changzhou exhibit frequent spikes within a short time frame, aligning with the characteristics of
391 concentrated emissions associated with open burning of straw during the autumn and winter seasons
392 in these regions. Such combustion under high temperatures and strong oxygen supply conditions
393 generates a higher proportion of levoglucosan (Chen et al., 2017; Cheng et al., 2013; Fabbri et al.,
394 2009). Additionally, Zibo, as a typical heavy industrial city, experiences biomass combustion
395 emissions that are influenced by the incomplete combustion processes of industrial burning and
396 residential heating with coal at low temperatures, leading to higher concentrations of galactosan
397 compared to mannosan (Haque et al., 2022; Yan et al., 2018). These factors collectively enhance the
398 chemical characteristic differences between Zibo and other sites. In contrast, Fig. S4(c) shows that
399 the time series from Hong Kong displays stable fluctuations in the concentrations of anhydro-
400 saccharides and K^+_{BB} , without frequent sudden peaks. This aligns with the biomass burning pattern
401 in this region, primarily relying on residential cooking and small-scale commercial activities (such
402 as wood and kitchen waste) (Lee et al., 2013; Leung et al., 2024). The biomass burning sources in
403 Hong Kong are characterized by low emission intensity and a sustained, dispersed release process.
404 The regional characteristics of these three cities provide important background and data support for

405 subsequent analyses of the daytime decay rates of anhydro-saccharides, aiding in the deeper
 406 understanding of their sources and variations under different environmental conditions, thereby
 407 laying the foundation for developing relevant pollution control strategies and optimization plans.




408
 409 **Fig. 2** Diurnal variations of levoglucosan, mannosan, and galactosan (left column); Lev/K⁺_{BB},
 410 Man/ K⁺_{BB}, and Gal/ K⁺_{BB} ratios (right column) at (a) Zibo, (b) Changzhou and (c) Hong Kong.

411 3.3 Daytime Decay Rate Calculation

412 Using equation (8), we calculated the daytime decay rates of anhydro-saccharides in the three
 413 cities. The decay rate of levoglucosan in Changzhou was $0.13 \pm 0.05 \text{ h}^{-1}$ (range: $0.01\sim 0.55 \text{ h}^{-1}$),
 414 ranking first among the three cities, indicating a significant decay rate. The decay rates of
 415 levoglucosan in Zibo and Hong Kong were similar, with values of $0.10 \pm 0.08 \text{ h}^{-1}$ (range: $0.001\sim 0.34$
 416 h^{-1}) and $0.10 \pm 0.05 \text{ h}^{-1}$ (range: $0.02\sim 0.25 \text{ h}^{-1}$), respectively. However, Hong Kong had the fastest
 417 decay rates for the other two anhydro-saccharides, especially galactosan, with a decay rate of 0.15

418 $\pm 0.06 \text{ h}^{-1}$ (range: $0.04\sim 0.33 \text{ h}^{-1}$), much higher than the other cities. In comparison, Changzhou's
 419 galactosan decay rate was $0.13 \pm 0.08 \text{ h}^{-1}$ (range: $0.01\sim 0.67 \text{ h}^{-1}$), and Zibo's galactosan decay rate
 420 was $0.10 \pm 0.03 \text{ h}^{-1}$ (range: $0.004\sim 0.31 \text{ h}^{-1}$), both of which were relatively smaller. Similarly, the
 421 decay rate of mannosan in Hong Kong was the highest, reaching $0.14 \pm 0.05 \text{ h}^{-1}$ (range: $0.04\sim 0.29$
 422 h^{-1}), followed by Changzhou at $0.13 \pm 0.07 \text{ h}^{-1}$ (range: $0.01\sim 0.60 \text{ h}^{-1}$), while Zibo had the lowest
 423 rate at only $0.09 \pm 0.03 \text{ h}^{-1}$ (range: $0.01\sim 0.33 \text{ h}^{-1}$). The detailed decay rates of anhydro-saccharides
 424 for the three cities are presented in Table S5. Based on the distribution of decay rates in three cities
 425 shown in Fig. 3, the average decay rate of levoglucosan in Changzhou is higher than in the other
 426 two cities, with 33.3% of the rates exceeding 0.15 h^{-1} . In contrast, the proportions for Zibo and Hong
 427 Kong are 12.9% and 13.0%, respectively. Moreover, Hong Kong shows a remarkable distribution
 428 of decay rates for mannosan and galactosan, with 37.7% of mannosan decay rates and 37.7% of
 429 galactosan decay rates exceeding 0.15 h^{-1} . These results indicate that in the three cities, Changzhou
 430 exhibits the highest average decay rate for levoglucosan, while Hong Kong shows the highest decay
 431 rates for mannosan and galactosan, and Zibo has the lowest decay rates for all three sugars.



432

433 **Fig. 3 Distribution of the decay rates of levoglucosan, mannosan and galactosan for (a) Zibo, (b)**


434

Changzhou and (c) Hong Kong.

435 Fig. 4 shows examples of linear fitting in the three cities, with the x-axis representing five time
436 points: 8:00, 10:00, 12:00, 14:00 and 16:00 each day, and the y-axis representing $\ln(\frac{C_i}{C_{KB}})$, with the
437 slope equal to k. In such a short time frame (approximately 8-hours), the combustion conditions of
438 burning plants and biomass sources are expected to remain relatively constant, making the influence
439 of source emissions on the changes in normalized sugar concentration negligible. The obtained k
440 values can be regarded as the average daytime decay rate for the day of observation. However, the
441 observational data for certain days do not conform to the linear decay pattern. This phenomenon is
442 primarily attributed to factors such as interference from fresh emissions and changes in air mass
443 origins, which cannot be directly ascertained based solely on ground-based observations. The
444 presence of invalid data during the fitting process does not contradict the research hypothesis; it
445 merely reflects the complexity of real ambient conditions. During the observation period, each day
446 was characterized by distinct biomass burning (BB) emission intensities (or the absence of BB
447 emissions), accompanied by varying meteorological conditions and oxidant levels. For days with
448 poor fitting performance or negative values of the rate constant k, this only indicates that the
449 combination of atmospheric conditions on those days violated the core assumptions underlying
450 Equation (8). The method employed in this study is only applicable to specific scenarios that satisfy
451 the fundamental assumptions, and the two-hourly measurement data enabled us to sample the target
452 compounds under a wide range of diurnal ambient conditions. The fitting results for the three cities
453 are as follows: Zibo sampled for a total of 67 days, with 31 days fitting the linear decay pattern
454 (46%); Changzhou sampled 45 days, with 21 days fitting the linear decay pattern (47%); and Hong
455 Kong sampled 106 days, with 69 days fitting the linear decay pattern (65%). The proportion of days
456 in Hong Kong that could be linearly fitted is significantly higher than that in Zibo and Changzhou.
457 This outcome may be closely related to the differences in climatic conditions and burning practices
458 across regions.

459 The analysis in Fig. S4 indicates that both Zibo and Changzhou experience frequent outdoor
460 straw burning activities in autumn and winter, where continuous fresh emissions release substantial
461 amounts of anhydro-saccharides and other biomass combustion tracers into the atmosphere. The
462 emission rates of these tracers within an 8-hour window far exceed the natural diffusion and
463 chemical degradation rates of pollutants, disrupting the stable concentration changes required for

464 linear decay, ultimately resulting in fitting failures. Furthermore, we compared the concentrations
 465 of atmospheric oxidants (O_x) on successful and unsuccessful fitting days. The results show that the
 466 mean concentrations of atmospheric oxidants on successful fitting days are higher across all three
 467 cities compared to unsuccessful fitting days. Specifically, the atmospheric oxidant concentration on
 468 successful fitting days in Zibo was 41.2 ± 8.5 ppb (range: 24.2~61.8 ppb), while it was 37.6 ± 7.8
 469 ppb (range: 17.9~65.7 ppb) on unsuccessful fitting days. In Changzhou, the concentration for
 470 successful days was 46.1 ± 13.0 ppb (range: 21.6~81.8 ppb), compared to 44.3 ± 10.6 ppb (range:
 471 14.6~60.8 ppb) on unsuccessful days. In Hong Kong, successful fitting days showed an atmospheric
 472 oxidant concentration of 31.9 ± 6.3 ppb (range: 19.7~53.4 ppb), while the concentration on
 473 unsuccessful days was 27.8 ± 6.8 ppb (range: 14.9~42.0 ppb). These results suggest that the relative
 474 deficiency of atmospheric oxidants may also be a significant factor preventing the concentration of
 475 anhydro-saccharides from exhibiting a clear linear decay pattern. To mitigate the bias arising from
 476 days with unsuccessful fitting, this study exclusively retained days characterized by positive decay
 477 rates and a coefficient of determination ($R^2 > 0.5$) for the estimation of decay rates and subsequent
 478 generalized additive model (GAM) analysis. The inclusion of such poorly fitted data points would
 479 artificially depress the average rate constant k , thereby potentially resulting in an underestimation
 480 of the authentic degradation rate



481
 482 **Fig. 4 Example calculation of the degradation rates of levoglucosan, mannosan and galactosan at**

483 **(a) Zibo, (b) Changzhou and (c) Hong Kong.**

484 The three anhydro-saccharides exhibit differences in their molecular structures, particularly
485 concerning the C-H bonds situated at various positions on the sugar rings. This results in variations
486 in the reaction potential of hydroxyl radicals ($\cdot\text{OH}$) with each sugar molecule. In the same city, all
487 three sugars undergo oxidation by $\cdot\text{OH}$ radicals. We conducted a correlation analysis of the decay
488 rates of these anhydro-saccharides across different cities. As illustrated in Fig. S5, there is a strong
489 correlation among the decay rates of anhydro-saccharides in the three cities. This suggests that
490 despite the differences in molecular structures, the oxidation mechanisms involving $\cdot\text{OH}$ are similar,
491 accounting for the high correlation between their decay rates. The differences in decay rates for the
492 three anhydro-saccharides in the same city may also relate to bond dissociation enthalpy (BDE).
493 According to John et al. (2020a and 2020b), the BDE of levoglucosan, mannosan, and galactosan
494 can be estimated using the accurate bond dissociation enthalpy tool (ALFABET) available online
495 (<https://bde.ml.nrel.gov/>, last accessed: October 21, 2025). The findings indicate that the C-H bonds
496 that are most susceptible to breakage in levoglucosan, mannosan, and galactosan are predominantly
497 located at positions 2 and 3, with corresponding BDE of 85.3–86.9 kcal mol⁻¹, 84.6–85.1 kcal mol⁻¹,
498 and 82.9–84.6 kcal mol⁻¹, respectively. This implies that the decay rates of the three anhydro-
499 saccharides should follow the order: galactosan > mannosan > levoglucosan (St John et al., 2020a;
500 St John et al., 2020b). However, this trend was only observed in Hong Kong and Changzhou. Hence,
501 besides bond dissociation enthalpy (BDE), there should be other influencing factors affecting the
502 decay rates. Section 3.4 provides a detailed exploration of the environmental factors that influence
503 the decay rates of anhydro-saccharides, with a particular focus on Zibo.

504 **3.4 Driving factors of the decreasing rate of anhydro-saccharides**

505 The three sampling sites represent cities with distinct meteorological conditions. Zibo has a
506 temperate monsoon climate, characterized by cold and dry winters. In contrast, Changzhou falls
507 under the subtropical monsoon climate, with winters being more humid than those in Zibo. Although
508 Hong Kong also has a subtropical monsoon climate, it is significantly influenced by the oceanic
509 climate, resulting in smaller temperature variations and a more humid winter. The differences in
510 climate conditions indirectly lead to variations in environmental factors, which in turn affect the
511 daytime degradation rate of anhydro-saccharides across different cities. Therefore, we compared the
512 environmental factors of the three cities with the calculated degradation rates of anhydro-
513 saccharides. These factors include ALWC related to liquid-phase reactions, the atmospheric

514 oxidative capacity indicator O_x , solar surface radiation (SSR), relative humidity (RH) and
515 temperature (T). Due to the lack of data on gas-phase anhydro-saccharides, the gas-phase oxidation
516 part was not discussed in this study.

517 As an indicator of the total amount of various oxidants in the atmosphere, O_x is used in this
518 study to explore its impact on the daytime degradation rate of levoglucosan. As shown in Fig. 5(b),
519 the O_x concentration in Changzhou is higher than in the other two cities, with an average value of
520 45.1 ± 11.8 ppb(range: 14.6~81.8 ppb). In contrast, O_x levels in Zibo (39.6 ± 8.3 ppb) are slightly
521 lower than Changzhou but much higher than that in Hong Kong (31.9 ± 6.3 ppb). Hoffmann et al.
522 (2010) reported that the reaction of levoglucosan with $\cdot OH$ is a major degradation pathway, and
523 model calculations indicate that levoglucosan is more readily oxidized by $\cdot OH$ during the daytime.
524 The average degradation flux during the winter daytime is $4.7 \text{ ng m}^{-3} \text{ h}^{-1}$ (Hennigan et al., 2010). On
525 the other hand, solar surface radiation (SSR), the key parameter for daytime $\cdot OH$ production,
526 showed similar levels in Changzhou and Hong Kong. The average solar radiation in Changzhou is
527 $381.7 \pm 165.6 \text{ W m}^{-2}$ (range: 147.3~779.0 W m^{-2}), while Hong Kong records $354.0 \pm 108.1 \text{ W m}^{-2}$
528 (range: 133.7~547.0 W m^{-2}). Zibo, by contrast, is markedly lower at $273.6 \pm 81.8 \text{ W m}^{-2}$ (range:
529 $79.8\sim 427.1 \text{ W m}^{-2}$). Slade and Knopf (2014) reported that the presence of water can reduce particle
530 viscosity, thereby enhancing $\cdot OH$ oxidation (Slade and Knopf, 2014). Therefore, the higher
531 degradation rate of levoglucosan in Changzhou compared to the other two cities may reflect
532 stronger $\cdot OH$ oxidation. As illustrated in Fig. 5(b), Changzhou exhibits the highest ALWC among
533 the three cities, averaging $15.6 \pm 15.5 \text{ } \mu\text{g m}^{-3}$ (range: 0.4~56.5 $\mu\text{g m}^{-3}$). Zibo follows closely at 13.4
534 $\pm 15.4 \text{ } \mu\text{g m}^{-3}$ (range: 0.7~55.7 $\mu\text{g m}^{-3}$), while Hong Kong records the lowest levels at $6.4 \pm 8.9 \text{ } \mu\text{g}$
535 m^{-3} (range: 0.2~39.7 $\mu\text{g m}^{-3}$). Furthermore, Slade and Knopf (2014) noted that an increase in relative
536 humidity accelerates the heterogeneous oxidation rate of levoglucosan (Slade and Knopf, 2014).
537 The relative humidity in Hong Kong is significantly higher than in the other two cities, with an
538 average of $52.8 \pm 13.0\%$ (range: 15.8~88.0%). Zibo has the lowest relative humidity, with an
539 average of $41.2 \pm 16.4\%$ (range: 18.4~89.8%), while Changzhou has an average relative humidity
540 of $48.9 \pm 19.2\%$ (range: 20.4~80.2%). When the relationship between the response variable and
541 explanatory variables is unclear, the generalized additive model (GAM) can be used to fit the
542 explanatory and response variables by plotting smooth functions, further assessing their linear or
543 nonlinear relationship (Shrestha, 2020; Stone, 1985; Xiao et al., 2018). In this study, we incorporated

544 the calculated effective decay days into the GAM model. The daytime degradation rate of anhydro-
545 saccharides calculated for the three cities was used as the response variable in the GAM model, and
546 the various influencing factors (O_x , ALWC, SSR, RH and T) were used as the corresponding
547 explanatory variables. Additionally, the model's accuracy was assessed by examining the R^2 , p-
548 values, and deviance explained (DE). The corresponding GAM is expressed as Equations (10)~(12).
549 Notably, this study used the natural logarithmic function of the ratio of anhydro-saccharides to K^+_{BB}
550 for linear fitting to analyze and calculate the degradation rate of anhydro-saccharides. This ratio
551 effectively mitigates the interference of boundary layer dynamical processes, such as diurnal
552 variations in mixing layer height and atmospheric vertical diffusion, which are meteorological
553 dilution effects. The underlying principle is that both anhydro-saccharides and K^+_{BB} originate from
554 biomass combustion and are subject to the same physical dilution effects driven by boundary layer
555 processes during atmospheric transport. Consequently, their ratio can cancel out these physical
556 effects on the concentration of individual species. Thus, the degradation rates included in the GAM
557 model have removed the contributions from meteorological physical dilution effects, reflecting
558 solely the intrinsic chemical degradation process of anhydro-saccharides.

$$559 \quad g(Lev) = s(ALWC) + s(T) + s(O_x) + s(RH) + s(SSR) + \beta \quad (10)$$

$$560 \quad g(Man) = s(ALWC) + s(T) + s(O_x) + s(RH) + s(SSR) + \beta \quad (11)$$

$$561 \quad g(Gal) = s(ALWC) + s(T) + s(O_x) + s(RH) + s(SSR) + \beta \quad (12)$$

562 In the equations, β represents the model intercept, while $s(i)$ denotes the smooth function
563 corresponding to each influencing factor. Table S7 presents the relevant parameters of the GAM
564 smooth functions, including effective degrees of freedom (Edf), reference degrees of freedom
565 (Ref.df), F-statistic (F), and p-value (p). These parameters are used to reflect the strength and
566 significance of the nonlinear associations between each influencing factor and the degradation rate
567 of anhydro-saccharides.





Fig. 5 Comparison of environmental factors and decay rates across the three cities: (a) decay rates of levoglucosan, mannosan and galactosan, (b) O_x, ALWC, SSR, RH and T.

To avoid the potential impact of multicollinearity among variables on the stability of model fitting and the accuracy of parameter estimates, this study performed a multicollinearity assessment on the explanatory variables using the variance inflation factor (VIF) prior to conducting the GAM analysis. A high VIF value points to strong multicollinearity between a given explanatory variable and the remaining explanatory variables; specifically, a VIF value exceeding 4 denotes the existence of significant multicollinearity in regression analysis (Shrestha, 2020; Xiao et al., 2018). The results of the multicollinearity test are presented in Table S6, where the VIF for all explanatory variables is less than 4, indicating that they successfully passed the multicollinearity assessment. The validation results of the GAM model are shown in Fig. S6. From the residual Q-Q plot (Fig. S6a), it can be observed that most of the data points approximately follow a straight line, indicating that the residuals of the GAM model generally follow a normal distribution. Fig. S6 (b) shows the scatter plot of the residuals versus the model's predicted values, which indicates that the residuals are randomly distributed. The residual histogram (Fig. S6c) demonstrates a rough symmetric distribution. Additionally, Fig. S6 (d) reveals that the observed values closely follow the “1:1” line, indicating a good fit between the observed and fitted values. These results confirm that the GAM model provides a good fit for the daytime attenuation rate of levoglucosan.

The analysis results from the GAM model for the decay rate of levoglucosan are shown in Table S7. The adjusted R² between the observed and estimated values was 0.66, with a bias explanation rate of 65.8%. The daytime decay rate of levoglucosan significantly increased with the rise in ALWC (p<0.05). As shown in Fig. 6(a), especially ALWC > 30 μg m⁻³, the decay rate increased as ALWC increased, indicating that ALWC is an important factor affecting the daytime


592 decay rate. This result is consistent with the findings of Slade and Knopf (2014), which suggest that
593 liquid-phase reactions reduce the viscosity of levoglucosan particles, allowing for easier absorption
594 of $\cdot\text{OH}$ and accelerating the daytime decay process (Slade and Knopf, 2014). We also find that the
595 decay rate significantly increased with temperature (T) ($p < 0.05$). However, temperature exerts a
596 modest positive contribution to the decay rate when $T < 10^\circ\text{C}$. In contrast, when $T > 10^\circ\text{C}$, the decay
597 rate increases markedly, indicating that the rise in temperature affects the decay rate. **An increase in**
598 **temperature generally accelerates the molecular thermal motion, elevates the collision frequency**
599 **between reactive radicals (e.g., OH radical and SO_4^- radical) with anhydro-saccharides molecules,**
600 **and thereby facilitates the occurrence of oxidative degradation reactions (Bai et al., 2013; Lai et al.,**
601 **2014). During the daytime, temperature increases is usually accompanied by high solar surface**
602 **radiation intensity; their synergistic effects can significantly promote atmospheric photochemical**
603 **reactions and the generation of reactive radicals, thus accelerating the degradation of anhydro-**
604 **saccharides (Bai et al., 2013; Wennberg, 2006). Additionally, the decay rate significantly increased**
605 **with the rise of O_x ($p < 0.05$). As shown in Fig. 6(a), the decay rate almost increased linearly with O_x .**
606 **When the $\text{O}_x > 80$ ppb, the decay rate is higher than 0.1 h^{-1} , suggesting that oxidants become the key**
607 **driving factor for the attenuation reaction under this condition. This study also found that relative**
608 **humidity ($p = 0.08$) and SSR ($p = 0.12$) did not significantly affect the decay rate. However, Fig.**
609 **6(a) shows that both are positively correlated with the daytime decay rate of levoglucosan,**
610 **especially $\text{RH} > 60\%$, which nearly increases linearly. Although SSR did not show a significant**
611 **upward trend when it was less than 400 W m^{-2} , when SSR exceeded 400 W m^{-2} , the decay rate**
612 **increased with the rise in SSR, suggesting that SSR still has some effect on the decay rate. Further**
613 **univariate GAM tests showed that, after excluding the interference of other variables, RH and SSR**
614 **were significantly positively correlated with the daytime decay rate of levoglucosan ($p < 0.05$). This**
615 **difference may arise from the weak correlation between RH and SSR and other variables, which**
616 **was partially masked in the multivariable model. Additionally, the multivariable model had lower**
617 **statistical power to test individual variables, potentially failing to identify effects that were close to**
618 **the significance level. The other two anhydro-saccharides, mannosan and galactosan, exhibited**
619 **similar characteristics to levoglucosan with respect to ALWC, RH, O_x , and SSR. As shown in Fig.**
620 **6(b) and Fig. 6(c), when temperature (T) $< 10^\circ\text{C}$, the positive contribution of temperature to the**
621 **decay rates of mannosan and galactosan was greater than levoglucosan. This indicates that although**

622 all three are anhydro-saccharides derived from biomass burning and their decay processes are
 623 generally regulated by similar environmental factors, mannosan and galactosan are more sensitive
 624 to temperature, likely attributed to differences in chemical stability induced by variations in their
 625 molecular structures. Hong Kong has the highest average temperature among the three cities, which
 626 may explain why the daytime degradation rates of mannosan and galactosan are higher in Hong
 627 Kong than in the other two cities.



628
 629 **Fig. 6 Influences of various factors on the daytime degradation rates of different saccharides**
 630 **analyzed using the GAM model: (a) levoglucosan, (b) mannosan, and (c) galactosan, as a function**
 631 **of ALWC, T, RH, O_x , and SSR. (The solid lines in each subplot represent the partial dependence**
 632 **trends of the saccharide degradation rates with respect to the corresponding factors. The shaded**
 633 **areas indicate the 95% confidence bands, reflecting the uncertainty of the model predictions at a**
 634 **95% confidence level)**

635 As shown in Fig. 6, ALWC, RH, and O_x are the three factors that contribute most to the
 636 degradation rate of anhydro-saccharides. Therefore, we plotted these factors against the degradation
 637 rate to better explore their relationship. As illustrated in Fig. 7, overall, all three factors show a
 638 positive correlation with the levoglucosan degradation rate. Specifically, as the concentration of O_x
 639 increases, relative humidity (RH) rises, and the liquid water content (ALWC) increases, the
 640 degradation rate of levoglucosan tends to increase. In contrast, when the levels of these three factors
 641 are relatively low, the degradation rate generally remains in a lower range, further confirming that
 642 these factors play a prominent role in the degradation process of levoglucosan.



643

644 **Fig. 7 Relationships between decay rates of levoglucosan (a), mannosan (b), and galactosan (c)**

645

with O_x, RH, and ALWC

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664


665

666

667

Based on the results from the GAM model, we separately examined the contributions of five factors to the degradation rates of dehydrated sugars in Zibo, as shown in Fig. 8. The results indicate that O_x contributes more significantly to the degradation rate of levoglucosan than the other two anhydro-saccharides, while aerosol liquid water content (ALWC) contributes similarly to the degradation rates of mannosan and galactosan, which is significantly higher than that of levoglucosan. These findings demonstrate that the degradation rates of anhydro-saccharides are not only correlated with their structural BDE but also regulated by other driving factors with distinct contribution patterns. Even within the macro atmospheric environment of the same city, microenvironmental differences in the aerosol particles hosting the three anhydro-saccharides can modulate their oxidation processes. For instance, variations in aerosol phase state (e.g., viscosity influenced by RH and T) may affect oxidant diffusion, leading to differential rates, as higher RH in Hong Kong could reduce viscosity and enhance heterogeneous oxidation for mannosan and galactosan (Slade and Knopf, 2014). Additionally, aqueous-phase oxidation (promoted by high ALWC) may dominate in Changzhou, while heterogeneous pathways prevail in lower-ALWC sites like Hong Kong, explaining pathway-specific sensitivities beyond BDE (Lai et al., 2014; Slade and Knopf, 2014). Differences in aerosol composition, such as inorganic ions (e.g., SO₄²⁻, NO₃⁻, NH₄⁺), further influence ALWC and viscosity; for example, higher sulfate in Zibo may acidify aerosols, potentially slowing certain pathways (Riva et al., 2016a; Riva et al., 2016b). Collectively, the combined effects of multiple driving factors and the differential sensitivities of the three anhydro-saccharides to these factors lead to inconsistent intensities of oxidation pathways (e.g., aqueous-phase oxidation and heterogeneous oxidation), thereby producing the divergent degradation rates observed. This may help explain why the degradation rates of the three dehydrated sugars in Zibo

668 do not follow the pattern predicted by BDE.



669

670 **Fig. 8 Contributions of ALWC, T, RH, SSR and O_x to the decay rates of levoglucosan, mannosan,**
671 **and galactosan in Zibo**

672 In conclusion, the analysis results from the GAM model suggest that the daytime decay rate of
673 anhydro-saccharides is influenced by multiple factors, with ALWC, RH, and O_x being the main
674 driving factors. Despite the lack of significance in the effects of Tand SSR, they still showed a
675 positive correlation with the decay rate. Given that the sampling times at the three sampling sites in
676 this study concentrated in the autumn and winter, it is expected that in summer, under conditions of
677 higher O_x, temperature, and SSR, the decay rate of anhydro-saccharides will significantly increase.
678 Therefore, further investigation into the degradation mechanisms of anhydro-saccharides is crucial
679 for accurately assessing the contribution of BB aerosols to global air quality, particularly in the
680 context of seasonal and environmental changes, and holds significant scientific and practical value.

681 **4. Conclusions**

682 This study employed TAG-GC/MS to obtain bihourly time resolution PM_{2.5}-bound anhydro-
683 saccharides (levoglucosan, mannosan, and galactosan) during the winter season in three typical
684 cities over three regions in China, including Zibo, Changzhou and Hong Kong, located in the NCP,
685 YRD and PRD regions, respectively. The decay rates of levoglucosan, mannosan, and galactosan in
686 the real atmosphere and the driving factors are investigated. Results indicate that levoglucosan had
687 the highest concentration among all the three anhydro-saccharides. In Zibo, the concentration of
688 levoglucosan was $45.5 \pm 32.3 \text{ ng m}^{-3}$, higher than Changzhou ($45.1 \pm 38.7 \text{ ng m}^{-3}$) and Hong Kong
689 ($27.5 \pm 15.6 \text{ ng m}^{-3}$). The diurnal variation of the three anhydro-saccharides showed a decreasing

690 trend during the daytime (8:00–16:00). We selected K^+_{BB} as a reference species and calculated the
691 daytime degradation rates of the three anhydro-saccharides in the three cities using the relative rate
692 constant method. The results indicated that the degradation rate of levoglucosan was highest in
693 Changzhou, at $0.13 \pm 0.05 \text{ h}^{-1}$ (range: $0.01\sim 0.55 \text{ h}^{-1}$), while mannosan and galactosan showed the
694 highest degradation rates in Hong Kong, at $0.14 \pm 0.05 \text{ h}^{-1}$ (range: $0.04\sim 0.29 \text{ h}^{-1}$) and $0.15 \pm 0.06 \text{ h}^{-1}$
695 (range: $0.04\sim 0.33 \text{ h}^{-1}$), respectively. Due to structural differences, particularly the varying positions
696 of the C-H bond within the sugar ring, the reaction mechanisms of hydroxyl radicals ($\cdot\text{OH}$) with the
697 three sugar molecules differ, leading to variations in the degradation rates of anhydrous sugars
698 within the same city. Environmental factors, such as air quality and climate type, in different cities
699 further contribute to variations in degradation rates. In addition, the differential sensitivity of various
700 anhydro-saccharides to these driving factors leads to differences in the decay rates of the three
701 sugars. The GAM model results indicate that the daytime decay rate of anhydro-saccharides is
702 primarily influenced by ALWC, RH, and O_x . Additionally, increases in T and SSR also contribute
703 to an enhanced decay rate. Our findings highlight that the degradation of anhydro-saccharides in
704 real atmospheric conditions occurs through various oxidative mechanisms. Further investigation
705 into the degradation mechanisms of anhydro-saccharides is crucial for accurately assessing the
706 contribution of BB aerosols to global air quality. The results of this study provide valuable data and
707 insights for future air quality management.

708 **Data availability.** Data will be available upon request to the corresponding authors.

709 **Author contributions.** **Conceptualization:** Li Li, Yu Jianzhen, Kun Zhang; **Formal analysis:** Biao
710 Zhou, Kun Zhang, Jiqi Zhu; **Methodology and Investigation:** Biao Zhou, Kun Zhang, Qiongqiong
711 Wang; **Writing - original draft:** Biao Zhou; **Writing - review & editing:** Kun Zhang, Li Li,
712 Jianzhen Yu, Qiongqiong Wang; **Funding acquisition and Supervision:** Li Li.

713

714 **Competing interests.** The authors declare no conflicts of interest.

715 **Acknowledgements.** This study is financially supported by Jing-Jin-Ji Regional Integrated
716 Environmental Improvement - National Science and Technology Major Project (2025ZD1202005,
717 2025ZD1202004), the National Natural Science Foundation of China (42375102). This work is
718 supported by the Shanghai Technical Service Center of Science and Engineering Computing,
719 Shanghai University.

720 References

- 721 Alvi, M.U., Kistler, M., Shahid, I., Alam, K., Chishtie, F., Mahmud, T., and Kasper-Giebl, A.:
722 Composition and source apportionment of saccharides in aerosol particles from an agro-industrial zone
723 in the Indo-Gangetic Plain, *Environ. Sci. Pollut. Res.*, 27, 14124-14137,
724 <http://dx.doi.org/10.1007/s11356-020-07905-2>, 2020.
- 725 Arangio, A.M., Slade, J.H., Berkemeier, T., Pöschl, U., Knopf, D.A., and Shiraiwa, M.: Multiphase
726 Chemical Kinetics of OH Radical Uptake by Molecular Organic Markers of Biomass Burning Aerosols:
727 Humidity and Temperature Dependence, Surface Reaction, and Bulk Diffusion, *The Journal of Physical*
728 *Chemistry A*, 119, 4533-4544, <http://dx.doi.org/10.1021/jp510489z>, 2015.
- 729 Bai, J., Sun, X., Zhang, C., Xu, Y., and Qi, C.: The OH-initiated atmospheric reaction mechanism and
730 kinetics for levoglucosan emitted in biomass burning, *Chemosphere*, 93, 2004-2010,
731 <http://dx.doi.org/10.1016/j.chemosphere.2013.07.021>, 2013.
- 732 Bao, M., Zhang, Y.-L., Cao, F., Lin, Y.-C., Wang, Y., Liu, X., Zhang, W., Fan, M., Xie, F., Cary, R., Dixon,
733 J., and Zhou, L.: Highly time-resolved characterization of carbonaceous aerosols using a two-wavelength
734 Sunset thermal-optical carbon analyzer, *Atmos. Meas. Tech.*, 14, 4053-4068,
735 <http://dx.doi.org/10.5194/amt-14-4053-2021>, 2021.
- 736 Chen, G., Canonaco, F., Tobler, A., Aas, W., Alastuey, A., Allan, J., Atabakhsh, S., Aurela, M.,
737 Baltensperger, U., Bougiatioti, A., De Brito, J.F., Ceburnis, D., Chazeau, B., Chebaicheb, H.,
738 Daellenbach, K.R., Ehn, M., El Haddad, I., Eleftheriadis, K., Favez, O., Flentje, H., Font, A., Fossum,
739 K., Freney, E., Gini, M., Green, D.C., Heikkinen, L., Herrmann, H., Kalogridis, A.C., Keernik, H., Lhotka,
740 R., Lin, C., Lunder, C., Maasikmets, M., Manousakas, M.I., Marchand, N., Marin, C., Marmureanu, L.,
741 Mihalopoulos, N., Mocnik, G., Necki, J., O'Dowd, C., Ovadnevaite, J., Peter, T., Petit, J.E., Pikridas, M.,
742 Matthew Platt, S., Pokorna, P., Poulain, L., Priestman, M., Riffault, V., Rinaldi, M., Rozanski, K.,
743 Schwarz, J., Sciare, J., Simon, L., Skiba, A., Slowik, J.G., Sosedova, Y., Stavroulas, I., Styszko, K.,
744 Teinmaa, E., Timonen, H., Tremper, A., Vasilescu, J., Via, M., Vodicka, P., Wiedensohler, A., Zografou,
745 O., Cruz Minguillon, M., and Prevot, A.S.H.: European aerosol phenomenology - 8: Harmonised source
746 apportionment of organic aerosol using 22 Year-long ACSM/AMS datasets, *Environ. Int.*, 166, 107325,
747 <http://dx.doi.org/10.1016/j.envint.2022.107325>, 2022.
- 748 Chen, J., Li, C., Ristovski, Z., Milic, A., Gu, Y., Islam, M.S., Wang, S., Hao, J., Zhang, H., He, C., Guo,
749 H., Fu, H., Miljevic, B., Morawska, L., Thai, P., Lam, Y.F., Pereira, G., Ding, A., Huang, X., and Dumka,
750 U.C.: A review of biomass burning: Emissions and impacts on air quality, health and climate in China,
751 *Sci. Total Environ.*, 579, 1000-1034, <http://dx.doi.org/10.1016/j.scitotenv.2016.11.025>, 2017.
- 752 Cheng, Y., Cao, X.B., Liu, J.M., Yu, Q.Q., Zhong, Y.J., Geng, G.N., Zhang, Q., and He, K.B.: New open
753 burning policy reshaped the aerosol characteristics of agricultural fire episodes in Northeast China, *Sci.*
754 *Total Environ.*, 810, 152272, <http://dx.doi.org/10.1016/j.scitotenv.2021.152272>, 2022.
- 755 Cheng, Y., Engling, G., He, K.B., Duan, F.K., Ma, Y.L., Du, Z.Y., Liu, J.M., Zheng, M., and Weber, R.J.:
756 Biomass burning contribution to Beijing aerosol, *Atmos. Chem. Phys.*, 13, 7765-7781,
757 <http://dx.doi.org/10.5194/acp-13-7765-2013>, 2013.
- 758 Donahue, N.M., Robinson, A.L., Hartz, K.E.H., Sage, A.M., and Weitkamp, E.A.: Competitive oxidation
759 in atmospheric aerosols: The case for relative kinetics, *Geophys. Res. Lett.*, 32,
760 <http://dx.doi.org/10.1029/2005gl022893>, 2005.
- 761 Engling, G., Lee, J.J., Tsai, Y.-W., Lung, S.-C.C., Chou, C.C.K., and Chan, C.-Y.: Size-Resolved
762 Anhydrosugar Composition in Smoke Aerosol from Controlled Field Burning of Rice Straw, *Aerosol Sci.*

763 Technol., 43, 662-672, <http://dx.doi.org/10.1080/02786820902825113>, 2009.

764 Fabbri, D., Torri, C., Simoneit, B.R.T., Marynowski, L., Rushdi, A.I., and Fabiańska, M.J.: Levoglucosan
765 and other cellulose and lignin markers in emissions from burning of Miocene lignites, *Atmos. Environ.*,
766 43, 2286-2295, <http://dx.doi.org/10.1016/j.atmosenv.2009.01.030>, 2009.

767 Fang, Z., Lai, A., Cai, D.M., Li, C.L., Carmieli, R., Chen, J.M., Wang, X.M., and Rudich, Y.: Secondary
768 Organic Aerosol Generated from Biomass Burning Emitted Phenolic Compounds: Oxidative Potential,
769 Reactive Oxygen Species, and Cytotoxicity, *Environ. Sci. Technol.*, 58, 8194-8206,
770 <http://dx.doi.org/10.1021/acs.est.3c09903>, 2024.

771 Fountoukis, C. and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium
772 model for K^+ - Ca^{2+} - Mg^{2+} - NH_4^+ - Na^+ - SO_4^{2-} - NO_3^- - Cl^- - H_2O aerosols, *Atmos. Chem. Phys.*, 7, 4639-4659,
773 <http://dx.doi.org/10.5194/acp-7-4639-2007>, 2007.

774 Fu, P.Q., Kawamura, K., Chen, J., Li, J., Sun, Y.L., Liu, Y., Tachibana, E., Aggarwal, S.G., Okuzawa, K.,
775 Tanimoto, H., Kanaya, Y., and Wang, Z.F.: Diurnal variations of organic molecular tracers and stable
776 carbon isotopic composition in atmospheric aerosols over Mt. Tai in the North China Plain: an influence
777 of biomass burning, *Atmos. Chem. Phys.*, 12, 8359-8375, <http://dx.doi.org/10.5194/acp-12-8359-2012>,
778 2012.

779 Haque, M.M., Zhang, Y., Bikkina, S., Lee, M., and Kawamura, K.: Regional heterogeneities in the
780 emission of airborne primary sugar compounds and biogenic secondary organic aerosols in the East Asian
781 outflow: evidence for coal combustion as a source of levoglucosan, *Atmos. Chem. Phys.*, 22, 1373-1393,
782 <http://dx.doi.org/10.5194/acp-22-1373-2022>, 2022.

783 He, X., Wang, Q., Huang, X.H.H., Huang, D.D., Zhou, M., Qiao, L., Zhu, S., Ma, Y., Wang, H., Li, L.,
784 Huang, C., Xu, W., Worsnop, D.R., Goldstein, A.H., and Yu, J.Z.: Hourly measurements of organic
785 molecular markers in urban Shanghai, China: Observation of enhanced formation of secondary organic
786 aerosol during particulate matter episodic periods, *Atmos. Environ.*, 240,
787 <http://dx.doi.org/10.1016/j.atmosenv.2020.117807>, 2020.

788 Hennigan, C.J., Izumi, J., Sullivan, A.P., Weber, R.J., and Nenes, A.: A critical evaluation of proxy
789 methods used to estimate the acidity of atmospheric particles, *Atmos. Chem. Phys.*, 15, 2775-2790,
790 <http://dx.doi.org/10.5194/acp-15-2775-2015>, 2015.

791 Hennigan, C.J., Sullivan, A.P., Collett, J.L., and Robinson, A.L.: Levoglucosan stability in biomass
792 burning particles exposed to hydroxyl radicals, *Geophys. Res. Lett.*, 37,
793 <http://dx.doi.org/10.1029/2010gl043088>, 2010.

794 Hoffman, D., Tilgner, A., Iinuma, Y. and Herrmann, H.: Atmospheric stability of levoglucosan A detailed
795 laboratory and modeling study, *Environ. Sci. Technol.*, 44, 694-699, <http://dx.doi.org/10.1021/es902476f>,
796 2010.

797 Hong, Y., Cao, F., Fan, M.-Y., Lin, Y.-C., Gul, C., Yu, M., Wu, X., Zhai, X., and Zhang, Y.-L.: Impacts
798 of chemical degradation of levoglucosan on quantifying biomass burning contribution to carbonaceous
799 aerosols: A case study in Northeast China, *Sci. Total Environ.*, 819,
800 <http://dx.doi.org/10.1016/j.scitotenv.2021.152007>, 2022.

801 Hu, C., Wei, Z., Zhan, H., Gu, W., Liu, H., Chen, A., Jiang, B., Yue, F., Zhang, R., Fan, S., He, P., Leung,
802 K.M.Y., Wang, X., and Xie, Z.: Molecular characteristics, sources and influencing factors of isoprene
803 and monoterpenes secondary organic aerosol tracers in the marine atmosphere over the Arctic Ocean,
804 *Sci. Total Environ.*, 853, 158645, <http://dx.doi.org/10.1016/j.scitotenv.2022.158645>, 2022.

805 Huff Hartz, K.E., Weitkamp, E.A., Sage, A.M., Donahue, N.M., and Robinson, A.L.: Laboratory
806 measurements of the oxidation kinetics of organic aerosol mixtures using a relative rate constants

807 approach, *J. Geophys. Res.:Atmos.*, 112, <http://dx.doi.org/10.1029/2006jd007526>, 2007.

808 Kang, M., Ren, L., Ren, H., Zhao, Y., Kawamura, K., Zhang, H., Wei, L., Sun, Y., Wang, Z., and Fu, P.:
809 Primary biogenic and anthropogenic sources of organic aerosols in Beijing, China: Insights from
810 saccharides and n-alkanes, *Environ. Pollut.*, 243, 1579-1587,
811 <http://dx.doi.org/10.1016/j.envpol.2018.09.118>, 2018.

812 Karavoltzos, S., Sakellari, A., Bakeas, E., Bekiaris, G., Plavsic, M., Proestos, C., Zinelis, S., Koukoulakis,
813 K., Diakos, I., Dassenakis, M., and Kalogeropoulos, N.: Trace elements, polycyclic aromatic
814 hydrocarbons, mineral composition, and FT-IR characterization of unrefined sea and rock salts:
815 environmental interactions, *Environ. Sci. Pollut. Res. Int.*, 27, 10857-10868,
816 <http://dx.doi.org/10.1007/s11356-020-07670-2>, 2020.

817 Kim, N., Park, M., Yum, S.S., Park, J.S., Song, I.H., Shin, H.J., Ahn, J.Y., Kwak, K.-H., Kim, H., Bae,
818 G.-N., and Lee, G.: Hygroscopic properties of urban aerosols and their cloud condensation nuclei
819 activities measured in Seoul during the MAPS-Seoul campaign, *Atmos. Environ.*, 153, 217-232,
820 <http://dx.doi.org/10.1016/j.atmosenv.2017.01.034>, 2017.

821 Kumar, V., Rajput, P., and Goel, A.: Atmospheric abundance of HULIS during wintertime in Indo-
822 Gangetic Plain: impact of biomass burning emissions, *J. Atmos. Chem.*, 75, 385-398,
823 <http://dx.doi.org/10.1007/s10874-018-9381-4>, 2018.

824 Kuo, L.J., Louchouart, P., and Herbert, B.E.: Influence of combustion conditions on yields of solvent-
825 extractable anhydrosugars and lignin phenols in chars: implications for characterizations of biomass
826 combustion residues, *Chemosphere*, 85, 797-805, <http://dx.doi.org/10.1016/j.chemosphere.2011.06.074>,
827 2011.

828 Lai, C., Liu, Y., Ma, J., Ma, Q., and He, H.: Degradation kinetics of levoglucosan initiated by hydroxyl
829 radical under different environmental conditions, *Atmos. Environ.*, 91, 32-39,
830 <http://dx.doi.org/10.1016/j.atmosenv.2014.03.054>, 2014.

831 Lee, Y.C., Lam, Y.F., Kuhlmann, G., Wenig, M.O., Chan, K.L., Hartl, A., and Ning, Z.: An integrated
832 approach to identify the biomass burning sources contributing to black carbon episodes in Hong Kong,
833 *Atmos. Environ.*, 80, 478-487, <http://dx.doi.org/10.1016/j.atmosenv.2013.08.030>, 2013.

834 Leung, C.W., Wang, X., and Hu, D.: Characteristics and source apportionment of water-soluble organic
835 nitrogen (WSON) in PM_{2.5} in Hong Kong: With focus on amines, urea, and nitroaromatic compounds,
836 *J. Hazard. Mater.*, 469, 133899, <http://dx.doi.org/https://doi.org/10.1016/j.jhazmat.2024.133899>, 2024.

837 Li, Q., Zhang, K., Li, R., Yang, L., Yi, Y., Liu, Z., Zhang, X., Feng, J., Wang, Q., Wang, W., Huang, L.,
838 Wang, Y., Wang, S., Chen, H., Chan, A., Latif, M.T., Ooi, M.C.G., Manomaiphiboon, K., Yu, J., and Li,
839 L.: Underestimation of biomass burning contribution to PM_{2.5} due to its chemical degradation based on
840 hourly measurements of organic tracers: A case study in the Yangtze River Delta (YRD) region, China,
841 *Sci. Total Environ.*, 872, <http://dx.doi.org/10.1016/j.scitotenv.2023.162071>, 2023a.

842 Li, R., Wang, Q., He, X., Zhu, S., Zhang, K., Duan, Y., Fu, Q., Qiao, L., Wang, Y., Huang, L., Li, L., and
843 Yu, J.Z.: Source apportionment of PM_{2.5} in Shanghai based on hourly organic molecular markers and
844 other source tracers, *Atmos. Chem. Phys.*, 20, 12047-12061, [http://dx.doi.org/10.5194/acp-20-12047-](http://dx.doi.org/10.5194/acp-20-12047-2020)
845 [2020](http://dx.doi.org/10.5194/acp-20-12047-2020), 2020.

846 Li, R., Zhang, K., Li, Q., Yang, L., Wang, S., Liu, Z., Zhang, X., Chen, H., Yi, Y., Feng, J., Wang, Q.,
847 Huang, L., Wang, W., Wang, Y., Yu, J.Z., and Li, L.: Characteristics and degradation of organic aerosols
848 from cooking sources based on hourly observations of organic molecular markers in urban environments,
849 *Atmos. Chem. Phys.*, 23, 3065-3081, <http://dx.doi.org/10.5194/acp-23-3065-2023>, 2023b.

850 Li, Y., Fu, T.-M., Yu, J.Z., Feng, X., Zhang, L., Chen, J., Boreddy, S.K.R., Kawamura, K., Fu, P., Yang,

851 X., Zhu, L., and Zeng, Z.: Impacts of Chemical Degradation on the Global Budget of Atmospheric
852 Levoglucosan and Its Use As a Biomass Burning Tracer, *Environ. Sci. Technol.*, 55, 5525-5536,
853 <http://dx.doi.org/10.1021/acs.est.0c07313>, 2021.

854 Liang, L., Engling, G., Du, Z., Cheng, Y., Duan, F., Liu, X., and He, K.: Seasonal variations and source
855 estimation of saccharides in atmospheric particulate matter in Beijing, China, *Chemosphere*, 150, 365-
856 377, <http://dx.doi.org/10.1016/j.chemosphere.2016.02.002>, 2016.

857 Liu, X., Zhang, Y.-L., Peng, Y., Xu, L., Zhu, C., Cao, F., Zhai, X., Haque, M.M., Yang, C., Chang, Y.,
858 Huang, T., Xu, Z., Bao, M., Zhang, W., Fan, M., and Lee, X.: Chemical and optical properties of
859 carbonaceous aerosols in Nanjing, eastern China: regionally transported biomass burning contribution,
860 *Atmos. Chem. Phys.*, 19, 11213-11233, <http://dx.doi.org/10.5194/acp-19-11213-2019>, 2019.

861 Mochida, M., Kawamura, K., Fu, P., and Takemura, T.: Seasonal variation of levoglucosan in aerosols
862 over the western North Pacific and its assessment as a biomass-burning tracer, *Atmos. Environ.*, 44, 3511-
863 3518, <http://dx.doi.org/10.1016/j.atmosenv.2010.06.017>, 2010.

864 Pan, R., Zhu, J., Chen, D., Cheng, H., Huang, L., Wang, Y., and Li, L.: Integrated analysis of air quality-
865 vegetation-health effects of near-future air pollution control strategies, *Environ. Pollut.*, 366, 125407,
866 <http://dx.doi.org/10.1016/j.envpol.2024.125407>, 2025.

867 Pio, C.A., Legrand, M., Alves, C.A., Oliveira, T., Afonso, J., Caseiro, A., Puxbaum, H., Sanchez-Ochoa,
868 A., and Gelencsér, A.: Chemical composition of atmospheric aerosols during the 2003 summer intense
869 forest fire period, *Atmos. Environ.*, 42, 7530-7543, <http://dx.doi.org/10.1016/j.atmosenv.2008.05.032>,
870 2008.

871 Pio, C.A., Legrand, M., Oliveira, T., Afonso, J., Santos, C., Caseiro, A., Fialho, P., Barata, F., Puxbaum,
872 H., Sanchez-Ochoa, A., Kasper-Giebl, A., Gelencsér, A., Preunkert, S., and Schock, M.: Climatology of
873 aerosol composition (organic versus inorganic) at nonurban sites on a west-east transect across Europe,
874 *J. Geophys. Res.:Atmos.*, 112, <http://dx.doi.org/10.1029/2006jd008038>, 2007.

875 Riva, M., Barbosa, T.D., Lin, Y.H., Stone, E.A., Gold, A., and Surratt, J.D.: Chemical characterization of
876 organosulfates in secondary organic aerosol derived from the photooxidation of alkanes, *Atmos. Chem.*
877 *Phys.*, 16, 11001-11018, <http://dx.doi.org/10.5194/acp-16-11001-2016>, 2016a.

878 Riva, M., Budisulistiorini, S.H., Zhang, Z.F., Gold, A., and Surratt, J.D.: Chemical characterization of
879 secondary organic aerosol constituents from isoprene ozonolysis in the presence of acidic aerosol, *Atmos.*
880 *Environ.*, 130, 5-13, <http://dx.doi.org/10.1016/j.atmosenv.2015.06.027>, 2016b.

881 Sang, X., Zhang, Z., Chan, C., and Engling, G.: Source categories and contribution of biomass smoke to
882 organic aerosol over the southeastern Tibetan Plateau, *Atmos. Environ.*, 78, 113-123,
883 <http://dx.doi.org/10.1016/j.atmosenv.2012.12.012>, 2013.

884 Shrestha, N.: Detecting Multicollinearity in Regression Analysis, *American Journal of Applied*
885 *Mathematics and Statistics*, 8, 39-42, <http://dx.doi.org/10.12691/ajams-8-2-1>, 2020.

886 Slade, J.H. and Knopf, D.A.: Multiphase OH oxidation kinetics of organic aerosol The role of particle
887 phase state and relative humidity, *Geophys. Res. Lett.*, 41, 5297-5306,
888 <http://dx.doi.org/10.1002/2014GL060582>, 2014.

889 St John, P.C., Guan, Y., Kim, Y., Etz, B.D., Kim, S., and Paton, R.S.: Quantum chemical calculations for
890 over 200,000 organic radical species and 40,000 associated closed-shell molecules, *Sci Data*, 7, 244,
891 <http://dx.doi.org/10.1038/s41597-020-00588-x>, 2020a.

892 St John, P.C., Guan, Y., Kim, Y., Kim, S., and Paton, R.S.: Prediction of organic homolytic bond
893 dissociation enthalpies at near chemical accuracy with sub-second computational cost, *Nat. Commun.*,
894 11, 2328, <http://dx.doi.org/10.1038/s41467-020-16201-z>, 2020b.

895 Stevens, H., Barmuta, L.A., Chase, Z., Saunders, K.M., Zawadzki, A., Bowie, A.R., Perron, M.M.G.,
896 Sanz Rodriguez, E., Paull, B., Child, D.P., Hotchkis, M.A.C., and Proemse, B.C.: Comparing
897 levoglucosan and mannosan ratios in sediments and corresponding aerosols from recent Australian fires,
898 *Sci. Total Environ.*, 945, 174068, <http://dx.doi.org/10.1016/j.scitotenv.2024.174068>, 2024.

899 Stone, C.J.: Additive regression and other nonparametric models, *Ann. Stat.*, 13, 689-705,
900 <http://dx.doi.org/10.1214/aos/1176349548>, 1985.

901 Vicente, E.D., Vicente, A., Evtuygina, M., Carvalho, R., Tarelho, L.A.C., Oduber, F.I., and Alves, C.:
902 Particulate and gaseous emissions from charcoal combustion in barbecue grills, *Fuel Process. Technol.*,
903 176, 296-306, <http://dx.doi.org/10.1016/j.fuproc.2018.03.004>, 2018.

904 Wang, Q., He, X., Zhou, M., Huang, D.D., Qiao, L., Zhu, S., Ma, Y.-g., Wang, H.-l., Li, L., Huang, C.,
905 Huang, X.H.H., Xu, W., Worsnop, D., Goldstein, A.H., Guo, H., and Yu, J.Z.: Hourly Measurements of
906 Organic Molecular Markers in Urban Shanghai, China: Primary Organic Aerosol Source Identification
907 and Observation of Cooking Aerosol Aging, *ACS Earth Space Chem.*, 4, 1670-1685,
908 <http://dx.doi.org/10.1021/acsearthspacechem.0c00205>, 2020.

909 Wang, Q., Wang, S., Chen, H., Zhang, Z., Yu, H., Chan, M.N., and Yu, J.Z.: Ambient Measurements of
910 Daytime Decay Rates of Levoglucosan, Mannosan, and Galactosan, *J. Geophys. Res.:Atmos.*, 130,
911 <http://dx.doi.org/10.1029/2024jd042423>, 2025.

912 Wang, Q. and Yu, J.Z.: Ambient Measurements of Heterogeneous Ozone Oxidation Rates of Oleic,
913 Elaidic, and Linoleic Acid Using a Relative Rate Constant Approach in an Urban Environment, *Geophys.*
914 *Res. Lett.*, 48, <http://dx.doi.org/10.1029/2021gl095130>, 2021.

915 Wennberg, P.O.: Radicals follow the Sun, *Nature*, 442, 145-146, <http://dx.doi.org/10.1038/442145a>, 2006.

916 White, W.H.: Chemical markers for sea salt in IMPROVE aerosol data, *Atmos. Environ.*, 42, 261-274,
917 <http://dx.doi.org/10.1016/j.atmosenv.2007.09.040>, 2008.

918 Xiao, L., Yu, L., and Hao, Y.: The burden associated with ambient PM_{2.5} and meteorological factors in
919 Guangzhou, China, 2012–2016: A generalized additive modeling of temporal years of life lost,
920 *Chemosphere*, 212, 705-714, <http://dx.doi.org/10.1016/j.chemosphere.2018.08.129>, 2018.

921 Xu, S., Ren, L., Lang, Y., Hou, S., Ren, H., Wei, L., Wu, L., Deng, J., Hu, W., Pan, X., Sun, Y., Wang, Z.,
922 Su, H., Cheng, Y., and Fu, P.: Molecular markers of biomass burning and primary biological aerosols in
923 urban Beijing: size distribution and seasonal variation, *Atmos. Chem. Phys.*, 20, 3623-3644,
924 <http://dx.doi.org/10.5194/acp-20-3623-2020>, 2020.

925 Yan, C., Sullivan, A.P., Cheng, Y., Zheng, M., Zhang, Y., Zhu, T., and Collett, J.L.: Characterization of
926 saccharides and associated usage in determining biogenic and biomass burning aerosols in atmospheric
927 fine particulate matter in the North China Plain, *Sci. Total Environ.*, 650, 2939-2950,
928 <http://dx.doi.org/10.1016/j.scitotenv.2018.09.325>, 2019.

929 Yan, C., Zheng, M., Sullivan, A.P., Shen, G., Chen, Y., Wang, S., Zhao, B., Cai, S., Desyaterik, Y., Li, X.,
930 Zhou, T., Gustafsson, Ö., and Collett, J.L.: Residential Coal Combustion as a Source of Levoglucosan in
931 China, *Environ. Sci. Technol.*, 52, 1665-1674, <http://dx.doi.org/10.1021/acs.est.7b05858>, 2018.

932 Yi, Y., Li, R., Zhang, K., Yang, X., Li, Q., Geng, C., Chen, H., Yang, W., Yu, J.Z., and Li, L.: Insights
933 Into the Influence of Anthropogenic Emissions on the Formation of Secondary Organic Aerosols Based
934 on Online Measurements, *J. Geophys. Res.:Atmos.*, 129, <http://dx.doi.org/10.1029/2024jd041479>, 2024.

935 Zhai, S., Jacob, D.J., Wang, X., Shen, L., Li, K., Zhang, Y., Gui, K., Zhao, T., and Liao, H.: Fine
936 particulate matter (PM_{2.5}) trends in China, 2013–2018: separating contributions from anthropogenic
937 emissions and meteorology, *Atmos. Chem. Phys.*, 19, 11031-11041, [http://dx.doi.org/10.5194/acp-19-](http://dx.doi.org/10.5194/acp-19-11031-2019)
938 [11031-2019](http://dx.doi.org/10.5194/acp-19-11031-2019), 2019.

939 Zhang, K., Yang, L., Li, Q., Li, R., Zhang, D., Xu, W., Feng, J., Wang, Q., Wang, W., Huang, L., Yaluk,
940 E.A., Wang, Y., Yu, J.Z., and Li, L.: Hourly measurement of PM_{2.5}-bound nonpolar organic compounds
941 in Shanghai: Characteristics, sources and health risk assessment, *Sci. Total Environ.*, 789,
942 <http://dx.doi.org/10.1016/j.scitotenv.2021.148070>, 2021a.

943 Zhang, L., Hu, B., Liu, X.L., Luo, Z.H., Xing, R., Li, Y.J., Xiong, R., Li, G., Cheng, H.F., Lu, Q., Shen,
944 G.F., and Tao, S.: Variabilities in Primary N-Containing Aromatic Compound Emissions from Residential
945 Solid Fuel Combustion and Implications for Source Tracers, *Environ. Sci. Technol.*,
946 <http://dx.doi.org/10.1021/acs.est.2c03000>, 2022.

947 Zhang, L., Zhou, P., Zhong, H., Zhao, Y., Dai, L., Wang, Q.g., Xi, M., Lu, Y., and Wang, Y.: Quantifying
948 the impacts of anthropogenic and natural perturbations on gaseous elemental mercury (GEM) at a
949 suburban site in eastern China using generalized additive models, *Atmos. Environ.*, 247,
950 <http://dx.doi.org/10.1016/j.atmosenv.2020.118181>, 2021b.

951 Zhang, Q., Jimenez, J.L., Canagaratna, M.R., Allan, J.D., Coe, H., Ulbrich, I., Alfarra, M.R., Takami, A.,
952 Middlebrook, A.M., Sun, Y.L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P.F., Salcedo, D., Onasch,
953 T., Jayne, J.T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J.,
954 Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell,
955 L., Griffin, R.J., Rautiainen, J., Sun, J.Y., Zhang, Y.M., and Worsnop, D.R.: Ubiquity and dominance of
956 oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere
957 midlatitudes, *Geophys. Res. Lett.*, 34, <http://dx.doi.org/10.1029/2007gl029979>, 2007.

958 Zhang, R., Jing, J., Tao, J., Hsu, S.C., Wang, G., Cao, J., Lee, C.S.L., Zhu, L., Chen, Z., Zhao, Y., and
959 Shen, Z.: Chemical characterization and source apportionment of PM_{2.5} in Beijing: seasonal perspective,
960 *Atmos. Chem. Phys.*, 13, 7053-7074, <http://dx.doi.org/10.5194/acp-13-7053-2013>, 2013.

961 Zhao, R., Mungall, E.L., Lee, A.K.Y., Aljawhary, D., and Abbatt, J.P.D.: Aqueous-phase photooxidation
962 of levoglucosan - a mechanistic study using aerosol time-of-flight chemical ionization mass spectrometry
963 (Aerosol ToF-CIMS), *Atmos. Chem. Phys.*, 14, 9695-9706, <http://dx.doi.org/10.5194/acp-14-9695-2014>,
964 2014.

965 Zhao, Y., Kreisberg, N.M., Worton, D.R., Isaacman, G., Weber, R.J., Liu, S., Day, D.A., Russell, L.M.,
966 Markovic, M.Z., VandenBoer, T.C., Murphy, J.G., Hering, S.V., and Goldstein, A.H.: Insights into
967 Secondary Organic Aerosol Formation Mechanisms from Measured Gas/Particle Partitioning of Specific
968 Organic Tracer Compounds, *Environ. Sci. Technol.*, 47, 3781-3787, <http://dx.doi.org/10.1021/es304587x>,
969 2013.

970 Zhu, S., Wang, Q., Qiao, L., Zhou, M., Wang, S., Lou, S., Huang, D., Wang, Q., Jing, S., Wang, H., Chen,
971 C., Huang, C., and Yu, J.Z.: Tracer-based characterization of source variations of PM_{2.5} and organic
972 carbon in Shanghai influenced by the COVID-19 lockdown, *Faraday Discuss.*, 226, 112-137,
973 <http://dx.doi.org/10.1039/d0fd00091d>, 2021.

974