

Process-Oriented Evaluation of ~~Stationary~~-Quasi-Stationary Rossby Waves and Their Impact on Surface Air Temperature Extremes in Dynamical Downscaling over North America

Koichi Sakaguchi¹, Seth A. McGinnis², L. Ruby Leung¹, Melissa S. Bukovsky³, Rachel R. McCrary², Ziming Chen¹, Chuan-Chieh Chang¹, and Yanjie Li⁴

¹Pacific Northwest National Laboratory, Richland, Washington, USA

²NSF National Center for Atmospheric Research, Boulder, Colorado, USA

³University of Wyoming, Laramie, WY, USA

⁴Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

Correspondence: Koichi Sakaguchi (Koichi.Sakaguchi@pnnl.gov)

Abstract. ~~Stationary~~-Quasi-stationary Rossby waves are a crucial component of the general circulation and play a significant role in regional water and energy cycles, as well as in extreme events. However, process-oriented evaluation for ~~stationary~~ Rossby waves is rarely performed for dynamical downscaling simulations. To close this gap, we evaluate three classes of dynamical downscaling approaches, with a focus on ~~stationary~~-quasi-stationary Rossby waves and their impact on surface air temperature over North America during Northern Hemisphere summer. The three classes of models differ in the way large-scale forcing is provided: a limited-area model (LAM) constrained only by lateral boundary conditions, represented by RegCM4 from the North American branch of the Coordinated Regional Downscaling Experiment (NA-CORDEX), a LAM with spectral nudging to maintain consistency in large-scale dynamics with the forcing data, represented by the Weather Research and Forecasting (WRF) model simulation in NA-CORDEX, and a global variable-resolution model with smoothly varying grid spacings, represented by the Community Atmosphere Model version 5.4, with the Model for Prediction Across Scales (MPAS) as its dynamical core (CAM-MPAS). With no constraints on the atmospheric dynamics, CAM-MPAS exhibits several mean biases in the upper-level circulations over the Pacific Coast region: a weaker subtropical jet, a northward-shifted mid-latitude jet, and an overestimated southerly flow. With the lateral boundary constraint alone, RegCM4 also exhibits weaker jets and overestimated southerly winds off the West Coast. Rossby ray theory reveals that those wind biases direct incoming Rossby waves northward. The erroneously routed Rossby waves distort the relationship between the accumulation of wave activity over the US West Coast and surface temperature anomalies over the Southern Great Plains, which emerges approximately four days after the convergence of ~~wave-activity~~-wave-activity flux in the ERA-Interim reanalysis. Furthermore, the response of heatwaves to the extreme wave activity flux is not reproduced by the two models, a serious drawback as a dynamical downscaling framework is expected to connect large-scale forcing to local-scale phenomena. The WRF model employing spectral nudging is largely free from the aforementioned problems. A pair of sensitivity simulations suggests that spectral nudging is the key to improving the dynamics of ~~stationary~~-quasi-stationary Rossby waves and their impact on surface air temperature. Our results also demonstrate the effectiveness of Rossby wave diagnostics that allow for realistic background

flows for assessing the credibility of dynamical downscaling over North America, where incoming Rossby waves propagate through complex circulation patterns before traveling across the continent. ~~Evaluating such process chains — from large-scale Rossby waves to local-scale extreme events — requires accounting for the region’s unique dynamical features.~~

1 Introduction

Rossby waves have the largest spatial scales among the atmospheric waves (1000s–10000s km). Their spatial extent makes it possible to connect tropical convection to mid-latitude weather (Wallace and Gutzler, 1981; Ambrizzi et al., 1995; Branstator, 2014). Rossby waves can be “quasi-stationary” by having a phase speed nearly equal to the background winds but in the opposite direction, thus their phase (maxima and minima) becomes fixed in space. ~~Since phase speed is inversely proportional to wavenumber, larger waves tend to have greater phase speed.~~ Some large waves become ~~stationary~~ quasi-stationary even within the atmospheric jet streams, where vorticity gradients and strong winds can trap and help the waves travel further (Manola et al., 2013; Branstator and Teng, 2017; Wirth, 2020). Such large, (quasi-)stationary Rossby waves are one of the important drivers for regional climate because their associated momentum and energy fluxes modify regional circulation and atmospheric stability (e.g., Weaver and Nigam, 2008; Hoskins and Woollings, 2015; Teng and Branstator, 2017, 2019; Wills et al., 2019; White et al., 2022). Rigorous evaluations of simulated Rossby waves are thus necessary for establishing confidence in regional climate projections. To this end, this study revisits and evaluates the large-scale circulations relevant to Rossby wave propagation to North America, as well as the physical connection between ~~stationary~~ quasi-stationary Rossby waves and regional climate, specifically near-surface air temperature (*tas*).

The heatwave over the Pacific Northwest (PNW) in July 2009 is a good example of a relationship between ~~stationary~~ quasi-stationary Rossby waves and *tas* anomaly. This event marked the highest maximum temperature in the record across the region (Bumbaco et al., 2013), until it was exceeded by a more recent heatwave in 2021 (White et al., 2023), which falls outside our study period. A spatiotemporal correlation between the upper-level geopotential height anomaly and the daily *tas* anomaly is evident during this month (Fig. 1d-i). Fig. 1a-c illustrate the flux of wave activity (WA), second-order variability of wind fields associated with Rossby waves (Takaya and Nakamura, 2001) (hereafter TN01). The WA flux delineates the flux of perturbation geopotential height in the direction of the group velocity, which is also associated with a negative momentum transport for the mean circulation (Takaya and Nakamura, 2001, section 4). In other words, ahead of the WA convergence, one sees an increase in the perturbation geopotential height and a reduction in mean wind speeds. The region behind the WA divergence experiences a decrease in perturbation geopotential height and an acceleration of the mean winds.

About two weeks before the most intense heatwave on July 29, the PNW region was under a weakly negative height anomaly (Fig. 1d), but the WA flux had already started converging over the region (Fig. 1a). The flux is dominantly meridionally oriented, flowing out northward from the subtropical eastern Pacific, where intense wave activity flux has been converging from the mid-latitude North Pacific. Some WA flux appears to originate from the tropical east Pacific region as well. ~~This southerly flow of WA is not obvious from the geopotential height pattern.~~ The WA flux convergence continued and became more intense over the next ten days, during which a positive geopotential anomaly built up over the PNW region (Fig. 1b,e). In response, a positive

60 *tas* anomaly has emerged (Fig. 1h). The WA flux convergence over the PNW continued, spreading the positive geopotential anomaly northward to cover Washington state in the United States and the entire Canadian West coast by July 29 (Fig. 1c,f), when a positive *tas* anomaly $> 6^{\circ}\text{C}$ has extended over most of the PNW region (Fig. 1i). The effect of WA flux divergence through geopotential changes to *tas* appears to take approximately six days based on the lead/lag correlation. Figure A1a shows that the linear correlation reaches a maximum value of 0.56 at a [negative lag of six days applied to the WA flux divergence](#). The evolution of the upper-level geopotential height anomalies follows the typical condition during heatwave events over the region, with the high anomaly centered over Vancouver Island near the Canada-US border (Fig. 1e,f) (Bumbaco et al., 2013). This circulation structure is a part of the East Pacific-North Pacific pattern that is characterized by a southward-shifted and more intense jet across the Pacific (Bell and Janowiak, 1995).




Figure 1. Evolution of [stationary-quasi-stationary](#) Rossby waves and surface air temperature (*tas*) during the 2009 heatwave event: (a-c) the flux (arrows) and divergence (color, blue = convergence, green = divergence) of daily-mean wave activity (WA) flux of derived from the 25-90 day band-passed geopotential height anomalies, (d-f) 200-hPa winds and geopotential height anomalies (25-90 day band-passed), and (g-i) daily *tas* anomaly, all variables from ERA-Interim.

65 The 2009 heatwave is just one example; significant connections between ~~stationary~~ quasi-stationary Rossby waves and regional climate and ~~extremes have been suggested for a long time~~ extreme events have long been suggested. Analyzing 30 years of reanalysis data, Schubert et al. (2011) found that meridional wind variabilities associated with stationary Rossby waves account for up to 60 % of surface temperature variabilities over large areas in North America. Teng et al. (2013) found that stationary waves with zonal wavenumber-5 patterns often appear 15-20 days before heatwave events in the United States
70 in 12,000 years of atmospheric general circulation model (GCM) simulations. Yuan et al. (2015) investigated the variability and trend of subtropical stationary waves during the NH summer. They found an increasing trend in wave amplitude over the 1979-2013 period, as well as changes in regional moisture fluxes ~~accompanying stationary waves, which affected hydroclimate~~ over-associated with stationary waves that affected hydroclimate across several regions, including the central United States. In ~~the last recent~~ decades, an increasing number of studies have investigated how quasi-stationary waves contribute to extreme
75 events (Coumou et al., 2014; Hoskins and Woollings, 2015; Kornhuber et al., 2017; Wolf et al., 2018).

Due to the significance of (quasi-)stationary waves on regional climate, several recent studies have evaluated Rossby waves in GCM simulations and further found connections between the model's skills in simulating Rossby waves and in simulating the surface climate (e.g., Holman et al., 2014; Luo et al., 2022). For example, Simpson et al. (2020) used standard performance metrics, such as spatial correlation and root-mean-square errors of the time-mean eddy streamfunction, to evaluate stationary
80 Rossby waves in two generations of model ensembles from the Coupled Model Intercomparison Project (CMIP) and a large ensemble of a single model. They found improved performance from the CMIP phase 5 (CMIP5) to CMIP6, and the model biases tend to be larger in JJA than in DJF. Other studies used metrics ~~developed based on~~ derived from linear wave theory and the vorticity budget to evaluate simulated Rossby waves. Nie et al. (2019) evaluated Rossby wave sources in the CMIP5 models, and Henderson et al. (2017) evaluated the teleconnection between North America and Madden-Julian Oscillation using the so-
85 called stationary wavenumbers. Some studies have taken a step further to use more complex diagnostics of Rossby waves, such as the WA flux and Rossby wave ray tracing, to find close connections between near-surface climate and Rossby wave propagation biases in GCMs (Garfinkel et al., 2022; Choi and Stan, 2024). However, few studies have evaluated large-scale stationary Rossby waves in regional, dynamical downscaling simulations.

For limited-area models (LAMs), previous studies have focused on atmospheric circulations with ~~smaller spatiotemporal~~ scales than stationary spatiotemporal scales smaller than those of quasi-stationary Rossby waves. Using the “Big-Brother Experiment” in which a smaller domain simulation is forced by the output from the larger-domain simulation using the same model, Denis et al. (2002b), Denis et al. (2003), and Dimitrijevic and Laprise (2005) evaluated the simulated atmospheric circulations on a monthly time scale. These studies found that lateral boundaries (LBs) do not significantly affect modeled sea-level pressure and relative vorticity; however, the vorticity fields exhibit some deviations from the driving model at higher
95 atmospheric levels. Using a similar experimental design but with an idealized dry test case, Park et al. (2014) found unphysical inertia-gravity waves excited at the LBs. The artificial waves become stronger with longer LB update time periods, particularly when they are substantially longer than the LAM timestep, which is usually the case in climate-scale model integration. Miguez-Macho et al. (2004) documented how the interactions between the simulated flow and specified flows at LBs distort large-scale circulations in regional simulations over North America, and also demonstrated the usefulness of spectral nudging

100 for the waves with synoptic and larger scales to remove the large-scale flow biases. Imberger et al. (2020) investigated the impact of the LB update frequency, size of the LB relaxation zone, and spectral nudging in the case study of a fast-propagating, strong mid-latitude storm. They found that the update frequency is most effective in mitigating reductions in storm intensity through LBs. Castro et al. (2007) and Chang et al. (2015) investigated how the modes of large-scale climate variabilities via Rossby waves are simulated in regional downscaling by Empirical Orthogonal Functions, focusing on the teleconnections
105 between tropical sea surface temperature (SST) and the North American Monsoon. They found that spectral nudging helps reproduce large-scale climate variabilities, but the dynamics and kinematics of Rossby waves were not their focus. Scarcity of Rossby wave evaluation in regional simulations may be related to an assumption that the large spatiotemporal scales of ~~stationary~~quasi-stationary Rossby waves are well resolved by the host GCM grid and sub-daily (e.g., six-hourly) frequency updates of LB conditions. However, this assumption is not necessarily valid.

110 A common numerical treatment of LB conditions is to blend the specified forcing with the state simulated by LAMs (e.g., Davies, 1976). Staniforth (1997) noted that such blending does not retain the balance within the flow, such as geostrophy. Deviation from the geostrophic balance excites inertia-gravity waves to restore the balance (Holton, 2004). The excitation of inertia-gravity waves would bring the state closer to geostrophic balance, but the LB treatment occurs at every time step; thus, the vicinity of the boundaries may always experience artificial imbalance. Such a disruption would distort the propagation of
115 incoming Rossby waves, and the persistent divergence produced by the unphysical inertia-gravity waves (Park et al., 2014) may also contaminate the amplitude of the incoming Rossby waves.

Another modeling framework for dynamical downscaling is global variable-resolution (VR) models. One such model, the Model for Prediction Across Scales (MPAS, Skamarock et al. (2012)), is developed on an unstructured grid that can smoothly change grid spacing over a specified region. This model has been shown not to have the aforementioned issues associated with
120 LBs (Park et al., 2014). However, the amplitude, pathways, and frequency of Rossby waves arriving in North America may be unrealistic. This is because for those waves originating from the tropics, the strength and spatial scales of the wave source are linked to the amplitude and profile of diabatic heating in the organized tropical convection, which is known to be difficult for GCMs to realistically simulate (~~Dai, 2006; Baemeister et al., 2014b; Bogenschutz et al., 2018; Park and Lee, 2021; Zhou et al., 2022; Chang et al., 2025a~~) (Dai, 2006; Baemeister et al., 2014a; Bogenschutz et al., 2018; Park and Lee, 2021; Zhou et al., 2022; Chang et al., 2025a). Fur-
125 thermore, GCMs have long-standing biases in the location and strength of the jet (Harvey et al., 2020; Simpson et al., 2020). For dynamical downscaling using LAMs, one can choose host GCMs with small biases in those aspects. For dynamical downscaling with a global VR model, the model must exhibit good skills in both global-scale and regional-scale processes.

There is thus a clear need to evaluate ~~stationary~~quasi-stationary Rossby waves in dynamical downscaling simulations; however, a process-oriented evaluation has not been ~~carried out to assess Rossby waves simulated by~~conducted to assess how
130 different modeling frameworks simulate them. To fill this gap, we evaluate three classes of dynamical downscaling approaches that have distinct representations of large-scale forcing. The first class is a standard regional climate simulation with a LAM, represented by the Regional Climate Model version 4 (RegCM4) simulation available from the North American branch of the Coordinated Regional Downscaling Experiment (Mearns et al., 2017) (NA-CORDEX). The second class is also ~~a~~an LAM simulation, but ~~employing~~employs spectral nudging to constrain ~~the~~ large-scale atmospheric dynamics; the WRF simulation

135 in NA-CORDEX is one such dataset. The third class is a global VR model that utilizes the MPAS dynamical core within the Community Atmosphere Model (CAM), referred to as CAM-MPAS. This model’s regional refinement and simulation design follow the NA-CORDEX protocol (Sakaguchi et al., 2023). We will demonstrate that these three classes of models exhibit distinct biases in the upper-level circulations and Rossby wave propagations. We also provide reviews and technical details of the diagnostics throughout the text and in the appendices for those interested in more background on Rossby wave theory.

140 2 Data and Methods

2.1 Downscaling and evaluation dataset

We use two simulations from the “Evaluation” experiment in NA-CORDEX (Diez-Sierra et al., 2022b), one using the RegCM4 model and the other using the WRF model. Both models are configured on 25-km grids following the NA-CORDEX protocol (Fig. 2b) (Diez-Sierra et al., 2022a). We also analyze another downscaling simulation conducted with the CAM-MPAS model
145 on a global VR grid with a 100-km coarse domain refined smoothly to 25-km grid spacing over North America (Fig. 2a).

The RegCM model is a widely used regional climate model with a long history (Giorgi and Anyah, 2012). Downscaled data from the fourth-generation RegCM4 are available from NA-CORDEX on both the 50 km and 25 km grids (Mearns et al., 2017; Bukovsky and Mearns, 2020; McGinnis and Mearns, 2021). This model version solves the primitive (hydrostatic) equations on a σ coordinate as described in Grell et al. (1994) and Elguindi et al. (2017). Multiple options are available for
150 the cumulus, ~~boundary-layer, and land-surface~~ boundary layer, and land-surface components (Giorgi et al., 2012). The physics parameterizations were selected based on the performance of test simulations over the CONUS region, particularly for warm-season precipitation (Arritt and Bukovsky, personal communication).

The WRF model is a regional model for weather and climate applications (Skamarock et al., 2008) and has been extensively used to study the present-day and future state of North American climate with a wide range of model resolutions and configurations (e.g., Chang et al., 2015; Liu et al., 2017; Chen et al., 2019; Srivastava et al., 2023). Version 3.5.1 was used for the
155 NA-CORDEX experiment (50 km and 25 km). The ~~model physics mostly follows those adopted by dynamical core solves the Euler equations without the hydrostatic assumption. The model physics largely follows that of~~ Castro et al. (2012), who focused on the warm-season climate of the western CONUS and the North American monsoon. Spectral nudging is applied to the temperature, winds, and geopotential height fields at the scales larger than approximately 1000 km to retain synoptic-scale
160 variability in the driving GCM or reanalysis data (Castro et al., 2005, 2012; Hu et al., 2018).

CAM-MPAS is an experimental model in which the dynamical core is ported from the MPAS-Atmosphere version 4 to the CAM model within a beta version of the CESM2. The technical description of the model and downscaling experiments are provided in Sakaguchi et al. (2023). Briefly, MPAS is a global dynamical core that solves the ~~fully compressible non-hydrostatic equations-of-motion~~ Euler equations on an unstructured grid (Skamarock et al., 2012). The unstructured grid can be configured
165 as a global quasi-uniform resolution grid or a VR grid ~~where-, in which~~ one or more regions of interest have finer grid spacing than the rest of the globe. Advantages of the global VR model over LAMs include the absence of LBs and the consistent

dynamical and physical schemes in ~~the downscaling (both the high-resolution) domain and the (downscaling) and~~ coarse-resolution ~~domain~~domains, which can avoid artificial shocks or gradients created by ~~the~~LBs in LAMs.

170 All models use the ERA-Interim reanalysis product for initial and boundary conditions, including six-hourly updates to the LBs and daily updates to SST and sea ice fraction (SIC) at the bottom (surface) boundary. RegCM4 does not have a specific sea ice scheme, so SIC from ERA-Interim is not used. CAM-MPAS uses SST and SIC only since there are no lateral boundaries; therefore, the large-scale circulations are not constrained. Table 1 lists the model characteristics and configurations. Table B1 compares the physics parameterizations used by the three models.




Figure 2. Model mesh examples: (a) global variable-resolution mesh for CAM-MPAS, (b) regional mesh for WRF. The mesh used by RegCM4 is visually similar to that of WRF (hence not shown), except it covers a slightly larger area.

The reference data we use is ERA-Interim, which drives the NA-CORDEX simulations for the “Evaluation” experiment. As
175 discussed by Laprise et al. (2008), we expect that dynamical downscaling adds value primarily in the small-scale processes while maintaining the large-scale flow provided from the driving data. If this tenet is true, incoming Rossby wave signals are not affected by LBs or other model details (see Appendix B1); Rossby wave metrics calculated from the driving data (ERA-Interim) and from downscaling simulations within the LAM domain should be very close to each other. On the other hand, if numerical aspects of the downscaling model affect the circulations, such as artificial sources of divergence over the
180 time scale of ~~stationary~~quasi-stationary Rossby waves, or the model exhibits mean biases in the general circulations (e.g., jet strength/width/positions), then we would see deviations in the Rossby waves between the driving data and the downscaling simulations. We are aware that this logic ignores a potential upscale effect from the downscaling simulation on the ~~stationary~~quasi-stationary Rossby waves. We will briefly discuss this assumption in Section 4; however, such upscaling signals cannot be easily quantified without a priori designed experiments (e.g., Denis et al., 2002b; Leung et al., 2013; Sakaguchi et al., 2016),
185 and this is left for future work.

Table 1. Characteristics of the three downscaling models. The sponge zone width in CAM-MPAS refers to the transition zone. All models solve compressible mass and momentum equations. SST: sea surface temperature, SIC: sea ice fraction.

Configuration	RegCM4	WRF	CAM-MPAS
	25 km	25 km	25 km
Model domain	Regional	Regional	Global
Horizontal grid	Cartesian, B-grid	Cartesian, C-grid	Unstructured, C-grid
Number of grid columns	123,825	96,036	137,218
Vertical grid	Sigma	Terrain-following hydrostatic pressure	Terrain-following height
Vertical levels	18	28	32
Domain top (hPa)	50	50	2
W momentum eqn.	Hydrostatic	Non-hydrostatic	Non-hydrostatic
Time step (s)	50	150	600 (Physics) 85 (dynamics)
Driving ocean BC variables	SST	SST,SIC	SST,SIC
Spectral nudging	No	Yes	No
Lateral boundary treatment	Nudging with exponential weights	Linear relaxation	Smoothly varying Δx
Horizontal sponge zone width	24 (grid points)	10 (grid points)	≈ 25 (degrees)

2.2 Data preparation

The NH summer season (JJA) in the 30-year period from 1980 to 2010 is analyzed, except for the CAM-MPAS simulation, which starts at 1990. Most analyses are performed using daily-mean variables at 200 hPa (zonal and meridional winds, geopotential height). This particular pressure level is chosen primarily because it is a standard pressure level available in the CORDEX archives (CORDEX, 2009). Note that grid-According to the CORDEX protocol’s model data requirements, daily mean quantities are calculated from three-hourly data (CORDEX, 2009). The CAM-MPAS data follow this requirement. The ERA-Interim data is available only every six hours, from which we calculated daily statistics. We compared the seasonal means and standard deviations of the monthly mean *tas* calculated from the six-hourly and three-hourly data, and found that the differences in these statistics are significantly smaller (less than 10 %) than the model biases against ERA-Interim (not shown).

Grid boxes adjacent to the LBs, or “sponge/buffer/relaxation zone”, where the external forcing and model-predicted variables are blended (Table 1, Appendix B1), have already been removed in the NA-CORDEX data (Table 1). This post-processing is designed for the usual-common use case of regional climate assessment within the model domain; for this study, it poses a challenge. This is because we *patch* the outside of the LAM domain with ERA-Interim data to produce spatially continuous fields, on which Rossby wave propagations are diagnosed. Without the relaxation zone that blends LAMs predictions and

ERA-Interim data, our patched diagnostic approach exhibits stronger gradients between the model and ERA-Interim data than with the relaxation zone. We used a brief WRF simulation to evaluate the impact of removing the buffer zone, which was found not to significantly alter the analysis results within the model domain (Fig. B2). However, within the blending zone, the strength and spatial pattern of derived quantities (e.g., vorticity, divergence, and WA fluxes) change, and overall, they are notably noisier without the blending zone (Fig. B2a,b,d,e). The noise and spurious WA fluxes can be reduced to some extent by spatial smoothing applied over the relaxation zone (Fig. B2c, f). We tested several smoothing methods and present the figures that utilized a Gaussian filter within the buffer zone when the noise is significant. We do not attempt to evaluate Rossby wave sources/sinks along the LBs; those crucial aspects will be assessed in future work.

Prior to the patched analyses, all the data are regridded to a global 0.7° latitude-longitude grid using the *patch* method available from the Earth System Modeling Framework (ESMF) library (Balaji et al., 2018). The 0.7° grid is nearly identical to the original ERA-Interim grid and is coarser than the downscaling datasets. We still remap the ERA-Interim data to this grid to more fairly compare variability and extremes with the models, since remapping can smooth the fields and affect those statistics (Sakaguchi et al., 2023). The patch method first estimates the grid corner values on the source grid using the second-order polynomials, then weight-averages the corner values to obtain the final estimate on a target point value on the destination grid; therefore, the computation is more expensive than the commonly used bilinear method (Zienkiewicz and Zhu, 1995). The patch method estimates the values and their derivatives more accurately than the bilinear method (Balaji et al., 2018), which is desirable for calculating Rossby wave diagnostics that involve spatial derivatives.

It is critical to rotate the grid-relative ~~u and v~~ u and v winds to the Earth-relative (eastward and northward) winds in the RegCM4 and WRF data before regridding. For the WRF model, the NCAR Command Language (NCL: NCAR (2017)) provides a function for wind rotation (`wrf_uvmet`). For RegCM4, we wrote an NCL function to rotate winds onto the Rotated Mercator projection, which is available in Sakaguchi (2025). It is often necessary to spatially smooth the variables, especially for winds at relatively high resolution. In most cases, we used a suite of spherical harmonic functions available in NCL: `vhaeC`, `tri_trunC`, and `shaeC`.

2.3 Diagnostic framework

2.3.1 Rossby wave ray theory

Wirth et al. (2018) reviewed diagnostics to study the dynamics of Rossby waves, particularly the frameworks to identify and track so-called Rossby wave packets. Wave energy, momentum, and other information propagate with the wave packets at the group velocity, not with individual wave crests/troughs (Vallis, 2017, Chapter 6). One ~~of the commonly used diagnostics is~~ the commonly used diagnostic is ray theory, which traces the trajectory of a wave packet from a specified source location. The potential or absolute vorticity equations are linearized by decomposing the variables into the base state (or background or reference ~~)-state~~), which does not vary in time during the lifetime of the wave packet, and the perturbation from the base state (wave motions). Assuming a wave-like solution to the linearized equation and scale separation between the wave motion (small) and base state (large), we can obtain an algebraic relationship among the wave frequency, wavenumbers,

and base states (the *wave dispersion relationship*) ~~-, which leads to at a given location.~~ Further assuming that the base state varies much more slowly than the waves do, we can get a set of ordinary differential equations for the time evolution of the wavenumbers and frequency. ~~Knowing this kinematic information at $t = 0$ and the base state~~ Combining these kinematic and dynamic relationships, we can ~~calculate the group velocity and~~ predict where the wave packets will travel ~~at from the source at $t = 0$ to another location at $t = 1$.~~ At the new location, we solve for the wavenumbers again ~~with the new environmental conditions~~, yielding a new group velocity. Repeating the process gives us the evolution of wavenumbers and group velocities across space and time (Li et al., 2015). Vallis (2017) provides a general introduction to ray theory for Rossby waves.

Hoskins and Karoly (1981) first applied the ray theory ~~to stationary~~ (Whitham, 1960) to quasi-stationary Rossby waves. Their ray theory assumes that the meridional wind in the base state (\bar{v}) is zero, and the zonal wind (\bar{u}) is a function of latitude only. Despite these strong assumptions, their result reproduced many aspects of the Rossby wave propagations inferred from statistical analyses and numerical model results. However, this assumption is ~~hard-difficult~~ to justify given our focus on regional climate over North America, as seen in the 2009 heatwave example in the Introduction ~~section.~~ ~~Therefore, we adapt~~ Karoly (1983) applied the ray theory ~~developed by Li et al. (2015) and Zhao et al. (2015) (hereafter LZ2015).~~ Unlike earlier Rossby wave ray-tracing methods, LZ2015 allows to a base state that varies in both the zonal and meridional directions, with non-zero \bar{u} and \bar{v} . ~~Their work was extended by Li et al. (2015) and Zhao et al. (2015) (hereafter LZ2015), which is adopted in our analysis.~~ The input for the LZ2015 ray theory consists of the ~~background winds $\bar{u}(x, y)$ and $\bar{v}(x, y)$,~~ the wave source location, ~~and~~ the initial zonal wavenumber k_0 , ~~and the background winds in the Mercator projection $(\bar{u}_M, \bar{v}_M) = (\bar{u}, \bar{v}) / \cos(\varphi)$ where φ is latitude.~~ Given those inputs, LZ2015 solves the following equations \div

(eqns. 11 and 12 in Li et al. (2015)):

$$\frac{d_g k}{dT} = -k \frac{\partial \bar{u}}{\partial X} \frac{\partial \bar{u}_M}{\partial X} - l \frac{\partial \bar{v}}{\partial X} \frac{\partial \bar{v}_M}{\partial X} - \frac{1}{K^2} \left(l \frac{\partial \bar{\eta}_x}{\partial X} - k \frac{\partial \bar{\eta}_y}{\partial X} \right) \quad (1)$$

$$\frac{d_g l}{dT} = -k \frac{\partial \bar{u}}{\partial Y} \frac{\partial \bar{u}_M}{\partial Y} - l \frac{\partial \bar{v}}{\partial Y} \frac{\partial \bar{v}_M}{\partial Y} - \frac{1}{K^2} \left(l \frac{\partial \bar{\eta}_x}{\partial Y} - k \frac{\partial \bar{\eta}_y}{\partial Y} \right) \quad (2)$$

where ~~k and l is the meridional wavenumber are the zonal and meridional wavenumbers~~ (m^{-1}), $K = \sqrt{k^2 + l^2}$ is the total wavenumber, $\bar{\eta}$ denotes the background absolute vorticity (s^{-1}), $\bar{\eta} = \bar{\zeta} + f$, f is the Coriolis parameter, and $\bar{\zeta}$ is the vertical component of the background relative vorticity. The ~~coordinate variables T , X , and Y are the time, zonal, and meridional coordinates for the mean state that has substantially larger scales than a local, wave-scale motion (t, x, y) .~~ Here, $x = a\lambda$ and $y = \ln[(1 + \sin \varphi) / \cos \varphi]$ in the Mercator projection. The operators $d_g/dT = \partial/\partial T + u_g \partial/\partial X + v_g \partial/\partial Y$ represent the total derivative describing the rate of change following the wave packet moving at the group velocity ~~(the subscript g denotes group, not geostrophic flow).~~ Expressions for the group velocity are given in the Appendix C. ~~The coordinate variables T , X , and Y are the zonal, meridional, and time coordinates for a local, wave-scale motion that has substantially smaller scales than those for the mean state (t, x, y) ; this scale separation allows us to treat wavenumbers as constant within the local coordinate (making eqns. 1 and 2 linear).~~ To maintain consistency with the assumed scale separation, climatological mean fields are smoothed by truncating wavenumbers ~~larger-greater~~ than 10 after spectral decomposition ~~using-in~~ spherical harmonics, before being passed

to the ~~ray tracing algorithm as the background state. The frequency must be specified in the ray tracing code, and ray tracing~~ algorithm. Also, to be consistent with our focus on quasi-stationary Rossby waves, the time frequency is set to zero for our analysis, ~~so that the ray theory solves the trajectories of stationary Rossby waves~~ (see eqns. C9, C10).

270 After running the ray tracing algorithm, we can visually compare wave ray trajectories in the background state from ERA-Interim and those from the model simulations. To make model evaluation more quantitative than relying on visual inspection of rays, we compare the probabilities of Rossby wave propagation ~~over at~~ each grid point, obtained by tracing a large number of rays. For example, if 5,000 Rossby wave rays are initiated in a source region with slightly different input parameters, and 50 of them pass a grid box, then the probability of 0.01 is assigned to the grid box. To create an ensemble ray tracing, we start rays
 275 every two grid boxes within a source region (≈ 20 to 40 degrees wide in the x and y directions), ~~which results in \approx~~ resulting in approx 150 to 300 source points per region. For each source point, we initiate Rossby waves with 12 different k_0 (1 – 12). We also consider three background states: the climatological winds for June, July, and August. Permuting the source points, 12 initial zonal wavenumbers, and three base states yields 5,000 to 10,000 ray trajectories from each source region.

We note that this is a rather arbitrary approach to creating an ensemble of Rossby rays, since our base state and choice of k_0
 280 may ignore important characteristics of Rossby waves in a particular region or time period. For instance, the preferred wavelengths of ~~stationary quasi-stationary~~ waves excited over the Indian Monsoon region and the Tibetan Plateau appear to differ [wavenumbers 6-7 for the former and 4-5 for the latter (Joseph and Srinivasan, 1999; Park et al., 2013)]. To more accurately quantify the probabilities of ray trajectories beyond model evaluations, one may consider a broader range of parameter space (Li et al., 2019) and/or specify the parameter ranges based on a priori knowledge of the wave sources and time period of interest
 285 (Garfinkel et al., 2022; Chang et al., 2023a). Here, our tenet is that, given the same set of parameters and specifications for the base state, dynamical downscaling models can reproduce the probability distributions of ~~stationary quasi-stationary~~ Rossby waves in the original forcing data if the relevant large-scale dynamics are faithfully retained.

2.3.2 Wave activity flux

In the introduction, we used the diagnostic derived by Takaya and Nakamura (1997, 2001) to visualize the ~~flux of WA WA~~
 290 flux, which is a linear combination of kinetic energy and enstrophy, ~~and~~ is also related to the momentum and energy exchange between the mean circulation and perturbations. Similar to LZ15, TN01 used a horizontally non-uniform background with non-zero meridional winds to derive their WA budget equation, making it an appealing tool for regional climate studies (Schneider et al., 2012; Sakaguchi et al., 2016; Chen et al., 2023; Zhang et al., 2024). TN01 obtained the following conservation equation for WA from the quasi-geostrophic (QG) potential vorticity equation:

$$\begin{aligned}
 \frac{\partial M}{\partial t} + \nabla \cdot \mathbf{W} &= D_T \\
 M &= \frac{1}{2}(A + E)
 \end{aligned}
 \tag{3}$$

where ~~M~~ M ($m \text{ s}^{-1}$) is the wave activity density, \mathbf{W} is WA flux ($m^2 \text{ s}^{-2}$), ~~A and E~~ A and E are the quantities proportional to perturbation vorticity and kinetic energy, and D_T represents non-conservative diabatic and friction terms ($m \text{ s}^{-2}$). Since

WA flux is denoted by W in TN01, we refer to their WA flux as the W -vector as well. All quantities are derived from the base-state and perturbation geopotential height. The actual expression for the W -vector is provided in the Appendix C3. The vertical components of the W -vector and ~~kinetic energy (hence M)~~ the wave activity density are not included in the analysis. This is primarily because they involve vertical derivatives; ~~thus,~~ but data at multiple pressure levels with sufficient resolution ~~are required.~~ ~~Such high-vertical-resolution data~~ at a daily frequency are not always available from model archives such as NA-CORDEX ~~and are therefore not included in the analysis.~~ As a result, we infer the source/sink of WA by the convergence/divergence of the horizontal components of the W -vector. With the complexity of realistic atmospheric fields, it can be challenging to identify the climatological sources of WA at a given location; one would need to systematically pre-process the perturbations to decompose Rossby waves into different spatiotemporal scales, or use idealized numerical experiments.

In this study, we apply a 25–90-day frequency band-pass filter to the perturbation geopotential height to extract the quasi-stationary Rossby wave signals. The phase velocity is set to zero in the W -vector terms (eqn. C19). The base state is the 30-year (20-year for CAM-MPAS) daily climatology. The W -vector is calculated for each day and then averaged to produce the 30-year JJA climatology for visualization purposes. We noted that the ~~W -vector is not sensitive to different~~ result is insensitive to varying levels of spatial smoothing ~~to~~ of the background state (not shown), presumably ~~because of~~ due to the underlying QG framework. Insensitivity to the background smoothness is an advantage for a diagnostic metric. On the other hand, the QG assumption appears to limit the validity of the W -vector in low-latitude regions, where we often observe unphysical variability in the W -vector. ~~Also, unless the model simulation is an idealized one with a specified wave source, the W -vector at a given time and~~ location includes contributions from all waves from different sources (at the specified phase velocity), which may complicate interpretation. On the other hand, the ray theory by

The LZ15 ray theory predicts wave-ray propagation based on relationships between wave kinematics and the background state (e.g., how background wind shear changes wave shapes), given the specified initial conditions. It is applicable over the tropics and deals with a single wave packet from a specified location, making the ~~dynamical interpretation more~~ source attribution straightforward. However, the barotropic, non-divergent vorticity equation underlying the LZ2015 does not consider ~~the~~ an influence of divergence on Rossby wave propagation (Li, 2020), ~~which is~~ and the wave amplitude is not diagnosed. These two are included in the W -vector ~~diagnosis.~~ ~~The ray theory is more computationally involved than the W -vector due to time~~ integration and does not diagnose the wave amplitude and momentum forcing, which diagnoses the wave characteristics directly from the perturbation geopotential height. Therefore, TN01 (the W -vector ~~and~~) and LZ15 (wave-ray) diagnostics complement each other, enabling a better understanding of model biases in Rossby wave dynamics.

3 Results

3.1 Evaluation of surface air temperature

We begin with the evaluation of tas in the dynamical downscaling simulations. Figure 3b-d shows the JJA-mean tas biases of the three models against ERA-Interim, showing rather distinct spatial patterns across the models. CAM-MPAS and WRF exhibit a warm bias over Canada, whereas RegCM4 tends to have a cold bias there. Over the western CONUS, CAM-MPAS




Figure 3. JJA-mean *tas* over North America in ERA-Interim (a) and *tas* difference between ERA-Interim and (b) CAM-MPAS, (c) RegCM4, and (d) WRF. The panel (e) shows the time series of JJA-mean *tas* anomaly in each year, averaged over the central North America (the black box in (b)-(d)). The mean bias against ERA-Interim is added to the anomaly time series and shown by the colored dashed lines. The legend text includes the linear correlation (r) between the model and the ERA-Interim time series.

tends to simulate higher *tas* while RegCM4 and WRF tend to simulate lower *tas* than ERA-Interim. An exception is central North America, where all three models exhibit warm biases to varying degrees, consistent with previous studies (Morcrette et al., 2018; Sy et al., 2024). CAM-MPAS has by far the worst bias centered around the U.S.-Canada border. The notably higher bias of CAM-MPAS implies the importance of LB constraint for simulating *tas*, assuming that physics parameterizations in each model perform equally well. In the RegCM4 simulation, the **highest-largest** bias over land occurs in the South Central region. The highest bias of WRF is over Canada and further south in SGP.

To assess the timing and magnitude of seasonal anomalies, we also plot the time series of JJA-mean *tas* anomalies relative to the all-year JJA climatology in each dataset (Fig. 3e), averaged over the central North America region (black box in Fig. 3b-d). The **model**-mean bias against ERA-Interim is added to the anomaly time series (also indicated by the horizontal dashed lines).

340 Without the LB constraint, the time evolution of *tas* anomaly in CAM-MPAS is not expected to precisely follow that of ERA-Interim, except for the years with substantially strong external forcing such as the cold anomaly in 1992 after the Pinatubo eruption in the previous year (Robock and Mao, 1995); the impact is felt by CAM-MPAS through the anomalously cold SST, but not through the aerosols since the CAM-MPAS model used prescribed aerosol forcing based on the year 2000 condition (the RegCM4 and WRF simulations do not consider aerosol effects either). RegCM4, with the LB constraint, produces a
345 reasonable correlation with ERA-Interim (0.75). In some years, however, *tas* anomaly in RegCM4 deviates significantly from that in ERA-Interim (e.g., 1995-1999). WRF with spectral nudging achieves the highest correlation of 0.94 with ERA-Interim, and also with the smallest mean bias over the central North America region (2.7, 0.5, and 0.3 °C for CAM-MPAS, RegCM4, and WRF, respectively).

Figure 4 compares simulated standard deviations (σ_{model}) of monthly mean *tas* of each grid box to those in ERA-Interim
350 (σ_{ERA-I}) as the ratio ($R_\sigma = \sigma_{model}/\sigma_{ERA-I}$). ERA-Interim shows the strongest variability in the PNW region in the United States (Fig. 4a). CAM-MPAS is able to capture this variability center as indicated by R_σ being close to one over the region (Fig. 4b). However, it overestimates the *tas* variability in western Canada and the central U.S. RegCM4 overestimates the variability over most of North America, particularly over western Canada, and northern and southern central U.S., and the east coast (Fig. 4c). The contrast in the RegCM4 skills between the mean and variability indicates that the LB forcing can constrain
355 the ~~time-mean-time mean~~ but not necessarily the temporal variability of *tas*. The WRF simulation again shows very good agreement with ERA-Interim (Fig. 4d).

3.2 JJA climatology of large-scale circulations

Acknowledging that not only the upper-level dynamics but also the local land-atmosphere interactions (Bukovsky et al., 2017; Ma et al., 2018) and the upscale growth of convective systems (Qin et al., 2023) play crucial roles in *tas* bias, we focus on
360 the role of the ~~seasonal-to-sub-seasonal~~ subseasonal to seasonal scale upper-level circulations through the lens of Rossby wave dynamics. This section reviews some key aspects of the JJA climatology of the upper-level circulations relevant to ~~stationary~~ quasi-stationary Rossby waves. The evaluation of the model-simulated upper-level circulations over North America follows it.

As in other seasons, the JJA-mean zonal winds are characterized by the extratropical and subtropical jets but with lower wind speeds and less zonally uniform structure (Fig. 5a). The extratropical jet is nearly circum-global except for the dis-
365 continuities over the eastern Pacific and Atlantic oceans, where the subtropical jet extends from $\approx 20^\circ\text{N}$ latitude to merge with the mid-latitude jet. Since jet streams serve as wave guides (Manola et al., 2013; Branstator and Teng, 2017; Wirth, 2020) (Manola et al., 2013; Branstator and Teng, 2017; Wirth, 2020; White and Mareshet Admasu, 2025), we expect that Rossby waves propagate from the Pacific Ocean to North America along the mid-latitude as well as the subtropical jets. When Rossby waves enter the East Pacific and the West Coast of North America, they encounter complex mean wind patterns, where the traditional
370 assumptions for the base state in Rossby wave dynamics — namely, zonally uniform flow with zero meridional winds — are not valid. Indeed, over the Western U.S., the mean zonal and meridional wind speeds are comparable; the former range from 12 to 20 m s^{-1} , while the latter can be as high as 8 m s^{-1} (Fig. 5b). Therefore, the ~~meridional-wind's role~~ role of the meridional wind in Rossby wave dynamics should not be ignored in this region.




Figure 4. JJA monthly standard deviations of tas in ERA-Interim (a) and the ratio of the standard deviations ($\sigma_{model}/\sigma_{ERA1}$) in (b) CAM-MPAS, (c) RegCM4, and (d) WRF.




Figure 5. The 30-year JJA-mean winds at the 200 hPa level in the ERA-Interim data: (a) zonal wind, (b) meridional wind, (c) divergence, (d) relative vorticity, (e) absolute vorticity, and (f) meridional gradient of absolute vorticity.

Vorticity and divergence are essential for Rossby wave dynamics and are also shown in the figure (Fig. 5c,d). One aspect of the jet being a waveguide's waveguide nature stems from the strong horizontal shear at its edges, which ~~creates relative vorticity and its enhances the vorticity~~ gradient. Also, the interaction between vorticity and divergence ~~creates vorticity ($\eta(\nabla \cdot \mathbf{v})$), alters the local vorticity balance (via vortex stretching, $-\eta(\nabla \cdot \mathbf{v})$), acting as a source of relative vorticity anomalies~~, often called Rossby Wave Sources (RWS) (Sardeshmukh and Hoskins, 1988). In JJA, local maxima and minima of the mean relative vorticity near the jet create the meridionally banded structure over the Pacific and Atlantic oceans. The relative vorticity maxima near the subtropical jets are also strong enough to produce zonal anomalies of the *absolute* vorticity (Fig. 5e). As a result, two sharp meridional gradients of absolute vorticity, or the regions of strong restoring force for Rossby waves, exist upstream of North America from the northern and tropical Pacific (Fig. 5f). ~~A notable divergence anomaly is found in the southern flank of the jet over the North Pacific, associated with the storm track.~~ In the tropics, strong 200 hPa divergence is co-located with ~~intense precipitation. Upper-level convergence exists~~ regions of intense deep convective precipitation, most notably in the Asian Monsoon. This massive latent heating drives a downstream dynamical response, maintaining a region of pronounced upper-level convergence over the Mediterranean and ~~Middle East regions in association with the Asian Monsoon (Rodwell and Hoskins, 1996)~~ the Middle East (Rodwell and Hoskins, 1996). Further north, a secondary divergence anomaly is observed along the southern flank of the North Pacific jet, associated with the midlatitude storm track. Those are potential source regions for Rossby waves propagating to North America.




Figure 6. The JJA-mean zonal winds (left column), meridional winds (middle column), and zonal anomaly geopotential height (right column) at the 200 hPa level over the NA-CORDEX domain, in ERA-Interim (a,e, i), CAM-MPAS (b,f,j), RegCM4 (c,g,k), and WRF (d,h,l). In the second row for CAM-MPAS, the gray markers denote the approximate boundaries between the high-resolution domain, transition zone, and low-resolution domain of the variable-resolution grid. In the bottom two rows for RegCM4 and WRF, the black dashed lines denote the original model domain boundary, and the gray dashed lines denote the boundaries of the post-processed NA-CORDEX data, which excludes the blending zone near the lateral boundaries (24 and 10 grid points for RegCM4 and WRF, respectively; see also Table 1 and Appendix B1). For RegCM4 and WRF, the regional model data are shown within the NA-CORDEX data domain, and ERA-Interim data are used outside the domain, including the blending zone.

390 For evaluating the upper-level circulations in the downscaling simulations, we focus on three variables: the mean 200-hPa zonal winds ($ua200$), meridional winds ($va200$), and zonal anomalies of geopotential heights ($zg200$)(Fig. 6). As explained in Section 2.2, the modeled fields from the two LAMs are patched with the same fields of ERA-Interim outside the model domain, a visualization also used by Denis et al. (2002b). We use the full fields here instead of the differences between the simulations and ERA-Interim to emphasize the overall spatial patterns and unphysical discontinuities (difference plots are provided in
395 Fig. B1). Ideally, for LAMs, the mean circulation across the LBs appears seamless. This is the case with the WRF simulation (Fig. 6d,h,l), where its spatial patterns are identical to those from the ERA-Interim even inside the model domain; the contour plot for the difference from ERA-Interim confirms negligible bias (Fig. B1). The geopotential height is slightly and uniformly higher within the WRF domain than ERA-Interim, but identifying the sources of $zg200$ bias in WRF is left for future work.

The overall patterns simulated by RegCM4 look reasonable, but discontinuities are apparent along the boundaries (Fig. 6c,
400 g, k). Inside the model domain, the subtropical jet entering California is weaker than ERA-Interim, and the $va200$ and $zg200$ patterns are shifted to the west. These patterns are time-invariant stationary waves that exist in the *mean* circulation, which we distinguish from quasi-stationary waves defined as the perturbation on the mean. CAM-MPAS captures the general structure of the upper-level circulations without artificial boundary effects (Fig. 6b, f, j); however, the jet core is weaker and located more northwestward than ERA-Interim, while the meridional wind speeds are overestimated (also see Figure B1a,d). Consistent
405 with the overestimated $va200$ speeds, the $zg200$ zonal anomaly over North America is too high compared to ERA-Interim (Fig. 6i,j). In other words, the amplitude of the time-mean stationary waves is too strong. The position of the positive maxima of $zg200$ anomaly coincides with the spatial structure of the mean warm bias in *tas* (Fig. 3b), indicating the contribution of the upper-level mean wind bias to the *tas* mean bias. On the other hand, in the case of RegCM4, the mean bias in the upper-level winds and warm bias in *tas* do not spatially overlap.

410 3.3 Model biases in Rossby wave propagations

The mean wind biases shown above imply that the waveguide structure for Rossby waves is also biased in the model simulations. ~~To verify, we use the ray theory~~ Before evaluating the model-simulated Rossby wave propagations, we first diagnose major waveguides in the ERA-Interim data. A commonly used diagnostic is the stationary wavenumber, K_s (Hoskins and Karoly, 1981), which is derived from the dispersion relationship of Rossby waves under a zonally uniform state with zero meridional winds.
415 The stationary wavenumber is defined as

$$K_s^2 = \frac{\bar{\eta}_y}{\bar{u}_M}. \quad (4)$$

It indicates where Rossby waves can propagate and where they are likely to be trapped or reflected; regions with real-valued K_s are conducive to Rossby wave propagation, while those with imaginary K_s are not. Over the regions where K_s is real-valued, it acts as a cut-off filter for stationary waves. When K_s is small (e.g., in strong westerlies), the total wavenumber allowed is
420 low, so only very long waves can exist as stationary waves. Stationary wavenumber also acts like the refractive index for the optical wave solution, such that Rossby wave rays (paths of group velocity vectors) bend toward regions of higher K_s [see Hoskins and Ambrizzi (1993), Li et al. (2018), and Appendix C].

In Figure 7a, we apply this metric to the JJA climatology of ERA-Interim at each grid point, assuming that the metric K_s is locally applicable (e.g., Hoskins and Ambrizzi, 1993; Henderson et al., 2017; Hoskins and Woollings, 2015). The grid boxes with imaginary values are shown in white in the figure. It depicts the two waveguides along the mid-latitude and subtropical jets into North America, consistent with the mean wind patterns. Interpretation of K_s as the refractive index suggests that a Rossby wave excited within the local maximum of K_s , associated with the mid-latitude jet, is trapped within the jet and propagates zonally since the strong vorticity gradients to the north and south refract back the wave. On the other hand, a Rossby wave excited in the subtropical jet would be refracted southward toward the equator with higher K_s (the red arrow in the figure) toward the critical latitude where waves cannot propagate further, rather than traveling into North America. South of the mid-latitude jet over the central Pacific, there is another prohibited region where the zonal winds are near zero, and the meridional gradient of absolute vorticity is slightly negative (see Fig. 5). Figure 7b shows the same diagnostic but calculated from the smoothed background state, which is more appropriate for the WKB approximation underlying the dispersion relationship (Appendix C). The overall waveguide structure is similar to that in Fig. 7a, but regional maxima (i.e., waveguides) are blurred, and the prohibited region over the central Pacific is replaced by small real values.

The assumptions of zonally uniform flow and zero meridional winds are not valid over the eastern Pacific and western North America, prompting us to perform ray tracing by LZ2015. ~~To trace Rossby wave rays to confirm the waveguide structure. To do this,~~ we need to specify the locations of the wave sources. Previous studies suggest several remote sources of Rossby waves reaching North America during the summer, including the East Asian and Indian Monsoon regions, the western Pacific, the Tibetan Plateau, and the Mediterranean (e.g., Ting, 1994; Ambrizzi et al., 1995; Trenberth et al., 1998; Wang et al., 2001; Lau and Weng, 2002; Ding and Wang, 2005; Wang et al., 2007; Lin, 2009; Li et al., 2015, 2019). Most of them found Rossby waves propagating along the mid-latitude jet, but ~~Rossby waves can travel~~ several studies suggested Rossby wave propagation along the subtropical jet from ~~wave sources in~~ the central and eastern tropical/subtropical Pacific to North America (Li et al., 2019) ~~Some of those waves are~~ (Li et al., 2019; Chang et al., 2023a; Chen et al., 2023; Lubis et al., 2024). Those waves can be initiated during the Madden-Julian Oscillation phases 5 and 6, travel across North America, and break over the Atlantic Ocean ~~which are shown to suppress tropical cyclone activities~~ (Chang et al., 2023a). A closely related subseasonal variability, the boreal summer intraseasonal oscillation (BSISO), is also found to enhance convective heating during particular phases, which triggers Rossby wave trains that tend to place a high-pressure ridge over the Pacific Northwest region (Lubis et al., 2024). ~~Chen et al. (2023) found that the CMIP6 models project wave sources in the tropical/subtropical eastern Pacific to intensify under the greenhouse gas forcing, exerting a more substantial influence on the surface temperature anomalies over western North America.~~

Ensemble ray tracing is performed as described in Section 2.3.1 for the source locations suggested by previous studies and by our preparatory analyses (Appendix C2). Results from ~~five-four~~ source locations are shown in Fig. 7 using the ERA-Interim climatological winds as the base state. ~~Stationary~~ Rossby waves excited in the northern North Pacific (NP) (Fig. 7ac) and eastern tropical/subtropical Pacific (EP) (Fig. 7bd) have significantly higher probabilities of propagating over North America than those originating from other areas. Waveguides extending from the eastern subtropical Pacific (20°-30°N) to North America are evident for the waves originating from ~~these two~~ both the NP and EP regions.




Figure 7. Probability-Waveguides for (quasi-)stationary Rossby waves, (a,b) diagnosed by stationary wavenumber, K_s , (eqn. 4) and (c-f) by the probability of Rossby wave propagation from obtained by the LZ15 ray tracing method. In (a selected), K_s is calculated from the JJA climatology from ERA-Interim on its native grid resolution (T255), while in (b) it is calculated from the smoothed climatology (T10), same as the background state for the ray tracing. The grid boxes with imaginary K_s are shown in white. The ray tracing results are presented separately for each source region: (ac) northern North Pacific, (bd) eastern subtropical Pacific, (ee) western tropical Pacific, (d) East-Asian Monsoon, (e) Indian Monsoon, and (f) Tibetan Plateau. Probability is calculated for each grid box as the fraction of rays reaching the grid box over the total number of rays traced from a source region. The base-state wind fields are the 1980-2010 monthly climatologies for June, July, and August from ERA-Interim.

Most waves excited in the West Pacific region travel southeast across the equator owing to the tropical easterly zonal wind and the monsoonal southerly meridional winds (Li et al., 2019). ~~Rays from the Indian Monsoon region travel west as equatorial Rossby waves; some rays travel further west across the Atlantic Ocean, as seen in Lin (2009).~~ The Tibetan Plateau generates Rossby waves that ~~travel west~~propagate westward; some of these waves arrive in North America from the east, while others turn ~~east~~eastward over North Africa and propagate along the jet stream. Some of those results may appear inconsistent with previous studies, and it is possible that wave activities originated from the other locations to reach North America, particularly in other seasons (Wang et al., 2020; Zhang et al., 2024), through non-linear processes such as Rossby wave breaking and associated wave reflection (Abatzoglou and Magnusdottir, 2006), or by the interactions of propagating Rossby waves and the background divergent circulation (Sardeshmukh and Hoskins, 1988; Li, 2020), which are not included in the linear ray theory. Nonetheless, one-point correlation maps for meridional winds are consistent with the ray tracing result, such that statistically significant lead/lag correlations ~~with the grid points~~ over North America are found only when the base points are specified in the NP and EP regions~~(and Caspian Sea but with negative lags, thus winds over North America leads the winds over Caspian Sea; see Appendix C2).~~ With Given those results, we consider it reasonable to focus on the upwind source regions of NP and EP ~~for evaluating to evaluate~~ regional downscaling simulations.

Sub-samples of individual ray trajectories from these two regions are shown in Fig. 8 to illustrate actual wave rays and their relationship to the initial zonal wavenumber (~~k_i~~ k_0) and background circulations. The figure uses climatological July winds as the base state, but the result is qualitatively similar to those obtained with June or August climatological flows (not shown). Rossby waves excited over the NP region with smaller ~~k_i~~ k_0 (i.e., longer wavelengths) tend to travel south/southeast toward the subtropical eastern Pacific, then turn east/northeast to reach North America. The climatological flow patterns immediately south of the NP source region have the northerly meridional winds of $\approx -5 \text{ m s}^{-1}$ with comparable or even weaker zonal winds (Fig. 5a,b). Also, the meridional group velocity is inversely proportional to the second power of the wavenumber; thus, smaller wavenumbers favor larger group velocity (eqn. C11). Those two aspects likely facilitate southward propagation from NP. On the other hand, those initiated with larger ~~k_i~~ k_0 tend to propagate along the mid-latitude jet and travel across North America near the U.S.-Canada border (Fig. 8c).

Located more southeastward, the EP region is situated within the northward meridional background winds (Fig. 5b). Consistently, the waves excited here with smaller ~~k_i~~ k_0 first propagate north, and turn around at the northern edge of the mid-latitude jet (Fig. 8e,f). The initial northward propagation is not obvious in the K_s diagnostics. Similar to the waves from the NP region, waves initiated with larger ~~k_i~~ k_0 tend to be trapped within the mid-latitude jet and propagate more zonally. For both source regions, waves with even larger ~~k_i~~ k_0 are not able to propagate across North America (Fig. 8c,d, g,h) (Li et al., 2019)). The result illustrates the sensitivity of Rossby ray propagation to the base state, ~~thereby~~ highlighting the profound impact of ~~mean circulation~~mean-circulation bias on modeled Rossby wave propagation.

The stationary wavenumber K_s and LZ15 ray tracing agree on the waveguide formed by the mid-latitude jet, which is more effective for waves initiated with $k_0 \approx 6$ in the LZ15 framework. For waves with smaller k_0 , LZ15 results diverge from the waveguide depicted by K_s on: 1) southward propagations over the central Pacific, where K_s prohibits wave propagation, 2) northeastward waveguides by the subtropical jet, where K_s implies equatorward propagation toward the critical latitude south of

the jet, and 3) northward propagations off the West Coast guided by southerly meridional winds. As shown below, wave activity flux patterns from TN01 are consistent with the LZ15 result, and meridional winds off the West Coast play an important role in understanding model biases in Rossby wave propagation and their downwind impact over North America. More sophisticated constructions of the background state for K_s have been suggested (e.g., White and Mareshet Admasu, 2025), which may produce waveguide structure that is more consistent with the LZ15 and TN01 diagnostics.

We evaluate the downscaling models by ~~the ratios of the comparing the~~ ray propagation probabilities ~~(obtained with the LZ15 framework, using the ratio of model to reanalysis probabilities,~~ $R_p = p_{model}/p_{ERA-Interim}$) (Fig. 9). The ray probabilities in the WRF simulation are almost identical to those in the ERA-Interim ($R_p \approx 1$ in Fig. 9 g,h), as expected from the small bias in the upper-level circulations. For the other two models, biases in jet and meridional wind speeds lead to significantly different wave-propagation patterns ~~compared with from those in~~ ERA-Interim. For the waves initiated in the NP region, CAM-MPAS overestimates the probabilities over Canada and northern CONUS, and underestimates them over the southern part of CONUS (Fig. 9c). This is likely the result of the northward-shifted mid-latitude jet and overestimated southerly winds over the West Coast (Fig. B1a,d), which promote more zonal propagations at higher latitudes instead of traveling to the south. Such propagations are seen for the waves initiated with relatively small zonal wavenumbers (Fig. 10a for $k_x = 4k_0 = 4$). For those waves, the mean circulation patterns in CAM-MPAS support longer-lived, circumglobal propagation that passes over North America twice, ~~contributing to higher thereby increasing~~ propagation probabilities. Such long-lived waves are rare for the same initial zonal wavenumbers with the ERA-Interim base state. For the waves from the EP region, stronger southerly winds over the West Coast region likely allow more waves to travel north, but the slightly weaker and wider jet in CAM-MPAS appears to be a less effective waveguide, spreading the rays more widely over North America, particularly to the south of the jet where CAM-MPAS simulates higher propagation probabilities (Fig. 9d).

The Rossby wave probabilities in RegCM4 ~~show some similarity with those in CAM-MPAS~~ (Fig. 9e,f) show some similarity with those in CAM-MPAS, likely due to the two models sharing the mean circulation biases over the western part of the NA-CORDEX domain (Fig. B1b,g,d,e). We can see more dense lines of wave rays emanating north from the EP source region (west of 120 °W) in the RegCM4 simulation-ray tracing than in ERA-Interim (Fig. 10b v.s. Fig. 8e), where stronger southerly winds are noted in the RegCM4 simulation (Fig. B1g,e). At the same time, the overestimated southerly winds appear to limit the southward wave propagation from the NP region, thus shifting the probabilities northward over North America (Fig. 9e). The mean wind patterns over North America in RegCM4 allow waves from the EP region with larger k_x/k_0 to travel farther than in the ERA-Interim base state, for example, for the initial zonal wavenumber of eight (Fig. 10c v.s. Fig. 8h). Those waves also contribute to the higher probabilities from the EP region. LB effects are not apparent in the RegCM4 ray-tracing results. This is due to the smoothing of the base state *after* the RegCM4 and ERA-Interim data are patched onto the global grid, thereby effectively weakening discontinuities at the lateral boundaries.

Overall, biases in the large-scale circulations in CAM-MPAS and RegCM4 tend to increase wave-propagation probabilities in the northern part of North America, particularly in the RegCM4 base state. Additionally, the probabilities for Rossby waves around 40°N over CONUS from the NP region are underestimated, whereas ~~both models overestimate those the waves~~ from the EP region are overestimated by both models. Those two biases would not simply cancel each other out, because Rossby




Figure 8. Samples of Rossby wave rays using the climatological July circulation from ERA-Interim as the base state. Each panel shows rays initiated from different the North Pacific source regions and region are shown on the left column, starting with a specific different initial zonal wavenumber (k) wavenumbers: (a) the North Pacific source region with initial $k = 3k_0 = 2$, (b) also from the North Pacific source region but with initial $k = 8k_0 = 4$, (c) the East Pacific source region with initial $k = 2k_0 = 6$, and (d) also $k_0 = 8$. The right column shows the rays from the tropical East Pacific source region but with initial $k = 6$: (e) the Tibet source region with initial $k = 1k_0 = 2$, (f) also from the Tibet source region but with initial $k = 9k_0 = 4$, (g) $k_0 = 6$, and (h) $k_0 = 8$. Line colors represent time-dependent total wavenumber K (equation C10), and red dots show the source point location. Rays are terminated when the total wavenumber reaches 40 (wavelength of $\approx 1,000$ km), assuming that they are not small-amplitude perturbations at the geostrophic scale anymore (wave-breaking).




Figure 9. Probability of Rossby wave propagation from the North Pacific and East Pacific source regions obtained from ERA-Interim (a, b), and the ratio of the probabilities as $R = P_{model}/P_{ERA-I}$ for CAM-MPAS (c, d), RegCM4 (e, f) and WRF (g, h). $R = 1 = 10^0$ means the equal probabilities of ray propagation in the model and ERA-Interim. A five-point running average is applied before plotting to reduce noise, primarily over the regions of low probabilities.

waves propagating from the NP region tend to have higher wavenumbers (shorter wavelengths) over North America than those from the EP region (Fig. 9e, g8). The smaller waves from the NP region may be more susceptible to breaking. In contrast, those from the EP region in wavenumbers four to six may have higher probabilities of resonating with Rossby waves of similar wavelengths but different frequencies (Petoukhov et al., 2013; Coumou et al., 2014). Those non-linear processes are not part of our diagnostics, though.

The shifted Rossby wave propagations in CAM-MPAS and RegCM4 may disrupt the spatiotemporal correlations between Rossby waves and surface climate, as seen in the 2009 heatwave example in the Introduction. In the two model simulations, the biases in Rossby wave probabilities and *tas* variabilities both exhibit large magnitudes variability are both large over the Pacific Northwest, suggesting a connection between the two biases them. We explore the connection in the following sections.

3.4 Wave activity flux and surface air temperature

~~Calculation of the WA flux by TN01, or W-vector, requires the base and perturbation (wave-) geopotential fields. We define stationary Rossby waves as the time-filtered daily-mean perturbation from the 30-year (20-year for CAM-MPAS) daily climatology (base state). A band-pass time filter is used to extract the variations within the 25–90-day frequency range. Also, the phase velocity is set to zero in the W-vector terms. The W-vector is calculated for each day and then averaged to produce the 30-year JJA climatology for visualization purposes.~~

~~From a global point of view, the~~ We begin with a global view of wave activity in ERA-Interim. The area with the most vigorous ~~stationary Rossby wave activity~~ quasi-stationary Rossby WA in the JJA season is the Pacific Ocean, followed by the Atlantic Ocean, both characterized by large flux and strong divergence of the W-vector (Fig. ??11a). We interpret the areas of divergence as indicating WA sources. Vigorous WA fluxes from the NP and EP wave sources converge on the West Coast of North America, then propagate across the continent to diverge out from the East Coast, in general agreement with the 2010 2009 heatwave case (Introduction). The pathways from the two source regions agree with the ray theory ray theory result, including the initial southward propagation from the NP region and waveguiding by the subtropical jet.

~~Figure ?? compares~~ The bottom four panels in Fig. 11 compare the W-vector patterns over North America in the downscaling models and ERA-Interim. The most notable feature is the bands of strong divergence/convergence pairs along the LBs in the RegCM4 simulation (Fig. ??e11d), which strongly suggests inconsistency between ERA-Interim and RegCM4 circulations, even considering the removed relaxation zone (Section 2.2). Although some spuriously large W-vectors emanating from LBs should be ignored, those downwind over the Pacific coastal area and the Pacific Northwest region are calculated fully from the model data, thus reliable. There, the W-vector in RegCM4 is oriented more zonally than in ERA-Interim, and some of the WA flux appear to originate at the LB rather than from the NP and EP regions. Not only the coastal region, but also the WA fluxes over the central U.S. differ between RegCM4 and ERAI; RegCM4 simulates a more northerly W-vector, while ERA-Interim suggests a more zonally propagating flux. The W-vectors in the WRF simulation are almost identical to those from ERA-Interim (Fig. ??d11e); subtle linear structures in the divergence pattern parallel to the lateral boundaries may be due to the removal of the relaxation zone in the W-vector calculation. The global VR simulation of the CAM-MPAS model does not suffer from such artifacts (Fig. ??b11c). However, the zonal propagation of the W-vector is shifted northward from the




Figure 10. Samples of Rossby wave rays using the climatological July circulation from the downscaling models as the base states. (a) CAM-MPAS for waves initiated in the North Pacific source region with initial $k=4$ zonal wavenumber $k_0=4$, (b) RegCM4 for waves from the East Pacific source region but with initial $k=2$ $k_0=2$, and (c) RegCM4 with initial $k=8$ $k_0=8$. Line colors represent time-dependent total wavenumber K (equation C10), and red dots show the source point location. Rays are terminated when the total wavenumber reaches 40 (wavelength of $\approx 1,000$ km, assuming that they are not small-amplitude perturbations at the geostrophic scale anymore (wave-breaking)).

U.S. to Canada, creating an anticyclonic rotation over the central U.S., possibly due to the overly strong positive geopotential anomaly (Fig. 6j). With this northward shift, the W-vector divergence over the East Coast of the U.S., as seen in ERA-Interim, is replaced with weak convergence in CAM-MPAS. In addition, the W-vector divergence in CAM-MPAS is overly strong near the coastlines and mountain ranges on the West Coast compared to other models and ERA-Interim. We have examined the variance spectra of surface topography, vertical velocity, and horizontal winds, but there is no indication that the topography and wind kinetic energy in CAM-MPAS differ significantly from those in other models (Fig. B3). Topography-related processes in the CAM-MPAS downscaling simulations will be investigated in future work, ~~which may help~~ potentially helping to explain the strong WA flux divergence over the mountainous region.

How are the differences in the WA flux patterns reflected in the regional climate? We first look at the lead/lag correlations between the five-day running mean $zg200$ anomalies across North America and the divergence of the W-vector averaged over the West Coast region, where we see strong convergence in the climatology of ERA-Interim (Fig. ??11a). ~~Since the W-vector is calculated from the geopotential anomalies band-passed for 25-90 day variabilities, this assessment essentially examines correlations across time scales (5 days vs. 25-90 days) and across space between the West Coast and the rest of North America.~~ Each time series consists of daily data spanning 30 years (20 years for CAM-MPAS) of JJA seasons. The statistical significance is determined following Li et al. (2019) using the two-tailed Student's t-test against the null hypothesis of zero correlation, taking into consideration the autocorrelation of each time series in determining the degrees of freedom ([eqn. 1 in Pyper and Peterman (1998)]).

In ERA-Interim, statistically significant negative correlations appear upstream over the Gulf of Alaska and positive correlation just off the U.S. West Coast around lag -2 (Fig. 12a), that is, $zg200$ anomalies off the U.S. West Coast for a given day is positively correlated with W-vector divergence over the West Coast happening two days later (or the higher $zg200$ anomalies off the U.S. West Coast are, the stronger WA flux divergence will be in two days later over the West Coast). CAM-MPAS can reproduce the correlation pattern, albeit weaker than ERA-Interim (Fig. 12b). RegCM4 misses the negative correlation over the Gulf of Alaska, extending the area with a positive correlation northwest toward the Gulf of Alaska (Fig. 12c). WRF with the spectral nudging can capture this lag-2 correlation pattern (Fig. 12d). At lag+4, statistically significant negative correlation appears over the SGP in ERA-Interim, creating a clear wave pattern (Fig. 12e). This negative correlation means that W-vector convergence (negative values) over the West Coast leads to a positive $zg200$ anomaly over SGP four days later. WRF generally captures this lag+4 correlation pattern, but without statistical significance over SGP (Fig. 12f). The other two models struggle to reproduce the lag+4 correlation (Fig. 12f,g)

The tas response to the W-vector divergence closely follows the $zg200$ response. Again, we calculate lead/lag correlations between five-day running mean tas anomalies at each grid point and the daily W-vector divergence averaged over the West Coast region (Fig. 13). In ERA-Interim, areas of statistically significant negative correlations appear over SGP around a lag of -2, with the maximum extent occurring when the tas anomaly is lagged by four days (lag+4). It suggests that the tas over the SGP tends to be higher than normal when the WA flux converges over the West Coast, particularly 4–8 days earlier. We noted that the significant lagged correlation remains when the tas anomaly is band-pass filtered for the periods




Figure 11. JJA climatology of horizontal components of the W-vector (arrows) and its divergence (color) at the 200 hPa level in (a, b) ERA-Interim, (c) CAM-MPAS, (d) RegCM4, and (e) WRF. The regions between 10°S and 10°N are masked because the Quasi-geostrophic assumption for the W-vector is not generally valid. The black boxes in (a,b) represent the source locations used for the ray tracing in Section 3.3. Note that the vector scale and color limits are different between (a) and the other panels. In the RegCM4 result, a Gaussian filter is applied to the relaxation zone.




Figure 12. The lead-lag correlation between W-vector divergence averaged over the West Coast (blue box) and the five-day running mean $zg200$ at each grid box. Negative lags mean that the $zg200$ time series leads and is shifted earlier by that amount —e.g., by 2 days in panel (a)— before the correlation is calculated against the time series of W-vector divergence. With positive lags, $zg200$ lags the W-vector divergence, i.e., the $zg200$ time series is shifted later by that amount. Yellow and black contours indicate areas with statistically significant correlations at $\alpha = 0.10$ and $\alpha = 0.05$ levels, respectively.

between 70 and 90 days (not shown). The long timescale may indicate a role for the land surface, particularly soil moisture (e.g., Dirmeyer and Halder, 2017). The actual physical processes underlying the correlation are left for future work.

Focusing on the lag+4 result, WRF is the only model to simulate the significant negative correlation over SGP and the overall structure of the lead-lag correlation (Fig. 13b,f). CAM-MPAS simulates a weak negative correlation over SGP but misses the statistical significance (Fig. 13d). Also, the correlation patterns over the Pacific Northwest, western Canada, and Alaska do not agree with those in ERA-Interim. Similarly, RegCM4 misses the negative correlation center over SGP, and also simulates unrealistic negative correlation over the eastern Pacific (Fig. 13e). The inability to reproduce the W-vector – *tas* correlation in these two models is likely one reason for the biases of the mean and/or variability of *tas* over SGP (Figs. 3 and 4).




Figure 13. Same as Fig. 12, but for the lead-lag correlations between the W-vector divergence averaged over the West Coast (blue box) and the five-day running mean (*tas*) anomaly at each grid box.

It is also possible to correlate the W-vector divergence and the *errors* in the simulated *tas*. Figure 14a presents the lead-lag correlations between the W-vector divergence averaged over the West Coast region in the RegCM4 simulation and the difference in the daily mean *tas* between RegCM4 and ERA-Interim at each grid point. In this case, we observe statistically significant correlations over SGP at lag +4. ~~This correlation and the lack of correlation between *tas* anomaly and the W-vector divergence (Fig. 13e) together~~ Those RegCM4 results suggest that accurately receiving Rossby wave signals through LBs and maintaining the large-scale ~~forcing-circulation~~ patterns are essential for LAMs to reproduce the cross-scale ~~forcing connections~~ from Rossby waves to *tas* ~~in the driving reanalysis or GCM data~~. Another example is based on the observation that the W-vector convergence is significantly overestimated by CAM-MPAS over British Columbia ~~in~~ Canada (Fig. ??11). The lead/lag correlations between the W-vector divergence averaged over British Columbia, and *tas* errors in CAM-MPAS exhibit significant positive correlation in the same region, which also overlaps the overestimated *tas* variability by the same model (Fig. 4b). ~~Across North America, the correlations show a wave-like structure resembling the Pacific-North America pattern, with significant correlations emerging over the Four Corners and Florida; however, those areas do not show systematic *tas* errors in the mean or standard deviations.~~ Part of these *tas* errors is attributable to out-of-sync temporal evolutions between ERA-Interim and global, free-running CAM-MPAS, which has its own internal variabilities. Nonetheless, this result ~~suggests error propagation~~ illustrates another example of how model error can propagate across scales, from the biased mean wind patterns through ~~erroneous propagations and convergence of Rossby waves WA to errors in the modeled *tas*~~ Rossby wave forcing to *tas* variability.




Figure 14. Similar to Fig. 13, but for the lead-lag correlations between the W-vector divergence averaged over the West Coast region in RegCM4 (a) and Canadian Pacific Northwest region in CAM-MPAS (b) and the simulation errors (model minus ERA-Interim) in the five-day running mean (*tas*) at each grid box.

3.5 Rossby wave and heatwaves

In this final subsection, we demonstrate a connection between ~~stationary-quasi-stationary~~ Rossby wave forcing and heatwave (HW) events, and assess how the downscaling simulations replicate the connection identified in ERA-Interim. We diagnose

HW events using the criteria outlined in Barriopedro et al. (2023) (their Appendix A2). Specifically, an HW event is a period of
625 three or more consecutive days with the daily maximum *tas* exceeding the 95th percentile of the reference period (1981-2010,
except for CAM-MPAS, for which we use 1990-2010). All seasons are included in HW identification, and the seasonal cycle
is not removed; thus, this criterion favors warm-season occurrences (Barriopedro et al., 2023). Figure 15a shows the spatial
distributions of the average fraction of HW days per JJA season (i.e., the number of HW days during one JJA season = 92
days). ERA-Interim indicates two local maxima, one over the southwestern U.S. and the other over the SGP, where $\approx 15\%$ of
630 JJA days, or about 14 HW days, are expected each summer.

The same HW definition is applied to the downscaling simulations, also shown in Fig. 15. CAM-MPAS ~~is able to~~ can
simulate the overall spatial patterns with two local maxima over the southwestern and south-central U.S., but overestimates
the number of HW days during the summer across most of North America, except in the eastern part, where it simulates fewer
HW ~~events~~ days. RegCM4 also simulates too many HWs in the JJA season across North America, except for SGP, where it
635 underestimates the number. The HW distributions simulated by WRF agree best with those in ERA-Interim.

How does the HW distribution change during the days with strong Rossby wave forcing? We examine the days when the
WA flux convergence ($-\nabla \cdot \mathbf{W}$) over the West Coast region (the same region used for the lead/lag correlations) exceeds the top
90th percentile of all years. The bar graphs in Fig. B4 show how those extreme days are distributed across years. Those top 10
percentiles are not uniformly distributed but instead exhibit variability on a 3–4 year timescale, according to the ERA-Interim
640 data. ~~A notable absence of extreme convergence days from 2004 to 2008 is followed by an abnormally high peak in 2009
(Fig. B4a), when unusually persistent upper-level circulations brought intense HWs to the Pacific Northwest, as illustrated
in the Introduction. We expect LAMs to reproduce~~ WRF reproduces this temporal distribution reasonably well, ~~since the
western inflow boundary conditions provide the WA flux variability. Again, it is the case with WRF, but not necessarily with
whereas~~ RegCM4 (Fig. B4c,d), ~~although RegCM4 is able to capture the 2009 peak.~~ does not. Free-running CAM-MPAS does
645 not simulate the 2009 peak or other clusters in sync with ERA-Interim (Fig. B4b), indicating the significant roles of biased
waveguide locations and/or the atmosphere's internal variability ~~on the WA flux variability, which is not well constrained by
the prescribed SST. The spatial structure of the W-vector during the extreme case is generally similar to the climatological
pattern (Fig. ??), so the model bias on the propagation pathways in the previous section applies to the extreme convergence
days (not shown). In terms of the magnitude of extreme~~ (. All models agree well with ERA-Interim on the magnitude of the top
650 10th percentile) W-vector convergence (indicated by the black markers in the figure), all models agree well with ERA-Interim:
the average magnitude of the extreme convergence is -27, -26, -27, and -25 ($10^{-8} \text{ m}^2 \text{ s}^{-1}$) in ERA-Interim, CAM-MPAS,
RegCM4, and WRF, respectively.

The bar graphs with the left y-axis show the distribution of the number of days with the top 10% strongest convergence
of WA flux convergence over the West Coast area (blue boxes in Figs. 12 and 13) in (a) ERA-Interim, (b) CAM-MPAS, (c)
655 RegCM4, and (d) WRF. The black circles represent the average magnitude of the extreme WA flux convergence in each JJA
season, with the right y-axis. In panel (b), the first 10 years are grayed-out since CAM-MPAS data are not available for this
period.




Figure 15. Fraction of the days identified as heatwaves (HWs) (or number of HW days per JJA (92) days) ~~identified as HWs~~, in (a) ERA-Interim, (b) CAM-MPAS, (c) RegCM4, and (d) WRF.

Calculating the fraction of HW days only on the days with extreme W-Vector convergence over the West Coast area in the ERA-Interim data, we see significantly higher HW fractions over the Midwest and the South Central U.S. and southern Canadian Prairies, and lower fractions in the northern Canadian Prairies, Quebec, and the Southwestern U.S. (Fig. 16a). That is, extremely rapid accumulations of WA over the West Coast region have a statistically significant impact on HW occurrences ~~over those broad regions across~~ across broad regions of North America. Note that the HW fraction is roughly doubled over the Central Plains from $\approx 0.12 - 0.15$ with all the samples to $\approx 0.2 - 0.3$ during the extreme WA flux convergence. Examining the composite means of the HW day fractions in the downscaling simulations, the WRF simulation yields the best agreement with ERA-Interim, although it does not accurately capture the reduced HW occurrences over the southwestern U.S. In CAM-MPAS, the higher HW fractions are seen over British Columbia and Quebec, the Southwest, and some parts of the Great Plains. Those responses differ from what ERA-Interim describes, and are somewhat similar to the areas with overestimated

670 variability of *tas* by CAM-MPAS this model (Fig. 4b). In-RegCM4, simulates HW surge with the extreme WA flux is more concentrated over the southern part of CONUS, possibly related to the more northerly WA flux downwind over the West Coast and Great Plains in RegCM4 this simulation compared to the easterly-westerly WA flux in ERA-Interim (those during the extreme convergence not shown, but similar to Fig. ??11). More in-depth analysis is required to conclude how the modeled HW response to WA flux is linked to the overall mean and/or variability biases of *tas*. Nonetheless, this diagnosis reveals a connection between extreme Rossby wave forcing and the occurrence of HWs over North America, which is accurately reproduced only by the WRF model with spectral nudging.




Figure 16. Difference in the fraction of the HW days, as the difference between the fraction calculated only during the extreme W-vector convergence over the Western Coast area and during the rest of the samples in (a) ERA-Interim, (b) CAM-MPAS, (c) RegCM4, and (d) WRF. The cross-hatch indicates that the difference is statistically significant at the 0.05 level.

Before summarizing our main findings, we report preliminary investigations into two outstanding questions arising from the presented results. First is the reason for the striking differences between the two LAMs, RegCM4 and WRF, in the upper-level circulations and ~~stationary~~ quasi-stationary Rossby waves presented above. The two differ in many ways: numerical grid discretizations, hydrostatic vs. non-hydrostatic dynamical core, width and weight functions of the LB buffer zones, and every component of physics parameterizations (Tables 1 and B1). Here, we focus on the impact of spectral nudging adopted in the WRF simulation for NA-CORDEX. We conducted two sensitivity simulations with WRF version 4.6.1. The only difference between the two sensitivity simulations is whether spectral nudging is used (“Nudge”) or not (“NoNudge”). The model configuration is identical to that used for the NA-CORDEX WRF simulations (Chang et al., 2015; Diez-Sierra et al., 2022a), except for using the hybrid sigma-pressure vertical coordinate that is the default option since WRF version 4.0, instead of the traditional sigma coordinate used in the NA-CORDEX simulation (Skamarock et al., 2019). It is not expected that the different vertical grid (with the same resolution) will impact the following result. Both simulations are initialized at 2010/03/01 00:00 UTC and run for nine months ~~to~~, ending at 2010/11/30 23:00 UTC. Only the results from the JJA months are presented. ~~We need the~~ The extended periods before and after JJA are used to apply the 25-90-day band-pass filters ~~to define the quasi-stationary perturbations~~, as done in the main result to calculate the W-vector for quasi-stationary Rossby waves.

~~Figure B5a,b compares the JJA-mean biases of the two simulations in $ua200$ against ERA-Interim.~~ Consistent with the main result, the Nudge experiment shows little difference from ERA-Interim on the upper-level circulations, while NoNudge simulates a weaker and shifted mid-latitude jet during this particular summer. ~~W-vector is calculated from the sensitivity simulations using the single-season JJA mean as the base state (Figure B5c,d(Fig. B5a,b)).~~ The NoNudge experiment produces artificial W-vector divergence/convergence pairs along the LBs, similar to the RegCM4 result, while the Nudge experiment shows no such artifacts. ~~The two simulations exhibit distinct patterns of the W-vector convergence over the West Coast region, likely the combined result of the biased mid-latitude jet and artificial LB effect. In the Nudge experiment, the W-vector converges over a larger area that encompasses the Southwest, the northern Rockies, and the Pacific Northwest. On the other hand, the NoNudge simulation directs WA fluxes into lower latitudes with little convergence over the Pacific Northwest. (Fig. B5c,d).~~ Although the simulation length is limited for a rigorous evaluation of Rossby waves, the result strongly supports the notion that spectral nudging is the dominant factor in the differences between RegCM4 and WRF. This interpretation is consistent with previous studies demonstrating the effectiveness of spectral nudging in maintaining large-scale circulations from the forcing data and their impact on the near-surface climate. (Miguez-Macho et al., 2004; Alexandru et al., 2009; Castro et al., 2012).

Another question is the possibility of upscale effects on the mean winds simulated by CAM-MPAS and RegRM4 (Figs. 6 and B1). Both models simulate stronger southerly winds and weaker mid-latitude and subtropical jets off the West Coast of CONUS than in ERA-Interim, which direct Rossby waves northward to higher latitudes (Figs. 9 and ~~??~~11). Since the two models have finer spatial resolutions than ERA-Interim and their dynamics are not constrained by nudging, momentum and vorticity sources that are not ~~well-resolved~~ well resolved by ERA-Interim may be better represented by the two models. The

710 impact of the Rockies and organized convection on Rossby waves over North America has been noted by previous studies (Ting, 1994; Stensrud, 2013; Rodwell et al., 2013).

The impact of different spatial resolutions ~~can be is~~ assessed by comparing the more recent reanalysis product, ERA5 (Hersbach et al., 2020), ~~to with~~ ERA-Interim, as ERA5 has a ~~similar grid spacing to grid spacing similar to that of~~ the models examined (≈ 25 km). ~~The We find that the resolution differences make negligible contributions to the~~ JJA-mean ua_{200} and zg_{200} ~~from ERA5 are compared to those from ERA-Interim in~~ (Fig. B6a,b, ~~from which it is clear that the resolution differences~~ ~~make negligible contributions to those mean fields~~). We also look at one-season average differences in va_{200} and zg_{200} between the NoNudge WRF simulation and ERA-Interim. If higher resolution is responsible for the upper-level wind differences, the NoNudge simulation might exhibit stronger southerly winds and a weaker jet, as in CAM-MPAS and RegCM4. ~~Figures~~ ~~This is not the case based on Figs. B5a and B6c,dtell us that this is not the case~~. The va_{200} difference shows weaker southerly winds off the West Coast in the WRF NoNudge experiment, which is the opposite of those by CAM-MPAS and RegCM4. The
720 seasonal means of ua_{200} and zg_{200} in NoNudge do not exhibit the same spatial patterns as those in CAM-MPAS or RegCM4. Therefore, it is unlikely that physically oriented upscale effects are the main reasons for the mean circulation differences between ERA-Interim and the two models.

In ~~our main results~~~~this study~~, quantifying large-scale biases is a common practice; we believe the novelty lies in ~~tying~~ ~~linking~~ circulation biases to ~~stationary-quasi-stationary~~ Rossby waves and ~~further, in turn,~~ to near-surface air temperature and
725 heatwaves. Some of the diagnostics we use could be helpful as part of climate model diagnostics packages (e.g., Eyring et al., 2020; Zhang et al., 2022; Stan et al., 2025). One use case is to identify GCMs with good skills in simulating wave sources and waveguides toward North America (e.g., ~~(Goldenson et al., 2023)~~~~Goldenson et al. (2023)~~). Another application is obviously the evaluation of a dynamical downscaling framework, and we expect metrics to depend on the specific regions. For North America, we recommend using ray theory to track Rossby waves from the North Pacific and the Eastern tropical/subtropical
730 regions, which “translate” the biases in the mean circulation into the likelihood of Rossby wave propagations over North America. Another useful diagnosis is the divergence/convergence of WA flux over the West Coast. Among available WA formulations, the WA budget equation derived by TN01 does not require a time average to define the perturbation component. This means we can evaluate time series of W-vector convergence, which can then be correlated with other variables, such as *tas*.

735 Despite the insights we can gain from those diagnostics, they also have limitations that ~~need to must~~ be overcome to ~~become~~ ~~part of~~ ~~be included in~~ such diagnostic packages. As stated earlier, ray tracing involves integrating ordinary differential equations over time, ~~which makes making~~ this technique computationally more intensive than typical evaluation ~~metries~~~~methods~~. W-vector diagnostics involve a more straightforward calculation, but their challenge lies in the complexity of the underlying theory rather than the numerical coding and data requirements for comprehensively analyzing three-dimensional budget terms.
740 The LZ2015 ray theory also shares the former challenge of complexity in ~~the its~~ underlying theory, but its publicly available repository includes documentation ~~that clarifies how the model data is used by the diagnostics~~ ~~explaining the source code, along~~ ~~with example calculations~~ and references to relevant literature (Yang, 2025; Yang and Li, 2025). ~~Other, simpler diagnostics~~ ~~exist for Rossby waves (e.g., Rossby wave sources and stationary wavenumber; see Appendix C), but they often involve more~~

745 ~~stringent assumptions, some of which may not be valid for studying regional climate. We hope this study provides valuable guidance, particularly for those interested in the North American climate.~~

Finally, all diagnostics we applied are based on the linear framework ~~, thus excluding that makes several assumptions and excludes~~ the effect of interactions with transient eddies (Teng et al., 2019) ~~on stationary wave propagation~~, waves with finite (larger) amplitude Huang and Nakamura (2017), and eventual wave-breaking (Zhang and Wang, 2018), despite the crucial roles they play in extreme events (Nakamura and Huang, 2018; Chang et al., 2023b). In Appendix C, we review the linear wave theory and other common diagnostics for Rossby waves, hoping to provide guidance for those focused on regional climate. We will continue to assess the robustness of Rossby wave diagnostics and their physical relationships with other climate variables at regional scales.

5 Conclusions

It is well known that not only global but also regional models exhibit warm *tas* bias during the summer over the central CONUS (Morcrette et al., 2018). While the role of physics parameterizations is significant on the surface warm bias, a recent study by Luo et al. (2022) found that biases in the upper-level stationary waves cause significant errors in the simulated surface temperature and precipitation. Investigating the sensitivity of WRF simulations to model resolution, convective parameterizations, and use of spectral nudging, ~~(Gao et al., 2017)~~ Gao et al. (2017) found that the SGP warm biases in the model are rather insensitive to the model resolution or the convective parameterization, but they were largely alleviated using spectral nudging that constrains the model to provide realistic moisture transport to the SGP region. Our results reiterate the importance of the model's ability to accurately simulate large-scale circulations, specifically the ~~stationary~~ quasi-stationary Rossby wave, for dynamical downscaling over North America. We evaluate three distinct dynamical downscaling approaches to incorporate large-scale forcing: 1) a standard regional climate simulation with a LAM, represented by the RegCM4 simulation, 2) a LAM simulation with spectral nudging to constrain the large-scale atmospheric dynamics, represented by the WRF simulation, and 760 3) a global VR model that simulates large-scale circulations on its global grid. The first two model data are obtained from the NA-CORDEX model archive (Mearns et al., 2017), while the CAM-MPAS data is produced by our previous work following the NA-CORDEX protocol (Sakaguchi et al., 2023). We evaluate the consistency of the large-scale circulations across the model domain boundaries by patching the LAM data and the driving data, ERA-Interim, onto a single global grid, on which the upper-level circulation and Rossby wave propagation are diagnosed.

770 We observe a striking contrast between RegCM4 and WRF in their consistency with the large-scale circulations of ERA-Interim. A pair of short WRF simulations with and without spectral nudging suggests that the spectral nudging applied in the NA-CORDEX WRF simulation is the primary reason for the difference between the two models. Specifically, in the RegCM4 simulation without spectral nudging, the mid-latitude and subtropical jets over the eastern North Pacific are weakened, and the time-mean geopotential patterns are shifted westward, resulting in stronger southerly meridional winds over the same region. 775 Furthermore, discontinuities in the time-mean circulation structure are apparent along LBs. The global model CAM-MPAS also suffers from mean circulation biases over the NA-CORDEX domain, featuring a weaker and northward-shifted mid-latitude

jet, a weaker subtropical jet over the eastern North Pacific, and an overly strong positive geopotential anomaly that is centered over the Pacific Northwest, which also results in stronger southerly meridional winds over the West Coast area. ~~CAM-MPAS's upper-level geopotential bias is found to overlap with the area of highest mean bias in *tas*.~~

780 A linear ray theory by LZ15 proves useful for linking those circulation biases to Rossby wave paths entering North America, owing to the relaxed assumption about the meridional winds in the mean background state. In the CAM-MPAS simulation, overestimated southerly winds and weaker zonal jets allow more Rossby waves to propagate northward from the low-latitude eastern Pacific to the Pacific Northwest, particularly to British Columbia. RegCM4 exhibits the same tendency as CAM-MPAS, overestimating the probability of Rossby wave passage over the Pacific Northwest and the eastern half of Canada. The WRF
785 model with spectral nudging reproduces the Rossby wave propagation patterns in ERA-Interim.

Another diagnostic to complement the ray theory is the flux of ~~kinetic~~ energy and momentum flux associated with Rossby wave packets, combined as wave activity (WA). The formulation by TN01 (W-vector) also allows a non-uniform background state and non-zero meridional winds, showing WA propagation patterns consistent with the LZ15 ray theory. ERA-Interim suggests that WA flux tends to converge over the West Coast of North America, and the temporal evolution of WA flux
790 convergence is correlated to that of *tas* anomaly over SGP. CAM-MPAS and RegCM4 cannot reproduce this correlation because of a biased mean circulation, leading to different Rossby wave propagation into North America: either to higher latitudes (CAM-MPAS) or at a different angle (RegCM4). We further find a relationship between extreme Rossby wave forcing (the top 10 percentile of WA flux convergence over the West Coast region) and HW occurrences across North America. This relationship is also not well simulated by CAM-MPAS and RegCM4.

795 The Rossby wave diagnostics used here translate large-scale circulation biases into ~~Rossby wave propagation~~ the propagation of Rossby waves and their influence on the surface climate. This physical connection across space and scales is disrupted by biases in the mean circulation patterns in the global VR model, or by the failure to faithfully retain large-scale forcing in the LAM dynamical downscaling. For the large spatiotemporal scales of ~~stationary-quasi-stationary~~ Rossby waves, spectral nudging that constrains only the larger spatial scales helps the model reproduce nearly all aspects of Rossby wave dynamics
800 and their impact on temperature anomalies across North America. ~~The implications for dynamical downscaling over North America~~ Their implications are: 1) the host GCMs and global VR models need to be able to simulate the wave sources in the eastern hemisphere (e.g., diabatic heating from organized convection) as well as the mean wind patterns over the Pacific and North America to provide correct waveguide into North America and 2) Spectral nudging is beneficial for dynamical downscaling using LAMs to avoid numerical artifacts from the LB treatment on incoming Rossby waves as well as to maintain
805 the large-scale circulation patterns on which Rossby waves propagate. Although our analysis focuses on the connection between Rossby waves and surface temperature, model biases in ~~wave activity~~ WA could also affect precipitation in the simulations, as seen in the common dry biases over the SGP. The latter can be further explored in the future using similar Rossby wave and ~~wave activity~~ WA diagnostics discussed in this study.

Code availability. The scripts used for post-processing, analysis, and visualization are available from the Zenodo archive at <https://doi.org/10.5281/zenodo.17458434> (Sakaguchi, 2025). The LZ15 ray tracing code is updated by Yang and Li (2025) and available from <https://github.com/yinan-codes/Rosby-wave-ray-tracing>.

Data availability. The ERA-Interim (for Medium-Range Weather Forecasts, 2009) is available at <https://doi.org/10.5065/D6CR5RD9> on the NSF NCAR's Geodata Science Exchange, so are the NA-CORDEX data at <https://doi.org/10.5065/D6SJ1JCH>. The CAM-MPAS data is described at <https://doi.org/10.25584/PNNL.data/1895153> and more easily downloadable through the National Energy Research Scientific Computing Center (NERSC) Science Gateway. The post-processed data used in the study is also available from the NERSC Science Gateway.

Appendix A: 2009 heatwave




Figure A1. (a) lead-lag-linear correlations between *tas* averaged over the Canada-U.S. Pacific Northwest [shown by a box with black dashed line Fig. 1g-h] and the lagged divergence of WA flux averaged over the U.S. Pacific Northwest [the red dashed-line box in 1a-c] and (b) time series of the regional-average *tas* anomaly (black), WA flux divergence (green), and the WA flux divergence shifted by six days earlier, corresponding the lag -6 at which the lag correlation reaches the absolute maximum (the lowest negative correlation).

Appendix B: Additional information for model evaluations




Figure B1. The JJA-mean zonal wind biases against ERA-Interim in (a) CAM-MPAS, (b) RegCM4, and (c) WRF. Corresponding meridional wind biases in (d, e, f) and geopotential biases in (g, h, i).




Figure B2. Illustration of the blending zone impact on derived diagnostics. The top row shows the JJA-mean vorticity calculated after remapping to the global grid and patching outside the domain with the ERA-Interim data, and bottom row shows W-vector and its divergence, using the model output with the buffer zone (a, d), without the buffer zone (b, d), and without buffer zone but Gaussian filter is applied to the (remapped) grid points located in the buffer zone (c, f).

Table B1. Physics parameterizations used in the three downscaling models. For the references for each parameterization, readers are referred to Diez-Sierra et al. (2022a) for RegCM4 and WRF, and Sakaguchi et al. (2023) for CAM-MPAS.

Component	RegCM4	WRF	CAM-MPAS
Land Surface	BATS	NOAH	CLM4
Subgrid land surface tiles	No	No	Yes
Boundary Layer	Modified Holtslag	MYJ	UW
Cloud microphysics	SUBEX	WSM3	MG2
Deep convection	Grell	Kain-Fritsch	Zhang-McFarlane
Shallow convection	-	Kain-Fritsch	UW
Longwave Radiation	CAM	RRTM	RRTMG
Shortwave Radiation	CAM	Goddard	RRTMG
Aerosols	no aerosols	no aerosols	prescribed




Figure B3. Power spectra of (a) pressure vertical velocity (ω , $Pa s^{-1}$) at the 200 hPa level, (b) ω at 500 hPa, (c) surface topography, and (d) horizontal winds at the 200 hPa level. All variables are regridded to the WRF native grid with 25 km grid spacing, then the Discrete Cosine Transform is used to calculate the spectra (Denis et al., 2002a).




Figure B4. The bar graphs with the left y-axis show the distribution of the number of days with the top 10 % strongest convergence of WA flux convergence over the West Coast area (blue boxes in Figs. 12 and 13) in (a) ERA-Interim, (b) CAM-MPAS, (c) RegCM4, and (d) WRF. The black circles represent the average magnitude of the extreme WA flux convergence in each JJA season, with the right y-axis. In panel (b), the first 10 years are grayed out since CAM-MPAS data are not available for this period.






Figure B5. The effect of spectral nudging with the WRF model in the JJA-mean zonal wind bias against ERA-Interim (left column) and horizontal component of W-vector and its divergence (right column), all at the 200 hPa level: (a,c) without spectral nudging, (b, d) with spectral nudging. [W-vector is calculated from the sensitivity simulations using the single-season JJA mean as the base state.](#)



(a) ERA5 - ERA-Interim, va_{200}



(b) ERA5 - ERA-Interim, zg_{200}



(c) WRF NoNudge - ERA-Interim, va_{200}



(d) WRF NoNudge - ERA-Interim, zg_{200}





Figure B6. Differences between ERA5 and ERA-Interim in the JJA-mean (a) va_{200} and (b) zg_{200} , as well as the differences between the WRF simulation without spectral nudging and ERA-Interim in (c) va_{200} and (d) zg_{200} . The 1980-2010 JJA climatology is used in (a) and (b), while the 2010 JJA season only is used in (c) and (d).

B1 Lateral boundary treatment in RegCM4 and WRF

This subsection introduces the configurations of the LB buffer zone in the RegCM4 and WRF simulations for NA-CORDEX. Both models follow the LB treatment proposed by Davies (1976), with several options for the shape and coefficients of the weighting functions to blend the model-predicted values and the large-scale forcing data. The RegCM4 and WRF simulations for NA-CORDEX differ in several of those options, as summarized below from Mearns et al. (2017) and Diez-Sierra et al. (2022a).

On the outermost grid point, referred to as the specified zone in Skamarock et al. (2008), all the prognostic variables are strictly those provided from the forcing data after interpolation in time and space (Fig. B7a). The specified zone is a single grid box in both RegCM4 and WRF simulations. The next n_{relax} grid points constitute the relaxation zone, where additional terms are included in the prognostic equations (eqn. 6 in Giorgi et al. (1993) and eqn. 6.1 in Skamarock et al. (2008)):

$$\left(\frac{\partial \alpha_m}{\partial t}\right)_n = (\text{advection, source/sinks, and other physical terms}) + \dots F(n)F_1(\alpha_f - \alpha_m) - F(n)F_2\nabla^2(\alpha_f - \alpha_m) \quad (\text{B1})$$

where α_m and α_f are the model-simulated and forcing values for a prognostic variable α , respectively, and n is the gridbox index from the boundary ($n = 1$ for the specified zone, and $n = n_{spec} + n_{relax}$ for the last grid box within the relaxation zone). In this formulation, model-predicted values are relaxed toward the external data by Newtonian relaxation (the second-to-last term) and the differences between the modeled and large-scale forcing values are smoothed by the diffusion-like term (the last term).

F_1 and F_2 are constants that depend on the timestep and grid spacing. F_1 is exactly the same in the two models as

$$F_1 = \frac{0.1}{\Delta t}$$

where Δt denotes the model time step (s). F_2 is slightly different; RegCM4 uses the following form

$$F_2 = \frac{(\Delta s)^2}{50\Delta t}$$

where Δs denotes the grid spacing (km). For WRF, it is given as

$$F_2 = \frac{1}{50\Delta t}$$

$F(n)$ is a weighting function that gradually reduces its magnitude from the outer-most to the inner-most grid boxes within the relaxation zone. Both RegCM4 and WRF offer the options of linear and exponential functions. The WRF simulations for NA-CORDEX use a linear weighting function with $n_{spec} = 1$ and $n_{relax} = 10$:


$$F(n) = \frac{n_{spec} + n_{relax} - n}{n_{relax} - 1} = -\frac{11}{9}n$$

, while the RegCM4 simulations use an exponential function:

$$F(n) = \exp\left(-\frac{(n-2)}{N_I}\right).$$

The coefficient $N_I = N_I(z)$ varies with height so that the model receives stronger large-scale forcing at higher altitudes (Giorgi et al., 1993). Fig. B7b shows three examples of RegCM4's exponential weight function for $N_I = 1, 3, 6$, along with the linear function in WRF.

(a) Lateral (western) boundary configuration



(b) Weighing coefficients, $F(n)$




Figure B7. Illustration of the lateral boundary conditions in the RegCM4 and WRF configurations for NA-CORDEX: (a) an example configuration of specified and relaxation zones in the western boundary with $n_{relax} = 10$, and (b) weighing coefficients ($F(n)$) in RegCM4 and WRF. Three curves are shown for RegCM4 corresponding to $N_I = 1, 3, 6$.

Appendix C: Rossby wave

850 C1 Linear wave theory

We provide a brief review of the linear wave theory to help readers without a strong background in this topic understand the Rossby wave diagnostics. The materials follow sections 7.7 and 10.5 in Holton (2004), chapter 6 in Vallis (2017), and more complex cases in [Karoly \(1983\)](#) and Li et al. (2015).

The dynamics of Rossby waves are studied in terms of the conservation law for quasi-geostrophic potential vorticity (e.g.,
855 Takaya and Nakamura, 2001; Li, 2020) since the restoring force for Rossby waves is the gradient of potential vorticity. For studying Rossby wave propagation in the atmosphere away from strong divergence, it is also common to use the vorticity equation in the barotropic atmosphere (i.e., a single layer with constant density in x and y , thus non-divergent circulations), which is:

$$\frac{\partial \eta}{\partial t} + u \frac{\partial \eta}{\partial x} + v \frac{\partial \eta}{\partial y} = 0 \quad (\text{C1})$$

860 where η denotes the absolute vorticity $\eta = \zeta + f$, $f = 2\Omega \sin \phi$ is the Coriolis parameter, and ζ is the vertical component of relative vorticity. The friction is ignored. [Also, with the assumption of a non-divergent system, \$u\$ and \$v\$ are the zonal and meridional components of rotational winds, respectively.](#) Strictly speaking, this [single-level-single-layer](#) (barotropic or shallow-water) vorticity equation is applicable for the middle ($< \approx 300 \text{ hPa}$) or near the top of the troposphere ($> \approx 150 \text{ hPa}$), where wave structure is nearly vertically uniform (equivalent barotropic) (Sardeshmukh and Hoskins, 1988; Jin and Hoskins, 1995;
865 Holton, 2004; Neduhai et al., 2024). At the same time, one needs to consider that outflows from the tropical convective systems to excite Rossby waves, and subsequent wave propagation to the middle latitudes, take place primarily in the upper troposphere ($\approx 100 - 300 \text{ hPa}$) (Jin and Hoskins, 1995; Neduhai et al., 2024). Our analysis of the 200-hPa level is chosen primarily for the availability of high-frequency outputs from NA-CORDEX at three pressure levels: 200, 500, and 850 hPa (CORDEX, 2009), but it seems to be a reasonable compromise.

870 It is also common to use streamfunction, instead of vorticity, to study Rossby waves (Chen and Chen, 1990). The vorticity budget terms are noisy and not straightforward to visually interpret (Kang and Held, 1986). Streamfunction is a scalar from which rotational winds are obtained by differentiation:

$$\begin{aligned} u &= -\frac{\partial \psi}{\partial y} \\ v &= \frac{\partial \psi}{\partial x} \end{aligned} \quad (\text{C2})$$

, and we can write vorticity in terms of streamfunction

$$\begin{aligned} \zeta &= \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \\ &= \frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} = \nabla^2 \psi \end{aligned} \quad (\text{C3})$$

875

Then equation C1 becomes:

$$\left(\frac{\partial}{\partial t} + u \frac{\partial}{\partial x} + v \frac{\partial}{\partial y} \right) (\nabla^2 \psi + f) = 0 \quad (C4)$$

All the diagnostics we use are based on a linear perturbation framework, in which Rossby waves are defined as small perturbations (a') from the background (mean) state (\bar{a}):

$$\begin{aligned} u &= \bar{u} + u' \\ v &= \bar{v} + v' \\ 880 \quad \psi &= \bar{\psi} + \psi' \\ \zeta &= \bar{\zeta} + \zeta' \\ \bar{\eta} &= \bar{\zeta} + f = \nabla^2 \bar{\psi} + f \end{aligned} \quad (C5)$$

then equation C4 becomes:

$$\begin{aligned} \left(\frac{\partial}{\partial t} + \bar{u} \frac{\partial}{\partial x} + \bar{v} \frac{\partial}{\partial y} \right) \nabla^2 \psi' + u' \frac{\partial \nabla^2 \bar{\psi} + f}{\partial x} + v' \frac{\partial \nabla^2 \bar{\psi} + f}{\partial y} &= 0 \\ \left(\frac{\partial}{\partial t} + \bar{u} \frac{\partial}{\partial x} + \bar{v} \frac{\partial}{\partial y} \right) \nabla^2 \psi' - \frac{\partial \psi'}{\partial y} \frac{\partial \bar{\eta}}{\partial x} + \frac{\partial \psi'}{\partial x} \frac{\partial \bar{\eta}}{\partial y} &= 0. \end{aligned} \quad (C6)$$

To study Rossby wave propagation from the source to remote locations, the above equation needs to be cast, those equations must be written in spherical coordinates or other map projections another coordinate system to account for the Earth's spherical geometry. Here, we keep the use simple Cartesian coordinates and refer the readers to previous studies for the equation in the appropriate coordinates (e.g., Hoskins and Karoly, 1981; Li et al., 2015). This partial differential equation for ψ' has parameters that vary in space, which prevents us from solving it analytically. To obtain an approximate analytical solution, we assume the mean state varies much more slowly than the wave disturbances, thus treating the mean state as constant locally (WKB approximation), and also assume plane wave solutions in the form of:

$$890 \quad \psi' = A(T, X, Y, \mathcal{F}) \exp[i(kx + ly - \omega t)] \quad (C7)$$

where T , X , and Y are the local or wave-scale base-state coordinates with substantially smaller larger scales of variations than those for the mean-state waves (t, x, y). A is the amplitude, ω is the angular frequency, k is the zonal wavenumber, and l is the meridional wavenumber, and ω is the angular frequency. (Fig. C1). A is a function of time and space. By substituting equation ??C7 to equation C6 while ignoring the variations of the base state, we can obtain the following dispersion relation [those and following equations can be found in Karoly (1983), Takaya and Nakamura (2001), and Li et al. (2015)]:

$$895 \quad \omega = \bar{u}k + \bar{v}l + \frac{\bar{\eta}_x l - \bar{\eta}_y k}{k^2 + l^2} \quad (C8)$$

where we have used subscripts x and y to denote partial differentiation with respect to x and y . Rearranging equation C8, we define the total wavenumber

$$K^2 = k^2 + l^2 = \frac{\bar{\eta}_y k - \bar{\eta}_x l}{\bar{u}k + \bar{v}l - \omega}. \quad (C9)$$

900 For stationary waves with $\omega = 0$, the total wavenumber is

$$K^2 = k^2 + l^2 = \frac{\bar{\eta}_y k - \bar{\eta}_x l}{\bar{u}k + \bar{v}l}. \quad (\text{C10})$$




Figure C1. Schematics to clarify the wavenumbers and wavelengths with two example cases: (a) zonal and meridional wavelengths are similar, and (b) meridional wavelength is longer than the zonal wavelength

~

The x- and y-components of group velocity are obtained from the dispersion relationship (with the quotient rule of differentiation):

$$u_g = \frac{\partial \omega}{\partial k} = \bar{u} + \frac{((k^2 - l^2)\bar{\eta}_y - 2kl\bar{\eta}_x)}{K^4} \quad (\text{C11})$$

$$v_g = \frac{\partial \omega}{\partial l} = \bar{v} + \frac{2kl\bar{\eta}_y + (k^2 - l^2)\bar{\eta}_x}{K^4}.$$

905 **Equation C10** tells that wavenumbers k and l depend on themselves and the mean state. Therefore, as a wave packet travels through a varying mean state, its wavenumbers also change. Their time evolution The time evolution of wavenumbers is governed by the conservation of the number of waves (Whitham, 1960): accounting for the spatial variation of the base state (rather than ignoring it, as in the previous derivation of the dispersion relationship). Accordingly, the derivatives of ω with respect to the large-scale coordinates (X, Y) determine the time rate of change of the wavenumbers k and l as the wave packet moves along the group velocity.

910

$$\frac{d_g k}{dT} = -\frac{\partial \omega}{\partial X} = -k \frac{\partial \bar{u}}{\partial X} - l \frac{\partial \bar{v}}{\partial X} - \frac{1}{K^2} \left(l \frac{\partial \bar{\eta}_x}{\partial X} - k \frac{\partial \bar{\eta}_y}{\partial X} \right) \quad (\text{C12})$$

$$\frac{d_g l}{dT} = -\frac{\partial \omega}{\partial Y} = -k \frac{\partial \bar{u}}{\partial Y} - l \frac{\partial \bar{v}}{\partial Y} - \frac{1}{K^2} \left(l \frac{\partial \bar{\eta}_x}{\partial Y} - k \frac{\partial \bar{\eta}_y}{\partial Y} \right) \quad (\text{C13})$$

(these are same as equations 1 and 2, but repeated here.) Equations C12 and C13 show that the base state wind shear changes
 915 the shape and scale of the wave (i.e., k and l) as the wave travels at the group velocity.

If we assume that the background (mean) circulations are represented by the zonal mean zonal winds (constant over longitudes) and zero meridional winds, then the dispersion relationship is

$$\omega = \bar{u}k - \frac{\bar{\eta}_y k}{k^2 + l^2} \quad (\text{C14})$$

and the group velocities are given by:

$$\begin{aligned} u_g &= \bar{u} + \frac{2\bar{\eta}_y k^2}{K^2} \\ v_g &= \frac{2\bar{\eta}_y kl}{K^2}. \end{aligned} \quad (\text{C15})$$

For stationary waves $\omega = 0$ in this zonally uniform background case, we have a stationary wavenumber as

$$K_s^2 = \frac{\bar{\eta}_y}{\bar{u}}$$

, which is widely used in literature to depict Rossby waveguides (e.g., Hoskins and Ambrizzi, 1993; Henderson et al., 2017; Hoskins and W
 -the same as eqn. 4 in the main text. In this simpler case, the total wavenumber depends on only two quantities: the meridional
 925 gradient of the background absolute vorticity and the background zonal wind. When either is negative, K_s is an imaginary number. In this case, instead of oscillating in space, the wave solution becomes evanescent, decaying exponentially with distance (eqn. C7); hence, Rossby waves do not propagate over such a region.

C2 ~~W~~-vector

Based on the equations C2, C3, and C5, the horizontal component of the wave activity flux by TN01, or the are written as: For
 930 a zonally uniform background state, eqn C12 suggests $\frac{\partial \omega}{\partial X} = 0 = \frac{d_g k}{dT}$, so k remains the same as given at the initial time k_0 , while l (hence K_s) evolves as the wavepacket travels through the background state. Changes in l along the path are related to the meridional propagation of wave packets (l is the y-component of the wavenumber vector, Fig. C1). Writing l as:

$$l = \pm \sqrt{K_s^2 - k_0^2} \quad (\text{C16})$$

where $|V| = \sqrt{\bar{u}^2 + \bar{v}^2}$. This is the form of W-vector on the pressure coordinate (eqn. C5 in TN01). Here, the mean winds (\bar{u}
 935 and \bar{v}) are the time-mean geostrophic winds, we can see: 1) where K_s is large and $K_s \gg k_0$, l is real and large, thus Rossby waves can propagate meridionally with smaller meridional wavelengths, 2) where $K_s \approx k_0$, l becomes small, the wave fronts become oriented North-South, and the wave energy travels almost purely zonally (Fig. C1b), and the perturbation winds (and

the associated streamfunction: u' , v' and ψ') are the deviation of *geostrophic* winds from the mean winds. 3) where $K_s = k_0$, $l = 0$, and the wave cannot propagate any further meridionally, and turn back toward higher K_s (turning latitude). If a Rossby wave is excited within the local maximum of K_s with a sufficient latitudinal width, then the wave is trapped within the latitude band and propagates zonally since the strong gradients to the north and south refract back the waves.

In Fig. 7a, we apply this metric to each grid point, assuming that the metric K_s is locally applicable; this is commonly done and able to provide a qualitative picture of the preferred wave pathways (Hoskins and Ambrizzi, 1993). An example of turning latitude is the north/south of the mid-latitude jet over the Pacific, where K_s is becoming smaller toward the inhibited region. To the south of the subtropical jet, zonal winds are tending to zero, changing from westerly to easterly (Fig. 5a). This makes K_s increase toward infinity, or the wavelength tends toward zero, implying that the waves break and mix into the background flow. Also, the group velocity decreases toward zero with increasing K_s (eqn C15), thereby prohibiting wave propagation. This is called critical latitude.

C2 Sources of Rossby waves propagating to North America

950 To apply ray tracing in section 3.3, it is necessary to specify the locations of the wave sources. Following previous studies, we used lead/lag correlation maps of the ~~daily-mean-perturbation-meridional-winds~~ daily-mean meridional perturbation winds, $va200'$, using the same background state and the perturbation winds for the W-vector diagnosis. The base point is placed in one of the eight source regions: North Pacific, East Pacific, West Pacific, East Asian Monsoon, Indian Monsoon, Tibetan Plateau, Caspian Sea, and Red Sea. The time series of $va200'$ after area-averaging over the base location (indicated by the box in Fig. 955 C2) is correlated to the same variables at all the other grid points, varying the lags from -15 to +15 days.

Among the eight source locations examined, statistically significant correlations in the grid points over North America are found for the North Pacific and East Pacific (Fig. C2 a,b), suggesting that those two are the critical wave sources for stationary quasi-stationary Rossby waves traveling to North America. However, the figure indicates other *indirect* wave sources from which wave signals reach those two sources. For example, from the East Asian Monsoon region (Fig. C2 d), statistically significant signals first appear to its west. The signal becomes stronger over three days, such that a statistically significant correlation links the East Asian region to Europe with a lag of -3. Then the signal in the downwind direction becomes strong and significant, reaching the North Pacific source region and almost the West Coast of North America (lags 0 and 3). Meanwhile, the wave signals from the Caspian Sea reach ~~the East Asian region~~ East Asia. A dipole wave pattern first establishes six days earlier (from day 0) over the Caspian Sea and Europe (Fig. C2 g, lag -6). Three days later (lag -3), another negative phase 960 appears to the east, then another positive phase over China, reaching the East Asian Monsoon source.

Sardeshmukh and Hoskins (1988) suggested another diagnostic for Rossby wave sources. They linearized equation C1 to solve for the time tendency of perturbation absolute vorticity to identify the source terms. Doing so, they suggested to partition the winds into rotational (\mathbf{v}_Ψ) and divergent (\mathbf{v}_χ) components when diagnosing Rossby wave sources, particularly in the tropics, so that one does not overlook the contribution of vorticity advection by the divergent winds:

$$970 \quad \frac{\partial \eta}{\partial t} + \mathbf{v}_\Psi \cdot \nabla \eta = -\mathbf{v}_\chi \cdot \nabla \eta - \eta(\nabla \cdot \mathbf{v}_\chi), \quad (\text{C17})$$

where advection by the rotational and divergent winds is separated, and the latter is moved to the right-hand side as a forcing. Linearizing the equation, we have:

$$\begin{aligned} \frac{\partial \eta'}{\partial t} + \bar{\mathbf{v}}_\Psi \cdot \nabla \eta' + \mathbf{v}'_\Psi \cdot \nabla \bar{\eta} &= -\nabla \cdot (\mathbf{v}'_\chi \bar{\eta}) - \nabla \cdot (\bar{\mathbf{v}}_\chi \eta') \\ &= -\bar{\eta}(\nabla \cdot \mathbf{v}'_\chi) - \mathbf{v}'_\chi \cdot \nabla \bar{\eta} - \eta'(\nabla \cdot \bar{\mathbf{v}}_\chi) - \bar{\mathbf{v}}_\chi \cdot \nabla \eta' \end{aligned} \quad (\text{C18})$$

The left-hand side is the change of the perturbation absolute vorticity following the rotational winds. The right-hand side now 975 has four terms: the first and third are stretching of the mean and perturbation vorticities by the perturbation and mean divergence, respectively. The second and fourth terms are the advection of the mean and perturbation vorticity by the perturbation and mean divergent winds, respectively. The right-hand-side terms are referred to as the Rossby Wave Source (RWS) (S'). Note that here RWS is defined for the perturbation vorticity, not the mean vorticity. Lin (2009) found the first two terms in the second line in eqn. C18 dominate other source terms. These are shown in Fig. C3, showing that all the source regions 980 mentioned above exhibit strong magnitudes of either source term.




Figure C2. Lead/lag correlation between the band-passed perturbation meridional winds at the 200 hPa level (va_{200}') averaged over the source regions (denoted by the blue rectangles) and all the other grid points. Each row represents different source locations: (a) North Pacific, (b) East Pacific, (c) West Pacific, (d) East Asian Monsoon, (e) Indian Monsoon, (f) Tibetan Plateau, (g) Caspian Sea, and (h) Red Sea.




Figure C3. The first (a) and second (b) terms in the Rossby wave source (equation C18) calculated from ERA Interim

C3 W vector

Based on the equations C2, C3, and C5, the horizontal component of the wave activity flux by TN01, or the “W-vector” are written as:

$$\begin{aligned} \mathbf{W}_x &= \frac{1}{2|\mathbf{V}|} \left[\bar{u} \left(\frac{\partial \psi'^2}{\partial x} - \psi' \frac{\partial^2 \psi'}{\partial x^2} \right) + \bar{v} \left(\frac{\partial \psi'}{\partial x} \frac{\partial \psi'}{\partial y} - \psi' \frac{\partial^2 \psi'}{\partial x \partial y} \right) \right] + \frac{\bar{u}}{|\mathbf{V}|} C_p M \\ \mathbf{W}_y &= \frac{1}{2|\mathbf{V}|} \left[\bar{u} \left(\frac{\partial \psi'}{\partial x} \frac{\partial \psi'}{\partial y} - \psi' \frac{\partial^2 \psi'}{\partial x \partial y} \right) + \bar{v} \left(\frac{\partial \psi'^2}{\partial y} - \psi' \frac{\partial^2 \psi'}{\partial y^2} \right) \right] + \frac{\bar{v}}{|\mathbf{V}|} C_p M \end{aligned} \quad (\text{C19})$$

985 where $|\mathbf{V}| = \sqrt{\bar{u}^2 + \bar{v}^2}$ and C_p is the wave phase speed. The last term represents the wave activity flux at the phase velocity along the background wind vector. This is the form of W-vector on the pressure coordinate (eqn. C5 in TN01). Here, the mean winds (\bar{u} and \bar{v}) are the time-mean *geostrophic* winds, and the perturbation winds (and the associated streamfunction: u' , v' and ψ') are the deviation of *geostrophic* winds from the mean winds. We set $C_p = 0$ for quasi-stationary waves.

990 *Author contributions.* KS performed the analysis/plots and wrote the article. SM processed some of the model data and helped KS with coding and analysis. LRL supervised the work by providing general guidance for the article’s direction and structure. MB provided technical information about the RegCM4 and WRF model configurations in the NA-CORDEX archive. SM, LRL, MB, and RM guided science questions and provided feedback on the analyses. ZC and CCC provided the technical background and literature on Rossby waves and their impact on extreme events. YL provided the ray tracing code and provided guidance on its use and interpretation. All authors reviewed and provided feedback on the article.

995 *Competing interests.* The authors declare that they have no conflict of interest.

1000 *Acknowledgements.* This work is supported by the Department of Energy Office of Science award number DE-SC0016605, “A Framework for Improving Analysis and Modeling of Earth System and Intersectoral Dynamics at Regional Scales.” [YL was supported by National Natural Science Foundation of China \(Grants 42175080\).](#) We acknowledge the general direction and programmatic support from Dr. Linda Mearns, who deceased on January 23, 2025. [We thank Dr. Sandro Lubis for the helpful discussion of wave activity flux. We also thank two anonymous referees for significantly improving the manuscript, in particular, the communication and accuracy of Rossby wave theory and diagnostics.](#) This research used resources of the National Energy Research Scientific Computing Center, a DOE Office of Science User Facility supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231 using NERSC awards BER-ERCAP0024296 and BER-ERCAP0032096.

References

- 1005 Abatzoglou, J. T. and Magnusdottir, G.: Planetary Wave Breaking and Nonlinear Reflection: Seasonal Cycle and Interannual Variability, *Journal of Climate*, 19, 6139–6152, <https://doi.org/10.1175/JCLI3968.1>, 2006.
- Alexandru, A., de Elia, R., Laprise, R., Separovic, L., and Biner, S.: Sensitivity Study of Regional Climate Model Simulations to Large-Scale Nudging Parameters, *Monthly Weather Review*, 137, 1666–1686, <https://doi.org/10.1175/2008MWR2620.1>, 2009.
- Ambrizzi, T., Hoskins, B. J., and Hsu, H.-H.: Rossby Wave Propagation and Teleconnection Patterns in the Austral Winter, *Journal of the Atmospheric Sciences*, 52, 3661–3672, [https://doi.org/10.1175/1520-0469\(1995\)052<3661:RWPATP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<3661:RWPATP>2.0.CO;2), 1995.
- 1010 Bacmeister, J. T., Wehner, M. F., Neale, R. B., Gettelman, A., Hannay, C., Lauritzen, P. H., Caron, J. M., and Truesdale, J. E.: Exploratory High-Resolution Climate Simulations using the Community Atmosphere Model (CAM), *Journal of Climate*, 27, 3073–3099, <https://doi.org/10.1175/JCLI-D-13-00387.1>, 2014a.
- Bacmeister, J. T., Wehner, M. F., Neale, R. B., Gettelman, A., Hannay, C., Lauritzen, P. H., Caron, J. M., and Truesdale, J. E.: Exploratory High-Resolution Climate Simulations using the Community Atmosphere Model (CAM), *Journal of Climate*, 27, 3073–3099, <https://doi.org/10.1175/JCLI-D-13-00387.1>, 2014b.
- 1015 Balaji, V., Boville, B., Cheung, S., Collins, N., Cruz, C., Silva, A., Deluca, C., Fainchtein, R. D., Eaton, B., Hallberg, B., Henderson, T., Hill, C., Iredell, M., Jacob, R., Jones, P., Kluzek, E., Kauffman, B., Larson, J., Li, P., Liu, F., Michalakes, J., Murphy, S., Neckels, D., Kuinghttons, R. O., Oehmke, B., Panaccione, C., Rosinski, J., Sawyer, W., Schwab, E., Smithline, S., Spector, W., Stark, D., Suarez, M., Swift, S., Theurich, G., Trayanov, A., Vasquez, S., Wolfe, J., Yang, W., Young, M., and Zaslavsky, L.: Earth System Modeling Framework ESMF Reference Manual for Fortran Version 7.1.0r, Tech. rep., The Earth System Modeling Framework, 2018.
- 1020 Barriopedro, D., García-Herrera, R., Ordóñez, C., Miralles, D. G., and Salcedo-Sanz, S.: Heat Waves: Physical Understanding and Scientific Challenges, *Reviews of Geophysics*, 61, e2022RG000780, <https://doi.org/10.1029/2022RG000780>, <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2022RG000780>, 2023.
- 1025 Bell, G. D. and Janowiak, J. E.: Atmospheric Circulation Associated With the Midwest Floods of 1993, *Bulletin of the American Meteorological Society*, 76, 681–696, https://journals.ametsoc.org/view/journals/bams/76/5/1520-0477_1995_076_0681_acawtm_2_0_co_2.xml, 1995.
- Bogenschutz, P. A., Gettelman, A., Hannay, C., Larson, V. E., Neale, R. B., Craig, C., and Chen, C.-C.: The path to CAM6: coupled simulations with CAM5.4 and CAM5.5, *Geoscientific Model Development*, 11, 235–255, <https://doi.org/10.5194/gmd-11-235-2018>, 2018.
- 1030 Branstator, G.: Long-Lived Response of the Midlatitude Circulation and Storm Tracks to Pulses of Tropical Heating, *Journal of Climate*, 27, 8809–8826, <https://doi.org/10.1175/JCLI-D-14-00312.1>, 2014.
- Branstator, G. and Teng, H.: Tropospheric Waveguide Teleconnections and Their Seasonality, *Journal of the Atmospheric Sciences*, 74, 1513–1532, <https://doi.org/10.1175/JAS-D-16-0305.1>, 2017.
- Bukovsky, M. S. and Mearns, L. O.: Regional climate change projections from NA-CORDEX and their relation to climate sensitivity, *Climatic Change*, 162, 645–665, <https://doi.org/10.1007/s10584-020-02835-x>, 2020.
- 1035 Bukovsky, M. S., McCrary, R. R., Seth, A., and Mearns, L. O.: A mechanistically credible, poleward shift in warm-season precipitation projected for the U.S. Southern Great Plains?, *Journal of Climate*, 30, 8275–8298, <https://doi.org/10.1175/JCLI-D-16-0316.1>, 2017.
- Bumbaco, K. A., Dello, K. D., and Bond, N. A.: History of Pacific Northwest Heat Waves: Synoptic Pattern and Trends, *Journal of Applied Meteorology and Climatology*, 52, 1618–1631, <https://doi.org/10.1175/JAMC-D-12-094.1>, 2013.

- 1040 Castro, C. L., Pielke, R. A., and Leoncini, G.: Dynamical downscaling: Assessment of value retained and added using the Regional Atmospheric Modeling System (RAMS), *Journal of Geophysical Research D: Atmospheres*, 110, 1–21, <https://doi.org/10.1029/2004JD004721>, 2005.
- Castro, C. L., Pielke, R. A., Adegoke, J. O., Schubert, S. D., and Pegion, P. J.: Investigation of the summer climate of the contiguous United States and Mexico using the Regional Atmospheric Modeling System (RAMS). Part II: Model climate variability, *Journal of Climate*, 20, 3866–3887, <https://doi.org/10.1175/JCLI4212.1>, 2007.
- 1045 Castro, C. L., Chang, H. I., Dominguez, F., Carrillo, C., Schemm, J. K., and Juang, H. M. H.: Can a regional climate model improve the ability to forecast the North American monsoon?, *Journal of Climate*, 25, 8212–8237, <https://doi.org/10.1175/JCLI-D-11-00441.1>, 2012.
- Chang, C.-C., Lubis, S. W., Balaguru, K., Leung, L. R., Hagos, S. M., and Klotzbach, P. J.: An Extratropical Pathway for the Madden–Julian Oscillation’s Influence on North Atlantic Tropical Cyclones, *Journal of Climate*, 36, 8539–8559, <https://doi.org/10.1175/JCLI-D-23-0251.1>, 2023a.
- 1050 Chang, C.-C., Wang, Z., Ting, M., and Ming, Z.: Summertime Subtropical Stationary Waves in the Northern Hemisphere: Variability, Forcing Mechanisms, and Impacts on Tropical Cyclone Activity, *Journal of Climate*, 36, 753–773, <https://doi.org/10.1175/JCLI-D-22-0233.1>, 2023b.
- Chang, C.-C., Zhao, M., Lubis, S. W., Chen, Z., Balaguru, K., Hagos, S., Leung, L. R., and Zhou, W.: Are North Atlantic Tropical Cyclones Modulated by the Madden–Julian Oscillation in HighResMIP AGCMs?, *Journal of Climate*, 38, 2891–2916, <https://doi.org/10.1175/JCLI-D-24-0509.1>, 2025a.
- 1055 Chang, C.-C., Zhao, M., Lubis, S. W., Chen, Z., Balaguru, K., Hagos, S., Leung, L. R., and Zhou, W.: Are North Atlantic Tropical Cyclones Modulated by the Madden–Julian Oscillation in HighResMIP AGCMs?, *Journal of Climate*, 38, 2891–2916, <https://doi.org/10.1175/JCLI-D-24-0509.1>, 2025b.
- 1060 Chang, H.-i., Castro, C. L., Carrillo, C. M., and Dominguez, F.: The more extreme nature of U.S. warm season climate in the recent observational record and two “well-performing” dynamically downscaled CMIP3 models, *Journal of Geophysical Research: Atmospheres*, 120, 8244–8263, <https://doi.org/10.1002/2015JD023333>, 2015.
- Chen, T.-C. and Chen, J.-M.: On the Maintenance of Stationary Eddies in Terms of the Streamfunction Budget Analysis, *Journal of the Atmospheric Sciences*, 47, 2818–2824, https://journals.ametsoc.org/view/journals/atsc/47/23/1520-0469_1990_047_2818_otmose_2_0_co_2.xml, 1990.
- 1065 Chen, X., Leung, L. R., Wigmosta, M., and Richmond, M.: Impact of Atmospheric Rivers on Surface Hydrological Processes in Western U.S. Watersheds, *Journal of Geophysical Research: Atmospheres*, 124, 8896–8916, <https://doi.org/10.1029/2019JD030468>, 2019.
- Chen, Z., Lu, J., Chang, C. C., Lubis, S. W., and Leung, L. R.: Projected increase in summer heat-dome-like stationary waves over North-western North America, *npj Climate and Atmospheric Science*, 6, 1–10, <https://doi.org/10.1038/s41612-023-00511-2>, 2023.
- 1070 Choi, N. and Stan, C.: Large-Scale Surface Air Temperature Bias in Summer over the CONUS and Its Relationship to Tropical Central Pacific Convection in the UFS Prototype 8, *Journal of Climate*, 38, 129, <https://doi.org/10.1175/JCLI-D-24-0078.1>, 2024.
- CORDEX: Experiment Protocol CORDEX-CMIP5 – RCMs, <https://cordex.org/experiment-guidelines/cordex-cmip5/experiment-protocol-cordex-cmip5-rcms/>, 2009.
- Coumou, D., Petoukhov, V., Rahmstorf, S., Petri, S., and Schellnhuber, H. J.: Quasi-resonant circulation regimes and hemispheric synchronization of extreme weather in boreal summer, *Proceedings of the National Academy of Sciences of the United States of America*, 111, 12 331–12 336, <https://doi.org/10.1073/pnas.1412797111>, 2014.
- Dai, A.: Precipitation Characteristics in Eighteen Coupled Climate Models, *Journal of Climate*, 19, 4605–4630, 2006.

- Davies, H. C.: A lateral boundary formulation for multi-level prediction models, *Quarterly Journal of the Royal Meteorological Society*, 102, 405–418, 1976.
- 1080 Denis, B., Côté, J., and Laprise, R.: Spectral decomposition of two-dimensional atmospheric fields on limited-area domains using the discrete cosine transform (DCT), *Monthly Weather Review*, 130, 1812–1829, [https://doi.org/10.1175/1520-0493\(2002\)130<1812:SDOTDA>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<1812:SDOTDA>2.0.CO;2), 2002a.
- Denis, B., Laprise, R., Caya, D., and Côté, J.: Downscaling ability of one-way nested regional climate models: the Big-Brother Experiment, *Climate Dynamics*, 18, 627–646, <https://doi.org/10.1007/s00382-001-0201-0>, 2002b.
- 1085 Denis, B., Laprise, R., and Caya, D.: Sensitivity of a regional climate model to the resolution of the lateral boundary conditions, *Climate Dynamics*, 20, 107–126, <https://doi.org/10.1007/s00382-002-0264-6>, 2003.
- Diez-Sierra, J., Iturbide, M., Gutiérrez, J. M., Fernández, J., Milovac, J., Cofiño, A. S., Cimadevilla, E., Nikulin, G., Levvasseur, G., Kjellström, E., Bülow, K., Horányi, A., Brookshaw, A., García-Díez, M., Pérez, A., Baño-Medina, J., Ahrens, B., Alias, A., Ashfaq, M., Bukovsky, M., Buonomo, E., Cabos, W. D., Caluwaerts, S., Chou, S. C., Christensen, O. B., Ciarlò, J. M., Coppola, E., Corre, L.,
- 1090 Demory, M.-E., Djurdjevic, V., Evans, J. P., Fealy, R., Feldmann, H., Jacob, D., Jayanarayanan, S., Katzfey, J., Keuler, K., Kittel, C., Kurnaz, M. L., Laprise, R., Lionello, P., McGinnis, S., Mercogliano, P., Nabat, P., Öñol, B., Ozturk, T., Panitz, H.-J., Paquin, D., Pieczka, I., Raffaele, F., Remedio, A. R., Scinocca, J., Sevault, F., Somot, S., Steger, C., Tangang, F., Teichmann, C., Termonia, P., Thatcher, M., Torma, C., van Meijgaard, E., Vautard, R., Warrach-Sagi, K., Winger, K., and Zittis, G.: CORDEX model component description, <https://zenodo.org/records/6553526>, 2022a.
- 1095 Diez-Sierra, J., Iturbide, M., Gutiérrez, J. M., Fernández, J., Milovac, J., Cofiño, A. S., Cimadevilla, E., Nikulin, G., Levvasseur, G., Kjellström, E., Bülow, K., Horányi, A., Brookshaw, A., García-Díez, M., Pérez, A., Baño-Medina, J., Ahrens, B., Alias, A., Ashfaq, M., Bukovsky, M., Buonomo, E., Caluwaerts, S., Chou, S. C., Christensen, O. B., Ciarlò, J. M., Coppola, E., Corre, L., Demory, M.-E., Djurdjevic, V., Evans, J. P., Fealy, R., Feldmann, H., Jacob, D., Jayanarayanan, S., Katzfey, J., Keuler, K., Kittel, C., Kurnaz, M. L., Laprise, R., Lionello, P., McGinnis, S., Mercogliano, P., Nabat, P., Öñol, B., Ozturk, T., Panitz, H.-J., Paquin, D., Pieczka, I., Raffaele, F., Remedio, A. R., Scinocca, J., Sevault, F., Somot, S., Steger, C., Tangang, F., Teichmann, C., Termonia, P., Thatcher, M., Torma, C.,
- 1100 Meijgaard, E. v., Vautard, R., Warrach-Sagi, K., Winger, K., and Zittis, G.: The Worldwide C3S CORDEX Grand Ensemble: A Major Contribution to Assess Regional Climate Change in the IPCC AR6 Atlas, *Bulletin of the American Meteorological Society*, 103, E2804–E2826, <https://doi.org/10.1175/BAMS-D-22-0111.1>, 2022b.
- Dimitrijevic, M. and Laprise, R.: Validation of the nesting technique in a regional climate model and sensitivity tests to the resolution of the lateral boundary conditions during summer, *Climate Dynamics*, 25, 555–580, <https://doi.org/10.1007/s00382-005-0023-6>, 2005.
- 1105 Ding, Q. and Wang, B.: Circumglobal Teleconnection in the Northern Hemisphere Summer, *Journal of Climate*, 18, 3483–3505, <https://doi.org/10.1175/JCLI3473.1>, 2005.
- Dirmeyer, P. A. and Halder, S.: Application of the Land–Atmosphere Coupling Paradigm to the Operational Coupled Forecast System, Version 2 (CFSv2), *Journal of Hydrometeorology*, 18, 85–108, <https://doi.org/10.1175/JHM-D-16-0064.1>, 2017.
- 1110 Elguindi, N., Bi, X., Giorgi, F., Nagarajan, B., Pal, J. S., Solmon, F., Rauscher, S. A., Zakey, A., O’Brien, T. A., Nogherotto, R., and Giuliani, G.: Regional Climate Model RegCM Reference Manual Version 4.7, Tech. rep., The Abdus Salam International Centre for Theoretical Physics, Trieste, Italy, issue: January, 2017.
- Eyring, V., Bock, L., Lauer, A., Righi, M., Schlund, M., Andela, B., Arnone, E., Bellprat, O., Brötz, B., Caron, L.-P., Carvalho, N., Cionni, I., Cortesi, N., Crezee, B., Davin, E. L., Davini, P., Debeire, K., de Mora, L., Deser, C., Docquier, D., Earnshaw, P., Ehbrecht, C., Gier, B. K., Gonzalez-Reviriego, N., Goodman, P., Hagemann, S., Hardiman, S., Hassler, B., Hunter, A., Kadow, C., Kindermann, S., Koirala,
- 1115

- S., Koldunov, N., Lejeune, Q., Lembo, V., Lovato, T., Lucarini, V., Massonnet, F., Müller, B., Pandde, A., Pérez-Zanón, N., Phillips, A., Predoi, V., Russell, J., Sellar, A., Serva, F., Stacke, T., Swaminathan, R., Torralba, V., Vegas-Regidor, J., von Hardenberg, J., Weigel, K., and Zimmermann, K.: Earth System Model Evaluation Tool (ESMValTool) v2.0 – an extended set of large-scale diagnostics for quasi-operational and comprehensive evaluation of Earth system models in CMIP, *Geoscientific Model Development*, 13, 3383–3438, <https://doi.org/10.5194/gmd-13-3383-2020>, 2020.
- 1120 for Medium-Range Weather Forecasts, E. C.: ERA-Interim Project, <https://doi.org/10.5065/D6CR5RD9>, 2009.
- Gao, Y., Leung, L. R., Zhao, C., and Hagos, S.: Sensitivity of U.S. summer precipitation to model resolution and convective parameterizations across gray zone resolutions, *Journal of Geophysical Research*, 122, 2714–2733, <https://doi.org/10.1002/2016JD025896>, 2017.
- Garfinkel, C. I., Chen, W., Li, Y., Schwartz, C., Yadav, P., and Domeisen, D.: The Winter North Pacific Teleconnection in Response to ENSO and the MJO in Operational Subseasonal Forecasting Models Is Too Weak, *Journal of Climate*, <https://doi.org/10.1175/JCLI-D-22-0179.1>, 2022.
- 1125 Giorgi, F. and Anyah, R. O.: The road towards RegCM4, *Climate Research*, 52, 3–6, <https://doi.org/10.3354/cr01089>, 2012.
- Giorgi, F., Marinucci, M. R., Bates, G. T., and De Canio, G.: Development of a Second-Generation Regional Climate Model (RegCM2). Part II: Convective Processes and Assimilation of Lateral Boundary Conditions, *Monthly Weather Review*, 121, 2814–2832, [https://doi.org/10.1175/1520-0493\(1993\)121<2814:DOASGR>2.0.CO;2](https://doi.org/10.1175/1520-0493(1993)121<2814:DOASGR>2.0.CO;2), 1993.
- 1130 Giorgi, F., Coppola, E., Solmon, F., Mariotti, L., Sylla, M. B., Bi, X., Elguindi, N., Diro, G. T. G., Nair, V., Giuliani, G., Turuncoglu, U. U. U., Cozzini, S., Güttler, I., O'Brien, T., Tawfik, A., Shalaby, A., Zakey, A., Steiner, A., Stordal, F., Sloan, L., and Brankovic, C.: RegCM4: model description and preliminary tests over multiple CORDEX domains, *Climate Research*, 52, 7–29, <https://doi.org/10.3354/cr01018>, 2012.
- 1135 Goldenson, N., Leung, L. R., Mearns, L. O., Pierce, D. W., Reed, K. A., Simpson, I. R., Ullrich, P., Krantz, W., Hall, A., Jones, A., and Rahimi, S.: Use-Inspired, Process-Oriented GCM Selection: Prioritizing Models for Regional Dynamical Downscaling, *Bulletin of the American Meteorological Society*, 104, E1619–E1629, <https://doi.org/10.1175/BAMS-D-23-0100.1>, 2023.
- Grell, G. A., Dudhia, J., and Stauffer, D. R.: A description of the fifth-generation Penn State/NCAR Mesoscale Model (MM5), Tech. rep., University Corporation for Atmospheric Research, Boulder, CO, ISSN 00424625, <https://doi.org/10.5065/D60Z716B>, publication Title: NCAR/TN-398+STR Volume: 115 Issue: 10, 1994.
- 1140 Harvey, B. J., Cook, P., Shaffrey, L. C., and Schiemann, R.: The Response of the Northern Hemisphere Storm Tracks and Jet Streams to Climate Change in the CMIP3, CMIP5, and CMIP6 Climate Models, *Journal of Geophysical Research: Atmospheres*, 125, 1–10, <https://doi.org/10.1029/2020JD032701>, 2020.
- Henderson, S. A., Maloney, E. D., and Son, S.-W.: Madden–Julian Oscillation Pacific Teleconnections: The Impact of the Basic State and MJO Representation in General Circulation Models, *Journal of Climate*, 30, 4567–4587, <https://doi.org/10.1175/JCLI-D-16-0789.1>, 2017.
- 1145 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 1150 Holman, K. D., Lorenz, D. J., and Notaro, M.: Influence of the Background State on Rossby Wave Propagation into the Great Lakes Region Based on Observations and Model Simulations, *Journal of Climate*, 27, 9302–9322, <https://doi.org/10.1175/JCLI-D-13-00758.1>, 2014.

- Holton, J. R.: An Introduction to Dynamic Meteorology, International Geophysics Series, Elsevier Academic Press, fourth edition edn., ISBN 978-0-12-354015-7, <https://shop.elsevier.com/books/an-introduction-to-dynamic-meteorology/holton/978-0-12-354015-7>, 2004.
- 1155 Hoskins, B. and Woollings, T.: Persistent Extratropical Regimes and Climate Extremes, *Current Climate Change Reports*, 1, 115–124, <https://doi.org/10.1007/s40641-015-0020-8>, 2015.
- Hoskins, B. J. and Ambrizzi, T.: Rossby wave propagation on a realistic longitudinally varying flow, *Journal of the Atmospheric Sciences*, 50, 1661–1671, 1993.
- 1160 Hoskins, B. J. and Karoly, D. J.: The steady linear response of a spherical atmosphere to thermal and orographic forcing, *Journal of the Atmospheric Sciences*, 38, 1179–1196, [http://journals.ametsoc.org/doi/abs/10.1175/1520-0469\(1981\)038<1179:TSLROA>2.0.CO;2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0469(1981)038<1179:TSLROA>2.0.CO;2), 1981.
- Hu, X.-M., Xue, M., McPherson, R. A., Martin, E., Rosendahl, D. H., and Qiao, L.: Precipitation Dynamical Downscaling Over the Great Plains, *Journal of Advances in Modeling Earth Systems*, 10, 421–447, <https://doi.org/10.1002/2017MS001154>, 2018.
- Huang, C. S. Y. and Nakamura, N.: Local wave activity budgets of the wintertime Northern Hemisphere: Implication for the Pacific and Atlantic storm tracks, *Geophysical Research Letters*, 44, 5673–5682, <https://doi.org/10.1002/2017GL073760>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/2017GL073760](https://onlinelibrary.wiley.com/doi/pdf/10.1002/2017GL073760), 2017.
- 1165 Imberger, M., Guo Larsén, X., Davis, N., and Du, J.: Approaches toward improving the modelling of midlatitude cyclones entering at the lateral boundary corner in the limited area WRF model, *Quarterly Journal of the Royal Meteorological Society*, 146, 3225–3244, <https://doi.org/10.1002/qj.3843>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/qj.3843](https://onlinelibrary.wiley.com/doi/pdf/10.1002/qj.3843), 2020.
- 1170 Jin, F. and Hoskins, B. J.: The direct response to tropical heating in a baroclinic atmosphere, *Journal of Atmospheric Sciences*, 52, 307 – 319, [https://doi.org/10.1175/1520-0469\(1995\)052<0307:TDRTH>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<0307:TDRTH>2.0.CO;2), place: Boston MA, USA, 1995.
- Joseph, P. V. and Srinivasan, J.: Rossby waves in May and the Indian summer monsoon rainfall, *Tellus A*, 51, 854–864, <https://doi.org/10.1034/j.1600-0870.1999.00021.x>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1034/j.1600-0870.1999.00021.x](https://onlinelibrary.wiley.com/doi/pdf/10.1034/j.1600-0870.1999.00021.x), 1999.
- 1175 Kang, I.-S. and Held, I. M.: Linear and Nonlinear Diagnostic Models of Stationary Eddies in the Upper Troposphere during Northern Summer, *Journal of the Atmospheric Sciences*, 43, 3045–3057, https://journals.ametsoc.org/view/journals/atsc/43/24/1520-0469_1986_043_3045_landmo_2_0_co_2.xml, 1986.
- Karoly, D. J.: Rossby wave propagation in a barotropic atmosphere, *Dynamics of Atmospheres and Oceans*, 7, 111–125, [https://doi.org/10.1016/0377-0265\(83\)90013-1](https://doi.org/10.1016/0377-0265(83)90013-1), 1983.
- 1180 Kornhuber, K., Petoukhov, V., Petri, S., Rahmstorf, S., and Coumou, D.: Evidence for wave resonance as a key mechanism for generating high-amplitude quasi-stationary waves in boreal summer, *Climate Dynamics*, 49, 1961–1979, <https://doi.org/10.1007/s00382-016-3399-6>, 2017.
- Laprise, R., de Elía, R., Caya, D., Biner, S., Lucas-Picher, P., Diaconescu, E., Leduc, M., Alexandru, a., and Separovic, L.: Challenging some tenets of Regional Climate Modelling, *Meteorology and Atmospheric Physics*, 100, 3–22, <https://doi.org/10.1007/s00703-008-0292-9>, 2008.
- 1185 Lau, K.-M. and Weng, H.: Recurrent Teleconnection Patterns Linking Summertime Precipitation Variability over East Asia and North America, *Journal of the Meteorological Society of Japan. Ser. II*, 80, 1309–1324, <https://doi.org/10.2151/jmsj.80.1309>, 2002.
- Leung, L. R., Ringer, T. D., Collins, W. D., Taylor, M. A., Ashfaq, M., and Framework, A. H. E.: A hierarchical evaluation of regional climate simulations, *EOS*, 94, 297–298, <https://doi.org/10.1002/2013EO340001>, 2013.
- 1190 Li, Y.: Enhanced Poleward Propagation of Barotropic Rossby Waves by the Free-Surface Divergent Effect, *Scientific Online Letters on the Atmosphere*, 16, 92–96, <https://doi.org/10.2151/sola.2020-016>, 2020.

- Li, Y., Li, J., Jin, F. F., and Zhao, S.: Interhemispheric propagation of stationary rossby waves in a horizontally nonuniform background flow, *Journal of the Atmospheric Sciences*, 72, 3233–3256, <https://doi.org/10.1175/JAS-D-14-0239.1>, 2015.
- 1195 Li, Y., Feng, J., Li, J., and Zhao, S.: The circle diagram in the group velocity domain for Rossby wave under the horizontally non-uniform flow, *Scientific Online Letters on the Atmosphere*, 14, 121–125, <https://doi.org/10.2151/SOLA.2018-021>, 2018.
- Li, Y., Feng, J., Li, J., and Hu, A.: Equatorial windows and barriers for stationary Rossby wave propagation, *Journal of Climate*, 32, 6117–6135, <https://doi.org/10.1175/JCLI-D-18-0722.1>, 2019.
- Lin, H.: Global Extratropical Response to Diabatic Heating Variability of the Asian Summer Monsoon, *Journal of the Atmospheric Sciences*, 66, 2697–2713, <https://doi.org/10.1175/2009JAS3008.1>, 2009.
- 1200 Liu, C., Ikeda, K., Rasmussen, R., Barlage, M., Newman, A. J., Prein, A. F., Chen, F., Chen, L., Clark, M., Dai, A., Dudhia, J., Eidhammer, T., Gochis, D., Gutmann, E., Kurkute, S., Li, Y., Thompson, G., and Yates, D.: Continental-scale convection-permitting modeling of the current and future climate of North America, *Climate Dynamics*, 49, 71–95, <https://doi.org/10.1007/s00382-016-3327-9>, 2017.
- Lubis, S. W., Chen, Z., Lu, J., Hagos, S., Chang, C.-C., and Leung, L. R.: Enhanced Pacific Northwest heat extremes and wildfire risks induced by the boreal summer intraseasonal oscillation, *npj Climate and Atmospheric Science*, 7, 232, <https://doi.org/10.1038/s41612-024-00766-3>, 2024.
- 1205 Luo, F., Selten, F., Wehrli, K., Kornhuber, K., Le Sager, P., May, W., Reerink, T., Seneviratne, S. I., Shiogama, H., Tokuda, D., Kim, H., and Coumou, D.: Summertime Rossby waves in climate models: substantial biases in surface imprint associated with small biases in upper-level circulation, *Weather and Climate Dynamics*, 3, 905–935, <https://doi.org/10.5194/wcd-3-905-2022>, 2022.
- Ma, H. Y., Klein, S. A., Xie, S., Zhang, C., Tang, S., Tang, Q., Morcrette, C. J., Van Weverberg, K., Petch, J., Ahlgrimm, M., Berg, L. K., 1210 Cheruy, F., Cole, J., Forbes, R., Gustafson, W. I., Huang, M., Liu, Y., Merryfield, W., Qian, Y., Roehrig, R., and Wang, Y. C.: CAUSES: On the Role of Surface Energy Budget Errors to the Warm Surface Air Temperature Error Over the Central United States, *Journal of Geophysical Research: Atmospheres*, 123, 2888–2909, <https://doi.org/10.1002/2017JD027194>, 2018.
- Manola, I., Selten, F., De Vries, H., and Hazeleger, W.: "Waveguidability" of idealized jets, *Journal of Geophysical Research Atmospheres*, 118, 10,432–10,440, <https://doi.org/10.1002/jgrd.50758>, 2013.
- 1215 McGinnis, S. and Mearns, L.: Building a climate service for North America based on the NA-CORDEX data archive, *Climate Services*, 22, 100233, <https://doi.org/10.1016/j.cliser.2021.100233>, 2021.
- Mearns, L. O., McGinnis, S., Korytina, D., Scinocca, J. F., Kharin, S., Jiao, Y., Qian, M., Lazare, M., Winger, K., Christensen, O. B., Nikulin, G., Arritt, R. W., Herzmann, D., Bukovsky, M. S., Chang, H.-I., Castro, C., Frigon, A., and Gutowski, W. J. J.: The NA-CORDEX dataset, version 1.0., <https://doi.org/https://doi.org/10.5065/D6SJ1JCH>, place: Boulder, Colorado, 2017.
- 1220 Miguez-Macho, G., Stenchikov, G. L., and Robock, A.: Spectral nudging to eliminate the effects of domain position and geometry in regional climate model simulations, *Journal of Geophysical Research D: Atmospheres*, 109, 1–15, <https://doi.org/10.1029/2003JD004495>, 2004.
- Morcrette, C. J., Van Weverberg, K., Ma, H. Y., Ahlgrimm, M., Bazile, E., Berg, L. K., Cheng, A., Cheruy, F., Cole, J., Forbes, R., Gustafson, W. I., Huang, M., Lee, W. S., Liu, Y., Mellul, L., Merryfield, W. J., Qian, Y., Roehrig, R., Wang, Y. C., Xie, S., Xu, K. M., 1225 Zhang, C., Klein, S., and Petch, J.: Introduction to CAUSES: Description of Weather and Climate Models and Their Near-Surface Temperature Errors in 5 day Hindcasts Near the Southern Great Plains, *Journal of Geophysical Research: Atmospheres*, 123, 2655–2683, <https://doi.org/10.1002/2017JD027199>, 2018.
- Nakamura, N. and Huang, C. S.: Atmospheric blocking as a traffic jam in the jet stream, *Science*, 361, 42–47, <https://doi.org/10.1126/science.aat0721>, 2018.
- NCAR: The NCAR Command Language, <https://doi.org/10.5065/D6WD3XH5>, 2017.

- 1230 Neduhal, V., Žagar, N., Lunkeit, F., Polichtchouk, I., and Zaplotnik, Ž.: Decomposition of the Horizontal Wind Divergence Associated With the Rossby, Mixed Rossby-Gravity, Inertia-Gravity, and Kelvin Waves on the Sphere, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD040427, <https://doi.org/10.1029/2023JD040427>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2023JD040427>, 2024.
- Nie, Y., Zhang, Y., Yang, X.-Q., and Ren, H.-L.: Winter and summer rossby wave sources in the CMIP5 models, *Earth and Space Science*, 6, 1831–1846, <https://doi.org/https://doi.org/10.1029/2019EA000674>, tex.eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019EA000674>, 2019.
- 1235 Park, H.-S., Xie, S.-P., and Son, S.-W.: Poleward Stationary Eddy Heat Transport by the Tibetan Plateau and Equatorward Shift of Westerlies during Northern Winter, *Journal of the Atmospheric Sciences*, 70, 3288–3301, <https://doi.org/10.1175/JAS-D-13-039.1>, 2013.
- Park, M. and Lee, S.: Is the Stationary Wave Bias in CMIP5 Simulations Driven by Latent Heating Biases?, *Geophysical Research Letters*, 48, e2020GL091678, <https://doi.org/10.1029/2020GL091678>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020GL091678>, 2021.
- 1240 Park, S.-H., Klemp, J. B., and Skamarock, W. C.: A Comparison of Mesh Refinement in the Global MPAS-A and WRF Models Using an Idealized Normal-Mode Baroclinic Wave Simulation, *Monthly Weather Review*, 142, 3614–3634, <https://doi.org/10.1175/MWR-D-14-00004.1>, 2014.
- Petoukhov, V., Rahmstorf, S., Petri, S., and Schellnhuber, H. J.: Quasiresonant amplification of planetary waves and recent Northern Hemisphere weather extremes, *Proceedings of the National Academy of Sciences*, 110, 5336–5341, <https://doi.org/10.1073/pnas.1222000110>, 2013.
- 1245 Pyper, B. J. and Peterman, R. M.: Comparison of methods to account for autocorrelation in correlation analyses of fish data, *Canadian Journal of Fisheries and Aquatic Sciences*, 55, 2127–2140, <https://doi.org/10.1139/f98-104>, 1998.
- Qin, H., Klein, S. A., Ma, H.-Y., Van Weverberg, K., Feng, Z., Chen, X., Best, M., Hu, H., Leung, L. R., Morcrette, C. J., Rumbold, H., and Webster, S.: Summertime Near-Surface Temperature Biases Over the Central United States in Convection-Permitting Simulations, *Journal of Geophysical Research: Atmospheres*, 128, e2023JD038624, <https://doi.org/10.1029/2023JD038624>, eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2023JD038624>, 2023.
- 1250 Robock, A. and Mao, J.: The Volcanic Signal in Surface Temperature Observations, *Journal of Climate*, 8, 1086–1103, https://journals.ametsoc.org/view/journals/clim/8/5/1520-0442_1995_008_1086_tvst_2_0_co_2.xml, 1995.
- 1255 Rodwell, M. J. and Hoskins, B. J.: Monsoons and the dynamics of deserts, *Quarterly Journal of the Royal Meteorological Society*, 122, 1385–1404, 1996.
- Rodwell, M. J., Magnusson, L., Bauer, P., Bechtold, P., Bonavita, M., Cardinali, C., Diamantakis, M., Earnshaw, P., Garcia-Mendez, A., Isaksen, L., Källén, E., Klocke, D., Lopez, P., McNally, T., Persson, A., Prates, F., and Wedi, N.: Characteristics of Occasional Poor Medium-Range Weather Forecasts for Europe, *Bulletin of the American Meteorological Society*, 94, 1393–1405, <https://doi.org/10.1175/BAMS-D-12-00099.1>, 2013.
- 1260 Sakaguchi, K.: Scripts for Rossby Wave Evaluation of Dynamical Downscaling over North America, <https://doi.org/10.5281/zenodo.17458434>, 2025.
- Sakaguchi, K., Lu, J., Leung, L. R., Zhao, C., Li, Y., and Hagos, S.: Sources and pathways of the upscale effects on the Southern Hemisphere jet in MPAS-CAM4 variable-Resolution simulations, *Journal of Advances in Modeling Earth Systems*, 8, 1786 – 1805, <https://doi.org/doi:10.1002/2016MS000743>, 2016.
- 1265 Sakaguchi, K., Leung, L. R., Zarzycki, C. M., Jang, J., McGinnis, S., Harrop, B. E., Skamarock, W. C., Gettelman, A., Zhao, C., Gutowski, W. J., Leak, S., and Mearns, L.: Technical descriptions of the experimental dynamical downscaling simulations over North America by the

- CAM–MPAS variable-resolution model, *Geoscientific Model Development*, 16, 3029–3081, <https://doi.org/10.5194/gmd-16-3029-2023>, 2023.
- 1270 Sardeshmukh, P. D. and Hoskins, B. J.: The generation of global rotational flow by steady idealized tropical divergence, *Journal of the Atmospheric Sciences*, 45, 1228–1251, 1988.
- Schneidereit, A., Schubert, S., Vargin, P., Lunkeit, F., Zhu, X., Peters, D. H. W., and Fraedrich, K.: Large-Scale Flow and the Long-Lasting Blocking High over Russia: Summer 2010, *Monthly Weather Review*, 140, 2967–2981, <https://doi.org/10.1175/MWR-D-11-00249.1>, 2012.
- 1275 Schubert, S., Wang, H., and Suarez, M.: Warm Season Subseasonal Variability and Climate Extremes in the Northern Hemisphere: The Role of Stationary Rossby Waves, *Journal of Climate*, 24, 4773–4792, <https://doi.org/10.1175/JCLI-D-10-05035.1>, 2011.
- Simpson, I. R., Bacmeister, J., Neale, R. B., Hannay, C., Gettelman, A., Garcia, R. R., Lauritzen, P. H., Marsh, D. R., Mills, M. J., Medeiros, B., and Richter, J. H.: An Evaluation of the Large-Scale Atmospheric Circulation and Its Variability in CESM2 and Other CMIP Models, *Journal of Geophysical Research: Atmospheres*, 125, 1–42, <https://doi.org/10.1029/2020JD032835>, 2020.
- 1280 Skamarock, W. C., Klemp, J. B., Dudhi, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X.-Y., Wang, W., and Powers, J. G.: A Description of the Advanced Research WRF Version 3, Tech. rep., University Corporation for Atmospheric Research, Boulder, CO, United States, ISBN NCAR/TN–468+STR, ISSN 1477870X, <https://doi.org/10.5065/D6DZ069T>, publication Title: NCAR Technical Note Issue: June, 2008.
- Skamarock, W. C., Klemp, J. B., Duda, M. G., Fowler, L. D., Park, S.-H., and Ringler, T. D.: A multiscale nonhydrostatic atmospheric model using Centroidal Voronoi Tessellations and C-grid staggering, *Monthly Weather Review*, 140, 3090–3105, <https://doi.org/10.1175/MWR-D-11-00215.1>, 2012.
- 1285 Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Liu, Z., Berner, J., Wang, W., Powers, J. G., Duda, M. G., Barker, D. M., and Huang, X.-Y.: A Description of the Advanced Research WRF Version 4, Tech. rep., National Center for Atmospheric Research, Boulder, Colo., publication Title: NCAR Technical Note, 2019.
- 1290 Srivastava, A. K., Ullrich, P. A., Rastogi, D., Vahmani, P., Jones, A., and Grotjahn, R.: Assessment of WRF (v 4.2.1) dynamically downscaled precipitation on subdaily and daily timescales over CONUS, *Geoscientific Model Development*, 16, 3699–3722, <https://doi.org/10.5194/gmd-16-3699-2023>, 2023.
- Stan, C., Kollapaneni, S., Jenney, A., Wang, J., Wu, Z., Zheng, C., Kim, H., Garfinkel, C., and Singh, A.: A Python diagnostics package for evaluation of MJO-Teleconnections in S2S forecast systems, *EGUsphere*, pp. 1–22, <https://doi.org/10.5194/egusphere-2025-1142>, 2025.
- 1295 Staniforth, A.: *Meteorology and Atmospheric Physics Regional Modeling : A Theoretical Discussion*, *Meteorology and Atmospheric Physics*, 63, 15–29, 1997.
- Stensrud, D. J.: Upscale Effects of Deep Convection during the North American Monsoon, *Journal of the Atmospheric Sciences*, 70, 2681–2695, <https://doi.org/10.1175/JAS-D-13-063.1>, 2013.
- Sy, S., Madonna, F., Serva, F., Diallo, I., and Quesada, B.: Assessment of NA-CORDEX regional climate models, reanalysis and in situ gridded-observational data sets against the U.S. Climate Reference Network, *International Journal of Climatology*, 44, 305–327, <https://doi.org/10.1002/joc.8331>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/joc.8331](https://onlinelibrary.wiley.com/doi/pdf/10.1002/joc.8331), 2024.
- 1300 Takaya, K. and Nakamura, H.: A formulation of a wave-activity flux for stationary Rossby waves on a zonally varying basic flow, *Geophysical Research Letters*, 24, 2985–2988, <https://doi.org/10.1029/97GL03094>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1029/97GL03094](https://onlinelibrary.wiley.com/doi/pdf/10.1029/97GL03094), 1997.

- 1305 Takaya, K. and Nakamura, H.: A formulation of a phase-independent wave-activity flux for stationary and migratory quasigeostrophic eddies on a zonally varying basic flow, *Journal of the Atmospheric Sciences*, 58, 608–627, 2001.
- Teng, H. and Branstator, G.: Connections Between Heat Waves and Circumglobal Teleconnection Patterns in the Northern Hemisphere Summer, in: *Climate Extremes*, pp. 177–193, American Geophysical Union (AGU), ISBN 978-1-119-06802-0, <https://doi.org/10.1002/9781119068020.ch11>, section: 11 _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/9781119068020.ch11>,
1310 2017.
- Teng, H. and Branstator, G.: Amplification of Waveguide Teleconnections in the Boreal Summer, *Current Climate Change Reports*, 5, 421–432, <https://doi.org/10.1007/s40641-019-00150-x>, 2019.
- Teng, H., Branstator, G., Wang, H., Meehl, G. A., and Washington, W. M.: Probability of US heat waves affected by a subseasonal planetary wave pattern, *Nature Geoscience*, 6, 1056–1061, <https://doi.org/10.1038/ngeo1988>, 2013.
- 1315 Teng, H., Branstator, G., Tawfik, A. B., and Callaghan, P.: Circumglobal Response to Prescribed Soil Moisture over North America, *Journal of Climate*, 32, 4525–4545, <https://doi.org/10.1175/JCLI-D-18-0823.1>, 2019.
- Ting, M.: Maintenance of Northern Summer Stationary Waves in a GCM, *Journal of the Atmospheric Sciences*, 51, 3286–3308, [https://doi.org/10.1175/1520-0469\(1994\)051<3286:MONSSW>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<3286:MONSSW>2.0.CO;2), 1994.
- Trenberth, K. E., Branstator, G. W., Karoly, D., Kumar, A., Lau, N.-C., and Ropelewski, C.: Progress during TOGA in understanding
1320 and modeling global teleconnections associated with tropical sea surface temperatures, *Journal of Geophysical Research: Oceans*, 103, 14 291–14 324, <https://doi.org/10.1029/97JC01444>, _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/97JC01444>, 1998.
- Vallis, G. K.: *Atmospheric and Oceanic Fluid Dynamics*, Cambridge University Press, 2nd edition edn., ISBN 978-1-107-06550-5, <https://doi.org/10.1017/9781107588417>, 2017.
- Wallace, J. M. and Gutzler, D. S.: Teleconnections in the Geopotential Height Field during the Northern Hemisphere Winter, *Monthly
1325 Weather Review*, 109, 784–812, [https://doi.org/10.1175/1520-0493\(1981\)109<0784:TITGHF>2.0.CO;2](https://doi.org/10.1175/1520-0493(1981)109<0784:TITGHF>2.0.CO;2), 1981.
- Wang, B., Wu, R., and Lau, K.-M.: Interannual Variability of the Asian Summer Monsoon: Contrasts between the Indian and the Western North Pacific–East Asian Monsoons, *Journal of Climate*, 14, 4073–4090, [https://doi.org/10.1175/1520-0442\(2001\)014<4073:IVOTAS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<4073:IVOTAS>2.0.CO;2), 2001.
- Wang, Z., Chang, C.-P., and Wang, B.: Impacts of El Niño and La Niña on the U.S. Climate during Northern Summer, *Journal of Climate*,
1330 20, 2165–2177, <https://doi.org/10.1175/JCLI4118.1>, 2007.
- Wang, Z., Wu, R., Duan, A., and Qu, X.: Influence of eastern tibetan plateau spring snow cover on north american air temperature and its interdecadal change, *Journal of Climate*, 33, 5123 – 5139, <https://doi.org/10.1175/JCLI-D-19-0455.1>, place: Boston MA, USA, 2020.
- Weaver, S. J. and Nigam, S.: Variability of the Great Plains Low-Level Jet: Large-Scale Circulation Context and Hydroclimate Impacts, *Journal of Climate*, 21, 1532–1551, <https://doi.org/10.1175/2007JCLI1586.1>, 2008.
- 1335 White, R. H. and Mareshet Admasu, L.: Temporally and zonally varying atmospheric waveguides – climatologies and connections to quasi-stationary waves, *Weather and Climate Dynamics*, 6, 549–570, <https://doi.org/10.5194/wcd-6-549-2025>, 2025.
- White, R. H., Kornhuber, K., Martius, O., and Wirth, V.: From Atmospheric Waves to Heatwaves: A Waveguide Perspective for Understanding and Predicting Concurrent, Persistent, and Extreme Extratropical Weather, *Bulletin of the American Meteorological Society*, 103, E923–E935, <https://doi.org/10.1175/BAMS-D-21-0170.1>, 2022.
- 1340 White, R. H., Anderson, S., Booth, J. F., Braich, G., Draeger, C., Fei, C., Harley, C. D. G., Henderson, S. B., Jakob, M., Lau, C.-A., Mareshet Admasu, L., Narinesingh, V., Rodell, C., Roocroft, E., Weinberger, K. R., and West, G.: The unprecedented Pacific Northwest heatwave of June 2021, *Nature Communications*, 14, 727, <https://doi.org/10.1038/s41467-023-36289-3>, 2023.

- Whitham, G. B.: A note on group velocity, *Journal of Fluid Mechanics*, 9, 347–352, <https://doi.org/10.1017/S0022112060001158>, 1960.
- Wills, R. C., White, R. H., and Levine, X. J.: Northern Hemisphere Stationary Waves in a Changing Climate, *Current Climate Change Reports*, 5, 372–389, <https://doi.org/10.1007/s40641-019-00147-6>, 2019.
- 1345 Wirth, V.: Waveguidability of idealized midlatitude jets and the limitations of ray tracing theory, *Weather and Climate Dynamics*, 1, 111–125, <https://doi.org/10.5194/wcd-1-111-2020>, 2020.
- Wirth, V., Riemer, M., Chang, E. K. M., and Martius, O.: Rossby Wave Packets on the Midlatitude Waveguide—A Review, *Monthly Weather Review*, 146, 1965–2001, <https://doi.org/10.1175/MWR-D-16-0483.1>, 2018.
- 1350 Wolf, G., Brayshaw, D. J., Klingaman, N. P., and Czaja, A.: Quasi-stationary waves and their impact on European weather and extreme events, *Quarterly Journal of the Royal Meteorological Society*, 144, 2431–2448, <https://doi.org/10.1002/qj.3310>, <https://onlinelibrary.wiley.com/doi/pdf/10.1002/qj.3310>, 2018.
- Yang, Y.: Rossby Wave Ray Tracing and Ray Flux, <https://github.com/yinan-codes/Rossby-wave-ray-tracing>, original-date: 2025-10-21T06:22:22Z, 2025.
- 1355 Yang, Y. and Li, J.: Novel monsoon indices based on vector projection and directed angle for measuring the East Asian summer monsoon, *Climate Dynamics*, 63, 210, <https://doi.org/10.1007/s00382-025-07696-7>, 2025.
- Yuan, J., Li, W., and Deng, Y.: Amplified subtropical stationary waves in boreal summer and their implications for regional water extremes, *Environmental Research Letters*, 10, 104009, <https://doi.org/10.1088/1748-9326/10/10/104009>, 2015.
- Zhang, C., Golaz, J.-C., Forsyth, R., Vo, T., Xie, S., Shaheen, Z., Potter, G. L., Asay-Davis, X. S., Zender, C. S., Lin, W., Chen, C.-C., Terai, C. R., Mahajan, S., Zhou, T., Balaguru, K., Tang, Q., Tao, C., Zhang, Y., Emmenegger, T., Burrows, S., and Ullrich, P. A.: The E3SM Diagnostics Package (E3SM Diags v2.7): a Python-based diagnostics package for Earth system model evaluation, *Geoscientific Model Development*, 15, 9031–9056, <https://doi.org/10.5194/gmd-15-9031-2022>, 2022.
- 1360 Zhang, G. and Wang, Z.: North Atlantic Extratropical Rossby Wave Breaking during the Warm Season: Wave Life Cycle and Role of Diabatic Heating, *Monthly Weather Review*, 146, 695–712, <https://doi.org/10.1175/MWR-D-17-0204.1>, 2018.
- 1365 Zhang, Y., Pan, Y., Xue, Y., Diallo, I., Zeng, X., Li, S., Neelin, J. D., Lau, W. K. M., Boone, A. A., Vitart, F., Yao, T., Tang, Q., Sato, T., Koo, M.-S., Ardilouze, C., Saha, S. K., Yang, J., Matera, S., Lin, Z., Qi, X., Qin, Y., Nakamura, T., Nobre, P., Peano, D., Senan, R., Takaya, Y., Wang, H., Zhang, H., Zhan, Y., Zhao, M., Mechoso, C. R., Bao, Q., Bottino, M. J., Hong, S., Lin, Y., Xie, S., Pan, X., Nayak, H. P., Chou, S. C., and Guo, W.: Near-global summer circulation response to the spring surface temperature anomaly in Tibetan Plateau — the GEWEX/LS4P first phase experiment, *Climate Dynamics*, 62, 2907–2924, <https://doi.org/10.1007/s00382-024-07210-5>, 2024.
- 1370 Zhao, S., Li, J., and Li, Y.: Dynamics of an interhemispheric teleconnection across the critical latitude through a southerly duct during boreal winter, *Journal of Climate*, 28, 7437–7456, <https://doi.org/10.1175/JCLI-D-14-00425.1>, 2015.
- Zhou, W., Leung, L. R., and Lu, J.: Linking Large-Scale Double-ITCZ Bias to Local-Scale Drizzling Bias in Climate Models, *Journal of Climate*, 35, 7965–7979, <https://doi.org/10.1175/JCLI-D-22-0336.1>, 2022.
- Zienkiewicz, O. C. and Zhu, J. Z.: Superconvergence and the superconvergent patch recovery, *Finite Elements in Analysis and Design*, 19, 11–23, [https://doi.org/10.1016/0168-874X\(94\)00054-J](https://doi.org/10.1016/0168-874X(94)00054-J), 1995.