

Managing glacial and periglacial hazards in the Alps: a geohistorical approach

Juliette Bazin^{*1,2}, Ludovic Ravanel^{1,3}, Sandrine Caroly²

¹EDYTEM, Savoie Mont-Blanc University, CNRS, 5 Boulevard de la Mer Caspienne 73370 Le Bourget-du-Lac, France

²PACTE, Grenoble Alpes University, INP, 46 Avenue Félix Viallet 38000 Grenoble, France

³Department of Geosciences, University of Oslo, 0371 Oslo, Norway

*Corresponding author: J. Bazin (juliette.bazin@univ-smb.fr)

Financial support : This research was funded by the EU ALCOTRA 2021-2027 Interreg VI-A *PrévRisk-CC* project.

Key words

Glacial and periglacial hazards, Risk management, Risk perception, Geo-history, European Alps

Abstract. Glacial and periglacial hazards such as ice or rock avalanches, glacial lake outburst floods, and debris flows have caused severe damage in populated Alpine mountain regions throughout history. The objective of this research is to investigate how risk management methods have evolved over time and across different countries in the European Alps. Through a geo-historical study, we compile an inventory of events that have triggered risk management actions over the past centuries. This management is analysed using eight guiding pillars: hazard understanding, preventive information, land-use planning, monitoring and surveillance, hazard-vulnerability-exposure mitigation, crisis management preparedness, response, and resilience.

Certain events, such as the discharge of the water pocket of the Tête Rousse glacier (FR) in 1892 or the collapse of the Allalin glacier (CH) in 1965, represent key turning points that mark a shift or renewal in how risks are perceived and addressed. Today, glacial and periglacial risk management benefits from a better understanding of both hazards and vulnerabilities. Our geo-historical analysis highlights that the inclusion of the local populations in crisis management has become an increasingly significant factor in decision-making processes. However, current risk management practices remain limited and would benefit from more participatory approaches, which in turn partly depend on the perception of hazards and their integration into management practices.

1 Introduction

Rock and ice mass movements in high mountain areas pose significant risks to populations and infrastructures. Numerous past events show how glacial and periglacial processes can deeply affect Alpine valleys through large-scale and/or cascading phenomena (e.g., Haeberli et al., 2016; Magnin et al., 2023). Their frequency and volume appear to be increasing (Ravanel and Deline, 2008, 2015; Jacquemart et al., 2024) as illustrated by recent disasters in Bondo (Graubünden, CH) in August 2017 (Walter et al., 2020), La Béarde (Isère, FR) in June 2024 (Mainieri et al., 2025), and Blatten (Valais, CH) in May 2025 (Büntgen et al., 2025).

This study focuses on major hazards affecting valley floors: ice, rock, and mixed avalanches, glacial lake outburst floods (GLOFs) and periglacial debris flows. These hazards are defined as natural phenomena impacting Alpine territories (Moles, 1972; Bisquet et al., 2025). A process becomes a risk

52 when it threatens lives, economic assets, or the environment. The notion of an *event* is useful for
53 historians as it marks “a cut, a discontinuity”, something “interesting”, sufficiently “important” or “new”
54 to be “told or enacted” (Dosse, 2010; Giacona et al., 2017). *Issues* are thus the exposed elements with
55 **varying** vulnerabilities to each hazard (Defossez et al., 2018).

56 **Permafrost warming and/or glacial debuttressing are contributing factors to rockfalls and rock**
57 **avalanches** (Fischer et al., 2006; Gruber and Haeberli, 2007; Huggel et al., 2010). They may directly
58 impact valley areas or provide material that triggers debris flows (Walter et al., 2020).

59 An ice avalanche occurs when a glacier section detaches and **moves quickly downwards** (Alean, 1985;
60 Richard, 2005; Faillettaz et al., 2015), destroying infrastructure, forests and lives, as **illustrated by the**
61 **collapse of the Allalin glacier (CH) onto worker accommodations during the construction of the**
62 **Mattmark dam in 1965** (Dalban Canassy et al., 2011). Indirectly, they can trigger snow avalanches
63 (Margreth and Funk, 1999) or debris flows when ice temporarily dams a **stream** (Richard, 2005).

64 GLOFs arise when a glacial lake or **another water body** suddenly drains, releasing large volumes of
65 water that can evolve into debris flows if sufficient **amounts of** sediment are available (Richard, 2005;
66 Huss et al., 2007; Carrivick and Tweed, 2016; Ancey et al., 2019). Their impacts can devastate entire
67 valleys, as seen in the Bagnes valley below the Giétro glacier (CH) in 1595 and 1818 (Vincent et al.,
68 2010; Ancey et al., 2019).

69 Integrated management of these risks (Prudent-Richard et al., 2008; Stoffel et al., 2014; Jacquemart et
70 al., 2024) is **essential for prevention because it allows for a holistic, coordinated approach that**
71 **anticipates, mitigates, and adapts to complex, interconnected climate and emerging risks before they**
72 **escalate into crises. We base our approach on the key components of risk management: hazard, exposure,**
73 **and vulnerability** (Niggli et al., 2024).

74 The database developed **in** this study documents glacial and periglacial events across the European Alps
75 over the past two centuries that required risk management. **Prevention measures from a political**
76 **perspective are not uniform across Alpine countries. We have chosen to primarily present examples from**
77 **France, Switzerland and Italy. Although Austria is also cited according to the sources we collected. The**
78 **authors acknowledge the limitations of this study, particularly the challenges associated with obtaining**
79 **comprehensive data; nevertheless, it is conceived as an exploratory approach. As such, it may serve as**
80 **an initial step toward the development of a broader collaborative framework. The purpose is not only to**
81 **provide a detailed account of the risk management measures implemented, but also to assess the**
82 **frequency and intensity of such events and to trace historical management trends.** An historical approach
83 enables the analysis of change processes in risk management (Giacona et al., 2017, 2019), beyond mere
84 event description (Girard and Rivière-Honegger, 2015).

85 We **thus** seek to understand how Alpine glacial and periglacial hazards have shaped scientific knowledge
86 and risk management practices.

87 The paper first outlines the theoretical and historical framework of risk management, followed by the
88 methods used to construct the database and apply the **eight** pillars of risk management **which we have**
89 **identified.** The results describe the spatial and temporal distribution of events and corresponding
90 management actions. Finally, we discuss the limitations of the data and the challenges of future
91 prevention **under** a changing climate.

92

93

94 **2 State of the art and theoretical positioning**

95

96 The concepts used here **require clarification to frame** our approach. A *risk* refers to the probability of
97 damage resulting from interactions between a physical process (the *hazard*), *exposure*, and *vulnerability*
98 (Pigeon, 2002). The hazards we study occur at high altitudes (> 2,000-2,500 m a.s.l.) and may reach
99 valley floors, threatening infrastructure, livelihoods, ecosystems, and populations (IPCC, 2023).
100 Mountain valleys are thus considered vulnerable due to their exposure to such hazards, **a condition**
101 **reinforced by the concentration of people and infrastructure, which increases susceptibility to cascading**
102 **effects** (IPCC, 2023; Niggli et al., 2024). Vulnerability also depends on *resilience* and *sensitivity*, *i.e.* the

103 system's ability to absorb shocks and the degree of damage it can sustain (Schneiderbauer and Ehrlich,
104 2004; Defosse et al., 2018). *Exposure* reflects the spatial and social **proximity** between hazard sources
105 and affected communities (IPCC, 2022).

106 The **continued** expansion of urban areas in high Alpine valleys, **driven by tourism and socioeconomic**
107 **development** (Hock and Rasul, 2022), **combined with climate-induced changes in the cryosphere,**
108 **increases potential losses** (Huggel et al., 2015) and underscores the need for effective *risk management*
109 (Allen et al., 2017). The latter encompasses strategies to anticipate, reduce, and respond to hazards
110 (Leone et al., 2021) and aims to reduce uncertainty through scientific **advances** (Margreth et al., 2011,
111 2017). While **comparable** to other Alpine hazards, glacial and periglacial **processes evolve rapidly under**
112 **climate change and can reach extreme magnitudes** (Haeberli et al., 1999; Richard, 2005). Anticipation
113 remains **difficult** because these hazards **co-evolve with a rapidly changing cryosphere:** glacier retreat,
114 de-icing of rock faces, **permafrost warming and associated slope movements,** formation or
115 disappearance of glacial lakes, and changes in glacier thermal regimes (Mainieri et al., 2025). **High**
116 **erosion rates in recently deglaciated or warming permafrost areas further increase debris-flow**
117 **susceptibility** (Huggel et al., 2015).

118 As historical experience shows, approaches based solely on hazard characterisation (Richard, 2005) are
119 necessary but not sufficient (Allen et al., 2017). Analysing past events improves understanding of
120 processes and supports better risk mitigation (Carrivick and Tweed, 2016). Rather than **compiling**
121 **exhaustive inventories, this study focuses is on lessons learned from major events, particularly how**
122 **societies have managed risks over time.**

123 Building a historical database requires a diachronic approach, inspired by diachronic monographs
124 (Girard and Rivière-Honegger, 2015). **This perspective (e.g. Desailly, 1990; Mendez, 2010) uses time**
125 **to explain current territorial and risk dynamics.** Such an approach allows the identification of
126 **geohistorical trajectories and rupture points in hazard evolution** (Valette and Carozza, 2019; Hugerot et
127 **al., 2021).** We adopt this framework to analyse long-term risk management, local practices, and the
128 **evolution of risk culture** (Dollfus and D'Ercole, 1996). Studies such as Favier (2006) and Fournier
129 (2010) on avalanches and floods in the Grenoble basin (FR), or Niggli et al. (2024) on GLOFs, illustrate
130 **how mountain societies have experienced and managed hazards.**

131 Adopting a diachronic perspective also requires careful source selection (**biases are discussed in § 5.2**).
132 Archival records complement narrative sources (Fournier, 2010). Since the 19th century, **historical**
133 **reconstructions have supported risk anticipation** (Dourlens, 2004). **However, exhaustive chronologies**
134 **remain unattainable, as many small or frequent events leave no archival trace** (Giacona et al., 2019).

135 To **address** this limitation, **historical approaches prioritise the analysis of processes and mechanisms**
136 **over completeness.** Our study thus provides a synthetic view of Alpine risk management. Similar
137 approaches, such as the temporal analysis of rockfalls in the Mont-Blanc massif (Ravel and Deline,
138 2008; Ravel et al., 2020), reveal long-term dynamics.

139 **Long-term monitoring initiatives also demonstrate the sensitivity of mountain environments to climate**
140 **change, such as the EU-PACE project (Permafrost and Climate in Europe), which highlights the role of**
141 **permafrost in slope stability and geohazards** (Harris et al., 2001; Etzelmüller et al., 2020).

142

143

144 **3 Construction of the database**

145

146 3.1 Inventory of major glacial and periglacial events in the Alps

147

148 3.1.1 Criteria used to identify events to be included in the database

149

150 Building a comprehensive long-term database is challenging as data availability before the 20th century
151 largely depends on societal vulnerability **and record-keeping** (Giacona, 2019).

152 We adopted a qualitative approach, retaining events that significantly marked local history or the
153 environment, living traces in archives or collective memory. Our criteria are grouped into four
154 categories.

155

156 *Extreme events*

157 We focused on rare, high-magnitude events exceeding several thousand cubic metres or causing serious
158 damage (Bourrelrier and Dunglas, 2009; Field et al., 2012). Quantitatively, these correspond to the upper
159 end of the frequency-intensity spectrum.

160

161 *Events with feedback*

162 We include events that led to in-depth post-event analyses aimed at improving process understanding,
163 beyond simple observation (e.g. the Giétro glacier collapse in 1818; Wiegandt and Lugon, 2008).

164

165 *Repeated events*

166 We consider events repeatedly occurring within the same watershed, focusing on major one. For
167 example, the Bockkarkees hanging glacier (AT) has produced >70 ice avalanches since 1933, including
168 events reaching 5×10^6 m³ in 1945 and $2-3 \times 10^6$ m³ in 1964 and 1975 (*Gridabase*; Kellerer-Pirklbauer
169 et al., 2012).

170

171 *Change in risk perception*

172 Events that significantly influenced risk perception are included, even if they do not meet the previous
173 criteria. Risk culture varies across Alpine countries and is shaped by historical events. Event memory
174 differs by context, making risk perception a social and political issue. The Mattmark disaster marked a
175 major policy shift, highlighting the role of unpredictability (Capozzi, 2011). Similarly, the Tête Rousse
176 disaster (FR) established in France the idea of glaciers as internal, hidden, yet predictable hazards. The
177 Marmolada glacier (IT) collapse in 2022 (Francese et al., 2024), although limited to high-altitude
178 impacts, caused 11 fatalities and strongly influenced public opinion, highlighting the need to better
179 identify glacial hazards (Chiarle et al., 2022). It also led to the creation of a national working group and
180 the publication of guidelines (Dipartimento della Protezione Civile, 2025).

181

182 We initially intended to start our inventory in 1985-1990, marking the first visible signs of the climate
183 crisis in the Alps (Ravanel, 2009; Ravanel et al., 2020), up to 2025. However, earlier key events such as
184 the 1892 Tête Rousse outburst (Mougin et Bernard, 1922) were too significant to omit. We therefore
185 extended the analysis back to the early 19th century, where reliable sources exist.

186 For each event, we collected the following data: location (country, region, mountain range, orientation,
187 glacier or summit name), rupture date(s), causes, mobilised volumes, runout distance, cascading effects,
188 damage and casualties, monitoring methods, crisis management, and sources.

189 This method was applied to three hazard types: ice avalanches, torrential hazards (including GLOFs and
190 debris flows), and rockfalls/rock avalanches.

191 For cascading processes, we record the most impactful downstream process. For example, if a serac fall
192 triggers lake overflow, the torrential process is retained. If permafrost degradation destabilises a moraine
193 dam, we document the resulting breach and downstream propagation.

194

195 3.1.2 Data sources

196

197 We used the *Gridabase* inventory produced as part of the European research project (« Gridabase -
198 Glaciorisk », s. d., 2001-2003). This database compiles 501 documented events across Austria, France,
199 Italy, and Switzerland, including floods, GLOFs, and ice avalanches (Richard, 2005). The aim of
200 *Gridabase* was to preserve as comprehensively as possible the memory and trace of glacial events
201 (Peissier and Courtray, 2012), including those of modest importance (excluded here according to our
202 criteria). However, investigations initiated by the Swiss Parliament highlighted that changing

203 environmental conditions exceeded historical empirical references, requiring broader system
204 assessments (Rickenmann and Zimmermann, 1993). *Gridabase* therefore presents a retrospective and
205 partial perspective, which we accounted for in our analysis. In addition, for Italian Alps, we used the
206 *Geoclimalp* cartographic inventory (Nigrelli et al., 2024; geoclimalp, 2026), which contains > 700 mass
207 movements > 1500 m a.s.l. between 2000 and 2022.

208 Other sources include local databases such as the *RTM Database* or the regional *Catasto Dissesti*. The
209 former is a French database developed by the *Restauration des Terrains de Montagne* (RTM) public
210 service of the *Office National des Forêts*. Since the late 19th century, it has recorded a wide range of
211 geomorphic processes, including avalanches, gully erosion, subsidence, landslides, rockfalls and debris
212 flows (Bisquert et al., 2025). The *Catasto Dissesti* is a regional mapped inventory of hydrogeological
213 and gravitational phenomena in Aosta Valley, enriched since the 1990s and regularly updated for risk
214 prevention (Giardino et al., 2010). We retained from both databases glacial and periglacial events,
215 including rock glacier ruptures, permafrost-related rockfalls and debris flows generated by cascading
216 processes.

217 The RTM has also produced numerous technical reports on glacial and periglacial events. These
218 documents, sometimes, unpublished, describes causes, trajectories, volumes, damage, and process
219 histories within catchment. They also document risk management actions, from modelling and early
220 warning to operational interventions (e.g. Patinoire lake in 1964; Étançons torrent in 2024; (Cathala et
221 al., 2021; Demolis et al., 2021; Mainieri et al., 2025) and crisis management through operational
222 measures (e.g. artificial draining of Rochemelon lake, FR, in 2005; Vincent et al., 2010; Cathala et al.,
223 2021).

224 Many events documented by Swiss authorities were also included, such as the flood protection measures
225 implemented at the Gruben glacier (Haeberli et al., 2001).

226 Scientific papers constitute a major part of our dataset, providing detailed analyses of hazard
227 mechanisms and management strategies. Individual events are often studied by multiple teams using
228 complementary approaches. The Whymper hanging glacier and the Planpincieux glacier (IT) illustrate
229 this: studies address monitoring systems, rupture scenarios, hazard assessment, and evacuation strategies
230 (Margreth and Funk, 1999; Margreth et al., 2011; Schindelegger, 2019; Dematteis et al., 2021; Troilo,
231 2025).

232 The inventories of Chiarle et al. (2007, 2022), Jacquemart et al. (2024), and Niggli et al. (2024) were
233 used to cross-check and validate our dataset.

234 Ministerial reports (e.g. Lacroix et al., 2022), the study of Magnin et al. (2023), and output from the
235 *Pôle Alpin des Risques Naturels* (PARN), provide insights into how Alpine countries manage glacial and
236 periglacial risks. They document research projects, monitoring strategies, prevention measures, and
237 crisis management practices (e.g. *Glaciorisk*, *PERMAdataROC*, *PermaNET*, *GlariskAlp*, *GLAMOS*,
238 *Prevrisk-CC*, *PermaRisk*, *SAMCO*, etc.). The *PAPROG* initiative fosters multidisciplinary collaboration
239 between researchers and practitioners to improve knowledge and management of glacial risks. Similarly,
240 *GEORESEARCH*, coordinates projects on monitoring, cryospheric dynamics, and quantitative risk
241 analysis in the Alps (*FROST.INI*, *Futurelakes*, *GlacierRocks*, *AlpSenseRely*, etc.).

242 The *World Glacier Monitoring Service* (WGMS) provides global data on glacier fluctuations (1959-
243 2010), including notable hazard events.

244 Finally, the local press provides insights into the societal impact of mass movements and associated
245 management responses. However, this source requires careful critical interpretation, as it may reflect
246 biases and lack technical accuracy (Joffe and Orfali, 2005). Despite these limitations, it remains valuable
247 for analysing societal risk perception.

248

249 3.2 Types of glacial and periglacial risk management

250

251 3.2.1 Definition of the eight pillars of risk management

252

253 Based on Alpine risk management systems (from local to national levels) and scientific literature,
254 Kienholz et al., (2004) and Link and Stötter (2015) developed a classification of risk prevention and
255 management methods. More recently, the Risk Management Cycle (RMC) – or Disaster Risk
256 Management (DRM) – has been conceptualised. It encompasses all policies, strategies, and actions
257 aimed at preventing, reducing, and managing disaster risks, as well as preparing for, responding to, and
258 recovering from their impacts (Tagarev et al., 2021). In Switzerland and Nordic countries, since the
259 1980s, natural hazard management has followed the principles of “Risk Minimisation” and “Integral
260 Risk Management” (Nyberg et al., 2026), which aim to reduce risks to acceptable levels through
261 assessment, prevention, mitigation, and preparedness, rather than eliminate them. Post-crisis phases
262 focus on learning to improve future measures.

263 In France, similar principles are embedded in public policy and summarized by the PARN for glacial
264 and periglacial risk prevention (Boudières and Peisser, 2013). In Italy, the *Dipartimento della Protezione*
265 *Civile* (2025) has produced a methodological framework defining objectives for understanding glacial
266 and periglacial phenomena. It outlines key pillars such as monitoring, public information, and crisis
267 resilience.

268 All these principles were used to classify the management strategies identified in our database.

269

270 ***Pillar 1 : Hazards, phenomena, and risks understanding***

271 This pillar improves knowledge through past-events analysis, archives, monitoring, modelling, and
272 experience feedback (Villeneuve et al., 2010; Faillettaz et al., 2015). It includes inventories and
273 vulnerability assessments (e.g. GlaRiskAlp project; (Lucchesi et al., 2014; Nigrelli et al., 2013). Within
274 DRM, this corresponds to risk identification and preparedness (Tagarev et al., 2021).

275

276 ***Pillar 2 : Preventive information***

277 This pillar promotes risk culture through public information and education (Beccera and Peltier, 2011),
278 enabling citizens to actively contribute to preparedness and crisis response (OECD, 2018).

279

280 ***Pillar 3 : Integration into land-use planning***

281 Risk is integrated into planning tools such as hazard maps and preventive plans to reduce exposure and
282 vulnerability (Schneiderbauer and Ehrlich, 2004). These tools combine modelling and spatial analysis
283 to support decision-making (Allen et al., 2022).

284

285 ***Pillar 4 : Monitoring and surveillance***

286 Monitoring at-risk sites (cf.: Pillar 1) enables early detection and warning (Giordan et al., 2020; Cathala
287 et al., 2024), supporting emergency preparedness. For example, Gormersee lake (CH) is monitored using
288 cameras, depth measurements, and seismic sensors to anticipate GLOF hazards (WGMS, 1993; Huss et
289 al., 2007). In Italy, glaciers such as Freney, Brenva, Grandes Jorasses, and Planpincieux are monitored
290 using time-lapse imagery to assess flow velocity (Dematteis et al., 2024). Satellite observations
291 complement field monitoring, especially for remote or large-scale events, though in the Alps they are
292 often used post-event to analyse processes (e.g. Marmolada glacier collapse in 2022; (Bondesan and
293 Francese, 2023; Olivieri and Bettanini, 2023) or to monitor supraglacial lakes hazards (e.g. Lys glacier
294 or the Forni glacier, IT; (Davide et al., 2021; Fugazza et al., 2018). However, field-based observation
295 remains central in the Alpine context. Field observations and human-based warning systems remain
296 central, including observer networks such as “*Regards d’Altitude*” (ONF-RTM, 2025).

297

298 ***Pillar 5 : Reducing hazard, exposure, and vulnerability***

299 This pillar relies on cooperation between scientists and authorities to limit damage when hazard cannot
300 be avoided (Marco et al., 2012). Mitigation measures include engineering works, early warning systems,
301 evacuation planning, and source interventions (Vincent et al., 2010b; Tagarev et al., 2021).

302 For GLOFs, measures include drainage, water-level regulation, and structural stabilisation (Niggli et al.,
303 2024). Exposure is reduced through channel management and protective structures, while vulnerability
304 is lowered by improving preparedness and response capacities.

306 ***Pillar 6 : Crisis management preparedness***

307 This pillar corresponds to measures taken to improve response capacity (Tagarev et al., 2021). It includes
308 planning, training, and coordination measures implemented before crises, such as contingency plans,
309 simulations, and risk integration into training (Einhorn and Peisser, 2011; Link and Stötter, 2015).

311 ***Pillar 7 : Response***

312 Response refers to immediate actions during or after an event to manage impacts, save lives, and ensure
313 safety (OECD, 2018). It includes emergency assistance and implementation of crisis plans (Tagarev et
314 al., 2021).

316 ***Pillar 8 : Resilience***

317 Resilience is defined as a system's ability to absorb shocks and recover, involving resistance, adaptation,
318 and reorganisation (Reghezza et al., 2006; Dauphiné and Provitolo, 2013). The objective is not full
319 recovery but achieving a new acceptable equilibrium (Einhorn, 2017). This phase also provides
320 opportunities to improve future risk reduction measures (Tagarev et al., 2021).

322 3.2.2 Distribution of events according to the eight pillars typology

324 Once the typology has been established, each event was assigned to the relevant management pillars
325 mobilised in response. Events were then grouped by decade (or by five-year intervals for rock hazards)
326 to improve temporal readability. This approach highlights changes in management practices over time
327 while limiting visual discontinuities associated with rare events.

330 4 Results

332 This study identifies 200 glacial and periglacial events across 109 Alpine catchment areas in France,
333 Switzerland, Italy, and Austria. These events were selected based on their magnitude, impacts, or
334 influence on the evolution of glacial and periglacial risk management practices.

336 4.1 Spatial and temporal distribution of the events

338 4.1.1 Distribution of the events across the Alps

340 The spatial distribution of events (Fig. 1) shows a marked concentration in certain areas, highlighting
341 particularly exposed or dynamically active zones. The Mont-Blanc massif (mainly its French and Italian
342 sides) clearly emerges as the most densely affected area, across all hazard types. It is followed by the
343 Valais Alps (CH). GLOFs are also more frequent in the western Alps.

344 The western Alps with high elevations, steep slopes, and extensive glacier cover, make them particularly
345 prone to mass movements. The Mont-Blanc massif and the Valais Alps are among the most monitored
346 and instrumented regions, which may introduce a 'visibility bias' (cf. 5.2). High population density and
347 tourism further enhance event detection and documentation, potentially amplifying their perceived
348 significance.

349 Figure 1 presents major events that occurred between the 17th century and today, revealing the long-term
350 dynamics of these hazards. They are not isolated events but part of long-term, sometimes recurring
351 trajectories. This representation shows that multiple hazard types coexist within the same Alpine
352 massifs, increasing the complexity of risk management.




Figure 1. Distribution of major glacial and periglacial events inventoried in the Alps (base map: OpenStreetMap).

354

4.2 Distribution of management pillars over time

356


4.2.1 Phenomena intensification

358

359

Keeping in mind potential source and event biases (cf. § 5.2), we nevertheless observe an intensification of glacial and periglacial hazards since the 1980s across all Alpine countries (Fig. 2). This trend is based solely on events that required risk management and contributed to hazard understanding.

361



362

Figure 2. Cumulative frequencies of inventoried glacial and periglacial events requiring risk management in the Alps.

363

364

365
366 4.2.2. Evolution over time of the pillars used

367
368 This section examines how risk management and prevention methods have evolved over time through
369 significant glacial or periglacial events. Events are analysed according to the management pillars
370 mobilised at the time.

371 For the three types of events, Pillars 2 (Preventive information) and 3 (Land use planning) appear only
372 from the 1990s onward, while Pillar 6 (Crisis preparedness) has been implemented more recently. Pillar
373 1 (Hazard understanding) remains the most consistently used. It also continues to be applied
374 retrospectively, as past events are reanalysed to improve risk knowledge.

375 The evolution over time of risk management associated with major *ice avalanche* events in the Alps is
376 presented in Figure 3. The graph shows two main phases: a long period (1597-1950) dominated almost
377 exclusively by Pillar 1 (Understanding hazard), followed by a gradual and accelerating diversification
378 of management approaches from the 1950s and 1960s onwards.

379 The Altels ice avalanche (CH) in 1895 marked a turning point in the recognition of the need to
380 understand high-altitude hazards. This avalanche was of exceptional size, with 4×10^6 m³ of ice breaking
381 away, killing six people and burying 170 cattle over an area of 1 km² (Forel, 1895; Du Pasquier, 1896).
382 For a long time, this event served as a benchmark for ice avalanches. "To what extent can we predict
383 such disasters, and to what degree will we be able to prevent them from happening again? Many Alpine
384 glaciers give rise to avalanches or ruinous debacles. (...) The same glaciers periodically give rise to the
385 same phenomena, so it is above all empirical knowledge of these phenomena that we need to acquire.
386 (...) An accurate knowledge of the dangerous points and the phenomena that occur there is absolutely
387 necessary. The era of independent personal research is over; it has produced all it could in this field (...).
388 What is needed now is more coordination, more method, more consistency in research" (Du Pasquier,
389 1896).

390 The predominance of Pillar 1 over several centuries should not be interpreted as a lack of institutional
391 response to glacial and periglacial hazards. It rather reflects the fact that, within the epistemic and
392 institutional context of the time, the production of knowledge and the documentation of phenomena
393 constituted the primary and legitimate form of risk management. At that time, management was largely
394 confined to the academic sphere (naturalists, cartographers, scientific academies), long before it became
395 integrated into engineering and spatial planning. This observation suggests that contemporary categories
396 of risk management should not be retrospectively applied to these periods. If preventive information
397 (Pillar 2) appears late, this indicates that decision-makers only gradually began to communicate
398 simplified risk information to exposed populations. This development is closely linked to the emergence
399 of crisis preparedness (Pillar 6). It also raises the question of the role assigned to local communities in
400 risk management, which has long remained expert-driven, with limited structured communication with
401 **residents**. The example of the Val Ferret (IT), where the terminal tongue of the Planpincieux glacier has
402 been monitored since 2014, illustrates this evolution. **Monitoring is coupled with rapid civil protection
403 responses and public information measures** (Giordan et al., 2020; Dematteis et al., 2024). The aim is to
404 continuously track glacier dynamics, and detect instability, and activate procedures such as road closures
405 or village evacuation.

406 This study also integrates avalanche hazard maps, such as the *Carte de Localisation des Phénomènes
407 d'Avalanche* (CLPA; Avalanche Phenomenon Location Maps) in France, *Catasto valanghe in Aosta
408 valley* (Italy), *Indication of avalanche hazards (ATH)* for Switzerland, *Wildbach und Lawinenverbauung
409 (WLV)* for Austria, into land-use planning (Pillar 3), even though they primarily address snow
410 avalanches. These maps first emerged in the 1970s, but events such as those at the Bourgeat glacier (FR)
411 in 1984 and 1993 show that ice avalanches can trigger snow avalanches (RTM-ONF, 2000, unpublished).

412 **We therefore consider these mapping tools as addressing combined snow-ice hazards.**

413




Figure 1. Temporal evolution of ice avalanche risk management in the Alps.


414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435

The evolution over time of risk management for major rockfall and rock avalanche events in the Alps is presented in Figure 4. It is divided into five-year intervals to better highlight recent dynamics, as most rockfall events have occurred since 1980.

The reduction in vulnerability, exposure, and hazard (Pillar 5) is particularly evident in these events. For example, the 1987 Val Pola rockslide (CH; $30 \times 10^6 \text{ m}^3$) involved preventive evacuation, illustrating proactive risk management (Chardon, 1990). In 2011, at Piz Cengalo / Bondo ($1.5 \times 10^6 \text{ m}^3$), preventive measures included public alerts, trail closures, evacuation of alpine buildings, and installation of protective structures (Bohnenblust, 2017).

Monitoring unstable rock faces is a key management tool (Pillar 4). Initially sporadic (1940–1944), it became more systematic from the mid-1980s. Its increasing prominence in recent years (especially 2020–2024) reflects growing investment in instrumental monitoring systems, driven by both the perceived intensification of hazards and advances in measurement technologies. Since 1986, the eastern face of Monte Rosa (IT) has been regularly monitored due to increasing mass movements (Fischer et al., 2006). Other emblematic sites have since been equipped, including Aiguille du Midi (FR) since 2005, the Matterhorn south face (IT) and Les Drus (FR) since 2007.

Finally, projects such as the EU AlpinSpace *PermaNET* program and the *PACE* project have contributed to structuring knowledge on Alpine permafrost, through harmonised datasets and coordinated monitoring across the Alps (Harris et al., 2001; Mair et al., 2011).



436
437 **Figure 2.** Temporal evolution of risk management for rockfalls / rock avalanches.
438

439 The evolution over time of risk management for major GLOF events in the Alps is presented in Figure
440 5. Unlike other hazards, GLOFs are documented as early as the 16th century, reflecting their high
441 visibility and destructing impacts in populated valleys. Their severity ensured systematic recording in
442 administrative and local archives, especially as their transformation into debris flows amplifies
443 downstream damage. The Bagnes valley and the Giétro glacier, for example, generated extensive records
444 from the 16th century onwards due to the scale of the destruction (Ancy et al., 2019).


445 The period 1850-1889 is absent from the graph, which is paradoxical from a climatic perspective.
446 Although post-Little Ice Age warming favoured lake formation and potentially increased GLOF
447 occurrence, this gap likely reflects missing data rather than reduced activity.

448 Efforts to reduce vulnerability and exposure appear earlier for GLOFs than for other hazards. Early
449 responses relied on direct engineering interventions (drainage, tunnels, dykes) in densely populated
450 valleys, preceding formalised risk management frameworks. This reflects a pragmatic, engineering-
451 driven approach shaped by local socio-economic constraints. Thus, from the late 16th century, recurrent
452 outbursts of the Margherita glacial lake (Rutor glacier, IT) prompted studies commissioned by the Duke
453 of Savoy (12 events between 1430 and 1680; discharge of 5×10^6 m³ in 1751). Technical solutions such
454 as dam and drainage tunnels were already considered in the 19th century (Vergnano et al., 2023).

455 Another notable example is the Giétro glacier event in 1818. A tunnel was constructed to drain the ice-
456 dammed lake, thereby reducing the magnitude of the floods and preventing major downstream damage
457 (Wiegandt and Lugon, 2008).

458 These recurrent events progressively led to diversified measures, including dykes, dams, storage basins,
459 subglacial drilling, siphoning, lake drainage, and moraine stabilisation, as well as warning systems. In
460 parallel, glacier monitoring became increasingly formalised, notably through photogrammetric surveys
461 of the Vernagtferner (AU), one of the best-documented glaciers due to its frequent GLOFs, the most
462 notable of which dates back to 1848 (Braun, 1995; *Gridabase - Glaciorisk*, s. d.).

463 However, the integration of populations through preventive information is more recent. For instance,
464 the 1986 Arsine proglacial lake case (FR) marked one of the first instances where scientific warning led
465 to coordinated crisis preparedness involving local populations (Lailly and Demolis, 2019).
466



467 **Figure 3.** Temporal evolution of risk management for Glacial Lake Outburst Floods.

468
469
470 4.2.3 Disruptive events that have improved the glacial and periglacial risk management and
471 understanding

472
473 The **construction** of a timeline tracing risk management responses to past events **has enabled the**
474 **identification of key turning points marking disruptions** and/or the emergence of new approaches.
475 Selected examples correspond to first documented implementations of new management practices
476 (**prediction, monitoring, mitigation, and coordinated crisis response**).

477
478 **Scale change**

479 Risk management has shifted from local (village-scale) responses to national frameworks. The Giétro
480 glacier disaster in 1818 illustrates this transition (Ancey et al., 2019). At that time, glaciology was still
481 emerging, yet the event demonstrated that improved process understanding could support more effective
482 interventions. Scientific advances progressively supported the development of preventive measures and
483 favoured the emergence of regional and national policies, replacing earlier reactive local approaches
484 (Wiegandt and Lugon, 2008). The disaster also marked a shift in risk perception, as hazards began to be
485 understood as natural processes rather than divine punishment (Carrivick and Tweed, 2016).

486
487 **Monitoring and surveillance**

488 The first known glacier inventory in the Alps dates back to the late 19th century (Vernagtferner), an area
489 characterised by frequent GLOFs over the past 400 years (Hoinkes, 1969). **Systematic photogrammetric**
490 **surveys over a century enabled the first detailed analyses of glacier dynamics** (Braun, 1995). Monitoring
491 intensified during the 20th century; the Giétro glacier has been monitored since the 1960s, including
492 measurements of flow velocity, structure, and mass balance. More recently, heavily instrumented sites
493 such as the Matterhorn have integrated multi-sensor monitoring systems (Weber et al., 2025) following
494 **recurrent instabilities** (Occhiena and Pirulli, 2012).

495
496 **Multidisciplinary network**

497 The 1892 Tête Rousse disaster was a foundational event in glacier risk management. The debris flow
498 has affected several villages downstream, causing 175 victims (Mougin and Bernard, 1922). It led to the

499 formation of multidisciplinary teams combining engineers, glaciologists, and foresters to investigate
500 causes and implement mitigation measures (Marco et al., 2012; Sirop et al., 2022).
501 Similarly, the 1965 Allalin glacier collapse at Mattmark (88 fatalities; Vivian, 1966) triggered
502 coordinated scientific and engineering responses (Dalban Canassy et al., 2011). This led to the creation
503 of a national working group on glacier hazards in Switzerland and the development of systematic hazard
504 assessment methods (Haeberli, 1983; Haeberli et al., 2004). A coalition of scientists has irrevocably
505 advanced the understanding of the mechanism of phased glacial rupture and sparked discussion on
506 practical methods of risk assessment at the international level (Huggel et al., 2004).
507 The Rochemelon lake case further illustrates this coordination. Cross-border collaboration enabled
508 monitoring, risk assessment, public warning, and controlled drainage of the lake (Vincent et al., 2010).
509 This case highlighted the need for continuity between scientific knowledge, decision-making, and
510 operational action (Marco et al., 2012, unpublished).

511 512 ***Reduction of exposure and vulnerability (evacuations, works, etc.)***

513 The first known organised evacuations for gravitational hazards occurred in 1987 (Val Pola rockslide;
514 $30 \times 10^6 \text{ m}^3$; Dramis et al., 1995). Early warnings based on slope instability enabled preventive
515 evacuation (Chardon, 1990). For glacial hazards, evacuations such as Planpincieux (1997) were based
516 on scenario modelling and monitoring data (Margreth and Funk, 1999; Margreth et al., 2011). These
517 approaches marked the emergence of anticipatory risk management. Engineering measures also
518 developed, as illustrated by the Belvedere glacier (IT) and the Effimero lake crisis (2002), where
519 combined strategies (monitoring, mapping, evacuation planning, and artificial drainage) were
520 implemented (Haeberli et al., 2002).

521 522 ***Predictive study***

523 In 1973, the first successful glacier collapse prediction was achieved at Weisshorn (CH) (Röthlisberger,
524 1981). Scientists observed an acceleration of the upper part of the glacier, accompanied by warnings
525 from local mountain guides (Faillettaz and Funk, 2013). Subsequent advances enabled short-term
526 predictions based on displacement monitoring, such as the Whympfer glacier collapse forecast 10 days
527 in advance (Faillettaz et al., 2016). Authorities adapted responses accordingly, restricting access while
528 avoiding unnecessary evacuation. These decisions increasingly relied on modelling and scenario-based
529 approaches (Schindelegger and Kanonier, 2019).
530 These developments are part of broader research programmes (e.g. *Glaciorisk*, *PERMAdataROC*,
531 *GlariskAlp*, *PAPROG*, etc.), aimed at improving hazard understanding and management.

532 533 ***Institutional risk understanding***

534 Swiss National Research Programmes (NRPs), significantly contributed to the institutionalisation of risk
535 knowledge. Projects under NRP 31, 48, and 61 advanced understanding of climate-driven hazards,
536 spatial planning tools, and emerging glacial lakes (Haeberli et al., 1999, 2016; Rothenbühler, 2006).
537 More broadly, Alpine risk management has evolved from “hazard protection” to “hazard management”
538 and ultimately to “risk management” (Link and Stötter, 2015).

539 Before 1950, strategies focused on structural protection (protective works) and exposure reduction
540 (legislation). There are numerous traces of such works in 19th century archives. For example, galleries
541 were excavated within the ice of the Allalingletscher glacier in 1834 to lower the level of its lake (a
542 glacial-moraine dam) and reduce the already high frequency of ice break-offs (Mariétan, 1953). Between
543 1589 and 1850, documents report 26 major sudden outburst events (Raymond et al., 2003).

544 After 1950, improved process understanding led to hazard mapping (Link and Stötter, 2015) and
545 probabilistic approaches; this was the age of *hazard management*.

546 Since the late 20th century, risk management has adopted a holistic perspective (Link and Stötter, 2015),
547 integrating vulnerability and risk acceptance. Before the 2000s, hazard management was considered *in*
548 *response* to an event. Recent approaches increasingly consider cascading processes (e.g. Evans et al.,

549 2009) and multiple scenarios, reflecting the growing complexity of cryospheric hazards under climate
550 change (Allen et al., 2022).

551 Historically, the scope of hazard assessment has progressively expanded from glacier-related processes
552 to include permafrost degradation and emerging landforms such as proglacial lakes. Current frameworks
553 now adopt integrated, forward-looking approaches that account for long-term cryosphere evolution and
554 cascading risks (Allen et al., 2017).

555

556 4.2.4 Proactive vs. reactive management

557

558 Our typology of glacial and periglacial risk management distinguishes between proactive and reactive
559 approaches. Management is *considered proactive* when it relies on anticipatory measures to prevent
560 potential risks. Within the Disaster Risk Management (DRM) framework, prevention, mitigation
561 (including spatial planning, protection measures), long-term adaptation to natural and socioeconomic
562 changes, and the development of response and recovery capacities are all proactive strategies (Tagarev
563 et al., 2021).

564 Management is *reactive* when it occurs during or after an event, aiming to manage the emergency,
565 analyse impacts, and improve future risk reduction measures (Tagarev et al., 2021).

566 The different principles of glacial and periglacial hazards management can operate independently but
567 are often combined. For instance, Monitoring and Surveillance (Pillar 4) is frequently coupled with
568 Exposure, Vulnerability or Hazard Mitigation (Pillar 5).

569 Regarding torrential hazards associated with glacial lakes, a proactive example is the 1985 warming on
570 the Arsine proglacial lake, which led to partial artificial drainage (Peissier and Courtray, 2012).
571 Similarly, an early warning system installed in 2008 at the Grindelwaldgletscher glacial lake (CH) based
572 on pressure sensors, enabled automatic alerts and proved effective during a drainage event (Bauder,
573 2017).

574 Conversely, the 1964 overflow of the Patinoire lake (FR), which caused damage downstream as far as
575 the village of Pralognan, prompted a reactive response. Post-event analyses included field surveys,
576 susceptibility studies, and impact modelling, leading to its later designation as a priority monitoring site
577 (Cathala et al., 2021).

578 The Whymper hanging glacier illustrates proactive management: in 1997, displacement measurements
579 enabled prediction of an imminent collapse, prompting evacuation of Planpincieux (Margreth and Funk,
580 1999). Continuous monitoring and preparedness measures have since been implemented (Faillettaz et
581 al., 2015, 2016). Similarly, the Forni glacier is monitored using satellite and terrestrial laser scanning to
582 detect instabilities (Fugazza et al., 2018).

583 In contrast, the 1965 Mattmark disaster (Allalin glacier) exemplifies reactive management. Authorities
584 focused on emergency response and post-event investigation. The event later triggered political debate
585 and highlighted the limits of predictability at the time (Ricciardi, 2016).

586 For rock avalanches, the Piz Cengalo case illustrates the transition from reactive to proactive
587 management. Following a first event in 2011, monitoring systems and protective infrastructure were
588 implemented (Walter et al., 2020). These included a storage basin, protective barriers, and early warning
589 systems enabling rapid evacuation (Bohnenblust, 2017).

590 In 2017, a major rock avalanche and subsequent debris flows ($3.1 \times 10^6 \text{ m}^3$) affected the Val Bondasca.
591 Despite underestimation of flow magnitude, early warning systems and protective measures
592 significantly reduced damage (OECD, 2018). This case demonstrates the effectiveness of combining
593 past-event analysis, monitoring, and preparedness, although some casualties still occurred (Walter et al.,
594 2020).

595

596

597 5 Discussions

598

599 5.1 Cascading processes affecting increasingly vulnerable populations

600
601 As shown by the events documented in this study, chain reactions in glacial and periglacial environments
602 tend to amplify beyond their initial triggering processes (OECD, 2018; Jacquemart et al., 2024), posing
603 increasing challenges for risk management. This trend is closely linked to rising exposure in Alpine
604 region. Mountain valleys have experienced significant urbanisation and infrastructure development in
605 recent decades, increasing pressure on historically exposed areas (Hock and Rasul, 2022). This growing
606 exposure is compounded by systemic vulnerability due to infrastructure interdependence: the destruction
607 of a road, bridge, or energy network during a cascading event can isolate communities and hinder
608 response capacities, as illustrated by La Bérarde in 2024 (Blanc et al., 2024). Risk management systems
609 have long treated hazards separately, but this segmented approach is increasingly inadequate for
610 complex, interacting phenomena affecting multiple components of populated territories. There is
611 therefore a need for integrated approaches that model process interactions while accounting for the
612 spatial distribution of populations and infrastructure (Tacnet et al., 2010; Nyberg et al., 2026). These
613 challenges are further intensified by climate change, which increases the likelihood of cascading
614 processes in densely populated Alpine valleys.

615

616 5.2 Biases linked to difficulties in compiling an exhaustive inventory of events

617

618 Compiling an inventory that aims to be exhaustive according to defined selection criteria entails
619 limitations and requires contextualisation of each event.

620

621 *The data source effect*

622 The apparent increase in events over time partly reflects biases related to data availability (e.g. Giacona
623 et al., 2017). This *data source effect* is closely linked to the nature and distribution of information
624 sources. As Giacona (2019) notes, the spatial and temporal distribution of events depends strongly on
625 the structure of the documentary corpus. Each recorded event therefore depends on the quality and
626 diversity of available sources. Scarcity of historical records limits traceability in earlier periods, whereas
627 recent data abundance increases event detection.

628 The data source effect thus requires careful consideration of variations in documentation volume. For
629 instance, technological advances such as webcam monitoring (e.g. Gornergletscher) generate continuous
630 and detailed observations, increasing the number of recorded events (Huss et al., 2007).

631

632 *Land use and the event prism*

633 The presence or absence of sources is also influenced by human occupation. The growing presence of
634 people in high mountains – through mountaineering (Mourey et al., 2023) and valley urbanisation
635 (Vannier et al., 2016) – has significantly increased observation capacity. Conversely, abandoned
636 settlements and pastures were once important witnesses to past hazards.

637 Even when hazards occur, their recognition depends on societal perception. An event may remain
638 unrecorded if it is not perceived as significant. As Giacona (2019) argues, an event gains social existence
639 only when it is perceived as a disruption. Risk perception evolves over time and is shaped by collective
640 experience, knowledge, and social context (Granet-Abisset, 2012). Increasing attention to high-
641 mountain hazards also reflects rising societal expectations for safety and accountability, which in turn
642 drive the production and diversification of information sources.

643

644 5.3 Resilience and response as late-developed concepts rather than absent practices

645

646 The emergence of Pillar 8 (Resilience) from the 1980s-90s onwards, and its consolidation after 2000,
647 should be interpreted cautiously. It does not indicate a prior absence of resilience from Alpine
648 communities, but rather the late institutionalisation of the concept in scientific and policy frameworks.
649 Post-event reconstruction dynamics existed well before this period, but were not conceptualised within
650 this framework, introducing an interpretative bias in diachronic analysis.

651 Similarly, Pillar 7 (Response) appears from the 1950s and remains present thereafter, without becoming
652 dominant. Its consistent but limited representation suggests that emergency response has always existed
653 but is less documented due to lower formalisation, leading to potential under-representation in the
654 database.

655 The use of a ninth separate pillar representing ‘*Experience feedback*’ was considered, since the
656 experience gained from an event enables a “structured process of capitalising on and utilising knowledge
657 resulting from the analysis of positive and/or negative events” (Villeneuve et al., 2010).

658 Feedback is the very condition for an event’s existence in archives: its narration and memory make it
659 historically visible. It therefore applies to every event, and would have created a bias in relation to the
660 other pillars. Therefore, it is implicitly included in Pillar 1.

661

662 5.4 National discrepancies in prevention

663

664 The study highlights that prevention measures are not uniform across Alpine countries and reveals a
665 geographical bias towards France, Switzerland, and Italy. This heterogeneity reflects significant
666 differences in legislative, institutional, and cultural frameworks for risk management.

667 In France, spatial planning relies on binding national regulations, and scientific action plans are also
668 defined at the national level. In Switzerland, management is largely devolved to the cantons,
669 encouraging locally adapted approaches but potentially generating inequalities depending on available
670 resources. In Italy, administrative fragmentation and regional disparities hinder the implementation of
671 coherent policies at the scale of catchments.

672 These governance differences affect both data comparability and the ability to derive cross-cutting
673 insights. More broadly, they highlight the need for stronger transnational Alpine governance. Given that
674 hazards transcend political borders, enhanced institutional coordination is essential for effective and
675 equitable risk prevention.

676

677

678 6 Conclusions and perspectives

679

680 This study **moves beyond** traditional approaches that focus primarily on post-event causes or hazards
681 characterization, and instead examines management **practices**. While event inventories **inform** on
682 frequency, they rarely capture organisational dynamics or adaptative capacities. Our analysis addresses
683 these dimensions.

684 By analysing 200 events since the early 19th century (some even earlier, though less frequent), we
685 observe both an intensification of events and a **progressive formalisation of management practices**,
686 reflected in archival records. **The growing number of simultaneously mobilised pillars in recent years**
687 **indicates a shift from single-discipline approaches to multi-actor, multi-scale governance. Beyond**
688 **increased event reporting, this reflects the maturation of institutional Alpine risk management**
689 **frameworks**. Although all high-altitude massifs are affected, our data highlights a concentration of major
690 events in France and western Switzerland, particularly GLOFs. **The completeness of this long-term**
691 **inventory remains constrained by sources and perception biases**.

692 Risk management has evolved in response to both major events and improved process understanding.
693 Forecasting, instrumentation, and monitoring of unstable zones have expanded, **alongside contingency**
694 **planning and evacuation strategies. Measures to reduce exposure and vulnerability increasingly account**
695 **for evolving hazards and urbanisation pressures**. Since the early 21st century, scientific initiatives aimed
696 at anticipating high-mountain mass movements that have multiplied. **Knowledge exchange across**
697 **borders has intensified, supporting more integrated and holistic risk management approaches**. Hazards
698 are now increasingly analysed as interconnected or cascading phenomena. As risk management grows
699 more complex, the social sciences emerge as an indispensable complement to natural sciences, bringing
700 to the fore dimensions – community vulnerability, risk perception, territorial identity, institutional
701 dynamics – that physical approaches alone cannot capture.

702 Switzerland appears as a leading actor in the institutionalisation of monitoring and hazard assessment,
703 particularly for glaciers and rock faces. Multidisciplinary collaboration and international knowledge
704 exchange are rapidly advancing.

705 Across many regions worldwide, glacial and permafrost environments now differ markedly from the
706 conditions under which settlements and infrastructure developed. Historical and empirical knowledge
707 alone is therefore insufficient to anticipate future glacier- and permafrost-related risks. This calls for
708 integrated, forward-looking approaches, based on continuous observation, repeated expert assessments,
709 and rapidly evolving technologies. Risk analysis must adopt a systemic perspective, covering the full
710 chain from hazard sources to socio-economic impacts.

711

712

713 *Author contributions.* JB, LR and SC designed the study and JB carried it out. JB prepared the
714 manuscript with contributions from all co-authors.

715

716 *Competing interests.* The authors declare that they have no conflict of interest.

717

718

719 **References**

720 Alean, J.: Ice Avalanches: Some Empirical Information about their Formation and Reach, *J. Glaciol.*,
721 31, 324–333, <https://doi.org/10.3189/S002214300006663>, 1985.

722 Allen, S., Frey, H., and Huggel, C.: Assessment of Glacier and Permafrost Hazards in Mountain
723 Regions. Technical Guidance Document, <https://doi.org/10.13140/RG.2.2.26332.90245>, 2017a.

724 Allen, S., Frey, H., and Huggel, C.: GAPHAZ 2017: Assessment of Glacier and Permafrost Hazards in
725 Mountain Regions – Technical Guidance Document., Glacier and Permafrost Hazards in Mountains
726 (GAPHAZ), International Association of Cryospheric Sciences (IACS) and the International
727 Permafrost Association (IPA) - Zurich, Switzerland / Lima, Peru, 2017b.

728 Allen, S., Frey, H., Haeberli, W., Huggel, C., Chiarle, M., and Geertsema, M.: Assessment Principles
729 for Glacier and Permafrost Hazards in Mountain Regions, in: *Oxford Research Encyclopedia of*
730 *Natural Hazard Science*, <https://doi.org/10.1093/acrefore/9780199389407.013.356>, 2022.

731 Ancy, C., Bardou, E., Funk, M., Huss, M., Werder, M. A., and Trehwela, T.: Hydraulic
732 Reconstruction of the 1818 Giétro Glacial Lake Outburst Flood, *Water Resour. Res.*, 55, 8840–8863,
733 <https://doi.org/10.1029/2019WR025274>, 2019.

734 **Geoclimalp Landslide Inventory:** <http://geoclimalp.to.cnr.it/landslide-inventory/#6/43.659/13.554>, last
735 access: 4 March 2026.

736 Gridabase - Glaciorisk: <https://www.nimbus.it/glaciorisk/gridabasemainmenu.asp>, last access: 27 May
737 2025.

738 Bauder, A.: The Swiss Glaciers 2013/14 and 2014/15 - Glaciological Report No. 135/136, 145pp,
739 https://doi.org/10.18752/GLREP_135-136, 2017.

740 Beccera, S. and Peltier, A.: L'information préventive pour réduire la vulnérabilité aux risques
741 d'inondation, élaboration et efficacité d'une réponse sociale, *Chang. Clim. Métarisque À Méta-Gouv.*,
742 pp.35-53, 2011.

743 Bisquert, A., Mainieri, R., Carlados, S., Robert, Y., Giacona, F., Verry, P., and Eckert, N.: La base de
744 données événementielles RTM pour la connaissance des risques naturels en montagne : L'exemple du
745 département de l'Isère (France), *J. Alp. Res. Rev. Géographie Alp.*, <https://doi.org/10.4000/137kf>,
746 2025.

- 747 Blanc, A., Misset, C., Mainieri, R., and Llamas, B.: Rétro-analyse de la crue du torrent des Etançons
748 du 21 juin 2024, RTM, 2024.
- 749 Bohnenblust, S.: Radar system monitors Pizzo Cengalo, Bondo, GEOPRÆVENT AG – Electron.
750 Monit. Nat. Hazards, 2017.
- 751 Bondesan, A. and Francese, R. G.: The climate-driven disaster of the Marmolada Glacier (Italy),
752 *Geomorphology*, 431, 108687, <https://doi.org/10.1016/j.geomorph.2023.108687>, 2023.
- 753 Boudières, V. and Peisser, C.: Prévention des Risques d’Origine Glaciaire et Périglaciaire (ROGP) -
754 Synthèse des connaissances et des pratiques, PARN, 2013.
- 755 Bourrelier, P.-H. and Dunglas, J.: Des événements naturels extrêmes aux figures de la catastrophe.,
756 *Ann. Mines - Responsab. Environ.*, N° 56, 41–47, <https://doi.org/10.3917/re.056.0041>, 2009.
- 757 Braun, L. N.: Les résultats principaux de 100 ans de recherche au Vernagtferner, Oetzal (Autriche),
758 *Houille Blanche*, 81, 109–110, <https://doi.org/10.1051/lhb/1995054>, 1995.
- 759 Büntgen, U., Oppenheimer, C., Farinotti, D., Nahtz, T., and Esper, J.: The 2025 Blatten disaster in the
760 Swiss Alps followed exceptional warming and highlights the vulnerability of people and heritage in
761 glaciated landscapes, *Commun. Earth Environ.*, 6, 994, <https://doi.org/10.1038/s43247-025-02994-8>,
762 2025.
- 763 Capozzi, C.: Mémoire de master en histoire - La catastrophe de Mattmark par la presse, Université de
764 Franche-Comté, 2011.
- 765 Carrivick, J. L. and Tweed, F. S.: A global assessment of the societal impacts of glacier outburst
766 floods, *Glob. Planet. Change*, 144, 1–16, <https://doi.org/10.1016/j.gloplacha.2016.07.001>, 2016.
- 767 Cathala, M., Magnin, F., Linsbauer, A., and Haeberli, W.: Modelling and characterizing glacier-bed
768 overdeepenings as sites for potential future lakes in the deglaciating French Alps, *Géomorphologie
769 Relief Process. Environ.*, 27, 19–36, <https://doi.org/10.4000/geomorphologie.15255>, 2021.
- 770 Cathala, M., Bock, J., Abdulsamad, F., Deline, P., Josnin, J.-Y., Ravanel, L., Revil, A., Richard, J.,
771 Verroust, F., and Magnin, F.: Assessing the role of permafrost in the preconditioning and triggering
772 factors of the September 2020 Crête des Grangettes rockfall (southern French Alps), *Géomorphologie
773 Relief Process. Environ.*, 30, <https://doi.org/10.4000/12yqn>, 2024.
- 774 Cauliez, R.: Mémoire M2 - Au-delà des eaux glacées. Pouvoirs gestionnaires et expériences populaires
775 sous le glacier de Tête-Rousse., Ecole Normale Supérieure, 229 pp., 2024.
- 776 Chardon, M.: Les catastrophes naturelles de l’été 1987 en Lombardie : crues, inondations, écoulement
777 de Val Pola, *Rev. Géographie Alp.*, 78, 59–87, <https://doi.org/10.3406/rga.1990.2767>, 1990.
- 778 Chiarle, M., Viani, C., Mortara, G., Deline, P., Tamburini, A., and Nigrelli, G.: Large glacier failures in
779 the Italian Alps over the last 90 years, *Geogr. Fis. E Din. Quat.*, 45, 19–40,
780 <https://doi.org/10.4461/GFDQ.2022.45.2>, 2022.
- 781 Dalban Canassy, P., Bauder, A., Dost, M., Fäh, R., Funk, M., Margreth, S., Müller, B., and Sugiyama,
782 S.: Hazard assessment investigations due to recent changes in Triftgletscher, Bernese Alps,
783 Switzerland, *Nat. Hazards Earth Syst. Sci.*, 11, 2149–2162, [https://doi.org/10.5194/nhess-11-2149-](https://doi.org/10.5194/nhess-11-2149-2011)
784 2011, 2011.
- 785 Dauphiné, A. and Provitolo, D.: Risques et catastrophes, Armand Colin,
786 <https://doi.org/10.3917/arco.dauph.2013.01>, 2013.

- 787 Davide, F., Antonella, S., Roberto Sergio, A., Carlo, D., Daniele, C. B., Luca, M., Fabiano, V.,
788 Claudio, S., and Guglielmina Adele, D.: Variations of Lys Glacier (Monte Rosa Massif, Italy) from the
789 Little Ice Age to the Present from Historical and Remote Sensing Datasets, in: *Glaciers and the Polar*
790 *Environment*, edited by: Kanao, M., Godone, D., and Dematteis, N., IntechOpen,
791 <https://doi.org/10.5772/intechopen.91202>, 2021.
- 792 Defossez, S., Vinet, F., and Leone, F.: Diagnostiquer la vulnérabilité face aux inondations : progrès et
793 limites, in: *Inondations 1: la connaissance juridique*, 2018.
- 794 Dematteis, N., Giordan, D., Troilo, F., Wrzesniak, A., and Godone, D.: Ten-Year Monitoring of the
795 Grandes Jorasses Glaciers Kinematics. Limits, Potentialities, and Possible Applications of Different
796 Monitoring Systems, *Remote Sens.*, 13, 3005, <https://doi.org/10.3390/rs13153005>, 2021.
- 797 Dematteis, N., Troilo, F., Scotti, R., Colombarolli, D., Giordan, D., and Maggi, V.: The use of
798 terrestrial monoscopic time-lapse cameras for surveying glacier flow velocity, *Cold Reg. Sci. Technol.*,
799 222, 104185, <https://doi.org/10.1016/j.coldregions.2024.104185>, 2024.
- 800 Demolis, B., Kuss, D., and Serrano, C.: Etude du lac de la Patinoire à Pralognan la Vanoise (73), ONF-
801 RTM INRAE, 2021.
- 802 Desailly, B.: Crues et inondations en Roussillon : le risque, le discours et l'aménagement, *Rev.*
803 *Géographique Pyrén. Sud-Ouest*, 61, 515–519, <https://doi.org/10.3406/rgpso.1990.3229>, 1990.
- 804 **Dipartimento della Protezione Civile: Rischio glaciale e periglaciale in ambiente alpino: un quadro**
805 **metodologico, Prima., 126 pp., 2025.**
- 806 Dollfus, O. and D'Ercole, R.: Les mémoires des catastrophes au service de la prévision et de la
807 prévention des risques naturels, 7–18, 1996.
- 808 Dosse, F.: Renaissance de l'événement, 1^{re} éd., Presses Universitaires de France,
809 <https://doi.org/10.3917/puf.dosse.2010.01>, 2010.
- 810 Dourlens, C.: La question des inondations au prisme des sciences sociales : un panorama de la
811 recherche publique, Ministère de l'Équipement, des Transports, du Logement, du Tourisme et de la
812 Mer, 112 pp., 2004.
- 813 **Dramis, F., Govi, M., Guglielmin, M., and Mortara, G.: Mountain Permafrost and Slope Instability in**
814 **the Italian Alps: the Val Pola Landslide, 6, 73–82, 1995.**
- 815 Du Pasquier, L.: L'avalanche du glacier de l'Altels le 11 septembre 1895, *Ann. Géographie*, 5, 458–
816 468, <https://doi.org/10.3406/geo.1896.5954>, 1896.
- 817 Einhorn, B.: Changement climatique et risques naturels - Prévention, Gestion intégrée, Adaptation,
818 2017.
- 819 Einhorn, B. and Peisser, C.: Gestion intégrée du risque généré par les poches d'eau du glacier de Tête
820 Rousse – De la tragédie de 1892 à la gestion de crise de 2010, Grenoble, 2011.
- 821 **Etzelmüller, B., Guglielmin, M., Hauck, C., Hilbich, C., Hoelzle, M., Isaksen, K., Noetzli, J., Oliva,**
822 **M., and Ramos, M.: Twenty years of European mountain permafrost dynamics—the PACE legacy,**
823 ***Environ. Res. Lett.*, 15, 104070, <https://doi.org/10.1088/1748-9326/abae9d>, 2020.**
- 824 **Evans, S. G., Tutubalina, O. V., Drobyshev, V. N., Chernomorets, S. S., McDougall, S., Petrakov, D.**
825 **A., and Hungr, O.: Catastrophic detachment and high-velocity long-runout flow of Kolka Glacier,**
826 **Caucasus Mountains, Russia in 2002, *Geomorphology*, 105, 314–321,**
827 **<https://doi.org/10.1016/j.geomorph.2008.10.008>, 2009.**

- 828 Faillettaz, J. and Funk, M.: Instabilités glaciaires et prédiction, *Société Vaudoise Sci. Nat.*, 159–174,
829 2013.
- 830 Faillettaz, J., Funk, M., and Vincent, C.: Avalanching glacier instabilities: Review on processes and
831 early warning perspectives, *Rev. Geophys.*, 53, 203–224, <https://doi.org/10.1002/2014RG000466>,
832 2015.
- 833 Faillettaz, J., Funk, M., and Vagliasindi, M.: Time forecast of a break-off event from a hanging glacier,
834 *The Cryosphere*, 10, 1191–1200, <https://doi.org/10.5194/tc-10-1191-2016>, 2016.
- 835 Favier, R.: Sociétés urbaines et culture du risque. Les inondations dans la France d’Ancien Régime, in:
836 *Les cultures du risque (XVIe-XXIe siècle)*, pp.49-86, 2006.
- 837 Field, C., Barros, V., Stocker, T., and Dahe, Q.: Managing the risks of extreme events and disasters to
838 advance climate change adaptation, Cambridge university press, New York, 582 pp., 2012.
- 839 Fischer, L., Käab, A., Huggel, C., and Noetzli, J.: Geology, glacier retreat and permafrost degradation
840 as controlling factors of slope instabilities in a high-mountain rock wall: the Monte Rosa east face,
841 *Nat. Hazards Earth Syst. Sci.*, 6, 761–772, <https://doi.org/10.5194/nhess-6-761-2006>, 2006.
- 842 Flotron, A.: Movement Studies on a Hanging Glacier in Relation with an Ice Avalanche, *J. Glaciol.*,
843 19, 671–672, <https://doi.org/10.3189/S0022143000029592>, 1977.
- 844 Forel, F. A.: *L’Eboulement du glacier des Altels*, Bureau des Archives, Genève, 513 pp., 1895.
- 845 Fournier, M.: Le riverain introuvable ! La gestion du risque d’inondation au défi d’une mise en
846 perspective diachronique : une analyse menée à partir de l’exemple de la Loire., *Aménagement de*
847 *l’Espace et Urbanisme*, Université François-Rabelais, Tours, 432 pp., 2010.
- 848 Francese, R. G., Valentino, R., Haeberli, W., Bondesan, A., Giorgi, M., Picotti, S., Pettenati, F.,
849 Sandron, D., Ramponi, G., and Valt, M.: Failure of Marmolada Glacier (Dolomites, Italy) in 2022:
850 Data-based back analysis of possible collapse mechanisms as related to recent morpho-climatic
851 evolution and possible trigger factors, <https://doi.org/10.5194/nhess-2024-212>, 19 November 2024.
- 852 Fugazza, D., Scaioni, M., Corti, M., D’Agata, C., Azzoni, R. S., Cernuschi, M., Smiraglia, C., and
853 Diolaiuti, G. A.: Combination of UAV and terrestrial photogrammetry to assess rapid glacier evolution
854 and map glacier hazards, *Nat. Hazards Earth Syst. Sci.*, 18, 1055–1071, <https://doi.org/10.5194/nhess-18-1055-2018>, 2018.
- 856 Giacona, F., Eckert, N., and Martin, B.: A 240-year history of avalanche risk in the Vosges Mountains
857 based on non-conventional (re)sources, *Nat. Hazards Earth Syst. Sci.*, 17, 887–904,
858 <https://doi.org/10.5194/nhess-17-887-2017>, 2017.
- 859 Giacona, F., Martin, B., Eckert, N., and Desarthe, J.: Une méthodologie de la modélisation en
860 géohistoire : de la chronologie (spatialisée) des événements au fonctionnement du système par la mise
861 en correspondance spatiale et temporelle, *Physio-Géo*, 171–199, <https://doi.org/10.4000/physio-geo.9186>, 2019.
- 863 Giardino, M., Ratto, S., Alberto, W., Armand, M., Cignetti, M., and Navillod, E.: Regional and local
864 risk assessments of alluvial fans by combination of historical and geomorphological data on debris
865 flows, the most damaging natural hazard in the Aosta Valley Region (NW-Italy), 2010.
- 866 Giordan, D., Dematteis, N., Allasia, P., and Motta, E.: Classification and kinematics of the
867 Planpincieux Glacier break-offs using photographic time-lapse analysis, *J. Glaciol.*, 66, 188–202,
868 <https://doi.org/10.1017/jog.2019.99>, 2020.

- 869 Girard, S. and Rivière-Honegger, A.: Le choix et la pratique de la monographie diachronique.
870 Contribution à l'étude de l'efficacité environnementale de la territorialisation de la politique de l'eau.,
871 in: Environnement, politiques publiques et pratiques locales, 359–384, 2015.
- 872 Granet-Abisset, A.-M.: L'historien, les risques et l'environnement : un regard sur la nature et les
873 hommes, in: 23èmes Journées Scientifiques de l'Environnement - Risques environnementaux :
874 détecter, comprendre, s'adapter, 2012.
- 875 Gruber, S. and Haeberli, W.: Permafrost in steep bedrock slopes and its temperature-related
876 destabilization following climate change, *J. Geophys. Res. Earth Surf.*, 112, 2006JF000547,
877 <https://doi.org/10.1029/2006JF000547>, 2007.
- 878 Haeberli, W.: Frequency and Characteristics of Glacier Floods in the Swiss Alps, *Ann. Glaciol.*, 4, 85–
879 90, <https://doi.org/10.3189/S0260305500005280>, 1983.
- 880 Haeberli, W., Käab, A., Hoelzle, M., Bösch, H., Funk, M., Mühl, D., and Keller, F.: Eisschwund und
881 Naturkatastrophen im Hochgebirge, vdf Hochschulverlag an der ETH Zürich, Zürich, 190pp pp., 1999.
- 882 Haeberli, W., Käab, A., Mühl, D. V., and Teysseire, P.: Prevention of outburst floods from periglacial
883 lakes at Grubengletscher, Valais, Swiss Alps, *J. Glaciol.*, 47, 111–122,
884 <https://doi.org/10.3189/172756501781832575>, 2001.
- 885 Haeberli, W., Käab, A., Paul, F., Chiarle, M., Mortara, G., Mazza, A., Deline, P., and Richardson, S.: A
886 surge-type movement at Ghiacciaio del Belvedere and a developing slope instability in the east face of
887 Monte Rosa, Macugnaga, Italian Alps, *Nor. Geogr. Tidsskr. - Nor. J. Geogr.*, 56, 104–111,
888 <https://doi.org/10.1080/002919502760056422>, 2002.
- 889 Haeberli, W., Huggel, C., Käab, A., Gruber, S., Noetzi, J., and Zraggen-Oswald, S.: Proceedings of
890 the International Conference „High mountain hazard prevention", 2004.
- 891 Haeberli, W., Buetler, M., Huggel, C., Friedli, T. L., Schaub, Y., and Schleiss, A. J.: New lakes in
892 deglaciating high-mountain regions – opportunities and risks, *Clim. Change*, 139, 201–214,
893 <https://doi.org/10.1007/s10584-016-1771-5>, 2016.
- 894 Harris, C., Haeberli, W., Vonder Mühl, D., and King, L.: Permafrost monitoring in the high mountains
895 of Europe: the PACE Project in its global context, *Permafr. Periglac. Process.*, 12, 3–11,
896 <https://doi.org/10.1002/ppp.377>, 2001.
- 897 Hock, R. and Rasul, G.: *The Ocean and Cryosphere in a Changing Climate: Special Report of the*
898 *Intergovernmental Panel on Climate Change*, 1st ed., Cambridge University Press,
899 <https://doi.org/10.1017/9781009157964>, 2022.
- 900 Hoinkes, H. C.: Surges of the Vernagtferner in the Ötztal Alps since 1599, *Can. J. Earth Sci.*, 6, 853–
901 861, <https://doi.org/10.1139/e69-086>, 1969.
- 902 Hugerot, T., Astrade, L., Gauchon, C., and Ployon, E.: De la carte diachronique à la trajectoire
903 géohistorique : modélisation du changement paysager d'un cône torrentiel, *Mappemonde*,
904 <https://doi.org/10.4000/mappemonde.6704>, 2021.
- 905 Huggel, C., Haeberli, W., Käab, A., Bieri, D., and Richardson, S.: An assessment procedure for glacial
906 hazards in the Swiss Alps, *Can. Geotech. J.*, 41, 1068–1083, <https://doi.org/10.1139/t04-053>, 2004.
- 907 Huggel, C., Fischer, L., Schneider, D., and Haeberli, W.: Research advances on climate-induced slope
908 instability in glacier and permafrost high-mountain environments, *Geogr. Helvetica*, 65, 146–156,
909 <https://doi.org/10.5194/gh-65-146-2010>, 2010.

- 910 Huggel, C., Carey, M., Clague, J. J., and Kääh, A.: The high-mountain cryosphere: environmental
911 changes and human risks, Cambridge university press, Cambridge, United Kingdom, 363 pp., 2015.
- 912 Huss, M., Bauder, A., Werder, M., Funk, M., and Hock, R.: Glacier-dammed lake outburst events of
913 Gornersee, Switzerland, *J. Glaciol.*, 53, 189–200, <https://doi.org/10.3189/172756507782202784>, 2007.
- 914 Intergovernmental Panel On Climate Change (Ipcce) (Ed.): Annex I: Glossary, in: *Climate Change 2022*
915 *- Mitigation of Climate Change*, Cambridge University Press, 1793–1820,
916 <https://doi.org/10.1017/9781009157926.020>, 2023.
- 917 **IPCC: Global Warming of 1.5°C: IPCC Special Report on Impacts of Global Warming of 1.5°C above**
918 **Pre-industrial Levels in Context of Strengthening Response to Climate Change, Sustainable**
919 **Development, and Efforts to Eradicate Poverty, 1st ed., Cambridge University Press,**
920 **<https://doi.org/10.1017/9781009157940>, 2022.**
- 921 **IPCC: Climate Change 2022 – Impacts, Adaptation and Vulnerability: Working Group II Contribution**
922 **to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, 1st ed., Cambridge**
923 **University Press, <https://doi.org/10.1017/9781009325844>, 2023.**
- 924 Jacquemart, M., Weber, S., Chiarle, M., Chmiel, M., Cicoira, A., Corona, C., Eckert, N., Gaume, J.,
925 Giacona, F., Hirschberg, J., Kaitna, R., Magnin, F., Mayer, S., Moos, C., Van Herwijnen, A., and
926 Stoffel, M.: Detecting the impact of climate change on alpine mass movements in observational
927 records from the European Alps, *Earth-Sci. Rev.*, 258, 29,
928 <https://doi.org/10.1016/j.earscirev.2024.104886>, 2024.
- 929 Joffe, H. and Orfali, B.: De la perception à la représentation du risque: le rôle des médias, *Hermès*, n°
930 41, 121, <https://doi.org/10.4267/2042/8962>, 2005.
- 931 Kellerer-Pirklbauer, A., Slupetzky, H., and Avian, M.: Ice-avalanche impact landforms: the event in
932 2003 at the glacier nördliches bockkarkees, hohe tauern range, austria, *Geogr. Ann. Ser. Phys. Geogr.*,
933 94, 97–115, <https://doi.org/10.1111/j.1468-0459.2011.00446.x>, 2012.
- 934 Kienholz, H., Krummenacher, B., Kipfer, A., and Perret, S.: Aspects of Integral Risk Management in
935 Practice - Considerations with Respect to Mountain Hazards in Switzerland, 45–50, 2004.
- 936 **Lacroix, D., Mayet, L., and Roche, P.-A.: Risques d’origine glaciaire et périglaciaire - Elements en**
937 **soutien à un plan d’action, IGEDD - IGA - IGESR, 2022.**
- 938 Laïly, B. and Demolis, B.: Glaciers blancs à risques - Retours d’expériences sur certaines crises, ONF-
939 RTM, 2019.
- 940 Leone, F., Meschinet de Richemond, N., and Freddy, V.: Aléas naturels et gestion des risques, Presses
941 Universitaires de France, 288 p. pp., 2021.
- 942 Link, S. and Stötter, J.: The development of mountain risk governance: challenges for application,
943 <https://doi.org/10.5194/nhessd-3-429-2015>, 16 January 2015.
- 944 **Lucchesi, S., Fioraso, G., Bertotto, S., and Chiarle, M.: Little Ice Age and contemporary glacier extent**
945 **in the Western and South-Western Piedmont Alps (North-Western Italy), *J. Maps*, 10, 409–423,**
946 **<https://doi.org/10.1080/17445647.2014.880226>, 2014.**
- 947 Magnin, F., Ravel, L., Ben-Asher, M., Bock, J., Cathala, M., Duvillard, P.-A., Jean, P., Josnin, J.-Y.,
948 Kaushik, S., Revil, A., and Deline, P.: De l’observation des écroulements aux solutions
949 opérationnelles : près de deux décennies d’études sur les risques cryo-gravitaires dans le massif du
950 Mont-Blanc, *Rev. Géographie Alp.*, 111–2, <https://doi.org/10.4000/rga.11644>, 2023.

- 951 Mainieri, R., Blanc, A., Astrade, L., Baratier, A., Berthet, J., Deline, P., Le Roy, M., Misset, C., De
 952 Montety, F., Robert, Y., and Schoeneich, P.: Les aspects géomorphologiques de la crue torrentielle du
 953 torrent des Étançons à la Bérarde du 21 juin 2024, *Géomorphologie Relief Process. Environ.*, 31,
 954 <https://doi.org/10.4000/14ipc>, 2025.
- 955 Mair, V., Zischg, A., Lang, K., Tonidandel, D., Krainer, K., Kellerer-Pirklbauer, A., Deline, P.,
 956 Schoeneich, P., Cremonese, E., Pogliotti, P., Gruber, S., and Böckli, L.: PermaNET - Réseau
 957 d'observation du permafrost sur le long terme - Rapport de synthèse, Interpraevent, Klagenfurt, 2011.
- 958 Marco, O., Lailly, B., and Charles, F.: Mesures de protection et contraintes d'ingénierie en milieu
 959 glaciaire : expérience des services intervenus sur les cas récents - ONF-RTM, 2012.
- 960 Margreth, S. and Funk, M.: Hazard mapping for ice and combined snow/ice avalanches — two case
 961 studies from the Swiss and Italian Alps, 159–173, 1999.
- 962 Margreth, S., Faillettaz, J., Funk, M., Vagliasindi, M., Diotri, F., and Broccolato, M.: Safety concept
 963 for hazards caused by ice avalanches from the Whymper hanging glacier in the Mont Blanc Massif,
 964 *Cold Reg. Sci. Technol.*, 69, 194–201, <https://doi.org/10.1016/j.coldregions.2011.03.006>, 2011.
- 965 Margreth, S., Funk, M., Tobler, D., Dalban, P., Meier, L., and Lauper, J.: Analysis of the hazard caused
 966 by ice avalanches from the hanging glacier on the Eiger west face, *Cold Reg. Sci. Technol.*, 144, 63–
 967 72, <https://doi.org/10.1016/j.coldregions.2017.05.012>, 2017.
- 968 Mariétan, I.: Note de sciences naturelles sur la vallée de Saas, *Bull. de la Murithienne* 76, 88–102,
 969 1959.
- 970 Moles, A.: Notes pour une typologie des événements, *Communications*, 18, 90–96,
 971 <https://doi.org/10.3406/comm.1972.1261>, 1972.
- 972 Mougin, P. and Bernard, C.: Étude sur le glacier de Tête-Rousse. Les avalanches en Savoie., Tome 4,
 973 3–90, 1922.
- 974 Mourey, J., Clivaz, C., and Bourdeau, P.: Analyser l'évolution des pratiques sportives en montagne peu
 975 aménagée à partir des données de fréquentation des cabanes. Application aux Alpes Valaisannes, *Rev.*
 976 *Géographie Alp.*, <https://doi.org/10.4000/rga.11110>, 2023.
- 977 Niggli, L., Allen, S., Frey, H., Huggel, C., Petrakov, D., Raimbekova, Z., Reynolds, J., and Wang, W.:
 978 GLOF Risk Management Experiences and Options: A Global Overview, in: *Oxford Research*
 979 *Encyclopedia of Natural Hazard Science*, Oxford University Press,
 980 <https://doi.org/10.1093/acrefore/9780199389407.013.540>, 2024.
- 981 Nigrelli, G., Chiarle, M., Nuzzi, A., Perotti, L., Torta, G., and Giardino, M.: A web-based, relational
 982 database for studying glaciers in the Italian Alps, *Comput. Geosci.*, 51, 101–107,
 983 <https://doi.org/10.1016/j.cageo.2012.07.027>, 2013.
- 984 Nigrelli, G., Paranunzio, R., Turconi, L., Luino, F., Mortara, G., Guerini, M., Giardino, M., and
 985 Chiarle, M.: First national inventory of high-elevation mass movements in the Italian Alps, *Comput.*
 986 *Geosci.*, 184, 105520, <https://doi.org/10.1016/j.cageo.2024.105520>, 2024.
- 987 Nyberg, L., Cedergren, A., De Herve, M. D. G., Gustavsson, J., Hassel, H., Tehler, H., and Wester, M.:
 988 What do we mean with integrated risk management? – A research profiling comparing DRM and other
 989 fields of risk management, *Int. J. Disaster Risk Reduct.*, 132, 105942,
 990 <https://doi.org/10.1016/j.ijdr.2025.105942>, 2026.

- 991 Occhiena, C. and Pirulli, M.: Analysis of Climatic Influences on Slope Microseismic Activity and
 992 Rockfalls: Case Study of the Matterhorn Peak (Northwestern Alps), *J. Geotech. Geoenvironmental*
 993 *Eng.*, 138, 1012–1021, [https://doi.org/10.1061/\(ASCE\)GT.1943-5606.0000662](https://doi.org/10.1061/(ASCE)GT.1943-5606.0000662), 2012.
- 994 **OECD: Progress towards an all-hazards approach to emergency preparedness and response, Nuclear**
 995 **Energy Agency Organisation for Economic Co-operation and Development, Paris, 2018.**
- 996 **Olivieri, L. and Bettanini, C.: Preliminary observation of Marmolada glacier collapse of July 2022**
 997 **with space-based cameras, *Remote Sens. Lett.*, 14, 21–29,**
 998 **<https://doi.org/10.1080/2150704X.2022.2152754>, 2023.**
- 999 **ONF-RTM: Revue annuelle des phénomènes d’origines glaciaire et périglaciaire de l’année 2024**
 1000 **Massif des Ecrins - Grandes Rousses, 2025.**
- 1001 Peissier, C. and Courtray, V.: Séminaire technique - Synthèse et conclusions : Gestion des risques
 1002 d’origine glaciaire et périglaciaire - PARN, Ministère de l’Ecologie, du Développement Durable, des
 1003 Transports et du Logement - Direction Générale de la Prévention des Risques, Grenoble - World Trade
 1004 Center, 2012.
- 1005 Pigeon, P.: Réflexions sur les notions et les méthodes en géographie des risques dits naturels, *Ann.*
 1006 *Géographie*, 111, 452–470, <https://doi.org/10.3406/geo.2002.21624>, 2002.
- 1007 Prudent-Richard, G., Gillet, M., Vengeon, J.-M., and Descotes-Genon, S.: Changements climatiques
 1008 dans les Alpes : Impacts et risques naturels, *ClimCHAlp - Interreg III B Alpine Space*, 2008.
- 1009 Ravel, L.: Évolution géomorphologique de la haute montagne alpine dans le contexte actuel de
 1010 réchauffement climatique, 113–124, <https://doi.org/10.3406/edyte.2009.1078>, 2009.
- 1011 Ravel, L. and Deline, P.: La face ouest des Drus (massif du Mont-Blanc) : évolution de l’instabilité
 1012 d’une paroi rocheuse dans la haute montagne alpine depuis la fin du petit âge glaciaire,
 1013 *Géomorphologie Relief Process. Environ.*, 14, 261–272,
 1014 <https://doi.org/10.4000/geomorphologie.7444>, 2008.
- 1015 **Ravel, L. and Deline, P.: Rockfall Hazard in the Mont Blanc Massif Increased by the Current**
 1016 **Atmospheric Warming, in: *Engineering Geology for Society and Territory - Volume 1*, edited by:**
 1017 **Lollino, G., Manconi, A., Clague, J., Shan, W., and Chiarle, M., Springer International Publishing,**
 1018 **Cham, 425–428, https://doi.org/10.1007/978-3-319-09300-0_81, 2015.**
- 1019 Ravel, L., Magnin, F., Gallach, X., and Deline, P.: Évolution des parois rocheuses gelées de haute
 1020 montagne sous forçage climatique. *La Météorologie.*, 34–40, [https://doi.org/10.37053/lameteorologie-](https://doi.org/10.37053/lameteorologie-2020-0090)
 1021 [2020-0090](https://doi.org/10.37053/lameteorologie-2020-0090), 2020.
- 1022 Raymond, M., Wegmann, M., and Funk, M.: Inventar gefährlicher Gletscher in der Schweiz,
 1023 Versuchsanstalt für Wasserbau Hydrologie und Glaziologie der Eidgenössischen Technischen
 1024 Hochschule Zürich, 2003.
- 1025 Reghezza, M., Leone, F., and Vinet, F.: La vulnérabilité: un concept problématique., in: *La*
 1026 *vulnérabilité des sociétés et des territoires face aux menaces naturelles - Analyses géographiques*, vol.
 1027 1, Montpellier III, 35–40, 2006.
- 1028 **Ricciardi, T.: Mattmark, 30 August 1965: A catastrophe that changed Switzerland’s perception of**
 1029 **Italian migrants, *Schweiz. Z. Für Gesch.*, 66, 401, 2016.**
- 1030 Richard, D.: Caractérisation et gestion des risques d’origine glaciaire, CEMAGREF - Unité de
 1031 recherche ETNA, Grenoble, 2005.

- 1032 Rickenmann, D. and Zimmermann, M.: The 1987 debris flows in Switzerland: documentation and
1033 analysis, *Geomorphology*, 8, 175–189, [https://doi.org/10.1016/0169-555X\(93\)90036-2](https://doi.org/10.1016/0169-555X(93)90036-2), 1993.
- 1034 Rothenbühler, C.: GISALP - räumlich-zeitliche Modellierung der klimasensitiven
1035 Hochgebirgslandschaft des Oberengadins, 3, 2006.
- 1036 Röthlisberger, H.: Eislawinen und Ausbrüche von Gletscherseen, *Gletsch. Klima - Glaciers Clim.*
1037 *Jahrb. Schweiz. Naturforschenden Ges. Wiss. Teil*, 170–212, 1981b.
- 1038 RTM-ONF: Révision partielle du Plan de Prévention des Risques Naturels Prévisibles, ONF-RTM,
1039 Chamonix-Mont-Blanc, 2000.
- 1040 Schindelegger, A.: Natural hazard risk governance - Report on the state of the Alps, Alpine
1041 Convention, 2019.
- 1042 Schindelegger, A. and Kanonier, A.: Natural Hazard Risk Governance: Status Quo in the EUSALP
1043 Region, EUSALP Action Group 8, Wien, 2019.
- 1044 Schneiderbauer, S. and Ehrlich, D.: Risk, hazard and people’s vulnerability to natural hazards - a
1045 review of definitions, concepts and data, European Commission, 2004.
- 1046 Gletschersee Grindelwald : Felssturz Schlossplatte: <http://www.gletschersee.ch/index.cfm/treeID/21>,
1047 last access: 12 February 2025.
- 1048 Sirop, M., Gagliardini, O., Serbource, P., and Bouvier, P.: La prévention des risques du glacier de Tête
1049 Rousse : une action pluri-acteurs, 27–29, 2022.
- 1050 Stoffel, M., Tiranti, D., and Huggel, C.: Climate change impacts on mass movements — Case studies
1051 from the European Alps, *Sci. Total Environ.*, 493, 1255–1266,
1052 <https://doi.org/10.1016/j.scitotenv.2014.02.102>, 2014.
- 1053 Tacnet, J. M., November, V., Richard, D., and Batton-Hubert, M.: Expertise, décision et incertitude :
1054 jusqu’où une approche interdisciplinaire est-elle possible dans le cadre de la gestion intégrée des
1055 risques naturels en montagne ?, in: *Conférence Outils Pour Décider Ensemble 2010 (OPDE)*, 18 p.,
1056 2010.
- 1057 Tagarev, T., Papadopoulos, G. A., Hagenlocher, M., Sliuzas, R., Ischiwatari, M., and Gallego, E.:
1058 Integrating the risk management cycle, in: *Science for disaster risk management 2020 :acting today,*
1059 *protecting tomorrow.*, Publications Office, LU, <https://doi.org/10.2760/571085>, 2021.
- 1060 Troilo, F., Lodigiani, M., Nicora, M., Perret, P., Christille, J.-M., Calabrese, M., Salvemini, C. B., and
1061 Sartor, S.: Monitoring Cryospheric Environment at a Regional Scale: Big Data from Sensor Networks
1062 and Experimental AI Applications in the Framework of the Glarisk-cc FESR Project, 12, 2025.
- 1063 Valette, P. and Carozza, J.-M.: *Géohistoire de l’environnement et des paysages*, CNRS éditions, Paris,
1064 2019.
- 1065 Vannier, C., Lefebvre, J., Longaretti, P.-Y., and Lavorel, S.: Patterns of landscape change in a rapidly
1066 urbanizing mountain region, *Cybergeo*, <https://doi.org/10.4000/cybergeo.27800>, 2016.
- 1067 Vergnano, A., Oggeri, C., and Godio, A.: Geophysical–geotechnical methodology for assessing the
1068 spatial distribution of glacio-lacustrine sediments: The case history of Lake Seracchi, *Earth Surf.*
1069 *Process. Landf.*, 48, 1374–1397, <https://doi.org/10.1002/esp.5555>, 2023.

- 1070 Villeneuve, E., Pérès, F., and Geneste, L.: An approach to improve risk assessment using experience
1071 feedback, in: Proceedings of IFIP Doctoral Spring Workshop "Product and Asset Lifecycle
1072 Management, 1–5, 2010.
- 1073 Vincent, C.: Fluctuations des bilans de masse des glaciers des Alpes françaises depuis le début du 20^e
1074 siècle au regard des variations climatiques, *Houille Blanche*, 88, 20–24,
1075 <https://doi.org/10.1051/lhb/2002100>, 2002.
- 1076 Vincent, C., Garambois, S., Thibert, E., Lefèbre, E., Le Meur, E., and Six, D.: Origin of the outburst
1077 flood from Glacier de Tête Rousse in 1892 (Mont Blanc area, France), *J. Glaciol.*, 56, 688–698,
1078 <https://doi.org/10.3189/002214310793146188>, 2010a.
- 1079 Vincent, C., Auclair, S., and Meur, E. L.: Outburst flood hazard for glacier-dammed Lac de
1080 Rochemelon, France, *J. Glaciol.*, 56, 91–100, <https://doi.org/10.3189/002214310791190857>, 2010b.
- 1081 Vivian, R.: La catastrophe du glacier Allalin, *Rev. Géographie Alp.*, 54, 97–112,
1082 <https://doi.org/10.3406/rga.1966.3248>, 1966.
- 1083 Walter, F., Amann, F., Kos, A., Kenner, R., Phillips, M., De Preux, A., Huss, M., Tognacca, C.,
1084 Clinton, J., Diehl, T., and Bonanomi, Y.: Direct observations of a three million cubic meter rock-slope
1085 collapse with almost immediate initiation of ensuing debris flows, *Geomorphology*, 351, 106933,
1086 <https://doi.org/10.1016/j.geomorph.2019.106933>, 2019.
- 1087 Weber, S., Beutel, J., Dietze, M., Bast, A., Kenner, R., Phillips, M., Leinauer, J., Mühlbauer, S.,
1088 Pfluger, F., and Krautblatter, M.: Progressive destabilization of a freestanding rock pillar in permafrost
1089 on the Matterhorn (Swiss Alps): Hydro-mechanical modeling and analysis, *Earth Surf. Dyn.*, 13,
1090 1157–1179, <https://doi.org/10.5194/esurf-13-1157-2025>, 2025.
- 1091 Wiegandt, E. and Lugon, R.: 3 Challenges of Living with Glaciers in the Swiss Alps, Past and Present,
1092 in: Darkening Peaks, edited by: Orlove, B., Wiegandt, E., and Luckman, B. H., University of
1093 California Press, 33–48, <https://doi.org/10.1525/9780520934245-004>, 2008.
- 1094 World Glacier Monitoring Service (WGMS): Fluctuations of glaciers 1985-1990, IAHS (ICSU) -
1095 UNEP - UNESCO, 1993.
- 1096