

## Managing glacial and periglacial hazards in the Alps: a geohistorical approach

Juliette Bazin\*<sup>1,2</sup>, Ludovic Ravel<sup>1,3</sup>, Sandrine Caroly<sup>2</sup>

<sup>1</sup>EDYTEM, Savoie Mont-Blanc University, CNRS, 5 Boulevard de la Mer Caspienne 73370 Le Bourget-du-Lac, France

<sup>2</sup>PACTE, Grenoble Alpes University, INP, 46 Avenue Félix Viallet 38000 Grenoble, France

<sup>3</sup>Department of Geosciences, University of Oslo, 0371 Oslo, Norway

\*Corresponding author: J. Bazin (juliette.bazin@univ-smb.fr)

Financial support : This research was funded by the EU ALCOTRA 2021-2027 Interreg VI-A *PrévRisk-CC* project.

### Key words

Glacial and periglacial hazards, Risk management, Risk perception, Geo-history, European Alps

**Abstract.** Glacial and periglacial hazards such as ice or rock avalanches, glacial lake outburst floods, and debris flows have caused severe damage in populated Alpine mountain regions throughout history. The objective of this research is to investigate how risk management methods have evolved over time and across different countries in the European Alps. Through a geo-historical study, we compile an inventory of events that have triggered risk management actions over the past centuries. This management is analysed using eight guiding pillars: hazard understanding, preventive information, land-use planning, monitoring and surveillance, hazard-vulnerability-exposure mitigation, crisis management preparedness, response, and resilience.

Certain events, such as the discharge of the water pocket of the Tête Rousse glacier (FR) in 1892 or the collapse of the Allalin glacier (CH) in 1965, represent key turning points that mark a shift or renewal in how risks are perceived and addressed. Today, glacial and periglacial risk management benefits from a better understanding of both hazards and vulnerabilities. Our geo-historical analysis highlights that the inclusion of the local populations in crisis management has become an increasingly significant factor in decision-making processes. However, current risk management practices remain limited and would benefit from more participatory approaches, which in turn partly depend on the perception of hazards and their integration into management practices.

### 1 Introduction

Rock and ice mass movements in high mountain areas pose significant risks to populations and infrastructure. Numerous past events show how glacial and periglacial processes can deeply affect Alpine valleys through large-scale and/or cascading phenomena (*e.g.*, Haeberli et al., 2016; Magnin et al., 2023). Their frequency and volume appear to be increasing (Ravel and Deline, 2008, 2015; Jacquemart et al., 2024) as illustrated by recent disasters in Bondo (Graubünden, CH) in August 2017 (Walter et al., 2020), La Béarde (Isère, FR) in June 2024 (Mainieri et al., 2025), and Blatten (Valais, CH) in May 2025 (Büntgen et al., 2025).

This study focuses on major hazards affecting valley floors: ice, rock, and mixed avalanches, glacial lake outburst floods (GLOFs) and periglacial debris flows. These hazards are defined as natural phenomena impacting Alpine territories (Moles, 1972; Bisquert et al., 2025). A process becomes a *risk*

52 when it threatens lives, economic assets, or the environment. The notion of an *event* is useful for  
53 historians as it marks “a cut, a discontinuity”, something “interesting”, sufficiently “important” or “new”  
54 to be “told or enacted” (Dosse, 2010; Giacona et al., 2017). *Issues* are thus the exposed elements with  
55 varying vulnerabilities to each hazard (Defossez et al., 2018).

56 Permafrost warming and/or glacial debuitressing are contributing factors to rockfalls and rock  
57 avalanches (Fischer et al., 2006; Gruber and Haerberli, 2007; Huggel et al., 2010; Fischer et al., 2012).  
58 They may directly impact valley areas or provide material that triggers debris flows (Walter et al., 2020).  
59 An ice avalanche occurs when a glacier section detaches and moves quickly downwards (Alean, 1985;  
60 Richard, 2005; Faillietaz et al., 2015), destroying infrastructure, forests and lives, as illustrated by the  
61 collapse of the Allalin glacier (CH) onto worker accommodations during the construction of the  
62 Mattmark dam in 1965 (Dalban Canassy et al., 2011). Indirectly, they can trigger snow avalanches  
63 (Margreth and Funk, 1999) or debris flows when ice temporarily dams a stream (Richard, 2005).  
64 GLOFs arise when a glacial lake or another water body suddenly drains, releasing large volumes of  
65 water that can evolve into debris flows if sufficient amounts of sediment are available (Richard, 2005;  
66 Huss et al., 2007; Carrivick and Tweed, 2016; Ancey et al., 2019). Their impacts can devastate entire  
67 valleys, as seen in the Bagnes valley below the Giétro glacier (CH) in 1595 and 1818 (Vincent et al.,  
68 2010; Ancey et al., 2019).

69 Integrated management of these risks (Prudent-Richard et al., 2008; Stoffel et al., 2014; Jacquemart et  
70 al., 2024) is essential for prevention because it allows for a holistic, coordinated approach that  
71 anticipates, mitigates, and adapts to complex, interconnected climate and emerging risks before they  
72 escalate into crises. We base our approach on the key components of risk management: hazard, exposure,  
73 and vulnerability (Niggli et al., 2024).

74 The database developed in this study documents glacial and periglacial events across the European Alps  
75 over the past two centuries that required risk management. Prevention measures from a political  
76 perspective are not uniform across Alpine countries. We have chosen to primarily present examples from  
77 France, Switzerland and Italy. Although Austria is also cited according to the sources we collected. The  
78 authors acknowledge the limitations of this study, particularly the challenges associated with obtaining  
79 comprehensive data; nevertheless, it is conceived as an exploratory approach. As such, it may serve as  
80 an initial step toward the development of a broader collaborative framework. The purpose is not only to  
81 provide a detailed account of the risk management measures implemented, but also to assess the  
82 frequency and intensity of such events and to trace historical management trends. An historical approach  
83 enables the analysis of change processes in risk management (Giacona et al., 2017, 2019), beyond mere  
84 event description (Girard and Rivière-Honegger, 2015).

85 We thus seek to understand how Alpine glacial and periglacial hazards have shaped scientific knowledge  
86 and risk management practices.

87 The paper first outlines the theoretical and historical framework of risk management, followed by the  
88 methods used to construct the database and apply the eight pillars of risk management which we have  
89 identified. The results describe the spatial and temporal distribution of events and corresponding  
90 management actions. Finally, we discuss the limitations of the data and the challenges of future  
91 prevention under a changing climate.

## 94 2 State of the art and theoretical positioning

95  
96 The concepts used here require clarification to frame our approach. A *risk* refers to the probability of  
97 damage resulting from interactions between a physical process (the *hazard*), *exposure*, and *vulnerability*  
98 (Pigeon, 2002). The hazards we study occur at high altitudes (> 2,000-2,500 m a.s.l.) and may reach  
99 valley floors, threatening infrastructure, livelihoods, ecosystems, and populations (IPCC, 2023).  
100 Mountain valleys are thus considered vulnerable due to their exposure to such hazards, a condition  
101 reinforced by the concentration of people and infrastructure, which increases susceptibility to cascading  
102 effects (IPCC, 2023; Niggli et al., 2024). Vulnerability also depends on *resilience* and *sensitivity*, i.e. the

103 system's ability to absorb shocks and the degree of damage it can sustain (Schneiderbauer and Ehrlich,  
104 2004; Defossez et al., 2018). *Exposure* reflects the spatial and social proximity between hazard sources  
105 and affected communities (IPCC, 2022).

106 The continued expansion of urban areas in high Alpine valleys, driven by tourism and socioeconomic  
107 development (Hock and Rasul, 2022), combined with climate-induced changes in the cryosphere,  
108 increases potential losses (Huggel et al., 2015) and underscores the need for effective *risk management*  
109 (Allen et al., 2017). The latter encompasses strategies to anticipate, reduce, and respond to hazards  
110 (Leone et al., 2021) and aims to reduce uncertainty through scientific advances (Margreth et al., 2011,  
111 2017). While comparable to other Alpine hazards, glacial and periglacial processes evolve rapidly under  
112 climate change and can reach extreme magnitudes (Haeberli et al., 1999; Richard, 2005). Anticipation  
113 remains difficult because these hazards co-evolve with a rapidly changing cryosphere: glacier retreat,  
114 de-icing of rock faces, permafrost warming and associated slope movements, formation or  
115 disappearance of glacial lakes, and changes in glacier thermal regimes (Mainieri et al., 2025). High  
116 erosion rates in recently deglaciated or warming permafrost areas further increase debris-flow  
117 susceptibility (Huggel et al., 2015).

118 As historical experience shows, approaches based solely on hazard characterisation (Richard, 2005) are  
119 necessary but not sufficient (Allen et al., 2017). Analysing past events improves understanding of  
120 processes and supports better risk mitigation (Carrivick and Tweed, 2016). Rather than compiling  
121 exhaustive inventories, this study focuses is on lessons learned from major events, particularly how  
122 societies have managed risks over time.

123 Building a historical database requires a diachronic approach, inspired by diachronic monographs  
124 (Girard and Rivière-Honegger, 2015). This perspective (*e.g.* Desailly, 1990; Mendez, 2010) uses time  
125 to explain current territorial and risk dynamics. Such an approach allows the identification of  
126 geohistorical trajectories and rupture points in hazard evolution (Valette and Carozza, 2019; Hugerot  
127 et al., 2021). We adopt this framework to analyse long-term risk management, local practices, and the  
128 evolution of risk culture (Dollfus and D'Ercole, 1996). Studies such as Favier (2006) and Fournier  
129 (2010) on avalanches and floods in the Grenoble basin (FR), or Niggli et al. (2024) on GLOFs, illustrate  
130 how mountain societies have experienced and managed hazards.

131 Adopting a diachronic perspective also requires careful source selection (biases are discussed in § 5.2).  
132 Archival records complement narrative sources (Fournier, 2010). Since the 19<sup>th</sup> century, historical  
133 reconstructions have supported risk anticipation (Dourlens, 2004). However, exhaustive chronologies  
134 remain unattainable, as many small or frequent events leave no archival trace (Giacona et al., 2019).

135 To address this limitation, historical approaches prioritise the analysis of processes and mechanisms  
136 over completeness. Our study thus provides a synthetic view of Alpine risk management. Similar  
137 approaches, such as the temporal analysis of rockfalls in the Mont-Blanc massif (Ravel and Deline,  
138 2008; Ravel et al., 2020), reveal long-term dynamics.

139 Long-term monitoring initiatives also demonstrate the sensitivity of mountain environments to climate  
140 change, such as the EU-PACE project (Permafrost and Climate in Europe), which highlights the role of  
141 permafrost in slope stability and geohazards (Harris et al., 2001; Etzelmüller et al., 2020).

142

143

### 144 **3 Construction of the database**

145

#### 146 3.1 Inventory of major glacial and periglacial events in the Alps

147

##### 148 3.1.1 Criteria used to identify events to be included in the database

149

150 Building a comprehensive long-term database is challenging as data availability before the 20<sup>th</sup> century  
151 largely depends on societal vulnerability and record-keeping (Giacona, 2019).

152 We adopted a qualitative approach, retaining events that significantly marked local history or the  
153 environment, living traces in archives or collective memory. Our criteria are grouped into four  
154 categories.

#### 155 **Extreme events**

156 We focused on rare, high-magnitude events exceeding several thousand cubic metres or causing serious  
157 damage (Bourrelier and Dunglas, 2009; Field et al., 2012). Quantitatively, these correspond to the upper  
158 end of the frequency–intensity spectrum.

#### 159 **Events with feedback**

160 We include events that led to in-depth post-event analyses aimed at improving process understanding,  
161 beyond simple observation (e.g. the Giétro glacier collapse in 1818; Wiegandt and Lugon, 2008).

#### 162 **Repeated events**

163 We consider events repeatedly occurring within the same watershed, focusing on major one. For  
164 example, the Bockkarkees hanging glacier (AT) has produced >70 ice avalanches since 1933, including  
165 events reaching  $5 \times 10^6$  m<sup>3</sup> in 1945 and  $2\text{--}3 \times 10^6$  m<sup>3</sup> in 1964 and 1975 (Gridabase; Kellerer-Pirklbauer  
166 et al., 2012).

#### 167 **Change in risk perception**

168 Events that significantly influenced risk perception are included, even if they do not meet the previous  
169 criteria. Risk culture varies across Alpine countries and is shaped by historical events. Event memory  
170 differs by context, making risk perception a social and political issue. **The Mattmark disaster marked a  
171 major policy shift, moving away from the legally criticized notion that natural hazards are inherently  
172 unpredictable toward a more predictive approach to hazard analysis, as reflected in the Federal  
173 Government's establishment of a glacier hazard working group** (Capozzi, 2011; Joris, 2025). Similarly,  
174 the Tête Rousse disaster (FR) established in France the idea of glaciers as internal, hidden, yet  
175 predictable hazards. The Marmolada glacier (IT) collapse in 2022 (Francesse et al., 2024), although  
176 limited to high-altitude impacts, caused 11 fatalities and strongly influenced public opinion, highlighting  
177 the need to better identify glacial hazards (Chiarle et al., 2022). It also led to the creation of a national  
178 working group and the publication of guidelines (Dipartimento della Protezione Civile, 2025).

179 We initially intended to start our inventory in 1985-1990, marking the first visible signs of the climate  
180 crisis in the Alps (Ravel, 2009; Ravel et al., 2020), up to 2025. However, earlier key events such as  
181 the 1892 Tête Rousse outburst (Mougin et Bernard, 1922) were too significant to omit. We therefore  
182 extended the analysis back to the early 19<sup>th</sup> century, where reliable sources exist.

183 For each event, we collected the following data: location (country, region, mountain range, orientation,  
184 glacier or summit name), rupture date(s), causes, mobilised volumes, runout distance, cascading effects,  
185 damage and casualties, monitoring methods, crisis management, and sources.

186 This method was applied to three hazard types: ice avalanches, torrential hazards (including GLOFs and  
187 debris flows), and rockfalls/rock-ice avalanches.

188 For cascading processes, we record the most impactful downstream process. For example, if a serac fall  
189 triggers lake overflow, the torrential process is retained. If permafrost degradation destabilises a moraine  
190 dam, we document the resulting breach and downstream propagation.

#### 191 **3.1.2 Data sources**

192 We used the *Gridabase* inventory produced as part of the European research project (« Gridabase -  
193 Glaciorisk », s. d., 2001-2003). This database compiles 501 documented events across Austria, France,  
194 Italy, and Switzerland, including floods, GLOFs, and ice avalanches (Richard, 2005). The aim of  
195 *Gridabase* was to preserve as comprehensively as possible the memory and trace of glacial events

**Commenté [JB1]:** W.Haeberli : The shift relates to the change from the heavily criticised “dogma of unpredictability” in the legal case towards predictive hazard analysis as initiated by the working group on glacier hazards established by the Federal Government.

203 (Peissier and Courtray, 2012), including those of modest importance (excluded here according to our  
204 criteria). However, investigations initiated by the Swiss Parliament highlighted that changing  
205 environmental conditions exceeded historical empirical references, requiring broader system  
206 assessments (Rickenmann and Zimmermann, 1993). *Gridabase* therefore presents a retrospective and  
207 partial perspective, which we accounted for in our analysis. In addition, for Italian Alps, we used the  
208 *Geoclimalp* cartographic inventory (Nigrelli et al., 2024; geoclimalp, 2026), which contains > 700 mass  
209 movements > 1500 m a.s.l. between 2000 and 2022.

210 Other sources include local databases such as the *RTM Database* or the regional *Catasto Dissesti*. The  
211 former is a French database developed by the *Restauration des Terrains de Montagne* (RTM) public  
212 service of the *Office National des Forêts*. Since the late 19<sup>th</sup> century, it has recorded a wide range of  
213 geomorphic processes, including avalanches, gully erosion, subsidence, landslides, rockfalls and debris  
214 flows (Bisquert et al., 2025). The *Catasto Dissesti* is a regional mapped inventory of hydrogeological  
215 and gravitational phenomena in Aosta Valley, enriched since the 1990s and regularly updated for risk  
216 prevention (Giardino et al., 2010). We retained from both databases glacial and periglacial events,  
217 including rock glacier ruptures, permafrost-related rockfalls and debris flows generated by cascading  
218 processes.

219 The RTM has also produced numerous technical reports on glacial and periglacial events. These  
220 documents, sometimes, unpublished, describes causes, trajectories, volumes, damage, and process  
221 histories within catchment. They also document risk management actions, from modelling and early  
222 warning to operational interventions (e.g. Patinoire lake in 1964; Étançons torrent in 2024; (Cathala et  
223 al., 2021; Demolis et al., 2021; Mainieri et al., 2025) and crisis management through operational  
224 measures (e.g. artificial draining of Rochemelon lake, FR, in 2005; Vincent et al., 2010; Cathala et al.,  
225 2021).

226 Many events documented by Swiss authorities were also included, such as the flood protection measures  
227 implemented at the Gruben glacier (Haeberli et al., 2001).

228 Scientific papers constitute a major part of our dataset, providing detailed analyses of hazard  
229 mechanisms and management strategies. Individual events are often studied by multiple teams using  
230 complementary approaches. The Whymper hanging glacier and the Planpincieux glacier (IT) illustrate  
231 this: studies address monitoring systems, rupture scenarios, hazard assessment, and evacuation strategies  
232 (Margreth and Funk, 1999; Margreth et al., 2011; Schindelegger, 2019; Dematteis et al., 2021; Troilo,  
233 2025).

234 The inventories of Chiarle et al. (2007, 2022), Jacquemart et al. (2024), and Niggli et al. (2024) were  
235 used to cross-check and validate our dataset.

236 Ministerial reports (e.g. Lacroix et al., 2022), the study of Magnin et al. (2023), and output from the  
237 *Pôle Alpin des Risques Naturels* (PARN), provide insights into how Alpine countries manage glacial and  
238 periglacial risks. They document research projects, monitoring strategies, prevention measures, and  
239 crisis management practices (e.g. *Glaciorisk*, *PERMAdataROC*, *PermaNET*, *GlariskAlp*, *GLAMOS*,  
240 *PERMOS*, *Prevrisk-CC*, *PermaRisk*, *SAMCO*, etc.). The *PAPROG* initiative fosters multidisciplinary  
241 collaboration between researchers and practitioners to improve knowledge and management of glacial  
242 risks. Similarly, *GEORESEARCH*, coordinates projects on monitoring, cryospheric dynamics, and  
243 quantitative risk analysis in the Alps (*FROST.INI*, *Futurelakes*, *GlacierRocks*, *AlpSenseRely*, etc.).

244 The *World Glacier Monitoring Service* (WGMS) provides global data on glacier fluctuations (1959-  
245 2010), including notable hazard events.

246 Finally, the local press provides insights into the societal impact of mass movements and associated  
247 management responses. However, this source requires careful critical interpretation, as it may reflect  
248 biases and lack technical accuracy (Joffe and Orfali, 2005). Despite these limitations, it remains valuable  
249 for analysing societal risk perception.

250

## 251 3.2 Types of glacial and periglacial risk management

252

### 253 3.2.1 Definition of the eight pillars of risk management

Commenté [JB2]: W.Haeberli : If GLAMOS is mentioned, PERMOS should also be mentioned.

254  
255 Based on Alpine risk management systems (from local to national levels) and scientific literature,  
256 Kienholz et al., (2004) and Link and Stötter (2015) developed a classification of risk prevention and  
257 management methods. More recently, the Risk Management Cycle (RMC) – or Disaster Risk  
258 Management (DRM) – has been conceptualised. It encompasses all policies, strategies, and actions  
259 aimed at preventing, reducing, and managing disaster risks, as well as preparing for, responding to, and  
260 recovering from their impacts (Tagarev et al., 2021). In Switzerland and Nordic countries, since the  
261 1980s, natural hazard management has followed the principles of “Risk Minimisation” and “Integral  
262 Risk Management” (Nyberg et al., 2026), which aim to reduce risks to acceptable levels through  
263 assessment, prevention, mitigation, and preparedness, rather than eliminate them. Post-crisis phases  
264 focus on learning to improve future measures.  
265 In France, similar principles are embedded in public policy and summarized by the PARN for glacial  
266 and periglacial risk prevention (Boudières and Peisser, 2013). In Italy, the *Dipartimento della Protezione  
267 Civile* (2025) has produced a methodological framework defining objectives for understanding glacial  
268 and periglacial phenomena. It outlines key pillars such as monitoring, public information, and crisis  
269 resilience.  
270 All these principles were used to classify the management strategies identified in our database.

271  
272 **Pillar 1 : Hazards, phenomena, and risks understanding**

273 This pillar improves knowledge through past-events analysis, archives, monitoring, modelling, and  
274 experience feedback (Villeneuve et al., 2010; Faillettaz et al., 2015). It includes inventories and  
275 vulnerability assessments (e.g. GlaRiskAlp project; Lucchesi et al., 2014; Nigrelli et al., 2013). Within  
276 DRM, this corresponds to risk identification and preparedness (Tagarev et al., 2021).

277  
278 **Pillar 2 : Preventive information**

279 This pillar promotes risk culture through public information and education (Beccera and Peltier, 2011),  
280 enabling citizens to actively contribute to preparedness and crisis response (OECD, 2018).

281  
282 **Pillar 3 : Integration into land-use planning**

283 Risk is integrated into planning tools such as hazard maps and preventive plans to reduce exposure and  
284 vulnerability (Schneiderbauer and Ehrlich, 2004). These tools combine modelling and spatial analysis  
285 to support decision-making (Allen et al., 2022).

286  
287 **Pillar 4 : Monitoring and surveillance**

288 Monitoring at-risk sites (cf.: Pillar 1) enables early detection and warning (Giordan et al., 2020; Cathala  
289 et al., 2024), supporting emergency preparedness. For example, Gornersee lake (CH) is monitored using  
290 cameras, depth measurements, and seismic sensors to anticipate GLOF hazards (WGMS, 1993; Huss et  
291 al., 2007). In Italy, glaciers such as Freney, Brenva, Grandes Jorasses, and Planpincieux are monitored  
292 using time-lapse imagery to assess flow velocity (Dematteis et al., 2024). Satellite observations  
293 complement field monitoring, especially for remote or large-scale events, though in the Alps they are  
294 often used post-event to analyse processes (e.g. Marmolada glacier collapse in 2022; (Bondesan and  
295 Francese, 2023; Olivieri and Bettanini, 2023; Francese et al., 2024) or to monitor supraglacial lakes  
296 hazards (e.g. Lys glacier or the Forni glacier, IT; (Davide et al., 2021; Fugazza et al., 2018). However,  
297 field-based observation remains central in the Alpine context. Field observations and human-based  
298 warning systems remain central, including observer networks such as “*Regards d'Altitude*” (ONF-RTM,  
299 2025).

300  
301 **Pillar 5 : Reducing hazard, exposure, and vulnerability**

302 This central pillar in risk / disaster risk management encompasses measures aimed at reducing three  
303 interconnected components of risk: hazard, exposure, and vulnerability. **These three components are  
304 addressed together here because, in practice, risk managers and decision-makers rarely operate by**

**Commenté [JB3]:** W. Haeberli : A newer and more complete analysis is: Francese, R.G., Valentino, R., Haeberli, W., Bondesan, A., Giorgi, M., Picotti, S., Pettenati, F., Sandron, D., Ramponi, G. and Valt, M. (2025): Failure of Marmolada Glacier (Dolomites, Italy) in 2022: data-based back analysis of possible collapse mechanisms. *Natural Hazards and Earth System Sciences* 25,3027-3053. <https://doi.org/10.5194/nhess-25-3027-2025>

**Commenté [JB4]:** C. Huggel : Pillar 5: this pillar is certainly the core of much of what we know as risk / disaster risk management. In principle it could make sense to separate hazard reduction from exposure and vulnerability reduction measures. This is typically done. Maybe add a sentence why you have taken all together here. The first sentence is a bit misleading, I think (limiting damage if hazards cannot be avoided). I think there is hardly any scenario in mountain environments where hazards can be avoided, it is always about limiting the frequency or magnitude of hazards (at the occurrence site or on their way downstream). Often, a distinction is made between structural and non-structural measures, this could also be mentioned. Overall I suggest that this pillar is explained in some more detail and better specification, especially because it is so central.

305 formally distinguishing between them. Mitigation strategies typically combine technical interventions  
306 and community capacity-building within integrated action plans. Artificially separating them would not  
307 reflect the reality of operational risk management. This is also consistent with the approach taken in this  
308 study, where the analysed policies and measures were designed and implemented in an integrated  
309 manner, making it difficult to disentangle the contribution of each component independently.  
310 In mountain contexts specifically, hazards can almost never be fully avoided; the focus therefore lies on  
311 limiting their frequency, magnitude, or spatial extent (at the source or along their path). Mitigation  
312 measures are commonly divided into two broad categories. Structural (or engineering) measures involve  
313 physical interventions. Non-structural measures include early warning systems, evacuation protocols,  
314 risk communication for examples. Together, these measures aim to reduce potential damage by acting  
315 on the hazard itself, on the elements exposed to it, or on the ability of communities and systems to cope  
316 with and recover from adverse events (Vincent et al., 2010b; Tagarev et al., 2021).  
317 For GLOFs, measures include drainage, water-level regulation, and structural stabilisation (Niggli et al.,  
318 2024). Exposure is reduced through channel management and protective structures, while vulnerability  
319 is lowered by improving preparedness and response capacities.

#### 320 ***Pillar 6 : Crisis management preparedness***

321 This pillar corresponds to measures taken to improve response capacity (Tagarev et al., 2021). It includes  
322 planning, training, and coordination measures implemented before crises, such as contingency plans,  
323 simulations, and risk integration into training (Einhorn and Peisser, 2011; Link and Stötter, 2015).

#### 324 ***Pillar 7 : Response***

325 Response refers to immediate actions during or after an event to manage impacts, save lives, and ensure  
326 safety (OECD, 2018). It includes emergency assistance and implementation of crisis plans (Tagarev et  
327 al., 2021).

#### 328 ***Pillar 8 : Resilience***

329 Resilience is defined as a system's ability to absorb shocks and recover, involving resistance, adaptation,  
330 and reorganisation (Reghezza et al., 2006; Dauphiné and Provitolo, 2013). The objective is not full  
331 recovery but achieving a new acceptable equilibrium (Einhorn, 2017). This phase also provides  
332 opportunities to improve future risk reduction measures (Tagarev et al., 2021).

### 333 3.2.2 Distribution of events according to the eight pillars typology

334 Once the typology has been established, each event was assigned to the relevant management pillars  
335 mobilised in response. Events were then grouped by decade (or by five-year intervals for rock hazards)  
336 to improve temporal readability. This approach highlights changes in management practices over time  
337 while limiting visual discontinuities associated with rare events.

## 338 **4 Results**

339 This study identifies 200 glacial and periglacial events across 109 Alpine catchment areas in France,  
340 Switzerland, Italy, and Austria. These events were selected based on their magnitude, impacts, or  
341 influence on the evolution of glacial and periglacial risk management practices.


### 342 4.1 Spatial and temporal distribution of the events

#### 343 4.1.1 Distribution of the events across the Alps

355 The spatial distribution of events (Fig. 1) shows a marked concentration in certain areas, highlighting  
 356 particularly exposed or dynamically active zones. The Mont-Blanc massif (mainly its French and Italian  
 357 sides) clearly emerges as the most densely affected area, across all hazard types. It is followed by the  
 358 Valais Alps (CH). GLOFs are also more frequent in the western Alps.

359 The western Alps high elevations, steep slopes, and extensive glacier cover, making them particularly  
 360 prone to mass movements. The Mont-Blanc massif and the Valais Alps are among the most monitored  
 361 and instrumented regions, which may introduce a ‘visibility bias’ (cf. 5.2). High population density and  
 362 tourism further enhance event detection and documentation, potentially amplifying their perceived  
 363 significance.

364 Figure 1 presents major events that occurred between the 17<sup>th</sup> century and today, revealing the long-term  
 365 dynamics of these hazards. They are not isolated events but part of long-term, sometimes recurring  
 366 trajectories. This representation shows that multiple hazard types coexist within the same Alpine  
 367 massifs, increasing the complexity of risk management.



**Figure 1.** Distribution of major glacial and periglacial events inventoried in the Alps. © European Space Agency (2024). Copernicus Global Digital Elevation Model. Distributed by OpenTopography


368 4.2 Distribution of management pillars over time

369

370 4.2.1 Phenomena intensification

371

372 Keeping in mind potential source and event biases (cf. § 5), we nevertheless observe an intensification  
 373 of **hazardous glacial and periglacial events** since the 1980s across all Alpine countries (Fig. 2). This  
 374 trend is based solely on events that required risk management and contributed to hazard understanding.



375 **Figure 2.** Cumulative frequencies of inventoried glacial and periglacial events requiring risk management in the Alps.

#### 377 4.2.2. Evolution over time of the pillars used

378 This section examines how risk management and prevention methods have evolved over time through  
 379 significant glacial or periglacial events. Events are analysed according to the management pillars  
 380 mobilised at the time.

381 For the three types of events, Pillars 2 (Preventive information) and 3 (Land use planning) appear only  
 382 from the 1990s onward, while Pillar 6 (Crisis preparedness) has been implemented more recently. Pillar  
 383 1 (Hazard understanding) remains the most consistently used. It also continues to be applied  
 384 retrospectively, as past events are reanalysed to improve risk knowledge.

385 The evolution over time of risk management associated with major *ice avalanche* events in the Alps is  
 386 presented in Figure 3. The graph shows two main phases: a long period (1597-1950) dominated almost  
 387 exclusively by Pillar 1 (Understanding hazard), followed by a gradual and accelerating diversification  
 388 of management approaches from the 1950s and 1960s onwards.

389 The Altels ice avalanche (CH) in 1895 marked a turning point in the recognition of the need to  
 390 understand high-altitude hazards. This avalanche was of exceptional size, with  $4 \times 10^6$  m<sup>3</sup> of ice breaking  
 391 away, killing six people and burying 170 cattle over an area of 1 km<sup>2</sup> (Forel, 1895; Du Pasquier, 1896).  
 392 For a long time, this event served as a benchmark for ice avalanches. "To what extent can we predict  
 393 such disasters, and to what degree will we be able to prevent them from happening again? Many Alpine  
 394 glaciers give rise to avalanches or ruinous debacles. (...) The same glaciers periodically give rise to the  
 395 same phenomena, so it is above all empirical knowledge of these phenomena that we need to acquire.  
 396 (...) An accurate knowledge of the dangerous points and the phenomena that occur there is absolutely  
 397 necessary. The era of independent personal research is over; it has produced all it could in this field (...).  
 398 What is needed now is more coordination, more method, more consistency in research" (Du Pasquier,  
 399 1896).


400 The predominance of Pillar 1 over several centuries should not be interpreted as a lack of institutional  
 401 response to glacial and periglacial hazards. It rather reflects the fact that, within the epistemic and  
 402 institutional context of the time, the production of knowledge and the documentation of phenomena  
 403 constituted the primary and legitimate form of risk management. At that time, management was largely  
 404 confined to the academic sphere (naturalists, cartographers, scientific academies), long before it became  
 405 integrated into engineering and spatial planning. This observation suggests that contemporary categories  
 406 of risk management should not be retrospectively applied to these periods. If preventive information  
 407 (Pillar 2) appears late, this indicates that decision-makers only gradually began to communicate  
 408 simplified risk information to exposed populations. This development is closely linked to the emergence  
 409 of crisis preparedness (Pillar 6). It also raises the question of the role assigned to local communities in

**Commenté [JB5]:** W.Haeberli : Better "hazardous glacial and periglacial events" (the term "hazard" is a mental construct relating to humans and to the future; this term should not be used in the sense of a (physical) "process" or a (past) "event". The formulation in the caption of Figure 2 is correct.

411 risk management, which has long remained expert-driven, with limited structured communication with  
 412 residents. The example of the Val Ferret (IT), where the terminal tongue of the Planpincieux glacier has  
 413 been monitored since 2014, illustrates this evolution. Monitoring is coupled with rapid civil protection  
 414 responses and public information measures (Giordan et al., 2020; Dematteis et al., 2024). The aim is to  
 415 continuously track glacier dynamics, and detect instability, and activate procedures such as road closures  
 416 or village evacuation.

417 This study also integrates avalanche hazard maps, such as the *Carte de Localisation des Phénomènes*  
 418 *d'Avalanche* (CLPA; Avalanche Phenomenon Location Maps) in France, *Catasto valanghe* in Aosta  
 419 valley (Italy), Indication of avalanche hazards (ATH) for Switzerland, *Wildbach und Lawinenverbauung*  
 420 (WLV) for Austria, into land-use planning (Pillar 3), even though they primarily address snow  
 421 avalanches. These maps first emerged in the 1970s, but events such as those at the Bourgeat glacier (FR)  
 422 in 1984 and 1993 show that ice avalanches can trigger snow avalanches (RTM-ONF, 2000, unpublished).  
 423 We therefore consider these mapping tools as addressing combined snow-ice hazards.

424



425

426

**Figure 3.** Temporal evolution of ice avalanche risk management in the Alps.

427 The evolution over time of risk management for major rockfall and rock avalanche events in the Alps is  
 428 presented in Figure 4. It is divided into five-year intervals to better highlight recent dynamics, as most  
 429 rockfall events have occurred since 1980.

430 The reduction in vulnerability, exposure, and hazard (Pillar 5) is particularly evident in these events. For  
 431 example, the 1987 Val Pola rockslide (CH;  $30 \times 10^6 \text{ m}^3$ ) involved preventive evacuation, illustrating  
 432 proactive risk management (Chardon, 1990). In 2011, at Piz Cengalo / Bondo ( $1.5 \times 10^6 \text{ m}^3$ ), preventive  
 433 measures included public alerts, trail closures, evacuation of alpine buildings, and installation of  
 434 protective structures (Bohnenblust, 2017).

435 Monitoring unstable rock faces is a key management tool (Pillar 4). Initially sporadic (1940–1944), it  
 436 became more systematic from the mid-1980s. Its increasing prominence in recent years (especially  
 437 2020-2024) reflects growing investment in instrumental monitoring systems, driven by both the  
 438 perceived intensification of hazards and advances in measurement technologies. Since 1986, the eastern  
 439 face of Monte Rosa (IT) has been regularly monitored due to increasing mass movements (Fischer et  
 440 al., 2006). Other emblematic sites have since been equipped, including Aiguille du Midi (FR) since  
 441 2005, the Matterhorn south face (IT) and Les Drus (FR) since 2007.

442 Finally, projects such as the EU AlpinSpace *PermaNET* program and the *PACE* project have contributed  
 443 to structuring knowledge on Alpine permafrost, through harmonised datasets and coordinated  
 444 monitoring across the Alps (Harris et al., 2001; Mair et al., 2011).

445




Figure 4. Temporal evolution of risk management for rockfalls / rock avalanches.

446  
447

448 The evolution over time of risk management for major GLOF events in the Alps is presented in Figure  
449 5. Unlike other hazards, GLOFs are documented as early as the 16<sup>th</sup> century, reflecting their high  
450 visibility and destructing impacts in populated valleys. Their severity ensured systematic recording in  
451 administrative and local archives, especially as their transformation into debris flows amplifies  
452 downstream damage. The Bagnes valley and the Giétro glacier, for example, generated extensive records  
453 from the 16<sup>th</sup> century onwards due to the scale of the destruction (Ancy et al., 2019).

454 The period 1850-1889 is absent from the graph, which is paradoxical from a climatic perspective.  
455 Although post-Little Ice Age warming favoured lake formation and potentially increased GLOF  
456 occurrence, this gap likely reflects missing data rather than reduced activity.

457 Efforts to reduce vulnerability and exposure appear earlier for GLOFs than for other hazards. Efforts to  
458 reduce vulnerability and exposure appear earlier for GLOFs than for other hazards. Early responses  
459 relied on direct engineering interventions (drainage, tunnels, dykes) in densely populated valleys,  
460 preceding formalised risk management frameworks. This reflects a pragmatic, engineering-driven  
461 approach shaped by local socio-economic constraints. Thus, from the late 16<sup>th</sup> century, recurrent  
462 outbursts of the Margherita glacial lake (Rutor glacier, IT) prompted studies commissioned by the Duke  
463 of Savoy (12 events between 1430 and 1680; discharge of  $5 \times 10^6 \text{ m}^3$  in 1751). Technical solutions such  
464 as dam and drainage tunnels were already considered in the 19<sup>th</sup> century (Vergnano et al., 2023).


465 Another notable example is the Giétro glacier event in 1818. A tunnel was constructed to drain the ice-  
466 dammed lake, thereby reducing the magnitude of the floods and preventing major downstream damage  
467 (Wiegandt and Lugon, 2008). **Also, at the foot of the Gruben Glacier, flood prevention work was carried  
468 out at two lakes after the 1970 outburst. For the proglacial lakes, the moraine dams were reinforced with  
469 concrete injections; and the thermokarst lake was drained through an artificial channel in 1995 (Haeberli  
470 et al., 2001).**

471 These recurrent events progressively led to diversified measures, including dykes, dams, storage basins,  
472 subglacial drilling, siphoning, lake drainage, and moraine stabilisation, as well as warning systems. In  
473 parallel, glacier monitoring became increasingly formalised, notably through photogrammetric surveys  
474 of the Vernagtferner (AU), one of the best-documented glaciers due to its frequent GLOFs, the most  
475 notable of which dates back to 1848 (Braun, 1995; *Gridabase - Glaciorisk*, s. d.).

476 However, the integration of populations through preventive information is more recent. For instance,  
477 the 1986 Arsine proglacial lake case (FR) marked one of the first instances where scientific warning led  
478 to coordinated crisis preparedness involving local populations (Lailly and Demolis, 2019).

**Commenté [JB6]:** W. Haeberli : In this context, it would be appropriate to mention the two multi-year, comprehensive hazard/risk-reducing work at Gruben and at Belvedere (references in the reference list)

**Commenté [JB7R6]:** Authors : As there are so many examples, we have chosen not to include them all, but only the most 'telling' ones that cover a wide range of areas.



480

481 **Figure 5.** Temporal evolution of risk management for Glacial Lake Outburst Floods.482 4.2.3 Disruptive events that have improved the glacial and periglacial risk management and  
483 understanding

484

485 The construction of a timeline tracing risk management responses to past events has enabled the  
486 identification of key turning points marking disruptions and/or the emergence of new approaches.  
487 Selected examples correspond to first documented implementations of new management practices  
488 (prediction, monitoring, mitigation, and coordinated crisis response).

489

490 **Scale change**

491 Risk management has shifted from local (village-scale) responses to national frameworks. The Giétro  
492 glacier disaster in 1818 illustrates this transition (Ancey et al., 2019). At that time, glaciology was still  
493 emerging, yet the event demonstrated that improved process understanding could support more effective  
494 interventions. Scientific advances progressively supported the development of preventive measures and  
495 favoured the emergence of regional and national policies, replacing earlier reactive local approaches  
496 (Wiegandt and Lugon, 2008). The disaster also marked a shift in risk perception, as hazards began to be  
497 understood as natural processes rather than divine punishment (Carrivick and Tweed, 2016).

498

499 **Monitoring and surveillance**

500 The first known glacier inventory in the Alps dates back to the late 19<sup>th</sup> century (Vernagtferner), an area  
501 characterised by frequent GLOFs over the past 400 years (Hoinkes, 1969). Systematic photogrammetric  
502 surveys over a century enabled the first detailed analyses of glacier dynamics (Braun, 1995). Monitoring  
503 intensified during the 20<sup>th</sup> century; the Giétro glacier has been monitored since the 1960s, including  
504 measurements of flow velocity, structure, and mass balance. More recently, heavily instrumented sites  
505 such as the Matterhorn have integrated multi-sensor monitoring systems (Weber et al., 2025) following  
506 recurrent instabilities (Occhiena and Pirulli, 2012).

507

508 **Multidisciplinary network**

509 The 1892 Tête Rousse disaster was a foundational event in glacier risk management. The debris flow  
510 has affected several villages downstream, causing 175 victims (Mougins and Bernard, 1922). It led to the

511 formation of multidisciplinary teams combining engineers, glaciologists, and foresters to investigate  
512 causes and implement mitigation measures (Marco et al., 2012; Sirop et al., 2022).  
513 Similarly, the 1965 Allalin glacier collapse at Mattmark (88 fatalities; Vivian, 1966) triggered  
514 coordinated scientific and engineering responses (Dalban Canassy et al., 2011). This led to the creation  
515 of a national working group on glacier hazards in Switzerland and the development of systematic hazard  
516 assessment methods (Haeberli, 1983; Haeberli et al., 2004). A coalition of scientists has irrevocably  
517 advanced the understanding of the mechanism of phased glacial rupture and sparked discussion on  
518 practical methods of risk assessment at the international level (Huggel et al., 2004).  
519 The Rochemelon lake case further illustrates this coordination. Cross-border collaboration enabled  
520 monitoring, risk assessment, public warning, and controlled drainage of the lake (Vincent et al., 2010).  
521 This case highlighted the need for continuity between scientific knowledge, decision-making, and  
522 operational action (Marco et al., 2012, unpublished).

523

#### 524 ***Reduction of exposure and vulnerability (evacuations, works, etc.)***

525 The first known organised evacuations for gravitational hazards occurred in 1987 (Val Pola rockslide;  
526  $30 \times 10^6$  m<sup>3</sup>; (Dramis et al., 1995). Early warnings based on slope instability enabled preventive  
527 evacuation (Chardon, 1990). For glacial hazards, evacuations such as Planpincieux (1997) were based  
528 on scenario modelling and monitoring data (Margreth and Funk, 1999; (Margreth et al., 2011). These  
529 approaches marked the emergence of anticipatory risk management. Engineering measures also  
530 developed, as illustrated by the Belvedere glacier (IT) and the Effimero lake crisis (2002), where  
531 combined strategies (monitoring, mapping, evacuation planning, and artificial drainage) were  
532 implemented (Haeberli et al., 2002).

533

#### 534 ***Predictive study***

535 In 1973, the first successful glacier collapse prediction was achieved at Weisshorn (CH) (Röthlisberger,  
536 1981b). Scientists observed an acceleration of the upper part of the glacier, accompanied by warnings  
537 from local mountain guides (Faillettaz and Funk, 2013). Subsequent advances enabled short-term  
538 predictions based on displacement monitoring, such as the Whymper glacier collapse forecast 10 days  
539 in advance (Faillettaz et al., 2016). Authorities adapted responses accordingly, restricting access while  
540 avoiding unnecessary evacuation. These decisions increasingly relied on modelling and scenario-based  
541 approaches (Schindelegger and Kanonier, 2019).

542 These developments are part of broader research programmes (e.g. *Glaciorisk*, *PERMAdataROC*,  
543 *GlariskAlp*, *PAPROG*, etc.), aimed at improving hazard understanding and management.

544

#### 545 ***Institutional risk understanding***

546 Swiss National Research Programmes (NRPs), significantly contributed to the institutionalisation of risk  
547 knowledge. Projects under NRP 31, 48, and 61 advanced understanding of climate-driven hazards,  
548 spatial planning tools, and emerging glacial lakes (Haeberli et al., 1999, 2016; Rothenbühler, 2006).  
549 More broadly, Alpine risk management has evolved from “hazard protection” to “hazard management”  
550 and ultimately to “risk management” (Link and Stötter, 2015).

551 Before 1950, strategies focused on structural protection (protective works) and exposure reduction  
552 (legislation). There are numerous traces of such works in 19<sup>th</sup> century archives. For example, galleries  
553 were excavated within the ice of the Allalingletscher glacier in 1834 to lower the level of its lake  
554 (dammed by the glacier tongue) and reduce the already high frequency of ice break-offs (Mariétan,  
555 1953). Between 1589 and 1850, documents report 26 major sudden outburst events (Raymond et al.,  
556 2003).

557 After 1950, improved process understanding led to hazard mapping (Link and Stötter, 2015) and  
558 probabilistic approaches; this was the age of *hazard management*.

559 Since the late 20<sup>th</sup> century, risk management has adopted a holistic perspective (Link and Stötter, 2015),  
560 integrating vulnerability and risk acceptance. Before the 2000s, hazard management was considered *in*  
561 *response* to an event. Recent approaches increasingly consider cascading processes (e.g. Evans et al.,

Commenté [JB8]: W. Haeberli : The lake was dammed by the glacier tongue, not a “glacial moraine dam”.

562 2009) and multiple scenarios, reflecting the growing complexity of cryospheric hazards under climate  
563 change (Allen et al., 2022).

564 Historically, the scope of hazard assessment has progressively expanded from glacier-related processes  
565 to include permafrost degradation and emerging landforms such as proglacial lakes. Current frameworks  
566 now adopt integrated, forward-looking approaches that account for long-term cryosphere evolution and  
567 cascading risks (Allen et al., 2017).

568

#### 569 4.2.4 Proactive vs. reactive management

570

571 Our typology of glacial and periglacial risk management distinguishes between proactive and reactive  
572 approaches. Management is considered *proactive* when it relies on anticipatory measures to prevent  
573 potential risks. Within the Disaster Risk Management (DRM) framework, prevention, mitigation  
574 (including spatial planning, protection measures), long-term adaptation to natural and socioeconomic  
575 changes, and the development of response and recovery capacities are all proactive strategies (Tagarev  
576 et al., 2021).

577 Management is *reactive* when it occurs during or after an event, aiming to manage the emergency,  
578 analyse impacts, and improve future risk reduction measures (Tagarev et al., 2021).

579 The different principles of glacial and periglacial hazards management can operate independently but  
580 are often combined. For instance, Monitoring and Surveillance (Pillar 4) is frequently coupled with  
581 Exposure, Vulnerability or Hazard Mitigation (Pillar 5).

582 Regarding torrential hazards associated with glacial lakes, a proactive example is the 1985 warming on  
583 the Arsine proglacial lake, which led to partial artificial drainage (Peissier and Courtray, 2012).  
584 Similarly, an early warning system installed in 2008 at the Grindelwaldgletscher glacial lake (CH) based  
585 on pressure sensors, enabled automatic alerts and proved effective during a drainage event (Bauder,  
586 2017). **Or the artificial lowering of the water level of Lake delle Locce and the construction of dams on  
587 the main torrent downstream of the Belvedere Glacier (Haeberli and Epifani, 1986).**

588 Conversely, the 1964 overflow of the Patinoire lake (FR), which caused damage downstream as far as  
589 the village of Pralognan, prompted a reactive response. Post-event analyses included field surveys,  
590 susceptibility studies, and impact modelling, leading to its later designation as a priority monitoring site  
591 (Cathala et al., 2021).

592 The Whymper hanging glacier illustrates proactive management: in 1997, displacement measurements  
593 enabled prediction of an imminent collapse, prompting evacuation of Planpincieux (Margreth and Funk,  
594 1999). Continuous monitoring and preparedness measures have since been implemented (Faillettaz et  
595 al., 2015, 2016). Similarly, the Forni glacier is monitored using satellite and terrestrial laser scanning to  
596 detect instabilities (Fugazza et al., 2018).

597 In contrast, the 1965 Mattmark disaster (Allalin glacier) exemplifies reactive management. Authorities  
598 focused on emergency response and post-event investigation. The event later triggered political debate  
599 and highlighted the limits of predictability at the time (Ricciardi, 2016).

600 For rock avalanches, the Piz Cengalo case illustrates the transition from reactive to proactive  
601 management. Following a first event in 2011, monitoring systems and protective infrastructure were  
602 implemented (Walter et al., 2020). These included a storage basin, protective barriers, and early warning  
603 systems enabling rapid evacuation (Bohnenblust, 2017). **Legal proceedings are still ongoing regarding  
604 the safety precautions taken prior to the event (Joris, 2025).**

605 In 2017, a major rock avalanche and subsequent debris flows ( $3.1 \times 10^6 \text{ m}^3$ ) affected the Val Bondasca.  
606 Despite underestimation of flow magnitude, early warning systems and protective measures  
607 significantly reduced damage (OECD, 2018). This case demonstrates the effectiveness of combining  
608 past-event analysis, monitoring, and preparedness, although some casualties still occurred (Walter et al.,  
609 2020).

610

611

## 612 5 Discussions

**Commenté [JB9]:** W. Haeberli : In the following section, the large amounts of reactive to proactive measures at Belvedere/Macugnaga und Gruben/Saas Balen should be mentioned.

**Commenté [JB10R9]:** Authors : As before: as there are many examples, we have chosen to include only a few.

**Commenté [JB11]:** W. Haeberli : Marti, K. (2025): Der steinige Weg vom Freispruch zum Fehlurteil. In: Joris, E. (Hg.): Mattmark 1965, Erinnerungen, Gerichtsurteile, italienisch-schweizerische Verflechtungen, 101 - 145. Rotpunktverlag.  
Note that the same author in the same book on pages 146-147 under the title «Das Dogma der Unvorhersehbarkeit, kein Phänomen der Vergangenheit» discusses the problematic legal situation after the Cengalo 2017 event. The court case is still ongoing

613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663

### 5.1 Cascading processes affecting increasingly vulnerable populations

As shown by the events documented in this study, chain reactions in glacial and periglacial environments tend to amplify beyond their initial triggering processes (OECD, 2018; Jacquemart et al., 2024), posing increasing challenges for risk management. This trend is closely linked to rising exposure in Alpine region. Mountain valleys have experienced significant urbanisation and infrastructure development in recent decades, increasing pressure on historically exposed areas (Hock and Rasul, 2022). This growing exposure is compounded by systemic vulnerability due to infrastructure interdependence: the destruction of a road, bridge, or energy network during a cascading event can isolate communities and hinder response capacities, as illustrated by La Bérarde in 2024 (Blanc et al., 2024). Risk management systems have long treated hazards separately, but this segmented approach is increasingly inadequate for complex, interacting phenomena affecting multiple components of populated territories. There is therefore a need for integrated and **future/scenario-based approaches** (Allen et al., 2017, 2022) that model process interactions while accounting for the spatial distribution of populations and infrastructure (Tacnet et al., 2010; Nyberg et al., 2026). These challenges are further intensified by climate change, which increases the likelihood of cascading processes in densely populated Alpine valleys.

### 5.2 Biases linked to difficulties in compiling an exhaustive inventory of events

Compiling an inventory that aims to be exhaustive according to defined selection criteria entails limitations and requires contextualisation of each event.

#### *The data source effect*

The apparent increase in events over time partly reflects biases related to data availability (e.g. Giacona et al., 2017). This *data source effect* is closely linked to the nature and distribution of information sources. As Giacona (2019) notes, the spatial and temporal distribution of events depends strongly on the structure of the documentary corpus. Each recorded event therefore depends on the quality and diversity of available sources. Scarcity of historical records limits traceability in earlier periods, whereas recent data abundance increases event detection.

The data source effect thus requires careful consideration of variations in documentation volume. For instance, technological advances such as webcam monitoring (e.g. Gornergletscher) generate continuous and detailed observations, increasing the number of recorded events (Huss et al., 2007).

#### *Land use and the event prism*

The presence or absence of sources is also influenced by human occupation. The growing presence of people in high mountains – through mountaineering (Mourey et al., 2023) and valley urbanisation (Vannier et al., 2016) – has significantly increased observation capacity. Conversely, abandoned settlements and pastures were once important witnesses to past hazards.

Even when hazards occur, their recognition depends on societal perception. An event may remain unrecorded if it is not perceived as significant. As Giacona (2019) argues, an event gains social existence only when it is perceived as a disruption. Risk perception evolves over time and is shaped by collective experience, knowledge, and social context (Granet-Abisset, 2012). Increasing attention to high-mountain hazards also reflects rising societal expectations for safety and accountability, which in turn drive the production and diversification of information sources.

### 5.3 Resilience and response as late-developed concepts rather than absent practices

The emergence of Pillar 8 (Resilience) from the 1980s-90s onwards, and its consolidation after 2000, should be interpreted cautiously. It does not indicate a prior absence of resilience from Alpine communities, but rather the late institutionalisation of the concept in scientific and policy frameworks.

**Commenté [JB12]:** W. Haeberli : Concerning integrative and – especially also – future/scenario-based approaches, GAPHAZ and in its updated version Allen et al. 2022 should be cited.

664 Post-event reconstruction dynamics existed well before this period, but were not conceptualised within  
665 this framework, introducing an interpretative bias in diachronic analysis.  
666 Similarly, Pillar 7 (Response) appears from the 1950s and remains present thereafter, without becoming  
667 dominant. Its consistent but limited representation suggests that emergency response has always existed  
668 but is less documented due to lower formalisation, leading to potential under-representation in the  
669 database.

670 The use of a ninth separate pillar representing ‘*Experience feedback*’ was considered, since the  
671 experience gained from an event enables a “structured process of capitalising on and utilising knowledge  
672 resulting from the analysis of positive and/or negative events” (Villeneuve et al., 2010).

673 Feedback is the very condition for an event’s existence in archives: its narration and memory make it  
674 historically visible. It therefore applies to every event, and would have created a bias in relation to the  
675 other pillars. Therefore, it is implicitly included in Pillar 1.

676

#### 677 5.4 National discrepancies in prevention

678

679 The study highlights that risk reduction measures are not uniform across Alpine countries and reveals a  
680 geographical bias towards France, Switzerland, and Italy. This heterogeneity reflects significant  
681 differences in legislative, institutional, and cultural frameworks for risk management. . International  
682 cooperation and governance initiatives such as GAPHAZ exist to try to bridge these differences (Allen  
683 et al., 2022).

684 In France, spatial planning relies on binding national regulations, and scientific action plans are also  
685 defined at the national level. In Switzerland, management is largely devolved to the cantons,  
686 encouraging locally adapted approaches but potentially generating inequalities depending on available  
687 resources. In Italy, administrative fragmentation and regional disparities hinder the implementation of  
688 coherent policies at the scale of catchments.

689 These governance differences affect both data comparability and the ability to derive cross-cutting  
690 insights. More broadly, they highlight the need for stronger transnational Alpine governance. Given that  
691 hazards transcend political borders, enhanced institutional coordination is essential for effective and  
692 equitable risk prevention.

693

694

## 695 6 Conclusions and perspectives

696

697 This study moves beyond traditional approaches that focus primarily on post-event causes or hazards  
698 characterization, and instead examines management practices. While event inventories inform on  
699 frequency, they rarely capture organisational dynamics or adaptative capacities. Our analysis addresses  
700 these dimensions.

701 By analysing 200 events since the early 19<sup>th</sup> century (some even earlier, though less frequent), we  
702 observe both an intensification of events and a progressive formalisation of management practices,  
703 reflected in archival records. The growing number of simultaneously mobilised pillars in recent years  
704 indicates a shift from single-discipline approaches to multi-actor, multi-scale governance. Beyond  
705 increased event reporting, this reflects the maturation of institutional Alpine risk management  
706 frameworks. Although all high-altitude massifs are affected, our data highlights a concentration of major  
707 events in France and western Switzerland, , particularly for glaciers, new lakes and permafrost-affected  
708 /glacially de-buttressed rock faces. The completeness of this long-term inventory remains constrained  
709 by sources and perception biases.

710 Risk management has evolved in response to both major events and improved process understanding. It  
711 continues to rapidly change far beyond historical and empirical antecedence. Forecasting,  
712 instrumentation, and monitoring of unstable zones have expanded, alongside contingency planning and  
713 evacuation strategies. Measures to reduce exposure and vulnerability increasingly account for evolving  
714 hazards and urbanisation pressures. Since the early 21<sup>st</sup> century, scientific initiatives aimed at

**Commenté [JB13]:** W. Haeberli : Better “risk reduction”.  
The term risk is a mental construct and as such cannot be  
“prevented”.

**Commenté [JB14]:** W. Haeberli : Such international  
cooperation and governance are especially also encouraged  
by GAPHAZ and Allen 2022.

**Commenté [JB15]:** W. Haeberli : Better “particularly for  
glaciers, new lakes and permafrost-affected/glacially de-  
buttressed rock faces.”

**Commenté [JB16]:** W. Haeberli : developed and continue  
to rapidly change far beyond historical/empirical  
precedence”

715 anticipating high-mountain mass movements that have multiplied. Knowledge exchange across borders  
716 has intensified, supporting more integrated and holistic risk management approaches. Hazards are now  
717 increasingly analysed as interconnected or cascading phenomena. As risk management grows more  
718 complex, the social sciences emerge as an indispensable complement to natural sciences, bringing to the  
719 fore dimensions – community vulnerability, risk perception, territorial identity, institutional dynamics –  
720 that physical approaches alone cannot capture.

721 Switzerland appears as a leading actor in the institutionalisation of monitoring and hazard assessment,  
722 particularly for glaciers and rock faces. Multidisciplinary collaboration and international knowledge  
723 exchange are rapidly advancing.

724 Across many regions worldwide, glacial and permafrost environments now differ markedly from the  
725 conditions under which settlements and infrastructure developed. Historical and empirical knowledge  
726 alone is therefore insufficient to anticipate future glacier- and permafrost-related risks. This calls for  
727 integrated, forward-looking approaches, based on continuous observation, repeated expert assessments,  
728 and rapidly evolving technologies. Risk analysis must adopt a systemic perspective, covering the full  
729 chain from hazard sources to socio-economic impacts.

730

731 *Data availability.* Raw inventory data is available on data.InDoRES

732 (<https://doi.org/10.48579/PRO/JA5PDT>)

733 *Author contributions.* JB, LR and SC designed the study and JB carried it out. JB prepared the  
734 manuscript with contributions from all co-authors.

735

736 *Competing interests.* The authors declare that they have no conflict of interest.

737

738

### 739 **References**

740 Alean, J.: Ice Avalanches: Some Empirical Information about their Formation and Reach, *J. Glaciol.*, 31,  
741 324–333, <https://doi.org/10.3189/S002214300006663>, 1985.

742 Allen, S., Frey, H., and Huggel, C.: Assessment of Glacier and Permafrost Hazards in Mountain Regions.  
743 Technical Guidance Document, <https://doi.org/10.13140/RG.2.2.26332.90245>, 2017.

744 Allen, S., Frey, H., Haerberli, W., Huggel, C., Chiarle, M., and Geertsema, M.: Assessment Principles for  
745 Glacier and Permafrost Hazards in Mountain Regions, in: *Oxford Research Encyclopedia of Natural  
746 Hazard Science*, <https://doi.org/10.1093/acrefore/9780199389407.013.356>, 2022.

747 Ancey, C., Bardou, E., Funk, M., Huss, M., Werder, M. A., and Trehwela, T.: Hydraulic Reconstruction of  
748 the 1818 Giétro Glacial Lake Outburst Flood, *Water Resour. Res.*, 55, 8840–8863,  
749 <https://doi.org/10.1029/2019WR025274>, 2019.

750 [geoclimalp.to.cnr.it/landslide-inventory](http://geoclimalp.to.cnr.it/landslide-inventory): [http://geoclimalp.to.cnr.it/landslide-](http://geoclimalp.to.cnr.it/landslide-inventory/#6/43.659/13.554)  
751 [inventory/#6/43.659/13.554](http://geoclimalp.to.cnr.it/landslide-inventory/#6/43.659/13.554), last access: 4 March 2026.

752 Gridabase - Glaciorisk: <https://www.nimbus.it/glaciorisk/gridabasemainmenu.asp>, last access: 27  
753 May 2025.

754 Bauder, A.: The Swiss Glaciers 2013/14 and 2014/15 - Glaciological Report No. 135/136, 145pp,  
755 [https://doi.org/10.18752/GLREP\\_135-136](https://doi.org/10.18752/GLREP_135-136), 2017.

756 Beccera, S. and Peltier, A.: L'information préventive pour réduire la vulnérabilité aux risques  
757 d'inondation, élaboration et efficacité d'une réponse sociale, *Chang. Clim. Métarisque À Méta-Gouv.*,  
758 pp.35-53, 2011.

759 Bisquert, A., Mainieri, R., Carladous, S., Robert, Y., Giacona, F., Verry, P., and Eckert, N.: La base de  
760 données événementielles RTM pour la connaissance des risques naturels en montagne : L'exemple du  
761 département de l'Isère (France), *J. Alp. Res. Rev. Géographie Alp.*, <https://doi.org/10.4000/137kf>,  
762 2025.

763 Blanc, A., Misset, C., Mainieri, R., and Llamas, B.: Rétro-analyse de la crue du torrent des Etançons du  
764 21 juin 2024, RTM, 2024.

765 Bohnenblust, S.: Radar system monitors Pizzo Cengalo, Bondo, GEOPRAEVENT AG – Electron. Monit.  
766 Nat. Hazards, 2017.

767 Bondesan, A. and Francese, R. G.: The climate-driven disaster of the Marmolada Glacier (Italy),  
768 *Geomorphology*, 431, 108687, <https://doi.org/10.1016/j.geomorph.2023.108687>, 2023.

769 Boudières, V. and Peisser, C.: Prévention des Risques d'Origine Glaciaire et Périglaciaire (ROGP) -  
770 Synthèse des connaissances et des pratiques, PARN, 2013.

771 Bourrelrier, P.-H. and Dunglas, J.: Des événements naturels extrêmes aux figures de la catastrophe;,  
772 *Ann. Mines - Responsab. Environ.*, N° 56, 41–47, <https://doi.org/10.3917/re.056.0041>, 2009.

773 Braun, L. N.: Les résultats principaux de 100 ans de recherche au Vernagtferner, Oetztal (Autriche),  
774 *Houille Blanche*, 81, 109–110, <https://doi.org/10.1051/lhb/1995054>, 1995.

775 Büntgen, U., Oppenheimer, C., Farinotti, D., Nahtz, T., and Esper, J.: The 2025 Blatten disaster in the  
776 Swiss Alps followed exceptional warming and highlights the vulnerability of people and heritage in  
777 glaciated landscapes, *Commun. Earth Environ.*, 6, 994, <https://doi.org/10.1038/s43247-025-02994-8>,  
778 2025.

779 Capozzi, C.: Mémoire de master en histoire - La catastrophe de Mattmark par la presse, Université de  
780 Franche-Comté, 2011.

781 Carrivick, J. L. and Tweed, F. S.: A global assessment of the societal impacts of glacier outburst floods,  
782 *Glob. Planet. Change*, 144, 1–16, <https://doi.org/10.1016/j.gloplacha.2016.07.001>, 2016.

783 Cathala, M., Magnin, F., Linsbauer, A., and Haerberli, W.: Modelling and characterizing glacier-bed  
784 overdeepenings as sites for potential future lakes in the deglaciating French Alps, *Géomorphologie  
785 Relief Process. Environ.*, 27, 19–36, <https://doi.org/10.4000/geomorphologie.15255>, 2021.

786 Cathala, M., Bock, J., Abdulsamad, F., Deline, P., Josnin, J.-Y., Ravanel, L., Revil, A., Richard, J., Verroust,  
787 F., and Magnin, F.: Assessing the role of permafrost in the preconditioning and triggering factors of  
788 the September 2020 Crête des Grangettes rockfall (southern French Alps), *Géomorphologie Relief  
789 Process. Environ.*, 30, <https://doi.org/10.4000/12yqn>, 2024.

790 Cauliez, R.: Mémoire M2 - Au-delà des eaux glacées. Pouvoirs gestionnaires et expériences populaires  
791 sous le glacier de Tête-Rousse., *Ecole Normale Supérieure*, 229 pp., 2024.

792 Chardon, M.: Les catastrophes naturelles de l'été 1987 en Lombardie : crues, inondations,  
793 écoulement de Val Pola, *Rev. Géographie Alp.*, 78, 59–87, <https://doi.org/10.3406/rga.1990.2767>,  
794 1990.

795 Chiarle, M., Iannotti, S., Mortara, G., and Deline, P.: Recent debris flow occurrences associated with  
796 glaciers in the Alps, *Glob. Planet. Change*, 56, 123–136,  
797 <https://doi.org/10.1016/j.gloplacha.2006.07.003>, 2007.

798 Chiarle, M., Viani, C., Mortara, G., Deline, P., Tamburini, A., and Nigrelli, G.: Large glacier failures in  
799 the Italian Alps over the last 90 years, *Geogr. Fis. E Din. Quat.*, 45, 19–40,  
800 <https://doi.org/10.4461/GFDQ.2022.45.2>, 2022.

801 Dalban Canassy, P., Bauder, A., Dost, M., Fäh, R., Funk, M., Margreth, S., Müller, B., and Sugiyama, S.:  
802 Hazard assessment investigations due to recent changes in Triftgletscher, Bernese Alps, Switzerland,  
803 *Nat. Hazards Earth Syst. Sci.*, 11, 2149–2162, <https://doi.org/10.5194/nhess-11-2149-2011>, 2011.

804 Dauphiné, A. and Provitolo, D.: Risques et catastrophes, Armand Colin,  
805 <https://doi.org/10.3917/arco.dauph.2013.01>, 2013.

806 Davide, F., Antonella, S., Roberto Sergio, A., Carlo, D., Daniele, C. B., Luca, M., Fabiano, V., Claudio, S.,  
807 and Guglielmina Adele, D.: Variations of Lys Glacier (Monte Rosa Massif, Italy) from the Little Ice Age  
808 to the Present from Historical and Remote Sensing Datasets, in: *Glaciers and the Polar Environment*,  
809 edited by: Kanao, M., Godone, D., and Dematteis, N., IntechOpen,  
810 <https://doi.org/10.5772/intechopen.91202>, 2021.

811 Defossez, S., Vinet, F., and Leone, F.: Diagnostiquer la vulnérabilité face aux inondations : progrès et  
812 limites, in: *Inondations 1 : la connaissance du risque*, 2018.

813 Dematteis, N., Giordan, D., Troilo, F., Wrzesniak, A., and Godone, D.: Ten-Year Monitoring of the  
814 Grandes Jorasses Glaciers Kinematics. Limits, Potentialities, and Possible Applications of Different  
815 Monitoring Systems, *Remote Sens.*, 13, 3005, <https://doi.org/10.3390/rs13153005>, 2021.

816 Dematteis, N., Troilo, F., Scotti, R., Colombarolli, D., Giordan, D., and Maggi, V.: The use of terrestrial  
817 monoscopic time-lapse cameras for surveying glacier flow velocity, *Cold Reg. Sci. Technol.*, 222,  
818 104185, <https://doi.org/10.1016/j.coldregions.2024.104185>, 2024.

819 Demolis, B., Kuss, D., and Serrano, C.: Etude du lac de la Patinoire à Pralognan la Vanoise (73), ONF-  
820 RTM INRAE, 2021.

821 Desailly, B.: Crues et inondations en Roussillon : le risque, le discours et l'aménagement, *Rev.*  
822 *Géographique Pyrén. Sud-Ouest*, 61, 515–519, <https://doi.org/10.3406/rgpso.1990.3229>, 1990.

823 Dipartimento della Protezione Civile: Rischio glaciale e periglaciale in ambiente alpino: un quadro  
824 metodologico, *Prima.*, 126 pp., 2025.

825 Dollfus, O. and D'Ercole, R.: Les mémoires des catastrophes au service de la prévision et de la  
826 prévention des risques naturels, 7–18, 1996.

827 Dosse, F.: Renaissance de l'événement, 1<sup>re</sup> éd., Presses Universitaires de France,  
828 <https://doi.org/10.3917/puf.dosse.2010.01>, 2010.

829 Dourlens, C.: La question des inondations au prisme des sciences sociales : un panorama de la  
830 recherche publique, Ministère de l'Équipement, des Transports, du Logement, du Tourisme et de la  
831 Mer, 112 pp., 2004.

832 Dramis, F., Govi, M., Guglielmin, M., and Mortara, G.: Mountain Permafrost and Slope Instability in  
833 the Italian Alps: the Val Pola Landslide, 6, 73–82, 1995.

834 Du Pasquier, L.: L'avalanche du glacier de l'Altels le 11 septembre 1895, *Ann. Géographie*, 5, 458–468,  
835 <https://doi.org/10.3406/geo.1896.5954>, 1896.

- 836 Einhorn, B.: Changement climatique et risques naturels - Prévention, Gestion intégrée, Adaptation,  
837 2017.
- 838 Einhorn, B. and Peisser, C.: Gestion intégrée du risque généré par les poches d'eau du glacier de Tête  
839 Rousse – De la tragédie de 1892 à la gestion de crise de 2010, Grenoble, 2011.
- 840 Etzelmüller, B., Guglielmin, M., Hauck, C., Hilbich, C., Hoelzle, M., Isaksen, K., Noetzli, J., Oliva, M., and  
841 Ramos, M.: Twenty years of European mountain permafrost dynamics—the PACE legacy, *Environ. Res.  
842 Lett.*, 15, 104070, <https://doi.org/10.1088/1748-9326/abae9d>, 2020.
- 843 Evans, S. G., Tutubalina, O. V., Drobyshev, V. N., Chernomorets, S. S., McDougall, S., Petrakov, D. A.,  
844 and Hungr, O.: Catastrophic detachment and high-velocity long-runout flow of Kolka Glacier, Caucasus  
845 Mountains, Russia in 2002, *Geomorphology*, 105, 314–321,  
846 <https://doi.org/10.1016/j.geomorph.2008.10.008>, 2009.
- 847 Faillietaz, J. and Funk, M.: Instabilités glaciaires et prédiction, *Société Vaudoise Sci. Nat.*, 159–174,  
848 2013.
- 849 Faillietaz, J., Funk, M., and Vincent, C.: Avalanching glacier instabilities: Review on processes and early  
850 warning perspectives, *Rev. Geophys.*, 53, 203–224, <https://doi.org/10.1002/2014RG000466>, 2015.
- 851 Faillietaz, J., Funk, M., and Vagliasindi, M.: Time forecast of a break-off event from a hanging glacier,  
852 *The Cryosphere*, 10, 1191–1200, <https://doi.org/10.5194/tc-10-1191-2016>, 2016.
- 853 Favier, R.: Sociétés urbaines et culture du risque. Les inondations dans la France d'Ancien Régime, in:  
854 *Les cultures du risque (XVIe-XXIe siècle)*, pp.49-86, 2006.
- 855 Field, C., Barros, V., Stocker, T., and Dahe, Q.: Managing the risks of extreme events and disasters to  
856 advance climate change adaption, Cambridge university press, New York, 582 pp., 2012.
- 857 Fischer, L., Kääb, A., Huggel, C., and Noetzli, J.: Geology, glacier retreat and permafrost degradation as  
858 controlling factors of slope instabilities in a high-mountain rock wall: the Monte Rosa east face, *Nat.  
859 Hazards Earth Syst. Sci.*, 6, 761–772, <https://doi.org/10.5194/nhess-6-761-2006>, 2006.
- 860 Fischer, L., Purves, R. S., Huggel, C., Noetzli, J., and Haeberli, W.: On the influence of topographic,  
861 geological and cryospheric factors on rock avalanches and rockfalls in high-mountain areas, *Nat.  
862 Hazards Earth Syst. Sci.*, 12, 241–254, <https://doi.org/10.5194/nhess-12-241-2012>, 2012.
- 863 Flotron, A.: Movement Studies on a Hanging Glacier in Relation with an Ice Avalanche, *J. Glaciol.*, 19,  
864 671–672, <https://doi.org/10.3189/S0022143000029592>, 1977.
- 865 Forel, F. A.: L'Eboulement du glacier des Altels, Bureau des Archives, Genève, 513 pp., 1895.
- 866 Fournier, M.: Le riverain introuvable ! La gestion du risque d'inondation au défi d'une mise en  
867 perspective diachronique : une analyse menée à partir de l'exemple de la Loire., *Aménagement de  
868 l'Espace et Urbanisme*, Université François-Rabelais, Tours, 432 pp., 2010.
- 869 Francese, R. G., Valentino, R., Haeberli, W., Bondesan, A., Giorgi, M., Picotti, S., Pettenati, F., Sandron,  
870 D., Ramponi, G., and Valt, M.: Failure of Marmolada Glacier (Dolomites, Italy) in 2022: Data-based  
871 back analysis of possible collapse mechanisms as related to recent morpho-climatic evolution and  
872 possible trigger factors, <https://doi.org/10.5194/nhess-2024-212>, 19 November 2024.

- 873 Fugazza, D., Scaioni, M., Corti, M., D'Agata, C., Azzoni, R. S., Cernuschi, M., Smiraglia, C., and Diolaiuti,  
874 G. A.: Combination of UAV and terrestrial photogrammetry to assess rapid glacier evolution and map  
875 glacier hazards, *Nat. Hazards Earth Syst. Sci.*, 18, 1055–1071, [https://doi.org/10.5194/nhess-18-1055-](https://doi.org/10.5194/nhess-18-1055-2018)  
876 2018, 2018.
- 877 Giacona, F., Eckert, N., and Martin, B.: A 240-year history of avalanche risk in the Vosges Mountains  
878 based on non-conventional (re)sources, *Nat. Hazards Earth Syst. Sci.*, 17, 887–904,  
879 <https://doi.org/10.5194/nhess-17-887-2017>, 2017.
- 880 Giacona, F., Martin, B., Eckert, N., and Desarthe, J.: Une méthodologie de la modélisation en  
881 géohistoire : de la chronologie (spatialisée) des événements au fonctionnement du système par la  
882 mise en correspondance spatiale et temporelle, *Physio-Géo*, 171–199,  
883 <https://doi.org/10.4000/physio-geo.9186>, 2019.
- 884 Giardino, M., Ratto, S., Alberto, W., Armand, M., Cignetti, M., and Navillod, E.: Regional and local risk  
885 assessments of alluvial fans by combination of historical and geomorphological data on debris flows,  
886 the most damaging natural hazard in the Aosta Valley Region (NW-Italy), 2010.
- 887 Giordan, D., Dematteis, N., Allasia, P., and Motta, E.: Classification and kinematics of the Planpincieux  
888 Glacier break-offs using photographic time-lapse analysis, *J. Glaciol.*, 66, 188–202,  
889 <https://doi.org/10.1017/jog.2019.99>, 2020.
- 890 Girard, S. and Rivière-Honegger, A.: Le choix et la pratique de la monographie diachronique.  
891 Contribution à l'étude de l'efficacité environnementale de la territorialisation de la politique de l'eau.,  
892 in: *Environnement, politiques publiques et pratiques locales*, 359–384, 2015.
- 893 Granet-Abisset, A.-M.: L'historien, les risques et l'environnement : un regard sur la nature et les  
894 hommes, in: *23èmes Journées Scientifiques de l'Environnement - Risques environnementaux :  
895 détecter, comprendre, s'adapter*, 2012.
- 896 Gruber, S. and Haeberli, W.: Permafrost in steep bedrock slopes and its temperature-related  
897 destabilization following climate change, *J. Geophys. Res. Earth Surf.*, 112, 2006JF000547,  
898 <https://doi.org/10.1029/2006JF000547>, 2007.
- 899 Haeberli, W.: Frequency and Characteristics of Glacier Floods in the Swiss Alps, *Ann. Glaciol.*, 4, 85–  
900 90, <https://doi.org/10.3189/S0260305500005280>, 1983.
- 901 Haeberli, W. and Epifani, F.: Mapping the Distribution of Buried Glacier Ice – An Example From Lago  
902 Delle Locce, Monte Rosa, Italian Alps, *Ann. Glaciol.*, 8, 78–81,  
903 <https://doi.org/10.3189/S026030550000118X>, 1986.
- 904 Haeberli, W., Käab, A., Hoelzle, M., Bösch, H., Funk, M., Mühl, D., and Keller, F.: *Eisschwund und  
905 Naturkatastrophen im Hochgebirge*, vdf Hochschulverlag an der ETH Zürich, Zürich, 190pp pp., 1999.
- 906 Haeberli, W., Käab, A., Mühl, D. V., and Teyssie, P.: Prevention of outburst floods from periglacial  
907 lakes at Grubengletscher, Valais, Swiss Alps, *J. Glaciol.*, 47, 111–122,  
908 <https://doi.org/10.3189/172756501781832575>, 2001.
- 909 Haeberli, W., Käab, A., Paul, F., Chiarle, M., Mortara, G., Mazza, A., Deline, P., and Richardson, S.: A  
910 surge-type movement at Ghiacciaio del Belvedere and a developing slope instability in the east face  
911 of Monte Rosa, Macugnaga, Italian Alps, *Nor. Geogr. Tidsskr. - Nor. J. Geogr.*, 56, 104–111,  
912 <https://doi.org/10.1080/002919502760056422>, 2002.

913 Haeberli, W., Huggel, C., Kääh, A., Gruber, S., Noetzli, J., and Zraggen-Oswald, S.: Proceedings of the  
914 International Conference „High mountain hazard prevention”, 2004.

915 Haeberli, W., Buetler, M., Huggel, C., Friedli, T. L., Schaub, Y., and Schleiss, A. J.: New lakes in  
916 deglaciating high-mountain regions – opportunities and risks, *Clim. Change*, 139, 201–214,  
917 <https://doi.org/10.1007/s10584-016-1771-5>, 2016.

918 Harris, C., Haeberli, W., Vonder Mühl, D., and King, L.: Permafrost monitoring in the high mountains  
919 of Europe: the PACE Project in its global context, *Permafr. Periglac. Process.*, 12, 3–11,  
920 <https://doi.org/10.1002/ppp.377>, 2001.

921 Hock, R. and Rasul, G.: The Ocean and Cryosphere in a Changing Climate: Special Report of the  
922 Intergovernmental Panel on Climate Change, 1st ed., Cambridge University Press,  
923 <https://doi.org/10.1017/9781009157964>, 2022.

924 Hoinkes, H. C.: Surges of the Vernagtferner in the Ötztal Alps since 1599, *Can. J. Earth Sci.*, 6, 853–861,  
925 <https://doi.org/10.1139/e69-086>, 1969.

926 Hugerot, T., Astrade, L., Gauchon, C., and Ployon, E.: De la carte diachronique à la trajectoire  
927 géohistorique : modélisation du changement paysager d’un cône torrentiel, *Mappemonde*,  
928 <https://doi.org/10.4000/mappemonde.6704>, 2021.

929 Huggel, C., Haeberli, W., Kääh, A., Bieri, D., and Richardson, S.: An assessment procedure for glacial  
930 hazards in the Swiss Alps, *Can. Geotech. J.*, 41, 1068–1083, <https://doi.org/10.1139/t04-053>, 2004.

931 Huggel, C., Fischer, L., Schneider, D., and Haeberli, W.: Research advances on climate-induced slope  
932 instability in glacier and permafrost high-mountain environments, *Geogr. Helvetica*, 65, 146–156,  
933 <https://doi.org/10.5194/gh-65-146-2010>, 2010.

934 Huggel, C., Carey, M., Clague, J. J., and Kääh, A.: The high-mountain cryosphere: environmental  
935 changes and human risks, Cambridge university press, Cambridge, United Kingdom, 363 pp., 2015.

936 Huss, M., Bauder, A., Werder, M., Funk, M., and Hock, R.: Glacier-dammed lake outburst events of  
937 Gornersee, Switzerland, *J. Glaciol.*, 53, 189–200, <https://doi.org/10.3189/172756507782202784>,  
938 2007.

939 Intergovernmental Panel On Climate Change (ipcc) (Ed.): Annex I: Glossary, in: *Climate Change 2022 -*  
940 *Mitigation of Climate Change*, Cambridge University Press, 1793–1820,  
941 <https://doi.org/10.1017/9781009157926.020>, 2023.

942 IPCC: Global Warming of 1.5°C: IPCC Special Report on Impacts of Global Warming of 1.5°C above  
943 Pre-industrial Levels in Context of Strengthening Response to Climate Change, Sustainable  
944 Development, and Efforts to Eradicate Poverty, 1st ed., Cambridge University Press,  
945 <https://doi.org/10.1017/9781009157940>, 2022.

946 IPCC: *Climate Change 2022 – Impacts, Adaptation and Vulnerability: Working Group II Contribution to*  
947 *the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, 1st ed., Cambridge  
948 University Press, <https://doi.org/10.1017/9781009325844>, 2023.

949 Jacquemart, M., Weber, S., Chiarle, M., Chmiel, M., Cicoira, A., Corona, C., Eckert, N., Gaume, J.,  
950 Giacona, F., Hirschberg, J., Kaitna, R., Magnin, F., Mayer, S., Moos, C., Van Herwijnen, A., and Stoffel,  
951 M.: Detecting the impact of climate change on alpine mass movements in observational records from  
952 the European Alps, *Earth-Sci. Rev.*, 258, 29, <https://doi.org/10.1016/j.earscirev.2024.104886>, 2024.

- 953 Joffe, H. and Orfali, B.: De la perception à la représentation du risque: le rôle des médias, *Hermès*, n°  
954 41, 121, <https://doi.org/10.4267/2042/8962>, 2005.
- 955 Joris, E.: *Mattmark 1965: Erinnerungen, Gerichtsurteile, italienisch-schweizerische Verflechtungen*, 1.  
956 Auflage., Rotpunktverlag, Zürich, 2025.
- 957 Kellerer-Pirklbauer, A., Slupetzky, H., and Avian, M.: Ice-avalanche impact landforms: the event in  
958 2003 at the glacier nördliches bockkarkees, hohe tauern range, austria, *Geogr. Ann. Ser. Phys. Geogr.*,  
959 94, 97–115, <https://doi.org/10.1111/j.1468-0459.2011.00446.x>, 2012.
- 960 Kienholz, H., Krummenacher, B., Kipfer, A., and Perret, S.: Aspects of Integral Risk Management in  
961 Practice - Considerations with Respect to Mountain Hazards in Switzerland, 45–50, 2004.
- 962 Lacroix, D., Mayet, L., and Roche, P.-A.: Risques d'origine glaciaire et périglaciaire - Elements en  
963 soutien à un plan d'action, IGEDD - IGA - IGESR, 2022.
- 964 Laïly, B. and Demolis, B.: *Glaciers blancs à risques - Retours d'expériences sur certaines crises*, ONF-  
965 RTM, 2019.
- 966 Leone, F., Meschinet de Richemond, N., and Freddy, V.: *Aléas naturels et gestion des risques*, Presses  
967 Universitaires de France, 288 p. pp., 2021.
- 968 Link, S. and Stötter, J.: The development of mountain risk governance: challenges for application,  
969 <https://doi.org/10.5194/nhessd-3-429-2015>, 16 January 2015.
- 970 Lucchesi, S., Fioraso, G., Bertotto, S., and Chiarle, M.: Little Ice Age and contemporary glacier extent in  
971 the Western and South-Western Piedmont Alps (North-Western Italy), *J. Maps*, 10, 409–423,  
972 <https://doi.org/10.1080/17445647.2014.880226>, 2014.
- 973 Magnin, F., Ravanel, L., Ben-Asher, M., Bock, J., Cathala, M., Duvillard, P.-A., Jean, P., Josnin, J.-Y.,  
974 Kaushik, S., Revil, A., and Deline, P.: De l'observation des écroulements aux solutions opérationnelles :  
975 près de deux décennies d'études sur les risques cryo-gravitaires dans le massif du Mont-Blanc, *Rev.*  
976 *Géographie Alp.*, 111–2, <https://doi.org/10.4000/rga.11644>, 2023.
- 977 Mainieri, R., Blanc, A., Astrade, L., Baratier, A., Berthet, J., Deline, P., Le Roy, M., Misset, C., De  
978 Montety, F., Robert, Y., and Schoeneich, P.: Les aspects géomorphologiques de la crue torrentielle du  
979 torrent des Étançons à la Bérarde du 21 juin 2024, *Géomorphologie Relief Process. Environ.*, 31,  
980 <https://doi.org/10.4000/14ipc>, 2025.
- 981 Mair, V., Zischg, A., Lang, K., Tonidandel, D., Krainer, K., Kellerer-Pirklbauer, A., Deline, P., Schoeneich,  
982 P., Cremonese, E., Pogliotti, P., Gruber, S., and Böckli, L.: PermaNET - Réseau d'observation du  
983 permafrost sur le long terme - Rapport de synthèse, Interpraevent, Klagenfurt, 2011.
- 984 Marco, O., Lailly, B., and Charles, F.: Mesures de protection et contraintes d'ingénierie en milieu  
985 glaciaire : expérience des services intervenus sur les cas récents - ONF-RTM, 2012.
- 986 Margreth, S. and Funk, M.: Hazard mapping for ice and combined snow/ice avalanches — two case  
987 studies from the Swiss and Italian Alps, 159–173, 1999.
- 988 Margreth, S., Faillettaz, J., Funk, M., Vagliasindi, M., Diotri, F., and Broccolato, M.: Safety concept for  
989 hazards caused by ice avalanches from the Whymper hanging glacier in the Mont Blanc Massif, *Cold*  
990 *Reg. Sci. Technol.*, 69, 194–201, <https://doi.org/10.1016/j.coldregions.2011.03.006>, 2011.

- 991 Margreth, S., Funk, M., Tobler, D., Dalban, P., Meier, L., and Lauper, J.: Analysis of the hazard caused  
 992 by ice avalanches from the hanging glacier on the Eiger west face, *Cold Reg. Sci. Technol.*, 144, 63–72,  
 993 <https://doi.org/10.1016/j.coldregions.2017.05.012>, 2017.
- 994 Mariétan, I.: Note de sciences naturelles sur la vallée de Saas, *Bull. de la Murithienne* 76, 88–102,  
 995 1959.
- 996 Moles, A.: Notes pour une typologie des événements, *Communications*, 18, 90–96,  
 997 <https://doi.org/10.3406/comm.1972.1261>, 1972.
- 998 Mougín, P. and Bernard, C.: Étude sur le glacier de Tête-Rousse. Les avalanches en Savoie., Tome 4, 3–  
 999 90, 1922.
- 1000 Mourey, J., Clivaz, C., and Bourdeau, P.: Analyser l'évolution des pratiques sportives en montagne peu  
 1001 aménagée à partir des données de fréquentation des cabanes. Application aux Alpes Valaisannes,  
 1002 *Rev. Géographie Alp.*, <https://doi.org/10.4000/rga.11110>, 2023.
- 1003 Niggli, L., Allen, S., Frey, H., Huggel, C., Petrakov, D., Raimbekova, Z., Reynolds, J., and Wang, W.: GLOF  
 1004 Risk Management Experiences and Options: A Global Overview, in: *Oxford Research Encyclopedia of*  
 1005 *Natural Hazard Science*, Oxford University Press,  
 1006 <https://doi.org/10.1093/acrefore/9780199389407.013.540>, 2024.
- 1007 Nigrelli, G., Chiarle, M., Nuzzi, A., Perotti, L., Torta, G., and Giardino, M.: A web-based, relational  
 1008 database for studying glaciers in the Italian Alps, *Comput. Geosci.*, 51, 101–107,  
 1009 <https://doi.org/10.1016/j.cageo.2012.07.027>, 2013.
- 1010 Nigrelli, G., Paranunzio, R., Turconi, L., Luino, F., Mortara, G., Guerini, M., Giardino, M., and Chiarle,  
 1011 M.: First national inventory of high-elevation mass movements in the Italian Alps, *Comput. Geosci.*,  
 1012 184, 105520, <https://doi.org/10.1016/j.cageo.2024.105520>, 2024.
- 1013 Nyberg, L., Cedergren, A., De Herve, M. D. G., Gustavsson, J., Hassel, H., Tehler, H., and Wester, M.:  
 1014 What do we mean with integrated risk management? – A research profiling comparing DRM and  
 1015 other fields of risk management, *Int. J. Disaster Risk Reduct.*, 132, 105942,  
 1016 <https://doi.org/10.1016/j.ijdrr.2025.105942>, 2026.
- 1017 Occhiena, C. and Pirulli, M.: Analysis of Climatic Influences on Slope Microseismic Activity and  
 1018 Rockfalls: Case Study of the Matterhorn Peak (Northwestern Alps), *J. Geotech. Geoenvironmental*  
 1019 *Eng.*, 138, 1012–1021, [https://doi.org/10.1061/\(ASCE\)GT.1943-5606.0000662](https://doi.org/10.1061/(ASCE)GT.1943-5606.0000662), 2012.
- 1020 OECD: Progress towards an all-hazards approach to emergency preparedness and response, Nuclear  
 1021 Energy Organisation for Economic Cooperation and Development, Paris, 2018.
- 1022 Olivieri, L. and Bettanini, C.: Preliminary observation of Marmolada glacier collapse of July 2022 with  
 1023 space-based cameras, *Remote Sens. Lett.*, 14, 21–29,  
 1024 <https://doi.org/10.1080/2150704X.2022.2152754>, 2023.
- 1025 ONF-RTM: Revue annuelle des phénomènes d'origines glaciaire et périglaciaire de l'année 2024  
 1026 Massif des Ecrins - Grandes Rousses, 2025.
- 1027 Peissier, C. and Courtray, V.: Séminaire technique - Synthèse et conclusions : Gestion des risques  
 1028 d'origine glaciaire et périglaciaire - PARN, Ministère de l'Écologie, du Développement Durable, des  
 1029 Transports et du Logement - Direction Générale de la Prévention des Risques, Grenoble - World Trade  
 1030 Center, 2012.

- 1031 Pigeon, P.: Réflexions sur les notions et les méthodes en géographie des risques dits naturels, *Ann. Géographie*, 111, 452–470, <https://doi.org/10.3406/geo.2002.21624>, 2002.
- 1032
- 1033 Prudent-Richard, G., Gillet, M., Vengeon, J.-M., and Descotes-Genon, S.: Changements climatiques dans les Alpes : Impacts et risques naturels, *ClimCHAlp - Interreg III B Alpine Space*, 2008.
- 1034
- 1035 Ravanel, L.: Évolution géomorphologique de la haute montagne alpine dans le contexte actuel de réchauffement climatique, 113–124, <https://doi.org/10.3406/edyte.2009.1078>, 2009.
- 1036
- 1037 Ravanel, L. and Deline, P.: La face ouest des Drus (massif du Mont-Blanc) : évolution de l’instabilité d’une paroi rocheuse dans la haute montagne alpine depuis la fin du petit âge glaciaire, *Géomorphologie Relief Process. Environ.*, 14, 261–272, <https://doi.org/10.4000/geomorphologie.7444>, 2008.
- 1038
- 1039
- 1040
- 1041 Ravanel, L. and Deline, P.: Rockfall Hazard in the Mont Blanc Massif Increased by the Current Atmospheric Warming, in: *Engineering Geology for Society and Territory - Volume 1*, edited by: Lollino, G., Manconi, A., Clague, J., Shan, W., and Chiarle, M., Springer International Publishing, Cham, 425–428, [https://doi.org/10.1007/978-3-319-09300-0\\_81](https://doi.org/10.1007/978-3-319-09300-0_81), 2015.
- 1042
- 1043
- 1044
- 1045 Ravanel, L., Magnin, F., Gallach, X., and Deline, P.: Évolution des parois rocheuses gelées de haute montagne sous forçage climatique. *La Météorologie.*, 34–40, <https://doi.org/10.37053/lameteorologie-2020-0090>, 2020.
- 1046
- 1047
- 1048 Raymond, M., Wegmann, M., and Funk, M.: Inventar gefährlicher Gletscher in der Schweiz, Versuchsanstalt für Wasserbau Hydrologie und Glaziologie der Eidgenössischen Technischen Hochschule Zürich, 2003.
- 1049
- 1050
- 1051 Reghezza, M., Leone, F., and Vinet, F.: La vulnérabilité: un concept problématique., in: *La vulnérabilité des sociétés et des territoires face aux menaces naturelles - Analyses géographiques*, vol. 1, Montpellier III, 35–40, 2006.
- 1052
- 1053
- 1054 Ricciardi, T.: Mattmark, 30 August 1965: A catastrophe that changed Switzerland’s perception of Italian migrants, *Schweiz. Z. Für Gesch.*, 66, 401, 2016.
- 1055
- 1056 Richard, D.: Caractérisation et gestion des risques d’origine glaciaire, CEMAGREF - Unité de recherche ETNA, Grenoble, 2005.
- 1057
- 1058 Rickenmann, D. and Zimmermann, M.: The 1987 debris flows in Switzerland: documentation and analysis, *Geomorphology*, 8, 175–189, [https://doi.org/10.1016/0169-555X\(93\)90036-2](https://doi.org/10.1016/0169-555X(93)90036-2), 1993.
- 1059
- 1060 Rothenbühler, C.: GISALP - räumlich-zeitliche Modellierung der klimasensitiven Hochgebirgslandschaft des Oberengadins, 3, 2006.
- 1061
- 1062 Röthlisberger, H.: Eislawinen und Ausbrüche von Gletscherseen, *Gletsch. Klima - Glaciers Clim. Jahrb. Schweiz. Naturforschenden Ges. Wiss. Teil*, 170–212, 1981b.
- 1063
- 1064 RTM-ONF: Révision partielle du Plan de Prévention des Risques Naturels Prévisibles, ONF-RTM, Chamonix-Mont-Blanc, 2000.
- 1065
- 1066 Schindelegger, A.: Natural hazard risk governance - Report on the state of the Alps, Alpine Convention, 2019.
- 1067

- 1068 Schindelegger, A. and Kanonier, A.: Natural Hazard Risk Governance: Status Quo in the EUSALP  
1069 Region, EUSALP Action Group 8, Wien, 2019.
- 1070 Schneiderbauer, S. and Ehrlich, D.: Risk, hazard and people's vulnerability to natural hazards - a  
1071 review of definitions, concepts and data, European Commission, 2004.
- 1072 Gletschersee Grindelwald : Felssturz Schlossplatte: <http://www.gletschersee.ch/index.cfm/treeID/21>,  
1073 last access: 12 February 2025.
- 1074 Sirop, M., Gagliardini, O., Serbource, P., and Bouvier, P.: La prévention des risques du glacier de Tête  
1075 Rousse : une action pluri-acteurs, 27–29, 2022.
- 1076 Stoffel, M., Tiranti, D., and Huggel, C.: Climate change impacts on mass movements — Case studies  
1077 from the European Alps, *Sci. Total Environ.*, 493, 1255–1266,  
1078 <https://doi.org/10.1016/j.scitotenv.2014.02.102>, 2014.
- 1079 Tacnet, J. M., November, V., Richard, D., and Batton-Hubert, M.: Expertise, décision et incertitude :  
1080 jusqu'où une approche interdisciplinaire est-elle possible dans le cadre de la gestion intégrée des  
1081 risques naturels en montagne ?, in: *Conférence Outils Pour Décider Ensemble 2010 (OPDE)*, 18 p.,  
1082 2010.
- 1083 Tagarev, T., Papadopoulos, G. A., Hagenlocher, M., Sliuzas, R., Ischiwatari, M., and Gallego, E.:  
1084 Integrating the risk management cycle, in: *Science for disaster risk management 2020 :acting today,*  
1085 *protecting tomorrow.*, Publications Office, LU, <https://doi.org/10.2760/571085>, 2021.
- 1086 Troilo, F., Lodigiani, M., Nicora, M., Perret, P., Christille, J.-M., Calabrese, M., Salvemini, C. B., and  
1087 Sartor, S.: Monitoring Cryospheric Environment at a Regional Scale: Big Data from Sensor Networks  
1088 and Experimental AI Applications in the Framework of the Glarisk-cc FESR Project, 12, 2025.
- 1089 Valette, P. and Carozza, J.-M.: *Géohistoire de l'environnement et des paysages*, CNRS éditions, Paris,  
1090 2019.
- 1091 Vannier, C., Lefebvre, J., Longaretti, P.-Y., and Lavorel, S.: Patterns of landscape change in a rapidly  
1092 urbanizing mountain region, *Cybergeo*, <https://doi.org/10.4000/cybergeo.27800>, 2016.
- 1093 Vergnano, A., Oggeri, C., and Godio, A.: Geophysical–geotechnical methodology for assessing the  
1094 spatial distribution of glacio-lacustrine sediments: The case history of Lake Seracchi, *Earth Surf.*  
1095 *Process. Landf.*, 48, 1374–1397, <https://doi.org/10.1002/esp.5555>, 2023.
- 1096 Villeneuve, E., Pérès, F., and Geneste, L.: An approach to improve risk assessment using experience  
1097 feedback, in: *Proceedings of IFIP Doctoral Spring Workshop "Product and Asset Lifecycle*  
1098 *Management*, 1–5, 2010.
- 1099 Vincent, C.: Fluctuations des bilans de masse des glaciers des Alpes françaises depuis le début du 20<sup>e</sup>  
1100 siècle au regard des variations climatiques, *Houille Blanche*, 88, 20–24,  
1101 <https://doi.org/10.1051/lhb/2002100>, 2002.
- 1102 Vincent, C., Garambois, S., Thibert, E., Lefebvre, E., Le Meur, E., and Six, D.: Origin of the outburst  
1103 flood from Glacier de Tête Rousse in 1892 (Mont Blanc area, France), *J. Glaciol.*, 56, 688–698,  
1104 <https://doi.org/10.3189/002214310793146188>, 2010a.
- 1105 Vincent, C., Auclair, S., and Meur, E. L.: Outburst flood hazard for glacier-dammed Lac de Rochemelon,  
1106 France, *J. Glaciol.*, 56, 91–100, <https://doi.org/10.3189/002214310791190857>, 2010b.

- 1107 Vivian, R.: La catastrophe du glacier Allalin, *Rev. Géographie Alp.*, 54, 97–112,  
1108 <https://doi.org/10.3406/rga.1966.3248>, 1966.
- 1109 Walter, F., Amann, F., Kos, A., Kenner, R., Phillips, M., De Preux, A., Huss, M., Tognacca, C., Clinton, J.,  
1110 Diehl, T., and Bonanomi, Y.: Direct observations of a three million cubic meter rock-slope collapse with  
1111 almost immediate initiation of ensuing debris flows, *Geomorphology*, 351, 106933,  
1112 <https://doi.org/10.1016/j.geomorph.2019.106933>, 2019.
- 1113 Weber, S., Beutel, J., Dietze, M., Bast, A., Kenner, R., Phillips, M., Leinauer, J., Mühlbauer, S., Pfluger,  
1114 F., and Krautblatter, M.: Progressive destabilization of a freestanding rock pillar in permafrost on the  
1115 Matterhorn (Swiss Alps): Hydro-mechanical modeling and analysis, *Earth Surf. Dyn.*, 13, 1157–1179,  
1116 <https://doi.org/10.5194/esurf-13-1157-2025>, 2025.
- 1117 Wiegandt, E. and Lugon, R.: 3 Challenges of Living with Glaciers in the Swiss Alps, Past and Present, in:  
1118 *Darkening Peaks*, edited by: Orlove, B., Wiegandt, E., and Luckman, B. H., University of California  
1119 Press, 33–48, <https://doi.org/10.1525/9780520934245-004>, 2008.
- 1120 World Glacier Monitoring Service (WGMS): *Fluctuations of glaciers 1985-1990*, IAHS (ICSJ) - UNEP -  
1121 UNESCO, 1993.
- 1122