



Exploring new EarthCARE observations for evaluating Greenland clouds in RACMO2.4

Thirza N. Feenstra¹, Willem Jan van de Berg¹, Gerd-Jan van Zadelhoff², David P. Donovan², Christiaan T. van Dalum², and Michiel R. van den Broeke¹

¹Institute for Marine and Atmospheric Research Utrecht, Utrecht University, The Netherlands

²Royal Netherlands Meteorological Institute (KNMI), de Bilt, the Netherlands

Correspondence: Thirza N. Feenstra (t.n.feenstra@uu.nl)

Abstract. Clouds present one of the major challenges for polar climate modeling and significantly contribute to uncertainties in climate and ice sheet mass balance projections, as their radiative effect can strongly impact ice and snow melt. Therefore, a reliable representation of clouds in polar climate models is essential, yet the observations necessary for their evaluation remain sparse. The launch of the Earth Cloud, Aerosol, and Radiation Explorer (EarthCARE) satellite in May 2024 helps bridge this gap by offering cloud observations in unprecedented detail using multiple instruments. Here, we demonstrate the potential of using these novel observations to evaluate cloud representation over the Greenland ice sheet in the regional climate model RACMO (version 2.4p1). To this end, we show along-track comparisons of co-located RACMO cloud profiles with EarthCARE lidar and radar observations. We compare both lidar backscatter and radar reflectivity observations, as well as retrieved cloud properties, with simulated RACMO profiles for two selected case studies. These first results indicate that RACMO simulates low- and mid-altitude ice clouds and snowfall at the correct locations, but fails to capture thinner high-altitude clouds. Additionally, RACMO typically underestimates cloud ice and snow water content, in particular in precipitating systems, where RACMO underestimates snowfall rates. Regarding supercooled liquid and mixed-phase clouds, RACMO does not always reproduce these, especially when they are located at higher altitudes. These first comparisons highlight the potential for using EarthCARE observations to evaluate regional climate models and provide directions for further development of RACMO.

1 Introduction

Over the last decades, Greenland ice sheet (GrIS) surface melt has been increasing, resulting from rising air temperatures (Hanna et al., 2021). The subsequent runoff of meltwater from the GrIS is now one of the main contributors to contemporary sea level rise (van den Broeke et al., 2016; Fox-Kemper et al., 2021; Otosaka et al., 2023) and is expected to keep increasing under future global warming (Bamber et al., 2019; Goelzer et al., 2020). Therefore, understanding the polar climate and its contemporary changes is critical for accurate projections of ice sheet mass loss and subsequent sea level rise.

Polar regional climate models (RCMs; e.g. Fettweis et al., 2017; Langen et al., 2017; Skamarock et al., 2019; Belušić et al., 2020; van Dalum et al., 2024) provide estimates of historical and projections of future Greenland ice sheet climate and surface



mass balance (SMB: accumulation minus ablation). One of the largest uncertainties in SMB estimates from RCM simulations arises from the representation of the microphysical structure of clouds (Hofer et al., 2019). Clouds govern accumulation through snow- and rainfall but also influence surface energy processes by cooling through reflection of shortwave radiation and warming through trapping of longwave radiation (Van Tricht et al., 2016; Niwano et al., 2019). Since surface melt is determined by the surface energy balance, correctly representing clouds and their interaction with the surface in RCMs is necessary to obtain reliable SMB estimates.

Polar clouds provide challenges for (regional) climate models. At high latitudes, mixed-phase clouds, in which supercooled liquid droplets and ice crystals coexist despite sub-zero temperatures, are frequently observed (Curry et al., 1996; Shupe, 2011). Accurately modeling their phase partitioning is of great importance, as both the sign and strength of the cloud radiative effect strongly depend on the cloud phase and water content (Shupe et al., 2015; Van Tricht et al., 2016; Hofer et al., 2019). Cloud phase changes are influenced by many poorly understood and competing processes, making them highly sensitive to the choice of parameterizations in atmospheric models (Forbes and Ahlgrimm, 2014; Taylor et al., 2019). Consequently, the simulated cloud phase and water content can vary considerably between models, depending on the microphysical parameterizations used (Hofer et al., 2019; Taylor et al., 2019). Additionally, models often struggle to capture optically thin ice clouds, which are common in polar regions (Tjernström et al., 2008; Taylor et al., 2019). These modeling challenges have led to contrasting conclusions regarding the role of clouds on GrIS surface melt. For instance, Van Tricht et al. (2016) and Hofer et al. (2019) both find that clouds enhance GrIS surface melt. However, Van Tricht et al. (2016) attribute this to an equal contribution from ice and liquid clouds, whereas Hofer et al. (2019) link higher melt rates to a larger fraction of liquid clouds. In contrast to these two studies, Niwano et al. (2019) report that, although clouds can increase the total integrated GrIS surface melt, mass loss in the ablation area can be reduced due to a reduction in solar and latent heat. In line with this, Wang et al. (2019) show that clouds enhance surface melt in the accumulation zone, but decrease melt in the ablation zone. This two-sided effect of clouds on surface mass loss, combined with the uncertainty regarding the impact of cloud phase on the radiative effect, stresses the need for observational constraints on clouds and their microphysical properties in models to improve our understanding of the interaction between clouds and ice sheet surface mass loss.

Evaluating cloud microphysical representation in climate models is particularly challenging for polar regions, as ground-based observations are limited (Shupe et al., 2013). Satellites can provide cloud observations over a larger spatial domain, but at the cost of lower temporal resolution. Specifically for a spectral imager, which can provide observations of microphysical processes, observations are limited to daylight hours. This can provide challenges in polar regions, as daylight is limited during polar winter. Particularly useful were the CloudSat and CALIPSO (Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation) satellites, equipped with radar and lidar systems, respectively, which were part of the so-called A-train constellation. This formation allowed CloudSat and CALIPSO to observe approximately the same ground track near-simultaneously with other Earth-observing satellites such as Aqua and Aura, which provided complementary measurements of radiation and atmospheric composition. These satellites have provided valuable observations used in a limited number of polar climate model evaluations. Lacour et al. (2018) present an assessment of cloud representation in eight CMIP5 models using CALIPSO observations and find that most models underestimate the cloud cover as well as the amount of liquid and mixed-phase clouds



over the GrIS. This results in an underestimation of the longwave radiative warming effect compared to ground-based radiation measurements at the Summit station in the centre of the GrIS. Similarly, they find an underestimation of the summer cloud radiative effect at the Summit. Conversely, both van Kampenhout et al. (2020) and Lenaerts et al. (2020) find that the atmospheric component of the Earth system model CESM2 overestimates the liquid water path and underestimates the ice water path over the GrIS, compared to CloudSat-CALIPSO observations.

While the aforementioned studies evaluate climatologies of vertically integrated cloud properties from global models with CloudSat and CALIPSO climatologies, model results can also be co-located in space and time with the satellite observations. Co-located evaluation has been done by Sankaré et al. (2022), who use CloudSat-CALIPSO observations to assess the representation of thin ice clouds in winter within the Canadian Regional Climate Model (CRCM6). They find a slight underestimation of ice water content during January 2007 in the Arctic region, but their analysis did not include clouds in the liquid phase.

Unfortunately, the CloudSat and CALIPSO satellites left the A-train constellation in 2018 (Zou et al., 2020) and ceased operation in 2023 (Skorokhodov and Kuryanovich, 2025). However, in May 2024, the newly developed Earth Cloud, Aerosol, and Radiation Explorer (EarthCARE; Wehr et al., 2023) was launched, which will bring the next generation of observations of atmospheric processes and will provide 3D profiles of clouds and aerosols. EarthCARE not only extends the CloudSat and CALIPSO observational record but also marks a big step forward by delivering the first exactly co-located measurements of clouds, aerosols, and radiation from space at a higher horizontal resolution than ever before. Without a lag in observation time between the instruments, no assumptions regarding temporal evolution have to be made, yielding more accurate atmospheric profiles. By combining observations of the four different instruments, an atmospheric lidar, a cloud profiling radar, a multi-spectral imager, and a broadband radiometer, EarthCARE provides observations of the vertical structure of clouds, aerosols, and radiation in unprecedented detail. These novel synergistic observations will be used to evaluate weather and climate models and improve their parameterizations. This is particularly valuable for the polar regions, where models frequently show radiation biases that may be linked to cloud processes (van Wessem et al., 2014; Lacour et al., 2018; Souverijns et al., 2019; Inoue et al., 2021; van Dalum et al., 2024). In this context, these high-resolution EarthCARE observations will be a much-needed addition to the sparse in-situ observational dataset currently available.

This study introduces a methodology for co-located comparison of cloud profiles simulated by RCMs or obtained from reanalyses to EarthCARE satellite observations. Here, we consider two representative EarthCARE overpasses to demonstrate the potential of these data and methods for model evaluation by applying this to model simulations of the polar RCM RACMO version 2.4p1. In section 2, we introduce RACMO version 2.4p1, the EarthCARE satellite data, and the methodology to compare the model and satellite data. Section 3 describes the first case study, considering an overpass on March 12, 2025. The second case study, on May 13, 2025, is described in section 4. In sections 5 and 6, we discuss our results and conclude, respectively.



90 2 Method

2.1 RACMO version 2.4p1 model description

RACMO (Regional Atmospheric Climate Model version 2.4p1, henceforth: RACMO) is a hydrostatic regional climate model, consisting of the High Resolution Limited Area Model (HIRLAM, version 5.03) dynamical core and the physics module of the Integrated Forecasting System (IFS, cycle 47r1) of the European Centre for Medium-Range Weather Forecasts (ECMWF)
 95 (van Meijgaard et al., 2008; van Dalum et al., 2024). Here, we will provide a detailed description of the cloud (micro)physics in version 2.4p1. For a description of other model components and improvements with respect to RACMO version 2.3p3, see van Dalum et al. (2024).

The cloud scheme in RACMO, which is based on the ECMWF IFS cloud physics (Fig. 1; ECMWF, 2020), is a single-moment scheme. In RACMO version 2.4p1, the prognostic treatment of cloud fraction and water content for cloud ice, liquid,
 100 rain and snow was introduced, allowing for several pathways for water phase changes and precipitation generation. The prognostic treatment of ice and liquid water yields a more physically realistic representation of supercooled liquid water and mixed-phase clouds compared to a diagnostic approach (Forbes and Tompkins, 2011). Treating rain and snow prognostically allows for modeling the precipitation fall speed and the horizontal advection of precipitation. Supersaturation of ice at temperatures below -38°C is parameterized using a threshold for the relative humidity with respect to ice, following Kärcher and
 105 Lohmann (2002) and Tompkins et al. (2007). For temperatures above -38°C , supercooled liquid water can be present. In mixed-phase clouds, which are assumed to be well-mixed, this supercooled liquid water can be converted to ice crystals through the Wegener-Bergeron-Findeisen process (Storelvmo and Tan, 2015). The Wegener-Bergeron-Findeisen process largely depends on the saturation ratio and the ice crystal number concentration (Pruppacher and Klett, 1997; Rotsteyn et al., 2000). The ice crystal number concentration is treated diagnostically and depends only on temperature, following Meyers et al. (1992). An-
 110 other sink for liquid droplets is the collection and freezing by falling snow particles, known as riming, which depends on the fall velocity and diameter of snow particles and the liquid water content (Wilson and Ballard, 1999). Both the snow and ice sedimentation velocities are set to a constant, whereas for rain, it is dependent on the particle size distribution. The autoconversion of liquid water to rain follows Khairoutdinov and Kogan (2000), where the droplet number concentration is kept constant within a grid box, at 300 cm^{-3} over land and 50 cm^{-3} over ocean. Autoconversion of ice water to snow follows Sundqvist
 115 (1978) and Lin et al. (1983), considering a tuned critical threshold of ice water content for snow to be aggregated by collision of ice particles. Snow sublimation is based on the Kessler (1969) formulation, and depends on the saturation deficit with respect to ice. Melting of ice and snow hydrometeors can occur when the wet-bulb temperature exceeds the melting point. When raindrops become supercooled, they might freeze, with larger drops more likely to freeze than smaller drops (Bigg, 1953). In the cloud scheme, secondary ice processes are not included.

120 Radiative fluxes are computed separately from the cloud scheme. The ECMWF radiation scheme (ecRAD) that is embedded in RACMO consists of the Rapid Radiation Transfer Model: Long-Wave (RRTMLW; Mlawer et al., 1997) and Rapid Radiation Transfer Model: Short-Wave (RRTMSW; Clough et al., 2005). The radiation scheme takes the temperature, cloud fraction, content, and phase, and the albedo as input from the IFS physics modules. For aerosols, carbon dioxide, ozone, and trace



gases, climatologies are used. To compute the radiative effects of clouds, the McICA (Monte Carlo Independent Column
 125 Approximation) method (McRad;) is used, which takes the cloud fraction, content, phase, and effective radius into account.
 The cloud effective radius is computed in the radiation scheme, following Martin et al. (1994) and Wood (2000) for the liquid
 effective radius and Sun and Rikus (1999) for the ice effective radius. Since the radiative transfer code is computationally very
 expensive, the radiation code is called only once every model hour for all grid points, rather than at every time step.

When implementing IFS cycle 47r1, several cloud parameterizations were tuned to better match precipitation and melt
 130 patterns over the GrIS (van Dalum et al., 2024). This includes speeding up the Wegener-Bergeron-Findeisen process by a
 factor of 2.5, doubling the fall speed of snow to 2 m s^{-1} and lowering the critical autoconversion threshold for snow from
 $3 \cdot 10^{-5} \text{ kg kg}^{-1}$ to $5 \cdot 10^{-7} \text{ kg kg}^{-1}$.




Figure 1. Schematic of processes and interactions within the cloud scheme of the ECMWF Integrated Forecasting System as is used in cycle47r1 (ECMWF, 2020).

2.2 EarthCARE satellite observations

EarthCARE carries four complementary instruments to obtain a comprehensive profile of clouds, aerosols, and radiation (Wehr
 135 et al., 2023). The ATmospheric LIDar (ATLID) uses 355 nm UV light to measure aerosols and optically thin clouds. The ATLID
 is a high-spectral-resolution (HSRL) lidar, allowing for the separation of molecular (air density) and particulate (aerosols and
 clouds) backscatter, referred to as Rayleigh and Mie backscatter, respectively. The former allows for an independent observation
 of the extinction and backscatter, which was not possible using the CALIOP lidar. The HSRL capability also enhances ATLID's
 capability to detect and quantify optically thin clouds and fine aerosols compared to CALIPSO. Use of a particulate polarization
 140 channel provides information on particle shapes, enabling more accurate retrievals of ice particle properties and aerosol types.
 Up to 20 km, the ATLID vertical sampling resolution is 103 m, while above 20 km, it is 500 m. The along-track sampling
 distance is 140 m, but the onboard summing of every two profiles results in an effective resolution of 280 m. The ATLID is



especially well-suited for the detection of thin ice clouds and small liquid droplets, but cannot observe optically thick clouds and precipitation, as these cause the lidar signal to be fully attenuated. Because the ALTID detects liquid droplets well, the signal attenuates rapidly in liquid layers; it is mainly the liquid cloud top that is detected. Thick clouds and precipitation are measured by the Cloud Profiling Radar (CPR). The CPR is a 94 GHz W-band radar and can fully penetrate through clouds, complementing the ATLID observations. Because the radar has a high sensitivity towards the larger particles, it cannot adequately detect smaller ice crystals or water droplets in precipitating systems. The CPR is the first space-borne W-band radar with Doppler capacity, bringing information on convective motions and precipitation fall speeds, which results in improved drizzle, rainfall, and snowfall rate observations. Compared to the CloudSat radar, the CPR has an increased sensitivity of about 5 dB, allowing for the detection of smaller ice crystals and low-altitude clouds. With a footprint of 750 m, compared to CloudSat's 1.4–1.7 km, EarthCARE's CPR has a significantly higher spatial resolution. The vertical sampling of both radars is 500 m. However, because the CPR oversamples the radar echoes at 100 m, compared to 250 m for CloudSat, the vertical resolution of the retrieved cloud profiles is higher for the CPR. Additionally, this allows the CPR to detect clouds closer to the surface, compared to CloudSat. The Multispectral Imager (MSI) provides observations in the four visible and near-infrared and three infrared channels over a 150 km wide swath for scene context and additional cloud and aerosol information. The synergistic retrievals based on these three instruments will yield the most accurate 3D profiles of clouds and aerosols to date. From these, radiative fluxes can be modeled, which can be compared to the top of atmosphere fluxes measured by the Broadband Radiometer (BBR).

As EarthCARE was launched in May 2024, retrieval algorithms are currently (October 2025) still being tested and finalized. Hence, not all EarthCARE products have been released yet. Currently, Level 1b (calibrated satellite measurements) and Level 2a (derived cloud and aerosol properties) single-instrument products and a few Level 2b combined instrument products are available. More multi-instrument products will become available at the end of 2025. Therefore, we use several Level 1b and Level 2 ATLID and CPR products that have been available since January and March 2025. As we are primarily interested in cloud properties, this study focuses on the Level 1b ATL-NOM and CPR-NOM (Eisinger et al., 2024), the Level 2a ATL-ICE (Donovan et al., 2024) and CPR-CLD (Mroz et al., 2023), and the Level 2b AC-TC (Irbah et al., 2023) products.

2.3 Evaluation approach

2.3.1 RACMO simulation description

The RACMO simulation used here covers the Greenland domain, which consists of the GrIS, Svalbard, Iceland, parts of the Canadian Arctic, and the surrounding oceans, as in van Dalum et al. (2024), and is carried out on a 5.5 km grid. In the vertical direction, the domain consists of 40 hybrid sigma atmospheric layers. At the lateral boundaries, RACMO is forced with 3-hourly ERA5 data (Hersbach et al., 2020). Additionally, ERA5 data is used to describe sea surface temperature and sea ice concentration at the sea surface boundary. We apply upper-air relaxation, in which the modeled temperature, wind speed, and moisture in the upper atmospheric layers are nudged towards the ERA5 fields (van de Berg and Medley, 2016).



175 2.3.2 Co-location procedure

To compare RACMO model results with the EarthCARE satellite observations, we obtain co-located RACMO profiles below the EarthCARE overpasses by extracting the RACMO grid points closest to the satellite trajectory. To obtain a fair comparison between the RACMO model results and EarthCARE observations, the timestamp of the model output should be as close as possible to the overpass time of the satellite. To achieve this, we write additional RACMO cloud output fields at the model time
 180 step that is closest to the central EarthCARE overpass time, instead of relying on hourly or multi-hourly output. Depending on the maximum modeled atmospheric wind speeds and simulated month, RACMO uses a time step between one and five minutes for the Greenland domain on 5.5 km resolution. Considering the time the satellite needs to pass over the ice sheet, typically around ten minutes, the time stamp of the RACMO output will not be more than ten minutes off with respect to the satellite overpass, even for locations at the boundaries of the domain. Since cloud processes are relatively slow at high latitudes, we
 185 consider this to be close enough in time not to influence the analysis.

When comparing RACMO and EarthCARE directly, the RACMO resolution is used. This means that the EarthCARE observations, having a higher resolution, are aggregated onto the RACMO grid in both the horizontal (latitude-longitude) and vertical (atmospheric height) directions. This is done by taking the mean of all EarthCARE grid cells nearest to a co-located RACMO grid cell. For the cloud and precipitation classification, the cloud (ice, mostly ice, mixed-phase, liquid, mostly liquid,
 190 or no cloud) and precipitation (rain, snow, or no precipitation) class occurring most often within this set of nearest EarthCARE grid points is selected. Since RACMO uses hybrid sigma levels, the heights of the vertical layers vary within the domain, depending on the surface topography, temperature, and humidity. Therefore, the vertical coordinates of the RACMO profiles are not uniform over the trajectory, as are the regridded EarthCARE profiles.

2.3.3 ATLID and CPR simulators

195 We compare both Level 1b backscatter and reflectivity profiles and Level 2 derived cloud properties. To compare RACMO model output with Level 1 EarthCARE observations, we simulate lidar Mie and Rayleigh backscatter and radar reflectivity based on the RACMO output. To simulate lidar backscatter, we use the Cardinal Campaign Tools ATLID simulator (Donovan and de Kloe, 2025). This simulator takes as input profiles of clouds, aerosols, temperature, pressure, and wind. Backscatter profiles of cloud water, ice, rain, and snow, as well as aerosols, are computed using the multiscatter module. For cloud cate-
 200 gories, the effective radius as computed in the radiation scheme in RACMO is used as input. For the aerosol categories, a fixed effective radius is used per aerosol category. The lidar ratio, linear depolarization ratio, and asymmetry factor are assumed to be constant for each category, as reported in Wandinger et al. (2023). In RACMO, aerosols are represented by climatological fields of 11 aerosol types that were produced by the Copernicus Atmosphere Monitoring Service (CAMS) (Bozzo et al., 2020). However, as input for the ATLID simulator, we use the four categories described by the Hybrid End-To-End Aerosol Classifi-
 205 cation (HETEAC; Wandinger et al., 2023). To convert the RACMO aerosol climatology to the HETEAC aerosol types (fine mode - weakly absorbing, fine mode - strongly absorbing, coarse mode - spherical, coarse mode - non-spherical), we use the same aerosol mapping approach as in Qu et al. (2023) and Donovan et al. (2023). Since the ATLID simulator is designed to



be used on a regular grid, we use a vertical resolution of 100 m for the simulated backscatter profiles instead of the RACMO's non-uniform vertical levels. This vertical resolution is comparable to the resolution of the ATLID backscatter observations.

210 Therefore, note that the lidar backscatter profiles are shown at this 100 m vertical resolution.

We simulate radar reflectivity using relationships between radar reflectivity and water content and correct for attenuation from precipitation, liquid water, and atmospheric gases. We neglect attenuation from ice crystals, as this is small for W-band radars (Hogan and Illingworth, 1999). For ice and snow water content, we use the relationship derived for W-band radar based on Protat et al. (2007):

$$215 \quad Z = \left(\frac{\log_{10}(\text{IWC}) + 0.0023T + 0.84}{0.000491T + 0.0939} \right), \quad (1)$$

where the IWC is the ice and snow water content in g m^{-3} , T is the temperature in $^{\circ}\text{C}$, and Z is the radar reflectivity in dBZ. For liquid and rain water content, we use the formulation from Matrosov et al. (2004):

$$Z = \left(\frac{\text{LWC}}{2.4} \right)^2, \quad (2)$$

220 where the LWC is the liquid and rain water content in g m^{-3} , and Z the radar reflectivity in $\text{mm}^6 \text{m}^{-3}$. We correct for attenuation from snowfall using the following relationship from Matrosov (2007):

$$\alpha_{\text{snow}}(h) = 0.12S^{1.1}, \quad (3)$$

where α_{snow} is the one-way attenuation in dB km^{-1} and S is the snowfall rate in mm hr^{-1} , derived from the snow water content and sedimentation velocity, which is constant at 2 m s^{-1} in RACMO. Attenuation through rainfall is corrected for using the relationship from Matrosov et al. (2008):

$$225 \quad \alpha_{\text{rain}}(h) = \frac{R \cdot \rho_a^{0.45}}{1.32}, \quad (4)$$

with α_{rain} the one-way attenuation in dB km^{-1} , ρ_a the air density and R the rainfall rate in mm hr^{-1} , derived from the snow water content and sedimentation velocity. The sedimentation velocity for rain depends on the particle size distribution as described in ECMWF (2020).

Attenuation from liquid water, water vapor, and oxygen is computed using relationships derived by Matrosov et al. (2004):

$$230 \quad A_{\text{liquid}}(h) = 7.56 \cdot \text{LWP}(h) \cdot (1 + 0.012(293 - T)), \quad (5)$$

$$A_{\text{H}_2\text{O}} = 0.077 \cdot \text{WVP}(h) \left(\frac{P_0}{1013} \right) \left(\frac{293}{T_0} \right)^{1.5} \cdot [1 - \exp(-0.42h)] \quad (6)$$

$$A_{\text{O}_2}(h) = \left(\frac{P_0}{1013} \right)^2 \left(\frac{293}{T_0} \right)^2 \cdot [(7.02 \cdot 10^{-2}h) - (4.81 \cdot 10^{-3}h^2) + (1.22 \cdot 10^{-4}h^3)] \quad (7)$$

with A_{liquid} , $A_{\text{H}_2\text{O}}$ and $A_{\text{O}_2}(h)$ the two-way attenuation from liquid water, water vapor and oxygen in dB, $\text{LWP}(h)$ and $\text{WVP}(h)$ the integrated liquid water and water vapor path from the top of the atmosphere until height h , T the temperature in Kelvin, T_0 the near-surface temperature in Kelvin and P_0 the near-surface pressure in Kelvin.



2.3.4 Treatment of ATLID, CPR, and RACMO water content estimates

Both the ATLID (ATL-ICE) and CPR (CPR-CLD) provide an estimate of ice water content (IWC). Where the ATLID provides a more reliable estimate of high and thin ice clouds, the CPR can yield estimates of thicker clouds and precipitation. Currently, a combined ATLID - CPR IWC product has not been released yet. Therefore, to obtain a complete cloud ice profile, we combine the ATLID and CPR IWC estimates using a criterion based on the normalized uncertainty, as described in Cole et al. (2023). When a grid cell has only one instrument providing an IWC estimate, this IWC estimate is used directly for the ATLID-CPR composite. When both instruments provide an IWC estimate, the IWC is chosen based on the normalized uncertainty, defined as,

$$\sigma_{A-ICE} = \sqrt{\left(\frac{\sigma_{IWC}^{A-ICE}}{IWC^{A-ICE}}\right)^2 + \left(\frac{\sigma_{r_{eff}}^{A-ICE}}{r_{eff}^{A-ICE}}\right)^2} \text{ and} \quad (8)$$

$$\sigma_{C-CLD} = \sqrt{\left(\frac{\sigma_{IWC}^{C-CLD}}{IWC^{C-CLD}}\right)^2 + \left(\frac{\sigma_{r_{eff}}^{C-CLD}}{r_{eff}^{C-CLD}}\right)^2},$$

where r_{eff} is the ice crystal effective radius, and σ_{IWC}^{A-ICE} , $\sigma_{r_{eff}}^{A-ICE}$, σ_{IWC}^{C-CLD} and $\sigma_{r_{eff}}^{C-CLD}$ are the 1σ uncertainties of the retrievals. The IWC estimate associated with the lowest normalized uncertainty is taken for the composite profile. Since in observations, there is no clear distinction between ice cloud and snow particles, we consider the IWC as a bulk quantity representing both ice and snow (Mason et al., 2024). Consequently, for comparing the IWC between RACMO and EarthCARE, we add the snow water content to the RACMO IWC. For all shown water content profiles for both EarthCARE and RACMO, we exclude grid cells with a water content lower than $10^{-7} \text{ kg m}^{-3}$. Regarding liquid water, Mason et al. (2024) showed that the CPR-only liquid water retrievals are considerably less reliable than those based on multiple instruments. Since the latter are not yet publicly available, we rely on the combined ATLID-CPR target classification (AC-TC) to identify the presence of liquid water, but do not consider the liquid water content. For a direct comparison between the target classification and RACMO model results, we make a classification based on the RACMO water content. We distinguish between the cloud classes ice, liquid and mixed-phase and the precipitation classes rain and snow. For all classes, the water content threshold is $10^{-7} \text{ kg m}^{-3}$. A cloud is only considered to be mixed phase if at least 10 % of the water content is in the ice phase and at least 10 % is in the liquid phase. A grid box can be assigned no class, only a cloud or precipitation class, or both a cloud and a precipitation class.

2.4 Case selection criteria

We consider two cases from the period after March 11th, 2025, since on this date, EarthCARE Level 2 data became available. The chosen cases should be sufficiently spaced in time to consider observations of different atmospheric conditions. A large part of the satellite overpass should be over the GrIS, since we want to investigate both the cloud representation over the ocean and the ice sheet. We specifically consider cases where there is overlap in cloudiness between the modeled and observed cloud scenes, which allows us to compare the two. Since models struggle to represent mixed-phase clouds in particular, we only choose cases in which these are present. Additionally, we are interested in precipitating particles, and therefore consider cases



265 in which snowfall occurs. As rainfall is limited to the late spring and summer and early fall in this region, we chose a scene
in which rainfall occurs for the second case. Considering these criteria, the chosen cases are on March 12th, 2025, 19:48 UTC
(EarthCARE frame 4479C) and May 13th, 2025, 03:10 UTC (EarthCARE frames 5433B and 5433C). For the chosen cases,
we use EarthCARE data of baseline BA for the ATLID Level 1b and all Level 2 products. For the CPR Level 1b products,
baseline CA (March case) and baseline CB (May case) are used.

270 3 Case study: March 12th, 2025

3.1 Case description

Here, we analyze an EarthCARE satellite overpass on March 12th, 2025, with the closest RACMO timestamp corresponding
to the overpass time of 19:48 UTC (Fig. 2a). Along almost the entire eastern ice sheet margin and along the western margin at
latitudes above 70° N, sea ice covers almost the entire coastal seas. The large-scale atmospheric flow is towards the southeast
275 in the northern half of the domain (Fig. 2a). A high-pressure system is present in the southeast. The near-surface temperatures
(dotted lines in Fig. 2e) over the ice sheet are sub-zero. Temperatures are only above the freezing point over a small area in the
Baffin Bay.

EarthCARE passes over the GrIS from the northeast to the southwest. In the northeast, over the Arctic Ocean, between Sval-
bard and Greenland, it encounters a cloudy region where RACMO simulates a relatively low cloud water path. In contrast, over
280 northwest Greenland and Baffin Bay, thicker clouds are modeled. The northwest area of Greenland is cloud-free. Partitioning
this into the different cloud types that RACMO simulates (i.e., ice, liquid, snow, and rain), most of the cloud water is in the
solid phase (Fig. 2b,d). Small amounts of supercooled liquid water (Fig. 2c) coexist with ice crystals (Fig. 2b) as mixed-phase
clouds. RACMO simulates relatively large snowfall amounts over northwest Greenland and Baffin Bay (Fig. 2d). Resulting
from the low temperatures in early spring at this high latitude, precipitation falling as rain is limited (Fig. 2e).




Figure 2. Modeled cloud scene on March 12th, 2025, 19:48 UTC. **(a)** Total cloud water path, vertically integrated [g m^{-2}], as simulated by RACMO. The thick black line shows the EarthCARE overpass. The contours of the 500 hPa geopotential height [m] levels are shown in dashed black lines. The hatched area indicates the presence of sea ice (sea ice extent larger than 15%). **(b-e)** Cloud water content [kg m^{-3}] as simulated by RACMO, for the co-located satellite overpass shown in **(a)**, for **(b)** cloud ice, **(c)** cloud liquid water, **(d)** cloud snow and **(e)** cloud rain. The dotted lines indicate the -50°C to 0°C temperature isotherms. Note that in **(b-e)** the x-axis follows the time coordinates. Hence, the latitude and longitude coordinates do not vary monotonically. In **(b-e)**, black areas correspond to the topography.

285 3.2 Comparison of simulated and observed Level 1 profiles

Based on the modeled profiles shown in Fig. 2a-d, we simulate Mie (Fig. 3b) and Rayleigh (Fig. 3d) attenuated backscatter and radar reflectivity profiles (Fig. 3f) to compare against the ATLID (Fig. 3a,c) and CPR (Fig. 3e) observations. In the absence of clouds or dense aerosol load, the ATLID Mie attenuated backscatter observations only show speckle, indicated in yellow and blue. High aerosol concentrations or optically very thin clouds appear in yellow to red, while thick ice clouds are shown in grey to white. Layers of liquid water show up as a clear white band. Below highly reflective layers, the ATLID signal is fully attenuated. There, the Mie attenuated backscatter signal indicates speckle again. In the Rayleigh channel, low attenuated



backscatter corresponds to full attenuation, which is shown in purple. In these areas, extinction and water content profiles can not be retrieved from the ATLID alone.

Considering the Mie attenuated backscatter profiles (Fig. 3a-b), RACMO captures a large part of the cloud structures over the ice sheet (black). In the southwest, RACMO misses high-altitude clouds. In this area, RACMO also shows a thicker and wider supercooled layer than can be identified from the ATLID observations. Contrastingly, it fails to capture several mid-altitude liquid layers near the western margin of the ice sheet, as well as a pronounced liquid layer detected by the lidar around 3 km altitude in the eastern part of the GrIS. However, RACMO does capture the lower-altitude supercooled layers observed by ATLID along the western margin of the ice sheet. Since RACMO uses a climatology instead of interactively simulating aerosols, it does not show strong backscatter signals from aerosol presence. However, some light blue areas appear, reflecting the smoothed aerosol field of the climatological input. Because aerosol loads in the Arctic are generally low (Hamilton et al., 2014), ATLID detects little aerosol as well, except for a small region in the northeast, corresponding to red and yellow areas below 2 km altitude. The yellow and light-blue speckling in Fig. 3a does not appear as strongly in the simulated profile, as it primarily represents measurement noise, which is not included in the ATLID simulator.

From Fig. 3a-b, it is clear that the ATLID signal rapidly extinguishes when it encounters thick ice clouds or liquid layers. These areas align with the low Rayleigh attenuated backscatter values in Fig. 3c-d. The top of these areas, therefore, indicates the top of a thick ice cloud or a liquid layer. For this overpass, both the Mie (Fig. 3a-b) and Rayleigh attenuated backscatter profiles (Fig. 3c-d) show that the top of the clouds over Baffin Bay, along the southwest of the overpass, are simulated at a lower height in RACMO. The height at which the signal is fully attenuated (purple) is located at lower altitudes in RACMO, and mainly corresponds to the supercooled liquid layers rather than the more extended ice cloud structures.

The previous findings can be confirmed by looking at the radar reflectivity (Fig. 3e-f). The CPR is not very sensitive to thin ice clouds, but captures larger and thicker cloud structures well (Fig. 3e). Looking again at the Baffin Bay area, RACMO underestimates the radar reflectivity (Fig. 3f), both in height and in the strength of the signal. This indicates that RACMO underestimates the cloud top and cloud water content of the clouds in this region. The CPR shows very high reflectivity values just above the surface due to surface backscatter (dark blue, Fig. 3e). This complicates the retrieval of near-surface clouds, particularly optically thin clouds such as fog. The blind zone of the CPR is, however, much smaller than the blind zone of CloudSat, which suffered from surface clutter in the lowest kilometer above the surface (Lamer et al., 2020).




Figure 3. Profiles of the March 12th, 2025, 19:48 UTC **(a,c,e)** observed (EarthCARE, **(a,c)** ATL-NOM, baseline BA and **(e)** CPR-NOM, baseline CA) and **(b,d,f)** modeled (RACMO) **(a-b)** Mie attenuated backscatter [$\text{sr}^{-1} \text{m}^{-1}$], **(c-d)** Rayleigh attenuated backscatter [$\text{sr}^{-1} \text{m}^{-1}$] and **(e-f)** radar reflectivity [dBZ]. Note that the x-axis follows the time coordinates. Hence, the latitude and longitude coordinates do not vary monotonically. Also note that in **(a-d)**, the vertical resolution is 100 m, while in **(e-f)** the vertical coordinates follow the RACMO hybrid-sigma levels. Black areas correspond to the topography.

3.3 Comparison of modeled and retrieved clouds and precipitation

The combined ATLID - CPR classification (AC-TC, Fig. 4a) provides information on cloud and precipitation types. In RACMO (Fig. 4b), clouds and snowfall often coexist. Since the CPR reflectivity is dominated by larger precipitation particles (Mason et al., 2024), in precipitating systems, it cannot be determined whether only snowflakes are present at a given location or



whether smaller cloud ice crystals are also present. As a result, snowfall in the AC-TC always co-occurs with cloud ice. This means that locations below a cloud where snow particles are precipitating (hatched regions with white background in Fig. 4b), will also be classified as cloud in the EarthCARE classification, which occurs over Baffin Bay. Here, we cannot determine whether the sole occurrence of snow, as modeled by RACMO, is correct or whether ice crystals are also present.

The classification confirms part of what can be identified in the backscatter and radar reflectivity profiles. Fig. 4 again shows that, for this overpass, RACMO did not capture the presence of high ice clouds over the Baffin Bay area. Over the ice sheet, RACMO generally agrees well with the observations in terms of ice cloud occurrence and snowfall, apart from missing some high-altitude ice clouds over the western part of the GrIS. Although the radar reflectivity profiles suggest that RACMO might underestimate snowfall, this is not evident from the classification. Despite missing some snowfall located at higher altitudes over Baffin Bay, RACMO captures most of the locations where snowfall is observed. Over the ice sheet, RACMO even models snowfall at higher altitudes than the EarthCARE observations indicate. This indicates that, rather than missing snowfall locations, the too low radar reflectivity might be explained by the snow water content being too low in RACMO. Furthermore, in RACMO, snow and ice almost always co-occur, whereas for EarthCARE, there are more locations where only ice is retrieved from the observations. This could be resulting from combining the ATLID and CPR classes, as for ATLID, there is no snowfall class because it cannot measure sedimentation velocities. Therefore, low snow water content might not be correctly classified in the ATLID-CPR classification. On the other hand, RACMO might generate snow too quickly when ice is present, which could lead to ice clouds dissipating too quickly, partly explaining the too low radar reflectivity (Fig. 3f).

Considering supercooled liquid and mixed-phase clouds, liquid and ice clouds typically coexist, with very few purely liquid clouds due to the low air temperatures during this season. For this overpass, RACMO shows fewer locations containing supercooled liquid water than EarthCARE, especially at mid-level altitudes. Near the ice sheet surface, however, there is relatively good agreement on the presence of liquid water, although RACMO produces mixed-phase layers that are too shallow. Over Baffin Bay, RACMO simulates a mixed-phase cloud around 2 km altitude that is not observed by the satellite. Contrastingly, the liquid water observed by EarthCARE at around 5 km height in this area is not captured by RACMO.




Figure 4. Cloud and precipitation classification for **(a)** EarthCARE (AC-TC, baseline BA) and **(b)** RACMO for March 12th, 2025, 19:48 UTC. Clouds are classified as ice (pink), liquid (blue), or mixed-phase (brown) clouds. Because the AC-TC is downsampled to the RACMO grid, some grid cells may fall between these categories and can then be classified as mostly ice (orange) or mostly liquid (green). The hatched areas indicate areas with snowfall or rainfall. Note that the x-axis follows the time coordinates. Hence, the latitude and longitude coordinates do not vary monotonically. Black areas correspond to the topography.

Both the ATLID (Fig. 5a) and the CPR (Fig. 5b) products provide an estimate of IWC. From Fig. 5a-b, it is evident that the ATLID and CPR provide complementary IWC profiles, with the ATLID effectively detecting the thinner, high-altitude clouds, and the CPR observing the lower and thicker clouds and precipitation. For a direct comparison with the RACMO IWC (Fig. 5e), we use the composite of the ATLID and CPR IWC profiles (Fig. 5d). Compared to the observations, RACMO performs well in simulating the mid-range IWC values, but does not fully capture the entire range of IWC values observed by EarthCARE. Specifically, RACMO fails to reproduce the highest IWC values. This is reflected in the IWC histograms (Fig. 5c), where RACMO shows fewer instances of high IWC and overestimates the number of locations with a mid-range IWC.




Figure 5. Ice water content [kg m^{-3}] from (a) ATLID (ATL-ICE, baseline BA), (b) CPR (CPR-CLD, baseline BA), (d) ATLID-CPR composite and (e) RACMO (including snow water content) for March 12th, 2025, 19:48 UTC. (c) shows the gridcell area-weighted histogram of ice water content for EarthCARE (pink, ATLID-CPR composite) and RACMO (blue, including snow water content). Note that in (a,b,d,e) the x-axis follows the time coordinates. Hence, the latitude and longitude coordinates do not vary monotonically. In (a,b,d,e), black areas correspond to the topography.

To evaluate snowfall estimates in RACMO, we use the precipitation retrievals from the CPR (Fig. 6a). The CPR can penetrate through precipitating clouds, allowing for precipitation retrievals, unlike the ATLID signals. The snowfall pattern in RACMO (Fig. 6b) agrees well with that observed over the GrIS, but is not fully captured over Baffin Bay, in line with what is indicated by the classification (Fig. 4). Towards the ice sheet interior, the near-surface snowfall rate is underestimated, as is the snowfall rate over Baffin Bay. Contrastingly, the snowfall rate directly at the western margin is overestimated in RACMO. This indicates that snow might precipitate fully out too quickly at the orographic barrier. This contrasting pattern indicates that the peak snowfall rates are not reached, and the snow likely precipitates out too fast. Hence, the width of the distribution of snowfall rates might be too small for this overpass. The snowfall rate depends on both the amount of snow generated by clouds and the sedimentation velocities of snow particles. In RACMO, a fixed sedimentation velocity of 2 m s^{-1} is applied, which lies at the upper end of the range observed by the radar (Fig. 6c). The lower sedimentation velocities observed by the CPR might



have resulted in snow particles remaining suspended in the atmosphere for a longer period than in the RACMO simulation. This could partly explain the discrepancies between RACMO and the observations in terms of snowfall and IWC (which also includes snow water content, Fig. 5).




Figure 6. (a-b) Snowfall rate [$\text{kg m}^{-2} \text{s}^{-1}$] from (a) CPR-CLD (baseline BA) and (b) RACMO for March 12th, 2025, 19:48 UTC. The RACMO snowfall rate is obtained by multiplying the snow water content by the sedimentation velocity. (c) Sedimentation velocity [m s^{-1}] from CPR-CLD. Note that the x-axis follows the time coordinates. Hence, the latitude and longitude coordinates do not vary monotonically. Black areas correspond to the topography.

365 4 Case study: May 13th, 2025

4.1 Case description

The second EarthCARE satellite overpass we consider is on May 13th, 2025, with the closest RACMO timestamp corresponding to the overpass time of 03:10 UTC (Fig. 7a). The sea ice cover during this period is similar to the March case, as the sea ice melting season has only just started. A high-pressure system is located south of Iceland. Thick clouds are present over the Atlantic Ocean, south of the GrIS, indicated by the large cloud water path. Moisture and clouds are transported from the Atlantic towards the south of the ice sheet and Baffin Bay. The Irminger Sea area is dominated by temperatures above the freezing point (dotted lines in Fig. 7e), while over the ice sheet, the near-surface temperatures are sub-zero.

This time, EarthCARE passes over the Greenland area from the southeast to the northwest. In RACMO, clouds are simulated over the southern half of the ice sheet and in its northernmost part. EarthCARE first encounters thick clouds over the Irminger Sea, then passes over the cloudy southern half of the ice sheet. After crossing the ice sheet's cloud-free region, it encounters another cloudy area over Ellesmere Island. Although most of the cloud water is again in the solid phase (Fig. 7b,d), RACMO simulates more water in the liquid phase compared to the March case (Fig. 7c). These liquid clouds sometimes coexist with



ice clouds. The higher temperatures over the Irminger Sea in May also lead to the occurrence of purely liquid clouds, and even with rainfall in a small region. Over the Irminger Sea, winds are mainly directed northward, approximately following the EarthCARE flight line. Hence, around 63° - 65° N, some of the generated snowfall (Fig. 7d) is advected onto the ice sheet.




Figure 7. Modelled cloud scene on May 13th, 2025, 03:10 UTC. **(a)** Total cloud water path, vertically integrated [g m^{-2}], as simulated by RACMO. The thick black line shows the EarthCARE overpass. The contours of the 500 hPa geopotential height [m] levels are shown in dashed black lines. The hatched area indicates the presence of sea ice (sea ice extent larger than 15%). **(b-e)** Cloud water content [kg m^{-3}] as simulated by RACMO, for the co-located satellite overpass shown in **(a)**, for **(b)** cloud ice, **(c)** cloud liquid water, **(d)** cloud snow and **(e)** cloud rain. The dotted lines indicate the -50°C to 0°C temperature isotherms. Note that in **(b-e)** the x-axis follows the time coordinates. Hence, the latitude and longitude coordinates do not vary monotonically. In **(b-e)**, black areas correspond to the topography.

4.2 Comparison of simulated and observed Level 1 profiles

The simulated ATLID backscatter profiles (Fig. 8b,d) indicate that RACMO captures the location of clouds reasonably well compared to the ATLID observations (Fig. 8a,c). Differences appear around 62° N, where ATLID detects high Mie attenuated backscatter values and is fully attenuated in the Rayleigh channel, indicating an optically thick ice cloud or snowfall, which RACMO does not reproduce. Over Ellesmere Island, RACMO simulates a larger area with high Mie attenuated backscatter



values, but appears to miss the rapid transition to full attenuation around 81.5° N, which suggests the presence of a supercooled liquid layer. The thin white band in Fig. 8a and the thin purple band in Fig. 8c over the non-cloudy part of the ice sheet represent the surface backscatter, as the signal has not reached full attenuation before reaching the surface yet. This does not show up in the modeled backscatter profiles (Fig. 8b,d), as surface scatter is not modeled in the lidar simulator. Over the Irminger Sea, a large red to yellow area is present in the RACMO profiles. This results from the high sea salt aerosol concentrations in the North Atlantic at this time of year in the CAMS aerosol climatology (Bozzo et al., 2020), which is consistent with in-situ observations (Saliba et al., 2019). However, in the EarthCARE observations, there is only limited backscatter resulting from aerosols in this region, indicating that there might be high variability in sea salt aerosol generation, which is not captured when using a climatology.

Considering the radar reflectivity (Fig. 8e-f), RACMO simulates cloudy regions at roughly the same locations as the CPR observations, but underestimates their reflectivity, which points to simulated ice and snow water contents that are too low. In line with the ATLID observations, RACMO misses the cloudy region around 62° N, as well as some thin clouds at the eastern margin.




Figure 8. Profiles of the May 13th, 2025, 03:10 UTC (a,c,e) observed (EarthCARE, (a,c) ATL-NOM, baseline BA and (e) CPR-NOM, baseline CB) and (b,d,f) modeled (RACMO) (a-b) Mie attenuated backscatter [$\text{sr}^{-1}\text{m}^{-1}$], (c-d) Rayleigh attenuated backscatter [$\text{sr}^{-1}\text{m}^{-1}$] and (e-f) radar reflectivity [dBZ]. Note that the x-axis follows the time coordinates. Hence, the latitude and longitude coordinates do not vary monotonically. Also note that in (a-d), the vertical resolution is 100 m, while in (e-f) the vertical coordinates follow the RACMO hybrid-sigma levels. Black areas correspond to the topography.

4.3 Comparison of modeled and retrieved clouds and precipitation

400 The cloud and precipitation classification (Fig. 9) indicates reasonably good agreement between the modeled and observed precipitation. The EarthCARE observations confirm the rainfall simulated by RACMO over the Irminger Sea. In terms of snowfall, RACMO captures most of the observed snowfall patterns, but simulates more snowfall at higher altitudes over the



ice sheet, as, again, ice and snow almost always co-occur. Along the eastern margin of the ice sheet, the model results and observations agree on the presence of liquid water, though not always on the altitude of the detected mixed-phase layers.

405 The precipitating low-altitude purely liquid cloud does not appear in the EarthCARE classification, likely because the CPR observations are obscured by surface clutter and attenuation by precipitation. Considering the detected rainfall in this area, it is likely that a liquid cloud is present here. Contrastingly, the observed liquid layers over the ice sheet interior and Ellesmere Island are not reproduced by RACMO. Regarding ice clouds, most of the ice clouds simulated by RACMO are also observed by EarthCARE. However, RACMO misses some clouds over the southern part of the Irminger Sea and the western part of the

410 GrIS. Looking at the ATLID Mie attenuated backscatter (Fig. 8a) and CPR radar reflectivity (Fig. 8e) profiles, these missing clouds are likely optically very thin. This is indicated by the very low backscatter and reflectivity values in these regions, which EarthCARE can observe because of the increased sensitivity of the lidar and radar instruments.




Figure 9. Cloud and precipitation classification for (a) EarthCARE (AC-TC, baseline BA) and (b) RACMO for May 13th, 2025, 03:10 UTC. Clouds are classified as ice (pink), liquid (blue), or mixed-phase (brown) clouds. Because the AC-TC is downsampled to the RACMO grid, some grid cells may fall between these categories and can then be classified as mostly ice (orange) or mostly liquid (green). The hatched areas indicate areas with snowfall or rainfall. Note that the x-axis follows the time coordinates. Hence, the latitude and longitude coordinates do not vary monotonically. Black areas correspond to the topography.

The observed IWC profile (Fig. 10a) confirms that the clouds over the southern part of the Irminger Sea and the western part of the GrIS are very thin, with concentrations well below $10^{-5} \text{ kg m}^{-3}$. In line with the March case, RACMO reproduces

415 locations with mid-range IWC values, but does not simulate sufficiently high IWC values at locations with snowfall, which are mainly located over the oceans. The IWC histograms (Fig. 10c) confirm this, showing too many grid cells with mid-range IWC values, whereas concentrations larger than $10^{-4} \text{ kg m}^{-3}$ hardly ever occur, showing a similar, but more extreme pattern than the March case.




Figure 10. Ice water content [kg m⁻³] from (a) ATLID-CPR (from ATL-ICE and CPR-CLD, baseline BA) composite and (b) RACMO (including snow water content) for May 13th, 2025, 03:10 UTC. (c) shows the gridcell area-weighted histogram of ice water content for EarthCARE (pink, ATLID-CPR composite) and RACMO (blue, including snow water content). Note that in (a-b) the x-axis follows the time coordinates. Hence, the latitude and longitude coordinates do not vary monotonically. In (a-b), black areas correspond to the topography.

Considering snowfall specifically, there is relatively good agreement on where snowfall occurs (Fig. 11a-b). However, the
 420 CPR detects very high snowfall rates, which RACMO does not reproduce for this overpass. Even though snowfall rates are
 highest directly at the eastern margin, similar to the March case, snowfall rates are not overestimated, and are too low in almost
 all locations. As in the March case, the sedimentation velocities obtained from the CPR (Fig. 11c) are much lower than the
 fixed sedimentation velocity of 2 m s⁻¹ in RACMO, which might explain part of the differences. It should be noted, however,
 that for locations with an order of magnitude difference in snowfall rate, e.g., around 66° N, even sedimentation velocities
 425 overestimated by a factor of two or three likely cannot explain all differences. Furthermore, RACMO simulates higher-altitude
 locations where low snowfall rates occur, which are not detected by the CPR. This is likely because the CPR is less sensitive
 to smaller particles. Mason et al. (2024) showed that CPR-only retrievals tend to miss lower snowfall rates, whereas multi-
 instrument synergistic retrievals, which include ATLID observations, can capture these due to the lidar's higher sensitivity to
 small particles. As for rain, RACMO simulates a larger area with rainfall than the CPR detects, but with similar rainfall rates
 430 (Fig. 11d-e). Because of the proximity to the surface, the CPR rainfall rates might be affected by surface clutter or attenuation,
 as can be seen in the radar reflectivity profile (Fig. 8e).




Figure 11. (a-b) Snowfall rate [$\text{kg m}^{-2} \text{s}^{-1}$] from (a) CPR-CLD (baseline BA) and (b) RACMO for May 13th, 2025, 03:10 UTC. (c) Sedimentation velocity [m s^{-1}] from CPR-CLD. (d-e) Rainfall rate [$\text{kg m}^{-2} \text{s}^{-1}$] from (d) CPR-CLD and (e) RACMO. The snowfall and rainfall rates are obtained by multiplying the snow and rain water content by the sedimentation velocity. Note that the x-axis follows the time coordinates. Hence, the latitude and longitude coordinates do not vary monotonically. Black areas correspond to the topography.

5 Discussion

This first EarthCARE-based evaluation of modeled polar clouds underlines the value of using satellite observations to evaluate regional climate models. Considering the recent launch of the EarthCARE satellite, this is the first study using these novel data for the evaluation of a polar climate model. Previous studies evaluating clouds over the GrIS in global models using CloudSat and CALIPSO data have reported contradictory results regarding ice and liquid cloud representation. Our results are most consistent with Lacour et al. (2018), indicating an underestimation of Greenland clouds in both phases. While our two case studies do not agree with the overestimated liquid water path, they are in line with the underestimation of ice clouds in CESM2, as found by van Kampenhout et al. (2020) and Lenaerts et al. (2020). Compared to these studies, which rely on climatologies of integrated quantities, our case study approach is more similar to that of Sankaré et al. (2022). Their focus, however, is on thin ice clouds during winter, whereas we include different types of clouds during early and late spring. Compared to CloudSat-



CALIPSO observations, they find an underestimation of ice water content in the Canadian Regional Climate Model (CRCM6) over the whole Arctic region, which is consistent with our results, although less pronounced than in our case. They attribute the underestimation of near-surface clouds to the modeled clouds precipitating out faster than the observed clouds. This agrees with the overly high snow sedimentation velocities we find with respect to the observations.

The underestimation of cloud water content likely contributes to the underestimation of the longwave downward flux in RACMO reported by van Dalum et al. (2024), since the cloud radiative effect is strongly correlated with the liquid and ice water path. Moreover, the longwave warming response typically dominates the shortwave cooling (Van Tricht et al., 2016). RACMO is not the only polar RCM in which such a radiation bias is found. Souverijns et al. (2019) report a similarly sized bias in the longwave downward flux in the COSMO-CLM² model over the Antarctic ice sheet, which they attribute to a lack of liquid clouds. Likewise, Inoue et al. (2021) find an underestimation of downward longwave radiation in the RCMs CAFS, CCLM, HIRHAM, MAR, and METUM over the sea ice-free Arctic, while WRF shows an overestimation. These biases are attributed, amongst others, to discrepancies in the partitioning between solid and liquid clouds, underestimated cloud occurrence, and excessive snowfall. Additionally, van Dalum et al. (2024) find an overestimation of precipitation over the GrIS compared to weather station data along the ice sheet margins, which might also be related to snow particles precipitating out quickly upon landfall over the ice sheet, like in our March case. Similarly, Ryan et al. (2020) find an overestimation of snowfall along the GrIS margins in a previous version of RACMO (version 2.3p2) and the RCM MAR compared to CloudSat snowfall rate retrievals. Together, these findings stress the importance of accurate precipitation and cloud microphysical representation in polar RCMs to obtain reliable surface melt and surface mass balance estimates.

Considering these first findings, we can work towards improved cloud microphysical representation in RACMO. These case studies suggest that some of our previous tuning choices should be reconsidered, such as the doubling of the snow sedimentation velocity, which now appears overestimated. Additionally, the process of conversion of ice to precipitating snow might also be overestimated, leading to overly rapid snow particle generation, resulting in ice clouds dissipating too quickly. Currently, the persistence of supercooled liquid layers might likely be suppressed by a too strong Wegener-Bergeron-Findeisen process, which converts too much liquid water into ice crystals. However, tuning this process will most likely result in lower simulated IWC, which is presumably already too low. This suggests that parameterizations describing the formation of clouds, such as supersaturation and ice crystal nucleation, should be considered as well. An improved cloud representation, achieved through tuning of these processes, will change the melt and precipitation patterns over the ice sheet, and consequently the SMB. Since the SMB is one of the key parameters of interest in polar RCMs, it provides an additional constraint during model tuning, as it should match the observed SMB as closely as possible.

While the ATLID and CPR retrievals used in this study provide high-resolution estimates of cloud properties, both instruments have their limitations. Although ATLID and CPR can provide a largely complete profile of the atmospheric structure, observations close to the surface will suffer from multiple scattering and surface clutter, making these observations less reliable. Therefore, fog layers and blowing snow are hard to detect. Furthermore, snowfall estimates relying only on CPR retrievals will be missing lower snowfall rates, as the CPR is not sensitive enough to observe the smaller ice crystals (Mason et al., 2024). At the same time, attenuation due to heavy precipitation can complicate the CPR retrievals. Regarding liquid clouds, we rely



mainly on the ATLID, as the CPR struggles to detect liquid layers (Mason et al., 2024). However, when liquid layers are located below thicker cirrus clouds, the ATLID will not be able to detect these liquid layers, as the signal will become fully attenuated because of the higher-altitude clouds. This is also the case for lower-located thin clouds, which might not be detected by the CPR, but cannot be observed by the ATLID when higher-altitude clouds are present. These limitations should be taken into account, especially when working with EarthCARE's single-instrument products. The use of lidar and radar simulators to be able to compare not only retrieved cloud profiles but also the observed backscatter and reflectivity profiles is therefore a valuable addition to the analysis.

As this analysis is based on some of the first available EarthCARE observations, calibration and validation efforts are still ongoing. This not only implies that newer, more reliable baselines of the EarthCARE products used in this study will become available, but also that additional multi-instrument synergistic products will be released by the end of 2025. These synergistic multi-instrument retrievals can partly overcome the limitations of each individual EarthCARE instrument and are, therefore, more reliable. An intercomparison of the EarthCARE cloud and precipitation retrieval products has shown that combining observations from the ATLID, CPR, and MSI yields the most accurate estimates of both the solid and liquid clouds and precipitation (Mason et al., 2024). Additionally, heating rates and radiation fluxes will be computed from the ATLID, CPR, and MSI retrieved cloud and aerosol profiles using radiative transfer modeling (Cole et al., 2023). Therefore, once available, these multi-instrument cloud and radiation products will be used to evaluate RACMO, using a similar methodology as described in this study. Although more reliable products will be available in the future, the current EarthCARE data already provide valuable observations of clouds, but their limitations should be considered when using these data for model evaluation. While these first case studies offer meaningful insights into cloud representation in RACMO, a more comprehensive evaluation based on multiple months of EarthCARE observations will be necessary for a reliable evaluation and will guide model development.

6 Conclusions

Using two selected case studies, we present the first comparison between EarthCARE observations and retrievals and simulated clouds and precipitation in a climate model, specifically the polar regional climate model RACMO. Our evaluation includes a comparison of simulated backscatter and reflectivity profiles against Level 1b ATLID and CPR observations, as well as an assessment of the modeled cloud and precipitation content and phase against the EarthCARE Level 2 cloud properties for two case studies. Comparing backscatter and reflectivity profiles provides a comprehensive overview of the atmospheric profile. These observations suggest that RACMO frequently misses thin, high-altitude clouds and likely underestimates the ice and snow water content, particularly in precipitating systems. These findings are supported by the ATLID and CPR water content retrievals and their combined classification. In particular, RACMO fails to reproduce the observed high snowfall rates, likely in part due to overly high sedimentation velocities in RACMO, causing snow particles to remain in the atmosphere for too short a time. Rainfall occurred only in the second case, where RACMO could reproduce the observed rainfall pattern. The EarthCARE observations provide information on the occurrence of supercooled liquid and mixed-phase clouds as well. These



reveal that RACMO captures most of the supercooled liquid layers near the surface, but struggles to simulate liquid clouds located at higher altitudes.

This study demonstrates the potential of using EarthCARE observations for evaluating regional climate models. Along-track comparisons provide insights into the vertical distribution of biases and the underlying processes, which is highly valuable for model development. EarthCARE passes over the GrIS around six times a day, providing sufficient spatial coverage for a comprehensive evaluation of clouds over the entire ice sheet and its surrounding oceans. In due time, additional EarthCARE multi-instrument products will become available, offering the best estimates of cloud and precipitation properties (Mason et al., 2024) as well as estimates of radiative fluxes and heating rates (Cole et al., 2023). These will, thereupon, be included for RACMO model evaluation and development. We expect that these observations will provide a reliable benchmark for evaluating and improving cloud microphysical processes, ultimately leading to improved cloud, radiation, and surface mass balance estimates in RACMO.

Code and data availability. The EarthCARE data can be downloaded from the ESA dissemination service. The frames used are 4479C for the March case and 5433B and 5433C for the May case. The products used are the ATL-NOM-1B product (baseline BA; European Space Agency, 2025a), the CPR-NOM-1B product (baseline CA and CB; Japan Aerospace Exploration Agency, 2025), the ATL-ICE-2A product (baseline BA; European Space Agency, 2025c), the CPR-CLD-2A product (baseline BA; European Space Agency, 2025d) and the AC-TC-2B product (baseline BA; European Space Agency, 2025b).

The RACMO model output and used EarthCARE data of the two timestamps can be downloaded from <https://doi.org/10.5281/zenodo.17590866> (Feenstra, 2025b).

The ATLID simulator can be accessed at <https://gitlab.com/KNMI-OSS/satellite-data-research-tools/cardinal-campaign-tools> (Donovan and de Kloe, 2025).

Software to compute co-located RACMO profiles, regrid EarthCARE data, simulate radar reflectivity and produce the figures shown in this manuscript can be found on <https://github.com/thirza-feenstra/EarthCARE4RCM> (Feenstra, 2025a).

In Fig. 2, Fig. 3e-f, Fig. 4, Fig. 5, Fig. 6, Fig. 7, Fig. 8e-f, Fig. 9, Fig. 10 and Fig. 11 the colormap Batlow is used (Crameri, 2023).

Author contributions. TF and WJvdB conceptualized the study. TF carried out the RACMO simulations, wrote the co-location software, and analysed the results. GJvZ and DD provided guidance on the use and interpretation of the EarthCARE observations and the simulation of backscatter and reflectivity profiles. CvD assisted with the use of RACMO2.4. TF wrote the manuscript with input from all authors.

Competing interests. DD is on the editorial board of the special issue: Early results from EarthCARE (AMT/ACP/GMD inter-journal SI).



Acknowledgements. We acknowledge the ECMWF for storage facilities and computational time on their supercomputer. We thank Shannon Mason for his input on the use of the different EarthCARE cloud and precipitation products.



References

- Bamber, J. L., Oppenheimer, M., Kopp, R. E., Aspinall, W. P., and Cooke, R. M.: Ice sheet contributions to future sea-level rise from structured expert judgment, *Proceedings of the National Academy of Sciences*, 116, 11 195–11 200, <https://doi.org/10.1073/pnas.1817205116>, 2019.
- 540 Belušić, D., De Vries, H., Dobler, A., Landgren, O., Lind, P., Lindstedt, D., Pedersen, R. A., Sánchez-Perrino, J. C., Toivonen, E., Van Ulft, B., Wang, F., Andrae, U., Batrak, Y., Kjellström, E., Lenderink, G., Nikulin, G., Pietikäinen, J.-P., Rodríguez-Camino, E., Samuelsson, P., Van Meijgaard, E., and Wu, M.: HCLIM38: a flexible regional climate model applicable for different climate zones from coarse to convection-permitting scales, *Geoscientific Model Development*, 13, 1311–1333, <https://doi.org/10.5194/gmd-13-1311-2020>, 2020.
- 545 Bigg, E. K.: The Supercooling of Water, *Proceedings of the Physical Society. Section B*, 66, 688–694, <https://doi.org/10.1088/0370-1301/66/8/309>, 1953.
- Bozzo, A., Benedetti, A., Flemming, J., Kipling, Z., and Rémy, S.: An aerosol climatology for global models based on the tropospheric aerosol scheme in the Integrated Forecasting System of ECMWF, *Geoscientific Model Development*, 13, 1007–1034, <https://doi.org/10.5194/gmd-13-1007-2020>, 2020.
- 550 Clough, S., Shephard, M., Mlawer, E., Delamere, J., Iacono, M., Cady-Pereira, K., Boukabara, S., and Brown, P.: Atmospheric radiative transfer modeling: a summary of the AER codes, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 91, 233–244, <https://doi.org/10.1016/j.jqsrt.2004.05.058>, 2005.
- Cole, J. N. S., Barker, H. W., Qu, Z., Villefranque, N., and Shephard, M. W.: Broadband radiative quantities for the EarthCARE mission: the ACM-COM and ACM-RT products, *Atmospheric Measurement Techniques*, 16, 4271–4288, <https://doi.org/10.5194/amt-16-4271-2023>, 2023.
- 555 Cramer, F.: Scientific colour maps, <https://doi.org/10.5281/zenodo.8409685>, 2023.
- Curry, J. A., Schramm, J. L., Rossow, W. B., and Randall, D.: Overview of Arctic Cloud and Radiation Characteristics, *Journal of Climate*, p. 1731–1764, 1996.
- Donovan, D. and de Kloe, J.: CARDINAL campaign tools ATLID simulator [software], <https://gitlab.com/KNMI-OSS/satellite-data-research-tools/cardinal-campaign-tools>, 2025.
- 560 Donovan, D. P., Kollias, P., Velázquez Blázquez, A., and van Zadelhoff, G.-J.: The generation of EarthCARE L1 test data sets using atmospheric model data sets, *Atmospheric Measurement Techniques*, 16, 5327–5356, <https://doi.org/10.5194/amt-16-5327-2023>, publisher: Copernicus GmbH, 2023.
- Donovan, D. P., van Zadelhoff, G.-J., and Wang, P.: The EarthCARE lidar cloud and aerosol profile processor (A-PRO): the A-AER, A-EBD, A-TC, and A-ICE products, *Atmospheric Measurement Techniques*, 17, 5301–5340, <https://doi.org/10.5194/amt-17-5301-2024>, 2024.
- 565 ECMWF: IFS documentation Cy47r1, Part IV: Physical Processes., <https://www.ecmwf.int/en/elibrary/81189-ifs-documentation-cy47r1-part-iv-physical-processes>, 2020.
- Eisinger, M., Marnas, F., Wallace, K., Kubota, T., Tomiyama, N., Ohno, Y., Tanaka, T., Tomita, E., Wehr, T., and Bernaerts, D.: The EarthCARE mission: science data processing chain overview, *Atmospheric Measurement Techniques*, 17, 839–862, <https://doi.org/10.5194/amt-17-839-2024>, 2024.
- 570 European Space Agency: EarthCARE ATLID NOM Level 1B, version BA, <https://doi.org/10.57780/ECA-BAD0EF3>, 2025a.
- European Space Agency: EarthCARE ATLID CPR TC Level 2B, version BA, <https://doi.org/10.57780/eca-7ba3052>, 2025b.
- European Space Agency: EarthCARE ATLID ICE Level 2A, version BA, <https://doi.org/10.57780/eca-e25465f>, 2025c.
- European Space Agency: EarthCARE CPR CLD Level 2A, version BA, <https://doi.org/10.57780/eca-7d84adf>, 2025d.



- 575 Feenstra, T.: EarthCARE4RCM [software], <https://github.com/thirza-feenstra/EarthCARE4RCM>, 2025a.
- Feenstra, T.: Dataset for "Exploring new EarthCARE observations for evaluating Greenland clouds in RACMO2.4" (Versie 1) [Data set], <https://doi.org/10.5281/zenodo.17590866>, 2025b.
- Fettweis, X., Box, J. E., Agosta, C., Amory, C., Kittel, C., Lang, C., van As, D., Machguth, H., and Gallée, H.: Reconstructions of the 1900–2015 Greenland ice sheet surface mass balance using the regional climate MAR model, *The Cryosphere*, 11, 1015–1033, <https://doi.org/10.5194/tc-11-1015-2017>, 2017.
- 580 Forbes, R. and Tompkins, A.: An improved representation of cloud and precipitation, <https://doi.org/10.21957/NFGULZHE>, publisher: ECMWF, 2011.
- Forbes, R. M. and Ahlgrimm, M.: On the Representation of High-Latitude Boundary Layer Mixed-Phase Cloud in the ECMWF Global Model, *Monthly Weather Review*, 142, 3425–3445, <https://doi.org/10.1175/MWR-D-13-00325.1>, 2014.
- 585 Fox-Kemper, B., Hewitt, H., Xiao, C., Aðalgeirsdóttir, G., Drijfhout, S., Edwards, T., Golledge, N., Hemer, M., Kopp, R., , Krinner, G., Mix, A., Notz, D., Nowicki, S., Nurhati, I., Ruiz, L., Sallée, J.-B., Slangen, A., , and Yu, Y.: Ocean, Cryosphere and Sea Level Change. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)], Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1211–1362, <https://doi.org/10.1017/9781009157896.011>, 2021.
- 590 Goelzer, H., Nowicki, S., Payne, A., Larour, E., Seroussi, H., Lipscomb, W. H., Gregory, J., Abe-Ouchi, A., Shepherd, A., Simon, E., Agosta, C., Alexander, P., Aschwanden, A., Barthel, A., Calov, R., Chambers, C., Choi, Y., Cuzzzone, J., Dumas, C., Edwards, T., Felikson, D., Fettweis, X., Golledge, N. R., Greve, R., Humbert, A., Huybrechts, P., Le clec'h, S., Lee, V., Leguy, G., Little, C., Lowry, D. P., Morlighem, M., Nias, I., Quiquet, A., Rückamp, M., Schlegel, N.-J., Slater, D. A., Smith, R. S., Straneo, F., Tarasov, L., van de Wal, R., and van den Broeke, M.: The future sea-level contribution of the Greenland ice sheet: a multi-model ensemble study of ISMIP6, *The Cryosphere*, 14, 3071–3096, <https://doi.org/10.5194/tc-14-3071-2020>, 2020.
- Hamilton, D. S., Lee, L. A., Pringle, K. J., Reddington, C. L., Spracklen, D. V., and Carslaw, K. S.: Occurrence of pristine aerosol environments on a polluted planet, *Proceedings of the National Academy of Sciences*, 111, 18466–18471, <https://doi.org/10.1073/pnas.1415440111>, 2014.
- 600 Hanna, E., Cappelen, J., Fettweis, X., Mernild, S. H., Mote, T. L., Mottram, R., Steffen, K., Ballinger, T. J., and Hall, R. J.: Greenland surface air temperature changes from 1981 to 2019 and implications for ice-sheet melt and mass-balance change, *International Journal of Climatology*, 41, E1336–E1352, <https://doi.org/10.1002/joc.6771>, 2021.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 610 Hofer, S., Tedstone, A. J., Fettweis, X., and Bamber, J. L.: Cloud microphysics and circulation anomalies control differences in future Greenland melt, *Nature Climate Change*, 9, 523–528, <https://doi.org/10.1038/s41558-019-0507-8>, 2019.



- Hogan, R. J. and Illingworth, A. J.: The Potential of Spaceborne Dual-Wavelength Radar to Make Global Measurements of Cirrus Clouds, *Journal of Atmospheric and Oceanic Technology*, pp. 1952–1969, <https://doi.org/10.1175/2011JAMC2646.1>, 1999.
- Inoue, J., Sato, K., Rinke, A., Cassano, J. J., Fettweis, X., Heinemann, G., Matthes, H., Orr, A., Phillips, T., Seefeldt, M., Solomon, A., and
 615 Webster, S.: Clouds and Radiation Processes in Regional Climate Models Evaluated Using Observations Over the Ice-free Arctic Ocean, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033 904, <https://doi.org/10.1029/2020JD033904>, 2021.
- Irbah, A., Delanoë, J., van Zadelhoff, G.-J., Donovan, D. P., Kollias, P., Puigdomènech Treserras, B., Mason, S., Hogan, R. J., and Tatarevic,
 A.: The classification of atmospheric hydrometeors and aerosols from the EarthCARE radar and lidar: the A-TC, C-TC and AC-TC prod-
 620 2023. *Atmospheric Measurement Techniques*, 16, 2795–2820, <https://doi.org/10.5194/amt-16-2795-2023>, publisher: Copernicus GmbH, 2023.
- Japan Aerospace Exploration Agency: EarthCARE/CPR L1B CPR one-sensor Received Echo Power Products and Doppler Product, <https://doi.org/10.57746/EO.01JDVCYDWJF63VPDBJP0VZ6V64>, 2025.
- Kessler, E.: On the Distribution and Continuity of Water Substance in Atmospheric Circulations, in: On the Distribution and Continuity
 of Water Substance in Atmospheric Circulations, pp. 1–84, American Meteorological Society, Boston, MA, ISBN 978-1-935704-36-2,
 625 https://doi.org/10.1007/978-1-935704-36-2_1, 1969.
- Khairoutdinov, M. and Kogan, Y.: A New Cloud Physics Parameterization in a Large-Eddy Simulation Model of Marine Stratocumulus, *Monthly Weather Review*, 128, 229–243, [https://doi.org/10.1175/1520-0493\(2000\)128<0229:ANCPPI>2.0.CO;2](https://doi.org/10.1175/1520-0493(2000)128<0229:ANCPPI>2.0.CO;2), 2000.
- Kärcher, B. and Lohmann, U.: A parameterization of cirrus cloud formation: Homogeneous freezing of supercooled aerosols, *Journal of
 Geophysical Research: Atmospheres*, 107, <https://doi.org/10.1029/2001JD000470>, 2002.
- Lacour, A., Chepfer, H., Miller, N. B., Shupe, M. D., Noel, V., Fettweis, X., Gallee, H., Kay, J. E., Guzman, R., and Cole, J.: How Well Are
 630 Clouds Simulated over Greenland in Climate Models? Consequences for the Surface Cloud Radiative Effect over the Ice Sheet, *Journal
 of Climate*, 31, 9293–9312, <https://doi.org/10.1175/JCLI-D-18-0023.1>, 2018.
- Lamer, K., Kollias, P., Battaglia, A., and Preval, S.: Mind the gap – Part 1: Accurately locating warm marine boundary layer clouds and
 precipitation using spaceborne radars, *Atmospheric Measurement Techniques*, 13, 2363–2379, <https://doi.org/10.5194/amt-13-2363-2020>,
 635 2020.
- Langen, P. L., Fausto, R. S., Vandecrux, B., Mottram, R. H., and Box, J. E.: Liquid Water Flow and Retention on the Green-
 land Ice Sheet in the Regional Climate Model HIRHAM5: Local and Large-Scale Impacts, *Frontiers in Earth Science*, 4,
<https://doi.org/10.3389/feart.2016.00110>, 2017.
- Lenaerts, J. T. M., Gettelman, A., Van Tricht, K., van Kampenhout, L., and Miller, N. B.: Impact of Cloud Physics on the Greenland Ice
 640 Sheet Near-Surface Climate: A Study With the Community Atmosphere Model, *Journal of Geophysical Research: Atmospheres*, 125,
 e2019JD031 470, <https://doi.org/10.1029/2019JD031470>, 2020.
- Lin, Y.-L., Farley, R. D., and Orville, H. D.: Bulk Parameterization of the Snow Field in a Cloud Model, *Journal of Climate and Applied
 Meteorology*, 22, 1065–1092, [https://doi.org/10.1175/1520-0450\(1983\)022<1065:BPOTSF>2.0.CO;2](https://doi.org/10.1175/1520-0450(1983)022<1065:BPOTSF>2.0.CO;2), 1983.
- Martin, G. M., Johnson, D. W., and Spice, A.: The Measurement and Parameterization of Effective Radius of Droplets in Warm Stratocumulus
 645 Clouds, *Journal of the Atmospheric Sciences*, 51, 1823–1842, [https://doi.org/10.1175/1520-0469\(1994\)051<1823:TMAPOE>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<1823:TMAPOE>2.0.CO;2),
 1994.
- Mason, S. L., Barker, H. W., Cole, J. N. S., Docter, N., Donovan, D. P., Hogan, R. J., Hünerbein, A., Kollias, P., Puigdomènech Treserras, B.,
 Qu, Z., Wandinger, U., and van Zadelhoff, G.-J.: An intercomparison of EarthCARE cloud, aerosol, and precipitation retrieval products,
Atmospheric Measurement Techniques, 17, 875–898, <https://doi.org/10.5194/amt-17-875-2024>, 2024.



- 650 Matrosov, S. Y.: Modeling Backscatter Properties of Snowfall at Millimeter Wavelengths, *Journal of the Atmospheric Sciences*, p. 1727–1736, <https://doi.org/10.1175/JAS3904.1>, 2007.
- Matrosov, S. Y., Uttal, T., and Hazen, D. A.: Evaluation of Radar Reflectivity–Based Estimates of Water Content in Stratiform Marine Clouds, *Journal of Applied Meteorology and Climatology*, pp. 405–419, [https://doi.org/https://doi.org/10.1175/1520-0450\(2004\)043<0405:EOREO>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0450(2004)043<0405:EOREO>2.0.CO;2), 2004.
- 655 Matrosov, S. Y., Battaglia, A., and Rodriguez, P.: Effects of Multiple Scattering on Attenuation-Based Retrievals of Stratiform Rainfall from CloudSat, *Journal of Atmospheric and Oceanic Technology*, p. 2199–2208, <https://doi.org/10.1175/2008JTECHA1095.1>, 2008.
- Meyers, M. P., DeMott, P. J., and Cotton, W. R.: New Primary Ice-Nucleation Parameterizations in an Explicit Cloud Model, *Journal of Applied Meteorology*, 31, 708–721, [https://doi.org/10.1175/1520-0450\(1992\)031<0708:NPINPI>2.0.CO;2](https://doi.org/10.1175/1520-0450(1992)031<0708:NPINPI>2.0.CO;2), 1992.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *Journal of Geophysical Research: Atmospheres*, 102, 16 663–16 682, <https://doi.org/10.1029/97JD00237>, 1997.
- 660 Mroz, K., Treserras, B. P., Battaglia, A., Kollias, P., Tatarevic, A., and Tridon, F.: Cloud and precipitation microphysical retrievals from the EarthCARE Cloud Profiling Radar: the C-CLD product, *Atmospheric Measurement Techniques*, 16, 2865–2888, <https://doi.org/10.5194/amt-16-2865-2023>, 2023.
- 665 Niwano, M., Hashimoto, A., and Aoki, T.: Cloud-driven modulations of Greenland ice sheet surface melt, *Scientific Reports*, 9, 10 380, <https://doi.org/10.1038/s41598-019-46152-5>, 2019.
- Otosaka, I. N., Shepherd, A., Ivins, E. R., Schlegel, N.-J., Amory, C., van den Broeke, M. R., Horwath, M., Joughin, I., King, M. D., Krinner, G., Nowicki, S., Payne, A. J., Rignot, E., Scambos, T., Simon, K. M., Smith, B. E., Sørensen, L. S., Velicogna, I., Whitehouse, P. L., A., G., Agosta, C., Ahlstrøm, A. P., Blazquez, A., Colgan, W., Engdahl, M. E., Fettweis, X., Forsberg, R., Gallée, H., Gardner, A., Gilbert, L., Gourmelen, N., Groh, A., Gunter, B. C., Harig, C., Helm, V., Khan, S. A., Kittel, C., Konrad, H., Langen, P. L., Lecavalier, B. S., Liang, C.-C., Loomis, B. D., McMillan, M., Melini, D., Mernild, S. H., Mottram, R., Mouginot, J., Nilsson, J., Noël, B., Pattie, M. E., Peltier, W. R., Pie, N., Roca, M., Sasgen, I., Save, H. V., Seo, K.-W., Scheuchl, B., Schrama, E. J. O., Schröder, L., Simonsen, S. B., Slater, T., Spada, G., Sutterley, T. C., Vishwakarma, B. D., van Wessel, J. M., Wiese, D., van der Wal, W., and Wouters, B.: Mass balance of the Greenland and Antarctic ice sheets from 1992 to 2020, *Earth System Science Data*, 15, 1597–1616, <https://doi.org/10.5194/essd-15-1597-2023>, 2023.
- 670 Protat, A., Delanoë, J., Bouniol, D., Heymsfield, A. J., Bansemmer, A., and Brown, P.: Evaluation of Ice Water Content Retrievals from Cloud Radar Reflectivity and Temperature Using a Large Airborne In Situ Microphysical Database, *Journal of Applied Meteorology and Climatology*, p. 557–572, <https://doi.org/https://doi.org/10.1175/JAM2488.1>, 2007.
- Pruppacher, H. R. and Klett, J. D.: Microphysics of clouds and precipitation, no. 18 in *Atmospheric and oceanographic sciences library*, Kluwer Academic Publishers, Dordrecht, 2nd rev. and enl. ed edn., ISBN 978-0-7923-4211-3, 1997.
- 680 Qu, Z., Donovan, D. P., Barker, H. W., Cole, J. N. S., Shephard, M. W., and Huijnen, V.: Numerical model generation of test frames for pre-launch studies of EarthCARE’s retrieval algorithms and data management system, *Atmospheric Measurement Techniques*, 16, 4927–4946, <https://doi.org/10.5194/amt-16-4927-2023>, 2023.
- Rotsteyn, L. D., Ryan, B. F., and Katzfey, J. J.: A Scheme for Calculation of the Liquid Fraction in Mixed-Phase Stratiform Clouds in Large-Scale Models, *Monthly Weather Review*, 128, 1070–1088, [https://doi.org/10.1175/1520-0493\(2000\)128<1070:ASFCOT>2.0.CO;2](https://doi.org/10.1175/1520-0493(2000)128<1070:ASFCOT>2.0.CO;2), 2000.
- 685 Ryan, J. C., Smith, L. C., Wu, M., Cooley, S. W., Miège, C., Montgomery, L. N., Koenig, L. S., Fettweis, X., Noel, B. P. Y., and van den Broeke, M. R.: Evaluation of CloudSat’s Cloud-Profiling Radar for Mapping Snowfall Rates Across the Greenland Ice Sheet, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031 411, <https://doi.org/10.1029/2019JD031411>, 2020.



- Saliba, G., Chen, C.-L., Lewis, S., Russell, L. M., Rivellini, L.-H., Lee, A. K. Y., Quinn, P. K., Bates, T. S., Haëntjens, N., Boss, E. S., Karp-Boss, L., Baetge, N., Carlson, C. A., and Behrenfeld, M. J.: Factors driving the seasonal and hourly variability of sea-spray aerosol number
 690 in the North Atlantic, *Proceedings of the National Academy of Sciences*, 116, 20 309–20 314, <https://doi.org/10.1073/pnas.1907574116>, 2019.
- Sankaré, H., Blanchet, J.-P., Laprise, R., and O'Neill, N. T.: Simulation of Arctic Thin Ice Clouds with Canadian Regional Climate Model Version 6: Verification against CloudSat-CALIPSO, *Atmosphere*, 13, 187, <https://doi.org/10.3390/atmos13020187>, 2022.
- Shupe, M. D.: Clouds at Arctic Atmospheric Observatories. Part II: Thermodynamic Phase Characteristics, *Journal of Applied Meteorology and Climatology*, p. 645–661, <https://doi.org/10.1175/2010JAMC2468.1>, 2011.
 695
- Shupe, M. D., Turner, D. D., Walden, V. P., Bennartz, R., Cadetdu, M. P., Castellani, B. B., Cox, C. J., Hudak, D. R., Kulie, M. S., Miller, N. B., Neely, R. R., Neff, W. D., and Rowe, P. M.: High and Dry: New Observations of Tropospheric and Cloud Properties above the Greenland Ice Sheet, *Bulletin of the American Meteorological Society*, <https://doi.org/10.1175/BAMS-D-11-00249.1>, 2013.
- Shupe, M. D., Turner, D. D., Zwink, A., Thieman, M. M., Mlawer, E. J., and Shippert, T.: Deriving Arctic Cloud Microphysics
 700 at Barrow, Alaska: Algorithms, Results, and Radiative Closure, *Journal of Applied Meteorology and Climatology*, 54, 1675–1689, <https://doi.org/10.1175/JAMC-D-15-0054.1>, 2015.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Liu, Z., Berner, J., Wang, W., Powers, J. G., Duda, M. G., Barker, D. M., and Huang, X.-Y.: A Description of the Advanced Research WRF Model Version 4, Tech. rep., UCAR/NCAR, <https://doi.org/10.5065/1DFH-6P97>, 2019.
- 705 Skorokhodov, A. V. and Kuryanovich, K. V.: Variability of Multilayer Cloud Field Structure over Western and Eastern Siberia in Summer and Winter in 2006–2023 Based on CALIPSO Data, *Atmospheric and Oceanic Optics*, 38, 290–299, <https://doi.org/10.1134/S1024856025700083>, 2025.
- Souverein, N., Gossart, A., Demuzere, M., Lenaerts, J. T. M., Medley, B., Gorodetskaya, I. V., Vanden Broucke, S., and van Lipzig, N. P. M.: A New Regional Climate Model for POLAR-CORDEX: Evaluation of a 30-Year Hindcast with COSMO-CLM2 Over Antarctica, *Journal of Geophysical Research: Atmospheres*, 124, 1405–1427, <https://doi.org/10.1029/2018JD028862>, 2019.
 710
- Storelvmo, T. and Tan, I.: The Wegener-Bergeron-Findeisen process – Its discovery and vital importance for weather and climate, *Meteorologische Zeitschrift*, 24, 455–461, <https://doi.org/10.1127/metz/2015/0626>, 2015.
- Sun, Z. and Rikus, L.: Parametrization of effective sizes of cirrus-cloud particles and its verification against observations, *Quarterly Journal of the Royal Meteorological Society*, 125, 3037–3055, <https://doi.org/10.1002/qj.49712556012>, 1999.
- 715 Sundqvist, H.: A parameterization scheme for non-convective condensation including prediction of cloud water content, *Quarterly Journal of the Royal Meteorological Society*, 104, 677–690, <https://doi.org/10.1002/qj.49710444110>, 1978.
- Taylor, P. C., Boeke, R. C., Li, Y., and Thompson, D. W. J.: Arctic cloud annual cycle biases in climate models, *Atmospheric Chemistry and Physics*, 19, 8759–8782, <https://doi.org/10.5194/acp-19-8759-2019>, 2019.
- Tjernström, M., Sedlar, J., and Shupe, M. D.: How Well Do Regional Climate Models Reproduce Radiation and Clouds in the Arctic? An
 720 Evaluation of ARCMIP Simulations, *Journal of Applied Meteorology and Climatology*, <https://doi.org/10.1175/2008JAMC1845.1>, 2008.
- Tompkins, A. M., Gierens, K., and Rädcl, G.: Ice supersaturation in the ECMWF integrated forecast system, *Quarterly Journal of the Royal Meteorological Society*, 133, 53–63, <https://doi.org/10.1002/qj.14>, 2007.
- van Dalum, C. T., van de Berg, W. J., Gadde, S. N., van Tiggelen, M., van der Drift, T., van Meijgaard, E., van Uft, L. H., and van den Broeke, M. R.: First results of the polar regional climate model RACMO2.4, *The Cryosphere*, 18, 4065–4088, <https://doi.org/10.5194/tc-18-4065-2024>, 2024.
 725



- van de Berg, W. J. and Medley, B.: Brief Communication: Upper-air relaxation in RACMO2 significantly improves modelled interannual surface mass balance variability in Antarctica, *The Cryosphere*, 10, 459–463, <https://doi.org/10.5194/tc-10-459-2016>, 2016.
- van den Broeke, M. R., Enderlin, E. M., Howat, I. M., Kuipers Munneke, P., Noël, B. P. Y., van de Berg, W. J., van Meijgaard, E., and Wouters, B.: On the recent contribution of the Greenland ice sheet to sea level change, *The Cryosphere*, 10, 1933–1946, <https://doi.org/10.5194/tc-10-1933-2016>, 2016.
- 730 van Kampenhout, L., Lenaerts, J. T. M., Lipscomb, W. H., Lhermitte, S., Noël, B., Vizcaíno, M., Sacks, W. J., and van den Broeke, M. R.: Present-Day Greenland Ice Sheet Climate and Surface Mass Balance in CESM2, *Journal of Geophysical Research: Earth Surface*, 125, <https://doi.org/10.1029/2019JF005318>, 2020.
- van Meijgaard, E., van Ulft, L., van de Berg, W. J., Bosveld, F., van den Hurk, B., Lenderink, G., and Siebesma, A.: The KNMI regional atmospheric climate model RACMO, version 2., KNMI Tech. Rep., <https://cdn.knmi.nl/knmi/pdf/bibliotheek/knmipubTR/TR302.pdf>, 2008.
- 735 Van Tricht, K., Lhermitte, S., Lenaerts, J. T. M., Gorodetskaya, I. V., L'Ecuyer, T. S., Noël, B., van den Broeke, M. R., Turner, D. D., and van Lipzig, N. P. M.: Clouds enhance Greenland ice sheet meltwater runoff, *Nature Communications*, 7, 10266, <https://doi.org/10.1038/ncomms10266>, 2016.
- 740 van Wessem, J. M., Reijmer, C. H., Lenaerts, J. T. M., van de Berg, W. J., van den Broeke, M. R., and van Meijgaard, E.: Updated cloud physics in a regional atmospheric climate model improves the modelled surface energy balance of Antarctica, *The Cryosphere*, 8, 125–135, <https://doi.org/10.5194/tc-8-125-2014>, 2014.
- Wandinger, U., Floutsi, A. A., Baars, H., Haarig, M., Ansmann, A., Hünerbein, A., Docter, N., Donovan, D., Van Zadelhoff, G.-J., Mason, S., and Cole, J.: HETEAC – the Hybrid End-To-End Aerosol Classification model for EarthCARE, *Atmospheric Measurement Techniques*, 16, 2485–2510, <https://doi.org/10.5194/amt-16-2485-2023>, 2023.
- 745 Wang, W., Zender, C. S., van As, D., and Miller, N. B.: Spatial Distribution of Melt Season Cloud Radiative Effects Over Greenland: Evaluating Satellite Observations, Reanalyses, and Model Simulations Against In Situ Measurements, *Journal of Geophysical Research: Atmospheres*, 124, 57–71, <https://doi.org/10.1029/2018JD028919>, 2019.
- Wehr, T., Kubota, T., Tzeremes, G., Wallace, K., Nakatsuka, H., Ohno, Y., Koopman, R., Rusli, S., Kikuchi, M., Eisinger, M., et al.: The EarthCARE mission–science and system overview, *Atmospheric Measurement Techniques*, 16, 3581–3608, 2023.
- 750 Wilson, D. R. and Ballard, S. P.: A microphysically based precipitation scheme for the UK meteorological office unified model, *Quarterly Journal of the Royal Meteorological Society*, 125, 1607–1636, <https://doi.org/10.1002/qj.49712555707>, 1999.
- Wood, R.: Parametrization of the effect of drizzle upon the droplet effective radius in stratocumulus clouds, *Quarterly Journal of the Royal Meteorological Society*, 126, 3309–3324, <https://doi.org/10.1002/qj.49712657015>, 2000.
- 755 Zou, L., Griessbach, S., Hoffmann, L., Gong, B., and Wang, L.: Revisiting global satellite observations of stratospheric cirrus clouds, *Atmospheric Chemistry and Physics*, 20, 9939–9959, <https://doi.org/10.5194/acp-20-9939-2020>, 2020.