

Hierarchical sedimentary architecture governs basin-scale solute dispersion: From pre-asymptotic dynamics to uncertainty propagation

5 Wanli Ren¹, Yue Fan², Anwen Pan¹, Heng Dai^{1*}, Jing Yang³, Mohamad Reza Soltanian⁴, Zhenxue Dai⁵,
6

¹ State Key Laboratory of Geomicrobiology and Environmental Changes, China University of Geosciences, Wuhan, Hubei, 430074, China.

10 ² Key Laboratory of Geotechnical Mechanics and Engineering of Ministry of Water Resources, Changjiang River Scientific Research Institute, Wuhan, Hubei, 430010, China.

³ School of Land Engineering, Chang'an University, Xi'an, Shanxi, 710064, China.

⁴ Departments of Geosciences and Environmental Engineering, University of Cincinnati, Cincinnati, Ohio, 45221, USA.

15 ⁵ College of Environmental and Municipal Engineering, Qingdao University of Technology, Qingdao, Shandong, 266033, China.

⁶ College of Construction Engineering, Jilin University, Changchun, Jilin, 130061, China.

Correspondence to: Heng Dai (daiheng@cug.edu.cn)

Abstract. Real aquifers are structured as hierarchical sedimentary systems, where multiscale heterogeneity and geometric connectivity jointly govern groundwater flow and solute migration. Although the general influence of heterogeneity has been extensively investigated, the scale-dependent effects of hierarchical organization, particularly under basin-scale flow conditions, remain inadequately quantified. In this study, we reconstructed a series of three-dimensional heterogeneous sedimentary architectures at the basin scale and performed numerical simulations to explore solute dispersion behaviour. The results reveal that the geometry and connectivity of dominant lithofacies at larger scale control macrodispersivity, while finer scale heterogeneity has only a secondary influence on plume evolution. Furthermore, the evolution of macrodispersivity is characterized by a prolonged pre-asymptotic phase, approaching a quasi-steady state after around 5000 days, with an asymptotic stability value of ~~170m~~ 170 m. This timescale is nearly 10 times longer than that inferred from the Borden site, where macrodispersivity stabilizes at around ~~0.4m~~ 0.4 m after approximately 400 days. This indicates that basin-scale solute transport remains non-ergodic over extended times and distances. Uncertainty analysis further identifies a distinct buffering

30 effect inherent to basin systems, in which the aggregation of numerous flow pathways dampens realization-to-realization variability caused by local heterogeneity. When integrated with previously reported laboratory- and sandbox-scale results from the same site, these findings establish a transferable framework linking hierarchical sedimentary architecture to multiscale dispersion and uncertainty. This framework advances theoretical understanding of non-Fickian transport and provides practical guidance for large-scale modeling and groundwater management in data-limited regions.

35 **1. Introduction**

The study of solute transport in heterogeneous porous media remains a central challenge in hydrogeology, with significant implications for groundwater science and engineering (Faroughi et al., 2023; Hansen and Berkowitz, 2020; Lee et al., 2018). Decades of field investigations, including the Borden, Cape Cod, MADE and Hanford tracer tests, and laboratory experiments have consistently demonstrated that spatial heterogeneity strongly governs plume migration, leading to non-
40 Fickian dispersion, anomalous breakthrough behaviour, and scale-dependent transport parameters (Sudicky, 1986; Garabedian et al., 1991; Boggs et al., 1992; Berkowitz et al., 2006; Botter et al., 2008; Chen et al., 2010, 2012; Agbotui et al., 2025; Tellam and Barker, 2006).

Scheibe and Freyberg (1995) systematically introduced the concept of hierarchical organization from sedimentology into hydrogeology to characterize aquifer heterogeneity. They pointed out that aquifer heterogeneity is not a single-scale problem
45 but rather results from the nested and superimposed geological units at different hierarchical levels (e.g., micro-, meso-, and macro-facies). Conceptually, this hierarchical-architecture view is consistent with sequence stratigraphy, which also organizes depositional heterogeneity in a nested manner through stratigraphic surfaces and stacking patterns. In practice, although stochastic theories and numerical upscaling frameworks have provided valuable tools to connect flow and mixing processes across spatial and temporal domains, many studies continue to compress heterogeneity into effective parameters (Dagan, 1984;
50 Gelhar, 1992; Dentz et al., 2011). Such simplification obscures the mechanistic role of multiscale sedimentary architecture in shaping flow pathways and controlling plume dynamics, and can bias spreading and mixing-relevant predictions (Fitts, 1996; Neuman and Tartakovsky, 2009; Dentz et al., 2023; Lester et al. 2016; Yin et al., 2023).

To elucidate the mechanisms underlying scale dependent transport in the heterogeneous porous media, a series of Lagrangian-based models has been developed within the hierarchical-architecture framework. By characterizing the spatial
55 organization of lithofacies across multiple scales, these models systematically integrate the multiscale heterogeneities into a unified representation, quantitatively linking sedimentary attributes (such as lithofacies volume proportion, mean length, and statistics of hydraulic conductivity) to transport metrics (such as dispersion and mixing). They have been successfully tested in laboratory and site-scale experiments (Soltanian et al., 2015b; Dai et al., 2020; Jia et al., 2023; Ma et al., 2025). Recent work further emphasizes that the role of lithofacies geometric attributes, particularly mean length and connectivity, control solute
60 dispersion and mixing in alluvial systems (Ramanathan et al., 2008; Soltanian et al., 2020; Ershadnia et al., 2021; Soltanian et al., 2017). High-resolution numerical simulations within this framework provide more intuitive and detailed information on

flow trajectories and interfacial reaction processes, thereby validating and supplementing the assumptions of the theoretical models (Ren et al., 2022).

65 ~~However, To date, the validation of most Lagrangian-based models, along with their associated simulation efforts, has primarily been conducted at the laboratory or site scale. For regional to basin scale solute transport at the regional scale or basin scale with processes involving long travel/migration distances and laterally extensive pathways still lacks flow paths, there remains a lack of quantitative characterization explicitly constrained by hierarchical sedimentary architecture. To date, most Lagrangian-based models' validation and related simulation work have been conducted at laboratory or site scales. Over the past decade, kilometer to basin-scale investigations have also begun to transition from simplified effective parameter~~
70 ~~descriptions toward structures. Although recent high-resolution, architecture-resolved analyses. Architectural attributes structure-resolving analytical approaches have shown that lithofacies geometry, connectivity, and conductivity (K) statistics have been shown to contrasts can~~ organize macroscopic dispersion and non-Fickian behaviour: high-resolution and architecture-resolved models built in the Llagas groundwater subbasin link lithofacies geometry and connectivity to behavior at kilometer scale, (Carle et al., 2006; Pauloo et al., 2021; Guo et al., 2019), it remains unclear which hierarchical level of
75 ~~heterogeneity remains transport-dominant under basin-scale flow conditions, how long pre-asymptotic behavior persists, and whether the superposition of numerous large-scale pathways amplifies or buffers predictive uncertainty.~~

~~Against this background, this study uses the Qiqihar aquifer system as a basin-scale testbed and combines hierarchical sedimentary architecture reconstruction with groundwater flow and solute transport simulations to investigate the following key questions: (i) Is basin-scale dispersion behavior primarily controlled by dominant large-scale lithofacies structures, or does it still require the explicit characterization of heterogeneity at finer scales? (ii) How do the geometry of solute sources, sedimentary architecture and hydraulic contrasts between lithofacies influence the evolution of macrodispersion and fast pathways/tailing contrasts (Carle et al., 2006); kilometer-scale simulations (Pauloo et al., 2021) demonstrate that aquifer anisotropy and seasonal recharge/pumping driven shifts in mean flow direction modulate the non-Fickian tails; and regional upscaling research under transient boundaries (Guo et al., 2019) shows that late time tails cannot be reproduced by the~~
85 ~~multirate mass transfer (MRMT) model calibrated under steady flow because mass transfer between high and low K units varies with boundary driven internal gradients. Even with these advances, architecture-resolved quantification of regional/basin scale plume evolution under its long-term pre-asymptotic behavior? (iii) Under field-representative basin-scale flow conditions, how do structural and hydraulic factors shape transport uncertainty and how uncertainties propagate and evolve? By comparing the results of this study with findings from previous laboratory- and field-scale studies conducted within~~
90 ~~the same conceptual framework (Dai et al., 2005; Ramanathan et al., 2010; Soltanian et al., heterogeneity and boundary driven gradients remains limited, particularly regarding the persistence of asymptotic behaviour and its uncertainty 2015a; Ren et al., 2023), this study clearly distinguishes which controlling factors possess scale-universality and which factors manifest specifically at the basin scale.~~

~~Accordingly, this study integrates hierarchical sedimentary structures characterization, numerical simulations, uncertainty analysis, using the Qiqihar aquifer system as a basin-scale testbed. Our objectives are to (i) quantify the relative~~
95

contributions of sedimentary architectural attributes, hydraulic statistics, and source size under field-representative aquifer and large-scale flow fields, (ii) identify which hierarchical scales of heterogeneity dominate solute dispersion, and (iii) evaluate how structural (model form) or parametric/data uncertainty propagate to transport predictions across scales. This work aims to provide guidance for more reliable predictions of solute transport in field-representative aquifer systems. Furthermore, as mentioned above, existing laboratory and site-scale studies have established mechanistic links between facies geometry/connectivity and solute dispersion (Dai et al., 2005; Ramanathan et al., 2010; Soltanian et al., 2015a). Recent global sensitivity analyses further demonstrate that limited aquifer structure parameters and hydraulic conductivity statistics exert first-order control on kilometer-scale dispersion (Ren et al., 2023). These findings enable comprehensive comparisons and discussions at experimental, field, and basin scales within the same research framework, thereby constructing a complete multiscale chain to identify which hierarchical sedimentary architecture and driving factors govern macroscopic spreading and dispersion across scales.

The paper is organized as follows: Section 2 introduces the geographic background of the study area, borehole data, and sedimentary architecture analysis methods; Section 3 describes the construction of the multiscale heterogeneous structural model and the simulation process of solute transport; Section 4 shows the simulation results and conducts uncertainty analysis to explore the influence of sedimentary architecture and permeability parameters on solute dispersion; Section 5 summarizes the main conclusions and outlines future research directions.

2. Methods

2.1 Overview of the study area

The study area is located on the east bank of the Nen River in Qiqihar City, Heilongjiang Province, China (Figure 1a). The Nen River defines the western boundary of the study area, and flows from the northeast to the southwest, with a total length of about 22 km and a width ranging from 400 m to 900 m. The regional terrain is gently inclined, sloping from north to south at less than 1‰, which exerts a primary control on groundwater flow direction and results in a general alignment of groundwater flow with that of the river. The aquifer system consists of a confined aquifer and a phreatic aquifer. The confined aquifer generally contains groundwater of relatively high quality, whereas groundwater contamination is largely concentrated in the phreatic aquifer. The present study therefore investigates solute transport processes within the phreatic aquifer to clarify the mechanisms of solute migration under the influence of sedimentary architecture.




Figure 1. (a) Geographic location map of the study area; (b) Boreholes and cross-sections

125 A total of 57 boreholes (see in Figure 1b) were collected for this study. According to the borehole data, the phreatic aquifer is composed of thick Quaternary unconsolidated deposits. The lithofacies of the aquifer is dominated by sandy gravel, gravelly medium- to coarse-grained sand, and medium- to fine-grained sand. The aquifer thickness varies from 22.1 m to 56.9 m, and the depth to the water table varies between 1.75 m and 5.86 m.

130 The phreatic aquifer is primarily recharged by infiltration from the Nen River, as river stage is consistently higher than the groundwater level. Other recharge includes precipitation infiltration and by lateral inflow from the northern and

northeastern boundaries. A relatively continuous aquitard, mainly composed of silty clay and clay, is located below this aquifer. However, it is locally absent near the Nen River and in the southern part of the study area, forming hydraulic windows that allow vertical connectivity between the two aquifer systems.

135 At the regional scale (kilometers), the phreatic aquifer exhibits a strongly layer-controlled, multiscale heterogeneity. Along the principal groundwater flow direction from upstream to downstream, the dominant lithofacies progressively transition from gravelly/medium-coarse sand to medium-fine sand. Away from the Nen River and into the floodplain, sediments become finer overall, with local development of low-permeability clay–silt lenses. Vertically, a pervasive “coarse-over-fine” organization is observed: the upper section is dominated by gravel and gravelly coarse/medium sand with moderate sorting, whereas the lower section consists mainly of medium-fine to fine sand with better sorting.

140 **2.2 Parameter determination**

2.2.1 Sediments heterogeneity parameters

Eight cross-sections (Figure 1b), oriented parallel to (defined as x -direction, sections 1-4) and perpendicular to (defined as y -direction, sections 5-8) the regional groundwater flow, were used to constrain the sedimentary structure characteristics. Following the multiscale hierarchical framework proposed by Dai et al. (2004) and Ritzi and Allen King (2007), this study
145 divided the lithofacies types of the aquifer into two hierarchical scales (Figure 2). Under this classification, Scale I and Scale II are used in a relative sense to denote two levels of heterogeneity representation within the same aquifer: Scale I resolves finer lithofacies variability, whereas Scale II represents a coarsened description in which Scale I facies are aggregated into composite units that preserve the dominant architectural organization. At Scale I, eight mutually exclusive lithofacies were defined based on permeability contrasts: gravel (G), coarse sand (CS), medium-coarse sand (MCS), medium sand (MS),
150 medium-fine sand (MFS), fine sand (FS), sub-sandy loam (SL), and clayey loam (C). At Scale II, these were aggregated into three composite units: gravel coarse sand (GCS), medium sand (MFS) and sandy clay (SC). Detailed delineations of the lithofacies at Scale I and Scale II are provided in Figures S1-S8 and S9-S16 in the Supplementary materials, respectively. From a sedimentological perspective, these sediments are interpreted as a river-alluvial-floodplain system associated with the Nen River. Coarse gravel/sand bodies represent channel zone (or paleochannel) deposits, while fine sand containing clay-silt
155 lenses represents floodplain and overflow deposits. Statistical analysis of borehole lithofacies characteristics also revealed a lateral grain size reduction towards downstream from the river, and a prevalent vertical structure of coarse sand overlaying fine sand in the aquifer.




Figure 2. Schematic diagram of lithofacies composition of the layered structural model of Qiqihar phreatic aquifer

160

The heterogeneous architecture models were produced via conditional stochastic simulation method under a transition-probability Markov-chain scheme, constrained by 57 boreholes and parameterized by lithofacies volume proportions and mean lengths. The ~~lithofacies~~ volume proportions (P) define the stationary occurrence probabilities ~~of each lithofacies~~, while the directional mean lengths (L_x, L_y, L_z) quantify ~~faeies~~ correlation scales of each lithofacies along the principal directions. To bias-correct parameter estimates affected by incomplete exposure of sections (see Figure S17 in Supplementary materials), a Bayesian updating scheme (Dai et al., 2005; White and Willis, 2000) and pixel-based image analysis were employed to calculate lithofacies lengths and volume proportions at both hierarchical scales. Comparison of the volume proportions computed from whole-sections (P_s) with those calculated from boreholes along the sections (P_D) revealed a close agreement (see Table S1 in Supplementary materials), indicating that these sections are sufficiently representative for estimating structural parameters.

170

The lithofacies volume proportions and mean vertical thicknesses (L_z) were derived from discrete borehole data, whereas mean horizontal lengths along x - and y - directions (L_x, L_y) were obtained from the cross-sections. Statistics of heterogeneous parameters are listed in Table 1. Define the horizontal anisotropy coefficient as $\alpha = L_y / L_x$, and the vertical anisotropy coefficient as $\varepsilon = L_z / [(L_x + L_y) / 2]$. According to Table 1, most lithofacies are horizontally isotropic at both scales, however, the MCS and SL lithofacies extend longer in the x - direction and the C lithofacies extend longer in the y - direction at Scale I. The mean anisotropy ratio decreases from ~ 0.005 (Scale I) to ~ 0.003 (Scale II), evidencing reduced apparent heterogeneity with model coarsening.

175

Table 1. Statistics of heterogeneous architecture parameters at Scale I and Scale II

Scale	Lithofacies	P	L_x (m)	L_y (m)	L_z (m)	α	ε
Scale I	G	0.34	2331.33	2495.40	8.94	1.070	0.004

	CS	0.06	2300.95	2310.10	11.41	1.004	0.005
	MCS	0.11	874.44	1187.37	9.43	1.358	0.009
	MS	0.13	1525.90	1496.85	6.45	0.981	0.004
	MFS	0.17	1516.81	1801.83	8.26	1.188	0.005
	FS	0.09	881.78	1009.77	3.99	1.145	0.004
	SL	0.04	640.32	791.79	2.55	1.236	0.004
	C	0.07	1098.27	763.32	2.48	0.695	0.003
Scale II	GCS	0.50	2205.99	2118.44	7.35	0.960	0.003
	MFS	0.39	1531.79	1412.87	5.06	0.922	0.003
	SC	0.11	817.75	842.73	2.60	1.030	0.003

In this study, the 57 boreholes provide hard conditioning data for faeieslithofacies occurrence and aquifer thickness, whereas the eight cross-sections supply additional structural constraints on lateral continuity and stratigraphic organization along and across the principal directions. This study isdoes not aim to deterministically reproduce specific in-situ plumes, but rather quantifies how hierarchical sedimentary architecture and associated parameter uncertainties govern basin-scale dispersion under field-representative flow conditions. The resulting heterogeneous model was intended to be statistically representative, while local connectivity in data-poor areas was treated as uncertain and quantified through a set of conditionallyconditional realizations. Although boreholes and corresponding cross-sections are more densely packed in the central and western parts of the study area and relatively sparse in the eastern area, this is sufficient to serve the objectives of this study.

2.2.2 Hydraulic conductivity

Hydraulic conductivity (K) was constrained using a combination of pumping-test analyses and grain-size-based empirical equations. A total of 45 pumping experiments were collected in the study area (including steady flow pumping tests and unsteady flow pumping tests) and the corresponding K values were obtained using analytical solutions and type-curve matching. However, the resulting K values are most representative for coarse-grained media. For fine-grained sediments, K was estimated from empirical equations. The soil samples collected from the shallow part of the study area were first analyzed for grain-size distribution, and K values were then estimated using the empirical equations summarized by Vuković and Soro (1992):

$$K = \frac{g}{\nu} C \varphi(n) d_e^2 \quad (1)$$

Where: K is the conductivity (m/d); ν is the kinematic viscosity (m^2/s); g is the gravity, taking the value of $9.81 m/s^2$; C is the empirical coefficient (provided in Table S2 in the Supplementary materials); $\varphi(n)$ is the dimensionless porosity function,

in which $n=0.225 \times (1+0.83)^{\eta}$, $\eta=d_{60}/d_{10}$, usually referred to as coefficient uniformity; d_e is the particle diameter corresponding to $e\%$ finer on the cumulative grain-size distribution curve.

200 Grain-size analyses showed that all samples had characteristic diameters of less than 0.2 mm (data from Dai et al., 2022).
 By comparing the K results calculated by various empirical formulas, the USBR method was finally adopted for the lithofacies SL. For the very low-permeability C unit, where grain-size methods are less reliable, K was inferred from a regional Plasticity Index- K ($PI-K$) empirical relationship, where the PI data was collected from clayey soils in the Qiqihar area. Through the
 205 on two scales were obtained, as shown in Table 2.

Table 2. Univariate statistics of conductivity for different hierarchical lithofacies at different scales

Scale	Lithofacies	$\bar{\varepsilon}$ (m/d)	σ_{ε}^2
Scale I	G	4.163	0.019
	CS	3.684	0.059
	MCS	3.165	0.126
	MS	2.775	0.177
	MFS	2.280	0.096
	FS	0.868	0.321
	SL	-0.251	0.116
Scale II	C	-4.212	0.118
	GCS	3.829	0.080
	MFS	2.336	0.402
	SC	-2.105	0.689

2.3 Construction of a heterogeneous structural model

Taking into account the aquifer structure and the topographic and geomorphological features of the study area, the simulation domain is set to be about 20 km in the x -direction, 22 km in the y -direction, and 50 m in the z -direction. Note that
 210 the purpose of this study is not ~~intend to aimed at~~ accurately ~~simulates~~ simulating groundwater dynamics or pollution plumes in the study area, but rather to discuss the influence of sedimentary structure and related parameters on solute dispersion. Therefore, for the convenience of modeling and parameter setting, the simulation area was simplified into an approximate square. As listed in Table 1, the SL facies type has the smallest mean horizontal extension (640.32 m), while the MS facies type has the smallest mean vertical extension (0.98 m). To capture lithological heterogeneity in detail, the grid cells were sized
 215 at $100 \text{ m} \times 100 \text{ m} \times 0.25 \text{ m}$, resulting in a total of $200 \times 220 \times 200$ cells.

Indicated kriging method based on Markov chain transfer probabilities was used to complete the modeling of heterogeneous architectures. Following Proce et al. (2004) and Ren et al. (2022), the sedimentary architecture was modeled in two stages. First, 50 realizations at Scale II and 50 realizations at Scale I were generated based on proportion and length statistics, respectively. This number of realizations is sufficient to obtain stable ensemble statistics (Zhou et al., 2018; Henri et al., 2020). Second, Scale I facies lithofacies were then mapped onto their corresponding Scale II domains to yield hierarchical models (hereafter referred to as the multiscale models). The resulting multiscale models thus embed fine-scale lithofacies heterogeneity within a framework that retains large-scale sedimentary architecture, enabling hierarchical coupling across scales. As an example, the three-dimensional and two-dimensional cross-sections of a particular heterogeneous model are shown in Figure 3. In all heterogeneous-architecture models, $x = 0$ is adjacent to the Nen River; larger x indicates greater distance from the river. Larger y denotes upstream, and smaller y denotes downstream.




Figure 3. An example of multiscale model: (a) 3-D facies, (b) 2-D sections; and Scale II model: (c) 3-D facies, (d) 2-D sections. Section lines are located at $x = 600\text{m}, 3000\text{m}, 7600\text{m}, 11000\text{m}$ and $y = 8000\text{m}, 12000\text{m}, 16000\text{m}, 18000\text{m}$


230 Facies structures in the simulated 2-D sections closely match field patterns and exhibit pronounced stratification. Along the groundwater flow direction, dominant faeieslithofacies transition from G and MCS to MS and MFS. Vertically, the succession fines downward, with coarser textures near the top and finer ones at the bottom.

2.4 Flow and solute transport simulation

2.4.1 Hydrogeologic conceptual model

235 Long-term groundwater level observations indicated that water levels had remained relatively stable over the years, suggesting a stable flow regime. According to multi-year groundwater level records, specified-head boundaries at $y = 0$ m and $y = 22$ km (corresponding to the upstream and downstream boundaries) with multi-year mean heads of 146 m and 140 m were set, respectively (Figure 4). The Nen River in the western part of the study area was also generalized as a steady head boundary (144 m). The eastern boundary was specified as a flux boundary, with flux values calculated from the natural hydraulic gradient.

240 The bottom of the phreatic aquifer is an aquitard with overflow discharge, so it was set as the flux boundary. The top boundary is the phreatic surface where recharge is primarily from precipitation, while discharge occurs through evaporation and artificial pumping.



245 **Figure 4. Conceptual hydrogeologic model of the study area and schematic diagram of solute plume planar source release conditions. The left side corresponds to the Nen River, and the pollution source is oriented approximately perpendicular to the regional groundwater flow direction.**

2.4.2 Groundwater flow simulation

In this study, MODFLOW-2005 (Harbaugh, 2005) was used to calculate the groundwater transport process in saturated porous media. After constructing the heterogeneous sedimentary architecture, stochastic K fields were generated by assigning

250 facies-conditioned K values to each cell based on the spatial distribution characteristics and statistics ($\bar{\mu}$ and σ_{μ}^2). Groundwater-flow simulations were run on the same grid as the architecture, ensuring a one-to-one correspondence between faeieslithofacies and K . Based on the recharge conditions of phreatic water, the lithology and thickness of the vadose-zone, city land and

farmland, four water-balance subregions were delineated (shown in Figure 1). Detailed water-balance equations and parameter values are provided in the source and sink calculation section of the Supplementary materials.

255 2.4.3 Solute transport simulation

The random walk particle tracer model program RWHet (LaBolle et al., 1996) was used to simulate the solute transport process. This study focuses on the transport characteristics of nonreactive solutes, and the governing equation for solute transport in saturated porous media defined as:

$$\begin{aligned} \frac{\partial}{\partial t} [\Theta(\mathbf{x}, t)c(\mathbf{x}, t)] = & - \sum_{i=1}^3 \frac{\partial}{\partial x_i} [v_i(\mathbf{x}, t)\Theta(\mathbf{x}, t)c(\mathbf{x}, t)] + \sum_{i,j=1}^3 \frac{\partial}{\partial x_i} [D_{ij}(\mathbf{x}, t)\Theta(\mathbf{x}, t) \frac{\partial c(\mathbf{x}, t)}{\partial x_j}] \\ & + \sum_k q_k(\mathbf{x}, t)c_k(\mathbf{x}, t)\delta_k(\mathbf{x} - \mathbf{x}_k) \end{aligned} \quad (2)$$

260 where c [M/L³] is the dissolved resident concentration; v [L/T] is the velocity; Θ [L³/L³] is the porosity; x_{ij} [L] is the distance along the respective Cartesian coordinate axis; c_k [M/L³] is the aqueous phase concentration in the flux q_k [L³/T] of water at x_k ; and δ_k is a Dirac function. D_{ij} [L²/T] is the local-scale dispersion tensor.

In this study, Θ was assumed to be stably isotropic and was set to the value of 0.35. Although setting a constant porosity may lead to deviations in the time required to reach a certain stage and the absolute value of the dispersion index plotted on the time axis, this study, however, emphasized the influence of aquifer structure and K -statistics under consistent settings, 265 where the spatial heterogeneity and connectivity of the corresponding velocity field were not significantly determined by subtle spatial variations in porosity. Another advantage of this choice is to avoid introducing other poorly constrained parameter fields into the model. For the same reason, the influence of local scale dispersion and molecular diffusion coefficient was not considered in this study, and therefore the corresponding dispersion and diffusion coefficients were taken as zero.

Solute transport was simulated under two distinct source scenarios: a point source and a planar source. The point source 270 represents localized releases (e.g., spills, leaks or point discharges), whereas the planar source represents an extended source zone that intersects multiple flow paths. These two scenarios are used to quantify how source dimensions control early-time sampling of heterogeneity and uncertainty in plume metrics. Using the bottom of the aquifer near the Nen River upstream as the origin of the coordinate system, the planar source (shown in Figure 4), oriented perpendicular to the groundwater flow, was centered at (10000, 3000, 25) m and extended 14000 m in the x -direction, 100 m in the y -direction, and 50 m in the z -direction. The point source was centered at (10000, 3000, 44) m, and extended 200 m in the x -direction, 200 m in the y -direction, and 2.5 m in the z -direction. A continuous NaCl source with a concentration of 800 mg/L was imposed based on 275 groundwater samples, with a background concentration of zero. Absorption type boundaries were defined at $y = 0$ m and $y = 22$ km to allow particles exit the simulation domain at the inlet and outlet boundaries, while reflection boundaries were used for all other boundaries to ensure particles remained within the domain. The solute transport simulation was performed over a 280 period of 10000 days, utilizing the same spatial discretization as the flow model.


285 The solute transport was measured by the solute concentration moments. According to the definitions of Freyberg (1986), the first moment, normalized by the mass in the solution, represents the position of the solute plume, expressed as the centroid coordinates (x_c, y_c, z_c) . The second spatial moment quantifies solute spreading around the centroid, given by the variances $(\sigma_{xx}^2, \sigma_{yy}^2, \sigma_{zz}^2)$. In the subsequent analysis, only the longitudinal component along the groundwater flow direction (y -direction) was retained.

3. Results

3.1 Numerical simulation results

3.1.1 Water flow simulation results

290 Model validation was performed using groundwater levels measured at 15 observation wells in 2020, whose spatial distribution is shown in Figure 5a. The average simulated heads from 50 realizations were compared with the observed values to evaluate model performance.




295 **Figure 5. (a) Distribution of groundwater level measurement points in 2020; (b) Fitting diagram between simulated water level and measured data**

The simulated water levels show good agreement with the observed values, closely following the 1:1 line. This visual consistency is supported by a relatively small error (RMSE = 0.507m), indicating that the water flow model reproduced the groundwater dynamics of the study area. This agreement further indicates that the hydraulic conductivity values derived from different methods are reasonable and consistent.

300 3.1.2 Solute transport simulation results

In stochastic transport modeling, aquifer heterogeneity is represented as independent random field with prescribed statistics. Consequently, transport characteristics are described using ensemble statistics over multiple realizations. ~~Two complementary metrics are commonly~~ Figure 6 shows the temporal evolution of plume center of mass $\gamma(t)$ and macro-dispersivity $\alpha(t)$ for multiscale (red) and Scale II (blue) models under point- and planar-source release conditions. Lines denote ensemble means, and shaded envelopes depict 10–90% confidence intervals from 50 stochastic realizations. Convergence tests indicate that 50 realizations are sufficient to achieve stable statistics. ~~Two complementary metrics are~~ employed to quantify macrodispersion (Dentz et al., 2000a, 2000b), which differ by their averaging procedure: the effective dispersion coefficient (D^{eff}) and the ensemble dispersion coefficient (D^{ens}). The former is computed by first calculating the moment-based dispersion for each realization and then averaging across realizations. In contrast, D^{ens} is derived directly from the spatial moments of the ensemble-averaged solute concentration field. ~~While both metrics characterize macrodispersion, D^{eff} more accurately represents the mean spreading in a typical field-scale realization and D^{ens} includes an “artificial” component from the spatial wandering of plume centroids between different realizations.~~ Under ergodic conditions or at sufficiently long times, these two measures are expected to converge, ~~but under finite-time and non-ergodic conditions D^{ens} may include an artificial contribution from realization-to-realization wandering of the plume centroid.~~ In this research, we report the corresponding macro-dispersivities, α_{eff} and α_{ens} , obtained by normalizing dispersion by the average groundwater velocity.

~~Figure 6 shows the temporal evolution of plume center of mass $\gamma(t)$ and macro-dispersivity $\alpha(t)$ for multiscale (red) and Scale II (blue) models under point- and planar source release conditions. Lines denote ensemble means, and shaded envelopes depict 10–90% confidence intervals from 50 stochastic realizations. Convergence tests indicate that 50 realizations are sufficient to achieve stable statistics. The close agreement between multiscale and Scale II models, especially under planar-source release (Figure 6c and 6d), demonstrates that at the basin scale, dispersion characteristics can be effectively captured by accurately representing the geometry of controlling lithofacies at larger scales, with diminished influence from smaller-scale heterogeneity. This finding is consistent with the findings of Ramanathan et al. (2010) and Soltanian et al. (2015b) at the site-scale research, who demonstrated similar behaviour using Lagrangian-based macrodispersion models. Comparable patterns were also observed in the site-scale simulations of Ren et al. (2022) at the Borden site, where $\alpha(t)$ exhibited a rapid increase and quickly converged to an asymptotic value.~~



330 **Figure 6. Results of Temporal evolution of longitudinal (a, c) solute transport distances distance $y_c(t)$ and (b, d) macro-dispersivity on two scales with time under two release conditions $\alpha(t)$ for multiscale (red) and Scale II (blue) models under point-source release (a, b) and planar-source release (c, d) conditions. Solid lines denote ensemble means, and shaded envelopes represent the 10th–90th percentile ranges.**

The time derivative of the mass transport distance corresponds to the mean velocity of solute migration in $y_c(t)$ provides a given direction. According to Figure 6a and 6c, the simple consistency check on the simulated transport behavior. The average transport velocity of the solute plume is approximately 0.058 m/d for a point-source release and 0.027 m/d for a planar-source release. Using the mean hydraulic conductivity of 12.05 m/d (Table 2), a regional hydraulic gradient of ~1%, and an effective porosity of ~0.35, (Figures 6a and 6c). The latter is close to the Darcy-based estimate of the mean groundwater velocity is, about 0.03 m/d. The consistency between this calculated velocity and the simulated plume migration suggests that the model adequately captures from the mean hydraulic conductivity of 12.05 m/d (Table 2), regional hydraulic gradient of ~1%, and effective porosity of ~0.35 used in this study. This consistency indicates that the flow and transport simulations reproduce the macroscopic solute transport characteristics in the study area. Comparisons with the Borden site highlight distinct timescales: $\alpha(t)$ at Borden increased rapidly and reached its asymptotic value within a relatively short time, whereas in this migration behavior of the study $\alpha(t)$ approached its asymptote more gradually (Ren et al., 2022) system.

345 A first-order result is the close agreement between the multiscale and Scale II simulations, particularly under planar-source release. This difference likely reflects the higher mean velocity at the Borden site (~0.091 m/d), which accelerates stabilization and indicates that basin-scale plume evolution can be captured primarily by the geometry and connectivity of the dominant large-scale lithofacies, whereas finer-scale heterogeneity mainly modulates the details of transport. The source geometry mainly affects early-time sampling of heterogeneity and thus the magnitude of realization-to-realization variability. Under point-source release, the plume initially samples only a limited portion of the K field, resulting in larger fluctuations in transport distance and solute dispersion. In contrast, the planar source intersects a broader set of flow paths from the outset, which narrows the uncertainty envelopes and yields a more stable ensemble-like response. This result is consistent with

350 previous theoretical and numerical studies showing that larger source areas enhance early sampling of heterogeneity, stabilize
large-scale transport statistics, and reduce uncertainty in plume behavior (Dagan, 2017; de Barros, 2018). This also confirms
that source dimensions remain important even at the basin scale.

At the release location, high proportions of gravel and medium to coarse sand lithofacies lead to strong local-scale
heterogeneity in the K field. Under point source release scenario, the plume initially samples a limited portion of the
355 heterogeneity, local scale flow velocity deviate significantly from average regional groundwater flow velocity, showing a large
fluctuation amplitude and significant uncertainty (shaded areas in Figures 6a and 6b). In contrast, planar source covers a
broader portion of the heterogeneous medium at early times. This “source area enlargement effect” smooths local-scale
velocity deviations and explains why confidence intervals are much narrower in the planar source scenarios. Similar trends
were reported by Cao et al. (2018) and de Barros (2018), who showed that enlarging the source area/width markedly narrows
360 uncertainty bands and reduces realization-to-realization spread of plume metrics in heterogeneous aquifers. Theoretically,
Dagan (2017) has explained why planar, large-area injections approach ergodic sampling of the conductivity field, by
smoothing local velocity fluctuations and stabilizing large-scale transport statistics.

The ensemble macro-dispersivity ($\alpha_{ens}(t)$) is more commonly used due to its simpler definition. Moreover, $\alpha_{ens}(t)$ often
overestimates the solute dispersion under non-ergodic or pre-asymptotic conditions. Figure 6b illustrates this overestimation
365 significantly and shows that the error associated with $\alpha_{ens}(t)$ increases progressively with transport time for both the multiscale
model and the larger-scale model. The persistence of this overestimation until the end of the 10000-day simulation suggests
that plume evolution remains largely governed by advection and has not yet reached a dispersion-dominated regime controlled
by local-scale dispersion. Once the solute plume samples a sufficiently large domain of the heterogeneous medium, realization
variability is substantially reduced and large-scale transport properties converge to the ensemble mean. Accordingly, $\alpha_{eff}(t)$
370 tends toward $\alpha_{ens}(t)$, consistent with theoretical predictions.

Figure 6 also shows that $\alpha_{eff}(t)$ and $\alpha_{ens}(t)$ do not fully converge within the 10000-day simulation period, although the
degree of separation varies with source geometry and transport stage. In general, the difference between the two metrics
remains more evident under point-source release condition. This incomplete convergence indicates that plume evolution
remains in a prolonged pre-asymptotic state under the simulated basin-scale transport conditions. At this stage, the plume's
375 evolution has not yet transitioned into the dispersion-dominated phase, which is characterized by the dominance of local-scale
dispersion processes. Compared with previously reported site-scale results at the Borden site, where macrodispersivity
approached a quasi-stationary value over a much shorter interval, the present basin-scale system exhibits a substantially slower
approach to convergence.

3.2 Uncertainty analysis of solute dispersion

380 A recent global sensitivity analysis indicated that a small set of geologically interpretable factors, most notably
faeieslithofacies volume proportions and in-facies mean hydraulic conductivity, exerts first-order influence on non-reactive
solute dispersion across regional to basin scales (Ren et al., 2023). Motivated by these findings, we next explore the individual

contribution of each factor to solute dispersion. To ensure a stable and consistent comparison while minimizing noise between realizations, all subsequent simulations were conducted under planar-source releases and with Scale II heterogeneous models. 385 This configuration focuses on the dominant lithofacies architecture, yielding an ensemble-like depiction of field-scale migration. This choice also provides a good chance to effectively narrow uncertainty bands and establish a clear baseline for systematically evaluating the impact of key parameters.

3.2.1 Effect of volume proportions

At Scale II, lithofacies are grouped into three types in decreasing order of permeability: GCS, MFS, and SC, with 390 volumetric proportions of 0.503, 0.385 and 0.112, respectively. Two scenarios are designed by changing the lithofacies volume proportions, while holding all other statistics fixed. In Group A, the proportions were set to 0.33, 0.34, and 0.33, approximating an aquifer with equal volumetric fractions of all lithofacies. In Group B, the mixture was skewed toward the low-permeability unit, with volume ratios of 0.2, 0.3, and 0.5, respectively. From a sedimentological perspective, in fluvial–alluvial systems the areal proportion of coarse deposits (e.g., gravel/sand bodies produced in paleochannel zones) versus floodplain fine deposits 395 can vary substantially at the basin scale, reflecting the coupled effects of stream power and sediment supply, channel migration, floodplain aggradation and development (Bridge, 2009). Accordingly, Group A (near-equal proportions) represents a more mixed and interbedded architecture consistent with frequent channel migration and ~~faeies~~lithofacies switching, whereas Group B (fine-dominated mixtures) represents a low-energy and/or distal floodplain setting where fine deposits are more prevalent and coarse bodies are more isolated.

400 Figure 7 shows the simulation results for different scenarios. With the increase of time, the solute transport distance increases proportionally, ~~and while~~ the $\alpha_{eff}(t)$ shows a power function growth trend ~~and approaches a quasi-stable value after roughly 5000 days. Across the tested scenarios, the asymptotic-scale $\alpha_{eff}(t)$ remains on the order of 170 m, indicating that the basin-scale system remains strongly dispersive over long travel times.~~ This behaviour is consistent with the well-established theory of large-scale dispersion in heterogeneous media, where transport is characterized by an initial non-Fickian regime 405 followed by a transition towards Fickian behaviour at late times (Dagan 1989; Neuman and Tartakovsky, 2009). ~~Across~~Increasing the ~~tested scenarios, the $\alpha_{eff}(t)$ stabilizes around 170 m after approximately 5000 days. High permeability lithofacies type (proportion of GCS) always provides preferential pathways that accelerate~~ enhances plume migration, ~~while low permeability type (SC) acts as barriers that restrict and leads to larger $\gamma_c(t)$ and $\alpha_{eff}(t)$ values, whereas increasing the proportion of SC suppresses both~~ plume ~~spreading and extent~~ solute residence times. This demonstrates that even at the basin 410 scale, migration and dispersivity. These trends indicate that modifying lithofacies proportions can still reshape ~~changes~~ the balance between connected preferential pathways and retarding controls. ~~Both high/low- K and low- K lithofacies exert strong controls on plume transport behaviours~~ domains, even though the overall basin-scale response remains relatively smooth. Previous theoretical and numerical studies (Fiori et al., 2010; Amooie et al., 2017; Soltanian and Ritzi, 2014; Puyguiraud et al., 2020), have also confirmed that large-scale solute dispersion is governed not by a single conductivity class, but by the 415 combined influence of extreme permeability contrasts.





Figure 7. **Results** Effect of lithofacies proportions on longitudinal (a) solute transport distances distance $y_c(t)$ and (b) macro-dispersivity $\alpha(t)$ under different lithofacies volume proportion scenarios planar-source release condition. Group A and Group B represent more mixed and fine-dominated architectures, respectively. Solid lines denote ensemble means, and shaded envelopes represent the 10th–90th percentile ranges.

A notable finding is that even when the proportion of GCS increases, the differences in plume metrics among scenarios remain modest, and the uncertainty bands widen only slightly. Actually, the plume travels only a short distance after 10000 days for this basin scale system (envelopes in Figure 7a), indicating that non ergodic conditions persist for extremely long times. This suggests that, changes in lithofacies proportions adjust transport characteristics, also show a systematic but the overall plume dynamics are buffered by the immensity of the flow domain. At regional basin scales, modifications moderate response to lithofacies proportions may act as secondary drivers of plume extent relative to other large-scale controls.

Realization to realization variability also reveals a clear scale dependent trend proportion. Increasing the proportion fraction of high- K lithofacies enhances solute mobility but simultaneously facies generally amplifies realization to realization the variability among realizations, whereas increasing the fraction of low- K lithofacies reduces both transport velocity and uncertainty, yielding facies produces more uniform outcomes. However, uncertainties do not grow indefinitely with conductive lithofacies dominance but instead plateau due to kilometer-scale the differences among scenarios remain much smaller than would typically be expected at the site scale. This suggests that, under basin-scale flow conditions, the effect of composition changes is present but moderated by large-scale spatial averaging. Such findings highlight the importance of explicitly accounting for scale dependent controls when extrapolating transport models from site to regional contexts, especially for risk assessment and large-scale groundwater management.

3.2.2 Influence of the mean value of the conductivity

The influence of conductivity on solute dispersion is primarily based on the difference in the mean K values among lithofacies. In the Qiqihar site, the mean K values are 46.02 m/d, 10.34 m/d, and 0.12 m/d for of GCS, MFS, and SC, respectively, indicating a pronounced permeability contrast. As is well known, K varies widely and is subject to considerable estimation and upscaling uncertainty at field scales. To isolate the role of individual lithofacies, three model groups were designed in which only the mean K of a single lithofacies was increased threefold, while the other two remained unchanged. The choice to expand by three times also takes into account the uncertainty of K at a medium to high level. In Group 1, the mean K of GCS was raised to 138.06 m/d. In Group 2, the mean K of MFS was increased to 31.02 m/d and in Group 3, the mean K of SC was increased to 0.36 m/d. In all cases, the variance of K was preserved, and the underlying heterogeneous sedimentary architecture remained unchanged. Thus, any changes in dispersion can be attributed to altered inter-facies K contrast and the resulting redistribution of flow among ~~faeies~~lithofacies. Solute transport simulations for these scenarios, conducted under planar-source release conditions, are presented in Figure 8.



450 **Figure 8. Results of (a) solute transport distances and (b) macro-dispersivity under the variations in lithofacies mean hydraulic conductivity. Effect of the mean K of individual Scale II lithofacies on longitudinal (a) solute transport distance $y_c(t)$ and (b) macro-dispersivity $\alpha(t)$ under planar-source release condition. In each scenario, the mean K is increased threefold while the lithofacies architecture and inter-facies variance remain unchanged. Solid lines denote ensemble means, and shaded envelopes represent the 10th–90th percentile ranges.**

455 Figures 8a-8c show the results for solute transport distance, and Figures 8d-8f are the $\alpha_{eff}(t)$ results. They all show a clear, asymmetric response when the mean K is tripled for a single lithofacies while keeping variance and architecture fixed. Increasing the mean K of the most permeable lithofacies (GCS) accelerates plume migration, raises the asymptotic α_{eff} , and slightly widens the uncertainty bands. Conversely, tripling the mean K of the medium permeability lithofacies (MFS) counterintuitively reduces migration distance and affects the asymptotic convergence of dispersion, whereas perturbing the K value of SC produces virtually no change. By calculating the global mean K of the 50 realizations, it can be obtained that the

460

global mean K value changes from 12.05 m/d of Qiqihar site to 22.20 m/d, 19.49m/d and 14.73 m/d after expanding the mean K values of the three lithofacies. Logically, MFS perturbation should have facilitated solute transport. Further analysis of the constructed heterogeneous sedimentary architecture model reveals that the dominant lithofacies progressively transition from gravelly/medium-coarse sand to medium-fine sand from upstream to downstream. MFS lithofacies is less distributed at the solute plume release location. As the conductivity in the model is generated based on lithofacies distribution, this may be a major reason for the anomalies in the transport distance results. Temporal variations in $\alpha_{eff}(t)$ also highlight the contribution of individual facies in shaping plume evolution. In the GCS perturbation, $\alpha_{eff}(t)$ increases sharply from the onset of solute release, and its asymptotic value remains well above the baseline case, emphasizing the dominance of coarse-type connectivity in channelling solute migration. In the MFS perturbation, $\alpha_{eff}(t)$ decreases in the early stages but then gradually converges to asymptotic equilibrium. This pattern reflects spatial averaging at the basin scale eventually restores ensemble-like behaviour. These early and late dynamics are (t) , and slightly widens the uncertainty bands. This indicates that the most permeable lithofacies has a strong influence on basin-scale plume migration. By contrast, increasing the mean K of MFS does not accelerate plume migration, even though the domain-scale mean conductivity increases (from 12.05 m/d of Qiqihar site to 19.49m/d). Instead, the transport distance is reduced relative to the baseline case, and the corresponding α_{eff} curve shows a different evolution toward quasi-stationary behavior. Increasing the mean K of SC produces only a very limited change in both transport distance and $\alpha_{eff}(t)$. The early and late dynamics of $\alpha_{eff}(t)$ are also consistent with observations of non-Fickian transport, where temporary suppression and delayed convergence are characteristic of strongly heterogeneous aquifers (Fiori and Dagan, 2000; Dentz et al., 2011). However, more importantly, the spatial averaging effect at the basin scale ultimately restores ensemble-like behaviour.

Collectively, the results indicate that macro-dispersion may not simply be controlled by the average conductivity of the system, but by the extent to which variations in facies properties reallocate groundwater fluxes across the underlying connected pathways. Specifically, enhancing GCS strengthens pre-existing preferential pathways, accelerating plume migration. The contrasting responses are likely related to the spatial roles of the three lithofacies in the modelled architecture. GCS is the most conductive lithofacies and is more influential near the release area, so increasing its mean K reinforces pre-existing fast routes and directly promotes plume migration. In contrast, increasing MFS is less dominant at the mean K of MFS release location and becomes more important farther along the transport domain. The mean K perturbation therefore reduces the contrast with GCS, and redistributes part of the groundwater flux into slower pathways, and thus away from the fastest paths, which suppresses early-time dispersion. This complex flux redistribution mechanism effectively suppresses the overall solute migration rate. For SC, even plume spreading. Even after a threefold increase leaves it far, SC remains much less conductive than the other facies, preventing it from contributing to the connected high- K transport network. Such asymmetric responses are consistent. These results show that basin-scale transport cannot be interpreted solely from the change in domain-average conductivity, because the effects of K perturbation depend strongly on which lithofacies is modified and where that lithofacies occurs within the connected architecture. These findings align with previous theoretical and numerical studies, which demonstrate/indicate that solute dispersion at large scales emerges from the interplay between

495 ~~faeieslithofacies~~ contrasts and connectivity, rather than from the mean conductivity values of single ~~faeieslithofacies~~ (Zinn and Harvey, 2003; Soltanian and Ritzi, 2014). ~~All these observations suggest that model calibration that focus only on asymptotic metrics may overlook early time transport features that are critical for risk assessment and monitoring system design.~~

4. Discussion

500 ~~Basin-scale simulations indicate that conservative solute dispersion and its uncertainty are principally organized by faeies conductivity contrasts and the geometry of the architectural elements at the larger scale, whereas modest perturbations to smaller-scale structure produce comparatively minor changes in transport metrics. This is consistent with the multiscale sedimentary architecture based theoretical analysis and with prior site-scale simulation studies. This study also distinguishes between $\alpha_{eff}(t)$ and $\alpha_{ens}(t)$: $\alpha_{ens}(t)$ increasingly overestimates the time-varying dispersion, while $\alpha_{eff}(t)$ measures spreading within a typical realization and approaches a stable value as sampling increases. The kilometer-scale $\alpha_{eff}(t)$ obtained in this study fall~~
505 ~~within the classical field-scale statistics compiled by Gelhar et al. (1992), who found $\alpha(t)$ typically in the tens to hundreds of meters at similar scales.~~

~~A central finding is the scale dependent nature of plume stabilization. Over a 10000 days simulation period, $\alpha_{eff}(t)$ initially increased slowly and then stabilized, but this stabilization occurs over a relatively prolonged period (around 5000 days), indicating that plume evolution remains in a pre-asymptotic (quasi-ergodic) state for thousands of days at the basin-scale.~~
510 ~~This behaviour contrasts with site scale results at the Borden site (Ren et al., 2022), where $\alpha_{eff}(t)$ was shown to reach an asymptotic value much more rapidly (after 400 days), plausibly facilitated by higher mean groundwater velocities. In the present basin-scale system, the solute explores only a small fraction of the heterogeneous flow field over a considerable period of time, the effective transport response is “buffered” by the domain immensity. Source size exerts a qualitatively consistent effect across scales: expanding the source promotes broader early time sampling of heterogeneity and thereby reduces inter-realization variability, yielding narrower uncertainty during transport process. By contrast, the uncertainty band for $\alpha(t)$ tends to stabilize or even decrease later at the basin scale, again reflecting domain-scale averaging that mutes source geometry effects far downstream. In the site scale tracer experiments (e.g., MADE, Borden), meter scale K variability may lead to apparent velocities differ by orders of magnitude between observation points, and strongly connected preferential pathways commonly yield pronounced early arrivals and heavy-tailed breakthrough curves (Zheng et al., 2011; Bianchi and Pedretti, 2017), thereby~~
515 ~~increasing predictive uncertainty in plume evolution and arrival time statistics. Simulations at the Borden site well demonstrate that the increased proportion of more permeable lithofacies significantly amplifies solute dispersion and output uncertainty. Basin-scale models, however, exhibit a weaker response, consistent with the buffering effect of long travel distance and multiple overlapping pathways.~~

~~This study further highlights that even at the basin scale, $\alpha(t)$ is not a simple constant. It is also closely related to the~~
525 ~~aquifer's heterogeneous structure and its hydraulic parameters. Scenario analysis reveals that the facies proportion and mean-conductivity perturbations illustrate two complementary mechanisms by which multiscale heterogeneity governs solute dispersion. Perturbations in facies proportions alter the statistical balance between preferential pathways and retarding domains,~~

whereas perturbations of mean conductivity reshape the contrast among facies types, redistributing fluxes across the connected network. At the basin scale, both mechanisms exert measurable influences on plume dynamics, but the divergence among scenarios remains modest due to kilometer scale spatial averaging and persistent non-ergodic conditions. Recent results clarify a central issue raised in the Introduction: under basin-scale flow conditions, the dominant controls on conservative solute dispersion are the geometry and connectivity of the larger-scale lithofacies framework, whereas finer-scale heterogeneity mainly modulates the details of plume evolution. Viewed together, the scenario analyses show that lithofacies proportions and mean K are not separate or competing explanations for this behavior, but two complementary ways in which the same architectural framework governs transport. Changing lithofacies proportions alters the statistical balance between preferential pathways and retarding domains, whereas changing the mean K of individual facies modifies how groundwater flux is partitioned among those pathways. What emerges from these results is that basin-scale dispersion is controlled less by domain-average hydraulic properties than by the way conductivity contrasts are embedded within the connected lithofacies architecture. This interpretation is consistent with previous regional and basin scale studies showing that aquifer-system heterogeneity, lithofacies architecture, and pathway organization remain influential controls on large-scale plume evolution (Carle et al., 2006; Pauloo et al., 2021). A recent global sensitivity analysis across multiscale heterogeneous media shows a robust ranking for non-reactive solute dispersion: (Ren et al., 2023): the ~~facies~~ lithofacies mean K is typically the most influential factor, followed by ~~facies~~-volume proportions and ~~facies~~-mean lengths; ~~whereas~~ variance and some correlation scales contribute less. ~~Importantly, when the heterogeneity integral scale reaches 100m to 1000m, the regional hydraulic gradient becomes non-negligible for non-reactive transport. Our~~ These basin-scale results are also broadly consistent with this ranking, especially in showing that lithofacies mean K and proportions exert the clearest influence on basin-scale dispersive behavior.

This study also indicates a markedly delayed approach to quasi-steady behavior at the basin scale. $\alpha_{eff}(t)$ approached a quasi-steady state only after roughly 5000 days, and $\alpha_{eff}(t)$ and $\alpha_{ens}(t)$ did not fully converge within the entire 10000-day simulation period. This persistent separation indicates that plume evolution remains strongly pre-asymptotic over long times and distances. Compared with the Borden tracer test study, where dispersivity approached quasi-stationary behavior after around 400 days (Ren et al., 2022), the present basin-scale system shows a substantially slower approach to convergence. Moreover, site-scale studies have also shown stronger sensitivity of plume behavior and predictive uncertainty to realization-specific pathways and source sampling (Zheng et al., 2011; de Barros and Dentz, 2016), whereas the basin-scale response here is more muted. These differences reflect not only domain-scale spatial averaging, but also the gradual way in which the plume samples the heterogeneous flow field and the delaying effect of multiple overlapping pathways on the emergence of ensemble-like transport. Within this setting, source geometry mainly influences early-time uncertainty by determining how much heterogeneity is sampled at the outset, while the long-term structure of plume evolution remains organized by the dominant large-scale architecture. These interpretations are also consistent with previous regional and basin scale studies showing that heterogeneous flow systems can sustain persistent non-Fickian transport over long distances and times, with a delayed approach to simplified or asymptotic transport behavior (Guo et al., 2019; Pauloo et al., 2021).

565 Combined with earlier laboratory sandbox experiments and site-scale studies developed within the same hierarchical framework (Ma et al., 2022, 2025), the present basin-scale results help refine how the role of heterogeneity should be understood across sedimentary scales. At local and site scales, conductivity contrasts, lithofacies geometry, and connectivity are often expressed more directly through realization-sensitive plume spreading, stronger source-size dependence, and more immediate pathway control on uncertainty (Dai et al., 2004; Ramanathan et al., 2010; Yin et al., 2020). At the basin scale, however, these rankings same controls remain operative, but their expression is increasingly filtered by long travel distances, pathway superposition, and spatial averaging, so that plume evolution becomes more slowly convergent and less sensitive to fine-scale structural differences. In this sense, the controlling mechanisms are not replaced at larger scales, rather, they are reorganized through scale-dependent averaging. In short, sedimentary heterogeneity exerts a scale-persistent but scale-reorganized control on solute transport.

575 This broader view also holds practical implications for model construction and interpretation. For problems focusing on bulk plume transport at regional to basin scales, it is not necessarily required to retain every detail of the fine-scale stratigraphic structure, provided that the dominant geological contrasts, lithofacies proportions, and directional organization of the connected flow framework are retained. This does not imply that finer-scale heterogeneity is irrelevant, on the contrary, it may remain important for early-time transport, near-source prediction, and problems sensitive to local concentration gradients. It must be acknowledged that neglecting porosity variations and molecular diffusion processes in this study may lead to an underestimation of early plume smoothing and lateral mixing, potentially delaying a significant convergence to Fick behaviour. However, at the basin scale and in the long-distance travel considered in this paper, structure-controlled velocity variations are expected to dominate the dispersion index; response, therefore, the main conclusions regarding relative the roles of lithofacies proportions, conductivity contrasts, and connectivity connected architecture remain unchanged. In that sense, the findings of this study support a scale-aware modeling strategy, in which the level of model complexity should be matched to the transport behavior of interest, rather than being increased indiscriminately across all scales. Such a scale-aware perspective provides a more practical theoretical basis for model simplification, monitoring design, and uncertainty assessment in data-limited regional aquifer systems.

585 Independent 3-D tank/column experiments conducted with sediments from the same Nen River setting (Ma et al., 2022, 2025) provide small-scale mechanistic insight into the scaling of $\alpha(t)$. By explicitly tying the growth of $\alpha(t)$ to velocity contrasts caused by K variability, to lithofacies geometry, and to cross-facies transition, a pronounced scale-dependence was observed: heterogeneous mixtures produced a much steeper growth of longitudinal $\alpha(t)$ than single-facies columns, with best-fit longitudinal $\alpha(t)$ on the order of 0.1 m. The consistency between the experimental evidence from this place and basin-scale model results supports our central conclusion that K -contrast and connectivity-governed mechanisms remain influential across multiple scales.

595 Taken together, the evidence from our study and a review of the broader literature supports three key points: (i) Solute dispersion and plume evolution are consistently shaped by coupled contrast and connectivity-driven processes across scales; (ii) the apparent influence of these mechanisms is scale-dependent and prolonged pre-asymptotic behaviour and pathway

superposition buffer bulk transport responses; and (iii) it is sufficient to preserve dominant geological contrasts, lithofacies proportions, and directional mean lengths for regional scale prediction, while problems near a release source or at early transport stage still require finer architectural resolution and careful treatment of source size. More broadly, this study emphasizes that reliable regional forecasts require not only calibrating mean hydraulic properties but also retaining the geological contrasts and connectivity that govern transport, together with explicit recognition of scale dependent averaging effects. These insights advance a more integrated understanding of basin-scale solute transport process and provide a conceptual basis for monitoring and management designs that remain robust under limited site characterization and persistent predictive uncertainty.

5. Conclusions

This study provides a comprehensive analysis of multiscale heterogeneity and its influence on non-reactive solute dispersion and modeling uncertainty within the Nen River Basin aquifer system. By integrating a hierarchical architectural model with numerical simulations and cross-scale validation, we have drawn several key conclusions that advance the understanding of transport in multiscale heterogeneous media.

First, our findings demonstrate that basin-scale solute dispersion is governed primarily by the geometry and connectivity of the dominant, large-scale lithofacies is sufficient to characterize solute dispersion at the basin scale, with smaller framework, whereas finer-scale architectural heterogeneity mainly modulates the details having a comparatively minor influence of transport. The evolution of dispersivity in our simulations is characterized by a long pre-asymptotic phase, which contrasts sharply with the rapid stabilization observed at sites like Borden. This unique behaviour highlights that at the basin scale, transport remains in a non-ergodic state for quite a long period of time, requiring a new perspective on long-term plume dynamics that accounts for this prolonged pre-asymptotic regime.

Second, we clarified the primary sources of modeling uncertainty. Our results show study identifies a “distinct buffering effect” inherent to in basin-scale systems, transport. where long travel distances and numerous superimposed pathways dampen the variability caused by local heterogeneity. This contrasts with local scale studies where preferential channels can lead to orders of magnitude variability in plume metrics. We found that the geometry of the contaminant source significantly impacts early time uncertainty, but this effect is muted at later stages by large scale spatial averaging. This underscores that for basin-scale problems, preserving dominant geological contrasts and connectivity is crucial, while the influence of local details diminishes over time. Although source geometry, lithofacies proportions, and inter-facies hydraulic contrasts all influence plume evolution, their effects on uncertainty are moderated by long travel distances, pathway superposition, and spatial averaging. Source dimensions remain important for early-time uncertainty, but their influence becomes less pronounced as transport integrates progressively larger parts of the heterogeneous domain.

Finally, we established a robust, multiscale framework for analyzing Third, the results support a scale-aware interpretation of solute transport in complex heterogeneous aquifers. This work highlights Sedimentary heterogeneity exerts a control on transport that reliable is persistent across scales but reorganized in its expression at the basin scale. For regional predictions

require not only accurate parameterization but also an explicit accounting for scale-dependent averaging effects- to basin-scale prediction, it is therefore most important to preserve dominant geological contrasts, lithofacies proportions, and the unique behaviour of dispersion in vast, non-ergodic systems. This approach is essential for designing effective connected architectural organization. This provides a more practical basis for model simplification, monitoring networks design, and robust risk uncertainty assessment models in areas with data-limited site characterization-regional aquifer systems.

Author contribution

635 Conceptualization: WLR and ZXD. Data curation: WLR. Formal analysis: WLR, AWP and YF. Funding acquisition: WLR and HD. Modeling and software: YF, MRS and JY. Visualization: AWP and WLR. Writing – original draft preparation: AWP. Writing – review & editing: WLR, YF, AWP, HD, JY, MRS, ~~ZXD~~ and ~~SHYZXD~~. All authors have read and agreed to the published version of the manuscript.

Competing Interests

640 One of the (co-)authors is a member of the editorial board of Hydrology and Earth System Sciences.

Acknowledgments

This work is funded by the National Natural Science Foundation of China (42302291, 42422208).

Reference

- 645 Agbotui, P. Y., Firouzbehi, F., & Medici, G.: Review of effective porosity in sandstone aquifers: insights for representation of contaminant transport. *Sustainability*, 17(14), 6469, <https://doi.org/10.3390/su17146469>, 2025.
- Amooie, M. A., Soltanian, M. R., and Moortgat, J.: Hydrothermodynamic mixing of fluids across phases in porous media. *Geophys. Res. Lett.*, 44(8), 3624-3634. <https://doi.org/10.1002/2016GL072491>, 2017.
- Berkowitz, B., Cortis, A., Dentz, M., and Scher, H.: Modeling non-Fickian transport in geological formations as a continuous time random walk. *Rev. Geophys.*, 44(2). <https://doi.org/10.1029/2005RG000178>, 2006.
- 650 ~~Bianchi, M., and Pedretti, D.: Geological entropy and solute transport in heterogeneous porous media. *Resour. Res.*, 53(6), 4691-4708. <https://doi.org/10.1002/2016WR020195>, 2017.~~
- Boggs, J. M., Young, S. C., Beard, L. M., Gelhar, L. W., Rehfeldt, K. R., and Adams, E. E.: Field study of dispersion in a heterogeneous aquifer: 1. Overview and site description. *Water. Resour. Res.*, 28(12), 3281-3291. <https://doi.org/10.1029/92WR01756>, 1992.
- 655 Botter, G., Peratoner, F., Putti, M., Zuliani, A., Zonta, R., Rinaldo, A., and Marani, M.: Observation and modeling of catchment-scale solute transport in the hydrologic response: A tracer study. *Water. Resour. Res.*, 44(5). <https://doi.org/10.1029/2007WR006611>, 2008.
- Bridge, J. S. (2009). *Rivers and floodplains: forms, processes, and sedimentary record*. John Wiley & Sons.

- 660 ~~Cao, G., Qin, R., Wu, Y., Wu, J., Xu, Z., and Zhang, C.: Effects of source size, monitoring distance and aquifer heterogeneity on contaminant mass discharge and plume spread uncertainty. Environ. Fluid. Mech., 18(2), 465-486. <https://doi.org/10.1007/s10652-017-9564-6>, 2018.~~
- Carle, S. F., Esser, B. K., and Moran, J. E. : High-resolution simulation of basin-scale nitrate transport considering aquifer system heterogeneity. Geosphere, 2(4), 195-209. <https://doi.org/10.1130/GES00032.1>, 2006.
- 665 Chen, C., Packman, A. I., Zhang, D., and Gaillard, J. F.: A multi-scale investigation of interfacial transport, pore fluid flow, and fine particle deposition in a sediment bed. Water. Resour. Res., 46(11). <https://doi.org/10.1029/2009WR009018>, 2010.
- ~~Chen, X., Murakami, H., Hahn, M. S., Hammond, G. E., Rockhold, M. L., Zachara, J. M., and Rubin, Y.: Three-dimensional Bayesian geostatistical aquifer characterization at the Hanford 300 Area using tracer test data. Water. Resour. Res., 48(6). <https://doi.org/10.1029/2011WR010675>, 2012.~~
- 670 Dagan, G.: Solute transport in heterogeneous porous formations. J. Fluid. Mech., 145, 151-177. <https://doi.org/10.1017/S0022112084002858>, 1984.
- Dagan, G.: Flow and Transport in Porous Formations. Springer Science and Business Media, 1989.
- Dagan, G.: Solute plumes mean velocity in aquifer transport: Impact of injection and detection modes. Adv. Water. Resour., 106, 6-10. <https://doi.org/10.1016/j.advwatres.2016.09.014>, 2017.
- 675 Dai, Z., Ma, Z., Zhang, X., Chen, J., Ershadnia, R., Luan, X., and Soltanian, M. R.: An integrated experimental design framework for optimizing solute transport monitoring locations in heterogeneous sedimentary media. J. Hydrol., 614, 128541. <https://doi.org/10.1016/j.jhydrol.2022.128541>, 2022.
- Dai, Z., Ritzi Jr, R. W., and Dominic, D. F.: Improving permeability semivariograms with transition probability models of hierarchical sedimentary architecture derived from outcrop analog studies. Water. Resour. Res., 41(7). <https://doi.org/10.1029/2004WR003515>, 2005.
- 680 Dai, Z., Ritzi Jr, R. W., Huang, C., Rubin, Y. N., and Dominic, D. F.: Transport in heterogeneous sediments with multimodal conductivity and hierarchical organization across scales. J. Hydrol., 294(1-3), 68-86. <https://doi.org/10.1016/j.jhydrol.2003.10.024>, 2004.
- 685 Dai, Z., Zhan, C., Dong, S., Yin, S., Zhang, X., and Soltanian, M. R.: How does resolution of sedimentary architecture data affect plume dispersion in multiscale and hierarchical systems?. J. Hydrol., 582, 124516. <https://doi.org/10.1016/j.jhydrol.2019.124516>, 2020.
- de Barros, F. P.: Evaluating the combined effects of source zone mass release rates and aquifer heterogeneity on solute discharge uncertainty. Adv. Water. Resour., 117, 140-150. <https://doi.org/10.1016/j.advwatres.2018.05.010>, 2018.
- ~~de Barros, F. P., & Dentz, M.: Pictures of blockscale transport: Effective versus ensemble dispersion and its uncertainty. Adv. Water. Resour., 91, 11-22. <https://doi.org/10.1016/j.advwatres.2016.03.004>, 2016.~~
- 690 Dentz, M., Hidalgo, J. J., and Lester, D.: Mixing in porous media: concepts and approaches across scales. Transport. Porous. Med., 146(1), 5-53. <https://doi.org/10.1007/s11242-022-01852-x>, 2023.

- Dentz, M., Kinzelbach, H., Attinger, S., and Kinzelbach, W.: Temporal behavior of a solute cloud in a heterogeneous porous medium: 2. Spatially extended injection. *Water. Resour. Res.*, 36(12), 3605-3614. <https://doi.org/10.1029/2000WR900211>, 2000a.
- 695 Dentz, M., Kinzelbach, H., Attinger, S., and Kinzelbach, W.: Temporal behavior of a solute cloud in a heterogeneous porous medium: 1. Point-like injection. *Water. Resour. Res.*, 36(12), 3591-3604. <https://doi.org/10.1029/2000WR900162>, 2000b.
- Dentz, M., Le Borgne, T., Englert, A., and Bijeljic, B.: Mixing, spreading and reaction in heterogeneous media: A brief review. *J. Contam. Hydrol.*, 120, 1–17. <https://doi.org/10.1016/j.jconhyd.2010.05.002>, 2011.
- Ershadnia, R., Hajirezaie, S., Amooie, A., Wallace, C. D., Gershenzon, N. I., Hosseini, S. A., Sturmer, D. M., Ritzi, R. W.,
700 and Soltanian, M. R.: CO₂ geological sequestration in multiscale heterogeneous aquifers: Effects of heterogeneity, connectivity, impurity, and hysteresis. *Adv. Water. Resour.*, 151, 103895. <https://doi.org/10.1016/j.advwatres.2021.103895>, 2021.
- Faroughi, S. A., Soltanmohammadi, R., Datta, P., Mahjour, S. K., and Faroughi, S.: Physics-informed neural networks with periodic activation functions for solute transport in heterogeneous porous media. *Mathematics*, 12(1), 63. <https://doi.org/10.3390/math12010063>, 2023.
- 705 Fiori, A., and Dagan, G.: Concentration fluctuations in aquifer transport: A rigorous first-order solution and applications. *J. Contam. Hydrol.*, 45(1-2), 139-163. [https://doi.org/10.1016/S0169-7722\(00\)00123-6](https://doi.org/10.1016/S0169-7722(00)00123-6), 2000.
- Fiori, A., Boso, F., de Barros, F. P., De Bartolo, S., Frampton, A., Severino, G., Suweis, S., and Dagan, G.: An indirect assessment on the impact of connectivity of conductivity classes upon longitudinal asymptotic macrodispersivity. *Water. Resour. Res.*, 46(8). <https://doi.org/10.1029/2009WR008590>, 2010.
- 710 Fitts, C. R.: Uncertainty in deterministic groundwater transport models due to the assumption of macrodispersive mixing: Evidence from the Cape Cod (Massachusetts, USA) and Borden (Ontario, Canada) tracer tests. *J. Contam. Hydrol.*, 23(1-2), 69-84. [https://doi.org/10.1016/0169-7722\(95\)00101-8](https://doi.org/10.1016/0169-7722(95)00101-8), 1996.
- Freyberg, D. L.: A natural gradient experiment on solute transport in a sand aquifer: 2. Spatial moments and the advection and dispersion of nonreactive tracers. *Water. Resour. Res.*, 22(13), 2031-2046, <https://doi.org/10.1029/WR022i013p02031>,
715 1986.
- Garabedian, S. P., LeBlanc, D. R., Gelhar, L. W., and Celia, M. A.: Large-scale natural gradient tracer test in sand and gravel, Cape Cod, Massachusetts: 2. Analysis of spatial moments for a nonreactive tracer. *Water. Resour. Res.*, 27(5), 911–924. <https://doi.org/10.1029/91WR00242>, 1991.
- 720 Gelhar, L. W., Welty, C., and Rehfeldt, K. R.: A critical review of data on field-scale dispersion in aquifers. *Water. Resour. Res.*, 28(7), 1955-1974. <https://doi.org/10.1029/92WR00607>, 1992.
- Guo, Z., Fogg, G. E., and Henri, C. V.: Upscaling of regional scale transport under transient conditions: Evaluation of the multirate mass transfer model. *Water. Resour. Res.*, 55(7), 5825–5841. <https://doi.org/10.1029/2019WR024953>, 2019.

- 725 Hansen, S. K., and Berkowitz, B.: Modeling non-Fickian solute transport due to mass transfer and physical heterogeneity on
arbitrary groundwater velocity fields. *Water. Resour. Res.*, 56(10), e2019WR026868.
<https://doi.org/10.1029/2019WR026868>, 2020.
- Harbaugh, A. W.: MODFLOW-2005, the US Geological Survey modular ground-water model: the ground-water flow process
(Vol. 6). Reston, VA, USA: US Department of the Interior, US Geological Survey, 2005.
- 730 Henri, C. V., Harter, T., & Diamantopoulos, E.: On the conceptual complexity of non-point source management: impact of
spatial variability. *Hydrol. Earth. Syst. Sc.*, 24(3), 1189-1209, <https://doi.org/10.5194/hess-24-1189-2020>, 2020.
- Jia, S., Dai, Z., Zhou, Z., Ling, H., Yang, Z., Qi, L., Wang A. H., Zhang, X. Y., Thanh H. V., and Soltanian, M. R.: Upscaling
dispersivity for conservative solute transport in naturally fractured media. *Water. Res.*, 235, 119844.
<https://doi.org/10.1016/j.watres.2023.119844>, 2023.
- 735 LaBolle, E. M., Fogg, G. E., and Tompson, A. F.: Random-walk simulation of transport in heterogeneous porous media: Local
mass-conservation problem and implementation methods. *Water. Resour. Res.*, 32(3), 583-593.
<https://doi.org/10.1029/95WR03528>, 1996.
- Lee, J., Rolle, M., and Kitanidis, P. K.: Longitudinal dispersion coefficients for numerical modeling of groundwater solute
transport in heterogeneous formations. *J. Contam. Hydrol.*, 212, 41-54. <https://doi.org/10.1016/j.jconhyd.2017.09.004>,
2018.
- 740 Lester, D. R., Trefry, M. G., and Metcalfe, G.: Chaotic advection at the pore scale: Mechanisms, upscaling and implications
for macroscopic transport. *Adv. Water. Resour.*, 97, 175-192. <https://doi.org/10.1016/j.advwatres.2016.09.007>, 2016.
- Ma, Z., Dai, Z., Zhang, X., Zhan, C., Gong, H., Zhu, L., Wallace C. D., and Soltanian, M. R.: Dispersivity variations of solute
transport in heterogeneous sediments: numerical and experimental study. *Stoch. Env. Res. Risk. A.*, 36(2), 661-677.
<https://doi.org/10.1007/s00477-021-02040-x>, 2022.
- 745 Ma, Z., Qi, L., Dai, Z., Yang, Z., Ma, Y., Wang, D., ... and Soltanian, M. R.: Impact of multiscale heterogeneous sediments
and boundary conditions on dispersivity spatial variations. *Water. Resour. Res.*, 61(6), e2024WR039151.
<https://doi.org/10.1029/2024WR039151>, 2025.
- Neuman, S. P., and Tartakovsky, D. M.: Perspective on theories of non-Fickian transport in heterogeneous media. *Adv. Water.
Resour.*, 32(5), 670-680. <https://doi.org/10.1016/j.advwatres.2008.08.005>, 2009.
- 750 Pauloo, R. A., Fogg, G. E., Guo, Z., and Henri, C. V.: Mean flow direction modulates non-Fickian transport in a heterogeneous
alluvial aquifer-aquitard system. *Water. Resour. Res.*, 57(3), e2020WR028655. <https://doi.org/10.1029/2020WR028655>,
2021.
- Proce, C. J., Ritzi, R. W., Dominic, D. F., and Dai, Z.: Modeling multi-scale heterogeneity and aquifer interconnectivity.
Groundwater, 42(5), 658-670. <https://doi.org/10.1111/j.1745-6584.2004.tb02720.x>, 2004.
- 755 Puyguraud, A., Perez, L. J., Hidalgo, J. J., and Dentz, M.: Effective dispersion coefficients for the upscaling of pore-scale
mixing and reaction. *Adv. Water. Resour.*, 146, 103782. <https://doi.org/10.1016/j.advwatres.2020.103782>, 2020.

- Ramanathan, R., Ritzi Jr, R. W., and Allen-King, R. M.: Linking hierarchical stratal architecture to plume spreading in a Lagrangian-based transport model: 2. Evaluation using new data from the Borden site. *Water. Resour. Res.*, 46(1). <https://doi.org/10.1029/2009WR007810>, 2010.
- 760 Ramanathan, R., Ritzi, R. W., Jr., and Huang, C.: Linking hierarchical stratal architecture to plume spreading in a Lagrangian-based transport model. *Water. Resour. Res.*, 44(4). <https://doi.org/10.1029/2007wr006282>, 2008.
- Ren, W., Dai, H., Yuan, S., Dai, Z., Ye, M., and Soltanian, M. R.: Global sensitivity study of non-reactive and sorptive solute dispersivity in multi-scale heterogeneous sediments. *J. Hydrol.*, 619, 129274. <https://doi.org/10.1016/j.jhydrol.2023.129274>, 2023.
- 765 Ren, W., Ershadnia, R., Wallace, C. D., LaBolle, E. M., Dai, Z., de Barros, F. P., and Soltanian, M. R.: Evaluating the effects of multiscale heterogeneous sediments on solute mixing and effective dispersion. *Water. Resour. Res.*, 58(9), e2021WR031886. <https://doi.org/10.1029/2021WR031886>, 2022.
- Ritzi Jr, R. W., and Allen-King, R. M.: Why did Sudicky [1986] find an exponential-like spatial correlation architecture for hydraulic conductivity at the Borden research site?. *Water. Resour. Res.*, 43(1). <https://doi.org/10.1029/2006WR004935>,
770 2007.
- Scheibe, T. D., and Freyberg, D. L.: Use of sedimentological information for geometric simulation of natural porous media structure. *Water. Resour. Res.*, 31(2), 325–337. <https://doi.org/10.1029/95WR02570>, 1995.
- Soltanian, M. R., and Ritzi, R. W.: A new method for analysis of variance of the hydraulic and reactive attributes of aquifers as linked to hierarchical and multi-scaled sedimentary architecture. *Water. Resour. Res.*, 50(12), 9766-9776. <https://doi.org/10.1002/2014WR015468>, 2014.
775
- Soltanian, M. R., Behzadi, F., and de Barros, F. P.: Dilution enhancement in hierarchical and multi-scale heterogeneous sediments. *J. Hydrol.*, 587, 125025. <https://doi.org/10.1016/j.jhydrol.2020.125025>, 2020.
- Soltanian, M. R., Ritzi, R. W., Huang, C. C., and Dai, Z.: Relating reactive solute transport to hierarchical and multi-scale sedimentary architecture in a Lagrangian-based transport model: 1. Time-dependent effective retardation factor. *Water. Resour. Res.*, 51(3), 1586-1600. <https://doi.org/10.1002/2014WR016353>, 2015a.
780
- Soltanian, M. R., Ritzi, R. W., Huang, C. C., and Dai, Z.: Relating reactive solute transport to hierarchical and multi-scale sedimentary architecture in a Lagrangian-based transport model: 2. Particle displacement variance. *Water. Resour. Res.*, 51(3), 1601-1618. <https://doi.org/10.1002/2014WR016354>, 2015b.
- Soltanian, M. R., Sun, A., and Dai, Z.: Reactive transport in the complex heterogeneous alluvial aquifer of Fortymile Wash, Nevada. *Chemosphere*, 179, 379-386, <https://doi.org/10.1016/j.chemosphere.2017.03.136>, 2017.
785
- Sudicky, E. A.: A natural gradient experiment on solute transport in a sand aquifer: Spatial variability of hydraulic conductivity and its role in the dispersion process. *Water. Resour. Res.*, 22(13), 2069-2082, <https://doi.org/10.1029/WR022i013p02069>, 1986

- 790 Tellam, J. H., & Barker, R. D.: Towards prediction of saturated-zone pollutant movement in fractured
permeable-matrix aquifers: the case of the UK Permo-Triassic sandstones.
<https://doi.org/10.1144/GSL.SP.2006.263.01.01>, 2006.
- Vuković, M., and Soro, A.: Determination of hydraulic conductivity of porous media from grain-size composition (pp. v+-83),
1992.
- 795 White, C. D., and Willis, B. J.: A method to estimate length distributions from outcrop data. *Math. Geol.*, 32(4), 389-419,
<https://doi.org/10.1023/A:1007510615051>, 2000.
- Yin, M., Ma, R., Zhang, Y., Lin, J., Guo, Z., and Zheng, C.: Competitive control of multi-scale aquifer heterogeneity on solute
transport in an alluvial aquifer. *J. Hydrol.*, 616, 128819, <https://doi.org/10.1016/j.jhydrol.2022.128819>, 2023.
- Yin, M., Zhang, Y., Ma, R., Tick, G. R., Bianchi, M., Zheng, C., Wei, W., Wei, S., and Liu, X.: Super-diffusion affected by
hydrofacies mean length and source geometry in alluvial settings. *J. Hydrol.*, 582, 124515,
800 <https://doi.org/10.1016/j.jhydrol.2019.124515>, 2020.
- Zheng, C., Bianchi, M., and Gorelick, S. M.: Lessons learned from 25 years of research at the MADE site. *Groundwater*, 49(5),
649-662, <https://doi.org/10.1111/j.1745-6584.2010.00753.x>, 2011.
- Zhou, D., Zhang, Y., Gianni, G., Lichtner, P., and Engelhardt, I.: Numerical modelling of stream-aquifer interaction:
Quantifying the impact of transient streambed permeability and aquifer heterogeneity. *Hydrol. Process*, 32(14), 2279-
805 2292, <https://doi.org/10.1002/hyp.13169>, 2018.
- Zinn, B., and Harvey, C. F.: When good statistical models of aquifer heterogeneity go bad: A comparison of flow, dispersion,
and mass transfer in connected and multivariate Gaussian hydraulic conductivity fields. *Water. Resour. Res.*, 39(3),
<https://doi.org/10.1029/2001WR001146>, 2003.