

The answer is blowing in the wind: seasonal hydrography and mixing of the inner sea of Tierra del Fuego, Southern Patagonia

Manuel I. Castillo^{2,8}, Constanza Zuñiga^{1,6}, Carmen Barrios-Guzmán^{1,9}, Natalia Cisternas, José Garcés-Vargas^{4,5}, Mauricio F. Landaeta^{3,7,8}, Andrea Piñones^{5,7}, Marcela Rojas^{2,6}, Alicia I. Guerrero¹, Maritza Sepúlveda¹

¹Laboratorio de Ecología y Conservación de Mamíferos Marinos (LECMMAR), Universidad de Valparaíso, Chile

²Laboratorio de Oceanografía Física y Satelital (LOFISAT), Facultad de Ciencias del Mar y de Recursos Naturales, Universidad de Valparaíso, Chile

³Laboratorio de Ictiología e Interacciones Biofísicas (LABITI), Instituto de Biología, Facultad de Ciencias, Universidad de Valparaíso, Chile.

⁴Instituto de Ciencias Marinas y Limnológicas, Facultad de Ciencias, Universidad Austral de Chile

⁵Centro FONDAP de Investigación en Dinámica de Ecosistemas Marinos de Altas Latitudes (IDEAL), Universidad Austral de Chile, Chile

⁶Programa de Magister en Oceanografía, PUCV/UV


⁷Millennium Institute Biodiversity of Antarctic and Sub-Antarctic Ecosystems (BASE), Santiago, Chile

⁸Centro de Observación y Análisis del Océano Costero, COSTAR-UV, Facultad de Ciencias del Mar y de Recursos Naturales, Universidad de Valparaíso, Chile

⁹Centro Bahía Lomas, Facultad de Ciencias, Universidad Santo Tomás, Chile

Correspondence to: Manuel I. Castillo (manuel.castillo@uv.cl)

Abstract. This study characterizes seasonal hydrography and mixing processes in Almirantazgo Fjord, a sensitive ecosystem in southern Chilean Patagonia. Although estuarine and tidal forcing conventionally explain fjord dynamics, wind stress effects remain less understood in this high-latitude region. The study analyses a comprehensive six-month dataset including a moored time-series of temperature, salinity, and dissolved oxygen, cross-fjord CTD transects, and hydrographic profiles derived from seal-deployed sensors. Observations indicate distinct seasonality, shifting from a stratified water column in summer—defined by low-salinity surface water from glacial melt—to a mixed winter state with significantly reduced vertical stability. The analysis identifies persistent, topographically channelled up-fjord winds as a primary physical driver. By applying the Wedderburn number (W_b) and mechanical energy balance calculations, we determined that strong wind stress perturbs the pycnocline ($W_b > 1$). During stratified summer periods, strong wind events (above the 90th percentile) generated wind power of the same order of magnitude as that of the estuarine circulation. Under such conditions, wind forcing amplifies vertical mixing, modulates the pressure gradient, and supports oxygenation in the upper and subsurface layers. First-order estimates indicate that the upper brackish layer is flushed in approximately one week, reflecting a dynamic surface exchange governed by the interplay of freshwater buoyancy and wind stress. These results indicate that wind constitutes a primary mechanism regulating the hydrographic structure and biogeochemical function of the Tierra del Fuego inner sea.



Graphical abstract: Schematic representation illustrating the key findings of the study on the Almirantazgo Fjord. The figure depicts typical summer (T_s, S_s, N^2_s) and winter (T_w, S_w, N^2_w) hydrographic profiles obtained via elephant seal-deployed CTD-SRLDs, alongside a conceptualization of the fjord's along-fjord and cross-fjord circulation patterns. Notice that the vertical scale of the upper layer (h_1) was exaggerated, to focus on the stratification layers. The outflow (Q_1) and inflow (Q_2) represent the estuarine exchange whereas Q_3 that could be tidal induced. The position of the instruments are also at the bottom of the head of the fjord at 30 m depth.

1 Introduction

Fjord basins originate from glacial advance and retreat, processes that produce elongated narrow basins characteristically containing one or several glacial moraines known as sills (Dyer, 2019; Farmer and Freeland, 1983; Geyer and MacCready, 2014; Inall and Gillibrand, 2010; Stigebrandt, 2012). These ecosystems play a key role in CO₂ sequestration, a function

50 resulting from the undersaturation of their surface waters (Aalto et al., 2021). Furthermore, the growing utilization and exposure of these systems has led to their designation as “aquatic critical zones” (Bianchi et al., 2020).

In these coastal systems, numerous processes operate on different temporal and spatial scales. This results in a complex combination of forces that determines the observed circulation patterns in regional and small-scale basins. Typically, residual
55 circulation in shallow water estuaries had been described as two-layered and density driven (Officer, 1976; Valle-Levinson et al., 2014). The density differences produce an along-fjord pressure gradient, due to freshwater input at the fjord head and authors like Officer (1976), Dyer (1997) and Valle-Levinson et al (2014), attributed as the main drivers of residual circulation. This gradient produces the two-layered, along-fjord gravitational circulation (Hansen and Rattray, 1965; Ribeiro et al., 2004). This pattern is manifested as a residual circulation pattern after several tidal cycles (MacCready and Banas, 2012) and can be
60 substantially modified by wind stress (Guo and Valle-Levinson, 2008), bathymetric and frictional effects, the Earth’s rotation (Yang et al., 2015), and drag (McCabe et al., 2006).

In deep-waters estuaries, like fjords, the two-layer circulation is often confined to a narrow portion of the water column, sequestered within a sharp pycnocline (Valle-Levinson et al., 2014). This fundamental structure is subject to modification by
65 several external forcings, including wind stress (e.g., Klinck et al., 1981; Castillo et al., 2017), remote density gradients (Stigebrandt, 2012), and tidal action (Ianniello, 1977; Valle-Levinson et al., 2007). Observations in Chilean fjords, particularly in northern Patagonia, have identified three-layer residual circulation patterns. Although these are primarily attributed to tidal forcing in deep fjords, the signal is relatively weak and often obscured by wind-driven effects (Valle-Levinson et al., 2014). Furthermore, research in Douglas Channel, British Columbia demonstrates a seasonal shift from three-layer circulation in
70 summer to a four-layer structure in winter. Such transitions are likely driven by the dynamic adjustment of estuarine flow, direct wind forcing, and the coupled barotropic and baroclinic responses to fluctuations in surface wind stress (Wan et al., 2017).

A complete description of fjord circulation must consider the cross-fjord dynamics. This cross-fjord pattern is generally weak
75 and consists of an overturning motion known as secondary circulation (Chant, 2002; Lacy and Monismith, 2001; MacCready and Geyer, 2010). The along-fjord wind stress can intensify the estuarine circulation during down-fjord winds or weaken the surface outflow during up-fjord winds (e.g. Valle-Levinson, 2010). The study of Jackson et al., (2025) shows that strong winds and weak stratification facilitate mixing of the water column promoting deep ventilation. In some fjords katabatic winter winds can lead cooling and reoxygenate subsurface waters for longer time like in Blue Inlet, British Columbia (Bianucci et al 2024).
80 This type of wind interaction, down-fjord winds, can drive upwelling that exposure deep water to the air-sea interface (Klymak et al., 2025).

Chilean Patagonia is one of the most extensive fjord regions of the world; the shape of this region was formed by the combined effect of glacial erosion since the Quaternary and the tectonic sinking of the central valley (Aracena et al., 2011). The region is formed by more than 100 thousand km of coastline and 40 thousand islands, which generate a complex system of fjords and channels as unique marine ecological hot spots (Hucke-Gaete et al., 2023; Landaeta et al., 2023). This system of channels and fjords in Chile between 41°S and 56°S has been geographically classified into three zones: the northern (41 - 46°S), the intermediate (46-50°S), and the southern (53-56°S) Patagonia (Pickard and Stanton, 1980). Early studies of the Chilean fjords indicated strong parallels with British Columbia fjords (Pickard, 1971; Farmer and Freeland, 1983).

Observations from the last four decades indicate that the predominant zonal (east-west) winds in Southern Patagonia have strengthened at a rate of 0.2-0.3 m s⁻¹ per decade (Garreaud et al., 2013; Giesecke et al., 2021). This strengthening is consistent with an increase in rainfall at a rate of 200 mm per decade in areas south of latitude 50°S (Garreaud et al., 2013). In the region, the Southern Annular Mode (SAM) is related with the generation of the westerlies (Aguayo et al., 2019). The SAM drives the formation of regional westerlies and has shifted toward its positive phase in response to climate change (Garreaud et al., 2013; Aguayo et al., 2019). This positive phase has been connected to an increase to 30% of the westerly winds since 1950 (Downes et al., 2017). El Niño-Southern Oscillation (ENSO) is another significant interannual climate pattern. During El Niño events, a reduction in freshwater inflow is associated with greater vertical mixing and the advection of oceanic waters into northern Patagonia's fjord systems (León-Muñoz et al., 2018).

The global increase in temperature (IPCC, 2021) suggests a significant alteration in the global heat budget and affects the retreat of ice sheets and mountain glaciers (Church et al., 2013). Interaction between glaciers and atmosphere can generate a local wind circulation known as katabatic winds (Spall et al., 2017; Bianucci et al 2024). These winds are a common characteristic of fjords adjacent to glaciers where they can exceed 20 m s⁻¹ (Farmer and Freeland, 1983; Stigebrandt, 2012; Spall et al., 2017). Along-channels winds may also advect heat and modulate the supply of warmer waters, thereby promoting glacial retreat (Moffat, 2014). The loss of glacial fields in southern Patagonia has been estimated at 10% (Rivera et al., 2017). In addition, within this framework, accelerated glacial melting and the subsequent freshening of surface waters directly impact vertical stratification (e.g. Boone et al., 2018). These alterations in freshwater input, combined with shifts in wind-stress intensity (which modulates mixing), could substantially modify the dynamics of sub-polar systems such as the Strait of Magellan and the inner sea of Tierra del Fuego. The southern Patagonia has two major ice-field glacial: the northern and the southern ice-field (Garces et al., 2026), a smaller glacial formation is located at the study region the Darwin Cordillera (Fig. 1). In these regions, katabatic flow plays an important role in defining on-glacier temperatures for valley glacier (Bravo et al., 2019). Thus, the increase of glacial melting could diminish the glacial area impacting in the katabatic flow intensity.

While the effects of density and tidal driven circulation on estuarine dynamics are well-documented (e.g. Farmer and Freeland, 1983; Stigebrandt, 2012; Geyer and MacCready, 2014), wind-driven circulation has received comparatively less attention

(Soto-Riquelme et al., 2023). Investigations of wind-driven circulation and its effects on the inner-sea of the Chilean Patagonia have been conducted in specific systems within northern and central Patagonia. These include the Reloncaví fjord (e.g. Valle-Levinson et al., 2007; Castillo et al., 2012, 2017), the Reloncaví sound (Letelier et al., 2011), the inner-sea of Chiloe (Soto-Riquelme et al., 2023; Lindford et al., 2024), the Guafo mouth (Ross et al., 2025), the Moraleda channel (Valle-Levinson and Blanco, 2004), the Aysen fjord (Caceres et al., 2002) and Jorge Montt Glacier (Moffat, 2014). Regional Patagonian wind patterns were analysed by Perez-Santos et al. (2019). At this spatial scale, atmospheric rivers have been identified as important factors controlling the regional wind effects (García-Santos et al., 2025). However, although this region is characterized by intense winds and strong tidal modulation (Antezana, 1999; Garreaud et al. 2013; Brun et al., 2020), the effects of the wind on the physical dynamics of the inner-sea of Tierra del Fuego remain poorly understood.

The intermediate zone (46-50°S) receives significant freshwater input within the Penas Gulf, originating principally from glacial discharge. Here, glacier is the primary source, and this freshwater contribution has been reinforced in recent years. The resulting freshwater interacts with the Cape Horn Current (CHC), causing its advection out of the Gulf. While the magnitude of these salinity anomalies decreases, they remain detectable, extending south of 50°S and north of 46°S (Cisternas et al., 2026).


The Southern Patagonian region is strongly affected by the Cape Horn Current (CHC), which is recognized as a component of the Antarctic Circumpolar Current (ACC) system (Lamy et al, 2015; Wu et al, 2019). Nevertheless, knowledge of the CHC's strength and variability remains limited (Zheng et al., 2023). Observational studies report current velocities exceeding 15 cm s⁻¹ (Chaigneau and Pizarro, 2005), while nearshore measurements show velocities greater than 30 cm s⁻¹ (Giesecke et al., 2021). Furthermore, an analysis combining model output and altimetry data calculated a CHC transport of 4.36 Sv. This value is comparable to the transport of the Humboldt Current System (HCS) between 5°S (1.8 Sv) and 15°S (5.2 Sv) (Chaigneau et al., 2013). More recently, research combining model output and in-situ observational data indicates that the CHC is largely geostrophic south of 49°S. This work also shows no clear seasonal variability with velocities of up to 0.3 m s⁻¹ in the upper 200 m of the water column (Garces-Vargas et al., 2026).

This study investigates the hydrographic response of the Almirantazgo Fjord—one of the southernmost fjords of South America, in Tierra del Fuego, Chilean Patagonia—to freshwater input from glacial melting and to along-fjord wind-stress. To describe the variability of the water column and its response to wind forcing, a six-month dataset was analysed. This dataset combines a time series of salinity, temperature, and dissolved oxygen with in-situ hydrographic data acquired from fixed CTD stations and animal-borne satellite CTD.

2 Material and methods

2.1 Study region

150 The Magellan Strait, situated in the southern Patagonia fjord region, connects the Atlantic and the Pacific oceans. Its internal dynamics govern the hydrography of the surrounding inland sea. Despite a sparse population, southern Patagonia possesses considerable economic significance for Chile. The growth of industries such as aquaculture, oil, green hydrogen, and tourism has altered land use, transport, and increased the ecological pressure on the Magellan Strait, Tierra del Fuego, and the Beagle Channel (Ariztía and Undurraga, 2025; Giesecke et al., 2024).



155 **Figure 1:** Study region. At the left side, a regional map of the interest region shows the main currents systems: the Humboldt Current System (HCS), the Cape Horn Current (CHC) and the Antarctic Circumpolar Current (ACC). At the right, the complexity of the Southern Patagonia it is shown, here bathymetry and topography are sowed in colorscale. Geographic features referenced in the text are identified, including Magellan Strait (MS), Inutil Bay (IB), Whiteside Channel (WC), Almirantazgo Fjord (AF), Brookes Fjord (BF), Ainsworth Bay (AB), Parry Fjord (PF), and Maria Cove (MC). The Darwin Cordillera is denoted by white-shaded high-altitude areas on the island of Tierra del Fuego. Depths and highs data were downloaded from ©GEBCO Compilation Group (2025).

Large freshwater inputs from riverine and glacial sources significantly affect the region (Sassi and Palma, 2017). The low salinity of the Magellan Strait results from a north-eastward flow from the Patagonian Current (Brun et al., 2020). The distribution of these waters is governed by the combined effects of wind, tides, and the Antarctic Circumpolar Current (Palma and Matano, 2012). Previous research indicates that waters from the Atlantic Ocean extend their effect westward through the Magellan Strait as far as Punta Arenas. In contrast, the effect of the Pacific Ocean extends eastward to Carlos III Island, where a shallow sill limits its influence on surface waters (Antezana, 1999; Brun et al., 2020).

170

An important geographical feature is the Darwin Cordillera (DC) mountain range which extends 200 km from west to east along the southwestern peninsula of Tierra del Fuego Island (Fig. 1). The DC is the core of the southern westerlies and experiences low precipitation. The region is characterized by a strong W-E temperature gradient (Carrasco et al., 2002; Garreaud et al., 2013; Meier et al., 2018; Izagirre et al., 2024). Near the study region (Fig. 1c), the glaciers of Parry fjord have been relatively stable since 1986, but the Darwin glacier retreated 3 km between 1986 and 2014. In this period, the region's glaciers have lost 10% of their area (Rivera et al., 2017).

The Almirantazgo Fjord (AF) is 75 km long, with a width of 16 km in the northern part and 5 km near its head with an averaged width of 12 km. The AF axis is 116° from the true north, an approximately East-Southeast (ESE) direction. It contains three major glacial connections: Brookes fjord (BF), the Ainsworth bay (AB), and Parry fjord (PF). At the head of the fjord, in Maria Cove, the fjord is shallow, with a mean depth of approximately 30 m for the first 7 km. The bottom then slopes steeply until the mouth of PF, deepening from 30 m to 200 m in less than 3 km. From PF to BF, the fjord depth is approximately 220 m. The deepest part of this basin is in the Whiteside channel (WC), where the bottom depth is approximately 450 m. This is one of the deepest parts of the inner sea of Tierra del Fuego (Fig. 1).

185 2.2 Atmospheric data

Meteorological data for the region were obtained from the Dirección General de Aeronáutica Civil (<https://climatologia.meteochile.gob.cl/>), sourced from airports at Punta Arenas (53.00°S, 70.84°W), Porvenir (53.25°S, 70.33°W), and Puerto Williams (54.93°S, 67.62°W). The dataset included wind magnitude and direction (meteorological convention), recorded at 1-hour time intervals between 1960 and 2025.

190

Additionally, to characterize the wind patterns, this study uses wind reanalysis data from ERA 5. The wind components at 10 m high (u_{10} , v_{10}) were downloaded in spatial grids of 0.25° x 0.25°. The ERA5 is the fifth-generation ECMWF reanalysis of global climate and weather data. Reanalysis integrates model data worldwide observations into a globally complete and consistent dataset on physical laws (Hersbach et al., 2023).

195

Wind-stress (τ) was calculated using the bulk formula by $\tau = \rho_a c_d |U_{10}| U_{10}$, where ρ_a is the air density (1.2 kg m^{-3}), U_{10} is the wind velocity vector (with components u_{10}, v_{10}), $|U_{10}|$ is wind speed (the scalar magnitude of U_{10}) and c_d is dimensionless wind-drag coefficient. The coefficient c_d was calculated for each wind component value in the time-series following Yelland and Taylor (1996), for $|U_{10}| < 6 \text{ m s}^{-1}$, $c_d = (0.29 + 3.1 |U_{10}|^{-1} + 7.7 |U_{10}|^{-2}) \times 10^{-3}$, whereas, for $6 \text{ m s}^{-1} \leq |U_{10}| \leq 26 \text{ m s}^{-1}$, $c_d = (0.60 + 0.07 U_{10}) \times 10^{-3}$.

2.3 Sea Level data and tides model

Sea level data were acquired from four tide gauge stations located within the inner sea of the Magellan Strait and the Almirantazgo Fjord (Fig. 1): Gregorio Cove, Punta Arenas, Puerto Williams, and Ushuaia (Table 1). The dataset was obtained from the Sea Level Station Monitoring Facility (<https://www.ioc-sealevelmonitoring.org/>) for the period of January to June 2024. At each station, measurements are recorded at a one-minute interval by a pressure sensor positioned near the seabed.

Additionally, a barotropic tidal global model TPX09 (Egbert and Erofeeva, 2002) was applied to determine the spatial distribution of amplitude and tidal current ellipses in the region (Fig. 3). The model provides complex tidal heights (h) and transport coefficients for 15 constituents at $1/30$ -degree resolution. The principal harmonic amplitudes (M_2 , N_2 , S_2 , O_1 , and K_1) were used to estimate the form factor (e.g. Pan et al. 2024) defined as $F = (O_1 + K_1)/(M_2 + S_2)$ was useful to characterize the regimes of the barotropic tide in the inner sea of Tierra del Fuego.

2.4 Hydrography data

The hydrography data were obtained using two different complementary methods: conventional CTDOF (AML Metrec X) deployments and CTDF (CTD-SRLD) casts from sensors affixed to Southern elephant seals.

The conventional instrumentation consisted of a CTD AML–Metrec X equipped with Temperature, Conductivity, Pressure, Fluorescence/Chlorophyll, and pH sensors, supplemented by an additional Aanderaa Optode 4831 infrared Dissolved Oxygen sensor. The CTDOF was configured for freshwater settings to permit measurements under low-conductivity conditions, with profiles acquired at a rate of four scans per second. At each oceanographic station, the instrument was activated on deck and subsequently lowered to 1 m below the surface, where it was held for one minute to ensure sensor stabilization. Data acquisition was then performed using an electrical winch to a depth up 200 m. To describe the oceanographic conditions, sampling was conducted at five cross-fjord oceanographic stations (Fig. 1). Data were collected during the austral summer (January 15th and 31st, 2024) and the austral winter (June 26th and 30th, 2024). A total of 60 cross-fjord profiles were acquired to characterize the oceanographic conditions near the head of the Almirantazgo Fjord. Post-processing of the AML-CTD data followed standard quality control procedures, consistent with other CTD instruments. This process included: visual inspection to remove spikes, exclusion of sensor stabilization time, selection of downward profiles, and vertical binning at 0.5 m intervals. Subsequently, mean profiles for January and June were calculated for each station and are presented in Figure 6.

Oceanographic conditions in the region were also assessed using CTD-SRLD (Bohemme et al., 2009) and CTD-Fluoro tags (Guinet et al., 2013). The CTD tags possess high accuracy ($\pm 0.005^{\circ}\text{C}$ for temperature, ± 0.01 mS/cm for conductivity, and $2 \text{ dBar} \pm (0.3 + 0.035\% \cdot \text{reading})/^{\circ}\text{K}$ for depth) and were calibrated by the manufacturer prior to deployment. These tags were affixed to Southern elephant seals (*Mirounga leonina*) from the colony located at Jackson Bay, located at the head of the Almirantazgo Fjord (Fig. 1c). *In-situ* temperature and salinity measurements were converted to Absolute Salinity (S_A), Conservative Temperature (Θ) following the TEOS-10 (IAPWS, 2010).

235

Water column stratification was characterized by the buoyancy frequency (N^2), calculated as $N^2 = -g/\rho_0 \partial\rho/\partial z$, where g is the gravitational acceleration, ρ_0 is the reference density, and $\partial\rho/\partial z$ is the vertical density gradient. Additionally, CTD data were used to estimate the energy required to homogenize the water column, expressed as the Potential Energy Anomaly (PEA), $PEA = 1/h \int_{-d}^0 g z (\rho_p - \rho) dz$, where h is the thickness for the calculation of average density (ρ_p). A high PEA value corresponds to high stratification (Simpson et al., 1981). The Freshwater Content (FWC), defined as the equivalent thickness of freshwater in the water column, was also calculated, $FWC = \int_{-d}^0 (s_0 - s) / s_0 dz$, where s_0 is the reference salinity at depth d (Blanton and Atkinson, 1983), here the study used $s_0 = 31 \text{ g kg}^{-1}$ which was the maximum salinity in the deeper zone of the Almirantazgo fjord. In all the profiles FWC was calculated by integrating over the entire water column down to the bottom depth.

245 To describe the hydrographic along-fjord patterns for January and June 2024, this study utilizes CTD-SRDL data collected between Inútil Bay (IB) and María Cove (MC). Although the elephant seals provided numerous hydrographic profiles across the region, the study defined an along-fjord route along the central axis between IB and MC (see Fig. 5a). This route consisted of points spaced at 0.1 km intervals over a 150 km span. All profiles within 2 km of each point, in a temporal window of 3-weeks centred at the CTD-AML dates (see Table 1), on the route were selected and averaged to obtain a single representative profile per point. For the vertical dimension, standard depths were established at every 1 m (0–20 m), 5 m (20–50 m), 10 m (60–100 m), and 50 m (150–500 m). Vertical interpolation was performed only at these standard depths, avoiding spatial (distance) along-fjord interpolation. This procedure was applied to the January and June 2024 data shown in Fig. 5c,d.

To assess the influence of different water masses, an Absolute Salinity of 31 g kg^{-1} was used as an index defining the upper boundary of the Modified Subantarctic Water (MSAAW). This water mass is produced by the combination of surface freshwaters with Subantarctic Water (SAAW), the dominant water mass within the region (Silva et al., 2009; Silva and Vargas, 2014).

2.5 Time series: observations and data analysis

Time series data of the water column properties were acquired using two distinct but complementary ways. The first method comprised a six-month (January-June 2024) near-bottom (30 m depth) record of sea level, salinity, temperature and dissolved

260

oxygen. The second method involved short-term (5-day) acquisitions of a fine vertical-scale temperature, salinity and dissolved oxygen during January 2025 (Fig. 2).




Figure 2: Location of the oceanographic stations at the head of the Almirantazgo fjord. The caption shows the location of oceanographic stations and mooring deployments within the innermost part of the Almirantazgo Fjord. In addition, the mooring system deployed at the sites and the instrumentation it is shown. Depths and highs data were downloaded from <https://www.gebco.net/data-products/gridded-bathymetry-data>. Bathymetric data ©GEBCO Compilation Group (2025).

The six-month time series was collected using a CTD WiSens NKE Instruments (<https://nke-instrumentation.com/>). This instrument was secured within a stainless steel, pyramid-shaped frame (1 m x 1 m; 0.7 m height) positioned on the seabed at 30 m depth near to 54.44°S, 69.05°W. The instrument specifications include a pressure range up to 300 m (0.1% accuracy), a temperature range of the -2°C to 35°C ($\pm 0.02^{\circ}\text{C}$), and salinity range 2 to 42 PSU (± 0.1 PSU). Dissolved oxygen (DO) concentration was recorded by a PMEL MiniDOT (<https://www.pme.com/products/minidot>), which functions via an optical sensor. This sensor has an accuracy of $\pm 0.2 \text{ mL L}^{-1}$ and a resolution of $7 \times 10^{-4} \text{ mL L}^{-1}$. The MiniDOT also contains temperature sensors with a range of 0°C and $35^{\circ}\text{C} \pm 0.1^{\circ}\text{C}$ and its housing is rated for 300 m depth. Both the Wisens NKE and MiniDOT sensors were affixed in individual stainless-steel frames attached to the main pyramid structure. This entire assembly was lowered to the bottom and remained moored until retrieval at the end of June 2024.

280 The 5-day time series was acquired from a mooring at A2 station (Fig. 2, Table 1) between January 22nd and 26th of 2025. This instrument array was designed to measure the vertical gradient of temperature, salinity, DO, and pressure. The array included CTD NKE WiSens and a miniDOT positioned at 0.3 m depth and at 30 m depth (near the bottom) to determine the vertical gradients of salinity ($\partial S/\partial z$), temperature ($\partial T/\partial z$), and dissolved Oxygen ($\partial DO/\partial z$).

285 To characterize the temporal variability and dominant oscillations observed in the data, a Morlet wavelet analysis (Morlet et al., 1982; Torrence and Compo, 1998; Grinsted et al., 2004) was applied to the time series of wind stress, sea level, Conservative Temperature, Absolute Salinity, and Dissolved Oxygen. This method permits the computations of both the dominant modes of variability and their temporal variations (Torrence and Compo, 1998). To quantify the relationship between the wind stress and the other variables, wavelet coherence was estimated following the methodology of Grinsted et al. (2004).

290

Table 1. Location and descriptions of the sea level stations, hydrographic (oceanographic) stations, and time series data used in the study.

	Station	Latitude (S)	Longitude (W)	Dates	Time interval
Winds	Punta Arenas	53° 00' 00"	70° 50' 24"	Jan 01/1960 – Jan 31/2025	1h
	Porvenir	53° 15' 00"	70° 19' 48"	Jan 01/1960 – Jan 31/2025	1h
	ERA 5 pixel	54° 11' 00"	70° 00' 40"	Jan 01/2024 – Jan 31/2025	1h
	Puerto Williams	54° 55' 48"	67° 37' 12"	Jan 01/2024 – Jan 31/2025	1h
Sealevel	Gregorio Cove	52°35'35.54"	70° 9'44.23"	Jan 01 – Jun 30 /2024	1 min.
	Punta Arenas	53°10'16.25"	70°54'17.54"		
	Puerto Williams	54°55'57.12"	67°36'29.05"		
	Ushuaia	54°49'1.2"	68°13'1.2"		
	María Cove	54°26'31.87"	69° 3'6.01"	Jan 15 – Jun 30/2024	10 min.
hydrography	CTD 1	54°23'51.9"	69°08'10.4"	summer:	0.25 s
	CTD 2	54°24'48.6"	69°09'38.9"	Jan 15 – 31/2024	
	CTD 3	54°25'45.6"	69°10'48.1"		
	CTD 4	54°26'32.6"	69°11'37.1"		
	CTD 5	54°26'50.7"	69°11'57.8"	Jun 26 – 30/2024	
Time	A1	54°26'31.87"	69° 3'6.01"	Jan 15 – Jun 26/2024	10 min.
	A2	54°27'54.11"	69° 0'18.36"	Jan 20 – 26 /2025	5 min.

series	Azopardo River discharge	54° 30' 10''	68° 49' 33''	Jan 01 /2024 – Jan 31/2025	30 min.
---------------	-----------------------------	--------------	--------------	----------------------------	---------

295 The potential for the wind stress to perturb the pycnocline was assessed using the dimensionless Wedderburn number (W_b) following Geyer (1997), Thorpe (2005) and Inall et al. (2015). The number is defined as: $W_b = (\tau L) (\Delta \rho g)^{-1} h_1^{-2}$, where L is the length of the fjord (or the horizontal scale of the wind), g is gravitational acceleration, and h_1 is the depth of the upper layer directly influenced by the wind stress τ . In a simplified two-layer model, the density difference $\Delta \rho = \rho_2 - \rho_1$ where ρ_1 and ρ_2 are the densities of the upper and deeper layer, respectively. According to this formulation, a value of $W_b < 1$ indicates that the

300 density gradient dominates. When $W_b = 1$, wind forcing and buoyancy effects are comparable, and horizontal mixing becomes significant. When $W_b > 1$, wind-driven effects are dominant (Inall et al., 2015).

3 Results

3.1 Wind pattern: seasonality of the zonal and meridional components

The 54-years hourly reanalysis dataset was used to establish the wind climatology for the study region (Fig. 2). The results

305 indicate that seasonal winds exhibit high directional consistency. Westerlies are the dominant pattern; south of 54°S in the Pacific Ocean, these winds acquire a slight southerly component, with the airflow generally following the coast. Wind magnitudes intensify near the coast, particularly during the warmer seasons (austral spring and summer), when speed exceeds 8 m s^{-1} (Fig. 2a, 2b). During the colder seasons (austral autumn and winter), regional winds are weaker ($< 6 \text{ m s}^{-1}$), particularly offshore in the Pacific Ocean. Within the inner sea of the Magellan Strait and Almirantazgo Fjord, wind magnitude was low

310 ($< 4 \text{ m s}^{-1}$), comparable to those observed in the open ocean. Regarding direction, winds within the Almirantazgo Fjord appear aligned with the fjord's axis and generally flow toward the fjord head throughout all the seasons (Fig. 2).




Figure 3: Climatology of winds in southern Patagonia. Mean wind vector are shown for the periods: a) JFM (January-March, 315 austral summer), b) AMJ (April-May, austral autumn), c) JAS (June-September, austral winter), and d) OND (October-December, austral spring). Data were derived from hourly $0.25 \times 0.25^\circ$ gridded ERA5 reanalysis data covering the period 1970–2024. ERA5 reanalysis data ©ECMWF/Copernicus.

The climatology from the meteorological stations indicates a high consistency with the regional winds described previously 320 (see Fig. A1). At Punta Arenas (PA), Porvenir (PV) and Puerto Williams (PW), eastward winds (meteorological westerlies) were predominant ($> 25\%$) during all the seasons. The highest wind speeds (c.a. 15 m s^{-1}) were observed at PA and PV during summer and spring. PV was an exception, as northward/southward winds were also recorded there during autumn and winter. For the Almirantazgo Fjord (AF) wind data were derived from a time series of an ERA5 reanalysis pixel (Fig. 3). At this location, the distribution of magnitude and direction was highly consistent with the regional patterns, showing a dominance of 325 eastward winds in all the seasons. The directional distribution indicates that up-fjord winds (directed towards the fjord's head) are prevailing conditions in AF (see Fig. A1).

330 **Table 2.** Main tidal harmonics amplitude and form parameter (F) in the sealevel stations of the study region.

Station	Amplitude (cm)					F
	M ₂	N ₂	S ₂	K ₁	O ₁	
Gregorio cove	132.67	39.08	26.45	29.23	25.66	0.33
Punta Arenas	48.06	9.24	25.24	27.41	22.39	0.67
Puerto Williams	55.13	16.41	7.04	18.68	16.30	0.56
Ushuaia	56.00	14.0	6.00	20.00	16.00	0.58
María cove	59.33	12.90	28.68	30.50	21.70	0.59

3.2 Sea level and Tidal influence on the region

335 Tidal amplitudes recorded by the region’s tide gauges (Fig. 1, Table 2) were primarily controlled by the M₂ semi-diurnal harmonics. The Gregorio station measured an M₂ amplitude of 132.7 cm, a magnitude at least double that observed at the other stations (which ranged from 48 cm to 56 cm) located south of the Magellan Strait. At the Punta Arenas station, the M₂ amplitude was 48.1 cm. In the Beagle Channel, south of the Almirantazgo Fjord, the Puerto Williams and Ushuaia stations recorded comparable M₂ amplitudes of approximately 55 cm. The lowest F value (0.33) was observed at Gregorio, whereas at the other stations, F> 0.6. The form factor (F) indicated that all the stations are characterized by a mixed, mainly semi-diurnal tidal regime.




Figure 4. Results based on the tidal model by Egbert and Erofeeva (2002) shown the regional distribution of the principal harmonic constituents: a) M_2 , b) N_2 , c) S_2 , and d) K_1 . Amplitude (m) is represented by color shading, while phase (°) is shown as contour lines. In the caption, the locations of the First (1stN) and the Second Narrows (2ndN) within Magellan Strait are indicated. Tidal model data © Oregon State University (TPX09; Egbert and Erofeeva, 2002).

To quantify the tidal influence in the study region, the barotropic tidal Model by Egbert and Erofeeva (2002) indicated that the northern part of the Magellan Strait and its connection with the Atlantic Ocean showed the highest M_2 amplitude (> 3 m). Amplitudes decreased southward, falling below 1 m off Punta Arenas and reaching approximately 0.5 m inside the Almirantazgo Fjord (Fig. 3). A marked change in amplitude (from high to lower) was observed at the Second Narrow, phase tends to increase from the Atlantic (10°) to the inner-sea reaching 90° into the Inutil Bay. The entire basin from the Inutil Bay to Maria Cove seems to be in phase at 90°. Other harmonic constituents were substantially smaller than M_2 accounting for < 10 % of the variability explained by the M_2 component (Fig. 4).

3.3 Temperature and Salinity patterns inside the Almirantazgo Fjord

The whole CTD-SRLD dataset (showed in Fig. A2) acquired from Southern elephant seals encompassed extensive areas, including the Pacific and Atlantic oceans and the inner sea of the Magellan Strait and Tierra del Fuego. To specifically address the dynamics of the Almirantazgo Fjord, the present analysis utilized only the CTD-SRLD data obtained from within the fjord.

The hydrographic data reveal two distinct regimes for the waters of southern Patagonia (see Fig. A1d). Within the inner channels of Tierra del Fuego, low absolute salinity (c.a. 31 g kg^{-1}) was observed in deeper waters (depths $> 100 \text{ m}$). Conversely, waters in the outer channels (in the Pacific and Atlantic oceans) were characterized by salinities exceeding 34 g kg^{-1} . These findings indicate that the inner channel waters are more stable (stratified) compared to the mixed conditions prevailing in the outer channels.




Figure 5. a) Locations of the hydrographic profiles of Absolute Salinity (S_A) and Conservative Temperature (CT) acquired via CTD-SRLD (Southern elephant seals) and CTD-AML. CTD-SRLD data points within Bahia Inutil, Whiteside Channel, and Almirantazgo Fjord are marked in red. b) CT/ S_A diagram the complete dataset (red dots) whereas in blue dots are the January and June elephants seals data showed in c) and d). Segmented black lines are isopycnals of conservative density anomaly (σ_{CT})

in kg m^{-3} . c) January and d) June vertical sections of CT and SA for each month. The along-fjord blue dots in panel (a) indicates the along-fjord transect. Average profiles of Brunt-Väisälä frequency (N^2), S_A , and CT for the entire transect are displayed to the right of the lower panels (c) and (d).

370

To compare and describe the seasonal patterns of the Almirantazgo Fjord, this study utilized CTD-SRLD data acquired from Southern elephant between January and June of 2024. The analysis was restricted to Conservative and Absolute Salinity measurements collected within the region extending from the Whiteside channel and Almirantazgo Fjord (Fig. 5).

375 The Conservative Temperature/Absolute Salinity (CT/ S_A) diagram (Fig. 5) acquired by elephant seals, illustrates the wide hydrographic variability within Almirantazgo Fjord. S_A ranged from a minimum of 26.28 g kg^{-1} near the surface (4 m depth) to a maximum of 31.13 g kg^{-1} near the fjord bottom. CT varied from 4.24°C to 10.94°C with a mean of $7.20 \pm 1.12^\circ\text{C}$. The conservative density anomaly (segmented lines, Fig. 4b) extended from 1020.0 kg m^{-3} in the surface waters during January to a maximum of 1024.5 kg m^{-3} in near-bottom waters during June. The mean CT and S_A profiles (Fig. 5c, d) show pronounced
380 upper-water-column stratification in summer, contrasting with the well-mixed conditions observed in winter.

The cross-fjord transect data were acquired using a CTD-AML Metrec during field measurements in January and July 2024. During the austral summer, S_A ranged from a minimum of 26.30 g kg^{-1} near the surface to a maximum of 30.99 g kg^{-1} near the bottom, with a seasonal average of $30.21 \pm 0.10 \text{ g kg}^{-1}$. Water temperature varies between 6.132°C and 10.81°C , with a mean
385 of $7.68 \pm 0.03^\circ\text{C}$.

During winter (June 2024), conditions in the Almirantazgo Fjord (AF) shifted. S_A ranged from 29.01 to 31.01 g kg^{-1} with a mean of $30.78 \pm 0.04 \text{ g kg}^{-1}$. The water column was colder, with CT between 2.97°C and 7.74°C (mean: $7.01 \pm 0.06^\circ\text{C}$). A thermal inversion was observed, with colder upper waters overlying warmer deeper waters. Density range was 1018.8 kg m^{-3}
390 to 1024.5 kg m^{-3} . Mean cross-fjord profiles indicated that stratification was salinity-dominated. In summer, continuously stratified conditions were present in the upper 30 m depth layer. In winter, however, the S_A profile was nearly homogeneous throughout the upper 100 m of the water column.

Based on the along-fjord and cross-fjord CTD sections and mean profiles (Figs. 5c,d and 6c,d), the study determined that
395 during January, the thickness of the upper brackish layer near the head of the fjord (at PF) was $h_1 = 20 \text{ m}$, compared to $h_1 = 5 \text{ m}$ during June. At the mouth of Almirantazgo Fjord, the upper layer depth was estimated to be $h_1 = 5 \text{ m}$ during both months. These observations are consistent with a salt-wedge structure in the upper layer during January that was absent during June (Figs. 5 and 6).




Figure 6. a) Location of the hydrographic profiles (Absolute Salinity, S_A ; Conservative Temperature, CT) acquired via CTD-AML. b) CT/ S_A diagram for the CTD-AML dataset. Segmented black lines represent conservative density anomaly (σ_{CT}) isopycnals in kg m^{-3} . Summer and d) Winter cross-fjord sections of S_A and CT, extending from station CTD1 (northern side) to CTD 5 (southern side) the cross-fjord stations are present in yellow circles in (a). Average vertical profiles of CT and S_A for the entire transect are displayed to the right of panels (c) and (d).

405

Both sets of hydrographic data (along-fjord and cross-fjord CT and S_A profiles) were used to estimate the Freshwater Content (FWC) and Potential Energy Anomaly (PEA) for January (summer) and June (winter) (Fig. 7). During January, the FWC exhibited a gradient, decreasing from the fjord head toward the Magellan Strait (MS). At 3.9 km from the head, which is near Parry Fjord (PF), the FWC was 1.37 m then FWC decrease trough the fjord's mouth presenting peaks events (c.a. 0.6 m) near of Brook's Fjord (BF) at 71 km from the head reaching a minimum FWC of 0.14 m near the connection with the Magellan

410

Strait (Fig. 7a). The along-fjord structure of PEA closely resembled that of the FWC, displaying a maximum of 157.2 J m^{-3} near PF and secondary maxima at 14 km (c.a. 152.5 J m^{-3}) and BF (c.a. 71.3 J m^{-3}). In June (winter), both FWC and PEA were lower and noisier with lower values (FWC $\sim 0.2 \text{ m}$, PEA $\sim 7 \text{ J m}^{-3}$) around BF location (70 km from the head) (Fig. 7b).




Figure 7. Distributions of Freshwater Content (FWC) and Potential Energy Anomaly (PEA) for January (dashed red line) and June (solid blue line) shown for the (a, b) along-fjord and (c, d) cross-fjord transects. Relative distance from the fjord head (along-fjord) and from the northern side (cross-fjord) is indicated with segmented thin black lines.

The cross-fjord FWC and PEA values were consistent with the along-fjord estimations, as summer values exceeded those recorded in winter. In summer, FWC ranged from 1.8 m (north side) to 2.1 m (south side), displaying a slight tilt toward the southern portion of the transect near PF (Fig. 6c). PEA values reached 200 J m^{-3} and exhibited a cross-fjord tilt analogous to that of the FWC. During the winter, FWC was less than 1.0 m and showed only a minimal cross-fjord tilt to the south. The winter cross-fjord PEA was $< 40 \text{ J m}^{-3}$ a value approximately five times lower than observed in summer (Fig. 7d).

3.4.1 Temporal variability: Summer to Winter Transition

During the austral summer 2024, instrumentation measuring sea level, temperature, salinity, and oxygen was deployed at a fixed station within the Almirantazgo Fjord, Tierra del Fuego. These sensors were subsequently retrieved during fieldwork conducted at the end of June 2024. The winds-stress time series presented in Fig. 8 covers from January to June, permitting a

comparison of its variability with the time series of sea level, temperature, dissolved oxygen, and salinity from the A1 mooring (Fig. 1). The following description is based on the data shown in Fig. 8.

435 The regional wind pattern indicated that winds persistently blew toward the fjord head, and the zonal (east-west) component was dominant across the seasons (Fig. 3, Fig. A1). Therefore, the wind-stress magnitude was analysed to characterize the variability and intensity of the winds within the fjord. The wind-stress magnitude recorded a minimum of $2 \times 10^{-4} \text{ N m}^{-2}$ (August 6th) and a maximum of $1.59 \times 10^{-1} \text{ N m}^{-2}$ (July 23rd); both extremes occurred during the austral winter. The mean wind-stress for the time-series was $3.09 \pm 2.26 \times 10^{-2} \text{ N m}^{-2}$ and the 90th percentile was $6.24 \times 10^{-2} \text{ N m}^{-2}$. Seasonal differences were apparent: the mean summer (December- February) wind-stress was $3.80 \pm 2.26 \times 10^{-2} \text{ N m}^{-2}$ which was higher than the mean
440 winter (June to August) wind-stress $2.28 \pm 2.16 \times 10^{-2} \text{ N m}^{-2}$.

The wind stress time series displayed a combination of high-frequency (c.a. 1 day) and synoptic (c.a. 5 days) oscillations from late January (summer) to mid-April (early autumn) 2024. During this summer and early autumn phase, wind stress events exceeding 0.1 N m^{-2} occurred in early February and late March. The wind pattern showed a seasonal transition from mid-April
445 to the end of June. During this late period, oscillations were predominantly synoptic (c.a. 5 to 7 days). No event surpassed 0.1 N m^{-2} , although notable events > 0.07 were recorded in the mid-May and mid-June 2024 (Fig. 8a).

The Azopardo river discharge time series for January-June contains a data gap between April 15th and May 7th (Fig. 7). The discharge was higher in summer ($48.23 \text{ m}^3 \text{ s}^{-1}$) and lower during winter ($32.76 \text{ m}^3 \text{ s}^{-1}$), with a mean of $40.50 \pm 4.47 \text{ m}^3 \text{ s}^{-1}$. The
450 data exhibited a pronounced negative seasonal trend from summer to winter, declining at a rate of $-0.10 \text{ m}^3 \text{ s}^{-1} \text{ d}^{-1}$. High frequency variability was evident throughout the record; like the wind stress, daily and synoptic oscillations seemed dominant (Fig. 8b).

Sea level (SL) was derived from the pressure sensor of the NKE WiSens, SL height was referenced to the minimum recorded
455 value to isolate amplitude and oscillations. Harmonic analysis indicated the dominance of the M_2 constituent (0.59 m), followed by K_1 (0.31 m), S_2 (0.29 m), O_1 (0.22 m), and N_2 (0.13 m). The form factor ($F = 0.593$) confirms a mixed, predominantly semi-diurnal tidal regime. This tidal forcing accounted for 76% of the total SL variability. An air pressure time series was not available to calculate adjusted sea level; the residual (non-tidal) variability was clearly of synoptic origin. The maximum tidal range was less than 3 m during spring tides and approximately 1 m during neap tides. Between February and June, 11 spring
460 and 10 neap periods were observed; however, the spring tides from February to late March were less pronounced than those in autumn and early winter (Fig. 8c).




Figure 8. Time series of: a) wind-stress magnitude; b) Azopardo River discharge; c) sea level; d) near-bottom (30 m depth) Conservative Temperature (red line) and Dissolved Oxygen (green line); and e) near-bottom (30 m depth) Absolute Salinity (blue line). Data for panels (c), (d), and (e) were recorded at A1 station in Maria Cove. Segmented lines on discharge, temperature, dissolved oxygen, and salinity indicate the seasonal trend for each time series. Notice that warmer events were marked with shading red, transition with shading grey and colder with shading blue.


470

475 Conservative Temperature (CT) displayed a pronounced negative seasonal trend of $-0.009^{\circ}\text{C d}^{-1}$ from summer to winter (Fig. 8d). The maximum CT (10.23°C) was observed during summer, while the minimum (4.98°C) occurred in winter. The time series exhibited a mean of $7.94 \pm 0.77^{\circ}\text{C}$. This seasonal signal was evident in the higher mean CT during summer ($8.08 \pm 0.73^{\circ}\text{C}$) compared to winter ($7.19 \pm 0.86^{\circ}\text{C}$). The record was characterized by short-duration thermal events: warmer during summer (February to end of March) and colder during autumn-winter (April to June). Similarly, Dissolved Oxygen DO, 480 showed a negative seasonal trend ($-0.001 \text{ mL L}^{-1} \text{ d}^{-1}$) over the same period (Fig. 7d). The mean DO was $6.9 \pm 0.2 \text{ mL L}^{-1}$ in summer and $6.8 \pm 0.5 \text{ mL L}^{-1}$ in winter. The time series featured high DO events that were brief (< 2 days) between February and mid-March but became more protracted (> 2 days) during autumn and winter, with extended events (> 1 week) observed from mid-May to June.

485 The S_A time series recorded a minimum of 27.56 g kg^{-1} (summer) and a maximum of 30.96 g kg^{-1} (winter) during the sampling period (Fig. 7e.). The mean S_A was $30.03 \pm 0.73 \text{ g kg}^{-1}$ and a distinct seasonal trend ($0.004 \text{ g kg}^{-1} \text{ d}^{-1}$) from fresher to saltier water conditions was evident between February to June. The mean summer value was $30.09 \pm 0.65 \text{ g kg}^{-1}$ compared to winter mean of $30.46 \pm 0.32 \text{ g kg}^{-1}$. Although the mean seasonal values were comparable, the variance in summer was four times greater than in winter. The low-salinity conditions in summer were associated with brief events (c.a. 1-2 days) of $< 30 \text{ g kg}^{-1}$ 490 waters, which occurred from late January to late March. From April to late June, these low-salinity events were more protracted (> 2 days), such as those in mid-May and mid-June, but remained above 29 g kg^{-1} (Fig. 8e).

3.4.2 Temporal Variability: Water Column Response to Wind Forcing

The summer-to-winter time series revealed a pronounced seasonal signal and provided evidence of a wind-driven response in the near-bottom (30 m) waters at the head of the Almirantazgo Fjord (Fig. 9). However, this single-depth dataset was 495 insufficient to resolve the vertical structure of the water column's response. Consequently, the short-term mooring was deployed in Maria Cove, near the Azopardo river outflow (Fig. 1). The mooring was instrumented to record the surface and bottom (30 m) water column response to variations during contrasting wind intensity.



500 **Figure 9.** Short time-series acquired at mooring site A2 in Maria Cove (January 20–26, 2025). Panels display, **a)** along-fjord (τ_u) and cross-fjord (τ_v) wind-stress components, **b)** Azopardo discharge, **c)** demeaned sealevel, **d)** density gradient ($\Delta \rho$), **e)** Wedderburn number (W_b) and **f)** Dissolved Oxygen gradient (ΔDO).

On January 20 and through most of January 22, the along-fjord (τ_u) and cross-fjord (τ_v) wind-stress components exhibited low
 505 magnitudes ($< |0.05| \text{ N m}^{-2}$). A τ_v event (c.a. -0.1 N m^{-2}) occurred at the end of January 22nd, marking a transition in the time

series. This event separated an initial period of weak winds (before 20:00 local time on Jan 22) from a subsequent period of increased wind forcing ($> 0.1 \text{ N m}^{-2}$) from January 23 to 25 (Fig. 8a). The Azopardo river discharge was relatively constant at approximately $44 \text{ m}^3 \text{ s}^{-1}$ during the weak wind period, decreasing slightly to $42 \text{ m}^3 \text{ s}^{-1}$ during the period of stronger winds (Fig. 8b). Regarding sea level, four distinct oscillations were observed during the weak wind period, presenting a clear semi-diurnal signal free of high-frequency noise. This pattern changed with the onset of strong winds, at which point the sea level record became characterized by high-frequency variability. During this latter period, the tidal oscillation was completely disrupted by the wind stress, especially between 00:00 to 12:00 of January 25 (Fig. 9c).


To characterize the changes in water column density, $\Delta\rho = \rho_2 - \rho_1$ was used, where ρ_2 is density at 30 m and ρ_1 is the surface density. On January 21 and early January 22, $\Delta\rho$ exceeded 10 kg m^{-3} . The gradient subsequently diminished to values near 1 kg m^{-3} and remained at this low level until the end of January 24. At that point, where $\Delta\rho$ increased and was maintained above 10 kg m^{-3} . It decreased again to approximately 1 kg m^{-3} during the end of January 25 and the beginning of January 26 (Fig. 9d).

The Wedderburn number (W_b) was calculated using $L = 75 \text{ km}$ (the basin length of AF) and $h_1 = 20 \text{ m}$, the upper depth of the brackish layer (h_1). For 96% of the deployment, events W_b remained < 1 . Events where $W_b > 1$ were observed on January 23 at 20:00. The highest values, which rose above 2 and reached a maximum of 3.77, occurred on January 24 between 20:00 and 23:00. A final event was recorded on January 25 at 02:00 (Fig. 9e).

At the beginning of the time-series (January 21 to end of January 23), the vertical Dissolved Oxygen gradient (ΔDO) was maintained with minor variability around -0.1 mL L^{-1} . Subsequently, the gradient increased to $> -0.3 \text{ mL L}^{-1}$ until mid-January 24. Following this, from mid-January 24 to mid-January 25, ΔDO was approximately 0 mL L^{-1} . The most negative gradient (c.a. -1 mL L^{-1}) was observed at the end of January 25 (Fig. 9f).

3.5 Spectral Analysis: pattern of variability near the bottom

The wind-stress magnitude exhibited high spectral amplitude for the oscillation with periods greater than 24 hours. This diurnal band showed its greatest amplitude from late January until the beginning of May, during the winter, this band was virtually absent (Fig. 10a). The subsequent period was observed in the 3-days band, which displayed high amplitude at the beginning of the time series (January 25 to February 24) and again around March 20 and mid-May. The 10-day band was the most significant period for the wind-stress magnitude. The first two weeks of February was the only time this band showed lower amplitudes; during that period, the variability was centred at 3 days. Furthermore, distinct events with high amplitudes spanning a broad range of periods (12h to 256h) were centred on specific dates: February 5, March 30, April 20, May 10, June 3, and June 14 (Fig. 10a).



540 **Figure 10.** Wavelet analysis for the time series acquired at the A1 mooring: a) wind stress, b) sea level, c) Conservative Temperature, d) Dissolved Oxygen and e) Absolute Salinity. For each variable, the wavelet power spectrum is shown (left panels), and the global wavelet spectrum is shown (right panels). In the power spectra, power is color-coded (red = maxima),

and the black contour represents the cone of influence where edge effects become important. Horizontal dashed lines denote periods of 4, 12, 24, and 72 hours.

545

The sea level (SL) data were dominated by tidal variability, with the highest spectral amplitudes occurring in the semi-diurnal (12 h) and diurnal (24 h) bands. Both bands exhibited a pronounced fortnightly modulation; this modulation became less distinct in the 12h band from mid-May until the end of the time series. Other evident variability was contained at 4 h band with low amplitude and bi-weekly modulation and the low-frequency (longer than 1-day periods) which showed marked high amplitudes at 3 days (around 72 h) at the beginning of the time-series until middle of February and beginning of April. The most important (and significant) period band was centred at 10 days (256 h) into the synoptic band. This band persists along the entire time-series but diminishes its amplitude between May and June (Fig. 10b).

550

The CT data indicated the significance of the 3-day band. High amplitudes were observed in this band during January and February; its amplitude subsequently decreased until June 3rd and 14th, when two distinct high-amplitude pulses were recorded (Fig. 9c). The global spectrum confirmed the relevance of the synoptic band, centred at 10 days (approx. 256 h). The wavelet power spectrum shows that this 10-day band was less prominent during March and April (Fig 10c).

555

The DO wavelet exhibited a pattern analogous to that of CT at the start of the time series, characterized by high variability within the 3-day band. However, unlike CT, the DO data displayed high amplitudes across the 3-day to 10-day bands between March and April; during this interval, the 5-day band appeared to intensify around March 20th and April 24th. Throughout the record, multiple high-amplitude events occurred across various periods (Fig. 10d).

560

For S_A , the time series from January 23rd to March 5th was characterized by low spectral amplitudes across all periods resolved by the wavelet analysis. Following this, from March to June, high amplitudes developed at periods greater than 3 days. Within the 5- to-10 day, S_A variability initially exhibited high amplitudes centred at 5 days (late April). From mid-May until the end of the record, the dominant variability shifted to the 10-day band. High-amplitude events spanning all periods > 24 h were observed centred on June 3rd and June 13th.

565

4 Discussion

The waters of the inner sea of Tierra del Fuego and the Magellan Strait are characterized by low-salinity (c.a. 31 g kg⁻¹) in comparison to the adjacent Pacific Ocean and Atlantic Ocean, where salinities are approximately 34 g kg⁻¹ (Panella et al., 1999; Palma and Silva, 2004; Brun et al., 2020). These oceanic waters enter the inner-sea and mix with regional freshwaters inputs. Strong tides (Medeiros and Kjerfve, 1988) and strong winds are the primary mechanisms driving mixing within the region.

570

The study of Brun et al. (2020) identifies the hydrographic condition of the Magellan Strait as the source of the low-salinity waters observed in the Atlantic; this study attributes the mixing primarily to tidal forcing. In the atmosphere, the Southern Annular Mode and other hemispheric-scale modes exert a substantial influence on Patagonia weather (Garreaud et al 2009). The region experiences an intensification of westerlies over Tierra del Fuego, which shows a significant positive trend during the austral summer (Garreaud et al., 2013). Local topography often channels these winds down-fjord (Oltmanns et al., 2014). This effect significantly impacts surface waters and ice, which can be driven down-fjord, subsequently promoting air-sea gas exchange and heat, and facilitating deep-water exchange within the fjords (Klymak et al., 2025).

The role of the wind promoting the deep exchange is an important dynamic component in semi-enclosed coastal systems, such as fjords, which are characterized by limited ventilation. Recent studies have documented warming and deoxygenation in the Puyuhuapi Fjord (Lindford et al., 2023) and in British Columbia (Jackson et al., 2021). Fjord circulation is driven by a combination of mechanism driven by density gradient, tides, and winds. In these systems, where the upper and intermediate layers are dominated by estuarine circulation (e.g. Stigebrandt, 2012). The role of wind has been understudied compared to density and tidal drivers in fjords (Soto-Riquelme et al. 2023) and this is also the case for the inner sea of the southern Patagonia fjords. In conditions of up-fjord winds, opposing to estuarine outflow, the transference of momentum could retain or even reverse the upper outflow (e.g. Caceres et al. 2002), this yield to a water column adjustment which drives in a three-layered residual pattern, particularly in fjords with weak tidal influence (Geyer, 1997). In systems with a strong tidally influence, deeper circulation could also generate a three-layer pattern observed in systems like Chilean Patagonia (e.g. Valle-Levinson et al., 2014). The mechanism of the formation of three or more layers in fjords has been documented by Farmer and Freeland (1983) and to verify the residual pattern of the circulation in the Almirantazgo fjord a dedicated study using ADCP data (as in Wan et al., 2017 or Castillo et al., 2012) will be required in future studies in the region.

4.1 Along-Fjord Exchange of Upper brackish Layer

The CTD-SRDL data was used in the estimations the upper outflows exchange and to determine flushing times of the same layer. The mean profiles from the along-fjord CTD-SRDL (Figs. 5c,d) and the mean profiles of the CTD-AML (Figs. 6c,d) were highly consistent between the seasons; highly stratified in January and mixed during June. Despite the fact, the study has several profiles in the region: inner the fjords and in the open ocean (see Fig. A2). In the Almirantazgo fjord, only few CTD-SRDL profiles were nearest to the cross-fjord stations (see Fig. 6a). Thus, the study was able to make a comparison for data of the 27 of January and between 6 and 9 of June of 2024. During those dates, CTD-SRDL data was taken nearest to CTD2 and CTD3 stations (Fig. 2, Fig. 6a). Considering that the study used a $h_1 = 20$ m during January, the comparison between upper layer (CTD depths ≤ 20 m) and deeper layer (CTD depths > 20 m) were used to quantify the agreement between CTD-SRDL and CTD-AML. The comparisons showed the best agreement for January temperatures ($r > 0.96$). During June, the relation

was lower with $r > 0.80$ in the upper layer. The summer Salinities presented $RSMD > 2.61 \text{ g kg}^{-1}$ with $r = 0.67$ at CTD2 in the upper layer. In June, RMSD was lower than January ($RSMD < 1.10 \text{ g kg}^{-1}$) with r between 0.85 (CTD2) and 0.94 (CTD3). In oceanic water, the performance change, the lowest errors and highest correlations were concentrated for profiles with depths $> 50 \text{ m}$ with $RSMD < 0.3$ and $r > 0.8$ for both temperature and salinity, indicating a good overall representation. Although the values in the surface layer were slightly lower, they remained consistently good for both variables. The low correlation in Salinity could be taken with the consideration that, **1)** few profiles were able to make the comparison, **2)** the dates selected were near under variable conditions of Salinity (see Fig. 8e), **3)** the processing of the traditional CTD take the downcast to be analyzed, whereas CTD-SRDL record data when elephant seals are coming to surface.

Utilizing the hydrographic data acquired from two complementary sources (CTD-SRLD and CTD-AML), within the Tierra del Fuego fjord system, and considering the basin geometry, a first-order estimation of the flushing time for the upper layer (T_F) of the Almirantazgo Fjord was estimated (from MC to BF). This approximation, based on the Knudsen theorem, this requires determining the upper Volume of the brackish layer. The study shows that during January the amount of freshwater was considerably higher than June (Fig. 7), additionally cross-fjord transect shows that during January the upper brackish layer (h_1) could be as deeper as 20 m but during June was shallower, $h_1 = 5 \text{ m}$ (Figs. 6c, d). Similarly, the along-fjord patterns showed that during January the fjord showed a salt-wedge pattern deeper near the head and shallower (c.a 5 m) near the mouth of the fjord (Figs. 5c) in contrary, during June the upper brackish layer has a constant and shallower (c.a 5 m) along the entire fjord. These cause geometry differences for the Volume between January (right trapezoidal prism) and June (right rectangular prism). Considering the average width of AF was nearly 12 km the upper Volume (V_1) of the brackish layer was $11.25 \times 10^9 \text{ m}^3$ in January whereas in June was $4.5 \times 10^9 \text{ m}^3$ yields that the upper brackish layer of January was 2.5 times of June.

At the mouth of the AF (Fig. 1) the study estimates the along-fjord exchange of the upper layer, $Q_1 = A_1 * v_1$, here A_1 was the transversal area at the mouth and v_1 is the advective along-fjord current. Expanding, $A_1 = b * h_1$, taking an averaged width of 12 km and $h_1 = 5 \text{ m}$ the transversal area in both January and June was $A_1 = 6 \times 10^4 \text{ m}^2$. This study do not take currents measurements but the Fisheries Chilean Institute (IFOP) is working to modelling the Magellan strait, they use a MIKE 3D model for modelling (<http://chonos.ifop.cl/>) the northern Patagonia (between Puerto Montt and Golfo de Penas, see Fig 1) and they using to estimate different operational results for decision makers (e.g. Reche et al., 2021; Ruiz et al., 2021). For the study region, the model lack in circulation validation (Ruiz, *pers. comm*) but the IFOP give us a weekly averages circulation patterns for January and June 2024 (see Fig. A3). Using that model output, the study obtained typical v_1 to determine the upper exchange Q_1 . The results showed that during, January $v_1 = 0.3 \text{ m s}^{-1}$ whereas in June $v_1 = 0.1 \text{ m s}^{-1}$ considering these values the output exchange in January was $Q_1 = 1.8 \times 10^4 \text{ m}^3 \text{ s}^{-1}$ and during June $Q_1 = 6 \times 10^3 \text{ m}^3 \text{ s}^{-1}$. Using that estimations to determine the flushing time ($T_F = V_1 * Q_1^{-1}$) of the upper layer, using this approximation T_F for January and June were similar order of magnitude of one week (Table 3). In this study high RMSD for surface Salinity could limited the precision of the Knudsen-based estimations. Despite that, the T_F quantification requires a second-order studies focusing into determine the water column

exchanges using a validated circulation model. As an example, Pinilla et al (2020) using a circulation validated model, determine the water age into the Puyuhuapi fjord obtaining longer than 1 year for intermediate and deeper waters of the fjord which could promote low DO in that region.

645

Table 3. Flushing times parameters for January and June. Upper (S_1/D_1) and deeper (S_2/D_2) average along-fjord Salinities/Densities, upper layer Volume (V_1), typical upper layer velocity (v_1), upper exchange (Q_1).

	S_1 [g kg ⁻¹]	S_2 [g kg ⁻¹]	D_1 [kg m ⁻³]	D_2 [kg m ⁻³]	V_1 [m ³]	v_1 [m s ⁻¹]	Q_1 [m ³ s ⁻¹]	T_F d
January	29.41 ± 1.13	30.75 ± 0.36	1022.6 ± 0.96	1024.1 ± 0.37	11.25 x10 ⁹	0.3	1.8x10 ⁴	~ 7
June	30.55 ± 0.46	30.58 ± 0.37	1023.9 ± 0.31	1024.0 ± 0.25	4.5 x10 ⁹	0.1	6 x10 ³	~ 9

650

The calculated upper layer flushing times were relatively longer than those reported for other highly stratified systems in northern Patagonia. For instance, Castillo et al. (2016) determined an upper-layer flushing time of 3 days for the Relocavi Fjord, and Calvete and Sobarzo (2011) found a similar 5-day flushing time for waters between the Guafo mouth and Elefantes Fjord. These results indicate that the upper layer of the Almirantazgo Fjord has flushing times of approximately one week, which may promote the concentration of materials within the basin, but the limitations of the Knudsen theorem must be taken into account and to serve as a tool that supports management and decision-makers, this first-order estimation of exchange times will require validation by a second-order estimation in future works.

655

4.2 Freshwater Dynamics and Stratification

Traditional CTD measurements acquired near Jackson Bay during both January and June recorded salinities below 31 g kg⁻¹. Data from the time series mooring (recovered in June) further indicated that near-bottom (c.a. 31 m depth) salinities reached a minimum of 27 g kg⁻¹ during the austral summer (January). This was followed by a seasonal increase, with values reaching a maximum of 30.5 g kg⁻¹ near the recovery dates in June 2024. The low salinities characteristic of the inner sea of Tierra del Fuego is attributed to the input of glacial meltwater, which exhibits strong seasonality. During summer, the along-fjord Freshwater Content (FWC) was higher in the first 20 km from the head of the fjord in the region where Parry fjord (PF) and the Azopardo river (AR) had the major influences on the freshwater input (Fig. 7a,b). Minor peaks of FWC and PEA were located at Brookes fjord (BF) suggesting the glacial melting influences on the FWC increase. Although a lack of comprehensive winter data prevents full confirmation, cross-fjord FWC measurements show that January values are at least twice as high as those observed during the June (Fig. 7a,b). This distribution is consistent with a surface slope toward the fjord head and an upper salt-wedge along-fjord structure (Fig. 5c). This typical structure of strong stratified estuaries, like fjords (Geyer and Ralston, 2011) has been documented previously in Chilean fjords like, the Reloncaví fjord (e.g. Castillo et al., 2012) and

665

670

Puyuhuapi fjord (e.g. Schneider et al., 2014). The FWC has been useful to quantify the amount of freshwater input in Chilean fjords system between 43.5°S to 46.5°S by Calvete and Sobarzo (2011) in this study the authors report FWC ~ 7 m near a region with tidal-glacial influence, here FWC was < 2 m suggesting a low amount of freshwater in the Almirantazgo fjord system. The summer upper-layer salinity from the seal-borne data was less well constrained than temperature, this uncertainty propagates into the along-fjord Freshwater Content estimations.

4.3 Patterns of Variability at 30 m depth near the fjord's head

Wavelet analysis of the time series indicated a dominance of low-frequency (periods > 24 h) variability in wind-stress magnitude, Conservative Temperature (CT), Dissolved Oxygen (DO) and Absolute Salinity (S_A). The sea level (SL) was the exception (Fig. 9b), being dominated by the semi-diurnal (12 h) and diurnal (24 h) signals. This tidal dominance is consistent with the harmonics analysis and the calculated form factor ($F = 0.593$).

In stratified systems, internal (baroclinic) seiches can be excited in the pycnocline, contributing to entrainment and enhancing heat and salt exchange between layers e.g. in the Gulmar fjord (Arneborg and Liljebladh, 2001), the Gullmaren fjord (Djurfeldt, 1987) and the Reloncavi fjord (Castillo et al., 2017). Up-fjord winds generate a surface slope ($\partial\eta/\partial x > 0$), highest at the fjord head. In stratified conditions ($\Delta\rho \gg 1$), this results in a deepening of the pycnocline at the head, creating an opposing pycnocline tilt ($\partial h_1/\partial x < 0$) (Farmer, 1976). In this study, the effect of the wind over the dynamics of the fjord is the main focus thus to determine the baroclinic adjustment scale (t_s), in the Almirantazgo fjord here the study consider the length of the AF which is 75 km to determine the time required for this pycnocline tilt to be established, is defined by $t_s \sim L / c_1$ (e.g. Klymak et al., 2025). Here, for January conditions $t_s = 3.34$ days whereas for June $t_s = 22$ days. Comparing t_s with T_F (Table 3), the results indicates that during January the baroclinic adjustment is lower than the flushing time but during June, $t_s > T_F$ suggesting that the wind-setup do not reach the quasi-steady state, probably due to the transient nature of the strong winds (Fig. 8) and the weak stratification of the water column during this month (Figs. 5 and 6) in the region.

Based on the established importance of the Earth's rotation, the 6-month mooring data, the short-time mooring data, and the hydrographic surveys, a schematic fjord dynamic is proposed. Winds in the region (Maria Cove, Fig. 2) are typically persistent and directed up-fjord (see Fig. A1). These forcing tilts the surface layer ($\partial\eta/\partial x$) and enhances the along-fjord barotropic pressure gradient. The response is a deepening of the pycnocline at the head and shallowing at the mouth (Fig. 5c). This salinity structure is concurrently modulated by summer glacial meltwater (Fig. 7a). These freshwater forms a buoyant plume. Under weak winds, this plume would likely be deflected by the Coriolis force and exit the fjord. However, moderate to-intense up-fjord winds—acting within the baroclinic adjustment time ($t_s = 3.34$ days)—appear to overcome this deflection, trapping the plume at the fjord head. During this process, the plume gains heat. Consequently, January up-fjord wind events are associated with pulses of warmer waters at 30 m depth (Fig. 8d). This is consistent to the observations made by Aravena et al., (2025) in

Punta Santa Ana at the south of the Magellan Strait. The associated turbulence also increases DO and promotes mixing of the upper freshwater, which reduces the salinity of the deeper waters (Fig. 8e). This process is markedly seasonal. A regime occurred in mid-March 2024, after which wind intensity decreased, and strong events became primarily synoptic. In this early autumn period, up-fjord wind events began to be associated with colder waters. In winter, the up-fjord wind events were related to colder waters. During June, surface water resides longer (see Table 3) at the surface than in January and thus loses heat to colder overlying air. These colder events are associated with high DO, resulting from enhanced turbulence (oxygen ingress) and the higher solubility of oxygen in colder water (Fig. 8d). Observations in the short-term are consistent with the synthesis. Following a strong up-fjord wind pulse ($W_b > 1$), the DO gradient approached zero ($\partial DO / \partial z \sim 0$), implying oxygenation of the layer at 30 m (Fig. 8e,f). The time-domain analysis (Fig. 8) retained the linear trends to reflect the seasonality. The decreasing river discharge trend and increasing salinity trend reflect the freshwater impact in the study region. However, the FWC estimations (Fig. 7a) suggest that freshwater inputs from glacial locations are more significant than riverine inputs. In the region, Izaguirre et al (2025) using aerial imagery between 1945-2024 showed that Darwin's Cordillera increase the number of glacial lakes and thus the freshwater input by Glacial Lake Outburst Floods (GLOFs) due to a warming progress. Although the specific volume of glacial inputs was not estimated, this study concludes that the combination of this substantial freshwater input with tidal and wind-driven mixing maintains the low-salinity conditions of the inner-sea of Tierra del Fuego.

4.3 Wind influence on the inner sea of Almirantazgo Fjord

While ERA5 reanalysis data are valuable for assessing the spatio-temporal variability of the wind forcing in southern Patagonia, comparisons with *in-situ* meteorological stations (e.g. Punta Arenas and Porvenir) reveal that ERA5 magnitudes are typically 20% lower than observations. This remarks that the wind-based estimations of the study were likely conservative and thus supporting the high influence of wind-driven on the dynamics of the Almirantazgo fjord.

To assess the potential for wind-driven mixing, the dimensionless Wedderburn number (W_b) was calculated. During January, intense wind-stress in AF could be as high as $\tau = 0.10 \text{ N m}^{-2}$ taken a mean density difference $\Delta\rho = 1.5 \text{ kg m}^{-3}$ (see Table 3) an upper layer $h_1 = 20 \text{ m}$ and considering the AF length $L = 75 \text{ km}$ the study obtained a $W_b = 1.28$. This value indicates that wind events can perturb the pycnocline during the stronger stratified condition of January. The June conditions were highly mixed with a $\Delta\rho = 0.1 \text{ kg m}^{-3}$ and $h_1 = 5 \text{ m}$, wind stress was able to perturb the pycnocline even in weakest winds conditions. For other hand, the observational data from the short-term deployment (Fig. 9e) confirm this: during highly stratified conditions, W_b repeatedly exceeded 1 and reached values as high as 3.7, demonstrating that wind is sufficient to drive dynamics at the fjord head. The h_1 depths for January and June were estimated by the cross-fjord and along-fjord maximum N^2 depths at the sites near of the CTD cross-fjord transects (Fig. 2). The sensitivity of the W_b parameter was assessed by calculating the wind-influenced depth $h_{1W} = \sqrt{\frac{\tau L}{\Delta\rho g}}$. In January, $h_{1W} = 22.6 \text{ m}$ closely aligns with the $h_1 = 20 \text{ m}$ used on the estimations. In June, however, $h_{1W} = 27.7 \text{ m}$ exceeding the seasonal depth for W_b by a factor of six. Although direct validation via ADCP data was

not available for this study – unlike the work of Wan et al (2017) in Douglas Channel – the fact that W_b meets or exceeds the prescribed layer depth indicates that wind-driven forcing is a major driver of the surface layer in this system.

The time series (Fig. 9) also reveals a feedback mechanism. Strong winds, mix the upper water column, which diminishes the density gradient ($\Delta\rho$) (Fig. 9d). This reduction in stratification, in turn, increases the W_b (as $\Delta\rho$ is in the denominator), enhancing the wind's mixing efficiency (Fig 9e). Conversely, when winds weaken (due to synoptic and daily oscillations), freshwater discharge strengthens the density gradient. This re-stratification requires greater wind energy to overcome, and in the absence of strong wind, W_b returns to near-zero values (Fig. 9e). Under $W_b > 1$ the water column DO yields to mixing conditions ($\Delta DO \sim 0$) showing a rapid response of the 30 m water column to strong winds (Fig. 9f).

745

To further quantify the wind's role, the power per unit area generated by the wind ($d\phi_W/dt = 9 \rho_a c_d W_{10}^3$) available for mixing was calculated, following methodologies from Denman and Miyake (1973) and Bowden (1981). Here, ρ_a is air density (1.23 kg m^{-3}), c_d is drag coefficient (1.1×10^{-3}) and W_{10} is the wind magnitude at 10 m height, here the study uses the mean magnitude (5.3 m s^{-1}) and percentile 90 (p90) of the magnitude (10 m s^{-1}) for the estimations. Additionally, the 9 coefficient was determined by $9 = \sqrt{\frac{\rho_a c_d}{\rho}}$ where ρ is the water density. The study determines for mean magnitude $d\phi_W/dt = 0.2 \times 10^{-3} \text{ W m}^{-2}$ whereas for p90, $d\phi_W/dt = 1.5 \times 10^{-3} \text{ W m}^{-2}$.

To determine if this wind power is sufficient for mixing, it must be compared to the power required to maintain stratification $d\phi_E/dt$ by the estuarine circulation (Simpson et al., 1990). This stratification power is given by: $d\phi_E/dt = (1/320) (g^2 h_1^5 / A_z \rho) (\partial\rho/\partial x)^2$. Following Osborn (1980), the vertical eddy diffusivity is parameterized as $K_z = 0.2(\epsilon/N^2)$. Based on microstructure measurements from winter 2024 (Rojas-Celis et al., 2025), we assume a representative dissipation rate of $\epsilon \sim O(10^{-6}) \text{ W kg}^{-1}$ and typical buoyancy frequency of $N^2 = 0.0028 \text{ s}^{-2}$ which yields $K_z \sim 7 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$.

The relationship between the vertical mixing of momentum and scalars is governed by the turbulent Prandtl number ($Pr_t = A_z/K_z$; Thorpe, 2005). Assuming the standard Reynolds analogy for neutral or weakly stratified flows ($Pr_t = 1$; Mellor and Yamada, 1982; Rodi, 1987), the resulting eddy viscosity ($A_z = K_z$) yields a stratification power of $d\phi_E/dt = 21 \times 10^{-3} \text{ W m}^{-2}$. This value is one order of magnitude higher than wind power ($d\phi_W/dt$), which would imply that wind forcing is insufficient for mixing. However, the $Pr_t = 1$ assumption breaks down under stratified environments where buoyancy forces suppress scalar mixing more efficiently than downward momentum transport (e.g. Munk and Anderson, 1948; Venayagamoorthy and Stretch, 2010). Consequently, in highly stable flows, Pr_t increases significantly, often reaching values $O(10)$ (e.g. Pacanowski and Philander, 1981; Peters et al., 1988). Applying a scaling of $Pr_t = 10$ to the summer conditions in the Almirantazgo fjord yields an effective eddy viscosity $A_z \sim 7 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$. This reduces the estuarine stratification power to $d\phi_E/dt = 2.1 \times 10^{-3} \text{ W m}^{-2}$.

Under these physically constrained parameterizations, first-order approximations indicate that estuarine stratification power becomes comparable to the wind power during intense episodic winds events of up to 9 m s^{-1} ($d\phi_w/dt = 1.2 \times 10^{-3} \text{ W m}^{-2}$). This scaling suggests that despite the high static stability of the water column, wind momentum can be sufficiently transferred downward to overcome stratification and induce mixing. Nonetheless, these steady-state assumptions warrant further validation through future observational and numerical studies to better constrain the spatiotemporal variability of mixing in the fjord.

775

Despite the dominant tidal forcing in the Magellan Strait (Fig. 3) and the strong salinity-driven stratification from glacial melt (Fig. 4c, 5c, 6a, 6c), the wind is a critical mixing agent. Both the Wedderburn number analysis and the mixing power comparison confirm the wind's capacity to mix the upper water column. This study provides the first quantification of the wind-driven effect in one of the southernmost fjord of Chile (Fig. 10).

780




Figure 11. Squared coherence wavelet spectrums, between τ (wind-stress) with a) Conservative Temperature (CT), Absolute Salinity (S_A) and Dissolved Oxygen (DO) for the time series acquired at the A1. Horizontal dashed lines denote periods of 4, 12, 24, and 72 hours.

785

Wind stress in the study area showed significant squared coherence with CT, SA, and DO at periods greater than 16 h, exhibiting marked synoptic variability at periods exceeding 70 h (Fig. 11). The most intense events ($> 0.07 \text{ N m}^{-2}$; Fig. 8) were highly coherent within the 24–72 h band, suggesting that synoptic wind forcing drives a uniform response in hydrographic properties at 30 m near the fjord head. Interestingly, CT coherence for periods $> 64 \text{ h}$ showed a distinct dip centered on April 4, 2024, despite remaining high during the surrounding periods (Fig. 11a). This indicates a temporary shift in the CT response to wind forcing between March 30 and April 9, which resulted in the observed breakdown in coherence between the variables.

790

In the context of climate change and the projected intensification of the Southern Hemisphere westerlies, these findings are critical. The Almirantazgo Fjord ecosystem, with a seasonal stratification of the upper layer and mixing processes dependent on both tides and wind, will be profoundly influenced by future atmospheric changes. Therefore, while tidal and buoyancy forces establish the baseline conditions, the answer to the fjord's physical and ecological future is unequivocally linked to the wind.

5 Conclusions

The hydrodynamics of the inner sea of Tierra del Fuego are primarily controlled by the seasonal cycle of glacial freshwater input, which establishes strong stratification during summer and modulates upper layer renewal rates. This buoyancy forcing drives a circulation characterized by rapid upper-layer flushing time in the order of one week. While M_2 tidal energy dominates the adjacent Magellan Strait, our analysis confirms that the specific dynamics of the Almirantazgo Fjord are governed by the interplay between this glacial freshwater buoyancy and along-fjord wind stress.

A key contribution of this study—supported by unprecedented spatial coverage from instrumented southern elephant seals—is the quantification of wind-driven mixing. The frequent occurrence of Wedderburn numbers > 1 (peaking at 3.7) provides direct evidence that synoptic wind events are sufficient to destabilize the pycnocline.

Ultimately, these physical processes have critical biogeochemical consequences. The observed oxygenation at 30 m depth of the fjord's head following strong up-fjord wind pulses indicates that atmospheric forcing is essential for ventilating the inner bay at the head of the fjord. As climate change projects an intensification of the Southern Hemisphere westerlies, the physical and ecological future of this fjord system will be increasingly defined by its sensitivity to wind-driven mixing.

Authors contribution

MS, CBG, AIG, MFL and MIC designed the study and wrote the initial manuscript draft. AP and JG-V contribute to improving the subsequent versions of the manuscript. NC, MR and CZ helped with data analysis. Discussions and iterative feedback from all co-authors significantly contributed to the revision of the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

825 The authors thank the researchers, students, and field personnel who assisted in data collection; their efforts made this study possible. We thank the two anonymous reviewers whose comments and suggestions helped to improve this manuscript. We also want to thanks to Cristian Ruiz from IFOP for their model output. Field measurements received funding from Anillo Seals ATE220033, with supplementary support provided by RED 21992 (MINEDUC, Chile). Logistic helps on Punta Arenas were carried out by IDEAL-FONDAP 151CTD-AML data acquisition utilized instrumentation funded by FONDEQUIP
830 EQM170115 (MIC). Furthermore, this research received financial support from FONDECYT 1231058, CIMAR 27F 24-11 (CONA, Chile), FONDEF ID22I10206. JG-V received support from FONDAP N° 15150003. NC received support from CCSS210020. The results contain modified Copernicus Climate Change Service information 2025. Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or data it contains. Bathymetry data was obtained from the General Bathymetric Chart of the Oceans (GEBCO, 2025). This work forms part of
835 the academic portfolio of MIC for full professorship at the University of Valparaíso (UV).

Code and Data Availability Statement

The datasets used open source and are available online: hourly ERA 5 wind data by Herbach et al. (2023), sealevel data at <https://www.ioc-sealevelmonitoring.org/>. The TPOX9 tidal model is available at OSU TPX (<https://www.tpxo.net/global>). The
840 wind data from meteorological stations along Chile is available at <https://climatologia.meteochile.gob.cl/application/requerimiento/producto/RE3008>. Bathymetry data was downloaded from GEBCO 2025 Grid. All data set used on this study are available at <https://doi.org/10.5281/zenodo.19433749>. In addition, all scripts used to obtain the results presented in this study could be shared upon request at the corresponding author.

References

845 Aalto, N. J., Campbell, K., Eilertsen, H. C., and Bernstein, H. C.: Drivers of Atmosphere-Ocean CO₂ Flux in Northern Norwegian Fjords, *Front Mar Sci*, 8, 692093, <https://doi.org/10.3389/FMARS.2021.692093/BIBTEX>, 2021.

Aguayo, R., León-Muñoz, J., Vargas-Baccheler, J., Montecinos, A., Garreaud, R., Urbina, M., Soto, D., and Iriarte, J. L.: The glass half-empty: climate change drives lower freshwater input in the coastal system of the Chilean Northern Patagonia.
850 *Climatic Change*, 155(3), 417–435. <https://doi.org/10.1007/S10584-019-02495-6/METRICS>, 2019.

Antezana, T.: Hydrographic features of Magellan and Fuegian inland passages and adjacent subantarctic waters, *Sci Mar*, 63, 23–34, <https://doi.org/10.3989/scimar.1999.63s123>, 1999.

- 855 Aracena, C., Lange, C. B., Luis Iriarte, J., Rebolledo, L., and Pantoja, S.: Latitudinal patterns of export production recorded in surface sediments of the Chilean Patagonian fjords (41–55°S) as a response to water column productivity, *Cont Shelf Res*, 31, 340–355, <https://doi.org/10.1016/J.CSR.2010.08.008>, 2011.
- 860 Aravena-Yáñez, V., Garcés-Vargas, J., & Benavides-Martínez, I. F.: Freshwater plume’s impact on the thermohaline structure of the water column in the Central Zone of the Strait of Magellan. *Regional Studies in Marine Science*, 83. <https://doi.org/10.1016/j.rsma.2025.104087>, 2025.
- Ariztía, T. and Undurraga, T.: Anticipating green hydrogen futures: exploring the socio-material production of a new clean fuel, *J Cult Econ*, <https://doi.org/10.1080/17530350.2025.2485999>, 2025.
- 865 Arneborg, L., and Liljebladh, B.: The internal seiches in Gullmar Fjord. Part II: Contribution to basin water mixing. In *Journal of Physical Oceanography* (Vol. 31, Issue 9). <https://doi.org/10.1175/1520-0485>, 2001.
- Bianchi, T. S., Arndt, S., Austin, W. E. N., Benn, D. I., Bertrand, S., Cui, X., Faust, J. C., Kozirowska-Makuch, K., Moy, C. M., Savage, C., Smeaton, C., Smith, R. W., and Syvitski, J.: Fjords as Aquatic Critical Zones (ACZs), <https://doi.org/10.1016/j.earscirev.2020.103145>, 2020.
- 870 Bianucci, L., Jackson, J. M., Allen, S. E., Krassovski, M. v., Giesbrecht, I. J. W., & Callendar, W. C. Fjord circulation permits a persistent subsurface water mass in a long, deep mid-latitude inlet. *Ocean Science*, 20(2), 293–306. <https://doi.org/10.5194/os-20-293-2024>, 2024.
- 875 Boehme, L., Lovell, P., Biuw, M., Roquet, F., Nicholson, J., Thorpe, S. E., Meredith, M. P., and Fedak, M.: Technical Note: Animal-borne CTD-Satellite Relay Data Loggers for real-time oceanographic data collection, *Ocean Sci.*, 5, 685–695, <https://doi.org/10.5194/os-5-685-2009>, 2009.
- 880 Boone, W., Rysgaard, S., Carlson, D. F., Meire, L., Kirillov, S., Mortensen, J., Dmitrenko, I., Vergeynst, L., Sejr, M. K. Coastal Freshening Prevents Fjord Bottom Water Renewal in Northeast Greenland: A Mooring Study From 2003 to 2015. *Geophysical Research Letters*, 45(6), 2726–2733. <https://doi.org/10.1002/2017GL076591>, 2018.
- Blanton, J. O., and Atkinson, L. P.: Transport and fate of river discharge on the continental shelf of the southeastern United States. *Journal of Geophysical Research*, 88(C8), 4730–4738. <https://doi.org/10.1029/JC088IC08P04730>, 1983.

- 885 Bravo, C., Quincey, D. J., Ross, A. N., Rivera, A., Brock, B., Miles, E., Silva, A. Air Temperature Characteristics, Distribution, and Impact on Modeled Ablation for the South Patagonia Icefield. *Journal of Geophysical Research: Atmospheres*, 124(2), 907–925. <https://doi.org/10.1029/2018JD028857>, 2019.
- Brun, A. A., Ramirez, N., Pizarro, O., and Piola, A. R.: The role of the Magellan Strait on the southwest South Atlantic shelf. *Estuarine, Coastal and Shelf Science*, 237, <https://doi.org/10.1016/j.ecss.2020.106661>, 2020
- 890 Cáceres, M., Valle-Levinson, A., Sepúlveda, H. H., and Holderied, K.: Transverse variability of flow and density in a Chilean fjord. In *Continental Shelf Research* (Vol. 22, Issues 11–13). [https://doi.org/10.1016/S0278-4343\(02\)00032-8](https://doi.org/10.1016/S0278-4343(02)00032-8), 2002.
- Calvete, C., and Sobarzo, M.: Quantification of the surface brackish water layer and frontal zones in southern Chilean fjords between Boca del Guafo (43°30'S) and Estero Elefantes (46°30'S). *Continental Shelf Research*, 31(3–4), 162–171.
- 895 <https://doi.org/10.1016/j.csr.2010.09.013>, 2011.
- Carrasco, J. F., Casassa, G., and Rivera, A.: Meteorological and climatological aspects of the Southern Patagonia Icefield. In G. Casassa, F. V. Sepúlveda, and R. M. Sinclair (Eds.), *The Patagonian Icefields: A unique natural laboratory for environmental and climate change studies* (pp. 29–41). Springer US. https://doi.org/10.1007/978-1-4615-0645-4_4, 2002.
- 900
- Castillo, M. I., Pizarro, O., Cifuentes, U., Ramirez, N., and Djurfeldt, L. : Subtidal dynamics in a deep fjord of southern Chile. *Continental Shelf Research*, 49, 73–89, 2012.
- Castillo, M. I., Pizarro, O., Ramírez, N., and Cáceres, M.: Seiche excitation in a highly stratified fjord of southern Chile: the
- 905 Reloncaví fjord. *Ocean Science*, 13(1), 145–160, 2017.
- Chaigneau, A., and Pizzaro, O.: Eddy characteristics in the eastern South Pacific. *Journal of Geophysical Research: Oceans*, 110(6), 1–12. <https://doi.org/10.1029/2004JC002815>, 2005.
- 910 Chaigneau, A., Dominguez, N., Eldin, G., Vasquez, L., Flores, R., Grados, C., and Echevin, V.: Near-coastal circulation in the Northern Humboldt Current System from shipboard ADCP data. *Journal of Geophysical Research: Oceans*, 118(10), 5251–5266. <https://doi.org/10.1002/JGRC.20328>, 2013.
- Chant, R. J.: Secondary circulation in a region of flow curvature: Relationship with tidal forcing and river discharge, *J Geophys Res Oceans*, 107, <https://doi.org/10.1029/2001jc001082>, 2002.
- 915

- Church, J. A., Monselesan, D., Gregory, J. M., and Marzeion, B.: Evaluating the ability of process based models to project sea-level change. *Environmental Research Letters*, 8(1), 014051. <https://doi.org/10.1088/1748-9326/8/1/014051>, 2013.
- 920 Cisternas, N., Garcés-Vargas, J., Castillo, M. I., Barrios-Guzmán, C., Barilari, F., Landaeta, M. F., Sepúlveda, M., Piñones, A. Salinity variability in the mixed layer off Chilean Patagonia: potential influence of Patagonian ice fields. *Progress in Oceanography*, 242, 103653. <https://doi.org/10.1016/j.pocean.2025.103653>, 2026.
- Deleersnijder, E., Campin, J. M., and Delhez, E. J. M.: The concept of age in marine modelling I. Theory and preliminary
925 model results. *Journal of Marine Systems*, 28(3–4), 229–267. [https://doi.org/10.1016/S0924-7963\(01\)00026-4](https://doi.org/10.1016/S0924-7963(01)00026-4), 2001.
- Denman, K. L., and Miyake, M.: Behavior of the mean wind, the drag coefficient, and the wave field in the open ocean. *Journal of Geophysical Research*, 78(12), 1917–1931. <https://doi.org/10.1029/jc078i012p01917>, 1973.
- 930 Downes, S. M., Langlais, C., Brook, J. P., and Spence, P.: Regional impacts of the westerly winds on Southern Ocean mode and intermediate water subduction. *Journal of Physical Oceanography*, 47(10), 2521–2530, 2017.
- Dyer, K. R. *Estuaries, a physical introduction* 2nd Edition, Wiley, 195 pp., 1997.
- 935 Dyer, K.: Estuarine Circulation, *Encyclopedia of Ocean Sciences*, Third Edition: Volume 1-5, 1–5, V6-67-V6-73, <https://doi.org/10.1016/B978-0-12-409548-9.11427-7>, 2019.
- Djurfeldt, L.: On the response of the fjord gullmaren under ice cover. *Journal of Geophysical Research: Oceans*, 92(C5), 5157–5167. <https://doi.org/10.1029/JC092iC05p05157>, 1987.
940
- Egbert, G. D., & Erofeeva, S. Y: Efficient inverse modeling of barotropic ocean tides. *Journal of Atmospheric and Oceanic technology*, 19(2), 183–204, 2002.
- Farmer, D. M. and Freeland, H. J.: The physical oceanography of Fjords, *Progress in Oceanography*, 147–220 pp.,
945 [https://doi.org/10.1016/0079-6611\(83\)90004-6](https://doi.org/10.1016/0079-6611(83)90004-6), 1983.
- Garreaud, R., Lopez, P., Minvielle, M., and Rojas, M.: Large-Scale control on the Patagonian Climate. *Journal of Climate*, 26(1), 215–230. <https://doi.org/10.1175/JCLI-D-12-00001.1>, 2013.

- 950 Garcés-Vargas, J., Piñones, A., Schneider, W., Landaeta, M. F., Castillo, M. I., Cisternas, N., Barrios-Guzmán, C., Barilari, F.
Seasonal dynamics and forcing mechanisms of the Cape Horn Current: insights from reanalysis data and hydrographic
observations. *Progress in Oceanography*, 242, 103665. <https://doi.org/10.1016/j.pocean.2025.103665>, 2026.
- Garcia-Santos, Y., Narváez, D. A., Jacques-Coper, M., Saldías, G. S., Bozkurt, D., and Alessio, B. M.: Dominant wind patterns
955 under the influence of atmospheric rivers: Implications for coastal upwelling off central-southern Chile. *Journal of Geophysical
Research: Oceans*, 130(3), e2024JC021444, 2025.
- GEBCO Compilation Group: GEBCO 2025 Grid (doi:10.5285/37c52e96-24ea-67ce-e063-7086abc05f29, 2025.
- 960 Geyer, W. R.: Influence of wind on dynamics and flushing of shallow estuaries. In *Estuarine, Coastal and Shelf Science* (Vol.
44, Issue 6). <https://doi.org/10.1006/ecss.1996.0140>, 1997.
- Geyer, W. R., and D. K. Ralston: The dynamics of strongly stratified estuaries." *Treatise on Estuarine and Coastal Science*.
Amsterdam: Elsevier (2011): 37-52, 2011.
- 965 Geyer, W. R. and MacCready, P.: The estuarine circulation, *Annu Rev Fluid Mech*, 46, 175–197,
<https://doi.org/10.1146/annurev-fluid-010313-141302>, 2014.
- Giesecke, R., Martín, J., Piñones, A., Höfer, J., Garcés-Vargas, J., Flores-Melo, X., Alarcón, E., Durrieu de Madron, X.,
970 Bourrin, F., and González, H. E.: General Hydrography of the Beagle Channel, a Subantarctic Interoceanic Passage at the
Southern Tip of South America. *Frontiers in Marine Science*, 8, 621822. <https://dx.doi.org/10.3389/fmars.2021.621822>, 2021.
- Giesecke, R., Galbán-Malagón, C., Salamanca, M., Chandia, C., Ruiz, C., Bahamondes, S., and Astorga-Gallano, D.:
Automated FerryBox monitoring reveals the first recorded river induced crude oil seep transport to the Strait of Magellan in
975 southern Patagonia. *Science of The Total Environment*, 920, 170706. <https://doi.org/10.1016/j.scitotenv.2024.170706>, 2024.
- Grinsted, A., Moore, J.C., Jevrejeva, S.: Application of the cross wavelet transform and wavelet coherence to geophysical time
series, *Nonlin. Processes Geophys.*, 11, 561–566, <https://doi.org/10.5194/npg-11-561-2004>, 2004.
- 980 Guinet, C., Xing, X., Walker, E., Monestiez, P., Marchand, S., Picard, B., Jaud, T., Authier, M., Cotté, C., Dragon, A. C.,
Diamond, E., Antoine, D., Lovell, P., Blain, S., D’Ortenzio, F., and Claustre, H.: Calibration procedures and first dataset of
Southern Ocean chlorophyll a profiles collected by elephant seals equipped with a newly developed CTD-fluorescence tags.
Earth System Science Data, 5(1), 15–29. <https://doi.org/10.5194/ESSD-5-15-2013>, 2013.

- 985 Guo, X. and Valle-Levinson, A.: Wind effects on the lateral structure of density-driven circulation in Chesapeake Bay, *Cont Shelf Res*, 28, 2450–2471, <https://doi.org/10.1016/j.csr.2008.06.008>, 2008.
- Hansen, D. V. and Rattray, M.: Gravitational circulation in straits and estuaries, *Journal of Marine Research*, 104–122 pp., 1965.
- 990 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS), DOI: 10.24381/cds.adbb2d47 (Accessed on 15-Nov-2025), 2023.
- 995 Hucke-Gaete, R., Viddi, F. A., and Simeone, A.: Marine Mammals and Seabirds of Chilean Patagonia: Focal Species for the Conservation of Marine Ecosystems, 233–261, https://doi.org/10.1007/978-3-031-39408-9_9, 2023.
- Ianniello, J. Tidally induced residual currents in estuaries of constant breadth and depth. *J Mar Res* 35(4):755–786, 1977.
- 1000 Inall, M. E., Nilsen, F., Cottier, F. R., and Daac, R. : Shelf/fjord exchange driven by coastal-trapped waves in the Arctic. *Journal of Geophysical Research: Oceans*, 120(12), 8283–8303. <https://doi.org/10.1002/2015JC011277>, 2015.
- Inall, M. E. and Gillibrand, P. A.: The physics of mid-latitude fjords: A review, *Geol Soc Spec Publ*, 344, 17–33, <https://doi.org/10.1144/SP344.3>, 2010.
- 1005 IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change[Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, In press, doi:10.1017/9781009157896, 2021
- 1010 Izagirre, E., Glasser, N., Menounos, B., Aravena, J. C., Faria, S., and Antiguiedad, I.: The glacial geomorphology of the Cordillera Darwin Icefield, Tierra del Fuego, southernmost South America. *Journal of Maps*, 20(1), 2378000, 2024.
- 1015

- Izagirre, E., Casassa, G., Dussailant, I., Miles, E. S., Wilson, R., Rada, C., Faria, S. and Antigüedad, I.: Evolution of glacial lakes and southernmost GLOFs in the Cordillera Darwin and Cloue Icefields (Tierra del Fuego) between 1945–2024. *Frontiers in Earth Science*, 13, 1641167, 2025.
- 1020 Jackson, J. M., Bianucci, L., Hannah, C. G., Carmack, E. C., and Barrette, J.: Deep Waters in British Columbia Mainland Fjords Show Rapid Warming and Deoxygenation From 1951 to 2020, *Geophys. Res. Lett.*, 48, e2020GL091094, <https://doi.org/10.1029/2020GL091094>, 2021.
- 1025 Jackson, J. M., Hare, A., Hannah, C. G., Hilborn, A., Zahner, S. J. V., Page, S., Rosen, S., Timmerman, A. H. V., Lee, L. Why Deep-Water Dissolved Oxygen Is Higher in Gandaawuu.ngaay Xyangs (Juan Perez Sound), Haida Gwaii, Than Other British Columbia Fjords. *Journal of Geophysical Research: Oceans*, 130(3). <https://doi.org/10.1029/2024JC021826>, 2025.
- 1030 Klinck JM, O'Brien JJ, Svendsen H. A simple model of fjord and coastal circulation interaction. *J Phys Oceanogr* 11:1612–1626, 1981.
- Klymak, J. M., Jackson, J. M., and Hannah, C. G.: Predicting Upwelling due to Down-Fjord Winds. *Journal of Physical Oceanography*, 55(9), 1553–1568. <https://doi.org/10.1175/jpo-d-24-0176.1>, 2025.
- 1035 Lacy, J. R. and Monismith, S. G.: Secondary currents in a curved, stratified, estuarine channel, *Journal of Geophysical Research: Oceans*, 31283–31302 pp., <https://doi.org/10.1029/2000jc000606>, 2001.
- 1040 Lamy, F., Arz, H. W., Kilian, R., Lange, C. B., Lembke-Jene, L., Wengler, M., Kaiser, J., Baeza-Urrea, O., Hall, I. R., Harada, N., and Tiedemann, R.: Glacial reduction and millennial-scale variations in Drake Passage throughflow. *Proceedings of the National Academy of Sciences*, 112(44), 13496–13501. <https://doi.org/10.1073/PNAS.1509203112>, 2015.
- Landaeta, M. F., Paredes, L. D., Castillo, M. I. and González, M. T.: Spatio-temporal patterns of ichthyoplankton in southern Chilean Patagonia: beta-diversity and associated environmental factors, *Fisheries Oceanography*, 32, 341–351, <https://doi.org/10.1111/fog.12633>, 2023
- 1045 Letelier, J., Soto-Mardones, L., Salinas, S., Osuna, P., López, D., Sepúlveda, H. H., Pinilla, E., and Rodrigo, C.: Variabilidad del viento, oleaje y corrientes en la región norte de los fiordos Patagónicos de Chile, *Revista de Biología Marina y Oceanografía*, 46, 363–377, 2011.

- 1050 Lavers, D. A., Hersbach, H., Rodwell, M. J., and Simmons, A.: An improved estimate of daily precipitation from the ERA5 reanalysis. *Atmospheric Science Letters*, 25(3). <https://doi.org/10.1002/asl.1200>, 2024.
- León-Munõz, J., Urbina, M. A., Garreaud, R., and Iriarte, J. L.: Hydroclimatic conditions trigger record harmful algal bloom in western Patagonia (summer 2016). *Scientific Reports*, 8(1). <https://doi.org/10.1038/s41598-018-19461-4>, 2018.
- 1055 Linford, P., Pérez-Santos, I., Montes, I., Dewitte, B., Buchan, S., Narváez, D., Saldías, G., Pinilla, E., Garreaud, R., Díaz, P., Schwerter, C., Montero, P., Rodríguez-Villegas, C., Cáceres-Soto, M., Mancilla-Gutiérrez, G., and Altamirano, R.: Recent Deoxygenation of Patagonian Fjord Subsurface Waters Connected to the Peru–Chile Undercurrent and Equatorial Subsurface Water Variability. *Global Biogeochemical Cycles*, 37(6), 1–25. <https://doi.org/10.1029/2022GB007688>, 2023.
- 1060 Linford, P., Pérez-Santos, I., Montero, P., Díaz, P. A., Aracena, C., Pinilla, E., Barrera, F., Castillo, M., Alvera-Azcárate, A., Alvarado, M., Soto, G., Pujol, C., Schwerter, C., Arenas-Urbe, S., Navarro, P., Mancilla-Gutiérrez, G., Altamirano, R., San Martín, J., and Soto-Riquelme, C.: Oceanographic processes driving low-oxygen conditions inside Patagonian fjords. *Biogeosciences*, 21(6), 1433–1459. <https://doi.org/10.5194/bg-21-1433-2024>, 2024.
- 1065 MacCready, P. and Banas, N. S.: Residual Circulation, Mixing, and Dispersion, *Treatise on Estuarine and Coastal Science*, 75–89 pp., <https://doi.org/10.1016/B978-0-12-374711-2.00205-9>, 2012.
- MacCready, P. and Geyer, W. R.: Advances in estuarine physics, *Ann Rev Mar Sci*, 2, 35–58, [https://doi.org/10.1146/annurev-](https://doi.org/10.1146/annurev-marine-120308-081015)
1070 [marine-120308-081015](https://doi.org/10.1146/annurev-marine-120308-081015), 2010.
- Medeiros, C., and Kjerfve, B.: Tidal characteristics of the Strait of Magellan. In *Continental Shelf Research* (Vol. 8, Issue 8). [https://doi.org/10.1016/0278-4343\(88\)90056-8](https://doi.org/10.1016/0278-4343(88)90056-8), 1988.
- 1075 Meier, W. J.-H., Griesinger, J., Hochreuther, P., and Braun, M. H.: An updated multi-temporal glacier inventory for the Patagonian Andes with changes between the Little Ice Age and 2016. *Frontiers in Earth Science*, 6, 62. <https://doi.org/10.3389/feart.2018.00062>, 2018.
- Mellor, G. L. and Yamada, T.: Development of a turbulence closure model for geophysical fluid problems, *Rev. Geophys.*,
1080 20, 851–875, <https://doi.org/10.1029/RG020i004p00851>, 1982.
- Moffat C.: Wind-driven modulation of warm water supply to a proglacial fjord, Jorge Montt Glacier, Patagonia. *Geophysical Research Letters*, 41, 3943–3950. <https://doi.org/10.1002/2014GL060071>, 2014

- Munk, W. H. and Anderson, E. R.: Notes on a theory of the thermocline, *J. Mar. Res.*, 7, 276–295, 1948.
- McCabe, R. M., MacCready, P., and Pawlak, G.: Form drag due to flow separation at a headland, *Journal of Physical Oceanography*, 2136–2152 pp., <https://doi.org/10.1175/JPO2966.1>, 2006.
- Officer, C.B. *Physical oceanography of estuaries*, Wiley, New York, 465 pp, 1976.
- Osborn, T. R. (1980). Estimates of the local rate of vertical diffusion from dissipation measurements. *Journal of physical oceanography*, 10(1), 83-89.
- Pacanowski, R. C., & Philander, S. G. H. : Parameterization of vertical mixing in numerical models of tropical oceans. *J. phys. Oceanogr*, 11(11), 1443-1451, 1981.
- Palma, E. D., and Matano, R. P.: A numerical study of the Magellan Plume. *Journal of Geophysical Research: Oceans*, 117(5), 5041. <https://doi.org/10.1029/2011JC007750> , 2012.
- Pan, H., Wei, Y., Xu, T., & Wei, Z. A time-varying tidal form factor considering the 18.61-year nodal cycle. *Estuarine, Coastal and Shelf Science*, 305, 108868. <https://doi.org/10.1016/j.ecss.2024.108868>, 2024.
- Pérez-Santos, I., Seguel, R., Schneider, W., Linford, P., Donoso, D., Navarro, E., Amaya-Cárcamo, C., Pinilla, E., and Daneri, G.: Synoptic-scale variability of surface winds and ocean response to atmospheric forcing in the eastern austral Pacific ocean. *Ocean Science*, 15(5), 1247–1266. <https://doi.org/10.5194/os-15-1247-2019>, 2019.
- Peters, H., Gregg, M. C., and Toole, J. M.: On the parameterization of equatorial turbulence, *J. Geophys. Res.-Oceans*, 93, 1199–1218, <https://doi.org/10.1029/JC093iC02p01199>, 1988.
- Pickard, G. L.: Some Physical Oceanographic Features of Inlets of Chile, *Journal of the Fisheries Research Board of Canada*, 28, 1077–1106, <https://doi.org/10.1139/F71-163>, 1971.
- Pickard, G. L., and Stanton, B. R.: Pacific Fjords - A Review of Their Water Characteristics. *Fjord Oceanography*, 1–51. https://doi.org/10.1007/978-1-4613-3105-6_1, 1980.
- Pinilla, E., Castillo, M. I., Pérez-Santos, I., Venegas, O., and Valle-Levinson, A.: Water age variability in a Patagonian fjord. *Journal of Marine Systems*, 210. <https://doi.org/10.1016/j.jmarsys.2020.103376>, 2020.

- Rabinovich, A. B.: Seiches and harbor oscillations. In Handbook of coastal and ocean engineering (pp. 243-286), 2018.
- 1120 Rodi, W.: Examples of calculation methods for flow and mixing in stratified fluids, J. Geophys. Res.-Oceans, 92, 5305–5328, <https://doi.org/10.1029/JC092iC05p05305>, 1987.
- Ribeiro, C. H. A., Waniek, J. J., and Sharples, J.: Observations of the spring-neap modulation of the gravitational circulation in a partially mixed estuary, in: Ocean Dynamics, 299–306, <https://doi.org/10.1007/s10236-003-0086-z>, 2004.
- 1125 Rivera, A., Bravo, C., and Buob, G.: Climate change and land Ice. International Encyclopedia of Geography: People, the Earth, Environment and Technology. <https://doi.org/10.1002/9781118786352>, 2017.
- Reche, P., Artal, O., Pinilla, E., Ruiz, C., Venegas, O., Arriagada, A., & Falvey, M. CHONOS: Oceanographic information website for Chilean Patagonia. *Ocean and Coastal Management*, 208. <https://doi.org/10.1016/j.ocecoaman.2021.105634>, 2021.
- 1130 2021.
- Ruiz, C., Artal, O., Pinilla, E., & Sepúlveda, H. H. Stratification and mixing in the Chilean Inland Sea using an operational model. *Ocean Modelling*, 158. <https://doi.org/10.1016/j.ocemod.2020.101750>, 2021.
- 1135 Rojas-Celis, M., Castillo, M. I., Pérez-Santos, I., Barrios-Guzmán, C., Garcés-Vargas, J., Guerrero, A., Landaeta, M. F., Piñones, A., and Sepúlveda, M.: Turbulent Mixing in Patagonian Fjords and Channels, EGUsphere [preprint], <https://doi.org/10.5194/egusphere-2025-5846>, 2025.
- Ross, L., Pérez-Santos, I., Linford, P., and Díaz, P. A.: Circulation in the Guafo Mouth: The gateway to northern Patagonia. *Science of the Total Environment*, 979(March). <https://doi.org/10.1016/j.scitotenv.2025.179512>, 2025.
- 1140 Science of the Total Environment, 979(March). <https://doi.org/10.1016/j.scitotenv.2025.179512>, 2025.
- Sassi, M.G., Palma, E.D.,: Modelo Hidrodinámico del Estrecho de Magallanes. Asociación Argentina de Mecánica Computacional. Mec. Comput. XXV (16), 1461–1477, 11. <https://cimec.org.ar/ojs/index.php/mc/article/view/577/550>, 2006.
- 1145 Schneider, W., Pérez-Santos, I., Ross, L., Bravo, L., Seguel, R., & Hernández, F.: On the hydrography of Puyuhuapi Channel, Chilean Patagonia. *Progress in Oceanography*, 129, 8–18. <https://doi.org/10.1016/j.pocean.2014.03.007>, 2014.
- Silva N and CA Vargas.: Hypoxia in Chilean Patagonian Fjords. *Progress in Oceanography*. 129(Part A): 62-74 pp, 2014.
- 1150 <https://doi.org/10.1016/j.pocean.2014.05.016>, 2014.

- Soto-Riquelme, C., Pinilla, E., and Ross, L.: Wind influence on residual circulation in Patagonian channels and fjords. *Continental Shelf Research*, 254. <https://doi.org/10.1016/j.csr.2022.104905>, 2023.
- 1155 Spall, M. A., Jackson, R. H., and Straneo, F.: Katabatic Wind-Driven Exchange in Fjords. *Journal of Geophysical Research: Oceans*, 122(10), 8246–8262. <https://doi.org/10.1002/2017JC013026>, 2017.
- Stigebrandt, A.: Barotropic and baroclinic response of a semi-enclosed basin to barotropic forcing from the sea. In *Fjord oceanography* (pp. 141-164). Boston, MA: Springer US, 1980.
- 1160 Stigebrandt, A.: Hydrodynamics and circulation of fjords, https://doi.org/10.1007/978-1-4020-4410-6_247, 2012.
- Thorpe, S. A.: *The turbulent ocean*, Cambridge University Press, Cambridge, UK, 439 pp., <https://doi.org/10.1017/CBO9780511819927>, 2005.
- 1165 Torrence, C. and Compo, G. P.: A Practical Guide to Wavelet Analysis, *Bull Am Meteorol Soc*, 79, 61–78, [https://doi.org/10.1175/1520-0477\(1998\)079<0061:APGTWA>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<0061:APGTWA>2.0.CO;2), 1998.
- Valle-Levinson, A., and Blanco, J. L.: Observations of wind influence on exchange flows in a strait of the Chilean Inland Sea. In *Journal of Marine Research* (Vol. 62, Issue 5). <https://doi.org/10.1357/0022240042387565>, 2004.
- 1170 Valle-Levinson, A., Sarkar, N., Sanay, R., Soto, D., and León, J.: Spatial structure of hydrography and flow in a Chilean fjord, *Estuario Reloncaví. Estuaries and coasts*, 30(1), 113-126, 2007.
- Valle-Levinson A.: Definition and classification of estuaries. In: Valle-Levinson A(ed). *Contemporary Issues in Estuarine Physics*, Cambridge University Press 1-11 pp. <https://doi.org/10.1017/CBO9780511676567.002>, 2010
- 1175 Valle-Levinson, A., Caceres, M. A., Pizarro, O. Variations of tidally driven three-layer residual circulation in fjords. *Ocean Dynamics*, 64(3), 459–469. <https://doi.org/10.1007/s10236-014-0694-9>, 2014.
- 1180 Venayagamoorthy, S. K. and Stretch, D. D.: On the turbulent Prandtl number in homogeneous stably stratified turbulence, *J. Fluid Mech.*, 644, 359–369, <https://doi.org/10.1017/S002211200999268X>, 2010.
- Wan, D., Hannah, C. G., Foreman, M. G. G., & Dosso, S. (2017). Subtidal circulation in a deep-silled fjord: Douglas Channel, British Columbia. *Journal of Geophysical Research: Oceans*, 122(5), 4163–4182. <https://doi.org/10.1002/2016JC012022>

Wu, Q., Zhang, X., Church, J. A., and Hu, J.: ENSO-Related global ocean heat content variations. *Journal of Climate*, 32(1), 45–68. <https://doi.org/10.1175/JCLI-D-17-0861.1>, 2019.

Yang, Z. Y., Cheng, H. Q., and Li, J. F.: Nonlinear advection, Coriolis force, and frictional influence in the South Channel of
1190 the Yangtze Estuary, China, *Sci China Earth Sci*, 58, 429–435, <https://doi.org/10.1007/s11430-014-4946-9>, 2015.

Yelland, M., and Taylor, P. K.: Wind stress measurements from the open ocean. *Journal of Physical Oceanography*, 26(4), 541–558. [https://doi.org/10.1175/1520-0485\(1996\)026<0541:WSMFTO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1996)026<0541:WSMFTO>2.0.CO;2), 1996.

1195 Zheng, Q., Bingham, R., and Andrews, O.: Using Sea Level to Determine the Strength, Structure and Variability of the Cape Horn Current. *Geophysical Research Letters*, 50(21), 1–10. <https://doi.org/10.1029/2023GL105033>, 2023.




Figure A1. Regional wind climatology for the 1970-2024 period. Wind direction follows the oceanographic convention (indicating the direction towards which the wind blows).




Figure A2. a) Elephant seal trajectories within the Patagonian fjords and channels during 2024 (red dots) and 2025 (yellow dots). Comparison of CTD-SRDL b) salinity and c) temperature data with is situ observations (CIMAR) and GLORYS model outputs for the defined by the rectangle in (a). d) Absolute Salinity of a transect between the Pacific Ocean, the Tierra del Fuego inner-sea, and the Atlantic Ocean. GLORYS ocean reanalysis data ©Copernicus Marine Service.




Figure A3. MIKE 3D output model of IFOP – CHONOS. Upper and deeper circulation in the study region to determine the exchange fluxes in the Almirantazgo fjord (AF).

1225