

Evaluating glycerol dialkyl glycerol tetraether (GDGT)-based reconstructions from varved lake sediments during the Holocene

Ashley M. Abrook^{1,2}, Gordon N. Inglis², Peter G. Langdon¹, McKenzie R. Bentley², Achim Brauer³, Ian Bull⁴, Daisy Fallows^{1,2}, Paul Lincoln⁵, Antti E. K. Ojala^{6,7}, Helen L. Whelton⁴, Celia Martin-Puertas⁸

¹ School of Geography and Environmental Science, University of Southampton, Southampton, UK

² School of Ocean and Earth Science, University of Southampton, Southampton, UK

³ GFZ Helmholtz Centre for Geosciences, Potsdam, Germany

⁴ School of Chemistry, University of Bristol, Bristol, UK

⁵ Department of Geography, Kings College London, London, UK

⁶ Department of Geography and Geology, University of Turku, Turku, Finland

⁷ Geological Survey of Finland, Espoo, Finland

⁸ Department of Geography, Royal Holloway University of London, Egham, UK

Correspondence to: Ashley M. Abrook (A.Abrook@soton.ac.uk)

Abstract. Advances in proxy development and proxy reconstructions within the Holocene increasingly reveal climatic complexity. Annually laminated (varved) lacustrine records provide an opportunity to assess this complexity at high temporal resolution. Organic geochemical proxies offer the potential for quantitative palaeoclimate reconstruction, however, their application to different varved lake settings remains limited. Here we explore the use of isoprenoid and branched glycerol dialkyl glycerol tetraethers (GDGTs) preserved in varved sediments as proxies for temperature. We analyse three Holocene-aged annually laminated lacustrine records spanning different climate regions and lake settings across Europe (Diss Mere, UK; Nautajärvi, Finland; Meerfelder Maar, Germany); including intervals at multi-decadal resolution within the mid-Holocene. We show that isoprenoid GDGT distributions in annually laminated sediment sequences are largely derived from methanogenic Euryarchaeota and yield unreliable lake surface temperature reconstructions. Conversely, branched GDGT reconstructions show good coherence with instrumental temperature data in mid- and high-latitude environments. Although we show that lake or catchment-specific processes, including differences in processes linked to varve sedimentation, hypoxia, sediment influx and landscape development, can influence brGDGT distributions in varved lakes, the trends and range of variability of our brGDGT derived Holocene temperature reconstructions broadly agree with regional European Holocene reconstructions. This suggests that temperature exerts a first-order control on the methylation of brGDGTs in varved lake sequences. Combined with precise varve chronologies, these biomarker records can be used to generate highly resolved climate data across the Holocene.

Keywords: isoprenoid and branched GDGTs, biomarkers, temperature reconstructions, varves, Holocene

1 Introduction

35 The Holocene (11.7 ka BP to present) has previously been considered as climatically stable when viewed alongside the Glacial-Interglacial cycles of the Quaternary. However, with increasing temporal and spatial resolution of proxy data, the Holocene reveals: a) rapid (multi-decadal) shifts in climate systems associated with abrupt climate events (e.g., Roland et al., 2015; Parker and Harrison, 2022; Candy et al., 2025), b) a disconnect between modelled climate evolution and proxy-based climate
40 of climatic variability through time (e.g., the North Atlantic Oscillation; Hernández et al., 2020). To disentangle this climatic complexity, highly resolved climate archives are required. Whilst internationally important archives with excellent temporal resolution exist, frequently these are limited in terrestrial palaeo- studies due to temporal limitations (e.g., tree rings; Briffa et al., 2004) and/or are restricted in their geographic location (e.g., ice-cores; Rasmussen et al., 2014, and speleothems; Wang et al., 2005). A complementary archive that offers the potential to disentangle climatic complexity across different regions over
45 long timescales are annually laminated (varved) lake sediments, with an absence of sedimentary mixing. These sediment sequences contain precise chronologies obtained *via* annual layer counting tied to radiocarbon and/or tephra frameworks (e.g., Lane et al., 2013; Zolitschka et al., 2015, Wulf et al., 2015). The analysis of climatic proxies within these sediments (e.g., Ojala et al., 2008; Larocque-Tobler et al., 2015; Luoto and Ojala et al., 2016; Lincoln et al., 2025) offers the potential for highly resolved (annual to multi-decadal) past climate reconstructions across wide spatial scales. Here, biomarkers are of
50 particular interest as they can provide quantified climatic and environmental observations from lakes (Rach et al., 2014) yet the degree to which biomarkers faithfully record past temperature in varved lake sediment archives remains an open question.

Isoprenoid and branched glycerol dialkyl glycerol tetraethers (GDGTs) are membrane-spanning lipids and are frequently used to reconstruct climate variability in lake sediments (e.g., Tierney and Russell, 2009; Woltering et al., 2014; Peuple et al., 2022;
55 Robles et al., 2023; Baxter et al., 2024; Zander et al., 2024). Isoprenoid GDGTs are archaeal lipids composed of two isoprenoid alkyl chains, with up to four cyclopentane moieties and one cyclohexane moiety identified in lakes (Schouten et al., 2013). The degree of isoGDGT cyclisation is captured *via* the TetraEther index of 86 carbons (TEX₈₆; Schouten et al., 2002) which is calibrated to surface water temperature in lacustrine settings (Powers et al., 2010). Branched GDGTs are bacterial lipids (Sinninghe Damsté et al., 2000) composed of two branched alkyl chains, varying in the number of methyl branches (4 to 6)
60 and cyclopentane moieties (0 to 2) (Weijers et al., 2007). The degree of methylation is captured *via* the methylation index of branched tetraethers (MBT'_{SME}; De Jonge et al., 2014), which in lakes is calibrated to either mean annual air temperature (MAAT; Russell et al., 2018), mean temperature of months above freezing (MAF; Martinez-Sosa et al., 2021; Raberg et al., 2021) and/or mean summer temperatures (MST; Pearson et al., 2011; 2025).

65 Whilst GDGTs have been applied extensively to Holocene-aged lake sediments, few studies have evaluated the environmental controls on GDGT distributions within annually laminated lake sediments in Europe (i.e., Weber et al., 2018; Zander et al.,

2024; Otiniano et al., 2024). These settings exhibit specific limnological processes, including hypolimnetic anoxia/hypoxia, thermal stratification and individual mixing regimes that may exert strong control upon the distribution of different GDGT compounds. Some of these features (e.g., seasonal anoxia), alongside lake morphometry and catchment size, are known to affect the ecology of archaeal and bacterial groups that synthesise isoprenoidal and branched GDGTs, respectively (Weber et al., 2018; van Bree et al., 2020; Baxter et al., 2024; Zander et al., 2024). As these proxies are becoming more common in varve-based palaeoclimate research, an evaluation is required to establish whether annually laminated lake records from different climatic settings and/or the processes that govern lamination sedimentology (e.g., clastic, organic, diatomaceous, carbonate lamina) impact GDGT distributions and associated metrics.

Here, we use three different annually laminated lake records from Europe to establish whether GDGT distributions in varved lake sediments are suitable for generating climate reconstructions during the Holocene. We assess whether processes that 1) are important for varve formation and preservation (e.g., stratification and anoxia) and 2) control varve composition, impact GDGT distributions and metrics. Isoprenoid and branched GDGTs are analysed at multi-decadal resolutions across key intervals of the Holocene, including the Holocene Thermal Maximum (ca. 6.5 – 5 ka BP) and the last 300 years. We compare our results with instrumental data and other independent proxies to provide a unique assessment of GDGT distributions in varved lake systems and evaluate the potential of using isoGDGTs and brGDGTs to reconstruct highly resolved temperature records across the Holocene and beyond.

2 Materials and Methods

2.1 Site descriptions

2.1.1 Diss Mere

Diss Mere (52° 22'N, 1° 6'E; Fig. 1) is a small (0.034 km²), 6 m deep, presently eutrophic lake with alkaline waters and a chalk bedrock (Martin-Puertas et al., 2021). The site occupies a region dominated by maritime climate. During lake stratification, pH is high in the epilimnion (values of ~9) and lower in the hypolimnion (~7) (Boyall et al., 2023). The lake contains no inflow or outflow, experiences minimal groundwater input (Boyall et al., 2023) and has an estimated catchment to lake area ratio (CA:LA) of 44. The DISS-16 record exhibits 15 m of sedimentation covering the last 10 thousand years (Martin-Puertas et al., 2021) with varves preserved between 10 and 2 ka BP. The varve couplets are calcite-organic (Fig. 1; including diatoms in the organic component) with some interannual variability (Martin-Puertas et al., 2021). The calcite layer is representative of summer sedimentation with organic constituents deposited during autumn to spring following resuspension of organic remains and wind-induced mixing of the water column (Martin-Puertas et al., 2021; Boyall et al., 2023).

2.2.2 Nautajärvi

Nautajärvi (61°48'N 24°24'E; Fig. 1) is a small (0.17 km²), ~20 m deep mesotrophic lake, exhibiting slightly acidic lake waters (modern pH of 5.8 – 6; Korkonen et al., 2017). The site occupies a region of sub-Arctic climate. The site sits in a tiered lake system within the Ävöntjärvi drainage basin and is fed by streams and Ristijärvi to the north and is drained by Lake Pitkävesi to the south (Ojala and Alenius, 2005) with a CA:LA of 71. The NAU-23 stratigraphy (Lincoln et al., 2025) contains 7.3 m of sedimentation, with 9,829 varves covering the upper 6.74 m (Ojala and Saarinen, 2002; Lincoln et al., 2025). The varve couplets are clastic-organic (Fig. 1), with the clastic layer forming in spring as a product of winter precipitation and snowmelt run off once the lake becomes ice-free, with the organic layer forming in the summer to winter months (Ojala and Alenius, 2005). Inverse stratification is observed during the winter months with anoxic bottom waters at this time (Ojala and Alenius, 2005; Ojala et al., 2013).

2.2.3 Meerfelder Maar

Meerfelder Maar (50°06'N, 6°45'E; Fig. 1) is a relatively small (0.25 km²), 18 m deep, eutrophic maar lake which occupies a predominantly maritime climate region (albeit with some continental influences) within mainland Europe. The lake has a CA:LA of 6. The 11.71 m long MFM-09 sequence contains 7.45 m of sediment across the Holocene (Martin-Puertas et al., 2012). Varves are preserved from 11.6 to ca. 1.5 ka BP (Martin-Puertas et al., 2012; 2017). The Holocene varve structures preserved at Meerfelder Maar are diatomaceous (Fig. 1), with a spring and summer layer composed of a monospecific diatom bloom and the autumn and winter layer composed of organic and minerogenic detritus (Brauer et al., 2000; Martin-Puertas et al., 2012). Hypolimnetic anoxia has previously been proposed at Meerfelder Maar, with treatments performed to increase bottom water oxygen conditions (Nürnberg et al., 2007). This process disturbed stratigraphy of the upper 500 years of the record.

Table 1: Lake descriptors from each site. * This is the assumed mixing regime for periods of varve formation across each site, at Diss Mere and Meerfelder Maar this is likely different to today: ¹ Martin-Puertas et al. (2021); ² Lincoln et al. (2025); ³ Nürnberg et al. (2007). ** The origin of Diss Mere is debated, with possible origins dating to the Anglian stage (Bailey, 2005).

Lake	Climatology	Lake origin	Mixing regime*	Varve composition
Diss Mere	Maritime	Thermokarst**	Meromictic ¹ (no lake water overturn)	Calcite – organic
Nautajärvi	Sub-Arctic	Glacial lake	Dimictic ² (two lake water overturn periods annually)	Clastic – organic
Meerfelder Maar	Maritime (with continental influences)	Volcanic (maar lake)	Meromictic ³ (no lake water overturn)	Diatomaceous – detrital




Figure 1: Location of each of the sites in Europe: a) Diss Mere, UK; b) Nautajärvi, Finland; c) Meerfelder Maar, Germany. Site boxes reveal lake bathymetry with depths and inflows/outflows. Varve depositional models are added to each box which demonstrate the main varve facies from each location, with three varve microfacies at Diss Mere and one each at Nautajärvi and Meerfelder Maar. Letters denote different cores from a) and b). Diss Mere adapted from Boyall et al. (2023); Nautajärvi adapted from Lincoln et al. (2025).

2.2.4 Chronology

Published chronologies are derived from annual layer counting and tied to either a radiocarbon and/or tephra framework (Martin-Puertas et al., 2012; 2021; Meerfelder Maar, Diss Mere respectively) or from the cross-correlation of marker layers between an annual layer-counted sequence and newly derived sediments (e.g., Ojala and Saarinen, 2002; Ojala and Alenius, 2005; Lincoln et al., 2025; Nautajärvi). For each sequence the chronological uncertainties are decadal in scale (see Martin-Puertas et al., 2012; 2021; Lincoln et al., 2025). Following Abrook et al. (2025), we present reconstructions as ka BP (as cal. ka BP at Diss Mere and Meerfelder Maar and ka varve years BP for Nautajärvi).

2.2 Biomarker analysis

135 Biomarker samples, 0.5 cm resolution at variable intervals (at an average of 8 – 10 varve years per sample over more densely
sampled core sections), were extracted from the across the Holocene and from modern sediments (covering the last ca. 300
years) from Diss Mere (n = 56) and Nautajärvi (n = 42). Due to historic hypolimnetic treatments at Meerfelder Maar (n = 31)
the last 500 years of sedimentation have been impacted. Analyses were therefore not performed on modern sediments at this
140 site. To obtain total lipid extracts (TLE), approximately 1 – 2g of wet sediment per sample was freeze dried and extracted
using Accelerated Solvent Extraction (Dionex ASE 350) with dichloromethane:methanol (9:1, v/v) at 100°C over 3 x 15 min
extraction cycles at 1,500 PSI. The TLE was dried under a stream of nitrogen (N₂) and separated into different compound
classes via silica gel column chromatography (F1; aliphatic, F2; aromatic, and F3; polar) using hexane:dichloromethane (9:1,
v/v; F1), hexane:dichloromethane (1:1, v/v; F2), and dichloromethane:methanol (1:1, v/v; F3) with a 4ml solvent volume for
each fraction. The polar fraction was re-dissolved in hexane:isopropanol (99:1, v/v) and passed through a 0.45 µm PTFE
145 (polytetrafluoroethylene) filter and analysed by high-performance liquid chromatography (HPLC) mass spectrometry (MS)
using an Agilent 1260 HPLC coupled to an Agilent 6130 single quadrupole MSD following the method outlined in Hopmans
et al. (2016) (supplementary information). Peak identification and GDGT integration followed a manual process with
identification aided using in-house GDGT laboratory standards after scanning for common mass ranges for GDGT analysis
(even values between m/z 1302 – 1292 for isoGDGTs and between (1022 – 1018, 1036 – 1032 and 1050 – 1046 for brGDGTs).

150 2.3 Biomarker analysis

2.3.1 GDGT-based temperature proxies

TEX₈₆ is defined following Schouten et al. (2002).

$$TEX_{86} = \frac{(GDGT-2 + GDGT-3 + \text{crenarchaeol isomer})}{(GDGT-1 + GDGT-2 + GDGT-3 + \text{crenarchaeol isomer})} \quad (1)$$

TEX₈₆ is calibrated to Lake Surface Temperature (LST) following Powers et al. (2010). See supplementary information.

155 $LST = 55.2 \cdot \text{TEX}_{86} - 14$ (2)

We use the MBT'_{SME} index following De Jonge et al. (2014).

$$MBT'_{SME} = \frac{(Ia + Ib + Ic)}{(Ia + Ib + Ic + IIa + IIb + IIc + IIIa)} \quad (3)$$

Where alphanumeric values relate to different brGDGT structures and the number of cyclopentane moieties and/or methyl
branches. MBT'_{SME} values are converted to mean temperature of months above freezing (MAF; Martinez-Sosa et al., 2021)
160 using a Bayesian calibration (BayMBT₀) developed for lakes (e.g., Martinez-Sosa et al., 2021). Present day mean annual
temperatures are used as prior means at each site: 10.5°C (Diss Mere); 5.3°C (Nautajärvi) and 9.9°C (Meerfelder Maar). Prior
standard deviation is set to 10°C.

We also use the multivariate calibration approach of Raberg et al. (2021).

$$MAF = 92.9 (\pm 15.98) + 63.84 (\pm 15.58) \cdot flb_{meth}^2 - 130.51 (\pm 30.73) \cdot flb_{meth} - 28.77 (\pm 5.44) \cdot flIa_{meth}^2 - 72.28 (\pm 17.38) \cdot flIb_{meth}^2 - 5.88 (\pm 1.36) \cdot flIc_{meth}^2 + 20.89 (\pm 7.69) \cdot flIIa_{meth}^2 - 40.54 (\pm 5.89) \cdot flIIa_{meth} - 80.47 (\pm 19.19) \cdot flIIb_{meth} \quad (4)$$

2.3.2 GDGT-based lake property reconstructions

The cyclisation of branched tetraethers (CBT) index (Weijers et al., 2007) that includes both 5- and 6-methyl isomers (De Jonge et al., 2014) is used to infer lake pH:

$$CBT' = {}^{10} \log \left(\frac{(Ic + IIa' + IIb' + IIc' + IIIa')}{(Ia + IIa + IIIa)} \right) \quad (5)$$

We subsequently convert CBT' values to pH using the calibration from Russell et al. (2018).

$$pH = 8.95 + 2.65 \cdot CBT' \quad (6)$$

The isomer ratio captures the relative abundance of 5-methyl vs 6-methyl brGDGTs (De Jonge et al., 2015) and is defined as:

$$IR_{6ME} = \frac{(IIa' + IIb' + IIc' + IIIa' + IIIb' + IIIc')}{(IIa + IIa' + IIb + IIb' + IIc + IIc' + IIIa + IIIa' + IIIb + IIIb' + IIIc + IIIc')} \quad (7)$$

Deviations from the expected temperature – MBT_{5ME} relationship are observed in lake sediments when the IR_{6ME} > 0.5 (Bauersachs et al., 2024).

We use the methane index (MI) to evaluate anaerobic methanotrophy within the lake water column and/or sediments, whereby sediments with high values (> 0.3 - 0.5) suggest intense anaerobic methane oxidation (Zhang et al., 2011).

$$MI = \frac{(GDGT-1 + GDGT-2 + GDGT-3)}{(GDGT-1 + GDGT-2 + GDGT-3 + crenarchaeol + crenarchaeol\ isomer)} \quad (8)$$

Crenarchaeol and its isomer are solely produced by *Nitrososphaeria* (Zhang et al., 2011; Schouten et al., 2013), whereas GDGT-0 can also be produced by methanogenic archaea (Euryarchaeota) (Auget et al., 2009). The relative abundance of GDGT-0 vs crenarchaeol (i.e. % GDGT-0 as Sinninghe Damsté et al. (2012); related to GDGT-0/cren) is thus used to evaluate methanogenesis across each of the different lakes.

$$\% GDGT - 0 = (GDGT - 0 / (GDGT - 0 + crenarchaeol)) \cdot 100 \quad (9)$$

Where values > 67% are considered to have a substantial methanogen component that could render TEX₈₆ reconstructions unreliable (e.g., Blaga et al., 2009; Inglis et al., 2015).

2.3.3 GDGT-based organic source proxies

- 190 The branched vs isoprenoid tetraether (BIT) index (Hopmans et al., 2004; Baxter et al., 2021) evaluates the relative proportion of terrestrial vs marine organic matter (OM) within individual samples and is defined as:

$$BIT = \frac{(Ia + IIa + IIa' + IIIa + IIIa')}{(Ia + IIa + IIa' + IIIa + IIIa' + \text{crenarchaeol})} \quad (10)$$

Where high values (e.g., 1) traditionally reflect high terrestrial input and low values (e.g., 0) indicate high marine/aquatic input.

- 195 The ratio of hexamethylated and pentamethylated compounds (Xiao et al., 2016) is used to assess input of terrestrial vs marine-derived brGDGTs, where $\Sigma IIIa / \Sigma IIa$ values < 0.59 suggest terrestrial provenance and high values > 0.92 represents a marine/aquatic provenance (Xiao et al., 2016).

- Finally, we use the branched and isoprenoid GDGT machine learning classification algorithm (BIGMaC) to constrain GDGT
200 source environments (Martinez-Sosa et al., 2023). The algorithm assesses the distribution of all isoGDGTs and brGDGTs and establishes (dis)similarity to GDGT distributions from a global marine, soil, lake, and peat calibration dataset.

2.4 Micro X-ray Fluorescence (μ -XRF) core scanning data

- To assist in disentangling environmental processes, geochemical element profiles of each sequence (DISS-16, NAU-23, MFM-09) have been acquired directly from split-core surfaces every 0.2 mm using an ITRAX μ -XRF core scanner at GFZ-Potsdam.
205 Element data has already been published (Martin-Puertas et al., 2012; Boyall et al., 2024; Lincoln et al., 2025), however, to aid interpretations we re-sample those data to match the ages of the GDGT data.

2.5 Data analyses

- Ternary diagrams were created in R through proportional abundances of tetra-, penta- and hexamethylated compounds using the 'ggtern' package (Hamilton and Ferry, 2018). Unconstrained and constrained ordination (principal components analysis;
210 PCA, and redundancy analysis; RDA) was performed on standardised brGDGT fractional abundance data. We used PCA and RDA owing to short gradient lengths observed within the GDGT data, which were determined in R. For RDA, explanatory variables consisted of the major elements from centered log ratio μ -XRF data (e.g., Fe_{clr}) from each lake. PCA and RDA analysis was performed in R using the 'factoextra' and 'Vegan' packages (Kassambara, 2023; Oksanen et al., 2024). Correlation plots are shown in supplementary information.

3.1 GDGT distributions and GDGT-based environmental metrics

IsoGDGTs were identified in all samples from Diss Mere, Meerfelder Maar and Nautajärvi. GDGT-0 dominates the isoGDGT assemblage at all three site locations, across modern and Holocene samples (mostly > 90% at Diss Mere and Meerfelder Maar, > 80% at Nautajärvi), with a lower relative abundance of GDGT-1 to -3, crenarchaeol and its isomer (Fig. 2). Each site exhibits
220 high MI values with values averaging 0.73 ± 0.21 , 0.70 ± 0.10 and 0.64 ± 0.09 , respectively (Fig. 5, 6, 7).

BrGDGTs were also identified in all samples. However, there are differences in the relative proportion of tetra-, penta- and hexamethylated brGDGTs between sites (Fig. 3). Nautajärvi and Meerfelder Maar have a higher contribution of 5-methyl brGDGTs than Diss Mere, with a higher relative proportion of penta- and hexamethylated compounds at Nautajärvi (e.g.,
225 brGDGT-IIa, IIIa). Diss Mere and Meerfelder Maar have a larger relative proportion of 6-methyl brGDGTs (e.g., brGDGT-IIa', IIIa') than Nautajärvi. This is reflected in low average IR_{6ME} values at Nautajärvi (0.15 ± 0.04), with higher values at Meerfelder Maar (0.41 ± 0.03) and Diss Mere (0.60 ± 0.06) (Figs. 5, 6, 7). The $\Sigma\text{IIIa} / \Sigma\text{IIa}$ ratio averages 0.74 ± 0.07 at Diss Mere, 0.64 ± 0.07 at Nautajärvi, and 0.55 ± 0.08 at Meerfelder Maar.

230 The BIGMaC algorithm classified 100% of GDGT distributions from Diss Mere and Meerfelder Maar and 67% of distributions from Nautajärvi as 'lake-type'. The remaining 33% of GDGTs from Nautajärvi (i.e., all modern [< 300 year] and two late-Holocene samples) are classified as 'peat-type'. Across all three sites the BIT index is very high and is > 0.92 (Fig 5, 6, 7).

3.2 Statistical analyses

PC1 and PC2 explain 39.3% and 25.7% of variation at Diss Mere, 56.1% and 23.9% of variation at Nautajärvi, and 38.8% and
235 23.2% of variation at Meerfelder Maar, respectively (Fig. 4). PCA shows that 5- and 6-methyl brGDGTs are separated at Diss Mere but not at Nautajärvi or Meerfelder Maar. At the latter two sites, however, the brGDGTs with the largest fractional abundance (i.e., brGDGT-Ia, IIa) are separated from the remainder of brGDGTs. PCA also reveals that at Diss Mere the modern and Holocene samples occupy a similar ordination space whilst at Nautajärvi the modern samples are adjunct from the remainder of the data.

240 RDA1 and RDA2 explain 52.2% and 32.6% of variation at Diss Mere, 84.1% and 9.1% of variation at Nautajärvi, and 51.9% and 20.8% of variation at Meerfelder Maar, respectively. At each site, redox sensitive elements appear important (e.g., Fe_{clr}, Mn_{clr}, S_{clr}) with additional importance given to the relationship between authigenic (e.g., Ca_{clr}) and allogenic components (K_{clr} and Ti_{clr}) between sites.




Figure 2: Fractional abundances of different GDGT molecules in the different lakes. a) isoprenoid GDGT distributions; b) branched GDGT distributions.




Figure 3: Ternary plots showing a) the relative proportions of tetramethylated, pentamethylated and hexamethylated compounds from Diss Mere, Meerfelder Maar and Nautajärvi, respectively against the same data from a global brGDGT peat and soil dataset (grey; Dearing Crampton-Flood et al., 2020); a global lake dataset (black; Martinez-Sosa et al., 2021); Arctic lakes (green; Raberg et al., 2021) and Alpine lakes (red; Bauersachs et al., 2024); and b) an inset figure of a).

3.3 Temperature and pH reconstructions

TEX₈₆ has a larger range of values at Diss Mere (average of 0.41 ± 0.13) compared to Meerfelder Maar (0.30 ± 0.04) and Nautajärvi (0.37 ± 0.08) (Figs. 5, 6, 7). When converted to LST the data from each site appears variable, with greatest range in LST observed at Diss Mere (-9.0°C to 25°C), Nautajärvi (1°C to 14°C) and Meerfelder Maar (-1°C to 7°C) (S1).

The lowest average MBT'_{SME} values are from Nautajärvi (0.36 ± 0.02) whilst Diss Mere (0.44 ± 0.04) and Meerfelder Maar (0.45 ± 0.02) exhibit higher values (Figs. 5, 6, 7). brGDGT temperatures using the Bayesian (Martinez-Sosa et al., 2021) and multivariate calibration (Raberg et al., 2021) approaches produce mostly similar trends across each site. Lowest absolute temperatures are generated from Nautajärvi with temperatures between 8°C and 12°C across all samples. Reconstructed temperatures from Diss Mere range from 9°C to 15°C . At Meerfelder Maar temperature estimates range between 11 to 14°C .

The CBT' index is highest at Diss Mere (0.05 ± 0.07) and lowest at Nautajärvi (-0.94 ± 0.15). This yields higher pH reconstructions at Diss Mere (9.08 ± 0.19) and lower pH estimates at Nautajärvi. (6.51 ± 0.39). Meerfelder Maar exhibits an intermediary pH reconstruction (8.11 ± 0.15) (Figs. 5, 6, 7).




Figure 4: Principal components analysis (PCA) left, and redundancy analysis (RDA) right. a) and b) detail the plots from Diss Mere; c) and d) from Nautajärvi; and e) and f) from Meerfelder Maar. a) and c) show modern (last 300-years) and Holocene sample distributions within the PCA plots. Explanatory variables in the RDA plots are major elements from μ -XRF data (from Martin-Puertas et al. (2012); Boyal et al. (2024); Lincoln et al. (2025)) with numerical points showing the median age of samples.

4.0 Discussion

4.1 IsoGDGTs influenced by methanogenesis in varved lake systems

In lacustrine environments, the TetraEther indeX of 86 carbons (TEX₈₆) has previously been used to infer Lake Surface Temperature (LST; Woltering et al., 2012). This assumes that isoGDGTs are mostly derived from ammonia-oxidizing archaea of the class *Nitrososphaeria*. However, methanogenic archaea may also synthesise smaller quantities of isoGDGTs with up to three cyclopentane moieties (Weijers et al., 2006), although this has not been confirmed in culture studies (Schouten et al., 2013). This implies that input of methanogenic archaea could bias lacustrine TEX₈₆ values (Blaga et al., 2009), especially in varved lakes where seasonal anoxia/hypoxia is a persistent feature. Here, the fractional abundance of GDGT-0 is elevated at all sites (Diss Mere > 0.84 [mostly > 0.9], Nautajärvi > 0.81 [excluding one sample of 0.62], and Meerfelder Maar > 0.94) and associated with very high % GDGT-0 values (> 67 %; see Inglis et al., 2015). These values suggest that isoGDGTs are not produced by ammonia-oxidizing archaea but instead synthesized mostly by methanogens.

Previous studies show that lakes that exhibit hypolimnetic anoxia have a higher abundance of GDGT-0 in sediments compared to the water column, implying that GDGT-0 is largely derived from methanogens present within the sediment (Blaga et al., 2009). However, in large lakes, Baxter et al. (2021) and Sinninghe Damsté et al. (2012) demonstrate that GDGT-0 can also be synthesised in high concentrations below the oxycline, whereas *Nitrososphaeria* synthesized abundant crenarchaeol in relatively oxygenated upper waters. No sediment trap or water column data is available here, but, as each lake exhibits hypolimnetic anoxia/hypoxia (e.g., Brauer et al., 2000; Ojala and Alenius, 2005; Martin-Puertas et al., 2021; Boyall et al., 2023), it is likely that isoGDGTs are largely produced by methanogenic Euryarchaeota. Given the inconsistent variability in LST between sites (Section 3.3), and, in some instances, an extreme range for Holocene climates (e.g., at Diss Mere, with reconstructions from -9.0°C to 25°C; S1), this suggests that a methanogen overprint is highly likely.

Anaerobic methanotrophs (ANME) can also synthesise GDGT-0 to GDGT-3 and may contribute to the observed high Methane Index (MI) values (Zhang et al., 2011; Kim and Zhang, 2023). These organisms are typically abundant in sites characterized by anaerobic oxidation of methane (AOM, e.g., cold seeps, mud volcanoes, gas hydrates) and require high concentrations of sulphate in the overlying water column (Zhao et al., 2024). Although sulphate is often variable in lakes, sulphate reduction can be maintained in freshwater systems (Segarra et al. 2015). At each location in our study, μ -XRF data reveals varying contributions of elemental sulphur (with relationships with GDGTs; S3 – S8) which, given hypoxic conditions, can lead to sulphate reduction (albeit strongly reducing conditions may not be maintained throughout the annual cycle; Lincoln et al., 2025). These features, alongside high MI values, may suggest AOM-impacted sediments. High MI and % GDGT-0 values are not surprising as the two proxies are linearly correlated in marine sediments (Inglis et al., 2015) and




Figure 5: Main GDGT reconstructions from Diss Mere. Shown are a) TEX₈₆; b) MBT'_{5ME} (De Jonge et al., 2014); c) CBT' (De Jonge et al., 2014); d) reconstructed pH (Russell et al., 2018); e) $\Sigma IIIa / \Sigma IIa$ ratios with cut offs from Xiao et al. (2016); f) IR_{6ME} (De Jonge et al., 2014; Bauersachs et al., 2024); g) the methane index (Zhang et al., 2011); h) % GDGT-0 and i) the branched vs isoprenoid tetraether index (BIT).




Figure 6: Main GDGT reconstructions from Nautajärvi. Shown are a) TEX_{86} ; b) $\text{MBT}'_{5\text{ME}}$ (De Jonge et al., 2014); c) CBT' (De Jonge et al., 2014); d) reconstructed pH (Russell et al., 2018); e) $\Sigma\text{IIIa} / \Sigma\text{IIa}$ ratios with a cut off from Xiao et al. (2016); f) $\text{IR}_{6\text{ME}}$ (De Jonge et al., 2014); g) the methane index (Zhang et al., 2011); h) % GDGT-0 and i) the branched vs isoprenoid tetraether index (BIT).




Figure 7: Main GDGT reconstructions from Meerfelder Maar across the Holocene. Shown are a) TEX₈₆; b) MBT'_{5ME} (de Jonge et al., 2014); c) CBT' (de Jonge et al., 2014); d) reconstructed pH (Russell et al., 2018); e) $\Sigma\text{IIIa} / \Sigma\text{IIa}$ ratios with a cut off from Xiao et al. (2016); f) IR_{6ME} (De Jonge et al., 2014; Bauersachs et al., 2024); g) the methane index (Zhang et al., 2011); h) % GDGT-0 and i) the branched vs isoprenoid tetraether index (BIT).

lakes (Collins et al., 2025). However, the relative contribution of methanogens and methanotrophs to the archaeal community is unknown and would require further isotopic analysis (e.g., Segarra et al., 2015). Overall, it is apparent that methane cycling can bias isoGDGT distributions in lakes (this study; Blaga et al., 2009; Weijers et al., 2011; Zhang et al., 2016; Baxter et al., 2021) and we caution against the use of TEX₈₆ in varved lake systems.

4.2 BrGDGTs likely derived from *in-situ* lacustrine production in varved lakes

BrGDGTs in lakes can have multiple sources including input *via* soils/peats (e.g., Peterse et al., 2014) and *in situ* production within the lake water column (e.g., Buckles et al., 2014). It is therefore important to understand the provenance of brGDGTs

350 in lacustrine environments as different soil, peat, and lake calibrations (e.g., Russell et al., 2018; Dearing Crampton-Flood et al., 2020; Martinez-Sosa et al., 2021; Raberg et al., 2021) can yield temperature estimates that can differ by up to 10°C (Tierney et al., 2010). This is likely to be important in annually laminated lakes where varve sediment composition may be reflective of changes in lacustrine or catchment productivity and changes in autochthonous process (e.g., primary productivity in the water column and suspension settling of mineral or organic particles vs in-wash).

355 Branched vs isoprenoid tetraether (BIT) values are historically interpreted to reflect changes in delivery of soil/peat derived organic matter into aquatic systems (e.g., Blaga et al., 2009; Sinninghe Damsté et al., 2009; Schouten et al., 2013). Across all three lakes, the BIT index is high for all samples (Holocene and modern) (> 0.9), which implies consistently high input of soil or peat organic matter into the lake. Each lake is relatively small ($< 0.25 \text{ km}^2$), and depositional models for detrital and clastic
360 lamination types indicate allochthonous input *via* catchment runoff limited to specific seasons (Meerfelder Maar, Brauer et al., 2000; Nautajärvi, Lincoln et al., 2025). Allied to this, each lake has different catchment to lake area (CA:LA) ratios which might impact BIT if viewed solely from a catchment provenance perspective. However, increasingly evidence suggests that higher BIT values can result from relatively low crenarachaeol concentrations in lakes (Loomis et al., 2011; Buckles et al., 2014; van Bree et al., 2020) and/or *in-situ* production of brGDGTs (Naeher et al., 2014; Buckles et al., 2014). Therefore, these
365 observations suggest caution when using the BIT to evaluate terrestrial vs aquatic input in annually laminated lake sediments.

To evaluate organic matter sources further, the $\Sigma\text{IIIa}/\Sigma\text{IIa}$ ratio (Xiao et al., 2016; Martin et al., 2019) can be used to help differentiate between soil vs aquatic input. This approach has mostly been applied in marine sediments (Xiao et al., 2016), but Martin et al. (2019) suggested that $\Sigma\text{IIIa}/\Sigma\text{IIa}$ ratios can also distinguish between soil vs aquatic input in lacustrine settings.
370 All Holocene samples from Diss Mere and Nautajärvi (except one sample from Nautajärvi at 1.0 ka BP) exhibit $\Sigma\text{IIIa}/\Sigma\text{IIa}$ ratios > 0.59 , which is higher than the proposed limit for soil production (Xiao et al., 2016). Modern (< 300 years) samples at Diss Mere exhibit similar values, but those from Nautajärvi range between 0.50 and 0.58, suggesting increasing input of soil-derived brGDGTs over the last few hundred years. The Meerfelder Maar record exhibits $\Sigma\text{IIIa}/\Sigma\text{IIa}$ ratios between 0.42 to 0.75, suggesting mixed contribution of soil and aquatic derived GDGTs. However, as Meerfelder Maar has the lowest CA:LA
375 ratio in this study, input solely *via* soil organic matter is unlikely. Albeit these sites likely exhibited a much larger hydrological catchment in periods of the Holocene. Nonetheless this highlights the limitations of $\Sigma\text{IIIa}/\Sigma\text{IIa}$ in varved (and typical) lake systems (O’Beirne et al., 2024).

An alternative approach involves comparing the proportion of tetra-, penta-, and hexamethylated GDGTs alongside those from
380 different environments, including soils, peats, and lakes (Dearing Crampton-Flood et al., 2020; Martinez-Sosa et al., 2021; Raberg et al., 2021; Bauersachs et al., 2024). For each site, our data reveals greater overlap with global lakes (as opposed to soils/peats), suggesting that brGDGTs in this study are largely produced within the lacustrine environment (Fig. 3). The BIGMaC machine-learning algorithm, on the basis of the fractional abundance of both isoGDGTs and brGDGTs in comparison

with a global dataset (Martinez-Sosa et al., 2023), also suggests that most samples from Diss Mere (100%), Nautajärvi (67%),
385 and Meerfelder Maar (100%) have GDGTs defined as ‘lake-type’. Only the modern samples and two late Holocene samples
(at 1.0 and 2.0 ka BP) from Nautajärvi are classified as ‘peat-type’. Taken together, this suggests that the predominant source
of brGDGTs in these laminated sediments is lacustrine.

4.3 Impact of lacustrine processes on brGDGT distributions

Seasonal stratification and hypolimnetic anoxia/hypoxia are considered key additional controls on brGDGT distributions in
390 lacustrine environments (Weber et al., 2018). In many stratified lakes, brGDGT production has been identified below the
thermocline (Buckles et al., 2014; Miller et al., 2018; Sinninghe Damsté et al., 2022) with enhanced brGDGT production in
low oxygen conditions (Weber et al., 2018; van Bree et al., 2020; Baxter et al., 2024; Zander et al., 2024). Changes in sediment
chemistry associated with individual laminations also reveals the influence of different seasonal processes in lake
environments, for example calcite precipitation in summer months and increased titanium in autumn and winter months
395 relating to changing catchment sediment flux (e.g., Marshall et al., 2012; Zolitschka et al., 2015), or varied organic carbon
content related productivity changes (e.g., Hällberg et al., 2023). To explore whether and/or how the brGDGT distributions in
lakes in this study are influenced by specific limnological processes (e.g., stratification, anoxia, in wash), we compare our data
to physical lake properties and μ -XRF measurements from the same sediment profiles within Holocene laminated core sections.

4.3.1 Redox influence upon brGDGT distributions

400 The lakes in this study are all thermally stratified in summer months. However, in terms of mixing regime, Diss Mere and
Meerfelder Maar were meromictic prior to 2 ka BP. Nautajärvi was dimictic in the early and late Holocene with episodic shifts
towards a more meromictic regime coinciding with episodes of reduced and strengthened overturning in the mid-Holocene
(Table 1; Lincoln et al., 2025). These mixing regimes influence the distribution of oxygen through the water column, with
each lake demonstrating hypoxic/anoxic bottom waters contributing to varve formation and preservation. In permanently
405 stratified (meromictic) lakes, pentamethylated and hexamethylated brGDGTs increase in abundance with depth and are
associated with anoxic conditions (Weber et al., 2018; Yao et al., 2020; van Bree et al., 2020; Baxter et al., 2024). As the lakes
in this study exhibit anoxic/hypoxic hypolimnetic conditions in the Holocene, as a result of their mixing regime (above; and
Table 1) high fractional abundances of pentamethylated (brGDGT-IIa, IIa') and hexamethylated brGDGTs (brGDGT-IIIa,
IIIa') suggests a potential redox influence on brGDGT distributions.

410

In addition, we identify relationships between brGDGTs and redox sensitive elements across each site location (i.e. sulphur
(S) and iron (Fe)). In lakes insoluble sulphides can form through conversion of sulphate by sulphate-reducing bacteria under
anoxic conditions (Luo, 2018), whilst precipitation of Fe-hydroxide is associated with oxic conditions (Davison, 1993). At
Nautajärvi, we identify a significant positive relationship between S_{clr} and brGDGT-Ia ($r = 0.45$, $p = 0.012$) and brGDGT-IIa
415 ($r = 0.54$; $p = 0.002$), and a significant negative relationship between Fe_{clr} and brGDGT-IIIa ($r = -0.52$; $p = 0.003$) (S7). Given

that we observe opposing relationships between these redox-sensitive elements (Fig. 4) alongside temperature-sensitive brGDGTs (i.e., lower relative brGDGT-IIIa with high Fe, derived from mineral precipitates linked to phases of relatively greater hypolimnetic oxidation during strengthened mixing phases; see Lincoln et al. (2025)), we suggest that hypoxia and shifts to relatively more oxygenated conditions may influence brGDGT production at Nautajärvi (e.g., Raberg et al., 2022; 420 2025).

In contrast, at Diss Mere, we observe the opposite relationship between S_{clr} and brGDGT-IIIa ($r = -0.46$; $p = 0.003$) (S3). This is also reflected in the redundancy analysis with S_{clr} being positively loaded with Fe_{clr} at Diss Mere (Fig. 4). There is also a significant correlation between $\ln(\text{Fe}/\text{Ti})$, which at Diss Mere captures the anoxic conditions of bottom waters (Boyall et al., 425 2024), and both brGDGT-IIa ($r = -0.4$; $p = 0.01$) and brGDGT-IIa' ($r = 0.46$; $p = 0.004$) (S3). Increased $\ln(\text{Fe}/\text{Ti})$ indicates increased bottom water anoxia (Boyall et al., 2024). As such, the negative relationship between brGDGT-IIa and $\ln(\text{Fe}/\text{Ti})$ suggests that anoxic conditions in the lake, due to incomplete mixing in a meromictic setting, result in lower relative brGDGT-IIa abundance. Whilst these patterns are observable at Diss Mere and Nautajärvi, we only observe a significant relationship between S_{clr} and brGDGT-IIa' at Meerfelder Maar ($r = 0.36$; $p = 0.045$) in terms of redox sensitive elements (S5). Therefore, 430 at Meerfelder Maar, like Diss Mere, anoxic conditions are considered less important than at Nautajärvi for brGDGTs, which is a pattern that has emerged across dimictic and meromictic lake settings globally (Raberg et al., 2025).

To further evaluate whether inferred redox sensitivity reflect changes in water column stratification, we compare our results with GDGT-based stratification indicators (e.g., Baxter et al. 2021). Baxter et al. (2021) use $f(\text{CREN}')$ (ratio of crenarchaeol 435 / crenarchaeol isomer) at Lake Chala to define periods of upper water column stratification, where higher values indicate greater stratification. As changes in mixing regime have been identified at Nautajärvi by Lincoln et al. (2025), we test GDGT approaches as stratification indicators here. Based on Baxter et al. (2021) increased $f(\text{CREN}')$ at Nautajärvi during the mid-Holocene suggests periods of enhanced lake water stratification (S9). Physical varve evidence suggests that Nautajärvi was sensitive during the mid-Holocene with periods of increased lake water stratification (greater meromixis) interspersed with 440 periods of periods of strengthened mixing (observed through Fe precipitation and colloid formation; Lincoln et al., 2025). Indeed, rapid fluctuations between higher and lower $f(\text{CREN}')$ in the mid-Holocene may align with these observations (S9). Whilst we acknowledge that interpretations from very different, albeit meromictic systems (in the mid-Holocene at Nautajärvi), may not be wholly transferable, the data appears to agree with physical varve evidence at this location reinforcing observations from Lincoln et al. (2025).

445

Therefore, the strength of redox influences on brGDGT distributions vary between sites. Nautajärvi appears to exhibit the greatest sensitivity to redox conditions and changes in mixing regime whereas this is weaker at Diss Mere and Meerfelder Maar. These results indicate that although anoxia and oxycline position can influence the distribution of certain 5-methyl brGDGTs, redox effects are site specific. However, the scatter observed in the correlations between brGDGTs and redox

450 sensitive elements, together with their contrasting behaviour across lakes, suggests that other controls (e.g., temperature and environmental factors) remain important drivers of brGDGT production in these systems.

4.3.2 BrGDGT response to limnological characteristics / processes within varved lakes

BrGDGT distributions, cyclisation of branched tetraethers (CBT')-derived pH reconstructions, and the isomer ratio (IR_{6ME}) differ across each site (Figs. 5, 6, 7). These differences are likely influenced by differences in water chemistry owing to the
455 global relationship between CBT' and IR_{6ME} and pH (De Jonge et al., 2014; Naafs et al., 2017) and therefore may be associated with varve formation.

At Diss Mere, calcite precipitation occurs predominantly within the summer leading to the formation of calcite laminae. This is a product of alkaline lake waters being supersaturated with $CaCO_3$ throughout the year, with a greater degree of precipitation
460 occurring during the summer months. Although our sampling strategy encompasses multiple varve laminations (i.e., both organic and calcite sub-layers in sub millimetre varve structures), brGDGT-derived pH values are relatively alkaline throughout the Holocene (~ 8.5 – 9.5). We do not observe a significant correlation between CBT' and Ca_{clr} or CBT' and $\ln(Ca/Ti)$ at Diss Mere (S4), likely due to this sampling approach. Nonetheless, our pH reconstructions are aligned to pH readings (~ 9) from epilimnetic waters (Boyall et al., 2023). At Nautajärvi the winter season is represented by clastic lamina
465 reflective of winter precipitation and snowmelt run off whereas the summer layer is related to lake productivity and allochthonous carbon yield from the catchment via surface runoff (Lincoln et al., 2025). This is consistent with more acidic conditions within the lake, as suggested by less alkaline pH reconstructions which is a product of the influx of acidic snowmelt during spring (e.g., Korkonen et al., 2017). However, we also observe a slight reduction in alkalinity from the early to late Holocene at Nautajärvi, which aligns with observations from other Scandinavian locations (Rosén and Hammarlund, 2007),
470 and may be attributed to a gradual increase in input of acidic soil or peat, as is consistent with a switch to the machine-learning-defined 'peat' source for late Holocene brGDGTs. We observe positive relationships between $\ln(Ca/Ti)$ ($r = 0.55$; $p = 0.001$), $\ln(Ca/K)$ ($r = 0.54$; $p = 0.002$) and CBT'/pH at Meerfelder Maar (S6), suggesting that variability in Ti_{clr} and K_{clr} impacts the 6-methyl isomerisation response (e.g., Peuple et al., 2022). Authigenic calcite only precipitates during a short period of the early Holocene in Meerfelder Maar (Martin-Puertas et al., 2017). The strong Ca_{clr} RDA loadings (Fig. 4) are driven by early
475 Holocene samples only and are not, therefore, considered a driver of the Holocene GDGT response.

The relative proportion of 6-methyl brGDGTs is variable between sites. This is reflected in differences in average IR_{6ME} with Diss Mere and Meerfelder Maar exhibiting greater values than Nautajärvi (Figs. 5, 6, 7). We observe a significant relationship between IR_{6ME} and $\ln(Ca/Ti)$ at both Diss Mere and Meerfelder Maar (S4, S6), which at the latter site appears predominantly
480 driven by the samples from within the early Holocene. This suggests that brGDGT isomerisation responds to annual $CaCO_3$ saturation within lake waters (Diss Mere) and changes to winter mineral flux (Meerfelder Maar) and therefore reflect site

dependent autochthonous and allochthonous processes. Given the low IR_{6ME} from Nautajärvi we do not discuss this further here.

485 Additionally, changes in organic carbon have been shown to drive brGDGT responses (Hällberg et al., 2023). Whilst no organic carbon data is available for comparison, we argue that changes in organic carbon content has a limited influence on temporal trends during the Holocene due to continued, rhythmic varve sedimentation. Such conditions suggest stable depositional and productivity environments without large shifts in organic carbon (e.g., Kril et al., 2025). Where carbon content likely increases (i.e. in the non-varved sediments < 2.0 ka BP at Diss Mere) organic carbon may influence brGDGTs. However, 490 disentangling this from processes controlling eutrophication remains difficult.

We show that differences in the distribution of brGDGTs and metrics can be related to processes that operate in seasonally stratified lakes. The data suggests that anoxia, limnological and catchment processes can influence brGDGTs in these environments but that different processes impact brGDGTs at each site. However, given the moderate correlations other factors 495 (e.g., climate) likely remain a key component (Section 4.4).

4.4 Reliability of brGDGT-derived temperature reconstructions from varved lake sediments

4.4.1 Modern brGDGT reconstructions consistent with instrumental observations

To ascertain whether brGDGT temperature reconstructions can be used to infer Holocene climate variability, we compare modern (< 300 year) brGDGT-based temperature reconstructions to instrumental data from Diss Mere and Nautajärvi (Fig. 8). 500 This same analysis is not possible from Meerfelder Maar owing to disrupted core top sediments.

BrGDGT-based mean temperature of months above freezing (MAF) reconstructions from Diss Mere are similar to long-term instrumental mean annual air temperature (MAAT) data, but are ~1°C to 2°C warmer than instrumental data between 1882 and 1950 AD but converge more closely after 1950 AD. As Diss Mere is not seasonally frozen, brGDGT-derived MAF 505 estimates *approximate* MAAT albeit, on average, appear slightly warmer in modern settings. At Nautajärvi, brGDGT temperature reconstructions overestimate instrumental MAAT by > 5°C when applying a global lacustrine MAF temperature calibration. The BIGMaC machine-learning algorithm suggests that brGDGTs from samples covering the last 300 years at this site could be derived from ‘peat’ rather than within the lake itself (see Section 4.2) and that a lacustrine calibration in these modern samples may be inappropriate. To explore this further, we recalculate temperatures at Nautajärvi using a combined 510 global peat and soil dataset (Dearing Crampton-Flood et al., 2020) calibrated to either MAAT or MAF. However, MAAT reconstructions yield temperatures that are consistently lower (3°C - 5°C) than instrumental MAAT.




Figure 8: Modern (< 300-year) temperature reconstructions from Diss Mere and Nautajärvi. Shown are a) GDGT-based MAF reconstructions at Diss Mere; b) this data alongside modern MAAT instrumental data from both the Lowestoft meteorological station (light blue) and from 5 km gridded temperatures using the HadUK dataset (light orange; Hollis et al., 2019); c) GDGT-based MAAT and MAF reconstructions at Nautajärvi; d) these GDGT-based MAF reconstructions alongside modern MAAT instrumental data from the Heinola Asemantaus (light blue) and Helsinki Kaisaniemi (light orange) meteorological stations; and e) Nautajärvi GDGT reconstructions against re-sampled May–November meteorological station air temperature data.

Instead, we resampled instrumental MAAT to MAF to only include months where the lake is sub-aerially exposed and find a stronger overlap between the brGDGT MAF temperatures and instrumental data. This suggests that brGDGTs at Nautajärvi are recording warm season temperatures and that a lake calibration is appropriate.

Whilst Nautajärvi has a very good overlap with instrumental data, Diss Mere only approximates instrumental data with occasional divergence between reconstructed and instrumental data. However, we do note that instrumental data are within reconstruction uncertainty of the GDGT calibration. Moreover, Diss Mere and Nautajärvi are different systems. Nautajärvi presently accumulates varves and is mesotrophic (Lincoln et al., 2025), whilst the upper sediments from Diss Mere are massive and the system is currently eutrophic (despite similar processes operating as during the Holocene; Boyall et al., 2023).

Eutrophication at Diss Mere is a result of increased detrital input via intensified human catchment activity over the last 1 kyr (Boyall et al., 2024). This shift in trophic state is realized through changes in diatom and blue-green algae communities (Boyall et al., 2024). We suggest that coeval shifts in the bacterial community, in a eutrophic system, may help explain why the overlap between reconstructed and instrumental data is weaker than at Nautajärvi. Shifting IR_{6ME} and $\Sigma IIIa / \Sigma IIa$ ratios also suggest bacterial community change. Similar patterns have been identified elsewhere (e.g. Miller et al., 2018; Zhao et al., 2021) associated with changing bacterial communities driven by eutrophication (Miller et al., 2018) and / or degradation and subsurface GDGT production (De Jonge et al., 2020). Given Diss Mere is presently eutrophic, this explanation seems plausible here.

This analysis shows that our data approximate modern instrumental data from these two locations yet consideration needs to be given to MAAT or MAF depending on location and present trophic conditions in the lake environment.

540 4.4.2 Effect of 6-methyl brGDGT isomers on Holocene MBT'_{5ME} values

Previous work has suggested that application of the methylation index of branched tetraethers (MBT'_{5ME}) in lakes with abundant 6-methyl brGDGTs can yield lower- or higher-than-expected temperature estimates (Dang et al., 2018; Russell et al., 2018; Bauersachs et al., 2024). Although the causes of this discrepancy remain unclear, it is considered to reflect a non-thermal response (Novak et al., 2025). There is no significant correlation between IR_{6ME} and MBT'_{5ME} at either Meerfelder Maar or Nautajärvi, respectively ($r = 0.01$; $p = 0.97$, $r = -0.36$; $p = 0.053$). At Diss Mere, there is a significant correlation between IR_{6ME} and MBT'_{5ME} ($r = 0.65$; $p < 0.001$) across the Holocene (S4), suggesting that input of 6-methyl brGDGTs may bias temperature reconstructions (Dang et al., 2018; Martinez-Sosa et al., 2020). However, this data incorporates sparsely sampled datapoints that track low-frequency climate variability (see Section 4.5). If the data is subset to remove the influence of low-frequency variability and includes only highly sampled mid-Holocene data, the IR_{6ME} and MBT'_{5ME} relationship is much weaker ($r = 0.27$; $p = 0.141$).

The observed similarities with regional reconstructions (Section 4.5) and instrumental data (Section 4.4.1), suggest that whilst there may be a minor influence, it is unlikely that the brGDGT distribution at Diss Mere is controlled by non-thermal factors associated with IR_{6ME} . Our results thus emphasize that previously identified 0.5 and 0.4 IR_{6ME} thresholds may vary on a site-by-site basis (e.g. Bauersachs et al., 2024; Novak et al., 2025).

4.5 New insights into terrestrial temperature evolution during the Holocene

Above we discussed the probable source of different GDGTs in annually laminated lake archives (Section 4.1 - 4.2), the influence of different lake processes on brGDGTs (Section 4.3) and the overall reliability of brGDGT reconstructions in modern and Holocene contexts (Section 4.4). Below we show the comparison of brGDGT reconstructions from our lakes to regional Holocene climate observations (Fig. 9). As reconstructions of MAF at Diss Mere (11-15°C) and Meerfelder Maar


(11-14°C) are both similar and warmer than at Nautajärvi (8-10°C) (Fig. 9), our brGDGT temperature reconstructions are, in the first order, performing as expected in light of site latitudinal differences. However, differences can be observed regionally.

4.5.1 Trends in temperature evolution across the Holocene

565 Diss Mere and Meerfelder Maar exhibit ~ 2°C - 3°C warming between 10 and 8 / 7 ka BP, respectively. From the available data, peak warmth occurs at Diss Mere between 8 and 4 ka BP, with the warmest temperature at 8 ka BP reconstructed from a single data point in a sparsely sampled section. Peak warmth at Meerfelder Maar appears between 5.6 and 3 ka BP with the observation at 3 ka again from a single datapoint. At Diss Mere temperatures generally follow a decreasing pattern from 8 ka BP (depending on calibration). In contrast, Nautajärvi reveals only a small temperature increase from the early to mid-Holocene from 9.8 ka BP, mid-Holocene peak warmth between 6 and 4.8 ka BP and rising temperatures across the late-Holocene (Fig. 570 9). Whilst caution is required when interpreting sparsely sampled datapoints, our reconstructions are broadly comparable with climate and physical proxy data obtained from these records (with the exception of warm temperatures during the late Holocene from Nautajärvi and Meerfelder Maar; S10, S11, S12), and comparative reconstructions from across Europe (Fig. 9) adding greater credence to our interpretations.

575 The general increase in temperature from the early Holocene as observed at Diss Mere and Meerfelder Maar, and decline in the late Holocene at Diss Mere is comparable to global temperature trends across the Holocene (e.g., Fig. 9; Kaufman et al., 2020; Salonen et al., 2024) and follows patterns of precessionally driven summer insolation (Loutre et al., 2004). However, increasing temperatures at Nautajärvi in the late Holocene appears at odds with both this pattern and Arctic neoglacial cooling (e.g., McKay et al., 2018). Orme et al. (2018) identify warming sea surface temperatures from 2 ka BP from the Greenland and Norwegian Seas and the Iceland Basin, which adds to evidence of warming from the south east Norwegian Sea, warm 580 climate anomalies in Scandinavia and a greater abundance of warm diatom taxa in the Baltic Sea (Seppä et al., 2009; Sejrup et al., 2016; van Wirdum et al., 2019). Despite this, it is challenging to reconcile opposing patterns in Europe, especially given that Nautajärvi is trending towards warmer climates than in the mid-Holocene. Therefore, at Nautajärvi ecological changes to bacterial communities, via changes in GDGT source in the mid – late Holocene, and changes in recorded seasonality of 585 temperature cannot be discounted.

Despite these proxy observations, the long-term evolution of temperature between regions (e.g., in the timing of peak warmth) reveal spatial differences with sites in mainland Europe showing different trends vs the UK. We suggest that the more maritime location of Diss Mere may be more sensitive to northern North Atlantic conditions, which, alongside certain sea surface 590 temperature records, depending on location and proxy, also reveal earlier peak warmth (e.g., Andersson et al., 2010; Berner et al., 2011; Cartapanis et al., 2022). However, why peak warmth is delayed at Meerfelder Maar, is not yet clear. At Nautajärvi mid-Holocene peak warmth is not surprising given similarities to a multi-ensemble pollen reconstruction (Salonen et al., 2024) and a brGDGT reconstruction (Otiniano et al., 2024) from northern Finland (Fig. 9). In Finland



595 **Figure 9: Comparative Holocene climate records. a) summer insolation at 65°N; b) Holocene global surface temperature anomalies (Kaufman et al., 2020); c) a pollen-based multi-method ensemble of Holocene July temperatures (Salonen et al., 2024); d) brGDGT**
600 **multiple linear regression MAF temperatures, northern Finland (Otiniano et al., 2024); e) brGDGT multiple linear regression MAAT temperatures, Massif-Central, France (Martin et al., 2020); f) brGDGT Bayesian MAF temperatures, eastern Carpathians, Romania (Ramos-Román et al., 2022); g) brGDGT Bayesian MAAT temperatures, Massif-Central, France (d'Olivera et al., 2023); and h, i, j) the Bayesian lake MAF (black; Martinez-Sosa et al., 2021) and methylation set multivariate regression MAF**
reconstructions (green; Raberg et al., 2021) from sites in this study.

delayed peak warmth may be related to the influence of the final wastage of the Fennoscandian ice sheet (Patton et al., 2017) and associated disruption of ocean circulation patterns (Cartapanis et al., 2022). Globally, at latitudes of $> 60^{\circ}\text{N}$, continental sites display peak warmth either prior to 10 ka BP or between 8 and 4 ka BP (Cartapanis et al., 2022). At 10 ka BP, Nautajärvi was a component of Lake Ancylus and connected to the Baltic Sea basin (Ojala et al., 2005) and therefore was not a competent lacustrine archive at this time. Therefore, regional temporal differences in temperature evolution appears variable between more maritime locations and mainland Europe.

Spatial differences in reconstructed temperatures also appear in the mid Holocene. Previous west-European GDGT-based reconstructions reveal a general increase in temperatures across the mid-Holocene (e.g., van den Bos et al., 2018; d'Oliveira et al., 2023; Fig. 9). Whilst reconstructions from a site in eastern Europe demonstrates cooling trends (e.g., Ramos-Román et al., 2022). Further reconstruction complexity is added with a western European site displaying warming in the mid-Holocene (Martin et al., 2020). However, this has been suggested to relate to differences in reconstructed temperature seasonality alongside localised site differences, for example in environmental contexts and altitude (d'Oliveira et al., 2023). Nonetheless, the trends in temperature reconstructions from both Diss Mere and Meerfelder Maar are comparable to European Holocene brGDGT-reconstructions (Fig. 9), albeit with local and/or spatial influences (e.g., site elevation, continental position) seemingly being important in the direction of temperature change during the mid-Holocene.

4.5.2 mid-Holocene variance

By exploiting varved lake sediments, we show that during the mid-Holocene, Diss Mere and Meerfelder Maar median reconstructions exhibit $\sim 2^{\circ}\text{C} - 3^{\circ}\text{C}$ of temperature variability at multi-decadal scales. Whilst this $\sim 2^{\circ}\text{C} - 3^{\circ}\text{C}$ temperature variability may be a product of calibration uncertainty, comparative climate reconstructions reveal similar patterns. Pollen and brGDGT-based reconstructions from the European mid-latitudes reveal a similar range of variability (e.g., Seppä et al., 2009; Martin et al., 2020); albeit with some records displaying increased variability ($\sim 4^{\circ}\text{C} - 5^{\circ}\text{C}$) at multi-decadal (Ramos-Román et al., 2022) and centennial scales (d'Oliveira et al., 2023) (Fig. 9). At Nautajärvi, temperatures are stable during the mid-Holocene (Fig. 9; median variability of 1°C). This is comparable to the brGDGT reconstruction from northern Finland between 7 and 4 ka BP (Otiniano et al., 2024). However, this is incongruent with non-brGDGT based temperature observations from Scandinavia (e.g., Shala et al., 2017; Salonen et al., 2024) where greater temperature variability ($2^{\circ}\text{C} - 3^{\circ}\text{C}$) is observed. Our reconstructions reveal a mostly comparable range of variability to other brGDGT reconstructions and suggest that higher latitudinal locations exhibit less variance than lower latitude locations. However, as reconstruction amplitude can be different between GDGT and traditional proxy reconstructions (pollen and/or chironomid) it is likely that some of the controls discussed above may be important in these lake settings (Section 4.3), or the seasonality of temperature recorded varies by proxy and by latitudinal setting.

4.5 Conclusion

635 We present iso- and brGDGT reconstructions from across the Holocene from three annually laminated records from Europe. We show that isoGDGTs are strongly influenced by methanogenesis linked to hypolimnetic hypoxia and that TEX₈₆ derived LST reconstructions are unreliable in annually laminated lake systems (or lakes that exhibit hypoxia/anoxia). In contrast, we show that brGDGTs are likely synthesised *in-situ* within each lake. Whilst there appears to be a redox influence at each location, the degree to which oxygen depletion affects brGDGT-temperatures appears minimal. This is similar to limnological
640 processes that influence lamination style and limnological characteristics (e.g., pH) in each lake.

We show that there is a strong correspondence between brGDGT temperatures and modern instrumental data, demonstrating suitability for brGDGT temperature reconstruction at those locations. Further, the reconstructed temperatures from Holocene-aged sediments at Diss Mere and Meerfelder Maar resemble regional climate reconstructions (partly at Meerfelder Maar) and
645 exhibit similar multi-decadal temperature variability. Nautajärvi exhibits limited temperature variability - this is comparable to another brGDGT reconstruction from Finland. However, there is mismatch between brGDGT and regional reconstructions during the late Holocene at this site (3 ka BP to present), suggesting other confounding factors may influence brGDGTs during the late Holocene at Nautajärvi.

650 Taken together, we show that annually laminated lake sequences are valuable archives for brGDGT-based climate reconstruction. However, we also show that processes that are specific to these unique laminated archives can lead to a brGDGT response. We therefore suggest that GDGT-based reconstructions be compared with additional site data or regional climate observations to disentangle the influence of lake processes. In doing so, GDGT-based reconstructions from annually laminated records have the potential to deconvolve both multi-decadal climate variability and if performed at higher temporal
655 resolutions, seasonal climate dynamics.

Data availability

All data associated with this article is available at Zenodo: <https://doi.org/10.5281/zenodo.17610783>

Author contributions

AMA, GNI and PGL conceptualised the study. AMA, GNI, and PGL formalised the methodology. AMA performed the
660 analysis with input from GNI, MRB, DF, IB and HW. AB, PL, AEKO and CMP were involved in data curation. AMA wrote the original manuscript and designed the figures with input from GNI and PGL. All authors contributed to the review and editing process.

Conflict of interest

The authors declare that there are no competing interests in this research.

665 Disclaimer

The authors declare that this work has not been published nor is under consideration anywhere else.

Acknowledgements

This study was funded by the UKRI Medical Research Council through a Future Leaders Fellowship held by C.M.-P.: DECADAL: Rethinking Palaeoclimatology for Society (MR/W009641/1) and further supported by a UKRI Natural
670 Environment Research Council National Environmental Isotope Facility (NEIF) grant (NEIF_2602.0423) to G.N.I, A.M.A and P.G.L. The authors wish to thank the UKRI Natural Environment Research Council (NERC) for partial funding of the National Environmental Isotope Facility (NEIF; contract no. NE/V003917/1). G.N.I. is supported by a GCRF Royal Society Dorothy Hodgkin Fellowship (DHF\R1\191178) with additional support via the Royal Society (RF\ERE\231019, RF\ERE\210068). The authors thank the Maarmuseum in Manderscheid for logistical support during coring of Meerfelder
675 Maar, with work at Nautajärvi conducted with the collaboration of Digital Waters Flagship (DIWA) (decision no. 359247) funded by the Research Council of Finland. The authors would like to thank Dr. Saija Saarni and Emilia Kosonen, who assisted with core extraction at Nautajärvi, Dr Sargent Bray and Dr Mark Peaple for discussions around analytical chemistry and GDGT-techniques and the DECADAL project team for discussions around the manuscript.

Financial support

680 This work was supported by the following funding sources:
UKRI Medical Research Council through a Future Leaders Fellowship (MR/W009641/1)
UKRI Natural Environment Research Council National Environmental Isotope Facility (NEIF) grant (NEIF_2602.0423)
UKRI Natural Environment Research Council partial funding of the National Environmental Isotope Facility (NEIF; NE/V003917/1).
685 GCRF Royal Society Dorothy Hodgkin Fellowship (DHF\R1\191178)
The Royal Society (RF\ERE\231019, RF\ERE\210068)
Digital Waters Flagship (DIWA) (no. 359247)

Short summary

We present lipid-based environmental and temperature reconstructions spanning the Holocene from three European varved
690 lakes. At each site archaea are impacted by methane which influence lake water temperature reconstructions, and bacteria appear to respond to oxygen conditions and specific lake and environmental processes. Nonetheless, bacterial temperature

reconstructions are comparable to regional climate data, showing the potential of varve archives for high resolution reconstructions.

References

- 695 Abrook, A. M., Langdon, P. G., Inglis, G. N., Brauer, A., Lincoln, P., Mayfield, R., Ojala, A. E. K., and Martin-Puertas, C.:
The role of summer temperature on aquatic insect diversity at multi-decadal scales within the Holocene, *Glob. Change Biol.*,
31, e70366, <https://doi.org/10.1111/gcb.70366>, 2025.
- Andersson, C., Pausata, F. S., Jansen, E., Risebrobakken, B., and Telford, R. J.: Holocene trends in the foraminifer record
700 from the Norwegian Sea and the North Atlantic Ocean, *Clim. Past*, 6, 179–193, <https://doi.org/10.5194/cp-6-179-2010>, 2010.
- Auguet, J. C., Barberan, A., and Casamayor, E. O.: Global ecological patterns in uncultured Archaea, *ISME J.*, 4, 182–190,
<https://doi.org/10.1038/ismej.2009.109> , 2009.
- 705 Bailey, I.: A high-resolution record of mid-Holocene climate change from Diss Mere, UK, Ph.D. thesis, University College
London, 2005.
- Bauersachs, T., Schubert, C. J., Mayr, C., Gilli, A., and Schwark, L.: Branched GDGT-based temperature calibrations from
Central European lakes, *Sci. Total Environ.*, 906, 167724, <https://doi.org/10.1016/j.scitotenv.2023.167724>, 2024.
- 710 Baxter, A. J., Peterse, F., Verschuren, D., Maitituerdi, A., Waldmann, N., and Sinninghe Damsté, J. S.: Disentangling
influences of climate variability and lake-system evolution on climate proxies derived from isoprenoid and branched
glycerol dialkyl glycerol tetraethers (GDGTs): the 250 kyr Lake Chala record, *Biogeosciences*, 21, 2877–2908,
<https://doi.org/10.5194/bg-21-2877-2024>, 2024.
- 715 Baxter, A. J., Van Bree, L. G. J., Peterse, F., Hopmans, E. C., Villanueva, L., Verschuren, D., and Sinninghe Damsté, J. S.:
Seasonal and multi-annual variation in the abundance of isoprenoid GDGT membrane lipids and their producers in the water
column of a meromictic equatorial crater lake (Lake Chala, East Africa), *Quat. Sci. Rev.*, 273, 107263,
<https://doi.org/10.1016/j.quascirev.2021.107263>, 2021.
- 720 Berner, K. S., Koç, N., Godtliebsen, F., and Divine, D.: Holocene climate variability of the Norwegian Atlantic Current
during high and low solar insolation forcing, *Paleoceanogr. Paleoclimatol.*, 26, PA2220,
<https://doi.org/10.1029/2010PA002002>, 2011.

- 725 Blaga, C. I., Reichart, G. J., Heiri, O., and Sinninghe Damsté, J. S.: Tetraether membrane lipid distributions in water-column
particulate matter and sediments: a study of 47 European lakes along a north–south transect, *J. Paleolimnol.*, 41, 523–540,
<https://doi.org/10.1007/s10933-008-9242-2>, 2009.
- Boyall, L., Martin-Puertas, C., Tjallingii, R., Milner, A. M., and Blockley, S. P.: Holocene climate evolution and human
730 activity as recorded by the sediment record of Lake Diss Mere, England, *J. Quat. Sci.*, 39, 972–986,
<https://doi.org/10.1002/jqs.3646>, 2024.
- Boyall, L., Valcárcel, J. I., Harding, P., Hernández, A., and Martin-Puertas, C.: Disentangling the environmental signals
recorded in Holocene calcite varves based on modern lake observations and annual sedimentary processes in Diss Mere,
735 England, *J. Paleolimnol.*, 70, 39–56, <https://doi.org/10.1007/s10933-023-00282-z>, 2023.
- Brauer, A., Endres, C., Zolitschka, B., and Negendank, J. F.: AMS radiocarbon and varve chronology from the annually
laminated sediment record of Lake Meerfelder Maar, Germany, *Radiocarbon*, 42, 355–368,
<https://doi.org/10.1017/S0033822200030307>, 2000.
740
- Briffa, K. R., Osborn, T. J., and Schweingruber, F. H.: Large-scale temperature inferences from tree rings: a review, *Glob.
Planet. Change*, 40, 11–26, [https://doi.org/10.1016/S0921-8181\(03\)00095-X](https://doi.org/10.1016/S0921-8181(03)00095-X), 2004.
- Buckles, L. K., Weijers, J. W., Verschuren, D., and Sinninghe Damsté, J. S.: Sources of core and intact branched tetraether
745 membrane lipids in the lacustrine environment: Anatomy of Lake Challa and its catchment, equatorial East Africa, *Geochim.
Cosmochim. Acta*, 140, 106–126, <https://doi.org/10.1016/j.gca.2014.04.042>, 2014.
- Candy, I., Boyall, L., Lincoln, P., Martin-Puertas, C., Matthews, I., Holt-Wilson, T., and Valcarcel, J.: A cold but stable 4.2
ka event in Britain and the northeastern Atlantic region, *Quat. Sci. Rev.*, 349, 109093,
750 <https://doi.org/10.1016/j.quascirev.2024.109093>, 2025.
- Cartapanis, O., Jonkers, L., Moffa-Sanchez, P., Jaccard, S. L., and de Vernal, A.: Complex spatio-temporal structure of the
Holocene Thermal Maximum, *Nat. Commun.*, 13, 5662, <https://doi.org/10.1038/s41467-022-33362-1>, 2022.
- 755 Collins, E. R., Ferland, T. M., Castañeda, I. S., Owen, R. B., Lowenstein, T. K., Cohen, A. S., Renaut, R. W., O’Beirne, M.
D., and Werne, J. P.: Hot-spring inputs and climate drive dynamic shifts in archaeal communities in Lake Magadi, Kenya
Rift Valley, *Biogeosciences*, 22, 3931–3948, <https://doi.org/10.5194/bg-22-3931-2025>, 2025.

- Crampton-Flood, E. D., Tierney, J. E., Peterse, F., Kirkels, F. M., and Sinninghe Damsté, J. S.: BayMBT: A Bayesian calibration model for branched glycerol dialkyl glycerol tetraethers in soils and peats, *Geochim. Cosmochim. Acta*, 268, 142–159, <https://doi.org/10.1016/j.gca.2019.09.043>, 2020.
- Dang, X., Ding, W., Yang, H., Pancost, R. D., Naafs, B. D. A., Xue, J., Lin, X., Lu, J., and Xie, S.: Different temperature dependence of the bacterial brGDGT isomers in 35 Chinese lake sediments compared to that in soils, *Org. Geochem.*, 119, 72–79, <https://doi.org/10.1016/j.orggeochem.2018.02.008>, 2018.
- Davison, W.: Iron and manganese in lakes, *Earth-Sci. Rev.*, 34, 119–163, [https://doi.org/10.1016/0012-8252\(93\)90029-7](https://doi.org/10.1016/0012-8252(93)90029-7), 1993.
- De Jonge, C., Fiskal, A., Han, X., and Lever, M.: Biomarker (brGDGT) degradation and production in lacustrine surface sediments: Implications for paleoclimate reconstructions, *EGU General Assembly 2020*, Online, EGU2020-12592, <https://doi.org/10.5194/egusphere-egu2020-12592>, 2020.
- De Jonge, C., Hopmans, E. C., Zell, C. I., Kim, J.-H., Schouten, S., and Sinninghe Damsté, J. S.: Occurrence and abundance of 6-methyl branched glycerol dialkyl glycerol tetraethers in soils: Implications for palaeoclimate reconstruction, *Geochim. Cosmochim. Acta*, 141, 97–112, <https://doi.org/10.1016/j.gca.2014.06.013>, 2014.
- De Jonge, C., Stadnitskaia, A., Hopmans, E. C., Cherkashov, G., Fedotov, A., Streletskaia, I. D., Vasiliev, A. A., and Sinninghe Damsté, J. S.: Drastic changes in the distribution of branched tetraether lipids in suspended matter and sediments from the Yenisei River and Kara Sea (Siberia): Implications for the use of brGDGT-based proxies in coastal marine sediments, *Geochim. Cosmochim. Acta*, 165, 200–225, <https://doi.org/10.1016/j.gca.2015.05.044>, 2015.
- d'Oliveira, L., Dugerdil, L., Ménot, G., Evin, A., Muller, S. D., Ansanay-Alex, S., Azuara, J., Bonnet, C., Bremond, L., Shah, M., and Peyron, O.: Reconstructing 15 kyr of southern France temperatures from coupled pollen and molecular (branched glycerol dialkyl glycerol tetraether) markers (Canroute, Massif Central), *Clim. Past*, 19, 2127–2156, <https://doi.org/10.5194/cp-19-2127-2023>, 2023.
- Hällberg, P.L., Schenk, F., Jarne-Bueno, G., Schankat, Y., Zhang, Q., Rifai, H., Phua, M., and Smittenberg, R.H.: Branched GDGT source shift identification allows improved reconstruction of an 8,000-year warming trend on Sumatra, *Org. Geochem.*, 186, p.104702, <https://doi.org/10.1016/j.orggeochem.2023.104702>, 2023.

- Hamilton, N. E. and Ferry, M.: ggtern: Ternary diagrams using ggplot2, *J. Stat. Softw.*, 87, 1–17, <http://doi.org/10.18637/jss.v087.c03>, 2018.
- Hernández, A., Martin-Puertas, C., Moffa-Sánchez, P., Moreno-Chamarro, E., Ortega, P., Blockley, S., Cobb, K. M., Comas-Bru, L., Giralt, S., Goosse, H., Luterbacher, J., Martrat, B., Muscheler, R., Parnell, A., Pla-Rabes, S., Sjolte, J., Scaife, A., Swingedouw, D., Wise, E., and Xu, G.: Modes of climate variability: synthesis and review of proxy-based reconstructions through the Holocene, *Earth-Sci. Rev.*, 209, 103286, <https://doi.org/10.1016/j.earscirev.2020.103286>, 2020.
- Hopmans, E. C., Schouten, S., and Sinninghe Damsté, J. S.: The effect of improved chromatography on GDGT-based palaeoproxies, *Org. Geochem.*, 93, 1–6, <https://doi.org/10.1016/j.orggeochem.2015.12.006>, 2016.
- Inglis, G. N., Farnsworth, A., Lunt, D., Foster, G. L., Hollis, C. J., Pagani, M., Jardine, P. E., Pearson, P. N., Markwick, P., Galsworthy, A. M., Raynham, L., Taylor, K. W. R., and Pancost, R. D.: Descent toward the Icehouse: Eocene sea surface cooling inferred from GDGT distributions, *Paleoceanography*, 30, 1000–1020, <https://doi.org/10.1002/2014PA002723>, 2015.
- Kassambara, A. and Mundt, F.: factoextra: Extract and visualize the results of multivariate data analyses, R package version 1.0.7, available at: <https://CRAN.R-project.org/package=factoextra>, 2020.
- Kaufman, D., McKay, N., Routson, C., Erb, M., Dätwyler, C., Sommer, P. S., Heiri, O., and Davis, B.: Holocene global mean surface temperature: a multi-method reconstruction approach, *Sci. Data*, 7, 201, <https://doi.org/10.1038/s41597-020-0530-7>, 2020.
- Kim, B. and Zhang, Y. G.: Methane Index: towards a quantitative archaeal lipid biomarker proxy for reconstructing marine sedimentary methane fluxes, *Geochim. Cosmochim. Acta*, 354, 74–87, <https://doi.org/10.1016/j.gca.2023.06.008>, 2023.
- Korkonen, S. T., Ojala, A. E. K., Kosonen, E., and Weckström, J.: Seasonality of chrysophyte cyst and diatom assemblages in varved Lake Nautajärvi – implications for palaeolimnological studies, *J. Limnol.*, 76, 366–379, <https://doi.org/10.4081/jlimnol.2017.1473>, 2017.
- Kril, M., Bonk, A., Żarczyński, M., Zolitschka, B., and Tylmann, W.: Varved sediments of Lake Gorzyńskie, western Poland: a new archive of climatic and environmental changes during the Late Glacial and the Holocene in central Europe, *JOPL*, 73, 4, pp.329-346, <https://doi.org/10.1007/s10933-025-00362-2>, 2025.

- Larocque-Tobler, I., Filipiak, J., Tylmann, W., Bonk, A., and Grosjean, M.: Comparison between chironomid-inferred mean-
825 August temperature from varved Lake Żabińskie (Poland) and instrumental data since 1896 AD, *Quat. Sci. Rev.*, 111, 35–50,
<https://doi.org/10.1016/j.quascirev.2015.01.001>, 2015.
- Lincoln, P., Tjallingii, R., Kosonen, E., Ojala, A. E. K., Abrook, A. M., and Martin-Puertas, C.: Disruption of boreal lake
circulation in response to mid-Holocene warmth: evidence from the varved sediments of Lake Nautajärvi, southern Finland,
830 *Sci. Total Environ.*, 964, 178519, <https://doi.org/10.1016/j.scitotenv.2025.178519>, 2025.
- Liu, Z., Cheng, J., Zheng, Y., Zhang, W., Liu, H., Wu, H., Zhu, J., and Xie, S.: The seasonal temperature conundrum for the
Holocene, *Sci. Adv.*, 11, e8950, [DOI: 10.1126/sciadv.adt8950](https://doi.org/10.1126/sciadv.adt8950), 2025.
- 835 Liu, Z., Zhu, J., Rosenthal, Y., Zhang, X., Otto-Bliesner, B. L., Timmermann, A., Smith, R. S., Lohmann, G., Zheng, W.,
and Elison Timm, O.: The Holocene temperature conundrum, *Proc. Natl. Acad. Sci. USA*, 111, E3501–E3505,
<https://doi.org/10.1073/pnas.1407229111>, 2014.
- Loomis, S. E., Russell, J. M., and Sinninghe Damsté, J. S.: Distributions of branched GDGTs in soils and lake sediments
840 from western Uganda: implications for a lacustrine paleothermometer, *Org. Geochem.*, 42, 739–751,
<https://doi.org/10.1016/j.orggeochem.2011.06.004>, 2011.
- Loutre, M. F., Paillard, D., Vimeux, F., and Cortijo, E.: Does mean annual insolation have the potential to change the
climate?, *Earth Planet. Sci. Lett.*, 221, 1–14, [https://doi.org/10.1016/S0012-821X\(04\)00108-6](https://doi.org/10.1016/S0012-821X(04)00108-6), 2004.
845
- Luo, Y.: Geochemical cycle and environmental effects of sulfur in lakes, *IOP Conf. Ser. Mater. Sci. Eng.*, 394, 052039,
<https://doi.org/10.1088/1757-899X/394/5/052039>, 2018.
- Luoto, T. P. and Ojala, A. E. K.: Controls of climate, catchment erosion and biological production on long-term community
850 and functional changes of chironomids in High Arctic lakes (Antczak-Orlewska Bard), *Palaeogeogr. Palaeoclimatol.*
Palaeoecol., 505, 63–72, <https://doi.org/10.1016/j.palaeo.2018.05.026>, 2018.
- Marshall, M., Schlolaut, G., Nakagawa, T., Lamb, H., Brauer, A., Staff, R., Ramsey, C. B., Tarasov, P., Gotanda, K.,
Haraguchi, T., Yokoyama, Y., Yonenobu, H., Tada, R., and Suigetsu 2006 Project Members: A novel approach to varve
855 counting using μ XRF and X-radiography in combination with thin-section microscopy, applied to the Late Glacial
chronology from Lake Suigetsu, Japan, *Quat. Geochronol.*, 13, 70–80, <https://doi.org/10.1016/j.quageo.2012.06.002>, 2012.

- 860 Martin, C., Ménot, G., Thouveny, N., Davtian, N., Andrieu-Ponel, V., Reille, M., and Bard, E.: Impact of human activities and vegetation changes on the tetraether sources in Lake St Front (Massif Central, France), *Org. Geochem.*, 135, 38–52, <https://doi.org/10.1016/j.orggeochem.2019.06.005>, 2019.
- 865 Martin, C., Ménot, G., Thouveny, N., Peyron, O., Andrieu-Ponel, V., Montade, V., Davtian, N., Reille, M., and Bard, E.: Early Holocene thermal maximum recorded by branched tetraethers and pollen in Western Europe (Massif Central, France), *Quat. Sci. Rev.*, 228, 106109, <https://doi.org/10.1016/j.quascirev.2019.106109>, 2020.
- Martínez-Sosa, P., Tierney, J. E., Pérez-Angel, L. C., Stefanescu, I. C., Guo, J., Kirkels, F., Sepúlveda, J., Peterse, F., Shuman, B. N., and Reyes, A. V.: Development and application of the branched and isoprenoid GDGT machine learning classification algorithm (BIGMaC) for paleoenvironmental reconstruction, *Paleoceanogr. Paleoclimatol.*, 38, e2023PA004611, <https://doi.org/10.1029/2023PA004611>, 2023.
- 870 Martínez-Sosa, P., Tierney, J. E., Stefanescu, I. C., Crampton-Flood, E. D., Shuman, B. N., and Routson, C.: A global Bayesian temperature calibration for lacustrine brGDGTs, *Geochim. Cosmochim. Acta*, 305, 87–105, <https://doi.org/10.1016/j.gca.2021.04.038>, 2021.
- 875 Martin-Puertas, C., Brauer, A., Dulski, P., and Brademann, B.: Testing climate–proxy stationarity throughout the Holocene: an example from the varved sediments of Lake Meerfelder Maar (Germany), *Quat. Sci. Rev.*, 58, 56–65, <https://doi.org/10.1016/j.quascirev.2012.10.023>, 2012.
- 880 Martin-Puertas, C., Tjallingii, R., Bloemsma, M., and Brauer, A.: Varved sediment responses to early Holocene climate and environmental changes in Lake Meerfelder Maar (Germany) obtained from multivariate analyses of micro X-ray fluorescence core scanning data, *J. Quat. Sci.*, 32, 427–436, <https://doi.org/10.1002/jqs.2935>, 2017.
- 885 Martin-Puertas, C., Walsh, A. A., Blockley, S. P., Harding, P., Biddulph, G. E., Palmer, A., Ramisch, A., and Brauer, A.: The first Holocene varve chronology for the UK: based on the integration of varve counting, radiocarbon dating and tephrostratigraphy from Diss Mere (UK), *Quat. Geochronol.*, 61, 101134, <https://doi.org/10.1016/j.quageo.2020.101134>, 2021.
- 890 McKay, N. P., Kaufman, D. S., Routson, C. C., Erb, M. P., and Zander, P. D.: The onset and rate of Holocene Neoglacial cooling in the Arctic, *Geophys. Res. Lett.*, 45, 12487–12496, <https://doi.org/10.1029/2018GL079773>, 2018.

- Miller, D. R., Habicht, M. H., Keisling, B. A., Castañeda, I. S., and Bradley, R. S.: A 900-year New England temperature reconstruction from in situ seasonally produced branched glycerol dialkyl glycerol tetraethers (brGDGTs), *Clim. Past*, 14, 1653–1667, <https://doi.org/10.5194/cp-14-1653-2018>, 2018.
- 895 Novak, J. B., Russell, J. M., Lindemuth, E. R., Prokopenko, A. A., Pérez-Angel, L., Zhao, B., Swann, G. E., and Polissar, P. J.: The branched GDGT isomer ratio refines lacustrine paleotemperature estimates, *Geochem. Geophys. Geosyst.*, 26, e2024GC012069, <https://doi.org/10.1029/2024GC012069>, 2025.
- Nürnberg, G. K.: Lake responses to long-term hypolimnetic withdrawal treatments, *Lake Reserv. Manag.*, 23, 388–409, <https://doi.org/10.1080/07438140709354026>, 2007.
- 900 <https://doi.org/10.1080/07438140709354026>, 2007.
- O’Beirne, M. D., Scott, W. P., Contreras, S., Araneda, A., Tejos, E., Moscoso, J., and Werne, J. P.: Distribution of branched glycerol dialkyl glycerol tetraether (brGDGT) lipids from soils and sediments from the same watershed are distinct regionally (central Chile) but not globally, *Front. Earth Sci.*, 12, 1383146, <https://doi.org/10.3389/feart.2024.1383146>, 2024.
- 905 Ojala, A. E., Alenius, T., Seppä, H., and Giesecke, T.: Integrated varve and pollen-based temperature reconstruction from Finland: evidence for Holocene seasonal temperature patterns at high latitudes, *Holocene*, 18, 529–538, <https://doi.org/10.1177/0959683608089207>, 2008.
- 910 Ojala, A. E. K. and Alenius, T.: 10 000 years of interannual sedimentation recorded in the Lake Nautajärvi (Finland) clastic–organic varves, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 219, 285–302, <https://doi.org/10.1016/j.palaeo.2005.01.002>, 2005.
- Ojala, A. E. K. and Saarinen, T.: Palaeosecular variation of the Earth's magnetic field during the last 10 000 years based on the annually laminated sediment of Lake Nautajärvi, central Finland, *Holocene*, 12, 391–400, <https://doi.org/10.1191/0959683602hl551rp>, 2002.
- 915 <https://doi.org/10.1191/0959683602hl551rp>, 2002.
- Ojala, A. E. K., Kosonen, E., Weckström, J., Korkonen, S., and Korhola, A.: Seasonal formation of clastic–biogenic varves: the potential for palaeoenvironmental interpretations, *GFF*, 135, 237–247, <https://doi.org/10.1080/11035897.2013.801925>, 2013.
- 920 Oksanen, J., Simpson, G., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O’Hara, R., Solymos, P., Stevens, M., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Carvalho, G., Chirico, M., De Caceres, M., Durand, S., Evangelista, H., FitzJohn, R., Friendly, M., Furneaux, B., Hannigan, G., Hill, M., Lahti, L., McGlinn, D., Ouellette, M.,

- Ribeiro Cunha, E., Smith, T., Stier, A., ter Braak, C., and Weedon, J.: Vegan: community ecology package, R package
925 version 2.6–6.1, available at: <https://CRAN.R-project.org/package=vegan>, 2024.
- Orme, L. C., Miettinen, A., Divine, D., Husum, K., Pearce, C., Van Nieuwenhove, N., Born, A., Mohan, R., and
Seidenkrantz, M. S.: Subpolar North Atlantic sea surface temperature since 6 ka BP: Indications of anomalous ocean–
atmosphere interactions at 4–2 ka BP, *Quat. Sci. Rev.*, 194, 128–142, <https://doi.org/10.1016/j.quascirev.2018.07.007>, 2018.
930
- Osman, M. B., Tierney, J. E., Zhu, J., Tardif, R., Hakim, G. J., King, J., and Poulsen, C. J.: Globally resolved surface
temperatures since the Last Glacial Maximum, *Nature*, 599, 239–244, <https://doi.org/10.1038/s41586-021-03984-4>, 2021.
- Otiniano, G. A., Porter, T. J., Phillips, M. A., Juutinen, S., Weckström, J. B., and Heikkilä, M. P.: Reconstructing warm-
935 season temperatures using brGDGTs and assessing biases in Holocene temperature records in northern Fennoscandia, *Quat.*
Sci. Rev., 329, 108555, <https://doi.org/10.1016/j.quascirev.2024.108555>, 2024.
- Parker, S. E. and Harrison, S. P.: The timing, duration and magnitude of the 8.2 ka event in global speleothem records, *Sci.*
Rep., 12, 10542, <https://doi.org/10.1038/s41598-022-14684-y>, 2022.
940
- Patton, H., Hubbard, A., Andreassen, K., Auriac, A., Whitehouse, P. L., Stroeven, A. P., Shackleton, C., Winsborrow, M.,
Heyman, J., and Hall, A. M.: Deglaciation of the Eurasian ice sheet complex, *Quat. Sci. Rev.*, 169, 148–172,
<https://doi.org/10.1016/j.quascirev.2017.05.019>, 2017.
- 945 Peaple, M. D., Bhattacharya, T., Lowenstein, T. K., McGee, D., Olson, K. J., Stroup, J. S., Tierney, J. E., and Feakins, S. J.:
Biomarker and pollen evidence for late Pleistocene pluvials in the Mojave Desert, *Paleoceanogr. Paleoclimatol.*, 37,
e2022PA004471, <https://doi.org/10.1029/2022PA004471>, 2022.
- Pearson, E. J., Juggins, S., Talbot, H. M., Weckström, J., Rosén, P., Ryves, D. B., Roberts, S. J., and Schmidt, R.: A
950 lacustrine GDGT–temperature calibration from the Scandinavian Arctic to Antarctic: Renewed potential for the application
of GDGT–paleothermometry in lakes, *Geochim. Cosmochim. Acta*, 75, 6225–6238,
<https://doi.org/10.1016/j.gca.2011.07.042>, 2011.
- Pearson, E. J., Juggins, S., Allbrook, H., Foster, L. C., Hodgson, D. A., Naafs, D. A., Phillips, T., and Roberts, S. J.:
955 Development of new global lake brGDGT-temperature calibrations: advances, applications, challenges, and
recommendations, *Quat. Sci. Rev.*, 369, 109615, <https://doi.org/10.1016/j.quascirev.2025.109615>, 2025.

- Peterse, F., Vonk, J. E., Holmes, R. M., Giosan, L., Zimov, N., and Eglinton, T. I.: Branched glycerol dialkyl glycerol tetraethers in Arctic lake sediments: Sources and implications for paleothermometry at high latitudes, *J. Geophys. Res. Biogeosci.*, 119, 1738–1754, <https://doi.org/10.1002/2014JG002639>, 2014.
- Powers, L., Werne, J. P., Vanderwoude, A. J., Sinninghe Damsté, J. S., Hopmans, E. C., and Schouten, S.: Applicability and calibration of the TEX₈₆ paleothermometer in lakes, *Org. Geochem.*, 41, 404–413, <https://doi.org/10.1016/j.orggeochem.2009.11.009>, 2010.
- Raberg, J. H., de Wet, G. A., Geirsdóttir, Á., Sepúlveda, J., and Miller, G. H.: Oxygen depletion in lake waters may skew brGDGT-inferred temperatures by more than 10 °C, *Geophys. Res. Lett.*, 52, e2024GL113562, <https://doi.org/10.1029/2024GL113562>, 2025.
- Raberg, J. H., Flores, E., Crump, S. E., de Wet, G., Dildar, N., Miller, G. H., Geirsdóttir, Á., and Sepúlveda, J.: Intact polar brGDGTs in Arctic lake catchments: Implications for lipid sources and paleoclimate applications, *J. Geophys. Res. Biogeosci.*, 127, e2022JG006969, <https://doi.org/10.1029/2022JG006969>, 2022.
- Raberg, J. H., Harning, D. J., Crump, S. E., de Wet, G., Blumm, A., Kopf, S., Geirsdóttir, Á., Miller, G. H., and Sepúlveda, J.: Revised fractional abundances and warm-season temperatures substantially improve brGDGT calibrations in lake sediments, *Biogeosciences*, 18, 3579–3603, <https://doi.org/10.5194/bg-18-3579-2021>, 2021.
- Rach, O., Brauer, A., Wilkes, H., and Sachse, D.: Delayed hydrological response to Greenland cooling at the onset of the Younger Dryas in western Europe, *Nat. Geosci.*, 7, 109–112, <https://doi.org/10.1038/ngeo2053>, 2014.
- Ramos-Roman, M. J., De Jonge, C., Magyari, E., Veres, D., Ilvonen, L., Develle, A. L., and Seppä, H.: Lipid biomarker (brGDGT)- and pollen-based reconstruction of temperature change during the Middle to Late Holocene transition in the Carpathians, *Glob. Planet. Change*, 215, 103859, <https://doi.org/10.1016/j.gloplacha.2022.103859>, 2022.
- Rasmussen, S. O., Bigler, M., Blockley, S. P., Blunier, T., Buchardt, S. L., Clausen, H. B., Cvijanovic, I., Dahl-Jensen, D., Johnsen, S. J., Fischer, H., Gkinis, V., Guillevic, M., Hoek, W. Z., Lowe, J. J., Pedro, J. B., Popp, T., Seierstad, I. K., Steffensen, J. P., Svensson, A. M., Vallenga, P., Vinther, B. M., Walker, M. J. C., Wheatley, J. J., and Winstrup, M.: A stratigraphic framework for abrupt climatic changes during the Last Glacial period based on three synchronized Greenland ice-core records: refining and extending the INTIMATE event stratigraphy, *Quat. Sci. Rev.*, 106, 14–28, <https://doi.org/10.1016/j.quascirev.2014.09.007>, 2014.

- Robles, M., Peyron, O., Ménot, G., Brugiapaglia, E., Wulf, S., Appelt, O., Blache, M., Vannière, B., Dugerdil, L., Paura, B., Ansanay-Alex, S., Cromartie, A., Charlet, L., Guédron, S., de Beaulieu, J.-L., and Joanin, S.: Climate changes during the Late Glacial in southern Europe: new insights based on pollen and brGDGTs of Lake Matese in Italy, *Clim. Past*, 19, 493–515, <https://doi.org/10.5194/cp-19-493-2023>, 2023.
- Roland, T. P., Daley, T. J., Caseldine, C. J., Charman, D. J., Turney, C. S. M., Amesbury, M. J., Thompson, G. J., and Woodley, E. J.: The 5.2 ka climate event: Evidence from stable isotope and multi-proxy palaeoecological peatland records in Ireland, *Quat. Sci. Rev.*, 124, 209–223, <https://doi.org/10.1016/j.quascirev.2015.07.026>, 2015.
- Rosén, P., and Hammarlund, D.: Effects of climate, fire and vegetation development on Holocene changes in total organic carbon concentration in three boreal forest lakes in northern Sweden, *Biogeosciences*, 4, 975–984, <https://doi.org/10.5194/bg-4-975-2007>, 2007.
- Russell, J. M., Hopmans, E. C., Loomis, S. E., Liang, J., and Sinninghe Damsté, J. S.: Distributions of 5- and 6-methyl branched glycerol dialkyl glycerol tetraethers (brGDGTs) in East African lake sediment: Effects of temperature, pH, and new lacustrine paleotemperature calibrations, *Org. Geochem.*, 117, 56–69, <https://doi.org/10.1016/j.orggeochem.2017.12.003>, 2018.
- Salonen, J. S., Kuosmanen, N., Alsos, I. G., Heintzman, P. D., Rijal, D. P., Schenk, F., Bogren, F., Luoto, M., Philip, A., Piilo, S., Trasune, L., Välranta, M., and Helmens, K. F.: Uncovering Holocene climate fluctuations and ancient conifer populations: insights from a high-resolution multi-proxy record from northern Finland, *Glob. Planet. Change*, 237, 104462, <https://doi.org/10.1016/j.gloplacha.2024.104462>, 2024.
- Schouten, S., Hopmans, E. C., and Sinninghe Damsté, J. S.: The organic geochemistry of glycerol dialkyl glycerol tetraether lipids: A review, *Org. Geochem.*, 54, 19–61, <https://doi.org/10.1016/j.orggeochem.2012.09.006>, 2013.
- Schouten, S., Hopmans, E. C., Schefuß, E., and Sinninghe Damsté, J. S.: Distributional variations in marine crenarchaeotal membrane lipids: a new tool for reconstructing ancient seawater temperatures?, *Earth Planet. Sci. Lett.*, 204, 265–274, [https://doi.org/10.1016/S0012-821X\(02\)00979-2](https://doi.org/10.1016/S0012-821X(02)00979-2), 2002.
- Segarra, K. E. A., Schubotz, F., Samarkin, V., Yoshinaga, M. Y., Hinrichs, K.-U., and Joye, S. B.: High rates of anaerobic methane oxidation in freshwater wetlands reduce potential atmospheric methane emissions, *Nat. Commun.*, 6, 7477, <https://doi.org/10.1038/ncomms8477>, 2015.

- Sejrup, H. P., Seppä, H., McKay, N. P., Kaufman, D. S., Geirsdóttir, Á., de Vernal, A., Renssen, H., Husum, K., Jennings, A., and Andrews, J. T.: North Atlantic–Fennoscandian Holocene climate trends and mechanisms, *Quat. Sci. Rev.*, 147, 365–378, <https://doi.org/10.1016/j.quascirev.2016.06.005>, 2016.
- 1030 Seppä, H., Bjune, A. E., Telford, R. J., Birks, H. J. B., and Veski, S.: Last nine-thousand years of temperature variability in northern Europe, *Clim. Past*, 5, 523–535, <https://doi.org/10.5194/cp-5-523-2009>, 2009.
- Shala, S., Helmens, K. F., Luoto, T. P., Salonen, J. S., Väiliranta, M., and Weckström, J.: Comparison of quantitative Holocene temperature reconstructions using multiple proxies from a northern boreal lake, *The Holocene*, 27, 1745–1755, 1035 <https://doi.org/10.1177/0959683617708442>, 2017.
- Sinninghe Damsté, J. S., Hopmans, E. C., Pancost, R. D., Schouten, S., and Geenevasen, J. A.: Newly discovered non-isoprenoid glycerol dialkyl glycerol tetraether lipids in sediments, *Chem. Commun.*, 17, 1683–1684, <https://doi.org/10.1039/B004517I>, 2000.
- 1040 Sinninghe Damsté, J. S., Ossebaar, J., Abbas, B., Schouten, S., and Verschuren, D.: Fluxes and distribution of tetraether lipids in an equatorial African lake: constraints on the application of the TEX₈₆ palaeothermometer and BIT index in lacustrine settings, *Geochim. Cosmochim. Acta*, 73, 4232–4249, <https://doi.org/10.1016/j.gca.2009.04.022>, 2009.
- 1045 Sinninghe Damsté, J. S., Ossebaar, J., Schouten, S., and Verschuren, D.: Distribution of tetraether lipids in the 25-ka sedimentary record of Lake Challa: extracting reliable TEX₈₆ and MBT/CBT palaeotemperatures from an equatorial African lake, *Quat. Sci. Rev.*, 50, 43–54, <https://doi.org/10.1016/j.quascirev.2012.07.001>, 2012.
- Sinninghe Damsté, J. S. S., Weber, Y., Zopfi, J., Lehmann, M. F., and Niemann, H.: Distributions and sources of 1050 isoprenoidal GDGTs in Lake Lugano and other central European (peri-)alpine lakes: Lessons for their use as paleotemperature proxies, *Quat. Sci. Rev.*, 277, 107352, <https://doi.org/10.1016/j.quascirev.2021.107352>, 2022.
- Tierney, J. E. and Russell, J. M.: Distributions of branched GDGTs in a tropical lake system: implications for lacustrine application of the MBT/CBT paleoproxy, *Org. Geochem.*, 40, 1032–1036, 1055 <https://doi.org/10.1016/j.orggeochem.2009.04.014>, 2009.
- Tierney, J. E., Russell, J. M., Eggermont, H., Hopmans, E. C., Verschuren, D., and Sinninghe Damsté, J. S.: Environmental controls on branched tetraether lipid distributions in tropical East African lake sediments, *Geochim. Cosmochim. Acta*, 74, 4902–4918, <https://doi.org/10.1016/j.gca.2010.06.002>, 2010.

- Van Bree, L. G., Peterse, F., Baxter, A. J., De Crop, W., Van Grinsven, S., Villanueva, L., Verschuren, D., and Sinninghe Damsté, J. S.: Seasonal variability and sources of in situ brGDGT production in a permanently stratified African crater lake, *Biogeosciences*, 17, 5443–5463, <https://doi.org/10.5194/bg-17-5443-2020>, 2020.
- 1065 Van den Bos, V., Engels, S., Bohncke, S. J. P., Cerli, C., Jansen, B., Kalbitz, K., Peterse, F., Renssen, H., and Sachse, D.: Late Holocene changes in vegetation and atmospheric circulation at Lake Uddelermeer (The Netherlands) reconstructed using lipid biomarkers and compound-specific δD analysis, *J. Quat. Sci.*, 33, 100–111, <https://doi.org/10.1002/jqs.3006>, 2018.
- 1070 Van Wirdum, F., Andrén, E., Wienholz, D., Kotthoff, U., Moros, M., Fanget, A. S., Seidenkrantz, M. S., and Andrén, T.: Middle to Late Holocene variations in salinity and primary productivity in the central Baltic Sea: a multiproxy study from the Landsort Deep, *Front. Mar. Sci.*, 6, 51, <https://doi.org/10.3389/fmars.2019.00051>, 2019.
- Wang, Y., Cheng, H., Edwards, R. L., He, Y., Kong, X., An, Z., Wu, J., Kelly, M. J., Dykoski, C. A., and Li, X.: The
 1075 Holocene Asian monsoon: links to solar changes and North Atlantic climate, *Science*, 308, 854–857, <https://doi.org/10.1126/science.1106296>, 2005.
- Weber, Y., Sinninghe Damsté, J. S., Zopfi, J., De Jonge, C., Gilli, A., Schubert, C. J., Lepori, F., Lehmann, M. F., and Niemann, H.: Redox-dependent niche differentiation provides evidence for multiple bacterial sources of glycerol tetraether
 1080 lipids in lakes, *Proc. Natl. Acad. Sci. USA*, 115, 10926–10931, <https://doi.org/10.1073/pnas.1805186115>, 2018.
- Weijers, J. W., Lim, K. L., Aquilina, A., Sinninghe Damsté, J. S., and Pancost, R. D.: Biogeochemical controls on glycerol dialkyl glycerol tetraether lipid distributions in sediments characterized by diffusive methane flux, *Geochem. Geophys. Geosyst.*, 12, Q10010, <https://doi.org/10.1029/2011GC003724>, 2011.
 1085
- Weijers, J. W., Schouten, S., Spaargaren, O. C., and Sinninghe Damsté, J. S.: Occurrence and distribution of tetraether membrane lipids in soils: Implications for the use of the TEX₈₆ proxy and the BIT index, *Org. Geochem.*, 37, 1680–1693, <https://doi.org/10.1016/j.orggeochem.2006.07.018>, 2006.
- 1090 Weijers, J. W. H., Schouten, S., van den Donker, J. C., Hopmans, E. C., and Sinninghe Damsté, J. S.: Environmental controls on bacterial tetraether membrane lipid distribution in soils, *Geochim. Cosmochim. Acta*, 71, 703–713, <https://doi.org/10.1016/j.gca.2006.10.003>, 2007.

- Woltering, M., Atahan, P., Grice, K., Heijnis, H., Taffs, K., and Dodson, J.: Glacial and Holocene terrestrial temperature variability in subtropical east Australia as inferred from branched GDGT distributions in a sediment core from Lake McKenzie, *Quat. Res.*, 82, 132–145, <https://doi.org/10.1016/j.yqres.2014.02.005>, 2014.
- Xiao, W., Wang, Y., Zhou, S., Hu, L., Yang, H., and Xu, Y.: Ubiquitous production of branched glycerol dialkyl glycerol tetraethers (brGDGTs) in global marine environments: a new source indicator for brGDGTs, *Biogeosciences*, 13, 5883–5894, <https://doi.org/10.5194/bg-13-5883-2016>, 2016.
- Yao, Y., Zhao, J., Vachula, R. S., Werne, J. P., Wu, J., Song, X., and Huang, Y.: Correlation between the ratio of 5-methyl hexamethylated to pentamethylated branched GDGTs (HP5) and water depth reflects redox variations in stratified lakes, *Org. Geochem.*, 147, 104076, <https://doi.org/10.1016/j.orggeochem.2020.104076>, 2020.
- Zander, P. D., Böhl, D., Sirocko, F., Auderset, A., Haug, G. H., and Martínez-García, A.: Reconstruction of warm-season temperatures in central Europe during the past 60,000 years from lacustrine branched glycerol dialkyl glycerol tetraethers (brGDGTs), *Clim. Past*, 20, 841–864, <https://doi.org/10.5194/cp-20-841-2024>, 2024.
- Zhang, Y.-G., Zhang, C. L., Liu, X.-L., Li, L., Hinrichs, K.-U., and Noakes, J. E.: Methane Index: A tetraether archaeal lipid biomarker indicator for detecting the instability of marine gas hydrates, *Earth Planet. Sci. Lett.*, 307, 525–534, <https://doi.org/10.1016/j.epsl.2011.05.031>, 2011.
- Zhang, Z., Smittenberg, R. H., and Bradley, R. S.: GDGT distribution in a stratified lake and implications for the application of TEX₈₆ in paleoenvironmental reconstructions, *Sci. Rep.*, 6, 34465, <https://doi.org/10.1038/srep34465>, 2016.
- Zhao, B., Russell, J. M., Tsai, V. C., Blaus, A., Parish, M. C., Liang, J., Wilk, A., Du, X., and Bush, M. B.: Evaluating global temperature calibrations for lacustrine branched GDGTs: Seasonal variability, paleoclimate implications, and future directions, *Quat. Sci. Rev.*, 310, 108124, <https://doi.org/10.1016/j.quascirev.2023.108124>, 2023.
- Zhao, Y., Liu, Y., Cao, S., Hao, Q., Liu, C., and Li, Y.: Anaerobic oxidation of methane driven by different electron acceptors: A review, *Sci. Total Environ.*, 946, 174287, <https://doi.org/10.1016/j.scitotenv.2024.174287>, 2024.
- Zolitschka, B., Francus, P., Ojala, A. E., and Schimmelmann, A.: Varves in lake sediments – a review, *Quat. Sci. Rev.*, 117, 1–41, <https://doi.org/10.1016/j.quascirev.2015.03.019>, 2015.