

Heat Increased ocean heat transport to the ~~Central~~-central Arctic is Reduced by the ~~despite~~ a well working Barents Sea Cooling Machine

Shaun A. Eisner¹, James A. Carton¹, Leon Chafik², and Lars H. Smedsrud³

¹Department of Atmospheric and Oceanic Science, University of Maryland, College Park, United States

²Department of Meteorology and Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

³Geophysical Institute & Bjerknes Centre for Climate Research, University of Bergen, Bergen, Norway

Correspondence: Shaun Eisner (seisner1@umd.edu)

Abstract. The Barents Sea is a primary gateway for Atlantic Water entering the ~~Central~~-central Arctic Ocean and ubiquitous water-mass transformation on the Barents shelf is key for mitigating increases in heat transport to the central Arctic through the St. Anna Trough. Using a mesoscale-permitting reanalysis spanning 40 years, we derive the first long-term estimate of heat transport through the St. Anna trough, finding that it has increased by ~~0.15~~ 0.11 TW/year since 1980. However, this is only
5 half of the ~~0.28~~ 0.23 TW/year trend in ~~increasing the~~ heat transport into the Barents Sea through the Barents Sea Opening. Decomposing the heat transports reveals that these trends are entirely due to warming temperatures at the sections with no discernible trend in the volume transports. We find that a northward migration of the largest heat fluxes from the ocean to the atmosphere have resulted in cooler and denser ~~Northern Barents Shelf Water~~ waters in the northeastern Barents Sea, mitigating the heat transported through the St. Anna trough. However, despite functioning properly, the "Barents Cooling Machine" has
10 been unable to keep pace with the dramatic warming of the Atlantic Water inflow, resulting in the residual trend in heat transport to the central Arctic. Finally, we present ~~the first observational evidence for~~ empirical evidence for a revised version of the "ocean feedback" hypothesis, ~~indicating that it~~ which modulates volume transport into and out of the Barents Sea on sub-decadal timescales.

1 Introduction

15 The Barents Sea is situated at the entrance to the deep ~~Central~~-central Arctic Ocean. Along with the ~~Bering and Fram Straits~~ Fram strait, this relatively shallow shelf sea is a primary gateway for warm Atlantic Water (AW) flowing northwards. Warm and salty AW enters the Barents Sea through the Barents Sea Opening (BSO), which serves as a major ~~source~~ moderator of heat and salt transport to the central Arctic (~~Skagseth et al., 2008~~) (Skagseth et al., 2008; Smedsrud et al., 2010) (See Figure 1). Despite exporting heat to the central Arctic, the Barents Sea has historically served as a "cooling machine", with strong
20 heat fluxes from the ocean to the atmosphere that cool the inflowing AW and transform it into denser, cooler ~~Barents Shelf Water~~ water masses (Smedsrud et al., 2010, 2013; Skagseth et al., 2020; Årthun et al., 2025). In recent decades, the Barents Sea has experienced a number of climatic changes which fall under the umbrella of "Atlantification", including loss of sea ice, rapid warming, and "Borealization" of its shelf ecosystems (Smedsrud et al., 2022; Ingvaldsen et al., 2021; Onarheim et al., 2018; Onarheim and Årthun, 2017). It has been suggested that ongoing warming and sea ice loss in the Barents has partially been

25 amplified by a slowdown of the "Barents Sea Cooling Machine" (Skagseth et al., 2020), while others suggest that the cooling machine may instead be migrating northward (Shu et al., 2021). Any changes in the efficiency of the Barents Sea cooling will have outsized downstream impacts on the future warming of the Arctic due to the outflow of ~~Barents Shelf Water~~ water from the Barents Sea into the Arctic Circumpolar Boundary Current (ACBC) which transports ~~Barents Shelf Water and AW~~ this water from the St. Anna Trough as far as the Amerasian Basin (Rudels et al., 1999; Woodgate et al., 2001; Aksenov et al., 30 2011). Correctly assessing long-term changes in the Barents Sea cooling system and the effect that these changes have on the properties of ~~Barents Shelf Water outflowing water exiting the Barents Sea~~ through the St. Anna Trough is therefore necessary to understanding the implications of Atlantification on the broader Arctic.

The Barents Sea is a complex system with decadal-scale air-ice-ocean oscillations (Ikeda, 1990) that requires feedback processes to operate. Smedsrud et al. (2013) evaluated two primary feedback mechanisms working to collectively maintain the 35 "Barents Sea Cooling Machine" - a "Wind Feedback" and an "Ocean Feedback". The wind feedback mechanism postulates that increases in the heat transport of AW entering through the BSO results in greater heat loss from the ocean to the atmosphere, resulting in lower surface pressures and stronger westerly winds which in turn continue to drive increased AW inflow. This mechanism has been addressed using models and observations (~~Bengtsson et al., 2004~~)(Bengtsson et al., 2004; Cai et al., 2022), but the increased cyclonic winds were not found to be causing increased AW inflow by Heukamp et al. (2023). The ocean 40 feedback mechanism postulates that the same increases in AW heat transport through the BSO which result in increased heat loss from the ocean to the atmosphere also drive an increase in the formation of dense water. This sets up an SSH gradient between the ~~northern-northeastern~~ Barents Sea and the Eurasian Basin that drives ~~an increase in transport out of the St. Anna Trough, which by mass continuity must increase the volume transport through the BSO~~ a barotropic flow through the Barents Sea that strengthens the inflow through BSO and outflow through STA.

45 Unlike the wind feedback mechanism, there has been little discussion and no direct observational evidence of the proposed ocean feedback mechanism, due primarily to a lack of sustained observations in the ~~Northern Barents-northeastern Barents Sea~~ Barents Sea in the vicinity of St. Anna Trough (~~although there are some observations, which have allowed for estimates of time mean quantities, such as in Schauer et al. (2002).~~). Recent discussions regarding the Barents Sea cooling machine have focused on the warming inflow through the BSO and a reduction in ocean heat loss associated with warmer ~~Barents Shelf Waters~~ (Skagseth et al., 2020). ~~However, there has been speculation that the cooling system has migrated northward, rather than reducing in efficiency~~ (Shu et al., 2021). ~~waters in the Barents Sea~~ (Skagseth et al., 2020). Again, sparse observations in the ~~Northern Barents-northeastern Barents Sea~~ Barents Sea limit the ability to assess whether changes to the location or efficiency of the cooling machine manifest in the transport of ~~Barents Shelf Water~~ water from the Barents Sea through the St. Anna trough and ultimately into the central Arctic, a key point we focus on in this study.

55 Reanalyses present a potential solution to the problem of assessing long-term changes in the export of ~~Barents Shelf Water~~ water from the Barents Sea and assessing the ocean feedback mechanism of Smedsrud et al. (2013). Historically, reanalyses have been used effectively to produce long-term estimates of ocean heat and volume transports (Carton and Santorelli, 2008; Karspeck et al., 2017). Comparison studies however, such as that of Uotila et al. (2019), have shown that many reanalyses lack sufficient vertical and horizontal resolution to adequately capture the structure and dynamics of the polar region. In this

60 study we use the new, mesoscale-permitting SODA4 reanalysis, which builds upon the Regional Arctic Reanalysis (RARE)
(Chepurin et al., 2025; Carton and Chepurin, 2023). SODA4 improves upon the vertical and horizontal resolution of existing
reanalyses and we assess its consistency with observations in the Barents Sea region. We use SODA4 to study trends in heat
transport through the Barents Sea Opening and St. Anna Trough from 1980-2020. Additionally, we assess how changes in the
Barents Sea cooling ~~system-processes~~ have affected the heat of ~~outflowing Barents Shelf Water~~ the Barents Sea outflow waters
65 and the evidence supporting the proposed ocean feedback hypothesis.

2 Methods

2.1 Reanalysis Construction

The SODA4 reanalysis is a global $\frac{1}{10}^\circ$, 75 vertical level ocean state estimate spanning the period from 1980 to the near present.
The base model is a coupled MOM5.1/SIS1 ice-ocean model on an Arakawa B-grid with displaced poles. The reanalysis
70 assimilates temperature and salinity profiles from the 2023 World Ocean Database (Mishonov, 2024) as well as ICOADS 3.0
(Freeman et al., 2017) and NOAA nighttime L3 Sea Surface Temperatures (~~Chepurin et al., 2025~~) (Jonasson et al., 2022b, a)
(Chepurin et al., 2025). In the Barents Sea, St. Anna Trough, and surrounding regions, SODA4 assimilates ~75,000 individual
profiles from the world ocean database over the period of coverage. Atmospheric forcing is provided by the ERA-5 atmospheric
reanalysis (Hersbach et al., 2019). ~~While the reanalysis~~ For more information on the construction of SODA, we refer the reader
75 to Carton and Giese (2008) or Chepurin et al. (2025).

While SODA4 is available at 5-daily temporal frequency, for this study, the monthly frequency output is used.

Temperature, salinity, velocity, sea surface height (SSH), and mixed layer depth (using the de Boyer Montégut et al. (2004)
density criterion) are all obtained directly from the SODA4 monthly output. While total surface heat fluxes are available from
the SODA4 reanalysis, in this study they are obtained directly from the ERA-5 reanalysis so that they can be separated into
80 turbulent (Latent & Sensible) and radiative (Short-wave & Long-wave) components.

In order to assess long-term changes in hydrographic properties, heat content, and atmospheric heat fluxes, time means of
each quantity are computed over each decade: 1980-1989, 1990-1999, 2000-2009, 2010-2019, and 2020-2024. Heat flux and
other rate of change quantities are first annually integrated and the annual integrals are then averaged over the appropriate
decadal period, while direct hydrographic properties such as temperature and salinity are simply averaged over the entire
85 decadal period.

In order to assess the coupling in the hypothesized ocean feedback mechanism, the SSH gradient is considered, which
supposedly drives the associated ~~geostrophic anomalies in flow through STA~~ barotropic flow through the Barents Sea. For
this assessment, we ~~construct a single numerical quantity which is referred to as the "SSH curl" or as " $\nabla \times$~~ consider the
90 spatial average of the magnitude of the SSH " in an abuse of notation. We define this quantity as the sum of the zonally and
~~meridionally-directed components of the SSH streamfunction. The "SSH curl" can be interpreted as measuring the strength of~~

~~the cyclonically-directed geostrophic flow generated by an SSH gradient anomaly.~~

$$\nabla \times SSH \equiv \frac{\partial SSH}{\partial x} - \frac{\partial SSH}{\partial y}$$

gradient, ∇SSH across the Barents Sea.

2.2 In Situ Observations

95 Annual time series of 0-200m depth averaged temperature and salinity are obtained from the Bear Island and Kola section time series (obtained from the ICES database: <https://ocean.ices.dk/iroc/>) and directly compared to annually and depth averaged temperature and salinity from the SODA4 output (Gonzalez-Pola et al., 2023). Time-mean hydrographic properties of the SODA4 output are also directly compared to the World Ocean Atlas 2023 climatology (Reagan et al., 2023).

2.3 Transport & Sections

100 Net heat and volume transports are evaluated over a number of key sections, including the BSO, the Fram Strait (FS), the St. Anna Trough (STA), and across the ACBC. The extent of these sections are indicated in Table 1 and Figure 1. All transports are depth-integrated to a maximum depth of 1000m. This cutoff depth is chosen as the data assimilation is poorly constrained in the Arctic at depths below 1000m. This yields a heat transport D_Q :

$$D_Q = \int_{z=-1000m}^{z=0m} \mathbf{u} \cdot \hat{\mathbf{n}} \rho C_p (T - T_{ref}) dA, \quad T_{ref} = -1.8^\circ C \quad (1)$$

105 where \mathbf{u} is the horizontal velocity vector and $\hat{\mathbf{n}}$ is the unit normal vector to the surface through which the transport is calculated. In the case of our sections, the unit normal is always either directly zonal or directly meridional, reducing the normal component of the horizontal velocity to either u or v .

While a reference temperature of $0^\circ C$ has traditionally been used for heat transport calculations in the Barents Sea region (Smedsrud et al., 2010), we adopt a reference temperature of $-1.8^\circ C$ in accordance with Woodgate et al. (2001) and Pnyushkov et al. (2018, 2021) so as to avoid any confusion in transport trends or direction resulting from temperatures in the vicinity of $0^\circ C$. $-1.8^\circ C$ is a natural lower limit for the reference temperature, as it is close to the freezing-point. To provide direct comparison with prior estimates regardless of reference temperature, we compute the true heat flux convergence of the Barents Sea, following Smedsrud et al. (2010). The true heat flux convergence is defined as the integrated divergence of heat over the entire volume, but can be rewritten in terms of the sum of heat transports across each of the boundaries using Green's Theorem:

115

$$\iiint \nabla \cdot \rho C_p (T - T_{ref}) dV = \sum_{i=1}^n D_Q^i \quad (2)$$

Therefore, the true heat flux convergence to the Barents Sea is independent of reference temperature as long as the volume budget is closed, but the absolute values at each section vary with this choice. In SODA4, the volume budget is approximately

closed when considering flow through BSO and the entire northern boundary (19-75°E, 79.5°N), so the true heat convergence
120 in SODA4 is approximately the sum of D_Q^{BSO} and D_Q^{NB} , where NB represents the northern boundary. For reference, we also
compute the heat convergence and time mean heat transports for the well-known GLORYS12 reanalysis (Lellouche et al.,
2021) (See Table A1).

Table 1. Transport Sections

Section	Longitudes	Latitudes
Barents Sea Opening (BSO)	19 E	70-75 N
St. Anna Trough (STA)	65-75 E	79.5 N
Fram Strait (FS)	3W-10E	79 N
Arctic Boundary Current (ACBC)	90E	81.5-83N

3 Results

3.1 Decadal Changes in Hydrography

125 We assess changes in the decadal-averaged vertical profiles of hydrographic properties as well as the change in the 2020s
decadally averaged heat content relative to the 1980s. Relative to the 1980s, the 2020s Barents-Kara sea heat content increased
by 1.1 ZJ. This is consistent with a steady $\sim 0.2^\circ\text{C}$ increase in potential temperature each decade. Notably, the increases
in potential temperature appear depth-amplified, ~~particularly in the 2020s, suggesting that increases in heat content are the~~
~~result of warmer inflows at greater than 50m depths (Figure 2a). While overall~~ While surface salinity and potential density
130 decreased by 0.5 psu and 1 kg/m³ respectively (Figure 2b), the trend is not steady and appears dominated by the signal in the
Kara Sea and ~~Northern-northeastern~~ Northern-northeastern Barents Sea. In the southern Barents Sea, both salinity and density have increased steadily
since the 1980s. ~~Increases in heat content are ubiquitous with the greatest increases in the vicinity of the Barents Sea Opening,~~
~~St. Anna Trough, and west of Novya Zemlya (Figure 4a).~~

To assess the reliability of SODA4, we provide comparison to the annual time series of temperature and salinity at the Bear
135 Island and Kola sections. In the vicinity of the BSO, SODA4 has a cold bias, while in the vicinity of Kola section, it has a slight
warm bias. The same pattern of temperature bias is evident when comparing the time mean SODA4 hydrography to the World
Ocean Atlas 2023, but is less pronounced in amplitude (see Figure S2S1). There is a 1-2°C warm bias in the Kara Sea and in
the vicinity of Novaya Zemlya, with intermittent regions of 1-1.5°C cold bias throughout the Southern Barents, near the BSO.
Despite the evident biases when compared with observations, the annual anomalies of temperature and salinity in SODA4 are
140 well correlated with those present in the Bear Island and Kola section time series (see Figure 3). Additionally, the same upward
positive trends in temperature ~~anomalies~~ that are evident in observations are reproduced in the SODA4 hydrography. This
supports the idea that despite the biases, the anomalies (and trends) in temperature are still consistent between the reanalysis

and observations, allowing us to assess trends in the heat transport. This is further supported by a realistic time mean heat convergence for the Barents Sea in SODA4, which is further discussed in the following section.

145 It is worth noting that there is a distinct ~~downward-negative~~ trend in salinity ~~anomalies~~ at both Bear Island and Kola section beginning in 2010. This appears consistent with the "great salinity anomaly" identified by Holliday et al. (2020). In SODA4, this anomaly is evident at Bear Island, but is not evident at the Kola section. For this reason, SODA4 deviates from observed salinity anomalies at Kola section by up to 0.2 psu beginning in 2015. This same pattern does not appear in the temperature anomalies.




Figure 1. The Barents Sea bathymetry and the analyzed sections; The Barents Opening (BSO), Fram Strait (FS), St. Anna Trough (STA), the Arctic Circumpolar Boundary Current (ACBC) and the Kola Section (KS). The northern boundary (19-75°E, 79.5°N) is also indicated as a dashed line. Colored red/purple arrows indicate the path of Atlantic Water as it traverses the shelfbreak, enters, and leaves the Barents Sea. Colored blue arrows indicate outflow of Arctic water via the East Greenland Current. Volume transports of the major inflows and outflows are used from Smedsrud et al. (2010) and Pnyushkov et al. (2018).

150 3.2 Heat Transport

~~Heat transport through both~~ Increases in heat content are ubiquitous in the Barents Sea with the greatest increases in the vicinity of the Barents Sea Opening, St. Anna Trough, and west of Novya Zemlaya (Figure 4a). Additionally, heat transport through the BSO and STA have increased over the 40-year period. However, the rate of increase of heat transport through the St. Anna trough (~~0.15-0.11~~ 0.15-0.11 TW/year) is roughly half that of the increase through the Barents Sea Opening (~~0.28-0.23~~ 0.28-0.23 TW/year), indicating that at least half of the additional heat advected into the Barents Sea is being lost (Figure ~~4b,c5~~ 4b,c5). The additional heat that does get transported through STA appears to be the primary source of increased heat in the core of the ACBC, which carries ~~Barents Shelf Water~~ water out of the Barents Sea, exiting through STA along the Eurasian shelf. There is a clear ~~increasing-positive~~ trend in the average temperature of water transported through the ACBC section (warming by almost 1°C over the entire 40 year period), but no discernible trend in heat transport through Fram Strait (Figure 5). This

160 indicates that the increases in heat transport through the STA are the primary advective source of increased heat transported to




Figure 2. Simulated Barents Sea properties over time. a) Vertical profiles of spatially averaged hydrographic properties over the entire Barents Sea domain for each decade. 1σ standard deviations for each decade are also indicated via shading. Potential temperature increases relatively uniformly each decade and is slightly depth intensified. Changes in salinity and density are largely dominated by decreases in the Eastern Kara Sea and Eurasian shelfbreak due to ice melt and increased river freshwater export.




Figure 3. Time Series of annual anomalies of Potential Temperature (LeftTop) and Salinity (RightBottom) at Bear Island (TopLeft) and Kola section (BottomRight) relative to the 1980-2023 time mean.

the central Arctic. It is worth noting however, that despite a lack of a trend in heat transport through Fram Strait, we do find a slight trend (~0.012°C/year) in the average temperature of water flowing through Fram Strait. This is similar to the findings of Beszczynska-Möller et al. (2012), who identified a 0.06°C/year trend. A potential reason for discrepancies in the magnitude

of the trend could be a mismatch in the sections considered, as we exclude the westernmost portion of Fram Strait so as not to count recirculating AW or the East Greenland Current.

The true heat convergence to the Barents Sea in SODA4 is 77.6 TW. This is nearly identical to the results of Smedsrud et al. (2010), which estimated a true heat convergence for the Barents Sea of ~ 73 TW. When referenced to a temperature of -1.8°C , 86.8 TW enters through the BSO and 15.9 TW exits through the northern boundary extending from Svalbard to the easternmost edge of STA in SODA4. The vast majority (11.2 TW) of this exiting heat leaves through the STA itself between $65\text{-}75^{\circ}\text{E}$. The close matching of the true heat convergence indicates that the temperature biases in SODA4 have little impact on the modelled heat convergence in the Barents Sea (Figure 4b).

Decomposing heat transport through the BSO and STA reveals that the increases in heat transport are entirely the result of changes in ~~heat content~~temperature, rather than changes in the volume transport (Figure 4c). The average potential temperature of the inflow through BSO has increased by 0.05 C/year , with a 0.02 C/year increase in the average temperature of the outflow through STA, despite no significant trends in the volume transport through either opening (Figure S1S2).

3.3 Heat Fluxes

The ~~reduced-smaller~~ increase in heat transport through STA relative to the BSO suggests that there must be a heat loss mechanism to ~~accomodate~~accommodate the reduction. We assess the change in decadal-averaged turbulent, radiative, and total heat fluxes in the 2020s relative to the 1980s. There is a clear pattern in both the turbulent and total heat fluxes of a reduction in the amount of cooling in the Southern Barents Sea alongside an increase in the amount of cooling in the central/~~northern~~northeastern Barents Sea (Figure 6a,c). Therefore, while atmospheric cooling efficiency is decreased in the vicinity of the BSO, there is an overcompensating increase in the ~~northern Barents~~northeastern Barents Sea. This dipolar pattern is largely the result of the wintertime (DJF) heat flux pattern and appears characteristic of ice edge retreat (Figure S3). When the ice edge retreats, water which was south of the ice edge now experiences warmer surface air temperatures and transfer less heat to the atmosphere, while newly ice-free waters remain warmer than surface air temperatures and are now exposed to intense atmospheric forcing, allowing for increased heat loss to the atmosphere. It is also consistent with the modelled pattern of heat flux changes in Shu et al. (2021).

3.4 Dense Water Formation

The ocean feedback mechanism of Smedsrud et al. (2013) suggests that increased heat loss from the ocean to the atmosphere drives increased dense water formation, that then, by way of an SSH gradient, generates ~~an increase in outflow through the St. Anna trough~~a barotropic flow through the Barents Sea, increasing transport through the BSO and STA. Skagseth et al. (2020) found that heat loss in the southern Barents had decreased, while the overall density had remained constant. We investigate the change in decadal-averaged density in the 2020s relative to the 1980s. Our results agree with the findings of Skagseth et al. (2020) that density in the Southern Barents has remained relatively constant over the previous 40 years. However, a hotspot of increasing density is evident in the ~~Northern Barents~~northeastern Barents Sea in the vicinity of the new ice edge (Figure 7a). This density change is collocated with a depression in SSH which establishes a strong SSH gradient between the ~~northern~~




Figure 4. (a) The change in decadal-averaged Ocean Heat Content in the 2020s relative to the 1980s. There are three "hotspots" located at the Barents Sea Opening, just west of Novaya Zemlya, and in the St. Anna trough. The mean ice edge for each decade is indicated. (b) Time series of the "true" heat convergence $\nabla_h \cdot Q$ of the Barents Sea in SODA4. The heat convergence of Smedsrud et al. (2010) is also indicated. (c) Time series of the temperature and volume decomposed heat transports ($\overline{T}v'$ and $v'\overline{v}$) through St. Anna Trough (STA) and Barents Sea Opening (BSO). Trend lines are indicated as well. There is no discernible trend in the volume component of heat transport through either opening, but trends in the temperature components of both the BSO and STA heat transports.

northeastern Barents Sea and the Eurasian Basin (Figure 7b). The increased density is consistent with an increase in mixed layer depth and decrease in base of ML stratification in the ~~Northern Barents~~ northeastern Barents Sea, indicating that the increase in cooling and dense water formation results in increased vertical mixing and erosion of near-surface stratification

200 (Figure 7c,d).




Figure 5. Time series of heat transport through the Barents Sea Opening (BSO), St. Anna Trough and Fram Strait. Average temperature along a section of the (bottom) Arctic Circumpolar Boundary Current (ACBC) is also indicated ([the extent of this section is given in Table 1](#)). Linear trends are indicated with dashed lines. There are increasing trends in heat transport through St. Anna Trough and the average temperature through the ACBC section but no discernible trend in heat transport through Fram Strait.

3.5 The Ocean Feedback Mechanism

We assess the correlations between the SSH ~~curl~~[gradient](#) and the outflow through St. Anna Trough over the 40-year time period. ~~Despite little to no trend in the volume transport through STA and a decreasing trend in the SSH curl in the vicinity of STA, there is a negative~~ [There is a positive](#) correlation between the two, with ~~$r = -0.87$~~ [\$r = 0.7\$](#) (Figure 8a). Additionally, there is a positive correlation between the outflow through STA and the inflow through BSO (~~$r = 0.84$~~ [\$r = 0.87\$](#)) (Figure 8b). To assess the time scales over which this relationship acts, the correlation and associated significance between STA and BSO transport and SSH curl and STA transport are assessed as a function of low-frequency cutoff (Figure 9). ~~STA-BSO~~ [SSH-STA](#) transport correlations fall off sharply after ~~1-2 years~~ [1 year](#), becoming insignificant beyond ~~the 3.5 year cutoff~~ [this point](#). Meanwhile, ~~SSH curl-STA~~ [BSO-STA](#) transport correlations fall ~~only slightly at 1-2 year cutoffs before steadily increasing to a maximum~~




Figure 6. Change in decadal-averaged Turbulent, Radiative, and total heat fluxes in the 2020s relative to the 1980s (positive values indicate upward heat flux out of the ocean). Changes in total heat fluxes resemble changes in turbulent heat flux, with increased heat leaving the ocean (resulting in increased cooling) in the northern-northeastern Barents sea, northwest of Novaya Zemlya, and decreased heat leaving the ocean in the southern Barents Sea (resulting in decreased cooling).




Figure 7. Change in decadal-averaged depth-averaged density (top-left) density, (b) mixed layer depth (top-right), curl of ssh (bottom-left) SSH gradient, and stratification at the base of the mixed layer (bottom-right) Brunt-Vaisala Frequency for the 2020s relative to the 1980s. SSH anomaly contours are shown in the left panels as well on (a) and (c). A region of increased density and SSH depression is evident in the Northern-northeastern Barents Sea, northwest of Novaya Zemlya. A region of anomalously low-strengthened SSH curl-gradient is evident in the St. Anna trough. Mixed layer depth and stratification both decrease in along the north-central northern boundary between the Barents Sea in the vicinity of ice-edge withdrawal and Eurasian Basin.

210 at a cutoff of 15 years. However, due to the reduced degrees of freedom, these correlations become insignificant after the 8 year cutoff. This indicates off at 4 years, becoming insignificant past this point. Together, these indicate that the relationships between STA outflow and SSH curl in the NE Barents remain consistent across most sub-decadal timescales and that the STA outflow-BSO inflow response only occurs at very short (1-3 year), interannual timescales. At timescales longer than 8 years,

215 it is possible that SSH curl and STA transport do have meaningful correlations but the time series is not long enough for such correlations to be statistically significant. Overall, this indicates that the feedback mechanism is not coherent at greater than decadal time scales the SSH gradient, STA outflow, and BSO inflow are only coherent at interannual to subdecadal timescales (Figure 9). The lack of correlation between the BSO inflow and STA outflow at greater-than-interannual timescales is also consistent with the findings that there is no significant trend in the volume transport through either opening over the 40-year period.

220 Collectively, this produces a revised picture of the ocean feedback mechanism whereby dense water formation has migrated northward towards the northeastern Barents Sea (consistent with Shu et al. (2021)) which has created an anomalous SSH gradient between the Barents Sea and central basin. Sub-decadal fluctuations in the SSH gradient then rapidly (over timescales less than 1 year) drive a barotropic flow through the Barents Sea, which adjusts transport through STA and BSO to reduce the SSH anomaly. The coupling between the BSO inflow and STA outflow is coherent over slightly longer timescales (up to 4 years), likely because of the multi-year flushing time which is experienced from when water masses enter the Barents Sea to when they leave (a back of the envelope calculation based on the average volume transport of 2.3 Sv from Smedsrud et al. (2010) would suggest this flushing time is ~ 4 years).




Figure 8. Scatter plots of $\nabla \times SSH$ in the NE Barents Sea vs. Volume Transport through STA (left) and Volume Transport through STA vs. Volume transport through BSO (right) with no low frequency cutoff. There is a clear linear relationship between SSH curl in the NE Barents and the outflow through STA, as well a linear relationship between the BSO inflow and STA outflow.

4 Discussion & Conclusions

230 The Barents Sea is a key moderator of heat and salt transport to the Central Arctic Ocean. Historically, the Barents Sea has served as a "cooling machine" where incoming warm and saline AW is cooled into denser Barents Shelf Water masses before emptying into the Eurasian Basin. However, recent studies have questioned the ongoing efficiency of the cooling machine. Despite its importance to the heat budget of the broader Arctic Ocean and understanding the mechanisms driving the




Figure 9. Pearson's r and associated significance (p -value) between the low frequency components of the BSO-STA outflows and the $\nabla \times SSH-STA-SSH-STA$ outflow as a function of the low frequency cutoff. A $p = 0.05$ threshold is also indicated. There are maxima for both values at the shortest timescales. $BSO-STA-SSH-STA$ correlations fall off sharply after 1 year, becoming insignificant by the 3-year cutoff, while $SSH-STA-BSO-STA$ correlations remain high at longer timescales but become insignificant at timescales greater than 8 years by the 4-year cutoff.




Figure 10. Diagram of the ~~observed~~ proposed ocean feedback loop consistent with the results of Smedsrud et al. (2013). The feedback is driven by dynamical processes that occur at sub-decadal timescales.

Barents Sea cooling machine, there remain no long-term observational records of heat transport exiting the Barents Sea. To address this lack of observational information, we use the SODA4 reanalysis to assess multi-decadal trends in transport and hydrography in the Barents Sea. We also provide ~~the first observational~~ evidence to support ~~the~~ a modified version of Smedsrud's original "Ocean-Feedback-Mechanismocean feedback" , ~~which has been proposed to constitute part of the broader cooling system (Smedsrud et al., 2013)~~ mechanism (Smedsrud et al., 2013), which we find to be restricted to sub-decadal timescales.

SODA4 depicts a clear ~~increasing~~ positive trend in the heat transport through both the Barents Sea Opening (~~0.28~~ 0.23 TW/year) and the St. Anna Trough (~~0.15~~ 0.11 TW/year), which are entirely the result of increases in the average temperature, with little to no change in volume transport over the 40 year period. Of particular note is that despite a decrease in heat loss to the atmosphere in the southern Barents, temperature changes in the 20s relative to the 80s are bottom amplified. This implies that the inflowing water from the Nordic Seas is the primary warming source. The temperature anomalies of SODA4 are consistent with those of the Bear Island and Kola section time series, suggesting that the increase in heat transport as a result of increasing water temperature is also consistent with observations. ~~This trend~~ While the increasing heat transport through BSO is consistent with prior studies, such as Skagseth et al. (2008, 2020); Årthun and Schrum (2010); Wang et al. (2019), it is difficult to compare the magnitude of the trend due to the limited temporal availability of observations which produces significant uncertainty in trend estimates. The primary time series of observations is the Bear Island section which does show the identified increase in mean temperature but does not span the entire BSO and thus cannot be used to estimate the time series of heat transport. While prior studies do find some trend in volume transport (notably the modelling experiments of Wang et al. (2019)), these studies also find that there is an even stronger trend in the temperature of the inflow, consistent with our finding that the trend in BSO heat transport is temperature, rather than volume dominated. The trend in heat transport has contributed to a significant warming of the water in the ACBC core, predominantly by reducing the capability of the Barents Shelf Water outflowing water from the Barents Sea to cool overlying AW transported from Fram Strait (which has remained fairly constant in its heat transport). Therefore, despite a mitigation in heat exiting the Barents Sea through the STA, the observed trend in STA heat transport still manifests in downstream increases in heat content, with significant implications for the warming of the central Arctic.

The ocean feedback mechanism proposed by Smedsrud et al. (2013) suggested that an increase in dense water formation in the Barents Sea sets up an SSH gradient that drives ~~increased volume transport through the STA and increased inflow through the BSO by mass continuity~~ barotropic flow through the Barents Sea, increasing inflow through BSO and outflow through STA. Our results support this hypothesis but only on sub-decadal timescales (see Figure 109). At longer timescales, the outflow through the STA is decoupled from both the BSO inflow and the SSH gradient. However, the increase in heat loss to the atmosphere, and dense water formation in the ~~Northern-northeastern~~ Barents Sea since the 1980s do suggest that the Barents Sea cooling ~~system processes~~ (including the ocean feedback mechanism) has migrated northward. This has resulted in less efficient cooling and less dense water formation in the southern Barents Sea. This is consistent with the findings of both Skagseth et al. (2020) and Shu et al. (2021) but suggests that there is more to the overall picture. While the poleward migration of the Barents Sea cooling machine has mitigated the increases in heat transport to the central Arctic by almost half, it has not been efficient enough to dissipate all of the increased heat entering through the ~~Barents-Sea-Opening~~ BSO. We suspect that the proposed Ocean-Feedback mechanism does play a role in dissipating the excess heat, but that it does so through dynamical processes that occur largely at sub-decadal timescales.

These results have significant implications for future understandings of heat transport into the central Arctic. Our results suggest that continued warming of inflowing AW will continue to be at least partially offset by the Barents ~~Cooling-Machine~~ Sea cooling machine but that at some point, the excess warming will become great enough so as to overwhelm the capacity of

the cooling machine. Therefore, continued modelling efforts from models capable of adequately resolving the Arctic mesoscale as well as observations in the Barents Sea and Eurasian Arctic are critical for determining the ultimate fate of the cooling machine.

Data availability. Time series from Kola and Bear Island Sections (S and T) are available at <https://ocean.ices.dk/iroc/>. SODA4 output is available at <https://dsrs.atmos.umd.edu/DATA>. GLORYS12 was obtained from the E.U. Copernicus Marine Service (<https://doi.org/10.48670/moi-327-00021>).

Author contributions. **SAE:** Conceptualization, Methodology, Software, Formal Analysis, Investigation, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization. **JAC:** Conceptualization, Methodology, Data Curation, Writing - Review & Editing, Resources, Supervision, Project Administration, Funding Acquisition. **LC:** Data Curation, Methodology, Writing - Review & Editing, Validation, Resources, Supervision, Funding Acquisition. **LHS:** Methodology and Heat Transport calculations, Writing - Review & Editing.

Competing interests. The authors declare there are no competing interests.

Acknowledgements. We are grateful to the Physical Oceanography Program of the National Science Foundation (OCE1948952) for providing financial support for this work.

L. Chafik was supported by the ECO2NORSE project funded by the Swedish National Space Agency (Dnr 2022-00172).

[L. H. Smedsrud was supported by the Overturning circulation in the new Arctic \(ArMOC\) project funded by the Norwegian Research Council \(Grant 335255\).](#)

References

- 290 Aksenov, Y., Ivanov, V. V., Nurser, A. G., Bacon, S., Polyakov, I. V., Coward, A. C., Naveira-Garabato, A. C., and Beszczynska-Moeller, A.: The Arctic circumpolar boundary current, *Journal of Geophysical Research: Oceans*, 116, 2011.
- Årthun, M. and Schrum, C.: Ocean surface heat flux variability in the Barents Sea, *Journal of Marine Systems*, 83, 88–98, 2010.
- Bengtsson, L., Semenov, V. A., and Johannessen, O. M.: The early twentieth-century warming in the Arctic—A possible mechanism, *Journal of Climate*, 17, 4045–4057, 2004.
- 295 Beszczynska-Möller, A., Fahrbach, E., Schauer, U., and Hansen, E.: Variability in Atlantic water temperature and transport at the entrance to the Arctic Ocean, 1997–2010, *ICES Journal of Marine Science*, 69, 852–863, <https://doi.org/10.1093/icesjms/fss056>, 2012.
- Cai, Z., You, Q., Chen, H. W., Zhang, R., Chen, D., Chen, J., Kang, S., and Cohen, J.: Amplified wintertime Barents Sea warming linked to intensified Barents oscillation, *Environmental research letters*, 17, 044 068, 2022.
- Carton, J. A. and Chepurin, G. A.: RARE: The Regional Arctic Reanalysis, *Journal of Climate*, 36, 2333 – 2348, <https://doi.org/10.1175/JCLI-D-22-0340.1>, 2023.
- 300 Carton, J. A. and Giese, B. S.: A Reanalysis of Ocean Climate Using Simple Ocean Data Assimilation (SODA), *Monthly Weather Review*, 136, 2999 – 3017, <https://doi.org/10.1175/2007MWR1978.1>, 2008.
- Carton, J. A. and Santorelli, A.: Global decadal upper-ocean heat content as viewed in nine analyses, *Journal of Climate*, 21, 6015–6035, 2008.
- 305 Chepurin, G. A., Carton, J. A., Sun, L., and Penny, S. G.: SODA4: a mesoscale ocean/sea ice reanalysis 1980–2024, <https://doi.org/10.5194/egusphere-2025-3810>, 2025.
- de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology, *Journal of Geophysical Research: Oceans*, 109, <https://doi.org/https://doi.org/10.1029/2004JC002378>, 2004.
- 310 Freeman, E., Woodruff, S. D., Worley, S. J., Lubker, S. J., Kent, E. C., Angel, W. E., Berry, D. I., Brohan, P., Eastman, R., Gates, L., Gloeden, W., Ji, Z., Lawrimore, J., Rayner, N. A., Rosenhagen, G., and Smith, S. R.: ICOADS Release 3.0: a major update to the historical marine climate record, *Int. J. Climatol.*, 37, 2211–2232, 2017.
- Gonzalez-Pola, C., Larsen, K. M. H., Fratantoni, P., and Beszczynska-Möller, A.: ICES Report on Ocean Climate 2021, <https://doi.org/10.17895/ices.pub.24755574.v1>, 2023.
- 315 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., et al.: ERA5 monthly averaged data on single levels from 1979 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS), 10, 252–266, <https://doi.org/10.24381/cds.f17050d7>, accessed on 14-05-2025, 2019.
- Heukamp, F. O., Kanzow, T., Wang, Q., Wekerle, C., and Gerdes, R.: Impact of cyclonic wind anomalies caused by massive winter sea ice retreat in the Barents Sea on Atlantic Water transport toward the Arctic: A model study, *Journal of Geophysical Research: Oceans*, 128, [e2022JC019045](https://doi.org/10.1029/2022JC019045), 2023.
- 320 Holliday, N. P., Bersch, M., Berx, B., Chafik, L., Cunningham, S., Florindo-López, C., Hátún, H., Johns, W., Josey, S. A., Larsen, K. M. H., et al.: Ocean circulation causes the largest freshening event for 120 years in eastern subpolar North Atlantic, *Nature communications*, 11, 585, 2020.
- Ikeda, M.: Decadal oscillations of the air-ice-ocean system in the Northern Hemisphere, *Atmosphere-Ocean*, 28, 106–139, 1990.

- 325 Ingvaldsen, R. B., Assmann, K. M., Primicerio, R., Fossheim, M., Polyakov, I. V., and Dolgov, A. V.: Physical manifestations and ecological implications of Arctic Atlantification, *Nature Reviews Earth & Environment*, 2, 874–889, 2021.
- Jonasson, O., Gladkova, I., and Ignatov, A.: Towards global daily gridded super-collated SST product from low earth orbiting satellites (L3S-LEO-Daily) at NOAA, in: *Ocean Sensing and Monitoring XIV*, vol. 12118, pp. 40–51, SPIE, 2022a.
- Jonasson, O., Ignatov, A., Pryamitsyn, V., Petrenko, B., and Kihai, Y.: JPSS VIIRS SST Reanalysis Version 3, *Remote Sensing*, 14, 3476,
330 2022b.
- Karspeck, A., Stammer, D., Köhl, A., Danabasoglu, G., Balmaseda, M., Smith, D., Fujii, Y., Zhang, S., Giese, B., Tsujino, H., et al.: Comparison of the Atlantic meridional overturning circulation between 1960 and 2007 in six ocean reanalysis products, *Climate Dynamics*, 49, 957–982, 2017.
- Lellouche, J.-M., Eric, G., Romain, B.-B., Gilles, G., Angélique, M., Marie, D., Clément, B., Mathieu, H., Olivier, L. G., Charly, R., et al.:
335 The Copernicus global 1/12 oceanic and sea ice GLORYS12 reanalysis, *Frontiers in Earth Science*, 9, 698 876, 2021.
- Mishonov, A. V.: *World Ocean Database 2023*, 2024.
- Onarheim, I. H. and Årthun, M.: Toward an ice-free Barents Sea, *Geophysical Research Letters*, 44, 8387–8395, 2017.
- Onarheim, I. H., Eldevik, T., Smedsrud, L. H., and Stroeve, J. C.: Seasonal and regional manifestation of Arctic sea ice loss, *Journal of Climate*, 31, 4917–4932, 2018.
- 340 Pnyushkov, A. V., Polyakov, I. V., Rember, R., Ivanov, V. V., Alkire, M. B., Ashik, I. M., Baumann, T. M., Alekseev, G. V., and Sundfjord, A.: Heat, salt, and volume transports in the eastern Eurasian Basin of the Arctic Ocean from 2 years of mooring observations, *Ocean Science*, 14, 1349–1371, 2018.
- Pnyushkov, A. V., Polyakov, I. V., Alekseev, G., Ashik, I., Baumann, T. M., Carmack, E., Ivanov, V., and Rember, R.: A steady regime of volume and heat transports in the eastern Arctic Ocean in the early 21st century, *Frontiers in Marine Science*, 8, 705 608, 2021.
- 345 Reagan, J. R., Boyer, T. P., García, H. E., Locarnini, R. A., Baranova, O. K., Bouchard, C., Cross, S. L., Mishonov, A. V., Paver, C. R., Seidov, D., Wang, Z., and Dukhovskoy, D.: *World Ocean Atlas 2023*, <https://www.ncei.noaa.gov/archive/accession/0270533>, 2023.
- Rudels, B., Friedrich, H. J., and Quadfasel, D.: The Arctic circumpolar boundary current, *Deep Sea Research Part II: Topical Studies in Oceanography*, 46, 1023–1062, 1999.
- Schauer, U., Loeng, H., Rudels, B., Ozhigin, V. K., and Dieck, W.: Atlantic water flow through the Barents and Kara Seas, *Deep Sea Research Part I: Oceanographic Research Papers*, 49, 2281–2298, 2002.
350
- Shu, Q., Wang, Q., Song, Z., and Qiao, F.: The poleward enhanced Arctic Ocean cooling machine in a warming climate, *Nature Communications*, 12, 2966, 2021.
- Skagseth, Ø., Furevik, T., Ingvaldsen, R., Loeng, H., Mork, K. A., Orvik, K. A., and Ozhigin, V.: Volume and heat transports to the Arctic Ocean via the Norwegian and Barents Seas, *Arctic–subarctic ocean fluxes: Defining the role of the northern seas in climate*, pp. 45–64,
355 2008.
- Skagseth, Ø., Eldevik, T., Årthun, M., Asbjørnsen, H., Lien, V. S., and Smedsrud, L. H.: Reduced efficiency of the Barents Sea cooling machine, *Nature Climate Change*, 10, 661–666, 2020.
- Smedsrud, L. H., Ingvaldsen, R., Nilsen, J. Ø., and Skagseth, Ø.: Heat in the Barents Sea: Transport, storage, and surface fluxes, *Ocean science*, 6, 219–234, 2010.
- 360 Smedsrud, L. H., Esau, I., Ingvaldsen, R. B., Eldevik, T., Haugan, P. M., Li, C., Lien, V. S., Olsen, A., Omar, A. M., Otterå, O. H., Risebrobakken, B., Sandø, A. B., Semenov, V. A., and Sorokina, S. A.: THE ROLE OF THE BARENTS SEA IN THE ARCTIC CLIMATE SYSTEM, *Reviews of Geophysics*, 51, 415–449, <https://doi.org/https://doi.org/10.1002/rog.20017>, 2013.

- 365 Smedsrud, L. H., Muilwijk, M., Brakstad, A., Madonna, E., Lauvset, S. K., Spensberger, C., Born, A., Eldevik, T., Drange, H., Jeansson, E.,
et al.: Nordic Seas heat loss, Atlantic inflow, and Arctic sea ice cover over the last century, *Reviews of Geophysics*, 60, e2020RG000725,
2022.
- Uotila, P., Goosse, H., Haines, K., Chevallier, M., Barthélemy, A., Bricaud, C., Carton, J., Fučkar, N., Garric, G., Iovino, D., et al.: An
assessment of ten ocean reanalyses in the polar regions, *Climate Dynamics*, 52, 1613–1650, 2019.
- 370 Wang, Q., Wang, X., Wekerle, C., Danilov, S., Jung, T., Koldunov, N., Lind, S., Sein, D., Shu, Q., and Sidorenko, D.: Ocean heat transport
into the Barents Sea: Distinct controls on the upward trend and interannual variability, *Geophysical Research Letters*, 46, 13 180–13 190,
2019.
- Woodgate, R. A., Aagaard, K., Muench, R. D., Gunn, J., Björk, G., Rudels, B., Roach, A., and Schauer, U.: The Arctic Ocean boundary
current along the Eurasian slope and the adjacent Lomonosov Ridge: Water mass properties, transports and transformations from moored
instruments, *Deep Sea Research Part I: Oceanographic Research Papers*, 48, 1757–1792, 2001.
- 375 Årthun, M., Brakstad, A., Dörr, J., Johnson, H. L., Mans, C., Semper, S., and Våge, K.: Atlantification drives recent strengthening of the
Arctic overturning circulation, *Science Advances*, 11, eadu1794, <https://doi.org/10.1126/sciadv.adu1794>, 2025.