

Roles of planktonic metabolism in CO₂ fluxes along the river-estuary continuum in a rapidly uplifting catchment of eastern Taiwan

Jhen-Nien Chen^{1,2}, Pei-En Chen³, Yu-Shiang Yen¹, Tzu-Hsuan Tu⁴, Lu-Yu Wang¹, Wan-Yin Lien¹, Yueh-Ting Lin^{2,3}, and Pei-Ling Wang^{3,5}

¹Department of Geosciences, National Taiwan University, Taipei 106, Taiwan

²Taiwan Ocean Research Institute, National Institutes of Applied Research, Taipei, Taiwan

³Institute of Oceanography, National Taiwan University, Taipei 106, Taiwan

⁴Department of Oceanography, National Sun Yat-sen University, Kaohsiung, Taiwan

⁵Science and Technology Research Institute for Decarbonization, National Taiwan University, Taipei 106, Taiwan

Correspondence to: Pei-Ling Wang (plwang@ntu.edu.tw)

Abstract. The contribution of river metabolism to carbon cycling remains poorly examined in catchments susceptible to the modulation of active tectonics. In these high-energy mountainous systems, chronically high turbidity and frequent sediment mobility create an unstable fluid regime that may impact planktonic metabolism and its contribution to aquatic carbon transformation. Using the Beinan River in southeastern Taiwan as a model system, this study aimed to quantify the potential rates of planktonic autotrophy and heterotrophy, and to identify the potential community members involved in these processes. To address this, river water samples were collected across wet and dry seasons for incubations with ¹³C-labeled dissolved inorganic carbon and amino acids. Our results revealed that metabolic rates were consistently higher in the wet season than in the dry season, with heterotrophic catabolic rates exceeding autotrophic carbon fixation by one to two orders of magnitude, indicating strong net heterotrophy. The obtained rates were further scaled up, resulting in a catchment-scale CO₂ evasion of $7.89 \times 10^7 \text{ mol yr}^{-1}$, which constitutes about 3.8% of the CO₂ flux derived from pyrite-induced carbonate weathering, oxidation of petrogenic carbon, and the river-air exchange. The microbial community structures varied seasonally at upstream sites, with more abundant lithoautotrophic sulfur- and ammonia/nitrite-oxidizing taxa in the wet season, as opposed to more abundant phototrophs or heterotrophs in the dry season. This study highlights that even in oligotrophic, physically dynamic mountainous systems prone to rapid uplift and erosion, planktonic metabolisms constitute a measurable and vital component of the riverine carbon cycle.

1. Introduction

Inland freshwater systems release CO₂ to the atmosphere at a flux (0.65–2.1 Pg C yr⁻¹) comparable with that of ocean CO₂ uptake (2.2 Pg C yr⁻¹) (Horgby et al., 2019; Lauerwald et al., 2015; Takahashi et al., 2009). The magnitude is enormous given their limited global land-surface coverage (0.3–0.56%) (Allen and Pavelsky, 2018; Aufdenkampe et al., 2011). Among a variety of inland sub-systems, mountainous catchments, which comprise 27% of the Earth's land surface (Ives et al., 1997) and drain more than 30% of the global runoff, receive far less attention when compared with low-altitude catchments in temperate regions (Bandyopadhyay et al., 1997; Meybeck et al., 2001; Viviroli and Weingartner, 2004). While the accessibility to sampling or measuring in such high-relief catchments is often limited, a few previous studies have projected that CO₂ evasion declines substantially by orders of magnitude as a function of distance from the headwater (Raymond et al., 2013; Schelker et al., 2016; Chiriboga and Borges, 2023). Such a change in flux is intrinsically tied to the summed effect of *p*CO₂ (CO₂ partial pressure) and the exchange coefficient. Even fewer studies have further demonstrated that mountainous rivers draining catchments composed of crystalline rocks are characterized by *p*CO₂ slightly exceeding those in equilibrium with the atmospheric CO₂ (Horgby et al., 2019). Therefore, the enhanced CO₂

evasion in the upstream region could be largely attributed to the steep topography, where turbulent flow and a rough riverbed enable rapid CO₂ exchange between air and water. As the river slope decreases toward the downstream region, the degree of topographically controlled CO₂ evasion gradually diminishes. The pattern highlights the role but also points out the deficiency in data coverage for mountainous catchments in configuring the global carbon inventory.

A number of factors have been suggested to contribute to the overall pattern of CO₂ evasion along the river corridor. Among these factors, groundwater input and *in situ* microbial processes are often considered the most important source mechanisms (Brunke and Gonser 1997; Cole et al., 2007; Marx et al., 2017). Measurements mostly using dissolved oxygen sensors have shown that aquatic metabolisms contributed to a sum of 28% of the CO₂ evasion for streams and rivers across the US continent (Butman and Raymond 2011; Hotchkiss et al., 2015). A few other studies have further demonstrated that benthic and hyporheic compartments contribute significantly to the whole stream metabolism in many upland systems, particularly when water clarity permits light to reach coarse streambed sediments (Hall et al., 2015; Battin et al., 2023; Bernhardt et al., 2022; Roche et al., 2025). For mountainous catchments, most studies intuitively assume that groundwater input would dominate over *in situ* aquatic microbial processes in a way that the catchments are composed of exposed bedrock rather than thick soils accumulated over a prolonged period of weathering and that the residence time of substrates is perhaps too short to be made available for microbial degradation (Marx et al., 2017; Duvert et al., 2018). Therefore, the quantities of the products derived from soil degradation (presumably attributed to dissolved organic carbon) would be scarce, thereby rendering the limited availability of substrates for microbial respiration. While the terrestrially derived fluxes (equivalent to lateral transport of groundwater) may dominate the CO₂ evasion in small streams, the percentage of aquatic metabolic flux to the overall evasion is postulated to increase with stream size (Hotchkiss et al., 2015). The exact fraction of metabolic flux to the overall evasion in mountainous rivers remains largely unconstrained.

The observations of diurnal variation in dissolved oxygen represent the net outcome combining primary production and ecosystem respiration. Therefore, the contribution of individual categories of metabolism and further extrapolation to the CO₂ variation could only be deconvoluted with the computational scheme inherited with assumptions (Appling et al., 2018; Saccardi and Winnick, 2021). Furthermore, whether observations based on the cratonic continent or large river systems could be applicable to other catchments around the world remains less constrained. In particular, small island catchments susceptible to active tectonics are constantly impacted by torrential precipitation associated with monsoons and typhoons, mass wasting processes, and structurally controlled transport of groundwater (McMillan and Srinivasan 2015). The event-based hydrological and geological impacts would have led to a pattern of nutrient and substrate availability (e.g., turbid river water during high water) and transport of microbial populations fundamentally distinct from the major river systems. As frequent sediment mobility, bed scouring, and episodic turbidity pulses may impact planktonic processes across the full range of hydrological conditions, dissecting the relative contributions of individual planktonic metabolic pathways (phototrophy, chemoautotrophy, and heterotrophy) provides a necessary first step toward resolving the role of the planktonic compartment within the broader river carbon-cycling network.

In this study, we aimed to quantify planktonic metabolic rates and characterize microbial community dynamics along the Beinan River. As a representative of high-standing island catchments, the Beinan River is characterized by high sediment exports and weathering rates that rank among the greatest in Taiwan (Dadson et al., 2003) and exceed the global average by orders of magnitude (Hilton and West, 2020). To quantify metabolic activity, we performed short-term incubations supplemented with ¹³C-labeled dissolved inorganic carbon (DIC) and amino acids (leucine and glycine) to constrain assimilation and respiration rates of different metabolic pathways, and complemented these with 16S rRNA gene amplicon sequencing to characterize microbial community composition. We tested three interrelated hypotheses: (i) the

seasonal transition to the wet season (characterized by higher temperatures, elevated nutrient availability, and increased particulate organic matter loading) would broadly shift planktonic metabolic activities; (ii) these seasonal hydrological changes would be reflected in shifts in microbial community composition; and (iii) in this physically unstable, chronically turbid catchment, frequent sediment mobility would drive temporal variability in planktonic metabolic rates and their relative contribution to the riverine CO₂ evasion. By testing these hypotheses with direct ¹³C-based rate measurements and community composition data, this study seeks to provide a mechanistic framework for understanding how physical disturbance, seasonal hydrology, and microbial structure interact to shape carbon cycling in tectonically active mountainous rivers.

2. Materials and methods

2.1 Field background

Taiwan was formed through dual subduction and interactions between the Eurasian and Philippine Sea Plates (Teng 1990). The northwestward impingement of the Luzon arc on the Eurasian continent since 5 Ma has created steep topographic changes with the unroofing of metamorphic complexes in the backbone range, from which major rivers originate (Chen et al., 2019). Across the island (particularly central and southern Taiwan), rapid uplift (up to several centimeters per year), river incision, event-based torrential precipitation, and frequent earthquakes have generated numerous landslides, with sediments transported along with flow or temporarily deposited in the riverbank/bed and subsequently remobilized during the high-water period (Dadson et al., 2003; Kao and Milliman, 2008). The Beinan River, being one of the largest rivers in Taiwan in terms of discharge and suspended sediment load (Dadson et al., 2003; Kao and Milliman, 2008; annual report of the Water Resource Agency, MOEA), flows eastward across the central mountain and drains southward along the Longitudinal Valley prior to reaching the ocean. The catchment covers an area of ~1600 km² and is primarily composed of schist and slate, with minor amounts of marble and meta-sandstone (Stanley et al., 1981). Soil development is limited and hindered by high erosion rates. Therefore, detrital materials that are susceptible to pulverization and abrasion are primarily composed of rock fragments. Land cover in the catchment is dominated by natural and semi-natural vegetation. Evergreen broadleaf and mixed montane forests cover the majority of the upper and middle catchment, with exposed bedrock, landslide scars, and riverbed gravel deposits accounting for a non-trivial fraction consistent with the high denudation regime of the region. Human activity and land use are limited and concentrated in the lower Longitudinal Valley and the coastal alluvial plain near the estuary. Population density in the catchment is lower than in western Taiwan, and no large impoundment exists on the main stem. A land-use and land-cover overview of the lower catchment is shown in Fig. 1b.

Sampling sites were selected to represent individual sub-basins of the Beinan catchment and to span the elevation gradient and landscape transition from metamorphic headwater tributaries to the estuary (Figs 1a and S1a). Among these sites, MLL01 is located in an upper tributary in the forested headwaters and is characterized by a steep, boulder- and cobble-dominated channel within a valley. CWL01 is on the upper main stem of the Beinan River, and DL is on an adjacent mid-catchment tributary that merges with the main stem near CWL01; both sites are characterized by coarse-grained beds and extensive exposure of gravel/sand bars during low flow. LY04 is on a lower tributary, where the channel broadens into a wider, cobble-bed reach. BNE is on the Beinan main stem at the river mouth. Field photographs of all five sites are provided in Fig. S2. *In situ* surface flow velocities measured during seven sampling campaigns ranged from 0.3 to 2.5 m s⁻¹ (Table S1), reflecting the high-energy characteristics of the catchment. At BNE, *in situ* salinity remained below 0.6 ppt in every campaign, and dissolved Cl⁻ and Na⁺ concentration (0.22–0.28 mM and 0.65–0.77 mM, respectively; Table S1) were several orders of magnitude lower than seawater values (~550 mM Cl⁻ and ~470 mM Na⁺) and comparable to

those at upstream freshwater sites, indicating that seawater intrusion was negligible during the sampled conditions and that BNE should be interpreted as the freshwater end of the river-estuary continuum rather than a fully mixed estuary.

Previous studies based on carbon isotopic compositions, radiocarbon contents, and Raman spectroscopic characteristics have inferred limited alteration of the source rocks and low contributions of soil to organic matter in suspended particulates (Hilton et al., 2008; Lien et al., 2025). Although substantial contributions of soil carbon have been reported (Lin et al., 2020), the analyzed samples were collected from the bank, which represents a mixture of sediments deposited over time. The isotopic and spectroscopic characteristics, combined with the rapid transit of river water, suggest that, during most periods, suspended particulates are primarily sourced from the lithogenic compartment and remain relatively recalcitrant to biological utilization. Monsoon combined with typhoons from summer to autumn contributes the majority of precipitation (generally >3000 mm) and runoff in the upstream region. For this study, the wet season was defined as the period between June and November 2020, during which precipitation accumulated substantially (Fig. S1). Notably, samples collected in May were referred to as the dry season, considering that the field campaign in May of 2020 was conducted prior to the significant increase in precipitation in late May (Fig. S1). It is also worth noting that the sampling period represents the driest year (with an annual precipitation of 1856 mm) over the last 20 years.

2.2 Sampling and field experiments

Seven field campaigns were conducted in March, May, July, August (twice), October 2020, and January 2021 (Table S1). This frequency was chosen to provide a systematic baseline that spans the full hydrologic cycle, encompassing the distinct dry (January – May) and wet (July – August) seasonal characteristics of southeastern Taiwan (Fig. S1). The intervals also enable consistent comparisons of metabolic rates across seasonal transitions, providing a framework to assess the long-term metabolic stability of these dynamic turbid systems. While *in situ* environmental parameters were characterized in all campaigns, metabolic rates were measured in March and May 2020 (for phototrophy and autotrophy only) and in August 2020 and January 2021 (for the full suite of tested metabolisms). At each site, *in situ* physicochemical characteristics (temperature, pH, conductivity, and river flow rate) were measured using the portable probes and meters (Hanna Instruments, HI-98130, RI, USA; Pro II SVR, Stalker, TX, USA). Samples were collected in different forms with different preservatives and preparations (Wang et al., 2024). In brief, river water was filtered using a sterilized 0.22 µm pore-size polyethersulfone (PES) membrane (Supor®) and collected for analyses of dissolved inorganic carbon (DIC), major cations and anions, dissolved organic carbon (DOC), and dissolved inorganic nutrients. To minimize carbon and trace-element contamination, all syringes and high-density polyethylene (HDPE) bottles for DOC and cation samples were pre-washed through a rigorous cleaning protocol consisting of an acid wash with concentrated nitric acid followed by triple-rinsing with both Milli-Q water and filtered river water before final collection. Cation samples were acidified with 1/10 volume of 20% nitric acid (Baseline grade, Seastar, BC, Canada). Samples were stored at room temperature (for DIC and inorganic solutes) or –20 °C (for DOC and nutrients). Particulate samples were collected on 0.3 µm pore-size glass fiber membranes pre-combusted at 500 °C for 6 hours to remove any potential organic contaminants. The total volume of filtrate for individual particulate samples was 1 L. The collected membranes were frozen at either –20 °C for analyses of chlorophyll *a* (Chl *a*) and abundances and isotopic compositions of particulate organic carbon (POC), or –80 °C for molecular analyses.





Figure 1: (a) Topographic map of the Beinan River catchment (eastern Taiwan) with sampling sites marked in red. (b) Land use and land cover of the lower Beinan catchment, adopted from the National Land Use Investigation dataset (National Land Surveying and Mapping Center, Ministry of the Interior, Taiwan).

For metabolic rate measurements, river water was collected into a water bag pre-washed with river water several times, and supplemented with ^{13}C -labeled bicarbonate or individual amino acids ($[1-^{13}\text{C}]$ glycine or $[1-^{13}\text{C}]$ L-leucine; ^{13}C labeled at the C-1 [carboxyl] position) at a final concentration of $50\ \mu\text{M}$ to constrain the rates of autotrophy/phototrophy (using

DIC) or heterotrophy/respiration (using amino acids). **DOC-based heterotrophy was given particular focus because DOC represents the substrate pool directly available to planktonic microorganisms, and particulate organic matter is considered recalcitrant for further degradation during river transit (Hilton et al., 2008; Lien et al., 2025).** The supplemented substrates with ^{13}C purity >99% were mixed with deionized water to prepare a 1 mM stock. The amino acid stock was acidified by titrating with HCl to pH <4 to remove the potential DIC associated with the dissolution of the amino acid powder. The stock of either DIC or amino acid was filter-sterilized and stored at 4 °C until further use in the field.

Six liters of river water were collected for each substrate type. All the incubation experiments were conducted in the field accommodation. The supplemented river water was dispensed into 1 L HDPE bottles, filled to capacity to eliminate headspace, and incubated for 4 to 6 hours. For ^{13}C -DIC autotrophy, three transparent (light) bottles and one opaque (dark) bottle were used per site per campaign. The light bottles were incubated under constant illumination (~18,000 lux, monitored by HOBO, Pendant MX) and placed in a flowing tap-water bath, by which heat from the lighting unit was dissipated to keep the bottles at near-ambient temperatures (ranging from 19.3 to 27.2°C). The dark bottle was incubated alongside. Heterotrophic incubations with ^{13}C -amino acids (glycine and leucine, three replicate bottles each) were conducted in opaque bottles at ambient temperature. The incubated samples were filtered through 0.3 µm pore-size glass fiber membranes to collect particulates comprising mineral/rock fragments and microorganisms. The membranes were rinsed with 2 N HCl immediately after filtration to remove carbonate minerals, then washed with sterilized saline solution. The collected membrane was stored at -80 °C for further isotopic and molecular analyses. Part of the filtrates from the amino acid incubations was preserved at room temperature for isotopic analyses of DIC. A conceptual diagram describing the experimental setup is shown in Fig. 2.




Figure 2: Conceptual diagram for the incubation setup in fields and following laboratory measurements.

2.3 Geochemical analyses

Anions were measured using an ICS-3000 ion chromatography (Thermo Fisher Scientific, USA). Major cations were analyzed using an inductively coupled plasma linked with a quadrupole mass spectrometry (ICP-MS; 7800, Agilent, Taiwan). Synthetic and river water standards (SLRS-5 from the National Research Council of Canada) were used to calibrate a range of cation concentrations. Concentrations of orthophosphate and ammonium were measured using molybdenum blue generated from the reductive reaction between phosphate complex and ascorbic acid (APHA Method 4500-P, 2005), and indophenol generated from the reaction between ammonium and thymol (APHA Method 4500-NH₃, 2005), respectively. The intensity of the color complex was quantified using a spectrophotometer (V-500, Jasco, MD, USA). Nitrite/nitrate concentrations were quantified using a commercial fluorometric assay (Cayman, MI, USA), which reacts with nitrite (original form or reduced from nitrate) with 2,3-diaminonaphthalene for the generation of 1(H)-

naphthotriazole (Miles et al., 1995). Chlorophyll *a* (Chl *a*) was extracted from membranes with acetone under dark conditions and quantified using a fluorometer (10-AU, Turner Designs, CA, USA). DOC concentration was measured using a Shimadzu TOC-L CPH/CPN system with high-temperature catalytic oxidation pre-calibrated with known concentrations of KHP (potassium hydrogen phthalate). The analytical uncertainty for these methods is less than 2%. Concentrations and carbon isotopic compositions of DIC for river water and incubated samples were analyzed by acidifying samples with concentrated phosphoric acid (85%) and purging the produced CO₂ into a total carbon analyzer (Aurora 1030, OI Analytical, USA) and an isotope ratio mass spectrometer (IRMS; MAT253, Thermo Fisher Scientific, USA) connected with a GC-IsoLink (Thermo Fisher Scientific, USA), respectively. Concentrations and carbon isotopic compositions of particulate organic carbon (POC) for river water and incubated samples were analyzed on freeze-dried, acid-rinsed membranes using an elemental analyzer (EA; MICRO cube, Elementar, Germany) and an IRMS (Delta V) coupled to a Flash EA (Thermo Fisher Scientific, USA), respectively. Carbon isotopic compositions were expressed as the delta notation:

$$\delta^{13}\text{C} = \left[\frac{R_{\text{sample}}}{R_{\text{standard}}} - 1 \right] \times 1000 \text{ ‰} \quad (1)$$

where *R* is the ratio of ¹³C to ¹²C, and the standard is Vienna Pee Dee Belemnite (VPDB). The uncertainties for DIC and POC concentrations and δ¹³C values were less than 2% and 0.2‰, respectively.

Metabolic rates were calculated using the following equations:

$$\rho_{\text{DIC_uptake}} = \frac{\Delta F^{13}\text{C}_{\text{POC}}}{F^{13}\text{C}_{\text{DIC}}} \times [\text{POC}] \times \frac{1}{t} \quad (2)$$

$$\rho_{\text{AA_uptake}} = \frac{\Delta F^{13}\text{C}_{\text{POC}}}{F^{13}\text{C}_{\text{DOC}}} \times [\text{POC}] \times \frac{1}{t} \quad (3)$$

$$\rho_{\text{AA_catabolic}} = \frac{\Delta F^{13}\text{C}_{\text{DIC}}}{F^{13}\text{C}_{\text{DOC}}} \times [\text{DIC}] \times \frac{1}{t} \quad (4)$$

where ρ_{DIC_uptake} is the rate of DIC assimilated into biomass (Eq.2), ρ_{AA_uptake} is the rate of amino acid assimilated into biomass (Eq. 3), and ρ_{AA_catabolic} is the rate of amino acid respired to DIC (Eq.4). *F*¹³C represents the fraction of ¹³C in POC, DIC, or DOC; Δ*F*¹³ is the change in *F*¹³C of POC or DIC during the incubation (post-incubation value minus the T0 reference value); [POC] and [DIC] are the measured POC and DIC concentrations, respectively; and *t* is the duration of incubation. The isotopic fractionation associated with the target metabolic reaction is assumed to be negligible. In this regard, the *F*¹³C_{DIC} in eq. 2 and *F*¹³C_{DOC} in Eqs. 3 and 4 represent the sum of the ¹³C fractions of added substrate and original forms (DIC or DOC). The δ¹³C values of *in situ* DOC were not available and, therefore, were assumed to be –27‰, considering that the isotopic compositions of soil organic matter are in a similar range (Wang et al., 2024; Lien et al., 2025). The rate for the incubation with DIC under light is considered to represent the summed rate of phototrophy and lithoautotrophy, and could be greater than the rate for the dark incubation. The difference is generally attributed to the phototrophic rate, assuming that the true lithoautotrophy is not affected by light illumination. For incubations with labeled DIC, the chosen concentration constituted a minor fraction of the environmental DIC pool (<5% of the environmental DIC; Wang et al., 2024). For incubations with labeled amino acid, the 50 μM addition represents a potential enrichment relative to ambient dissolved free amino acid concentration in oligotrophic systems (typically < 0.1 μM; Jørgensen and Søndergaard 1984; Raymond and Spencer 2015) and is comparable to the total ambient DOC concentration (20–80 μM). In this context, the measured autotrophic and phototrophic rates based on DIC labeling are considered representative of *in situ* rates, whereas the measured amino acid assimilation and respiration rates represent potential heterotrophic activity.

Seasonal differences in environmental parameters were assessed using linear mixed-effects models (R packages lme4 and lmerTest), with season as a fixed effect and site as a random intercept to account for repeated sampling. Strongly skewed variables (TSM, POC, Chl *a*, NH₄⁺) were log₁₀-transformed prior to analysis to satisfy the normality assumption on residuals. Among-site differences in metabolic rates were assessed using Kruskal-Wallis tests followed by Dunn's post-hoc tests. Within-site seasonal variations were evaluated using Kruskal-Wallis tests (multiple time points) or Mann-Whitney U tests (pairwise comparisons). Relationships between rates and environmental variables were examined using Pearson correlations. All the statistical analyses were performed using R version 4.4.1.

2.4 Molecular analyses of community compositions

Crude DNA was extracted and purified using the DNeasy PowerSoil kit (Qiagen, USA) following the manufacturer's protocol, and stored at -80 °C for further analyses. Polymerase chain reaction (PCR) amplification was conducted on all extracts using the primers 515F (GTG CCA GCM GCC GCG GTA A) and 806R (CCC GTC AAT TCM TTT RAG T), which target the V4 region of the 16S rRNA gene (Kozich et al., 2013). The amplicons were purified, pooled, barcoded, and sequenced using the Illumina MiSeq platform (Illumina, USA). The detailed methodology and reagents employed for PCR have been described in Tu et al. (2017, 2022). Due to the insufficient quantity of DNA (<50 ng in total), extracts from incubated samples were not further processed for stable isotope probing analysis, which targets the population members actively assimilating the fed ¹³C-labeled substrate.

Quantitative PCR (qPCR) was performed to determine the copy numbers of bacterial and archaeal 16S rRNA genes and genes encoding ribulose-1,5-bisphosphate carboxylase/oxygenase form I (*cbbL*) (Alfreider et al., 2003) using the QuantStudio™ 3 Real-Time PCR System (Applied Biosystems, ABI, USA). The standard preparations and PCR conditions were as described in Tu et al. (2017) and Wang et al. (2014). The primers and annealing temperatures for the different genes are summarized in Table 1. The copy number was calculated using threshold cycles calibrated against standards with known concentrations of gene amplicons, assuming 650 g mol⁻¹ for one base pair of DNA.

Raw sequences of 16S rRNA gene amplicons were processed using Mothur ver. 1.48.0 (Schloss et al., 2009) and QIIME2 (Bolyen et al., 2019). The individual reads were quality-filtered, denoised, merged, and dechimerized to generate Amplicon Sequence Variants (ASVs). Taxonomic classification was performed using the SILVA database (version 138) (Quast et al., 2012). Sequence reads were rarefied to a depth of 100 rounds of random resampling (without replacement) to account for differences in sequencing depth. Alpha diversity indices, including observed ASVs and Chao1 for richness (Chao, 1987) and the Shannon index for evenness (Shannon and Weaver, 1949), were calculated. Differences in alpha diversity among sites and between seasons (wet and dry) were tested by the Kruskal-Wallis test. For beta diversity, the entire ASV table was normalized, and Bray-Curtis dissimilarities were computed. Differences among sites and seasons were tested by permutational multivariate analysis of variance (PERMANOVA; adonis2 in package vegan, 999 permutations). Relationships between microbial community composition and environmental variables were examined by canonical correspondence analysis (CCA) (González et al., 2008), with the significance of each environmental vector assessed by envfit (vegan, 9999 permutations; *p* < 0.05). All statistical analyses were performed using R version 4.4.1 with the vegan, ggplot2, and phyloseq packages. The obtained sequences were deposited in GenBank (BioProject ID: PRJNA1270665).

271

Table 1: Primer information in this study.

Primer name	Sequence (5'-3')	Target group/gene	Annealing temp (°C)	Reference
B27F	AGAGTTTGATCMTGGCTCAG	Bacteria/16S rRNA	55	Lane (1991)
Eub338R	GCTGCCTCCCGTAGGAGT	Bacteria/16S rRNA	55	Amann et al. (1990)
U1492R	GGTTACCTTGTTACGACTT	Bacteria/16S rRNA	55	Lane (1991)
A8F	TCYGGTTGATCCTGCC	Archaea/16S rRNA	55	Huber et al. (2002)
A806F	ATTAGATACCCSBGTAGTCC	Archaea/16S rRNA	55	Takai and Horikoshi (2000)
A958R	YCCGGCGTTGAMTCCAATT	Archaea/16S rRNA	55	Hinrichs et al. (1999)
U1513R	ACGGHTACCTTGTTACGACTT	Archaea/16S rRNA	55	Huber et al. (2002)
cbbLR1F	AAGGAYGCGAGAACATC	autotroph/ <i>cbbL</i>	58	Selesi et al. (2007)
cbbLR1intR	TGCAGSATCATGTCTTT	autotroph/ <i>cbbL</i>	58	Selesi et al. (2007)

272

2.5 Computation for catchment-scale CO₂ fluxes

To scale planktonic metabolic rates to the catchment level, the five sampling sites were treated as the outlets of five sub-catchments, with boundaries and areas adopted from the Water Resources Agency (WRA) hydrological atlas of the Beinan basin (Table S6). Measured rates were multiplied by the average water depth of each investigated reach (50 cm for tributaries, 70 cm for main stem) to obtain areal fluxes. These depth estimates reflect typical conditions observed during field campaigns, excluding flash flood periods. DIC uptake rates under light and dark conditions were assumed to represent daytime and nighttime rates, respectively, with total daily DIC uptake calculated as the sum of equal contributions from each. Using leucine incorporation rates as a conservative estimate of heterotrophic activity, net CO₂ production for each sub-catchment was calculated by multiplying the per-site area flux with the sub-catchment river surface area, taken as 0.47% of its sub-catchment area (Raymond et al., 2013). The total catchment yield was obtained as the sum of the five sub-catchment yields.

To compare the metabolic fluxes with the CO₂ evasion, CO₂ exchange fluxes between river water and the atmosphere were calculated for each sub-catchment using the following equation:

$$\text{Flux} = k_{\text{CO}_2} \times \Delta (C_{\text{measured}} - C_{\text{eq}}) \quad (5)$$

where k_{CO_2} is the gas exchange velocity, and C represents the partial pressure of CO₂ either from measurement or in equilibrium with the atmospheric CO₂ concentration (assumed to be 410 ppm). C_{measured} was derived from the DIC concentration, pH, salinity, temperature, and total atmospheric pressure measured in the same field campaign using co2sys (Lewis and Wallace, 1998). k_{CO_2} was computed using the relationships reported by Ulseth et al. (2019), where k_{600} values vary positively with eD (energy dissipation rate and is equal to the product of g [gravitational acceleration], v [flow velocity], and s [slope of the reach]). Depending on whether the gas exchange is controlled by turbulence or bubbles, the relationships between k_{600} and eD are different (Ulseth et al., 2019):

$$\ln k_{600} = 6.43 + 1.18 * \ln eD \quad (\text{for } eD > 0.02) \quad (6)$$

$$\ln k_{600} = 3.10 + 0.35 * \ln eD \quad (\text{for } eD < 0.02) \quad (7)$$

The flow velocity was adopted from the measured ones obtained in the field, which represented the maximum velocity on the river surface. Because friction would be substantial near the channel bed, the measured surface velocity would be higher than the average channel velocity integrated over the channel cross section. However, considering the water depth is generally small, the measured flow velocity may not substantially deviate from the true one. Nevertheless, the calculated eD values would represent an overestimate. Slope was calculated for the investigated reach (~1000 m in length) using the DEM (digital elevation model) available to the public (data.gov.tw). All the parameters used for calculation were summarized in Table S7. The obtained k_{600} value was further converted to k_{CO_2} using the following equation (Wanninkhof, 2014):

$$k_{\text{CO}_2} = k_{600} / [(600 / Sc_{\text{CO}_2})]^{(-n)} \quad (8)$$

where Sc is the Schmidt number of CO₂ at the *in situ* temperature and n is assumed to be 0.5 for turbulent water (Jähne et al., 1987).

The catchment-scale CO₂ evasion for target metabolisms or the air-river exchange was susceptible to the uncertainty in both the assumed river surface area and the measured metabolic or flux rates at each site. River surface area fluctuates substantially between events or seasons, and the surface area of low-order reaches may be partly shaded by the trees or

310 too narrow (< 1 m) to be resolved by satellite imaging. It becomes technically challenging to fully recover the variations
311 in surface area across the investigated tributaries at different temporal and spatial scales. Therefore, we arbitrarily assigned
312 a conservative uncertainty of $\pm 50\%$ to river surface area. For each site, this area uncertainty was combined with the
313 measured variability in the corresponding rate or flux (Tables S6 and S7). Site-level uncertainties were then similarly
314 propagated to the catchment scale.

315 3. Results

316 Physicochemical characteristics for each field campaign were categorized into the wet and dry seasons in accordance with
317 the precipitation records (Tables 2 and S1; Fig. S1). Water temperatures varied between 12 and 30 °C, with consistently
318 higher values in the wet than dry seasons (mean wet-dry difference = $+5.9$ °C, $p < 0.001$; Table S2). pH was lower in the
319 wet than dry seasons ($p = 0.003$). TSM contents ranged from several to hundreds of mg L^{-1} and were higher in the wet
320 than dry seasons, with an average ~ 11 -fold increase in the wet season ($p = 0.002$). The $\delta^{13}\text{C}_{\text{POC}}$ values ranged from -28
321 to -20‰ and were more negative in the dry than in the wet season ($p < 0.001$). Ammonium concentrations ranged between
322 0.3 and 16.5 μM and were higher in the wet than in the dry season ($p < 0.001$). In contrast, nitrate/nitrite and phosphate
323 concentrations ranged from 1.7 to 48.3 μM and 0.01 to 0.31 μM , respectively, and neither of them differed significantly
324 between seasons ($p = 0.690$ and $p = 0.504$, respectively; Table S2). Chl *a* concentrations ranged from below the detection
325 limit to 2.1 mg m^{-3} and were higher in the dry season than in the wet season (except for DL; $p = 0.007$). DIC concentrations
326 and their $\delta^{13}\text{C}$ values varied between 1.26 and 2.88 mM and between -6.8 and $+0.6\text{‰}$, respectively, with no significant
327 difference between seasons ($p = 0.852$ and $p = 0.421$, respectively). Sulfate concentrations ranged between 1.03 and 2.06
328 mM for most sites (except for MLL01 at 2.92 – 3.59 mM) and were higher in the wet than dry seasons ($p = 0.004$). Sodium
329 and calcium concentrations were lower in the wet than dry seasons ($p = 0.005$ and $p < 0.001$, respectively).

330

Table 2: Physicochemical characteristics among sites and seasons.

Tributary (altitude)	n (sample numb. for wet or dry season)	T ¹ (°C)	pH	TSM ¹ (mg L ⁻¹)	POC (mg m ⁻³)	δ ¹³ C- POC (‰)	DOC (μM)	DIC (mM)	δ ¹³ C-DIC (‰)	Chl a (mg m ⁻³)	TIN ² (μM)	NH ₄ ⁺ (μM)	PO ₄ ²⁻ (μM)	SO ₄ ²⁻ (mM)
		n= 4, 3	n= 4, 3	n= 2, 3	n= 2, 3	n= 2, 3	n= 4, 3	n= 3, 3	n= 4, 3	n= 3, 2	n= 4, 3	n= 3, 3	n= 4, 3	n= 3, 3
MLL01 (699 m)	Wet seasons	23.4 ± 2.8	8.4 ± 0.4	157 ± 113	2432 ± 1868	-24.3 ± 0.3	30.4 ± 12.9	1.92 ± 0.05	-1.8 ± 2.1	0.07 ± 0.06	10.0 ± 2.5	4.6 ± 4.3	0.04 ± 0.05	3.36 ± 0.21
	Dry seasons	17.3 ± 4.7	8.1 ± 0.5	11 ± 16	152 ± 32	-27.4 ± 0.3	20.6 ± 1.7	1.91 ± 0.34	-2.7 ± 1.3	0.44 ± 0.08	6.8 ± 6.2	0.6 ± 0.3	0.05 ± 0.04	3.02 ± 0.13
	Total	20.8 ± 4.7	8.2 ± 0.4	70 ± 98	1064 ± 1560	-26.1 ± 1.7	26.2 ± 10.5	1.92 ± 0.21	-2.2 ± 1.8	0.22 ± 0.21	8.7 ± 4.3	2.6 ± 3.5	0.05 ± 0.04	3.19 ± 0.24
CWL01 (375 m)	Wet seasons	26.1 ± 2.4	7.8 ± 0.1	27 ± 32	1477 ± 1381	-23.4 ± 2.7	31.2 ± 16.0	1.81 ± 0.48	-2.8 ± 1.3	0.58 ± 0.38	6.0 ± 0.8	4.9 ± 1.9	0.06 ± 0.04	1.53 ± 0.04
	Dry seasons	20.2 ± 6.8	8.5 ± 0.3	12 ± 18	236 ± 129	-26.8 ± 1.0	23.9 ± 3.3	2.18 ± 0.37	-2.1 ± 1.9	1.33 ± 1.02	5.2 ± 3.9	2.5 ± 0.6	0.04 ± 0.03	1.61 ± 0.04
	Total	23.1 ± 5.6	8.2 ± 0.4	18 ± 22	732 ± 973	-25.4 ± 2.4	28.1 ± 12.1	1.99 ± 0.43	-2.5 ± 1.5	0.83 ± 0.67	5.7 ± 2.3	3.7 ± 1.8	0.05 ± 0.03	1.57 ± 0.06
DL (382 m)	Wet seasons	26.6 ± 2.2	7.9 ± 0.6	4 ± 3	291 ± 24	-25.9 ± 0.2	35.3 ± 19.9	2.16 ± 0.05	-1.9 ± 0.9	1.14 ± 0.54	16.0 ± 4.2	6.4 ± 6.5	0.14 ± 0.02	1.08 ± 0.04
	Dry seasons	21.9 ± 7.3	8.4 ± 0.2	11 ± 18	160 ± 63	-26.4 ± 0.4	20.2 ± 3.4	2.43 ± 0.41	-1.9 ± 0.6	0.64 ± 0.24	18.0 ± 3.0	1.5 ± 1.3	0.15 ± 0.06	1.13 ± 0.12
	Total	24.6 ± 5.2	8.2 ± 0.5	8 ± 13	212 ± 86	-26.2 ± 0.4	28.9 ± 16.3	2.29 ± 0.30	-1.9 ± 0.7	0.94 ± 0.48	16.8 ± 3.6	4.3 ± 5.4	0.14 ± 0.04	1.11 ± 0.08
LY04 (238 m)	Wet seasons	27.4 ± 2.3	8.0 ± 0.2	499 ± 172	9361 ± 4358	-20.4 ± 0.3	22.9 ± 4.0	2.30 ± 0.38	-2.9 ± 1.4	0.05 ± 0.09	9.6 ± 4.5	7.1 ± 3.2	0.08 ± 0.03	1.70 ± 0.24
	Dry seasons	21.1 ± 6.0	8.4 ± 0.3	15 ± 15	174 ± 48	-27.5 ± 0.4	17.4 ± 3.9	2.34 ± 0.43	-3.9 ± 1.0	0.37 ± 0.09	8.3 ± 2.8	2.2 ± 0.7	0.07 ± 0.00	1.41 ± 0.06
	Total	24.7 ± 5.1	8.2 ± 0.3	208 ± 280	3849 ± 5484	-24.7 ± 3.9	20.5 ± 4.7	2.41 ± 0.32	-3.3 ± 1.2	0.18 ± 0.19	9.0 ± 3.6	4.7 ± 3.4	0.07 ± 0.02	1.56 ± 0.22
BNE (7 m)	Wet seasons	28.0 ± 2.6	7.8 ± 0.1	106 ± 131	2282 ± 2718	-25.0 ± 1.4	45.1 ± 22.4	2.58 ± 0.45	-4.9 ± 2.5	0.78 ± 1.03	31.0 ± 14.1	7.3 ± 6.2	0.19 ± 0.09	1.84 ± 0.20
	Dry seasons	21.5 ± 6.8	8.4 ± 0.3	12 ± 17	245 ± 73	-26.4 ± 0.6	35.6 ± 10.0	2.02 ± 0.28	-5.7 ± 0.2	1.73 ± 0.43	30.1 ± 10.5	1.7 ± 0.2	0.25 ± 0.07	1.46 ± 0.09
	Total	25.2 ± 5.5	8.4 ± 0.2	49 ± 84	1060 ± 1759	-25.8 ± 1.1	41.0 ± 17.6	2.30 ± 0.45	-5.2 ± 1.8	1.16 ± 0.92	30.6 ± 11.7	4.9 ± 5.3	0.21 ± 0.08	1.65 ± 0.25

* Parameters shown in bold are significantly different between two seasons (Wilcoxon test, *p* < 0.05). Detailed data are provided in Table S1.

¹ T: water temperature; TSM: total suspended matter.

² TIN includes nitrate and nitrite (nitrite was below the detection limit).

331
332
333
334
335

336 $\rho_{\text{DIC_uptake}}$ under light conditions varied from 0.03 to 1.98 mg C m⁻³ hr⁻¹ (Fig. 3a; Table S3). The rates measured in August
337 were either significantly higher than (for MLL01, LY04, and BNE) or comparable (for CWL01 and DL) with those in
338 other months (Fig. 3a). Within-site temporal variation was significant for all five sites (Kruskal-Wallis: MLL01, H = 9.97,
339 $p = 0.019$; CWL01, H = 9.97, $p = 0.019$; DL, H = 7.2, $p = 0.027$; LY04, H = 9.15, $p = 0.027$; BNE, H = 9.46, $p = 0.024$).
340 The rates were also higher for BNE than for MLL01, CWL01, and DL (Kruskal-Wallis, H=25.07, $p < 0.0001$; Dunn's
341 post-hoc, $p < 0.05$; Fig.3a). In contrast, incubations with ¹³C-labeled DIC under dark conditions exhibited a drastically
342 different pattern (Fig. 3b). Most rates were either below the detection limit or less than 0.05 mg C m⁻³ hr⁻¹. Exceptions
343 occurred for sites MLL01 (0.7 mg C m⁻³ hr⁻¹), LY04 (1.6 mg C m⁻³ hr⁻¹), and BNE (0.19 mg C m⁻³ hr⁻¹) in August.
344




Figure 3: Carbon transformation rates along the Beinan River. DIC uptake rates under (a) light and (b) dark incubation conditions across four sampling dates. The inset in panel (b) shows the same data on a logarithmic scale. (c) Amino acid uptake rates and (d) catabolic rates for glycine and leucine across four sampling dates. The inset in panel (c) shows the same data on a logarithmic scale. Error bars represent the standard deviation of triplicate measurements (n=3) for the light DIC, glycine, and leucine treatments. The dark DIC incubation was performed on a single set (n=1) and is shown without error bars. No bar is shown for replicates that yielded negative rates, which were excluded from analysis. Letters above bars indicate statistically significant differences among sites based on the Kruskal-Wallis test followed by Dunn's pairwise post-hoc test (panel a: $H = 25.07$, $p < 0.0001$; panel d: $H = 15.47$, $p = 0.004$). No significant among-site differences were detected for amino acid uptake rates in panel (c) ($H = 5.680$, $p = 0.224$). Asterisks following site names in panel (a) indicate significant within-site temporal variation (Kruskal-Wallis, $p < 0.05$).

ρ_{AA_uptake} ranged from below the detection limit to $0.49 \text{ mg C m}^{-3} \text{ hr}^{-1}$ (Fig. 3c; Table S3). These rates were generally higher in August than in January at all sites. Uptake rates with glycine were generally higher than those with leucine (except at LY04 in August) (Fig. 3c). No significant differences were detected among sites for amino acid uptake rates (Kruskal-Wallis, $H = 5.68$, $p = 0.224$). For comparison, catabolic rates ranged from 4.5 to $154.4 \text{ mg C m}^{-3} \text{ hr}^{-1}$ and varied spatially and temporally (Fig. 3d; Table S3). $\rho_{AA_catabolic}$ with glycine were also higher than those with leucine across most incubations (except for DL and LY04 in January) (Fig. 3d). The catabolic rates appeared to be higher in August than in January for sites CWL01, DL, and LY04, but were comparable between the two months for site BNE. No significant among-site differences for catabolic rates were detected (Kruskal-Wallis, $H = 15.470$, $p = 0.004$). Within-site differences between two time points were not tested, because a two-sample rank test with only three replicates per group ($n = 3$) cannot yield a statistically significant result. By contrast, the catabolic rates for MLL01 were higher in January than in August. Most of the added amino acids were utilized for respiration (as the product of total CO_2 ; $>99\%$) rather than biosynthesis.

To identify the environmental drivers of the metabolic rates, Pearson correlation analyses were performed between the rates and the measured environmental parameters (Fig. S4). DIC uptake rates under light conditions were positively correlated ($p < 0.01$) with multiple parameters, including DOC, POC, $\delta^{13}\text{C}$ -POC, nitrate/nitrite, phosphate, TSM, water temperature, and NH_4^+ . The phototrophic rates, defined as the difference between rates under light and dark, were positively correlated with DOC, nitrate/nitrite, and phosphate. In contrast, DIC uptake rates under dark conditions were most strongly positively correlated with sediment-associated parameters (TSM, POC, $\delta^{13}\text{C}$ -POC, and NH_4^+). Glycine uptake rates were positively correlated with DOC, nitrate/nitrite, and phosphate, whereas leucine uptake was positively correlated with TSM, POC, $\delta^{13}\text{C}$ -POC, and NH_4^+ . Glycine catabolic rates showed a strong negative correlation with DOC concentration, while leucine catabolic rates showed no significant correlation with any environmental parameter.

Scaling measured rates to the catchment level yielded net metabolic CO_2 production fluxes of $5.16 \text{ mol m}^{-2} \text{ yr}^{-1}$ for MLL01, $21.11 \text{ mol m}^{-2} \text{ yr}^{-1}$ for CWL01, $4.22 \text{ mol m}^{-2} \text{ yr}^{-1}$ for DL, $5.35 \text{ mol m}^{-2} \text{ yr}^{-1}$ for LY04, and $24.44 \text{ mol m}^{-2} \text{ yr}^{-1}$ for BNE (Table S6). The total yield of planktonic metabolic CO_2 for individual sites ranged from 10^5 to 10^7 mol yr^{-1} , with a catchment-wise total of $7.89 \times 10^7 \text{ mol yr}^{-1}$ ($4.56 \times 10^7 - 1.14 \times 10^8 \text{ mol yr}^{-1}$). For comparison, CO_2 evasion fluxes through the river-air exchange ranged from 0.20 – $1.76 \text{ mol m}^{-2} \text{ d}^{-1}$ for CWL01, 0.76 – $17.3 \text{ mol m}^{-2} \text{ d}^{-1}$ for MLL01, 0.27 – $0.73 \text{ mol m}^{-2} \text{ d}^{-1}$ for DL, 0.39 – $1.56 \text{ mol m}^{-2} \text{ d}^{-1}$ for LY04, and 0.20 – $0.90 \text{ mol m}^{-2} \text{ d}^{-1}$ for BNE (Table S7). Summing exchange rates across the catchment yielded a total annual emission of 2.08×10^9 ($0.96 - 3.2 \times 10^9 \text{ mol yr}^{-1}$). Planktonic microbial processes contributed approximately 3.8% of the total annual CO_2 evasion rate (Table S6 and S7).

Bacterial and archaeal 16S rRNA gene abundances for river water ranged from 2.3×10^3 to $3.9 \times 10^7 \text{ copies L}^{-1}$ and from 5.6×10^4 to $5.5 \times 10^6 \text{ copies L}^{-1}$, respectively (Fig. S3). The abundances of *cbbL* ranged from 1.8×10^3 to $7.2 \times 10^5 \text{ copies L}^{-1}$. All the gene abundances were generally higher in January than those in other months (mostly by more than one order of magnitude, except for BNE). The variation in gene abundance between January and August was most pronounced at

LY04, where bacterial 16S rRNA gene copy numbers were ~2 orders of magnitude smaller in August than in January (Fig. S3). By comparison, archaeal 16S rRNA gene abundances at BNE were relatively stable across the four sampling months. Statistical analyses by season were not conducted due to insufficient biological replication in the gene abundance measurements.

A total of 532,787 reads of 16S rRNA gene amplicons were obtained and classified into 13,005 ASVs. The observed ASVs and Chao1 estimates based on the rarefied dataset (n=5220) ranged from 200 to 1600 and from 150 to 2300, respectively (Table S3). The Shannon index ranged from 4.4 to 6.9. Alpha diversity varied among sites and sampling intervals (Fig. 4a). Kruskal-Wallis tests did not detect statistically significant differences among sites (Observed ASVs: $H = 6.99, p = 0.14$; Chao1: $H = 6.35, p = 0.17$; Shannon: $H = 8.36, p = 0.08$) or between seasons (all $p > 0.72$). Nonetheless, Shannon diversity tended to be highest at the upstream tributary CWL01 (mean = 6.4) and lowest at the river mouth BNE (mean = 5.0). At MLL01 and LY04, observed ASVs and Chao1 dropped markedly during the early wet season (July), whereas CWL01, DL, and BNE showed more stable values across the study period.





Figure 4: (a) Alpha diversity indices (observed ASVs, Chao1, Shannon index) and (b) community compositions at the phylum level, with Proteobacteria resolved at the class level. Error bars on Chao1 represent ± 1 SE estimated from rarefaction. Time intervals with blue background represent the wet season.

Community compositions varied temporally and spatially to different degrees, with the major phyla including Bacteroidota, Cyanobacteria, Firmicutes, Campilobacterota, Desulfobacterota, and Proteobacteria (Alpha and Gamma divisions) (2–20%; Fig. 4b). Gamma-Proteobacteria appeared to be more abundant in most time intervals and at most sites (except for BNE). Cyanobacteria was more abundant in the dry season than in the wet season, and vice versa for

411 Campilobacterota and Desulfobacterota (Fig.4b). For BNE, Bacteroidota constituted the major phylum for two out of
 412 three dry season samples. Firmicutes combined with Gamma-Proteobacteria outnumbered (>54%) the others during the
 413 wet season and January. Among all detected ASVs, ASVs with sequences related to Thiotrichaceae, Erysipelotrichaceae,
 414 and Phormidiaceae represented the most abundant ones. However, their abundances varied dramatically among sites
 415 (Table S8).

416 Beta diversity analyses showed that community composition was structured primarily by site (PERMANOVA: $F_{4,19} =$
 417 2.01 , $R^2 = 0.30$, $p = 0.001$), while the seasonal effect was marginal ($F_{1,22} = 1.29$, $R^2 = 0.06$, $p = 0.087$). The joint site-and-
 418 season model explained 35% of the variance ($p = 0.001$). In the CCA ordination (Fig. 5), BNE samples deviated from the
 419 cluster formed by the other sites and were associated with elevated DOC. For the remaining sites, dry-season samples
 420 clustered, whereas wet-season samples were more dispersed and showed positive correlations with TSM, POC, and ^{13}C -
 421 POC. All four environmental vectors were significant (envfit, $p < 0.05$; $r^2 = 0.62$ – 0.96 ; Fig. 5).

422



423

424 **Figure 5. Canonical correspondence analysis (CCA) ordination of microbial community composition of 16S rRNA gene**
 425 **amplicons (ASV proportion). Significant environmental vectors (envfit, 9999 permutations, $p < 0.05$) are shown as solid arrows:**
 426 **TSM ($r^2 = 0.94$, $p = 0.006$), POC ($r^2 = 0.96$, $p = 0.007$), DOC ($r^2 = 0.62$, $p = 0.011$), and $\delta^{13}\text{C}$ -POC ($r^2 = 0.65$, $p = 0.035$).** Symbols
 427 **indicate site (color) and season (circle = dry, triangle = wet).**

428

429 4. Discussion

430

430 4.1 Solute and nutrient status, and river metabolism

431 The general physicochemical characteristics of river water (pH, major ions, TOC, TSM, and ^{13}C -DIC) were comparable
 432 to those reported previously (Wang et al., 2024). These solute characteristics are mostly derived from carbonate
 433 dissolution (>80% of the total solute load) driven by sulfuric acid produced by microbially mediated pyrite oxidation
 434 (Wang et al., 2024). Considering that a great portion of discharge is attributed to monsoons or typhoons, solute yields

from carbonate dissolution during these extreme events dominate over the quiescent period and could be translated into a significant flux of CO₂ emissions that constitutes a disproportionally large fraction of global flux (~1% from 0.002% of three major catchments in Taiwan). The high contribution pattern of carbonate weathering to the solute budget has also been observed for other active orogens (e.g., New Zealand and the Himalayas) (Emberson et al., 2016; Kemeny et al., 2021). Furthermore, a high-water regime during summer monsoon or typhoons drains erodible materials (sediments accumulated in banks or hillslopes) into channels. As a result, TSM concentrations increased to a range of hundreds of mg L⁻¹, up to a factor of 40 (at LY04) between the wet and dry seasons (Table 2). This magnitude is far less than previous observations during the extreme events (up to 10 g L⁻¹; Dadson et al. 2003; Kao and Milliman 2008; Wang et al., 2024), given that the annual precipitation during the sampling period was much less than previous years (1856 mm for this study period versus ten-year average of 3402 mm for the most upstream Siangyang weather station; CWB annual report), but could still be sufficient to influence the expression of investigated metabolisms (see the next section). Similar increases in POC concentrations were also observed (Table 2). For comparison, dissolved cation concentrations were characterized by varying degrees of dilution attributable to increased discharge (linear mixed-effects model, $p < 0.05$; Tables S1 and S2). In contrast, sulfate is significantly more abundant in the wet than dry seasons (linear mixed-effects model, $p = 0.004$), a pattern consistent with enhanced pyrite oxidation during sediment mobilization in this catchment (Wang et al., 2024) and the elevated wet-season abundance of sulfur-oxidizing taxa (e.g., *Thiobacillus* and *Sulfuricurvum*) (Section 4.6).

While nitrate/nitrite and phosphate concentrations did not differ significantly between seasons, Chl *a* concentrations were significantly higher in the dry season, and ammonium concentrations were significantly higher in the wet season (linear mixed-effects model, both $p < 0.05$; Table S2). As ammonium is the degradative product of organic particulates, the higher ammonium concentrations in the wet season suggest either greater organic degradation during river transit or more extensive purging of soil porewater during the high-water regime (Calmels et al., 2011). It is also worth noting that the comparable nitrate/nitrite and phosphate concentrations between seasons, or higher ammonium concentrations in the wet season, could translate into higher yields of these nutrients given much higher discharge. Part of the ammonium production may even have been channeled from the degradation of amino acids, which can also be sourced from the degradation of suspended organic particulates. As indicated by our incubation results, amino acid mineralization occurred across all investigated sites distributed from the upstream to the estuary. Their rates were also higher in the wet than in the dry season, coinciding with the increased input of TSM. Alternatively, soil development is limited in the investigated catchment. High erosion and heavy precipitation associated with extreme hydrologic events could have rapidly drained stagnant soil porewater along with soil particulates and groundwater residing in the fracture network into the river. As these water bodies retain the degradative products of organic matter and their transport is slow (Alley et al., 2002; Calmels et al., 2011), these nutrient constituents would accumulate to high levels during the quiescent period and be rapidly purged only when erosion and high hydraulic head are imposed. As ammonium concentrations tended to increase from the upstream headwater (MLL01: 2.6 μM) toward the river mouth (BNE: 4.9 μM) (3.7–4.7 μM; Table 2), the concentration gradient was even more pronounced during the wet season than the dry season. The data pattern seems consistent with progressive accumulation of soil-derived inputs along the catchment under high flow conditions, and may be accounted for by the increasing agricultural land use in the lower catchment near the estuary (Fig. 1b). Finally, water temperatures during the wet season were systematically ~6°C higher than during the dry season (Table 2). Enhanced microbial degradation of suspended or soil organic matter driven by this temperature increase may have contributed to increased ammonium release. Overall, current data do not allow us to identify the primary cause driving enhanced ammonium concentrations in the wet season. High-frequency monitoring of river water and groundwater across the event period is

warranted to gauge the mechanism of nutrient release from the soil. Such a data pattern is also comparable to the nutrient-deprived oligotrophic state commonly observed in the open ocean (Overholt et al., 2022).

Chl *a* concentrations were significantly higher during the dry season than during the wet season (linear mixed-effects model, $p = 0.007$), by a factor of at least 2.2 (except for site DL). Pearson correlation analysis revealed a significant negative correlation between Chl *a* and TSM ($r = -0.46$, $p < 0.05$) and no significant correlation between Chl *a* and POC ($r = -0.41$, $p = 0.06$) (Table S5). Chl *a* concentrations at sites MLL01 and LY04 during the wet season were barely detectable. While nutrient concentrations (nitrate/nitrite and phosphate) do not vary considerably between MLL01 and LY04, nutrients do not appear to be a deterministic factor in controlling phototrophic activity. Instead, the load of suspended sediment can affect the expression of phototrophy even at a scale of hundreds of mg L^{-1} , a level that is not particularly unusual for mountainous rivers. Potential mechanisms inhibiting phototrophic activity include the shading created by dense sediments, grain pulverization, and abrasion (Stanley et al., 2010). The results are consistent with O'Donnell and Hotchkiss (2022), who found that high flow events drove gross primary production (GPP) and ecosystem respiration (ER) to deviate from baseline metabolism. Their results also indicate that both GPP and ER rates were reduced by flash flow, with GPP rates varying at a greater magnitude than ER (Błaszczak et al., 2019; Hall et al., 2015).

4.2 Autotrophic carbon fixation: rates and controlling factors

DIC uptake rates were significantly correlated with TSM, POC, $\delta^{13}\text{C}$ -POC, and NH_4^+ regardless of light availability (Fig. S4). Under light conditions, DIC uptake rates were also correlated with temperature, DOC, and nutrients. DIC rates for the wet season could be qualitatively categorized into two patterns. The first category is represented by sites CWL01, DL, and BNE, where rates with light were higher than those without light (Fig. 3a and 3b). The largest difference at $43 \text{ mg C m}^{-3} \text{ d}^{-1}$ was observed at the most downstream site, BNE (Table S3). The second category is represented by sites MLL01 and LY04, where rates with light were either comparable to or slightly less than those without light. For incubations with DIC for the dry season, rates with light were generally much larger than those without light. In fact, rates without light were below the detection limit for sites MLL01, DL, and BNE. The only exception occurs for site CWL01, where DIC uptake rates with light were slightly higher than those without light. The results in part corroborate the Chl *a* pattern but offer an additional dimension for constraining river metabolism.

The contrast pattern described above is apparently consistent with light limitation imposed by suspended sediment, which was far higher at MLL01 and LY04 (wet-season TSM ~ 157 and $\sim 499 \text{ mg L}^{-1}$) than at CWL01, DL, and BNE ($\sim 4 - 106 \text{ mg L}^{-1}$). Furthermore, differences in rates with and without light could be interpreted as the phototrophic rates, assuming that lithoautotrophic rates (obtained from dark conditions) are constant regardless of whether light is available. In this regard, phototrophic rates for sites MLL01 and LY04 were lower in the wet season and were even completely inhibited with the replacement of lithoautotrophy. In contrast, phototrophic rates were greater in the wet season than in the dry season for the other three sites, even though lithoautotrophy was substantially reduced during the dry season. Additionally, the significantly positive correlations between DIC rates under light conditions and TSM, POC, and $\delta^{13}\text{C}$ -POC contradict the expression of phototrophy. This pattern could be largely accounted for by the fact that a large fraction of the DIC rates with light (115-145%) were attributed to the lithoautotrophic rates during the wet season (Table S3). Therefore, the significant correlation for light conditions resembles that for dark conditions. By comparison, the significant correlation between the difference in DIC rates and nutrient concentration is compatible with the fact that primary production is highly dependent on nutrient availability in aquatic environments (Cloern, 1999; Domingues et al., 2005). As the nutrients generated from the degradation of organic matter accumulate along the river flow, BNE was characterized by the highest concentrations of nutrients and phototrophic rates in the wet season (Fig. 3). Taken together, these patterns suggest that

515 phototrophic rates were primarily limited by light availability at MLL01 and LY04, whereas nutrient availability was the
516 more relevant limiting factor at CWL01, DL, and BNE. Notably, the Chl *a* concentration did not show a significant
517 correlation with the phototrophic rate (Fig. S4).

518 4.3 Particulate-mediated lithoautotrophy in the wet season

519 The incubation with DIC under dark conditions yielded DIC uptake rates spanning from below the detection limit to 1.6
520 mg C m⁻³ hr⁻¹ (Fig. 3; Table S3). These rates constitute only a small fraction (<5%) of the rates under light conditions but
521 become generally high and even comparable with the rates under light conditions for samples collected during the period
522 of high sediment load (LY04 and MLL01 in August). As these two sub-catchments are susceptible to high erosion of slate
523 formations, such a spatial pattern of dark DIC uptake is consistent with geological control rather than the upstream-
524 downstream position along the catchment. The rate pattern is intriguing and suggests intimate relationships with the
525 potential substrates supplied from the suspended particulates or reduced ions from the groundwater influx.

526 For particulate matter, potential substrates include reduced sulfur or iron-bearing minerals (e.g., pyrite) and ammonium
527 encapsulated within the interlayer of clay minerals (Petit et al., 2006). Pyrite is prevalently distributed in metasedimentary
528 rocks of the catchment and has been identified as the main solute and CO₂ contributor of weathering (Blattmann et al.,
529 2019; Bufe et al., 2021). A previous study conducted on the same catchment has shown a highly positive correlation of
530 the yields between riverine sulfate, suspended sediment, and sulfur oxidizers (mainly composed of *Thiobacillus* and
531 *Sulfuricurvum* members) (Wang et al., 2024). The data pattern suggests that sulfate/sulfuric acid is primarily generated
532 by the microbially mediated oxidation of pyrite in erodible materials in riparian zones or hillslopes. As the torrential
533 precipitation continues, these materials, along with the sulfur oxidizers, are drained into the river channels and recovered
534 by sampling and analysis. The data corroborate that the erosion of bedrocks and subsequent pulverization on hillslopes
535 or in river channels facilitate the exposure of pyrite to the atmosphere or to air-infiltrated pore space, thereby rendering
536 the oxidative reaction that produces sulfuric acid for dissolution of carbonate or other minerals (Hilton and West, 2020).
537 Furthermore, particulate pulverization and abrasion can also disrupt the cleavage or interlayer structure of clay minerals,
538 thereby releasing ammonium into the surrounding water (Yu et al., 2023). While river water is presumably saturated with
539 atmospheric oxygen, pyrite and ammonium associated with suspended particulates are readily available for
540 lithoautotrophic microbial exploitation and colonization.

541 For groundwater influx into rivers, potential substrates for lithoautotrophy include ferrous iron, ammonium, and sulfide,
542 all of which could be produced by either microbial reduction of oxidants (e.g., iron oxyhydroxide, nitrate, and sulfate) or
543 degradation of organic matter (for the release of ammonium). Considering that sulfate minerals have not been identified
544 or recovered from the catchment to date, ferrous iron and ammonium produced in subsurface aquifers could be drained
545 and discharged into rivers. Indeed, ammonium concentrations were significantly higher in the wet than in the dry seasons
546 (Table 2). Our current analyses also revealed that potential sulfur oxidizers related to these two genera (*Thiobacillus* and
547 *Sulfuricurvum* members) and others (e.g., *Sulfurovum*, *Sulfurifustis*) and ammonium oxidizers related to *Nitrosomonas*
548 and *Nitrosopumilus* constituted a significant proportion of communities (>10%) in the wet season as compared to 5% in
549 the dry season. A detailed experimental design is required to pinpoint the exact substrate involved in the aquatic
550 lithoautotrophy.

551 4.4 Heterotrophic amino acid metabolism and controlling factors

552 In contrast to autotrophic carbon fixation, heterotrophic metabolism showed distinct seasonal patterns and substrate
553 preferences. Glycine and leucine uptake rates were correlated with nutrients and particulate characteristics (concentration

and isotopic composition), respectively (Fig. S4). As the nutrients represent the degradative products of organic particulates, these relationships seem to suggest that the direct uptake of amino acids by riverine community members is linked to the metabolites generated by particulate degradation (for glycine) or particulate-associated populations (for leucine). For the former scenario, metabolites or nutrients released from the particulate degradation accumulated along the flow, peaking at the estuary site (BNE) when compared with upstream sites. Therefore, the correlation relationships suggest that enhanced nutrient abundances facilitate the boosting of activity in a riverine population, preferentially for glycine uptake. Such a relationship is also analogous to the priming effect on soil microbiota, where the addition of exogenous substrate or substance promotes further degradation of organic matter (Blagodatskaya and Kuzyakov 2008; Chen et al., 2014). On the other hand, the latter scenario suggests that leucine uptake is linked to the size of the population attached to particulate organic matter. With a larger influx of TSM in the wet season, the size of the attached population could have been correspondingly enhanced, providing a larger population of microorganisms with functional preferences for leucine uptake. The inference also suggests a divergent modulation of physiological preferences for glycine and leucine uptake mediated by two distinct populations. The inference may be better constrained by expanding the incubations to various concentrations of particulates and simultaneous analyses of stable isotope probing of cellular components (proteins, lipids, or nucleic acids) to identify the plausible populations actively incorporating either of these amino acids. Despite these substrate-specific correlations with site-varying parameters, amino acid uptake rates did not differ significantly among sites (Kruskal-Wallis, $H = 5.68$, $p = 0.224$), suggesting that the correlative patterns described above reflect catchment-scale trends rather than localized among-site contrasts.

The exploitability of organic substrates for heterotrophy is worth noting. Glycine uptake rates were generally higher than leucine uptake rates for the wet season (except for LY04, where both rates were comparable) (Fig. 3c). In the dry season, these two uptake rates were comparable but noticeably much lower than those in the wet season. For comparison, glycine catabolic rates were higher than leucine catabolic rates for CWL01, DL, and LY04 in the wet season. Both rates were comparable to each other at other sites (Fig. 3d). Overall, the selectivity of glycine over leucine for the wet season was prevalent (at four out of five sites for uptake and three out of five sites for catabolism). The cause of substrate selectivity may be attributed to the preferential utilization of simply structured substrates by specific community members or enzymatic expression (Geisseler and Horwath 2014; Palenik and Morel 1990). The selectivity for glycine likely reflects higher metabolic efficiency of glycine-centered pathways. The reductive glycine pathway (rGlyP) and glycine cleavage system (GCS) offer highly ATP-efficient routes for both biosynthesis and energy generation (Dronsella et al., 2025; Kikuchi et al., 2008), providing a more consistent mechanistic basis for glycine metabolism across seasons. In contrast, leucine catabolism requires a complex multi-step pathway involving at least six enzymatic reactions to yield acetyl-CoA and acetoacetate (Brosnan and Brosnan, 2006; Massey et al., 1976), making respiratory oxidation costly. Therefore, microorganisms would preferentially incorporate leucine into proteins (Kirchman, 2001; Kirchman et al., 1985), such that leucine-derived rates provide a conservative estimate for heterotrophic activity. As both amino acids are required for biosynthesis, further experiments are needed to clarify the expression of specific metabolic pathways.

The results for heterotrophic incubations in this study revealed that catabolic rates exceeded assimilation rates by at least two orders of magnitude (Fig. S5). The great rate contrast was partially accounted for by the fact that ^{13}C was labeled at the carboxyl position of amino acids in our incubations. The carboxyl carbon in amino acids would be primarily cleaved and converted into CO_2 , whereas all the other unlabeled carbon would be assimilated and partially converted into CO_2 at the cost of energy expenditure (Hill et al., 2013; Suttle et al., 1991). Therefore, carbon uptake would be primarily sourced from non-carboxyl carbon, with limited contribution from carboxyl carbon. Other than the site preference associated with the labeled position, which would result in underestimation of both rates, the ratios of catabolic to uptake rates remain

related to community activity and composition modulated by environmental parameters. Except for January incubation with leucine for CWL01, all the other incubations with amino acids yielded a ratio of catabolic to uptake rates for January greater than for August. The ratios even reached an infinite level (uptake rate below the detection limit) for MLL01 and DL in January. The overwhelmingly large ratios were primarily controlled by heterotrophic assimilation at extremely low levels during the dry season when sediment supply was low. Although the incubations were provided with readily available amino acids, the slow assimilation suggests that the communities in the dry season may be customized to a deprived supply of nutrients and small organic molecules generated from the degradation of particulate organic matter. Therefore, any solid or soluble substrate available for microbial utilization would be initially diverted to catabolism for energy generation to sustain certain maintenance activities (McInerney et al., 2010). Thus, even when the amino acids were provided during incubations, their assimilation was not immediately enhanced under such energy-limited conditions for heterotrophs. Conversely, communities in the wet season had access to an abundant supply of particulate organic matter to catalyze metabolism for energy generation or assimilation, which resulted in a relatively larger fraction of the fed amino acids being diverted to assimilation or biosynthesis (Table S3, Fig. S5). Nevertheless, the heterotrophy in such oligotrophic mountainous rivers yields excessive CO₂ that is susceptible to net emission through the air-water exchange.

4.5 Implications for river heterotrophy and carbon cycling

The rate comparison between autotrophy and heterotrophy (assimilation plus catabolism) indicates that the planktonic compartment of the Beinan River is net heterotrophic, producing excess CO₂ through planktonic microbial processes. As the incubations were performed over a short period of time (4–6 hrs), the physiological status of community members likely resembles *in situ* conditions. For DIC-based measurements provided with ¹³C at a concentration less than 5% of the ambient DIC pool, the derived rates were considered to bracket the range of true rates. For amino acid-based measurements, the 50 μM addition of amino acid exceeds the ambient concentration and half-saturation constants (K_m) for oligotrophic systems (11–79 nM in Boysen et al., 2022; 1.41 μM in Brailsford et al., 2019). Therefore, our experimental setup may represent the possible stimulation of uptake and respiration for our river systems. Furthermore, part of the unlabeled carbon in amino acids that was assimilated and eventually respired was not quantified in our experiments; the rates obtained were mostly derived from the carboxyl ¹³C in amended amino acids. Therefore, our measurements represent an underestimate of respiration under the experimental setup. To provide a first-order and conservative quantification of possible heterotrophic CO₂ production, the leucine catabolic rates were arbitrarily chosen to estimate the carbon export at the catchment scale.

We further note that the experimental design captures only part of metabolisms on dissolved organic substrates and does not directly probe the decomposition of pre-existing suspended POC. It is also worth noting that the current estimates constrain only the planktonic compartment of river metabolisms. The benthic and hyporheic zones, which are connected to the river water and contribute to the cycling of organic carbon, were not measured in this study. In addition, this catchment-scale estimate is sensitive to the assumed river surface area fraction (0.47% of sub-catchment area; Raymond et al., 2013), derived from a global average reported for a range of 0.30–0.56%. Given the challenge of independently constraining this fraction for the present catchment, we arbitrarily adopted a conservative ± 50% uncertainty for river surface area, and combined this with the measured variability in rate or flux for each site (Section 2.5). This approach yields a catchment-wise planktonic CO₂ production estimate of $7.99 \times 10^7 \text{ mol yr}^{-1}$ ($4.56 \times 10^7 - 1.14 \times 10^8 \text{ mol yr}^{-1}$), and a total river-air CO₂ evasion flux of 2.08×10^9 ($0.96 - 3.20 \times 10^9 \text{ mol yr}^{-1}$).

Our planktonic metabolic CO₂ production ($7.89 \times 10^7 \text{ mol yr}^{-1}$) accounts for less than 5% of river-atmosphere CO₂ exchange flux ($\sim 2.08 \times 10^9 \text{ mol yr}^{-1}$) and CO₂ production from chemical weathering ($2.8 \times 10^9 \text{ mol yr}^{-1}$; Wang et al., 2024) and petrogenic carbon oxidation ($1.7\text{-}2.9 \times 10^9 \text{ mol yr}^{-1}$; Lien et al., 2025). This pattern indicates that net CO₂ emissions through the river-air interface are overwhelmingly driven by the weathering of minerals and petrogenic carbon. While benthic and hyporheic contributions may have been reduced during the high-water period (Battin et al. 2023; Hall et al., 2015), they are likely substantial during dry-season low-TSM conditions when the water column is clear, and light can reach the stream bed. To place the planktonic contribution in the broader context of stream metabolism, our rates were compared with rates reported for morphologically comparable upland streams. Our measured DIC uptake rates ($0.03\text{-}1.98 \text{ mg C m}^{-3} \text{ h}^{-1}$) were transformed into a flux of 4×10^{-4} to $2 \times 10^{-2} \text{ g C m}^{-2} \text{ d}^{-1}$ assuming an average water depth of 0.5 or 0.7 m (Table S6). By comparison, benthic gross primary production fluxes for clear upland streams typically range from $\sim 10^{-1}$ to several $\text{g C m}^{-2} \text{ d}^{-1}$ (Hall et al., 2015; Bernhardt et al., 2022). Furthermore, Rovelli et al. (2017) isolated water-column metabolism from benthic metabolism in UK headwater streams and reported that water-column metabolism contributed 25% to annual ecosystem respiration and primary production, with peak rates of $2.5\text{-}3.5 \text{ g C m}^{-2} \text{ d}^{-1}$. Therefore, planktonic autotrophic fluxes for the Beinan River are one to two orders of magnitude lower than benthic primary production fluxes reported elsewhere. In contrast, planktonic heterotrophic catabolic fluxes ($0.055\text{-}1.85 \text{ g C m}^{-2} \text{ d}^{-1}$) for the Beinan catchment fall within the typical range of total ecosystem respiration fluxes reported for upland streams (~ 0.5 to several $\text{g C m}^{-2} \text{ d}^{-1}$; Battin et al., 2008; Hall et al., 2015; Rovelli et al., 2017; Bernhardt et al., 2022). Because ecosystem respiration in most of these studies encompasses planktonic, benthic, and hyporheic contributions rather than separating them, our planktonic-only rates suggest that the planktonic compartment can be a significant fraction of total stream respiration. Overall, these comparisons indicate that the low fractional contribution of planktonic metabolism to total CO₂ evasion in the present study reflects the dominance of weathering-derived CO₂ in this catchment, rather than anomalously low planktonic metabolic rates. More detailed measurements are warranted to validate whether different compartments of the Beinan river systems contribute to the net carbon export in a pattern resembling other similar catchments.

Previous studies based on dissolved oxygen data demonstrate that both fluxes of net ecosystem production (NEP) and CO₂ emissions derived from the river-air exchange vary over several orders of magnitude as the discharge or stream order increases for the catchments distributed over contiguous US and European boreal regions (Dodds et al., 2013; Cory et al., 2014; Hotchkiss et al., 2015; Rasilo et al., 2017; Lupon et al., 2019; Bernal et al., 2022). Using a discharge-weighted regression, Hotchkiss et al. (2015) found that NEP fluxes varied much less. Such a regional data pattern also suggests that the contribution fraction of NEP to the total emission increases from $\sim 15\%$ to $\sim 50\%$ as the discharge increases to hundreds of cms (cubic meter per second). Only limited data simultaneously covering NEP, CO₂ emissions, and discharge are available for individual catchments. These studies show that the fractions range from 17% to 75% for rivers in Canada (Rasilo et al., 2017), from 17% to 51% in Sweden (Lupon et al., 2019), $\sim 22\%$ in southeastern US (Dodds et al., 2013), and from 51% to 57% in northeastern Spain (Bernal et al., 2022). Recent modeling for catchments in contiguous US suggests that the NEP accounts for $\sim 7\%$ of the total evasion (Maavara et al., 2025). The net planktonic CO₂/exchange fractions ($<5\%$) reported herein are even smaller than or comparable to any fraction obtained in previous studies. Such a low range of net planktonic CO₂ exchange fluxes is in great contrast to most of the examples obtained elsewhere, even for the site (BNE) located nearest to the shoreline or with the greatest stream order (SO = 5). The pattern suggests that steep topography and rapid erosion associated with active mountain building limit the development of soils and enable a scarce supply of nutrients and rapid transit of river water, all of which apparently restrain the expression of planktonic metabolisms. In contrast, voluminous amounts of metamorphic rocks are fractured and eroded, providing unlimited

674 materials readily for the oxidation of pyrite and petrogenic carbon mediated by microbial processes in the subsurface and
675 riverbanks (Wang et al., 2024; Lien et al., 2025). Consequently, CO₂ generated by these weathering reactions is drained
676 into the river along with groundwater flow and dominates over the contribution from other sources.

677 4.6 Seasonal and spatial dynamics of microbial communities

678 Variations in community diversity and composition were evident across different sites and sampling periods. Beta
679 diversity analysis indicated that microbial community compositions generally varied over time across most sites (Fig. 5).
680 Samples collected during the dry season were generally clustered (with the exception of BNE), and moderately deviated
681 from those collected during the wet season. Based on these ordination patterns and a detailed taxonomic analysis, the sites
682 can be categorized into three distinct groups: those exhibiting clear seasonal patterns (MLL01, CWL01, LY04), site with
683 complex temporal dynamics that do not strictly follow seasonal variation (DL), and the most downstream site (BNE)
684 where communities for different months are scattered to a great extent.

685 MLL01 and LY04 exhibited the most pronounced seasonal trends in community diversity and composition (Figs. 4 and
686 5). Community diversity was lower during the wet season than the dry season, except for the one in January at MLL01
687 (Fig. 4a). During the dry season, communities at both sites were predominantly composed of lithoautotrophs (Table S8).
688 More specifically, MLL01 and LY04 were dominated by iron-oxidizing *Sideroxydans*, and sulfur-oxidizing *Thiothrix*,
689 respectively. Both genera utilize the Calvin-Benson-Bassham cycle for carbon fixation (Emerson et al., 2010, 2013;
690 Grabovich et al., 2023; Ravin et al., 2021). As dark DIC uptake rates were low at these sites during the dry season (Fig.
691 3b; Section 4.3), the abundant lithoautotrophic taxa suggest weak expression of their functions. Major members in
692 communities appeared to shift towards different compositions in the wet season, such as autotrophic *Sulfuricurvum* and
693 *Thiobacillus* (sulfur-oxidizing) at MLL01 and a combination of autotrophic and heterotrophic taxa at LY04 (*Thiobacillus*,
694 *Arenimonas*, and Coxiellaceae). The wet-season shifts coincided with the highest dark DIC uptake rates measured in this
695 study (Section 4.3), indicating that the wet-season lithoautotrophic communities were both abundant and functionally
696 active. The presence of heterotrophic Coxiellaceae during the wet season coincided with increased TSM and POC levels,
697 suggesting their introduction through terrestrial runoff rather than *in situ* growth and the impact of hydrological changes
698 on community compositions.

699 CWL01 was characterized by high community diversity during the sampling period (except in January, Fig. 4a), and
700 showed a distinct seasonal pattern in a way that lithoautotrophic genera (*Sulfuricurvum* and *Thiobacillus*) became more
701 abundant in August, and Cyanobacteria dominated throughout the dry season (Fig. 4b). This interpretation partially aligns
702 with the low dark DIC uptake rates (Fig. 3). The patterns also suggest that abundances of lithoautotrophic and
703 photosynthetic communities are complementary across different seasons, demonstrating significant metabolic versatility
704 within the community.

705 DL was characterized by an intricate temporal pattern that did not follow the distinct dry-wet seasonal variations.
706 Community diversity remained relatively stable across the seasons (Fig. 4). The communities during the dry season
707 (January, March, and May) clustered more tightly than the others (Fig. 5) and were predominantly composed of
708 Cyanobacteria and *Thiothrix*, along with various heterotrophs (e.g., Comamonadaceae and Microscillaceae). The
709 community in August also resembled those in the dry season and was dominated by *Thiothrix* and *Nitrospira*, highlighting
710 the importance of sulfur and nitrogen metabolisms. However, the community in July deviated from this cluster and was
711 dominated by the heterotrophic *Bacillus*. The temporal variability was correlated with DOC concentration, suggesting
712 that organic matter availability and hydrological circulation rather than season per se drive community turnover at this
713 site.

The most downstream site, BNE, was notably distinct from all other mountainous sites. It was characterized by relatively low community diversity throughout the sampling period (except for January) (Fig. 4) and high community variation between campaigns. (Fig. 5). The communities for March and May were dominated by *Fluviicola*, a genus specializing in degrading complex organic compounds (Woyke et al., 2011). For July and August, the community compositions shifted to the predominance of *Turicibacter* and *Polynucleobacter*, both of which are capable of decomposing dissolved organic matter (Bosshard et al., 2002; Watanabe et al., 2012). The prevalence of heterotrophic taxa throughout all seasons was consistent with a high glycine catabolic rate (Fig. 3) and a significant correlation between community variation and DOC concentration (Fig. 5). The pattern also reflects its downstream role, where organic matter accumulated along the drainage path facilitates heterotrophic respiration.

The variation in community composition likely reflects the combined influence of catchment position and hydrological seasonality on substrate and energy availability. The upstream sites (MLL01, CWL01, LY04) are underlain predominantly by metamorphic rocks that can supply reduced iron and sulfur species. The substrate availability is consistent with the detection of abundant lithoautotrophic taxa. At these sites, dominant lithoautotrophic taxa also shifted between dry and wet periods (e.g., iron-oxidizing Sideroxydans and sulfur-oxidizing Thiobacillus in the dry season at MLL01 and LY04 versus *Sulfuricurvum* and *Thiobacillus* in the wet season), suggesting that hydrological seasonality reorganizes the availability or accessibility of specific taxa rather than simply favoring or disfavoring lithoautotrophy as a whole. At DL, community composition instead tracked variation in DOC concentration rather than season directly, pointing to hydrological transport of dissolved organic matter as the more relevant driver at this site. At the most downstream site, BNE, communities were consistently dominated by heterotrophic taxa (*Fluviicola*, *Turicibacter* and *Polynucleobacter*) and were highly variable between campaigns regardless of season – a pattern consistent with the nature of this site, which integrates organic matter and nutrients accumulated from the entire upstream catchment. This downstream transition from rock-driven to organic-matter-driven metabolism is broadly consistent with longitudinal patterns of substrate processing described for river networks elsewhere (e.g., the River Continuum Concept, Vannote et al., 1980). Together, these patterns suggest that hydrological seasonality shapes microbial communities through several site-dependent drivers, including substrate and redox turnover, transport and production of organic matter and nutrients, and integration of upstream inputs, rather than a single mechanism operating uniformly across the catchment. We also noted that abundance and activity were not always tightly coupled across the catchment. For example, lithoautotrophs were numerically abundant at MLL01 and LY04 during the dry season, yet dark DIC uptake rates were low at these sites in dry months. The pattern suggests that the most abundant taxa were not always the most metabolically active. This decoupling underscores that 16S rRNA gene-based abundance is a static metric and does not directly resolve activity. Further studies based on RNA are warranted to identify actively metabolizing community members.

Conclusions

This study elucidates the dynamic nature of planktonic river metabolisms in the Beinan River system. By supplementing ¹³C-labelled substrates, our incubations showed that all categories of metabolic rates were consistently higher during the wet season than the dry season, and that heterotrophic rates were higher than autotrophic rates. The deduced net planktonic metabolic CO₂ rates account for less than 5% of CO₂ exchange at the air-water interface, pyrite-induced weathering, and petrogenic carbon oxidation. Microbial community composition shifted from lithoautotrophic sulfur- and ammonia/nitrite-oxidizing taxa in the wet season to a mixture of phototrophs and heterotrophs in the dry season, reflecting the dynamic response of biological processes to variations in river hydrology and particle loading. Although the study is inherited with several methodological limitations, future work should pursue high-frequency event-driven monitoring,

754 kinetic experiments for *in situ* substrate uptake, RNA-based approaches for actively metabolizing community members,
755 and parallel benthic and hyporheic measurements. Nevertheless, characterization of the water-column compartment
756 remains essential for tracking upstream microbial and geochemical signals and for predicting downstream fluxes to
757 estuarine environments. Our findings highlight the complexity and dynamics of planktonic river metabolisms and
758 microbial communities in tectonically active mountainous catchments prone to intense hydrological events and rapid
759 landscape changes.

760 **Author contribution**

761 Jhen-Nien Chen: conceptualization; formal analysis; investigation; methodology; writing—original draft. Pei-En Chen:
762 conceptualization; methodology; formal analysis; writing—original draft; Yu-Shiang Yen: methodology; formal analysis;
763 investigation. Tzu-Hsuan Tu: conceptualization; methodology; writing - review & editing. Lu-Yu Wang:
764 conceptualization; methodology; data curation. Wan-Yin Lien: conceptualization; formal analysis; writing - review &
765 editing. Yueh-Ting Lin: conceptualization; data curation; writing - review & editing. Pei-Ling Wang: Conceptualization;
766 funding acquisition; investigation; methodology; project administration; resources; supervision; validation; writing—
767 review and editing.

768 **Competing interests**

769 The authors declare that they have no conflict of interest.

770 **Acknowledgements**

771 We are grateful for the funding support from the Taiwanese MOE and NSTC to THT (NSTC 113-2116-M-110-002) and
772 PLW (NSTC 114-2116-M-002-026, NSTC 113-2116-M-002-027, MOE 113L901005). We also acknowledge the
773 logistical support and comments provided by Li-Hung Lin and the assistance in sampling and geochemical analyses by
774 Ya-Fang Cheng, Tzu-Jung Cheng, Yun-Hsuan Lee, Ruei-Feng Tsai, Bo-Yu Chen, Jing-Yi Tseng, and Ling-Wen Liu.

775 **References**

- 776 Alfreider, A., Vogt, C., Hoffmann, D., and Babel, W.: Diversity of Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase
777 Large-Subunit Genes from Groundwater and Aquifer Microorganisms, *Microb. Ecol.*, 45, 317–328,
778 <https://doi.org/10.1007/s00248-003-2004-9>, 2003.
- 779 Allen, G. H. and Pavelsky, T. M.: Global extent of rivers and streams, *Science*, 361, 585–588,
780 <https://doi.org/10.1126/science.aat0636>, 2018.
- 781 Alley, W. M., Healy, R. W., LaBaugh, J. W., and Reilly, T. E.: Flow and Storage in Groundwater Systems, *Science*,
782 296, 1985–1990, <https://doi.org/10.1126/science.1067123>, 2002.
- 783 Appling, A. P., Hall, R. O., Yackulic, C. B., and Arroita, M.: Overcoming Equifinality: Leveraging Long Time Series
784 for Stream Metabolism Estimation, *J. Geophys. Res. Biogeosciences*, 123, 624–645,
785 <https://doi.org/10.1002/2017JG004140>, 2018.
- 786 Aufdenkampe, A. K., Mayorga, E., Raymond, P. A., Melack, J. M., Doney, S. C., Alin, S. R., Aalto, R. E., and Yoo, K.:
787 Riverine coupling of biogeochemical cycles between land, oceans, and atmosphere, *Front. Ecol. Environ.*, 9, 53–60,
788 <https://doi.org/10.1890/100014>, 2011.
- 789 Amann, R. I., Binder, B. J., Olson, R. J., Chisholm, S. W., Devereux, R., and Stahl, D. A.: Combination of 16S rRNA-
790 targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations, *Appl Environ*
791 *Microbiol.*, 56, 1919–1925, <https://doi.org/10.1128/aem.56.6.1919-1925.1990>, 1990.

792 Bandyopadhyay, J., Kraemer, D., Kattelman, R. and Kundzewicz, Z. W.: Highland waters: A resource of global
793 significance, in *Mountains of the World: A Global Priority*, edited by B. Messerli and J. Ives, pp. 131 – 155, Parthenon,
794 New York. 1997.

795 Battin, T. J., Louis A. Kaplan, Stuart Findlay, Charles S. Hopkinson, Eugenia Marti, Aaron I. Packman, J. Denis
796 Newbold, and Francesc Sabater: Biophysical controls on organic carbon fluxes in fluvial networks, *Nature Geoscience*,
797 1, 95–100, 2008.

798 Battin, T. J., Lauerwald, R., Bernhardt, E. S., Bertuzzo, E., Gener, L. G., Hall, R. O., Hotchkiss, E. R., Maavara, T.,
799 Pavelsky, T. M., Ran, L., Raymond, P., Rosentreter, J. A., and Regnier, P.: River ecosystem metabolism and carbon
800 biogeochemistry in a changing world, *Nature*, 613, 449–459, <https://doi.org/10.1038/s41586-022-05500-8>, 2023.

801 Bernhardt, E. S., Savoy, P., Vlah, M. J., Appling, A. P., Koenig, L. E., Hall, R. O., Arroita, M., Blaszcak, J. R., Carter,
802 A. M., Cohen, M., Harvey, J. W., Heffernan, J. B., Helton, A. M., Hosen, J. D., Kirk, L., McDowell, W. H., Stanley, E.
803 H., Yackulic, C. B., and Grimm, N. B.: Light and flow regimes regulate the metabolism of rivers, *Proc. Natl. Acad. Sci.*
804 U.S.A., 119, e2121976119, <https://doi.org/10.1073/pnas.2121976119>, 2022.

805 Bernal, S., Cohen, M. J., Ledesma, J. L. J., Kirk, L., Martí, E., and Lupon, A.: Stream metabolism sources a large
806 fraction of carbon dioxide to the atmosphere in two hydrologically contrasting headwater streams, *Limnol. Oceanogr.*,
807 67, 2621–2634, <https://doi.org/10.1002/lno.12226>, 2022.

808 Blagodatskaya, E. and Kuzyakov, Y.: Mechanisms of real and apparent priming effects and their dependence on soil
809 microbial biomass and community structure: critical review, *Biol. Fertil. Soils*, 45, 115–131,
810 <https://doi.org/10.1007/s00374-008-0334-y>, 2008.

811 Blaszcak, J. R., Delesantro, J. M., Urban, D. L., Doyle, M. W., and Bernhardt, E. S.: Scoured or suffocated: Urban
812 stream ecosystems oscillate between hydrologic and dissolved oxygen extremes, *Limnol. Oceanogr.*, 64, 877–894,
813 <https://doi.org/10.1002/lno.11081>, 2019.

814 Blattmann, T. M., Wang, S.-L., Lupker, M., Märki, L., Haghipour, N., Wacker, L., Chung, L.-H., Bernasconi, S. M.,
815 Plötze, M., and Eglinton, T. I.: Sulphuric acid-mediated weathering on Taiwan buffers geological atmospheric carbon
816 sinks, *Sci. Rep.*, 9, 2945, <https://doi.org/10.1038/s41598-019-39272-5>, 2019.

817 Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., Alexander, H., Alm, E. J.,
818 Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K., Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan,
819 B. J., Caraballo-Rodríguez, A. M., Chase, J., Cope, E. K., Da Silva, R., Diener, C., Dorrestein, P. C., Douglas, G. M.,
820 Durall, D. M., Duvallet, C., Edwardson, C. F., Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J. M., Gibbons, S. M.,
821 Gibson, D. L., Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S., Holste, H., Huttenhower, C., Huttley, G.
822 A., Janssen, S., Jarmusch, A. K., Jiang, L., Kaehler, B. D., Kang, K. B., Keefe, C. R., Keim, P., Kelley, S. T., Knights,
823 D., Koester, I., Kosciulek, T., Kreps, J., Langille, M. G. I., Lee, J., Ley, R., Liu, Y.-X., Loftfield, E., Lozupone, C.,
824 Maher, M., Marotz, C., Martin, B. D., McDonald, D., McIver, L. J., Melnik, A. V., Metcalf, J. L., Morgan, S. C.,
825 Morton, J. T., Naimey, A. T., Navas-Molina, J. A., Nothias, L. F., Orchanian, S. B., Pearson, T., Peoples, S. L., Petras,
826 D., Preuss, M. L., Priesse, E., Rasmussen, L. B., Rivers, A., Robeson, M. S., Rosenthal, P., Segata, N., Shaffer, M.,
827 Shiffer, A., Sinha, R., Song, S. J., Spear, J. R., Swafford, A. D., Thompson, L. R., Torres, P. J., Trinh, P., Tripathi, A.,
828 Turnbaugh, P. J., Ul-Hasan, S., Van Der Hooft, J. J. J., Vargas, F., Vázquez-Baeza, Y., Vogtmann, E., Von Hippel, M.,
829 et al.: Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2, *Nat Biotechnol*, 37,
830 852–857, <https://doi.org/10.1038/s41587-019-0209-9>, 2019.

831 Bosshard, P. P., Zbinden, R., and Altwegg, M.: *Turicibacter sanguinis* gen. nov., sp. nov., a novel anaerobic, Gram-
832 positive bacterium., *Int. J. Syst. Evol. Microbiol.*, 52, 1263–1266, <https://doi.org/10.1099/00207713-52-4-1263>, 2002.

833
834 Boysen, A. K., Durham, B. P., Kumler, W., Key, R. S., Heal, K. R., Carlson, L. T., Groussman, R. D., Armbrust, E. V.,
835 and Ingalls, A. E.: Glycine betaine uptake and metabolism in marine microbial communities, *Environmental*
836 *Microbiology*, 24, 2380–2403, <https://doi.org/10.1111/1462-2920.16020>, 2022.

837

838 Brailsford, F. L., Glanville, H. C., Golyshin, P. N., Johnes, P. J., Yates, C. A., and Jones, D. L.: Microbial uptake
839 kinetics of dissolved organic carbon (DOC) compound groups from river water and sediments, *Sci Rep*, 9, 11229,
840 <https://doi.org/10.1038/s41598-019-47749-6>, 2019.

841
842 Brunke, M. and Gonser, T.: The ecological significance of exchange processes between rivers and groundwater,
843 *Freshw. Biol.*, 37, 1–33, <https://doi.org/10.1046/j.1365-2427.1997.00143.x>, 1997.

844
845 Bufer, A., Hovius, N., Emberson, R., Rugenstein, J. K. C., Galy, A., Hassenruck-Gudipati, H. J., and Chang, J.-M.: Co-
846 variation of silicate, carbonate and sulfide weathering drives CO₂ release with erosion, *Nat. Geosci.*, 14, 211–216,
847 <https://doi.org/10.1038/s41561-021-00714-3>, 2021.

848
849 Butman, D. and Raymond, P. A.: Significant efflux of carbon dioxide from streams and rivers in the United States, *Nat. Geosci.*, 4, 839–842, <https://doi.org/10.1038/ngeo1294>, 2011.

850
851 Calmels, D., Galy, A., Hovius, N., Bickle, M., West, A. J., Chen, M.-C., and Chapman, H.: Contribution of deep
852 groundwater to the weathering budget in a rapidly eroding mountain belt, Taiwan, *Earth Planet. Sci. Lett.*, 303, 48–58,
<https://doi.org/10.1016/j.epsl.2010.12.032>, 2011.

853
854 Chao, A.: Estimating the Population Size for Capture-Recapture Data with Unequal Catchability, *Biometrics*, 43, 783–
791, <https://doi.org/10.2307/2531532>, 1987.

855
856 Chen, R., Senbayram, M., Blagodatsky, S., Myachina, O., Dittert, K., Lin, X., Blagodatskaya, E., and Kuzyakov, Y.:
857 Soil C and N availability determine the priming effect: microbial N mining and stoichiometric decomposition theories,
Glob. Change Biol., 20, 2356–2367, <https://doi.org/10.1111/gcb.12475>, 2014.

858
859 Chen, W.-S., Yeh, J.-J., and Syu, S.-J.: Late Cenozoic exhumation and erosion of the Taiwan orogenic belt: New
860 insights from petrographic analysis of foreland basin sediments and thermochronological dating on the metamorphic
orogenic wedge, *Tectonophysics*, 750, 56–69, <https://doi.org/10.1016/j.tecto.2018.09.003>, 2019.

861
862 Chiriboga, G. and Borges, A. V.: Andean headwater and piedmont streams are hot spots of carbon dioxide and methane
emissions in the Amazon basin, *Commun. Earth Environ.*, 4, 76, <https://doi.org/10.1038/s43247-023-00745-1>, 2023.

863
864 Cloern, J. E.: The relative importance of light and nutrient limitation of phytoplankton growth: a simple index of coastal
ecosystem sensitivity to nutrient enrichment, *Aquat. Ecol.*, 33, 3–16, 1999.

865
866 Cole, J. J., Prairie, Y. T., Caraco, N. F., McDowell, W. H., Tranvik, L. J., Striegl, R. G., Duarte, C. M., Kortelainen, P.,
867 Downing, J. A., Middelburg, J. J., and Melack, J.: Plumbing the Global Carbon Cycle: Integrating Inland Waters into
the Terrestrial Carbon Budget, *Ecosystems*, 10, 172–185, <https://doi.org/10.1007/s10021-006-9013-8>, 2007.

868
869 Cory, R. M., Ward, C. P., Crump, B. C., and Kling, G. W.: Sunlight controls water column processing of carbon in
arctic fresh waters, *Science*, 345, 925–928, <https://doi.org/10.1126/science.1253119>, 2014.

870
871 Dadson, S. J., Hovius, N., Chen, H., Dade, W. B., Hsieh, M.-L., Willett, S. D., Hu, J.-C., Horng, M.-J., Chen, M.-C.,
872 Stark, C. P., Lague, D., and Lin, J.-C.: Links between erosion, runoff variability and seismicity in the Taiwan orogen,
Nature, 426, 648–651, <https://doi.org/10.1038/nature02150>, 2003.

873
874 Dodds, W. K., Veach, A. M., Ruffing, C. M., Larson, D. M., Fischer, J. L., and Costigan, K. H.: Abiotic controls and
875 temporal variability of river metabolism: multiyear analyses of Mississippi and Chattahoochee River data, *Freshw. Sci.*,
32, 1073–1087, <https://doi.org/10.1899/13-018.1>, 2013.

876
877 Domingues, R. B., Barbosa, A., and Galvão, H.: Nutrients, light and phytoplankton succession in a temperate estuary
878 (the Guadiana, south-western Iberia), *Estuar. Coast. Shelf Sci.*, 64, 249–260, <https://doi.org/10.1016/j.ecss.2005.02.017>,
2005.

879
880 Dronsella, B., Orsi, E., Schulz-Mirbach, H., Benito-Vaquerizo, S., Yilmaz, S., Glatter, T., Bar-Even, A., Erb, T. J., and
881 Claassens, N. J.: One-carbon fixation via the synthetic reductive glycine pathway exceeds yield of the Calvin cycle, *Nat Microbiol*, 10, 646–653, <https://doi.org/10.1038/s41564-025-01941-9>, 2025.

882 Duvert, C., Butman, D. E., Marx, A., Ribolzi, O., and Hutley, L. B.: CO₂ evasion along streams driven by groundwater
883 inputs and geomorphic controls, *Nat. Geosci.*, 11, 813–818, <https://doi.org/10.1038/s41561-018-0245-y>, 2018.

884 Emberson, R., Hovius, N., Galy, A., and Marc, O.: Chemical weathering in active mountain belts controlled by
885 stochastic bedrock landsliding, *Nat. Geosci.*, 9, 42–45, <https://doi.org/10.1038/ngeo2600>, 2016.

886 Emerson, D., Fleming, E. J., and McBeth, J. M.: Iron-Oxidizing Bacteria: An Environmental and Genomic Perspective,
887 *Annu. Rev. Microbiol.*, 64, 561–583, <https://doi.org/10.1146/annurev.micro.112408.134208>, 2010.

888 Emerson, D., Field, E. K., Chertkov, O., Davenport, K. W., Goodwin, L., Munk, C., Nolan, M., and Woyke, T.:
889 Comparative genomics of freshwater Fe-oxidizing bacteria: implications for physiology, ecology, and systematics,
890 *Front. Microbiol.*, 4, <https://doi.org/10.3389/fmicb.2013.00254>, 2013.

891 Geisseler, D. and Horwath, W. R.: Investigating amino acid utilization by soil microorganisms using compound specific
892 stable isotope analysis, *Soil Biol. Biochem.*, 74, 100–105, <https://doi.org/10.1016/j.soilbio.2014.02.024>, 2014.

893 González, I., Déjean, S., Martin, P., and Baccini, A.: **CCA** : An R Package to Extend Canonical Correlation Analysis, *J.*
894 *Stat. Softw.*, 23, <https://doi.org/10.18637/jss.v023.i12>, 2008.

895 Grabovich, M. Yu., Ravin, N. V., and Boden, R.: *Thiothrix*, in: *Bergey's Manual of Systematics of Archaea and*
896 *Bacteria*, edited by: Whitman, W. B., Wiley, 1–21, <https://doi.org/10.1002/9781118960608.gbm01229.pub2>, 2023.

897 Hall, R. O., Yackulic, C. B., Kennedy, T. A., Yard, M. D., Rosi-Marshall, E. J., Voichick, N., and Behn, K. E.:
898 Turbidity, light, temperature, and hydropeaking control primary productivity in the Colorado River, Grand Canyon:
899 Semimechanistic modeling of daily GPP, *Limnol. Oceanogr.*, 60, 512–526, <https://doi.org/10.1002/lno.10031>, 2015.

900 Hill, P. G., Warwick, P. E., and Zubkov, M. V.: Low microbial respiration of leucine at ambient oceanic concentration
901 in the mixed layer of the central Atlantic Ocean, *Limnology & Oceanography*, 58, 1597–1604,
902 <https://doi.org/10.4319/lo.2013.58.5.1597>, 2013.

903 Hilton, R. G. and West, A. J.: Mountains, erosion and the carbon cycle, *Nat. Rev. Earth Environ.*, 1, 284–299,
904 <https://doi.org/10.1038/s43017-020-0058-6>, 2020.

905 Hilton, R. G., Galy, A., Hovius, N., Chen, M.-C., Horng, M.-J., and Chen, H.: Tropical-cyclone-driven erosion of the
906 terrestrial biosphere from mountains, *Nat. Geosci.*, 1, 759–762, <https://doi.org/10.1038/ngeo333>, 2008.

907 Hinrichs, K.-U., Hayes, J. M., Sylva, S. P., Brewer, P. G., and DeLong, E. F.: Methane-consuming archaeobacteria in
908 marine sediments, *Nature*, 398, 802–805, <https://doi.org/10.1038/19751>, 1999.

909 Horgby, Å., Segatto, P. L., Bertuzzo, E., Lauerwald, R., Lehner, B., Ulseth, A. J., Vennemann, T. W., and Battin, T. J.:
910 Unexpected large evasion fluxes of carbon dioxide from turbulent streams draining the world's mountains, *Nat.*
911 *Commun.*, 10, 4888, <https://doi.org/10.1038/s41467-019-12905-z>, 2019.

912 Hotchkiss, E. R., Hall Jr, R. O., Sponseller, R. A., Butman, D., Klaminder, J., Laudon, H., Rosvall, M., and Karlsson, J.:
913 Sources of and processes controlling CO₂ emissions change with the size of streams and rivers, *Nat. Geosci.*, 8, 696–
914 699, <https://doi.org/10.1038/ngeo2507>, 2015.

915 Huber, H., Hohn, M. J., Rachel, R., Fuchs, T., Wimmer, V. C., and Stetter, K. O.: A new phylum of Archaea
916 represented by a nanosized hyperthermophilic symbiont, *Nature*, 417, 63–67, <https://doi.org/10.1038/417063a>, 2002.

917 Ives, J.D., Messerli, B. and Spiess, E., Introduction. In: *Mountains of the World: A Global Priority*, B. Messerli and J.D.
918 Ives (Eds.), Parthenon, New York and London, 115, 1997.

919 Jähne, B., Heinz, G., and Dietrich, W.: Measurement of the diffusion coefficients of sparingly soluble gases in water, *J.*
920 *Geophys. Res. Oceans*, 92, 10767–10776, <https://doi.org/10.1029/JC092iC10p10767>, 1987.

921 Jørgensen, N. O. G. and Søndergaard, M.: Are dissolved free amino acids free?, *Microb. Ecol.*, 10, 301-316,
922 <https://link.springer.com/article/10.1007/BF02015556>, 1984.

923 Kao, S. J. and Milliman, J. D.: Water and Sediment Discharge from Small Mountainous Rivers, Taiwan: The Roles of
924 Lithology, Episodic Events, and Human Activities, *J. Geol.*, 116, 431–448, <https://doi.org/10.1086/590921>, 2008.

925 Kemeny, P. C., Lopez, G. I., Dalleska, N. F., Torres, M., Burke, A., Bhatt, M. P., West, A. J., Hartmann, J., and Adkins,
926 J. F.: Sulfate sulfur isotopes and major ion chemistry reveal that pyrite oxidation counteracts CO₂ drawdown from
927 silicate weathering in the Langtang-Trisuli-Narayani River system, Nepal Himalaya, *Geochim. Cosmochim. Acta*, 294,
928 43–69, <https://doi.org/10.1016/j.gca.2020.11.009>, 2021.

929 Kikuchi, G., Motokawa, Y., Yoshida, T., and Hiraga, K.: Glycine cleavage system: reaction mechanism, physiological
930 significance, and hyperglycinemia, *Proc. Jpn. Acad., Ser. B*, 84, 246–263, <https://doi.org/10.2183/pjab.84.246>, 2008.

931 Kirchman, D.: Measuring bacterial biomass production and growth rates from leucine incorporation in natural aquatic
932 environments, in: *Methods in Microbiology*, vol. 30, Elsevier, 227–237, [https://doi.org/10.1016/S0580-9517\(01\)30047-](https://doi.org/10.1016/S0580-9517(01)30047-8)
933 8, 2001.

934 Kirchman, D., K’nees, E., and Hodson, R.: Leucine incorporation and its potential as a measure of protein synthesis by
935 bacteria in natural aquatic systems, *Appl Environ Microbiol.*, 49, 599–607, [https://doi.org/10.1128/aem.49.3.599-](https://doi.org/10.1128/aem.49.3.599-607.1985)
936 607.1985, 1985.

937 Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., and Schloss, P. D.: Development of a Dual-Index
938 Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence Data on the MiSeq Illumina Sequencing
939 Platform, *Appl. Environ. Microbiol.*, 79, 5112–5120, <https://doi.org/10.1128/AEM.01043-13>, 2013.

940 Lane, D.J., 16S/23S rRNA sequencing. *Nucleic acid techniques in bacterial systematics*. E. Stackebrandt and M.
941 Goodfellow, eds. New York, NY, John Wiley and Sons: 115-175, 1991.

942 Lauerwald, R., Laruelle, G. G., Hartmann, J., Ciais, P., and Regnier, P. A. G.: Spatial patterns in CO₂ evasion from the
943 global river network, *Glob. Biogeochem. Cycles*, 29, 534–554, <https://doi.org/10.1002/2014GB004941>, 2015.

944 Lewis, E., and Wallace, D.: Program developed for CO₂ system calculations. Ornl/Cdiac-105 1–21, 1998.

945 Lien, W.-Y., Chen, C.-T., Lee, Y.-H., Su, C.-C., Wang, P.-L., and Lin, L.-H.: Two-stage oxidation of petrogenic
946 organic carbon in a rapidly exhuming small mountainous catchment, *Commun. Earth Environ.*, 6, 45,
947 <https://doi.org/10.1038/s43247-025-02015-8>, 2025.

948 Lin, B., Liu, Z., Eglinton, T. I., Kandasamy, S., Blattmann, T. M., Haghipour, N., Huang, K.-F., and You, C.-F.: Island-
949 wide variation in provenance of riverine sedimentary organic carbon: A case study from Taiwan, *Earth Planet. Sci.*
950 *Lett.*, 539, 116238, <https://doi.org/10.1016/j.epsl.2020.116238>, 2020.

951 Lupon, A., Denfeld, B. A., Laudon, H., Leach, J., Karlsson, J., and Sponseller, R. A.: Groundwater inflows control
952 patterns and sources of greenhouse gas emissions from streams, *Limnol. Oceanogr.*, 64, 1545–1557,
953 <https://doi.org/10.1002/lno.11134>, 2019.

954 Maavara, T., Yuan, Z., Johnson, A. M., Zhang, S., Aho, K. S., Brinkerhoff, C. B., Logozzo, L. A., and Raymond, P.:
955 River metabolism in the contiguous United States: A West of extremes, *Science*, 390, 622–627, 2025.

956 Marx, A., Dusek, J., Jankovec, J., Sanda, M., Vogel, T., Van Geldern, R., Hartmann, J., and Barth, J. A. C.: A review of
957 CO₂ and associated carbon dynamics in headwater streams: A global perspective, *Rev. Geophys.*, 55, 560–585,
958 <https://doi.org/10.1002/2016RG000547>, 2017.

959 Massey, L. K., Sokatch, J. R., and Conrad, R. S.: Branched-Chain Amino Acid Catabolism in Bacteria, *BACTERIOL.*
960 *REV.*, 40, 1976.

961 McInerney, M. J., Hoehler, T., Gunsalus, R. P., and Schink, B.: Introduction to Microbial Hydrocarbon Production:
 962 Bioenergetics, in: Handbook of Hydrocarbon and Lipid Microbiology, edited by: Timmis, K. N., Springer Berlin
 963 Heidelberg, Berlin, Heidelberg, 319–335, https://doi.org/10.1007/978-3-540-77587-4_21, 2010.

964 McMillan, H. K. and Srinivasan, M. S.: Characteristics and controls of variability in soil moisture and groundwater in a
 965 headwater catchment, Hydrol. Earth Syst. Sci., 19, 1767–1786, <https://doi.org/10.5194/hess-19-1767-2015>, 2015.

966 Meybeck, M., Green, P., and Vörösmarty, C.: A New Typology for Mountains and Other Relief Classes: An
 967 Application to Global Continental Water Resources and Population Distribution, Mt. Res. Dev., 21, 34–45,
 968 [https://doi.org/10.1659/0276-4741\(2001\)021%255B0034:ANTFMA%255D2.0.CO;2](https://doi.org/10.1659/0276-4741(2001)021%255B0034:ANTFMA%255D2.0.CO;2), 2001.

969 Miles, A. M., Chen, Y., Owens, M. W. and Grisham, M. B.: Fluorometric determination of nitric oxide. Methods A
 970 Companion to Methods Enzymol. 7: 40–47. doi:10.1006/meth.1995.1006, 1995.

971 O'Donnell, B. and Hotchkiss, E. R.: Resistance and resilience of stream metabolism to high flow disturbances,
 972 Biogeosciences, 19, 1111–1134, <https://doi.org/10.5194/bg-19-1111-2022>, 2022.

973 Overholt, W. A., Trumbore, S., Xu, X., Bornemann, T. L. V., Probst, A. J., Krüger, M., Herrmann, M., Thamdrup, B.,
 974 Bristow, L. A., Taubert, M., Schwab, V. F., Hölzer, M., Marz, M., and Küsel, K.: Carbon fixation rates in groundwater
 975 similar to those in oligotrophic marine systems, Nat. Geosci., 15, 561–567, [https://doi.org/10.1038/s41561-022-00968-](https://doi.org/10.1038/s41561-022-00968-5)
 976 5, 2022.

977 Palenik, B. and Morel, F.: Comparison of cell-surface L-amino acid oxidases from several marine phytoplankton, Mar.
 978 Ecol. Prog. Ser., 59, 195–201, <https://doi.org/10.3354/meps059195>, 1990.

979 Petit, S., Righi, D. and Madejová, J.: Infrared spectroscopy of NH₄⁺-bearing and saturated clay minerals: A review of
 980 the study of layer charge. Appl. Clay Sci. 34: 22–30. doi:10.1016/j.clay.2006.02.007, 2006

981 Quast, C., Priesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O.: The SILVA
 982 ribosomal RNA gene database project: improved data processing and web-based tools, Nucleic Acids Res., 41, D590–
 983 D596, <https://doi.org/10.1093/nar/gks1219>, 2012.

984 Rasilo, T., Hutchins, R. H. S., Ruiz-González, C., and Del Giorgio, P. A.: Transport and transformation of soil-derived
 985 CO₂, CH₄ and DOC sustain CO₂ supersaturation in small boreal streams, Sci. Total Environ., 579, 902–912,
 986 <https://doi.org/10.1016/j.scitotenv.2016.10.187>, 2017.

987 Ravin, N. V., Rudenko, T. S., Smolyakov, D. D., Beletsky, A. V., Rakitin, A. L., Markov, N. D., Fomenkov, A., Sun,
 988 L., Roberts, R. J., Novikov, A. A., Karnachuk, O. V., and Grabovich, M. Y.: Comparative Genome Analysis of the
 989 Genus *Thiothrix* Involving Three Novel Species, *Thiothrix subterranea* sp. nov. Ku-5, *Thiothrix litoralis* sp. nov. AS
 990 and “*Candidatus Thiothrix anitrata*” sp. nov. A52, Revealed the Conservation of the Pathways of Dissimilatory Sulfur
 991 Metabolism and Variations in the Genetic Inventory for Nitrogen Metabolism and Autotrophic Carbon Fixation, Front.
 992 Microbiol., 12, 760289, <https://doi.org/10.3389/fmicb.2021.760289>, 2021.

993 Raymond, P. A., Hartmann, J., Lauerwald, R., Sobek, S., McDonald, C., Hoover, M., Butman, D., Striegl, R., Mayorga,
 994 E., Humborg, C., Kortelainen, P., Dürr, H., Meybeck, M., Ciais, P., and Guth, P.: Global carbon dioxide emissions from
 995 inland waters, Nature, 503, 355–359, <https://doi.org/10.1038/nature12760>, 2013.

996 Raymond, P. A. and Spencer, R. G. M.: Riverine DOM, in: Biogeochemistry of Marine Dissolved Organic Matter,
 997 Elsevier, 509–533, <https://doi.org/10.1016/B978-0-12-405940-5.00011-X>, 2015.

998 Roche, K., Dentz, M., Samolis, R., Hixson, J. L., González-Pinzón, R., Herzog, S. P., Kim, J., Ward, A. S., and Bolster,
 999 D.: Quantifying benthic biolayer contributions to whole-stream reactivity by unifying surface and subsurface
 1000 observations: field evidence and theory, <https://doi.org/10.22541/essoar.173170006.67575057/v1>, 2025.

1001 Rovelli, L., Attard, K. M., Binley, A., Heppell, C. M., Stahl, H., Trimmer, M., and Glud, R. N.: Reach-scale river
1002 metabolism across contrasting sub-catchment geologies: Effect of light and hydrology, *Limnology & Oceanography*,
1003 62, <https://doi.org/10.1002/lno.10619>, 2017.

1004 Saccardi, B. and Winnick, M.: Improving Predictions of Stream CO₂ Concentrations and Fluxes Using a Stream
1005 Network Model: A Case Study in the East River Watershed, CO, USA, *Glob. Biogeochem. Cycles*, 35,
1006 e2021GB006972, <https://doi.org/10.1029/2021GB006972>, 2021.

1007 Schelker, J., Singer, G. A., Ulseth, A. J., Hengsberger, S., and Battin, T. J.: CO₂ evasion from a steep, high gradient
1008 stream network: importance of seasonal and diurnal variation in aquatic pCO₂ and gas transfer: CO₂ evasion from a
1009 steep, high gradient stream network, *Limnol. Oceanogr.*, 61, 1826–1838, <https://doi.org/10.1002/lno.10339>, 2016.

1010 Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., Lesniewski, R. A., Oakley, B. B.,
1011 Parks, D. H., Robinson, C. J., Sahl, J. W., Stres, B., Thallinger, G. G., Van Horn, D. J., and Weber, C. F.: Introducing
1012 mothur: Open-Source, Platform-Independent, Community-Supported Software for Describing and Comparing
1013 Microbial Communities, *Appl. Environ. Microbiol.*, 75, 7537–7541, <https://doi.org/10.1128/AEM.01541-09>, 2009.

1014 Selesi, D., Pattis, I., Schmid, M., Kandeler, E., and Hartmann, A.: Quantification of bacterial RubisCO genes in soils by
1015 cbbL targeted real-time PCR, *Journal of Microbiological Methods*, 69, 497–503,
1016 <https://doi.org/10.1016/j.mimet.2007.03.002>, 2007.

1017 Shannon, C. E., and Weaver, W.: The mathematical theory of communication., University of Illinois Press, 1949.

1018 Stanley, E. H., Powers, S. M., and Lottig, N. R.: The evolving legacy of disturbance in stream ecology: concepts,
1019 contributions, and coming challenges, *J. North Am. Benthol. Soc.*, 29, 67–83, <https://doi.org/10.1899/08-027.1>, 2010.

1020 Stanley R.S., Hill L.B., Chang H.C., and Hu H.N.: A transect through the metamorphic core of the Central Mountains,
1021 southern Taiwan, in: *Memoir of the Geological Society of China (Taiwan)*, vol. 4, 443–473, 1981.

1022 Suttle, C. A., Chan, A. M., and Fuhrman, J. A.: Dissolved free amino acids in the Sargasso Sea: uptake and respiration
1023 rates, turnover times, and concentrations, *Mar. Ecol. Prog. Ser.*, 70, 189–199, <https://doi.org/10.3354/meps070189>,
1024 1991.

1025 Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G.,
1026 Chavez, F., Sabine, C., Watson, A., Bakker, D. C. E., Schuster, U., Metzl, N., Yoshikawa-Inoue, H., Ishii, M.,
1027 Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T. S., Tilbrook, B.,
1028 Johannessen, T., Olsen, A., Bellerby, R., Wong, C. S., Delille, B., Bates, N. R., and De Baar, H. J. W.: Climatological
1029 mean and decadal change in surface ocean pCO₂, and net sea–air CO₂ flux over the global oceans, *Deep Sea Res. Part II*
1030 *Top. Stud. Oceanogr.*, 56, 554–577, <https://doi.org/10.1016/j.dsr2.2008.12.009>, 2009.

1031 Takai, K. and Horikoshi, K.: Rapid Detection and Quantification of Members of the Archaeal Community by
1032 Quantitative PCR Using Fluorogenic Probes, *Appl Environ Microbiol*, 66, 5066–5072,
1033 <https://doi.org/10.1128/AEM.66.11.5066-5072.2000>, 2000.

1034 Teng, L. S.: Geotectonic evolution of late Cenozoic arc-continent collision in Taiwan, *Tectonophysics*, 183, 57–76,
1035 [https://doi.org/10.1016/0040-1951\(90\)90188-E](https://doi.org/10.1016/0040-1951(90)90188-E), 1990.

1036 Tu, T.-H., Wu, L.-W., Lin, Y.-S., Imachi, H., Lin, L.-H., and Wang, P.-L.: Microbial Community Composition and
1037 Functional Capacity in a Terrestrial Ferruginous, Sulfate-Depleted Mud Volcano, *Front. Microbiol.*, 8, 2137,
1038 <https://doi.org/10.3389/fmicb.2017.02137>, 2017.

1039 Tu, T.-H., Chen, L.-L., Chiu, Y.-P., Lin, L.-H., Wu, L.-W., Italiano, F., Shyu, J. B. H., Raisossadat, S. N., and Wang,
1040 P.-L.: The biogeographic pattern of microbial communities inhabiting terrestrial mud volcanoes across the Eurasian
1041 continent, *Biogeosciences*, 19, 831–843, <https://doi.org/10.5194/bg-19-831-2022>, 2022.

- 1042 Ulseth, A. J., Hall, R. O., Boix Canadell, M., Madinger, H. L., Niayifar, A., and Battin, T. J.: Distinct air–water gas
1043 exchange regimes in low- and high-energy streams, *Nat. Geosci.*, 12, 259–263, [https://doi.org/10.1038/s41561-019-](https://doi.org/10.1038/s41561-019-0324-8)
1044 0324-8, 2019.
- 1045 Vannote, R. L., Minshall, G. W., Cummins, K. W., Sedell, J. R., Cushing, C. E.: The River Continuum Concept, *Can. J.*
1046 *Fish. Aquat. Sci.*, 37, 130-137, <http://doi.org/10.1139/f80-017>, 1980.
- 1047 Viviroli, D. and Weingartner, R.: The hydrological significance of mountains: from regional to global scale, *Hydrol.*
1048 *Earth Syst. Sci.*, 8, 1017–1030, <https://doi.org/10.5194/hess-8-1017-2004>, 2004.
- 1049 Wang, P.-L., Tu, T.-H., Lin, L.-H., Chou, H.-L., Wang, Y.-J., Chen, J.-N., Wang, L.-Y., Chang, J.-M., Chu, M.-F., Hsu,
1050 Y.-C., Chang, C.-P., Wu, Y.-M., Lin, Y.-T., and Ke, C.-C.: Microbial communities modulate chemical weathering and
1051 carbon dioxide cycling in an active orogen in Taiwan, *Commun. Earth Environ.*, 5, 174, [https://doi.org/10.1038/s43247-](https://doi.org/10.1038/s43247-024-01345-3)
1052 024-01345-3, 2024.
- 1053 Wang, Y., Tian, R. M., Gao, Z. M., Bougouffa, S., and Qian, P. Y.: Optimal eukaryotic 18S and universal 16S/18S
1054 ribosomal RNA primers and their application in a study of symbiosis, *PLoS ONE*, 9,
1055 <https://doi.org/10.1371/journal.pone.0090053>, 2014.
- 1056 Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnol. Oceanogr.*
1057 *Methods*, 12, 351–362, <https://doi.org/10.4319/lom.2014.12.351>, 2014.
- 1058 Watanabe, K., Komatsu, N., Kitamura, T., Ishii, Y., Park, H., Miyata, R., Noda, N., Sekiguchi, Y., Satou, T., Watanabe,
1059 M., Yamamura, S., Imai, A., and Hayashi, S.: Ecological niche separation in the *Polynucleobacter* subclusters linked to
1060 quality of dissolved organic matter: a demonstration using a high sensitivity cultivation-based approach, *Environ.*
1061 *Microbiol.*, 14, 2511–2525, <https://doi.org/10.1111/j.1462-2920.2012.02815.x>, 2012.
- 1062 Woyke, T., Chertkov, O., Lapidus, A., Nolan, M., Lucas, S., Del Rio, T. G., Tice, H., Cheng, J.-F., Tapia, R., Han, C.,
1063 Goodwin, L., Pitluck, S., Liolios, K., Pagani, I., Ivanova, N., Huntemann, M., Mavromatis, K., Mikhailova, N., Pati, A.,
1064 Chen, A., Palaniappan, K., Land, M., Hauser, L., Brambilla, E.-M., Rohde, M., Mwirichia, R., Sikorski, J., Tindall, B.
1065 J., Göker, M., Bristow, J., Eisen, J. A., Markowitz, V., Hugenholtz, P., Klenk, H.-P., and Kyrpides, N. C.: Complete
1066 genome sequence of the gliding freshwater bacterium *Fluviicola taffensis* type strain (RW262T), *Stand. Genomic Sci.*,
1067 5, 21–29, <https://doi.org/10.4056/sigs.2124912>, 2011.
- 1068 Yu, A. J., Lin, X., Zhu, J., He, H., and Li, L.: Environmental effects on ammonium adsorption onto clay minerals:
1069 Experimental constraints and applications, *Appl. Clay Sci.*, 246, 107165, <https://doi.org/10.1016/j.clay.2023.107165>,
1070 2023.