

Impacts of Free-Tropospheric Thermodynamics and Ice Nuclei Recycling on Arctic Mixed-Phase Cloud Properties

Benjamin Ascher^{1,2} and Fabian Hoffmann²

¹Meteorological Institute Munich, Ludwig Maximilian University, Munich, Germany

²Institute for Meteorology, Free University of Berlin, Berlin, Germany

Correspondence: Benjamin Ascher (ben.ascher@lmu.de) and Fabian Hoffmann (f.hoffmann@fu-berlin.de)

Abstract. Radiatively driven Arctic stratocumulus clouds have important climactic impacts due to their effects on surface radiative balance. The presence of both liquid and ice within Arctic stratocumulus, and their interaction through the Wegener-Bergeron-Findeisen (WBF) process, strongly affects the properties and lifetimes of these clouds. To assess the impacts of mixed-phase microphysical processes in Arctic Stratocumulus, we use a Lagrangian cloud microphysical model within a large eddy simulation framework to simulate a single-layer cloud under varying free-tropospheric humidity and above-cloud inversion strength. We also run two simulations in which precipitating ice crystals have their ice nuclei (IN) re-injected into the model domain, rather than removed. We find that IN recycling plays a critical role in maintaining the presence of ice in the mixed-phase cloud. The simulations with drier free-tropospheric air experience greater sublimation of ice crystals below cloud, recycling of ice crystals, and a higher ice water path than simulations with more humid free-tropospheric air. We also find that the impact of inversion strength on cloud microphysical characteristics is strongly modulated by free-tropospheric relative humidity, with decreased inversion strength resulting in both increased and decreased liquid water path under high and low free-tropospheric relative humidity, respectively.

1 Introduction

Stratocumulus clouds, weakly convective shallow cloud systems, occur frequently over large areas of the Earth's surface across many different latitudinal and climate regimes (Shi et al., 2018; Wood, 2012). These long-lived cloud fields play a critical role in the climate system, as their high albedo reflects a large portion of incoming solar radiation, resulting in a cooling effect during the day. Their impact on Earth's radiative balance, and investigation into how this impact may change due to anthropogenic climate and aerosol forcing, has been extensively studied (Caldwell and Bretherton, 2009; Tan and Storelvmo, 2019).

Compared to stratocumulus clouds in the subtropics, Arctic stratocumulus clouds have important impacts on the climate system not just as a result of their reflection of solar radiation back into space, but also from their emission of longwave radiation downward to the surface (Eirund et al., 2020; Garrett et al., 2002; Tan and Storelvmo, 2019; Tan et al., 2023). With the rapid rate of Arctic warming as a result of climate change, a better understanding of Arctic stratocumulus, and their influence on Arctic energy balance, is thus a research area of significant importance.

While subtropical stratocumulus clouds are entirely composed of liquid, Arctic stratocumulus clouds are complicated by the presence of ice (Mioche et al., 2017; McFarquhar et al., 2011; Shupe et al., 2006; Verlinde et al., 2007). Due to the lower equilibrium vapor pressure over ice compared to liquid at the same temperature, ice crystals within Arctic stratocumulus clouds grow much more rapidly than liquid cloud droplets, effectively competing for water vapor and reducing the liquid water path (LWP) of these clouds, an effect known as the Wegener-Bergeron-Findeisen (WBF) process (Harrington et al., 1999).

While many stratocumulus clouds in the Arctic are maintained by surface fluxes, particularly under Cold-Air Outbreak conditions, those Arctic stratocumulus clouds which form atop decoupled boundary layers are typically maintained by cloud-top radiative cooling, which is itself dependent on the LWP (de Boer et al., 2009; Griesche et al., 2021; Klein and Coauthors, 2009; McFarquhar et al., 2011; Sotiropoulou et al., 2014). Therefore, the reduction in LWP due to the WBF process, and concomitant reduction in cloud-top radiative cooling, can weaken the boundary-layer circulations which maintain the cloud. If enough ice is present, this can lead to a destructive feedback loop and the eventual dissipation of the cloud entirely. Even if the cloud persists, the reduction of LWP which is a consequence of the presence of ice alters the downward flux of longwave radiation to the surface beneath the cloud. The ice crystal number concentration, which strongly affects LWP and ice water path (IWP) within the clouds, is heavily dependent on the number of ice nuclei (IN) within the cloud. Larger IN concentrations have been shown both in modeling and observational studies to increase IWP at the expense of LWP, sometimes leading to complete glaciation and dissipation of the cloud (Ahola et al., 2020; Das et al., 2025; Ovchinnikov et al., 2014; Solomon et al., 2018).

Arctic stratocumulus clouds also often precipitate ice, which stabilizes the boundary layer by cooling as falling ice crystals sublimate below the cloud. This boundary layer stabilization, as well as the production of cold pools, plays an important role in determining the morphology and organization of Arctic stratocumulus clouds (Eirund et al., 2019; Sotiropoulou et al., 2014).

Due to the importance of ice processes within Arctic stratocumulus to the morphology and lifetimes of these clouds, as well as their radiative impacts on the surface, these clouds have been the subject of both modeling and observational studies (Ahola et al., 2020; Avramov and Harrington, 2010; de Boer et al., 2009; Eirund et al., 2019; Fan et al., 2009; Harrington et al., 1999; McFarquhar et al., 2007, 2011; Ovchinnikov et al., 2014; Paukert and Hoose, 2014; Raatikainen et al., 2022; Savre and Ekman, 2015; Shupe et al., 2006, 2011; Solomon et al., 2015, 2018; Verlinde et al., 2007). However, despite the insights gained from these projects, numerous unknowns remain regarding ice growth within Arctic stratocumulus and its effects on the cloud as a whole. While many processes affect the growth rate and number concentration of ice crystals in Arctic stratocumulus, including aggregation, riming, secondary ice production from ice-ice collisions, shattering of cloud droplets upon heterogeneous freezing, and rime splintering, we will focus primarily on ice growth through vapor deposition (Gayet et al., 2009-10-16; Jackson, 2012; Pasquier et al., 2022-12-12; Sotiropoulou et al., 2020). In particular, we will examine the roles of environmental humidity, entrainment strength, and IN recycling on vapor depositional growth of ice.

The role of entrainment and mixing on ice crystal growth within Arctic stratocumulus clouds is another outstanding source of uncertainty in the prediction of microphysical properties of these clouds. Arctic stratocumulus clouds typically occur in shallow boundary layers beneath a drier and warmer free atmosphere above (McFarquhar et al., 2007, 2011; Mioche et al., 2017; Shupe et al., 2006; Verlinde et al., 2007). At the top of the cloud, entrainment of this warm, dry air into the cloud can occur. As a result

of the WBF process, dry air entrainment in a mixed-phase cloud impacts liquid and ice particles quite differently. Hoffmann (2020) suggests that such dry air entrainment could enhance the growth of ice crystals, as rapid evaporation of cloud droplets exposed to subsaturated air provides a ready source of water vapor which can then be consumed by ice crystals. Both in-situ aircraft measurements and remote sensing of single-layer Arctic stratocumulus, however, consistently find a predominantly-liquid layer near cloud top, suggesting that WBF processes may be less important at cloud top than sedimentation of ice crystals and the production of supersaturation through longwave radiative cooling (de Boer et al., 2009; Griesche et al., 2021; McFarquhar et al., 2007, 2011; Mioche et al., 2017; Shupe et al., 2006; Verlinde et al., 2007). The rate of entrainment of air from the entrainment layer into the cloud is highly dependent on the magnitude of the temperature difference across the entrainment layer, with smaller temperature differences (weaker inversion strengths) being conducive to greater entrainment rates (Randall, 1984; Wood, 2012; Xu and Xue, 2015). The effects of entrainment on cloud microphysical properties are also strongly influenced by the relative humidity above the cloud, meaning that entrainment can alternately act to contribute to or deplete from the total water budget of Arctic stratocumulus clouds (Egerer et al., 2021; Solomon et al., 2014). How these different inversion strengths and relative humidities interact, particularly in the presence of ice, has not been well studied despite its potentially significant impacts on cloud properties and radiative balance.

The goal of this study is therefore to examine the following two science questions relating to microphysical processes within Arctic stratocumulus clouds. 1) How does the relative humidity above the cloud layer affect the microphysical properties within Arctic stratocumulus clouds? 2) How does the strength of the above-cloud temperature inversion affect the entrainment rate and microphysical processes within Arctic Stratocumulus clouds?

2 Methods

To answer the above questions, we chose to simulate a single-layer mixed-phase Arctic stratocumulus cloud field with the System for Atmospheric Modeling (Khairoutdinov and Randall, 2003), an anelastic, nonhydrostatic Large Eddy Simulation (LES) model. This model was run with a Lagrangian cloud microphysics model (Hoffmann et al., 2015), similar to the Super-Droplet Method (SDM) of Shima et al. (2009, 2020). In the SDM approach, aerosol particles and hydrometeors are represented as Super-Droplets (SDs), collections of many identical particles. A single gridbox may contain anywhere from dozens to hundreds of these SDs. Each SD has a weighting factor, indicating how many physical aerosol particles/hydrometeors are represented by the SD; extensive quantities, including total water mass, total aerosol mass, total ice nucleating particle (INP) surface area; and intensive quantities such as dimensions (if an ice crystal), freezing temperature, and time spent within ice- or liquid-supersaturated conditions. Each SD also has a unique ID assigned either at the start of the simulation or upon emission (if a particle emission source is present in the domain).

The growth of ice crystals follows the approach of Chen and Lamb (1994). This method treats ice crystals as spheroids with oblate and prolate axes (denoted as a - and c -axes, respectively). This parameterization uses an Inherent Growth Function (IGF) to determine which shape ice crystals growing by vapor deposition should develop. This IGF, based primarily on laboratory studies of ice crystal growth in free-fall chambers by Takahashi et al. (1991) are purely a function of temperature. This has the

consequence that, at a given temperature, two ice particles growing by vapor deposition under ice supersaturated conditions will develop the same habits (though at different rates, depending on the magnitude of ice supersaturation). This parameterization also calculates an effective deposition density and ventilation coefficient for ice crystals in an attempt to capture the impacts of
95 faceting and hollowing during vapor depositional growth and fall speed through vapor, respectively.

As we are interested in examining the behavior of ice crystals under vapor deposition with little impact from collisional processes, we chose to simulate a single-layer precipitating mixed-phase Arctic stratocumulus cloud observed on 26 April 2008 during Flight 31 of the Indirect and Semi-Direct Aerosol Campaign (ISDAC, McFarquhar et al. (2011)). This cloud was observed both by ground-level remote sensing instruments and in situ aircraft observations. Observations of this day indicated
100 that the majority of ice crystals were pristine, with little indications of riming or aggregation (Jackson, 2012). Additionally, the single-layer nature of the cloud precludes seeder-feeder effects and reduces the chances of ice-ice collisions. Drizzle was also absent, indicating a lack of collisional growth among cloud droplets (Ovchinnikov et al., 2014). Therefore, droplet collisional growth, riming, aggregation, and secondary ice production were disabled in our simulations. This particular cloud has been examined extensively by other modeling studies, making it a good case to simulate as our results can be compared with those
105 of previous work (Ahola et al., 2020; Das et al., 2025; Fu and Xue, 2017; Paukert and Hoose, 2014; Raatikainen et al., 2022; Savre and Ekman, 2015; Sulia and Kumjian, 2017).

Initial environmental conditions are identical to those used in Ovchinnikov et al. (2014) and depicted here in Skew-T format in Figure 1. The boundary layer is initially decoupled from the surface, as evidenced by the stable lapse rate between the surface and 400 m. Between 400–650 m, there is well-mixed (dry adiabatic) layer, while between 650 and 820 m, the atmosphere
110 follows a moist adiabatic lapse rate. We also initialize all simulations with condensate following an adiabatic LWP profile between 650–820 m (not shown). The maximum initial value of liquid water mixing ratio is 0.163 g kg^{-1} at 820 m. Zonal wind is initially set to -7 m s^{-1} throughout the domain, while meridional wind varies linearly with height from -2 m s^{-1} at the surface to 4.735 m s^{-1} at 2250 m (here, negative values for zonal and meridional wind indicate easterly and northerly winds, respectively). Initial temperatures are approximately -8.7°C at the surface, between -13°C and -15°C within the
115 cloud, and -10.5°C at the maximum of the above-cloud inversion.

All simulations use 10 meter vertical and 20 meter horizontal grid spacing, with 144 vertical levels and 64 grid points in each horizontal dimension. Horizontal boundaries are periodic, with a rigid model top at 1.44 km. Simulations are run with a 0.5 second timestep for 8 hours. Subgrid-scale turbulence is parameterized with a 1.5 order TKE-based closure, while radiation is parameterized using a Liquid Water Content (LWC)-profile approach as in Ovchinnikov et al. (2014). Surface energy fluxes
120 are neglected as in Ovchinnikov et al. (2014). Initial CCN number mixing ratio is 200 mg^{-1} ($\sim 267 \text{ cm}^{-3}$ at the lowest model level) and initial IN number concentration is 1 L^{-1} . To reduce computation time, we only add SDs up to 1200 m of altitude.

We do not permit ice formation at the start of the simulation, but rather restrict it to occur beginning one hour after the start of the simulation. This is to permit a radiatively driven cloud circulation and STBL to begin to form without competition from ice crystals for water vapor. After one hour, SDs containing IN which are in liquid-supersaturated conditions undergo probabilistic
125 immersion freezing. The probability of a given SD freezing is calculated using a parameterization based on the behavior of SNOWMAX bacteria (Wex et al., 2015). SNOWMAX bacteria become active INP at approximately -5°C and maintain




Figure 1. Initial thermodynamic conditions of the CONTROL simulation. Initial temperatures of the CONTROL and WEAKINV simulations are indicated by the red and maroon solid lines, respectively. Initial dewpoints of CONTROL and DRY are indicated in blue and light blue solid lines, respectively. Dry and moist adiabats are indicated by the red and blue dashed lines, respectively. Isolines of constant water vapor mixing ratio are depicted from 0.8 g kg^{-1} to 1.6 g kg^{-1} , at intervals of 0.2 g kg^{-1} , in the green dashed lines.

approximately the same ice nucleating activity between -10 and -20 °C. Because temperatures within the cloud during this case are between -13 and -15 °C, the use of a SNOWMAX parameterization for ice nucleating activity therefore ensures that most IN-containing particles freeze within only a couple of timesteps after they enter liquid-supersaturated conditions. Once a particle has nucleated, its SD type is instantly set to "ice" rather than "liquid". The ice nucleation parameterization used in this study takes no account of the amount of IN contained within an SD, but rather uses a simple "IN/No IN" binary classification for SDs. Future modeling work will explore the sensitivity of cloud microphysical evolution to different species or sizes of IN as described in Fu and Xue (2017).

To test the sensitivity of particle growth and cloud properties to both environmental conditions and microphysical parameterization, we present results from seven simulations. Initially, these were a CONTROL case using the same initial atmospheric profile as that used in Ovchinnikov et al. (2014), a DRY case in which the water vapor mixing ratio above 825 m was reduced from 1.2 to 0.8 g kg^{-1} , a WEAKINV case in which the inversion strength was reduced from 3.5 K to 1.5 K, and a DRY_WEAKINV case combining the weaker inversion strength of WEAKINV and the reduced above-cloud water vapor mix-

ing ratio of DRY. Note that in WEAKINV, above-cloud water vapor mixing ratio is unchanged, meaning that the above-cloud
140 relative humidity increases relative to CONTROL due to the reduced inversion strength.

During the course of our analysis, we also discovered that depletion of IN through particle sedimentation was an important factor in controlling both ice and liquid growth rates in all simulations. To examine the impact of perfectly-efficient IN recycling on cloud microphysical and boundary-layer evolution, we ran an additional two simulations, one with CONTROL environmental conditions and one with DRY conditions, in which any particle which reached the surface was re-injected into
145 the below-cloud boundary layer as a dry CCN/IN particle. Note that in this manuscript, the "efficiency" of IN recycling refers to the fraction of ice crystals which return their IN to the boundary layer through sublimation rather than precipitate to the surface. "Perfectly efficient" IN recycling thus indicates that all IN are returned to the boundary layer, rather than removed through precipitation. These simulations are denoted as INFLUX and DRY_INFLUX. ~~Finally, to~~ We should emphasize that the INFLUX and DRY_INFLUX simulations are not meant to represent a specific process such as surface emission of INP. Rather, they
150 serve as a counterfactual to the CONTROL and DRY simulations, allowing us to disentangle the effects of free-tropospheric thermodynamics from INP recycling on the cloud microphysical properties.

To assess the sensitivity of cloud properties to increased free-tropospheric IN number concentration, we conducted a simulation, HIGHIN, in which initial IN number concentrations above the inversion layer are increased by a factor of 5.6 (reflecting the high-IN conditions in Hiranuma et al. (2013)). The results of HIGHIN are discussed very briefly in section 3.2. Table 1
155 illustrates the full set of simulations.

We also ran one simulation to assess the sensitivity of our results to the model horizontal domain extent, though we do not discuss it in detail below. This simulation, denoted as LARGEDOMAIN, was run with 256 grid points in each horizontal dimension and 144 vertical levels. Vertical and horizontal grid spacing were the same as in CONTROL, meaning that the domain in LARGEDOMAIN was 4 times as wide as CONTROL along each horizontal dimension. Due to computational
160 restraints, we were only able to conduct this simulation for 5 hours. During this time, LWP IWP, cloud-top radiative cooling, and TKE were of similar magnitude in both CONTROL and LARGEDOMAIN (Fig. A1). Likewise, the horizontal extent of updrafts, downdrafts, and regions of ice water were also broadly consistent between CONTROL and LARGEDOMAIN (Fig. A2). This indicates that our use of a smaller domain for the other six simulations is unlikely to have significantly influenced our results.

165 3 Results

3.1 CONTROL

In the CONTROL simulation, a cloud is present from the start of the model run, but in an unrealistic state, with very little vertical motion. This initial state takes some time to spin up, as observed by the reduction in cloud liquid water mixing ratio at approximately time $t = 10$ minutes into the simulation. After $t = 30$ minutes, LWP steadily increases up to a value of
170 approximately 15.0 g m^{-2} at $t = 1$ hour (Fig. 2a).

Table 1. Simulation parameters. "Above-Cloud Water Vapor Mixing Ratio" refers to the above-cloud water [vapor](#) mixing ratio at initialization. "Inversion Strength" refers to the magnitude of the initial above-cloud temperature inversion. "IN Re-Injection" indicates whether IN-containing superparticles are removed from the simulation (None) or placed back in the simulation domain (Active) when an IN-containing superparticle reaches the surface.

Simulation Name	Above-Cloud Water Vapor Mixing Ratio (g kg^{-1})	Inversion Strength (K)	IN Re-Injection
CONTROL	1.2	3.5	None
DRY	0.8	3.5	None
WEAKINV	1.2	1.5	None
DRY_WEAKINV	0.8	1.5	None
INFLUX	1.2	3.5	Active
DRY_INFLUX	0.8	3.5	Active
HIGHIN	1.2	3.5	None




Figure 2. Time series of LWP (a) and IWP (b) in six simulations. The initiation of ice formation at $t = 1$ hour is indicated by the vertical dotted line.




Figure 3. As in Figure 2, but showing ice crystal number concentration averaged within cloud (defined as regions with liquid water mixing ratio $> 0.01 \text{ g kg}^{-1}$) (a) and vertically-integrated IN number concentration below cloud (b).

At 1 hour into the simulation, ice processes are allowed to begin occurring within the model. This results in the rapid nucleation of virtually every IN-containing superdroplet within the cloud and a peak ice crystal number concentration of approximately $\sim 1.2 \text{ L}^{-1}$ (Fig. 3a). The relatively high ice crystal number concentration and rapid growth of the ice crystals by vapor deposition leads to a classic WBF evolution and the net evaporation of cloud water, resulting in a decrease of LWP from $\sim 15.0 \text{ g m}^{-2}$ at $t = 1$ hour to $\sim 12.6 \text{ g m}^{-2}$ at $t = 1.7$ hours (Fig. 2a). Simultaneously, the IWP increases from 0.0 g m^{-2} at $t = 1$ hour to $\sim 7.6 \text{ g m}^{-2}$ at $t = 1.7$ hours (Fig. 2b).

As the ice crystals grow larger, they begin to sediment out of the cloud, resulting in a rapid decrease in ice crystal number concentration within the cloud. Indeed, by $t = 2.3$ hours, the average ice crystal number concentration within the cloud has plummeted to only $\sim 0.2 \text{ L}^{-1}$ (Fig. 3a), a value which is somewhat below the observed ice crystal number concentration of $\sim 0.35 \text{ L}^{-1}$ (Savre and Ekman, 2015). The depletion of ice crystal number concentration within the cloud is discussed in more detail below.

Though each individual ice crystal grows much more rapidly than the cloud droplets, the reduction in ice crystal number concentration means that the total rate of vapor deposition within the cloud decreases. So much so, in fact, that it decreases below the production of supersaturation through radiative and adiabatic cooling within the cloud, allowing the liquid cloud droplets to begin growing once more. After reaching a minimum at $t = 1.7$ hours, the LWP steadily increases throughout the rest of the simulation, reaching $\sim 35.8 \text{ g m}^{-2}$ at $t = 8$ hours (Fig. 2a).




Figure 4. As in Figure 2, but displaying average radiative cooling rates within the uppermost 50 m of the cloud (a) and the vertically integrated turbulent kinetic energy up to cloud top (b).

This increase in cloud water is aided by a positive feedback loop within the cloud; liquid condensate near the cloud top emits infrared (longwave) radiation, leading to strong longwave cooling and the formation of a strong downdraft region of negatively buoyant air. This sinking air within this downdraft region, by conservation of momentum, leads to a surrounding region of ascent and supersaturation. Within this ascending, supersaturated region, even more liquid water content is produced, leading to greater longwave cooling at cloud top, and a stronger sinking motion within the downdraft (Petters et al., 2012). Cloud-top radiative cooling rates increase from $\sim 55 \text{ K day}^{-1}$ at $t = 1.7$ hours to $\sim 80 \text{ K day}^{-1}$ at $t = 8$ hours (Fig 4a), while the vertically integrated Turbulent Kinetic Energy (TKE), hereafter referred to as the TKE path, increases from $\sim 80 \text{ m}^3 \text{ s}^{-2}$ at $t = 1.7$ hours to $\sim 160 \text{ m}^3 \text{ s}^{-2}$ by $t = 8$ hours (Fig. 4b). This increase in TKE path is a consequence not just of the strengthening of vertical motion and mixing within the cloud, but a deepening of the subcloud mixed layer. Importantly, because water vapor mixing ratios are higher in the decoupled near-surface layer, this means that the expanding below-cloud mixed layer entrains greater and greater quantities of water vapor, also increasing the amount of liquid water within the cloud. At approximately 6.5 hours into the simulation, the mixed layer reaches the surface (not shown), meaning that the entire atmosphere up to the cloud top is well-mixed. The cloud itself also deepens during this time; though cloud top height remains nearly constant due to cloud-top entrainment balancing large-scale subsidence, the cloud base sinks from $\sim 659 \text{ m}$ at $t = 2$ hours to $\sim 551 \text{ m}$ at $t = 8$ hours (Figs. 5, 6a).




Figure 5. Time-series of cloud-top entrainment rate (a) and cloud depth (b) for six simulations. Cloud-top entrainment rate is calculated the rate of change of cloud top height minus the value of large-scale subsidence. Ice initiation at 1 hour is denoted by the vertical black dotted line.

The average size of ice crystals within the cloud increases throughout the simulation, from $\sim 500 \mu\text{m}$ at $t = 2$ hours to $\sim 700 \mu\text{m}$ at $t = 8$ hours (Fig. 6). The temperatures within the cloud fall between -13°C and -15°C , resulting in predominantly plate-like growth of the crystals (Fig. A1A3). As mentioned briefly above, these large ice crystals fall through and below the cloud. Due to the lower equilibrium water vapor pressure over ice compared to liquid, the ice crystals continue to grow by vapor deposition even slightly below the cloud base, down to ~ 450 m at $t = 1$ hour. Below this altitude, however, the relative humidity with respect to ice falls below 100%, resulting in sublimation. This sublimation means that the number of ice particles reaching the surface is only $\sim 59.2\%$ of those which are present at the lowest ice-supersaturated altitude, though this proportion has considerable variability with time (Fig. A2A4).

This sublimation of ice crystals below cloud base is extremely important to the microphysical properties within the cloud itself. In the CONTROL simulation, there is no source of IN emission from the surface, while ice particles which reach the surface without sublimating are removed from the model domain, taking their IN with them. In contrast, ice particles which completely sublimate at some point below cloud base become dry IN. (We will refer to this complete sublimation, where a particle’s model classification changes from that of an ice crystal to that of a dry IN particle, as “desiccation”.) This results in two effects: an overall depletion of IN from the domain, and a buildup of recycled IN below cloud base where ice crystals have desiccated.

A “band” of elevated IN concentrations below cloud base emerges approximately 1.5 hours into the simulation, initially at an altitude of ~ 250 m. Note that this band does not form exactly where the majority of ice crystals desiccate, but rather




Figure 6. A time-height comparison of mean ice crystal radius (measured along the maximum particle dimension). (a,b) display radius in blue-purple shading, while (c) shows the difference in radius between DRY and CONTROL in red and blue colors. Cloud boundaries are shown in solid black lines in (a) and (b). Ice initiation at 1 hour is denoted by the vertical black dotted line.

slightly below (Fig. 7). This is because the majority of ice crystals desiccate within the well-mixed region beneath cloud base, meaning that their IN are quickly spread throughout the mixed layer. The ice crystals which desiccate below the mixed layer, on the other hand, release their INs into a stable environment where they remain concentrated within a relatively narrow altitude range (the “band” visible in Fig. 7). As the mixed-layer deepens toward the surface, the altitude of this band sinks over time. At approximately 6.5 hours into the simulation, the “band” disappears as it intersects the surface. This indicates that there is no longer a stable layer which desiccating ice particles can release their IN into. Instead, IN from desiccated ice crystals spread vertically throughout the mixed layer and into the cloud.

Nucleation rates at the cloud top are more than an order of magnitude less than those which occur at cloud base (Fig. 8a). Because the CONTROL simulation lacks an IN source at the surface, this means that the nucleation of new ice crystals within the cloud is thus almost entirely determined by the rate at which these recycled below-cloud IN are ingested into the cloud base. This leads to ice crystal nucleation and growth exclusively in updraft regions where recycled IN are lifted into the cloud from below, while downdraft regions are almost entirely liquid-phase (Figs. 8b, 9a,b). This total dependence of ice nucleation on recycled IN also means that the total number of IN in the domain decreases over time, as each round of ice nucleation, growth, sedimentation, sublimation, recycling, and re-nucleation always results in a loss of IN as some particles reach the surface without completely sublimating. Surprisingly, the actual number concentration of ice crystals within the cloud remains in a roughly steady-state condition between 4 and 8 hours in the simulation. However, the mixed-phase nature of the cloud is moribund; with no way to replenish IN, the vertically-integrated IN concentration below cloud drops precipitously from $\sim 6 \times 10^5 \text{ m}^{-2}$ at $t = 2$ hours to $\sim 2 \times 10^5 \text{ m}^{-2}$ by $t = 8$ hours, continuing to fall rapidly as the simulation ends (Fig. 7).




Figure 7. A time-height plot of IN number concentration (shaded) and ice crystal desiccation rate (in contours) in the CONTROL simulation. Contour values range from $5 \times 10^{-5} \text{ L}^{-1} \text{ s}^{-1}$ (light blue) to $2.5 \times 10^{-4} \text{ L}^{-1} \text{ s}^{-1}$ (dark blue). Cloud boundaries are indicated by solid black lines, while ice initiation at $t = 1$ hour is shown in the vertical black dotted line.

This indicates that the cloud, already liquid-dominated, will become almost entirely warm-phase with little ice or mixed-phase characteristics.

Another interesting consequence of heterogeneous ice nucleation occurring almost exclusively at cloud base is that ice formation and growth within the cloud occurs quite nonuniformly, colocated with updrafts which bring IN into liquid-saturated conditions. This means that ice within the cloud is almost entirely restricted to updrafts, with very little ice water content in downdraft regions, similar to the modeling results of Raatikainen et al. (2022) (Figs. 8, 9a,b). The almost complete confinement of ice particles to updraft regions means that there is essentially no ice growth by vapor deposition in downdraft regions. Since downdraft regions are entirely liquid, with no competition from ice, this leads to the rather surprising result that downdraft regions of the cloud actually have a slightly higher liquid water content than updraft regions (Fig. 9c)!

3.2 Sensitivity to Humidity - DRY

The DRY simulation displays greatly reduced LWP and cloud depths compared to CONTROL. LWP just before the onset of ice formation at $t = 1$ hour is only $\sim 10.2 \text{ g m}^{-2}$, compared to $\sim 15.0 \text{ g m}^{-2}$ in CONTROL (Fig. 1a). This reduction in LWP




Figure 8. (a) A time series of heterogeneous ice nucleation rate at cloud base (blue) and cloud top (green) in the CONTROL simulation. Ice initiation at $t = 1$ hour is denoted by the black vertical dotted line. (b) A time-average vertical profile of ice phase fraction in updrafts (red, dashed), downdrafts (blue, dashed), and over the entire cloud (black, solid) between hours 2 and 8 in CONTROL.




Figure 9. (a) A horizontal x-z cross-section of the CONTROL simulation at approximately 4.5 hours elapsed time. Ice water mixing ratio is denoted by blue shading, while contours of vertical velocity from -0.2 to 0.2 m s^{-1} are displayed in the blue and red contours (for downdrafts and updrafts, respectively). Cloud boundaries are denoted by solid black lines. (b) Time-average vertical profile of ice water mixing ratio in updrafts (solid) and downdrafts (dashed) between hours 2 and 8. (c) same as (b), but for liquid water mixing ratio.

compared to CONTROL persists and widens as the simulation progresses. Although LWP does still increase steadily in DRY from hours 2 to 8, it does so at a slower rate than in CONTROL. Between hours 2 and 8, the average LWP in DRY is $\sim 14.3 \text{ g m}^{-2}$, $\sim 40.6\%$ less than the average of $\sim 24.1 \text{ g m}^{-2}$ in CONTROL (Fig. 1a). This reduction in LWP is due to the entrainment of drier above-cloud air than in CONTROL, reducing cloud depth and condensational growth within the cloud (Figs. 5b, 10a,c). As discussed in Section 2, the initial water vapor mixing ratio above cloud is set to 0.8 g kg^{-1} in DRY, compared to 1.2 g kg^{-1} in CONTROL. This corresponds to an initial above-cloud relative humidity of $\sim 41.3\%$ in DRY and $\sim 61.9\%$ in CONTROL.




Figure 10. Time-height plots of total condensation rate for six simulations (in green shading). Cloud boundaries for each simulation are indicated by the solid black lines, while ice initialization is indicated by the vertical black dotted line.

This reduction in LWP also slows down (but does not eliminate) the positive feedback loop which deepens the cloud in CONTROL. With a lower initial LWP, radiative cooling at cloud top is reduced in DRY compared to CONTROL. Between hours 2 and 8, the average longwave radiative cooling rates within the top 50 meters of the cloud are $\sim 65.2 \text{ K day}^{-1}$ and $\sim 48.6 \text{ K day}^{-1}$ in CONTROL and DRY, respectively (Fig. 4a).

However, the most interesting aspect of DRY is the evolution of ice within and below the cloud. The initial “pulse” of ice formation which begins at $t = 1$ hour and continues until approximately 2 hours is weaker in DRY than CONTROL (Fig. 2b). Even after the “pulse”, IWP remains depressed in the DRY simulation compared to the CONTROL simulation until approximately $t = 5$ hours, after which IWP increases above the value in CONTROL (Fig. 2b). Between $t = 2$ -5 hours, average




Figure 11. A time-height comparison of net deposition rate per particle. Deposition rates per particle are shown in green and brown shading in (a,b), while differences between DRY and CONTROL are shown in blue and red shading in (c). Cloud contours for respective simulations are indicated by solid black lines in (a,b), while the initiation of ice processes at $t = 1$ hour is shown by the vertical dotted black lines.

IWP in CONTROL is $\sim 3.65 \text{ g m}^{-2}$, compared to $\sim 2.87 \text{ g m}^{-2}$ in DRY. However, between $t = 5$ -8 hours, average IWP in CONTROL has fallen to $\sim 2.73 \text{ g m}^{-2}$ but has risen to 3.65 g m^{-2} in DRY.

265 These results were unexpected, and, at first glance, seem to corroborate those of Hoffmann (2020), that entrainment of dry air and evaporation of cloud droplets lead to an enhancement of ice crystal growth through the WBF process. However, this is not in fact the case in our experiments; rather, the increase in IWP is due to differences in IN recycling between the CONTROL and DRY simulations. As discussed above, recycling of IN below cloud base by the desiccation of sedimenting ice crystals is the only way in which the number concentration of below-cloud IN can be increased. Without this recycling, IN which were
 270 consumed by nucleation would always be removed from the domain as ice crystals reached the surface, quickly starving the simulation of IN and leading to a purely liquid cloud. This therefore means that the amount of recycled IN is therefore a critical factor in controlling the rate of new ice nucleation and growth within the cloud and a major determinant of total IWP.

In DRY, two important factors lead to much more efficient IN recycling than in CONTROL. The first is that ice crystals simply don't experience as much vapor depositional growth in DRY as they do in CONTROL. Per-particle deposition rates are
 275 reduced in DRY compared to CONTROL, as is the vertical depth of ice-supersaturated air (Fig. 11). As a result, ice crystals in DRY experience reduced maximum growth rates and spend less time within ice-supersaturated conditions than those in CONTROL. This leads to smaller ice crystals when they begin to sublimate. At the altitude corresponding to 100% RH with respect to ice (below which falling ice crystals will sublimate), the mean ice radius (averaged over the long axis of each particle) is $\sim 872 \text{ } \mu\text{m}$ in CONTROL and only $\sim 705 \text{ } \mu\text{m}$ in DRY (Figs. 6, A3A5).

280 Compounding the fact that ice crystals in DRY are smaller when they begin to sublimate, the below-cloud mixed layer is also drier in DRY. Between hours 2 and 8, the average relative humidity over ice in the ice-subaturated region below cloud




Figure 12. (a) Profiles of relative humidity with respect to ice averaged over time between hours 2 and 8. (b) Time series of surface precipitation (in mm of water equivalent) in six simulations. Ice initiation at $t = 1$ hour is indicated by the vertical dotted black line in (b).

is $\sim 91.3\%$ in DRY, compared with $\sim 92.7\%$ in CONTROL. Compounding the greater dryness is the fact that the depth of this subsaturated layer is also approximately 30 meters deeper in the DRY compared to CONTROL (Fig. 12a).

Together, the combination of the smaller size of ice crystals when they begin to sublime, and the deeper and drier layer in which they sublime, leads to a greater fraction of ice crystals in DRY desiccating rather than reaching the surface. Indeed, between hours 2 and 8, the average number concentration of ice crystals at the lowest model level in DRY is only $\sim 37.5\%$ of the number concentration at the lowest ice-supersaturated altitude (not shown). By contrast, $\sim 59.2\%$ of sedimenting ice crystals at the lowest ice-supersaturated altitude reach the surface in CONTROL (Fig. A2A4b). This greater rate of desiccation in DRY in turn increases the recycling of IN below cloud (Figs 3b, 14). Since ice crystal number concentrations within the cloud are primarily determined by the rate of uptake of recycled IN from below the cloud, DRY, with higher recycled IN concentrations, has $\sim 44.1\%$ greater in-cloud ice crystal number concentrations than CONTROL averaged between hours and 8 (Fig. 3a). Although each individual ice crystal is, on average, smaller in DRY, the fact that there are more of them means that the total deposition and IWP eventually become larger than in CONTROL, as discussed above (Figs 13b, 2b, 6, A3A5). This suggests that the increase in IWP in DRY is driven by differences in ice crystal number concentration, rather than any manifestation of entrainment-enhanced ice growth as reported in Hoffmann (2020). The effects of IN recycling on ice crystal number concentration and IWP are discussed further in Sections 3.5 and 3.6.

Notably, the increase in in-cloud ice crystal number concentration in DRY is much greater than that in HIGHIN. Between hours 2 and 8, average in-cloud ice crystal number concentration in HIGHIN is $\sim 18.1\%$ greater than that in CONTROL,




Figure 13. Time series of average in-cloud deposition rate (a) and condensation rate (b). Note that this is not the deposition rate per particle, but rather per mass of air. Ice initiation at $t = 1$ hour is indicated by the vertical dotted black lines. Quantities are smoothed with a Gaussian time filter with a standard deviation of 5 minutes for clarity.




Figure 14. A time-height comparison of IN number concentration and nucleation rate. (a,b) show IN number concentration in pink-purple shading and ice nucleation rate in blue contours. The contour range is from $2 \times 10^{-4} \text{ particle L}^{-1}$ to $1 \times 10^{-3} \text{ particle L}^{-1}$ with a contour interval of $2 \times 10^{-4} \text{ particle L}^{-1}$. Contours are smoothed in time with a gaussian kernel with a 2.5 minute standard deviation for clarity. (c) shows the difference in IN number concentration between DRY and CONTROL in red and blue shading.

compared with a $\sim 44.1\%$ increase in DRY (Fig. A4A6). This indicates that the strong cloud-top inversion inhibits cloud-top
 300 entrainment and prevents the cloud in HIGHIN from accessing many of the IN in the free-troposphere. In contrast, the greater
 IN recycling below cloud base in DRY increases the number of IN in the below-cloud mixed layer, which can be advected back
 into and nucleated within the cloud.

3.3 Sensitivity to Inversion Strength - WEAKINV

Cloud top height increases steadily over the course of the WEAKINV simulation, unlike CONTROL, in which cloud top
 305 height remains almost constant. This is, unsurprisingly, due to an increase in the entrainment rate at cloud top in WEAKINV
 compared to CONTROL. The average entrainment rate between hours 2 and 8 in CONTROL is $\sim 4.20 \text{ mm s}^{-1}$, just barely
 larger than the prescribed large-scale subsidence velocity of 4.12 mm s^{-1} (Fig. 5a). By contrast, the average entrainment rate
 in WEAKINV is $\sim 54.4\%$ higher, at $\sim 6.48 \text{ mm s}^{-1}$. While cloud base height lowers over time in both simulations (as the
 cloud deepens), it does so at a slower rate in WEAKINV than CONTROL. Cloud base height sinks at an average rate of ~ 5.03
 310 mm s^{-1} in CONTROL, compared to only $\sim 3.30 \text{ mm s}^{-1}$ in WEAKINV (not shown). This slower rate of cloud base decrease
 in WEAKINV is due to the greater entrainment of warm air from the entrainment interfacial layer (EIL) into and beneath the
 cloud, which raises the temperature of the boundary layer compared to CONTROL. This warming of the boundary layer in
 WEAKINV also raises the height of the cloud base compared to CONTROL. Together, the combination of higher cloud base
 and higher cloud top in WEAKINV mean that the average cloud depth between hours 2 and 8 is $\sim 249 \text{ m}$, a $\sim 8.35\%$ increase
 315 compared to the average cloud thickness of $\sim 229 \text{ m}$ in CONTROL (Fig. 5b).

The deeper cloud in WEAKINV leads to a larger LWP compared to CONTROL, with an average value of $\sim 27.4 \text{ g m}^{-2}$
 between hours 2 and 8, a $\sim 13.5\%$ increase over the average LWP of $\sim 24.2 \text{ g m}^{-2}$ in CONTROL (Fig. 2a). This increase in
 LWP is due to a combination of thicker cloud and greater condensate mixing ratio compared to CONTROL. As discussed
 previously, the cloud thickness is $\sim 8.35\%$ greater in WEAKINV than CONTROL. At the same time, average in-cloud liquid
 320 water mixing ratio is $\sim 5.2\%$ greater in WEAKINV compared to CONTROL (not shown). This is primarily because the greater
 cloud depth in WEAKINV results in a deeper region of moist adiabatic ascent, and therefore greater total supersaturation
 production, compared to CONTROL. Though one would expect the greater entrainment rate in WEAKINV to mix more dry
 air into the cloud, slowing the rate of condensation, this does not occur to any great extent. Indeed, the average condensation
 rate within the cloud is just $\sim 0.85\%$ less in WEAKINV than it is in CONTROL (Figs. 10, 13b). The relative lack of drying in
 325 WEAKINV, despite the greater entrainment rate, is due to the fact that the above-cloud air in WEAKINV is at a higher relative
 humidity than that in CONTROL, as a consequence of reducing the inversion strength while leaving the above-cloud [water](#)
[vapor](#) mixing ratio unchanged. The average relative humidity of air 50 meters above cloud top (above the entrainment interfacial
 layer) between hours 2 and 8 in CONTROL is $\sim 61.9\%$, considerably drier than the $\sim 75.8\%$ average relative humidity of above-
 cloud air in WEAKINV. Though not shown directly, differences in above-cloud relative humidity between simulations can be
 330 inferred from Figs. 1 and 12a.

The greater cloud depth in WEAKINV also increases the lifetime of ice crystals ascending and falling through the cloud
 compared to CONTROL. Between hours 2 and 8, the average in-cloud lifetime of ice crystals in WEAKINV is $\sim 4.5\%$ longer

than that in CONTROL (not shown). Coupled with a per-particle deposition rate which is $\sim 7.2\%$ greater in WEAKINV than CONTROL, this leads to an $\sim 11.2\%$ increase in average in-cloud ice water mixing ratio between hours 2 and 8 in WEAKINV compared to CONTROL. This increased deposition rate in WEAKINV is the result of multiple factors, including a greater average vertical velocity variance in WEAKINV and greater lifetime of ice crystals (Fig. A5A7). The greater vertical velocity variance in WEAKINV translates to higher updraft (and downdraft) speeds, resulting in a greater production of supersaturation compared to CONTROL. The greater lifetime of ice, meanwhile, also results in greater average per-particle deposition rates as older and larger ice particles experience faster depositional growth. The greater depositional growth rate and deeper cloud depth in WEAKINV means that the average IWP between hours 2 and 8 is $\sim 12.1\%$ greater than that in CONTROL, at $\sim 3.53 \text{ g m}^{-2}$ compared to $\sim 3.15 \text{ g m}^{-2}$ in CONTROL (Fig. 2b). Ice crystals at the lowest ice-supersaturated altitude are also larger in WEAKINV than CONTROL, with an average major-axis diameter of $\sim 913 \text{ }\mu\text{m}$ compared to $\sim 871 \text{ }\mu\text{m}$ in CONTROL (Fig. A3A5).

The greater entrainment rate of free-tropospheric air in WEAKINV has important consequences for the thermodynamics of the sub-cloud boundary layer. The ingestion of warm air downward into the cloud results in a higher cloud base in WEAKINV compared to CONTROL and a warmer, drier sub-cloud boundary layer. The average relative humidity with respect to ice throughout the sub-cloud boundary layer is $\sim 1.08\%$ smaller in WEAKINV compared to CONTROL (Fig. 12a). Interestingly, however, the fraction of ice crystals present at the lowest ice-supersaturated altitude in WEAKINV which reach the surface is $\sim 57.5\%$, barely different from the average value of $\sim 59.3\%$ in CONTROL. Similarly, both CONTROL and WEAKINV show similar amounts of ice precipitation at the surface (Fig. 12b). This suggests that the greater ice crystal size in WEAKINV and the drier sub-cloud boundary layer compared to CONTROL have counteracting effects on ice crystal desiccation rate, leading to a similar overall fraction of ice crystals which desiccate.

3.4 Combining Humidity and Inversion Strength - DRY_WEAKINV

DRY_WEAKINV displays the lowest LWP of any of the four sensitivity simulations. Average LWP between hours 2 and 8 in DRY_WEAKINV is only $\sim 10.5 \text{ g m}^{-2}$, a reduction of $\sim 56.4\%$, $\sim 26.6\%$, and $\sim 61.6\%$ from CONTROL, DRY, and WEAKINV, respectively (Fig. 2a). This reduction in LWP is driven by a reduction in both condensational growth and cloud depth compared to CONTROL, DRY, or WEAKINV. Between hours 2 and 8, the average in-cloud condensation rate in DRY_WEAKINV is $\sim 17.3\%$, $\sim 10.1\%$, and $\sim 16.6\%$ smaller than those in CONTROL, DRY, and WEAKINV, respectively (Figs. 13b, 10). DRY_WEAKINV also has the thinnest cloud of the 4 simulations; the average cloud depth of $\sim 160.7 \text{ m}$ between hours 2 and 8 is a $\sim 30\%$ decrease compared to CONTROL, a $\sim 11.2\%$ decrease compared to DRY, and a $\sim 35.4\%$ decrease from WEAKINV (Fig. 5b). The fact that the cloud in DRY_WEAKINV is thinner than that in DRY is particularly surprising given the fact that the cloud top height increases slowly throughout the course of the simulation in DRY_WEAKINV while it falls slowly during DRY (not shown). However, the cloud base in DRY_WEAKINV is the highest of all simulations. These results present a sharp contrast with WEAKINV, which displayed a higher LWP and deeper cloud than CONTROL. The differences between WEAKINV and DRY_WEAKINV clearly illustrates that not just the magnitude, but also the sign of the impacts of entrainment on cloud microphysical properties is strongly affected by the relative humidity of the free troposphere.

The reduction in LWP in DRY_WEAKINV inhibits the rate of longwave cooling at cloud top. Between hours 2 and 8, the average longwave cooling rate within the uppermost 50 meters of the cloud is $\sim 38.4 \text{ K day}^{-1}$, a $\sim 41.1\%$, $\sim 21.0\%$, and $\sim 42.1\%$ reduction compared to CONTROL, DRY, and WEAKINV, respectively (Fig. 4a). However, even in DRY_WEAKINV, this reduced longwave cooling rate is sufficient to promote a positive feedback loop of cloud development as in CONTROL, DRY, and WEAKINV. Longwave cooling rates increase from approximately $\sim 29.1 \text{ K day}^{-1}$ at hour 2 to $\sim 45.2 \text{ K day}^{-1}$ at hour 8, as LWP likewise increases from $\sim 6.2 \text{ g m}^{-2}$ to $\sim 16.3 \text{ g m}^{-2}$ over the same time period. This reduced longwave cooling also suppresses turbulence in DRY_WEAKINV compared to the other simulations, with a TKE path $\sim 18.6\%$, $\sim 11.9\%$, and $\sim 23.8\%$ smaller than those in CONTROL, DRY, and WEAKINV, respectively (Fig. 4b). This reduction in longwave cooling and TKE path may act to limit cloud-top entrainment; mean cloud-top entrainment rate between hours 2 and 8 in DRY_WEAKINV is $\sim 4.74 \text{ mm s}^{-1}$, a $\sim 26.7\%$ decrease compared to CONTROL (Fig. 5a). This suggests the presence of a negative feedback in the presence of weak inversion strength and dry above-cloud air, in which initial dry air entrainment decreases liquid water content and radiative cooling at cloud top, reducing turbulence and cloud-top entrainment, helping to insulate the cloud from further drying.

IWP in DRY_WEAKINV, as in all simulations, is highly variable over time. The initial "pulse" of ice formation is strongly reduced compared to DRY, CONTROL, or WEAKINV, with a peak IWP of $\sim 4.9 \text{ g m}^{-2}$ (Fig. 2b). This constitutes a $\sim 35.9\%$, $\sim 24.6\%$, and $\sim 45.0\%$ reduction compared to CONTROL, DRY, and WEAKINV, respectively. Between hours 2 and 8, IWP in DRY_WEAKINV is $\sim 3.7\%$, $\sim 7.0\%$, and $\sim 14.1\%$ reduced compared to CONTROL, DRY, and WEAKINV, respectively. However, this average difference masks the fact that IWP in DRY_WEAKINV increases over time relative to the other three simulations (Fig. 2b). As with DRY, this relative increase in IWP is due to the more efficient recycling of IN beneath the cloud compared to the CONTROL or WEAKINV simulations. Examining Figure 3b, the DRY_WEAKINV simulation stands out as having the highest below-cloud IN quantities of any simulation without IN re-injection.

The very efficient IN recycling in DRY_WEAKINV is due to the same factors as those which led to greater IN recycling in DRY: reduced ice crystal depositional growth rates resulting in smaller ice crystals, and a deep and dry ice-subsaturated layer. Between hours 2 and 8, the average depth of the ice-subsaturated layer below cloud is $\sim 493 \text{ m}$ in DRY_WEAKINV, compared to $\sim 414 \text{ m}$, $\sim 442 \text{ m}$, and $\sim 445 \text{ m}$ in CONTROL, DRY, and WEAKINV, respectively. Simultaneously, the average relative humidity with respect to ice in the subsaturated region of the sub-cloud boundary layer in DRY_WEAKINV is $\sim 3.2\%$, $\sim 1.8\%$, and $\sim 1.7\%$ less than that in CONTROL, DRY, and WEAKINV, respectively (Fig. 12a). Ice crystals sedimenting into the ice-subsaturated layer are also smaller than those in the other three simulations, with a major axis diameter of only $\sim 625 \text{ }\mu\text{m}$ compared to $\sim 871 \text{ }\mu\text{m}$, $\sim 705 \text{ }\mu\text{m}$, and $\sim 913 \text{ }\mu\text{m}$ in CONTROL, DRY, and WEAKINV, respectively (Fig. A3A5). Taken together, it's therefore unsurprising that a greater fraction of ice crystals in DRY_WEAKINV desiccate before reaching the surface than in the other three sensitivity simulations. In fact, between hours 2 and 8, only $\sim 29.7\%$ of ice crystals present at the lowest ice-supersaturated altitude in DRY_WEAKINV reach the surface, compared to $\sim 59.3\%$ in CONTROL, $\sim 37.5\%$ in DRY, and $\sim 57.5\%$ in WEAKINV (not shown). Due to both fewer and smaller ice crystals reaching the surface in DRY_WEAKINV compared to the other three simulations, ice precipitation at the surface is also lower in DRY_WEAKINV compared to CONTROL, DRY, or WEAKINV (Fig. 12b).

3.5 Sensitivity to IN - INFLUX

INFLUX has much-reduced cloud condensate compared to CONTROL, particularly in the latter half of the simulation. Unlike in CONTROL, where both the cloud depth and LWP steadily increase between hours 2 and 8, in INFLUX these quantities both remain fairly stable with time. Between hours 2 and 8, the average LWP in INFLUX is $\sim 13.5 \text{ g m}^{-2}$, $\sim 44.0\%$ less than the average value of $\sim 24.1 \text{ g m}^{-2}$ in CONTROL (Fig. 2a). Interestingly, this is a similar magnitude of LWP reduction as seen in DRY. For the same time period, average IWP in INFLUX is $\sim 102\%$ higher than that in CONTROL, at $\sim 6.36 \text{ g m}^{-2}$ compared to $\sim 3.15 \text{ g m}^{-2}$ (Fig. 2b).

Unsurprisingly, the reduction in LWP and increase in IWP compared to CONTROL are a result of the greater number of ice crystals in INFLUX depleting more water vapor from the cloud through vapor deposition. Though deposition rates per ice particle are higher in CONTROL simulation compared to INFLUX, the much greater ice crystal number concentration in INFLUX simulation leads to greater total deposition rates. In INFLUX, the average vapor deposition rate per particle between hours 2 and 8 is $\sim 25.4 \text{ ng particle}^{-1} \text{ s}^{-1}$, a $\sim 27.9\%$ decrease from the average value of $\sim 35.2 \text{ ng particle}^{-1} \text{ s}^{-1}$ in CONTROL (not shown). However, because the average in-cloud ice crystal number concentration at the same time is $\sim 197\%$ higher in INFLUX than CONTROL, at $\sim 0.51 \text{ particles L}^{-1}$ compared to $\sim 0.17 \text{ particles L}^{-1}$, the total in-cloud deposition rate in INFLUX is $1.18 \times 10^{-5} \text{ g kg}^{-1} \text{ s}^{-1}$, $\sim 121\%$ higher than the average value of $5.35 \times 10^{-6} \text{ g kg}^{-1} \text{ s}^{-1}$ in CONTROL (Figs. 3 13a).

This greater consumption of water vapor through vapor deposition on ice crystals in INFLUX thus reduces the amount of condensational growth compared to CONTROL. Between hours 2 and 8, average in-cloud condensation rates in INFLUX are $\sim 81.6\%$ of those in CONTROL, at $1.12 \times 10^{-4} \text{ g kg}^{-1} \text{ s}^{-1}$ compared to $1.38 \times 10^{-4} \text{ g kg}^{-1} \text{ s}^{-1}$ (Fig. 13b). Comparing only condensation rates actually understates the difference between the two simulations, since, as can be seen in Fig. 10, not only are condensation rates higher in CONTROL than in INFLUX, but they occur over a greater depth as the cloud is thicker. In fact, while the cloud continuously grows deeper in both CONTROL and DRY, it remains at an almost constant thickness in INFLUX. This suggests that the feedback loops present in the CONTROL and DRY simulations between cloud condensate, radiative cooling, production of turbulence, and supersaturation production may be inhibited with certain concentrations of IN, in line with previous research finding that large enough IN concentrations will cause complete cloud glaciation and/or dissipation (Das et al., 2025; Loewe et al., 2017; Morrison et al., 2005; Paukert and Hoose, 2014). Interestingly, the results from INFLUX disagree quantitatively with those of Ovchinnikov et al. (2014), who found that a background IN concentration of 1 L^{-1} supported a steady increase in LWP with time from $\sim 10 \text{ g m}^{-2}$ to between $\sim 40\text{-}50 \text{ g m}^{-2}$. Our INFLUX simulation, by contrast, agrees better with their simulations using fixed IN concentrations of 4 L^{-1} , in which LWP changes little with time, remaining between 10 and 20 g m^{-2} . This may be because Ovchinnikov et al. (2014) utilized an ice crystal growth parameterization with a specified mass-diameter relationship which did not allow for variable habit development as in our model.

The reduction in LWP in the INFLUX simulation, similarly to the DRY simulation, acts to inhibit cloud-top longwave radiative cooling compared to CONTROL. Average longwave cooling rates between hours 2 and 8 in the uppermost 50 meters

of the cloud are $\sim 53.3 \text{ K day}^{-1}$ in INFLUX, $\sim 18.3\%$ less than the $\sim 65.2 \text{ K day}^{-1}$ in CONTROL (Fig. 4a). Coupled with the decrease in condensational latent heating, this leads the INFLUX simulation to have an average TKE path between hours 2 and 8 which is $\sim 17.1\%$ smaller than that in CONTROL (Fig. 4).

3.6 Combining Humidity and IN - DRY_INFLUX

440 The DRY_INFLUX simulation makes for a very interesting comparison, allowing us to analyze how the interaction of thermodynamic and IN modification affect the development of the cloud and boundary layer differently than either modification in isolation. The DRY_INFLUX has the lowest LWP or cloud depth compared to either CONTROL, DRY, or INFLUX. Indeed, its average LWP between hours 2 and 8 is a mere $\sim 8.8 \text{ g m}^{-2}$, $\sim 38.3\%$ less than in DRY, $\sim 34.6\%$ less than in INFLUX, and $\sim 63.4\%$ less than CONTROL (Fig. 2). This is because water vapor within the cloud is depleted both by the enhanced vapor
445 deposition rates from higher ice crystal number concentrations and by increased evaporation through mixing with the dry air above the cloud. As discussed above in Section 3.5, the decrease in LWP associated with higher ice crystal number concentrations in the INFLUX simulation is of a similar magnitude to that associated with enhanced evaporation in the DRY simulation, when compared to LWP in CONTROL. This suggests that, under the environmental conditions of the ISDAC simulation, the effects of perfectly efficient IN recycling and the magnitude of the [water vapor](#) mixing ratio decrease in DRY cause a similar
450 magnitude of supersaturation depletion within the cloud.

This reduction in LWP also leads to a large reduction in longwave radiative cooling in the DRY_INFLUX simulation. Between hours 2 and 8, the average cooling rate in the highest 50 meters of the cloud is only $\sim 39.6 \text{ K day}^{-1}$, compared with $\sim 65.2 \text{ K day}^{-1}$ in CONTROL, a $\sim 39.3\%$ reduction. This is also $\sim 28.2\%$ less than INFLUX and $\sim 18.4\%$ less than DRY (Fig. 4a). This decrease in cloud-top radiative cooling results in a $\sim 26.6\%$ reduction of TKE path from $\sim 153 \text{ m}^3 \text{ s}^{-2}$ in CONTROL
455 to $\sim 112 \text{ m}^3 \text{ s}^{-2}$ in DRY_INFLUX (TKE path in DRY_INFLUX is also $\sim 12.1\%$ lower than that in INFLUX and $\sim 21.1\%$ less than in DRY) (Fig. 4b). However, the cloud appears to be in no danger of glaciating or dissipating in DRY_INFLUX, as LWP increases slowly but steadily from $\sim 7.9 \text{ g m}^{-2}$ at hour 4 to $\sim 9.5 \text{ g m}^{-2}$ at hour 8 (Fig. 2a).

IWP in DRY_INFLUX is lower than that in INFLUX, while higher than that in either DRY or CONTROL. Between hours 2 and 8, the average IWP in DRY_INFLUX is $\sim 4.55 \text{ g m}^{-2}$, $\sim 44.6\%$ and $\sim 39.6\%$ higher than CONTROL and DRY, re-
460 spectively, while also reduced by $\sim 28.5\%$ compared with INFLUX (Fig. 2b). Similar to our comparison of INFLUX and CONTROL, in-cloud deposition rates per particle in DRY_INFLUX are smaller than those in the CONTROL or DRY simulations, at approximately $\sim 20.1 \text{ ng particle}^{-1} \text{ s}^{-1}$ in DRY_INFLUX, compared to $\sim 35.2 \text{ ng particle}^{-1} \text{ s}^{-1}$ in CONTROL and $\sim 26.6 \text{ ng particle}^{-1} \text{ s}^{-1}$ in DRY (not shown). Notably, the average ice radii are smaller both within and below cloud base in the DRY_INFLUX simulation compared with CONTROL, DRY, or INFLUX; Average ice particle radius at the lowest ice-
465 supersaturated altitude is $\sim 610 \mu\text{m}$, compared to $\sim 728 \mu\text{m}$ in INFLUX, $\sim 705 \mu\text{m}$ in DRY, and $\sim 872 \mu\text{m}$ in CONTROL (Fig. [A3A5](#)). However, the much greater ice crystal number concentration in DRY_INFLUX means that both the total deposition rate and the IWP are higher than those in either CONTROL or DRY (Fig. 13a).

As discussed above in the comparison of DRY and CONTROL, the smaller size of ice crystals when they begin to sublimate and the drier sub-cloud sublimation layer within DRY_INFLUX both contribute to increased desiccation rates of ice crystals

470 and a reduction in the number and mass of ice crystals reaching the surface compared to CONTROL. This is most strikingly seen in a comparison of accumulated precipitation over the course of the six simulations, which illustrates that, despite the much greater IWP in DRY_INFLUX compared to CONTROL, the total accumulated snowfall at the surface is higher in CONTROL throughout almost the entire simulation is higher than in DRY_INFLUX, due to higher sub-cloud sublimation rates in the latter simulation (though, by the end of the simulation, CONTROL has become so IN-depleted that it is likely that running both 475 simulations for another 8 hours would result in DRY_INFLUX overtaking CONTROL in terms of total surface precipitation) (Fig. 12b). Unlike DRY and CONTROL, however, the increased sublimation and desiccation rates within DRY_INFLUX result neither in a higher IN number concentration below cloud nor a higher in-cloud ice crystal number concentration compared to INFLUX, as the total amount of IN-containing particles (both “dry” INs and ice crystals containing INs) are held constant in both simulations.

480 4 Discussion and Conclusions

To examine the sensitivity of Arctic mixed-phase clouds to environmental humidity, inversion strength, and IN depletion, we ran seven simulations using an LES model with Lagrangian cloud microphysics. Our results demonstrate the clear feedback between the microphysical and boundary-layer processes in Arctic mixed-phase stratocumulus clouds, and the role of ice in determining this feedback.

485 Reduced above-cloud relative humidity in DRY results in both expected and unexpected impacts on cloud and boundary-layer properties. Due to increased cloud-top evaporation, LWP within the cloud is reduced compared to CONTROL, as is, initially, IWP. By the end of the DRY simulation, however, IWP is higher than in the CONTROL simulation, though LWP remains much lower (Fig. 2). This is due to enhanced IN recycling driven by desiccation of sublimating ice crystals. The entrainment of drier above-cloud air into the cloud reduces the available supersaturation for ice crystal growth by vapor de- 490 position, resulting in smaller ice crystals (Fig. 6). Simultaneously, turbulent transport of this drier above-cloud air through and below the cloud reduces water vapor mixing ratios below cloud base, increasing the water vapor deficit experienced by ice crystals below the cloud. Both of these factors result in a greater number of sedimenting ice crystals desiccating before reaching the surface and returning their IN to the boundary layer, which are then nucleated once more at cloud base (Fig. 14). The higher number concentration of ice crystals in the DRY simulation compared to CONTROL therefore results in a higher 495 final IWP, even though the average size of each individual ice crystal is smaller in DRY than CONTROL.

We do not find any evidence to support the hypothesis of dry air entrainment enhancing ice crystal growth, as hypothesized in Hoffmann (2020). The increased IWP present in the DRY simulation would seem to lend credence to this hypothesis, but, as described above, the increase in IWP in the DRY simulation is solely a result of more efficient IN recycling. Within the DRY simulation, deposition rates per particle are reduced compared to CONTROL, but the greater number concentration of 500 ice crystals within cloud results in a higher total deposition rate and IWP (Figs. 11, 13a). Examination of the INFLUX and DRY_INFLUX simulations, both of which run with perfectly efficient IN recycling, clearly shows a reduction in IWP in the

latter simulation (Fig. 2b). Depositional growth rates, both per particle and total, are lower in DRY_INFLUX than INFLUX, as entrainment of dry air reduces the available supersaturation for ice crystal growth.

The WEAKINV and DRY_WEAKINV simulations support the finding of several previous studies of subtropical stratocumulus that above-cloud relative humidity plays a critical role in modulating the effects of cloud-top entrainment on cloud properties (Ackerman et al., 2004; Eastman and Wood, 2018; Xu and Xue, 2015). In particular, the fact that LWP increases in WEAKINV compared to CONTROL, while simultaneously decreasing in DRY_WEAKINV compared to CONTROL, indicates that not only the magnitude, but the *sign* of microphysical impacts of increased entrainment is dependent on the above-cloud environmental humidity, in line with the results of Xu and Xue (2015). In WEAKINV, the inversion strength above the cloud top is reduced while the [water](#) vapor mixing ratio is unchanged. In effect, this increases the above-cloud relative humidity compared to CONTROL. Despite an overall greater cloud-top entrainment rate in WEAKINV compared to control, the entrainment of higher-humidity air leads to a deepening of the cloud and boundary layer, a larger LWP, and greater cloud-top longwave radiational cooling (Figs. 2a, 4a). The deeper cloud also promotes longer-lived and larger ice particles, leading to an increase in IWP compared to CONTROL (Fig. 2b). Interestingly, while cloud depth is greater and LWP is increased in WEAKINV compared to CONTROL, below-cloud relative humidity is reduced by a similar amount as in DRY (Fig. 12a).

In DRY_WEAKINV, by contrast, the combination of decreased inversion strength and drier above-cloud air strongly depletes cloud water compared to either CONTROL, DRY, or WEAKINV. While above-cloud relative humidity in DRY_WEAKINV is greater than in DRY, it is still reduced compared to CONTROL, while the reduced inversion strength means that this dry above-cloud air is entrained at a much greater rate than either CONTROL or DRY. Unlike WEAKINV, this increased entrainment of dry air results in a dramatic reduction of cloud depth, LWP, and cloud-top longwave radiative cooling rate (Figs. 2a, 4). The entrainment of dry above-cloud air in DRY_WEAKINV also reduces the relative humidity with respect to ice in the sub-cloud boundary layer while reducing both the lifetime and depositional growth rate of ice crystals within the cloud (Fig. 12a). These effects are similar to, but more pronounced than those seen in DRY. As such, it is unsurprising that a larger proportion of sedimenting ice crystals are desiccated in DRY_WEAKINV than in either CONTROL, DRY, or WEAKINV. This increased desiccation results in more efficient IN recycling and a greater in-cloud ice crystal number concentration in DRY_WEAKINV compared to CONTROL, DRY, or WEAKINV (Fig. 3). As a result, by the end of the simulation, IWP in DRY_WEAKINV is similar to that in DRY and greater than that in CONTROL or WEAKINV, both of which have much less efficient below-cloud IN recycling (Fig. 2b).

The imposition of perfectly efficient IN recycling in the INFLUX simulation results in a dramatically increased IWP compared to CONTROL as a much greater number of ice crystals are present in the cloud (Fig. 2b). However, this greater ice crystal number concentration also suppresses LWP, as more supersaturation is consumed by growing ice crystals rather than by cloud droplets (Figs. 2, 13). This suppression of LWP also lessens the degree of mixing in the boundary layer; a lower LWP leads to a reduction in cloud-top radiative cooling and generation of turbulent kinetic energy, a finding supported by previous modeling studies of Arctic stratocumulus clouds atop decoupled boundary layers (Klein and Coauthors, 2009; Ovchinnikov et al., 2014; Solomon et al., 2018; Stevens et al., 2018) (Fig. 4). INFLUX also shows no increase in LWP or thickening of the

cloud over time, unlike CONTROL, indicating that the higher ice crystal number concentration in this simulation compared to CONTROL interrupts the LWP-radiative feedback which acts to increase LWP and deepen the cloud (Figs. 2a, 3a).

Our results indicate that the recycling of IN is of key importance to the microphysical properties of the cloud and is influenced both by the rate of growth of ice crystals within the cloud as well as the characteristics of the sub-cloud boundary layer. While numerous modeling studies have previously highlighted the importance of IN recycling to the maintenance of Arctic mixed-phase clouds, the influence of above-cloud relative humidity on IN recycling has not been specifically examined (Ahola et al., 2020; Fan et al., 2009; Raatikainen et al., 2022; Solomon et al., 2015).

While surface emission rates of IN in the wintertime Arctic over sea ice appear to be quite low (Bigg, 1996; Creamean et al., 2018; Tan et al., 2023; Wex et al., 2019), the lack of any surface source of IN in our simulations may not be entirely reflective of the aerosol environment off the coast of Alaska in late April; more recent research has highlighted the role of small leads of open water as a source of marine IN (Creamean et al., 2022; Griesche et al., 2021). Previous modeling of the ISDAC Flight 31 cloud also highlighted the importance of marine INP emissions to maintaining mixed-phase cloud properties over a 16 hour simulation (Raatikainen et al., 2022), as IN recycling was insufficient to preserve a steady-state IN or ice crystal number concentration, similar to our results. Further constraints on the rate of terrestrial and marine INP emissions in Arctic spring are necessary to accurately capture the influence of these IN sources in modeling of Arctic mixed-phase stratocumulus. Interestingly, our findings also indicate that recycled IN from desiccating ice crystals can be trapped in a "band" of stably stratified air below the sub-cloud mixed layer and just above the surface. Once the sub-cloud mixed layer reaches the surface, these IN are mixed throughout the boundary layer and back into the cloud. A similar phenomenon could occur with surface IN emissions under conditions of low surface wind speeds and surface sensible heat fluxes. In such an environment, the well-mixed layer stretching down from the cloud might not initially extend all the way to the surface, allowing IN to build up near the surface. Once the sub-cloud mixed layer reaches the surface, this elevated concentration of marine IN could be advected into the cloud and increase the rate of ice nucleation, potentially increasing IWP within the cloud.

This study also did not account for multiple types of IN; while nucleation rates in the LCM are probabilistic, the parameterization used is that of Wex et al. (2015) for fragments of SNOWMAX bacteria, which nucleate at high temperatures. Under long-range transport of other aerosols such as mineral dust or volcanic ash, however, the omission of these INP sources may underestimate ice crystal number concentrations within the cloud (Prenni et al., 2009). Additionally, Fu and Xue (2017) found that cloud-top radiative cooling can act to nucleate colder-temperature INPs over time, helping counteract the effect of IN depletion through precipitation scavenging. Future simulations with multiple IN species would help illustrate the importance of this effect under different background aerosol and environmental conditions.

As we noted in the Introduction, several ice growth mechanisms which affect mixed-phase Arctic stratocumulus are not represented in these simulations, particularly riming, aggregation, and SIP, meaning that the results of our simulations may not be exactly representative of real-world conditions. Our simulations also only examined two very specific environments, while Arctic stratocumulus clouds are known to occur under a wide range of atmospheric conditions (de Boer et al., 2009; McFarquhar et al., 2011; Griesche et al., 2021; Shupe et al., 2006, 2011; Verlinde et al., 2007). How IN recycling affects Arctic stratocumulus clouds under such environmental conditions is therefore left unanswered by the present analysis, but may have

significant effects on the structure and longevity of these clouds, as well as their radiative impacts at the surface. To address these limitations of our analysis, future modeling research with different microphysical schemes and with different initial environmental conditions, as well as more observational studies measuring IN and ice crystal number concentration within and below Arctic stratocumulus, are therefore essential to improve knowledge of the processes affecting and behavior of Arctic
575 mixed-phase stratocumulus.

Appendix A: Supplemental Figures




Figure A1. Time series of LWP (a), IWP (b), cloud-top longwave radiative cooling rate (c), and vertically-integrated TKE (d) in the CONTROL (solid blue) and LARGEDOMAIN (dotted blue) simulations.




Figure A2. Plan view comparison of the CONTROL (a) and LARGEDOMAIN (b) simulations at $z = 500$ m and $t = 5$ hours. Vertical velocity is shown in the red and blue dashed lines, while ice water mixing ratio is indicated with blue shading.




Figure A3. Time-height diagram of air temperature (a) and number concentration-weighted mean particle aspect ratio (b). Cloud boundaries are denoted by black lines.




Figure A4. Time series of ice crystal number concentration (a) and fraction of ice crystals at the lowest ice-supersaturated altitude which reach the surface (b) in CONTROL. In (a), ice crystal number concentration at the lowest ice-supersaturated altitude is shown in red, while the ice crystal number concentration at the surface is shown in blue. In (b), the fraction of ice crystals reaching the surface is calculated using time-shifted arrays, with the shift equal to the maximum temporal correlation between the arrays of ice crystal number concentration at the lowest ice-supersaturated altitude and ice crystal number concentration at the surface.






Figure A5. Time series of mean ice crystal radius (calculated along the maximum dimension of each ice crystal) at the lowest ice-supersaturated altitude in six simulations.



simulations.

Figure A6. As in Fig. 3, but for the CONTROL (blue), DRY (orange), and HIGHIN (black) simulations.

Time-height plots of the square root of vertical velocity variance for six simulations. Vertical velocity variance is indicated in shading, while cloud boundaries (defined as regions where cloud water mixing ratio is $> 0.01 \text{ g kg}^{-1}$) are marked in black



lines.

Figure A7. Time-height plots of the square root of vertical velocity variance for six simulations. Vertical velocity variance is indicated in shading, while cloud boundaries (defined as regions where cloud water mixing ratio is $> 0.01 \text{ g kg}^{-1}$) are marked in black lines.

Code and data availability. The model data used in the production of this paper is shared at the following DOI: <https://doi.org/10.5282/ubm/data.720>.

A Jupyter Notebook file to reproduce the figures and quantitative analysis seen in this paper is available at the following DOI: <https://doi.org/10.5281/zenodo.1411111>.

(Ascher and Hoffmann, 2025; Ascher, 2025). Ascher and Hoffmann (2025) also includes a short guide on how to set up a Python environment

580 and use the Jupyter notebooks of Ascher (2025).

Author contributions. Benjamin Ascher contributed research and the main writing component of this manuscript. Fabian Hoffmann was the primary architect of the Lagrangian cloud microphysics model used in this research project, offered helpful edits and contributions to the text, and served as PhD advisor to the first author.

Competing interests. The authors declare that they have no conflicts of interest.

585 *Acknowledgements.* This work was supported by the Emmy Noether program of the German Research Foundation (DFG) under Grant HO 6588/1-1. The authors gratefully acknowledge the Gauss Centre for Supercomputing e.V. (<http://www.gauss-centre.eu>) for helping this project by providing computing time on the GCS supercomputer SuperMUC-NG at the Leibniz Supercomputing Centre (<http://www.lrz.de>). We would also like to thank Dr. Jessie Creamean for her role as editor during the peer-review process, as well as Dr. Étienne Vignon and one anonymous reviewer for their insightful feedback which has improved this manuscript.

- Ackerman, A. S., Kirkpatrick, M. P., Stevens, D. E., and Toon, O. B.: The Impact of Humidity above Stratiform Clouds on Indirect Aerosol Climate Forcing, *Nature*, 432, 1014–1017, <https://doi.org/10.1038/nature03174>, 2004.
- Ahola, J., Korhonen, H., Tonttila, J., Romakkaniemi, S., Kokkola, H., and Raatikainen, T.: Modelling Mixed-Phase Clouds with the Large-Eddy Model UCLALES–SALSA, *Atmospheric Chemistry and Physics*, 20, 11 639–11 654, <https://doi.org/10.5194/acp-20-11639-2020>, 595 2020.
- Ascher, B.: Benapplus1/ISDAC_2025: Version 0.5, <https://doi.org/10.5281/zenodo.17735740>, 2025.
- Ascher, B. and Hoffmann, F.: Model Data to Recreate Figures Seen in the Manuscript "Impacts of Environmental Conditions and Ice Nuclei Recycling on Arctic Mixed-Phase Cloud Properties.", <https://doi.org/10.5282/ubm/data.720>, 2025.
- Avramov, A. and Harrington, J. Y.: Influence of Parameterized Ice Habit on Simulated Mixed Phase Arctic Clouds, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD012108>, 2010. 600
- Bigg, E. K.: Ice Forming Nuclei in the High Arctic, *Tellus B: Chemical and Physical Meteorology*, 48, 223–233, <https://doi.org/10.3402/tellusb.v48i2.15888>, 1996.
- Caldwell, P. and Bretherton, C. S.: Response of a Subtropical Stratocumulus-Capped Mixed Layer to Climate and Aerosol Changes, *Journal of Climate*, 22, 20–38, <https://doi.org/10.1175/2008JCLI1967.1>, 2009.
- Chen, J.-P. and Lamb, D.: The Theoretical Basis for the Parameterization of Ice Crystal Habits: Growth by Vapor Deposition, *Journal of the Atmospheric Sciences*, 51, 1206–1222, [https://doi.org/10.1175/1520-0469\(1994\)051<1206:TTBFTP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<1206:TTBFTP>2.0.CO;2), 1994. 605
- Creamean, J. M., Kirpes, R. M., Pratt, K. A., Spada, N. J., Maahn, M., de Boer, G., Schnell, R. C., and China, S.: Marine and Terrestrial Influences on Ice Nucleating Particles during Continuous Springtime Measurements in an Arctic Oilfield Location, *Atmospheric Chemistry and Physics*, 18, 18 023–18 042, <https://doi.org/10.5194/acp-18-18023-2018>, 2018.
- Creamean, J. M., Barry, K., Hill, T. C. J., Hume, C., DeMott, P. J., Shupe, M. D., Dahlke, S., Willmes, S., Schmale, J., Beck, I., Hoppe, C. J. M., Fong, A., Chamberlain, E., Bowman, J., Scharien, R., and Persson, O.: Annual Cycle Observations of Aerosols Capable of Ice Formation in Central Arctic Clouds, *Nature Communications*, 13, 3537, <https://doi.org/10.1038/s41467-022-31182-x>, 2022. 610
- Das, A., Clothiaux, E. E., and Harrington, J. Y.: Ice Crystal Habit Effects on the Resilience of Arctic Mixed-Phase Stratus Clouds in a One-Dimensional Model, *Journal of Geophysical Research: Atmospheres*, 130, e2024JD041 305, <https://doi.org/10.1029/2024JD041305>, 615 2025.
- de Boer, G., Eloranta, E. W., and Shupe, M. D.: Arctic Mixed-Phase Stratiform Cloud Properties from Multiple Years of Surface-Based Measurements at Two High-Latitude Locations, *Journal of the Atmospheric Sciences*, 66, 2874–2887, <https://doi.org/10.1175/2009JAS3029.1>, 2009.
- Eastman, R. and Wood, R.: The Competing Effects of Stability and Humidity on Subtropical Stratocumulus Entrainment and Cloud Evolution from a Lagrangian Perspective, *Journal of the Atmospheric Sciences*, 75, 2563–2578, <https://doi.org/10.1175/JAS-D-18-0030.1>, 2018. 620
- Egerer, U., Ehrlich, A., Gottschalk, M., Griesche, H., Neggers, R. A. J., Siebert, H., and Wendisch, M.: Case Study of a Humidity Layer above Arctic Stratocumulus and Potential Turbulent Coupling with the Cloud Top, *Atmospheric Chemistry and Physics*, 21, 6347–6364, <https://doi.org/10.5194/acp-21-6347-2021>, 2021.
- Eirund, G. K., Lohmann, U., and Possner, A.: Cloud Ice Processes Enhance Spatial Scales of Organization in Arctic Stratocumulus, *Geophysical Research Letters*, 46, 14 109–14 117, <https://doi.org/10.1029/2019GL084959>, 2019. 625

Eirund, G. K., Possner, A., and Lohmann, U.: The Impact of Warm and Moist Airmass Perturbations on Arctic Mixed-Phase Stratocumulus, *Journal of Climate*, 33, 9615–9628, <https://doi.org/10.1175/JCLI-D-20-0163.1>, 2020.

Fan, J., Ovtchinnikov, M., Comstock, J. M., McFarlane, S. A., and Khain, A.: Ice Formation in Arctic Mixed-Phase Clouds: Insights from a 3-D Cloud-Resolving Model with Size-Resolved Aerosol and Cloud Microphysics, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/10.1029/2008JD010782>, 2009.

Fu, S. and Xue, H.: The Effect of Ice Nuclei Efficiency on Arctic Mixed-Phase Clouds from Large-Eddy Simulations, *Journal of the Atmospheric Sciences*, 74, 3901–3913, <https://doi.org/10.1175/JAS-D-17-0112.1>, 2017.

Garrett, T. J., Radke, L. F., and Hobbs, P. V.: Aerosol Effects on Cloud Emissivity and Surface Longwave Heating in the Arctic, *Journal of the Atmospheric Sciences*, 59, 769–778, [https://doi.org/10.1175/1520-0469\(2002\)059<0769:AEOCEA>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<0769:AEOCEA>2.0.CO;2), 2002.

Gayet, J.-F., Treffeisen, R., Helbig, A., Bareiss, J., Matsuki, A., Herber, A., and Schwarzenboeck, A.: On the Onset of the Ice Phase in Boundary Layer Arctic Clouds, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/10.1029/2008JD011348>, 2009-10-16.

Griesche, H. J., Ohneiser, K., Seifert, P., Radenz, M., Engelmann, R., and Ansmann, A.: Contrasting Ice Formation in Arctic Clouds: Surface-Coupled vs. Surface-Decoupled Clouds, *Atmospheric Chemistry and Physics*, 21, 10 357–10 374, <https://doi.org/10.5194/acp-21-10357-2021>, 2021.

Harrington, J. Y., Reisin, T., Cotton, W. R., and Kreidenweis, S. M.: Cloud Resolving Simulations of Arctic Stratus: Part II: Transition-season Clouds, *Atmospheric Research*, 51, 45–75, [https://doi.org/10.1016/S0169-8095\(98\)00098-2](https://doi.org/10.1016/S0169-8095(98)00098-2), 1999.

Hiranuma, N., Brooks, S. D., Moffet, R. C., Glen, A., Laskin, A., Gilles, M. K., Liu, P., Macdonald, A. M., Strapp, J. W., and McFarquhar, G. M.: Chemical Characterization of Individual Particles and Residuals of Cloud Droplets and Ice Crystals Collected on Board Research Aircraft in the ISDAC 2008 Study, *Journal of Geophysical Research: Atmospheres*, 118, 6564–6579, <https://doi.org/10.1002/jgrd.50484>, 2013.

Hoffmann, F.: Effects of Entrainment and Mixing on the Wegener–Bergeron–Findeisen Process, *Journal of the Atmospheric Sciences*, 77, 2279–2296, <https://doi.org/10.1175/JAS-D-19-0289.1>, 2020.

Hoffmann, F., Raasch, S., and Noh, Y.: Entrainment of Aerosols and Their Activation in a Shallow Cumulus Cloud Studied with a Coupled LCM–LES Approach, *Atmospheric Research*, 156, 43–57, <https://doi.org/10.1016/j.atmosres.2014.12.008>, 2015.

Jackson, R.: The Dependence of Arctic Mixed Phase Stratus Ice Cloud Microphysics on Aerosol Concentration Using Observations Acquired during ISDAC and M-PACE, Ph.D. thesis, University of Illinois at Urbana-Champaign, 2012.

Khairoutdinov, M. F. and Randall, D. A.: Cloud Resolving Modeling of the ARM Summer 1997 IOP: Model Formulation, Results, Uncertainties, and Sensitivities, *Journal of the Atmospheric Sciences*, 60, 607–625, [https://doi.org/10.1175/1520-0469\(2003\)060<0607:CRMOTA>2.0.CO;2](https://doi.org/10.1175/1520-0469(2003)060<0607:CRMOTA>2.0.CO;2), 2003.

Klein, S. A. and Coauthors: Intercomparison of Model Simulations of Mixed-Phase Clouds Observed during the ARM Mixed-Phase Arctic Cloud Experiment. I: Single-Layer Cloud, *Quarterly Journal of the Royal Meteorological Society*, 135, 979–1002, <https://doi.org/10.1002/qj.416>, 2009.

Loewe, K., Ekman, A. M. L., Paukert, M., Sedlar, J., Tjernström, M., and Hoose, C.: Modelling Micro- and Macrophysical Contributors to the Dissipation of an Arctic Mixed-Phase Cloud during the Arctic Summer Cloud Ocean Study (ASCOS), *Atmospheric Chemistry and Physics*, 17, 6693–6704, <https://doi.org/10.5194/acp-17-6693-2017>, 2017.

- McFarquhar, G. M., Zhang, G., Poellot, M. R., Kok, G. L., McCoy, R., Tooman, T., Fridlind, A., and Heymsfield, A. J.: Ice Properties of Single-Layer Stratocumulus during the Mixed-Phase Arctic Cloud Experiment: 1. Observations, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2007JD008633>, 2007.
- 665 McFarquhar, G. M., Ghan, S., Verlinde, J., Korolev, A., Strapp, J. W., Schmid, B., Tomlinson, J. M., Wolde, M., Brooks, S. D., Cziczo, D., Dubey, M. K., Fan, J., Flynn, C., Gultepe, I., Hubbe, J., Gilles, M. K., Laskin, A., Lawson, P., Leaitch, W. R., Liu, P., Liu, X., Lubin, D., Mazzoleni, C., Macdonald, A.-M., Moffet, R. C., Morrison, H., Ovchinnikov, M., Shupe, M. D., Turner, D. D., Xie, S., Zelenyuk, A., Bae, K., Freer, M., and Glen, A.: Indirect and Semi-direct Aerosol Campaign: The Impact of Arctic Aerosols on Clouds, *Bulletin of the American Meteorological Society*, 92, 183–201, <https://doi.org/10.1175/2010BAMS2935.1>, 2011.
- 670 Mioche, G., Jourdan, O., Delanoë, J., Gourbeyre, C., Febvre, G., Dupuy, R., Monier, M., Szczap, F., Schwarzenboeck, A., and Gayet, J.-F.: Vertical Distribution of Microphysical Properties of Arctic Springtime Low-Level Mixed-Phase Clouds over the Greenland and Norwegian Seas, *Atmospheric Chemistry and Physics*, 17, 12 845–12 869, <https://doi.org/10.5194/acp-17-12845-2017>, 2017.
- Morrison, H., Shupe, M. D., Pinto, J. O., and Curry, J. A.: Possible Roles of Ice Nucleation Mode and Ice Nuclei Depletion in the Extended Lifetime of Arctic Mixed-Phase Clouds, *Geophysical Research Letters*, 32, <https://doi.org/10.1029/2005GL023614>, 2005.
- 675 Ovchinnikov, M., Ackerman, A. S., Avramov, A., Cheng, A., Fan, J., Fridlind, A. M., Ghan, S., Harrington, J., Hoose, C., Korolev, A., McFarquhar, G. M., Morrison, H., Paukert, M., Savre, J., Shipway, B. J., Shupe, M. D., Solomon, A., and Sulia, K.: Intercomparison of Large-Eddy Simulations of Arctic Mixed-Phase Clouds: Importance of Ice Size Distribution Assumptions, *Journal of Advances in Modeling Earth Systems*, 6, 223–248, <https://doi.org/10.1002/2013MS000282>, 2014.
- Pasquier, J. T., Henneberger, J., Ramelli, F., Lauber, A., David, R. O., Wieder, J., Carlsen, T., Gierens, R., Maturilli, M., and Lohmann, U.: Conditions Favorable for Secondary Ice Production in Arctic Mixed-Phase Clouds, *Atmospheric Chemistry and Physics*, 22, 15 579–15 601, <https://doi.org/10.5194/acp-22-15579-2022>, 2022-12-12.
- 680 Paukert, M. and Hoose, C.: Modeling Immersion Freezing with Aerosol-Dependent Prognostic Ice Nuclei in Arctic Mixed-Phase Clouds, *Journal of Geophysical Research: Atmospheres*, 119, 9073–9092, <https://doi.org/10.1002/2014JD021917>, 2014.
- Petters, J. L., Harrington, J. Y., and Clothiaux, E. E.: Radiative–Dynamical Feedbacks in Low Liquid Water Path Stratiform Clouds, *Journal of the Atmospheric Sciences*, 69, 1498–1512, <https://doi.org/10.1175/JAS-D-11-0169.1>, 2012.
- 685 Prenni, A. J., Demott, P. J., Rogers, D. C., Kreidenweis, S. M., Mcfarquhar, G. M., Zhang, G., and Poellot, M. R.: Ice Nuclei Characteristics from M-PACE and Their Relation to Ice Formation in Clouds, *Tellus B: Chemical and Physical Meteorology*, 61, 436–448, <https://doi.org/10.1111/j.1600-0889.2008.00415.x>, 2009.
- Raatikainen, T., Prank, M., Ahola, J., Kokkola, H., Tonttila, J., and Romakkaniemi, S.: The Effect of Marine Ice-Nucleating Particles on Mixed-Phase Clouds, *Atmospheric Chemistry and Physics*, 22, 3763–3778, <https://doi.org/10.5194/acp-22-3763-2022>, 2022.
- 690 Randall, D. A.: Stratocumulus Cloud Deepening through Entrainment, *Tellus A: Dynamic Meteorology and Oceanography*, 36, 446–457, <https://doi.org/10.3402/tellusa.v36i5.11646>, 1984.
- Savre, J. and Ekman, A. M. L.: Large-Eddy Simulation of Three Mixed-Phase Cloud Events during ISDAC: Conditions for Persistent Heterogeneous Ice Formation, *Journal of Geophysical Research: Atmospheres*, 120, 7699–7725, <https://doi.org/10.1002/2014JD023006>, 2015.
- 695 Shi, X., Hagen, H. L., Chow, F. K., Bryan, G. H., and Street, R. L.: Large-Eddy Simulation of the Stratocumulus-Capped Boundary Layer with Explicit Filtering and Reconstruction Turbulence Modeling, *Journal of the Atmospheric Sciences*, 75, 611–637, <https://doi.org/10.1175/JAS-D-17-0162.1>, 2018.

- Shima, S., Kusano, K., Kawano, A., Sugiyama, T., and Kawahara, S.: The Super-Droplet Method for the Numerical Simulation of Clouds and Precipitation: A Particle-Based and Probabilistic Microphysics Model Coupled with a Non-Hydrostatic Model, *Quarterly Journal of the Royal Meteorological Society*, 135, 1307–1320, <https://doi.org/10.1002/qj.441>, 2009.
- Shima, S.-i., Sato, Y., Hashimoto, A., and Misumi, R.: Predicting the Morphology of Ice Particles in Deep Convection Using the Super-Droplet Method: Development and Evaluation of SCALE-SDM 0.2.5-2.2.0, -2.2.1, and -2.2.2, *Geoscientific Model Development*, 13, 4107–4157, <https://doi.org/10.5194/gmd-13-4107-2020>, 2020.
- Shupe, M. D., Matrosov, S. Y., and Uttal, T.: Arctic Mixed-Phase Cloud Properties Derived from Surface-Based Sensors at SHEBA, *Journal of the Atmospheric Sciences*, 63, 697–711, <https://doi.org/10.1175/JAS3659.1>, 2006.
- Shupe, M. D., Walden, V. P., Eloranta, E., Uttal, T., Campbell, J. R., Starkweather, S. M., and Shiobara, M.: Clouds at Arctic Atmospheric Observatories. Part I: Occurrence and Macrophysical Properties, *Journal of Applied Meteorology and Climatology*, 50, 626–644, <https://doi.org/10.1175/2010JAMC2467.1>, 2011.
- Solomon, A., Shupe, M. D., Persson, O., Morrison, H., Yamaguchi, T., Caldwell, P. M., and Boer, G. d.: The Sensitivity of Springtime Arctic Mixed-Phase Stratocumulus Clouds to Surface-Layer and Cloud-Top Inversion-Layer Moisture Sources, *Journal of the Atmospheric Sciences*, 71, 574–595, <https://doi.org/10.1175/JAS-D-13-0179.1>, 2014.
- Solomon, A., Feingold, G., and Shupe, M. D.: The Role of Ice Nuclei Recycling in the Maintenance of Cloud Ice in Arctic Mixed-Phase Stratocumulus, *Atmospheric Chemistry and Physics*, 15, 10 631–10 643, <https://doi.org/10.5194/acp-15-10631-2015>, 2015.
- Solomon, A., de Boer, G., Creamean, J. M., McComiskey, A., Shupe, M. D., Maahn, M., and Cox, C.: The Relative Impact of Cloud Condensation Nuclei and Ice Nucleating Particle Concentrations on Phase Partitioning in Arctic Mixed-Phase Stratocumulus Clouds, *Atmospheric Chemistry and Physics*, 18, 17 047–17 059, <https://doi.org/10.5194/acp-18-17047-2018>, 2018.
- Sotiropoulou, G., Sedlar, J., Tjernström, M., Shupe, M. D., Brooks, I. M., and Persson, P. O. G.: The Thermodynamic Structure of Summer Arctic Stratocumulus and the Dynamic Coupling to the Surface, *Atmospheric Chemistry and Physics*, 14, 12 573–12 592, <https://doi.org/10.5194/acp-14-12573-2014>, 2014.
- Sotiropoulou, G., Sullivan, S., Savre, J., Lloyd, G., Lachlan-Cope, T., Ekman, A. M. L., and Nenes, A.: The Impact of Secondary Ice Production on Arctic Stratocumulus, *Atmospheric Chemistry and Physics*, 20, 1301–1316, <https://doi.org/10.5194/acp-20-1301-2020>, 2020.
- Stevens, R. G., Loewe, K., Dearden, C., Dimitrelos, A., Possner, A., Eirund, G. K., Raatikainen, T., Hill, A. A., Shipway, B. J., Wilkinson, J., Romakkaniemi, S., Tonttila, J., Laaksonen, A., Korhonen, H., Connolly, P., Lohmann, U., Hoose, C., Ekman, A. M. L., Carslaw, K. S., and Field, P. R.: A Model Intercomparison of CCN-limited Tenuous Clouds in the High Arctic, *Atmospheric Chemistry and Physics*, 18, 11 041–11 071, <https://doi.org/10.5194/acp-18-11041-2018>, 2018.
- Sulia, K. J. and Kumjian, M. R.: Simulated Polarimetric Fields of Ice Vapor Growth Using the Adaptive Habit Model. Part I: Large-Eddy Simulations, *Monthly Weather Review*, 145, 2281–2302, <https://doi.org/10.1175/MWR-D-16-0061.1>, 2017.
- Takahashi, T., Endoh, T., Wakahama, G., and Fukuta, N.: Vapor Diffusional Growth of Free-Falling Snow Crystals between -3 and -23°C, *Journal of the Meteorological Society of Japan*, 69, 15–30, https://doi.org/10.2151/jmsj1965.69.1_15, 1991.
- Tan, I. and Storelvmo, T.: Evidence of Strong Contributions From Mixed-Phase Clouds to Arctic Climate Change, *Geophysical Research Letters*, 46, 2894–2902, <https://doi.org/10.1029/2018GL081871>, 2019.
- Tan, I., Sotiropoulou, G., Taylor, P. C., Zamora, L., and Wendisch, M.: A Review of the Factors Influencing Arctic Mixed-Phase Clouds: Progress and Outlook, in: *Clouds and Their Climatic Impacts, Geophysical Monograph Series*, pp. 103–132, ISBN 978-1-119-70035-7, <https://doi.org/10.1002/9781119700357.ch5>, 2023.

- Verlinde, J., Harrington, J. Y., McFarquhar, G. M., Yannuzzi, V. T., Avramov, A., Greenberg, S., Johnson, N., Zhang, G., Poellot, M. R., Mather, J. H., Turner, D. D., Eloranta, E. W., Zak, B. D., Prenni, A. J., Daniel, J. S., Kok, G. L., Tobin, D. C., Holz, R., Sassen, K., Spangenberg, D., Minnis, P., Tooman, T. P., Ivey, M. D., Richardson, S. J., Bahrmann, C. P., Shupe, M., DeMott, P. J., Heymsfield, A. J., and Schofield, R.: The Mixed-Phase Arctic Cloud Experiment, *Bulletin of the American Meteorological Society*, 88, 205–222, <https://doi.org/10.1175/BAMS-88-2-205>, 2007.
- Wex, H., Augustin-Bauditz, S., Boose, Y., Budke, C., Curtius, J., Diehl, K., Dreyer, A., Frank, F., Hartmann, S., Hiranuma, N., Jantsch, E., Kanji, Z. A., Kiselev, A., Koop, T., Möhler, O., Niedermeier, D., Nillius, B., Rösch, M., Rose, D., Schmidt, C., Steinke, I., and Stratmann, F.: Intercomparing Different Devices for the Investigation of Ice Nucleating Particles Using Snomax[®] as Test Substance, *Atmospheric Chemistry and Physics*, 15, 1463–1485, <https://doi.org/10.5194/acp-15-1463-2015>, 2015.
- Wex, H., Huang, L., Zhang, W., Hung, H., Traversi, R., Becagli, S., Sheesley, R. J., Moffett, C. E., Barrett, T. E., Bossi, R., Skov, H., Hünnerbein, A., Lubitz, J., Löffler, M., Linke, O., Hartmann, M., Herenz, P., and Stratmann, F.: Annual Variability of Ice-Nucleating Particle Concentrations at Different Arctic Locations, *Atmospheric Chemistry and Physics*, 19, 5293–5311, <https://doi.org/10.5194/acp-19-5293-2019>, 2019.
- Wood, R.: Stratocumulus Clouds, *Monthly Weather Review*, 140, 2373–2423, <https://doi.org/10.1175/MWR-D-11-00121.1>, 2012.
- Xu, X. and Xue, H.: Impacts of Free-Tropospheric Temperature and Humidity on Nocturnal Nonprecipitating Marine Stratocumulus, *Journal of the Atmospheric Sciences*, 72, 2853–2864, <https://doi.org/10.1175/JAS-D-14-0387.1>, 2015.