

How ice apron loss and permafrost degradation ~~promote~~promoted the Platteikogel rock ~~slides~~slope failure: A thermo-mechanical reconstruction

Felix Pfluger¹, Samuel Weber^{2,3}, Natalie Barbosa^{1,4}, Florentin Hofmeister⁵, Johannes Leinauer¹, Peter Wegmann⁶, and Michael Krautblatter¹

¹Landslide Research Group, TUM School of Engineering and Design, Technical University of Munich, Munich, Germany

²WSL Institute for Snow and Avalanche Research, SLF, Davos Dorf, Switzerland

³Climate Change, Extremes and Natural Hazards in Alpine Regions Research Center CERC, Davos Dorf, Switzerland

⁴Department of Earth and Environmental Sciences, Faculty of Earth Sciences, GeoBio Center, Ludwig Maximilians University, Munich, Germany

⁵Bavarian Academy of Sciences and Humanities, Geodesy and Glaciology, Alfons-Goppel Str. 11, 80539 Munich, Germany.

⁶Department of Computer Science, TUM School of Computation, Information and Technology, Technical University of Munich, Munich, Germany

Correspondence: Felix Pfluger (felix.pfluger@tum.de)

Abstract.

The Alpine cryosphere changes at unprecedented speed, affecting the thermal, hydrological, and mechanical state and behaviour of rock slopes. While numerous studies investigated singular drivers for progressive rock slope failures, the knowledge of hydro-thermo-mechanically coupled processes remains scarce. In this manuscript, we investigate the 2024 permafrost rock

5 ~~slide~~slope failure at Platteikogel with a volume of 50,000 m³ (3,395 m a.s.l., above Vernagtferner, Austria). We aim to assess how observed ice apron loss and related permafrost warming promote the release mechanism. We reconstructed multidecadal thermal evolution accounting for the thermal impact of ice apron loss. Based on field observations, we derived a conceptual model on how ice apron loss potentially affects rock slope destabilization. Integrating the outcome of the preceding steps, we performed a mechanical stability analysis assuming that the rock ~~slide~~slope failed along ice-filled discontinuities. The me-

10 chanical model indicates that the ~~rock slide~~failure can not be solely explained by a warming-driven decrease in shear strength of ice-filled discontinuities, suggesting that other failure processes superimpose or even dominate. The implemented system feedback related to ice apron loss suggests that hydrostatic pressure buildup due to water infiltration and rockfall-induced unloading thereby promoted the Platteikogel rock ~~slide~~releaseslope failure. In summary, we demonstrate that ice apron loss not only leads to increased rockfall activity but also accelerates progressive ~~rock slope~~failure, promoting the ~~release of rock~~

15 ~~slides~~detachment event. In upcoming decades, ice aprons on steep rock slopes above 3000 m in the European Alps are expected to experience drastic area loss, exposing potential source zones for future rock ~~slides~~slope failures.

1 Introduction

Recent rock slope failures in the cryosphere of the European Alps underscore both the anticipated increase in frequency under a warming climate (Intergovernmental Panel on Climate Change, 2022) and the complexity of the failure processes: Notable examples include the permafrost rock slide at Fluchthorn (Austria; 2023; Krautblatter et al., 2024), the rock slide beneath glacier ice at Piz Scerscen (Switzerland; 2024; Pierhöfer et al., 2024), and the complex glacier failure at Blatten which was preceded by permafrost-related rockfalls (Switzerland; 2025; Islam et al., 2025). All of these failures evolved into multi-kilometer, highly mobile rock–ice avalanches, illustrating the potential — or, in the case of Blatten, the actual — threat posed to valley populations by cascading processes.

While rock slope destabilization mechanisms have been investigated for changes in glaciers (Fischer et al., 2010; Rechberger and Zangerl, 2022; Walden et al., 2025) or permafrost (Gruber and Haeberli, 2007; Phillips et al., 2017; Etzelmüller et al., 2022), less attention has been drawn towards ice aprons. Ice aprons - defined as irregularly shaped, small ice bodies - typically less than 0.1 km² in area and located on slopes steeper than 40° - undergo rapid area loss in the 21st century, and have almost exclusively been studied in the Western Alps (Ravanel et al., 2023). While glaciers retreat from the bottom up, the upper boundary of ice aprons shifts top-downwards, as observed during continuous area loss. Over a 70-year observation period (1952–2019) in the Mont Blanc Massif, the area of ice aprons has declined by 47% (Kaushik et al., 2022). Ice apron loss is hypothesized to negatively impact the stability of permafrost rock slopes by inducing thermo-mechanical alteration upon their disappearance (Guillet and Ravanel, 2020), however, the actual processes remain poorly constrained.

In this paper, we aim to decipher the failure mechanism of the Plateikogel rock ~~slide-slope failure~~ (3,395 m a.s.l., release volume of 50,000 m³, spring 2024, situated in the Vernagtferner Basin, Tyrol, Austria). The rock ~~slide-mass~~ detached from a ridge flanked by glaciers. Ice aprons reached the detachment area in 1970 but have since lost substantial elevation. We investigate the destabilization of the rock slope combining three complementary steps: (i) Analysis of visible cryospheric and geomorphic changes, (ii) modeling decadal permafrost evolution using the conductive heat flow model (CryoGrid 2D; Czekirka et al., 2023), and (iii) modeling rock slope mechanics with distinct element code (UDEC; Itasca Consulting Group, 2019), integrating (i) and (ii) to assess the mechanical impacts associated with the paraglacial transition (Fig. 1).

This ~~cross-disciplinary-multi-method~~ study is the first to apply a laboratory-derived rock–ice mechanical shear model to an actual rock slope failure at slope scale, challenging its applicability to larger scales. Furthermore, we demonstrate the thermal impact of ice apron loss on permafrost warming and propose a conceptual model describing the feedbacks between ice apron loss and rock slope destabilization. This study aims to provide an integrated understanding of the coupled thermo-hydro-mechanical processes that promote rock ~~slides-slope failures~~ in permafrost.

We address the following questions:

- (i) How does the area loss of the ice apron affect the thermal state of permafrost at the Plateikogel?
- (ii) Assuming ice-filled fractures, did climate-driven permafrost warming contribute to the destabilization and release of the rock slope?

50 (iii) Which other mechanisms were relevant in the final phase of slope failure?




Figure 1. Conceptualized workflow of this study, which iterates from (i) observing surface ice changes and geomorphological changes to (ii) permafrost modeling to (iii) mechanical investigations of rock slope stability. The expected changes Δ within the observation period of more than four decades are illustrated. Abbreviation: A.. Area, T.. Temperature, FoS.. Factor of safety. Arrows indicate result transfer used as input for subsequent modeling.

In this paper, we follow the ~~process- and material-based~~ landslide terminology of [Hungr et al. \(2014\)](#). ~~Pre-failure refers to the period of initial deformation and progressive rock mass damage, typically at millimeters to centimeters per year, which may culminate in the rockslide release (sudden failure), marked by accelerated displacement and full detachment along a distinct shear plane, potentially evolving into a rock avalanche. Post-failure refers to any geomorphic activity in the area affected by the sudden failure~~ [Hermanns et al. \(2022\)](#) and use the term *rock slope failure* to describe landslides with detachment in rock, occurring days, months, or even years afterward ([Leroueil et al., 1996](#)) independent of failure kinematics. This term encompasses the pre-failure deformation and associated geomorphic activity, the main failure event (sudden failure), and the post-failure deformation or related geomorphic activity. The spatial data and time series used for the analysis presented in this publication are summarized in the Tables A1 and A2.

60 2 Characterization of the field site

2.1 The Platteikogel rock ~~slide~~ slope failure

The Platteikogel rock ~~slide~~ slope failure (46°52'14.2"N 10°50'46.2"E) ~~detached from~~ occurred on a NE-SW-oriented mountain ridge at 3 400 m a.s.l. surrounding the Vernagtferner glacier, Ötztal, Tyrol, Austria ([Fig. 2a](#); Glacier ID: 489; ~~World Glacier Monitoring Service~~ [WGMS.2025](#)). ~~After detachment, the rock slide transitioned into a rock avalanche.~~ The affected area and deposits are shown ~~in the aerial photograph of the post-failure state~~ on aerial photographs and the slope map in Figure 2b. The

exact timing of the event remains unclear and can only be narrowed down to the period between 25th April and 6th June 2025 (Sentinel-2 L2A; European Space Agency, 2024). No seismic or hydrological recordings in the near surroundings indicate any unambiguous signals related to the event. Landslide-related metrics are given in Table 1.

Table 1. Classification and metrics regarding the Platteikogel rock ~~slide/rock-avalanche~~slope failure.

Landslide Metrics	
Pre-failure elevation of ridge	3395 m
Elevation of frontal runout deposits	3120 m
Total elevation difference H	275 m
Length of total runout L	600 m (Basal contact: 100 m on bedrock, 500 m on glacier surface)
H/L ratio	0.45
Fahrböschungswinkel (overall-slope-reach angle) β	24.62°
<u>Affected total area including deposits A</u>	<u>48,000 m²</u>
Minimum estimated volume loss at detachment area* V	50,000 m ³
	1.04
Affected-total-area A 48,000-m² -V/A ratio	
<u>Estimated average depth of basal shear plane**</u>	<u>20 m (range: 15–25 m)</u>
Center of Gravity: ΔCoG for pre-/post-failure state	dx=85, dy=100 m
Potential Energy of event** $E_{pot} = \rho * V * g * dy$	127 * 10 ⁹ J (TNT equivalent 30,000 kg)

*Change detection was conducted for the point clouds of 2023 and 2024 (each 2 cm resolution) at the detachment area ~~of the rock-slide~~ only. The volume error is smaller than 100 m³. **assumed $\rho = 2600 \text{ kg m}^{-3}$. **The location and shape of the basal shear plane are shown in the profile in Figure 6a.

The geology at the detachment area consists of ~~Schistose~~-schistose gneiss with feldspar augen and staurolite minerals,
 70 enclosing a local intersection of ~~Muscovite~~-muscovite schist in the direct vicinity of the South of the detachment area (Kreuss et al., 2012). Structural geology was likely to favour the destabilization of the Platteikogel rock slope (Fig. A1 - analysis of rock outcrops): The pronounced foliation of the metamorphic rock is visible from afar, marking the many incised ~~steep~~-gullies in the steep wall - Fig. 2**a**). The foliation exhibits a general dip direction toward S/SSE and steep dip angles ranging from 60° to 85°. Moreover, two prominent, non-foliation parallel joint sets were identified, both striking nearly perpendicular to the direction
 75 of rock slope displacement. Their dip directions are northwest and southeast. General joint spacing ranges from centimeters to several decameters up to meters and varies locally. Although the dip angles of both joint sets (ranging between 60° to 80°) are generally steeper than the slope angle, suggesting that sliding is inhibited by geometry, the narrow joint spacing likely favored a step-path shear failure (Eberhardt et al., 2004). Slab-shaped, disintegrated, angular blocks characterize the detachment area at post-failure state (Fig. 2**b**).

80 2.2 Cryospheric changes and pre-failure rockfall activity

Glaciers surround both sides of the ridge of the detachment area, with ice aprons extending steeply upwards below the ridge crest. Given that a dynamic cryosphere affects rock slope mechanics through changes in water availability, temperature, and local stress field (ice loss/rockfalls), the subsequent paragraphs focus on the evolution of the cryosphere in the area of the detachment.

85 The detachment area lies within permafrost (100% probability of permafrost occurrence according to permafrost distribution map; Otto et al., 2020). Moreover, the presence of ice aprons indicates subzero rock surface temperatures (Benn and Evans, 2014). Figure 3 (a-c) demonstrates the visible changes of ice aprons at the detachment area in past decades (ice apron typology: *steep ice apron above glacier*, cf. Fig. 1 (4) in Ravel et al., 2023): Between 1970 and 1999, the upper limit of ice aprons decreased to the position of the former Bergschrund, losing more than 50 m in elevation at the southeastern flank of the ridge.
90 In the period 1999 to 2019, the recent ice-free bedrock exhibits incised gullies, while debris cones accumulated on the glacier below. In contrast, the ice apron at the northwestern flank exhibits less loss in area, but shows overall derogation, with widening of the berschrund, and opening of crevasses underneath in 1999. For the period analyzed (1970-2019), the Bergschrund at the northwestern flank remained almost in a stationary position (± 5 m).

Analyzing surface ice elevation changes of Kleiner Vernagtferner located northwest of the ridge in the years before failure
95 from 2015 onwards, a rate of elevation change $> 3\text{ m yr}^{-1}$ was calculated for the lower part of the glacier, while a rate of elevation change of $\approx 0.3\text{ m yr}^{-1}$ was calculated for the area of ice aprons. Similar values were derived for the Platteiferner located southeast of the ridge (Fig. 3 d-f). While ice aprons were in direct contact with the rock in the detachment area, both glaciers were situated more than 50 m below. Focusing only on the uppermost area of ice aprons, the analyzed data suggests a loss of at least 5 m of ice apron thickness in the decade before the failure in 2024.

100 Together with observed changes in surface ice, rockfall activity was evident in the detachment area since 2015 onwards (Fig. 3g-i): Between 2015–2018, seven rockfalls occurred (561 m² total; 5–311 m² each). No events were detected from 2018–2021. From 2021–2023, fourteen rockfalls occurred (412 m² total; 2–236 m² each).




Figure 2. (a) Overview of the location and extent of the Platteikogel rock slope failure situated in the cirque of the Kleiner Vernagtferner. (b) Aerial view from NW showing the full dimensions of the affected area at post-failure state, and (eb) zoom-in to the detachment area. Both photographs (ba and eb) were captured by a UAV on 8 August 2024. (ba) From the macro perspective, the steep and vertical gullies incised in the headwall of the Vorderer Brochkogel summit clearly demonstrate the tectonic structure of the foliated metamorphic rock mass. (eb) The shape of the individual boulders and blocks of the disintegrated rock mass is slab-like according to the foliation structure. Distinct zones with a high concentration of fine grains are visible in the detachment zone. (c) Overview of the location and extent of the Platteikogel rock slope failure situated in the cirque of the Kleiner Vernagtferner. (d) Slope map at pre-failure state, including the profiles used for modeling studies. (e) Slope map at post-failure state showing the morphology of deposition, and the change detection (threshold of detection: 5 m). Data source: (ac) Orthophotography acquired by Land Tirol - <https://www.tirol.gv.at/data/> (last access: 7 January 2025).




Figure 3. (a-c) Changes in ice cover 5 decades before failure, (d-f) surface ice elevation change, and (g-i) affected rockfall area years before failure shown for the area of Plattekogel rock slope failure (for location of analyzed area see Fig. 2a). The spatio-temporal surface ice changes (d-f) and rockfall inventory (g-i) were processed following the approach for multi-temporal quantification of surface changes described in the supplement of Barbosa et al. (2024). (d-f) Glacier outlines were mapped manually on orthophotos with 20 cm resolution for the respective years. The limit of detection was set to -0.6 m. Note: Due to changes made to the camera system in the 2021 campaign, (e) shows artifacts due to the suboptimal model alignment, such as the pronounced scattering at the SE exposed slopes in the lower right. (g-i) Spatial distribution of rockfall activity at the ridges surrounding the Kleiner Vernagtferner. Rockfalls are manually mapped using geomorphic change detection from a DSM derived from aerial imagery at 20 cm spatial resolution. We used a 100 m grid to display the cumulative rockfall area in m² per time interval. The glacier extent was manually mapped. Data sources: (a-c) orthophotography acquired by Land Tirol - <https://www.tirol.gv.at/data/> (last access: 7 January 2025), (d-i) large-format aerial imagery acquired by 3D RealityMaps GmbH.

3 Methodological approach

105 An extended map view, including the positions of all meteorological stations from which data were used for the thermal modeling conducted subsequently, is shown in Figure SM1 - the label SM denotes supplementary material. The cross-section selected for the modeling studies follows the direction of mass movements and runs perpendicular to the mountain's ridge. Its topography was inferred from the pre-failure digital elevation model of 2023 (Table A1, [i, for location see Fig. 2d](#)) and was smoothed to a step width of 5 m.

3.1 Modeling the thermal evolution of the mountain ridge

110 We use the [transient](#) conductive heat flux model CryoGrid 2D, which was applied in other studies on permafrost evolution in steep rock walls (Myhra et al., 2017, 2019; Czekirda et al., 2023), in order to reconstruct permafrost conditions in the decades before the rock ~~slide-release~~ [slope failure event](#) in 2024. [With Cryogrid 2D, the subsurface temperature field is calculated by solving the heat diffusion equation following Fourier's law of heat conduction according to defined material- and temperature-dependent parameters. The finite element solver MILAMIN package \(Dabrowski et al., 2008\) was employed](#)
115 [to numerically solve complex geometries on unstructured grids, based on specified boundary conditions and the imposed temperature forcing along the model topography. Time discretisation follows a finite-difference backward Euler scheme.](#) A detailed description of [the](#) CryoGrid 2D [model](#) is provided by Myhra et al. (2017). Here, we use the Cryogrid 2D version as applied by Czekirda et al. (2023). The modeling strategy follows their approach and is outlined below.

3.1.1 Model calibration and setup

120 First, we calibrated the thermal parameters with measured borehole data: We assumed uniform rock mass properties (single lithology according to field observations). The volumetric fraction of rock and water was set to 0.95 and 0.05, accounting for fractured and jointed gneissic rock mass (Myhra et al., 2017, 2019). We calibrated the parameters thermal conductivity k , and volumetric heat capacity of c_v with measured temperatures of the borehole at Matterhorn (PERMOS, 2024), which has similar lithology and altitude to the Platteikogel detachment area. Therefore, we utilized a simplified column mesh and
125 horizontal model topography, and applied a uniform heat flux of 50 mW m^{-1} at 6000 m depth. The model was then forced using measured temperatures at 0.1 m depth. Measured borehole temperatures at greater depths were compared with the simulated temperature profiles (Fig. A2,a1-a4). The configuration of parameters resulting in the best model-fit (compare Fig. A2a1-a4) was selected for all further simulations: $k = 2 \text{ W K}^{-1} \text{ m}^{-1}$, $c_v = 3 \cdot 10^6 \text{ J m}^{-3} \text{ K}^{-1}$. For the simulation regarding the Platteikogel rock ~~slide~~ [slope failure](#), we constructed an unstructured triangular mesh using the smoothed cross-section profile
130 of the mountain ridge by using the Triangle library (Shewchuk, 1996). The node density was decreased gradually with higher depth (see Table SM1). Aforementioned calibrated parameters were prescribed to the entire Platteikogel mountain.

3.1.2 Applied forcings

Figure 4 illustrates the cross-section with varying surface types, indicating their thermal functionality regarding permafrost in the rock mass below. To simulate the thermal evolution of the Plattekogel ridge, we forced the model using mean monthly rock surface temperatures (RST), which were projected along the rock topography. RSTs are obtained from air temperatures (AT); their difference defines the surface offset (SO). First, we created a long-term dataset of monthly AT for the site of Plattekogel: Mean monthly lapse rates were calculated using the temperature records (2003-2024) from two nearby meteorological stations (i), at 2863.9 m a.s.l., and (ii), at 3437 m a.s.l. (GeoSphere Austria, 2024). Second, linear regression models were applied to extrapolate the temperature time series for station (ii), using (iii) historical mean monthly air temperature records from long-term monitoring station Obergurgl (Auer et al., 2007), which is located at 1938 m a.s.l. in the neighboring valley. This resulted in a mean monthly AT time series for the period 1900 to 2024, corresponding to the approximate elevation of the ~~rock slide's~~ detachment area. Thirdly, the calculated lapse rates were used to adjust the monthly AT values to the corresponding elevations along the profile (for station details see Table A2, i-iii).

On the basis of AT, we inferred RST for varying surface types along the profile - bedrock, snow cover, ice aprons, and glacier ice - by using temperature transfer functions. Accounting for natural variability and uncertainty in SO, we specified a plausible range of values for the sampling of ensemble simulations. These functions are explained in the following. Examples for the application of these functions are demonstrated in Figure SM2a-e.

(i) Seasonal snow cover: Seasonal snow cover acts as a thermal insulator, buffering cold AT signals (Haberkorn et al., 2015). Snow reduction factors nF [-] (Smith and Riseborough, 2002; Gislén et al., 2013) were multiplied with negative AT for months with snow cover only (assumed period of snow cover: 1st November to ~~31st~~ 31 May, Fig. SM2b). For positive AT in this period, we enforced an isothermal snow cover with 0°C . ~~We~~ Following (Czekirda et al., 2023), we used the slope of the profile to assess the nF -factors ranging from 0.5 for slope $< 30^{\circ}$ to 1 for slope $> 60^{\circ}$ along the profile topography (see Fig. SM3; analogous to Czekirda et al., 2023). and calculated RST below the snow cover using the empirical transfer function (see Fig. SM3):

$$\text{RST}_{\text{snow cover}} = \begin{cases} AT \cdot (nF(\text{slope}) + x) [^{\circ}\text{C}], & \text{if } AT < 0^{\circ}\text{C}, x \in [-0.1, 0.1], \\ 0^{\circ}\text{C}, & \text{if } AT \geq 0^{\circ}\text{C}. \end{cases} \quad (1)$$

(ii) Bedrock: Snow-free, sun-exposed rock surfaces undergo significant radiative warming (Magnin et al., 2019). For snow-free locations with either slope angles $> 60^{\circ}$ or during the snow-free months, we calculated the RST using fixed temperature offsets accounting only for the aspect of the mountain flank (Fig. SM2b). The here defined surface offset range chosen for radiative warming of rock surfaces is consistent with measured data (Czekirda et al., 2023, rock walls in Norway), and calculated RST-AT offsets on the basis of measurements from the Matterhorn Hörnli ridge, Switzerland, 3500 m a.s.l. (see Fig. SM4, data source: Weber et al., 2024).

$$RST_{\text{exposed rock}} = \begin{cases} AT + SO_{NW} [^{\circ}\text{C}], & SO_{NW} \in [-1, +1], \\ AT + SO_{SE} [^{\circ}\text{C}], & SO_{SE} \in [+2, +4]. \end{cases} \quad (2)$$

(iii) Ice aprons: Ice aprons insulate the rock in summer due to the lower thermal diffusivity of warm ice, and limit warming above 0°C by latent heat consumption during the surface melt. In winter, they enhance the cooling of the bedrock by the relative increase of diffusivity of colder ice (James, 1968). Temperatures measured at depths between 2.5 and 8.8 m of a NNE-facing ice apron at 3470 m a.s.l. at Tour Ronde (France) show that mean annual temperatures strongly converge at $-5 \pm 0.5^{\circ}\text{C}$ regardless of the depth in the mentioned range (Ravel et al., 2023). While the seasonal air temperature signal is pronounced at shallow depths, it exhibits only a marginal amplitude at 8.8 m depth. For the modeling strategy, we assume that ice aprons, for their entire vertical extension, consist of uniform thickness. To account for the thermal effect of ice aprons on the rock surface underneath, we implemented a low-pass filter on the basis of the analytical solution of the 1D conductive heat equation (Carslaw and Jaeger, 1980), which dampens monthly fluctuations of AT with increasing ice thickness, while annual signals penetrate deeper (Fig. SM2c; approximating the observed results of Ravel et al., 2023). Using properties of ice: $\rho_{ice} = 917 \text{ kg m}^{-3}$, $k_{ice} = 2.1 \text{ W m}^{-1} \text{ K}^{-1}$, and $c_{ice} = 2009 \text{ J kg}^{-1} \text{ K}^{-1}$ and assuming uniform thickness of ice $H_{ice} \in [1, 9] \text{ m}$ throughout the entire simulation time, we calculate the damping factor: $e^{-H_{ice}/\delta}$, with penetration depth $\delta = \sqrt{2\kappa/\omega_y}$, $\omega_y = 2\pi/(365 \cdot 24 \cdot 3600) \text{ s}$ and diffusivity $\kappa = k_{ice}/(\rho_{ice} \cdot c_{ice})$. The formula used to calculate the monthly temperature at the rock-ice apron interface is

$$RST_{\text{ice aprons}} = \min \left(TIAS_{\text{mean}} + e^{-H_{ice}/\delta} [TIAS_m - TIAS_{\text{mean}}], 0^{\circ}\text{C} \right) \quad (3)$$

$$H_{ice} \in [1, 9] \text{ m}$$

Note that the minimum statement introduced in the formula limits the monthly temperature at the rock-ice apron interface to a maximum of 0°C. Here, TIAS refers to the Temperature at the Ice Apron Surface specified for a location of 0.1 m below the current ice surface (analogous to the standard depth specified for RST measurements). It serves to calculate the $RST_{\text{ice aprons}}$ and is calculated as follows.

Eq. 3a: Monthly temperature at ice apron surface...

$$TIAS_m = \begin{cases} RST_{\text{snow}}(m), & \text{if } m \in \{1, 2, 3, 4, 5, 11, 12\}, \text{ snow cover on top of ice apron} \rightarrow \text{Eq. 1,} \\ AT(m), & \text{otherwise, ice apron surface exposed} \rightarrow \text{no surface offset;} \end{cases}$$

and Eq. 3b: Mean annual temperature at ice apron surface calculated for each year respectively...

$$TIAS_{\text{mean}} = \frac{1}{12} \sum_{m=1}^{12} TIAS_m.$$

(iv) For the glacier below the ice aprons - the Bergschrund marks the transition - we assumed rather thin glacier ice in the order of a few decameters, with marginal movement. However, the glacier may distinctly differ from the ice apron by its greater thickness of the ice body. In the areas near the headwalls of the cirque, the ice might still be frozen to bedrock (Benn and Evans, 2014). For the simulation, we assume thick perennial snow cover on top of the glacier, using a fixed nF -factor throughout the entire simulation time, and suggest that the ice thickness is well beyond the depth of the seasonal penetration signal (Fig. SM2d). Monthly glacier-bed temperatures were calculated from the mean annual air temperature (MAAT) as:

$$RST_{glacier} = MAAT \cdot nF^{\circ C}, \quad nF \in [+0.4, +0.6] \quad (4)$$

Apart from AT, we considered the gradual retreat in ice apron since 1970 (Fig. 2a-c and Fig. 4) as a dynamic variable in our model, while snow cover, solar radiation, and glacier extent were treated as static throughout the simulation period. We therefore implemented a linearly decreasing ice-bedrock boundary for the southeastern flank from 1970 at an elevation of 3355 m a.s.l. to 1999 at an elevation of 3290 m a.s.l. For the northwestern flank, the ice-bedrock boundary decreased from 1970 at an elevation of 3365 m a.s.l. to 2024 at an elevation of 3310 m a.s.l. (Fig. SM5). In our simulation, only two states are considered: Ice aprons of constant thickness throughout their existence and extent, or ice-free surfaces following top-down retreat. For the newly exposed ice-free surfaces, RSTs were calculated respecting snow cover and radiative bedrock warming (Eqs. 1 and 2; Fig. SM2e). In contrast to the varying ice apron extent, the glacier in the area of the cirque, delimited by the observed stationary bergschrund (± 5 m in horizontal direction between 1970 and 2023), was considered stationary, and its temporal variations in thickness were ignored. Before 1970, we assumed the glacier and ice apron extent to be identical to the 1970 extent.

3.1.3 Simulation strategy

The model was initialized with the mean annual RST of 1900 (calculated with the mean of the values in defined ranges in Eqs. 1 to 4) along the model topography and the geothermal heat flux at the bottom boundary until a steady state was reached within the model domain (marginal difference of modeled temperature between consecutive years of less than $10^{-4}^{\circ C}$). Starting from the initialized state in 1900, we ran 100 individual simulations until 2024. Each with randomized offset parameters chosen from specified ranges in order to account for uncertainties through the use of temperature transfer functions. Each simulation was forced with RST calculated on basis of randomly sampled offset parameters within specified ranges: Eq. 1: $x \in [-0.1, 0.1]$; Eq. 2: $SO_{NW} \in [-1, +1]$, $SO_{SE} \in [+2, +4]^{\circ C}$; Eq. 3: $H_{ice} \in [1, 9]$ m; Eq. 4: $nF \in [+0.4, +0.6]$. The full workflow following (a) model calibration, (b) setup, (c) surface forcings, to (d) simulation strategy is comprised in Figure A2.

3.2 Mechanical modeling of the failure mechanism

3.2.1 The mechanical implications of ice apron loss

Based on our observations of ice apron retreat (Section 2.2), we derive a conceptual model emphasizing the coupled effects on (i) permafrost degradation, (ii) hydrogeology, and (iii) topographic modification through rockfall activity (Fig. 5), which in turn




Figure 4. Cross-section through the ridge of Platteikogel, demonstrating varying surface types and the observed downslope retreat of ice apron since 1970 onwards - Given dates/elevation marks are inferred from historic orthophotographs. The topography (upper boundary) and lower boundary (implied, at 6000 m depth) mark the frame for the meshed model. Note: The vertical scale of the cross-section is exaggerated by a factor of 2.

influences rock slope stability. The concept explained here serves to define the simulation scenarios for the rock mechanical analysis of the Platteikogel rock slideslope failure.

We briefly address the consequences of ice apron loss on the various systems: (i) Upon ice apron retreat, newly exposed surfaces are subject to increased radiative heating and sensible heat exchange (Deline et al., 2015). From now on, an active layer might seasonally be formed, enhancing permafrost degradation. (ii) Ice that once sealed the bedrock, preventing rain or meltwater from infiltration, vanishes and makes the uppermost bedrock more permeable. As a result, water infiltration can lead to the buildup of hydrostatic water pressure within the rock mass (Offer et al., 2025; Scandroglio et al., 2025). (iii) The upper meters of exposed bedrock experience a thermal shock by regular freeze-thaw cycles, which lead to accelerated fatigue of rock and weathering processes (Jia et al., 2015), resulting in increased rockfall events (Draebing and Mayer, 2021).

The system exhibits a strong feedback loop with coupled interdependencies: In fractured permafrost rock, both conductive and advective thermal transport processes are relevant (i ↔ ii). The latter typically channels energy transport by water flow paths along fractures, forming local thaw corridors resulting in heterogeneous permafrost zones (Hasler et al., 2011; Magnin and Josnin, 2021). Hydrostatic pressure (ii) mechanically widens joint walls (Witherspoon et al., 1980; Ji et al., 2013), en-

hancing flow paths and concentrating thermal energy transport (i) or releasing rockfalls (iii) (Krautblatter and Moser, 2009) (Krautblatter and Moser, 2009; Kuhn et al., 2025). (iii) Rockfalls modify the surface, exposing deeper rock to atmospheric conditions and reinforcing thaw (i) or channel infiltration of surface water (ii).

230 The aforementioned consequences of ice apron loss - enhanced warming of permafrost, induced hydrostatic pressure to previously frozen area or changes of local stress state by rockfalls, decrease the mechanical stability of rock slopes by reducing the rock slope's strength (i) or enhancing driving forces (ii & iii; Krautblatter et al., 2013). For the purpose of analyzing the failure mechanism of the Platteikogel rock slideslope failure, we integrated these concepts into a mechanical modeling study.




Figure 5. Conceptual model indicating the impact of ice apron loss on (i) permafrost, (ii) hydrogeology, (iii) geomorphic processes. The glacier and ice apron thickness is sketched arbitrarily. The lower right comprises the processes that are investigated by mechanical modeling in this paper. Note: (i) Contrast in albedo (snow/ice $\approx 0.2\text{--}0.9$; rock $\approx 0.05\text{--}0.2$) and thermal response (ice limited to 0°C ; rock potentially warms beyond melting point). (ii) Very low ice permeability ($k_{ice} \approx 10^{-18}\text{m}^{-2}$). (iii) Topographic change and mass loss may raise effective stress σ_{eff} , beyond critical stress σ_{crti} .

3.2.2 Model setup

235 We use the 2D mechanical modeling framework UDEC (Universal Distinct Element Code) by Itasca Consulting Group (2019) to analyze the mechanics promoting the ~~release of the Platteikogel rock slide~~major rock detachment. UDEC employs the distinct element method to simulate rock masses as discrete blocks, defined by discontinuities such as joints or faults. These discontinuities act as contact boundaries during simulations, allowing sliding, toppling, or rotation of individual blocks. We used the same cross-section as in the CryoGrid 2D simulation and projected the geometry of joint sets, inferred from the 2024
240 point cloud analysis, into the model domain (see Fig. A1 and Section 2.1). The joint spacing was upscaled to 10 m. On this basis, we created two model setups accounting for different structural geometry (Fig. 6a). For setup A, the location of the

basal shear plane inferred from post-failure DSM was explicitly integrated in the model, while for setup B, we adopted an implicit approach: To account for natural irregularities, the spread of the varying joint angles, we introduced a joint set based on squeezed Voronoi polygons (i.e., see Gerstner et al., 2023). This configuration facilitated free deformation and sliding of the model without prescribing a specific basal shear plane. By integrating these structures, the model domain was subdivided into more than 5800 discrete blocks. The discrete blocks were meshed with a maximum edge length of 5 m, creating triangular zones, and were assigned linear elastic block models. The rounding length of block corners was set to 0.5 m in order to minimize computation time and allow for the rotation of blocks. We fixed the left, bottom, and right model boundaries with no-velocity conditions and assigned a gravitational acceleration of 9.81 ms^{-2} . The elastic blocks were assigned parameters for density, bulk- and shear-modulus of $\rho_{rock\ mass} = 2600 \text{ kg m}^{-3}$, $K_{rock\ mass} = 4 \text{ GPa}$, and $G_{rock\ mass} = 1 \text{ GPa}$, which were estimated after Hoek and Brown (2019), as shown in Table A3. The block parameters were kept constant for all simulations, as shallow rock slides with a basal shear plane in the upper decameters below the surface are mainly dominated by shearing along discontinuities, rather than deformation of brittle intact material. The contact surfaces of the blocks were governed by a Mohr-Coulomb shear model, with varying shear parameters accounting for different conditions of the contact areas, such as surface roughness, joint-infillings, or intact rock bridges (Gischig et al., 2011; Mamot et al., 2021; Rechberger and Zangerl, 2022).

3.2.3 Simulation strategy

We initialized a base model ~~with defining an in-situ stress ratio of $k = 0.5$ ($0.5 \sigma_y = \sigma_x = \sigma_z$) prior to the simulation start~~ (cf. Fischer et al., 2010; Rechberger and Zangerl, 2022), assigned shear parameters of $\phi = 40^\circ$ and $c = 0.5 \text{ MPa}$, and ran the simulation until reaching mechanical equilibrium. For all further scenarios simulated (Fig. 6b), this base model was used as the starting point. To assess stability, we cycled 30,000 ~~mechanical time-model~~ steps, studying overall model deformation and monitoring the displacement of specified locations within the rock mass throughout the simulation time.

3.2.4 Scenarios and implementation

In a first step, referred to as scenario S0, we tested the rock slopes' predisposition to failure by accounting for ~~varying-model geometry-different structural model configurations~~ (setup A & B) and back-calculated the rock slopes' hypothetical pre-failure condition by conducting a sensitivity test on varying pairs of shear parameters applied to all discontinuities throughout the model domain (i.e., analogous to Rechberger and Zangerl, 2022). The applied Mohr-Coulomb shear criterion relates shear stress τ [Pa] to normal stress σ [Pa] multiplied by the tangent of the friction angle ϕ [$^\circ$] and adding cohesion c [Pa] as an intercept.

$$\tau = \sigma \cdot \tan(\phi) + c, \quad \phi \in [20, 40]^\circ, c \in [0.0, 0.5] \text{ MPa} \quad (5)$$

Based on considerations of possible promoting factors or triggers (Section 3.2.1, Fig. 5), we examine the following scenarios, aiming to investigate the mechanical response to a changing cryosphere. ~~These scenarios represent quasi-static realizations of~~

the rock slope under specific states or external forcings (Fig. 6b) and therefore illustrate mechanical system responses to isolated effects rather than complex interwoven system dynamics.

275 S1 Permafrost degradation

Discontinuities in permafrost rock are often filled with ice (Gruber and Haeberli, 2007; Krautblatter et al., 2013; Zangerl et al., 2019). The temperature of these ice-filled fractures or joints controls the shear strength of contact surfaces (Mamot et al., 2018, 2020; Huang et al., 2023). To test the mechanical response to permafrost degradation, we performed a unidirectional thermo-mechanical simulation, where the output of the thermal model (CryoGrid 2D) was translated into the mechanical model (UDEC). We assigned temperature-dependent shear parameters $\phi(T), c(T)$ to discontinuities in the corresponding temperature regions resulting from the CryoGrid 2D simulation. Temperature zones from the CryoGrid 2D model output were replicated with regions in UDEC, whose discontinuities were assigned the corresponding temperature-dependent shear criterion. The herby used temperature-dependent Mohr-Coulomb shear criterion was derived from shear tests on rock-ice-rock sandwich samples and was proposed for ice-filled joints in permafrost rock by (Mamot et al., 2018). In their equtation τ and σ are expressed in [kPa]:

$$\tau(T) = \sigma \cdot ((0.42 \pm 0.15) - (0.21 \pm 0.06) \cdot T) + (53.3 \pm 20.6 - (73.5 \pm 8.8) \cdot T) \quad (6)$$

The equation incorporates temperature-independent (subscript $_{rock}$) and temperature-dependent (T) parts of the friction coefficient μ and cohesion c, which is demonstrated in a general form below:

$$\tau(T) = \sigma \cdot (\mu_{rock} - \mu(T)) + (c_{rock} - c(T)), \quad with \mu = \tan(\phi)$$

290 The limit of applicability of the temperature-dependent shear criterion is given by the test settings in the laboratory, ranging from normal stresses between 100 and 400 kPa and temperatures between -8 and -0.5°C. The $\pm$ sign reflects the spread of results stemming from the laboratory tests used to construct the shear criterion. In a simplified approach, we assume ice-filled discontinuities throughout the full model domain, while neglecting irregularities such as rock bridges or surface roughness of discontinuities.

295 S2 Transient buildup of hydrostatic pressure

The buildup of hydrostatic pressure within permafrost rock is a central trigger for releasing permafrost rock ~~slides~~ slope failure (Gruber and Haeberli, 2007; Fischer et al., 2010; Pfluger et al., 2025). Although observations and measurements of permafrost hydrogeology in rock slopes are scarce and typically site specific, a field study suggests plausible values for transient water columns in fracture systems of several decameters upon peak snow melt or rain-water infiltration periods (Scandroglio et al., 2025, back calculation from water discharge measured at fracture outlet). In addition, piezometric heads of more than 10 m were recorded in ~~fractured permafrost rock in a one-year measurement interval of 2024 (Offer et al., 2025)~~ boreholes within fractured rock in permafrost (Offer et al., 2025), and

sporadic permafrost (Aspaas et al., 2026). Moreover, failure scarps of larger ~~rock-slides-failures~~ often exhibit wet areas, observed directly after detachment, which point to locally ponded water within the rock mass short before sudden-failure conditions (i.e., Fluchthorn, Austria, 2023 event, Krautblatter et al., 2024; Piz Scerscen, Switzerland, 2024 event, PERMOS, 2024). With this scenario, we test the mechanical response to applied hydrostatic ~~water-pressure-of-pressure~~ equivalent to a 30 m water column~~and-a-~~ repreting hydrogeological conditions upon peak surface water infiltration - compare to reported hydrostatic heads of 27 ± 6 m during average snowmelt and 40 ± 10 m for extreme events (Scandroglio et al., 2025). The pressurized zone is assigned a lateral width of 30 m ~~within-varying-areas-and~~ applied to different regions of the rock mass. Throughout the mechanical cycling, static water pressure is applied within discontinuities only, exerting normal stress on joint walls, while pore pressure within blocks is neglected. Temporal hydrogeological evolution is not explicitly modeled. Instead, spatially variable pressurized zones are used to represent conceptually inferred, locally ponded groundwater conditions within permafrost rock slopes, potentially promoted by channelized flow along fractures (Hasler et al., 2011; Magnin and Josnin, 2021). Unlike the analysis of observed rockfalls (Sect. 2) and permafrost warming (Sect. 3.1), changing groundwater conditions were not directly captured within the scope of this study, but their implications were inferred conceptually. This setup does not aim to reproduce actual groundwater flow conditions but isolates the mechanical effect of water pressure, emphasizing the sensitivity of slope stability to the spatial availability of water within the rock mass.

S3 Rockfalls and ice apron loss

Rockfalls or surface ice loss considerably alter the stress field of the rock slope in sudden moments or within relatively short periods, such as a decade (Deline et al., 2015). These processes have been found to substantially impact the morphology of Platteikogel rock slope before failure (Fig. 3g-i), and thus will be investigated in regard to mechanical response for preparing the ~~rock-slide~~detachment. Therefore, rock slope topography was altered by removing individual blocks, simulating the mechanical response adapting to the new stress field and topography.

Here, we apply the factor of safety (FoS) concept in a modified way. Traditionally, FoS quantifies slope stability by comparing resisting and driving forces (Wyllie and Mah, 2004). However, in our UDEC model with several thousand contact surfaces, this concept is challenging to adapt. As our focus lies on the final detachment phase, we interpret FoS through the evolution of displacement functions instead (cf. Pfluger et al., 2025). The amount of displacement may serve as a proxy to quantify failure. A displacement plateau towards the end of the simulation suggests that the rock slope remains stable approaching mechanical equilibrium (FoS > 1), whereas a continuously increasing displacement indicates progressive failure and the initiation of the ~~rock-slide-release~~detachment (FoS $\approx$ 1 or below).




Figure 6. Simulation strategy for applying the mechanical modeling framework. (a) Model topography and boundary conditions, and two setups A & B varying in structural geometry, as shown in the circled overviews. Joint sets 1 and 2 are included throughout the full model domain. In setup A, the basal shear plane is sketched as a specified shear plane defining the structural failure path. In B, the basal shear plane is missing explicitly, but the model domain was added with a squeezed Voronoi structure, creating many possible failure paths. (b) Overview of the four modeling scenarios and their implementation.

4 Results

4.1 Thermal evolution of the subsurface

From 1980 to 2024, air temperature warming of $\approx 2^{\circ}\text{C}$ is evident at the location of Platteikogel (Fig. SM8a). As permafrost in depths of several decameters evolves over decades, we display the mean annual temperature of mountain permafrost for the year 1980, at the start of the warming trend, and for 2023, shortly before the Platteikogel rock ~~slide detached~~ slope failure (Fig. 7a,b). Throughout this period, warming is evident within the full mountain and especially pronounced on the southeastern flank, while on the steep sections of the northwestern flank, ~~cold~~ low temperatures were better maintained. The temperature asymmetry mainly stems from the radiative warming of the sun-exposed bedrock, which is more pronounced for southeastern aspects, and is a result of the implemented temperature transfer function (Eq. 2). The vanishing ice aprons, which uncovered bedrock down to the elevation of the glacier surface at the southeastern flank between 1970 and 2000, controlled surface exposure and therefore enforced permafrost warming from the southeastern flank towards the northwestern flank. Comparing modeled temperatures along a 20 m deep monitoring profile running parallel to the rock topography, the simulations reveal mean annual temperatures below -3°C , with a minimum of -5.5°C modeled for the northwestern flank in 1980 (Fig. 7c). For 2023, only 15% of the monitoring profile remained below -3°C - marking the lense-shaped cold permafrost body at the northwestern flank - while at the southeastern flank temperatures warmed approximately to -2°C (Fig. 7d). Table 2 displays modeled temperature and its increase at 20 m below the surface for specified locations. Overall, the modeled temperatures along the monitoring profile display a similar shape of temperature progression for 1980 and 2023, reflecting the impact of rising AT throughout the period. However, the ice apron loss on the southeastern flank, between a profile distance of 80 and 180 m, indicates a flip in temperature progression from a local depression (1980) towards ~~an extrusion-like form~~ a local peak (2023).

Table 2. Modeled median annual permafrost temperatures at 20 m below surface picked from Fig. 7b,d for specified locations along the profile distance. Note that the vertical location above 150 m profile distance became ice-free in 1978, and above 300 m in 1987 (see Fig. SM5). ΔT (10m) is given as an additional reference and not explicitly shown. Note that the difference of the 10-year-moving average air temperature between 1980 and 2020 is $+2^{\circ}\text{C}$.

Distance along x-axis [m]	1980 Temperature (20 m)	2023 Temperature (20 m)	ΔT (20 m)	ΔT (10 m)*	Location below
50	-3.2	-2.6	+0.6°C	+1.1°C	SE Glacier
150	-3.6	-1.6	+2°C	+2.7°C	Ice-free, SE flank
250	-3.3	-1.7	+1.6°C	+1.7°C	Ridge top
300	-5.2	-3.8	+1.4°C	+1.7°C	NW Ice apron/ice-free

To estimate the thermal impact of vanishing ice aprons, we compared a model with gradual ice apron retreat to a model assuming a fixed ice apron extent since 1970 onwards (Fig. 8). The observed retreat of ice aprons on the southeastern flank from 1970 to 2000, and its absence since 2000 onwards, indicates a surplus in permafrost temperature of $\approx 1^{\circ}\text{C}$ at 20-meter




Figure 7. Modeled mean annual temperature distribution for (a) 1980 and (c) 2023. The glacier's location and ice apron extent are shown for reference. The plotted temperature distribution shows the median temperatures of all the simulated models (n=100). (b) & (d) Probability density plot of modeled temperatures shown at 20 m below the rock surface, accounting for parameter variations within the given ranges (Eqs. 1 to 4). Note: All models were run from 1900 onwards. The ice apron extent was fixed from 1900 to 1970, while the gradual top-down retreat was simulated from 1970 onwards.

355 depth in 2023. In contrast, retreat of the ice apron on the northwestern flank affected permafrost temperature only marginally, owing to the neglected radiative heating on the northwestern flank in our model.




Figure 8. Impact of ice aprons on the evolution of mountain temperature. Difference in mean annual temperatures between a model assuming retreating ice aprons from 1970 onwards and a model assuming fixed ice apron extent from 1900 to 2023 according to the extent in 1970. Both models were forced since 1900 onwards, with implementing forcings calculated assuming the mean values specified in the defined ranges (Eqs. 1 to 4: (1) $x = 0$, (2) $SO_{NW} = 0^\circ\text{C}$ & $SO_{SE} = 3^\circ\text{C}$, (3) $H_{ice} = 5\text{ m}$, (4) $nF_{glacier} = 0.5$). Note: The SE flank indicates higher temperatures for absent ice aprons (radiative warming implemented with a $+3^\circ\text{C}$ -air temperature offset for snow-free months only - Eq. 2). The NW flank indicates slightly lower temperatures as a result of the attenuated signal of AT (Eq. 3), and the assumed no-offset condition accounting for the absence of radiative warming at the NW flank. The black area in the model marks regions where $|\Delta T| < 0.01^\circ\text{C}$.

4.2 Mechanical investigations on failure processes

Mechanical simulations of setups A and B yielded similar results despite differences in structural detail (explicit basal shear plane vs. implicit shear plane developing along multiple substructures). In the following, we concentrate on setup A, displaying results for scenarios S0 to S3, while presenting results of setup B in the appendix (Figs. A3-~~12~~12). We briefly address the differences between setups A and B at the end of this section.

4.2.1 Model with basal shear plane - explicit failure path - setup A

S0 In order to back-calculate the theoretical pre-failure conditions of critical discontinuities short before the ~~rock-slide~~ release detachment, we ran sensitivity tests of varying shear parameters, which revealed the following situation (Fig. 9a): Of the conducted 21 simulations, ~~7-9~~9 resulted in horizontal displacement of the tracked block of more than 0.01 m (Fig. 9d). While models with $\phi = 40^\circ$ remained in stable conditions, indicated by the displacement functions approaching a plateau at the end of cycling, models with $\phi = 30$ or 20° and cohesion below 0.1 or 0.2 MPa, respectively, display continuously propagating displacement at the end of cycling. The full activation of the basal shear plane is presented in the sequence of Figures 9b-d, where (b) ~~indicates the first activation of shear planes at the rock slope's toe~~ first indicates

370 activation of the central discontinuity set. Assuming a reduction in cohesion, i.e., through progressive weathering or rock fatigue, in (c) ~~the shear path extends to deeper regions, and in~~, the shear plane is activated starting from the rock slope's toe and from there extending towards deeper regions. In (d), the full shear plane is activated, and the displaced blocks do not regain stable positions.

375 S1 Assuming the presence of ice-filled discontinuities and their temperature-dependent shear strength according to the permafrost temperature short before ~~rock-slide-release~~ the detachment (Fig. 10a), the simulation revealed stable slope conditions (Fig. 10b). In scenario S0, failure occurs as a result of progressive destabilization of the rock slope's toe. These kinematics were prevented here, as permafrost temperature is ~~coldest~~ lowest and shear parameters are highest at the steeper parts of the northwestern flank. In contrast, warmer temperatures and lower shear strength dominate the mechanically less relevant southeastern flank. It is worth noting that even a more pessimistic perspective, with 380 higher temperatures simulated (approx. +1°C compared to the 2023 median temperature state shown in Fig. 7c) and incorporating the lower bound of spread from the conversion of temperature-dependent shear parameters (Eq. 6, lower values of shear parameters), results in stable rock slope conditions.

To test the mechanical response to hydrostatic pressure or rockfalls (scenarios S2 and S3), we use the simulated results from the models of scenario S0 (the back-calculated pre-failure state, with a factor of safety (FoS) slightly above 1, i.e., $\phi = 30^\circ$ and 385 $c = 0.1$ MPa, as shown in Fig. 9) and of scenario S1, which represents the temperature-dependent shear model according to 2023 temperatures (Fig. 10b), as the basis.

Base model S0 - Pre-failure conditions $\phi = 30^\circ$ and $c = 0.1$ MPa:

390 S2_{S0} Applied hydrostatic pressure leads to varying magnitudes of shear displacement, mainly depending on where the pressure acts in regard to the monitoring location (Fig. ??11a-c). For the defined hydrostatic pressure, the displacement functions of the models (a-c) do not approach a plateau, indicating unstable slope conditions. In (a,b), the basal shear plane is fully activated. In (c), the basal shear plane is partly activated, only in the areas at the toe of the rock slope. Yet, if that region detaches, the stress is redistributed to other parts at the basal shear plane above, gradually shaping a failure path which evolves towards the peak of the mountain, similar to what is demonstrated in Figure 9b-d with the gradual decrease in 395 cohesion.

S3_{S0} Rockfalls affect the stress field and change the local topography. The removal of a block at the lower slope part results in an increase in x-displacement, as indicated in the displacement function. Yet the form of function appears to approach a plateau, indicating the model is likely to regain equilibrium (Figure ??11d). For the removal of blocks at the upper part (e), the model exhibits an elastic response (see displacement function), suggesting the overall model kinematics were 400 governed by the base model S0, which was used as input. Through local unloading, less stress was exerted on the steeper section of the basal shear plane, stopping the ongoing gradual deformation (compare the displacement function at S0 vs S3).




Figure 9. Rock mechanical back analysis characterizing pre-failure contact surfaces (scenario S0,A) - Results of the UDEC simulation. (a) Compilation of sensitivity tests and the corresponding displacement functions for the location of the black square shown in (b-d). Note that displacements from the initialization of the model were excluded from the graph. (b-d) exhibiting model states after cycling 30,000 ~~mechanical timesteps~~ model steps in the order of a gradual decrease in cohesion. The coloured patches indicate the absolute shear displacement along the contact surfaces of blocks at the end of cycling (dual-coded with the size to display the most prominent areas of shearing within the model). Blue vectors mark the displacement direction and relative magnitude. The black square marks the location of the monitoring point, while the graph on top shows the displacement in the horizontal direction along mechanical cycling time. For intercomparison, (b-d) share the same scale for shear displacement.




Figure 10. Coupling of the temperature-dependent shear criterion to the thermal model state (scenario S1,A). (a) Displaying thermal model state of 2023 as calculated with Cryogrid (Fig. 7c) and corresponding temperature-dependent shear parameters for ice-filled discontinuities acc. to Eq. 6, mean values used. (b) UDEC results of the unidirectional coupled model at the end of cycling. Explanation of the chart as shown in Figure 9.

Base model S1 - Temperature-dependent shear model of 2023 temperatures:

- $S2_{S1}$ Applied hydrostatic pressure leads to a relatively distinct increase in shear displacement (Fig. ??f-h12a-c). However, modeled absolute displacement remains below ~~0.01~~ 0.046 m, and activated shear planes are locally within the rock mass, leaving the entire basal shear plane unaffected. Hydrostatic pressure exerted in the area of the southeastern flank or at a centered position within the mountain (f,g) indicates displacement functions approaching a plateau. In contrast, hydrostatic pressure in proximity of the rock slide's toe (h or northeastern flank (a,c) shows how the uppermost blocks become unstable and detach from the rock slope - simulating a local rockfall release rather than the full rock-slide release. major release. Hydrostatic pressure exerted at a centered position within the mountain (b) shows the modeled highest displacement according to the displacement function. Yet, this is a local effect as the monitor point lies within the area of applied water pressure, while the basal shear zone at its margins remains unaffected.
- $S3_{S1}$ The effect of a rockfall in the area of the slope's toe, i.e., the removal of a block (Fig. ??i12d), shows a direct response by the activation of shearing along contacts of neighboring blocks, but leaves regions more distant unaffected. As shown with In contrast to the base model S0, block removal in the upper slope (Fig. ??j12e) reveals an elastic stress-relief response that counteracts downslope displacement (blue arrows), enhancing local shear activation at shallow discontinuities above the basal shear plane.




Figure 11. The impact of hydrostatic pressure and rockfalls on rock slope stability calculated with UDEC for model setup A ~~-(a-e)~~ using the model of back-calculated pre-failure state from scenario S0 ($\phi = 30^\circ$ and $c = 0.1$ MPa; see Fig. 10c) ~~or (f-j) the temperature-dependent shear criterion S1 (Fig. 10b) as base a basis.~~ (a-c/f-h) Impact of hydrostatic water pressure on slope mechanics, illustrated for three different locations of assumed water pressure. (d,e/f-j) Impact of rockfalls (removal of individual blocks) on slope mechanics. Explanation of charts as shown in Figure 9. Note that the state of the model is displayed for the end of the simulation, and the illustrated shear displacement shows the cumulative displacement of S0 ~~/S1~~ and S2 or S3, respectively.




Figure 12. The impact of hydrostatic pressure and rockfalls on rock slope stability calculated with UDEC for model setup A using the model with the temperature-dependent shear criterion S1 acc. to 2023 (Fig. 10b) as a basis. (a-c) Impact of hydrostatic water pressure on slope mechanics, illustrated for three different locations of assumed water pressure. (d,e) Impact of rockfalls (removal of individual blocks) on slope mechanics. Explanation of charts as shown in Figure 9. Note that the state of the model is displayed for the end of the simulation, and the illustrated shear displacement shows the cumulative displacement of S1 and S2 or S3, respectively.

4.2.2 Model with multiple shear planes - implicit failure path - setup B

S0 Comparing the results for the different model geometries of model of the structural model setup A (defined basal shear plane) and with B (Voronoi structures, without explicitly defined shear plane, but higher degree of structural detail),

we briefly describe the difference. For the back-calculation of the pre-failure state of discontinuities (scenario S0), both setups reveal the same ~~displacement magnitudes (Fig. A3a-e), and the same~~ parameter combinations that exceed the 0.01 m displacement benchmark, ~~except $\phi = 40^\circ$ & $c = 0.025\text{MPa}$, which reaches 0.009 m (Fig. A3d). The rock slide detaching a).~~ Simulated displacement magnitudes resemble both setups (Fig. A3b-d), yet due to the many substructures, ~~shearing occurs along many discontinuities rather than one central structure.~~ The rock slope failure under setup B was formed by multiple parallel shear planes, which are slightly below the pre-defined basal shear plane in setup A. Despite the implicit approach with the Voronoi structures, almost the same failure volume could be replicated (Fig. A3e-d).

S1 The coupled temperature-dependent shear criterion demonstrates marginal shear displacement on the order of 10^{-5}m for 2023 temperature, while setup A reaches the order of 10^{-4}m . Shear displacements in critical regions are not observed, while localized ~~enhanced-marginal~~ shearing areas originate ~~from extra-small blocks within deeper areas~~ (Fig. A4).

S2 Applied hydrostatic pressure demonstrates a distinct impact on overall shear displacement for base model S0 with back-calculated pre-failure shear parameters (Fig. ~~??A5~~a-c), ~~similar to setup A~~. In contrast, the model state according to the temperature-dependent shear criterion of 2023 (S1) was marginally affected, ~~and displacement remained low with recorded displacement functions an order of magnitude lower than for setup A, but in general showing a similar model response~~ (Fig. ~~??f-hA6~~a-c).

S3 Rockfalls affecting the slope topography and the stress ~~field led to the same results~~ redistribution led to enhanced displacement of the failure volume at setup B, for both locations of rockfalls simulated (base model S0, Fig.A5 d,e). In contrast to setup A, where shearing decreases as a result of rockfalls from the ridge area (Fig.11e), the rockfall from the same area indicated the opposite effect for setup B(Fig.??, with enhanced shearing at similar areas. Regarding the base model S1, both setups A and B resulted in similar model responses (cf. Fig.12 & A6 d,e,i, j, ~~as demonstrated for setupA above-).~~

To quantify the mechanical impact of the individual processes, we directly compare scenario results by using the modeled absolute x-displacement recorded at the end of cycling 30000 model steps for the specified monitoring location as the benchmark metric (see displacement functions in the figures). The comparison in Figure 13 demonstrates that hydrostatic pressure (as modeled here) had the most pronounced impact on the kinematic response of the rock slope. Permafrost, as modeled with the implementation of the temperature-dependent shear criterion, had a minor impact on the destabilization process. The impact of rockfalls depends on the structural model setup: While setup A showed minor effects, setup B exhibited x-displacements comparable to those suggested by hydrostatic pressure (cf. S3: a and b in Figure 13). Small block sizes facilitated shearing along many substructures and favoured locally the sliding of neighbouring blocks affected by the void at the post-rockfall state. In general, a smaller block size (setup B) led to lower total x-displacement at the monitoring location, compared to a larger block model with a defined shear basal shear plane (setup A). Within a single setup, however, displacement patterns remain internally consistent and comparable between scenarios. Both model setups indicate that at higher pre-failure shear strength (base model S1), the kinematic rock slope response is less susceptible to hydrostatic pressure applied and rockfalls.




Figure 13. [Compilation of results listed for each scenario shown for the structural model configuration of \(a\) setup A and \(b\) setup B. The bars display the absolute x-displacement at the end of cycling 30,000 model steps for the location of the monitoring point shown in the reference figures. Results stem from individual simulations shown for \(a\) in Figures 9 - 12 and for \(b\) in Figures A3 - A6. The color scale indicates the relative share of the basal shear plane / critical shear planes that is activated at this model state. Note: In UDEC, “model steps” refer to numerical iterations. One model step corresponds to 0.00492 s \(setup A\) and 0.00342 s \(setup B\) of physical time, regardless of the scenarios simulated. Model steps scale linearly with time.](#)

5 Discussion

455 5.1 Does ice apron loss promote the Platteikogel rock ~~slideslope~~ failure?

The Platteikogel rock ~~slide detached from slope~~ failure originated in heavily weathered gneiss with closely spaced foliation and joints in steep terrain, providing favorable preconditions for gravitational mass movements (Fischer et al., 2006). To discuss the mechanism leading to the ~~release of the rock slide~~ detachment, we distinguish between *promoting drivers* which act on a rock slope [system](#) over months to millions of years (Dietze et al., 2017), preparing the rock slope system towards future failure

460 (Prager et al., 2008), and *trigger* referring to an event which directly leads to the ~~detachment-of-a-rock-slide~~main detachment in sec, min, hours, or days after the event - i.e., rainfall or seismic shaking (Leinauer et al., 2024). The patterns shifting a rock slope system towards instability are typically determined by nonlinear key controls in space and time (Krautblatter and Moore, 2014). While we did not identify a clear trigger for the Platteikogel rock ~~slides~~slope failure, we observed processes like ice apron loss over 50 years. The loss of ice aprons and their impact on permafrost, hydrogeology, and rockfall processes
465 (Fig. 5), illustrates how multiple drivers interact during ~~paraglacial transition~~, highlighting its the transition from paraglacial to periglacial conditions, highlighting the nonlinear influence on rock instability.

The destabilizing effect of permafrost degradation on rock slopes has been cross-confirmed by various methods: Laboratory tests simulating permafrost degradation underpin the change of physical rock (Mellor, 1973; Dwivedi et al., 2000; Davies et al., 2001; Krautblatter and Hauck, 2007; Draebing and Krautblatter, 2012) and mixed rock-ice material properties (Mamot et al.,
470 2021; Han et al., 2023; Huang et al., 2024). Field studies demonstrate that observed kinematics and permafrost dynamics are strongly interwoven, impacting small-scale objects such as rock pillars (Weber et al., 2025, volume 100 m³), and full rock slope systems on the slope-scale (Etzelmüller et al., 2022). Dated prehistoric slip surfaces suggest that rock slide formation coincided with permafrost degradation phases, leading Hilger et al. (2021) to conclude that permafrost degradation was likely the primary driver for the failure of the studied post-glacial rock slides. Moreover, empirical evidence confirms enhanced ge-
475 omorphological activity during the paraglacial transition, from small-scale rockfalls (Hartmeyer et al., 2020; Draebing and Mayer, 2021) to large-scale rock ~~slides~~slope failures (Gruber and Haeberli, 2007; Ballantyne et al., 2014; McColl and Draebing, 2019). Whether rock slopes can withstand this transition depends on their pre-failure condition - that is, whether their overall factor of safety was already close to failure.

The concept of progressive failure mechanism and the promotion of the Platteikogel rock slope failure is explained theoretically (Fig. 14a) and on the basis of our simulation results (Fig. 14b): Cold permafrost since the Little Ice Age stabilizes the slope by adding cohesion through ice in fractures, and suppressing crack propagation (Krautblatter and Leith, 2015). However, in recent decades, ice apron loss and warming air temperatures have accelerated permafrost warming. The observed rockfall activity, together with ice apron loss, modified the topography. Consequently, elastic rock mass adaptation leads to widening of joints (Leith et al., 2014; Grämiger et al., 2017). Ice apron loss, joint widening, and heterogeneous permafrost conditions, which are
485 typically found in complex topographic terrain (Noetzli et al., 2007), favour the infiltration of water into the rock slope and therefore potentially enable hydrostatic pressure buildup within stability-relevant (scenario S2). Given sufficient accumulated pre-failure damage (i.e., base model S0), the rock slope is prone to failure upon hydrostatic pressure buildup during peak water infiltration events (snowmelt/rainfall event, S2) or as a response to rockfalls (S3, setup B: small block-size model). Yet, shear strength degradation as a result of permafrost warming (S1) had a minor impact on the destabilization (Fig. 14b). The
490 implications of permafrost warming and strength degradation are discussed in the following section in detail.

5.2 Beyond thermal warming: Superimposed processes accelerate failure

The coupling of the temperature-dependent shear criterion for ice-filled discontinuities to modeled permafrost temperatures of 2023 (Scenario S1, Fig. 10) reveals stable slope conditions for the Platteikogel rock slope, suggesting that the degradation of




Figure 14. (a) Conceptually inferred stability function of the Platteikogel rock slope, highlighting the non-linear influence during the peri-paraglacial transition. (b) Dominant promoting factors investigated in this study that are relevant in the period shortly before failure. Results from the thermal and mechanical simulations are included to illustrate the corresponding impact.

shear strength of ice-filled discontinuities through warming temperatures alone can not explain the release of the observed rock
 495 ~~slide~~ detachment. Therefore, we draw the following conclusions.

1. Other mechanisms superimpose to promote slope failure - i.e., hydrostatic pressure as described by (Gruber and Haeberli, 2007; Cathala et al., 2024; Pfluger et al., 2025) and/or the effects of mass unloading and topographic modification (through glacier ice retreat; Fischer et al., 2010; Leith et al., 2014), which is the case of Platteikogel, explored through rockfalls or ice apron loss. With the mechanical modeling study, we could demonstrate the effect of both process, while
 500 the impact on the destabilization, here measured in terms of shear displacement - was greater for a model which had accumulated more ~~fatigue~~ fatigue and was closer to a state of sudden-failure (=rock ~~slide-release~~ detachment, scenario S0), than for a model suggesting less pre-failure damage (S1, compare Fig. ??11 & 12). The difference in shear parameters suggests that the temperature-dependent shear criterion used in S1 potentially overestimates the actual pre-failure shear strength ($\Delta_{S1-S0}\phi > +14^\circ$, $\Delta_{S1-S0}c > +0.14MPa$). The modeling results indicate that in the case of scenario S0,
 505 the imposed factors may act as a trigger for the ~~rock-slide-release~~ major detachment, while in scenario S1, they may lead to ~~the~~ accumulation of rock mass damage, promoting failure in the future.

2. The discontinuity surfaces are heterogeneous and more complex in nature. Yet, the presence of ice-filled discontinuities according to (Eq. 5; Mamot et al., 2018) was prescribed throughout the full model domain in scenario S1, assuming similar shear charaterisits under same temperature. The applied temperature-dependent shear criterion explains the final stage
 510 of shear failure along ice-filled contact surfaces of rocks, given that rock bridges are eroded, asperities are smoothed, and shear strength is controlled only by the compound material of ice and rock. It accounts for failure timing linked to temperature, where cold delays failure. Given the theoretical pre-failure conditions, as created in the laboratory for tested rock-ice-rock sandwich samples with diameters of 15 cm and artificially smoothed rock surfaces, warming above

–1 °C would lead to the rock ~~slide-release~~slope failure, assuming a model with uniform temperatures. Compared to the simulated heterogeneous temperatures, with average permafrost temperatures of ≈ -3 °C at stability-relevant areas at the rock slopes' toe (Fig. 7d), this corresponds to a temperature discrepancy of more than 2°C. In any case, extrapolating the laboratory constraint conditions of ice-filled contact surfaces to a natural shear plane spanning 100 m in length is a strong simplification, neglecting other shear plane features and irregularities. For intact rock, serving as a proxy for rock bridges, a model explicitly capturing temperature-dependent shear strength degradation in the range from -5 to 0 °C has not yet been established. Nevertheless, existing studies clearly indicate that rock mechanical properties are sensitive to warming at these temperatures, demonstrating a general warming-related weakening (see Section 5.1).”

Although our thermo-mechanical modelling approach involves several uncertainties (see Supplementary Discussion – Sect. Scope of interpretation and limitations to modeling), it provides a first-order assessment of the central processes that promote permafrost rock slope failures (i.e., compare to Gruber and Haeberli, 2007; Krautblatter et al., 2013; Grämiger et al., 2020)(cf. Gruber and

The mechanical model indicates that the failure is strongly controlled by the stability of the toe of the slope. Simulated rockfalls (scenario S3) yield in additional displacement of the failed rock mass above - especially for the model with small block sizes (setup B) and a shear parametrization close to failure (Fig. A5d,e). These findings suggest that the Platteikogel rock slope failure could have occurred as a rock collapse (sequential smaller detachments). However, morphological observations suggest that detachment happened as one major, single push event. The form of deposits (bifurcation - see Fig. 2a,e), the reach angle of 24.6°, and H/L ratio of 0.45 suggest that the failure can kinematically be classified as a rock avalanche, likely resulting from a single push event (cf. kinematic model in Figure SM6). The runout length can be explained by the low contact friction and lubrication effect (basal water film) resulting from the basal contact to the glacier surface and present snow cover. The failed volume of 50,000 m³ is at the lower end of cubatures reported for rock avalanches (cf. rock collapse vs. rock avalanche in Hermanns et al., 2020). To conclude, the Platteikogel rock slope failure likely occurred as one major failure event, which was preceded by rockfalls observed in the decade before (Fig. 3) and by detachments shortly after. Post-failure activity was indicated by proximal deposits that are generally larger than distal deposits (Detailed map in Fig. SM7), suggesting secondary failure events.

5.3 Ice apron loss likely enhances thermal and hydrogeological changes

Ice aprons represent a glacial heritage, often preserving ice several thousand years old (Guillet et al., 2021). Their existence indicates the presence of permafrost underneath (Benn and Evans, 2014). The observed area decline in the Western Alps in the last decades (Guillet and Ravel, 2020; Kaushik et al., 2022; Ravel et al., 2023) was also evident at the Vernagtferner Basin in general (compare time series of historic orthophotos; Table A1, ii), which is located in the Eastern Alps of Europe, and in detail in direct proximity to the Platteikogel detachment area (Fig. 3a-c).

The observed ice apron loss at Vernagtferner, including the Platteikogel site, is primarily attributed to continuously rising air temperatures since 1980 (Fig. SM8a). Winter precipitation has remained largely stable since 1970, with no significant trend (winter mass balance of Vernagtferner used as proxy; Fig. SM8b). In contrast, summer precipitation has increased since 1980,

with the trend intensifying after 2010 (10-year moving average: 500 mm in 1980, 650 mm in 2010, and 800 mm in 2020; Fig. SM8c). However, it remains unclear and largely site-specific if ice aprons respond to a general increase in precipitation by growth or thickness loss. Yet, for the case of Platteikogel, the pronounced area loss of ice apron at the southern flank occurred
550 before 2000, suggesting a strong control by rising air temperatures.

In our thermal model, ice aprons were represented as static bodies with temporally varying extents, forced by linear retreat rates inferred from orthophotos. This approach did not capture their dynamic interaction with atmospheric variables. Current knowledge on ice aprons remains limited, with scarce temperature data for the ice or underlying rock (Ravanel et al., 2023), and no information on heat fluxes between the rock, ice, and atmosphere. To approximate the thermal characteristics of ice aprons
555 in our thermal model, we applied a low-pass filter on air temperatures mimicking buffered seasonal signals with ice apron thickness (Eq. 3). This reproduced the general pattern observed in temperature profiles of glacier ice with depth (Jacquemart et al., 2025) and incorporated the elimination of clear seasonal signals beyond 10 m depth.

Our model highlights the thermal contrast between previously ice-covered and now ice-free rock surfaces, which have since been exposed to radiative warming (Fig. 8). The results clearly demonstrate the impact on surfaces receiving high amounts of
560 solar radiation, whereas for northwest-facing rock slopes, the effect is less pronounced and cannot be reliably assessed with given information and no constraints on rock–ice–atmosphere heat fluxes.

With these abstractions of ice aprons, we capture only part of the cryospheric dynamics. Rather than reproducing full system complexity, we demonstrate how ice apron loss can influence permafrost degradation over multiple decades through conductive processes and simplified atmospheric coupling, while neglecting precipitation and snow dynamics. The conceptual model of ice
565 aprons (Fig. 5), illustrating interactions with the hydro- (Francese et al., 2025; Pfluger et al., 2025) and geosphere (Hartmeyer et al., 2020; Fey et al., 2025) concomitant with atmospheric warming, builds on empirical relationships and inferred processes from studies conducted on glaciers. Due to similarities and the scarce knowledge about ice aprons, analogies were drawn between glaciers and ice aprons. For the Platteikogel rock [slideslope failure](#), the modeled ice aprons did not thermally affect the mechanically critical shear planes. However, the ongoing rockfall activity and hydrogeological changes are likely associated
570 with the retreat of ice aprons.

6 Conclusion

The 2024 Platteikogel rock ~~slide~~-slope failure (Vernagtferner Basin, Austria) provides a benchmark case demonstrating how ice apron loss and permafrost warming promote the failure of high-alpine rock slopes. Multi-decadal ice apron area loss and pre-failure rockfall activity preceded the studied ~~rock-slide~~major rock detachment. Based on field observations and conceptual reasoning, we developed a numerical modeling study drawing the following conclusions:

- **Climate warming is the main driver for rising permafrost temperatures between 1980 and 2023.** Conductive thermal simulations indicate decadal warming at 20 m depth before rock ~~slide~~-detachment: $+0.14^{\circ}\text{C decade}^{-1}$ below the cold-based glacier in the area of the bergschrund, $+0.37^{\circ}\text{C decade}^{-1}$ below the ridge top at the location of the basal shear plane, $+0.47^{\circ}\text{C decade}^{-1}$ at the southeast-, and $+0.33^{\circ}\text{C decade}^{-1}$ at the northwest-exposed, ice-apron free slopes (median values of 100 twin simulations).
- **Ice apron loss accelerates permafrost warming through atmospheric exposure of rock surfaces, as shown by conductive thermal modeling.** Forced with a monthly air temperature offset for solar radiative warming of $+3^{\circ}\text{C}$ for southeast-exposed slopes during snow-free months, the model suggests that the observed loss of the ice aprons on the southeastern flank between 1970 and 2000 caused approximately 1°C of additional permafrost warming at 20 m depth by 2023, relative to simulations where the ice aprons remained intact from 1970 to 2023. With radiative warming as the dominant factor in warming rock upon becoming ice-free, the effect is less pronounced for aspects with little incoming radiation.
- **Modeled mean 2023 temperatures along the basal shear plane (-4 to -2°C) indicate stable slope conditions, as supported by the mechanical model coupled to the thermal simulation.** Herby, we linked the shear strength, as the dominant parameter of rock slide control, to a temperature-dependent shear model, valid for ice-filled discontinuities. The finding suggests that other mechanisms apart from permafrost warming acted in driving the observed rock ~~slide-release~~detachment, the slope did not fail along ice-filled discontinuities, or the application of laboratory-derived temperature-dependent shear parameters does not match real-world shear surface conditions.
- **Degrading ice aprons feature geomorphic change by enhancing frost-weathering activity through atmospheric recoupling and by enabling water infiltration into rock slopes, which in turn facilitates the generation of rock slope failures.** Besides atmospheric recoupling, hydrostatic pressure buildup, and stress ~~Stress~~ alterations from ice apron loss and rockfalls, ~~drive progressive as well as hydrostatic pressure buildup, drive progressive rock slope~~ failure, as demonstrated by the mechanical simulation.
- **Given the major ice apron loss and structural predisposition at the Platteikogel, similar rock slope failures may be anticipated if sufficient-observations-kinematic monitoring data are available.** However, precise failure time predictions without kinematic data remain unrealistic.

Code availability. The scripts used to reproduce the UDEC modeling study are made available via the following repository:
<https://zenodo.org/records/19028752>.

Data availability. The input data and results of the CryoGrid 2D modeling studies were made available via the following repository:
605 <https://zenodo.org/records/19028752>. A video of the modeled temperature evolution can be found therein.
Spatial data used for pre- and post-failure characterisation are referenced in Table A1 and accessible via the provided sources or upon request. Meteorological and hydrological time series used to characterize pre- and post-failure conditions are referenced in Table A2 and can be downloaded from the original sources.






Figure A1. Structural analysis of two rock outcrops in proximity to the ~~rock-slide's~~ detachment area. In total, 11339 planes were reconstructed and analyzed using the FACET plugin in Cloud Compare software (Dewez et al., 2016). Parameters for FACET plugin: Fusion algorithm *Kd-tree*, *max-anlge* 20, *max relative distance* 1.00. (a) Stereographic projection of the geological planes using CLAR-method for illustration and 'Kamb exponential smoothing': Units are in numbers of standard deviations by which the density estimate differs from uniform (produced with mplstereonet Kington, 2024)). The two identified clusters are joint planes. (b) Orthophotographic view of the ridge exhibiting the detachment area of the rock ~~slideslope failure~~, including the location of the two rock outcrops and the whitish highlighted structural features of morphology (iii) that intersect the ridge: Steep dipping, perpendicular oriented foliation planes. The UAV point cloud model recorded in August 2024 was used as the basis for analysis in Cloud Compare (2 cm model resolution).

(a) Model calibration of subsurface properties


- (i) Volumetric fraction of rock and water acc. to Myhra et al. (2017,2019) for gneissic rock slopes
- (ii) Use of Matterhorn borehole temperatures (0.1 - 52.5 m, Permos (2025)) to fit thermal parameters in a pseudo 1D model (simplified column mesh and horizontal model topography)
- Forcing with measured borehole temperatures at 0.1 m depths (daily mean)
Best-fit of thermal conductivity k , and volumetric heat capacity of c_v chosen for all further simulations:
- $$k = 2 \text{ W K}^{-1} \text{ m}^{-1}, c_v = 3 \cdot 10^6 \text{ J m}^{-3} \text{ K}^{-1}$$
- (iii) Single lithology at Platteikogel assuming homogenous subsurface properties



(b) Setup and geometry



(c) Forcing applied to rock topography



(d) Ensemble simulation

- (i) **Model initialization**
Run until reaching thermal equilibrium with mean annual GST of 1900
- (ii) **Random sampling** of parameters in defined ranges (c) and calculation of GST along rock topography
- (iii) **Ensemble simulations: n = 100**
Starting 1900 until 2023
- Ice apron retreat modeled since 1970 onwards, from 1900 to 1969 fixed extension to the observed elevation in 1970

Figure A2. Simulation strategy applying the permafrost model Cryogrid 2D: (a) Calibration of thermal parameters with measured borehole temperature showing best-fit simulation results (a1-a4). (b) Cross-section through the ridge of Platteikogel, demonstrating varying surface types and the observed downslope retreat of ice apron since 1970 onwards - Given dates/elevation marks are inferred from historic orthophotographs. The Topography (upper boundary) and lower boundary (implied, at 6000 m depth) mark the frame for the meshed model. Note: The vertical scale of the cross-section is exaggerated by a factor of 2. (c) Processing of atmospheric forcing using temperature transfer functions to derive RST on the basis of AT, considering varying surface types. (d) Workflow for conducting the ensemble simulations.




Figure A3. Rock mechanical back analysis characterizing pre-failure joint surfaces (scenario S0,B) - Results of the UDEC simulation. (a-c) exhibiting model after cycling 30,000 mechanical timesteps model steps in the order of a gradual decrease in cohesion. The coloured patches indicate the absolute shear displacement along the contact surfaces of blocks at the end of cycling (dual-coded with the size to display the most prominent areas of shearing within the model). Blue vectors mark the displacement direction and relative magnitude. The black square marks the location of the monitoring point, while the graph on top shows the displacement in the horizontal direction along mechanical cycling time. (d) Compilation of sensitivity tests and the corresponding displacement functions for the location of the black square. Note that displacements from the initialization of the model were excluded from the graph.




Figure A4. Coupling of the temperature-dependent shear criterion to the thermal model state (scenario S1,B). (a) Displaying thermal model state of 2023 as calculated with Cryogrid (Fig. 7c) and corresponding temperature-dependent shear parameters for ice-filled discontinuities acc. to Eq. 6, mean values used. (b) UDEC results of the unidirectional coupled model at the end of cycling. Explanation of chart as shown in Figure 9. Note that displacement vectors illustrate the settlement of the overall mountain, resulting of marginal shear along steep south dipping joint sets. Displacement vectors are here exxagerated by a factor 10^5 to make them visible.




Figure A5. The impact of hydrostatic pressure and rockfalls on rock slope stability calculated with UDEC for model setup B using the model of back-calculated pre-failure state from scenario S0 ($\phi = 30^\circ$ and $c = 0.1$ MPa; see Fig. A4c) as a basis. (a-c) Impact of hydrostatic water pressure on slope mechanics, illustrated for three different locations of assumed water pressure. (d,e) Impact of rockfalls (removal of individual blocks) on slope mechanics. Explanation of charts as shown in Figure A3. Note that the state of the model is displayed for the end of the simulation, and the illustrated shear displacement shows the cumulative displacement of S0 and S2 or S3, respectively. (b,c) Small surficial blocks appear to fly freely due to numerical instability caused by hydrostatic pressure applied beneath the surface.




Figure A6. The impact of hydrostatic pressure and rockfalls on rock slope stability calculated with UDEC for model setup B ~~-(a-e)~~ using the model ~~of back-calculated pre-failure state from scenario S0 ($\phi = 30^\circ$ and $c = 0.1$ MPa; see Fig. 10c) or (f-j)~~ with the temperature-dependent shear criterion S1 ~~acc. to 2023~~ (Fig. ~~10A4b~~) as ~~base~~ basis. (a-c/~~f-h~~) Impact of hydrostatic water pressure on slope mechanics, illustrated for three different ~~location-locations~~ of assumed water pressure. (d,e/~~f-j~~) Impact of rockfalls (removal of individual blocks) on slope mechanics. Explanation of charts as shown in Figure 9A3. Note that the state of the model is displayed for the end of the simulation, and the illustrated shear displacement shows the ~~cummulative~~ cumulative displacement of ~~S0/S1~~ and S2 or S3, respectively.

Table A1. Spatial data used to for pre- and post-failure characterisation.

Index	Spatial data	Year	File type/reso- lution	Source	Usage
ii	DSM and orthophoto of Platteikogel post-failure state	2024	0.2 m	RGB images recorded by UAV-campaign on 24 July 2024, this study; DSM and orthophoto processed using Agisoft Metashape professional (https://www.agisoft.com/)	Overview photos, processed DEM for (a) structural analysis of rock surfaces: mapping of joints, (b) rock-slide <u>Failure</u> volume detection
ii	Orthophotos	1969, 1999, 2003, 2009, 2013, 2019	0.2 - 0.25 m	Provided by Land Tirol - https://www.tirol.gv.at/data/ (last access: 7 January 2025)	Visual ice apron change
iii	Large-format aerial imagery	2015, 2018, 2021, 2023	0.2 m	Recorded by 3D RealityMaps GmbH, DSM processed acc. to Barbosa et al. (2024), this study	Surface ice elevation change, rockfall inventory, pre-failure topography for 2023 cross-section

Table A2. Meteorological and hydrological time series used to characterize pre- and post-failure conditions.

Index	Time series	Parameters	Observation period	Measurement interval	Location	Source	Usage
i	Pitztal glacier meteorological station	AT	1994 - 2024	hourly	< 5 km distance to site, at 2863.9 m a.s.l.	GeoSphere Austria (2024), station id: 17315; 46°55'37.0"N 10°52'45.0"E	Calculation of monthly AT lapse rate
ii	Brunnenkogel meteorological stations	AT	2003 - 2024	hourly	< 5 km distance to site, at 3437 m a.s.l.	GeoSphere Austria (2024), station id: 173200; 46°54'46.0"N 10°51'42.0"E	Calculation of monthly AT lapse rate
iii	Obergurgl-Vent historical air temperature record	AT	1851 - 2024	monthly homogenized data	< 14 km distance to site, 1938 m a.s.l	Auer et al. (2007), HISTALP station mode, dataset accessible via https://www.zamg.ac.at/histalp/ (last access: 21 January 2025), station name: OBG, Austria; 46°52'01.0"N 11°01'28.0"E	Long-term AT as basis for CryoGrid 2D model forcing
iv	Matterhorn borehole temperatures (MAT 0205)	T between 0.1 and 40 m depth	2019 - 2024	daily	Swizz, 3343.96 m a.s.l.	PERMOS data portal (https://www.permos.ch/de/data-portal , last access 11.02.2025); 45°58'55.3"N 7°40'33.8"E	Parameter calibration of the CryoGrid 2D model
v	Matterhorn air temperature (MH30) and rock surface temperature (MH27: south-, MH30: north-exposed face)**	AT, RST at 0.1 m depth	2010 - 2023	hourly	Swizz, approx. 3500 m a.s.l.	Weber et al. (2024); 45°58'48.0"N 7°40'12.0"E	RST conversion accounting for solar incoming radiation (Eq. 2)

*We hereby refer to the detachment area of the Platteikogel rock ~~slide~~slope failure. Abbreviations: T.. temperature, AT.. air temperature, P.. air pressure, RR..precipitation sum, SWR.. short wave radiation, T_{water} .. water temperature, RST.. rock surface temperature. The given coordinates are based on the WGS84 datum. **MH27: aspect = 90°, slope = 70°; MH30: aspect = 340°, slope = 90°.

Table A3. Characterization of the rock mass at Platteikogel by using the Geological Strength Index - parametrization (1). Intact rock properties (2) are tested in the laboratory. The material properties representing the rock mass (3) are derived from (1 & 2) to be subsequently assigned to the linear elastic blocks within the UDEC model. Specification for deriving parameters: 'estimated' values are derived from categorical relations or from given graphs. 'calculated' values are calculated according to the suggested formula.

Parameter	Abbrev.	Value	Unit	Source
(1) Geological Strength Index	GSI	25	-	estimated according to Hoek and Brown (2019, Fig. 7)
Material constant intact rock	m_i	20	-	estimated acc. to (Marinos and Hoek, 2000, Table S1)
Disturbance factor	D	0	-	acc. to Hoek and Brown (2019)
Material constants rock mass (m_b, s, a)	m_b	1.37322	-	calculated acc. to Hoek et al. (2002, eq. 3)
	s	0.00024	-	calc. acc. to Hoek et al. (2002, eq. 4)
	a	0.53127	-	calc. acc. to Hoek et al. (2002, eq. 5)
(2) Uniaxial Compressiv Strength	UCS	101	MPa	UCS test**, unfrozen, $\bar{x}, n = 5$
Young's modulus intact rock	E_i	39	GPa	UCS test**, unfrozen, $\bar{x}, n = 5$
Poisson ratio intact rock	ν_i	0.25	-	UCS test**, unfrozen, $\bar{x}, n = 5$
(3) Young' modulus rock mass	E_{rm}	2.383	GPa	calc. acc. to Hoek et al. (2002, eq. 11a)
Poisson ratio rock mass	ν_{rm}	0.4	-	estimated acc. to Vászrhelyi (2009, eq. 13)
Compressive modulus rock mass	K_{rm}	3.972*	GPa	calc. acc. to $K_{rm} = \frac{E_{rm}}{3(1 - 2\nu_{rm})}$
Shear modulus rock mass	G_{rm}	0.851*	GPa	calc. acc. to $G_{rm} = \frac{E_{rm}}{2(1 + \nu_{rm})}$
Tensile strength of rock mass	σ_t	0.017	MPa	calc. acc. to Hoek et al. (2002, eq. 7)
Friction angle of rock mass	ϕ_{rm}	40	°	estimated after Cai et al. (2004, Fig. 5)
Cohesion of rock mass	c_t	0.8	MPa	estimated after Cai et al. (2004, Fig. 5)

*These values were rounded to $K_{rm} = 4 \text{ GPa}$ and $G_{rm} = 1 \text{ GPa}$ and assigned to the blocks for all UDEC simulations.

** UCS tests conducted according to recommendations of Mutschler (2004) under constant strain for gneissic rock samples collected in regard to [studying](#) the Bliggspitze rock [slide-study-slope failure](#) in Kaunertal, Austria (Pfluger et al., 2025), which are of similar lithology to the rocks at the Platteikogel rock [slideslope failure](#).

Author contributions. FP initiated the idea, created the concept of both modeling studies, conducted the mechanical and thermal modeling studies, and wrote the manuscript. SW revised the manuscript and directed the focus of the storyline and the thermal modeling. NB conducted the change detection/rockfall inventory and the pre-processing of the data. FH conducted the meteorological and hydrological analysis. JL conducted the seismic analysis and revised the manuscript. PW revised the CryoGrid 2D code, conducted data preprocessing, and calibration analysis. MK assisted with discussions.

Competing interests. At least one of the (co-)authors is a member of the editorial board of Earth Surface Dynamics.

Acknowledgements. The authors thank Christoph Mayer for sharing his observation on the Platteikogel rock ~~slide~~slope failure. We thank Justyna Czekirda and Bernd Etzelmüller for sharing the CryoGrid 2D code (Czekirda et al., 2023), and express special thanks to Sebastian Westermann, who was open to helping us with issues when implementing the model. Furthermore, we thank 3D RealityMaps GmbH for the production of Orthophotos and DSM from high-resolution aerial imagery in the frame of the AlpsenseRely project (Teilprojekt for LMU TUSO1UFS-77318), used for the detection of rockfall activity before the ~~rock slide~~major failure and glacier mapping since 2010. FP expresses his gratitude to Maximilian Reinhard for his invaluable assistance in conducting the UAV survey in 2024 and for his companionship during fieldwork. FP expresses further gratitude to the individuals who contributed to data sharing or processing, fieldwork assistance, or their open dialogue: Christine Fey, Robert Kenner, Manuel Saigger, Alex Fröhlich, Theresa Hayeck, Matthias Siebers, and Christian Sommer. AI - statement: The authors used Grammarly and ChatGPT exclusively to assist with grammar and language editing. No content or interpretations were generated by AI. This research has been supported by the Bayerisches Staatsministerium für Wissenschaft und Kunst (grant no. M3OCCA).

References

- Aspaas, A., Bievre, G., Lacroix, P., Langet, N., Aga, J., Skrede, I., Kristensen, L., Etzelmüller, B., and Renard, F.: Seasonal and Inter-Annual Evolution of the Deformation of Two Arctic Landslides, *EGU*sphere, pp. 1–53, <https://doi.org/10.5194/egusphere-2026-62>, 2026.
- Auer, I., Böhm, R., Jurkovic, A., Lipa, W., Orlik, A., Potzmann, R., Schöner, W., Ungersböck, M., Matulla, C., Briffa, K., Jones, P., Efthymi-
630 adis, D., Brunetti, M., Nanni, T., Maugeri, M., Mercalli, L., Mestre, O., Moisselin, J.-M., Begert, M., Müller-Westermeier, G., Kveton, V.,
Bochnicek, O., Stastny, P., Lapin, M., Szalai, S., Szentimrey, T., Cegnar, T., Dolinar, M., Gajic-Capka, M., Zaninovic, K., Majstorovic,
Z., and Nieplova, E.: HISTALP—Historical Instrumental Climatological Surface Time Series of the Greater Alpine Region, *International
Journal of Climatology*, 27, 17–46, <https://doi.org/10.1002/joc.1377>, 2007.
- Ballantyne, C. K., Sandeman, G. F., Stone, J. O., and Wilson, P.: Rock-Slope Failure Following Late Pleistocene Deglaciation on Tectonically
635 Stable Mountainous Terrain, *Quaternary Science Reviews*, 86, 144–157, <https://doi.org/10.1016/j.quascirev.2013.12.021>, 2014.
- Barbosa, N., Leinauer, J., Jubanski, J., Dietze, M., Münzer, U., Siegert, F., and Krautblatter, M.: Massive Sediment Pulses Triggered by a
Multi-Stage 1303 Alpine Cliff Fall (Hochvogel, DE–AT), *Earth Surface Dynamics*, 12, 249–269, [https://doi.org/10.5194/esurf-12-249-
2024](https://doi.org/10.5194/esurf-12-249-2024), 2024.
- Benn, D. and Evans, D. J. A.: *Glaciers and Glaciation*, 2nd Edition, Routledge, London, 2 edn., ISBN 978-0-203-78501-0,
640 <https://doi.org/10.4324/9780203785010>, 2014.
- Cai, M., Kaiser, P. K., Uno, H., Tasaka, Y., and Minami, M.: Estimation of Rock Mass Deformation Modulus and Strength of
Jointed Hard Rock Masses Using the GSI System, *International Journal of Rock Mechanics and Mining Sciences*, 41, 3–19,
[https://doi.org/10.1016/S1365-1609\(03\)00025-X](https://doi.org/10.1016/S1365-1609(03)00025-X), 2004.
- Carslaw, H. S. and Jaeger, J. C.: *Conduction of Heat in Solids*, Clarendon Pr, Oxford, 2. ed., repr edn., ISBN 978-0-19-853303-0, 1980.
- Cathala, M., Bock, J., Abdulsamad, F., Deline, P., Josnin, J.-Y., Ravel, L., Revil, A., Richard, J., Verroust, F., and Magnin, F.: Assessing
645 the Role of Permafrost in the Preconditioning and Triggering Factors of the September 2020 Crête Des Granges Rockfall (Southern
French Alps), *Géomorphologie : relief, processus, environnement*, 30, <https://doi.org/10.4000/12yqn>, 2024.
- Czekirda, J., Etzelmüller, B., Westermann, S., Isaksen, K., and Magnin, F.: Post-Little Ice Age Rock Wall Permafrost Evolution in Norway,
The Cryosphere, 17, 2725–2754, <https://doi.org/10.5194/tc-17-2725-2023>, 2023.
- Dabrowski, M., Krotkiewski, M., and Schmid, D. W.: MILAMIN: MATLAB-based Finite Element Method Solver for Large Problems,
650 *Geochemistry, Geophysics, Geosystems*, 9, <https://doi.org/10.1029/2007GC001719>, 2008.
- Davies, M. C., Hamza, O., and Harris, C.: The Effect of Rise in Mean Annual Temperature on the Stability of Rock Slopes Containing
Ice-Filled Discontinuities, *Permafrost and periglacial processes*, 12, 137–144, 2001.
- Deline, P., Gruber, S., Delaloye, R., Fischer, L., Geertsema, M., Giardino, M., Hasler, A., Kirkbride, M., Krautblatter, M., Magnin, F., McColl,
655 S., Ravel, L., and Schoeneich, P.: Chapter 15 - Ice Loss and Slope Stability in High-Mountain Regions, in: *Snow and Ice-Related
Hazards, Risks, and Disasters*, edited by Shroder, J. F., Haeberli, W., and Whiteman, C., *Hazards and Disasters Series*, pp. 521–561,
Academic Press, Boston, ISBN 978-0-12-394849-6, <https://doi.org/10.1016/B978-0-12-394849-6.00015-9>, 2015.
- Dewez, T. J. B., Girardeau-Montaut, D., Allanic, C., and Rohmer, J.: FACETS: A CloudCompare Plugin to Extract Geological Planes from
Unstructured 3D Point Clouds, *Int. Arch. Photogramm. Remote Sens. Spatial Inf. Sci.*, XLI-B5, 799–804, [https://doi.org/10.5194/isprs-
archives-XLI-B5-799-2016](https://doi.org/10.5194/isprs-

660 archives-XLI-B5-799-2016), 2016.
- Dietze, M., Turowski, J. M., Cook, K. L., and Hovius, N.: Spatiotemporal Patterns, Triggers and Anatomies of Seismically Detected Rock-
falls, *Earth Surface Dynamics*, 5, 757–779, <https://doi.org/10.5194/esurf-5-757-2017>, 2017.

- Draebing, D. and Krautblatter, M.: P-Wave Velocity Changes in Freezing Hard Low-Porosity Rocks: A Laboratory-Based Time-Average Model, *The Cryosphere*, 6, 1163–1174, 2012.
- 665 Draebing, D. and Mayer, T.: Topographic and Geologic Controls on Frost Cracking in Alpine Rockwalls, *Journal of Geophysical Research: Earth Surface*, 126, e2021JF006163, 2021.
- Dwivedi, R. D., Soni, A. K., Goel, R. K., and Dube, A. K.: Fracture Toughness of Rocks under Sub-Zero Temperature Conditions, *International Journal of Rock Mechanics and Mining Sciences*, 37, 1267–1275, [https://doi.org/10.1016/S1365-1609\(00\)00051-4](https://doi.org/10.1016/S1365-1609(00)00051-4), 2000.
- Eberhardt, E., Stead, D., and Coggan, J.S.: Numerical Analysis of Initiation and Progressive Failure in Natural Rock Slopes—the 1991 Randa
670 Rockslide, *International Journal of Rock Mechanics and Mining Sciences*, 41, 69–87, 2004.
- Etzelmüller, B., Czekirka, J., Magnin, F., Duvillard, P.-A., Ravel, L., Malet, E., Aspaas, A., Kristensen, L., Skrede, I., Majala, G. D., Jacobs, B., Leinauer, J., Hauck, C., Hilbich, C., Böhme, M., Hermanns, R., Eriksen, H. Ø., Lauknes, T. R., Krautblatter, M., and Westermann, S.: Permafrost in Monitored Unstable Rock Slopes in Norway – New Insights from Temperature and Surface Velocity Measurements, *Geophysical Surveying, and Ground Temperature Modelling, Earth Surface Dynamics*, 10, 97–129, [https://doi.org/10.5194/esurf-10-97-](https://doi.org/10.5194/esurf-10-97-2022)
675 2022, 2022.
- European Space Agency: Sentinel-2 L2A image, tile ID T33TWN, acquired on 10 May 2024, European Space Agency, 2024.
- Fey, C., Wichmann, V., and Zangerl, C.: Influence of Permafrost Degradation and Glacier Retreat on Recent High Mountain Rockfall Distribution in the Eastern European Alps, *Earth Surface Processes and Landforms*, 50, e70063, <https://doi.org/10.1002/esp.70063>, 2025.
- Fischer, L., Käb, A., Huggel, C., and Noetzli, J.: Geology, Glacier Retreat and Permafrost Degradation as Controlling Factors of Slope
680 Instabilities in a High-Mountain Rock Wall: The Monte Rosa East Face, *Natural Hazards and Earth System Sciences*, 6, 761–772, <https://doi.org/10.5194/nhess-6-761-2006>, 2006.
- Fischer, L., Amann, F., Moore, J. R., and Huggel, C.: Assessment of Periglacial Slope Stability for the 1988 Tschierwa Rock Avalanche (Piz Morteratsch, Switzerland), *Engineering Geology*, 116, 32–43, <https://doi.org/10.1016/j.enggeo.2010.07.005>, 2010.
- Francesse, R. G., Valentino, R., Haeblerli, W., Bondesan, A., Giorgi, M., Picotti, S., Pettenati, F., Sandron, D., Ramponi, G., and Valt, M.:
685 Failure of Marmolada Glacier (Dolomites, Italy) in 2022: Data-Based Back Analysis of Possible Collapse Mechanisms, *Natural Hazards and Earth System Sciences*, 25, 3027–3053, <https://doi.org/10.5194/nhess-25-3027-2025>, 2025.
- GeoSphere Austria: Bundesanstalt Für Geologie, Geophysik, Klimatologie Und Meteorologie - Data Hub, <https://data.hub.geosphere.at/>, 2024.
- Gerstner, R., Fey, C., Kuschel, E., Valentin, G., Voit, K., and Zangerl, C.: Polyphase Rock Slope Failure Controlled by Pre-Existing Geological Structures and Rock Bridges, *Bulletin of Engineering Geology and the Environment*, 82, 363, [https://doi.org/10.1007/s10064-023-](https://doi.org/10.1007/s10064-023-03382-2)
690 03382-2, 2023.
- Gischig, V., Amann, F., Moore, J. R., Loew, S., Eisenbeiss, H., and Stempfhuber, W.: Composite Rock Slope Kinematics at the Current Randa Instability, Switzerland, Based on Remote Sensing and Numerical Modeling, *Engineering Geology*, 118, 37–53, <https://doi.org/10.1016/j.enggeo.2010.11.006>, 2011.
- 695 Gislås, K., Etzelmüller, B., Farbrot, H., Schuler, T. V., and Westermann, S.: CryoGRID 1.0: Permafrost Distribution in Norway Estimated by a Spatial Numerical Model, *Permafrost and Periglacial Processes*, 24, 2–19, <https://doi.org/10.1002/ppp.1765>, 2013.
- Grämiger, L. M., Moore, J. R., Gischig, V. S., Ivy-Ochs, S., and Loew, S.: Beyond Debuitressing: Mechanics of Paraglacial Rock Slope Damage during Repeat Glacial Cycles, *Journal of Geophysical Research: Earth Surface*, 122, 1004–1036, 2017.

- Grämiger, L. M., Moore, J. R., Gischig, V. S., Loew, S., Funk, M., and Limpach, P.: Hydromechanical Rock Slope Damage During Late Pleistocene and Holocene Glacial Cycles in an Alpine Valley, *Journal of Geophysical Research: Earth Surface*, 125, <https://doi.org/10.1029/2019JF005494>, 2020.
- Gruber, S. and Haeberli, W.: Permafrost in Steep Bedrock Slopes and Its Temperature-Related Destabilization Following Climate Change, *Journal of Geophysical Research: Earth Surface*, 112, https://www.zora.uzh.ch/id/eprint/3936/2/Gruber_Haeberli_Permafrost_Steep_2007.pdf, 2007.
- Guillet, G. and Ravanel, L.: Variations in Surface Area of Six Ice Aprons in the Mont-Blanc Massif since the Little Ice Age, *Journal of Glaciology*, 66, 777–789, <https://doi.org/10.1017/jog.2020.46>, 2020.
- Guillet, G., Preunkert, S., Ravanel, L., Montagnat, M., and Friedrich, R.: Investigation of a Cold-Based Ice Apron on a High-Mountain Permafrost Rock Wall Using Ice Texture Analysis and Micro-14C Dating: A Case Study of the Triangle Du Tacul Ice Apron (Mont Blanc Massif, France), *Journal of Glaciology*, 67, 1205–1212, <https://doi.org/10.1017/jog.2021.65>, 2021.
- Haberkorn, A., Hoelzle, M., Phillips, M., and Kenner, R.: Snow as a Driving Factor of Rock Surface Temperatures in Steep Rough Rock Walls, *Cold Regions Science and Technology*, 118, 64–75, 2015.
- Han, Y., Du, L., and Shen, S.: Study on Shear Test and Shear Displacement of Frozen Joints with Different Opening Degrees, *Natural Hazards*, 115, 289–307, <https://doi.org/10.1007/s11069-022-05555-w>, 2023.
- Hartmeyer, I., Delleske, R., Keuschnig, M., Krautblatter, M., Lang, A., Schrott, L., and Otto, J.-C.: Current Glacier Recession Causes Significant Rockfall Increase: The Immediate Paraglacial Response of Deglaciating Cirque Walls, *Earth Surface Dynamics*, 8, 729–751, 2020.
- Hasler, A., Gruber, S., Font, M., and Dubois, A.: Advective Heat Transport in Frozen Rock Clefts: Conceptual Model, Laboratory Experiments and Numerical Simulation, *Permafrost and Periglacial Processes*, 22, 378–389, 2011.
- Hermanns, R. L., Penna, I. M., Oppikofer, T., Noël, F., and Velardi, G.: Rock Avalanche, in: *Treatise on Geomorphology*, pp. 85–105, Elsevier, ISBN 978-0-12-818235-2, <https://doi.org/10.1016/B978-0-12-818234-5.00183-8>, 2022.
- Hilger, P., Hermanns, R. L., Czokir, J., Myhra, K. S., Gosse, J. C., and Etzelmueller, B.: Permafrost as a First Order Control on Long-Term Rock-Slope Deformation in (Sub-)Arctic Norway, *Quaternary Science Reviews*, 251, 106718, <https://doi.org/10.1016/j.quascirev.2020.106718>, 2021.
- Hoek, E. and Brown, E. T.: The Hoek–Brown Failure Criterion and GSI – 2018 Edition, *Journal of Rock Mechanics and Geotechnical Engineering*, 11, 445–463, <https://doi.org/10.1016/j.jrmge.2018.08.001>, 2019.
- Hoek, E., Carranza-Torres, C., Corkum, B., et al.: Hoek-Brown Failure Criterion-2002 Edition, *Proceedings of NARMS-Tac*, 1, 267–273, 2002.
- Huang, S., Cai, H., Xin, Z., and Liu, G.: The Temperature-Dependent Shear Strength of Ice-Filled Joints in Rock Mass Considering the Effect of Joint Roughness, Opening and Shear Rates, *The Cryosphere*, 17, 1205–1223, <https://doi.org/10.5194/tc-17-1205-2023>, 2023.
- Huang, S., Liu, G., Cheng, A., Zheng, L., Liu, F., Tian, J., and Musonda, K.: Experimental Study on the Influence of Water Saturation and Temperature Condition on the Basic Friction Angle, <https://doi.org/10.21203/rs.3.rs-4439361/v1>, 2024.
- Hungr, O., Leroueil, S., and Picarelli, L.: The Varnes Classification of Landslide Types, an Update, *Landslides*, 11, 167–194, <https://doi.org/10.1007/s10346-013-0436-y>, 2014.
- Intergovernmental Panel on Climate Change: The Ocean and Cryosphere in a Changing Climate: Special Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, 1 edn., ISBN 978-1-009-15796-4 978-1-009-15797-1, <https://doi.org/10.1017/9781009157964>, 2022.

- Islam, N., Carrivick, J. L., Coulthard, T., Westoby, M., Dunning, S., and Gindraux, S.: A Growing Threat of Multi-Hazard Cascades Highlighted by the Birch Glacier Collapse and Blatten Landslide in the Swiss Alps, *Geology Today*, 41, 200–205, <https://doi.org/10.1111/gto.12526>, 2025.
- 740 Itasca Consulting Group: UDEC – Universal Distinct Element Code, User’s Manual, Minneapolis, United States, 2019.
- Jacquemart, M., Welty, E., Gastaldello, M., and Carcanade, G.: Glenglat: A Database of Global Englacial Temperatures, *Earth System Science Data*, 17, 1627–1666, <https://doi.org/10.5194/essd-17-1627-2025>, 2025.
- James, D. W.: The Thermal Diffusivity of Ice and Water between -40 and + 60° C, *Journal of Materials Science*, 3, 540–543, <https://doi.org/10.1007/BF00549738>, 1968.
- 745 Ji, S.-H., Koh, Y.-K., Kuhlman, K. L., Lee, M. Y., and Choi, J. W.: Influence of Pressure Change During Hydraulic Tests on Fracture Aperture, *Groundwater*, 51, 298–304, <https://doi.org/10.1111/j.1745-6584.2012.00968.x>, 2013.
- Jia, H., Xiang, W., and Krautblatter, M.: Quantifying Rock Fatigue and Decreasing Compressive and Tensile Strength after Repeated Freeze–Thaw Cycles, *Permafrost and Periglacial Processes*, 26, 368–377, 2015.
- Kaushik, S., Ravelle, L., Magnin, F., Yan, Y., Trouve, E., and Cusicanqui, D.: Effects of Topographic and Meteorological Parameters on the
- 750 Surface Area Loss of Ice Aprons in the Mont Blanc Massif (European Alps), *The Cryosphere*, 16, 4251–4271, <https://doi.org/10.5194/tc-16-4251-2022>, 2022.
- Kington: Mplstereonet: A Python Package for Creating Stereonets in Matplotlib, <https://pypi.org/project/mplstereonet/>, 2024.
- Krautblatter, M. and Hauck, C.: Electrical Resistivity Tomography Monitoring of Permafrost in Solid Rock Walls, *Journal of Geophysical Research: Earth Surface*, 112, <https://doi.org/10.1029/2006JF000546>, 2007.
- 755 Krautblatter, M. and Leith, K.: Glacier- and Permafrost-Related Slope Instabilities, in: *The High-Mountain Cryosphere*, edited by Huggel, C., pp. 147–165, Cambridge University Press, 2015.
- Krautblatter, M. and Moore, J. R.: Rock Slope Instability and Erosion: Toward Improved Process Understanding, *Earth Surface Processes and Landforms*, 39, 1273–1278, <https://doi.org/10.1002/esp.3578>, 2014.
- Krautblatter, M. and Moser, M.: A Nonlinear Model Coupling Rockfall and Rainfall Intensity Based on a Four Year Measurement in a High
- 760 Alpine Rock Wall (Reintal, German Alps), *Natural Hazards and Earth System Sciences*, 9, 1425–1432, <https://doi.org/10.5194/nhess-9-1425-2009>, 2009.
- Krautblatter, M., Funk, D., and Günzel, F. K.: Why Permafrost Rocks Become Unstable: A Rock-Ice-Mechanical Model in Time and Space, *Earth Surface Processes and Landforms*, 38, 876–887, <https://doi.org/10.1002/esp.3374>, 2013.
- Krautblatter, M., Weber, S., Dietze, M., Keuschnig, M., Stockinger, G., Brückner, L., Beutel, J., Figl, T., Trepmann, C., Hofmann, R., Rau,
- 765 M., Pfluger, F., Barbosa Mejia, L., and Siegert, F.: The 2023 Fluchthorn Massive Permafrost Rock Slope Failure Analysed, *EGU General Assembly 2024*, Vienna, Austria, 14–19 Apr 2024, <https://doi.org/10.5194/egusphere-egu24-20989>, 2024.
- Kreuss, O., Krenmayr, H. G., Pavlik, W., Bayer, I., and Hirschhofer, J.: GEOFAST 1:50 000 Geological Map of 173 - Sölden, <https://www.geologie.ac.at/>, 2012.
- Kuhn, D., Hermanns, R. L., Fuchs, M., Schüßler, N., Torizin, J., Aga, J., Bendle, J., Eiken, T., and Balzer, D.: Warming-Induced Destabi-
- 770 lization of Polar Coastal Rock Cliffs and the Role of Thermokarst: A Case Study of Forkastningsfjellet on Svalbard, *Science of The Total Environment*, 968, 178 807, <https://doi.org/10.1016/j.scitotenv.2025.178807>, 2025.
- Leinauer, J., Dietze, M., Knapp, S., Scandroglio, R., Jokel, M., and Krautblatter, M.: How Water, Temperature, and Seismicity Control the Preconditioning of Massive Rock Slope Failure (Hochvogel), *Earth Surface Dynamics*, 12, 1027–1048, <https://doi.org/10.5194/esurf-12-1027-2024>, 2024.

- 775 Leith, K., Moore, J. R., Amann, F., and Loew, S.: Subglacial Extensional Fracture Development and Implications for Alpine Valley Evolution, *Journal of Geophysical Research: Earth Surface*, 119, 62–81, 2014.
- Leroueil, S., Locat, J., Vaunat, J., Picarelli, L., Lee, HOMA., and Faure, R. M.: Geotechnical Characterization of Slope Movements, in: *Proceedings of the Seventh International Symposium on Landslides*, (Ed. K. Senneset) Trondheim, Norway, Balkema, Rotterdam., pp. 53–74, 1996.
- 780 Magnin, F. and Josnin, J.-Y.: Water Flows in Rock Wall Permafrost: A Numerical Approach Coupling Hydrological and Thermal Processes, *Journal of Geophysical Research: Earth Surface*, 126, e2021JF006394, <https://doi.org/10.1029/2021JF006394>, 2021.
- Magnin, F., Etzelmüller, B., Westermann, S., Isaksen, K., Hilger, P., and Hermanns, R. L.: Permafrost Distribution in Steep Rock Slopes in Norway: Measurements, Statistical Modelling and Implications for Geomorphological Processes, *Earth Surface Dynamics*, 7, 1019–1040, <https://doi.org/10.5194/esurf-7-1019-2019>, 2019.
- 785 Mamot, P., Weber, S., Schröder, T., and Krautblatter, M.: A Temperature- and Stress-Controlled Failure Criterion for Ice-Filled Permafrost Rock Joints, *The Cryosphere*, 12, 3333–3353, <https://doi.org/10.5194/tc-12-3333-2018>, 2018.
- Mamot, P., Weber, S., Lanz, M., and Krautblatter, M.: Brief Communication: The Influence of Mica-Rich Rocks on the Shear Strength of Ice-Filled Discontinuities, *The Cryosphere*, 14, 1849–1855, <https://doi.org/10.5194/tc-14-1849-2020>, 2020.
- Mamot, P., Weber, S., Eppinger, S., and Krautblatter, M.: A Temperature-Dependent Mechanical Model to Assess the Stability of Degrading
- 790 Permafrost Rock Slopes, *Earth Surface Dynamics*, 9, 1125–1151, 2021.
- Marinos, P. and Hoek, E., eds.: *GSI: A Geologically Friendly Tool for Rock Mass Strength Estimation*, OnePetro, 2000.
- McColl, S. T. and Draebing, D.: Rock Slope Instability in the Proglacial Zone: State of the Art, *Geomorphology of Proglacial Systems: Landform and Sediment Dynamics in Recently Deglaciated Alpine Landscapes*, pp. 119–141, 2019.
- Mellor, M.: Mechanical Properties of Rocks at Low Temperatures, in: *2nd International Conference on Permafrost.*, pp. 334–344, 1973.
- 795 Mutschler, T.: Neufassung Der Empfehlung Nr. 1 Des Arbeitskreises “Versuchstechnik Fels” Der Deutschen Gesellschaft Für Geotechnik e. V.: Einaxiale Druckversuche an Zylindrischen Gesteinsprüfkörpern, *Bautechnik*, 81, 825–834, <https://doi.org/10.1002/bate.200490194>, 2004.
- Myhra, K. S., Westermann, S., and Etzelmüller, B.: Modelled Distribution and Temporal Evolution of Permafrost in Steep Rock Walls Along a Latitudinal Transect in Norway by CryoGrid 2D, *Permafrost and Periglacial Processes*, 28, 172–182, <https://doi.org/10.1002/ppp.1884>,
- 800 2017.
- Myhra, K. S., Westermann, S., and Etzelmüller, B.: Modeling Conductive Heat Flow Between Steep Rock Walls and Talus Slopes – Thermal Processes and Geomorphological Implications, *Frontiers in Earth Science*, 7, <https://doi.org/10.3389/feart.2019.00192>, 2019.
- Noetzli, J., Gruber, S., Kohl, T., Salzmann, N., and Haeberli, W.: Three-Dimensional Distribution and Evolution of Permafrost Temperatures in Idealized High-Mountain Topography, *Journal of Geophysical Research: Earth Surface*, 112, 2007.
- 805 Offer, M., Weber, S., Krautblatter, M., Hartmeyer, I., and Keuschnig, M.: Pressurised Water Flow in Fractured Permafrost Rocks Revealed by Borehole Temperature, Electrical Resistivity Tomography, and Piezometric Pressure, *The Cryosphere*, 19, 485–506, <https://doi.org/10.5194/tc-19-485-2025>, 2025.
- Otto, J.-C., Schrott, L., and Keller, F.: Map of Permafrost Distribution for Austria, Europe, <https://doi.org/10.1594/PANGAEA.917719>, 2020.
- PERMOS: Rock Falls Data Portal, <https://www.permos.ch/data-portal/rock-falls>, 2024.
- 810 Pfluger, F., Weber, S., Steinhauser, J., Zangerl, C., Fey, C., Fürst, J., and Krautblatter, M.: Massive Permafrost Rock Slide under a Warming Polythermal Glacier Deciphered through Mechanical Modeling (Bliggspitze, Austria), *Earth Surface Dynamics*, 13, 41–70, <https://doi.org/10.5194/esurf-13-41-2025>, 2025.

- Phillips, M., Wolter, A., Lüthi, R., Amann, F., Kenner, R., and Bühler, Y.: Rock Slope Failure in a Recently Deglaciated Permafrost Rock Wall at Piz Kesch (Eastern Swiss Alps), February 2014, *Earth Surface Processes and Landforms*, 42, 426–438, <https://doi.org/10.1002/esp.3992>, 2017.
- Pierhöfer, L., Bartelt, P., Bühler, Y., Hafner, E., Kenner, R., Walter, F., and Phillips, M.: Bergsturz Vom 14. April 2024 Am Piz Scerscen, Graubünden, <https://doi.org/10.55419/wsl:38392>, 2024.
- Prager, C., Zangerl, C., Patzelt, G., and Brandner, R.: Age Distribution of Fossil Landslides in the Tyrol (Austria) and Its Surrounding Areas, *Natural Hazards and Earth System Sciences*, 8, 377–407, <https://doi.org/10.5194/nhess-8-377-2008>, 2008.
- Ravel, L., Guillet, G., Kaushik, S., Preunkert, S., Malet, E., Magnin, F., Trouvé, E., Montagnat, M., Yan, Y., and Deline, P.: Ice Aprons on Steep High-Alpine Slopes: Insights from the Mont-Blanc Massif, Western Alps, *Journal of Glaciology*, 69, 1275–1291, <https://doi.org/10.1017/jog.2023.15>, 2023.
- Rechberger, C. and Zangerl, C.: Rock Mass Characterisation and Distinct Element Modelling of a Deep-Seated Rock Slide Influenced by Glacier Retreat, *Engineering Geology*, 300, 106584, 2022.
- Scandroglio, R., Weber, S., Rehm, T., and Krautblatter, M.: Decadal in Situ Hydrological Observations and Empirical Modeling of Pressure Head in a High-Alpine, Fractured Calcareous Rock Slope, *Earth Surface Dynamics*, 13, 295–314, <https://doi.org/10.5194/esurf-13-295-2025>, 2025.
- Shewchuk, J. R.: Triangle: Engineering a 2D Quality Mesh Generator and Delaunay Triangulator, in: *Applied Computational Geometry: Towards Geometric Engineering*, edited by Lin, M. C. and Manocha, D., vol. 1148 of *Lecture Notes in Computer Science*, pp. 203–222, Springer-Verlag, 1996.
- Smith, M. W. and Riseborough, D. W.: Climate and the Limits of Permafrost: A Zonal Analysis, *Permafrost and Periglacial Processes*, 13, 1–15, <https://doi.org/10.1002/ppp.410>, 2002.
- Vásárhelyi, B.: A Possible Method for Estimating the Poisson's Rate Values of the Rock Masses, *Acta Geodætica et Geophysica Hungarica*, 44, 313–322, <https://doi.org/10.1556/AGeod.44.2009.3.4>, 2009.
- Walden, J., Jacquemart, M., Higman, B., Hugonnet, R., Manconi, A., and Farinotti, D.: Landslide Activation during Deglaciation in a Fjord-Dominated Landscape: Observations from Southern Alaska (1984–2022), *Natural Hazards and Earth System Sciences*, 25, 2045–2073, <https://doi.org/10.5194/nhess-25-2045-2025>, 2025.
- Weber, S., Beutel, J., and Cicoira, A.: In-Situ Measurements in Steep Bedrock Permafrost in an Alpine Environment on the Matterhorn Hörnligrat, Zermatt Switzerland: 2008-2023, <https://doi.org/10.1594/PANGAEA.967586>, 2024.
- Weber, S., Beutel, J., Dietze, M., Bast, A., Kenner, R., Phillips, M., Leinauer, J., Mühlbauer, S., Pfluger, F., and Krautblatter, M.: Progressive Destabilization of a Freestanding Rock Pillar in Permafrost on the Matterhorn (Swiss Alps): Field Observations, Laboratory Experiments and Mechanical Modeling, *EGU sphere*, 2025, 1–31, <https://doi.org/10.5194/egusphere-2025-1151>, 2025.
- Witherspoon, P. A., Wang, J. S. Y., Iwai, K., and Gale, J. E.: Validity of Cubic Law for Fluid Flow in a Deformable Rock Fracture, *Water Resources Research*, 16, 1016–1024, <https://doi.org/10.1029/WR016i006p01016>, 1980.
- World Glacier Monitoring Service: Fluctuations of Glaciers (FoG) Database, <https://doi.org/10.5904/WGMS-FOG-2025-02B>, 2025.
- Wyllie, D. C. and Mah, C.: *Rock slope engineering*, CRC Press, 2004.
- Zangerl, C., Fey, C., and Prager, C.: Deformation Characteristics and Multi-Slab Formation of a Deep-Seated Rock Slide in a High Alpine Environment (Bliggspitze, Austria), *Bulletin of Engineering Geology and the Environment*, 78, 6111–6130, <https://doi.org/10.1007/s10064-019-01516-z>, 2019.