

Measurement Report: Unraveling PM₁₀ Sources and Oxidative Potential Across Chinese Regions Based on CNN-LSTM Data Preprocessing and Receptor Model

Qinghe Cai¹, Dongqing Fang^{2,3}, Junli Jin³, Xiaoyu Hu⁴, Yuxuan Cao¹, Tianyi Zhao^{1,5}, Yang Bai¹, Yang Zhang^{1,6}

¹College of Resources and Environment, University of Chinese Academy of Sciences, Beijing, 100049, China

²Key Laboratory of Urban Meteorology, China Meteorological Administration, Beijing 100089, China

³Meteorological Observation Center, China Meteorological Administration, Beijing, 100081, China

⁴City University of Hong Kong, Hongkong, 999077, China

⁵Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou, 510640, China

⁶Beijing Yanshan Earth Critical Zone National Research Station, Beijing, 101408, China

Correspondence to: Dongqing Fang (fangdongqing@163.com) and Yang Zhang (zhangyang@ucas.ac.cn)

Abstract. The oxidative potential (OP) of particulate matter is a key driver of PM₁₀-induced adverse health effects, triggering oxidative stress and inflammatory responses that increase respiratory and cardiovascular disease risks. To evaluate PM₁₀ and its OP characteristics across China, samples were collected from twelve representative monitoring stations from June 2022 to May 2023. A deep learning model combining Convolutional Neural Networks and Long Short-Term Memory networks (CNN-LSTM) was employed to reconstruct anomalous PM₁₀ data, achieving R² values of 0.8840 for test sets. Significant spatial variations in PM₁₀ were observed, with highest concentrations in the northwestern regions (Xi'an: $98.20 \pm 52.92 \mu\text{g}\cdot\text{m}^{-3}$, Dunhuang: $90.36 \pm 54.72 \mu\text{g}\cdot\text{m}^{-3}$), the lowest in the northeast (Longfengshan: $40.04 \pm 24.04 \mu\text{g}\cdot\text{m}^{-3}$, Dalian: $40.35 \pm 15.66 \mu\text{g}\cdot\text{m}^{-3}$), and elevated levels in suburban areas (average: $85.43 \pm 46.69 \mu\text{g}\cdot\text{m}^{-3}$). Urban sites showed the highest OP values ($0.61 \pm 0.21 \text{ nmol H}_2\text{O}_2\cdot\text{m}^{-3}$), with significantly higher PM₁₀ concentrations in northern regions compared to southern ones ($p < 0.05$). Most sites exhibited peak PM₁₀ and OP levels in winter and lowest in summer. Source apportionment using Positive Matrix Factorization (PMF) revealed dust (13.2-27.4%), biomass burning (16.2-39.3%), traffic (16.6-21.4%), and agricultural activities (13-22%) as main contributors to PM₁₀. Mass-normalized OP (OP_m) analysis revealed traffic sources showing consistently high values ($0.013\text{-}0.022 \text{ nmol H}_2\text{O}_2\cdot\mu\text{g}^{-1}$). These findings highlight the need to control traffic emissions and other major sources to reduce OP and protect public health.

1 Introduction

Particulate matter (PM) is one of the main pollutants affecting air quality and human health. Among these, PM₁₀, which refers to suspended particles with an aerodynamic diameter of 10 μm or less, has received considerable attention due to its complex sources, extensive environmental and health effects. The sources of PM₁₀ are both complex and diverse, including anthropogenic activities such as fossil fuel combustion, industrial production, traffic emissions and dust, as well as natural sources such as dust storms and volcanic eruptions (Xue et al., 2010). PM₁₀ can remain suspended in the atmosphere for

extended periods of time, significantly affecting atmospheric visibility while potentially exerting profound effects on regional and global climate change through both direct and indirect mechanism (Slanina and Zhang, 2004). More critically, PM₁₀ poses a serious threat to human health. Upon entering the human body via the respiratory system, it can be deposited in the airways and lungs, triggering respiratory diseases such as asthma, chronic obstructive pulmonary disease (COPD) and even lung cancer (Cao et al., 2016). Furthermore, PM₁₀ can penetrate the alveolar barrier and enter the circulatory system, inducing systemic diseases such as cardiovascular disease and diabetes (Huang, 2023).

In the context of accelerating global industrialization and urbanization, PM₁₀ pollution has emerged as a critical environmental concern. Research conducted by the World Health Organization (WHO) indicates that air pollution is responsible for millions of premature deaths worldwide each year, with PM₁₀ being a major contributor (Cohen et al., 2005). The mechanisms by which PM₁₀ affects human health are diverse and complex, one of the primary mechanisms being its ability to induce excessive production of reactive oxygen species (ROS), subsequently triggering oxidative stress (OS) effects. Components within PM₁₀, such as transition metals and polycyclic aromatic hydrocarbons (PAHs), can directly or indirectly promote ROS generation, leading to cell membrane lipid peroxidation, protein denaturation, and DNA damage (Chirino et al., 2010). Furthermore, ROS can activate inflammatory signaling pathways, including nuclear factor κ B (NF- κ B), which amplify inflammatory responses and further leading to cellular dysfunction and tissue damage (Wang et al., 2017). This interplay between oxidative stress and inflammatory responses is considered a critical pathophysiological basis for various PM₁₀-induced diseases. Several studies suggest that oxidative potential (OP) may be a more accurate indicator of PM health effects than its mass concentration, providing a new perspective for assessing PM health risks (Gao et al., 2020; Bates et al., 2019).

The OP of PM serves as a critical indicator for assessing its toxicity and is closely related to the generation of ROS. Research indicates that the OP of PM is strongly correlated with its physicochemical properties and sources (He and Zhang, 2023). In particular, PM of smaller size typically exhibits higher OP, possibly due to its larger specific surface area and enhanced bioavailability (Saffari et al., 2014; Yao et al., 2024). Water-soluble transition metals (e.g., iron and copper) and organic carbon (e.g., PAHs) in PM are considered to be the primary chemical components that influence OP. These components can induce ROS generation either by catalyzing Fenton reactions or by directly participating in redox processes (Saffari et al., 2014; Guo et al., 2020). Sources of OP in PM are varied and include primarily traffic emissions, fossil fuel combustion, and secondary organic aerosol formation (Bates et al., 2019; Saffari et al., 2014). Significantly, photochemical aging of PM in the atmosphere further alters its OP, possibly related to the formation of secondary organic aerosols, changes in oxidation states of metallic components during the aging process, and the oxidation degree of reactive organic compounds (An et al., 2022; Ma et al., 2025). In addition, the oxygen content in the fuel has been shown to be a critical factor affecting OP, as exemplified by the typically high OP of PM generated from biomass combustion (Hedayat et al., 2016).

However, an accurate assessment of the health risks associated with PM₁₀ requires an accurate analysis of its sources and chemical compositions. High-quality, complete datasets are essential for reliable source apportionment and subsequent risk assessment. Environmental monitoring data often contain missing values and anomalies due to instrument malfunction, maintenance periods, or extreme weather conditions, which can significantly affect the accuracy of subsequent analyses. In

recent years, with the rapid development of deep learning technology, its application in handling environmental data quality issues has received increasing attention. Deep learning models, particularly the combination of Convolutional Neural Networks (CNN) and Long Short-Term Memory networks (LSTM), have demonstrated significant advantages in identifying and
70 correcting anomalies and filling missing values in time series environmental data. CNNs effectively extract spatial features, while LSTMs excel at capturing long-term dependencies in time series (Huang and Kuo, 2018; Li et al., 2020). This hybrid model not only identifies anomalies, but also improves data completeness and reliability by predicting and replacing anomalous or missing values (Lee et al., 2019; Qin et al., 2019). Compared with traditional machine learning methods, CNN-LSTM models show superior performance in several evaluation metrics, such as Mean Absolute Error (MAE), Root Mean Square
75 Error (RMSE) (Huang and Kuo, 2018; Yang et al., 2020a; Li et al., 2020). CNN-LSTM models retain significant value in processing atmospheric particulate matter data for data quality improvement. Their spatial feature extraction capabilities effectively identify and correct anomalies caused by instrument malfunction or local pollution events, thereby improving data quality (Zhang and Zhou, 2023). Through training and learning, CNN-LSTM models can effectively predict and fill missing data, providing a high-quality data foundation for subsequent source apportionment and risk assessment analyses (Li et al.,
80 2020; Yang et al., 2020a).

After data pre-processing, the Positive Matrix Factorization (PMF) model was used to analyse PM_{10} sources in this study. The PMF model can identify major pollution sources and their contribution rates by decomposing the observation data matrix without requiring prior information (Paatero and Tapper, 1994). In recent years, PMF models have been extensively applied in PM_{10} and $PM_{2.5}$ source apportionment, often in combination with other techniques such as multiple linear regression (MLR)
85 (Weber et al., 2018). Based on the source contribution results from PMF analysis, MLR models can further quantify the contributions of different sources to the OP of PM, providing crucial evidence to reveal the association between PM sources and their health effects. Recent studies have innovatively introduced machine learning methods, such as multilayer perceptron (MLP), to model OP based on source contribution results from PMF analysis, significantly improving model predictive accuracy and explanatory power (Borlaza et al., 2022).

90 In this study, we adopted a comprehensive approach to process PM_{10} data and evaluate its OP. First, we removed anomalies from PM_{10} data and used a deep learning model combining CNN and LSTM to predict and replace anomalous values. This method effectively captures spatial and temporal features in time-series data, thereby improving data completeness and prediction accuracy. Then, we employed the PMF model for PM_{10} source apportionment to identify its major sources. Finally, based on the PMF results, we calculated the OP per unit mass of PM_{10} (OP_m) to investigate the intrinsic toxicity of different
95 emission sources. Through this series of methods, this study aims to reveal the OP characteristics and sources of PM_{10} in typical regions of China.

2 Materials and Methods

2.1 Sample Collections

Daily ambient PM₁₀ samples were collected every three days from June 2022 to May 2023 at the twelve stations of the CMA
100 Atmosphere Watch Network (CAWNET), with their distribution shown in **Figure 1** and detailed information provided in
Table 1. Remote sites were selected in areas far from anthropogenic pollution sources to ensure the representativeness of the
background monitoring data. Rural sites were selected in typical areas, with sampling points located away from local pollution
sources and elevated above the surrounding ground to minimize local disturbances. At urban sites, sampling points were
typically located 50-100 m above the average urban elevation in order to collect mixed aerosol samples rather than aerosols
105 from single sources. Suburban sites were located in transition zones between urban and rural areas to reflect aerosol
characteristics under different environmental conditions. All aerosol samples were collected using MiniVol™ air samplers
(Airmetrics, Oregon, USA) operating continuously for 24 hours from 9:00 AM to 9:00 AM the following day (Beijing time)
at a flow rate of 5 L·min⁻¹. Whatman 47 mm quartz fiber filters (QM/A) were used for sampling. To prevent contamination
from affecting the experimental results, all filters were heated at 800°C for 3 hours prior to use to remove potential organic
110 contaminants.




Figure 1. Locations of 12 CAWNET stations. The map base is from the Ministry of Natural Resources' Standard Map Service, review number GS (2019)1822.

Table 1. Information for twelve CAWNET stations.

Station name	Province	Lat.&Long.	Elev.(m)	Type
Changde (CHD)	Hunan	29° 10.2' N, 111° 42.6' E	150.6	Rural
Chengdu (CD)	Sichuan	30° 39 'N, 104° 2.4' E	587.0	Urban
Dalian (DL)	Liaoning	38° 54' N, 121° 37.8' E	91.5	Urban
Dunhuang (DH)	Gansu	40° 9' N, 94° 40.8' E	1137.5	Suburban
Gucheng (GC)	Hebei	39° 7.8' N, 115° 48' E	15.2	Rural
Jinsha (JS)	Hubei	29° 37.8' N, 114° 12' E	751.4	Remote
Lhasa (LS)	Tibet	29° 40.2' N, 91° 7.8' E	3660.0	Urban
Lin'An (LA)	Zhejiang	30° 18' N, 119° 44' E	138.6	Remote
Longfengshan (LFS)	Heilongjiang	44° 43.8' N, 127° 36' E	331.0	Remote
Nanning (NN)	Guangxi	22° 49.2' N, 108° 21' E	159.0	Urban
Xi'An (XA)	Shaanxi	34° 25.8' N, 108° 58.2' E	363.0	Urban
Zhengzhou (ZZ)	Henan	34° 46.8' N, 113° 40.8' E	110.4	Suburban

115 **2.2 Chemical and OP analysis**

2.2.1 Chemical compositions analysis

Quantitative measurements of OC and EC were performed using the DRI Model 2015A thermal/optical carbon analyzer developed by the Desert Research Institute, USA. After OC and EC analysis, ion chromatography (Dionex 600 series, USA) was used to analyze and determine various ions, including Na⁺, NH₄⁺, K⁺, Ca²⁺, Mg²⁺, F⁻, Cl⁻, NO₃⁻, and SO₄²⁻. This method
120 has been widely used as a highly efficient and sensitive analytical technique for the determination of water-soluble ions in PM₁₀ and PM_{2.5} (Domingos et al., 2012; Cui et al., 2008; Yan et al., 2006).

2.2.2 OP analysis

The 2',7'-Dichlorodihydrofluorescein (DCFH) method is widely used for detecting particle-bound ROS, mainly due to its lack of specificity and selectivity for various ROS species (Antonini et al., 1998; Cohn et al., 2008; Huang et al., 2016). In this
125 study, the 2',7'-Dichlorodihydrofluorescein diacetate (DCFH-DA) probe method was employed to measure ROS levels induced by PM₁₀. First, DCFH-DA (97%, Sigma-Aldrich, USA) was prepared as a 1 mmol·mL⁻¹ stock solution using anhydrous ethanol and mixed with 0.01 mol·L⁻¹ NaOH solution in a 1:4 (v/v) ratio. The mixture was kept at room temperature in the dark for 30 min to ensure complete alkaline hydrolysis of DCFH-DA to DCFH. Phosphate buffer solution (PBS, 0.0067 mol·L⁻¹, pH 7.2) was then added to adjust the pH to 7.0-7.4. The hydrolyzed DCFH solution was stored at 4°C in the dark and used within 2
130 hours. Horseradish peroxidase (HRP) was dissolved in phosphate buffer to prepare a 10 unit·mL⁻¹ HRP stock solution. It was mixed with the DCFH solution prior to use to achieve final concentrations of 10 μmol·L⁻¹ DCFH and 0.5 units·mL⁻¹ HRP in

the reaction system. To generate a standard curve, a $1000 \mu\text{g}\cdot\text{mL}^{-1} \text{H}_2\text{O}_2$ solution was diluted with ultrapure water to generate H_2O_2 standard solutions at concentrations of 20, 40, 80, 160, 200, 240, 320, 400, and $800 \text{ nmol}\cdot\text{L}^{-1}$. In a 96-well plate, $20 \mu\text{L}$ standard solution and $60 \mu\text{L}$ DCFH-HRP mixture were added, with three replicates for each concentration. After 15 minutes of dark incubation at 37°C , fluorescence intensity was measured using a multifunctional microplate reader (SynergyTMH1, BioTek America) at an excitation wavelength of 485 nm and an emission wavelength of 535 nm. Quantification was done through converting the sample's fluorescent intensity to the equivalent quantity of H_2O_2 ($\text{nmol H}_2\text{O}_2\cdot\text{m}^{-3}$). Before sample analysis, PM_{10} samples were extracted in phosphate buffer solution by sonication for 30 minutes. After centrifugation, the supernatant was collected for testing. In the 96-well plate, $20 \mu\text{L}$ of sample solution and $60 \mu\text{L}$ of DCFH-HRP mixture were added, with three replicates per sample. Water blanks, filter blanks, and DCFH-HRP background blanks were included to eliminate background interference. After 15 minutes of dark incubation at 37°C , fluorescence intensity was measured and converted to H_2O_2 equivalent concentrations using the H_2O_2 standard curve to characterize PM_{10} -induced ROS levels. Throughout the experimental procedure, the microplate reader was preheated for 30 minutes before measurement to ensure a stable incubation temperature of 37°C . Background fluorescence values were subtracted from each measurement, and the relative standard deviation (RSD) of the replicates was controlled within 5% to ensure the accuracy and reliability of the data.

2.3 Data analysis

2.3.1 CNN

One-dimensional convolutional neural networks (1D-CNN) have significant theoretical advantages and practical value in processing time series data. The core mechanism relies on local connectivity and weight sharing, where each neuron is connected only to a local region of the input data, while the convolution kernel weights are shared across the entire input sequence. This design significantly reduces the number of model parameters, improving computational efficiency while effectively mitigating overfitting problems. Moreover, 1D-CNN achieves translational invariance through convolution and pooling operations, ensuring robustness to input data translations and enabling stable capture of key patterns in time-series data. Crucially, 1D-CNN possesses automatic feature extraction capabilities, allowing the model to independently learn and extract multi-level feature representations from raw data through end-to-end training, thus reducing dependence on manual feature engineering. As illustrated in **Figure 2**, the input sequence $x_1 \sim x_6$ undergoes convolution operations to generate feature mappings $y_1 \sim y_4$, with purple, green, and yellow connections linking the input layer to the convolution layer. Each connection maintains its distinct weight value, with connections of the same color sharing identical weights. By stacking multiple convolutional layers, the model progressively learns higher-level feature representations, offering robust expressive capabilities for time-series data modeling and prediction.




Figure 2. The one-dimensional (1D) convolution operation process.

2.3.2 LSTM

Long Short-Term Memory (LSTM) networks are specialized recurrent neural networks that effectively address the long-term dependency problems inherent in traditional RNN (Hochreiter and Schmidhuber, 1997). LSTM introduces memory cells and gating mechanisms that selectively remember or forget information, enabling the capture of long-term dependencies in sequences (Okut, 2021). The network uses three primary gating mechanisms: the forgetting gate, the input gate, and the output gate. The operating principle of the LSTM is illustrated in **Figure 3**, where σ represents the sigmoid function as shown in Eq. (1). Compared to traditional RNNs, LSTM networks exhibit superior handling of the vanishing gradient problem and can learn dependencies over longer time steps (Sherstinsky, 2020). These capabilities have led to the widespread application of LSTM in various domains, including time series prediction and natural language processing (Vennerød et al., 2021). The specific mathematical formulations of LSTM are detailed in Eqs. (1) - (6):

$$f_t = \sigma(W_f \cdot [h_{t-1}, x_t] + b_f) \quad (1)$$

$$i_t = \sigma(W_i \cdot [h_{t-1}, x_t] + b_i) \quad (2)$$

$$\tilde{c}_t = \tanh(W_c \cdot [h_{t-1}, x_t] + b_c) \quad (3)$$

$$c_t = f_t \times c_{t-1} + i_t \times \tilde{c}_t \quad (4)$$

$$o_t = \sigma(W_o \cdot [h_{t-1}, x_t] + b_o) \quad (5)$$

$$h_t = o_t \times \tanh(c_t) \quad (6)$$

Where W_f represents the weight matrix of the forget gate, and b_f denotes its bias term. h_{t-1} is the previous hidden state and x_t is the current input. The sigmoid activation function σ controls the proportion of information retention. i_t represents the output of the input gate, while $\tilde{c}_t$ indicates the candidate memory value. W_i and W_c represent the weight matrices for the input gate and candidate memory respectively, while b_i and b_c denote their corresponding bias terms. c_t represents the memory cell state at the current time step. W_o denotes the weight matrix of the output gate, b_o represents its bias term, and o_t indicates the output of the output gate.





Figure 3. The schematic diagram of the Long Short-Term Memory (LSTM).

2.3.3 CNN-LSTM

Several studies have shown that CNN-LSTM models have excellent performance in PM prediction, with low error rates and reduced training times (Li et al., 2020; Huang and Kuo, 2018). In this study, PM₁₀ concentration data were preprocessed for 11 chemical components: OM, EC, Na⁺, NH₄⁺, K⁺, Ca²⁺, Mg²⁺, F⁻, Cl⁻, NO₃⁻, and SO₄²⁻. Specifically, if the sum of the chemical components in a data set exceeded the PM₁₀ mass concentration or fell below 50% of the PM₁₀ mass concentration, the PM₁₀ concentration in that data set was considered anomalous and removed. After screening, the remaining data were retained and included in the training set. A hybrid model combining CNN and LSTM was used to predict PM₁₀ concentrations based on the training set. The model first extracts local features from the data through two CNN layers: the first CNN layer uses 16 channels and a kernel size of 2, while the second CNN layer uses 32 channels and the same kernel size, capturing local feature patterns through a sliding window with a stride of 1. Each CNN layer is followed by a ReLU activation function to introduce non-linearity, and a Dropout layer with a probability of 0.2 to enhance generalization capability. Subsequently, a 2-layer LSTM network (with 64 hidden units) captures long-term dependencies in the time series, with the LSTM layers also applying the same Dropout mechanism; finally, the prediction results are output through a fully connected layer. During the training process, Mean Squared Error (MSE) was used as the loss function, and the Adam optimizer was employed for parameter optimization, with an initial learning rate set to 0.0005. When training the model, a total of 3000 training epochs were set, while dynamically monitoring the loss value, with early stopping when the loss value fell below a preset threshold of 0.0007. Upon completion of training, the model was evaluated on both training and test sets by calculating MAE, RMSE, and Coefficient of Determination (R²) to comprehensively evaluate the predictive performance of the model.



Conv1D: One Dimensional Convolution Layer LSTM: Long Short Term Memory Neural Network
 FC: Fully Connected Neural Network ReLU: Rectified Linear Unit
 Dropout: A regularization technique to prevent overfitting

Figure 4. The architecture of the CNN-LSTM in this study.

2.4 Source apportionment

In this study, the US Environmental Protection Agency (US-EPA) EPA PMF 5.0 software (US-EPA, 2017) was used to perform source apportionment of PM₁₀. PMF is a multivariate statistical method based on factor analysis that has been widely applied in source apportionment studies of atmospheric particulate matter (Paatero and Tapper, 1994). The PMF model identifies pollution sources and their contribution rates by decomposing the observed data matrix into two non-negative matrices - the factor contribution matrix (G) and the factor profile matrix (F). The mathematical model can be expressed as:

$$X = GF + E \quad (7)$$

Where X is the observation data matrix ($n \times m$), G is the factor contribution matrix ($n \times p$), F is the factor profile matrix ($p \times m$), and E is the residual matrix. Here, n represents the number of samples, m represents the number of chemical species, and p represents the number of factors. The PMF model optimizes the decomposition results by minimizing the objective function Q :

$$Q = \sum_{i=1}^n \sum_{j=1}^m \left(\frac{x_{ij} - \sum_{k=1}^p g_{ik} f_{kj}}{u_{ij}} \right) \quad (8)$$

Where x_{ij} is the concentration of chemical component j in sample i , u_{ij} is the corresponding uncertainty, g_{ik} is the contribution of factor k in sample i , and f_{kj} is the proportion of chemical component j in factor k . By introducing non-negative

constraints, the PMF model can more reasonably explain the physical significance of pollution sources (Paatero, 1997). The uncertainty (Unc) of the sample data is calculated using Eqs. (9) and (10):

$$Unc = \frac{5}{6} \times MDL(x_{ij} < MDL) \quad (9)$$

$$Unc = \sqrt{(EF_{ij} \times x_{ij})^2 + (0.5 \times MDL)^2(x_{ij} \geq MDL)} \quad (10)$$

225 Where MDL represents the method detection limit, and EF_{ij} denotes the error fraction of component j in sample i . In this study, the EF values for OP_v were set as the standard deviation during analysis (Verma et al., 2015), while the other components were set at 10%.

3 Results and discussion

3.1 CNN-LSTM prediction results

230 The CNN-LSTM model was trained using non-outlier datasets consisting of PM_{10} concentration measurements and their corresponding eleven chemical constituents, including OM, EC, Na^+ , NH_4^+ , K^+ , Ca^{2+} , Mg^{2+} , F^- , Cl^- , NO_3^- , and SO_4^{2-} . To ensure the integrity of the data quality, outlier elimination was performed based on the sum of the chemical components. After the outlier screening process described in Section 2.3.3, 471 non-outlier datasets meeting the quality criteria were retained for model training and evaluation, with 85% allocated to the training set and 15% to the test set. The trained CNN-LSTM model
235 was then used to predict PM_{10} concentrations for the 766 outlier datasets by using their eleven chemical constituent concentrations as input features, with the predicted values replacing the original outlier measurements to maintain data completeness. Model performance was evaluated independently on both the training and test sets using three metrics: MAE, RMSE, and R^2 . These performance metrics are mathematically expressed in Eqs. (11) - (13). In these equations, y_i represents the actual value, $\hat{y}_i$ denotes the predicted value, n indicates the sample size, and $\bar{y}$ represents the mean of the actual values.

$$240 \quad MAE = \frac{1}{n} \sum_{i=1}^n |y_i - \hat{y}_i| \quad (11)$$

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2} \quad (12)$$

$$R^2 = 1 - \frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{\sum_{i=1}^n (y_i - \bar{y})^2} \quad (13)$$

3.1.1 Comparison with Conventional Gap-filling Techniques

The model was evaluated on both the training and test sets after completion of training, with results presented in **Table 2** and
245 **Figure 5**. As shown in **Figure 5 (a)**, the training process converged effectively, with the loss function decreasing steadily and

250

stabilizing at approximately 0.0007, indicating successful model optimization without overfitting. For the training set, the CNN-LSTM model achieved a MAE of $6.6614 \mu\text{g}\cdot\text{m}^{-3}$, a RMSE of $8.7162 \mu\text{g}\cdot\text{m}^{-3}$, and a R^2 of 0.9670. When evaluated on the test set, the model demonstrated an MAE of $12.6705 \mu\text{g}\cdot\text{m}^{-3}$, a RMSE of $17.4965 \mu\text{g}\cdot\text{m}^{-3}$, and an R^2 of 0.8840.

Table 2. Comparison of MAE, RMSE, and R^2 among different models.

Model Type	MAE ($\mu\text{g}\cdot\text{m}^{-3}$)	RMSE ($\mu\text{g}\cdot\text{m}^{-3}$)	R^2
Linear Regression	12.6852	17.8804	0.8028
RF	14.6494	20.0135	0.8482
KNN	15.6263	24.2398	0.8135
CNN-LSTM	12.6705	17.4935	0.8840




Figure 5. (a) LOSS trends for the training sets; comparison of predictions and observations for the (b) training and (c) test sets by the CNN-LSTM.

255

Traditional gap-filling techniques in air quality data reconstruction commonly include Linear Regression, Random Forest (RF), and k-nearest neighbors (KNN) methods, which have been widely applied in environmental data analysis (Méndez et al., 2023). To evaluate the effectiveness of our CNN-LSTM approach, we conducted a comprehensive comparison with these conventional methods and individual deep learning components, with detailed results presented in **Table 2**.

260

In the ranking of MAE performance, from low to high, the results are: CNN-LSTM ($12.6705 \mu\text{g}\cdot\text{m}^{-3}$), Linear Regression ($12.6852 \mu\text{g}\cdot\text{m}^{-3}$), RF ($14.6494 \mu\text{g}\cdot\text{m}^{-3}$), and KNN ($15.6263 \mu\text{g}\cdot\text{m}^{-3}$). While in the ranking of RMSE, from low to high, the performance follows: CNN-LSTM ($17.4935 \mu\text{g}\cdot\text{m}^{-3}$), Linear Regression ($17.8804 \mu\text{g}\cdot\text{m}^{-3}$), RF ($20.0135 \mu\text{g}\cdot\text{m}^{-3}$), and KNN ($24.2398 \mu\text{g}\cdot\text{m}^{-3}$). In terms of R^2 , from high to low, the algorithms perform as follows: CNN-LSTM (0.8840), RF (0.8482), KNN (0.8135), and Linear Regression (0.8028).

265

This study shows that the CNN-LSTM model proposed in this paper demonstrates good performance across all evaluation metrics. Linear regression, despite achieving competitive MAE values, shows limitations in capturing complex non-linear relationships, as evidenced by its lower R^2 value. This can be attributed to the linear model's relative inability to capture the

complex non-linear relationships inherent in atmospheric particulate matter dynamics (Singh et al., 2012; Dragomir and Oprea, 2014). RF demonstrated moderate performance with balanced metrics across MAE, RMSE, and R^2 . KNN showed the least effective overall performance, particularly evident in its highest RMSE value of $24.2398 \mu\text{g}\cdot\text{m}^{-3}$.

Overall, CNN-LSTM is the strongest performer among all tested models. This result confirms that the combination of CNN and LSTM is quite effective for PM_{10} gap-filling applications. We conducted ablation experiments in **Section S1**. As shown in **Table S1**, the performances of CNN and LSTM are both good, but that of CNN-LSTM is even more outstanding. It is indicated that for PM_{10} data reconstruction, it is beneficial to first perform feature extraction using CNN, and then input the feature values into the LSTM architecture for temporal pattern recognition.

3.1.2 Leave-One-Site-Out Cross-Validation

To validate the model's generalization capability and ensure it captures physically meaningful variability rather than site-specific biases, we employed a site-type-based Leave-One-Site-Out (LOSO) cross-validation strategy. The monitoring sites within the study area were categorized into four types based on their functional characteristics and geographical environment: urban sites, rural sites, suburban sites, and remote sites. LOSO cross-validation was performed separately for each site type. For each iteration, one site of a particular type was withheld during model training. The model was then trained using data from the remaining sites and tested on the withheld site.

Table 3. Results of Leave-One-Site-Out (LOSO) cross-validation for different site types.

Site Type	MAE ($\mu\text{g}\cdot\text{m}^{-3}$)	RMSE ($\mu\text{g}\cdot\text{m}^{-3}$)	R^2
Urban	10.9436	15.2544	0.9235
Rural	12.5448	17.6598	0.8297
Suburban	14.5934	19.0105	0.8450
Remote	7.8523	9.3182	0.7892

Table 3 presents the model performance metrics for different site types in the LOSO cross-validation, including MAE, RMSE, and R^2 . The cross-validation results demonstrate that the established neural network model exhibits good predictive performance across different site types, with R^2 values exceeding 0.78 for all site types. This confirms that the model can effectively reproduce the physically meaningful variability in pollutant concentrations rather than merely fitting site-specific characteristics.

Remote stations exhibit the lowest MAE ($7.8523 \mu\text{g}\cdot\text{m}^{-3}$) and RMSE ($9.3182 \mu\text{g}\cdot\text{m}^{-3}$), indicating minimal prediction errors at these site types. This is primarily attributed to remote stations being located far from major pollution sources, resulting in relatively low pollutant concentrations with gradual variations that reduce absolute prediction errors. However, Remote stations show a relatively low R^2 value (0.7892), which may be related to their smaller concentration variation range, making the correlation coefficient more sensitive to minor prediction deviations. Urban stations demonstrate the highest R^2 value

(0.9235), while maintaining moderate MAE ($10.9436 \mu\text{g}\cdot\text{m}^{-3}$) and RMSE ($15.2544 \mu\text{g}\cdot\text{m}^{-3}$) levels. This result indicates that the model can effectively capture pollutant concentration trends in urban environments. Urban areas feature dense and diverse pollution sources, including vehicle emissions and industrial emissions, which increase prediction complexity but provide larger concentration variation ranges that facilitate the establishment of robust predictive relationships. Rural stations show moderate performance across all metrics, with MAE of $12.5448 \mu\text{g}\cdot\text{m}^{-3}$, RMSE of $17.6598 \mu\text{g}\cdot\text{m}^{-3}$, and R^2 of 0.8297. Rural areas are primarily influenced by relatively simple pollution sources such as regional transport and agricultural activities, resulting in more regular spatiotemporal variation patterns of pollutant concentrations that the model can effectively learn. Suburban stations display the highest MAE ($14.5934 \mu\text{g}\cdot\text{m}^{-3}$) and RMSE ($19.0105 \mu\text{g}\cdot\text{m}^{-3}$), but with an R^2 value (0.8450) similar to rural stations. As transitional zones between urban and rural areas, suburban regions are simultaneously influenced by urban pollution dispersion and rural pollution sources, exhibiting distinct transitional and complex pollution characteristics. This composite pollution environment increases the difficulty of model prediction and may result in higher prediction errors.

3.2 PM₁₀ mass and chemical composition concentrations

3.2.1 Annual average

The analysis of PM₁₀ concentrations across diverse locations in China shows a remarkable spatial variation in the annual mean concentrations of PM₁₀ and its chemical constituents from June 2022 to May 2023, as shown in **Table 3**. Significantly elevated PM₁₀ levels were observed at northwestern sites, with Xi'An (XA) and Dunhuang (DH) recording concentrations of $98.20 \mu\text{g}\cdot\text{m}^{-3}$ and $90.36 \mu\text{g}\cdot\text{m}^{-3}$, respectively, while other sites had concentrations ranging from 40 to $80 \mu\text{g}\cdot\text{m}^{-3}$. These spatial patterns suggest complex interactions between natural and anthropogenic factors. The elevated PM₁₀ concentrations observed in XA, a major industrial city and densely populated metropolitan area, are primarily due to industrial emissions and substantial high traffic volumes. Due to its location in an arid region, DH is likely influenced by dust storm events, as evidenced by higher concentrations of crustal elements such as Ca^{2+} (Yu et al., 2020). While Na^+ is typically associated with sea salt spray, its presence at inland sites such as DH may indicate contributions from crustal material or other local sources (Zhang et al., 2014b). In contrast, the lowest PM₁₀ concentrations were observed at Longfengshan (LFS) and Dalian (DL) in the northeastern region, with values of $40.04 \mu\text{g}\cdot\text{m}^{-3}$ and $40.35 \mu\text{g}\cdot\text{m}^{-3}$, respectively. These relatively lower concentrations may be due to relatively less anthropogenic activities and better air quality in these regions. LFS, located at the interface of agricultural and forested landscapes, primarily receives PM₁₀ contributions from natural sources, such as soil dust resuspension and biomass burning reported in previous research (Yu et al., 2012). Meanwhile, Dalian's coastal location likely contributes to its lower PM₁₀ concentrations. The observed Na^+ concentration of $2.36 \mu\text{g}\cdot\text{m}^{-3}$ in DL may reflect the influence of marine aerosols (Shi et al., 2022). In addition, air quality in DL is likely modulated by meteorological conditions, especially sea breezes, which facilitate the dispersion and dilution of pollutants, thereby reducing PM₁₀ concentrations (Wang et al., 2002). In the densely populated regions of Gucheng (GC) and Zhengzhou (ZZ), where anthropogenic pollution sources are abundant, the annual mean PM₁₀ concentrations were $79.18 \mu\text{g}\cdot\text{m}^{-3}$ and $80.50 \mu\text{g}\cdot\text{m}^{-3}$, respectively. These elevated PM₁₀ levels are

strongly correlated with intensive anthropogenic sources in these regions, including industrial activities, traffic emissions, and construction dust. As major industrial and transportation hubs, GC and ZZ have particularly high concentrations of organic matter ($OM=1.2*OC$) and elemental carbon (EC), specifically $19.67 \mu\text{g}\cdot\text{m}^{-3}$ and $4.89 \mu\text{g}\cdot\text{m}^{-3}$ in GC, $17.35 \mu\text{g}\cdot\text{m}^{-3}$ and $4.12 \mu\text{g}\cdot\text{m}^{-3}$ in ZZ. Additionally, the concentrations of sulfate (SO_4^{2-}) and nitrate (NO_3^-) concentrations in ZZ and GC were measured to be $8.70 \mu\text{g}\cdot\text{m}^{-3}$, $13.71 \mu\text{g}\cdot\text{m}^{-3}$ and $6.00 \mu\text{g}\cdot\text{m}^{-3}$, $10.94 \mu\text{g}\cdot\text{m}^{-3}$, respectively. These values, which are significantly higher than in other regions, indicate particularly active secondary aerosol formation processes in these areas (Yang et al., 2020b).

In the southwestern region, Chengdu (CD), located in the Sichuan Basin, recorded an annual mean PM_{10} concentration of $59.56 \mu\text{g}\cdot\text{m}^{-3}$. This region is characterized by high aerosol optical depth and reduced visibility, attributed to poor dispersion conditions and significant local industrial emissions (Li et al., 2003; Zhang et al., 2012).

The central Chinese sites of Jinsha (JS), Changde (CHD), and Lin'An (LA) showed relatively lower annual mean concentrations of PM_{10} which are $47.17 \mu\text{g}\cdot\text{m}^{-3}$, $46.59 \mu\text{g}\cdot\text{m}^{-3}$, and $48.16 \mu\text{g}\cdot\text{m}^{-3}$, respectively. Despite these lower concentrations, the chemical composition shows distinct regional characteristics. Ca^{2+} concentrations of $2.48 \mu\text{g}\cdot\text{m}^{-3}$ and $2.19 \mu\text{g}\cdot\text{m}^{-3}$ in JS and LA, respectively, likely reflect contributions from soil dust resuspension (Shen, 2016). K^+ concentration of $0.44 \mu\text{g}\cdot\text{m}^{-3}$ observed in CHD may be related to agricultural activities in the region (Liu et al., 2016).


Lhasa (LS), located in the center of the Tibetan Plateau at an elevation of 3,663 meters, has PM_{10} concentrations that are primarily influenced by natural factors due to its relatively sparse population and limited industrial emissions. Nevertheless, LS maintained an average PM_{10} concentration of $47.82 \mu\text{g}\cdot\text{m}^{-3}$, mainly due to extensive dust resuspension from arid and exposed terrain, coupled with regional dust storm events. The plateau's climatic conditions, characterized by particularly strong winds and low humidity, enhance the dispersal of soil dust and maintain relatively high PM_{10} levels despite the absence of significant anthropogenic sources.

The annual mean PM_{10} concentrations for urban, rural, suburban, and remote sites were $59.99 \pm 29.38 \mu\text{g}\cdot\text{m}^{-3}$, $62.88 \pm 27.58 \mu\text{g}\cdot\text{m}^{-3}$, $85.43 \pm 39.43 \mu\text{g}\cdot\text{m}^{-3}$, and $45.12 \pm 14.67 \mu\text{g}\cdot\text{m}^{-3}$, respectively. These data show that urban-rural transition zones had the highest PM_{10} concentrations, which may be due to the simultaneous influence of multiple pollution sources from both urban and rural areas, including industrial emissions, traffic pollution, and agricultural activities (Li et al., 2014). In contrast, remote sites had the lowest PM_{10} concentrations, reflecting minimal anthropogenic influence in these regions, with primary pollution sources consisting of natural dust resuspension and long-range transported pollutants (Jiao et al., 2021).

Table 3. Annual average concentrations ($\mu\text{g}\cdot\text{m}^{-3}$) of PM_{10} and its chemical composition in different regions of China from June 2022 to May 2023.

Station	Type	PM_{10}	OM	EC	Na^+	NH_4^+	K^+	Mg^{2+}	Ca^{2+}	F^-	Cl^-	SO_4^{2-}	NO_3^-
Chengdu	Urban	59.56	17.09	3.97	2.18	2.11	0.30	0.23	2.67	0.15	0.58	6.29	9.36
Dalian	Urban	40.35	9.35	2.30	2.36	0.74	0.25	0.25	1.89	0.04	0.69	3.19	5.00
Lhasa	Urban	47.82	16.85	4.16	2.51	0.07	0.32	0.26	1.75	0.05	1.18	1.55	1.12
Nanning	Urban	54.23	12.87	3.50	2.03	1.20	0.37	0.21	2.89	0.07	0.64	7.21	5.09
Xi'An	Urban	98.20	19.13	4.87	2.50	2.64	0.76	0.37	4.97	0.15	1.67	8.67	12.82
Changde	Rural	46.59	9.05	2.17	0.44	2.76	0.44	0.08	1.02	0.03	0.27	6.16	6.18
Gucheng	Rural	79.18	19.67	4.89	2.08	1.78	0.35	0.46	4.01	0.09	1.21	6.00	10.94
Dunhuang	Suburban	90.36	23.24	4.78	4.43	0.16	0.36	0.46	6.31	0.06	2.57	5.90	2.29
Zhengzhou	Suburban	80.50	17.35	4.12	1.71	3.43	0.45	0.32	3.03	0.21	0.86	8.70	13.71
Jinsha	Remote	47.17	12.14	2.07	1.52	1.45	0.40	0.20	2.48	0.08	0.58	5.82	6.89
Lin'An	Remote	48.16	13.02	2.92	1.37	1.46	0.34	0.22	2.19	0.04	0.66	5.37	7.42
Longfengshan	Remote	40.04	12.31	2.52	1.21	1.14	0.36	0.15	1.61	0.06	0.50	4.04	4.25

3.2.2 Seasonal variation



355 **Figure 6. Stacked Representation of Monthly averaged PM₁₀ Concentrations and Chemical Composition (µg·m⁻³) across Chinese Regions, Including Unknown Components from June 2022 to May 2023 (n.d: Unknown Components). The map base is from the Ministry of Natural Resources' Standard Map Service, review number GS (2019)1822.**

Monthly variations in PM₁₀ concentrations are shown in **Figure 6**. Overall, the study area shows a significant seasonal differentiation of PM₁₀ concentrations, characterized by minimum levels in summer (June-August), maximum levels in winter (December-February), and a secondary peak in spring (March-May). Multiple studies have also identified distinct seasonal patterns in PM₁₀ concentrations, with minimal concentrations in summer and maximal concentrations in winter (Yang, 2009; Qu et al., 2010; Li et al., 2009). The lower PM₁₀ concentrations observed in summer may be attributed to increased precipitation, which effectively scavenges atmospheric particulate matter (Yang, 2009). In addition, research has shown significant negative correlations between PM₁₀ concentrations and temperature, as well as positive correlations with atmospheric pressure (Han et al., 2015; Li et al., 2019). Elevated PM₁₀ concentrations in winter are primarily associated with increased solid fuel consumption during the heating season (Tsvetanova et al., 2017). Additionally, unfavorable meteorological conditions in winter, including high atmospheric stability, reduced atmospheric boundary layer height, and frequent temperature inversions,

exacerbate the accumulation of pollutants (Zhao et al., 2014). Five monitoring stations including GC, LFS, DH, LA, and Nanning (NN) exhibited significantly elevated concentrations during spring, which can be attributed to multiple factors. Firstly, the frequent occurrence of dust events during spring increases atmospheric particulate matter concentrations. Secondly, weak wind conditions and local circulation patterns establish local emissions as the primary source of PM₁₀ (Park et al., 2019). Moreover, regional transport represents a significant influencing factor, with studies indicating substantial contributions to PM₁₀ concentrations from dust transport from northwestern regions and pollutant transport from surrounding urban agglomerations in spring (Ham et al., 2017).

The results indicate significant seasonal variations in monthly mean concentrations of OM and EC in urban, rural, and suburban sites. All three functional site types showed the lowest concentrations in summer and the highest in winter, consistent with previous studies confirming the widespread winter-high and summer-low seasonal pattern of carbonaceous components in PM₁₀ across China (Tian et al., 2013). The elevated concentrations of OM and EC in winter correlate primarily with increased fossil fuel and biomass combustion emissions during the heating season, coupled with unfavorable meteorological dispersion conditions. Conversely, the decrease concentrations in summer are attributed to increased precipitation, increased mixing layer height, and reduced stationary source emissions due to higher temperatures. However, remote sites showed different seasonal patterns than other sites, with OM and EC concentration peaks occurring in spring and fall. This phenomenon may be associated with regional-scale dust transport, biomass burning activities, and increased open-source emissions, while also reflecting minimal local anthropogenic influence at remote sites, better representing regional background concentration variations.

We observed generally higher concentrations of SO₄²⁻ and NO₃⁻ in winter compared to lower concentrations in summer. This seasonal pattern is primarily due to increased SO₂ and NO_x emissions from extensive fossil fuel combustion, especially coal, during the winter heating season, which provides abundant precursors for the formation of sulfate and nitrate. In addition, stable atmospheric stratification and frequent temperature inversions in winter inhibit the dispersion of pollutants, leading to near-surface accumulation of these secondary inorganic ions. Furthermore, the relatively lower temperatures in winter facilitate the gas-to-particle conversion of gaseous precursors, promoting the partitioning of semi-volatiles such as ammonium sulfate and ammonium nitrate to the particulate phase (Wang et al., 2020). In contrast, higher summer temperatures favor the gaseous state of these semi-volatile substances, while frequent convection and stronger atmospheric dispersion conditions significantly reduce sulfate and nitrate concentrations in PM₁₀ (Simonich and Hites, 1994). This seasonal pattern is consistent with observations from other regional studies and reflects the close relationship between secondary inorganic ion formation mechanisms and meteorological conditions (Liu et al., 2017a; Wang et al., 2023a) .

3.3 OP concentrations

As shown in **Figure 7**, OP measurements conducted at twelve different sampling sites across China from June 2022 to May 2023 revealed significant temporal and spatial variability in OP_v. Further analysis revealed a strong correlation between OP_v and the degree of urbanization at the sampling sites. During the sampling period, the urban site in Chengdu had significantly

higher OP_v levels compared to the other sites, while the rural site in CHD had the lowest OP_v levels. However, the study revealed unexpectedly high average OP_v levels at the rural site in GC, ranking second highest among all sites. This finding is consistent with the high PM_{10} mass concentrations observed at this site, suggesting a strong correlation between particulate matter loading and OP_v levels. GC, located in the Beijing-Tianjin-Hebei region characterized by high population density and typical pollution concentration, experiences elevated OP_v levels likely due to the combined influence of high PM_{10} concentrations, pollutant transport from surrounding urban areas, and local emissions (Han et al., 2015). In contrast, the urban site in DL demonstrated relatively low average OP_v levels, ranking second lowest. This phenomenon may be attributed to the coastal location of DL, which benefits from strong marine air mass modulation and favorable atmospheric dispersion conditions (Meng et al., 2019), resulting in comparatively lower OP_v levels.




Figure 7. Seasonal variations of (a) PM_{10} concentrations ($\mu g \cdot m^{-3}$) and (b) OP_v ($nmol H_2O_2 \cdot m^{-3}$) across different regions of China. The map bases are from the Ministry of Natural Resources' Standard Map Service, review number GS (2019)1822.

As shown in **Figure 7 (a) and (b)**, sites located in northern China exhibited significantly elevated PM_{10} concentrations and OP_v levels during the autumn and winter seasons. This phenomenon in northern Chinese sites can be attributed to several factors unique to northern China's regional characteristics. Firstly, the widespread reliance on coal-based central heating systems and biomass burning for residential heating in northern China during the heating season (typically from November to March) (Liu et al., 2017b; Li et al., 2017) sharply contrasts with southern China where heating demand is minimal due to milder winter temperatures. In addition, northern China's continental climate creates more severe winter meteorological conditions, including prolonged periods of low wind speeds, frequent temperature inversions, and significantly reduced atmospheric boundary layer heights compared to the more temperate conditions in southern regions, which severely inhibited pollutant dispersion (Li et al., 2017). Despite lower levels of urbanization in rural areas, PM_{10} concentrations were comparable to urban areas due to the widespread use of solid fuels (Li et al., 2014). **Figure 7 (b)** shows that nine of the twelve sites had

lower OP_v values in summer. This may be due to more frequent rainfall, which reduces PM_{10} concentrations and subsequently leads to lower OP_v levels. However, sites such as LS and CD maintained relatively high OP_v levels during the summer. This phenomenon may be related to the enhanced of photochemical reactions during summer, especially under conditions of high temperature and strong solar radiation, resulting in a significant increase in secondary organic aerosol (SOA) formation (Zhou et al., 2019; Saffari et al., 2014). In particular, Lhasa's high-altitude location, characterized by minimal precipitation and intense solar radiation, further promoted photochemical reactions, resulting in elevated OP_v levels.

We observed elevated OP_v levels at remote stations (such as LFS, JS, and LA stations) in spring. This phenomenon may be attributed to the minimal influence of anthropogenic pollution sources at remote stations, which typically exhibit more homogeneous mixing states and consequently have relatively lower and more stable OP_v levels during other seasons. However, the frequent occurrence of dust storms and increased temperature inversion events during spring can lead to elevated particulate matter concentrations. In addition, the potential metal components carried by dust particles and the formation of secondary aerosols further enhance OP_v levels (Saffari et al., 2014), resulting in significantly elevated OP_v levels during spring.

Table 4. Annual averaged OP_v ($nmol\ H_2O_2\cdot m^{-3}$) for PM_{10} across different regions of China from June 2022 to May 2023.

Station	OP_v	
	Average	Median
Chengdu	0.85	0.57
Dalian	0.30	0.14
Lhasa	0.60	0.57
Nanning	0.56	0.50
Xi'An	0.73	0.74
Changde	0.22	0.21
Gucheng	0.83	0.75
Dunhuang	0.76	0.50
Zhengzhou	0.42	0.40
Jinsha	0.54	0.40
Lin'An	0.46	0.45
Longfengshan	0.57	0.52

We conducted a discussion on the differences between northern and southern sites across these 11 stations (excluding LS) in China. The geographical division corresponding to the station is shown in **Table S2**. The distinction between northern and southern sites and the specific analytical methods are detailed in **S2**. As shown in **Figure 8(a)**, OP_v concentrations in northern regions exhibited higher levels during the winter, primarily due to increased pollutant emissions associated with coal-based heating activities. In contrast, southern regions exhibited peak OP_v concentrations in June, possibly due to enhanced

photochemical reactions facilitated by stronger solar radiation intensity. However, a significant decrease was observed in July and August, which may be attributed to the increased frequency of precipitation events leading to enhanced wet deposition and the removal of particulate matter. **Figure 8 (b)** shows that the annual mean OP_v concentrations in northern regions were significantly higher than those in southern regions ($p < 0.05$). This spatial variation can be attributed to several factors, including lower precipitation rates, frequent dust weather events, and emissions of coal combustion emissions in northern regions.




Figure 8. Comparison of (a) monthly and (b) annual average OP_v between sites in northern and southern China.

3.4 Source apportionment

3.4.1 Source apportionment of PM_{10}

This study employed the PMF model to conduct a detailed analysis of PM_{10} sources at four representative sites selected based on distinct geographical and environmental characteristics. The selection criteria considered regional representativeness, pollution characteristics, and geographical diversity across China. The selected sites include: NN, an urban site in southern China with coastal proximity; Longfengshan (LFS), a remote site located in the northeastern region of Heilongjiang Province; ZZ, a suburban site serving as a major transportation hub in central China; and GC, a rural site situated in the heavily polluted Beijing-Tianjin-Hebei region. These four sites collectively represent different pollution source characteristics and regional environmental conditions, enabling a comprehensive understanding of PM_{10} source apportionment across diverse geographical and climatic zones in China. The optimal number of factors for PMF analysis was determined based on Q_{true}/Q_{robust} values and BS mapping evaluation, as illustrated in **Figure S1**. The PMF results in this study were subjected to BS, DISP, and BS-DISP error estimation analyses. Summary of error estimation diagnostics with PMF at NN, LFS, ZZ and GC stations are shown in **Table S3**. Results indicate that PM_{10} in NN likely originates primarily from biomass burning, traffic, dust, secondary aerosols,

and sea salt emissions. Sources of PM₁₀ in LFS may include biomass burning, traffic, dust, agricultural activities, and secondary aerosols. The ZZ site showed industry, agricultural activities, traffic, coal combustion, dust, and secondary aerosols as the main sources. PM₁₀ sources in GC are biomass burning, traffic, dust, agricultural activity emissions, secondary aerosols, and coal combustion. **Figure 9** summarizes the distribution of PM₁₀ mass concentrations among the major sources at the four sites.





Figure 9. The contributions of Biomass burning, Traffic, Dust, Secondary aerosol, Sea salt, Agricultural activities, Coal combustion, and Industry to the atmospheric concentration of PM₁₀ mass (%) as derived by PMF modelling at NN, LFS, ZZ, and GC.



470 **Figure 10. Chemical profiles of the source factors identified at NN, LFS, ZZ and GC. The bars represent the chemical composition profiles (left y-axis) and the dots indicate the contribution values (right y-axis).**

As a typical urban site, the PM₁₀ source apportionment results at the NN site indicate that biomass burning, dust, and traffic are likely the main contributors, accounting for 39.3%, 27.4%, and 21.4% of total sources, respectively. As shown in **Figure 10**, the first factor contained high levels of Cl⁻ (76.9%), Mg²⁺ (27.6%), and Na⁺ (14.3%), elements typically associated with sea salt (Viana et al., 2008), contributing approximately 5% to PM₁₀. Sea salt as a source of PM₁₀ in NN likely enters urban areas primarily through coastal air mass transport. NN is about 110 kilometers from the Beibu Gulf, and when prevailing southerly winds occur, sea salt aerosols from the South China Sea may migrate to inland cities through atmospheric circulation. The second factor contained high levels of Na⁺ (66.7%), Ca²⁺ (48.1%) and Mg²⁺ (17.5%), contributing approximately 26% to PM₁₀. This likely represents dust sources (Sharma et al., 2016), indicating that human activities such as urban construction may have some impact on particulate emissions. The third factor had high levels of NH₄⁺ (84.2%), SO₄²⁻ (45.7%), and NO₃⁻ (44.4%), contributing approximately 6.9% to PM₁₀, possibly representing secondary aerosols. This suggests that the process of gaseous precursors (such as SO₂, NO_x, and VOCs) in the atmosphere forming secondary particles through photochemical reactions may have a certain impact on PM₁₀ concentrations (Yue et al., 2015). The fourth factor contained high levels of Mg²⁺ (54.8%), EC (50.7%) and OC (38.2%), contributing approximately 20.2% to PM₁₀, possibly related to traffic. EC and OC have long been considered the main tracer elements for traffic emission sources, particularly vehicle exhaust emissions (Saarikoski et al., 2008; Sowlat et al., 2016; Esmaeilirad et al., 2020). Research has shown that Mg is one of the elements present in high concentrations in brake pad materials. Mg is typically used as a filler material in brake pads, and along with Fe, Ba and Cu, serves as a characteristic element of brake wear (Mckenzie et al., 2009). At the NN urban site, which is heavily influenced by traffic, brake wear is likely the primary source of these elements. The fifth factor had high levels of K⁺ (74.9%), OC (37.2%), and EC (35.1%), substances typically associated with biomass burning (Stracquadiano et al., 2019). This factor made a significant contribution to urban PM₁₀ in NN, approximately 39.3%, indicating that biomass burning may be one of the important sources of atmospheric particulate pollution in NN. Although the observation point is located in the urban area of NN, which may be at some distance from areas where straw burning occurs, studies have shown that particulate matter produced by biomass burning may undergo long-distance transport (Uranishi et al., 2019).

The PM₁₀ source apportionment results for LFS indicate that secondary aerosols may be the main contributor, accounting for 36.2% of total sources. Source analysis identified five potential major factors: In the first factor, NH₄⁺ (71.0%), Mg²⁺ (26.5%), and NO₃⁻ (18.0%) were present in high concentrations. NH₄⁺ and NO₃⁻ are the main nitrogen components in agricultural fertilizers (Cao et al., 2018), while Mg²⁺ is commonly added to fertilizers as a supplementary element (Sun et al., 2018). This factor may be related to agricultural activities, particularly fertilizer application processes. The second factor contained high levels of Na⁺ (74.6%), Mg²⁺ (46.2%), and Ca²⁺ (50.8%), elements typically associated with dust sources (Zhang et al., 2014a; Sharma et al., 2016), contributing approximately 16.9% to PM₁₀. The third factor had high levels of EC (74.0%) and OC (38.3%), components typically associated with traffic (Esmaeilirad et al., 2020), contributing approximately 17.6%. The fourth factor contained high levels of Cl⁻ (79.2%), Mg²⁺ (22.7%), OC (22.5%), and K⁺ (15.2%), among them, K⁺ and Cl⁻ have been identified as reliable indicators of biomass burning (Saggu and Mittal, 2020), contributing approximately 13% to PM₁₀. The

505 fifth factor had high levels of SO_4^{2-} (70.7%) and NO_3^- (71.9%), with NH_4^+ (27.3%) also making a considerable contribution, these components are typically associated with secondary aerosol formation processes (Yue et al., 2015).

ZZ is located in a suburban area, and the diversity of its PM_{10} sources may reflect the complex environmental characteristics of this region. Source apportionment results suggest that there may be six major pollution sources in this area, with their respective contribution proportions as follows: The first factor had high contribution of K^+ (21.7%) and Cl^- (83.9%), but low
510 contribution of OC (4.6%) and EC (2.8%), possibly indicating the influence of industrial emissions, such as food manufacturing, cement manufacturing, salt production, or industrial activities involving potassium chloride compounds (Yin et al., 2019; Seo et al., 2019), with a contribution proportion of approximately 9.5%. The second factor contained high levels of Na^+ (77.6%), Mg^{2+} (35.3%), and Ca^{2+} (43.2%), elements typically associated with dust sources (Sharma et al., 2016), contributing approximately 13.2% to PM_{10} . In the third factor, Mg^{2+} (42.1%) and SO_4^{2-} (46.9%) had relatively high
515 concentrations. Since SO_4^{2-} primarily originates from fuel combustion (Schwartz, 1993), and Mg is specifically mentioned as an element enriched in the magnetic separation of coal fly ash (Strzałkowska, 2021), this factor is associated with emissions from coal combustion when regional characteristics are considered. Coal combustion accounts for around 15.5% of PM_{10} emissions and is likely to be associated with combined heat and power facilities in the surrounding area. The fourth factor had high levels of EC (49.2%) and OC (22.1%), components typically associated with traffic (Esmaeilirad et al., 2020), contributing
520 approximately 16.6%. The fifth factor contained high levels of NH_4^+ (80.1%), SO_4^{2-} (33.0%), and NO_3^- (52.6%), components typically associated with secondary aerosol formation processes (Yue et al., 2015), accounting for approximately 23.2% of total PM_{10} sources. The sixth factor had high levels of K^+ (60.6%), Ca^{2+} (38.4%), EC (39.3%), OC (28.8%), and NO_3^- (25.0%), based on comprehensive analysis of these characteristic species, this factor may be related to agricultural activity emissions, contributing approximately 22% to PM_{10} . Ca^{2+} , OC and EC may be related to surface soil dust (Yu and Cao, 2023), Jung et al.
525 found elevated K^+ concentrations at schools near corn farms, supporting the agricultural source attribution (Jung et al., 2024), while NO_3^- would be related to fertilizer application (Cao et al., 2018).


The PM_{10} source apportionment results for the GC show that agricultural activities, traffic emissions, secondary aerosols, and biomass burning are the main contributors, accounting for 20.5%, 20%, 18.5%, and 18.1% of total sources, respectively. The factor with K^+ (42.7%), NO_3^- (38.4%), and Ca^{2+} (29.2%) as primary characteristic species may be related to agricultural
530 activities, accounting for 20.5%. This likely reflects the contribution of corn, wheat, and other farming activities around the site to PM_{10} , potentially associated with the agricultural-dominant economic structure of this rural area. Ca^{2+} and NO_3^- may originate from agricultural soil dust during tillage and other agricultural processes, and NO_3^- could be related to fertilizer application (Yu and Cao, 2023; Cao et al., 2018). Similar to the ZZ site, this agricultural source attribution is supported by Jung et al., who found elevated K^+ concentrations at schools near corn farms (Jung et al., 2024). The factor characterized by
535 EC (65.1%) and OC (48.1%) likely comes from traffic (Esmaeilirad et al., 2020), representing the second-largest contributor to PM_{10} at 20%. This indicates that transportation activities in rural areas may have a significant impact on PM_{10} concentrations. The GC is relatively close to National Highway 107, and traffic emissions from the highway may contribute to the site's PM_{10} concentration through transport. Additionally, the increasing vehicle ownership in rural areas may be a contributing factor.

Secondary aerosols, characterized by NH_4^+ (93.0%), SO_4^{2-} (45.6%), and NO_3^- (52.3%), account for 18.5%, indicating the important role of atmospheric secondary transformation processes in PM_{10} formation in this region (Yue et al., 2015). The factor characterized by K^+ (22.3%) and Cl^- (76.2%) may be related to biomass burning (Saggu and Mittal, 2020), accounting for 18.1%. This could be associated with activities such as straw burning and residential fuel use, particularly during crop harvest seasons and winter heating periods when such activities may increase. The factor characterized by Na^+ (75.2%) and Ca^{2+} (44.6%) may be related to dust (Sharma et al., 2016), accounting for 14.7%, potentially reflecting the impact of agricultural cultivation and road dust on PM_{10} . The factor characterized by Mg^{2+} (43.0%) and SO_4^{2-} (47.8%) may be related to coal combustion emissions, accounting for 8.3%. This suggests that industrial activities and residential coal use in rural areas may have some impact on PM_{10} , especially during the winter heating season when such emissions may become more prominent. The contrasting OC/EC loadings in the agricultural activities factor between the suburban ZZ site and rural GC site reveal important insights into the spatial heterogeneity of agricultural emissions. The suburban ZZ site, located in the intensively cultivated Central Plains, experiences higher carbonaceous aerosol loadings from mechanized farming operations, which contribute significantly to EC emissions through diesel exhaust from agricultural machinery (Liu et al., 2018). In contrast, the rural GC site in Baoding represents areas with traditional, less mechanized farming practices, resulting in minimal EC contributions from agricultural activities.

A notable pattern observed among the three sites with agricultural activities (LFS, ZZ, and GC) is the differential contribution of NH_4^+ within agricultural emission factors, with NH_4^+ being exclusively associated with agricultural activities at the LFS site. This spatial variation reflects the complex interplay between regional meteorological conditions, agricultural practices, and atmospheric chemistry processes. At the LFS site in northeastern China, cooler climate conditions favor the stability of particulate NH_4^+ , allowing its direct retention within agricultural emission factors (Wang et al., 2020). The concentrated fertilizer application during the spring planting season, combined with lower ambient temperatures that minimize NH_4^+ volatilization, preserves the distinct agricultural source signature at this remote location (Huo et al., 2025). Conversely, at the warmer ZZ and GC sites in central and northern China, NH_4^+ undergoes more extensive atmospheric processing due to higher ambient temperatures. These conditions promote the volatilization of NH_4^+ to gaseous NH_3 , which subsequently undergoes secondary reactions with acidic species (SO_4^{2-} and NO_3^-) to form ammonium-containing secondary aerosols (Stelson and Seinfeld, 1982; Wang et al., 2015).

3.4.2 Source apportionment of OP in PM_{10}

In this study, the PMF model was applied to identify the sources of OP_v in PM_{10} samples collected from four sites. Through comparison of OP_m across different pollution sources, insights into the differential toxic efficiencies of various sources were obtained.



570 **Figure 11. Comparison of OP_m contributions from different emission sources across NN, LFS, ZZ and GC sites.**

Traffic

Traffic emission is a common important contributing source to OP_v at the four sites NN, LFS, ZZ and GC. The high OP of traffic emissions is mainly attributed to the oxidative components in their particulate matter emissions, including organic carbon as well as potentially present PAHs and transition metals (TMs) (Valavanidis et al., 2008). Traffic sources showed high OP_m (0.013-0.022 nmol $H_2O_2 \cdot \mu g^{-1}$) at all monitoring sites, indicating that the particulate matter they generate has significant toxic efficiency. The high oxidative activity of traffic-emitted particulate matter may be related to its complex chemical composition. PAHs emitted from traffic sources can form quinone compounds and other oxygen-containing organic compounds, which can promote reactive oxygen species generation through redox reactions (Nielsen, 1996; Libalova et al., 2018).

580 Biomass burning

Biomass burning sources were only detected to contribute to OP at NN and GC sites, with OP_m of 0.019 and 0.006 nmol $H_2O_2 \cdot \mu g^{-1}$, respectively. Notably, biomass burning has the highest OP_m in NN. This is closely related to the frequent crop straw burning activities in the region. Particulate matter emitted from biomass burning has complex chemical compositions, and its OP primarily originates from various organic compounds produced during the combustion process. The higher toxic

585 efficiency of biomass burning at the NN site may be related to multiple oxidatively active components generated from incomplete combustion. Additionally, biomass burning produces substantial amounts of PAHs, which can form quinone intermediates through metabolic activation within cells, subsequently participating in ROS generation processes (Zhu et al., 2024; Libalova et al., 2018). The biomass burning process also releases water-soluble organic carbon (WSOC), which contains humic-like substances (HULIS) with significant oxidative activity (Yan et al., 2015; Salma et al., 2010). These macromolecular
590 organic compounds contain abundant functional groups such as hydroxyl, carbonyl, and carboxyl groups, which can generate reactive oxygen species like hydroxyl radicals and hydrogen peroxide through photochemical reactions and metal-catalyzed reactions (Verma et al., 2015; Lin and Yu, 2011). The biomass burning at LFS sites, despite contributing to PM₁₀, lacks oxidative activity, which may be related to different combustion conditions or degrees of combustion completeness leading to reduced generation of oxidatively active components. Differences in combustion temperature and oxygen supply conditions
595 significantly affect the formation and transformation of organic compounds, under high-temperature complete combustion conditions, most organic compounds are oxidatively decomposed, thereby reducing the oxidative activity of particulate matter (Tuet et al., 2017).

Secondary aerosols

Secondary aerosols exhibited distinctly different patterns at different sites. At the LFS and GC sites, OP_m values of secondary
600 aerosols are 0.005 and 0.014 nmol H₂O₂·μg⁻¹, respectively, while the OP contributions of secondary aerosols were completely absent at NN and ZZ sites. This spatial heterogeneity may be related to the formation mechanisms, aging degree, and precursor composition of secondary aerosols. The photochemical oxidation processes of volatile organic compounds (VOCs) in the atmosphere play an important role in the oxidative activity of secondary aerosols (Kong et al., 2023). In atmospheric photochemical reactions, precursors such as benzene compounds and terpenes undergo complex chemical transformations
605 under the action of oxidants like ·OH and ozone, forming products with strong oxidative activity (Chen et al., 2022). These reaction products include various oxygen-containing organic compounds, such as aldehydes, ketones, carboxylic acids, and peroxides, which can participate in ROS generation through multiple pathways like photochemical reactions and Fenton reactions (Wei et al., 2022; Lin and Yu, 2011). The absence of secondary aerosol OP contributions at NN and ZZ sites may be related to multiple factors. First, the concentrations and compositions of VOC precursors at these two sites may not be favorable
610 for forming secondary organic aerosols with high oxidative activity, or the concentrations of atmospheric oxidants may be low, limiting the generation of oxidatively active products (Chen et al., 2022; Kalbande et al., 2022). Additionally, differences in meteorological conditions such as relative humidity, temperature, and solar radiation intensity significantly affect the formation and aging processes of secondary aerosols, and unfavorable meteorological conditions may lead to the formation of products with weaker oxidative activity (Xu et al., 2017; Wu et al., 2022).

615 Dust

The minerals contained in dust can participate in the ROS generation process (Nishita-Hara et al., 2019; Lodovici and Bigagli, 2011). The OP_m values of dust at NN, LFS, and GC, are 0.004, 0.009, and 0.016 $nmol\ H_2O_2\cdot\mu g^{-1}$, respectively. The dust source at the GC site exhibited the highest toxic efficiency, whereas the NN site showed relatively lower OP_m indicating weaker intrinsic toxicity of local dust. Notably, the ZZ site demonstrated zero OP contribution from dust sources, which may be attributed to distinct chemical characteristics of dust particles resulting from regional soil composition, geological background, or urbanization level. The oxidative activity of dust particles is primarily associated with their complex mineral composition and surface chemical properties. Due to varying geological backgrounds, dust from different regions exhibits significant differences in mineral composition, consequently affecting its OP. Quartz particles demonstrate certain oxidative activity through surface-catalyzed reactions that generate hydroxyl radicals, facilitated by surface silanol groups and defect sites (Konecny et al., 2001). Meteorological conditions also significantly influence the oxidative activity of dust (Joshi et al., 2017; Ma et al., 2023). The NN site is located in southern China, where higher precipitation levels compared to northern regions may contribute to reduced oxidative reactivity.

Coal combustion

PM emitted from coal combustion may contain numerous TMs and PAHs, which can promote ROS generation through pathways such as the Fenton reaction, thereby enhancing the OP of the particles (Pardo et al., 2020). Notably, coal combustion sources exhibit significant differences in OP_m contributions at the ZZ and GC sites (0.008 and 0.017 $nmol\ H_2O_2\cdot\mu g^{-1}$, respectively). This disparity in mass-specific toxicity may reflect variations in coal types or pollution control technologies across different regions, ultimately influencing the chemical composition and toxicological characteristics of the emitted particles. The organic carbon fraction of coal combustion emissions contains substantial amounts of oxygenated organic compounds, such as aldehydes and ketones, which can directly participate in oxidative stress reactions or act as precursors for generating stronger oxidants (Wang et al., 2023b). Different coal types produce particles with distinct oxidative properties. For instance, lignite combustion, due to its higher volatile matter content, tends to generate more organic compounds, resulting in relatively higher oxidative activity (Martens et al., 2021). Combustion technologies and pollution control measures also significantly influence the OP of coal-derived particles. Modern coal-fired power plants equipped with desulfurization (FGD), denitrification (SCR/SNCR), and particulate removal systems can effectively reduce certain redox-active components, thereby lowering the toxic efficiency of the emitted particles (Tao et al., 2020; Asif et al., 2022). The observed differences in oxidative activity between the ZZ and GC sites may be closely linked to local coal quality, combustion facility technologies, and the implementation level of environmental protection measures.

Agricultural activity

The OP of agricultural activities was only detected at the ZZ and GC sites, with OP_m of 0.012 and 0.004 $nmol\ H_2O_2\cdot\mu g^{-1}$, respectively, while no OP contribution from agricultural activities was observed at the LFS site. The source of agricultural

activities at ZZ exhibited relatively higher toxic efficiency, indicating that agricultural-derived particulate matter in this region possesses stronger oxidative activity. In contrast, the toxic efficiency of agricultural sources at GC was comparatively weaker. This regional disparity may be attributed to variations in agricultural practices, types of fertilizers or pesticides used, and agricultural waste management methods across different areas, leading to significant differences in the chemical composition and oxidative activity of agricultural-derived particles. NO_3^- formed during fertilizer application can influence particle oxidative properties through ionic strength effects and acidification processes (Lodovici and Bigagli, 2011). Pesticide usage constitutes another critical source of agricultural OP. Degradation products of organophosphorus pesticides often demonstrate strong oxidative activity (Lukaszewicz-Hussain, 2010). Fine particles generated during pesticide spraying may carry these reactive components into the atmosphere, enhancing the overall OP of agricultural-derived particulate matter. The absence of agricultural OP at LFS could be related to the intensity and type of farming practices in this northern urban area. The surrounding agriculture likely adopts extensive farming methods with lower fertilizer and pesticide application rates, resulting in insufficient concentrations of oxidative components in agricultural-sourced particles.

Sea salt

Although NN is located inland, it is influenced by air masses originating from the South China Sea, enabling long-range transport of sea salt aerosols that affect local atmospheric OP. Halogen compounds (e.g., Cl^- , Br^-) in sea salt can catalyze the generation of radicals such as $\cdot\text{OH}$ and $\text{Cl}\cdot$, thereby participating in atmospheric oxidation processes (Cao et al., 2024; Knipping et al., 2000). At the NN site, the OP_m of sea salt aerosols was measured at $0.016 \text{ nmol H}_2\text{O}_2 \cdot \mu\text{g}^{-1}$.

Industry

The PMF results revealed that industrial emissions at ZZ were dominated by Cl^- and K^+ with negligible contributions from OC and EC. OC typically serves as the primary contributor to particle oxidative activity through redox-active organic species, including quinones and phenolic compounds that can participate in electron transfer reactions and generate reactive oxygen species (Libalova et al., 2018; Jiang and Jang, 2018). The absence of organic carbon compounds provides a mechanistic explanation for the zero OP observed in this source profile.

4 Conclusions

This study utilized a comprehensive approach to analyze the characteristics and sources of PM_{10} and its OP at 12 representative sites in China. The main findings are summarized as follows:

1. Performance of CNN-LSTM deep learning model

675 The CNN-LSTM deep learning model exhibited robust performance in reconstructing missing data for PM₁₀ mass concentrations and outliers in chemical components. The model achieved R² values of 0.9670 and 0.8840 for the training and testing sets, respectively. These results highlight the potential of the model to address missing data issues in PM₁₀ research.

2. Spatiotemporal variations in PM₁₀ and OP levels

PM₁₀ and OP concentrations showed remarkable spatial and temporal variations:

- PM₁₀ concentrations were relatively higher in XA and DH in the northwestern region, while lower in LFS and DL in the 680 northeastern region.
- Suburban sites generally exhibited higher PM₁₀ concentrations compared to other site types.
- OP levels were relatively higher in CD and GC, with urban sites having higher OP_v values than other sites.
- Annual average PM₁₀ concentrations in northern regions were typically higher than in southern regions.
- Seasonally, PM₁₀ and OP levels were higher in winter and lower in summer, suggesting the potential benefits of implementing 685 targeted control measures during high-risk periods to mitigate adverse health impacts.

3. Source Apportionment Findings

Source apportionment using PMF indicated that dust, biomass burning, traffic emissions, and agricultural activities were likely the main contributing sources to PM₁₀ mass concentrations at the study sites. Understanding the contributions of these sources is crucial for developing more effective PM₁₀ reduction strategies.

690 4. OP Source Analysis

OP_m analysis revealed significant spatial variations in toxic efficiency across different sources. Traffic sources demonstrated consistently high OP_m (0.013-0.022 nmol H₂O₂·μg⁻¹) at four sites, indicating significant toxic efficiency of traffic-generated particulate matter. Biomass burning showed the highest OP_m at NN (0.019 nmol H₂O₂·μg⁻¹), closely related to frequent crop straw burning activities. Secondary aerosols exhibited the highest OP_m value at GC (0.014 nmol H₂O₂·μg⁻¹). Dust sources 695 showed regional differences, with highest toxic efficiency at GC (0.016 nmol H₂O₂·μg⁻¹) and zero contribution at ZZ. Coal combustion sources exhibited significant differences between ZZ and GC (0.008 and 0.017 H₂O₂·μg⁻¹), reflecting variations in coal types and pollution control technologies. Sea salt aerosols at NN demonstrated OP_m of 0.016 nmol H₂O₂·μg⁻¹ through long-range transport from coastal regions. Industrial emissions at ZZ showed zero OP due to the absence of organic carbon compounds in the source profile.

700 The study results underscore the importance of identifying and quantifying OP sources to assess and mitigate health risks associated with PM₁₀ exposure. The source apportionment findings suggest that emission reduction measures targeting traffic, biomass burning, dust, agricultural activities, and coal combustion may help lower OP levels and protect public health. This research employed deep learning techniques to analyze the spatiotemporal distribution characteristics, source apportionment, and influencing factors of PM₁₀ and its OP in different typical regions of China from multiple perspectives. The findings 705 provide a scientific basis for better understanding the causes of PM₁₀ pollution, formulating control strategies, and mitigating health risks. Future studies should focus on further investigating the identification and health risk assessment of toxic and

harmful components in PM₁₀, exploring the toxicological mechanisms of OP, and developing integrated indicators that combine chemical components and toxicity for characterizing and evaluating PM₁₀ pollution.

Code availability. The software code is available upon request.

710 **Data availability.** The data used in this study are available on the Zenodo data repository platform at <https://doi.org/10.5281/zenodo.15420768>.

Author contribution. Q.C.: Software, Writing – original draft preparation. D.F. and J.J.: Data curation, Conceptualization, Writing – review & editing. X. H.: Management of OP measurements. Y.C.: Methodology, Investigation. T.Z. and Y.B.: Software, Methodology. Y. Z.: Conceptualization, Writing – review & editing.

715 **Competing interests.** The authors declare that they have no conflict of interest.

Disclaimer. Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Financial support. This work was supported by the National Natural Science Foundation of China (No. 21677145), the Fundamental Research Funds for the Central Universities (No. E0E48927X2) and the Foundation of CMA Meteorological
720 Observation Center (#MOCQN202409).

Acknowledgments: The authors are grateful to the study participants and field staff in the twelve stations of the CMA Atmosphere Watch Network (CAWNET).

References

- 725 An, J. Q., Liu, H. B., Wang, X. M., Talifu, D., Abulizi, A., Maihemuti, M., Li, K. J., Bai, H. F., Luo, P., and Xie, X. X.: Oxidative potential of size-segregated particulate matter in the dust-storm impacted Hotan, northwest China, *Atmos. Environ.*, 280, <https://doi.org/10.1016/j.atmosenv.2022.119142>, 2022.
- Antonini, J. M., Clarke, R. W., Murthy, G. G. K., Sreekanthan, P., Jenkins, N., Eagar, T. W., and Brain, J. D.: Freshly generated stainless steel welding fume induces greater lung inflammation in rats as compared to aged fume, *Toxicol. Lett.*, 98, 77-86, 730 [https://doi.org/10.1016/s0378-4274\(98\)00103-9](https://doi.org/10.1016/s0378-4274(98)00103-9), 1998.
- Asif, Z., Chen, Z., Wang, H., and Zhu, Y.: Update on air pollution control strategies for coal-fired power plants, *Clean Technol. Environ. Policy*, 24, 2329-2347, <https://doi.org/10.1007/s10098-022-02328-8>, 2022.
- Bates, J. T., Fang, T., Verma, V., Zeng, L. H., Weber, R. J., Tolbert, P. E., Abrams, J. Y., Sarnat, S. E., Klein, M., Mulholland, J. A., and Russell, A. G.: Review of Acellular Assays of Ambient Particulate Matter Oxidative Potential: Methods and 735 Relationships with Composition, Sources, and Health Effects, *Environ. Sci. Technol.*, 53, 4003-4019, <https://doi.org/10.1021/acs.est.8b03430>, 2019.
- Borlaza, L. J., Weber, S., Marsal, A., Uzu, G., Jacob, V., Besombes, J. L., Chatain, M., Conil, S., and Jaffrezo, J. L.: Nine-year trends of PM₁₀ sources and oxidative potential in a rural background site in France, *Atmos. Chem. Phys.*, 22, 8701-8723, <https://doi.org/10.5194/acp-22-8701-2022>, 2022.
- 740 Cao, L. M., Zhou, Y., Zhang, Z., Sun, W. W., and Chen, W. H.: Impacts of airborne particulate matter and its components on respiratory system health, *Chinese journal of preventive medicine*, 50, 1114-1118, <http://dx.doi.org/10.3760/cma.j.issn.0253-9624.2016.12.018>, 2016.
- Cao, P., Lu, C., and Yu, Z.: Historical nitrogen fertilizer use in agricultural ecosystems of the contiguous United States during 1850–2015: application rate, timing, and fertilizer types, *Earth Syst. Sci. Data*, 10, 969-984, [https://doi.org/10.5194/essd-10-](https://doi.org/10.5194/essd-10-969-2018)
- 745 969-2018, 2018.
- Cao, Y. Q., Wang, Z., Liu, J. R., Ma, Q. X., Li, S. Y., Liu, J., Li, H., Zhang, P., Chen, T. Z., Wang, Y. H., Chu, B. W., Zhang, X. H., Saiz-Lopez, A., Francisco, J. S., and He, H.: Spontaneous Molecular Bromine Production in Sea-Salt Aerosols, *Angewandte Chemie-International Edition*, 63, <https://doi.org/10.1002/anie.202409779>, 2024.
- Chen, T., Zhang, P., Chu, B., Ma, Q., Ge, Y., Liu, J., and He, H.: Secondary organic aerosol formation from mixed volatile 750 organic compounds: Effect of RO₂ chemistry and precursor concentration, *npj Clim. Atmos. Sci.*, 5, 95, <https://doi.org/10.1038/s41612-022-00321-y>, 2022.
- Chirino, Y. I., Sánchez-Pérez, Y., Osornio-Vargas, á. R., Morales-Bárcenas, R., Gutiérrez-Ruíz, M. C., Segura-García, Y., Rosas, I., Pedraza-Chaverri, J., and Garcia-Cuellar, C. M.: PM₁₀ impairs the antioxidant defense system and exacerbates oxidative stress driven cell death, *Toxicol. Lett.*, 193, 209-216, <http://doi.org/10.1016/j.toxlet.2010.01.009>, 2010.
- 755 Cohen, A. J., Ross Anderson, H., Ostro, B., Pandey, K. D., Krzyzanowski, M., Künzli, N., Gutschmidt, K., Pope, A., Romieu, I., Samet, J. M., Smith, K.: The global burden of disease due to outdoor air pollution, *Journal of Toxicology and Environmental Health Part A Current Issues*, 68, 1301-1307, <http://doi.org/10.1080/15287390590936166>, 2005.

- Cohn, C. A., Simon, S. R., and Schoonen, M. A. A.: Comparison of fluorescence-based techniques for the quantification of particle-induced hydroxyl radicals, *Part. Fibre Toxicol.*, 5, <http://doi.org/10.1186/1743-8977-5-2>, 2008.
- 760 Dragomir, E. G. and Oprea, M.: Air Quality Forecasting by Using Nonlinear Modeling Methods, *Nonlinear Dynamics of Electronic Systems*, 438, 387-394, 2014.
- Esmailirad, S., Lai, A., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., Uzu, G., Daellenbach, K., Canonaco, F., Hassankhany, H., Arhami, M., Baltensperger, U., Prévôt, A. S. H., Schauer, J. J., Jaffrezo, J. L., Hosseini, V., and El Haddad, I.: Source apportionment of fine particulate matter in a Middle Eastern Metropolis, Tehran-Iran, using PMF with organic and
- 765 inorganic markers, *Sci. Total Environ.*, 705, <https://doi.org/10.1016/j.scitotenv.2019.135330>, 2020.
- Gao, D., Ripley, S., Weichenthal, S., and Pollitt, K. J. G.: Ambient particulate matter oxidative potential: Chemical determinants, associated health effects, and strategies for risk management, *Free Radical Biology and Medicine*, 151, 7-25, <http://dx.doi.org/10.1016/j.freeradbiomed.2020.04.028>, 2020.
- Guo, H. B., Jin, L., and Huang, S. J.: Effect of PM characterization on PM oxidative potential by acellular assays: a review,
- 770 *Rev. Environ. Health*, 35, 461-470, <https://doi.org/10.1515/reveh-2020-0003>, 2020.
- Ham, J., Lee, H. J., Cha, J. W., and Ryoo, S. B.: Potential Source of PM₁₀, PM_{2.5}, and OC and EC in Seoul During Spring 2016, *Atmosphere-Korea*, 27, 41-54, <http://dx.doi.org/10.14191/Atmos.2017.27.1.041>, 2017.
- Han, L. J., Zhou, W. Q., and Li, W. F.: Increasing impact of urban fine particles (PM_{2.5}) on areas surrounding Chinese cities, *Scientific Reports*, 5, <http://dx.doi.org/10.1038/srep12467>, 2015.
- 775 He, L. C. and Zhang, J. F.: Particulate matter (PM) oxidative potential: Measurement methods and links to PM physicochemical characteristics and health effects, *Crit. Rev. Environ. Sci. Technol.*, 53, 177-197, <https://doi.org/10.1080/10643389.2022.2050148>, 2023.
- Hedayat, F., Stevanovic, S., Milic, A., Miljevic, B., Nabi, M. N., Zare, A., Bottle, S. E., Brown, R. J., and Ristovski, Z. D.: Influence of oxygen content of the certain types of biodiesels on particulate oxidative potential, *Sci. Total Environ.*, 545, 381-
- 780 388, <http://dx.doi.org/10.1016/j.scitotenv.2015.12.036>, 2016.
- Hochreiter, S. and Schmidhuber, J.: Long Short-term Memory, *Neural Comput.*, 9, 1735-1780, <https://doi.org/10.1162/neco.1997.9.8.1735>, 1997.
- Huang, C. J. and Kuo, P. H.: A Deep CNN-LSTM Model for Particulate Matter (PM_{2.5}) Forecasting in Smart Cities, *Sensors*, 18, <https://doi.org/10.3390/s18072220>, 2018.
- 785 Huang, W., Zhang, Y., Zhang, Y., Fang, D., and Schauer, J. J.: Optimization of the Measurement of Particle-Bound Reactive Oxygen Species with 2',7'-dichlorofluorescein (DCFH), *Water, Air, & Soil Pollution*, 227, 164, <https://doi.org/10.1007/s11270-016-2860-9>, 2016.
- Huang, X.: The Impact of PM₁₀ and Other Airborne Particulate Matter on the Cardiopulmonary and Respiratory Systems of Sports Personnel under Atmospheric Exposure, *Atmosphere*, 14, <https://doi.org/10.3390/atmos14111697>, 2023.

- 790 Huo, H. X., Gao, Y. T., Sun, L., Gao, Y., Gao, H. W., and Yao, X. H.: Investigating Dual Character of Atmospheric Ammonia on Particulate NH_4NO_3 : Reducing Evaporation Versus Promoting Formation, *Atmosphere*, 16, <https://doi.org/10.3390/atmos16060685>, 2025.
- Jiang, H. and Jang, M.: Dynamic Oxidative Potential of Atmospheric Organic Aerosol under Ambient Sunlight, *Environ. Sci. Technol.*, 52, 7496-7504, <https://doi.org/10.1021/acs.est.8b00148>, 2018.
- 795 Jiao, J., Jia, X. F., Yan, P., Cao, F., Fang, D. Q., Ma, Q. L., Yu, D. J., Chu, J. H.: Chemical characterization of PM_{10} in the east-central background region of China, 2016-2017. *Journal of Applied Meteorological Science*, 32, 65-77, <https://doi.org/10.11898/1001-7313.20210106>, 2021.
- Joshi, N., Romanias, M. N., Riffault, V., and Thevenet, F.: Investigating water adsorption onto natural mineral dust particles: Linking DRIFTS experiments and BET theory, *Aeolian Research*, 27, 35-45, <https://doi.org/10.1016/j.aeolia.2017.06.001>,
800 2017.
- Jung, C., Huang, C., Su, H., Chen, N., and Yeh, C.: Impact of agricultural activity on $\text{PM}_{2.5}$ and its compositions in elementary schools near corn and rice farms, *Sci. Total Environ.*, 906, 167496, <https://doi.org/10.1016/j.scitotenv.2023.167496>, 2024.
- Kalbande, R., Yadav, R., Maji, S., Rathore, D. S., and Beig, G.: Characteristics of VOCs and their contribution to O_3 and SOA formation across seasons over a metropolitan region in India, *Atmos. Pollut. Res.*, 13, 101515, <https://doi.org/10.1016/j.apr.2022.101515>, 2022.
805
- Knipping, E. M., Lakin, M. J., Foster, K. L., Jungwirth, P., Tobias, D. J., Gerber, R. B., Dabdub, D., and Finlayson-Pitts, B. J.: Experiments and simulations of ion-enhanced interfacial chemistry on aqueous NaCl aerosols, *Science*, 288, 301-306, <https://doi.org/10.1126/science.288.5464.301>, 2000.
- Konecny, R., Leonard, S., Shi, X., Robinson, V., and Castranova, V.: Reactivity of free radicals on hydroxylated quartz surface and its implications for pathogenicity experimental and quantum mechanical study, *Journal of environmental pathology, toxicology and oncology : official organ of the International Society for Environmental Toxicology and Cancer*, 20 Suppl 1, 119-132, <https://doi.org/10.1615/JEnvironPatholToxicolOncol.v20.iSuppl.1.110>, 2001.
810
- Kong, L., Zhou, L., Chen, D., Luo, L., Xiao, K., Chen, Y., Liu, H., Tan, Q., and Yang, F.: Atmospheric oxidation capacity and secondary pollutant formation potentials based on photochemical loss of VOCs in a megacity of the Sichuan Basin, China, *Sci. Total Environ.*, 901, 166259, <https://doi.org/10.1016/j.scitotenv.2023.166259>, 2023.
815
- Lee, S., Shin, J.: Hybrid Model of Convolutional LSTM and CNN to Predict Particulate Matter, *International Journal of Information and Electronics Engineering*, <https://doi.org/10.18178/IJIEE.2019.9.1.701>, 2019.
- Li, H., Zhang, Q., Zhang, Q., Chen, C., Wang, L., Wei, Z., Zhou, S., Parworth, C., Zheng, B., Canonaco, F. J.: Wintertime aerosol chemistry and haze evolution in an extremely polluted city of the North China Plain: significant contribution from coal and biomass combustion, *Atmos. Chem. Phys.*, 17, 1-31, <https://doi.org/10.5194/acp-17-4751-2017>, 2017.
820
- Li, N., Hao, M., Phalen, R. F., Hinds, W. C., and Nel, A. E.: Particulate air pollutants and asthma: A paradigm for the role of oxidative stress in PM-induced adverse health effects, *Clin. Immunol.*, 109, 250, <https://doi.org/10.1016/j.clim.2003.08.006>, 2003.

- Li, Q., Wyatt, A., and Kamens, R. M.: Oxidant generation and toxicity enhancement of aged-diesel exhaust, *Atmos. Environ.*, 43, 1037-1042, <https://doi.org/10.1016/j.atmosenv.2008.11.018>, 2009.
- Li, T., Hua, M., and Wu, X.: A Hybrid CNN-LSTM Model for Forecasting Particulate Matter (PM_{2.5}), *Ieee Access*, 8, 26933-26940, <https://doi.org/10.1109/access.2020.2971348>, 2020.
- Li, W., Wang, C., Wang, H. Q. J., Chen, J. W., Yuan, C. Y., Li, T. C., Wang, W. T., Shen, H. Z., Huang, Y., Wang, R., Wang, B., Zhang, Y. Y., Chen, H., Chen, Y. C., Tang, J. H., Wang, X. L., Liu, J. F., Coveney, R. M., and Tao, S.: Distribution of atmospheric particulate matter (PM) in rural field, rural village and urban areas of northern China, *Environ. Pollut.*, 185, 134-140, <https://doi.org/10.1016/j.envpol.2013.10.042>, 2014.
- Li, X., Song, H., Zhai, S., Lu, S., Kong, Y., Xia, H., and Zhao, H.: Particulate matter pollution in Chinese cities: Areal-temporal variations and their relationships with meteorological conditions (2015-2017), *Environ. Pollut.*, 246, <https://doi.org/10.1016/j.envpol.2018.11.103>, 2019.
- Libalova, H., Milcova, A., Cervena, T., Vrbova, K., Rossnerova, A., Novakova, Z., Topinka, J., and Rossner, P.: Kinetics of ROS generation induced by polycyclic aromatic hydrocarbons and organic extracts from ambient air particulate matter in model human lung cell lines, *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 827, 50-58, <https://doi.org/10.1016/j.mrgentox.2018.01.006>, 2018.
- Lin, P. and Yu, J. Z.: Generation of Reactive Oxygen Species Mediated by Humic-like Substances in Atmospheric Aerosols, *Environ. Sci. Technol.*, 45, 10362-10368, <https://doi.org/10.1021/es2028229>, 2011.
- Liu, H., Zheng, J., Qu, C., Zhang, J., Wang, Y., Zhan, C., Yao, R., and Cao, J.: Characteristics and Source Analysis of Water-Soluble Inorganic Ions in PM₁₀ in a Typical Mining City, Central China, *Atmosphere*, 8, <https://doi.org/10.3390/atmos8040074>, 2017a.
- Liu, P., Zhang, C., Mu, Y., Liu, C., Xue, C., Ye, C., Liu, J., Zhang, Y., Zhang, H.: The possible contribution of the periodic emissions from farmers' activities in the North China Plain to atmospheric water-soluble ions in Beijing, *Atmos. Chem. Phys.*, 16, <https://doi.org/10.5194/acp-16-10097-2016>, 2016.
- Liu, P., Zhang, C., Xue, C., Mu, Y., Liu, J., Zhang, Y., Tian, D., Ye, C., Zhang, H., Guan, J.: The contribution of residential coal combustion to atmospheric PM_{2.5} in northern China during winter, *Atmos. Chem. Phys.*, 17, 1-37, <https://doi.org/10.5194/acp-17-11503-2017>, 2017b.
- Liu, Y., Zhang, W., Yang, W., Bai, Z., and Zhao, X.: Chemical Compositions of PM_{2.5} Emitted from Diesel Trucks and Construction Equipment, *Aerosol Science and Engineering*, 2, 51-60, <https://doi.org/10.1007/s41810-017-0020-2>, 2018.
- Lodovici, M. and Bigagli, E.: Oxidative Stress and Air Pollution Exposure, *Journal of Toxicology*, 2011, <https://doi.org/10.1155/2011/487074>, 2011.
- Lukaszewicz-Hussain, A.: Role of oxidative stress in organophosphate insecticide toxicity – Short review, *Pestic. Biochem. Physiol.*, 98, 145-150, <https://doi.org/10.1016/j.pestbp.2010.07.006>, 2010.

- Ma, S., Cheng, D., Tang, Y., Fan, Y., Li, Q., He, C., Zhao, Z., and Xu, T.: Investigation of oxidative potential of fresh and O₃-aging PM_{2.5} from various emission sources across urban and rural regions, *J. Environ. Sci.*, 151, 608-615, <https://doi.org/10.1016/j.jes.2024.04.023>, 2025.
- Ma, T., Niuhe, J., Lu, S., Zhang, L., Zhou, S., Liu, J., Zhang, W., Liu, X., Ebere, E. C., Wang, Q., and Wang, W.: Comparison of the heterogeneous reaction of NO₂ on the surface of clay minerals and desert dust particles, *Environ. Pollut.*, 334, 122134, <https://doi.org/10.1016/j.envpol.2023.122134>, 2023.
- Martens, P., Czech, H., Tissari, J., Ihalainen, M., Suhonen, H., Sklorz, M., Jokiniemi, J., Sippula, O., and Zimmermann, R.: Emissions of Gases and Volatile Organic Compounds from Residential Heating: A Comparison of Brown Coal Briquettes and Logwood Combustion, *Energy & Fuels*, 35, 14010-14022, <https://doi.org/10.1021/acs.energyfuels.1c01667>, 2021.
- McKenzie, E. R., Money, J. E., Green, P. G., and Young, T. M.: Metals associated with stormwater-relevant brake and tire samples, *Sci. Total Environ.*, 407, 5855-5860, <https://doi.org/10.1016/j.scitotenv.2009.07.018>, 2009.
- Méndez, M., Merayo, M. G., and Núñez, M.: Machine learning algorithms to forecast air quality: a survey, *Artificial intelligence review*, 1-36, <https://doi.org/10.1007/s10462-023-10424-4>, 2023.
- Meng, L., Li, Y., Han, S., and Hao, T.: Influence of Sea-Land Breeze on the Concentrations of PM_{2.5} and O₃ in Tianjin City, *Research of Environmental Sciences*, 32, 390-398, <http://dx.doi.org/10.13198/j.issn.1001-6929>, 2018. 12. 10, 2019.
- Nielsen, T.: Traffic contribution of polycyclic aromatic hydrocarbons in the center of a large city, *Atmos. Environ.*, 30, 3481-3490, [https://doi.org/10.1016/1352-2310\(96\)00096-9](https://doi.org/10.1016/1352-2310(96)00096-9), 1996.
- Nishita-Hara, C., Hirabayashi, M., Hara, K., Yamazaki, A., and Hayashi, M.: Dithiothreitol-Measured Oxidative Potential of Size-Segregated Particulate Matter in Fukuoka, Japan: Effects of Asian Dust Events, *Geohealth*, 3, 160-173, <https://doi.org/10.1029/2019gh000189>, 2019.
- Okut, H.: Deep Learning for Subtyping and Prediction of Diseases: Long-Short Term Memory, *Deep Learning Applications*, <https://doi.org/10.5772/intechopen.96180>, 2021.
- Paatero, P.: Least squares formulation of robust non-negative factor analysis, *Chemom. Intell. Lab. Syst.*, 37, 23-35, [https://doi.org/10.1016/s0169-7439\(96\)00044-5](https://doi.org/10.1016/s0169-7439(96)00044-5), 1997.
- Paatero, P. and Tapper, U. J. E.: Positive matrix factorization: A nonnegative factor model with optimal utilization of error estimates of data values, *Environmetrics*, 5, <https://doi.org/10.1002/env.3170050203>, 1994.
- Pardo, M., Qiu, X. H., Zimmermann, R., and Rudich, Y.: Particulate Matter Toxicity Is Nrf₂ and Mitochondria Dependent: The Roles of Metals and Polycyclic Aromatic Hydrocarbons, *Chem. Res. Toxicol.*, 33, 1110-1120, <https://doi.org/10.1021/acs.chemrestox.0c00007>, 2020.
- Park, I. S., Kim, H. K., Song, C. K., Jang, Y. W., Kim, S. H., Cho, C. R., Owen, J. S., Kim, C. H., Chung, K. W., and Park, M. S.: Meteorological Characteristics and Assessment of the Effect of Local Emissions during High PM₁₀ Concentration in the Seoul Metropolitan Area, *Asian J. Atmos. Environ.*, 13, 117-135, <https://doi.org/10.5572/ajae.2019.13.2.117>, 2019.
- Qin, D. M., Yu, J., Zou, G. J., Yong, R. H., Zhao, Q., and Zhang, B.: A Novel Combined Prediction Scheme Based on CNN and LSTM for Urban PM_{2.5} Concentration, *Ieee Access*, 7, 20050-20059, <https://doi.org/10.1109/access.2019.2897028>, 2019.

- 890 Qu, W. J., Arimoto, R., Zhang, X. Y., Zhao, C. H., Wang, Y. Q., Sheng, L. F., Fu, G.: Spatial distribution and interannual variation of surface PM₁₀ concentrations over eighty-six Chinese cities, *Atmos. Chem. Phys.*, 10, 5641-5662, <https://doi.org/10.5194/acp-10-5641-2010>, 2010.
- Saarikoski, S. K., Sillanpää, M. K., Saarnio, K. M., Hillamo, R. E., Pennanen, A. S., and Salonen, R. O.: Impact of biomass combustion on urban fine particulate matter in Central and Northern Europe, *Water, Air, Soil Pollut.*, 191, 265-277, <https://doi.org/10.1007/s11270-008-9623-1>, 2008.
- 895 Saffari, A., Daher, N., Shafer, M. M., Schauer, J. J., and Sioutas, C.: Global perspective on the oxidative potential of airborne particulate matter: A synthesis of research findings, *Environ. Sci. Technol.*, 48, 7576, <https://doi.org/10.1021/es500937x>, 2014.
- Saggi, G. S. and Mittal, S. K.: Source apportionment of PM₁₀ by positive matrix factorization model at a source region of biomass burning, *J. Environ. Manage.*, 266, <https://doi.org/10.1016/j.jenvman.2020.110545>, 2020.
- 900 Salma, I., Mészáros, T., Maenhaut, W., Vass, E., and Majer, Z.: Chirality and the origin of atmospheric humic-like substances, *Atmos. Chem. Phys.*, 10, 1315-1327, <https://doi.org/10.5194/acp-10-1315-2010>, 2010.
- Schwartz, S. E.: Does fossil fuel combustion lead to global warming?, *Energy*, 18, 1229-1248, [https://doi.org/10.1016/0360-5442\(93\)90012-3](https://doi.org/10.1016/0360-5442(93)90012-3), 1993.
- Seo, M., Lee, S. Y., Lee, C., and Cho, S. S.: Recycling of Cement Kiln Dust as a Raw Material for Cement, *Environments*, 6, <https://doi.org/10.3390/environments6100113>, 2019.
- 905 Sharma, S. K., Mandal, T. K., Jain, S., Saraswati, Sharma, A., and Saxena, M.: Source Apportionment of PM_{2.5} in Delhi, India Using PMF Model, *Bull. Environ. Contam. Toxicol.*, 97, 286-293, <https://doi.org/10.1007/s00128-016-1836-1>, 2016.
- Shen, Z. X., Sun, J., Cao, J. J., Zhang, L. M., Zhang, Q., Lei, Y. L., Gao, J. J., Huang, R. J., Liu, S. X., Huang, Y., Zhu, C. S., Xu, H. M., Zheng, C. L., Liu, P. P., Xue, Z. G.: Chemical profiles of urban fugitive dust PM_{2.5} samples in Northern Chinese cities, *Sci. Total Environ.* 569-570, 619-626, <https://doi.org/10.1016/j.scitotenv.2016.06.156>, 2016.
- 910 Sherstinsky, A.: Fundamentals of Recurrent Neural Network (RNN) and Long Short-Term Memory (LSTM) network, *Physica D-Nonlinear Phenomena*, 404, <https://doi.org/10.1016/j.physd.2019.132306>, 2020.
- Shi, J., Yan, J. P., Wang, S. S., Zhao, S. H., Zhang, M. M., Xu, S. Q., Lin, Q., and Yang, H.: Determinant of Sea Salt Aerosol Emission in the Southern Hemisphere in Summer Time, *Earth and Space Science*, 9, <https://doi.org/10.1029/2022ea002529>, 2022.
- 915 Simonich, S. L. and Hites, R. A.: Vegetation-Atmosphere Partitioning of Polycyclic Aromatic Hydrocarbons, *Environ. Sci. Technol.*, 28, 939-943, <https://doi.org/10.1021/es00054a028>, 1994.
- Singh, K. P., Gupta, S., Kumar, A., and Shukla, S. P.: Linear and nonlinear modeling approaches for urban air quality prediction, *Sci. Total Environ.*, 426, 244-255, <https://doi.org/10.1016/j.scitotenv.2012.03.076>, 2012.
- 920 Slanina, S. and Zhang, Y.: Aerosols: Connection between regional climate change and air quality (IUPAC Technical Report), 76, 1241-1253, <https://doi.org/10.1351/pac200476061241>, 2004.

- Sowlat, M. H., Hasheminassab, S., and Sioutas, C.: Source apportionment of ambient particle number concentrations in central Los Angeles using positive matrix factorization (PMF), *Atmos. Chem. Phys.*, 16, 4849-4866, <http://doi.org/10.5194/acp-16-4849-2016>, 2016.
- 925 Stelson, A. W. and Seinfeld, J. H.: Relative humidity and temperature dependence of the ammonium nitrate dissociation constant, *Atmos. Environ.*, 16, 983-992, [https://doi.org/10.1016/0004-6981\(82\)90184-6](https://doi.org/10.1016/0004-6981(82)90184-6), 1982.
- Stracquadanio, M., Petralia, E., Berico, M., La Torretta, T. M. G., Malaguti, A., Mircea, M., Gualtieri, M., and Ciancarella, L.: Source Apportionment and Macro Tracer: Integration of Independent Methods for Quantification of Woody Biomass Burning Contribution to PM₁₀, *Aerosol Air Qual. Res.*, 19, 711-723, <http://doi.org/10.4209/aaqr.2018.05.0186>, 2019.
- 930 Strzałkowska, E.: Morphology, chemical and mineralogical composition of magnetic fraction of coal fly ash, *Int. J. Coal Geol.*, 240, 103746, <https://doi.org/10.1016/j.coal.2021.103746>, 2021.
- Sun, X., Chen, J., Liu, L., Rosanoff, A., Xiong, X., Zhang, Y., and Pei, T.: Effects of Magnesium Fertilizer on the Forage Crude Protein Content Depend upon Available Soil Nitrogen, *J. Agric. Food Chem.*, 66, 1743-1750, <https://doi.org/10.1021/acs.jafc.7b04028>, 2018.
- 935 Tao, L., Wang, Y., Yue, C., Shen, Z., Liu, Z., Liu, Q., Lu, J., Ding, H., and Pan, W.: Synergistic Emission Reduction of Particulate Pollutants in Coal-fired Power Plants Using Ultra-low Emission Technology, *Aerosol Air Qual. Res.*, 20, 2529-2535, <https://doi.org/10.4209/aaqr.2020.01.0004>, 2020.
- Tian, Y. Z., Xiao, Z. M., Han, B., Shi, G.L., Wang, W., Hao, H. Z., Li, X., Feng, Y. C., and Zhu, T.: Seasonal Study of Primary and Secondary Sources of Carbonaceous Species in PM₁₀ from Five Northern Chinese Cities, *Aerosol Air Qual. Res.*, 13, 148-161, <https://doi.org/10.4209/aaqr.2012.01.0010>, 2013.
- 940 Tsvetanova, I., Zheleva, I., Filipova, M., and Stefanova, A.: Statistical analysis of ambient air PM₁₀ contamination during winter periods for Ruse region, Bulgaria, 13th National Congress on Theoretical and Applied Mechanics (NCTAM), Bulgarian Acad Sci, Inst Mech Campus I, Sofia, BULGARIA, Sep 06-10, <http://doi.org/10.1051/mateconf/201814501007>, 2018.
- Tuet, W. Y., Chen, Y., Xu, L., Fok, S., Gao, D., Weber, R. J., and Ng, N. L.: Chemical oxidative potential of secondary organic aerosol (SOA) generated from the photooxidation of biogenic and anthropogenic volatile organic compounds, *Atmos. Chem. Phys.*, 17, 839-853, <http://doi.org/10.5194/acp-17-839-2017>, 2017.
- Uranishi, K., Ikemori, F., Shimadera, H., Kondo, A., and Sugata, S.: Impact of field biomass burning on local pollution and long-range transport of PM_{2.5} in Northeast Asia, *Environ. Pollut.*, 244, 414-422, <http://doi.org/10.1016/j.envpol.2018.09.061>, 2019.
- 950 Valavanidis, A., Fiotakis, K., and Vlachogianni, T.: Airborne Particulate Matter and Human Health: Toxicological Assessment and Importance of Size and Composition of Particles for Oxidative Damage and Carcinogenic Mechanisms, *Journal of Environmental Science and Health Part C-Environmental Carcinogenesis & Ecotoxicology Reviews*, 26, 339-362, <https://doi.org/10.1080/10590500802494538>, 2008.
- Vennerød, C. B., Kjærø, A., and Bugge, E. S. J. A.: Long Short-term Memory RNN, <http://arxiv.org/abs/2105.06756>, 2021.

- 955 Verma, V., Fang, T., Xu, L., Peltier, R. E., Russell, A. G., Ng, N. L., and Weber, R. J.: Organic Aerosols Associated with the Generation of Reactive Oxygen Species (ROS) by Water-Soluble PM_{2.5}, *Environ. Sci. Technol.*, 49, 4646-4656, <https://doi.org/10.1021/es505577w>, 2015.
- Viana, M., Kuhlbusch, T. A., Querol, X., Alastuey, A., Harrison, R. M., Hopke, P. K., Winiwarter, W., Vallius, M., Szidat, S., and Prevot, A. S.: Source apportionment of particulate matter in Europe: A review of methods and results, *J. Aerosol Sci.*, 39, 827-849, <https://doi.org/10.1016/j.jaerosci.2008.05.007>, 2008.
- 960 Wang, J., Huang, J., Wang, L., Chen, C., Yang, D., Jin, M., Bai, C., and Song, Y.: Urban particulate matter triggers lung inflammation via the ROS-MAPK-NF- κ B signaling pathway, *J. Thorac. Dis.*, 9, 4398-4412, <https://doi.org/10.21037/jtd.2017.09.135>, 2017.
- Wang, M., Kong, W., Marten, R., He, X.-C., Chen, D., Pfeifer, J., Heitto, A., Kontkanen, J., Dada, L., Kürten, A., Yli-Juuti, T., Manninen, H. E., Amanatidis, S., Amorim, A., Baalbaki, R., Baccarini, A., Bell, D. M., Bertozzi, B., Bräkling, S., Brilke, S., Murillo, L. C., Chiu, R., Chu, B., De Menezes, L.-P., Duplissy, J., Finkenzeller, H., Carracedo, L. G., Granzin, M., Guida, R., Hansel, A., Hofbauer, V., Krechmer, J., Lehtipalo, K., Lamkaddam, H., Lampimäki, M., Lee, C. P., Makhmutov, V., Marie, G., Mathot, S., Mauldin, R. L., Mentler, B., Müller, T., Onnela, A., Partoll, E., Petäjä, T., Philippov, M., Pospisilova, V., Ranjithkumar, A., Rissanen, M., Rörup, B., Scholz, W., Shen, J., Simon, M., Sipilä, M., Steiner, G., Stolzenburg, D., Tham, Y. J., Tomé, A., Wagner, A. C., Wang, D. S., Wang, Y., Weber, S. K., Winkler, P. M., Wlasits, P. J., Wu, Y., Xiao, M., Ye, Q., Zauner-Wieczorek, M., Zhou, X., Volkamer, R., Riipinen, I., Dommen, J., Curtius, J., Baltensperger, U., Kulmala, M., Worsnop, D. R., Kirkby, J., Seinfeld, J. H., El-Haddad, I., Flagan, R. C., and Donahue, N. M.: Rapid growth of new atmospheric particles by nitric acid and ammonia condensation, *Nature*, 581, 184-189, <https://doi.org/10.1038/s41586-020-2270-4>, 2020.
- 975 Wang, N. L., Yao, T. D., Thompson, L. G., and Davis, M. E.: Indian monsoon and North Atlantic Oscillation signals reflected by Cl⁻ and Na⁺ in a shallow ice core from Dasuopu Glacier, Xixabangma, Himalaya, *Ann. Glaciol.*, 273-277, <https://doi.org/10.3189/172756402781816825>, 2002.
- Wang, Q. Z., Ding, H., Yu, F. W., Chao, N., Li, Y., Jiang, Q. Q., Huang, Y., Duan, L., Ji, Z. Q., Zhou, R., Yang, Z. P., Zheng, K. Y., and Miao, X. P.: The Characteristics and Impact Factors of Sulfate and Nitrate in Urban PM_{2.5} over Typical Cities of Hangzhou Bay Area, China, *Atmosphere*, 14, <https://doi.org/10.3390/atmos14121799>, 2023a.
- 980 Wang, S. S., Nan, J. L., Shi, C. Z., Fu, Q. Y., Gao, S., Wang, D. F., Cui, H. X., Saiz-Lopez, A., and Zhou, B.: Atmospheric ammonia and its impacts on regional air quality over the megacity of Shanghai, China, *Sci. Rep.* 5, <https://doi.org/10.1038/srep15842>, 2015.
- Wang, T., Huang, R.-J., Yang, L., Dai, W., Ni, H., Gong, Y., Guo, J., Zhong, H., Lin, C., and Xu, W.: Direct emissions of particulate glyoxal and methylglyoxal from biomass burning and coal combustion, *Sci. Total Environ.*, 862, 160757, <https://doi.org/10.1016/j.scitotenv.2022.160757>, 2023b.

- Weber, S., Uzu, G., Calas, A., Chevrier, F., Besombes, J. L., Charron, A., Salameh, D., Jezek, I., Mocnik, G., and Jaffrezo, J. L.: An apportionment method for the oxidative potential of atmospheric particulate matter sources: application to a one-year study in Chamonix, France, *Atmos. Chem. Phys.*, 18, 9617-9629, <https://doi.org/10.5194/acp-18-9617-2018>, 2018.
- 990 Wei, J., Fang, T., and Shiraiwa, M.: Effects of Acidity on Reactive Oxygen Species Formation from Secondary Organic Aerosols, *ACS Environ. Au*, 2, 336-345, <https://doi.org/10.1021/acsenvironau.2c00018>, 2022.
- Wu, Y., Liu, D., Tian, P., Sheng, J., Liu, Q., Li, R., Hu, K., Jiang, X., Li, S., Bi, K., Zhao, D., Huang, M., Ding, D., and Wang, J.: Tracing the Formation of Secondary Aerosols Influenced by Solar Radiation and Relative Humidity in Suburban Environment, *J. Geophys. Res.:Atmos.*, 127, e2022JD036913, <https://doi.org/10.1029/2022JD036913>, 2022.
- 995 Xu, W., Han, T., Du, W., Wang, Q., Chen, C., Zhao, J., Zhang, Y., Li, J., Fu, P., Wang, Z., Worsnop, D. R., and Sun, Y.: Effects of Aqueous-Phase and Photochemical Processing on Secondary Organic Aerosol Formation and Evolution in Beijing, China, *Environ. Sci. Technol.*, 51, 762-770, <https://doi.org/10.1021/acs.est.6b04498>, 2017.
- Xue, Y. H., Wu, J. H., Feng, Y. C., Dai, L., Bi, X. H., Li, X. A., Zhu, T., Tang, S. B., and Chen, M. F.: Source Characterization and Apportionment of PM₁₀ in Panzhuhua, China, *Aerosol Air Qual. Res.*, 10, 367-377, <https://doi.org/10.4209/aaqr.2010.01.0002>, 2010.
- 1000 Yan, C., Zheng, M., Sullivan, A. P., Bosch, C., Desyaterik, Y., Andersson, A., Li, X., Guo, X., Zhou, T., Gustafsson, Ö., and Collett, J. L.: Chemical characteristics and light-absorbing property of water-soluble organic carbon in Beijing: Biomass burning contributions, *Atmos. Environ.*, 121, 4-12, <https://doi.org/10.1016/j.atmosenv.2015.05.005>, 2015.
- Yang, G., Lee, H., and Lee, G.: A Hybrid Deep Learning Model to Forecast Particulate Matter Concentration Levels in Seoul, South Korea, *Atmosphere*, 11, <https://doi.org/10.3390/atmos11040348>, 2020a.
- 1005 Yang, L.: The space-time variations of PM₁₀ concentration in major cities of China during 2000-2007, *Journal of Arid Land Resources and Environment*, 23, 51-54, <https://www.cnndoi.org/Resolution/Handler?doi=10.13448/j.cnki.jalre.2009.09.017>, 2009.
- Yang, L. M., Wang, S. B., Duan, S. G., Yan, Q. S., Jiang, N., Zhang, R. Q., and Li, S. L.: Characteristics and formation mechanisms of secondary inorganic ions in PM_{2.5} during winter in a central city of China: Based on a high time resolution data, *Atmos. Res.*, 233, <https://doi.org/10.1016/j.atmosres.2019.104696>, 2020b.
- 1010 Yao, K. X., Xu, Y. H., Zheng, H., Zhang, X. J., Song, Y. X., and Guo, H. B.: Oxidative potential associated with reactive oxygen species of size-resolved particles: The important role of the specific sources, *J. Environ. Manage.*, 360, <https://doi.org/10.1016/j.jenvman.2024.121122>, 2024.
- 1015 Yin, H., Guo, L., Xie, M., Xu, Z., Zhao, D., and Shi, Z.: Source Tracking of Food Industry Wastewater Discharge into Sewers Using Marker Species, *J. Tongji Univ. Nat. Sci.*, 47, 1168-1174, <http://dx.doi.org/10.11908/j.issn.0253-374x.2019.08.013>, 2019.
- Yu, D. J., Wu, Y. L., Song, Q. L., Dai, X., Lin, W. L.: Environmental Characteristics and Its Observations at Longfengshan WMO Regional Atmospheric Background Station, *Climate Change Research Letters*, 01, 65-73, <http://dx.doi.org/10.12677/ccrl.2012.12008>, 2012.
- 1020

- Yu, H., Zhao, X. Y., Wang, J., Yin, B. H., Geng, C. M., Wang, X. H., Gu, C., Huang, L. H., Yang, W., and Bai, Z. P.: Chemical characteristics of road dust PM_{2.5} fraction in oasis cities at the margin of Tarim Basin, *J. Environ. Sci.*, 95, 217-224, <https://doi.org/10.1016/j.jes.2020.03.030>, 2020.
- Yu, Y. and Cao, J.: Chemical Fingerprints and Source Profiles of PM₁₀ and PM_{2.5} from Agricultural Soil in a Typical Polluted
1025 Region of Northwest China, *Aerosol Air Qual. Res.*, 23, 220419, <https://doi.org/10.4209/aaqr.220419>, 2023.
- Yue, D. L., Zhong, L. J., Zhang, T., Shen, J., Zhou, Y., Zeng, L. M., Dong, H. B., and Ye, S. Q.: Pollution Properties of Water-Soluble Secondary Inorganic Ions in Atmospheric PM_{2.5} in the Pearl River Delta Region, *Aerosol Air Qual. Res.*, 15, 1737-1747, <https://doi.org/10.4209/aaqr.2014.12.0333>, 2015.
- Zhang, R., Cao, J., Tang, Y., Arimoto, R., Shen, Z., Wu, F., Han, Y., Wang, G., Zhang, J., and Li, G.: Elemental profiles and
1030 signatures of fugitive dusts from Chinese deserts, *Sci. Total Environ.* 472, 1121-1129, <https://doi.org/10.1016/j.scitotenv.2013.11.011>, 2014a.
- Zhang, T. T., Shao, Y., Gong, H. Z., Li, L., and Wang, L. F.: Salt Content Distribution and Paleoclimatic Significance of the Lop Nur "Ear" Feature: Results from Analysis of EO-1 Hyperion Imagery, *Remote Sens.*, 6, 7783-7799, <https://doi.org/10.3390/rs6087783>, 2014b.
- Zhang, X. L. and Zhou, W. S.: Structural Vibration Data Anomaly Detection Based on Multiple Feature Information Using CNN-LSTM Model, *Structural Control & Health Monitoring*, 2023, <http://dx.doi.org/10.1155/2023/3906180>, 2023.
- Zhang, X. Y., Wang, Y. Q., Niu, T., Zhang, X. C., Gong, S. L., Zhang, Y. M., and Sun, J. Y.: Atmospheric aerosol compositions in China: spatial/temporal variability, chemical signature, regional haze distribution and comparisons with global aerosols, *Atmos. Chem. Phys.*, 12, 779-799, <https://doi.org/10.5194/acp-12-779-2012>, 2012.
- Zhao, C. X., Wang, Y. Q., Wang, Y. J., Zhang, H. L., and Zhao, B. Q.: Temporal and Spatial Distribution of PM_{2.5} and PM₁₀
1040 Pollution Status and the Correlation of Particulate Matters and Meteorological Factors During Winter and Spring in Beijing, *Huanjing Kexue*, 35, 418-427, <https://doi.org/10.13227/j.hjkk.2014.02.013>, 2014.
- Zhou, J., Elser, M., Huang, R. J., Krapf, M., Fröhlich, R., Bhattu, D., Stefenelli, G., Zotter, P., Bruns, E. A., Pieber, S. M., Ni, H. Y., Wang, Q. Y., Wang, Y. C., Zhou, Y. Q., Chen, C. Y., Xiao, M., Slowik, J. G., Brown, S., Cassagnes, L. E., Daellenbach, K. R., Nussbaumer, T., Geiser, M., Prevot, A. S. H., El-Haddad, I., Cao, J. J., Baltensperger, U., and Dommen, J.:
1045 Predominance of secondary organic aerosol to particle-bound reactive oxygen species activity in fine ambient aerosol, *Atmos. Chem. Phys.*, 19, 14703-14720, <http://doi.org/10.5194/acp-19-14703-2019>, 2019.
- Zhu, X., Ho, K. F., Yang, T. T., Laiman, V., Sun, J., Shen, Z., and Chuang, H. C.: Emission Factors of PAHs Components and Bioreactivity in PM_{2.5} from Biomass Burning, *Aerosol Air Qual. Res.*, 24, 230068, <http://doi.org/10.4209/aaqr.230068>, 2024.

1050