

# **Egusphere-2025-657 – Answer to reviewers**

## **REPORT#2**

This paper describes a method for adjusting operational surface fields, specifically wind speed and current speed, to minimise forecast errors in the coastal zone. Artificial neural networks trained with radar data from land stations (HFR) and satellites (SAR) adjust forcing fields. Overall, the paper reads well and contains all relevant references. However, the vast amount of acronyms limits readability to some extent. The methodology is reasonably well documented, and the verification of surface winds and currents appears sound. I appreciate the attempt to separate errors in the forcings that drive the wave model.

We thank Reviewer #2 for the positive feedback and for taking the time to review our manuscript. We have tried to address all your suggestions in this revised version, and we believe they have significantly improved the manuscript. We hope this revision meets with your approval.

Verification for "short-term wave forecasting" is missing. Perhaps this is out of scope, then the title would need to be edited accordingly. Wave verification is too brief and could benefit from a more detailed analysis, other than a case study. Time series plots presented in Figure 13-16 did not convince me of an improved forecast system. Numerical weather prediction models feature some variability in skill over a complete seasonal cycle. Training and verification for a particular season can introduce seasonal bias in the correction method.

We thank Reviewer #2 for this important remark. Our aim was to focus on the Neural Networks for correcting wind and currents, and we acknowledge that a full verification of short-term wave forecasting is indeed beyond the scope of this study.

Our testing with the IBI-WAV model was primarily intended to demonstrate the overall potential and feasibility of the methodology. We consider that a proper, comprehensive validation of the forecast skill warrants dedicated attention and a specific follow-up study.

We are aware that training and verifying data from a particular season can introduce seasonal biases. To mitigate this, we intentionally selected a verification period that was as out-of-sample as possible, which partially explains the modest relative performance of the methodology shown in the figures.

We agree with the reviewer that two isolated extreme events are not fully representative of the system's performance under all extreme conditions. This is why we have highlighted

throughout the revised text that any results and conclusions regarding extreme events should be considered as preliminary.

We are currently working on a follow-up study to test the feasibility of this methodology using more comprehensive benchmarks, including a more detailed analysis of extreme event data.

The introduction could benefit from a paragraph on the expected impact on the wave model skill, not just for the wind generation part, but also for the effect of currents on wave predictability.

We thank the Reviewer#2 for this remark. We have added the following text in the Introduction:

*Another main forcing at the coastal zone are the surface currents. Including currents in spectral wave models can reduce the errors on significant wave heights by more than 30% in some macrotidal environments, such as at the coast of Brittany (France) (Ardhuin et al., 2012). Wave-current interaction can affect (i) refraction due to currents, (ii) shoaling, and (iii) current-driven frequency shifting (Staneva et al., 2017; Cavalieri et al., 2018; Law Chune and Aouf, 2018; Bruciaferri et al., 2021; Calvino et al., 2022). While numerical wave models have demonstrated considerable skill in predicting wave conditions, the predictive skill of coastal circulation models remains comparatively lower, particularly in complex coastal regions (Fringer et al., 2019, García-León et al., 2022).*

### **Specific comments**

Referencing in text uses inconsistent punctuation and contains et al. year, et al., year and author, year.

We thank the Reviewer#2 for this remark. We have revised the whole manuscript and we have standardized citation formatting and Figure references.

Line 29-30: "Third generation spectral wave models ..., as they address wave generation and propagation". Third-generation models explicitly represent the nonlinear wave-wave interactions. Generation and propagation refer to second-generation models.

We thank the Reviewer#2 for this remark. Please find the revised version of the text:

*Third-generation spectral wave models are suitable for the regional scale because they address wave generation, propagation, and non-linear interactions (WISE group, 2007).*

Line 31-32: "three factors". What about the representation error? In the coastal zone, many processes are often overlooked, such as wetting and drying, river discharge, variable water depth, changes in beach profile, and seagrass, among others.

We thank the Reviewer #2 for this remark. Please find the revised version of the text:

*They are computationally affordable and reliable for many applications (Capet et al., 2020). However, errors tend to increase at the coast, due to the compound effect of many factors: (i) accuracy in the forcings; (ii) limitations in the physical parameterizations; (iii) overlooked physical processes (e.g. river discharge, coastal morphodynamics, wave-biota interaction, wet-and-dry), (iv) model discretization and numerical schemes, etc.*

Line 34: "(ii) reduce biases and errors in the inputs (Durrant et al 2013) ...". Complementary to Durrant et al., Zieger (2025) assessed forecast errors in an operational wave model. (<https://doi.org/10.1071/ES25010>).

We thank the Reviewer #2 for this reference. We have added Zieger (2025) in the Introduction.

Line 38-39: "Reducing biases and errors in the forcings have certain advantages...it does not require changing the physics." The parameterisations that represent the physics will naturally contain some form of bias that stems from model development. The underlying tuning parameters are bound to the frequency of model fields (daily, 6 hours, 3 hours, hourly) and the underlying statistic (i.e., cumulative, average, instantaneous). As a result, one has to change the physics kind of.

We thank the Reviewer #2 for this remark. Please find the revised version of the text:

*Reducing biases and errors in the forcings has certain advantages over other strategies, because it does not increase computational time by changing the model resolution. Lower forcing errors also bounds a source of uncertainty, assisting in the implementation of the other two strategies. Main inputs in the spectral wave models are (i) wind fields, (ii) surface ocean currents and (iii) bathymetry. At regional scale models, the bathymetry is considered as a static forcing, because the rate of change for short-term forecasting (i.e. a few days) is only significant at shallow waters. Hence, for the forecast window considered, winds and currents can be considered as the main dynamic forcings.*

Line 84: "spectacular growth" does not sound well. Perhaps "exponential growth".

We thank the reviewer#2 for this remark. We have added "exponential growth" in the revised version.

Line 90: "concrete". This word does not feel right and does not add clarity to the sentence (omit). There is a slight overuse of the word 'concrete' in the manuscript.

We thank the Reviewer#2 for this remark. All mentions to “concrete” have been removed.

Line 128: Could you please provide more specific details on the ST4 configuration? Could you state the value for BETAMAX used?

We thank the Reviewer#2 for this remark. Please find the revised version of the text:


*Wave physics within IBI-WAV are parameterized using the ST4 formulation (Ardhuin et al., 2010), accounting for energy dissipation due to wave breaking and swell decay. The MFWAM implementation of ST4 is further enhanced by incorporating a Phillips tail spectrum to accurately represent the high-frequency portion of the wave spectrum. The  $\beta_{max}$  tuning parameter, that adjusts the transfer of energy and momentum from the wind to the waves, is 1.48. This is consistent with other applications in the literature (i.e. 1.39 in Valiente et al., 2023; 1.52 in Ardhuin et al., 2010 or 1.75 in Alday et al., 2021).*

Line 186-187: "It has been discarded images" is gibberish. Please rephrase the sentence.

We thank the Reviewer#2 for this remark. We have rewritten this sentence.

Line 360-361: "So, it can be concluded that the ANN is able to predict winds that are closer to the SAR data than IFS." Would one not verify against an independent dataset (also Figure 5)? The ANN training would be well aware of the characteristics and structure of SAR data.

We thank the Reviewer#2 for this remark. Our main aim with the Wind ANN was to show that the GAN can effectively mimic the SAR data. The SAR data, when compared with an independent data source (in-situ buoy), presented better model skill (please refer to Figure 1). Note also the data scarcity in the area (575 valid time points in almost 5 years). This is one of the reasons that we required to keep SAR data from the Mediterranean and the NE Atlantic within the same training dataset. Another reason for this choice of training dataset was to obtain a homogeneous response and avoid disparate wind predictions across different basins. In current research, we are trying to add more SAR data, but further assessment is needed.



**Figure 1.** (up) Results of comparing SAR data with the in-situ buoy, and (low) comparing ECMWF-IFS with in-situ buoy. Note that the Wind ANN mimics well the SAR data, and the error metrics of the SAR are lower than ECMWF-IFS.

Line 479: "Benefits from the ANN forcing are more remarkable during extreme events. During storm Arwen, strong Northern winds at the Northern Iberian Peninsula (Figure 12a) ..." What is the rationale for showing daily-average marine wind speed to depict extreme events? The most pronounced feature in wind speed difference (Fig. 12b) is the north-west south-east shift around coastlines. In addition, wind speed verification at the GAL station (Fig. 13 and Fig. 14) indicates that surface wind speed is consistently overpredicted. At the same time, the responding wave field exhibits lower magnitudes for the duration of the storms. On the Mediterranean side, GCA station, is not able to capture the double peak on 19<sup>th</sup> and 20<sup>th</sup> January. Can you elaborate on this?

We thank the Reviewer#2 for this remark. We have added the following text in the revised version:

*Overall improvement is also suggested during the two extreme events. However, because the sample of storms is limited, the results and interpretations must be considered as preliminary. Figure 12 illustrates the expected order of magnitude of the differences between*

*the ECMWF-IFS and the Wind ANN products. These daily-averaged differences are representative of how the Wind ANN corrected the wind forcings during the peak of Storm Arwen. Furthermore, they provide qualitative insight into the reasons for the over- or under-performance observed in the wave results for both the TOT and WND experiments. During storm Arwen, strong Northern winds at the Northern Iberian Peninsula (**Fig. 12[a]**) generated Northern wind-sea waves, that were reinforced with NW swell waves. The integrated  $H_{m0}$  of this mixed sea state can be seen in **Fig. 13**. The storm peak was better handled (**Fig. 13[a]**), because at the wind-sea generation area, the ANN Winds predicted more intensity than ECMWF-IFS. The wind speed time series at GAL-D does not show much difference (**Fig. 13[c]**), but **Fig. 12[b]** shows clearly the extra wind energy at the area (red marked region), that lead the TOT and WND simulations to capture better the storm peak at 27<sup>th</sup> November 2021. Note also that the NW Mediterranean has less energy in KAILANI than in IFS (i.e. the blue area in **Fig. 12[b]**), suggesting the same issues mentioned in the general performance.*

The Generative Adversarial Network (GAN) corrects wind speed but does not generate atmospheric patterns that strongly differ from ECMWF-IFS, leading to similar spatial structures.

We also note the specific case of the Gran Canaria (GCA) station. In this area, even the control simulation (ECMWF-IFS winds) was indeed unable to reproduce the observed moderate waves. The presence of the double peak at GCA is an interesting case. While the GAN successfully corrects systematic biases, it currently appears unable to reproduce highly localized wind effects. Since the waves were primarily wind-sea, the discrepancy could stem either from the local winds not being properly modeled or from the spectral wave model not accurately handling wave growth. As the reviewer correctly pointed out, the ANN's enhancement seemed to be a minor improvement that primarily involved increasing the wind speed in that zone.

However, any enhancements achieved under extreme events should be considered preliminary, and further research is needed for proper validation. This statement has been added throughout the text.

Line 671: "The ANN forcings have positive impact on the wave forecast, especially under extreme events." To be honest, the results for significant wave height seem to reduce the variability in the signal. Peaks are less pronounced in Figure 15, and almost no signal in the time series in Figure 16. In general, a diurnal signal appears to be present in the observed wave field, which is lost in the simulation.

We thank the Reviewer#2 for this remark. We agree that the variability is not the same as in the in-situ buoy, but this low variability is also present in the control simulation.

This statement has been removed in the revised version.

Line 674: "Wind ANN tends to decrease the overestimation of ECMWF-IFS wind speed". Is this correct? Time series plots in Figure 13-16 indicate that ANN winds are frequently higher than ECMWF and observations.

We thank the Reviewer#2 for this remark. This statement is correct regarding the Tarragona site, where the predicted ANN winds tend to be lower. Specifically, Figure 15 shows that the ANN winds are lower at the buoy location. Additionally, Figure 12 illustrates that the NW Mediterranean is an area where ECMWF has consistently higher values than IFS (indicated by the blue zones). The reasons are the same as those noted in the Line 360-361 specific comment: SAR data has better agreement than the IFS but exhibits lower overestimation in the NW Med. Furthermore, as was also mentioned, the IBI-WAV parameters are tuned for the NE Atlantic and therefore do not perform properly in the NW Mediterranean.