

1 **Review article: Harnessing data driven methods for climate multi-**  
2 **hazard and multi-risk assessment**

3  
4 Davide Mauro Ferrario<sup>1,2,3</sup>, Marcello Sano<sup>1,2,4</sup>, Margherita Maraschini<sup>1,2</sup>, Andrea Critto<sup>1,2</sup>, Silvia  
5 Torresan<sup>1,2</sup>

6 <sup>1</sup>Department of environmental sciences, informatics and statistics, Ca' Foscari University Venice, 30170, Venice, Italy

7 <sup>2</sup>CMCC Foundation, EURO Mediterranean Center on Climate change, 73100, Lecce, Italy

8 <sup>3</sup> Institute for Advanced Study IUSS Pavia, 27100, Pavia, Italy

9 <sup>4</sup>Griffith University, 4222, Gold Coast, Australia

10 *Correspondence to:* Andrea Critto ([critto@unive.it](mailto:critto@unive.it))

11 • Davide Mauro Ferrario, [davide.ferrario@cmcc.it](mailto:davide.ferrario@cmcc.it)

12 • Marcello Sano [marcello.sano@unive.it](mailto:marcello.sano@unive.it)

13 • Margherita Maraschini, [margherita.maraschini@cmcc.it](mailto:margherita.maraschini@cmcc.it)

14 • Andrea Critto, [critto@unive.it](mailto:critto@unive.it)

15 • Silvia Torresan, [silvia.torresan@cmcc.it](mailto:silvia.torresan@cmcc.it)

16

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

17 **Abstract.** In recent years, interest in data-driven methods, such as machine learning and multivariate statistics for multi-hazard  
18 and multi-risk assessment has surged, due to their ability to integrate vast amounts of data in modelling complex non-linear  
19 relationships between hazard and risk factors. This review explores data-driven methods in climate multi-hazard and risk  
20 analysis, focusing on four themes: (i) data processing and collection; (ii) hazard identification, prediction and analysis; (iii)  
21 risk analysis; and (iv) future risk scenarios under climate change. Key findings highlight the extensive use of machine learning  
22 to combine Earth observations and climate data for downscaling and land use and land cover characterisation; the application  
23 of deep learning for hazard prediction; the use of ensemble methods for risk analysis; and the growing emphasis on explainable  
24 AI frameworks. Supervised machine learning approaches trained on historical impact data to project future climate risks have  
25 also emerged as a significant research area. Future research in this area should focus on modelling multi-hazard interactions,  
26 particularly triggering and cascading effects, integrate dynamic vulnerability and exposure factors, and address uncertainties  
27 associated with using machine learning for extrapolation. Advancements in Earth observations and textual data integration,  
28 alongside the development of open-access disaster catalogues, will also be crucial for improving multi-risk analyses and  
29 supporting AI-driven early warning systems tailored to regional needs.

30

31

33 **Table of Contents**

34	<b><u>1 Introduction.....</u></b>	<b><u>5</u></b>
35	<b><u>2 Methodology.....</u></b>	<b><u>7</u></b>
36	<b><u>2.1 Research questions.....</u></b>	<b><u>7</u></b>
37	<b><u>2.2 Methodological framework: search methodology, screening, reporting and interpreting.....</u></b>	<b><u>9</u></b>
38	<b><u>2.3 Limitations and scope of the review.....</u></b>	<b><u>10</u></b>
39	<b><u>3 Results and discussions.....</u></b>	<b><u>12</u></b>
40	<b><u>3.1 Data.....</u></b>	<b><u>12</u></b>
41	3.1.1 Climate datasets.....	12
42	3.1.2 Earth observations.....	15
43	3.1.3 Textual data.....	16
44	<b><u>3.2 Multi-hazard.....</u></b>	<b><u>17</u></b>
45	3.2.1 Identify, classify and cluster.....	17
46	3.2.2 Hazard forecasting and prediction.....	19
47	3.2.2 Modelling hazard interaction.....	21
48	<b><u>3.3 Multi-risk.....</u></b>	<b><u>25</u></b>
49	3.3.1 Modelling risk combining susceptibility, exposure and vulnerability.....	25
50	3.3.2 Modelling risk predicting impacts.....	27
51	<b><u>3.4 Future.....</u></b>	<b><u>31</u></b>
52	3.4.1 Predicting future hazards.....	31
53	3.4.2 Modelling future impacts.....	33
54	<b><u>3.5 Limitations and future research directions.....</u></b>	<b><u>35</u></b>
55	<b><u>4 Conclusion.....</u></b>	<b><u>37</u></b>
56	<b><u>Appendix A: Abbreviations.....</u></b>	<b><u>38</u></b>

57	<b><u>Appendix B: Summary tables of the collected studies</u></b> .....	<b>40</b>
58	<b><u>Author contribution</u></b> .....	<b>67</b>
59	<b><u>Competing interest</u></b> .....	<b>67</b>
60	<b><u>Financial support</u></b> .....	<b>67</b>
61	<b><u>References</u></b> .....	<b>68</b>
62	▼ .....	
63		

<b>Deleted: 1 Introduction</b> → 5*
<b>2 Methodology</b> → 7*
<b>2.1 Research questions</b> → 7*
<b>2.2 Methodological framework: search methodology, screening, reporting and interpreting</b> → 9*
<b>2.3 Limitations and scope of the review</b> → 10*
<b>3 Results and discussions</b> → 12*
<b>3.1 Data</b> → 12*
3.1.1 Climate datasets → 12*
3.1.2 Earth observations → 15*
3.1.3 Textual data → 16*
<b>3.2 Multi-hazard</b> → 17*
3.2.1 Identify, classify and cluster → 17*
3.2.2 Hazard forecasting and prediction → 19*
3.2.2 Modelling hazard interaction → 21*
<b>3.3 Multi-risk</b> → 25*
3.3.1 Modelling risk combining susceptibility, exposure and vulnerability → 25*
3.3.2 Modelling risk predicting impacts → 27*
<b>3.4 Future</b> → 31*
3.4.1 Predicting future hazards → 31*
3.4.2 Modelling future impacts → 33*
3.5 Limitations and future research directions → 35*
<b>4 Conclusion</b> → 37*
<b>Appendix A: Abbreviations</b> → 38*
<b>Appendix B: Summary tables of the collected studies</b> → 40*
<b>Author contribution</b> → 67*
<b>Competing interest</b> → 67*
<b>Financial support</b> → 67*
<b>References</b> → 68*
<b>1 Introduction</b> → 5*
<b>2 Methodology</b> → 7*
<b>2.1 Research questions</b> → 7*
<b>2.2 Methodological framework: search methodology, screening, reporting and interpreting</b> → 9*
<b>3 Results and discussions</b> → 10*
<b>3.1 Data</b> → 10*
3.1.1 Climate datasets → 10*
3.1.2 Earth observations → 13*
3.1.3 Textual data → 15*
<b>3.2 Multi-hazard</b> → 16*
3.2.1 Identify, classify and cluster → 16*
3.2.2 Hazard forecasting and prediction → 18*
3.2.2 Modelling hazard interaction → 19*
<b>3.3 Multi-risk</b> → 23*
3.3.1 Modelling risk combining susceptibility, exposure and vulnerability → 23*
3.3.2 Modelling risk predicting impacts → 25*
<b>3.4 Future</b> → 30*
3.4.1 Predicting future hazards → 30*
3.4.2 Modelling future impacts → 31*
<b>4 Conclusion</b> → 33*
<b>Appendix A: Abbreviations</b> → 36*
<b>Appendix B: Summary tables of the collected studies</b> → 37*
<b>Author contribution</b> → 61*
<b>Competing interest</b> → 61*
<b>Financial support</b> → 61*
<b>References</b> → 62*

## 122 **1 Introduction**

123 The growing interconnectedness between socio-economic and natural systems, coupled with the escalating challenges  
124 presented by climate change, has led to increased complexities in climate risk analysis. At the same time, a wider availability  
125 of data on multiple risk drivers, including weather observations, Earth observations (EO), climate reanalyses and projections,  
126 socio-economic indicators, and social media, coupled with advances in machine learning (ML) and statistical methods, are  
127 increasing the potential of data-driven methodologies, which promise to revolutionise climate risk assessment (Kashinath et  
128 al., 2021a; Reichstein et al., 2019). To unlock the full potential of this data, it is crucial to develop and apply advanced methods  
129 for processing, harmonizing, and integrating heterogeneous datasets. These efforts enable the generation of actionable insights  
130 essential for effective multi-hazard and multi-risk assessments, by leveraging the accessibility of large datasets to be explored  
131 with advanced ML and statistical techniques.

132 Complex dynamics characterize socio-environmental and climate risk: applications may underestimate impacts if they do not  
133 take into account the compounding, cascading and amplifying interactions of hazards and their effect on vulnerability and  
134 exposure factors. In fact, (i) compounding hazards (co-occurring in the same location and at the same time) can lead to impacts  
135 which may be substantially higher than the sum of the single events taken in isolation (Arosio et al., 2020; Zscheischler et al.,  
136 2018), (ii) the occurrence of one hazard itself can modify vulnerability or resilience of the system, exposing assets or  
137 communities to higher risks, such as in the case of consecutive hazards (de Ruiter & van Loon, 2022), and (iii) impacts and  
138 risks can propagate across multiple scales and sectors, extending far beyond the area initially hit and affecting whole systems  
139 (Arosio et al., 2021; Pescaroli & Alexander, 2018), such as in the case of high-impact and low-probability events (Linkov et  
140 al., 2022). For these reasons, the international community (Intergovernmental Panel on Climate Change (IPCC), 2023;  
141 UNDRR, 2020) has recently pledged for a paradigm shift from single hazard towards a more comprehensive representation of  
142 multiple and interconnected climatic risks (AghaKouchak et al., 2020; De Angeli et al., 2022; Gallina et al., 2020; Šakić  
143 Trogrlić et al., 2024; Terzi et al., 2019; Tilloy et al., 2019; Ward et al., 2022). To achieve this shift, it is essential to develop  
144 data-driven methodologies that can analyse and predict the interactions and dependencies between multiple hazards, enabling  
145 a more accurate characterisation of their compounding and cascading effects.

146 To better navigate the many definitions surrounding multi-hazard risk concepts, this paper adopts the terminology used in  
147 Zschau (2017), where *multi-layer single hazards* refers to applications focussing on more than one hazards, without  
148 considering hazard interactions; *multi-hazard* focuses on hazards interaction; *multi-hazard risk* refers to applications  
149 considering risks in a multi-hazard framework, without discussing interactions at vulnerability level, and finally *multi-risk*  
150 refers to the most complex analysis comprising interactions at both hazard and vulnerability level.

151 The complex nature of multi-hazard events presents significant challenges to existing risk assessment methodologies, which  
152 treat hazards and risks singularly and often struggle to handle the non-linear interactions and feedback loops between multiple  
153 risk drivers (Tilloy et al., 2019). ML techniques have recently gained traction in climate science and risk analysis for their  
154 ability to process and integrate large, heterogeneous datasets from sources such as weather observations, Earth observations,

155 climate reanalyses and projections, socio-economic indicators, and even social media. By learning from historical data, they  
156 can uncover non-linear risk patterns and detect correlations across spatial and temporal scales, driving their growing use in  
157 climate risk assessment (Reichstein et al., 2019; Zennaro et al., 2021).

158 Integrating these heterogeneous data sources can help in capturing multi-hazard interactions and characterise their impacts on  
159 social, economic, and natural systems, especially thanks to the introduction of new Deep Learning (DL) architectures and  
160 models, specialized in capturing both spatial and temporal non-linear interactions (S. Park et al., 2023).

161 As ML models have become more complex, attention has shifted toward making these models more interpretable and  
162 explainable (Carvalho et al., 2019). This is especially important for applications focussing on risk, where it is crucial to quantify  
163 the contribution of each input feature to the model's prediction, making it easier to assess how different risk variables impact  
164 the overall risk. In this context, explainability frameworks improve the robustness of risk assessments and enhance trust in the  
165 model's outputs by providing insights into how the model arrives at specific conclusions (S. Jiang et al., 2024; McGovern et  
166 al., 2019), supporting transparency and accountability for stakeholders.

167 In addition to ML methods, this review briefly considers the role of copulas as multivariate statistical tools in multi-risk  
168 assessment. Copulas enable explicit modelling of the dependence structure between variables, making them particularly  
169 valuable for analysing compound events in which multiple hazards occur simultaneously or sequentially (see, for example,  
170 Agrawal, 2022; Hochrainer-Stigler et al., 2019). They have, for instance, been used to characterise the joint occurrence of  
171 droughts and heatwaves, yielding insights into their combined impacts on agriculture and water resources (see e.g. Ribeiro et  
172 al., 2020). Although their application is more specialised than most ML approaches, copulas provide critical information about  
173 inter-hazard dependencies, supporting a deeper understanding of compounding and interacting risks. Their inclusion in this  
174 review therefore highlights their importance in contexts requiring precise statistical modelling of hazard interactions and  
175 underscores how they complement broader ML-based strategies in climate-risk analysis. To advance this field, there is a critical  
176 need for predictive frameworks that can leverage these advanced methods to forecast long-term future multi-hazard and multi-  
177 risk scenarios, addressing uncertainties and guiding adaptive risk management strategies under changing climatic conditions.

178 To support implementation, the development of a wide range of open-source libraries (e.g., *scikit-learn*, *TensorFlow*, *Keras*,  
179 *PyTorch*, *VineCopulas* (Claassen et al., 2024), etc.), allows users to implement, train, validate, and deploy models with  
180 minimal programming expertise, making it possible for non-experts or domain specialists with limited knowledge to efficiently  
181 apply advanced techniques to risk modelling. This democratization of tools reduces the technical barriers for researchers and  
182 practitioners, enabling more interdisciplinary collaborations and accelerating the adoption of data-driven methods in climate  
183 risk management (Rolnick et al., 2019).

184 This paper aims to provide a comprehensive review of data-driven methods, with a specific focus on ML approaches, for multi-  
185 hazard and multi-risk assessment, exploring ongoing applications, current limitations and future perspectives, while also  
186 addressing the use of copulas, a non-ML statistical method, to highlight its role in modelling dependencies in compound hazard  
187 events. Unlike other recent reviews that have focused on ML (particularly DL) for specific hazards or sectors – such as extreme  
188 events (Salcedo-Sanz et al., 2022), hydrology (Tripathy & Mishra, 2024), geophysics (S. Yu & Ma, 2021), wildfires (Jain et

189 al., 2020), and climate risk (Zennaro et al., 2021) – this paper takes a cross-cutting perspective on multi-hazard and multi-risk  
190 assessment. By structuring the discussion around successive stages of risk analysis – data processing, hazard prediction, risk  
191 assessment, and future scenarios – we connect climate risk and data-driven methods while also identifying critical gaps,  
192 particularly in linking hazard interactions with vulnerability.

193  
194 The paper is structured as follows: Section 2 Methodology outlines the research questions, and the search methodology  
195 employed for the review. Section 3 Results and discussions summarises the literature review findings and discusses key  
196 insights related to each of the research questions. Section 4 Conclusion provides a summary of the key insights and outlines  
197 the next steps for research in this field. The Appendices provide an abbreviation dictionary ([Appendix A: Abbreviations](#)), as  
198 well as the summary tables of main articles collected for each research question ([Appendix B: Summary tables of the collected  
199 studies](#)).

**Deleted:** Appendix A: Abbreviations

**Deleted:** Appendix B: Summary tables of the collected studies

## 200 **2 Methodology**

201 This paper follows a systematic review process based on the Preferred Reporting Items for Systematic Reviews and Meta-  
202 Analyses (PRISMA) methodology, which ensures a standardized, systematic, and transparent framework for analysing and  
203 synthesizing existing literature (O’Dea et al., 2021). The method involves several steps, among which the main ones are:  
204 defining of the research questions; developing a protocol detailing the search methodology (including database to search,  
205 keywords, timeframe and selection criteria); collecting and screening relevant literature; synthesizing and interpreting the  
206 findings. Such a stepwise process ensures a thorough search for relevant studies, consistent criteria for the selection of papers,  
207 and clear documentation of the review process, therefore reducing the risk of bias and enhancing the robustness and  
208 replicability of the analysis (Sarkis-Onofre et al., 2021).

### 209 **2.1 Research questions**

210 Each of the four research questions (Figure 1) is focussed on a specific topic and presents several sub-topics, offering a  
211 structured framework to explore current applications, address challenges, and pinpoint future opportunities. These research  
212 questions are:

- 213 1. Data: How can Machine Learning improve data collection and processing?
- 214 2. Multi-Hazard: How can Machine Learning and statistical tools be used to analyse extreme events, and model hazard  
215 interactions?
- 216 3. Multi-Risk: How can Machine Learning applications integrate vulnerability and exposure in multi-risk analysis?
- 217 4. Future: How can Machine Learning and statistical tools be used to predict long-term future multi-hazard and multi-  
218 risk?
- 219



Figure 1: Research questions and sub-themes

222  
223

224 The first research question examines how ML can help process diverse types of data, extracting and harmonising the  
225 information needed to analyse multi-hazard and multi-risk by addressing current gaps such as data sparsity, inconsistency  
226 across sources, and the lack of harmonised formats. This contributes to improving the quality and comparability of risk  
227 assessments by enabling integrated use of climate, EO, and textual datasets. In particular, the sub-themes are divided based on  
228 the type of data analysed:

- 229 I. Climate data (time series of geospatial climate data), which describe the characteristics of climate-related hazards  
230 across space and time. Preparing this data for multi-hazard and multi-risk applications often requires ML methods  
231 (i.e. feature engineering) to increase spatial and temporal resolution, harmonise and extend the time coverage of the  
232 datasets or correct for biases (Schneider et al., 2023).
- 233 II. EO, which can be used to characterise exposure and vulnerability layers and extract information on impacts  
234 (Ghaffarian & Emtchani, 2021; Novellino et al., 2024).
- 235 III. Textual data, such as newspapers or social media, which in the last years have been leveraged for extracting  
236 information on diverse impacts (Sodoge et al., 2023).

237

238 The second research question investigates how ML and statistical tools improve the identification and modelling of hazard  
239 dynamics by capturing complex spatio-temporal patterns, compounding effects, and non-linear interactions that traditional  
240 approaches often overlook. This helps advance multi-hazard and multi-risk analysis by providing more accurate detection,  
241 classification, and modelling of extreme events. In particular, the key sub-themes are:

- 242 I. Analyse which methods can be used to identify, classify and cluster extreme events, producing spatio-temporal  
243 footprints of multi-hazard events (H. Yu et al., 2022).
- 244 II. The prediction of (multi-)hazard events, for example through early warning systems or seasonal predictions  
245 (Bhowmik et al., 2023).
- 246 III. The analysis of hazard interactions, for example characterising joint distributions through copulas (Bevacqua et al.,  
247 2021) or multi-hazard susceptibility maps (Pourghasemi et al., 2019).

248 The third research question concerns the application of ML for the integration of vulnerability and exposure into multi-risk  
249 analysis addressing the current gap where vulnerability and exposure are often treated as static or secondary layers rather than  
250 dynamic drivers of risk. This integration strengthens the ability of multi-risk assessments to capture how socio-economic  
251 conditions and adaptation measures interact with hazards to shape overall risk. In particular, the key themes are:


- 252 I. Multi-hazard exposure and vulnerability on assessments, integrating susceptibility mapping with information on  
253 specific exposure layers, such as buildings and population (Rusk et al., 2022).
- 254 II. Modelling risk from past impacts data, often through supervised ML approaches that use hazard, vulnerability and  
255 exposure indicators as predictors (Dal Barco et al., 2024).

256 The fourth research question investigates the possible contribution of ML and statistical tools into the analysis of (long-term)  
257 future multi-hazard and multi-risk, where uncertainty associated with the representation of future extremes in climate  
258 projections further complicates risk modelling, highlighting a critical gap in existing approaches, which often fail to adequately  
259 capture compound and cascading extremes under changing climate conditions. This research question clarifies how ML can  
260 enhance scenario building, improve uncertainty quantification, and support more robust long-term multi-risk assessments. In  
261 particular, the key sub-themes are:

- 262 I. Modelling future multi-hazard trends and spatial patterns using statistical methods, in particular for compound and  
263 consecutive events (Zscheischler et al., 2018).
- 264 II. Assessing future impacts based on climate change projections, often using methods trained on historical data and  
265 applied to ensembles of RCP projections (S. J. Park & Lee, 2020).

## 266 **2.2 Methodological framework: search methodology, screening, reporting and interpreting**

267 The search was performed on Scopus, focusing on articles published in English. Since the analysis focuses on ML applications  
268 and multi-risk, the timeframe 2010 – 2024 was chosen because both areas of research are recent and other reviews have  
269 addressed earlier periods, highlighting that most applications in ML and climate risk have been published only in the last few  
270 years (Zennaro et al., 2021). For each research question, a dedicated search was performed. Each search string was generated  
271 by the combination of a set of method-related keywords (e.g. those related to ML or statistical methods), common across all  
272 questions, and a set of thematic keywords, specific to each research question (Figure 2).



273  
274 **Figure 2: Literature review methodology**

275 After collecting articles for each research question, the papers were first filtered by following typologies: 'journal articles',  
276 'conference papers', and 'book chapters'. Afterwards, for each research question, the papers were screened by title, then by  
277 abstract, and finally by full text. The final screening selected 153 key papers to be analysed in the literature review. This  
278 information was then summarised into tables, identifying the type of applications, the type of data used, the case study and the  
279 methods used. Finally, for each research questions, the results were discussed to understand the maturity level of the  
280 applications, their limitations and possible future developments.

281 **2.3 Limitations and scope of the review**

282 While this review follows the PRISMA guidelines for search strategy, screening, and reporting, a formal numerical quality  
283 scoring of individual studies was not applied, consistent with standard practice in PRISMA-based reviews of computational  
284 methods in geoscience and climate risk (e.g., Zennaro et al., 2021; Salcedo-Sanz et al., 2022; Ghaffarian et al., 2023). Instead,  
285 quality and relevance were assessed qualitatively during full-text screening based on three criteria: methodological rigour  
286 (evaluated through the presence and type of model validation, e.g., cross-validation, independent test sets, or benchmark  
287 comparisons), relevance to the research questions, and diversity across data sources, geographical coverage, hazard types, and  
288 ML approaches. The latter criterion was applied explicitly to avoid over-representation of any single method or region in the  
289 final corpus, and is documented in Appendix B.

Formatted: Font:

Formatted: Heading 2

290 Another limitation concerns terminological consistency. Although this review adopts the Zschau (2017) framework to  
291 reclassify papers during full-text screening, the terms multi-hazard, multi-risk, compound, and cascading are used with  
292 considerable inconsistency across the reviewed literature, a well-documented feature of the field (Gill & Malamud, 2014;  
293 Tilloy et al., 2019). Because paper selection was based on keyword matching against author-assigned terminology, the corpus  
294 necessarily reflects this heterogeneity, and the thematic categories used in the synthesis should be understood as analytical  
295 conveniences rather than sharp taxonomic boundaries.

296 Moreover, while this review focuses on ML and copula-based methods as the primary data-driven approaches for multi-hazard  
297 and multi-risk assessment, it is important to acknowledge that several complementary quantitative frameworks exist and have  
298 been the subject of dedicated reviews that fall outside the scope of the present work. Bayesian networks (BNs) provide a  
299 probabilistic framework for multi-hazard causal modelling, capturing conditional dependencies between risk drivers through  
300 directed acyclic graphs and propagating uncertainty in a transparent, interpretable way; they are particularly valuable in data-  
301 sparse contexts where causal structure can be informed by expert knowledge, and their application to climate multi-risk  
302 assessment has been reviewed in depth by Sperotto et al. (2017). Agent-based models (ABMs) simulate the adaptive behaviour  
303 of individuals and institutions under hazard scenarios, making them suited to capturing dynamic vulnerability, evacuation  
304 dynamics, and community resilience processes that purely data-driven models cannot represent; comprehensive reviews of  
305 their application in disaster management are provided by Zhuo & Han (2020) and Anshuka et al. (2022). More broadly, the  
306 full landscape of quantitative methods for modelling hazard interrelationships, including stochastic, empirical, and mechanistic  
307 approaches, is systematically covered by Tilloy et al. (2019), providing a valuable complement to the ML-focused perspective  
308 of the present review.

309 Finally, some considerations need to be taken on the geographical distribution of the 152 papers included in this review, which  
310 reveal a marked concentration in a small number of regions. In terms of lead authorship, Europe (35.5%) and East Asia (27.0%)  
311 together account for nearly two thirds of the corpus, followed by North America (21.1%), while the Global South is  
312 substantially underrepresented: Africa, South America, and Oceania collectively contribute less than 5% of lead authors. A  
313 similar pattern holds for co-authorship, though with a slight broadening of participation: South/SE Asia rises to 6.4% and  
314 Middle East to 7.3%, suggesting that researchers from these regions participate more frequently as collaborators than as lead  
315 investigators. The most pronounced shift occurs in the case study column: Global studies account for 14.9% of the corpus, and  
316 South/SE Asia (10.6%), Middle East (6.8%), and Africa (5.0%) are more represented as study areas than as sources of  
317 authorship, indicating that data-driven methods developed in high-income regions are frequently applied to, rather than  
318 developed within, lower-income contexts. The full breakdown of lead author institution country, co-author countries, and case  
319 study regions, together with a Sankey diagram illustrating the flows between these three dimensions, is provided in Appendix  
320 B. These geographical imbalances should be borne in mind when interpreting the findings of this review, as the methods,  
321 datasets, and risk priorities that dominate the literature inevitably reflect the institutional contexts in which the research was  
322 produced.

323

## 324 **3 Results and discussions**

### 325 **3.1 Data**

#### 326 **3.1.1 Climate datasets**

327 The application of ML methods to produce new, complete, or high-resolution hazard datasets (either from meteorological  
328 observations, climate reanalyses or future projection) is quite established, and mainly focuses on data with sparse and irregular  
329 measurements. A typical indicator which is derived with ML methodologies is soil moisture: in-situ measurements are usually  
330 scarce and not uniformly distributed, satellite images (which will be discussed later) often presents temporal gaps and can only  
331 provide information on the first layer and struggles in complex topographies and it presents a complex dynamic that is  
332 influenced by many different drivers (similarly to multi-risk prediction) such as precipitation, temperature, evaporation,  
333 topography and land use. For example, Kang et al. (2018) and O. & Orth, (2021) investigate the complex interactions at  
334 different soil levels and temporal scales with a Long-Short Term Memory (LSTM) model that takes as inputs the topography,  
335 vegetation and atmospheric conditions and predicts each soil moisture layer in succession, using ERA-5 reanalysis as  
336 assessment endpoint. LSTM is widely applied to model the behaviour of other hydrological variables, such as snow, run-off  
337 and river catchments. Entity-Aware LSTM was used for rainfall-runoff modelling by Kratzert et al., (2018); Kratzert, Klotz,  
338 Shalev, et al., (2019), to include both static and dynamic inputs allowing the algorithm to explicitly differentiate the two  
339 different types. Ghiggi et al. (2019) applies Random Forest (RF) regression to predict monthly runoff rates in the timeframe  
340 1902-2020, based on antecedent precipitation and temperature from an atmospheric reanalysis, validating the results with in-  
341 situ streamflow observations. Other research focuses on different variables and in particular investigate the irregular  
342 distribution of sensors: Amato et al. (2020) introduces a multi-step methodology to interpolate irregularly distributed spatio-  
343 temporal timeseries, first decomposing the signal and then learning stochastic spatial coefficients which can be spatially  
344 modelled and mapped on a regular grid with Artificial Neural Networks (ANN), allowing the reconstruction of the complete  
345 spatio-temporal signal.

346 ML methods have been applied also to climate reanalyses and models. Early applications, such as He et al. (2016), tested RF  
347 regression to statistically downscale spatially precipitation data, using few covariates and demonstrating how this approach is  
348 able to catch the non-linear relations between variables, minimising overfitting and collinearity issues between predictors.  
349 However, the algorithm struggled with skewed datasets and even the final model, which is the combination of two different  
350 RF models, trained respectively on high-precipitation and low-precipitation values, fails to detect the complex spatial and  
351 temporal complexity of precipitation data, overestimating the intensity and spatial distribution of low precipitation and  
352 underestimating high precipitation. Other applications are focussing on Deep Learning models: CNNs are used to downscale  
353 many variables from future climate models (among which, air temperature, precipitation, 10-m wind speed, 2-m relative  
354 humidity, downward shortwave radiation) (Lin et al., 2023). Generative models particularly Generative Adversarial Networks  
355 (GAN) and diffusion models, are widely used for this task. GANs consist of two neural networks – a generator and a  
356 discriminator - that are trained simultaneously in a competitive process. The generator attempts to create realistic fake data

357 that can fool the discriminator, while the discriminator works to distinguish between real and fake data. For example, specific  
358 GANs based on Convolutional Neural Networks (CNNs) have been applied to post-process weather forecast outputs. These  
359 models can enhance the resolution of precipitation data by a factor of ten, producing more realistic and spatially coherent  
360 forecasts compared to the original input data (Harris et al., 2022). Diffusion models, on the other hand, learn to reverse a noise  
361 process: first the model adds sequentially noise to input data, then the model learns how to predict the noise at each step, and  
362 once trained, it can start with noisy data and work backwards, progressively removing the noise to generate a new, realistic  
363 dataset. Diffusion models are related to variational inference, where the forward process defines a probabilistic trajectory from  
364 data to noise, and the reverse process defines a generative path from noise back to data. Unlike other generative models like  
365 GANs, which learn through a "discriminative" process (trying to fool a discriminator network), diffusion models learn through  
366 this smooth diffusion and denoising process (Yeğin & Amasyalı, 2024). For example, diffusion models are applied to  
367 downscale multiple climate models, also providing information on the uncertainty downscaling, by generating a large number  
368 of ensemble members based on probability distribution sampling (Ling et al., 2024a). DL approaches are often used to  
369 downscale low-resolution future models to Convection Permitting (CP) climate models, where the main advantage of these  
370 techniques is their reduced computational costs compared to the development of a CP climate models (Bretherton et al., 2022;  
371 Clark et al., 2022). The role of Artificial Intelligence (AI) in climate predictions is discussed in Schneider et al. (2023). This  
372 study advocates for the development of global models at 10–50 km resolution, harnessing AI and EO for the calibration and  
373 development of higher-resolution regional simulations.

374 In recent years, there has been growing interest in hybrid modelling: approaches that combine data-driven ML methods with  
375 physical or process-based models or constraints, as a way to benefit from both high flexibility and physical realism. Such  
376 hybrid / physics-informed ML methods help address several limitations of pure data-driven models: they can enforce  
377 conservation laws, reduce overfitting to noise, improve generalization especially under conditions outside the training domain,  
378 and provide more interpretable insights into underlying drivers. For instance, He et al., (2023) integrates ML corrections into  
379 a land-surface / atmospheric model using data assimilation, remote sensing LAI and soil moisture to improve climate  
380 simulations. Similarly, Huynh et al., (2025) combines process-based hydrological flux models with neural networks to correct  
381 for scale mismatches and to better capture spatial heterogeneity. Also, (S. Yu et al., 2024) provides benchmarks for ML  
382 emulators that mimic nested high-resolution physical simulations. Despite their promise, hybrid models also face important  
383 limitations. They often require substantial domain and physical knowledge to be formulated appropriately and to ensure  
384 physical consistency (Willard et al., 2022). Moreover, coupling ML architectures with numerical process models can remain  
385 computationally demanding, particularly for high-resolution simulations or large spatio-temporal domains (Reichstein et al.,  
386 2019). Calibration and validation can also be complex, as balancing the contributions of the physical and data-driven  
387 components often involves ad hoc or case-specific tuning (Read et al., 2019). Finally, interpretability may still be reduced  
388 when the ML component acts as a black box, obscuring how physical constraints shape predictions (Kashinath et al., 2021a).  
389 These challenges are also relevant for hazard prediction, where process dynamics such as land–atmosphere feedback play  
390 central roles and require models that are both physically credible and statistically robust. Thus, hybrid models represent an

391 emerging frontier at the interface of ML, process-based modeling, and data assimilation, particularly relevant for both climate  
392 data reconstruction and hazard modelling and deserve explicit consideration in future reviews and benchmarking efforts.

393 Machine learning applications for climate and environmental datasets have greatly improved the reconstruction and  
394 downscaling of variables from sparse and irregular observations. However, a critical yet often under-addressed aspect in this  
395 field is uncertainty quantification (UQ), which is particularly relevant when these datasets are later used for hazard or risk  
396 assessments (Beven, 2018). Uncertainty in ML-based arises from multiple sources: Aleatoric uncertainty stems from the  
397 intrinsic variability and noise in the underlying measurements, such as sensor errors, missing satellite observations, or  
398 inconsistent temporal coverage; epistemic uncertainty originates from limited or biased training data and model structural  
399 choices (Xu et al., 2022). Several probabilistic approaches have been explicitly designed to represent spatial data uncertainty  
400 by learning distributions rather than deterministic predictions, mainly involving Bayesian Networks (BN) and Gaussian  
401 Processes (GP) (Siddique et al., 2022). For example, Multi-fidelity Gaussian Processes with a 5/2 Matern kernel in particular,  
402 were used to downscale precipitation data from ERA-5 over high mountain terrain. Multi fidelity models combine low-fidelity  
403 observations (which are usually more numerous and less expensive to obtain) with high-fidelity ones. This makes the model  
404 more suited than other state-of-the-art machine learning methods for smaller datasets and able to quantify and narrow the  
405 uncertainty associated with the precipitation estimates, which is especially needed over ungauged areas and can be used to  
406 estimate the likelihood of extreme events that lead to floods or droughts (Tazi et al., 2024). Andersson et al., (2023) applies  
407 Convolutional Neural Processes (ConvNPs), to suggest informative sensor placements by finding sites that maximally reduce  
408 prediction uncertainty, testing it for air temperature anomalies measurements in Antarctica. Convolutional Neural Processes  
409 (ConvNPs) extend the probabilistic framework of Gaussian Processes by learning flexible, data-driven covariance structures  
410 through neural networks. While traditional GPs provide robust uncertainty estimates but suffer from scalability and stationarity  
411 constraints (M. Jiang et al., 2022), ConvNPs maintain a probabilistic foundation while scaling linearly with data size and  
412 accommodating irregular spatial inputs (Garnelo et al., 2018). DeepSensor<sup>1</sup>, a specific GitHub python package, was developed  
413 to facilitate the application of Neural Processes in environmental sciences, especially for downscaling, interpolation, sensor  
414 placement and data imputation. Monte Carlo Dropout (MCD) enhances epistemic uncertainty quantification in climate data  
415 and was tested on neural networks for probabilistic medium-range weather forecasting (Garg et al., 2022). Deep generative  
416 models such as diffusion or GAN frameworks can further approximate uncertainty by generating ensembles of plausible  
417 realisations that sample the predictive probability space (Ling et al., 2024b; Saha & Ravela, 2022). Despite these advances,  
418 most studies still focus primarily on improving resolution and accuracy, while systematic approaches to quantifying and  
419 propagating uncertainty through the modelling chain, from data to hazard and risk estimates, remain limited (Beven, 2018).  
420 Addressing this challenge is crucial, as downstream risk assessments rely heavily on the reliability of the climate inputs that  
421 feed them.

---

<sup>1</sup> <https://github.com/alan-turing-institute/deepsensor>

### 422 3.1.2 Earth observations

423 EO data, when combined with ML is increasingly recognised for its critical role in supporting actionable multi-hazard and  
424 multi-risk assessment, as evidenced by new initiatives from ESA and NOAA's Centre for AI, where particular attention is  
425 devoted to the use of EO for discovering impacts in remote areas and developing early warning systems.

426 Remote sensing images are used to improve climate datasets, for example increasing the spatial coverage in areas with sparse  
427 measurements or providing real data to bias-correct/downscale modelled data. Multiple AI methods, such as Support Vector  
428 Machine (SVM) (Ahmad et al., 2010; Jing et al., 2016a), Ridge Regression (Kang et al., 2018), RF (Han et al., 2023; Jing et  
429 al., 2016b) and LSTM (Fang et al., 2017) are applied for developing soil moisture datasets.

430 EO provides consistent, near-real time observations of environmental conditions that are critical for early warning and hazard  
431 characterisation. For instance, indicators such as vegetation stress (Miyoshi et al., 2020; Schiefer et al., 2020; Veras et al.,  
432 2022), surface temperature anomalies can enable the early detection of droughts (Barrett et al., 2020), floods (Dasgupta et al.,  
433 2022) or wildfires (Jain et al., 2020) especially in remote and data scarce areas. DL and Physics Informed Neural Networks  
434 can leverage radar (e.g., Sentinel-1 SAR), to estimate water levels for flood forecasting (Dasgupta et al., 2022; Gierszewska  
435 & Berezowski, 2024) or fused into predictive models that refine hazard forecasts for severe weather and anticipate cascading  
436 impacts (Flora et al., 2021). Remote sensing plays a crucial role in hazard dataset development by helping mitigate bias that  
437 may be inherited by ML-based risk models. These models are often trained on datasets calibrated with data from resource-rich  
438 regions, where the majority of weather stations are located. As a result, they may struggle to generalize effectively to  
439 underdeveloped areas, which are frequently the most vulnerable to extreme events (McGovern et al., 2019, 2022).

440 EO combined with ML is also used in assessing environmental quality, such as water quality (Sagan et al., 2020; Sit et al.,  
441 2020). These applications mainly showcase simpler models, such as short neural networks and SVM (Nazeer et al., 2017),  
442 Decision Trees (DT), RF, Cubist Regression and Extreme Gradient Boosting (XGBoost), due to their ease of implementation  
443 and relative scarcity of ground measurement data (J. Liu et al., 2023). They focus on optically parameters, such as chlorophyll-  
444 a, turbidity and suspended solids, but also others such as of nutrients and other non-optical parameter) can be predicted relying  
445 on models integrating meteorological and hydrological variables (S. Chen et al. 2022).

446 A central application of EO is in supporting impact and damage assessments: change detection techniques that compare pre-  
447 and post- event imagery are used to estimate physical impacts (T. Bai et al., 2023). This includes building damage (Y. Bai et  
448 al. 2018), infrastructure collapse (Sublime & Kalinicheva 2019) due to earthquakes or tsunamis (Ji et al. 2018), but also flood  
449 extent (Munawar et al., 2021), landslides (Lei et al., 2019) and wildfire scars (Bo et al., 2022; Tran et al., 2020). The main  
450 challenges encountered in these applications are due to the return periods of satellites, which may limit their ability to detect  
451 fast changing impacts; to the presence of clouds, which can hamper visibility especially during the occurrence of extreme  
452 events likely to cause damages; and to changes in luminosity or season (Faiza et al., 2012).

453 Moreover, EO enables long-term recovery tracking and vulnerability/exposure monitoring, with applications using proxies  
454 such as night-time lights to measure recovery trajectories (Kabiru et al., 2023; Qiang et al., 2020). For examples, studies have

455 used EO and ML to track how rapidly services return to urban slums post disaster, highlighting which population remain  
 456 exposed and underserved (Ghaffarian & Emtehani, 2021). Similarly, UNET-based CNNs are used to identify deprivation  
 457 pockets from satellite images and track during their recovery process (J. Wang et al., 2019), or to derive proxy indicators for  
 458 poverty from satellite night lights (Jean et al., 2016), in combination with transfer learning to overcome scarcity of labelled  
 459 data (S. J. Pan & Yang, 2010). At longer timescales, techniques like K-Nearest Neighbour (KNN), SVM, ANN and RF are  
 460 used to classify urban and rural land cover, detect land use changes or informal settlements (Adam et al., 2014; Yuh et al.,  
 461 2023; Zerrouki et al., 2019).

462 In summary, the integration of EO with ML and statistical techniques offers a powerful toolkit for multi-hazard and multi-risk  
 463 assessment, supporting early warning, targeted preparedness, rapid impact estimation, and recovery monitoring.

### 464 3.1.3 Textual data

465 In addition to remote sensing, textual data from sources such as social media and newspapers offer valuable information for  
 466 impact assessment. Natural Language Processing (NLP) algorithms can harness this textual data, facilitating applications  
 467 across various hazard types, including landslides, volcanoes, drought, earthquakes, floods, and wildfires. In general, the  
 468 procedure typically consists in several steps, in which the textual sources are first screened based on metadata (such as location  
 469 or the presence of disaster-related words in titles); then NLP or semantic algorithms (Angelov, 2020) are used to extract  
 470 keywords from the main text and convert the textual data into tabular/numeric; then a classification algorithm is applied to  
 471 choose between impact/no impact data or link the impacts to a specific sector or hazard. Additional steps may also involve the  
 472 retrieval of spatial information from textual data. Many different algorithms can be employed, with logistic/lasso regression  
 473 (Genkin et al., 2007), Naïve Bayes Classifiers (L. Jiang et al., 2016), KNNs (Shah et al., 2020) and ANNs (Nam et al., 2014),  
 474 being the most common. In the field of disaster mapping, SVM are tested by Asinthara et al. (2022), while Powers et al. (2023)  
 475 compares CNN and specific pre-trained language models; Koshy & Elango (2023) tests a multi-modal method leveraging text  
 476 and images from social media, employing the language models BERT; Mehrotra et al., (2022) test SVM, DT, RF, Adaboost,  
 477 Gradient Boosting, XGBoost, LSTM in combination with language models. Twitter (now X) was the main social media that  
 478 has been used to detect impacts, while newspaper articles have also been used, in particular for slow onset hazards, such as  
 479 droughts. For example, Sodge et al. (2023) apply several NLP and ML methods to automatize the detection of drought impacts  
 480 from newspaper articles; the procedure classifies impacts into 25 classes, based on the sector (e.g., forestry, livestock, forestry,  
 481 transport etc.) by using different Supervised ML models (Naïve Bayes, Lasso Regression, RF, ANN). In general, rule-based  
 482 methods are preferred to ML models when the number of samples is limited (X. Liu et al., 2018).

483

484 **Table 1. Data-related methods, gaps and opportunities.**

Section	Methods	Gaps	Opportunities
---------	---------	------	---------------

<b>3.1.1 Climate datasets</b>	LSTM for soil moisture & hydrology; RF for runoff/precipitation; ConvNPs for sparse sensors; ANNs for interpolation; CNNs, GANs, diffusion models for downscaling; GPs for uncertainty quantification	Struggles with sparse/irregular data; poor scalability (GPs); extremes misrepresented; limited uncertainty treatment	Hybrid ML–physics models; scalable probabilistic methods; better uncertainty quantification; generative models for ensembles
<b>3.1.2 Earth observations (EO)</b>	SVM, RF, LSTM for soil moisture; CNNs/autoencoders for land cover, impacts, disaster recovery; transfer learning for data-scarce regions; ML for water quality (RF, ANN, XGBoost)	Bias toward data-rich regions; gaps/clouds limit detection; false positives; weak multi-hazard integration	Robust models for missing/noisy data; near-real-time EO pipelines; integrate EO with socio-economic data; transfer learning for vulnerable regions
<b>3.1.3 Textual data</b>	NLP + ML (Naïve Bayes, RF, SVM, CNN, BERT, LSTM); multimodal (text + images); rule-based for small datasets	Few labelled datasets; language/cultural bias; imprecise spatial info; noisy social media inputs	Multilingual/transfer learning; improved geolocation extraction; integrate with EO/sensor data; robust methods for noisy/misinformation-prone data

485

486

487 This section contributes to the field of multi-hazard and multi-risk analysis by showing how ML applications to climate  
488 datasets, Earth observations, and textual data can overcome data sparsity and heterogeneity, thereby enabling the generation  
489 of more complete, high-resolution, and multi-source datasets that are essential for capturing hazard interactions and cascading  
490 risks.

## 491 3.2 Multi-hazard

### 492 3.2.1 Identify, classify and cluster

493 The initial step in conducting a comprehensive multi-risk assessment involves a thorough analysis of hazard factors, which is  
494 critical for effective climate risk evaluation and enhancing disaster preparedness. In this context, identifying various hazards,  
495 classifying them into distinct categories, and extracting their spatio-temporal footprints through clustering techniques are  
496 fundamental processes.

497 The identification of impacts from satellite images to discover hazard footprints, such as for landslides, earthquakes, floods  
498 was discussed in the previous section because it is mainly an image processing task, where the goal is to identify differences  
499 between two images. This section focuses on the identification of extreme events from climate datasets, which require specific

500 considerations on the typology of hazards and risk considered and is subject to different definitions and multiple interpretations.  
501 The most common approach to identify multiple hazards from climate datasets is to use thresholds to identify univariate  
502 extreme events and then combine them at a later stage into a multi-hazard database. In order to identify the thresholds, two  
503 methods are applied: empirical thresholds (e.g., defining a max temperature over which an event is considered extreme) or  
504 statistical thresholds (e.g., calculating a pixel-wise and/or day-wise percentile to identify events that exceeds a threshold that  
505 can vary spatially and temporally). Empirical thresholds are usually fine-tuned to link extreme events to impacts on specific  
506 sectors or local applications, and many applications focus on temperature extremes and health (Ray et al., 2021; X. Sun et al.,  
507 2014). Statistical thresholds are preferred when analysing global trends and merging multi-hazard extremes because they allow  
508 a more consistent and probabilistic robust comparison between different hazards. Percentiles can be easily adapted to model  
509 spatial and temporal variations in data and are ideal for global application that cover multiple landscapes where a unique  
510 empirical threshold cannot be univocally determined. For example, in Ionita et al. (2021), specific percentiles are used to  
511 identify heatwaves and drought from temperature and SPI indicators respectively, before applying Empirical Orthogonal  
512 Functions to investigate their drivers and their centre of actions over Europe; Similarly, Sutanto et al. (2020) is using  
513 percentiles to identify heatwaves, droughts and wildfires from temperature, soil moisture and Fire Weather Index (FWI),  
514 analysing spatial overlaps of the daily binary hazard maps to identify simultaneously occurrences of dry hazards and then  
515 investigating cascading events by looking at different combinations of hazard sequences. Claassen et al. (2023), proposes a  
516 methodology to identify multi-hazard events combining static footprints derived from the processing of satellite images (e.g.  
517 for landslides, floods, tsunamis) with dynamic footprints (based on statistical percentiles) of climate hazards (e.g., heatwaves,  
518 droughts, extreme precipitation, extreme wind, etc.), proposing a methodology to identify consecutive events using a specific  
519 time lag and analysing the global distribution of the various multi-hazard events.

520 Return periods are another statistical technique used to identify extreme events, studying the likelihood of an event of a certain  
521 magnitude occurring in a chosen timeframe (Liao et al., 2021). Return periods are most often applied in hydrology, when  
522 dealing with flooding and storm surge events (G. Liu et al., 2020, 2023; Mattei et al., 2021; Zanini et al., 2020). These  
523 applications fit a probability distribution (typically a Generalised Extreme Value Distribution, calculated over the number  
524 exceeding of a threshold or over maxima) which allow for an estimation of the uncertainty of the threshold. Percentile  
525 thresholds, returns periods and Generalised Extreme Value (GEV) distributions are also used conjunctly, such as in Orth et al.  
526 (2022), where different hydrological hazards (floods, frost, heat waves, droughts, and storms) and their contrasting impacts  
527 are analysed against multiple sectoral assessment endpoints (Gross Primary Productivity for vegetation, crop yields, human  
528 mortality, damages to properties and public attention).

529 It is important to note that these approaches focus initially on univariate extremes, and only at a second stage, the identified  
530 events are merged to produce multi-hazard events, checking for overlapping in time and space. This can lead to the  
531 underestimation of compound joint-extreme events which arise as a combination of multiple indicators not individually  
532 extreme.

533 Other approaches focus on identifying and classifying extreme events from climate reanalyses using DL, especially in case of  
534 cyclones or other hazards that are characterised by the interaction of multiple atmospheric drivers. Y. Liu et al. (2016) was one  
535 of the first to apply CNN based on AlexNet to detect and classify tropical cyclones, atmospheric rivers and weather fronts  
536 from climate datasets, such as ERA-5, CAM5.1. One of the main challenges in this domain is the scarcity of labelled data for  
537 training supervised ML models. This is discussed by Racah et al. (2016), who expanded the previous approach, developing a  
538 semi-supervised CNN model to overcome the lack of labelled data and created an extreme weather dataset as benchmark. In  
539 general, the skewness of datasets is another common challenge for identifying climate anomalies with supervised approaches:  
540 often data on which the ML models are trained on present very few samples of conditions leading to impacts (Dal Barco et al.,  
541 2024).

542 Other studies focus on the identification of the spatio-temporal footprints of the climate hazards, in particular with algorithms  
543 such as Density Based Spatial Clustering Applications with Noise (DBSCAN, Ester et al., 1996), grouping single point  
544 anomalies into clusters in time and space. These approaches are applied in single hazards, such as droughts (Cammalleri &  
545 Toreti, 2023), heatwaves (J. Wang & Yan, 2021) or earthquakes (Di Martino et al., 2018a). With regard to multi-hazards  
546 applications, DBSCAN is used by Tilloy et al. (2022) to cluster compound precipitation and wind compound extreme events  
547 in Great Britain and by (H. Yu et al., 2022) to investigate droughts, heatwaves, cold-waves, extreme wind and extreme  
548 precipitation in Eurasian Drylands, studying how the coordinates of the centroid of the clusters are shifting hot and dry events  
549 to northern latitudes due to climate change.

### 550 3.2.2 Hazard forecasting and prediction

551 Before delving into more risk-based applications, it is worth noting that in the last few years, the application of DL models  
552 such as Transformers (Vaswani et al., 2017), Graph Neural Networks (GNN) (Veličković et al., 2017) and Physics Informed  
553 Neural Networks (Kashinath et al., 2021b; Lütjens et al., 2021) has prompted a revolution in weather forecasting. Early  
554 applications of AI models, primarily using RF and SVM, were largely aimed at replacing specific steps within numerical  
555 weather forecasts. More recently, DL tools have gained prominence due to their ability to capture long-range dependencies,  
556 handle complex and irregular data structures and integrate the solutions of equations of physical systems into a unified  
557 framework, enabling DL to be successfully employed for modelling the whole medium range weather forecasting process (Bi  
558 et al., 2022; Chen et al., 2023; Keisler, 2022).

559 Applications that focus on predicting or forecasting hazards are still mainly focussed on single hazard approaches. However,  
560 some single hazard approaches were included in this review because their multi-variate approach includes the combination of  
561 different static (as land use, topography, socio-economic data) and dynamic (e.g., atmospheric and marine data) parameters  
562 and implicitly deal with multi-hazard interactions (e.g., a wildfire may be more probable when dry and hot conditions are  
563 present, a drought can be influenced by temperature and soil moisture, etc.). For example, Haggag et al. (2021) propose an  
564 ANN prediction model in a multi-hazard perspective, but then test it on past disaster records to predict floods in Ontario using

565 indices for climate extremes inputs. Monte Carlo dropout techniques have been employed to quantify epistemic uncertainty,  
566 for example in surge forecasts (Macdonald et al., 2025) and flood modelling (M. Nguyen et al., 2024).

567 One of the main algorithms applied to forecast hazards is LSTM: Kratzert, Klotz, Brandstetter, et al. (2019) apply adapted  
568 LSTM to disentangle static and dynamic inputs and analyse both high and low extremes in river flows, considering climate  
569 susceptibility and integrating static and dynamic inputs. Tiggeloven et al. (2021) propose a LSTM/CNN architecture to predict  
570 global storm surge residuals based on atmospheric conditions, investigating how the model's performance varied based on  
571 changes of the spatial area input into the convolutional model. With regard to vegetation, long-range temporal dependencies  
572 from several climate variables are investigated with a LSTM model (Kraft et al., 2019). Many applications focus on forecasting  
573 of air quality hazards, especially in urban areas: compared to other types of environmental impacts, such as water quality, the  
574 network of air quality monitoring stations offers hourly data at a high spatial resolution, enabling the training of AI models to  
575 dynamically forecast at short lead times. Applications include the short-term prediction of ozone levels in Kuwait (Freeman et  
576 al., 2018), the development of a daily air quality index in Beijing and Guilin (Q. Wu & Lin, 2019), or the prediction of  
577 concentration of micro particular matter in the air of Seoul (Chang-Hoi et al., 2021).

578 Another popular DL architecture is GNN, applied in weather forecasting (Keisler, 2022; Lam et al., 2022) and river  
579 networks/flooding predictions (Bentivoglio et al., 2023; Kazadi et al., 2024; A. Y. Sun et al., 2021). The key advantage of  
580 GNNs over CNNs is their ability to capture complex relationships in non-Euclidean data. While CNNs are limited by fixed  
581 sliding windows and may miss correlations between adjacent pixels or non-adjacent zones, GNNs excel in modelling graph-  
582 structured data, allowing for more accurate representations (Kipf & Welling, 2016). In particular, Kazadi et al. (2024) apply a  
583 combination of GNN and Gated Recurrent Unit (GRU, a type of recurrent neural network), for spatio-temporal flood  
584 prediction, accounting for spatially distributed precipitation data, as well as static features such as bathymetry and topography,  
585 comparing its performances against a LISFLOOD-FP simulation of Hurricane Harvey (2017) in Houston, Texas and showing  
586 improvements in terms of accuracy and faster training (100x) and testing (1000x) times. Similarly, Transformers are applied  
587 for river flood prediction, outperforming other RNNs in terms of computational costs and performances, also increasing the  
588 interpretability of the model (Castangia et al., 2023).

589 CNN, ANN, LSTM are still popular for drought and heat events, which are characterised by longer scale spatio-temporal  
590 dynamics. For example, Bonino et al. (2024) compare the performances of CNN, LSTM and RF for the prediction of marine  
591 heatwaves; Patil et al. (2023) employ CNN to predict drought in East Africa 3 or 4 season ahead, analysing the contribution  
592 of different climate drivers at multiple spatial and temporal scales; ANN are used for forecasting drought risk at near real time  
593 in India, using Artificial Neural Network models (Singh et al., 2021). Other algorithms (SVM, Random Forest, XGBoost,  
594 Extra Trees) are still often applied to analyse low probability extreme events in specific locations, where the lack of data  
595 constrains the training of Deep Neural Networks, such as the storm surge height caused by tropical cyclones in New York  
596 (Ayyad et al., 2022).

### 597 3.2.2 Modelling hazard interaction

598 Recent work has applied interpretable ML frameworks to hazard modelling, aiming not only at prediction but also at identifying  
599 key drivers. For instance, S. Jiang, Bevacqua, et al. (2022) and S. Jiang, Zheng, et al. (2022) used LSTMs to study river  
600 flooding in Europe, combining feature attribution methods such as Expected Gradients (Erion et al., 2021) and Additive  
601 Decomposition (Du et al., 2019) to disentangle the roles of snowmelt and precipitation. By running models across decades,  
602 they revealed shifts in dominant flood drivers, with precipitation becoming increasingly important. Other studies have applied  
603 gradient-based methods (A. Y. Sun et al., 2021), CNN heatmaps (Patil et al., 2023), attention mechanisms (Castangia et al.,  
604 2023), and sensitivity analysis (Bentivoglio et al., 2023; Bonino et al., 2024; Kratzert, Klotz, Shalev, et al., 2019). These  
605 advances improve interpretability, yet ML approaches remain limited by high data demands, sensitivity to training biases, and  
606 the difficulty of generalising beyond observed conditions (Bentivoglio et al., 2023). Their strength lies in prediction and  
607 uncovering nonlinear relationships, but the black-box nature of many models complicates causal modelling (Freeman et al.,  
608 2018).

609 While most ML studies focus on univariate hazards, compound events require methods that capture joint extremes. Copulas  
610 offer a flexible statistical framework to model dependence structures between variables, such as the co-occurrence of high  
611 river discharge, intense rainfall, and coastal surges (Hao & Singh, 2016; Nelsen, 2006). By decoupling marginal distributions  
612 from their dependence structure, copulas can assess joint probabilities of rare events with more precision than traditional  
613 multivariate models (Tilloy et al., 2019). Applications include pair copulas for compound flooding in Italy (Bevacqua et al.,  
614 2017a), Joe copulas for concurrent river–coastal extremes (Sadegh et al., 2017), and copula-based Bayesian networks for  
615 flood–drought interactions (Couasnon et al., 2018). However, several challenges remain: selecting appropriate copula families  
616 is non-trivial (since different families imply different tail dependencies, yet many common families assume simplistic  
617 dependency or exchangeability) (Oh & Patton, 2015); capturing joint tail dependence becomes increasingly difficult in high  
618 dimensions (vines, mixtures, or hierarchical copulas may help but bring computational and inference burdens) (Simpson et al.,  
619 2020); physical drivers (e.g. precipitation skew, changing climate forcings, watershed characteristics) are often only indirectly  
620 represented through marginal or covariate models (Hochrainer-Stigler et al., 2019b). Therefore, while copulas are powerful  
621 for probabilistic risk quantification, they are less suited to dynamic forecasting or process-based understanding without  
622 additional model structure or ensembles (Tootoonchi et al., 2022).

623 Comparison and complementarities.

624 ML and copula methods approach hazard interactions from distinct perspectives. ML excels at prediction and feature  
625 discovery but struggles with transparency and extrapolation, while copulas provide interpretable dependence structures and  
626 joint probability estimates but scale poorly with dimensionality and lack causal interpretability. ML can identify critical  
627 hazard predictors and generate inputs, while copulas rigorously quantify their joint occurrence. Yet, few studies combine  
628 these strengths; most rely on either predictive ML or probabilistic copulas in isolation. For example, an LSTM may forecast  
629 river discharge under given precipitation and snowmelt conditions, while a copula model can then quantify the probability

630 that extreme discharge co-occurs with extreme rainfall or sea-level rise. Together, ML and copulas can provide a more  
631 complete picture: ML enables forecasting and driver attribution, while copulas ensure rigorous treatment of dependence  
632 structures and joint extremes (Sadegh et al., 2017; Tilloy et al., 2019). Combining both approaches offers a promising  
633 pathway for advancing compound risk assessments. Some approaches, such as, T. Jiang et al., (2023) used a hybrid ML-  
634 copula method to estimate the probability of consecutive drought events (in particular from meteorological to ecological  
635 droughts), combining several ML classifiers (KNN, RF, SVM, ...) to estimate the propagation probability of meteorological  
636 drought given its characteristics, and C-vine copulas to model conditional probability model of the paired meteorological  
637 and ecological drought events. Closing this gap, for instance, by integrating ML-derived drivers into copula frameworks, or  
638 benchmarking ML-learned dependencies against copula-based models, represents a promising but underexplored direction  
639 for compound risk assessment.

#### 640 **Susceptibility mapping**

641 Susceptibility in the context of natural hazards refers to the predisposition of an area to experience a specific hazard and  
642 considers different factors (usually categorised into hazard or vulnerability in risk assessment), such as topography, geology,  
643 hydrology, land use and vegetation and highlights “territorial characteristics”, disregarding the more dynamic and time-  
644 dependent component of risks (Wubalem, 2022). The methodology for creating multi-hazard susceptibility maps using ML  
645 usually consists in three steps: first, for each hazard, the susceptibility factors are identified; then, supervised ML techniques  
646 are employed to create single hazards susceptibility maps, considering the different conditioning factors as predictors and the  
647 areas impacted by the analysed hazards in the past as assessment endpoints; finally, the single hazard maps are combined to  
648 produce the final multi-hazard susceptibility map. Eventually, feature importance techniques are applied as a fourth step to  
649 extract the most susceptible factors for each hazard or multi-hazard combination.

650 ML has been applied extensively to derive multi-hazard susceptibility maps, which can identify areas prone to multiple disaster  
651 and help disaster management planning. However, these applications are typically trained on average, static climatic conditions  
652 and do not consider temporal interactions between risk factors (such as the cumulative impacts of a series of successive extreme  
653 rain events, the duration of a heatwave or changes in vulnerability caused by wildfires). Moreover, the type of multi-hazard  
654 events for which they are applied is often limited to wildfires, landslides, floods, and earthquakes (Abu El-Magd et al., 2021;  
655 Ahmadi et al., 2021; Cao et al., 2020): in fact, these methods rely on the presence of catalogues of past clearly defined  
656 hazard spatial footprint: for other climate hazards, such as extreme winds, hails, or heatwaves susceptibility is not investigated.  
657 Furthermore, input data for susceptibility mapping are aggregated over long time frames, in order to ensure robustness of the  
658 analysis. However, changes in vulnerability and exposure parameters occurring in the analysed periods, for example due to  
659 newly implemented adaptation measures, are overlooked, potentially leading to overestimation (or underestimation) of areas  
660 at risks.

661 The most common approach for integrating susceptibility parameters into multi-risk assessment is by producing multi-hazard  
662 susceptibility mapping, where susceptibility to multiple hazard (including factors for hazard, such as yearly precipitation, but  
663 also vulnerability parameters, such as slope) can provide a valuable point of reference for decision makers in sustainable land-

664 use planning or infrastructure development. A number of studies are focusing on mountainous regions, using a range of ML  
665 models, including Logistic Regression, ANN, DT, SVM, RF, Boosted Regression Trees (BRT), or Generalised Linear Models  
666 (GLM) (Javidan et al., 2021; Karakas et al., 2023; Kariminejad et al., 2022; H. D. Nguyen et al., 2023; Pourghasemi et al.,  
667 2019, 2020; Pouyan et al., 2021; Yousefi et al., 2020) The multi-hazard combination usually covers floods, landslides,  
668 avalanches and forest fires, which have clear footprints that can be used to train single hazard susceptibility, and integrate other  
669 risks which can be assessed through already available risk maps, such as seismic risk maps at a later stage (Bordbar et al.,  
670 2022). Different hazards are included by Piao et al. (2022), who test BRT, RF and Classification And Regression Trees (CART)  
671 in the Gangwon-do region in South Korea (an area rich in forests and ecological diversity) to establish a multi-hazard  
672 probability map for forest fires and droughts; in this study the multi-hazard interactions are investigated, considering drought  
673 as an amplifying hazard for forest fires. Mandal et al. (2022) focus instead on coastal areas, in particular in West Bengal  
674 (India), considering tropical cyclones, embankment breaching, storm and tidal surge, inundations, extreme rainfall, salinization  
675 and erosion; RF and ANN are applied to produce multi-hazard susceptibility maps. Ullah et al. (2022) test a CNN to produce  
676 flash floods, landslides and debris flow multi-hazard susceptibility mapping, comparing its performances with Logistic  
677 Regression and KNN methods in terms of accuracy, coefficient of determination, Mean Absolute Error and Root Mean Squared  
678 Error. The input data consist of field surveys, topography, hydrology, and environmental data, while the locations of historical  
679 flash flood, debris flow and landslide locations are extracted from Google Earth images. The feature importance scores are  
680 derived using a Random Forest model and are used to enhance the analysis of the multi-hazard maps. It is interesting to note  
681 that in this case, the CNN layer is 1-dimensional and is not used to analyse the spatial context of the pixels, but it runs across  
682 the 14 layers of predicting variables, producing an independent output pixel by pixel.

683 While the literature on this topic is quite established, most of these applications propose a multi-layer single hazard risk, rather  
684 than a full multi-hazard or multi-risk approach: in fact, the single hazard maps are often combined linearly or via a matrix  
685 considering combined risk categories, without elaborating further on the hazard interactions. Another common challenge in  
686 the development of susceptibility maps is the skewness of the training dataset, which are characterized by a predominance of  
687 areas with no damage. These greatly affects the training and testing of the models, and specific sampling procedures are often  
688 applied, rather than relying on balancing weights when training the ML model. Most often, all the positive samples (e.g., where  
689 some impact was recorded) are included; a buffer area is applied to the positive samples and subtracted from the whole dataset  
690 to exclude areas near recorded impacts; a number of points of comparable magnitude to the positive ones is sampled from the  
691 difference dataset to ensure that the final training dataset includes a balanced representation of impacted and non-impacted  
692 areas. This is a key step of the susceptibility mapping and can potentially add biases to the model, if the selected samples are  
693 not representative of the whole dataset or if there is a high autocorrelation. Spatial or temporal autocorrelation needs to be  
694 considered when splitting between training, validation and test data: random splitting methods assume data is independent and  
695 identically distributed. Specific techniques, such as spatio-temporal block cross validation (Zanetti et al., 2022) need to be  
696 considered to account for this. For example, a recent paper by Sweet et al. (2023) shows the impact of different validation

697 techniques in a RF model for the prediction of agricultural yield, and their implications on performances and robustness of the  
 698 interpretation of the model.

699 **Table 2: Multi-hazard related methods, gaps and opportunities.**

Section	Methods	Gaps	Opportunities
<b>3.2.1 Identify, classify &amp; cluster</b>	Thresholding (empirical & percentiles) to build multi-hazard catalogs; return periods & GEV; EOFs; CNNs (semi-/supervised) for extreme-weather object detection in reanalyses; DBSCAN for spatio-temporal footprints and compound clusters.	Under-detection of joint (non-univariate) extremes when hazards are merged post-hoc; label scarcity & class imbalance for supervised DL; skewed datasets; sensitivity to spatial/temporal non-stationarity.	Unified pipelines that detect <b>compound</b> signatures directly (multivariate thresholds + clustering); semi-/self-supervised DL to mitigate label scarcity; robust cluster tracking of compound hotspots under change.
<b>3.2.2 Hazard forecasting &amp; prediction</b>	LSTM/CNN for hydrology, storm surge, drought/heat; Transformers for floods; GNN/GRU for river-network dynamics; classical ML (RF/SVM/XGB) for local extremes when data are limited.	High data demands; generalisation beyond observed regimes; limited interpretability; performance varies with spatial context and input windowing.	Physics-informed/graph-aware DL for better extrapolation; attention/attribution to expose drivers; global-to-local transfer learning; benchmarking vs. process models for trust.
<b>3.2.2 Modelling hazard interactions</b>	Copulas (pair/vine/Joe) for joint extremes; copula-BNs for river-coastal compounding; XAI on LSTMs/CNNs/Transformers (gradients, attention, sensitivity) to reveal drivers and shifts.	Copula family selection & tail-dependence in high dimensions; ML black-box limits causal insight; difficulty linking physical drivers to dependence structures.	<b>Hybrid ML-copula</b> stacks (ML to predict/characterise events, copulas to quantify joint probabilities); benchmarking ML-learned dependencies against copula baselines; conditional vines for cascades.
<b>Susceptibility mapping (multi-hazard)</b>	Supervised ML (LR/GLM, RF, SVM, BRT, CART, ANN, CNN) to build single-hazard susceptibility then combine into multi-hazard maps; feature importance to rank drivers.	Often “multi-layer single-hazard” (weak interaction modelling); training skew (few positives); sampling bias & autocorrelation; static inputs ignore evolving exposure/vulnerability.	Spatio-temporal CV (block) to curb leakage; dynamic susceptibility that updates with sequences/adaptation; explicit interaction terms or graph-based fusion; extend beyond the usual

Deleted: ¶

Formatted: Caption, Keep with next

Formatted Table

			quartet (fire/landslide/flood/quake).
--	--	--	------------------------------------------

701

702 This section contributes to the field of multi-hazard and multi-risk analysis by reviewing methods for identifying, classifying,  
703 and clustering hazard events from diverse datasets, highlighting how threshold-based approaches, clustering algorithms, deep  
704 learning models, and copulas can capture the spatio-temporal footprints and interactions of hazards, thereby advancing the  
705 ability to detect, forecast, and model compound and cascading events.

### 706 3.3 Multi-risk

#### 707 3.3.1 Modelling risk combining susceptibility, exposure and vulnerability

708 Many studies are found to focus on modelling risk by combining hazard maps produced via susceptibility mapping with ML  
709 and vulnerability and exposure layers. Single hazards such as wildfires, floods and landslides are the often considered, and  
710 buildings, population and infrastructures are the ~~Kotaridis & Lazaridou (2022) consider flooding risk in Tuscany and applied~~  
711 a 2D CNN to produce an urban flooding susceptibility map. Differently from Ullah et al. (2022) the CNN applied here makes  
712 use of the spatial context of each pixel, considering a 5x5 patch centred on a specific pixel (an area of 50 x 50 m<sup>2</sup> since the  
713 pixel size is 10m), creating 20000 different samples from the initial map, each one with a 5x5x9 size, where the last number  
714 corresponds to the different predictors of the susceptibility mapping that are considered as channels in the CNN architecture.  
715 Thus, not only the selection of the initial samples, but also the selection of the size of the patch is a key hyperparameter to be  
716 considered: in this case, a cross validation is used to choose the best patch size. The vulnerability maps are created dividing  
717 the land use into 5 classes, which are then multiplied with the hazard layer to calculate the final risk map. Convolutional Neural  
718 Networks (CNNs) offer significant advantages over traditional algorithms in spatial analysis due to their ability to process  
719 areas as 2D maps. This enables the model to leverage Max Pooling layers to capture and simplify the spatial context of events.  
720 Unlike models that focus on individual point characteristics, CNNs can better model and integrate the broader spatial  
721 relationships. For example, Zhao et al. (2020) test CNN for urban flood susceptibility too but instead of producing separate  
722 maps for hazard and vulnerability, anthropogenic factors were used as predictors for the susceptibility map. The study  
723 compares the performances of different ML models: a simple (with 1 convolutional layer) CNN architecture, LeNet5 (Lecun  
724 et al., 1998), a slightly deeper CNN (with 2 convolutional layers), SVM and RF models. Different input strategies are tested:  
725 a point based strategy that only considers input at a given site; a partial spatial strategy that considers the surrounding pixels,  
726 flattening the 2D image to a 1D vector, thus losing partially the spatial context, but allowing the neighbouring pixels to be  
727 fed to SVM and RF models as additional predictors; a patch strategy, similar to the one described before for the CNN models,  
728 which granted the best performances. This study also discusses the use of Deep CNNs, which is discouraged since the typical  
729 sample size and model is too small to tune the high number of parameters required by Deep CNNs.

Deleted: moKotaridis & Lazaridou (2022)

731 Rusk et al. (2022) analyse population risk in the Hindu-Kush and Himalaya region, producing a multi-hazard map for  
732 landslides, floods and wildfire with the MaxEnt (Maximum Entropy) algorithm, which is then overlaid with population  
733 distribution. The paper also produces a matrix of multi-hazard interactions, dividing them into three types: when hazards are  
734 directly linked (e.g., flooding causing a landslide), when their linkage is mediated by an environmental condition (e.g., land  
735 use changes caused by wildfires increasing the probability of a landslide), or when their linkage is mediated by infrastructure  
736 or urban processes (e.g., a landslide damaging a dam, triggering a flood). However, a quantitative assessment of these multi-  
737 hazard interactions is not provided and only the records of these events are used to complement the multi-risk map. A similar  
738 approach is used in Austria, (Fuchs et al., 2015) considering river flooding, torrential flooding and snow avalanches as hazards  
739 and buildings as assets. In this case, buildings vulnerability is investigated, categorising them based on location, size, building  
740 category and the construction period. The different urbanisation patterns, very high in mountainous terrain of the Hindu-Kush-  
741 Himalaya (HKH) and quite low for Austria, influenced the final risk score assessment, with the HKH showing more areas at  
742 higher risk (Rusk et al., 2022). Sammonds et al. (2023) analyse hurricane, flood and landslide risk on population, producing  
743 single hazard susceptibility maps with statistical methods and discussing the vulnerability of population, considering gender,  
744 age, and population density; the final multi-hazard hurricane risk is obtained as a product of the single hazard susceptibility  
745 scores, overlaid with weights determined with Analytic Hierarchy Process (AHP), and the vulnerability score. Other  
746 applications focus on Vietnam, where RF is applied to derive risk for buildings and population against multi-hazard  
747 susceptibility maps for floods and wildfires (Luu et al., 2024). RF is applied to calculate single and multi-hazard susceptibility  
748 maps for China for flooding, landslides, and debris flows and the railway infrastructure was overlaid to analyse present and  
749 future risk, considering newly planned railway links (K. Liu et al., 2018). In general, a number of studies are found to apply  
750 non-ML approaches, including multi-criteria decision-making and expert judgements methods to calculate susceptibility and  
751 vulnerability layers, such as in Arvin et al. (2023), that focuses on infrastructure resilience in Iran, considering flooding,  
752 landslides and earthquake as hazards, and 25 indicators at the county level and Khatakho et al. (2021), focussing on population  
753 exposed to flooding, earthquakes and wildfires near Kathmandu (Nepal).

754 A critical limitation of the studies reviewed in this section is the static treatment of vulnerability. Most applications use fixed  
755 proxies – building footprints, land-use classifications, census-derived population density – that do not evolve in response to  
756 hazard occurrence, adaptation measures, or broader socio-economic change (Haer et al., 2019, de Ruiter & van Loon, 2022).  
757 This static framing can substantially underestimate risk in contexts where vulnerability is shaped by governance failures,  
758 structural inequalities, or rapid urban expansion (Ward et al., 2022; Šakić Trogrlić et al., 2024). A particularly underexplored  
759 challenge in multi-hazard risk assessment is that vulnerabilities do not simply add up across hazards: they interact. Synergies  
760 and asynergies between vulnerabilities mean that the combination of hazards can fundamentally alter how exposed elements  
761 are affected. For instance, adaptation measures designed to reduce risk from one hazard may increase vulnerability to another,  
762 and damage caused by a first hazard event can leave a system more vulnerable to a subsequent one (Albulescu & Armas, 2024;  
763 de Ruiter & van Loon, 2022). Stolte et al. (2024) further demonstrate through a global systematic review of urban vulnerability  
764 that the drivers of vulnerability differ substantially across hazard types, and explicitly call for research into multi-hazard

765 vulnerability dynamics as a necessary step beyond the current dominant paradigm of treating multiple hazards in parallel rather  
766 than in interaction. Despite growing conceptual recognition of this problem, it remains essentially unaddressed in the data-  
767 driven literature reviewed in this study, where vulnerability interactions are neither modelled nor discussed. Social justice  
768 dimensions also remain largely absent from the reviewed multi-risk literature: only few of the papers analysed explicitly  
769 consider vulnerability dimensions such as gender, while the question of how ML-based risk maps might inherit biases from  
770 historically underinvested impact datasets remains largely unaddressed (McGovern et al., 2022).

771 Another aspect to consider is uncertainty and its propagation across the risk modelling chain: attempts to propagate it formally  
772 across the hazard–exposure–vulnerability–risk chain are rare even in single-hazard contexts: Kropf et al., (2022) introduced a  
773 sensitivity and uncertainty analysis framework within the CLIMADA platform that varies hazard, exposure, and vulnerability  
774 inputs simultaneously, and Dawkins et al. (2023) extended this to formally quantify uncertainty contributions from each  
775 component, with an application using GAM for heat-stress risk assessment, but neither study addresses multi-hazard  
776 interactions. However, no study in the reviewed corpus achieves end-to-end UQ in a multi-hazard risk context, propagating  
777 uncertainty from input data through hazard modelling and ML or statistical methods to the final risk estimate.

### 778 3.3.2 Modelling risk predicting impacts

779 Another popular approach to model multi-risk with ML is to use impacts as a proxy and training supervised ML models on  
780 past impacts. Examples of possible impacts are excess mortality for health risks, economic damages and monetary losses,  
781 number of emergency signals or specific environmental indicators, such as ecological status. With regard to ML methodology,  
782 approaches are similar to the ones applied for predicting hazard values, considering multiple predictors covering climate,  
783 topography, land use and anthropogenic factors, but the final assessment endpoint, impact data, is very different from typically  
784 hazard data, having a coarser resolution in time and space and resulting in much smaller datasets. Thus, most of the studies  
785 focus on simpler and more interpretable ML methods like ensemble methods, rather than the DL approaches which are popular  
786 for hazard prediction. Moreover, more attention is dedicated to the interpretation of the factors and the explainability of  
787 methods (Ghaffarian et al., 2023), with most applications presenting some form of feature importance analysis, either as a  
788 built-in feature of the model, such as for RF, or as a a-posteriori analysis with SHAP values. In this section, studies are grouped  
789 based on the sectors and type of impact considered, considering health, food security and crops, environmental quality &  
790 biodiversity, physical damages and economic losses.

#### 791 Health

792 Studies focussing on environmental-health risks often analyse the combination of heat and air quality stressors and use excess  
793 mortality as predicant variable. These applications aim at disentangling complex temporal patterns, consisting of a long-term  
794 trend, driven by multiple (and often unknown) factors, and short-term peaks, mainly driven by summer heatwaves; moreover,  
795 time-lags needs to be considered. Thus, statistical methods, such as Distributed Lag non-linear models have been widely  
796 applied (Gasparrini, 2014) to model exposure lag-response of mortality to environmental stressors. More recently, RF has been  
797 applied, analysing the role of humidity in urban mortality during heatwaves at the global scale (Guo et al., 2024) or predicting

**Deleted:** Compared to the publications focussing only on the multi-hazard aspect, the applications that also consider exposure and vulnerability do not test multiple ML algorithms, often rely on expert-based judgments or simple frequency analysis. Moreover, while these applications focus on multiple hazards, the analysis of vulnerability factors is often overlooked: in most cases, only exposure layers are used to produce risk maps. Even when vulnerability is explicitly considered, it is calculated only for single hazards and as a static parameter. Recent publications, such as de Ruyter & van Loon, (2022) highlight the importance of considering dynamic vulnerability factors, especially in multi-hazard and multi-risk contexts, where vulnerability can vary because of the changes cause on the system caused by the occurrence of the hazards of extreme events, or because of specific adaptation and responses. Even though the use of EO can help to inform the models of changes in the system (such as land cover changes due to wildfires and landslides, or new urbanisation patterns after reconstructions), these indicators are not yet integrated in multi-risk mappings.

**Deleted:** (Kropf et al. 2022)

**Deleted:** ¶

818 heat-stroke occurrence in China (Y. Wang et al., 2019), while SVM is applied for analysing previous diseases, population  
819 density and urbanisation (X. Wang et al., 2021). One of the most interesting papers, Boudreault et al., (2023) test 9 different  
820 ML, DL and statistical methods (such as Generalised Additive Models – GAMs) in the Metropolitan City of Montreal,  
821 considering weekly all-cause mortality as predictand and air temperature, humidity, wind, Particle Matter (PM) 2.5, Ozone  
822 (O<sub>3</sub>), Nitrogen Dioxide (NO<sub>2</sub>), Sulphur Dioxide (SO<sub>2</sub>), Carbon Monoxide (CO) as predictors. Among the methods tested,  
823 Tree based methods (RF, XGBoost) usually perform better overall, while statistical methods (and GAM in particular) are more  
824 accurate in predicting the mortality peaks; Deep Learning approaches, such as MLP and LSTM have instead the worst  
825 performances. This is partially explained by the limited size of the dataset and the inclusion of non-climate causes in the  
826 predictand, likely to cause overfitting in the DL models. Another study also focussing on Canada proposes an AI-based  
827 framework to extrapolate vulnerability from health-heat relationship: Côté et al. (2024) test this approach considering two  
828 steps: first, a model to predict daily mortality from mean temperature for 3 days, age, income and period of the year as  
829 predictors and then a second model predicting annual mortality over aggregated areas with specific socio-economic and  
830 environmental (air quality, vegetation, ...) characteristics. The model tested are AutoGluon (an automatic ML framework  
831 allowing to train and test ML models without expert knowledge<sup>2</sup>), GP and Deep Gaussian Process (Deep GP). The results  
832 shows that GP are able to model better the daily mortality trends, especially during extreme temperature, while AutoGluon is  
833 slightly better for the annual analysis. GP with non-linear (e.g., 5/2 Matern Kernel (Y. Pan et al., 2021)) are in fact able to  
834 better handle noise and small data samples (J. Wang, 2023), and their limit is their computational costs (M. Jiang et al., 2022);  
835 on the other hand, the more complex Deep GP handed the worst outcomes, highlighting the challenges in tuning more complex  
836 Deep GPs (Tazi et al., 2023). Other studies focus on predicting the influence of water quality parameters, such as turbidity, on  
837 the risk of cholera disease outbreaks in Indian Coastal municipalities using a RF predictor (Campbell et al., 2020).

### 838 **Food security and crops**

839 The second group of reviewed studies focus on the nexus between food production, food security and migrations. For instance,  
840 Busker et al., (2024) apply XGBoost to predict food insecurity in the Horn of Africa. This model, takes as input several factors,  
841 integrating climatological variables, biological hazards, food and fuel prices, macroeconomic indicators, conflicts and  
842 humanitarian assistance, aggregating data on the administrative units for which the assessment endpoint variable (food  
843 security) was available. The model is tested for its ability to predict the onset of crises up to 12 months in advance,  
844 demonstrating superior performance in agro-pastoral areas compared to croplands. SHAP values are employed to analyse the  
845 key risk drivers. The findings of this study highlight its potential application in operational early warning systems, such as  
846 FEWS NET.

847 Tàrraga et al. (2024) also investigate the dynamic relationships between droughts, conflicts and food security, focussing on  
848 their impact on population displacement. In this case, ML is not used to predict displacement, but causal discovery methods  
849 are tested to retrieve its drivers within Somalia from 2016 to 2023. In particular, Granger Causality and Peter and Clark

---

<sup>2</sup> <https://auto.gluon.ai/stable/index.html>

850 Momentary Conditional Independence (PCMRI) are tested to generate plausible causal graphs of drought displacement,  
851 showing limitations for Granger causality due to the high dimensionality and autocorrelation of the time series, while the  
852 PCMRI method is able to disentangle the intertwined vulnerabilities and different leading times connecting drought impacts,  
853 water and food security systems along with episodes of violent conflict. The reliability of the causal model depends on the  
854 quality of training data and several assumptions are required, such as causal sufficiency (i.e., all possible driving variables of  
855 drought displacement need to be considered in the analysis), no contemporaneous causal effects and causal stationarity. Note  
856 that although causal sufficiency is valid, the associations between the other variables (e.g., SPEI, market prices, fatalities) may  
857 be influenced by confounding factors rather than direct causality.

858 Different types of copulas (Normal, Student's t, Archimedean with different distributions) are tested to model risk by linking  
859 bivariate return periods of temperature and precipitation to crop yields, analysing the impact of dry and hot, dry and cold, wet  
860 and hot, wet and cold conditions (Zscheischler et al., 2017). Nested Archimedean copulas were used to model the tri-variate  
861 dependence between maximum temperature and spring precipitation on crop yields, estimating the impact differences between  
862 single and compound hazards, using combinations of heat and precipitation stress (Ribeiro et al., 2020).

#### 863 **Environmental quality and biodiversity**

864 Numerous studies focus directly on environmental impacts, such as the influence of land use and urban planning on water  
865 quality. For example, R. Wang et al., (2021) apply RF with SHAP values to model stream water quality and specific pollutants  
866 based on four different urban planning scenarios in Texas. The model allows to correlate urban sprawl to water quality  
867 degradation and was used to forecast environmental impacts under different urban development pattern scenarios. In Li et al.  
868 (2022) the ensemble model XGBoost is used to predict water quality in beach locations in lake Eyre, paired with SHAP for  
869 increased explainability. Other studies focus on ecosystem and biodiversity: for example, RF and Logistic regressions are  
870 tested to predict forest loss in Borneo from topographical and anthropogenic variables (distance to urban areas, population,  
871 etc.), highlighting the advantages of RF for modelling multi-scale spatial relationships between risk drivers (Cushman et al.,  
872 2017). Similarly, in Islam et al. (2021), the spatio-temporal dynamics of wetlands in Bangladesh and their negative effects on  
873 biodiversity are analysed using Decision trees, RF and SVM. RF and SVM are the best performing algorithms and in general,  
874 the papers highlighted the role of remote sensing, for mapping wetlands variations in time. Species distributions is also  
875 investigated, with many applications discussing the different spatial approaches for river network modelling. For example,  
876 Schmidt et al. (2020) test the MaxEnt algorithm with two representations of rivers, highlighting how a high-resolution model  
877 based on river reaches is better at discovering individual local habitat features, whereas lower resolution sub-catchment scale  
878 models better account for more general drivers in fish distribution. Teichert et al. (2016) apply a RF model to identify the  
879 dominant stressors for fish presence in estuaries, investigating the interactions among stressors evaluating ecological benefits  
880 expected from reducing pressure. In particular, an RF model is trained to predict ecological status in 90 locations using 17  
881 predictors describing the different stressors (urbanisation, flow changes, water pollution, oxygen depletion, etc.). Then,  
882 simulations are run to analyse the benefit of restorations comparing the difference between the baseline model and a model


883 where the intensity of stressors was varied. The difference between single and multiple restoration action is analysed,  
 884 highlighting the importance of combined restoration schemes and the non-linearity of their effects.

885 **Economic losses and physical damages**

886 This final category focus on studies modelling economic losses or physical impacts: Dal Barco et al. (2024) model the  
 887 occurrence of impacts due to extreme weather events in the Veneto coastal municipalities, with a combination of two ML  
 888 models: first a classifier (RF, SVM, ANN) is trained to predict the probability of daily impacts in coastal municipalities using  
 889 meteorological data as predictors and a Boolean variable based on impact reports from the Regional Authorities as predictand;  
 890 then a Linear Regression is used to predict the yearly occurrences of damages based on the outcome of the first model.  
 891 However, the coarse resolution of the impact data, the biases in human collected impact catalogues, and the skewedness of the  
 892 dataset can pose significant challenges to the training of a ML-model predicting direct physical impacts. Other studies focus  
 893 on modelling tropical cyclones along the East Coast of the US with ANN: Pilkington & Mahmoud (2017) investigate the  
 894 complex connections between all meteorological factors (wind, pressure, storm surge, and precipitation resulting in inland  
 895 flooding) of a tropical cyclone and how those interact with the location of landfalls to produce a certain level of economic  
 896 damage. The vulnerability and resilience of the different coastal locations are investigated essentially using the model to predict  
 897 losses with varying meteorological factors taken from past historical events but switching their landfall location. Other  
 898 approaches, such as Mukherjee et al. (2018) test SVM and RF to analyse impacts on the energy sector in the US caused by  
 899 extreme weather events, leveraging the records of disruptions from outage data of the Department of Energy in the US and  
 900 using as predictors a set of climatic and socio-economic variables aggregated at state level. In this study, two different models  
 901 are trained, in order to account for the differences in the risk drivers between the more frequent energy disruptions and the  
 902 extreme events, which are separated based on their quantile. Finally, other studies focus on the impacts on specific economic  
 903 sectors, such as finance and tourism: Carannante et al. (2024) propose a pricing model for climate change risk, particularly  
 904 physical risk, developing a type of climate risk-insured loan, based on a bioclimatic composite indicator developed with ML.  
 905 In particular, a temporal dynamic RF (considering variables at different lag-times) is used to produce a monthly risk index,  
 906 based on atmospheric variables (wind, precipitation, temperature) obtained mainly from remote sensing datasets, which is used  
 907 to model impacts on beach resorts in Italy and inform the subsequent climate-risk loan mechanism.

908 **Table 3: Multi risk related methods, gaps and opportunities.**

Section	Methods	Gaps	Opportunities
<b>3.3.1 Risk via susceptibility + exposure + vulnerability</b>	Overlay of single-hazard susceptibility (RF, SVM, ANN, BRT, CART, MaxEnt, CNN with patch context) with exposure (buildings, population,	Often “multi-layer single-hazard” (weak/implicit interaction modelling); vulnerability treated as static; label imbalance and sampling bias; spatial/temporal autocorrelation leakage; limited	Dynamic vulnerability/exposure updates using EO and time-sequenced hazards; spatio-temporal block cross-validation; interaction-aware fusion (graphs, learned weights); extend to wind, hail, heat,

Deleted: 

Formatted: Caption, Keep with next

Formatted Table

	infrastructure) and simple vulnerability layers; AHP/MCDM weighting; feature importance/SHAP to rank drivers.	hazard breadth beyond fire/landslide/flood/quake.	surge; probabilistic risk surfaces with uncertainty bands.
<b>3.3.2 Predicting impacts — Health</b>	Ensemble and hybrid ML approaches (RF, XGBoost, SVM, DL, copulas, causal ML) applied to health, food, environmental, and economic impacts; explainable AI (SHAP) and probabilistic modeling for driver attribution.	Impact labels are sparse, coarse, biased, and confounded; scale mismatches and aggregation blur signals; extremes and tails poorly represented; DL tends to overfit and transfer poorly across cities/regions/climates; uncertainty quantification and causal attribution often limited.	(1) <b>Data &amp; catalogs:</b> build geocoded, event-level, cross-sector impact datasets and standardized labels (health, yields, biodiversity, losses). (2) <b>Causal &amp; lag-aware stacks:</b> combine DLNM / explicit-lag models with ML and causal discovery to capture delayed and causal pathways. (3) <b>Multi-source fusion &amp; transfer:</b> integrate EO, in-situ, socio-economic and market data; use domain-adaptation/transfer learning for cross-region generalization.

910

911 This section contributes to the field of multi-hazard and multi-risk analysis by examining how ML and statistical approaches  
912 combine hazard, exposure, and vulnerability layers or directly predict impacts, highlighting opportunities and challenges in  
913 capturing dynamic vulnerability, addressing data limitations, and improving the interpretability of risk models across health,  
914 food security, environmental, economic, and infrastructure sectors.

915 **3.4 Future**

916 **3.4.1 Predicting future hazards**

917 Several studies focus on data-driven applications to predict long-term future multi-hazard and multi-risk scenarios.  
918 Zscheischler et al. (2018) discuss the importance of compound events for future risk assessment and presents several  
919 approaches and discusses the main challenges related to the use of future climate projections and weather simulations to analyse  
920 future compound events. The role of bias correction and its connection to multi-hazard events and impact models is analysed:  
921 future projections are often bias corrected to align the distribution of the modelled variables to the distribution of the observed  
922 ones, in the reference timeframe. However, some issues can arise: the simplest approaches focus on adjusting the averages of

923 the variables and do not correct the tails of the distributions, thus modifying the behaviour of extreme events. Methods such  
924 as quantile mapping, are needed to align the historical and future datasets before the application of any statistical or ML  
925 methods. Sensitivity analysis can be performed to analyse how the model reacts to changes in inputs and the robustness of  
926 future scenarios (Kim et al., 2023). Moreover, bias corrections are often univariate, and do not consider the effects on joint tail  
927 distributions and consequently impact models based on these inputs are affected; multivariate bias correction models are then  
928 encouraged (Sippel et al., 2016).

929 When dealing with the future of multi-hazard events, statistical methods are most often applied to identify hotspots and test  
930 trends, similarly to the applications focussing on historical data. For example, Ridder et al. (2022) consider hot, dry, wet and  
931 windy compound events by selecting cells which exceed the 99<sup>th</sup> percentile for wind and precipitation in the same day. Then  
932 results are presented in changes in return period and annual event density, where the latter is a measure for how often an event  
933 affects a region and how much of the region is affected, calculated from the number of grid cells affected. Similarly, Zhu et al.  
934 (2023) investigate future compound wind and precipitation extreme at the global scale, analysing 14 CMIP6 models,  
935 identifying compound events through the 95<sup>th</sup> percentile and discussing the sources of uncertainties via the HS09 statistical  
936 method (Hawkins & Sutton, 2009) splitting between internal variability, model uncertainty and scenario uncertainty. Further  
937 analyses discuss the spatial and temporal performances of future projections: Ridder et al., (2021) find good performances in  
938 CMIP6 simulation for precipitation and wind compound extremes over North America, Europe and Asia, but poor  
939 performances over Australia, probably linked to the limits in the modelling of tropical and extratropical cyclones and local  
940 convection systems. Also, copulas are used to analyse spatial complementary patterns of compound events, such as in Ghanbari  
941 et al. (2021), which analyse the joint return period of compound floods along the US coast, incorporating sea level rise and  
942 peak river flows for future climate change risk scenarios with copulas. H. Wu et al., (2023, 2024), employ Vine copulas to  
943 analyse hot & dry and pluvial & hot events in future scenarios, using the Single Model Initial Conditions Large Ensemble  
944 (SMILE) approach.

945 Bevacqua et al., (2023) stress the importance of SMILE for a robust analysis of future compound climate events. In fact, a  
946 SMILE consists of many simulations from a single climate model, each starting from slightly different initial states (differently  
947 from classical model ensembles, like CMIP6, which consists of many different runs from different models). Each realization  
948 differs solely due to internal climate variability and ensures a better quantification of future uncertainties, and at the same time  
949 it provides a much larger dataset to analyse statistically compound events. Multiple SMILEs can then be combined to identify  
950 model differences and distinguish between internal climate variability and structural model differences. Sometimes, especially  
951 when dealing with unprecedented, High-Impact, Low-Probability events, climate projections or even SMILE or statistical  
952 weather generation are not sufficient: in these cases, storyline approaches are often used as alternative to explore future multi-  
953 risk patterns (Moezzi et al., 2017; Shepherd et al., 2018). These approaches fit well within common practices in disaster risk  
954 management, which consider event-based scenarios for emergency preparedness, allowing for interaction with local  
955 stakeholders to evaluate the effectiveness of selected measures (Sillmann et al., 2021) and to explore low-likelihood and high  
956 impact plausibility events (Bevacqua et al., 2021).

### 957 3.4.2 Modelling future impacts

958 A common approach to estimate future risks involves using future climate projections as input data for ML models that have  
959 been trained on historical data of past impacts, similar to applications that focus on assessing current risks by leveraging past  
960 impacts. For example, the study of future cyclone impacts in New York and New Jersey, is feeding four General Circulation  
961 models as input for a SVM / AdaBoost risk model (Ayyad et al. Park & Lee (2020)rk & Lee (2020) test the performances of  
962 three algorithms, K-NN, RF and SVM to analyse coastal risks in South Korea, considering rainfall, tides, topography and land  
963 use, training the model on past floodings and then predicting future risks using monthly averages of rainfall and tidal values  
964 from RCP 4.5 and 8.5 ensembles. Future risk scenarios are calculated aggregating the risk model outcomes for each decade  
965 from 2030s to 2080s. In a successive publication, Park et al. (2023) apply a similar ML methodology to investigate adaptation  
966 strategies for coastal flooding: in this case, the ML model is trained on historical data with two different adaptation strategies,  
967 seawalls or green spaces, and then the future adaptation models are implemented, either maintaining current adaptation  
968 infrastructures or increasing one specific strategy. To ensure comparability between the adaptation scenarios, infrastructure  
969 construction costs are standardized, guaranteeing that the two distinct adaptation pathways incurred equal expenses.

970 In general, it is considered good practice to use ensemble projections and values calculated over multiple years, in order to  
971 increase the robustness of the future scenarios; however, some risk analyses focus on just a few selected years: Lim & Kim  
972 (2022) test RF for future rainfall induced landslides, also analysing different adaptation pathways and considering an increase  
973 in forested or urban areas. Instead of using monthly or daily values for the ML model, yearly values are used in the model, for  
974 specific years (2050, 2092), which are considered significant for representing future scenarios. This approach is valuable for  
975 analysing specific extreme events that may be overlooked when averaging across multiple models or years, and it reduces  
976 computational demands. However, it carries the risk of biasing the analysis, as the selection of specific years may result in  
977 outcomes that are not fully representative of the broader range of future scenarios. Bayesian Networks were tested by Pham et  
978 al., (2023) in a multi-model chain approach combining ocean hydrodynamics models, wind-wave models, and shoreline  
979 extraction models to analyse sea water quality impacts and shoreline erosion under different RCP projections (4.5 and 8.5).  
980 Bayesian Networks are applied due to their ability to integrate heterogeneous data sources, including quantitative and  
981 qualitative inputs and several data fusion steps to harmonise different spatial coverage, temporal resolutions and data formats,  
982 with a final risk assessment conducted at municipality level and yearly/ decadal scale.

983 With regard to the water-food nexus, ML is being progressively employed as an alternative to process or statistical methods  
984 for future crop yield estimation, showing increased performances and higher computational efficiency: Leng & Hall (2020)  
985 test a RF model for annual yield prediction in the US for a 2° C global warming scenario; while Khan et al. (2024) select  
986 Gradient Boosting to model the relationships between daily climate variables, hazard indicators, such as Consecutive dry days  
987 (CDD) and crop production with CMIP6 data. Tabari & Willems (2023) carry out a global risk assessment from hot and dry  
988 events, employing Copulas and integrating data from Shared Socio-economic Pathways (SSP) scenarios, future land use  
989 patterns population and governance. ML methods are used also to predict the risk of increased conflicts due to climate stressors:

990 a RF classifier is applied by Hoch et al., (2021) to predict water-related conflicts in Africa using different SSP future  
 991 projections, integrating socio-economic predictors (population, education, GDP, governance) and climate predictors  
 992 (precipitation, evaporation, flood volume, soil water). The model is trained on historical data up to 2015 and tested with  
 993 projections from 2016 to 2050. Future temperature-related mortality in different European regions is analysed by García-León  
 994 et al., (2024) considering 4 scenarios of global warming (1.5 °C, 2°C, 3°C, 4°C) with an ensemble of CMIP5 models, analysing  
 995 disparities between cold-related deaths and heat-related deaths and analysing the role of age, health infrastructure and climate  
 996 change with a Distributed Lag Non-Linear model. In particular, different scenarios are discussed: present climate and present  
 997 population, present climate with future population from EUROPOP 2019; future climate under different warming level with  
 998 future population exposure.

999 Future risk patterns are also calculated implementing future multi-hazard susceptibility maps: for example, Rahman et al.,  
 1000 (2024) analyse future coastal multi-hazard risks in Bangladesh, implementing an LSTM algorithm, in combination with RF  
 1001 feature selection and a Genetic Algorithm (GA) optimiser. In particular, GA is used to identify optimal or near-optimal  
 1002 solutions, searching the space of LSTM parameters through a process of selection, crossover and mutation. The combination  
 1003 of the LSTM's ability to capture sequential patterns and long-term dependencies and GA's efficiency in navigating complex  
 1004 search spaces, is proved to achieve better convergence, avoid local minima, and optimise both the architecture and parameters  
 1005 of the LSTM model (Zamani et al., 2022). Other future multi-hazard susceptibility approaches include Ya et al., (2023), who  
 1006 analysed future risks in the Tibetan plateau considering climate and land use changes. Logistic Regression is used to produce  
 1007 susceptibility maps, while future climate scenarios were taken from CMIP6 future projections. In order to create future land  
 1008 use, this paper focus on PLUS, a RF-based model analysing the relationship between influencing factors and land use changes  
 1009 (Liang et al., 2021). Another approach for future land use is applied by Saha et al., (2021), which focus on modelling cultural  
 1010 heritage site future multi-hazard susceptibility in the Sikkim state in India, considering different climate scenarios from CMIP5  
 1011 and land use from an empirical model (Dyna-CLUE) incorporating spatial logistic regression (W. Jiang et al., 2015). Bayesian  
 1012 Additive Regression Trees and Bayesian Generalised Linear models are applied to produce multi-hazard susceptibility maps,  
 1013 considering extreme rainfall, landslides and earthquakes. Another dynamical model, a Cellular Automata- Markov model  
 1014 (Clarke et al., 1997) is used to predict future land use changes in Iran to investigate flood risks, testing RF, XGBoost and  
 1015 Gradient Boosting as algorithms for producing susceptibility maps (Janizadeh et al., 2021).

1016 **Table 4: Future related methods, gaps and opportunities.**

Section	Methods	Gaps	Opportunities
<b>3.4.1 Predicting future hazards</b>	Bias correction for projections (incl. <b>quantile mapping</b> ); hotspot/trend detection via percentile thresholds (e.g., 95th–99th), return periods; uncertainty sources and propagation;	Univariate bias correction can distort extremes/joint tails; regional skill varies; limited direct detection of <b>compound</b>	Adopt <b>multivariate</b> bias correction; combine SMILEs to separate internal variability vs. structural model differences; scale up <b>vine</b> copulas for

Deleted: ¶

Formatted: Caption, Keep with next

Formatted Table

	<b>vine copulas</b> for joint tails; <b>SMILE</b> large ensembles; <b>storyline</b> event-based scenarios for HILP analysis.	signals; uncertainty treatment often partial.	compound events; embed <b>storylines</b> for preparedness.
<b>3.4.2 Modelling future impacts</b>	Trained on historical <b>impacts</b> and applied to future ensembles; ensemble ML methods (RF, XGBoost, ...) for coastal risk, conflict risks, crop yield and adaptation scenarios; <b>Bayesian Networks</b> for multi-model chains (hydrodynamics–waves–shoreline); distributed-lag models for future health impacts; future susceptibility integrating land use changes	Impact data often coarse, biased, and sparse; studies often rely on <b>few years</b> → low representativeness; causal discovery hinges on strong assumptions; biases due to scale mismatch in climate–exposure–impact data.	Use <b>multi-model, multi-year ensembles</b> and probabilistic models to quantify and propagate uncertainty.; evaluate <b>adaptation portfolios</b> with standardized costs; integrate dynamic <b>land use + socio-economic</b> pathways (SSPs); uncertainty-aware learning, and open, geocoded impact datasets.

1018

1019 This section contributes to the field of multi-hazard and multi-risk analysis by reviewing how ML and statistical methods are  
1020 applied to predict future hazards and impacts, highlighting the importance of bias correction, ensemble and SMILE approaches,  
1021 and storyline methods, as well as the integration of socio-economic and land use projections. It emphasizes how these  
1022 approaches can improve the robustness of long-term risk scenarios, support adaptation planning, and guide strategies to address  
1023 uncertainties in future multi-risk patterns.

1024 **3.5 Limitations and future research directions**

1025 Beyond the methodological advances documented in the preceding sections, this review also identifies a set of persistent  
1026 limitations and structural gaps in the field that are directly relevant to the operational uptake of data-driven approaches in  
1027 multi-hazard risk management. A first and fundamental limitation is climate non-stationarity, as highlighted in Section 3.4  
1028 Future: ML models trained on historical hazard records implicitly assume that the statistical relationships between predictors  
1029 and outcomes will remain stable into the future. As Reichstein et al. (2025) argue in the context of early warning systems,  
1030 relying on past norms and training distributions will prove inappropriate under non-stationary risk conditions, where projected  
1031 increases in hazard frequency and severity, combined with shifting exposure and vulnerability, create conditions that fall  
1032 outside the range of any historical training set. This is particularly acute for compound and cascading events, which are by  
1033 definition rare in the historical record yet are precisely the configurations that climate change is projected to intensify.  
1034 A second limitation is the gap between hazard prediction and impact prediction. Most ML applications reviewed optimise for  
1035 hazard or susceptibility metrics, but impact prediction requires integrating physical hazard outputs with dynamic exposure and  
1036 vulnerability data at sub-kilometre scales, a challenge that the reviewed papers largely sidestep by using static proxies. This  
1037 gap between technical model performance and actionable risk information represents one of the most important unresolved

Formatted: Font color: Auto

Formatted: Heading 2

038 challenges in translating ML-based risk assessment into operational decision-making (Tiggeloven et al., 2025, Reichstein et  
039 al., 2025).

040 A third set of concerns relates to interpretability and trust. The black-box nature of deep learning models creates well-  
041 recognised barriers to adoption in high-stakes regulatory and emergency management contexts, where stakeholders need not  
042 only a prediction but a justification they can interrogate and contest. However, current XAI applications remain predominantly  
043 proof-of-concept and are rarely integrated into operational early warning or risk assessment workflows (Ghaffarian et al.,  
044 2023). Moreover, reproducibility and validation remain a persistent concern. In geoscientific applications, spatial  
045 autocorrelation means that random train-test splits routinely inflate apparent model skill relative to genuinely independent  
046 spatial holdouts (Sweet et al., 2023), and the reviewed literature shows limited adoption of spatially blocked cross-validation  
047 or independent regional test sets. These limitations do not invalidate the contributions reviewed here, but they do underscore  
048 the need for more rigorous validation protocols, realistic appraisal of out-of-sample performance, and explicit discussion of  
049 the conditions under which ML approaches can be expected to generalise beyond their training contexts.

050 Another methodological gap identified by this review is the absence of end-to-end uncertainty quantification frameworks for  
051 multi-hazard risk assessment. Current practice addresses UQ in fragments: aleatory uncertainty in input data is handled at the  
052 start of the chain, epistemic uncertainty in ML models is occasionally addressed through Bayesian or ensemble methods at the  
053 hazard stage, and copula-based approaches characterise joint uncertainty for statistically correlated hazard pairs, but these  
054 efforts are rarely connected, and they do not extend to the full multi-hazard concept, which encompasses cascading and  
055 triggered hazards beyond the reach of shared statistical distributions. A genuinely integrated framework would propagate both  
056 aleatory and epistemic uncertainty continuously from input data through multi-hazard interactions, ML and statistical model  
057 outputs, and exposure and vulnerability components, to the final risk estimate (Beven et al., 2018).

058 A further methodological consideration for future research is the development of data-driven frameworks that move beyond  
059 static representations of vulnerability and exposure. The reviewed literature overwhelmingly treats these components as fixed  
060 spatial layers, with limited engagement with their dynamic, socially differentiated, and governance-mediated dimensions.  
061 Addressing this gap will require closer integration of ML and statistical methods with approaches capable of representing how  
062 vulnerability evolves over time, including agent-based modelling, participatory data collection, and socially-informed  
063 frameworks that explicitly account for adaptive behaviour, equity, and governance processes. Progress in this direction would  
064 not only improve the realism of multi-risk assessments but also strengthen their relevance for policy and decision-making in  
065 contexts where social vulnerability is itself a driver of risk (Cannon 2017; Bankoff and Hilhorst 2022).

066 Finally, the geographical distribution of the reviewed studies, visualised in the Sankey diagram in Appendix B, points to an  
067 imbalance that is worth acknowledging explicitly. Europe, North America and East Asia together account for more than 80%  
068 of lead authorships, while Africa and South America contribute less than 5%. This pattern partly reflects the Scopus, English-  
069 only, 2010–2024 scope of the search strategy, which may systematically underrepresent research published in other languages  
070 or in regional journals not indexed by Scopus. At the same time, the decoupling between authorship geography and case study  
071 geography, with South/SE Asia, the Middle East, and Africa appearing more frequently as study areas than as sources of

1072 authorship, suggests that data-driven methods are in several cases developed in data-rich institutional contexts and  
1073 subsequently applied to regions with different risk dynamics, data availability, and governance structures (Tiggeloven et al.,  
1074 2025). While drawing strong conclusions about data colonialism or algorithmic bias from a bibliometric analysis alone would  
1075 go beyond the scope of this review, these patterns do raise questions that the community should engage with: whether training  
1076 datasets and validation benchmarks are representative of the contexts in which models are ultimately deployed, and whether  
1077 the priorities shaping methodological innovation reflect the needs of the most exposed populations. Future work in this area  
1078 should pay closer attention to the transferability of data-driven multi-risk frameworks across different socio-economic and  
1079 data environments (Tiggeloven et al., 2025), and collaborative initiatives fostering locally-grounded research in currently  
1080 underrepresented regions would strengthen both the scientific robustness and the equity dimensions of the field (Naudé &  
1081 Viluesa, 2021).

Formatted: Font: Font color: Auto

#### 1082 **4 Conclusion**

1083 This paper presents a comprehensive review of data-driven applications aimed at modelling and enhancing our understanding  
1084 of climate-related multi-hazard and multi-risk events. Based on the selection of over 1,400 studies and an in-depth analysis of  
1085 136 key papers, the review addresses four research areas: (i) data processing and collection, (ii) hazard analysis, (iii) risk  
1086 analysis, and (iv) future risk scenarios, each divided in several sub-topics. summarises the main methods used in each research  
1087 question, illustrating the different approaches for each sub-topic. In particular, the figure highlights the strong connections  
1088 between Earth observations processing and ML techniques like CNN; on the other hand, RF, other ensemble methods and  
1089 GAM are mostly applied for risk impacts and future risk assessment, while LSTM, ANN and other DL approaches are most  
1090 common for hazard prediction, reflecting a growing trend toward leveraging sophisticated AI architectures for climate and  
1091 hazards applications, and a focus on simpler, more interpretable models for risk applications. Despite the current prevalence  
1092 of single-hazard applications in ML research, there is growing recognition of the importance of multi-risk strategies. Notable  
1093 advancements include copula-based compound event analyses and ML-driven multi-hazard susceptibility maps. Future  
1094 research should prioritize a more comprehensive understanding of multi-risk interactions – such as triggering, cascading, or  
1095 amplifying effects – by considering the interplay between hazard factors, vulnerability, and exposure dynamics, which are  
1096 often overlooked or treated independently in current studies. DL methods, with their capacity to capture complex, non-linear  
1097 interactions across spatio-temporal dimensions, offer promising avenues for progress, yet remain underexplored in operational  
1098 multi-risk contexts. However, these methods require high-resolution impact data, which remains a significant challenge due  
1099 to limited availability, inconsistency across regions, and issues of data quality and standardization. While EO and textual data  
1100 can aid in generating new multi-risk disaster catalogues, traditional sensor-based and human-curated disaster catalogues remain  
1101 essential for validation, representing a major bottleneck that constrains model validation, transferability, and ultimately the  
1102 uptake of these methods in practice. By addressing these methodological and data gaps, the field can move toward more robust,

Field Code Changed

1103 interpretable, and actionable multi-risk assessments, ultimately strengthening the integration of machine learning into climate  
 1104 services that support adaptation, resilience, and disaster risk reduction.

1105 The gap between hazard prediction and impact prediction remains largely unresolved and bridging it will demand closer  
 1106 integration of data driven model outputs with dynamic representations of exposure and vulnerability, including human  
 1107 behaviour, adaptive responses, and the social and governance dimensions that determine how risk is distributed across  
 1108 communities. Explainability is a further priority: XAI methods need to move beyond their current role as exploratory tools and  
 1109 be embedded into operational early warning and risk assessment workflows, where their ability to illuminate driver interactions  
 1110 and build stakeholder trust is most consequential. End-to-end uncertainty quantification across the full modelling chain remains  
 1111 absent and developing integrated frameworks that propagate both aleatory and epistemic uncertainty from inputs through  
 1112 multi-hazard interactions to the final risk estimate is one of the most important open methodological challenges for the field.  
 1113 Underlying all of these challenges is the problem of non-stationarity: as climate change intensifies hazard frequency and  
 1114 severity, shifts exposure and vulnerability, and increases the likelihood of compound and cascading configurations that fall  
 1115 outside any historical training set, the assumption that past conditions are a reliable guide to future risk becomes increasingly  
 1116 untenable, with direct consequences for the validity of ML-based projections of multi-risk evolution.  
 1117 Addressing these gaps, alongside the geographic and equity imbalances documented in this review, will require not only  
 1118 methodological innovation but also more inclusive research practices: collaborative frameworks that bring together physical  
 1119 scientists, social scientists, and communities in currently underrepresented regions, co-producing knowledge that is robust,  
 1120 transferable, and genuinely relevant to those most exposed to the evolving risks of a changing climate.

1121 ▼

1122 **Appendix A: Abbreviations**

1123 **Table A1: Acronyms of methods (in alphabetical order)**

Acronym	Full Name
AI	Artificial Intelligence
ANN	Artificial Neural Network
BRT	Boosted Regression Trees
CART	Classification and Regression Trees
CNN	Convolutional Neural Network
ConvNP	Convolutional Neural Process
DBSCAN	Density Based Spatial Clustering Application with Noise
DeepGP	Deep Gaussian Process
DL	Deep Learning
DT	Decision Tree

**Deleted:** Finally, this review highlights the importance of adopting a multidisciplinary approach that combines expertise in multivariate statistics, machine learning, big data, climate science, and risk assessment. Such collaboration is essential for leveraging new climate projections tailored to extreme events and their societal and ecological impacts. Advancements in AI and ML will further facilitate synthetic data generation, advanced pattern analysis, and AI-driven early warning systems. Stakeholder engagement – including policymakers, communities, and industries – is essential to ensure actionable and regionally tailored strategies for risk reduction and climate adaptation.

... [1]

**Deleted:** Finally, this review highlights the importance of adopting a multidisciplinary approach that combines expertise in multivariate statistics, machine learning, big data, climate science, and risk assessment. Such collaboration is essential for leveraging new climate projections tailored to extreme events and their societal and ecological impacts. Advancements in AI and ML will further facilitate synthetic data generation, advanced pattern analysis, and AI-driven early warning systems. Stakeholder engagement – including policymakers, communities, and industries – is essential to ensure actionable and regionally tailored strategies for risk reduction and climate adaptation.

**Deleted: A2**

EG	Expected Gradient
GA	Genetic Algorithm
GAM	Generalised Additive Models
GAN	Generative Adversarial Network
GLM	Generalised Linear Models
GNN	Graph Neural Network
GP	Gaussian Process
GRU	Gated Recurrent Unit
IG	Integrated Gradient
KNN	K Nearest Neighbour
LSTM	Long Short Term Memory
MaxEnt	Maximum Entropy
ML	Machine Learning
NLP	Natural Language Processing
PCMCI	Peter and Clark Momentary Conditional Independence
RF	Random Forest
SHAP	Shapley Values
SVM	Support Vector Machine
XGBoost	Extreme Gradient Boosting

1163

1164 **Table A2: Other acronyms (in alphabetical order)**

Acronym	Full Name
AHP	Analytical Hierarchy Processes
CO	Carbon Monoxide
CDD	Consecutive Dry Days
CMIP	Coupled Model Intercomparison Project
DynaCLUE	Dynamic Conversion of Land Use and its Effect
EO	Earth observations
FWI	Fire Weather Index
GEV	Generalised Extreme Value (distributions)
HKH	Hindu-Kush and Himalaya (Region)
NO2	Nitrogen Dioxide
O3	Ozone
RCP	Representative Concentration Pathways

Deleted: A3

PLUS	Patch-generating Land Use Simulation
PM	Particle Matter
SO2	Sulphur dioxide
SMILE	Single Model Initial-condition Large Ensemble
SPEI	Standardised Precipitation and Evapotranspiration Index
SPI	Standardised Precipitation Index

1166

1167 **Appendix B: Summary tables of the collected studies**

1168 The literature review followed the PRISMA guidelines to ensure transparency and reproducibility in the identification,  
 1169 screening, and selection of studies. The process is summarized in the PRISMA flow diagram and detailed as follows.  
 1170 First, records were retrieved from major scientific databases (Scopus) and filtered by type, retaining only *articles, conference*  
 1171 *papers, and book chapters* and language (English). Next, documents were filtered by subject area and keyword, selecting only  
 1172 those classified under *Environmental Science* and *Earth and Planetary Science* as subjects and considering machine learning,  
 1173 climate change, risk assessments (and their synonyms and variations) as keywords.

1174 In the third step (title screening stage), studies not focusing on *natural hazards, multi-hazard, or risk assessment* were  
 1175 excluded. During the abstract screening stage, each paper was evaluated for its relevance to the review’s research questions,  
 1176 focusing particularly on the use of machine learning (ML) techniques and their application to multi-hazard or multi-risk  
 1177 contexts. Studies were retained if they explicitly applied ML, AI, or statistical learning methods to the modelling,  
 1178 characterization, or assessment of natural hazards, or if they addressed interactions between multiple hazard types (e.g.,  
 1179 cascading or compound events) and their associated risks. Papers focusing solely on single hazards without methodological  
 1180 innovation or on unrelated environmental modelling were excluded. This step ensured that the final selection captured studies  
 1181 advancing methodological understanding of ML-driven hazard analysis, as well as those integrating multiple hazard processes  
 1182 or risk dimensions. Finally, the full-text review identified the most relevant and representative papers, ensuring balanced  
 1183 coverage across different hazard types and AI methodologies. The final selections were based on diversity in data sources,  
 1184 geographical coverage, hazard types and machine learning methods used. This process ensured that the resulting corpus reflects  
 1185 the breadth of current research at the intersection of AI, Earth observation, and multi-hazard risk assessment.


1186 The number of studies retained at each step is summarized below (numbers correspond to the four main research questions):

1187 Table B1: summary of the screening step results

Screening step	RQ1	RQ2	RQ3	RQ4
Initial retrieval	24,335	9,542	22,054	1,961
After type filtering	17,676	8,731	3,548	344
After title screening	6,999	801	2,215	104
After abstract screening	376	107	214	67


Full text screening	52	50	29	22
---------------------	----	----	----	----

188  
 189  
 190 Figures B1, B2 and B3 provide an overview of the methodological and geographic characteristics of the reviewed literature  
 191 and together contextualise some of the limitations discussed in the main text. Figure B1 shows the distribution of ML methods  
 192 across application domains: CNNs dominate EO-processing applications, reflecting their architectural suitability for gridded  
 193 spatial data. Random Forest and other ensemble methods are the predominant choice for risk impact modelling and future risk  
 194 assessment, consistent with their interpretability, robustness to overfitting, and compatibility with tabular socio-environmental  
 195 predictor sets. LSTM and other sequence-to-sequence deep learning architectures concentrate in hazard prediction tasks where  
 196 temporal dynamics and memory effects are critical. Notably, methods specifically designed for multi-hazard interaction  
 197 modelling – such as graph neural networks or multi-output probabilistic models – are conspicuously absent from the reviewed  
 198 literature, confirming that the methodological frontier lies in adapting or designing architectures that natively represent inter-  
 199 hazard dependencies rather than applying single-hazard models independently.  
 200 Figure B2 illustrates the geographic distribution of reviewed papers across lead author country, all author countries, and case  
 201 study region, revealing that research output is heavily concentrated in China, the USA, and Western Europe. Figure B3 further  
 202 summarises this regional imbalance: Europe and North America together account for over 55% of lead authorships, while  
 203 regions such as Africa, South/SE Asia, and South America remain substantially underrepresented both as producers and  
 204 subjects of research. This geographic skew has direct methodological implications: the predominance of country-level  
 205 aggregated indicators, and the limited availability of sub-national spatially resolved datasets, partly reflects the data  
 206 infrastructure of the regions where most studies are conducted and may systematically underrepresent the vulnerability  
 207 dynamics of lower-income contexts where disaster impacts are most severe.  
 208



**Figure B1: Main methods used for each research topic**


1209  
1210



211

212 **Figure B2: Geographic distribution of reviewed papers: Sankey diagrams between main authors countries, co-author countries and**  
 213 **analysed case studies**

214



**Figure B3: Summary statistics of geographic distribution of reviewed papers**

215  
 216  
 217

1218

1219

1220 **Table B2: Summary of the research questions and their keywords**

Topic	Research question	Thematic keywords	Method-keywords
Data	How can data-driven applications improve data collection and processing?	Climate, model, observations, reanalysis, remote sensing, earth observations, social media, newspapers, downscaling, bias, impacts	ML (ML), AI (Artificial Intelligence), DL (Deep Learning), NN (neural networks), multivariate statistics, regression, prediction, forecast, classification, anomaly detection, copulas, interpretability, explainability
Hazard	How can data-driven applications be used to identify, classify, and cluster extreme events, and model hazard interactions?	Multi-hazard, drought, flood, heatwave, wildfire, landslide, storm, hurricane, volcanic, earthquake, wind, compound, consecutive, extremes	
Risk	How can data-driven applications integrate vulnerability and exposure in multi-risk analysis?	Multi-risk, climate-risk, multi-sector, environment(al), energy, health, infrastructure, susceptibility, vulnerability, exposure	
Future	How can data-driven applications be used to predict long-term future multi-hazard and multi-risk?	Climate change, tipping points, uncertainty, projections, future risk, RCP, storylines	

1221

1222

1223 **Table B3: Final selection of studies for RQ1: Data**

Reference	Title	Hazards/ Main variable	ML methods

Topic 1: Data - Climate			
(Orth et al., 2022)	Global soil moisture data derived through machine learning trained with in-situ measurements	Soil moisture	LSTM
(Ghiggi et al., 2019)	GRUN: an observation-based global gridded runoff dataset from 1902 to 2014	run-off	RF
(Anderson et al., 2019)	Environmental sensor placement with convolutional Gaussian neural processes	air temperature	CONVNP
(Tazi et al., 2024)	Downscaling precipitation over High-mountain Asia using multi-fidelity Gaussian processes: improved estimates from ERA5	precipitation	GP
(He et al., 2016)	Spatial downscaling of precipitation using adaptable random forests	precipitation	RF
(Lin et al., 2023)	Deep learning downscaled high-resolution daily near surface meteorological datasets over East Asia	Air temperature, humidity, wind, radiation	CNN
(Harris et al., 2022)	A Generative Deep Learning Approach to Stochastic Downscaling of Precipitation Forecasts	precipitation	GAN

Deleted: (Ling et al., 2024a)

... [2]

(Bretherton et al., 2022)	Correcting Coarse-Grid Weather and Climate Models by Machine Learning From Global Storm-Resolving Simulations	atmospheric variables	RF, ANN
(Clark et al., 2022)	Correcting a 200 km Resolution Climate Model in Multiple Climates by Machine Learning From 25 km Resolution Simulations	atmospheric variables	RF, ANN
(He et al., 2023)	Improving regional climate simulations based on a hybrid data assimilation and machine learning method	Atmospheric, vegetation, soil	Hybrid physics - XGBoost
(Huynh et al., 2025)	A distributed hybrid physics-AI framework for learning corrections of internal hydrological fluxes and enhancing high-resolution regionalized flood modeling	Hydrological	Hybrid physics -ANN
(S. Yu et al., 2024)	ClimSim-Online: A Large Multi-scale Dataset and Framework for Hybrid ML-physics Climate Emulation	Climate/atmospheric	CNN, Encoder/decoder, Heteroskedastic regression, MLP, randomized Prior Network, Conditional Variation Autoencoder
(Willard et al., 2022)	Integrating Scientific Knowledge with Machine Learning for Engineering and Environmental Systems	Literature review	Literature review

(Read et al., 2019)	Process-Guided Deep Learning Predictions of Lake Water Temperature	Water temperature	Hybrid physical / LSTM
(Xu et al., 2022)	Quantifying the uncertainty of precipitation forecasting using probabilistic deep learning	Precipitation	Probabilistic Deep learning
(Siddique et al., 2022)	A Survey of Uncertainty Quantification in Machine Learning for Space Weather Prediction	Space weather, uncertainty quantification	Gaussian Processes, Physics informed Neural Networks
(Ling et al., 2024)	Diffusion model-based probabilistic downscaling for 180-year East Asian climate reconstruction	Climate indices, hot and dry compound events, wind	Diffusion probabilistic downscaling model
(Saha & Ravela, 2022)	Downscaling Extreme Rainfall Using Physical-Statistical Generative Adversarial Learning	Extreme precipitation	Physical/ Generative Adversarial Network
Topic 2: Data - Earth observations			
(Ahmad et al., 2010)	Estimating soil moisture using remote sensing data: A machine learning approach	soil moisture	SVM, ANN, Linear regression

(Kang et al., 2018)	Spatial Upscaling of Sparse Soil Moisture Observations Based on Ridge Regression	soil moisture	Ridge Regression
(Han et al., 2023)	Global long term daily 1 km surface soil moisture dataset with physics informed machine learning	soil moisture	RF
(Jing et al., 2016a)	A Comparison of Different Regression Algorithms for Downscaling Monthly Satellite-Based Precipitation over North China	precipitation	CART, KNN, RF, SVM
(Jing et al., 2016b)	A Spatial Downscaling Algorithm for Satellite-Based Precipitation over the Tibetan Plateau Based on NDVI, DEM, and Land Surface Temperature	precipitation	RF, SVM
(Fang et al., 2017)	Prolongation of SMAP to Spatiotemporally Seamless Coverage of Continental U.S. Using a Deep Learning Neural Network	Soil Moisture	LSTM
(Adam et al., 2014)	Land-use/cover classification in a heterogeneous coastal landscape using RapidEye imagery: evaluating the performance of random forest and support vector machines classifiers	LULC classification (coastal)	RF, SVM
(Yuh et al., 2023)	Application of machine learning approaches for land cover monitoring in northern Cameroon	LULC monitoring	RF, SVM, KNN, ANN

(Zerrouki et al., 2019)	A Machine Learning-Based Approach for Land Cover Change Detection Using Remote Sensing and Radiometric Measurements	LULC change detection	RF, SVM, KNN, ANN
(Miyoshi et al., 2020)	A Novel Deep Learning Method to Identify Single Tree Species in UAV-Based Hyperspectral Images	Tree species mapping	CNN
(Schiefer et al., 2020)	Mapping forest tree species in high resolution UAV-based RGB-imagery by means of convolutional neural networks	Tree species mapping	CNN
(Veras et al., 2022)	Fusing multi-season UAS images with convolutional neural networks to map tree species in Amazonian forests	Tree species mapping	CNN
(J. Wang et al., 2019)	Deprivation pockets through the lens of convolutional neural networks	Identify deprived urban areas	CNN
(Ghaffarian et al., 2021)	Monitoring Urban Deprived Areas with Remote Sensing and Machine Learning in Case of Disaster Recovery	Track disaster recovery in urban deprived areas	SVM
(Nazeer et al., 2017)	Evaluation of Empirical and Machine Learning Algorithms for Estimation of Coastal Water Quality Parameters	Water quality	ANN
(J. Liu et al., 2023)	Monitoring Total Suspended Solids and Chlorophyll-a Concentrations in Turbid Waters: A Case Study of the Pearl River Estuary and Coast Using Machine Learning	Water quality (Turbidity)	ANN, RF, XGBoost, SVM

(S. Chen et al., 2022)	Machine learning-based estimation of riverine nutrient concentrations and associated uncertainties caused by sampling frequencies	Water Quality (River Nutrients)	SVM, RF, ANN
(Sublime & Kalinicheva, 2019)	Automatic Post-Disaster Damage Mapping Using Deep-Learning Techniques for Change Detection: Case Study of the Tohoku Tsunami	Change detection after disaster (earthquake/ tsunami)	CNN based autoencoder
(Ji et al., 2018)	Earthquake/Tsunami Damage Assessment for Urban Areas Using Post-Event PolSAR Data	Change detection after disaster (earthquake/ tsunami)	SVM
(Y. Bai et al., 2018)	Towards Operational Satellite-Based Damage-Mapping Using U-Net Convolutional Network: A Case Study of 2011 Tohoku Earthquake-Tsunami	Change detection after disaster (earthquake/ tsunami)	CNN
(Lei et al., 2019)	End-to-end Change Detection Using a Symmetric Fully Convolutional Network for Landslide Mapping	Change detection (landslide mapping)	CNN
(Bo et al., 2022)	BASNet: Burned Area Segmentation Network for Real-Time Detection of Damage Maps in Remote Sensing Images	Change detection (wildfire mapping)	CNN
(Tran et al., 2020)	Damage-Map Estimation Using UAV Images and Deep Learning Algorithms for Disaster Management System	Change detection (wildfire mapping)	CNN

(Munawar et al., 2021)	UAVs in Disaster Management: Application of Integrated Aerial Imagery and Convolutional Neural Network for Flood Detection	Change detection (flood mapping)	CNN
(Kabiru et al., 2023)	The relationship between multiple hazards and deprivation using open geospatial data and machine learning	Hydrological (floods, landslides), geophysical (earthquake, volcanic), biological, meteorological (temperature), human (transport, industrial, miscellaneous)	Random Forest
(Qiang et al., 2020)	Observing community resilience from space: Using nighttime lights to model economic disturbance and recovery pattern in natural disaster	Hurricane impacts	Univariate/multivariate regression
(Dasgupta et al., 2022)	Towards Daily High-resolution Inundation Observations using Deep Learning and EO	Floods (inundation maps)	Convolutional Neural Network
(Gierszewska & Berezowski, 2024)	A physics-guided neural network for flooding area detection using SAR imagery and local river gauge observations	Floods (inundation maps)	Physics/Neural Network
Topic 3: Data - Texts			
(Asinthara et al., 2022)	Classification of Disaster Tweets using Machine Learning and Deep Learning Techniques	Classifying disaster tweets	SVM, Naïve Bayes

Deleted: c

(Powers et al., 2023)	Using artificial intelligence to identify emergency messages on social media during a natural disaster: A deep learning approach	Classifying disaster tweets	BERT, XLNet, SVM
(Koshy & Elango, 2023)	Multimodal tweet classification in disaster response systems using transformer-based bidirectional attention model	Classifying disaster tweets and images	BERT, Transformers, LSTM
(Mehrotra et al., 2022)	A Multi-stage Classification Framework for Disaster-Specific Tweets	Classifying disaster tweets	SVM, DT, RF, ADABOOST, GBM, XGB, LSTM, BERT, XLNET
(Sodoge et al., 2023)	Automatized spatio-temporal detection of drought impacts from newspaper articles using natural language processing and machine learning	Classifying drought impacts from newspapers	Naïve Bayes, Lasso Regression, RF, ANN

1226

1227

1228 **Table B4: Final selection of studies for RQ2: Hazard**

Reference	Title	Hazards/ Main variable	ML methods
Topic 1: Hazard – identify, classify, cluster			
(Ionita et al., 2021)	Compound Hot and Dry Events in Europe: Variability and Large-Scale Drivers	Hot and Dry compound events	Percentile based thresholds, Empirical Orthogonal Functions

(Sutanto et al., 2020)	Heatwaves, droughts, and fires: Exploring compound and cascading dry hazards at the pan-European scale	Heatwave, drought, wildfire	Percentile based thresholds
(Claassen et al., 2023)	A new method to compile global multi-hazard event sets	Heatwave, coldwave, drought, wildfire, floods, earthquakes, wind, tsunami, tropical cyclone, volcano, landslide	Percentile based thresholds
(Liao et al., 2021)	Growing Threats From Unprecedented Sequential Flood-Hot Extremes Across China	consecutive flood - heatwave	Return periods
(Sfetsos et al., 2023)	Multi-Hazard Extreme Scenario Quantification Using Intensity, Duration, and Return Period Characteristics	Heatwave, coldwave, precipitation, snowfall, wind extremes	Return periods
(Orth et al., 2022)	Contrasting biophysical and societal impacts of hydro-meteorological extremes	Heatwave, Drought, Floods, Wildfire	Return periods, percentiles
(Y. Liu et al., 2016)	Application of Deep Convolutional Neural Networks for Detecting Extreme Weather in Climate Datasets	Extreme weather (Tropical cyclones, atmospheric rivers, weather fronts)	CNN
(Racah et al., 2016)	ExtremeWeather: A large-scale climate dataset for semi-supervised detection, localization, and understanding of extreme weather events	Extreme weather (Tropical cyclones, atmospheric rivers, weather fronts)	CNN (semi-supervised)
(Cammalleri & Toreti, 2023)	A Generalized Density-Based Algorithm for the Spatiotemporal Tracking of Drought Events	Drought	DBSCAN, Percentile based thresholds

(J. Wang & Yan, 2021)	Rapid rises in the magnitude and risk of extreme regional heat wave events in China	heatwaves	DBSCAN, Percentile based thresholds
(Di Martino et al., 2018b)	Spatiotemporal extended fuzzy C-means clustering algorithm for hotspots detection and prediction	earthquakes	DBSCAN
(Tilloy et al., 2022)	A methodology for the spatiotemporal identification of compound hazards: wind and precipitation extremes in Great Britain (1979–2019)	Wind and precipitation	DBSCAN, Percentile based thresholds
(H. Yu et al., 2022)	Hotspots, co-occurrence, and shifts of compound and cascading extreme climate events in Eurasian drylands	Drought, heatwave, coldwave, precipitation, wind	DBSCAN, Percentile based thresholds
Topic 2: Hazard - Predict			
(Haggag et al., 2021)	A deep learning model for predicting climate-induced disasters	Multi-Hazard (flood tested)	ANN
(Kratzert, Klotz, Shalev, et al., 2019)	Towards learning universal, regional, and local hydrological behaviors via machine learning applied to large-sample datasets	Floods	LSTM
(Kratzert, Klotz, Brandstetter, et al., 2019)	Using LSTMs for climate change assessment studies on droughts and floods	Floods, droughts	LSTM
(Tiggeloven et al., 2021)	Exploring deep learning capabilities for surge predictions in coastal areas	Storm Surge	LSTM, CNN, ANN

(S. Jiang, Bevacqua, et al., 2022)	River flooding mechanisms and their changes in Europe revealed by explainable machine learning	River floods, pluvial floods, snowmelt floods	LSTM
(Kraft et al., 2019)	Identifying Dynamic Memory Effects on Vegetation State Using Recurrent Neural Networks	Hot and dry events (impacts on vegetation)	LSTM
(Freeman et al., 2018)	Forecasting air quality time series using deep learning	LSTM	
(Q. Wu & Lin, 2019)	A novel optimal-hybrid model for daily air quality index prediction considering air pollutant factors	Air quality (various pollutants)	LSTM
(Chang-Hoi et al., 2021)	Development of a PM2.5 prediction model using a recurrent neural network algorithm for the Seoul metropolitan area, Republic of Korea	Air quality (PM 2.5)	RNN
(Bentivoglio et al., 2023)	Rapid spatio-temporal flood modelling via hydraulics-based graph neural networks	Floods	GNN
(Kazadi et al., 2024)	FloodGNN-GRU: a spatio-temporal graph neural network for flood prediction	Floods	GNN-GRU
(A. Y. Sun et al., 2021)	Explore Spatio-Temporal Learning of Large Sample Hydrology Using Graph Neural Networks	Floods	GNN
(Castangia et al., 2023)	Transformer neural networks for interpretable flood forecasting	Floods	Transformers

Deleted: (Q. Wu & Lin, 2019)zone

(Bonino et al., 2024)	Machine learning methods to predict sea surface temperature and marine heatwave occurrence: a case study of the Mediterranean Sea	marine heatwaves	CNN, LSTM, RF
(Patil et al., 2023)	Predicting extreme floods and droughts in East Africa using a deep learning approach	drought	CNN
(Singh et al., 2021)	Drought risk assessment and prediction using artificial intelligence over the southern Maharashtra state of India	drought	ANN
(Ayyad et al., 2022)	Machine learning-based assessment of storm surge in the New York metropolitan area	storm surge	RF, XGBoost, Extra Trees, SVM
(Macdonald et al., 2025)	Robust storm surge forecasts for early warning system: a machine learning approach using Monte Carlo Bayesian model selection algorithm	Storm surge	Monte Carlo dropout + Bayesian NN
(M. Nguyen et al., 2024)	Estimating uncertainty in flood model outputs using machine learning informed by Monte Carlo analysis	Flooding	Monte Carlo dropout + Bayesian NN
Topic 3: Hazard - Interactions			
(Couasnon et al., 2018)	A Copula-Based Bayesian Network for Modeling Compound Flood Hazard from Riverine and Coastal Interactions at the Catchment Scale: An Application to the Houston Ship Channel, Texas	Compound river and coastal flood	Copulas, Bayesian Networks

(Sadegh et al., 2017)	Multivariate Copula Analysis Toolbox (MvCAT): Describing dependence and underlying uncertainty using a Bayesian framework	droughts, floods	Copulas
(Bevacqua et al., 2017b)	Multivariate statistical modelling of compound events via pair-copula constructions: analysis of floods in Ravenna (Italy)	River floods, precipitation, coastal floods	Copulas
(Bevacqua et al., 2021)	Guidelines for Studying Diverse Types of Compound Weather and Climate Events	compound flooding, precipitation/landslide	Copulas, regressions, percentile thresholds, clustering
(Hochrainer et al., 2019)	Large scale extreme risk assessment using copulas: an application to drought events under climate change for Austria	drought	copulas
(Tootoonchi et al., 2022)	Copulas for hydroclimatic analysis: A practice-oriented overview	Temperature, precipitation	copulas
(Jiang et al., 2023)	Estimating propagation probability from meteorological to ecological droughts using a hybrid machine learning copula method	Droughts	Copulas, 3D clustering, 11 ML methods (KNN, SVM, GP, DT, MLP, AdaBoost, Naive Bayes, quadratic discriminant analysis, Gradient Boosting, XGBoost, Random Forest)
(Cao et al., 2020)	Multi-geohazards susceptibility mapping based on machine learning—a case study in Jiuzhaigou, China	rockfall, landslide, debris flow	RF, SVM, XGBoost
(Javidan et al., 2021)	Evaluation of multi-hazard map produced using MaxEnt machine learning technique	flood, landslide, gully erosion	MaxEnt

(Karakas et al., 2023)	A Hybrid Multi-Hazard Susceptibility Assessment Model for a Basin in Elazig Province, Türkiye	Landslide, Flood, Earthquake	RF
(Kariminejad et al., 2022)	Analytical techniques for mapping multi-hazard with geo-environmental modeling approaches and UAV images	collapsed pipe, gully erosion, landslide	BRT, Flexible discriminant analysis, Multivariate adaptive regression spline, Mixture discriminant analysis, RF, GLM and SVM
(H. D. Nguyen et al., 2023)	Multi-hazard assessment using machine learning and remote sensing in the North Central region of Vietnam	Flood, landslide	SVM, RF, AdaBoost
(Pourghasemi et al., 2020)	Assessing and mapping multi-hazard risk susceptibility using a machine learning technique	Flood, landslide, wildfire	RF
(Pouyan et al., 2021)	A multi-hazard map-based flooding, gully erosion, forest fires, and earthquakes in Iran	gully erosion, wildfire, earthquake	RF, SVM, BRT
(Yousefi et al., 2020)	A machine learning framework for multi-hazards modeling and mapping in a mountainous area	avalanche, landslide, wildfire, subsidence, flood	SVM, BRT, GLM, FDA
(Piao et al., 2022)	Multi-hazard mapping of droughts and forest fires using a multi-layer hazards approach with machine learning algorithms	drought, wildfire	CART, RF, BRT
(Ullah et al., 2022)	Multi-hazard susceptibility mapping based on Convolutional Neural Networks	flash flood, debris flow, landslide	CNN, RF

(Mandal et al., 2022)	Mapping the multi-hazards risk index for coastal block of Sundarban, India using AHP and machine learning algorithms	cyclones, storm surge, coastal erosion	ANN, RF
-----------------------	----------------------------------------------------------------------------------------------------------------------	----------------------------------------	---------

1230

1231

1232 Table B5: Final selection of studies for RQ3: Risk

Deleted: ¶

... [3]

Reference	Title	Hazards/ Main variable	ML methods
Topic 1: Risk - Combining hazard, exposure and vulnerability			
(Kotaridis & Lazaridou, 2022)	Integration of convolutional neural networks for flood risk mapping in Tuscany, Italy	flood	CNN
(Zhao et al., 2020)	Urban flood susceptibility assessment based on convolutional neural networks	flood	CNN
(Rusk et al., 2022)	Multi-hazard susceptibility and exposure assessment of the Hindu Kush Himalaya	flood, landslide, wildfire	MaxEnt
(Fuchs et al., 2015)	A spatiotemporal multi-hazard exposure assessment based on property data	river flood, snow avalanche, torrential flood	Frequency ratio
(Sammonds et al., 2023)	Hurricane risk assessment in a multi-hazard context for Dominica in the Caribbean	hurricane, landslides, floods	Frequency ratio, analytical hierarchy process
(Luu et al., 2024)	Integrating multi-hazard susceptibility and building exposure: A case study for Quang Nam province, Vietnam	flood, wildfire	RF, CART

(K. Liu et al., 2018)	Susceptibility of existing and planned Chinese railway system subjected to rainfall-induced multi-hazards	flood, landslide, debris flow	RF
(Arvin et al., 2023)	Assessment of infrastructure resilience in multi-hazard regions: A case study of Khuzestan Province	flood, landslide, earthquake	analytical hierarchy process
(Khatakho et al., 2021)	Multi-Hazard Risk Assessment of Kathmandu Valley, Nepal	flood, earthquake, wildfire	analytical hierarchy process
Topic 2: Risk – Predicting impacts			
(Gasparrini, 2014)	Modeling exposure–lag–response associations with distributed lag non-linear models	heatwave, air pollution	Distributed Lag Non-Linear Models
(Guo et al., 2024)	Regional variation in the role of humidity on city-level heat-related mortality	heatwave, humidity	RF
(Y. Wang et al., 2019)	A random forest model to predict heatstroke occurrence for heatwave in China	heatwave, humidity	RF
(X. Wang et al., 2021)	Quantitative Impact Analysis of Climate Change on Residents' Health Conditions with Improving Eco-Efficiency in China: A Machine Learning Perspective	heatwave, humidity, previous diseases	SVM
(Boudreault et al., 2023)	Machine and deep learning for modelling heat-health relationships	heatwave, air pollution	DT, RF, GBM, SLP, MLP, LSTM, GLM, GAM, DLNM
(Côté et al., 2024)	Vulnerability assessment of heat waves within a risk framework using artificial intelligence	heatwave, air pollution	Auto-Gluon, GP, Deep GP

(Busker et al., 2024)	Predicting Food-Security Crises in the Horn of Africa Using Machine Learning	Heatwaves, droughts, precipitation, conflict	XGB
(Tárraga et al., 2024)	Causal discovery reveals complex patterns of drought-induced displacement	drought, precipitation, conflict	Granger Causality, PCMCI
(Zscheischler et al., 2017)	Bivariate return periods of temperature and precipitation explain a large fraction of European crop yields	drought, heatwave, precipitation	Copulas
(Ribeiro et al., 2020)	Risk of crop failure due to compound dry and hot extremes estimated with nested copulas	drought, heatwave	Copulas
(R. Wang et al., 2021)	Predicting stream water quality under different urban development pattern scenarios with an interpretable machine learning approach	water quality, land use planning	RF
(Li et al., 2022)	Interpretable tree-based ensemble model for predicting beach water quality	water quality	DT, RF, CatBoost, GBM, XGBoost
(Cushman et al., 2017)	Multiple-scale prediction of forest loss risk across Borneo	forest loss	RF, logistic regression
(Islam et al., 2021)	Machine learning algorithm-based risk assessment of riparian wetlands in Padma River Basin of Northwest Bangladesh	drought, topography, environmental and antropogenic stressors	RF, SVM, DT, ANN

(Schmidt et al., 2020)	The role of spatial units in modelling freshwater fish distributions: Comparing a subcatchment and river network approach using MaxEnt	topography, environmental and antropogenic stressors	MaxEnt
(Teichert et al., 2016)	Restoring fish ecological quality in estuaries: Implication of interactive and cumulative effects among anthropogenic stressors	topography, environmental and antropogenic stressors	RF
(Dal Barco et al., 2024)	A machine learning approach to evaluate coastal risks related to extreme weather events in the Veneto region (Italy)	precipitation, wind, sea level rise, storm surges	AN(Pilkington & Mahmoud, 2017)n
(Pilkington & Mahmoud, 2017)	Spatial and temporal variations in resilience to tropical cyclones along the United States coastline as determined by the multi-hazard hurricane impact level model	wind, storm surge, precipitation, flooding	ANN
(Mukherjee et al., 2018)	A multi-hazard approach to assess severe weather-induced major power outage risks in the U.S.	heatwave, wildfire, hurricane, coldwave, wind, precipitation	SVM, RF
(Carranante et al., 2024)	Machine learning-based climate risk sharing for an insured loan in the tourism industry	wind, precipitation, heatwave	RF

1236

1237

1238

1239 **Table B5: Final selection of studies for RQ4: future**

Reference	Title	Hazards/ variable	Main	ML methods
Topic 1: Future: hazard				
(Zscheischler et al., 2018)	Future climate risk from compound events	compound events		copulas, storylines
(Ridder et al., 2022)	Increased occurrence of high impact compound events under climate change	drought, heatwaves, precipitation, wind		percentile threshold, return period
(Zhu et al., 2023)	Compound wind and precipitation extremes at a global scale based on CMIP6 models: Evaluation, projection and uncertainty	wind, precipitation		percentile threshold, return period
(Ridder et al., 2021)	Do CMIP6 Climate Models Simulate Global or Regional Compound Events Skillfully?	wind, precipitation		percentile threshold, return period
(Ghanbari et al., 2021)	Climate Change and Changes in Compound Coastal-Riverine Flooding Hazard Along the U.S. Coasts	coastal flood, river flood, sea level rise		copulas
(H. Wu et al., 2023)	Increasing Risks of Future Compound Climate Extremes with Warming Over Global Land Masses	drought, heatwave, precipitation		copulas
(H. Wu et al., 2024)	Predicting compound agricultural drought and hot events using a Cascade Modeling framework combining Bayesian Model Averaging ensemble with Vine Copula (CaMBMAViC)	drought, heatwave		copulas

(Bevacqua et al., 2021)	Guidelines for Studying Diverse Types of Compound Weather and Climate Events	High-Impact Low-Probability Events	storylines
Topic 2: Future – Risk			
(Ayyad et al., 2023)	Climate change impact on hurricane storm surge hazards in New York/New Jersey Coastlines using machine-learning	hurricane, storm surge	SVM, AdaBoost
(S. J. Park & Lee, 2020)	Prediction of coastal flooding risk under climate change impacts in South Korea using machine learning algorithms	precipitation, storm surge, sea level rise	KNN, RF, SVM
(S. Park et al., 2023)	Adaptation strategies for future coastal flooding: Performance evaluation of green and grey infrastructure in South Korea	precipitation, storm surge, sea level rise	KNN, RF, SVM
(Lim & Kim, 2022)	Can Forest-Related Adaptive Capacity Reduce Landslide Risk Attributable to Climate Change? -Case of Republic of Korea	precipitation, landslide	RF
(Pham et al., 2023)	Multi-model chain for climate change scenario analysis to support coastal erosion and water quality risk management for the Metropolitan city of Venice	coastal erosion, water quality, storm surge	Bayesian Network
(García-León et al., 2024)	Temperature-related mortality burden and projected change in 1368 European regions: a modelling study	heatwave, future population, economic factors	weighted averages

(Rahman et al., 2024)	Multi-hazard could exacerbate in coastal Bangladesh in the context of climate change	flash floods, river floods, coastal floods, landslide	LSTM
(Ya et al., 2023)	Increased flood susceptibility in the Tibetan Plateau with climate and land use changes	flood	logistic regression
(Liang et al., 2021)	Understanding the drivers of sustainable land expansion using a patch-generating land use simulation (PLUS) model: A case study in Wuhan, China	Future land use	RF
(Saha et al., 2021)	Modelling multi-hazard threats to cultural heritage sites and environmental sustainability: The present and future scenarios	earthquake, landslide, precipitation	BRT, BART, BGLM
(Janizadeh et al., 2021)	Mapping the spatial and temporal variability of flood hazard affected by climate and land-use changes	Floods	GBM, XGB
(Leng & Hall, 2020)	Predicting spatial and temporal variability in crop yields: an inter-comparison of machine learning, regression and process-based models	precipitation, drought, heatwave	RF
(Khan et al., 2024)	Association of precipitation extremes and crops production and projecting future extremes using machine learning approaches with CMIP6 data	Precipitation	XGB

(Tabari & Willems, 2023)	Global risk assessment of compound hot-dry events in the context of future climate change and socioeconomic factors	drought, heatwaves	Copulas
--------------------------	---------------------------------------------------------------------------------------------------------------------	--------------------	---------

1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252

1253 **Author contribution**

1254 DMF: Conceptualisation, Methodology, Formal analysis, Investigation, Data curation, Visualisation, Writing – original draft.  
1255 MS: Conceptualisation, Methodology, Validation, Writing – review and editing.  
1256 MM: Conceptualisation, Data curation, Writing – review and editing.  
1257 AC: Funding acquisition, Supervision, Conceptualisation, Writing – review and editing.  
1258 ST: Funding acquisition, Supervision, Conceptualisation, Project administration.

1259 **Competing interest**

1260 The authors declare that they have no conflict of interest.

1261 **Financial support**

1262 The research was carried out within the frame of *Myriad\_EU project* (<https://www.myriadproject.eu/>), which has received  
1263 fundings from the European Union’s Horizon 2020 research and innovation programme call H2020-LC-CLA-2018-2019-2020  
1264 under grant agreement number 101003276.

1265  
1266

Deleted: .....Page Break.....

1269 **References**

- 1270 Abu El-Magd, S. A., Ali, S. A., & Pham, Q. B. (2021). Spatial modeling and susceptibility zonation of landslides using random  
1271 forest, naïve bayes and K-nearest neighbor in a complicated terrain. *Earth Science Informatics*, *14*(3), 1227–1243.  
1272 <https://doi.org/10.1007/s12145-021-00653-y>
- 1273 Adam, E., Mutanga, O., Odindi, J., & Abdel-Rahman, E. M. (2014). Land-use/cover classification in a heterogeneous coastal  
1274 landscape using RapidEye imagery: evaluating the performance of random forest and support vector machines  
1275 classifiers. *International Journal of Remote Sensing*, *35*(10), 3440–3458.  
1276 <https://doi.org/10.1080/01431161.2014.903435>
- 1277 AghaKouchak, A., Chiang, F., Huning, L. S., Love, C. A., Mallakpour, I., Mazdiyasi, O., Moftakhari, H., Papalexou, S. M.,  
1278 Ragno, E., & Sadegh, M. (2020). Climate Extremes and Compound Hazards in a Warming World. *Annual Review of*  
1279 *Earth and Planetary Sciences*, *48*(1), 519–548. <https://doi.org/10.1146/annurev-earth-071719-055228>
- 1280 Agrawal, S. (UCLA). (2022). *The Effectiveness of Copulas for Modeling Compound Climate Extreme Events in Boulder*  
1281 *County, Colorado*. [UCLA]. <https://doi.org/ProQuest> ID: Agrawal\_ucla\_0031N\_21275. Merritt ID:  
1282 <ark:/13030/m59m1m6w>.
- 1283 Ahmad, S., Kalra, A., & Stephen, H. (2010). Estimating soil moisture using remote sensing data: A machine learning approach.  
1284 *Advances in Water Resources*, *33*(1), 69–80. <https://doi.org/10.1016/j.advwatres.2009.10.008>
- 1285 Ahmadlou, M., Al-Fugara, A., Al-Shabeeb, A. R., Arora, A., Al-Adamat, R., Pham, Q. B., Al-Ansari, N., Linh, N. T. T., &  
1286 Sajedi, H. (2021). Flood susceptibility mapping and assessment using a novel deep learning model combining multilayer  
1287 perceptron and autoencoder neural networks. *Journal of Flood Risk Management*, *14*(1).  
1288 <https://doi.org/10.1111/jfr3.12683>
- 1289 Albulescu, A.-C. and Armaş, I.: An impact-chain-based exploration of multi-hazard vulnerability dynamics: the multi-hazard  
1290 of floods and the COVID-19 pandemic in Romania, *Nat. Hazards Earth Syst. Sci.*, *24*, 2895–2922,  
1291 <https://doi.org/10.5194/nhess-24-2895-2024>,
- 1292 Amato, F., Guignard, F., Robert, S., & Kanevski, M. (2020). A novel framework for spatio-temporal prediction of  
1293 environmental data using deep learning. *Scientific Reports*, *10*(1), 22243. <https://doi.org/10.1038/s41598-020-79148-7>
- 1294 Anderson, M. J., de Valpine, P., Punnett, A., & Miller, A. E. (2019). A pathway for multivariate analysis of ecological  
1295 communities using copulas. *Ecology and Evolution*, *9*(6), 3276–3294. <https://doi.org/10.1002/ece3.4948>
- 1296 Andersson, T. R., Bruinsma, W. P., Markou, S., Requeima, J., Coca-Castro, A., Vaughan, A., Ellis, A.-L., Lazzara, M. A.,  
1297 Jones, D., Hosking, S., & Turner, R. E. (2023). Environmental sensor placement with convolutional Gaussian neural  
1298 processes. *Environmental Data Science*, *2*, e32. <https://doi.org/10.1017/eds.2023.22>
- 1299 Angelov, D. (2020). *Top2Vec: Distributed Representations of Topics*. <http://arxiv.org/abs/2008.09470>
- 1300 Arosio, M., Cesarini, L., & Martina, M. L. V. (2021). Assessment of the Disaster Resilience of Complex Systems: The Case  
1301 of the Flood Resilience of a Densely Populated City. *Water*, *13*(20), 2830. <https://doi.org/10.3390/w13202830>

Formatted: English (US)

Deleted: ¶

- 1303 Arosio, M., Martina, M. L. V., & Figueiredo, R. (2020). The whole is greater than the sum of its parts: a holistic graph-based  
1304 assessment approach for natural hazard risk of complex systems. *Natural Hazards and Earth System Sciences*, 20(2),  
1305 521–547. <https://doi.org/10.5194/nhess-20-521-2020>
- 1306 Arvin, M., Beiki, P., Hejazi, S. J., Sharifi, A., & Atashafrooz, N. (2023). Assessment of infrastructure resilience in multi-  
1307 hazard regions: A case study of Khuzestan Province. *International Journal of Disaster Risk Reduction*, 88, 103601.  
1308 <https://doi.org/10.1016/j.ijdr.2023.103601>
- 1309 Asinthara, K., Jayan, M., & Jacob, L. (2022). Classification of Disaster Tweets using Machine Learning and Deep Learning  
1310 Techniques. *2022 International Conference on Trends in Quantum Computing and Emerging Business Technologies*  
1311 *(TQCEBT)*, 1–5. <https://doi.org/10.1109/TQCEBT54229.2022.10041629>
- 1312 Ayyad, M., Hajj, M. R., & Marsooli, R. (2022). Machine learning-based assessment of storm surge in the New York  
1313 metropolitan area. *Scientific Reports*, 12(1), 19215. <https://doi.org/10.1038/s41598-022-23627-6>
- 1314 Ayyad, M., Hajj, M. R., & Marsooli, R. (2023). Climate change impact on hurricane storm surge hazards in New York/New  
1315 Jersey Coastlines using machine-learning. *Npj Climate and Atmospheric Science*, 6(1), 88.  
1316 <https://doi.org/10.1038/s41612-023-00420-4>
- 1317 Bai, T., Wang, L., Yin, D., Sun, K., Chen, Y., Li, W., & Li, D. (2023). Deep learning for change detection in remote sensing:  
1318 a review. *Geo-Spatial Information Science*, 26(3), 262–288. <https://doi.org/10.1080/10095020.2022.2085633>
- 1319 Bai, Y., Mas, E., & Koshimura, S. (2018). Towards Operational Satellite-Based Damage-Mapping Using U-Net Convolutional  
1320 Network: A Case Study of 2011 Tohoku Earthquake-Tsunami. *Remote Sensing*, 10(10), 1626.  
1321 <https://doi.org/10.3390/rs10101626>
- 1322 Bankoff, G., & Hilhorst, H. (2022). Why Vulnerability Still Matters: The Politics of Disaster Risk Creation. *Routledge*.  
1323 <https://doi.org/10.4324/9781003219453>.
- 1324 Barrett, A. B., Duivenvoorden, S., Salakpi, E. E., Muthoka, J. M., Mwangi, J., Oliver, S., & Rowhani, P. (2020). *Forecasting*  
1325 *vegetation condition for drought early warning systems in pastoral communities in Kenya*.  
1326 <http://arxiv.org/abs/1911.10339>
- 1327 Bentivoglio, R., Isufi, E., Jonkman, S. N., & Taormina, R. (2023). Rapid spatio-temporal flood modelling via hydraulics-based  
1328 graph neural networks. *Hydrology and Earth System Sciences*, 27(23), 4227–4246. <https://doi.org/10.5194/hess-27-4227-2023>
- 1330 Bevacqua, E., De Michele, C., Manning, C., Couasnon, A., Ribeiro, A. F. S., Ramos, A. M., Vignotto, E., Bastos, A., Blesić,  
1331 S., Durante, F., Hillier, J., Oliveira, S. C., Pinto, J. G., Ragno, E., Rivoire, P., Saunders, K., Wiel, K., Wu, W., Zhang,  
1332 T., & Zscheischler, J. (2021). Guidelines for Studying Diverse Types of Compound Weather and Climate Events. *Earth's*  
1333 *Future*, 9(11). <https://doi.org/10.1029/2021EF002340>
- 1334 Bevacqua, E., Maraun, D., Hobæk Haff, I., Widmann, M., & Vrac, M. (2017a). Multivariate statistical modelling of compound  
1335 events via pair-copula constructions: analysis of floods in Ravenna (Italy). *Hydrology and Earth System Sciences*, 21(6),  
1336 2701–2723. <https://doi.org/10.5194/hess-21-2701-2017>

Formatted: Font: Italic

- 1337 Bevacqua, E., Maraun, D., Hobæk Haff, I., Widmann, M., & Vrac, M. (2017b). Multivariate statistical modelling of compound  
1338 events via pair-copula constructions: analysis of floods in Ravenna (Italy). *Hydrology and Earth System Sciences*, *21*(6),  
1339 2701–2723. <https://doi.org/10.5194/hess-21-2701-2017>
- 1340 Bevacqua, E., Suarez-Gutierrez, L., Jézéquel, A., Lehner, F., Vrac, M., Yiou, P., & Zscheischler, J. (2023). Advancing research  
1341 on compound weather and climate events via large ensemble model simulations. *Nature Communications*, *14*(1), 2145.  
1342 <https://doi.org/10.1038/s41467-023-37847-5>
- 1343 Beven, K. (2018). *Environmental Modelling*. CRC Press. <https://doi.org/10.1201/9781482288575>
- 1344 Bhowmik, R. T., Jung, Y. S., Aguilera, J. A., Prunicki, M., & Nadeau, K. (2023). A multi-modal wildfire prediction and early-  
1345 warning system based on a novel machine learning framework. *Journal of Environmental Management*, *341*, 117908.  
1346 <https://doi.org/10.1016/j.jenvman.2023.117908>
- 1347 Bi, K., Xie, L., Zhang, H., Chen, X., Gu, X., & Tian, Q. (2022). *Pangu-Weather: A 3D High-Resolution Model for Fast and*  
1348 *Accurate Global Weather Forecast*. <http://arxiv.org/abs/2211.02556>
- 1349 Bo, W., Liu, J., Fan, X., Tjahjadi, T., Ye, Q., & Fu, L. (2022). BASNet: Burned Area Segmentation Network for Real-Time  
1350 Detection of Damage Maps in Remote Sensing Images. *IEEE Transactions on Geoscience and Remote Sensing*, *60*, 1–  
1351 13. <https://doi.org/10.1109/TGRS.2022.3197647>
- 1352 Bonino, G., Galimberti, G., Masina, S., McAdam, R., & Clementi, E. (2024). Machine learning methods to predict sea surface  
1353 temperature and marine heatwave occurrence: a case study of the Mediterranean Sea. *Ocean Science*, *20*(2), 417–432.  
1354 <https://doi.org/10.5194/os-20-417-2024>
- 1355 Bordbar, M., Aghamohammadi, H., Pourghasemi, H. R., & Azizi, Z. (2022). Multi-hazard spatial modeling via ensembles of  
1356 machine learning and meta-heuristic techniques. *Scientific Reports*, *12*(1), 1451. [https://doi.org/10.1038/s41598-022-](https://doi.org/10.1038/s41598-022-05364-y)  
1357 [05364-y](https://doi.org/10.1038/s41598-022-05364-y)
- 1358 Boudreault, J., Campagna, C., & Chebana, F. (2023). Machine and deep learning for modelling heat-health relationships.  
1359 *Science of The Total Environment*, *892*, 164660. <https://doi.org/10.1016/j.scitotenv.2023.164660>
- 1360 Bretherton, C. S., Henn, B., Kwa, A., Brenowitz, N. D., Watt-Meyer, O., McGibbon, J., Perkins, W. A., Clark, S. K., & Harris,  
1361 L. (2022). Correcting Coarse-Grid Weather and Climate Models by Machine Learning From Global Storm-Resolving  
1362 Simulations. *Journal of Advances in Modeling Earth Systems*, *14*(2). <https://doi.org/10.1029/2021MS002794>
- 1363 Busker, T., van den Hurk, B., de Moel, H., van den Homberg, M., van Straaten, C., Odongo, R. A., & Aerts, J. C. J. H. (2024).  
1364 Predicting Food-Security Crises in the Horn of Africa Using Machine Learning. *Earth's Future*, *12*(8).  
1365 <https://doi.org/10.1029/2023EF004211>
- 1366 Cammalleri, C., & Toreti, A. (2023). A Generalized Density-Based Algorithm for the Spatiotemporal Tracking of Drought  
1367 Events. *Journal of Hydrometeorology*, *24*(3), 537–548. <https://doi.org/10.1175/JHM-D-22-0115.1>
- 1368 Campbell, A. M., Racault, M.-F., Goult, S., & Laurenson, A. (2020). Cholera Risk: A Machine Learning Approach Applied  
1369 to Essential Climate Variables. *International Journal of Environmental Research and Public Health*, *17*(24), 9378.  
1370 <https://doi.org/10.3390/ijerph17249378>

1371 Cannon, T. (2017). Social Vulnerability and Environmental Hazards. *International Encyclopedia of Geography*, Wiley.  
1372 <https://doi.org/10.1002/9781118786352.wbieg0845>.

1373 Cao, J., Zhang, Z., Du, J., Zhang, L., Song, Y., & Sun, G. (2020). Multi-geohazards susceptibility mapping based on machine  
1374 learning—a case study in Jiuzhaigou, China. *Natural Hazards*, *102*(3), 851–871. [https://doi.org/10.1007/s11069-020-](https://doi.org/10.1007/s11069-020-03927-8)  
1375 [03927-8](https://doi.org/10.1007/s11069-020-03927-8)

1376 Carannante, M., D’amato, V., Fersini, P., & Forte, S. (2024). Machine learning-based climate risk sharing for an insured loan  
1377 in the tourism industry. *Quality & Quantity*. <https://doi.org/10.1007/s11135-024-01958-y>

1378 Carvalho, D. V., Pereira, E. M., & Cardoso, J. S. (2019). Machine Learning Interpretability: A Survey on Methods and Metrics.  
1379 *Electronics*, *8*(8), 832. <https://doi.org/10.3390/electronics8080832>

1380 Castangia, M., Grajales, L. M. M., Aliberti, A., Rossi, C., Macii, A., Macii, E., & Patti, E. (2023). Transformer neural networks  
1381 for interpretable flood forecasting. *Environmental Modelling & Software*, *160*, 105581.  
1382 <https://doi.org/10.1016/j.envsoft.2022.105581>

1383 Chang-Hoi, H., Park, I., Oh, H.-R., Gim, H.-J., Hur, S.-K., Kim, J., & Choi, D.-R. (2021). Development of a PM2.5 prediction  
1384 model using a recurrent neural network algorithm for the Seoul metropolitan area, Republic of Korea. *Atmospheric*  
1385 *Environment*, *245*, 118021. <https://doi.org/10.1016/j.atmosenv.2020.118021>

1386 Chen, K., Han, T., Gong, J., Bai, L., Ling, F., Luo, J.-J., Chen, X., Ma, L., Zhang, T., Su, R., Ci, Y., Li, B., Yang, X., &  
1387 Ouyang, W. (2023). *FengWu: Pushing the Skillful Global Medium-range Weather Forecast beyond 10 Days Lead*.  
1388 <http://arxiv.org/abs/2304.02948>

1389 Chen, S., Zhang, Z., Lin, J., & Huang, J. (2022). Machine learning-based estimation of riverine nutrient concentrations and  
1390 associated uncertainties caused by sampling frequencies. *PLOS ONE*, *17*(7), e0271458.  
1391 <https://doi.org/10.1371/journal.pone.0271458>

1392 Claassen, J. N., Koks, E. E., de Ruiter, M. C., Ward, P. J., & Jäger, W. S. (2024). VineCopulas: an open-source Python package  
1393 for vine copula modelling. *Journal of Open Source Software*, *9*(101), 6728. <https://doi.org/10.21105/joss.06728>

1394 Claassen, J. N., Ward, P. J., Daniell, J., Koks, E. E., Tiggeloven, T., & de Ruiter, M. C. (2023). A new method to compile  
1395 global multi-hazard event sets. *Scientific Reports*, *13*(1), 13808. <https://doi.org/10.1038/s41598-023-40400-5>

1396 Clark, S. K., Brenowitz, N. D., Henn, B., Kwa, A., McGibbon, J., Perkins, W. A., Watt-Meyer, O., Bretherton, C. S., & Harris,  
1397 L. M. (2022). Correcting a 200 km Resolution Climate Model in Multiple Climates by Machine Learning From 25 km  
1398 Resolution Simulations. *Journal of Advances in Modeling Earth Systems*, *14*(9). <https://doi.org/10.1029/2022MS003219>

1399 Clarke, K. C., Hoppen, S., & Gaydos, L. (1997). A self-modifying cellular automaton model of historical urbanization in the  
1400 San Francisco Bay area. *Environment and Planning B: Planning and Design*, *24*(2), 247–261.  
1401 <https://doi.org/10.1068/b240247>

1402 Côté, J.-N., Germain, M., Levac, E., & Lavigne, E. (2024). Vulnerability assessment of heat waves within a risk framework  
1403 using artificial intelligence. *Science of The Total Environment*, *912*, 169355.  
1404 <https://doi.org/10.1016/j.scitotenv.2023.169355>

Formatted: Font: Italic

1405 Couason, A., Sebastian, A., & Morales-Nápoles, O. (2018). A Copula-Based Bayesian Network for Modeling Compound  
1406 Flood Hazard from Riverine and Coastal Interactions at the Catchment Scale: An Application to the Houston Ship  
1407 Channel, Texas. *Water*, 10(9), 1190. <https://doi.org/10.3390/w10091190>

1408 Cushman, S. A., Macdonald, E. A., Landguth, E. L., Malhi, Y., & Macdonald, D. W. (2017). Multiple-scale prediction of  
1409 forest loss risk across Borneo. *Landscape Ecology*, 32(8), 1581–1598. <https://doi.org/10.1007/s10980-017-0520-0>

1410 Dal Barco, M. K., Maraschini, M., Ferrario, D. M., Nguyen, N. D., Torresan, S., Vascon, S., & Critto, A. (2024). A machine  
1411 learning approach to evaluate coastal risks related to extreme weather events in the Veneto region (Italy). *International  
1412 Journal of Disaster Risk Reduction*, 108, 104526. <https://doi.org/10.1016/j.ijdr.2024.104526>

1413 Dasgupta, A., Hybbeneth, L., & Waske, B. (2022). *Towards Daily High-resolution Inundation Observations using Deep  
1414 Learning and EO*. <http://arxiv.org/abs/2208.09135>.

1415 Dawkins, L. C., Bernie, D. J., Pianosi, F., Lowe, J. A., & Economou T., (2023). Quantifying Uncertainty and Sensitivity in  
1416 Climate Risk Assessments: Varying Hazard, Exposure and Vulnerability Modelling Choices. *Climate Risk Management  
1417 40: 100511*. <https://doi.org/10.1016/j.crm.2023.100511>.

1418 De Angeli, S., Malamud, B. D., Rossi, L., Taylor, F. E., Trasforini, E., & Rudari, R. (2022). A multi-hazard framework for  
1419 spatial-temporal impact analysis. *International Journal of Disaster Risk Reduction*, 73, 102829.  
1420 <https://doi.org/10.1016/j.ijdr.2022.102829>

1421 de Ruiter, M. C., & van Loon, A. F. (2022). The challenges of dynamic vulnerability and how to assess it. *IScience*, 25(8),  
1422 104720. <https://doi.org/10.1016/j.isci.2022.104720>

1423 Di Martino, F., Pedrycz, W., & Sessa, S. (2018a). Spatiotemporal extended fuzzy C-means clustering algorithm for hotspots  
1424 detection and prediction. *Fuzzy Sets and Systems*, 340, 109–126. <https://doi.org/10.1016/j.fss.2017.11.011>

1425 Di Martino, F., Pedrycz, W., & Sessa, S. (2018b). Spatiotemporal extended fuzzy C-means clustering algorithm for hotspots  
1426 detection and prediction. *Fuzzy Sets and Systems*, 340, 109–126. <https://doi.org/10.1016/j.fss.2017.11.011>

1427 Du, M., Liu, N., Yang, F., Ji, S., & Hu, X. (2019). On Attribution of Recurrent Neural Network Predictions via Additive  
1428 Decomposition. *The World Wide Web Conference*, 383–393. <https://doi.org/10.1145/3308558.3313545>

1429 Erion, G., Janizek, J. D., Sturmfels, P., Lundberg, S. M., & Lee, S.-I. (2021). Improving performance of deep learning models  
1430 with axiomatic attribution priors and expected gradients. *Nature Machine Intelligence*, 3(7), 620–631.  
1431 <https://doi.org/10.1038/s42256-021-00343-w>

1432 Ester, M., Kriegel, H.-P., Sander, J., & Xu, X. (1996). A Density-Based Algorithm for Discovering Clusters in Large Spatial  
1433 Databases with Noise. *KDD-96 Proceedings*.

1434 Faiza, B., Yuhaniz, S. S., Hashim, S. Z. M., & AbdulRahman, K. K. (2012). A Review and Analysis of Image Misalignment  
1435 Problem in Remote Sensing. *International Journal of Scientific and Engineering Research*, 3(3), 82–86.

1436 Fang, K., Shen, C., Kifer, D., & Yang, X. (2017). Prolongation of SMAP to Spatiotemporally Seamless Coverage of  
1437 Continental U.S. Using a Deep Learning Neural Network. *Geophysical Research Letters*, 44(21).  
1438 <https://doi.org/10.1002/2017GL075619>

Formatted: Italian

Formatted: Italian

Formatted: Font: Italic, Italian

Formatted: Italian

1439 Flora, M. L., Potvin, C. K., Skinner, P. S., Handler, S., & McGovern, A. (2021). Using Machine Learning to Generate Storm-  
1440 Scale Probabilistic Guidance of Severe Weather Hazards in the Warn-on-Forecast System. *Monthly Weather Review*,  
1441 *149*(5), 1535–1557. <https://doi.org/10.1175/MWR-D-20-0194.1>

1442 Freeman, B. S., Taylor, G., Gharabaghi, B., & Thé, J. (2018). Forecasting air quality time series using deep learning. *Journal*  
1443 *of the Air & Waste Management Association*, *68*(8), 866–886. <https://doi.org/10.1080/10962247.2018.1459956>

1444 Fuchs, S., Keiler, M., & Zischg, A. (2015). A spatiotemporal multi-hazard exposure assessment based on property data.  
1445 *Natural Hazards and Earth System Sciences*, *15*(9), 2127–2142. <https://doi.org/10.5194/nhess-15-2127-2015>

1446 Gallina, V., Torresan, S., Zabeo, A., Critto, A., Glade, T., & Marcomini, A. (2020). A multi-risk methodology for the  
1447 assessment of climate change impacts in coastal zones. *Sustainability (Switzerland)*, *12*(9).  
1448 <https://doi.org/10.3390/su12093697>

1449 García-León, D., Masselot, P., Mistry, M. N., Gasparrini, A., Motta, C., Feyen, L., & Ciscar, J.-C. (2024). Temperature-related  
1450 mortality burden and projected change in 1368 European regions: a modelling study. *The Lancet Public Health*, *9*(9),  
1451 e644–e653. [https://doi.org/10.1016/S2468-2667\(24\)00179-8](https://doi.org/10.1016/S2468-2667(24)00179-8)

1452 Garg, S., Rasp, S., & Thuerey, N. (2022). *WeatherBench Probability: A benchmark dataset for probabilistic medium-range*  
1453 *weather forecasting along with deep learning baseline models*. <http://arxiv.org/abs/2205.00865>

1454 Garnelo, M., Schwarz, J., Rosenbaum, D., Viola, F., Rezende, D. J., Eslami, S. M. A., & Teh, Y. W. (2018). *Neural Processes*.  
1455 <http://arxiv.org/abs/1807.01622>

1456 Gasparrini, A. (2014). Modeling exposure–lag–response associations with distributed lag non-linear models. *Statistics in*  
1457 *Medicine*, *33*(5), 881–899. <https://doi.org/10.1002/sim.5963>

1458 Genkin, A., Lewis, D. D., & Madigan, D. (2007). Large-Scale Bayesian Logistic Regression for Text Categorization.  
1459 *Technometrics*, *49*(3), 291–304. <https://doi.org/10.1198/004017007000000245>

1460 Ghaffarian, S., & Emtehani, S. (2021). Monitoring Urban Deprived Areas with Remote Sensing and Machine Learning in Case  
1461 of Disaster Recovery. *Climate*, *9*(4), 58. <https://doi.org/10.3390/cli9040058>

1462 Ghaffarian, S., Taghikhah, F. R., & Maier, H. R. (2023). Explainable artificial intelligence in disaster risk management:  
1463 Achievements and prospective futures. *International Journal of Disaster Risk Reduction*, *98*, 104123.  
1464 <https://doi.org/10.1016/j.ijdr.2023.104123>

1465 Ghanbari, M., Arabi, M., Kao, S., Obeysekera, J., & Sweet, W. (2021). Climate Change and Changes in Compound Coastal-  
1466 Riverine Flooding Hazard Along the U.S. Coasts. *Earth's Future*, *9*(5). <https://doi.org/10.1029/2021EF002055>

1467 Ghiggi, G., Humphrey, V., Seneviratne, S. I., & Gudmundsson, L. (2019). GRUN: an observation-based global gridded runoff  
1468 dataset from 1902 to 2014. *Earth System Science Data*, *11*(4), 1655–1674. <https://doi.org/10.5194/essd-11-1655-2019>

1469 Gierszewska, M., & Berezowski, T. (2024). *A physics-guided neural network for flooding area detection using SAR imagery*  
1470 *and local river gauge observations*. <http://arxiv.org/abs/2410.08837>

1471 Guo, Q., Mistry, M. N., Zhou, X., Zhao, G., Kino, K., Wen, B., Yoshimura, K., Satoh, Y., Cvijanovic, I., Kim, Y., Ng, C. F.  
1472 S., Vicedo-Cabrera, A. M., Armstrong, B., Urban, A., Katsouyanni, K., Masselot, P., Tong, S., Sera, F., Huber, V., ...

- 1473 Honda, Y. (2024). Regional variation in the role of humidity on city-level heat-related mortality. *PNAS Nexus*, 3(8).  
1474 <https://doi.org/10.1093/pnasnexus/pgae290>
- 1475 Haer, T., Botzen, W.J.W., & Aerts, J.C.J.H. (2019). Advancing disaster policies by integrating dynamic adaptive behaviour in  
1476 risk assessments using an agent-based modelling approach. *Environmental Research Letters*, 14(4), 044022.  
1477 <https://doi.org/10.1088/1748-9326/ab077>
- 1478 Haggag, M., Siam, A. S., El-Dakhkhni, W., Coulibaly, P., & Hassini, E. (2021). A deep learning model for predicting climate-  
1479 induced disasters. *Natural Hazards*, 107(1), 1009–1034. <https://doi.org/10.1007/s11069-021-04620-0>
- 1480 Han, Q., Zeng, Y., Zhang, L., Wang, C., Prikaziuk, E., Niu, Z., & Su, B. (2023). Global long term daily 1 km surface soil  
1481 moisture dataset with physics informed machine learning. *Scientific Data*, 10(1), 101. <https://doi.org/10.1038/s41597-023-02011-7>
- 1482
- 1483 Hao, Z., & Singh, V. P. (2016). Review of dependence modeling in hydrology and water resources. *Progress in Physical  
1484 Geography: Earth and Environment*, 40(4), 549–578. <https://doi.org/10.1177/0309133316632460>
- 1485 Harris, L., McRae, A. T. T., Chantry, M., Dueben, P. D., & Palmer, T. N. (2022). A Generative Deep Learning Approach to  
1486 Stochastic Downscaling of Precipitation Forecasts. *Journal of Advances in Modeling Earth Systems*, 14(10).  
1487 <https://doi.org/10.1029/2022MS003120>
- 1488 Hawkins, E., & Sutton, R. (2009). The Potential to Narrow Uncertainty in Regional Climate Predictions. *Bulletin of the  
1489 American Meteorological Society*, 90(8), 1095–1108. <https://doi.org/10.1175/2009BAMS2607.1>
- 1490 He, X., Chaney, N. W., Schleiss, M., & Sheffield, J. (2016). Spatial downscaling of precipitation using adaptable random  
1491 forests. *Water Resources Research*, 52(10), 8217–8237. <https://doi.org/10.1002/2016WR019034>
- 1492 He, X., Li, Y., Liu, S., Xu, T., Chen, F., Li, Z., Zhang, Z., Liu, R., Song, L., Xu, Z., Peng, Z., & Zheng, C. (2023). Improving  
1493 regional climate simulations based on a hybrid data assimilation and machine learning method. *Hydrology and Earth  
1494 System Sciences*, 27(7), 1583–1606. <https://doi.org/10.5194/hess-27-1583-2023>
- 1495 Hoch, J. M., de Bruin, S. P., Buhaug, H., Von Uexkull, N., van Beek, R., & Wanders, N. (2021). Projecting armed conflict  
1496 risk in Africa towards 2050 along the SSP-RCP scenarios: a machine learning approach. *Environmental Research  
1497 Letters*, 16(12), 124068. <https://doi.org/10.1088/1748-9326/ac3db2>
- 1498 Hochrainer-Stigler, S., Balkovič, J., Silm, K., & Timonina-Farkas, A. (2019a). Large scale extreme risk assessment using  
1499 copulas: an application to drought events under climate change for Austria. *Computational Management Science*, 16(4),  
1500 651–669. <https://doi.org/10.1007/s10287-018-0339-4>
- 1501 Hochrainer-Stigler, S., Balkovič, J., Silm, K., & Timonina-Farkas, A. (2019b). Large scale extreme risk assessment using  
1502 copulas: an application to drought events under climate change for Austria. *Computational Management Science*, 16(4),  
1503 651–669. <https://doi.org/10.1007/s10287-018-0339-4>
- 1504 Huynh, N. N. T., Garambois, P.-A., Renard, B., Colleoni, F., Monnier, J., & Roux, H. (2025). A distributed hybrid physics-  
1505 AI framework for learning corrections of internal hydrological fluxes and enhancing high-resolution regionalized flood  
1506 modeling. *Hydrology and Earth System Sciences*, 29(15), 3589–3613. <https://doi.org/10.5194/hess-29-3589-2025>

1507 Ionita, M., Caldarescu, D. E., & Nagavciuc, V. (2021). Compound Hot and Dry Events in Europe: Variability and Large-Scale  
1508 Drivers. *Frontiers in Climate*, 3. <https://doi.org/10.3389/fclim.2021.688991>

1509 Islam, A. R. Md. T., Talukdar, S., Mahato, S., Ziaul, S., Eibek, K. U., Akhter, S., Pham, Q. B., Mohammadi, B., Karimi, F., &  
1510 Linh, N. T. T. (2021). Machine learning algorithm-based risk assessment of riparian wetlands in Padma River Basin of  
1511 Northwest Bangladesh. *Environmental Science and Pollution Research*, 28(26), 34450–34471.  
1512 <https://doi.org/10.1007/s11356-021-12806-z>

1513 Jain, P., Coogan, S. C. P., Subramanian, S. G., Crowley, M., Taylor, S., & Flannigan, M. D. (2020). *A review of machine*  
1514 *learning applications in wildfire science and management*. <https://doi.org/10.1139/er-2020-0019>

1515 Janizadeh, S., Chandra Pal, S., Saha, A., Chowdhuri, I., Ahmadi, K., Mirzaei, S., Mosavi, A. H., & Tiefenbacher, J. P. (2021).  
1516 Mapping the spatial and temporal variability of flood hazard affected by climate and land-use changes in the future.  
1517 *Journal of Environmental Management*, 298, 113551. <https://doi.org/10.1016/j.jenvman.2021.113551>

1518 Javidan, N., Kavian, A., Pourghasemi, H. R., Conoscenti, C., Jafarian, Z., & Rodrigo-Comino, J. (2021). Evaluation of multi-  
1519 hazard map produced using MaxEnt machine learning technique. *Scientific Reports*, 11(1), 6496.  
1520 <https://doi.org/10.1038/s41598-021-85862-7>

1521 Jean, N., Burke, M., Xie, M., Davis, W. M., Lobell, D. B., & Ermon, S. (2016). Combining satellite imagery and machine  
1522 learning to predict poverty. *Science*, 353(6301), 790–794. <https://doi.org/10.1126/science.aaf7894>

1523 Ji, Y., Sri Sumantyo, J., Chua, M., & Waqar, M. (2018). Earthquake/Tsunami Damage Assessment for Urban Areas Using  
1524 Post-Event PolSAR Data. *Remote Sensing*, 10(7), 1088. <https://doi.org/10.3390/rs10071088>

1525 Jiang, L., Li, C., Wang, S., & Zhang, L. (2016). Deep feature weighting for naive Bayes and its application to text classification.  
1526 *Engineering Applications of Artificial Intelligence*, 52, 26–39. <https://doi.org/10.1016/j.engappai.2016.02.002>

1527 Jiang, M., Pedrielli, G., & Ng, S. H. (2022). Gaussian Processes for High-Dimensional, Large Data Sets: A Review. 2022  
1528 *Winter Simulation Conference (WSC)*, 49–60. <https://doi.org/10.1109/WSC57314.2022.10015416>

1529 Jiang, S., Bevacqua, E., & Zscheischler, J. (2022). River flooding mechanisms and their changes in Europe revealed by  
1530 explainable machine learning. *Hydrology and Earth System Sciences*, 26(24), 6339–6359. [https://doi.org/10.5194/hess-](https://doi.org/10.5194/hess-26-6339-2022)  
1531 [26-6339-2022](https://doi.org/10.5194/hess-26-6339-2022)

1532 Jiang, S., Sweet, L., Blougouras, G., Brenning, A., Li, W., Reichstein, M., Denzler, J., Shanguan, W., Yu, G., Huang, F., &  
1533 Zscheischler, J. (2024). How Interpretable Machine Learning Can Benefit Process Understanding in the Geosciences.  
1534 *Earth's Future*, 12(7). <https://doi.org/10.1029/2024EF004540>

1535 Jiang, S., Zheng, Y., Wang, C., & Babovic, V. (2022). Uncovering Flooding Mechanisms Across the Contiguous United States  
1536 Through Interpretive Deep Learning on Representative Catchments. *Water Resources Research*, 58(1).  
1537 <https://doi.org/10.1029/2021WR030185>

1538 Jiang, T., Su, X., Zhang, G., Zhang, T., & Wu, H. (2023). Estimating propagation probability from meteorological to ecological  
1539 droughts using a hybrid machine learning copula method. *Hydrology and Earth System Sciences*, 27(2), 559–576.  
1540 <https://doi.org/10.5194/hess-27-559-2023>

1541 Jiang, W., Chen, Z., Lei, X., Jia, K., & Wu, Y. (2015). Simulating urban land use change by incorporating an autologistic  
1542 regression model into a CLUE-S model. *Journal of Geographical Sciences*, 25(7), 836–850.  
1543 <https://doi.org/10.1007/s11442-015-1205-8>

1544 Jing, W., Yang, Y., Yue, X., & Zhao, X. (2016a). A Comparison of Different Regression Algorithms for Downscaling Monthly  
1545 Satellite-Based Precipitation over North China. *Remote Sensing*, 8(10), 835. <https://doi.org/10.3390/rs8100835>

1546 Jing, W., Yang, Y., Yue, X., & Zhao, X. (2016b). A Spatial Downscaling Algorithm for Satellite-Based Precipitation over the  
1547 Tibetan Plateau Based on NDVI, DEM, and Land Surface Temperature. *Remote Sensing*, 8(8), 655.  
1548 <https://doi.org/10.3390/rs8080655>

1549 Kabiru, P., Kuffler, M., Sliuzas, R., & Vanhuyse, S. (2023). The relationship between multiple hazards and deprivation using  
1550 open geospatial data and machine learning. *Natural Hazards*, 119(2), 907–941. <https://doi.org/10.1007/s11069-023-1551-05897-z>

1552 Kang, J., Jin, R., Li, X., Zhang, Y., & Zhu, Z. (2018). Spatial Upscaling of Sparse Soil Moisture Observations Based on Ridge  
1553 Regression. *Remote Sensing*, 10(2), 192. <https://doi.org/10.3390/rs10020192>

1554 Karakas, G., Kocaman, S., & Gokceoglu, C. (2023). A Hybrid Multi-Hazard Susceptibility Assessment Model for a Basin in  
1555 Elazig Province, Türkiye. *International Journal of Disaster Risk Science*, 14(2), 326–341.  
1556 <https://doi.org/10.1007/s13753-023-00477-y>

1557 Kariminejad, N., Pourghasemi, H. R., & Hosseinalizadeh, M. (2022). Analytical techniques for mapping multi-hazard with  
1558 geo-environmental modeling approaches and UAV images. *Scientific Reports*, 12(1), 14946.  
1559 <https://doi.org/10.1038/s41598-022-18757-w>

1560 Kashinath, K., Mustafa, M., Albert, A., Wu, J.-L., Jiang, C., Esmacilzadeh, S., Azizzadenesheli, K., Wang, R., Chattopadhyay,  
1561 A., Singh, A., Manepalli, A., Chirila, D., Yu, R., Walters, R., White, B., Xiao, H., Tchelepi, H. A., Marcus, P.,  
1562 Anandkumar, A., ... Prabhat. (2021a). Physics-informed machine learning: case studies for weather and climate  
1563 modelling. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*,  
1564 379(2194), 20200093. <https://doi.org/10.1098/rsta.2020.0093>

1565 Kashinath, K., Mustafa, M., Albert, A., Wu, J.-L., Jiang, C., Esmacilzadeh, S., Azizzadenesheli, K., Wang, R., Chattopadhyay,  
1566 A., Singh, A., Manepalli, A., Chirila, D., Yu, R., Walters, R., White, B., Xiao, H., Tchelepi, H. A., Marcus, P.,  
1567 Anandkumar, A., ... Prabhat. (2021b). Physics-informed machine learning: case studies for weather and climate  
1568 modelling. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*,  
1569 379(2194), 20200093. <https://doi.org/10.1098/rsta.2020.0093>

1570 Kazadi, A., Doss-Gollin, J., Sebastian, A., & Silva, A. (2024). FloodGNN-GRU: a spatio-temporal graph neural network for  
1571 flood prediction. *Environmental Data Science*, 3, e21. <https://doi.org/10.1017/eds.2024.19>

1572 Keisler, R. (2022). *Forecasting Global Weather with Graph Neural Networks*. <https://doi.org/arXiv:2202.07575v1>

1573 Khan, F., Spöck, G., Liou, Y.-A., & Ali, S. (2024). Association of precipitation extremes and crops production and projecting  
1574 future extremes using machine learning approaches with CMIP6 data. *Environmental Science and Pollution Research*,  
1575 31(42), 54979–54999. <https://doi.org/10.1007/s11356-024-34652-5>

1576 Khatakho, R., Gautam, D., Aryal, K. R., Pandey, V. P., Rupakhety, R., Lamichhane, S., Liu, Y.-C., Abdouli, K., Talchabhadel,  
1577 R., Thapa, B. R., & Adhikari, R. (2021). Multi-Hazard Risk Assessment of Kathmandu Valley, Nepal. *Sustainability*,  
1578 13(10), 5369. <https://doi.org/10.3390/su13105369>

1579 Kim, Y., Evans, J. P., & Sharma, A. (2023). Correcting biases in regional climate model boundary variables for improved  
1580 simulation of high-impact compound events. *IScience*, 26(9), 107696. <https://doi.org/10.1016/j.isci.2023.107696>

1581 Kipf, T. N., & Welling, M. (2016). *Semi-Supervised Classification with Graph Convolutional Networks*.

1582 Koshy, R., & Elango, S. (2023). Multimodal tweet classification in disaster response systems using transformer-based  
1583 bidirectional attention model. *Neural Computing and Applications*, 35(2), 1607–1627. [https://doi.org/10.1007/s00521-](https://doi.org/10.1007/s00521-022-07790-5)  
1584 022-07790-5

1585 Kotaridis, I., & Lazaridou, M. (2022). Integration of convolutional neural networks for flood risk mapping in Tuscany, Italy.  
1586 *Natural Hazards*, 114(3), 3409–3424. <https://doi.org/10.1007/s11069-022-05525-2>

1587 Kraft, B., Jung, M., Körner, M., Requena Mesa, C., Cortés, J., & Reichstein, M. (2019). Identifying Dynamic Memory Effects  
1588 on Vegetation State Using Recurrent Neural Networks. *Frontiers in Big Data*, 2.  
1589 <https://doi.org/10.3389/fdata.2019.00031>

1590 Kratzert, F., Klotz, D., Brandstetter, J., Hoedt, P.-J., Nearing, G., & Hochreiter, S. (2019). *Using LSTMs for climate change*  
1591 *assessment studies on droughts and floods*. <https://doi.org/https://doi.org/10.48550/arXiv.1911.03941>

1592 Kratzert, F., Klotz, D., Brenner, C., Schulz, K., & Herrnegger, M. (2018). Rainfall–runoff modelling using Long Short-Term  
1593 Memory (LSTM) networks. *Hydrology and Earth System Sciences*, 22(11), 6005–6022. [https://doi.org/10.5194/hess-22-](https://doi.org/10.5194/hess-22-6005-2018)  
1594 6005-2018

1595 Kratzert, F., Klotz, D., Shalev, G., Klambauer, G., Hochreiter, S., & Nearing, G. (2019). Towards learning universal, regional,  
1596 and local hydrological behaviors via machine learning applied to large-sample datasets. *Hydrology and Earth System*  
1597 *Sciences*, 23(12), 5089–5110. <https://doi.org/10.5194/hess-23-5089-2019>

1598 Kropf, C. M., Ciullo, A., Otth, L., Meiler, S., Rana, A., Schmid, E., McCaughey, J. W., & Bresch, D. N. (2022). Uncertainty  
1599 and Sensitivity Analysis for Probabilistic Weather and Climate-Risk Modelling: An Implementation in CLIMADA  
1600 v.3.1.0. *Geoscientific Model Development* 15 (18): 7177–201. <https://doi.org/10.5194/gmd-15-7177-2022>.

1601 Lam, R., Sanchez-Gonzalez, A., Willson, M., Wirnsberger, P., Fortunato, M., Pritzel, A., Ravuri, S., Ewalds, T., Alet, F.,  
1602 Eaton-Rosen, Z., Hu, W., Merose, A., Hoyer, S., Holland, G., Stott, J., Vinyals, O., Mohamed, S., Battaglia, P., &  
1603 contribution, equal. (2022). *GraphCast: Learning skillful medium-range global weather forecasting*.

1604 Lecun, Y., Bottou, L., Bengio, Y., & Haffner, P. (1998). Gradient-based learning applied to document recognition. *Proceedings*  
1605 *of the IEEE*, 86(11), 2278–2324. <https://doi.org/10.1109/5.726791>

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Formatted: Font: Italic

1606 Lei, T., Zhang, Q., Xue, D., Chen, T., Meng, H., & Nandi, A. K. (2019). End-to-end Change Detection Using a Symmetric  
1607 Fully Convolutional Network for Landslide Mapping. *ICASSP 2019 - 2019 IEEE International Conference on Acoustics,  
1608 Speech and Signal Processing (ICASSP)*, 3027–3031. <https://doi.org/10.1109/ICASSP.2019.8682802>

1609 Leng, G., & Hall, J. W. (2020). Predicting spatial and temporal variability in crop yields: an inter-comparison of machine  
1610 learning, regression and process-based models. *Environmental Research Letters*, *15*(4), 044027.  
1611 <https://doi.org/10.1088/1748-9326/ab7b24>

1612 Li, L., Qiao, J., Yu, G., Wang, L., Li, H.-Y., Liao, C., & Zhu, Z. (2022). Interpretable tree-based ensemble model for predicting  
1613 beach water quality. *Water Research*, *211*, 118078. <https://doi.org/10.1016/j.watres.2022.118078>

1614 Liang, X., Guan, Q., Clarke, K. C., Liu, S., Wang, B., & Yao, Y. (2021). Understanding the drivers of sustainable land  
1615 expansion using a patch-generating land use simulation (PLUS) model: A case study in Wuhan, China. *Computers,  
1616 Environment and Urban Systems*, *85*, 101569. <https://doi.org/10.1016/j.compenvurbsys.2020.101569>

1617 Liao, Z., Chen, Y., Li, W., & Zhai, P. (2021). Growing Threats From Unprecedented Sequential Flood-Hot Extremes Across  
1618 China. *Geophysical Research Letters*, *48*(18). <https://doi.org/10.1029/2021GL094505>

1619 Lim, C.-H., & Kim, H.-J. (2022). Can Forest-Related Adaptive Capacity Reduce Landslide Risk Attributable to Climate  
1620 Change?—Case of Republic of Korea. *Forests*, *13*(1), 49. <https://doi.org/10.3390/f13010049>

1621 Lin, H., Tang, J., Wang, S., Wang, S., & Dong, G. (2023). Deep learning downscaled high-resolution daily near surface  
1622 meteorological datasets over East Asia. *Scientific Data*, *10*(1), 890. <https://doi.org/10.1038/s41597-023-02805-9>

1623 Ling, F., Lu, Z., Luo, J.-J., Bai, L., Behera, S. K., Jin, D., Pan, B., Jiang, H., & Yamagata, T. (2024a). Diffusion model-based  
1624 probabilistic downscaling for 180-year East Asian climate reconstruction. *Npj Climate and Atmospheric Science*, *7*(1),  
1625 131. <https://doi.org/10.1038/s41612-024-00679-1>

1626 Ling, F., Lu, Z., Luo, J.-J., Bai, L., Behera, S. K., Jin, D., Pan, B., Jiang, H., & Yamagata, T. (2024b). Diffusion model-based  
1627 probabilistic downscaling for 180-year East Asian climate reconstruction. *Npj Climate and Atmospheric Science*, *7*(1),  
1628 131. <https://doi.org/10.1038/s41612-024-00679-1>

1629 Liu, G., Gao, Z., Chen, B., Fu, H., Jiang, S., Wang, L., Wang, G., & Chen, Z. (2020). Extreme values of storm surge elevation  
1630 in Hangzhou Bay. *Ships and Offshore Structures*, *15*(4), 431–442. <https://doi.org/10.1080/17445302.2019.1661618>

1631 Liu, G., Yang, B., Nong, X., Kou, Y., Wu, F., Zhao, D., & Yu, P. (2023). Risk Level Assessment of Typhoon Hazard Based  
1632 on Loss Utility. *Journal of Marine Science and Engineering*, *11*(11), 2177. <https://doi.org/10.3390/jmse11112177>

1633 Liu, J., Qiu, Z., Feng, J., Wong, K. P., Tsou, J. Y., Wang, Y., & Zhang, Y. (2023). Monitoring Total Suspended Solids and  
1634 Chlorophyll-a Concentrations in Turbid Waters: A Case Study of the Pearl River Estuary and Coast Using Machine  
1635 Learning. *Remote Sensing*, *15*(23), 5559. <https://doi.org/10.3390/rs15235559>

1636 Liu, K., Wang, M., Cao, Y., Zhu, W., & Yang, G. (2018). Susceptibility of existing and planned Chinese railway system  
1637 subjected to rainfall-induced multi-hazards. *Transportation Research Part A: Policy and Practice*, *117*, 214–226.  
1638 <https://doi.org/10.1016/j.tra.2018.08.030>

- 1639 Liu, X., Guo, H., Lin, Y., Li, Y., & Hou, J. (2018). Analyzing Spatial-Temporal Distribution of Natural Hazards in China by  
 1640 Mining News Sources. *Natural Hazards Review*, 19(3). [https://doi.org/10.1061/\(ASCE\)NH.1527-6996.0000291](https://doi.org/10.1061/(ASCE)NH.1527-6996.0000291)
- 1641 Liu, Y., Racah, E., Prabhat, Correa, J., Khosrowshahi, A., Lavers, D., Kunkel, K., Wehner, M., & Collins, W. (2016).  
 1642 *Application of Deep Convolutional Neural Networks for Detecting Extreme Weather in Climate Datasets*.  
 1643 <https://doi.org/10.1101/011156>
- 1644 Lütjens, B., Crawford, C. H., Veillette, M., & Newman, D. (2021). *PCE-PINNs: Physics-Informed Neural Networks for  
 1645 Uncertainty Propagation in Ocean Modeling*.
- 1646 Luu, C., Forino, G., Yorke, L., Ha, H., Bui, Q. D., Tran, H. H., Nguyen, D. Q., Duong, H. C., & Kervyn, M. (2024). *Integrating  
 1647 multi-hazard susceptibility and building exposure: A case study for Quang Nam province, Vietnam*.  
 1648 <https://doi.org/10.5194/egusphere-2024-57>
- 1649 Macdonald, E., Tubaldi, E., & Patelli, E. (2025). Robust storm surge forecasts for early warning system: a machine learning  
 1650 approach using Monte Carlo Bayesian model selection algorithm. *Stochastic Environmental Research and Risk  
 1651 Assessment*, 39(7), 2789–2816. <https://doi.org/10.1007/s00477-025-02993-3>
- 1652 Mandal, P., Maiti, A., Paul, S., Bhattacharya, S., & Paul, S. (2022). Mapping the multi-hazards risk index for coastal block of  
 1653 Sundarban, India using AHP and machine learning algorithms. *Tropical Cyclone Research and Review*, 11(4), 225–243.  
 1654 <https://doi.org/10.1016/j.terr.2023.03.001>
- 1655 Mattei, G., Di Luccio, D., Benassai, G., Anfuso, G., Budillon, G., & Aucelli, P. (2021). Characteristics and coastal effects of  
 1656 a destructive marine storm in the Gulf of Naples (southern Italy). *Natural Hazards and Earth System Sciences*, 21(12),  
 1657 3809–3825. <https://doi.org/10.5194/nhess-21-3809-2021>
- 1658 McGovern, A., Ebert-Uphoff, I., Gagne, D. J., & Bostrom, A. (2022). Why we need to focus on developing ethical, responsible,  
 1659 and trustworthy artificial intelligence approaches for environmental science. *Environmental Data Science*, 1, e6.  
 1660 <https://doi.org/10.1017/eds.2022.5>
- 1661 McGovern, A., Lagerquist, R., John Gagne, D., Jergensen, G. E., Elmore, K. L., Homeyer, C. R., & Smith, T. (2019). Making  
 1662 the Black Box More Transparent: Understanding the Physical Implications of Machine Learning. *Bulletin of the  
 1663 American Meteorological Society*, 100(11), 2175–2199. <https://doi.org/10.1175/BAMS-D-18-0195.1>
- 1664 Mehrotra, H., Mishra, A., & Pal, S. (2022). A Multi-stage Classification Framework for Disaster-Specific Tweets. *SN  
 1665 Computer Science*, 3(1), 24. <https://doi.org/10.1007/s42979-021-00930-z>
- 1666 Miyoshi, G. T., Arruda, M. dos S., Osco, L. P., Marcato Junior, J., Gonçalves, D. N., Imai, N. N., Tommaselli, A. M. G.,  
 1667 Honkavaara, E., & Gonçalves, W. N. (2020). A Novel Deep Learning Method to Identify Single Tree Species in UAV-  
 1668 Based Hyperspectral Images. *Remote Sensing*, 12(8), 1294. <https://doi.org/10.3390/rs12081294>
- 1669
- 1670 Mls K, Kořinek M, Štekerová K, Tučník P, Bureš V, Čech P, Husáková M, Mikulecký P, Nacházel T, Ponce D, Zanker M,  
 1671 Babič F, Triantafyllou I. (2023). Agent-based models of human response to natural hazards: systematic review of tsunami  
 1672 evacuation. *Nat Hazards (Dordr)*, 115(3), 1887-1908. <https://doi.org/10.1007/s11069-022-05643-x>.

Formatted: Font: Italic

- 1673 Moezzi, M., Janda, K. B., & Rotmann, S. (2017). Using stories, narratives, and storytelling in energy and climate change  
1674 research. *Energy Research & Social Science*, 31, 1–10. <https://doi.org/10.1016/j.erss.2017.06.034>
- 1675 Mukherjee, S., Nateghi, R., & Hastak, M. (2018). A multi-hazard approach to assess severe weather-induced major power  
1676 outage risks in the U.S. *Reliability Engineering & System Safety*, 175, 283–305.  
1677 <https://doi.org/10.1016/j.res.2018.03.015>
- 1678 Munawar, H. S., Ullah, F., Qayyum, S., Khan, S. I., & Mojtahedi, M. (2021). UAVs in Disaster Management: Application of  
1679 Integrated Aerial Imagery and Convolutional Neural Network for Flood Detection. *Sustainability*, 13(14), 7547.  
1680 <https://doi.org/10.3390/su13147547>
- 1681 Nam, J., Kim, J., Loza Mencía, E., Gurevych, I., & Fürnkranz, J. (2014). *Large-Scale Multi-label Text Classification —*  
1682 *Revisiting Neural Networks* (pp. 437–452). [https://doi.org/10.1007/978-3-662-44851-9\\_28](https://doi.org/10.1007/978-3-662-44851-9_28)
- 1683 Naudé, W., & Vinuesa, R. (2021). Data Deprivations, Data Gaps and Digital Divides: Lessons from the COVID-19 Pandemic.  
1684 *Big Data & Society* 8 (2); 20539517211025545. <https://doi.org/10.1177/20539517211025545>.
- 1685 Nazeer, M., Bilal, M., Alsahli, M., Shahzad, M., & Waqas, A. (2017). Evaluation of Empirical and Machine Learning  
1686 Algorithms for Estimation of Coastal Water Quality Parameters. *ISPRS International Journal of Geo-Information*, 6(11),  
1687 360. <https://doi.org/10.3390/ijgi6110360>
- 1688 Nelsen, R. (2006). *An Introduction to Copulas*. Springer New York. <https://doi.org/10.1007/0-387-28678-0>
- 1689 Nguyen, H. D., Dang, D., Bui, Q., & Petrisor, A. (2023). Multi-hazard assessment using machine learning and remote sensing  
1690 in the North Central region of Vietnam. *Transactions in GIS*, 27(5), 1614–1640. <https://doi.org/10.1111/tgis.13091>
- 1691 Nguyen, M., Wilson, M., Lane, E., Brasington, J., & Pearson, R. (2024). Estimating uncertainty in flood model outputs using  
1692 machine learning informed by Monte Carlo analysis. *2024 International Conference on Machine Intelligence for*  
1693 *GeoAnalytics and Remote Sensing (MIGARS)*, 1–3. <https://doi.org/10.1109/MIGARS61408.2024.10544837>
- 1694 Novellino, A., Pennington, C., Leeming, K., Taylor, S., Alvarez, I. G., McAllister, E., Amhardt, C., & Winson, A. (2024).  
1695 Mapping landslides from space: A review. *Landslides*, 21(5), 1041–1052. <https://doi.org/10.1007/s10346-024-02215-x>
- 1696 O., S., & Orth, R. (2021). Global soil moisture data derived through machine learning trained with in-situ measurements.  
1697 *Scientific Data*, 8(1), 170. <https://doi.org/10.1038/s41597-021-00964-1>
- 1698 O’Dea, R. E., Lagisz, M., Jennions, M. D., Koricheva, J., Noble, D. W. A., Parker, T. H., Gurevitch, J., Page, M. J., Stewart,  
1699 G., Moher, D., & Nakagawa, S. (2021). Preferred reporting items for systematic reviews and meta-analyses in ecology  
1700 and evolutionary biology: a <sc>PRISMA</sc> extension. *Biological Reviews*, 96(5), 1695–1722.  
1701 <https://doi.org/10.1111/brv.12721>
- 1702 Oh, D. H., & Patton, A. J. (2015). Modelling Dependence in High Dimensions with Factor Copulas. *Finance and Economics*  
1703 *Discussion Series*, 2015.0(51), 1–41. <https://doi.org/10.17016/feds.2015.051>
- 1704 Orth, R., O, S., Zscheischler, J., Mahecha, M. D., & Reichstein, M. (2022). Contrasting biophysical and societal impacts of  
1705 hydro-meteorological extremes. *Environmental Research Letters*, 17(1), 014044. <https://doi.org/10.1088/1748-9326/ac4139>

Formatted: English (US)

Formatted: Font: Italic

1707 Pan, S. J., & Yang, Q. (2010). A Survey on Transfer Learning. *IEEE Transactions on Knowledge and Data Engineering*,  
1708 22(10), 1345–1359. <https://doi.org/10.1109/TKDE.2009.191>

1709 Pan, Y., Zeng, X., Xu, H., Sun, Y., Wang, D., & Wu, J. (2021). Evaluation of Gaussian process regression kernel functions  
1710 for improving groundwater prediction. *Journal of Hydrology*, 603, 126960.  
1711 <https://doi.org/10.1016/j.jhydrol.2021.126960>

1712 Park, S. J., & Lee, D. K. (2020). Prediction of coastal flooding risk under climate change impacts in South Korea using machine  
1713 learning algorithms. *Environmental Research Letters*, 15(9). <https://doi.org/10.1088/1748-9326/ABA5B3>

1714 Park, S., Sohn, W., Piao, Y., & Lee, D. (2023). Adaptation strategies for future coastal flooding: Performance evaluation of  
1715 green and grey infrastructure in South Korea. *Journal of Environmental Management*, 334, 117495.  
1716 <https://doi.org/10.1016/j.jenvman.2023.117495>

1717 Patil, K. R., Doi, T., & Behera, S. K. (2023). Predicting extreme floods and droughts in East Africa using a deep learning  
1718 approach. *Npj Climate and Atmospheric Science*, 6(1), 108. <https://doi.org/10.1038/s41612-023-00435-x>

1719 Pescaroli, G., & Alexander, D. (2018). Understanding Compound, Interconnected, Interacting, and Cascading Risks: A  
1720 Holistic Framework. *Risk Analysis*, 38(11), 2245–2257. <https://doi.org/10.1111/risa.13128>

1721 Pham, H. V., Dal Barco, M. K., Cadau, M., Harris, R., Furlan, E., Torresan, S., Rubinetti, S., Zanchettin, D., Rubino, A.,  
1722 Kuznetsov, I., Barbarioli, F., Benetazzo, A., Sclavo, M., & Critto, A. (2023). Multi-model chain for climate change  
1723 scenario analysis to support coastal erosion and water quality risk management for the Metropolitan city of Venice.  
1724 *Science of The Total Environment*, 904, 166310. <https://doi.org/10.1016/j.scitotenv.2023.166310>

1725 Piao, Y., Lee, D., Park, S., Kim, H. G., & Jin, Y. (2022). Multi-hazard mapping of droughts and forest fires using a multi-layer  
1726 hazards approach with machine learning algorithms. *Geomatics, Natural Hazards and Risk*, 13(1), 2649–2673.  
1727 <https://doi.org/10.1080/19475705.2022.2128440>

1728 Pilkington, S., & Mahmoud, H. (2017). Spatial and temporal variations in resilience to tropical cyclones along the United  
1729 States coastline as determined by the multi-hazard hurricane impact level model. *Palgrave Communications*, 3(1), 14.  
1730 <https://doi.org/10.1057/s41599-017-0016-1>

1731 Pourghasemi, H. R., Gayen, A., Panahi, M., Rezaie, F., & Blaschke, T. (2019). Multi-hazard probability assessment and  
1732 mapping in Iran. *Science of The Total Environment*, 692, 556–571. <https://doi.org/10.1016/j.scitotenv.2019.07.203>

1733 Pourghasemi, H. R., Kariminejad, N., Amiri, M., Edalat, M., Zarafshar, M., Blaschke, T., & Cerda, A. (2020). Assessing and  
1734 mapping multi-hazard risk susceptibility using a machine learning technique. *Scientific Reports*, 10(1), 3203.  
1735 <https://doi.org/10.1038/s41598-020-60191-3>

1736 Pouyan, S., Pourghasemi, H. R., Bordbar, M., Rahmanian, S., & Clague, J. J. (2021). A multi-hazard map-based flooding,  
1737 gully erosion, forest fires, and earthquakes in Iran. *Scientific Reports*, 11(1), 14889. <https://doi.org/10.1038/s41598-021-94266-6>

1738

1739 Powers, C. J., Devaraj, A., Ashqeen, K., Dontula, A., Joshi, A., Shenoy, J., & Murthy, D. (2023). Using artificial intelligence  
1740 to identify emergency messages on social media during a natural disaster: A deep learning approach. *International*  
1741 *Journal of Information Management Data Insights*, 3(1), 100164. <https://doi.org/10.1016/j.jjimei.2023.100164>

1742 Qiang, Y., Huang, Q., & Xu, J. (2020). Observing community resilience from space: Using nighttime lights to model economic  
1743 disturbance and recovery pattern in natural disaster. *Sustainable Cities and Society*, 57, 102115.  
1744 <https://doi.org/10.1016/j.scs.2020.102115>

1745 Racah, E., Beckham, C., Maharaj, T., Kahou, S. E., Prabhat, & Pal, C. (2016). *ExtremeWeather: A large-scale climate dataset*  
1746 *for semi-supervised detection, localization, and understanding of extreme weather events*.  
1747 <https://doi.org/https://doi.org/10.48550/arXiv.1612.02095>

1748 Rahman, M., Shufeng, T., Tumon, M. S. H., Hossain, M. A., Kim, H.-J., Islam, M. M., Alam, M., Sadiq, S., Ningsheng, C.,  
1749 Ullah, K., Zafar, M. A., & Raju, M. R. (2024). Multi-hazard could exacerbate in coastal Bangladesh in the context of  
1750 climate change. *Journal of Cleaner Production*, 457, 142289. <https://doi.org/10.1016/j.jclepro.2024.142289>

1751 Ray, K., Giri, R. K., Ray, S. S., Dimri, A. P., & Rajeevan, M. (2021). An assessment of long-term changes in mortalities due  
1752 to extreme weather events in India: A study of 50 years' data, 1970–2019. *Weather and Climate Extremes*, 32, 100315.  
1753 <https://doi.org/10.1016/j.wace.2021.100315>

1754 Read, J. S., Jia, X., Willard, J., Appling, A. P., Zwart, J. A., Oliver, S. K., Karpatne, A., Hansen, G. J. A., Hanson, P. C.,  
1755 Watkins, W., Steinbach, M., & Kumar, V. (2019). Process-Guided Deep Learning Predictions of Lake Water  
1756 Temperature. *Water Resources Research*, 55(11), 9173–9190. <https://doi.org/10.1029/2019WR024922>

1757 Reichstein, M., Camps-Valls, G., Stevens, B., Jung, M., Denzler, J., Carvalhais, N., & Prabhat. (2019). Deep learning and  
1758 process understanding for data-driven Earth system science. *Nature*, 566(7743), 195–204.  
1759 <https://doi.org/10.1038/s41586-019-0912-1>

1760 Reichstein, M., Benson, V., Blunk, J., Camps-Valls, G., Creutzig, F., Fearnley, C. J., Han, B., Kornhuber, K., Rahaman, N.,  
1761 Schölkopf, B., Tárraga, J. M., Vinuesa, R., Dall, K., Denzler, J., Frank, D., Martini, G., Nganga, N., Maddix, D. C., &  
1762 Weldemariam, K. (2025). Early warning of complex climate risk with integrated artificial intelligence. *Nature*  
1763 *Communications*, 16(1), 2564. <https://doi.org/10.1038/s41467-025-57640-w>

1764 Ribeiro, A. F. S., Russo, A., Gouveia, C. M., Páscoa, P., & Zscheischler, J. (2020). Risk of crop failure due to compound dry  
1765 and hot extremes estimated with nested copulas. *Biogeosciences*, 17(19), 4815–4830. [https://doi.org/10.5194/bg-17-](https://doi.org/10.5194/bg-17-4815-2020)  
1766 4815-2020

1767 Ridder, N. N., Pitman, A. J., & Ukkola, A. M. (2021). Do CMIP6 Climate Models Simulate Global or Regional Compound  
1768 Events Skillfully? *Geophysical Research Letters*, 48(2). <https://doi.org/10.1029/2020GL091152>

1769 Ridder, N. N., Ukkola, A. M., Pitman, A. J., & Perkins-Kirkpatrick, S. E. (2022). Increased occurrence of high impact  
1770 compound events under climate change. *Npj Climate and Atmospheric Science*, 5(1), 3. [https://doi.org/10.1038/s41612-](https://doi.org/10.1038/s41612-021-00224-4)  
1771 021-00224-4

Formatted: English (UK)

Formatted: English (UK)

Formatted: English (UK)

Field Code Changed

Formatted: English (US)

1772 Rolnick, D., Donti, P. L., Kaack, L. H., Kochanski, K., Lacoste, A., Sankaran, K., Ross, A. S., Milojevic-Dupont, N., Jaques,  
1773 N., Waldman-Brown, A., Luccioni, A., Maharaj, T., Sherwin, E. D., Mukkavilli, S. K., Kording, K. P., Gomes, C., Ng,  
1774 A. Y., Hassabis, D., Platt, J. C., ... Bengio, Y. (2019). *Tackling Climate Change with Machine Learning*.  
1775 <https://doi.org/https://doi.org/10.48550/arXiv.1906.05433>

1776 Rusk, J., Maharjan, A., Tiwari, P., Chen, T.-H. K., Shneiderman, S., Turin, M., & Seto, K. C. (2022). Multi-hazard  
1777 susceptibility and exposure assessment of the Hindu Kush Himalaya. *Science of The Total Environment*, *804*, 150039.  
1778 <https://doi.org/10.1016/j.scitotenv.2021.150039>

1779 Sadegh, M., Ragno, E., & AghaKouchak, A. (2017). Multivariate Copula Analysis Toolbox (MvCAT): Describing dependence  
1780 and underlying uncertainty using a Bayesian framework. *Water Resources Research*, *53*(6), 5166–5183.  
1781 <https://doi.org/10.1002/2016WR020242>

1782 Sagan, V., Peterson, K. T., Maimaitijiang, M., Sidike, P., Sloan, J., Greeling, B. A., Maalouf, S., & Adams, C. (2020).  
1783 Monitoring inland water quality using remote sensing: potential and limitations of spectral indices, bio-optical  
1784 simulations, machine learning, and cloud computing. *Earth-Science Reviews*, *205*, 103187.  
1785 <https://doi.org/10.1016/j.earscirev.2020.103187>

1786 Saha, A., Pal, S. C., Santosh, M., Janizadeh, S., Chowdhuri, I., Norouzi, A., Roy, P., & Chakraborty, R. (2021). Modelling  
1787 multi-hazard threats to cultural heritage sites and environmental sustainability: The present and future scenarios. *Journal*  
1788 *of Cleaner Production*, *320*, 128713. <https://doi.org/10.1016/j.jclepro.2021.128713>

1789 Saha, A., & Ravela, S. (2022). *Downscaling Extreme Rainfall Using Physical-Statistical Generative Adversarial Learning*.  
1790 <https://doi.org/https://doi.org/10.48550/arXiv.2212.01446>

1791 Šakić Trogrlić, R., Reiter, K., Ciurean, R. L., Gottardo, S., Torresan, S., Daloz, A. S., Ma, L., Padrón Fumero, N., Tatman, S.,  
1792 Hochrainer-Stigler, S., de Ruiter, M. C., Schlumberger, J., Harris, R., García-Gonzalez, S., García-Vaquero, M., Arévalo,  
1793 T. L. F., Hernandez-Martin, R., Mendoza-Jimenez, J., Ferrario, D. M., ... Ward, P. J. (2024). Challenges in assessing  
1794 and managing multi-hazard risks: A European stakeholders perspective. *Environmental Science & Policy*, *157*, 103774.  
1795 <https://doi.org/10.1016/j.envsci.2024.103774>

1796 Salcedo-Sanz, S., Pérez-Aracil, J., Ascenso, G., Del Ser, J., Casillas-Pérez, D., Kadow, C., Fister, D., Barriopedro, D., García-  
1797 Herrera, R., Restelli, M., Giuliani, M., & Castelletti, A. (2022). *Analysis, Characterization, Prediction and Attribution*  
1798 *of Extreme Atmospheric Events with Machine Learning: a Review*.

1799 Sammonds, P., Alam, A., Day, S., Stavrianaki, K., & Kelman, I. (2023). Hurricane risk assessment in a multi-hazard context  
1800 for Dominica in the Caribbean. *Scientific Reports*, *13*(1), 20565. <https://doi.org/10.1038/s41598-023-47527-5>

1801 Sarkis-Onofre, R., Catalá-López, F., Aromataris, E., & Lockwood, C. (2021). How to properly use the PRISMA Statement.  
1802 *Systematic Reviews*, *10*(1), 117. <https://doi.org/10.1186/s13643-021-01671-z>

1803 Schiefer, F., Kattenborn, T., Frick, A., Frey, J., Schall, P., Koch, B., & Schmidtlein, S. (2020). Mapping forest tree species in  
1804 high resolution UAV-based RGB-imagery by means of convolutional neural networks. *ISPRS Journal of*  
1805 *Photogrammetry and Remote Sensing*, *170*, 205–215. <https://doi.org/10.1016/j.isprsjprs.2020.10.015>

1806 Schmidt, H., Radinger, J., Teschlade, D., & Stoll, S. (2020). The role of spatial units in modelling freshwater fish distributions:  
1807 Comparing a subcatchment and river network approach using MaxEnt. *Ecological Modelling*, *418*, 108937.  
1808 <https://doi.org/10.1016/j.ecolmodel.2020.108937>

1809 Schneider, T., Behera, S., Boccaletti, G., Deser, C., Emanuel, K., Ferrari, R., Leung, L. R., Lin, N., Müller, T., Navarra, A.,  
1810 Ndiaye, O., Stuart, A., Tribbia, J., & Yamagata, T. (2023). Harnessing AI and computing to advance climate modelling  
1811 and prediction. *Nature Climate Change*, *13*(9), 887–889. <https://doi.org/10.1038/s41558-023-01769-3>

1812 Sfetsos, A., Politi, N., & Vlachogiannis, D. (2023). Multi-Hazard Extreme Scenario Quantification Using Intensity, Duration,  
1813 and Return Period Characteristics. *Climate*, *11*(12), 242. <https://doi.org/10.3390/cli11120242>

1814 Shah, K., Patel, H., Sanghvi, D., & Shah, M. (2020). A Comparative Analysis of Logistic Regression, Random Forest and  
1815 KNN Models for the Text Classification. *Augmented Human Research*, *5*(1), 12. <https://doi.org/10.1007/s41133-020-00032-0>

1816

1817 Shepherd, T. G., Boyd, E., Calel, R. A., Chapman, S. C., Dessai, S., Dima-West, I. M., Fowler, H. J., James, R., Maraun, D.,  
1818 Martius, O., Senior, C. A., Sobel, A. H., Stainforth, D. A., Tett, S. F. B., Trenberth, K. E., van den Hurk, B. J. J. M.,  
1819 Watkins, N. W., Wilby, R. L., & Zenghelis, D. A. (2018). Storylines: an alternative approach to representing uncertainty  
1820 in physical aspects of climate change. *Climatic Change*, *151*(3–4), 555–571. [https://doi.org/10.1007/s10584-018-2317-](https://doi.org/10.1007/s10584-018-2317-9)  
1821 [9](https://doi.org/10.1007/s10584-018-2317-9)

1822 Sillmann, J., Shepherd, T. G., van den Hurk, B., Hazeleger, W., Martius, O., Slingo, J., & Zscheischler, J. (2021). Event-Based  
1823 Storylines to Address Climate Risk. *Earth's Future*, *9*(2). <https://doi.org/10.1029/2020EF001783>

1824 Simpson, E. S., Wadsworth, J. L., & Tawn, J. A. (2020). *A geometric investigation into the tail dependence of vine copulas*.  
1825 <https://doi.org/10.1016/j.jmva.2021.104736>

1826 Singh, T. P., Nandimath, P., Kumbhar, V., Das, S., & Barne, P. (2021). Drought risk assessment and prediction using artificial  
1827 intelligence over the southern Maharashtra state of India. *Modeling Earth Systems and Environment*, *7*(3), 2005–2013.  
1828 <https://doi.org/10.1007/s40808-020-00947-y>

1829 Sippel, S., Otto, F. E. L., Forkel, M., Allen, M. R., Guillod, B. P., Heimann, M., Reichstein, M., Seneviratne, S. I., Thonicke,  
1830 K., & Mahecha, M. D. (2016). A novel bias correction methodology for climate impact simulations. *Earth System*  
1831 *Dynamics*, *7*(1), 71–88. <https://doi.org/10.5194/esd-7-71-2016>

1832 Sit, M., Demiray, B. Z., Xiang, Z., Ewing, G. J., Sermet, Y., & Demir, I. (2020). A comprehensive review of deep learning  
1833 applications in hydrology and water resources. *Water Science and Technology*, *82*(12), 2635–2670.  
1834 <https://doi.org/10.2166/wst.2020.369>

1835 Sodoge, J., Kuhlicke, C., & de Brito, M. M. (2023). Automated spatio-temporal detection of drought impacts from newspaper  
1836 articles using natural language processing and machine learning. *Weather and Climate Extremes*, *41*, 100574.  
1837 <https://doi.org/10.1016/j.wace.2023.100574>

Formatted: English (US)

Formatted: English (US)

838 Sperotto, A., Molina, J.L., Torresan, S., Critto, A., Marcomini, A. (2017). Reviewing Bayesian Networks potentials for climate  
839 change impacts assessment and management: A multi-risk perspective. *Journal of Environmental Management*, 202 (1),  
840 320-331. <https://doi.org/10.1016/j.jenvman.2017.07.044>.

1841 Sublime, J., & Kalinicheva, E. (2019). Automatic Post-Disaster Damage Mapping Using Deep-Learning Techniques for  
1842 Change Detection: Case Study of the Tohoku Tsunami. *Remote Sensing*, 11(9), 1123.  
1843 <https://doi.org/10.3390/rs11091123>

1844 Sun, A. Y., Jiang, P., Mudunuru, M. K., & Chen, X. (2021). Explore Spatio-Temporal Learning of Large Sample Hydrology  
1845 Using Graph Neural Networks. *Water Resources Research*, 57(12). <https://doi.org/10.1029/2021WR030394>

1846 Sun, X., Sun, Q., Zhou, X., Li, X., Yang, M., Yu, A., & Geng, F. (2014). Heat wave impact on mortality in Pudong New Area,  
1847 China in 2013. *Science of The Total Environment*, 493, 789–794. <https://doi.org/10.1016/j.scitotenv.2014.06.042>

1848 Sutanto, S. J., Vitolo, C., Di Napoli, C., D'Andrea, M., & Van Lanen, H. A. J. (2020). Heatwaves, droughts, and fires:  
1849 Exploring compound and cascading dry hazards at the pan-European scale. *Environment International*, 134, 105276.  
1850 <https://doi.org/10.1016/j.envint.2019.105276>

1851 Sweet, L., Müller, C., Anand, M., & Zscheischler, J. (2023). Cross-Validation Strategy Impacts the Performance and  
1852 Interpretation of Machine Learning Models. *Artificial Intelligence for the Earth Systems*, 2(4).  
853 <https://doi.org/10.1175/AIES-D-23-0026.1>

854 Stolte, T.R., Koks, E.E., de Moel, H., Reimann, L., van Vliet, J., de Ruijter, M.C., & Ward, P.J. (2024). VulneraCity – drivers  
855 and dynamics of urban vulnerability based on a global systematic literature review. *International Journal of Disaster*  
856 *Risk Reduction*, 108, 104535. <https://doi.org/10.1016/j.ijdr.2024.104535>

1857 Tabari, H., & Willems, P. (2023). Global risk assessment of compound hot-dry events in the context of future climate change  
1858 and socioeconomic factors. *Npj Climate and Atmospheric Science*, 6(1), 74. [https://doi.org/10.1038/s41612-023-00401-](https://doi.org/10.1038/s41612-023-00401-7)  
1859 [7](https://doi.org/10.1038/s41612-023-00401-7)

1860 Tárraga, J. M., Sevillano-Marco, E., Muñoz-Marí, J., Piles, M., Sitokonstantinou, V., Ronco, M., Miranda, M. T., Cerdà, J., &  
1861 Camps-Valls, G. (2024). Causal discovery reveals complex patterns of drought-induced displacement. *IScience*, 27(9),  
1862 110628. <https://doi.org/10.1016/j.isci.2024.110628>

1863 Tazi, K., Lin, J. A., Viljoen, R., Gardner, A., John, S., Ge, H., & Turner, R. E. (2023). Beyond intuition, a Framework for  
1864 Applying GPs to Real-World Data. *ICML Workshop on Structured Probabilistic Inference and Generative Modelling*.

1865 Tazi, K., Orr, A., Hernandez-González, J., Hosking, S., & Turner, R. E. (2024). Downscaling precipitation over High-mountain  
1866 Asia using multi-fidelity Gaussian processes: improved estimates from ERA5. *Hydrology and Earth System Sciences*,  
1867 28(22), 4903–4925. <https://doi.org/10.5194/hess-28-4903-2024>

1868 Teichert, N., Borja, A., Chust, G., Uriarte, A., & Lepage, M. (2016). Restoring fish ecological quality in estuaries: Implication  
1869 of interactive and cumulative effects among anthropogenic stressors. *Science of The Total Environment*, 542(Part A),  
1870 383–393. <https://doi.org/10.1016/j.scitotenv.2015.10.068>

Formatted: Font: Italic

Formatted: English (US)

1871 Terzi, S., Torresan, S., Schneiderbauer, S., Critto, A., Zebisch, M., & Marcomini, A. (2019). Multi-risk assessment in mountain  
1872 regions: A review of modelling approaches for climate change adaptation. *Journal of Environmental Management*,  
1873 232(February), 759–771. <https://doi.org/10.1016/j.jenvman.2018.11.100>

1874 Tiggeloven, T., Couason, A., van Straaten, C., Muis, S., & Ward, P. J. (2021). Exploring deep learning capabilities for surge  
1875 predictions in coastal areas. *Scientific Reports*, 11(1), 17224. <https://doi.org/10.1038/s41598-021-96674-0>

1876 Tiggeloven, T., Pfeiffer, S., Matanó, A., van den Honberg, Thalheimer, L., Reichstein, M., Torresan, S. (2025). The Role of  
1877 Artificial Intelligence for Early Warning Systems: Status, Applicability, Guardrails, and Ways Forward. *iScience* 28  
1878 (11). [https://www.cell.com/iscience/fulltext/S2589-0042\(25\)01950-9](https://www.cell.com/iscience/fulltext/S2589-0042(25)01950-9).

1879 Tilloy, A., Malamud, B. D., & Joly-Laugel, A. (2022). A methodology for the spatiotemporal identification of compound  
1880 hazards: wind and precipitation extremes in Great Britain (1979–2019). *Earth System Dynamics*, 13(2), 993–1020.  
1881 <https://doi.org/10.5194/esd-13-993-2022>

1882 Tilloy, A., Malamud, B. D., Winter, H., & Joly-Laugel, A. (2019). A review of quantification methodologies for multi-hazard  
1883 interrelationships. *Earth-Science Reviews*, 196, 102881. <https://doi.org/10.1016/j.earscirev.2019.102881>

1884 Tootoonchi, F., Sadegh, M., Haerter, J. O., Rätty, O., Grabs, T., & Teutschbein, C. (2022). Copulas for hydroclimatic analysis:  
1885 A practice-oriented overview. *WIREs Water*, 9(2). <https://doi.org/10.1002/wat2.1579>

1886 Tran, D. Q., Park, M., Jung, D., & Park, S. (2020). Damage-Map Estimation Using UAV Images and Deep Learning  
1887 Algorithms for Disaster Management System. *Remote Sensing*, 12(24), 4169. <https://doi.org/10.3390/rs12244169>

1888 Ullah, K., Wang, Y., Fang, Z., Wang, L., & Rahman, M. (2022). Multi-hazard susceptibility mapping based on Convolutional  
1889 Neural Networks. *Geoscience Frontiers*, 13(5), 101425. <https://doi.org/10.1016/j.gsf.2022.101425>

1890 Vaswani, A., Shazeer, N., Parmar, N., Uszkoreit, J., Jones, L., Gomez, A. N., Kaiser, L., & Polosukhin, I. (2017). *Attention Is*  
1891 *All You Need*. <https://doi.org/10.1016/j.arXiv.1706.03762v7>

1892 Veličković, P., Cucurull, G., Casanova, A., Romero, A., Liò, P., & Bengio, Y. (2017). *Graph Attention Networks*.  
1893 <https://doi.org/10.1016/j.arXiv.1706.03762v7>

1894 Veras, H. F. P., Ferreira, M. P., da Cunha Neto, E. M., Figueiredo, E. O., Corte, A. P. D., & Sanquetta, C. R. (2022). Fusing  
1895 multi-season UAS images with convolutional neural networks to map tree species in Amazonian forests. *Ecological*  
1896 *Informatics*, 71, 101815. <https://doi.org/10.1016/j.ecoinf.2022.101815>

1897 Wang, J. (2023). An Intuitive Tutorial to Gaussian Process Regression. *Computing in Science & Engineering*, 25(4), 4–11.  
1898 <https://doi.org/10.1109/MCSE.2023.3342149>

1899 Wang, J., Kuffer, M., Roy, D., & Pfeffer, K. (2019). Deprivation pockets through the lens of convolutional neural networks.  
1900 *Remote Sensing of Environment*, 234, 111448. <https://doi.org/10.1016/j.rse.2019.111448>

1901 Wang, J., & Yan, Z. (2021). Rapid rises in the magnitude and risk of extreme regional heat wave events in China. *Weather*  
1902 *and Climate Extremes*, 34, 100379. <https://doi.org/10.1016/j.wace.2021.100379>

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Formatted: Font: Italic

Formatted: English (US)

- 1903 Wang, Q., Zhang, X., Chen, G., Dai, F., Gong, Y., & Zhu, K. (2018). Change detection based on Faster R-CNN for high-  
1904 resolution remote sensing images. *Remote Sensing Letters*, 9(10), 923–932.  
1905 <https://doi.org/10.1080/2150704X.2018.1492172>
- 1906 Wang, R., Kim, J.-H., & Li, M.-H. (2021). Predicting stream water quality under different urban development pattern scenarios  
1907 with an interpretable machine learning approach. *Science of The Total Environment*, 761, 144057.  
1908 <https://doi.org/10.1016/j.scitotenv.2020.144057>
- 1909 Wang, X., Ma, Z., & Dong, J. (2021). Quantitative Impact Analysis of Climate Change on Residents' Health Conditions with  
1910 Improving Eco-Efficiency in China: A Machine Learning Perspective. *International Journal of Environmental Research  
1911 and Public Health*, 18(23), 12842. <https://doi.org/10.3390/ijerph182312842>
- 1912 Wang, Y., Song, Q., Du, Y., Wang, J., Zhou, J., Du, Z., & Li, T. (2019). A random forest model to predict heatstroke occurrence  
1913 for heatwave in China. *Science of The Total Environment*, 650, 3048–3053.  
1914 <https://doi.org/10.1016/j.scitotenv.2018.09.369>
- 1915 Ward, P. J., Daniell, J., Duncan, M., Dunne, A., Hananel, C., Hochrainer-Stigler, S., Tijssen, A., Torresan, S., Ciurean, R.,  
1916 Gill, J. C., Sillmann, J., Couasnon, A., Koks, E., Padrón-Fumero, N., Tatman, S., Tronstad Lund, M., Adesiyun, A.,  
1917 Aerts, J. C. J. H., Alabaster, A., ... de Ruiter, M. C. (2022). Invited perspectives: A research agenda towards disaster  
1918 risk management pathways in multi-(hazard-)risk assessment. *Natural Hazards and Earth System Sciences*, 22(4), 1487–  
1919 1497. <https://doi.org/10.5194/nhess-22-1487-2022>
- 1920 Willard, J., Jia, X., Xu, S., Steinbach, M., & Kumar, V. (2022). *Integrating Scientific Knowledge with Machine Learning for  
1921 Engineering and Environmental Systems*. <http://arxiv.org/abs/2003.04919>
- 1922 Wu, H., Su, X., & Singh, V. P. (2023). Increasing Risks of Future Compound Climate Extremes With Warming Over Global  
1923 Land Masses. *Earth's Future*, 11(9). <https://doi.org/10.1029/2022EF003466>
- 1924 Wu, H., Su, X., Singh, V. P., & Niu, J. (2024). Predicting compound agricultural drought and hot events using a Cascade  
1925 Modeling framework combining Bayesian Model Averaging ensemble with Vine Copula (CaMBMAViC). *Journal of  
1926 Hydrology*, 642, 131901. <https://doi.org/10.1016/j.jhydrol.2024.131901>
- 1927 Wu, Q., & Lin, H. (2019). A novel optimal-hybrid model for daily air quality index prediction considering air pollutant factors.  
1928 *Science of The Total Environment*, 683, 808–821. <https://doi.org/10.1016/j.scitotenv.2019.05.288>
- 1929 Wubalem, A. (2022). Landslide Inventory, Susceptibility, Hazard and Risk Mapping. In *Landslides*. IntechOpen.  
1930 <https://doi.org/10.5772/intechopen.100504>
- 1931 Xu, L., Chen, N., Yang, C., Yu, H., & Chen, Z. (2022). Quantifying the uncertainty of precipitation forecasting using  
1932 probabilistic deep learning. *Hydrology and Earth System Sciences*, 26(11), 2923–2938. <https://doi.org/10.5194/hess-26-2923-2022>
- 1934 Ya, R., Wu, J., Tang, R., & Zhou, Q. (2023). Increased flood susceptibility in the Tibetan Plateau with climate and land use  
1935 changes. *Ecological Indicators*, 156, 111086. <https://doi.org/10.1016/j.ecolind.2023.111086>

- 1936 Yeğın, M. N., & Amasyalı, M. F. (2024). *Theoretical research on generative diffusion models: an overview*.  
 1937 <https://doi.org/10.1016/j.neucom.2024.128373>
- 1938 Yousefi, S., Pourghasemi, H. R., Emami, S. N., Pouyan, S., Eskandari, S., & Tiefenbacher, J. P. (2020). A machine learning  
 1939 framework for multi-hazards modeling and mapping in a mountainous area. *Scientific Reports*, *10*(1), 1–14.  
 1940 <https://doi.org/10.1038/s41598-020-69233-2>
- 1941 Yu, H., Lu, N., Fu, B., Zhang, L., Wang, M., & Tian, H. (2022). Hotspots, co-occurrence, and shifts of compound and cascading  
 1942 extreme climate events in Eurasian drylands. *Environment International*, *169*, 107509.  
 1943 <https://doi.org/10.1016/j.envint.2022.107509>
- 1944 Yu, S., Hu, Z., Subramaniam, A., Hannah, W., Peng, L., Lin, J., Bhouri, M. A., Gupta, R., Lütjens, B., Will, J. C., Behrens,  
 1945 G., Busecke, J. J. M., Loose, N., Stern, C. I., Beucler, T., Harrop, B., Heuer, H., Hillman, B. R., Jenney, A., ... Pritchard,  
 1946 M. (2024). *ClimSim-Online: A Large Multi-scale Dataset and Framework for Hybrid ML-physics Climate Emulation*.  
 1947 <http://arxiv.org/abs/2306.08754>
- 1948 Yuh, Y. G., Tracz, W., Matthews, H. D., & Turner, S. E. (2023). Application of machine learning approaches for land cover  
 1949 monitoring in northern Cameroon. *Ecological Informatics*, *74*, 101955. <https://doi.org/10.1016/j.ecoinf.2022.101955>
- 1950 Zanetti, M., Allegri, E., Sperotto, A., Torresan, S., & Critto, A. (2022). Spatio-temporal cross-validation to predict pluvial  
 1951 flood events in the Metropolitan City of Venice. *Journal of Hydrology*, *612*, 128150.  
 1952 <https://doi.org/10.1016/j.jhydrol.2022.128150>
- 1953 Zanini, E., Eastoe, E., Jones, M. J., Randell, D., & Jonathan, P. (2020). Flexible covariate representations for extremes.  
 1954 *Environmetrics*, *31*(5). <https://doi.org/10.1002/env.2624>
- 1955 Zennaro, F., Furlan, E., Simeoni, C., Torresan, S., Aslan, S., Critto, A., & Marcomini, A. (2021). Exploring machine learning  
 1956 potential for climate change risk assessment. *Earth-Science Reviews*, *220*, 103752.  
 1957 <https://doi.org/10.1016/j.earscirev.2021.103752>
- 1958 Zerrouki, N., Harrou, F., Sun, Y., & Hocini, L. (2019). A Machine Learning-Based Approach for Land Cover Change  
 1959 Detection Using Remote Sensing and Radiometric Measurements. *IEEE Sensors Journal*, *19*(14), 5843–5850.  
 1960 <https://doi.org/10.1109/JSEN.2019.2904137>
- 1961 Zhao, G., Pang, B., Xu, Z., Peng, D., & Zuo, D. (2020). Urban flood susceptibility assessment based on convolutional neural  
 1962 networks. *Journal of Hydrology*, *590*, 125235. <https://doi.org/10.1016/j.jhydrol.2020.125235>
- 1963 Zhu, X., Yang, Y., & Tang, J. (2023). Compound wind and precipitation extremes at a global scale based on CMIP6 models:  
 1964 Evaluation, projection and uncertainty. *International Journal of Climatology*, *43*(16), 7588–7605.  
 1965 <https://doi.org/10.1002/joc.8281>
- 1966 Zhuo, L., Han, D. (2020). Agent-based modelling and flood risk management: A compendious literature review. *Journal of*  
 1967 *Hydrology*, *591*, 125600. <https://doi.org/10.1016/j.jhydrol.2020.125600>
- 1968 Zschau. (2017). *Where are we with multihazards, multirisks assessment capacities?*, in: *Science for disaster risk management*  
 1969 *2017: knowing better and losing less*, edited by: Poljansek, K., Marin Ferrer, M., De Groeve, T., and Clark, I., European

Formatted: Font: Italic

1970        *Union, Brussels, Belgium.* [https://drmkc.jrc.ec.europa.eu/knowledge/science-for-drm/science-for-disaster-risk-](https://drmkc.jrc.ec.europa.eu/knowledge/science-for-drm/science-for-disaster-risk-management-2017)  
1971        [management-2017](https://drmkc.jrc.ec.europa.eu/knowledge/science-for-drm/science-for-disaster-risk-management-2017)  
1972        Zscheischler, J., Orth, R., & Seneviratne, S. I. (2017). Bivariate return periods of temperature and precipitation explain a large  
1973        fraction of European crop yields. *Biogeosciences*, *14*(13), 3309–3320. <https://doi.org/10.5194/bg-14-3309-2017>  
1974        Zscheischler, J., Westra, S., van den Hurk, B. J. J. M., Seneviratne, S. I., Ward, P. J., Pitman, A., AghaKouchak, A., Bresch,  
1975        D. N., Leonard, M., Wahl, T., & Zhang, X. (2018). Future climate risk from compound events. *Nature Climate Change*,  
1976        *8*(6), 469–477. <https://doi.org/10.1038/s41558-018-0156-3>  
1977

**Page 38: [1] Deleted** **Davide Ferrario** **3/11/26 2:12:00 PM**

x.....  
**Page 46: [2] Deleted** **Davide Ferrario** **3/9/26 4:06:00 PM**

▲.....  
**Page 60: [3] Deleted** **Davide Ferrario** **3/12/26 11:58:00 AM**

x.....