

Soil signals of key mechanisms driving greater protection of organic carbon under aspen compared to spruce forests in a North American montane ecosystem

Authors: Lena Wang¹, Sharon A. Billings², Li Li³., Daniel R. Hirmas⁴, Keira Johnson¹, Devon Kerins³, Julio Pachon⁵, Curtis Beutler⁶, Karla M. Jarecke¹, Vaishnavi Varikuti⁴, Micah Unruh², Hoori Ajami⁷, Holly Barnard⁸, Alejandro N. Flores⁹, Kenneth Williams⁶, Pamela L. Sullivan¹

¹ College of Earth Ocean and Atmospheric Science, Oregon State University. Address: 101 SW 26th St, Corvallis, OR, 97331, USA

²Department of Ecology and Evolutionary Biology and Kansas Biological Survey & Center for Ecological Research, University of Kansas. Address: 2101 Constant Ave., Lawrence, KS, 66047, USA

³Department of Civil and Environmental Engineering, Pennsylvania State University, Address: Sackett, #212, University Park, PA 16802, USA.

⁴Department of Plant and Soil Science, Texas Tech University. Address: 2500 Broadway Lubbock, Texas 7940 USA.

⁵Sydney Institute of Agriculture, School of Life and Environmental Sciences, University of Sydney, New South Wales, Australia. Address: The University of Sydney, NSW 2006, Australia.

⁶Lawrence Berkeley National Laboratory, Berkeley, CA USA Rocky Mountain Biological Laboratory, Gothic, CO USA. Address: 1 Cyclotron Road, Berkeley, CA 94720, USA.

⁷Department of Environmental Sciences, University of California Riverside. Address: 2460B Geology Building, Riverside, CA 92521USA.

⁸Department of Geography, Institute of Arctic and Alpine Research, University of Colorado – Boulder. Address: Guggenheim 110, 260 UCB Boulder, Colorado 80309-0260, USA.

⁹Department of Geosciences, Boise State University, Boise Address: 1295 University Drive, Boise, ID 83706 USA.

Correspondence to: Pamela L. Sullivan(Sullipam@oregonstate.edu)

Highlights:

1. Soil organic carbon concentrations are consistently greater under aspen compared to spruce
2. Microbial Efficiency - Matrix Stabilization model helps explain SOC differences
3. Smaller aggregate sizes under aspen further help explain SOC concentrations
4. A lower probability of SOC destabilization likely persists under aspen stands

Abstract

Soil organic carbon (SOC) is often retained more effectively in aspen-dominated forests compared to coniferous forests in North America, yet the reasons why are unclear. A potential driver could be differences in SOC protection mechanisms. Over decades to centuries, chemical (e.g., mineral association) and physical (e.g., aggregation) processes can work to preserve SOC stocks, which can vary across cover types. To investigate this hypothesis, we evaluate controls on SOC concentrations in the Coal Creek watershed (CO, USA), a montane ecosystem dominated by quaking aspen and Engelmann spruce and underlain by granite and sandstone. We examined a combination of biological, chemical, physical, and environmental conditions to evaluate potential abiotic and biotic mechanisms of SOC preservation at multiple depths. As expected, we observed greater SOC concentrations under aspen compared to spruce. Growing season soil moisture, temperature, and CO₂ and O₂ varied with slope position and aspect, and thus forest cover type. Dissolved organic carbon (DOC) was lower under aspen compared to spruce. Exo-enzyme data indicate that aspen soil microbes exhibited greater effort to seek organically-bound resources; consistent with this, soil organic N exhibited higher $\delta^{15}\text{N}$ values, hinting at a greater degree of organic matter processing. Finally, aspen soils exhibited greater root abundance, and aspen mineral soils revealed smaller mean aggregate diameters compared to conifer sites. Our data suggest enhanced biotic activities in aspen-dominated forest soils that promote both chemical and physical protection of SOC in aspen- relative to spruce-dominated forests, and associated limitations on potential DOC export.

Keywords (1-7 words): Critical Zone, Ecohydrology, Montane Ecosystems, Soil Organic Carbon, Climate Change

1 Introduction

The distribution and composition of temperate montane forests are changing (Alexander et al., 1987; Anderegg et al., 2013), driven by increasing air temperature, earlier snowmelt, earlier onset and extent of the growing season (Godsey et al., 2014; Mote et al., 2018; Rhoades et al., 2018), and increasing frequency and intensity of disturbance (e.g., drought, fire, logging, and insect infestation; Canelles et al., 2021). For example, aspen stands have lost substantial live density and basal area to Englemann spruce, sub-alpine fir, and Douglas Fir since 1964 with an

increasing rate of decline since 1994 (Alexander, 1987; Coop et al., 2014). Changes in montane forest cover can directly impact soil organic carbon (SOC) stability. Given that SOC [influences](#) the availability of nutrients, soil stability, ecosystem water fluxes, and biosphere-atmosphere exchange of greenhouse gases (Jackson et al., 2017), and that global SOC reservoirs represent far more C than plant biomass and the atmosphere (Scharlemann et al., 2014), unraveling drivers of SOC stability remains an important research goal (Billings et al., 2021). Between paired aspen and conifer stands at numerous sites throughout North America, SOC pools differ substantially (review in Langanieri et al., 2017). Studies consistently show that C under conifers is more readily destabilized than under aspen (Woldeselassie et al., 2012; Langanieri et al., 2013; [Boča et al., 2020](#); [Román Dobarco et al., 2021](#)). Further, SOC pools in aspen-dominated environments tend to be composed of larger stocks of mineral-associated organic C (MAOC), which is a relatively stable SOC fraction, than those under conifers ([66% compared to 48% MAOC to SOC, respectively](#); [Román Dobarco and Van Miegroet, 2014](#); [Román Dobarco et al., 2021](#)). Yet, the mechanisms driving such differences in SOC stability under aspen and conifer remain elusive.

Examining soil physical attributes and how they can differ with plant cover type may help us understand differences in MAOC fate in aspen vs. conifer forests. For example, soil aggregation is a key process promoting SOC protection in many soil types (Blanco-Canqui and Lal, 2004). Soil aggregation refers to the clustering or binding of soil particles into larger units. This process is promoted by interactions among colloidal material and binding compounds (microaggregates; Six et al., 2004; Blanco-Canqui and Lal, 2005; Weil and Brady, 2017; Araya and Ghezzehei, 2019) and among particulate organic C (POC; Cotrufo et al., 2019), and clay minerals or clay-sized particles (Six et al., 2000). The collapse and formation of aggregates influence the protection of SOC. For example, the breakdown of macroaggregates into microaggregates often leads to the release of dissolved organic C (DOC) (Cincotta et al., 2019), some of which can undergo microbial uptake and mineralization to CO₂. In contrast, aggregate formation can limit soil microbial access to SOC on aggregates' interiors, helping to shield it from exo-enzymatic attack (Jastrow 1996; Six et al., 2000; Woolf et al., 2019).

Multiple mechanisms may drive differences in soil aggregation across aspen and conifer soils. First, soils beneath conifers are often more acidic (PoPONoe et al., 1992; Buck and St. Clair, 2012) and thus may promote a greater abundance of relatively small aggregates, given that increases in soil solution [H⁺] can weaken soil aggregation processes (Stătescu et al., 2013). Second, differences in rooting abundance among aspen and conifers may drive differences in aggregate formation across these cover types. Aspen tends to produce shallow roots that generally extend to ~ 30 cm deep (Sheppard et al., 2006), while conifers tend to develop both lateral and tap roots, the latter of which can extend relatively deep into the soil and bedrock (Mauer et al., 2012). Spruce tends to exhibit lower fine root biomass compared to aspen

(Mekontchou et al., 2020). Fine roots may promote aggregate formation through enmeshment processes, while coarse roots may promote aggregate collapse because of roots perforating aggregates (Bronick and Lal, 2005). Differences in soil moisture between aspen and conifers driven by differences in aspect, foliar cover, and transpiration rates (Buck and St. Clair, 2012) may also influence aggregate stability, as rapid changes in soil moisture can cause aggregates to burst while a gradual increase in moisture can stabilize aggregates (Amezketta, 1999).

Combined, these concurrent and competing processes may drive differences in soil aggregation between aspen- vs. conifer-dominated soils in ways that are difficult to predict due to complex and non-linear interactions, and require the synthesis of findings across biological and pedological disciplines to understand.

Soil moisture and temperature not only influence physical aggregation processes, and thus the protection and preservation of SOC, but also the degree to which microbes transform SOC into CO₂ or alter the transport of organic C pools to depth. Where soil moisture is higher, greater transport of organic C pools into the subsurface may be feasible, potentially increasing the amount of organic matter sorbed to minerals at greater depths (Mikutta et al., 2019). Conversely, DOC leaching may increase, and subsequent DOC export could reduce SOC concentrations (Roulet and Moore, 2006; Monteith et al., 2007). Soil temperature also may drive differences in SOC transformations across aspen and conifer sites, given that aspect exerts strong control on aspen distribution. Soil temperatures tend to be warmer under the sunnier, aspen-dominated stands compared to conifer stands (Buck and St. Clair, 2012). In a temperature-limited montane system, warmer temperatures under aspen stands may increase microbial metabolic activity and turnover, and thus accelerate microbial necromass formation, a process linked to greater stocks of relatively persistent SOC (Liang et al., 2019), perhaps due to necromass-promoting aggregate formation and stabilization (Sae-Tun et al., 2022). Thus, understanding soil hydrologic behaviors as well as solute transport down-profile, traditionally the realms of hydrologists and soil biogeochemists, in addition to biological and pedological processes, is important for understanding patterns of SOC transformations.

Finally, differences in the chemical composition of aspen and conifer biomass and their root exudates may explain differences in MAOC stocks between the two stand types (Boča et al., 2020). For example, aspen litter tends to exhibit lower lignin concentrations than coniferous litterfall (Moore et al., 2006). The Microbial Efficiency - Matrix Stabilization (MEMS) framework (Cutrofo et al., 2013) would suggest this more labile plant material may be easier for soil microbes to assimilate and transform into microbial necromass, which can become more physically or chemically protected through aggregation and chemical bonding (Kleber et al., 2007; von Lützow et al., 2008; Cutrofo et al., 2013) and lead to relatively more persistent stocks of SOC (Liang et al., 2019; Buckridge et al., 2022). Differences in microbial activities between aspen and conifer may further be exacerbated by differences in root exudation between these

species. For example, Norway spruce can exhibit lower exudation rates than silver birch (Sadnes et al., 2005), and deciduous trees appear to experience greater exudation rates than pines (Wang et al., 2021). Though many studies explore the biotic, chemical, physical, and hydrologic processes that can influence SOC transformations and preservation, these processes are rarely examined at the same time. Thus, it remains unclear why conifer-dominated forests consistently harbor smaller amounts of SOC, and why aspen-dominated forests exhibit greater SOC stabilization.

Here, we use a holistic, critical-zone approach (Chorover et al., 2007) to understand SOC dynamics and drivers. We explore a suite of abiotic and biotic factors as they relate to SOC pool sizes across two forest cover types at Coal Creek, a watershed in central Colorado, USA, dominated by Englemann spruce (*Picea engelmanni*) on the north-facing hillslopes, and aspen (*Populus tremuloides*) on the south-facing hillslopes. Coal Creek has experienced relatively high variability in stream water DOC concentrations in recent years (2005-2019; Leonard et al., 2022). The mysterious, almost tripling of stream DOC concentrations in some years (2018-2019) may indicate recent shifts in upslope biogeochemical processes such as greater forest stress associated with climate change (Leonard et al., 2022) and subsequent changes in hydrologic flow paths (Zhi et al., 2020; Kerins et al., 2023) that influence C transport from soil profiles to stream water. We test the hypothesis that oft-observed greater stocks of SOC in soils under aspen stands compared to conifer-dominated soils are linked to greater soil microbial activities in aspen stand soils. We further hypothesize that relatively stable microaggregates are more abundant in aspen-dominated soils than spruce-dominated soils, resulting from higher microbial activities, and thus higher necromass production rates. Differences in rooting strategies and ecohydrology (e.g., evapotranspiration, soil moisture) between aspen and spruce stands will likely exert a secondary control on C stability. We finally hypothesize that the proliferation of fine roots in aspen-dominated soils are linked to smaller water-stable aggregates, and the generally coarser and deeper roots under spruce facilitate the transport of moisture and dissolved C down through the soil profile to a greater extent than under aspens, a process directly linked to DOC export to streams.

To test these hypotheses, we quantified multiple metrics describing basic abiotic conditions, SOC pools, soil microbial activities, soil aggregate-size distributions, and rooting distributions on five hillslopes dominated by either spruce or aspen, underlain by two contrasting lithologies and located at two hillslope positions (i.e., backslope and footslope). We aim to clarify some of the mechanisms governing aspen- and conifer-dominated forest soil microbial activity, soil aggregation, and soil moisture dynamics and their impact on SOC protection and potential DOC transport into surface water, illuminating the possible trajectories of SOC and DOC in rapidly changing, montane forest watersheds.

2 Study Area




Figure 1: A map of the Coal Creek catchment. Colors represent land cover types, where aspen (orange) are dominantly at lower south-facing slopes while conifer (green) are on both north and south facing slopes. Shapes represent lithology type where granite sites (blue triangles) are in the western part of the catchment and sandstone sites (pink circles) are in the eastern part of the catchment. AS is aspen sandstone, and SS is spruce sandstone. ESG and EAG are spruce granite and aspen granite, respectively. They are in Elk Creek, a sub catchment of the Coal Creek catchment. While ESG is on a dominantly south facing slope, it is north facing within the Elk Creek catchment. SG is also a spruce granite site. Note that all sites reside at contrasting hillslope positions: backslope = AS and SG, and footslopes = SS, ESG, and EAG.

Coal Creek (53 km²) is a high-elevation (2715 m), headwater tributary of the Upper Colorado River Basin located in the Colorado Rocky Mountains near the town of Crested Butte (Fig. 1). Coal Creek is a sub-catchment of the larger East River watershed (300 km²) and falls within the research domains of the U.S. Department of Energy funded Watershed Function Science Focus Area and Rocky Mountain Biological Laboratory (RMBL). The watershed is seasonally snow-covered from November through June. The area has a continental, subarctic climate with long, cold winters and short, cool summers. The mean annual temperature is 0.9 °C and the mean annual precipitation is 670 mm (Carroll et al., 2018), with approximately 60% falling as snow between October and May. This area has been warming since the 1980s and the fraction of snow has been decreasing roughly at 1% per year (Zhi et al., 2020). Due to these warming temperatures, the growing season in Crested Butte appears to be extending (Wadgymar et al., 2018).

The geology of Coal Creek is underlain by sandstone, siltstone, shale, and coal units from the Mesa Verde Formation, variegated claystone and shale from the Wasatch Formation, and some intrusive granite diorite, granite, quartz, and monzonite that are Middle Tertiary aged (Gaskill et al., 1991). Soils are predominantly mapped as carbonate free Alfisols, Mollisols, and Inceptisols (Soil Survey Staff, 2023).

Spruce, aspens, and alpine meadows can be found in the Coal Creek watershed. North-facing slopes are dominated by Engelmann Spruce, while aspen and Engelmann spruce can be found on south-facing slopes. We focused on five sites during this study. Three of our sites lie within the main drainage of Coal Creek including two spruce sites (spruce sandstone, SS; spruce granite, SG) and one aspen (aspen sandstone (AS)). The last two sites are located in Elk Creek, a sub-catchment of Coal Creek, which includes one spruce site and one aspen site, both underlain by granite (Elk spruce granite (ESG) and Elk aspen granite (EAG)). While ESG is on a dominantly south facing slope, it is north facing within the Elk Creek catchment.

3 Methods

To quantify the impact of aspen vs. conifer land cover on soil organic C dynamics at Coal Creek, we dug two pits roughly one meter deep at all five sites. The first series of pits were dug in the summer 2020 and 2021 (Table 1). The second series of pits were dug in the summer of 2022. Aspen was dominant at two sites (AS and EAG), while spruces were dominant at three sites (SS, ESG, and SG). Because aspen- and conifer-dominated forests in this region tend to occur on hillslopes of contrasting aspects, it was not possible to isolate land cover from aspect effects (e.g., temperature, radiation). While the sites were selected based on their land cover, other key ecosystem features underlying lithology (either granite or sandstone), and hillslope position (either backslopes or footslopes) (Fig. 1) also differed across the sites. We address these site features as potential sources of variation in our response variables in the discussion.

Soil pits were described following Schoeneberger et al. (2012), then each pit face was photographed with a high-resolution, digital single-lens reflex camera (D5600, Nikon, Minato City, Tokyo, Japan) to quantify rooting depth distributions following Billings et al. (2018). Bulk soil samples were collected by depth every 10 cm for the first set of pits (2020-2021), and by horizon for the second set of pits (2022). Samples were then immediately stored in a refrigerator or freezer until they could be ground, sieved to 2 mm and analyzed. Twice in the summer of 2022 (late June and mid-August), soil was collected from 3 auger sampling locations within ~100 meters of each pit to characterize soil chemistry. Soils were augured at 10 cm intervals to 110 cm (or deepest possible depth), and samples were stored in coolers with ice packs in the field and transported back to the lab and stored at 4 °C (most analyses) or frozen (DOC and microbial biomass C, exo-enzyme assays, nitrate).

Table 1. Sampling design and analysis for the soil pits and augers samples.

	Soil Pits	Soil Pits	Auger Samples
Timing	2020-2021	2022 <u>July</u>	2022

	(July and September)		(June and August)
Total Depth (cm)	~110	~110	~110
Soil Sampling Intervals	10 cm	Horizon	10 cm
Soil moisture and gas sensors	3 depths (15, 45, 110 cm)		
Root Distributions	X	X	
%C and %N	X	X	X
Extractable nitrate concentrations		X	
$\delta^{15}\text{N}$		X	
pH	X	X	X
Effective cation exchange capacity (ECEC)	X		
Texture	X		
Wet aggregate size distribution (ASD)		X	
Dissolved organic carbon (DOC)		X	X
Microbial biomass carbon		X	
β -glucosidase and N-acetyl- β -D-glucosaminidase		X	

3.1 Measuring Soil Organic C and Nitrogen Dynamics

We assessed SOC and SON concentrations and stocks and the likelihood of SOC and SON degradation by microbes by analyzing bulk soil samples at 10-cm intervals. We determined SOC and SON on subsamples (~75 mg) via an elemental analyzer (Vario Macro Cube, Elementar, Ronkonkoma, NY). We used SOC and SON concentration measurements to calculate each subsample's C:N ratio. To determine stocks of SOC in each horizon, we multiplied SOC concentrations by soil bulk density obtained in each horizon. Bulk density was measured using a three-dimensional laser scanner (3D Scanner Ultra HD, NextEngine, Inc., Santa Monica, CA) following Rossi et al. (2008).

We measured extractable, dissolved organic C (DOC) to estimate organic C that can be relatively easily mobilized and transported out of the soil profiles; note that this differs from DOC measured in soil porewater using lysimeters, and instead represents a salt-extractable pool. We analyzed soil samples at 10-cm intervals to auger refusal collected at each site during the growing season. Soil samples were extracted within three months of collection date. A total of 7.5 g of soil at field moisture was extracted with 30 ml of simulated rainwater (Laegdsmand et al., 1999). The rainwater extracts were comprised of 47.9 μM NaNO_3 , 4.69 μM KCl, 23.81 μM

CaCl₂ x 2H₂O, 12.09 μM MgSO₄ x 7H₂O , and 18.24 μM (NH₄)₂SO₄ and adjusted to a pH of 4.2 ± 0.5 using HCl. Samples were placed on a shaker table for 30 minutes and centrifuged at 80 Hz (s⁻¹) for 15 minutes. Samples were filtered through 0.45 μm nylon syringe filters and 50 ml acid washed syringes. Filtered samples were stored in 10 ml centrifuge tubes, frozen and shipped overnight in a cooler with dry ice to the University of Kansas. DOC was analyzed from the thawed samples using a Violet-pink Mn (III)-pyrophosphate solution and a microplate reader (Biotek, UT).

To better understand the potential for microbial activity in these soils, we quantified microbial biomass C by horizon from pits dug in the summer of 2022 (Brooks et al. 1985). We exposed 5 g of each soil sample to chloroform for 24 h. To these fumigated sub-samples and to 5 g of unfumigated sub-samples, we added 20 ml of 0.5 M K₂SO₄ and shook for 30-40 minutes at 220 rpm. These samples were filtered through a 0.45 μm polyethersulfone (PES) filter and their DOC concentration was determined via colorimetry (Bartlett and Ross 1988) on a Synergy HT microplate reader (Agilent, USA). To assess the degree to which soil microbial communities were generating exo-enzymes that catalyze soil organic matter decay and thus can provide assimilable C- and N-rich compounds, we quantified potential activity rates of two such enzymes. We measured activity of β-glucosidase and N-acetyl-β-D-glucosaminidase, herein referred to as BGase and NAGase, which are linked to microbial C (BGase) and N and C (NAGase) acquisition (Sinsabaugh and Moorhead, 1994; Allison et al., 2011, Stone et al., 2014), using 4-methylumbelliferyl β-D-glucopyranoside (for BGase) and 4-methylumbelliferyl N-acetyl-β-D-glucosaminide (for NAGase) fluorescent tags. These tags were added to slurries made from approximately 1 gram of soil and pH-adjusted 50 mM sodium acetate. We pipetted the blended sample into the desired substrate and incubated all plates at 25 °C for 18 hours. Fluorescence from a Synergy HT plate reader (Agilent, USA) was used as a proxy for each enzyme's capacity to cleave monomers from the respective molecules undergoing decay (DeForest, 2009; German et al., 2011).

We quantified salt-extractable NO₃⁻ because of its importance as a biotically-available form of N, and also because of its status as a readily leachable ion. As such, it can serve as an indicator of each soil's capacity to undergo elemental loss in surface soil with hydrologic fluxes, and provides a valuable point of comparison to DOC values. We extracted ~10 g (fresh weight) of each soil sample with 0.5M K₂SO₄ and repeated the shaking and filtering steps described above for MBC. Extracts were analyzed for NO₃⁻ (Synergy HT, Agilent, USA) using Shand et al. (2008), a microplate-based approach that relies on hydrazine sulphate and sulphanilamide to generate a color intensity directly related to NO₃⁻ concentration.

We also quantified soil organic matter δ¹⁵N, given these signatures' value as an indicator of the degree to which soil microbes have processed soil organic matter (Nadelhoffer and Fry 1988; Billings and Richter 2006). Sub-samples of each soil were dried, ground to fine powder, and

weighed into a tin capsule for analysis. Values of $\delta^{15}\text{N}$ were obtained at the Kansas State University Stable Isotope Lab, where an Elementar EA Vario Pyrocube linked to an Elementar Geovision Isotope Ratio Mass Spectrometer determine N concentration and $\delta^{15}\text{N}$, respectively.

3.2 Measuring Soil Chemical and Physical Properties

To better assess possible differences in the chemical and physical controls on SOC stability we also measured pH, effective cation exchange capacity (ECEC), soil texture, and wet aggregate size distribution (ASD). We focused on pH as it is known to strongly control microbial communities and mineral associated organic C (MAOC) (Kleber et al., 2015). The soil pH was determined in a 1:1 H_2O soil slurry (Soil Survey Staff, 2022). We focused on ECEC because ECEC has a high positive correlation with SOC, clay content, and aluminum and iron oxides (Solly et al., 2020), which are highly correlated with the formation of MAOC (Kleber et al., 2015). ECEC was determined by summing Ca, Mg, and K extracted using a Mehlich-3 solution (Culman et al., 2019). Mehlich-3 extraction was used instead of an ammonium acetate extraction, because the soils samples had a pH of <7.5 and there is very little to no calcium carbonate. In these conditions Mehlich-3 and ammonium acetate extractions yield similar ECEC values (Rutter et al., 2021).

We examined soil texture at each pit for several reasons. First, the total amount of clay is important to MAOC, and second, texture is known to impact the distribution and connectivity of pores. This connectivity influences how easily oxygen can diffuse into a soil profile and thus processes such as microbial respiration (Schjønning et al., 1999; Moldrup et al., 2001), and further regulates water and solute transport down-profile. Soil texture was analyzed on pit samples collected from 2020-2021 using a laser diffraction (LD) unit (Bettersizer S3, Bettersize Instruments, Dandong, Liaoning, China). Five grams of soil was sieved to 2 mm, and organic matter was removed by treating samples with 30% hydrogen peroxide. Ten ml of 10% sodium hexametaphosphate (HMP) was added to the solution to prevent flocculation. The soil solution was pipetted into the Bettersizer until obscuration levels were between 14-20. We set clay- silt and silt-sand boundaries to be 6.6 and 60.33 μm , respectively (Makó et al., 2017).

We quantified aggregate size distributions as one key metric of soil structure. Aggregate-size distributions were measured on each soil horizon following Nimmo and Perkins (2002). Briefly, around 25 g of the largest air-dried aggregates were fully saturated with a Dickson apparatus (Dickson et al., 1991), and placed on a Yoder device where sieves (#4, 10, 17, 70) and soil samples were raised and lowered in the water 2.8 cm per stroke at a rate of 36 strokes per a minute for 10 minutes. Following this agitation in water, the sieves with their respective aggregates were placed in a drying oven at 105 °C for 12 hours. The soil material remaining on each sieve was dispersed with 200 ml of 2 g L⁻¹ HMP, mixed for 10 minutes, passed through the sieve again, and oven-dried at 105 °C for 2 hours. Weights were recorded and mass fractions of water-stable aggregates were then calculated. Sieves divided aggregates into 5 classes:

aggregates > 4.76 mm, aggregates between 2–4.76 mm, aggregates between 0.21–1 mm, and aggregates less than 0.21 mm. To simplify our analysis, we agglomerated these into 3 classes following Souza et al. (2023): fine aggregates (< 0.21 mm), intermediate aggregates (0.21–4.76 mm), and coarse aggregates (> 4.76 mm). A weighted geometric mean aggregate diameter (GMD) was calculated for each triplicate using the mass fractions of each aggregate-size class; the mean and standard deviation were calculated from these triplicate values to represent the aggregate diameter of each sample. The GMD values were divided by SOC content and the resulting values were used to characterize the propensity of C to form aggregates.

3.3 Measuring Rooting Distributions

To determine the relationship between roots, and C stability and transport, we measured the fraction of soil volume containing fine and coarse roots throughout the soil profiles using images collected from all 10 pits (e.g., 2020/2021 and 2022). We used ImageJ (Schneider et al., 2012) to overlay each image with a 1x1 cm grid. We then manually checked each 1x1 cm grid cell for the presence of a fine root (diameter < 1 mm) or coarse root (diameter > 1mm) and noted these presence/absence scores for each grid cell. Our focus is the soil volume containing roots and thus directly influenced by roots. As such, only presence/absence and not count data were recorded, and in any cell containing both fine and coarse roots the presence of only the coarse root(s) was recorded given their greater volume (Billings et al., 2018). These measures are thus a conservative measure of direct root influence on soil volumes, derived at the cm scale for soil pedons. Centimeter-scale cell presence/absence data were transformed into the fraction of each 1-cm thick layer containing roots.

3.4 Sensor Data

Soil sensor arrays were installed in the first set of pits (2020/2021) at the completion of sampling. Sensors were installed at depths of 15 cm, 45 cm, and 110 cm (or deepest depth) to monitor soil temperature (°C) and volumetric water content (VWC ; EC-5, Meter Group, Pullman, WA), matric potential (kPa) (Teros 21, Meter Group, Pullman, WA), O₂ concentration (%) (IB201806, Apogee Instruments, Logan, UT) and CO₂ concentration (ppm) (F0275476, Eosense, Dartmouth, Canada). Data were collected every 30 minutes for moisture, matric potential, and temperature and hourly for O₂ and CO₂ given the power requirements. We focus on CO₂ and O₂ as they are indicators of soil microbial and root biotic activities including heterotrophic respiration. Microbial activity directly and indirectly affects the formation of MAOC, SOC stabilization, and microaggregation (Dohnalkova et al., 2022). We used sensor data to investigate additional environmental controls on C dynamics. We converted O₂ from millivolt readings to % by adding calibrated values to the millivolt value of O₂. Each calibrated value was specific to the sensor installed and determined using atmospheric concentrations prior to installation. To focus on the growing season, we selected data from June 15-August 29, which was 14 days before the first sample was collected (June 29th) and ending 14 days after the last

sample was collected (August 15th). AS reflects 2021 data, SS reflects average daily 2021 and 2022, and ESG, EAG, and SG reflects 2022 data. These differences were because pits were installed with sensors in different years and some of the instrumentation had power outages and other unforeseen issues. We averaged daily temperature and VWC by week and examined average and standard deviation of the O₂ and CO₂ over the growing season.

3.5 Data Analysis

Spatial replicates controlling for all ecosystem-scale factors were not feasible in this study. Instead, we advance our understanding of SOC stability by examining a more diverse suite of biotic and abiotic ecosystem characteristics than is often the case in SOC-focused work. Our work begins to unravel the complex interactions among cover type characteristics, soil properties, and hydrologic settings in SOC dynamics. We used Wilcoxon Rank sum tests to determine if differences between aspen and spruce concentrations of SOC, DOC, total soil nitrogen and nitrate, and of ratios of DOC to SOC were significant. We used linear mixed effects (LME) methods via the R package *lme4* (Bates et al., 2014) to assess the influence of vegetation type, depth, and their interaction (N=5) on soil abiotic conditions, various forms of soil nitrogen and C and $\delta^{15}\text{N}$, ASD, and root abundances. We tested if variables were normally distributed using the Shapiro-Wilks test and transformed the data to achieve a normal distribution if they were non-normal. The soil chemical properties of SOC, EOC, EOC:SOC, ECEC were log transformed, while C:N data were transformed with the function $x^{1/3}$. Root fractions and soil solution pH did not require transformation to meet model assumptions. We assessed if vegetation type exerted a meaningful influence on the previously mentioned variables by constructing four models. The two simplest models included only vegetation type or depth, both as fixed effects. A third model included those fixed effects additively (e.g., Vegetation + Depth), and a fourth model included their interaction. We resolved the lack of independence of soil depth within each pedon by incorporating site identifiers as a random effect term in the model. We then tested the normality of the model residuals using Shapiro-Wilk test. For all models that passed this test, we compared the model fits using analysis of variance (ANOVA) and visually examined model residual errors for homogeneity of variance; the best model fit was selected based on the lowest Akaike information criterion (AIC) following Hauser et al. (2020). We interpret the results of these LME models conservatively, given the low number of replicate sites for each land cover type. We could not perform a LME model on microbial biomass and enzyme data due to the relatively limited number of samples. This limited our ability to include vegetation*depth interactions in those models.

4 Results

4.1 Soil Properties and Development

Clay, silt, and sand content at the aspen sites (AS and EAG; Fig. S1) and one of the conifer sites (ESG) exhibited little variation with depth (average 33.1% clay and 18.8% sand), while the two other conifer sites had a greater sand and lower silt and clay content, particularly at depths greater than 25 cm (SS and SG; Fig. S1). Cation exchange capacity was similar among the aspen and conifer sites with averages of 7.2 ± 4.9 and 8.2 ± 6.8 (meq/100 g soil), respectively, with elevated values at the surface that declined with depth (Fig S2).

We were able to access and describe soil profiles to approximately 100 cm (Fig. 2, Table. S1). All sites had weak to moderately strong subangular blocky structure throughout the soil profile, and most sites had weak to moderately strong granular structure in A and upper B horizons. Dendritic tubular pores, interpreted to be abandoned root channels, were present throughout the soil profile of aspen sites, while they were less common in the conifer soil profiles. Aspen sites exhibited faint organic stains and organoargillans (i.e., dark, organic stained clay films) throughout the soil profile, while conifer sites had clay bridges and krotovina throughout the soil profile. The krotovina suggest greater bioturbation under conifer than aspen. Both vegetation types exhibited ferriargillans (i.e., clay coats that include Fe oxides), clay films, and charcoal, although ferriargillans and clay films were more prominent under conifer. Clay bridges, organoargillans, and ferriargillans indicate illuviation. Lithologic discontinuities were identified in SS, ESG, and EAG indicating colluvial inputs into these footslope pedons.




Figure 2: Soil profiles described at each site. Horizon colors represent the moist color of the soils as matched to the soil-color or Munsell chart.

Soils at both aspen sites (AS and EAG) are Ustic Haplocryolls with thick, SOC-enriched surface horizons (mollic epipedons) and showing evidence of incipient subsoil development in the form of moderately thick cambic horizons. Soils under conifer sites are Typic Haplocrypts (SS and SG) and Eutric Haplocryalfs (ESG). Although surface horizons under conifer were not as well-developed (ochric epipedons), the subsurface showed similar incipient pedogenesis in the form

of cambic horizons for SS and SG and greater development in the case of ESG where an argillic horizon was identified between 19-90 cm below the mineral surface.

4.2 Soil abiotic conditions

To understand how soil abiotic conditions are linked to SOC forms and processing pathways, we focused our analysis of soil temperature and moisture during the growing season (June – August; Fig. 3). As expected, soil temperature increased at all sites as the growing season progressed peaking in mid to late July, with the warmest temperatures observed near the surface and lower variability observed at depth. We also observed that the aspen sites (AS & EAG), which are on south-facing slopes, are warmer than conifer sites with an average surface soil (15 cm deep) temperature of 14.3 ± 1.2 and 10.4 ± 1.0 °C, respectively, during the growing season. Aspen sites were generally drier than conifer. The average volumetric water content in the surface soils (15 cm deep) at aspen sites was 0.15 ± 0.05 and the average volumetric water content at spruce sites was 0.24 ± 0.05 cm³ cm⁻³.

We also examined soil pH. Across the entire soil profile, pH was similar at the aspen and spruce sites (5.6 ± 0.3 and 5.3 ± 0.4 , respectively) but their depth trends differed with spruce soils having slightly more acidic pH near the surface compared to the aspen (Fig. S2). This trend reversed at approximately 60 cm, where the aspen soils became slightly more acidic compared to the conifer soils.




Figure 3: Temperature (a-e; °C) and volumetric water content (VWC (cm³ cm⁻³); f-i) data for aspen sandstone (AS; a, f), spruce sandstone (SS; b, g), spruce granite (ESG; c, h), aspen granite (EAG; d, i), and spruce granite (SG; e, j). AS reflects 2021 data, SS reflects averaged 2021 and 2022, ESG, EAG, and SG reflects 2022 data.

4.3 Soil Organic C and Nitrogen

Across all sites, SOC concentrations ranged from 46.0-62.6 mg g⁻¹ near the surface (5 cm deep) to 4.8 to 29.0 mg g⁻¹ at depth (Fig. 4a). SOC concentrations were generally higher under aspen compared to spruce sites ($p < 0.0001$; Fig. 4a), but LME models also suggest that the best fit model included a significant interaction between vegetation and depth ($p < 0.001$), suggesting that SOC declines with depth for both vegetation types but to a greater extent under spruce compared to aspen. Stocks of SOC for all depth intervals ranged between 0.01 and 1.31 kg m⁻² (Fig. 4b); these values did not exhibit consistent declines with depth or clear differences across cover type.

In contrast to SOC concentrations, DOC was generally higher under the spruce stands compared to aspen. Similar to SOC, a model including a significant interaction between vegetation and depth was the best predictor of DOC values ($p < 0.001$), likely reflecting variable DOC values at different depths in both vegetation types (Fig. 4c). The DOC:SOC ratio also exhibited a significant interaction between vegetation type and depth (Fig. 4d; $p = 0.0007$). As with DOC, this significant interaction likely reflects variable ratio values for each cover type across depths.




Figure 4: The mean and standard deviation of (a) soil organic carbon (SOC) concentrations, (b) SOC stock [by horizon per pit, solid lines indicate sites underlain by granite and dotted lines aspen], (c) dissolved organic carbon (DOC), and (d) the ratio of dissolved organic carbon to soil organic carbon (DOC:SOC) with depth under two different vegetation types, aspen (orange) and spruce (green).

Total soil nitrogen ranged from 0.2 mg g⁻¹ at depth to 4.63 mg g⁻¹ near the surface. A model including an interaction between vegetation type and depth was the best fit ($p = 0.003$; Fig. 5a). Aspen values were greater than those in spruce-dominated soils at all depths; the significant

interaction implies that the decline with depth was greater in spruce soils. Nitrate concentrations averaged $214 \pm 323 \text{ ng g}^{-1}$ near the surface and $2.3 \pm 3.8 \text{ ng g}^{-1}$ at depth. Nitrate was elevated under the aspen compared to the spruce sites (Fig. 5b), and values under both vegetation types varied with depth. A model that included both depth and vegetation type with no interaction was a meaningfully better predictor of nitrate concentrations than either depth or vegetation alone ($p = 0.035$), and including a depth-vegetation interaction did not improve model fit. Aspen soil C:N averaged 10.9 ± 1.1 and remained fairly constant with depth (Fig. 5c). The spruce sites showed greater variation with depth with a similar mean value of 19.3 in the top 20 cm but widely variable values at the deepest points, ranging from 4.6 to 28.7 (Fig. 5c). A model including a vegetation and depth interaction was a meaningfully better fit than all simpler models ($p = 0.0008$), suggesting that the visibly greater variation in C:N with depth in spruce soils was a significantly different depth trend from the fairly constant aspen values. The lowest value measured, at depth in one of the spruce forests, is suggestive of soil organic matter highly-processed by microbial communities (Ziegler et al. 2017).

$\delta^{15}\text{N}$ signatures showed less distinct depth trends compared to the total nitrogen and nitrate, mirroring the relative lack of clear depth trends in C:N. Though variation across sites limited our ability to find statistically-significant differences across vegetation types or a significant influence of depth, $\delta^{15}\text{N}$ of soil organic matter in spruce plots tended to be lower than that of the aspen (Fig. 5d), hinting that soil N has undergone more microbial processing (Nadelhoffer and Fry 1988; Billings and Richter 2006) under aspens compared to under conifers. This interpretation is consistent with the mean spruce C:N values being greater than those in aspen forests (Fig. 5c).




Figure 5: The mean and standard deviation of (a) soil nitrogen, (b) soil nitrate, (c) carbon to nitrogen ratio (C:N), and $\delta^{15}\text{N}$ with depth under two different vegetation types, aspen (orange) and spruce (green). For each mean and standard deviation, where error bars are not visible the deviation is smaller than the point.

4.4 Biotic activity

4.4.1 Roots

The LME models indicate that models including vegetation-depth interactions were the most effective at describing total and coarse root fractions ($p < 0.001$), with generally greater root abundances in the aspen compared to the spruce (Fig. 6a & c). In contrast, vegetation type offered no additional explanatory power to the depth-dependent fine root abundance ($p > 0.05$; Fig. 6b), suggesting that the greater total root abundance under aspen was driven by the coarse root fraction. The difference between aspen and spruce root abundances were continuous with depth for the total root fraction but more punctuated with coarse root fraction. For example, higher coarse root fractions were observed from 30-60 cm and greater than 90 cm for the aspen as compared to the spruce. Interestingly, overall spruce root fractions decreased faster with depth than aspen root fractions. When we standardized DOC with rooting abundance, we found generally greater concentrations of DOC per unit root abundance under spruce soils, particularly with respect to total and fine roots (Fig. 7).




Figure 6: The mean (points) and standard deviation(shading) of (a) total, (b) fine, and (c) coarse root fractions quantified at 1-cm depth interval under two different vegetation types, aspen (orange) and spruce (green).




Figure 7: The mean (points) and standard deviation (bars) of DOC divided by mean (a) total, (b) fine, and (c) coarse root fractions every 10 cm under two different vegetation types, aspen (orange) and spruce (green). Root fractions represent the count of fine (<1 mm) or coarse (≥ 1 mm) in 10 cm depth increments.

4.4.2 Enzyme activity and microbial biomass

Exo-enzyme activity, their ratios, and microbial biomass C decreased from the surface with depth (Fig. 8a.-d.). Exo-enzyme activity standardized by microbial biomass lacked distinct depth trends (data not shown). Beta values of exponential decay curves fit to these data, merged for each cover type, were larger (more negative) for the spruce sites compared to aspen, indicating steeper declines in exo-enzymatic activities, microbial biomass C, and BGase activity relative to NAGase activity in spruce-dominated forest soils.




Figure 8: Exponential curves fitting enzyme and microbial biomass data. Panels (a) β -glucosidase (BGase) (b) β -N-acetyl glucosaminidase (NAGase) (c) microbial biomass and (d) the ratio of BGase to NAGase. Each point represents one site and one depth, with each curve thus defined by multiple spruce and aspen sites.

4.4.3 Soil O₂ and CO₂

We examined soil O₂ and CO₂ concentrations during the growing season to better understand patterns of respiration (Fig. 9). Soil CO₂ concentrations increased with depth across all sites, while O₂ concentrations were more variable. Soil O₂ concentrations remained relatively stable at aspen sites and at spruce granite sites (AS, EAG, and SG). However, O₂ concentrations decreased with depth at the remaining two spruce sites—one sandstone and one granite (SS and ESG).




Figure 9: Average and standard error (obscured by symbols) of the soil O₂ (%) and CO₂ (%) concentrations during at aspen (orange) and conifer (green) sites during the growing season at depths 15 cm (light), 45 cm (medium), and 110 cm (dark), with lines connecting depths within each profile. AS, Aspen Sandstone; EAG, Elk Aspen Granite; ESG, Elk Spruce Granite; SG, Spruce Granite; SS, Spruce Sandstone.

4.4 Soil Aggregates

The mean geometric diameter of soil aggregates was generally smaller under aspen compared to spruce (Fig. 10a). Aspen aggregates tended to be finer, with fewer intermediate and coarse aggregates compared to the spruce soil (Fig. 10b-d). LME models indicated that vegetation-depth interactions were the most meaningful in driving all three aggregate size classes ($p < 0.001$). To explore the propensity of SOC to promote and stabilize aggregation in these soils, we standardized aggregates by SOC concentrations (Fig. 11; Souza et al. 2023). LME models indicate that an interaction between vegetation type and depth contributes to variation in the fine aggregate:SOC data (Fig. 11a; $p < 0.0001$), suggesting that in the bottom third of the profile, SOC in spruce soils also contributes to fine aggregate formation. Adding an interaction between vegetation and depth to the model improved model fit for coarse aggregate:SOC

values ($p < 0.0001$), highlighting a higher coarse aggregate:SOC ratio in the spruce sites that varied little with depth compared to aspen soils (Fig. 11b) and suggesting that SOC in spruce-dominated soils exhibits a higher propensity to form coarse aggregates throughout the soil profile.




Figure 10: The geometric mean and standard deviation of (a) all aggregates, and fraction of (b) fine aggregates (< 0.21 mm), (c) intermediate aggregates (0.21 - 4.76 mm), and (d) coarse aggregates (> 4.76 mm). Colors are associated with vegetation, aspen (orange) and spruce (green).




Figure 11: The mean and standard deviation of the fraction of (a) fine aggregates (< 0.21 mm) and (b) coarse aggregates (> 4.76 mm) divided by the fraction of SOC. Colors are associated with vegetation, aspen (orange) and spruce (green). Please note each aggregate size class is divided by the total SOC, not the C associated with each size class.

5 Discussion

By integrating knowledge from biology, pedology, hydrology, and soil chemistry we were better able to understand how multiple factors interact to drive observed SOC patterns in aspen and conifer montane forests. Our data indicate that differences in SOC protection give rise to often observed patterns of elevated SOC storage in soils under aspen compared to those in conifer stands (Woldeselassie et al., 2012, Laganieri et al., 2013, Boča et al., 2020, Román Dobarco et al., 2021). Furthermore, our study suggests that aspen-dominated soils may experience enhanced degrees of microbial transformation of SOC, with the products of those transformations exhibiting a greater tendency to reside in relatively small aggregates and thus protect C to a greater degree (Fig. 12). Consistent with this idea, we also observed less DOC loss in aspen soils compared to soil under spruce stands and slightly higher concentrations of DOC per unit root abundance under the spruce stands. These differences suggest greater infiltration of DOC to deeper horizons in spruce soils compared to those in aspen stands. It is important to highlight that spatial replicates controlling for all factors of interest at an ecosystem scale were not feasible, but that our work moves beyond considerations of vegetation biomass characteristics that often dominate investigations of contrasting SOC dynamics. Instead we begin to unravel the complex interactions among cover type characteristics, soil properties, and hydrologic settings (e.g., hillslope position). Below, we discuss the drivers of SOC form and fate in greater detail and interpret these findings in light of recent increases in stream water DOC concentration in this spruce-dominated watershed.




Figure 12. Summary of observations across aspen and spruce sites at Coal Creek, CO (USA) that are interpreted to indicate a greater amount of chemical and physical protection of SOC under aspen sites.

5.1 Microbial data are consistent with the Microbial Efficiency - Matrix Stabilization framework

Our data provide multiple lines of evidence that SOC protection, and thus fate, in these montane forests is largely controlled by biotic action linked to soil mineral material. Here greater values of total N, nitrate, BGase and $\delta^{15}\text{N}$ and lower C:N under aspen compared to spruce (Fig. 5, 8), suggest a greater degree of microbially processed organic matter under the aspen stands where greater SOC contents were measured (Fig. 4). These data hint that the microbial community under aspen stands functions in a manner consistent with the Microbial Efficiency - Matrix Stabilization (MEMS) framework (Cotrufo et al., 2012), transforming relatively labile leaf litter (e.g., under aspen) into byproducts more readily stabilized within soil profiles to a greater extent than appears to occur with slower-turnover litterfall (e.g., spruce). Differences in litterfall composition and thus decay rates across aspen and conifer species are typical, with generally lower lignin and higher nitrogen content in aspen litter (Moore et al., 2006). Our inference about litterfall differences promoting microbial byproduct stabilization is consistent with findings from across western Canada, where investigators observe relatively more active microbial communities under aspen compared to paired spruce stands throughout a growing season (Norris et al., 2016). Specifically, one interpretation of these C:N, $\delta^{15}\text{N}$, exo-enzyme, SOC, and DOC data at our sites is that tree species-specific composition of litterfall appears to have prompted greater microbial activities (Fig. 8), likely promoting greater contributions of microbial necromass to the SOC pool. This, in turn, may promote greater SOC retention in aspen-dominated soils; though investigation of specific necromass-derived compounds in these soils (e.g., Liang et al., 2019) is beyond the scope of this work, it represents a valuable way forward to testing this inference.

5.2 SOC transformations likely influence aggregate sizes and the probability of destabilization

The smaller aggregate sizes in aspen-dominated soils further support the notion that SOC stability is enhanced by higher microbial activity and increased necromass production rates. SOC is better protected and has generally longer mean residence times in smaller aggregates than larger aggregates (Six and Jastrow, 2002; Six et al., 2004). Literature hints that the larger size aggregates (Fig. 10c) and greater propensity for C to form large aggregates (Fig. 11b) observed in the spruce-dominated soils at our sites may be due to a greater abundance of particulate organic matter (POM) in spruce compared to aspen forest soils (Cotrufo et al., 2015; Cotrufo et al., 2019); this may be the case if spruce litterfall is more difficult to decompose. Taken together, these lines of evidence are consistent with aspen-dominated forests harboring SOC pools that tend to promote relatively small aggregate formation that can preserve SOC to a greater extent, particularly deeper in the profile where MAOC tends to dominate SOC pools (Jackson et al. 2017).

We also observed patterns in soil data suggesting that aspen soil SOC pools are more dominated by MAOC than those under spruce. The greater abundances of smaller aggregates and total soil nitrogen and nitrate concentrations, and lower C:N ratios (Fig. 5a-c), in aspen compared to conifer soils are consistent with relatively greater MAOC than POC concentrations (Kögel-Knaper et al., 2008; Ye et al., 2018; Sokal et al., 2022). Combined with the lower DOC:SOC ratio in aspen-dominated soils, these data suggest that a greater fraction of SOC in aspen-dominated soils is mineral-bound and relatively difficult to transform into microbially-available pools of DOC. We interpret these data to suggest that microbially-mediated

transformations of SOC promote differences in the abundance of MAOC and the physical structure of soil aggregates that leads to differences in the SOC protection.

5.3 Roots may indirectly regulate depth profiles of EOC losses

Roots can influence SOC stability through their promotion of both physical and chemical protection. Specifically, roots can play an important role in the formation and breakdown of soil aggregates (Oades, 1984; Singer et al., 1992; Le Bissonnais 1996; Attou et al., 1998), they can create biopores that can support the transport of DOC to depth (Sigen et al., 1997; Angers and Caron, 1998; Boger et al., 2010; Zhang et al., 2015; Lucas et al., 2019), and root exudates can prime microbial activity, enhance decomposition, and support the formation of MAOC (Jilling et al., 2021; Fossum et al., 2022). Our data revealed little direct correspondence of root abundance with SOC. However, per unit root abundance, spruce soils appear to harbor more DOC compared to aspen. This was particularly evident per unit total and fine roots, and is suggestive of greater movement of DOC through spruce soil profiles with potential greater losses of DOC to stream water compared to aspen-dominated soils. A complementary explanation would be that there are differences in the amount of DOC exudation by roots between the two species, and indeed such difference in exudation rates have been hypothesized in the literature (Buck and St. Clair, 2012; Boča et al., 2020). We might expect that greater exudation would lead to a greater increase in the MAOC pool and enhanced C stability (Even and Cotrufo, 2024), which could explain the lower values of DOC relative to SOC observed under aspen.

5.4 Aspect exerts some control on Coal Creek SOC dynamics.

South-facing slopes tend to be warmer and drier than north-facing slopes in the northern hemisphere (Burnett et al., 2008), and thus they can prompt more microbial decomposition of SOC. In our study aspen cover co-occurs where soil temperatures are warmer (Fig. 2). As such, it is possible that the exo-enzymatic signals of generally greater microbial activity in aspen-dominated soils compared to spruce-dominated soils (Fig. 8) is prompted more so by enhanced soil temperatures than by differences in aspen and spruce organic matter characteristics, and that enhanced soil temperatures also contribute to smaller soil aggregates, perhaps also due to greater microbial activities. We note that the volumetric fraction of soil moisture was also lower in the aspen, particularly at the shallowest soils, but that aspen soils appear to stay sufficiently moist (0.10-0.20 under aspen vs. 0.20-0.30 under the spruce) to support microbial activity responses to the higher temperatures. Consistent with this idea, soil CO₂ and O₂ concentrations generally suggest that microbial activities in the warmer, aspen-dominated soils are greater than in the cooler, spruce-dominated soils. Cooler, wetter conditions of the spruce-dominated soils, particularly following snow melt may prompt a deeper infiltration of moisture and DOC down profile, leading to the elevated DOC/root biomass observed under the spruce stands. While disentangling the impact of elevated soil temperatures from that of the chemical composition of organic inputs from aspen trees within the soil profile is difficult, soil nitrogen and $\delta^{15}\text{N}$ data are consistent with the idea that litterfall chemistry, and not just temperatures, promoted greater microbial activities in the aspen-dominated soils. We suggest that investigating the comprehensive, integrated effects of warmer, aspen-dominated sites on SOC dynamics compared to cooler, spruce-dominated sites offer a straightforward approach to assessing landscape-scale transitions in watershed C dynamics.

6 Changes in SOC destabilization and release have implications for stream water quality

Widespread increases in stream water DOC concentrations have been reported around the world in recent decades (Evans et al., 2005; Alvarez-Cobelas, 2012; Stanley et al., 2012; Pagano et al., 2014). Increases in stream water DOC concentration can harm global water quality by altering light and thermal regimes, nutrient cycling (e.g., Morris et al., 1995; Cory et al., 2015), the transport and bioavailability of heavy metals (e.g., Dupré et al., 1999; Trostle et al., 2016), and creating harmful disinfection byproducts (Leonard et al., 2022). Consistent with these global trends, recent findings at Coal Creek also report increasing DOC concentrations (Leonard et al., 2022; Kerins et al., 2024). As such, our research may help to shed light on drivers of stream water DOC, and thus has implications for changing drinking water quality in the region. Specifically, our work hints that differences in aggregate-size distributions may play an underappreciated role in influencing stream water C chemistry. Aggregate size can be modulated by vegetation type (i.e., smaller aggregates associated with Aspen) (Fig. 11; Jiménez et al., 2012; Zhao et al., 2017), and aggregation and disaggregation both represent mechanisms that can influence the transport of DOC to streams (Fan et al., 2022). Larger aggregates appear more prone to induce DOC transport into streams due to their relatively greater propensity to undergo fragmentation and associated loss of DOC (Cincotta et al., 2019; Fan et al., 2022).

Understanding how these different types of vegetation affect the chemical and physical properties of soil, and how this influences C release, is further complicated by climate change. Increasing temperature, a phenomenon evident in many Rocky Mountain environments including Coal Creek (Zhi et al., 2020), can cause aggregates to become less stable (Lavee et al., 1996, Wang et al., 2016), soil microbes to increase their C demand (Belay-Tedla et al., 2009, Hu et al., 2017), and recalcitrant C to undergo decay more rapidly (Luo et al., 2009). Dry soil conditions, which are often prompted by warming (Lakshmi et al., 2003), can induce a decrease in microbial biomass, which is often incorporated into stable aggregates (Gillballi et al., 2007). In addition to warming induced changes to subsurface properties and function, changing stand composition prompted by warming and drying can alter C dynamics. Some research indicates a high mortality rate among aspen stands and the expansion of conifer stands associated with increases in drought (Anderegg et al., 2013, Brewen et al., 2021), while others indicate the expansion of bark beetles and wildfires may promote the encroachment of aspen into conifer stands (Andrus et al., 2021). Our work suggests that the distribution of spruce and aspen in a watershed may influence soil release of DOC and its subsequent transport into streams, given that spruce vegetation appears to be associated with larger aggregates (Fig. 10), a potential for greater DOC loss per unit SOC (Fig. 4c), greater sand content (depth >25 cm; Fig. S1) and thus likely greater values of hydraulic conductivity, and generally higher soil moisture content (Fig. 3) compared to aspen-dominated soils. Thus, shifts in stand composition associated with perturbations linked to large-scale global changes have the potential to influence DOC transport from the hillslope to the stream.

7 Conclusions

Our work explores the interplay of different forest cover types and abiotic conditions in governing soil microbial activities, which then influence the propensity of SOC pools to form and stabilize soil aggregates of different sizes. In turn, these processes appear to promote varying capacities of a soil to protect SOC from destabilization. Our work contributes to the ongoing process of examining suites of biotic and abiotic whole-ecosystem (i.e., the critical zone, Richter and Billings, 2015) features to understand SOC dynamics (e.g., Keller, 2019; Mainka et al., 2022; Wasner et al., 2024) and offers a springboard for subsequent studies in which a greater number of spatially replicated sites across all gradients of interest may be feasible. Specifically, our data suggest that aspen-derived organic matter is linked to greater transformation rates by soil microbes and greater stabilization of SOC stocks, prompting a lower probability of relatively labile pools of SOC undergoing transport down-profile. If so, this suggests that aspen-dominated stands may experience a lower probability of promoting DOC transport across landscapes into streams. This phenomenon may be driven by greater rates of microbial necromass formation and generation of relatively smaller aggregates, and highlights how models like MEMS (Cotrufo et al., 2013) can be important for projecting not just CO₂ release to the atmosphere and SOC stabilization, but down-profile and downstream C transport as well. Though soil temperature differences likely played a role in the greater soil microbial activities in aspen, the generally higher nitrogen in aspen soils lends credence to the idea that litterfall chemistry itself played a key role in the higher rates of soil microbial activities. As such, the patterns that emerge in our data suggest that processes that control landcover ultimately also control SOC dynamics and soil structure in ways that may directly impact the delivery of organic C pools deep within soil profiles and stream water quality, and be sensitive to changing climatic conditions. Here, we demonstrate how the critical zone paradigm offers a valuable approach for examining, interrogating, and understanding watersheds, linking vegetation dynamics to subsurface processes and ultimately to the flux of water and C from hillslopes to streams.

Data statement

Soil sensor and soil properties data can be obtained at HydroShare, <http://www.hydroshare.org/resource/9948ad04a9a74246ad9bd5f8decb40b9>

Author Contributions (CRediT):

Wang, L: *Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Visualization Writing – original draft, Writing – review and editing*

Billings, S.: *Conceptualization, Funding acquisition, Investigation, Methodology, Writing – original draft, Writing – review and editing*

Li, L.: *Conceptualization, Investigation, Funding acquisition, Writing – review and editing*

Hirmas, D.R.: *Data Curation, Funding acquisition, Methodology, Investigation, Writing – review and editing*

Johnson, K.: *Data Curation, Investigation, Writing – review and editing*

Kerins, D. : *Investigation, Writing – review and editing*

Pachon, J: *Data Curation, Investigation, Writing – review and editing*

Curtis Beutler: *Investigation, Writing – review and editing*

Jarecke, K.M.: *Data Curation, Investigation, Writing – review and editing*

Varikuti, V: *Data Curation, Investigation, Writing – review and editing*

Unruh, Micah,: *Writing – review and editing*

Ajami, H: *Data Curation, Investigation, Funding acquisition, Methodology, Writing – review and editing*

Barnard, H.R.: *Investigation, Funding acquisition, Writing – review and editing*

Flores, A.N: *Funding acquisition, Writing – review and editing*

Williams, K. H.: *Funding acquisition, Investigation, Writing – review and editing*

Sullivan, P.L.: *Conceptualization, Funding acquisition, Methodology, Project administration, Supervision, Visualization, Writing – original draft, Writing – review and editing*

Competing Interests

The authors declare that they have no conflict of interest

Acknowledgements

We would like to thank Reece Gregory, Nicole Hornslein, Ariel Mollhagen, and Michael Mackenzie. This material is based upon work supported by the National Science Foundation under Grants NSF 2121694 (P. L. Sullivan) and 2012796 (P. L. Sullivan); NSF 2012669 (H. R. Barnard), the Department of Energy under Grant DE-SC0020146 (L. Li; P. L. Sullivan), NSF 2121639 (S.A. Billings), and NSF 2121760 (H. Ajami; D. Hirmas). This material is partially based upon work supported as part of the Watershed Function Scientific Focus Area funded by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research under Contract No. DE-AC02-05CH11231. Finally, our understanding of these sites benefited from data provided for project doi.org/10.46936/mone.proj.2023.60933/60008945 awarded to

SAB and PLS by the Molecular Observation Network (MONet) at the Environmental Molecular Sciences Laboratory (<https://ror.org/04rc0xn13>), a DOE Office of Science user facility sponsored by the Biological and Environmental Research program under Contract No. DE-AC05-76RL01830.

References

- Alaghmand, S., Beecham, S. and Hassanli, A., 2014. Impacts of vegetation cover on surface-groundwater flows and solute interactions in a semi-arid saline floodplain: a case study of the Lower Murray River, Australia. *Environmental Processes*, 1, pp.59-71. <https://doi.org/10.1007/s40710-014-0003-0>
- Alban, D.H., 1982. Effects of nutrient accumulation by aspen, spruce, and pine on soil properties. *Soil Science Society of America Journal*, 46(4), pp.853-861 <https://doi.org/10.2136/sssaj1982.03615995004600040037x>
- Alexander, R.R., 1987. *Ecology, silviculture, and management of the Engelmann spruce--subalpine fir type in the central and southern Rocky Mountains* (No. 659). US Department of Agriculture, Forest Service.
- Allison, S.D., 2014. Modeling adaptation of carbon use efficiency in microbial communities. *Frontiers in Microbiology*, 5, p.571. <https://doi.org/10.3389/fmicb.2014.00571>.
- Alvarez-Cobelas, M., Angeler, D. G., Sánchez-Carrillo, S., Almendros, G., 2012. A worldwide view of organic carbon export from catchments. *Biogeochemistry*, 107, 275-293. <https://doi.org/10.1007/s10533-010-9553-z>
- Amézketa, E., 1999. Soil aggregate stability: a review. *Journal of sustainable agriculture*, 14(2-3), pp.83-151. https://doi.org/10.1300/J064v14n02_08.
- Anderegg, L.D., Anderegg, W.R., Abatzoglou, J., Hausladen, A.M. and Berry, J.A., 2013. Drought characteristics' role in widespread aspen forest mortality across Colorado, USA. *Global Change Biology*, 19(5), pp.1526-1537. <https://doi.org/10.1111/gcb.12146>.
- Anderson, M.A., Graham, R.C., Alyanakian, G.J. and Martynn, D.Z., 1995. Late summer water status of soils and weathered bedrock in a giant sequoia grove. *Soil Science*, 160(6), pp.415-422. <https://doi.org/10.1097/00010694-199512000-00007>.
- Anderson, S.P., Hinckley, E.L., Kelly, P. and Langston, A., 2014. Variation in critical zone processes and architecture across slope aspects. *Procedia Earth and Planetary Science*, 10, pp.28-33. <https://doi.org/10.1016/j.proeps.2014.08.006>
- Andrus, R.A., Hart, S.J., Tutland, N. and Veblen, T.T., 2021. Future dominance by quaking aspen expected following short-interval, compounded disturbance interaction. *Ecosphere*, 12(1), p.e03345. <https://doi.org/10.1002/ecs2.3345>.

831 Angers, D.A. and Caron, J., 1998. Plant-induced changes in soil structure: processes and
832 feedbacks. *Biogeochemistry*, 42, pp.55-72. <https://doi.org/10.1023/A:1005944025343>

833 Araya, S.N. and Ghezzehei, T.A., 2019. Using machine learning for prediction of saturated
834 hydraulic conductivity and its sensitivity to soil structural perturbations. *Water Resources*
835 *Research*, 55(7), pp.5715-5737. <https://doi.org/10.1029/2018WR024357>

836 Attou, F., Bruand, A. and Le Bissonnais, Y., 1998. Effect of clay content and silt—clay fabric on
837 stability of artificial aggregates. *European Journal of Soil Science*, 49(4), pp.569-577.
838 <https://doi.org/10.1046/j.1365-2389.1998.4940569.x>

839 Averill, C., Turner, B.L. and Finzi, A.C., 2014. Mycorrhiza-mediated competition between plants
840 and decomposers drives soil carbon storage. *Nature*, 505(7484), pp.543-545.
841 <https://doi.org/10.1038/nature12901>

842 Banwart, S., Bernasconi, S.M., Bloem, J., Blum, W., Brandao, M., Brantley, S., Chabaux, F., Duffy,
843 C., Kram, P., Lair, G. and Lundin, L., 2011. Soil processes and functions in critical zone
844 observatories: hypotheses and experimental design. *Vadose Zone Journal*, 10(3), pp.974-987.
845 <https://doi.org/10.2136/vzj2010.0136>

846 Belay-Tedla, A., Zhou, X., Su, B., Wan, S. and Luo, Y., 2009. Labile, recalcitrant, and microbial
847 carbon and nitrogen pools of a tallgrass prairie soil in the US Great Plains subjected to
848 experimental warming and clipping. *Soil Biology and Biochemistry*, 41(1), pp.110-116.
849 <https://doi.org/10.1016/j.soilbio.2008.10.003>

850 Bergstrom, A., Jencso, K. and McGlynn, B., 2016. Spatiotemporal processes that contribute to
851 hydrologic exchange between hillslopes, valley bottoms, and streams. *Water Resources*
852 *Research*, 52(6), pp.4628-4645. <https://doi.org/10.1002/2015WR017972>

853 Berner, R.A., 1992. Weathering, plants, and the long-term carbon cycle. *Geochimica et*
854 *Cosmochimica Acta*, 56(8), pp.3225-3231. [https://doi.org/10.1016/0016-7037\(92\)90300-8](https://doi.org/10.1016/0016-7037(92)90300-8)

855 Besnard, E., Chenu, C., Balesdent, J., Puget, P. and Arrouays, D., 1996. Fate of particulate
856 organic matter in soil aggregates during cultivation. *European Journal of Soil Science*, 47(4),
857 pp.495-503. <https://doi.org/10.1111/j.1365-2389.1996.tb01849.x>

858 Billings, S.A., and Richter, D.D., 2006. Changes in stable isotopic signatures of soil nitrogen and
859 carbon during forty years of forest development. *Oecologia* 148: 325–333; 10.1007/s00442-
860 006-0366-7. <https://doi.org/10.1007/s00442-006-0366-7>

861 Billings, S.A., Hirmas, D., Sullivan, P.L., Lehmeier, C.A., Bagchi, S., Min, K., Brecheisen, Z., Hauser,
862 E., Stair, R., Flournoy, R. and deB. Richter, D., 2018. Loss of deep roots limits biogenic agents of
863 soil development that are only partially restored by decades of forest regeneration. *Elem Sci*
864 *Anth*, 6, p.34. <https://doi.org/10.1525/elementa.287>.

865 Billings, S.A., Lajtha, K., Malhotra, A., Berhe, A.A., de Graaff, M.-A., Earl, S., Fraterrigo, J.,
866 Georgiou, K., Grandy, S., Hobbie, S.E., Moore, J.A.M., Nadelhoffer, K., Pierson, D., Rasmussen,
867 C., Silver, W.L., Sulman, B.N., Weintraub, S., and Wieder, W., 2021. Soil organic carbon is not

868 just for soil scientists: Measurement recommendations for diverse practitioners. *Ecological*
869 *Applications* 31 doi: 10.1002/eap.2290

870 Boča, A., Jacobson, A.R. and Van Miegroet, H., 2020. Aspen soils retain more dissolved organic
871 carbon than conifer soils in a sorption experiment. *Frontiers in Forests and Global Change*, 3,
872 p.594473. <https://doi.org/10.3389/ffgc.2020.594473>.

873 Bogner, C., Gaul, D., Kolb, A., Schmiedinger, I. and Huwe, B., 2010. Investigating flow
874 mechanisms in a forest soil by mixed-effects modelling. *European Journal of Soil Science*, 61(6),
875 pp.1079-1090. <https://doi.org/10.1111/j.1365-2389.2010.01300.x>

876 Bornyasz, M.A., Graham, R.C. and Allen, M.F., 2005. Ectomycorrhizae in a soil-weathered
877 granitic bedrock regolith: linking matrix resources to plants. *Geoderma*, 126(1-2), pp.141-160.
878 <https://doi.org/10.1016/j.geoderma.2004.11.023>

879 Brewen, C.J., Berrill, J.P., Ritchie, M.W., Boston, K., Dagley, C.M., Jones, B., Coppoletta, M. and
880 Burnett, C.L., 2021. 76-year decline and recovery of aspen mediated by contrasting fire regimes:
881 Long-unburned, infrequent and frequent mixed-severity wildfire. *Plos one*, 16(2),
882 p.e0232995. <https://doi.org/10.1371/journal.pone.0232995>.

883 Brzostek, E.R., Greco, A., Drake, J.E. and Finzi, A.C., 2013. Root carbon inputs to the rhizosphere
884 stimulate extracellular enzyme activity and increase nitrogen availability in temperate forest
885 soils. *Biogeochemistry*, 115, pp.65-76. <https://doi.org/10.1007/s10533-012-9818-9>

886 Bronick, C.J. and Lal, R., 2005. Soil structure and management: a review. *Geoderma*, 124(1-2),
887 pp.3-22. <https://doi.org/10.1016/j.geoderma.2004.03.005>

888 Brusseau, M.L. and Rao, P.S.C., 1990. Modeling solute transport in structured soils: A
889 review. *Geoderma*, 46(1-3), pp.169-192. [https://doi.org/10.1016/0016-7061\(90\)90014-Z](https://doi.org/10.1016/0016-7061(90)90014-Z)

890 Buck, J.R. and St. Clair, S.B., 2012. Aspen increase soil moisture, nutrients, organic matter and
891 respiration in Rocky Mountain forest communities. *PLoS One*, 7(12),
892 p.e52369. <https://doi.org/10.1371/journal.pone.0052369>.

893 Buckeridge, K.M., Creamer, C. and Whitaker, J., 2022. Deconstructing the microbial necromass
894 continuum to inform soil carbon sequestration. *Functional Ecology*, 36(6), pp.1396-1410.
895 <https://doi.org/10.1111/1365-2435.14014>

896 Burnett, B.N., Meyer, G.A. and McFadden, L.D., 2008. 'Aspect-related microclimatic influences
897 on slope forms and processes, northeastern Arizona', *Journal of Geophysical Research*, 113(F3),
898 p. F03002. Available at: <https://doi.org/10.1029/2007JF000789>.

899 Brzostek, E.R., Greco, A., Drake, J.E. and Finzi, A.C., 2013. Root carbon inputs to the rhizosphere
900 stimulate extracellular enzyme activity and increase nitrogen availability in temperate forest
901 soils. *Biogeochemistry*, 115, pp.65-76. <https://doi.org/10.1007/s10533-012-9818-9>

902 Canelles, Q., Aquilué, N., James, P. M., Lawler, J., & Brotons, L. (2021). Global review on
903 interactions between insect pests and other forest disturbances. *Landscape Ecology*, 36, 945-
904 972. <https://doi.org/10.1007/s10980-021-01209-7>

- Carbone, M.S., Still, C.J., Ambrose, A.R., Dawson, T.E., Williams, A.P., Boot, C.M., Schaeffer, S.M. and Schimel, J.P., 2011. Seasonal and episodic moisture controls on plant and microbial contributions to soil respiration. *Oecologia*, 167, pp.265-278. <https://doi.org/10.1007/s00442-011-1975-3>
- Chen, S., Franklin, R.E. and Johnson, A.D., 1997. Clay film effects on ion transport in soil. *Soil science*, 162(2), pp.91-96.
- Chorover, J., Kretzschmar, R., Garcia-Pichel, F., and Sparks, D. L, 2007. Soil biogeochemical processes within the critical zone. *Elements*, 3(5), 321-326. <https://doi.org/10.2113/gselements.3.5.321>
- Cincotta, M. M., Perdrual, J.N., Shavitz, A., Libenson, A., Landsman-Gerjoi, M., Perdrual, N., Armfield, J., Adler, T., and Shanley, J.B., 2019. Soil aggregates as a source of dissolved organic carbon to streams: an experimental study on the effect of solution chemistry on water extractable carbon." *Frontiers in Environmental Science*: 172. <https://doi.org/10.3389/fenvs.2019.00172>.
- Clark, A.L. and Clair, S.B.S., 2011. Mycorrhizas and secondary succession in aspen–conifer forests: Light limitation differentially affects a dominant early and late successional species. *Forest Ecology and Management*, 262(2), pp.203 207. <https://doi.org/10.1016/j.foreco.2011.03.024>.
- Coop, J.D., Barker, K.J., Knight, A.D. and Pecharich, J.S., 2014. Aspen (*Populus tremuloides*) stand dynamics and understory plant community changes over 46 years near Crested Butte, Colorado, USA. *Forest Ecology and Management*, 318, pp.1 12. <https://doi.org/10.1016/j.foreco.2014.01.019>.
- Cory, R. M., Harrold, K. H., Neilson, B. T., and Kling, G. W., 2015. Controls on dissolved organic matter (DOM) degradation in a headwater stream: the influence of photochemical and hydrological conditions in determining light-limitation or substrate-limitation of photo-degradation. *Biogeosciences*, 12(22), 6669-6685. <https://doi.org/10.5194/bg-12-6669-2015>
- Cotrufo, M.F., Wallenstein, M.D., Boot, C.M., Denef, K. and Paul, E., 2013. The Microbial Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: Do labile plant inputs form stable soil organic matter?. *Global change biology*, 19(4), pp.988-995. <https://doi.org/10.1111/gcb.12113>
- Cotrufo, M.F., Soong, J.L., Horton, A.J., Campbell, E.E., Haddix, M.L., Wall, D.H. and Parton, W.J., 2015. Formation of soil organic matter via biochemical and physical pathways of litter mass loss. *Nature Geoscience*, 8(10), pp.776-779. <https://doi.org/10.1038/ngeo2520>.
- Cotrufo, M.F., Ranalli, M.G., Haddix, M.L., Six, J. and Lugato, E., 2019. Soil carbon storage informed by particulate and mineral-associated organic matter. *Nature Geoscience*, 12(12), pp.989-994. <https://doi.org/10.1038/s41561-019-0484-6>.

943 Cruz-Paredes, C., Tájmel, D. and Rousk, J., 2021. Can moisture affect temperature dependences
 944 of microbial growth and respiration?. *Soil Biology and Biochemistry*, 156, p.108223.
 945 <https://doi.org/10.1016/j.soilbio.2021.108223>

946 Cryer, D.H. and Murray, J.E., 1992. Aspen regeneration and soils. *Rangelands Archives*, 14(4),
 947 pp.223-226.

948 Culman, S., 2019. Calculating Cation Exchange Capacity, Base Saturation, and Calcium
 949 Saturation. Ohioline, 22 Aug., ohioline.osu.edu/factsheet/anr-81.

950 Dapples, E.C., 1947. Sandstone types and their associated depositional environments. *Journal of*
 951 *Sedimentary Research*, 17(3), pp.91-100. [https://doi.org/10.1306/D42692BA-2B26-11D7-](https://doi.org/10.1306/D42692BA-2B26-11D7-8648000102C1865D)
 952 [8648000102C1865D](https://doi.org/10.1306/D42692BA-2B26-11D7-8648000102C1865D)

953

954 DeForest, J.L., 2009. The influence of time, storage temperature, and substrate age on potential
 955 soil enzyme activity in acidic forest soils using MUB-linked substrates and L-DOPA. *Soil Biology*
 956 *and Biochemistry*, 41(6), pp.1180-1186. <https://doi.org/10.1016/j.soilbio.2009.02.029>
 957 [Get rights and content](https://doi.org/10.1016/j.soilbio.2009.02.029)

958 de Wit, H.A., Bryn, A., Hofgaard, A., Karstensen, J., Kvilevåg, M.M. and Peters, G.P., 2014.
 959 Climate warming feedback from mountain birch forest expansion: reduced albedo dominates
 960 carbon uptake. *Global Change Biology*, 20(7), pp.2344-2355.
 961 <https://doi.org/10.1111/gcb.12483>

962 Dickson, E.L., Rasiah, V. and Groenevelt, P.H., 1991. Comparison of four prewetting techniques
 963 in wet aggregate stability determination. *Canadian journal of soil science*, 71(1), pp.67-72.
 964 <https://doi.org/10.4141/cjss91-006>

965 Dohnalkova, A.C., Tfaily, M.M., Chu, R.K., Smith, A.P., Brislawn, C.J., Varga, T., Crump, A.R.,
 966 Kovarik, L., Thomashow, L.S., Harsh, J.B. and Keller, C.K., 2022. Effects of Microbial-Mineral
 967 Interactions on Organic Carbon Stabilization in a Ponderosa Pine Root Zone: A Micro-Scale
 968 Approach. *Frontiers in Earth Science*, 10, p.799694. <https://doi.org/10.3389/feart.2022.799694>

969 Dupré, B., Viers, J., Dandurand, J. L., Polve, M., Bénézech, P., Vervier, P., & Braun, J. J. (1999).
 970 Major and trace elements associated with colloids in organic-rich river waters: ultrafiltration of
 971 natural and spiked solutions. *Chemical Geology*, 160(1-2), 63-80.
 972 [https://doi.org/10.1016/S0009-2541\(99\)00060-1](https://doi.org/10.1016/S0009-2541(99)00060-1)

973 Eusterhues, K., Rumpel, C. and Kögel-Knabner, I., 2005. Organo-mineral associations in sandy
 974 acid forest soils: Importance of specific surface area, iron oxides and micropores. *European*
 975 *Journal of Soil Science*, 56(6), pp.753-763. <https://doi.org/10.1111/j.1365-2389.2005.00710.x>

976 Evans, C. D., Monteith, D. T., and Cooper, D. M., 2005. Long-term increases in surface water
 977 dissolved organic carbon: observations, possible causes and environmental
 978 impacts. *Environmental pollution*, 137(1), 55-71. <https://doi.org/10.1016/j.envpol.2004.12.031>

979 Even, R. J., & Cotrufo, M. F., 2024. The ability of soils to aggregate, more than the state of
 980 aggregation, promotes protected soil organic matter formation. *Geoderma*, 442, 116760.

981 Fang, C. and Moncrieff, J.B., 2001. The dependence of soil CO₂ efflux on temperature. *Soil*
 982 *Biology and Biochemistry*, 33(2), pp.155-165. [https://doi.org/10.1016/S0038-0717\(00\)00125-5](https://doi.org/10.1016/S0038-0717(00)00125-5)

983 Fossum, C., Estera-Molina, K.Y., Yuan, M., Herman, D.J., Chu-Jacoby, I., Nico, P.S., Morrison,
 984 K.D., Pett-Ridge, J. and Firestone, M.K., 2022. Belowground allocation and dynamics of recently
 985 fixed plant carbon in a California annual grassland. *Soil Biology and Biochemistry*, 165,
 986 p.108519. <https://doi.org/10.1016/j.soilbio.2021.108519>

987 Geological Survey (US) and Gaskill, D.L., 1991. *Geologic map of the gothic quadrangle, Gunnison*
 988 *County, Colorado*. The Survey. <https://doi.org/10.3133/gq1689>.

989 German, D.P., Weintraub, M.N., Grandy, A.S., Lauber, C.L., Rinkes, Z.L. and Allison, S.D., 2011.
 990 Optimization of hydrolytic and oxidative enzyme methods for ecosystem studies. *Soil Biology*
 991 *and Biochemistry*, 43(7), pp.1387-1397. <https://doi.org/10.1016/j.soilbio.2011.03.017>

992 Ghotsa Mekontchou, C., Houle, D., Bergeron, Y., Roy, M., Gardes, M., Séguin, A. and Drobyshv,
 993 I., 2022. Contrasting structure of root mycorrhizal communities of black spruce and trembling
 994 aspen in different layers of the soil profile in the boreal mixedwoods of eastern Canada. *Plant*
 995 *and Soil*, 479(1-2), pp.85-105. <https://doi.org/10.1007/s11104-022-05410-8>

996 Gilllabel, J., Denef, K., Brenner, J., Merckx, R. and Paustian, K., 2007. Carbon sequestration and
 997 soil aggregation in center-pivot irrigated and dryland cultivated farming systems. *Soil Science*
 998 *Society of America Journal*, 71(3), pp.1020-1028. <https://doi.org/10.2136/sssaj2006.0215>

999 Gislason, S.R., Oelkers, E.H., Eiriksdottir, E.S., Kardjilov, M.I., Gisladottir, G., Sigfusson, B.,
 1000 Snorrason, A., Elefsen, S., Hardardottir, J., Torssander, P. and Oskarsson, N., 2009. Direct
 1001 evidence of the feedback between climate and weathering. *Earth and Planetary Science*
 1002 *Letters*, 277(1-2), pp.213-222. <https://doi.org/10.1016/j.epsl.2008.10.018>

1003 Godsey, S.E., Kirchner, J.W. and Tague, C.L., 2014. Effects of changes in winter snowpacks on
 1004 summer low flows: case studies in the Sierra Nevada, California, USA. *Hydrological Processes*,
 1005 28(19), pp.5048-5064. <https://doi.org/10.1038/s41612-018-0012-1>.

1006 Graham, R., Rossi, A. and Hubbert, R., 2010. Rock to regolith conversion: Producing hospitable
 1007 substrates for terrestrial ecosystems. *GSA today*, 20, pp.4-9.

1008 Hemingway, J.D., Rothman, D.H., Grant, K.E. et al., 2019. Mineral protection regulates long-
 1009 term global preservation of natural organic carbon. *Nature* 570, 228–231.
 1010 <https://doi.org/10.1038/s41586-019-1280-6>

1011 Hodge, A., Berta, G., Doussan, C., Merchan, F. and Crespi, M., 2009. Plant root growth,
 1012 architecture and function. <https://doi.org/10.1007/s11104-009-9929-9>

1013 Hodges, C., Kim, H., Brantley, S.L. and Kaye, J., 2019. Soil CO₂ and O₂ concentrations illuminate
 1014 the relative importance of weathering and respiration to seasonal soil gas fluctuations. *Soil*

1015 *Science Society of America Journal*, 83(4), pp.1167-
 1016 1180. <https://doi.org/10.2136/sssaj2019.02.0049>.

1017 Hoff, C.C., 1957. A comparison of soil, climate, and biota of conifer and aspen communities in
 1018 the central Rocky Mountains. *The American Midland Naturalist*, 58(1), pp.115-
 1019 140. <https://doi.org/10.2307/2422365>

1020 Homer, C. H., Fry, J. A., and Barnes, C. A., 2012. The National land cover database, US geological
 1021 survey fact sheet 2012–3020. *US Geological Survey: Reston, VA, USA*.

1022 Hu, Y., Wang, Z., Wang, Q., Wang, S., Zhang, Z., Zhang, Z. and Zhao, Y., 2017. Climate change
 1023 affects soil labile organic carbon fractions in a Tibetan alpine meadow. *Journal of Soils and*
 1024 *Sediments*, 17, pp.326-339. <https://doi.org/10.1007/s11368-016-1565-4>

1025 Hubbard, Susan S., Kenneth Hurst Williams, Deb Agarwal, Jillian Banfield, Harry Beller, Nicholas
 1026 Bouskill, Eoin Brodie et al., 2019. The East River, Colorado, Watershed: A mountainous
 1027 community testbed for improving predictive understanding of multiscale hydrological–
 1028 biogeochemical dynamics. *Vadose Zone Journal* 17, no. 1: 1-25.
 1029 <https://doi.org/10.2136/vzj2018.03.0061>.

1030 Jackson, R. B., K. Lajtha, S. E. Crow, G. Hugelius, M. G. Kramer, and G. Pineiro. 2017. The ecology
 1031 of soil carbon: Pools, vulnerabilities, and biotic and abiotic controls. *Annual Review of Ecology,*
 1032 *Evolution, and Systematics* 48:419–445. [https://doi.org/10.1146/annurev-ecolsys-112414-](https://doi.org/10.1146/annurev-ecolsys-112414-054234)
 1033 [054234](https://doi.org/10.1146/annurev-ecolsys-112414-054234)

1034 Jastrow, J.D. 1996. Soil aggregate formation and the accrual of particulate and mineral-
 1035 associated organic matter. *Soil Biology and Biochemistry* 28:665-676.
 1036 [https://doi.org/10.1016/0038-0717\(95\)00159-X](https://doi.org/10.1016/0038-0717(95)00159-X)

1037 Jastrow, J.D., Miller, R.M. and Boutton, T.W., 1996. Carbon dynamics of aggregate-associated
 1038 organic matter estimated by carbon-13 natural abundance. *Soil Science Society of America*
 1039 *Journal*, 60(3), pp.801-807. <https://doi.org/10.2136/sssaj1996.03615995006000030017x>

1040 Jencso, K.G., McGlynn, B.L., Gooseff, M.N., Bencala, K.E. and Wondzell, S.M., 2010. Hillslope
 1041 hydrologic connectivity controls riparian groundwater turnover: Implications of catchment
 1042 structure for riparian buffering and stream water sources. *Water Resources Research*, 46(10).
 1043 <https://doi.org/10.1029/2009WR008818>

1044 Jiang, P., Shuai, P., Sun, A., Mudunuru, M. K., and Chen, X., 2023. Knowledge-informed deep
 1045 learning for hydrological model calibration: an application to Coal Creek Watershed in
 1046 Colorado. *Hydrology and Earth System Sciences*, 27(14), 2621-2644.
 1047 <https://doi.org/10.5194/hess-27-2621-2023>

1048 Jilling, A., Keiluweit, M., Gutknecht, J.L. and Grandy, A.S., 2021. Priming mechanisms providing
 1049 plants and microbes access to mineral-associated organic matter. *Soil Biology and Biochemistry*,
 1050 158, p.108265. <https://doi.org/10.1016/j.soilbio.2021.108265>

1051 Jin, L., Andrews, D. M., Holmes, G. H., Lin, H., and Brantley, S. L., 2011. Opening the “black box”:
 1052 Water chemistry reveals hydrological controls on weathering in the Susquehanna Shale Hills

1053 Critical Zone Observatory. *Vadose Zone Journal*, 10(3), 928– 942.
 1054 <https://doi.org/10.2136/vzj2010.0133>

1055 Johnson, K., Harpold, A., Carroll, R. W., Barnard, H., Raleigh, M. S., Segura, C., ... and Sullivan, P.
 1056 L., 2023. Leveraging Groundwater Dynamics to Improve Predictions of Summer Low-Flow
 1057 Discharges. *Water Resources Research*, 59(8), e2023WR035126.
 1058 <https://doi.org/10.1029/2023WR035126>

1059 Keller, C.K., 2019. Carbon exports from terrestrial ecosystems: A Critical-Zone framework.
 1060 *Ecosystems* 22:1691-1705. <https://doi.org/10.1007/s10021-019-00375-9>

1061 Kerins, D., and Li, L., 2023. High dissolved carbon concentration in arid rocky mountain streams.
 1062 *Environmental Science & Technology*, 57(11), 4656-4667.
 1063 <https://doi.org/10.1021/acs.est.2c06675>

1064 Kerins, D., Sadayappan, K., Zhi, W., Sullivan, P. L., Williams, K. H., Carroll, R. W., Barnard H.,
 1065 Sprenger M., Wenming D., Williams K., Pedrial J., and Li, L., 2024. Hydrology outweighs
 1066 temperature in driving production and export of dissolved carbon in a snowy mountain
 1067 catchment. *Water Resources Research*, 60(7), e2023WR036077.
 1068 <https://doi.org/10.1029/2023WR036077>

1069 Kerins D., Sullivan, P.L., Kayalvizhi S., Wei, Z., Rosemary C., Barnard H., Sprenger M., Wenming
 1070 D., Williams K., Pedrial J., Li L., in review. Dry Conditions Decrease Carbon Production and
 1071 Export but Amplify the Importance of Deep Carbon Respiration in a Water-Limited Mountain
 1072 Catchment.

1073 Kleber, M., Eusterhues, K., Keiluweit, M., Mikutta, C., Mikutta, R. and Nico, P.S., 2015. Mineral–
 1074 organic associations: formation, properties, and relevance in soil environments. *Advances in*
 1075 *agronomy*, 130, pp.1-140. <https://doi.org/10.1016/bs.agron.2014.10.005>.

1076 Kleber, M., Sollins, P. and Sutton, R., 2007. A conceptual model of organo-mineral interactions
 1077 in soils: self-assembly of organic molecular fragments into zonal structures on mineral surfaces.
 1078 *Biogeochemistry*, 85, pp.9-24. <https://doi.org/10.1007/s10533-007-9103-5>

1079 Kochenderfer, J.N., 1973. Root distribution under some forest types native to West Virginia.
 1080 *Ecology*, 54(2), pp.445-448. <https://doi.org/10.2307/1934355>

1081 Kögel-Knabner, I., Guggenberger, G., Kleber, M., Kandeler, E., Kalbitz, K., Scheu, S., Eusterhues,
 1082 K. and Leinweber, P., 2008. Organo-mineral associations in temperate soils: Integrating biology,
 1083 mineralogy, and organic matter chemistry. *Journal of Plant Nutrition and Soil Science*, 171(1),
 1084 pp.61-82. <https://doi.org/10.1002/jpln.200700048>

1085 Laegdsmand, M., Villholth, K.G., Ullum, M. and Jensen, K.H., 1999. Processes of colloid
 1086 mobilization and transport in macroporous soil monoliths. *Geoderma*, 93(1-2), pp.33-
 1087 59. [https://doi.org/10.1016/S0016-7061\(99\)00041-5](https://doi.org/10.1016/S0016-7061(99)00041-5)

1088 Laganiere, J., Paré, D., Bergeron, Y., Chen, H.Y., Brassard, B.W. and Cavard, X., 2013. Stability of
 1089 soil carbon stocks varies with forest composition in the Canadian boreal biome. *Ecosystems*, 16,
 1090 pp.852-865. <https://doi.org/10.1007/s10021-013-9658-z>.

1091 Laganière, J., Boča, A., Van Miegroet, H. and Paré, D., 2017. A tree species effect on soil that is
 1092 consistent across the species' range: the case of aspen and soil carbon in North America.
 1093 *Forests*, 8(4), p.113. <https://doi.org/10.3390/f8040113>.

1094 Lakshmi, V., Jackson, T.J. and Zehrhuhs, D., 2003. Soil moisture–temperature relationships:
 1095 results from two field experiments. *Hydrological processes*, 17(15), pp.3041-3057.
 1096 <https://doi.org/10.1002/hyp.1275>

1097 Lal, R., 2004. Mechanisms of Carbon Sequestration in Soil Aggregates AU-Blanco-Canqui,
 1098 Humberto. *Crit. Rev. Plant Sci*, 23(6), pp.481-504. <https://doi.org/10.1080/07352680490886842>

1099 Langston, A.L. *et al.* (2015) 'Evidence for climatic and hillslope-aspect controls on vadose zone
 1100 hydrology and implications for saprolite weathering: CLIMATIC CONTROL ON VADOSE ZONE
 1101 MOISTURE', *Earth Surface Processes and Landforms*, 40(9), pp. 1254–1269. Available at:
 1102 <https://doi.org/10.1002/esp.3718>.

1103 Larson, J. H., Frost, P. C., Lodge, D. M., & Lamberti, G. A. (2007). Photodegradation of dissolved
 1104 organic matter in forested streams of the northern Great Lakes region. *Journal of the North
 1105 American Benthological Society*, 26(3), 416-425.

1106 Lavelle, P., Spain, A., Fonte, S., Bedano, J. C., Blanchart, E., Galindo, V., ... and Zangerlé, A., 2020.
 1107 Soil aggregation, ecosystem engineers and the C cycle. *Acta Oecologica*, 105, 103561.
 1108 <https://doi.org/10.1016/j.actao.2020.103561>

1109 Leonard, Laura T., Gary F. Vanzin, Vanessa A. Garayburu-Caruso, Stephanie S. Lau, Curtis A.
 1110 Beutler, Alexander W. Newman, William A. Mitch, James C. Stegen, Kenneth H. Williams, and
 1111 Jonathan O. Sharp. Disinfection byproducts formed during drinking water treatment reveal an
 1112 export control point for dissolved organic matter in a subalpine headwater stream." *Water
 1113 Research X* 15 (2022): 100144. <https://doi.org/10.1016/j.wroa.2022.100144>.

1114 Leonard, L. T., Vanzin, G. F., Garayburu-Caruso, V. A., Lau, S. S., Beutler, C. A., Newman, A. W.,
 1115 ... & Sharp, J. O. (2022). Disinfection byproducts formed during drinking water treatment reveal
 1116 an export control point for dissolved organic matter in a subalpine headwater stream. *Water
 1117 Research X*, 15, 100144. <https://doi.org/10.1016/j.wroa.2022.100144>

1118 Liang, C., Amelung, W., Lehmann, J. and Kästner, M., 2019. Quantitative assessment of
 1119 microbial necromass contribution to soil organic matter. *Global change biology*, 25(11),
 1120 pp.3578-3590. <https://doi.org/10.1111/gcb.14781>

1121 Liu, X. and Biondi, F., 2021. Inter-specific transpiration differences between aspen, spruce, and
 1122 pine in a sky-island ecosystem of the North American Great Basin. *Forest Ecology and
 1123 Management*, 491, p.119157. <https://doi.org/10.1016/j.foreco.2021.119157>.

1124 Lucas, M., Schlüter, S., Vogel, H.J. and Vetterlein, D., 2019. Roots compact the surrounding soil
 1125 depending on the structures they encounter. *Scientific reports*, 9(1), p.16236.
 1126 <https://doi.org/10.1038/s41598-019-52665-w>

1127 Luo, C., Xu, G., Wang, Y., Wang, S., Lin, X., Hu, Y., Zhang, Z., Chang, X., Duan, J., Su, A. and Zhao,
 1128 X., 2009. Effects of grazing and experimental warming on DOC concentrations in the soil
 1129 solution on the Qinghai-Tibet plateau. *Soil Biology and Biochemistry*, 41(12), pp.2493-2500.
 1130 <https://doi.org/10.1016/j.soilbio.2009.09.006>

1131 Mambelli, S., Bird, J.A., Gleixner, G., Dawson, T.E. and Torn, M.S., 2011. Relative contribution of
 1132 foliar and fine root pine litter to the molecular composition of soil organic matter after in situ
 1133 degradation. *Organic Geochemistry*, 42(9), pp.1099-1108.
 1134 <https://doi.org/10.1016/j.orggeochem.2011.06.008>

1135 Mainka, M., Summerauer, L., Wasner, D., Garland, G., Gripenrog, M., Berhe, A.A., Doetterl, S.
 1136 2022. Soil geochemistry as a driver of soil organic matter composition: insights from a soil
 1137 chronosequence. *Biogeosciences* 19:1675-1689. <https://doi.org/10.5194/bg-19-1675-2022>

1138 Makiel, M. *et al.* (2022) 'Formation of iron oxyhydroxides as a result of glauconite weathering in
 1139 soils of temperate climate', *Geoderma*, 416, p. 115780. Available at:
 1140 <https://doi.org/10.1016/j.geoderma.2022.115780..>

1141 Mataix-Solera, J., Cerdà, A., Arcenegui, V., Jordán, A. and Zavala, L.M., 2011. Fire effects on soil
 1142 aggregation: a review. *Earth-Science Reviews*, 109(1-2), pp.44-60.
 1143 <https://doi.org/10.1016/j.earscirev.2011.08.002>

1144 Mauer, O. and Palátová, E., 2012. Root system development in Douglas fir (*Pseudotsuga*
 1145 *menziesii* [Mirb.] Franco) on fertile sites. *Journal of Forest Science*, 58(9), pp.400-
 1146 409. <https://doi.org/10.17221/94/2011-JFS>.

1147 Mekontchou, C.G., Houle, D., Bergeron, Y., Drobyshev, I. 2020. Contrasting root system
 1148 structure and belowground interactions between black spruce (*Picea mariana* (Mill.) B.S.P) and
 1149 trembling aspen (*Populus tremuloides* Michx) in boreal mixed woods of eastern Canada.
 1150 *Forests*. 11:127. <https://doi.org/10.3390/f11020127>

1151 Menzel, A. and Fabian, P., 1999. Growing season extended in Europe. *Nature*, 397(6721),
 1152 pp.659-659. <https://doi.org/10.1038/17709>

1153 Mikutta, R., Turner, S., Schippers, A., Gentsch, N., Meyer-Stüve, S., Condrón, L.M., Peltzer, D.A.,
 1154 Richardson, S.J., Eger, A., Hempel, G. and Kaiser, K., 2019. Microbial and abiotic controls on
 1155 mineral-associated organic matter in soil profiles along an ecosystem gradient. *Scientific*
 1156 *reports*, 9(1), p.10294. <https://doi.org/10.1038/s41598-019-46501-4>.

1157 Mitchell, P.J., Lane, P.N. and Benyon, R.G., 2012. Capturing within catchment variation in
 1158 evapotranspiration from montane forests using LiDAR canopy profiles with measured and
 1159 modelled fluxes of water. *Ecohydrology*, 5(6), pp.708-720. <https://doi.org/10.1002/eco.255>

1160 Moldrup, P., Deepagoda, T.C., Hamamoto, S., Komatsu, T., Kawamoto, K., Rolston, D.E. and de
 1161 Jonge, L.W., 2013. Structure-dependent water-induced linear reduction model for predicting

1162 gas diffusivity and tortuosity in repacked and intact soil. *Vadose Zone Journal*, 12(3).
 1163 <https://doi.org/10.2136/vzj2013.01.0026>

1164 Monteith, D.T., Stoddard, J.L., Evans, C.D., De Wit, H.A., Forsius, M., Høgåsen, T., Wilander, A.,
 1165 Skjelkvåle, B.L., Jeffries, D.S., Vuorenmaa, J. and Keller, B., 2007. Dissolved organic carbon
 1166 trends resulting from changes in atmospheric deposition chemistry. *Nature*, 450(7169), pp.537-
 1167 540. <https://doi.org/10.1038/nature06316>

1168 Moore, T.R., Trofymow, J.A., Prescott, C.E., Fyles, J. and Titus, B.D., 2006. Patterns of carbon,
 1169 nitrogen and phosphorus dynamics in decomposing foliar litter in Canadian forests. *Ecosystems*,
 1170 9, pp.46-62. <https://doi.org/10.1007/s10021-004-0026-x>.

1171 Morbidelli, R., Saltalippi, C., Flammini, A. and Govindaraju, R.S., 2018. Role of slope on
 1172 infiltration: A review. *Journal of hydrology*, 557, pp.878-886.
 1173 <https://doi.org/10.1016/j.jhydrol.2018.01.019>

1174 Morgan, M.D., 1969. Ecology of aspen in Gunnison County, Colorado. *American Midland*
 1175 *Naturalist*, pp.204-228. <https://doi.org/10.2307/2423831>

1176 Morris, D. P., Zagarese, H., Williamson, C. E., Balseiro, E. G., Hargreaves, B. R., Modenutti, B., ...
 1177 & Queimalinos, C. (1995). The attenuation of solar UV radiation in lakes and the role of
 1178 dissolved organic carbon. *Limnology and Oceanography*, 40(8), 1381-1391.
 1179 <https://doi.org/10.4319/lo.1995.40.8.1381>

1180 Nadelhoffer, K.J., Fry, B., 1988. Nitrogen-15 and carbon-13 abundances in forest soil organic
 1181 matter. *Soil Science Society of America Journal* 52:1633-1640.
 1182 <https://doi.org/10.2136/sssaj1988.03615995005200060024x>

1183 Nédeltcheva, T.H., Piedallu, C., Gégout, J.C., Stussi, J.M., Boudot, J.P., Angeli, N. and Dambrine,
 1184 E., 2006. Influence of granite mineralogy, rainfall, vegetation and relief on stream water
 1185 chemistry (Vosges Mountains, north-eastern France). *Chemical Geology*, 231(1-2), pp.1-15.
 1186 <https://doi.org/10.1016/j.chemgeo.2005.12.012>

1187 Neris, J., Jiménez, C., Fuentes, J., Morillas, G. and Tejedor, M., 2012. Vegetation and land-use
 1188 effects on soil properties and water infiltration of Andisols in Tenerife (Canary Islands, Spain).
 1189 *Catena*, 98, pp.55-62. <https://doi.org/10.1016/j.catena.2012.06.006>

1190 Neville, J., Tessier, J.L., Morrison, I., Scarratt, J., Canning, B. and Klironomos, J.N., 2002. Soil
 1191 depth distribution of ecto-and arbuscular mycorrhizal fungi associated with *Populus*
 1192 *tremuloides* within a 3-year-old boreal forest clear-cut. *Applied Soil Ecology*, 19(3), pp.209-216..
 1193 [https://doi.org/10.1016/S0929-1393\(01\)00193-7](https://doi.org/10.1016/S0929-1393(01)00193-7).

1194 Nicoll, B.C., Berthier, S., Achim, A., Gouskou, K., Danjon, F. and Van Beek, L.P.H., 2006. The
 1195 architecture of *Picea sitchensis* structural root systems on horizontal and sloping terrain. *Trees*,
 1196 20, pp.701-712. <https://doi.org/10.1007/s00468-006-0085-z>.

1197 Nikolakopoulou, M., Argerich, A., Drummond, J.D., Gacia, E., Martí, E., Sorolla, A. and Sabater,
 1198 F., 2018. Emergent macrophyte root architecture controls subsurface solute transport. *Water*
 1199 *Resources Research*, 54(9), pp.5958-5972. <https://doi.org/10.1029/2017WR022381>

1200 Nimmo, J.R. and Perkins, K.S., 2002. 2.6 Aggregate stability and size distribution. *Methods of soil*
 1201 *analysis: part 4 physical methods*, 5, pp.317-328. <https://doi.org/10.2136/sssabookser5.4.c14>

1202 Nye, P.H., 1981. Changes of pH across the rhizosphere induced by roots. *Plant and soil*, 61,
 1203 pp.7-26. <https://doi.org/10.1007/BF02277359>

1204 Norris, C. E., Quideau, S. A., & Oh, S. W. (2016). Microbial utilization of double-labeled aspen
 1205 litter in boreal aspen and spruce soils. *Soil Biology and Biochemistry*, 100, 9-20.
 1206 <https://doi.org/10.1016/j.soilbio.2016.05.013>

1207 Oades, J.M., 1984. Soil organic matter and structural stability: mechanisms and implications for
 1208 management. *Plant and soil*, 76, pp.319-337. <https://doi.org/10.1007/BF02205590>

1209 Okada, H. (1971). Classification of sandstone: analysis and proposal. *The Journal of Geology*,
 1210 79(5), 509-525. <https://doi.org/10.1306/D42692BA-2B26-11D7-8648000102C1865D>

1211 Ouyang, N., Zhang, Y., Sheng, H., Zhou, Q., Huang, Y. and Yu, Z., 2021. Clay mineral composition
 1212 of upland soils and its implication for pedogenesis and soil taxonomy in subtropical China.
 1213 *Scientific Reports*, 11(1), p.9707. <https://doi.org/10.1038/s41598-021-89049-y>.

1214 Pagano, T., Bida, M., & Kenny, J. E. (2014). Trends in levels of allochthonous dissolved organic
 1215 carbon in natural water: a review of potential mechanisms under a changing
 1216 climate. *Water*, 6(10), 2862-2897. <https://doi.org/10.3390/w6102862>

1217 Panhwar, Q.A., Naher, U.A., Shamshuddin, J., Othman, R. and Ismail, M.R., 2016. Applying
 1218 limestone or basalt in combination with bio-fertilizer to sustain rice production on an acid
 1219 sulfate soil in Malaysia. *Sustainability*, 8(7), p.700. <https://doi.org/10.3390/su8070700>

1220 Paré, D. and Bergeron, Y., 1996. Effect of colonizing tree species on soil nutrient availability in a
 1221 clay soil of the boreal mixedwood. *Canadian Journal of Forest Research*, 26(6), pp.1022-1031.
 1222 <https://doi.org/10.1139/x26-113>

1223 Popenoe, J.H., Bevis, K.A., Gordon, B.R., Sturhan, N.K. and Hauxwell, D.L., 1992. Soil-vegetation
 1224 relationships in Franciscan terrain of northwestern California. *Soil Science Society of America*
 1225 *Journal*, 56(6), pp.1951-1959. <https://doi.org/10.2136/sssaj1992.03615995005600060050x>

1226 Ramesh, T., Bolan, N.S., Kirkham, M.B., Wijesekara, H., Kanchikerimath, M., Rao, C.S., Sandeep,
 1227 S., Rinklebe, J., Ok, Y.S., Choudhury, B.U. and Wang, H., 2019. Soil organic carbon dynamics:
 1228 Impact of land use changes and management practices: A review. *Advances in agronomy*, 156,
 1229 pp.1-107. <https://doi.org/10.1016/bs.agron.2019.02.001>.

1230 Reisman, D., Rutkowski, T., Smart, P., Gusek, J. and Sieczkowski, M., 2009. Passive treatment
 1231 and monitoring at the standard mine superfund site Crested Butte, CO. *Proceedings America*
 1232 *Society of Mining and Reclamation*, pp.1107-1128.

1233 Rhoades, A.M., Ullrich, P.A. and Zarzycki, C.M., 2018. Projecting 21st century snowpack trends
1234 in western USA mountains using variable-resolution CESM. *Climate Dynamics*, 50(1-2), pp.261-
1235 288. <https://doi.org/10.1007/s00382-017-3606-0>.

1236 Riveros-Iregui, D.A., McGlynn, B.L., Marshall, L.A., Welsch, D.L., Emanuel, R.E. and Epstein, H.E.,
1237 2011. A watershed-scale assessment of a process soil CO₂ production and efflux model. *Water*
1238 *Resources Research*, 47(10). <https://doi.org/10.1029/2010WR009941>

1239 Roulet, N. and Moore, T.R., 2006. Browning the waters. *Nature*, 444(7117), pp.283-284.
1240 <https://doi.org/10.1038/444283a>

1241 Burnett, B. N., Meyer, G. A., & McFadden, L. D. (2008). Aspect-related microclimatic influences
1242 on slope forms and processes, northeastern Arizona. *Journal of Geophysical Research: Earth*
1243 *Surface*, 113(F3). <https://doi.org/10.1029/2010WR009941>

1244 Dobarco, M. R., & Van Miegroet, H. (2014). Soil Organic Carbon Storage and Stability in the
1245 Aspen-Conifer Ecotone in Montane Forests in Utah, USA. *Forests*, 5(4), 666-688.
1246 <https://doi.org/10.3390/f5040666>

1247 Román Dobarco, M., Jacobson, A.R. and Van Miegroet, H., 2021. Chemical composition of soil
1248 organic carbon from mixed aspen-conifer forests characterized with Fourier transform infrared
1249 spectroscopy. *European Journal of Soil Science*, 72(3), pp.1410-1430.
1250 <https://doi.org/10.1111/ejss.13065>.

1251 Rossi, A.M., Hirmas, D.R., Graham, R.C. and Sternberg, P.D., 2008. Bulk density determination
1252 by automated three-dimensional laser scanning. *Soil Science Society of America Journal*, 72(6),
1253 pp.1591-1593. <https://doi.org/10.2136/sssaj2008.0072N>

1254 Rutter, E.B., Ruiz Diaz, D. and Hargrave, L.M., 2022. Evaluation of Mehlich-3 for determination
1255 of cation exchange capacity in Kansas soils. *Soil Science Society of America Journal*, 86(1),
1256 pp.146-156. <https://doi.org/10.1002/saj2.20354>

1257 Sadnes, A., Eldhuset, T.D., Wollebaek, G. 2005. Organic acids in root exudates and soil solution
1258 of Norway spruce and silver birch. *Soil Biology and Biochemistry* 37:259-k269.
1259 <https://doi.org/10.1016/j.soilbio.2004.07.036>

1260 Sae-Tun, O., Bodner, G., Rosinger, C., Zechmeister-Boltenstern, S., Mentler, A. and Keiblinger,
1261 K., 2022. Fungal biomass and microbial necromass facilitate soil carbon sequestration and
1262 aggregate stability under different soil tillage intensities. *Applied Soil Ecology*, 179, p.104599.
1263 <https://doi.org/10.1016/j.apsoil.2022.104599>

1264 Scharlemann, J.P., Tanner, E.V., Hiederer, R. and Kapos, V., 2014. Global soil carbon:
1265 understanding and managing the largest terrestrial carbon pool. *Carbon management*, 5(1),
1266 pp.81-91. <https://doi.org/10.4155/cmt.13.77>

1267 Schjønning, P., Thomsen, I.K., Møberg, J.P., de Jonge, H., Kristensen, K. and Christensen, B.T.,
1268 1999. Turnover of organic matter in differently textured soils: I. Physical characteristics of
1269 structurally disturbed and intact soils. *Geoderma*, 89(3-4), pp.177-198.
1270 [https://doi.org/10.1016/S0016-7061\(98\)00083-4](https://doi.org/10.1016/S0016-7061(98)00083-4)

1271 Schneider, C.A., Rasband, W.S. and Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image
1272 analysis. *Nature methods*, 9(7), pp.671-675. <https://doi.org/10.1038/nmeth.2089>

1273 Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and Soil Survey Staff. 2012. Field book for
1274 describing and sampling soils, Version 3.0. Natural Resources Conservation Service, National
1275 Soil Survey Center, Lincoln, NE.

1276 Seyfried, G.S., Canham, C.D., Dalling, J.W. and Yang, W.H., 2021. The effects of tree-mycorrhizal
1277 type on soil organic matter properties from neighborhood to watershed scales. *Soil Biology and*
1278 *Biochemistry*, 161, p.108385. <https://doi.org/10.1016/j.soilbio.2021.108385>.

1279 Shand, C.A., Williams, B.L., Coutts, G., 2008. Determination of N-species in soil extracts using
1280 microplate techniques. *Talanta* 74 ,648-654; [10.1016/j.talanta.2007.06.039](https://doi.org/10.1016/j.talanta.2007.06.039)

1281 Shen, R., Pennell, K.G. and Suuberg, E.M., 2013. Influence of soil moisture on soil gas vapor
1282 concentration for vapor intrusion. *Environmental Engineering Science*, 30(10), pp.628-637.
1283 <https://doi.org/10.1089/ees.2013.0133>

1284 Shepperd, W., Rogers, P.C. and Bartos, D., 2006. Ecology, management, and restoration of
1285 aspen in the Sierra Nevada. <https://doi.org/10.2737/RMRS-GTR-178>.

1286 Silver, W.L., Lugo, A.E. and Keller, M., 1999. Soil oxygen availability and biogeochemistry along
1287 rainfall and topographic gradients in upland wet tropical forest soils. *Biogeochemistry*, 44,
1288 pp.301-328. <https://doi.org/10.1007/BF00996995>.

1289 Singer, M.J., Southard, R.J., Warrington, D.N. and Janitzky, P., 1992. Stability of synthetic sand-
1290 clay aggregates after wetting and drying cycles. *Soil Science Society of America Journal*, 56(6),
1291 pp.1843-1848. Le Bissonnais, Y., 1996. Aggregate stability and assessment of soil crustability and
1292 erodibility: I. Theory and methodology. *Eur. J. Soil Sci.* 47 (4), 425–437.
1293 <https://doi.org/10.2136/sssaj1992.03615995005600060032x>

1294 Singh, S.H.I.P.R.A., 2018. Understanding the role of slope aspect in shaping the
1295 vegetation attributes and soil properties in Montane ecosystems. *Tropical Ecology*, 59(3),
1296 pp.417-430.

1297 Sinsabaugh, R.L. and Moorhead, D.L., 1994. Resource allocation to extracellular enzyme
1298 production: a model for nitrogen and phosphorus control of litter decomposition. *Soil biology*
1299 *and biochemistry*, 26(10), pp.1305-1311. [https://doi.org/10.1016/0038-0717\(94\)90211-9](https://doi.org/10.1016/0038-0717(94)90211-9).

1300 Six, J., Bossuyt, H., Degryze, S. and Denef, K., 2004. A history of research on the link between
1301 (micro) aggregates, soil biota, and soil organic matter dynamics. *Soil and tillage research*, 79(1),
1302 pp.7-31. <https://doi.org/10.1016/j.still.2004.03.008>

1303 Six, J., Conant, R.T., Paul, E.A. and Paustian, K., 2002. Stabilization mechanisms of soil organic
1304 matter: implications for C-saturation of soils. *Plant and soil*, 241, pp.155
1305 176. <https://doi.org/10.1023/A:1016125726789>

1306 Six, J. and Jastrow, J.D., 2002. Organic matter turnover. *Encyclopedia of soil science*, 10.

1307 Soil Survey Staff, Natural Resources Conservation Service, United States Department of
 1308 Agriculture. Web Soil Survey. Available online at the following link:
 1309 <http://websoilsurvey.sc.egov.usda.gov/>. Accessed [5/21/2023].

1310 Soil Survey, S., 2014,. Keys to soil taxonomy.

1311 Sokol, N.W., Whalen, E.D., Jilling, A., Kallenbach, C., Pett-Ridge, J. and Georgiou, K., 2022. Global
 1312 distribution, formation and fate of mineral-associated soil organic matter under a changing
 1313 climate: A trait-based perspective. *Functional Ecology*, 36(6), pp.1411-
 1314 1429. <https://doi.org/10.1111/1365-2435.14040>

1315 Solly, E.F., Weber, V., Zimmermann, S., Walthert, L., Hagedorn, F. and Schmidt, M.W., 2020. A
 1316 critical evaluation of the relationship between the effective cation exchange capacity and soil
 1317 organic carbon content in Swiss forest soils. *Frontiers in Forests and Global Change*, 3,
 1318 p.98. <https://doi.org/10.3389/ffgc.2020.00098>.

1319 Souza, L.F., Hirmas, D.R., Sullivan, P.L., Reuman, D.C., Kirk, M.F., Li, L., Ajami, H., Wen, H., Sarto,
 1320 M.V., Loecke, T.D. and Rudick, A.K., 2023. Root distributions, precipitation, and soil structure
 1321 converge to govern soil organic carbon depth distributions. *Geoderma*, 437, p.116569.
 1322 <https://doi.org/10.1016/j.geoderma.2023.116569>

1323 Staff, S. S., 1999,. Soil taxonomy: a basic system of soil classification for making and interpreting
 1324 soil surveys. *Agriculture handbook*, 436.

1325 Stanley, E. H., Powers, S. M., Lottig, N. R., Buffam, I., & Crawford, J. T. (2012). Contemporary
 1326 changes in dissolved organic carbon (DOC) in human-dominated rivers: is there a role for DOC
 1327 management?. *Freshwater Biology*, 57, 26-42.
 1328 <https://doi.org/10.1111/j.1365-2427.2011.02613.x>

1329

1330 Stătescu, F., Zaucă, D.C. and Pavel, L.V., 2013. Soil structure and water-stable aggregates.
 1331 *Environmental Engineering & Management Journal (EEMJ)*,
 1332 12(4). <https://doi.org/10.30638/eemj.2013.091>.

1333 Sternberg, P.D., Anderson, M.A., Graham, R.C., Beyers, J.L. and Tice, K.R., 1996. Root
 1334 distribution and seasonal water status in weathered granitic bedrock under chaparral.
 1335 *Geoderma*, 72(1-2), pp.89-98. [https://doi.org/10.1016/0016-7061\(96\)00019-5](https://doi.org/10.1016/0016-7061(96)00019-5).

1336 Stone, E.L. and Kalisz, P.J., 1991. On the maximum extent of tree roots. *Forest ecology and*
 1337 *management*, 46(1-2), pp.59-102. [https://doi.org/10.1016/0378-1127\(91\)90245-Q](https://doi.org/10.1016/0378-1127(91)90245-Q)

1338 Stone, M.M., DeForest, J.L. and Plante, A.F., 2014. Changes in extracellular enzyme activity and
 1339 microbial community structure with soil depth at the Luquillo Critical Zone Observatory. *Soil*
 1340 *Biology and Biochemistry*, 75, pp.237-247. <https://doi.org/10.1016/j.soilbio.2014.04.017>.

1341 Stump, L.M. and Binkley, D., 1993. Relationships between litter quality and nitrogen availability
 1342 in Rocky Mountain forests. *Canadian Journal of Forest Research*, 23(3), pp.492-
 1343 502. <https://doi.org/10.1139/x93-067>

1344 Sullivan, P. L., Godd  ris, Y., Shi, Y., Gu, X., Schott, J., Hasenmueller, E. A., ... & Brantley, S. L.
 1345 (2019). Exploring the effect of aspect to inform future earthcasts of climate-driven changes in
 1346 weathering of shale. *Journal of Geophysical Research: Earth Surface*, 124(4), 974-993.
 1347 <https://doi.org/10.1029/2017JF004556>

1348 Sullivan, P.L., Billings, S.A., Hirmas, D., Li, L., Zhang, X., Ziegler, S., Murenbeeld, K., Ajami, H.,
 1349 Guthrie, A., Singha, K. and Gim  nez, D., 2022. Embracing the dynamic nature of soil structure: A
 1350 paradigm illuminating the role of life in critical zones of the Anthropocene. *Earth-Science*
 1351 *Reviews*, 225, p.103873.<https://doi.org/10.1016/j.earscirev.2021.103873>.

1352 Tew, R.K., 1968. *Properties of soil under aspen and herb-shrub cover* (Vol. 78). US Department
 1353 of Agriculture, Forest Service, Intermountain Forest & Range Experiment Station.

1354 Tisdall, J.M. and OADES, J.M., 1982. Organic matter and water-stable aggregates in soils. *Journal*
 1355 *of soil science*, 33(2), pp.141-163. <https://doi.org/10.1111/j.1365-2389.1982.tb01755.x>
 1356

1357 von L  tzow, M., K  gel-Knabner, I., Ludwig, B., Matzner, E., Flessa, H., Ekschmitt, K.,
 1358 Guggenberger, G., Marschner, B. and Kalbitz, K., 2008. Stabilization mechanisms of organic
 1359 matter in four temperate soils: Development and application of a conceptual model. *Journal of*
 1360 *Plant Nutrition and Soil Science*, 171(1), pp.111-124. <https://doi.org/10.1002/jpln.200700047>

1361

1362 Wadgyamar, S.M., Ogilvie, J.E., Inouye, D.W., Weis, A.E. and Anderson, J.T., 2018. Phenological
 1363 responses to multiple environmental drivers under climate change: insights from a long-term
 1364 observational study and a manipulative field experiment. *New Phytologist*, 218(2), pp.517-
 1365 529.<https://doi.org/10.1111/nph.15029>.

1366 Wainwright, Haruko M., Sebastian Uhlemann, Maya Franklin, Nicola Falco, Nicholas J. Bouskill,
 1367 Michelle E. Newcomer, Baptiste Dafflon et al. Watershed zonation through hillslope clustering
 1368 for tractably quantifying above-and below-ground watershed heterogeneity and functions.
 1369 *Hydrology and Earth System Sciences* 26, no. 2 (2022): 429-444. [https://doi.org/10.5194/hess-](https://doi.org/10.5194/hess-26-429-2022)
 1370 [26-429-2022, 2022](https://doi.org/10.5194/hess-26-429-2022)

1371 Wang, Y., Gao, S., Li, C., Zhang, J. and Wang, L., 2016. Effects of temperature on soil organic
 1372 carbon fractions contents, aggregate stability and structural characteristics of humic substances
 1373 in a Mollisol. *Journal of Soils and Sediments*, 16, pp.1849-1857.[https://doi.org/10.1007/s11368-](https://doi.org/10.1007/s11368-016-1379-4)
 1374 [016-1379-4](https://doi.org/10.1007/s11368-016-1379-4)

1375 Wang, Q., Xiao, J., Ding, J., Zou, T. 2021. Differences in root exudate inputs and rhizosphere
1376 effects on soil N transformation between deciduous and evergreen trees. *Plant and Soil* 458
1377 <https://doi.org/10.1007/s11104-019-04156-0>

1378 Wasner,D., Abramoff,R., Griepentrog,M., Venegas,E.Z., Boeckx,P., & Doetterl,S. (2024) The role
1379 of climate, mineralogy and stable aggregates for soil organic carbon dynamics along a
1380 geoclimatic gradient. *Global Biogeochemical Cycles*, 38,e2023GB007934.
1381 <https://doi.org/10.1029/2023GB007934>

1382 Weil, R.R. and Brady, N.C., 2017. *The nature and properties of soils* (Fifteen Ed).

1383 White, T., Brantley, S., Banwart, S., Chorover, J., Dietrich, W., Derry, L., et al., 2015. The role of
1384 critical zone observatories in critical zone science. In: *Developments in Earth Surface Processes*,
1385 vol. 19. Elsevier, pp. 15–78. <https://doi.org/10.1016/B978-0-444-63369-9.00002-1>

1386 Witty, J.H., Graham, R.C., Hubbert, K.R., Doolittle, J.A. and Wald, J.A., 2003. Contributions of
1387 water supply from the weathered bedrock zone to forest soil quality. *Geoderma*, 114(3-4),
1388 pp.389-400.[https://doi.org/10.1016/S0016-7061\(03\)00051-X](https://doi.org/10.1016/S0016-7061(03)00051-X).

1389 Woldeeslassie, M., Van Miegroet, H., Gruselle, M.C. and Hambly, N., 2012. Storage and stability
1390 of soil organic carbon in aspen and conifer forest soils of northern Utah. *Soil Science Society of*
1391 *America Journal*, 76(6), pp.2230-2240..<https://doi.org/10.2136/sssaj2011.0364>.

1392 Woolf, D., and Lehmann, J. 2019. Microbial models with minimal mineral protection can explain
1393 long-term soil organic carbon persistence. *Sci Rep* 9, 6522. [https://doi.org/10.1038/s41598-](https://doi.org/10.1038/s41598-019-43026-8)
1394 [019-43026-8](https://doi.org/10.1038/s41598-019-43026-8)

1395 Yang, J. Q., Zhang, X., Bourg, I. C., and Stone, H. A., 2021. 4D imaging reveals mechanisms of
1396 clay-carbon protection and release. *Nature communications*, 12(1), 622.

1397 Ye, C., Chen, D., Hall, S.J., Pan, S., Yan, X., Bai, T., Guo, H., Zhang, Y., Bai, Y. and Hu, S., 2018.
1398 Reconciling multiple impacts of nitrogen enrichment on soil carbon: plant, microbial and
1399 geochemical controls. *Ecology Letters*, 21(8), pp.1162-1173.<https://doi.org/10.1111/ele.13083>.

1400 Zhai, J., Liu, R., Liu, J., Zhao, G. and Huang, L., 2014. Radiative forcing over China due to albedo
1401 change caused by land cover change during 1990–2010. *Journal of Geographical Sciences*, 24,
1402 pp.789-801. <https://doi.org/10.1007/s11442-014-1120-4>

1403 Zhang, Y., Niu, J., Yu, X., Zhu, W. and Du, X., 2015. Effects of fine root length density and root
1404 biomass on soil preferential flow in forest ecosystems. *Forest Systems*, 24(1),
1405 p.12.<https://doi.org/10.5424/fs/2015241-06048>.

1406 Zhao, D., Xu, M., Liu, G., Ma, L., Zhang, S., Xiao, T. and Peng, G., 2017. Effect of vegetation type
1407 on microstructure of soil aggregates on the Loess Plateau, China. *Agriculture, ecosystems &*
1408 *environment*, 242, pp.1-8. <https://doi.org/10.1016/j.agee.2017.03.014>

1409 Zhi, W., Williams, K.H., Carroll, R.W., Brown, W., Dong, W., Kerins, D. and Li, L., 2020. Significant
1410 stream chemistry response to temperature variations in a high-elevation mountain watershed.
1411 *Communications Earth & Environment*, 1(1), p.43.[https://doi.org/10.1038/s43247-020-00039-](https://doi.org/10.1038/s43247-020-00039-w)
1412 [w](https://doi.org/10.1038/s43247-020-00039-w).
1413 Ziegler, S.E., Benner, R., Billings, S.A., Edwards, K.A., Philben, M., Zhu, X., Laganière, J. 2017.
1414 Climate warming can accelerate carbon fluxes without changing soil carbon stocks. *Frontiers in*
1415 *Earth Science* 5:2 doi: 10.3389/feart.2017.00002.