

The Diurnal Susceptibility of Subtropical Clouds to Aerosols

Marcin J. Kurowski¹, Matthew D. Lebsock¹, and Kevin M. Smalley²

¹Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA.

²Lawrence Livermore National Laboratory, Livermore, California, USA.

Abstract. The diurnal susceptibility of clouds and their radiative properties to aerosols is examined during ~~the~~their Lagrangian transition from subtropical stratocumulus to shallow cumulus regimes. Using large-eddy simulations, ~~a~~we analyze the six-day ~~air-mass evolution along a 3800-km~~evolution of an air mass along a 3,800-km observed trajectory from the coast of Peru toward the equator ~~is analyzed~~. Pristine and polluted scenarios are simulated with forcing imposed from weather reanalysis. The polluted scenario exhibits stronger diurnal variations in cloud water, cloud fraction, and albedo, with enhanced nighttime entrainment and suppressed precipitation. The overall response of cloud properties and outgoing shortwave radiation to droplet number concentration follows a distinct diurnal pattern: strong positive cloud adjustments dominate at night and in the morning, while weak negative adjustments prevail in the afternoon. This cycle is driven by the competition between precipitation suppression, which enhances cloud water and coverage, and entrainment drying, which depletes them. In polluted conditions, enhanced entrainment leads to a deeper and more decoupled boundary layer that cannot be sustained by surface fluxes in the afternoon, resulting in negative cloud adjustments. ~~This entrainment enhancement is mediated by the~~The enhanced entrainment rate under polluted conditions is caused by the reduced sedimentation of cloud and precipitation water from the entrainment zone. While the Twomey effect dominates the diurnal average albedo response, the diurnal variation in the competing cloud adjustments lead to a near-neutral net adjustment effect in the afternoon, highlighting the critical role of ~~diurnal~~diurnally varying processes in aerosol-cloud interactions.

. The authors' copyright for this publication is transferred to Jet Propulsion Laboratory, California Institute of Technology.

1 Introduction

The interactions between aerosol and clouds represent one of the largest sources of uncertainties in the anthropogenic radiative forcing of Earth's climate (IPCC, 2021, 2022). The radiative effect of the collective set of changes to cloud morphology by aerosol is known as the Effective Radiative Forcing due to Aerosol Cloud Interactions (~~ERF_{ACI}~~) (Wall et al., 2022) (ERF_{ACI}; Boucher et al., 2013), which is composed of a number of different cloud changes (Twomey, 1977; Albrecht, 1989; Boucher and Lohmann, 1995; Lohmann and Lohmann, 2000). The first order effect, often referred to as the Twomey effect (Twomey, 1977) posits that an increase in cloud droplet number (N_c) for fixed cloud liquid water path (LWP_c) results in a greater integrated water droplet cross sectional area and thus an increase in cloud optical depth (τ_c) and cloud albedo (A_c). The magnitude of the Twomey effect is thought to

be relatively well understood (Quaas et al., 2020) (Fan et al., 2016; Bellouin et al., 2020; Quaas et al., 2020). However, second-order indirect effects, or cloud adjustments, result from changes to the cloud liquid water path (LWP_c) and cloud cover fraction (f_c), where the domain mean liquid water path is $LWP = f_c LWP_c$. These cloud adjustments are less well understood. It was first thought that increases in N_c would inhibit the formation of precipitation and thus increase cloud lifetime (Albrecht, 1989; Pincus and Baker, 1994). More recently, it was suggested that increasing N_c can reduce LWP_c through a decreased sedimentation efficiency causing an increase in liquid near the cloud top which enhances the efficiency of the entrainment of dry free-tropospheric air into the cloud layer (Bretherton et al., 2007; Ackerman et al., 2004). (Ackerman et al., 2004; Bretherton et al., 2007)

To quantify the various aerosol-cloud interactions, the sensitivity of the reflected shortwave flux ($F^\uparrow$) is often decomposed into three terms (Bellouin et al., 2020) representing changes in τ_c (cloud optical depth) at fixed LWP_c (S_N), additional changes in τ_c resulting from changes in LWP_c at fixed N_c (S_{LWP}) and changes in f_c at fixed τ_c (S_f):

$$\frac{d \ln F^\uparrow}{d \ln N_c} \frac{d F^\uparrow}{d \ln N_c} = \underbrace{\text{Twomey Effect}(S_N)}_{\frac{\partial F^\uparrow}{\partial \ln N_c}} + \underbrace{\text{LWP adjustment}(S_{LWP})}_{\frac{\partial F^\uparrow}{\partial \ln LWP_c} \cdot \frac{d \ln LWP_c}{d \ln N_c}} + \underbrace{\text{fraction adjustment}(S_f)}_{\frac{\partial F^\uparrow}{\partial \ln f_c} \cdot \frac{d \ln f_c}{d \ln N_c}} \quad (1)$$

There is observational evidence for both increases and decreases in the LWP_c . For example, Han et al. (2002) use satellite data to show that clouds have positive, negative, and neutral sensitivity to aerosol in roughly equal proportions. It is also clear that the sign of the response is dependent on the cloud state. Lebsock et al. (2008) find that the LWP_c tends to increase with increased N_c for precipitating clouds and decrease with increasing N_c for non-precipitating clouds. Evidence from ship-tracks show both positive and negative sensitivity (Ackerman et al., 2000; Coakley and Walsh, 2002), with the observation that the sign of the response is associated with the mesoscale cellular structure with open-celled regimes tending to have a positive response and closed-cells tending to have a negative response, presumably due to their differential propensity to precipitate (Christensen and Stephens, 2012). A recent review of polluted clouds down-wind of anthropogenic pollution sources finds a weak albeit slightly negative average response of LWP_c to aerosol perturbations (Toll et al., 2019). To the contrary, Manshausen et al. (2022) recently find a large positive increases in LWP_c by using ship location data to find a large number of ‘invisible’ ship tracks, which are not readily identifiable in satellite imagery. Regional variability and observational uncertainties, such as cloud regime differences, further complicate LWP responses (Wood, 2012). We note that cloud adjustments are dependent on the background N_c , which can explain the presence of both positive and negative adjustments without contradiction. This state dependence in the cloud adjustment is manifest as the ‘inverted V’ relationship between N_c and LWP_c implying positive adjustment at low N_c and negative adjustment at high N_c (Gryspeerd et al., 2022).

Observed positive correlations between aerosol optical depth and f_c have long been considered suspect due to the tendency to observe enhanced clear sky reflectance in the vicinity of clouds due to three dimensional radiative effects (Várnai and Marshak, 2009). For example, carefully controlling for the distance of an aerosol retrieval to the nearest cloud nearly halves

the magnitude of the relationship between f_c and aerosol optical depth (Christensen et al., 2017). To entirely avoid the influence of artificial correlations, more recent observational studies have used either the observed N_c or a model derived aerosol field in place of the aerosol optical depth to derive the slope $df_c/d\ln N_c$. Although the magnitude is highly uncertain, studies tend to find a positive correlation (Gryspeerd et al., 2016; Wall et al., 2023).

Most observational satellite studies are based on visible and near infrared imager data with fixed diurnal sampling time therefore there are few hints as to the observed diurnal cycle of the cloud adjustments. A study of a South Atlantic shipping lane shows that Terra MODIS shows a larger positive LWP adjustment than Aqua MODIS, and the Terra/Aqua show positive/negative f_c adjustments (Diamond et al., 2020). The recent observational study of Smalley et al. (2024) uses a combination of geostationary and microwave imager data to find a strong diurnal cycle in the response of the domain mean LWP to variation in N_c . Decreases in LWP are observed during the day and neutral or positive responses of LWP during the night time hours. They speculated that this diurnal cycle in LWP sensitivity was driven primarily by the diurnal variation in precipitation sensitivity, however there is no way to confirm or refute the causation with observations. The discovery of this large diurnal cycle in the cloud adjustments presents yet another significant uncertainty in our current knowledge because the ERF_{ACI} is weighted by the diurnally varying incoming solar radiation.

Many Large-Eddy Simulation (LES) studies ~~use~~ employ idealized scenarios with constant forcings to extract key controls of the cloud system and simplify the interpretation of the results. This ~~has generally been the case for the many studies examining approach has been foundational in studies of~~ aerosol indirect effects ~~as well (e.g., Hoffmann et al., 2020), where aerosols modify cloud albedo and lifetime through changes in droplet number and precipitation processes (e.g., Moeng et al., 1996; Feingold et al.,~~ . While this approach has the advantage of simplicity, it neglects two important modes of variability in the subtropical cloudy boundary layer: (1) the large diurnal cycle, and (2) the multi-day transition of stratocumulus to cumulus boundary layers. A handful of studies have touched on these modes of variability in the context of aerosol indirect effects. For example, the study of Sandu et al. (2008) shows that increases in N_c increase the amplitude of the diurnal cycle of LWP_c in simulated stratocumulus. Furthermore, Sandu and Stevens (2011) show that transitions from Stratocumulus to Cumulus are a response to increasing ~~Sea Surface Temperatures~~ sea surface temperature (SST) through Lagrangian LES in the North East Pacific. However, Yamaguchi et al. (2017) find that aerosol number concentration influences the timing of the transition through its mediation of drizzle. ~~Prabhakaran et al. (2024) perform Lagrangian simulations of Stratocumulus perturbed by localized pulses of aerosol perturbations designed to simulate intentional~~ Prabhakaran et al. (2024) perform Lagrangian LESs of stratocumulus clouds transitioning to cumulus, perturbed by intermittent aerosol injections to simulate marine cloud brightening. They find ~~a distinct~~ that aerosol perturbations suppress precipitation and enhance cloud reflectivity, with greater sensitivity in pristine conditions due to precipitation-driven transverse circulations, and note diurnal variations in radiative forcing due to the solar cycle. Zhang et al. (2024) use LES with a conditional Monte-Carlo subsampling approach to study non-precipitating marine stratocumulus, finding a diurnal cycle in ~~the~~ cloud property sensitivity ~~with negative cloud adjustments occurring where aerosol-induced liquid water path adjustments are more negative at night due to enhanced entrainment, but less negative in the afternoon and positive adjustments in the early morning, which they attribute to differences in absorbed solar radiation in the cloud layer.~~ Erfani et al. (2022), buffered by shortwave absorption dependent on cloud liquid water path. Erfani et al. (2022)

perform Lagrangian LES ~~along a subtropical cloud transition and demonstrate that the LWP adjustment depends of a stratocumulus-to-cumulus transition in a subtropical marine environment, demonstrating that aerosol-induced LWP adjustments depend on the cloud state, where the cloud adjustments are more pronounced along a pristine precipitating trajectory regime.~~ In pristine conditions with active precipitation, aerosol perturbations suppress drizzle, leading to larger LWP increases in stratocumulus clouds compared to polluted conditions, where precipitation is already limited.

This study addresses the susceptibility of clouds and their properties to aerosol concentrations along their realistic multi-day Lagrangian transition from the subtropics to the tropics, with a focus on their diurnal variability. Furthermore, we decompose the susceptibility into three main components: the Twomey effect, LWP adjustment, and cloud fraction adjustment, showing that the Twomey effect is the primary factor controlling this susceptibility, while the other two contribute notably to the diurnal variability. The combined effects result in a significant susceptibility during the morning hours, with a diminishing net effect in the afternoon and evening. Our methodology is outlined in Section 2, the simulation results and their analysis are presented in Section 3, and Section 4 summarizes the study and presents the conclusions.

2 Methodology

2.1 Lagrangian Trajectory

The Lagrangian trajectory ~~selected for used in~~ this study was ~~chosen from the many produced in Smalley et al. (2024)~~ produced using the methodology outlined in ~~Smalley et al. (2022)~~ Smalley et al. (2022), and was then selected from the ensemble generated in Smalley et al. (2024). The trajectory is propagated forward in time using a 10 minute time step with the 3-hourly 925 hPa winds from the Modern-Era Retrospective analysis for Research and Applications, version 2 (~~MERRA-2 Gelaro et al., 2017~~) (MERRA-2; Gelaro et al., 2017). The selected trajectory west of Peru spans about ~~3800~~ 3,800 km and extends from the subtropics to the tropics over the Pacific Ocean (Fig. 1). It represents a classical example of the stratocumulus-to-cumulus transition (~~SCT~~) for an air mass propagating over the ocean upon increasing sea surface temperature (~~SST~~) and reduced large-scale subsidence. It starts at 20°S and 80°W on 2019-10-06 00:00:00 UTC (i.e., around 18:00 local time) and follows the mean ~~PBL~~ planetary boundary layer (PBL) flow during its six-day evolution. Note that the calculated trajectory provides only an approximate reconstruction of the real air mass movement due to both the presence of wind shear and the limited accuracy and resolution of reanalysis data.

Several observed cloud properties are matched to the trajectory where they are available. These include LWP from the fleet of passive microwave imagers (Wentz and Spencer, 1998). Higher frequency LWP observations are taken from the corrected geostationary data of Smalley and Lebsock (2023). Additional geostationary data products derived from the Advanced Baseline Imager (ABI) on GOES-16 include the cloud fraction, cloud top height, cloud optical depth, and cloud effective radius (Walther and Straka, 2019–2021). Finally, the profiles of several MERRA-2 variables are collocated along the trajectory to provide forcing data for the LES. These variables include horizontal wind components, water vapor, potential temperature, and large-scale subsidence, in addition to sea surface temperature.

2.2 Large-Eddy Simulations

125 We use the System for Atmospheric Modeling (Khairoutdinov and Randall, 2003) to simulate the transition. The domain size is $40.92 \times 40.92 \text{ km}^2$. The horizontal grid spacing is 40 m, while the vertical grid spacing is 8 m in the PBL, gradually increasing with altitude. The initial and boundary conditions are based on MERRA-2 reanalysis data interpolated to the trajectory points. However, adjustments to the initial atmospheric state were necessary to reproduce the thick stratocumulus layer observed on that day. The ~~most important change is sharpening original MERRA-2 profiles, due to their coarse vertical resolution~~
130 ~~and smoothed inversion structure, only support shallow convection when used in LES. To enable stratocumulus formation,~~ the inversion layer ~~thickness was reduced~~ to around 40m ~~instead of the original MERRA-2 resolution, which only produces~~
~~shallow convection m, providing a sharper capping inversion more consistent with stratocumulus-topped boundary layers~~
(Stevens et al., 2005; Berner et al., 2011).

The free-tropospheric temperature and moisture profiles are nudged with 1-hr timescale starting 500-m above the PBL
135 height defined as the top of inversion layer. Because the model cannot directly follow changes in the mesoscale pressure gradient that controls boundary-layer winds, we apply weak nudging of the mean PBL winds with a timescale of 12 hours. Furthermore, to suppress the development of spurious circulations within the domain during longer simulations, we apply weak ~~homogenization~~ horizontal homogenization of temperature and water vapor mixing ratio with a 48-hour timescale. Microphysics is parameterized using the ~~one-moment~~ scheme of Khairoutdinov and Kogan (2000), ~~and~~ Instead of prognosing
140 cloud droplet number concentrations, four different aerosol-related scenarios are prescribed along the trajectory in terms of ~~droplet number fixed time-dependent~~ concentrations (Fig. 1 e). All scenarios begin with high coastal droplet number concentrations typical of polluted continental air, gradually decreasing to $25 \text{ ee}^{-1} \text{ cm}^{-3}$ for pristine air, $50 \text{ ee}^{-1} \text{ cm}^{-3}$ and $100 \text{ ee}^{-1} \text{ cm}^{-3}$ for intermediate conditions, and $200 \text{ ee}^{-1} \text{ cm}^{-3}$ for polluted air. These scenarios represent ~~the uncertainty in~~
realistic variability in number concentrations and the associated aerosol-cloud interactions ~~and including~~ their impact on cloud
145 microphysics and radiative properties, ~~as well as observational uncertainties. The~~ The 25 cm^{-3} asymptotic case best agrees
with the satellite observations and should be considered the baseline simulation. The sea surface temperature changes from approximately 290 K to nearly 297 K, with surface fluxes interactively calculated based on local atmospheric conditions near the surface. Interactive short-wave and long-wave radiation effects are also included. A similar Lagrangian perspective and modeling setup was applied in many other studies (e.g., van der Dussen et al., 2013; Sandu and Stevens, 2011; Yamaguchi
150 et al., 2017). Note that while the boundary conditions follow observations, the PBL development is determined by the processes occurring within it.

Finally, we comment that in this case, the impact of changing subcloud atmospheric conditions on surface moisture supply
across different scenarios is relatively small, as latent surface heat fluxes increase by only several percent in the polluted
scenario compared to the pristine one, with 6-day averages of 137 and 145 W m^{-2} , respectively (see supplemental material).
155 All other simulation results, including sensitivity scenarios analyzed further, fall within that envelope determined by the pristine
and polluted scenarios.

2.3 Diurnal Controls of Indirect Radiative Effect

To understand the relative diurnal contributions of the Twomey effect and the cloud adjustments to the indirect effect, offline radiative transfer calculations are performed. The reflected shortwave flux is given by

$$160 \quad F^\uparrow = F_o \mu_o A, \quad (2)$$

where F_o is the solar constant, μ_o is the cosine of the solar zenith angle, and A is the all-sky albedo. The all-sky albedo is calculated as the sum of a clear and cloud sky components

$$A = (1 - f_c) \alpha_{surf} + f_c A_c, \quad (3)$$

where α_{surf} is the ocean surface albedo assumed to be 0.06, and A_c is the albedo of the cloudy part of the domain. We neglect clear sky absorption of the radiation. Accounting for multiple reflections between a cloud layer with albedo (α_{cld}) and the reflecting surface this is (Stephens, 1984) gives the combined albedo for the cloudy part of the domain as

$$A_c = \alpha_{cld} + \frac{\alpha_{surf}(1 - \alpha_{cld})^2}{1 - \alpha_{surf}\alpha_{cld}}. \quad (4)$$

Appendix A describes the offline calculations of α_{cld} , including a proper accounting of the solar zenith angle, which is a critical factor when addressing the diurnal cycle. Finally, the Cloud Radiative Effect (CRE) is calculated as

$$170 \quad CRE = F_o \mu_o f_c (A_{cld} - \alpha_{surf}) \quad (5)$$

The cloud optical depth is calculated at each time step from the domain mean time-dependent modeled LWP_c and N_c assuming an adiabatic cloud vertical structure following Hoffmann et al. (2023) (Brenguier et al., 2000) following the specific implementation of Hoffmann et al. (2023)

$$\tau_c = 0.2 N_c^{1/3} LWP_c^{5/6}. \quad (6)$$

175 The offline radiation calculations are used to decompose the ERF_{ACI} into the three indirect sensitivity terms defined in Eq. 1. Knowing that $F^\uparrow = F^\uparrow(N_c, LWP_c, f_c)$, the sensitivity can be estimated using the pristine and polluted simulation results as follows:

$$S_N = \frac{\partial F^\uparrow}{\partial \ln N_c} \approx \frac{F^\uparrow(N_{c200}, \overline{LWP}_c, \overline{f_c}) - F^\uparrow(N_{c25}, \overline{LWP}_c, \overline{f_c})}{\ln N_{c200} - \ln N_{c25}}, \quad (7)$$

$$S_{LWP} = \frac{\partial F^\uparrow}{\partial \ln LWP_c} \cdot \frac{d \ln LWP_c}{d \ln N_c} \approx \frac{F^\uparrow(\overline{N}_c, LWP_c(N_{c200}), \overline{f_c}) - F^\uparrow(\overline{N}_c, LWP_c(N_{c25}), \overline{f_c})}{\ln N_{c200} - \ln N_{c25}}, \quad (8)$$

$$180 \quad S_f = \frac{\partial F^\uparrow}{\partial \ln N_c} = \frac{\partial F^\uparrow}{\partial f_c} \cdot \frac{df_c}{d \ln N_c} \approx \frac{F^\uparrow(\overline{N}_c, \overline{LWP}_c, f_c(N_{c200})) - F^\uparrow(\overline{N}_c, \overline{LWP}_c, f_c(N_{c25}))}{\ln N_{c200} - \ln N_{c25}}. \quad (9)$$

Here, the overbar ~~represents~~ denotes the mean of the polluted and pristine ~~simulations~~ values along the trajectory, ~~meaning~~ as a function of time. This means that for each of the three terms, i.e., Eqs. 7-9, we estimate the sensitivity of ~~$F^\uparrow$~~ $F^\uparrow$ in only one direction within the three-dimensional parameter space, ~~while keeping~~ (N_c, LWP_c, f_c) , while holding the other two parameters fixed at their mean values. Note that ~~;~~ because these sensitivities are expressed in terms of reflected fluxes rather than albedo, they inherently account for the diurnal variation in incoming solar flux. ~~Furthermore, since we analyze the susceptibility of $\ln F^\uparrow$ rather than $F^\uparrow$ itself, the results are intended to be more universal across seasons, as they are scaled by solar insolation~~ radiation and therefore drop to zero at night.

2.4 Results

3 Results

185 3.0.1 Evaluation of LES evolution against observations

3.1 Evaluation of LES evolution against observations

We begin by evaluating the diurnal evolution of the simulated clouds and the realism of the LES against the observations. Figure 1 provides a summary of the evolution of the clouds for four N_c scenarios over the six-day simulation. Panel *a* shows the path of the trajectory while panel *f* shows the ~~Sea Surface Temperature (SST)~~ sea surface temperature along the trajectory. Panel *e* shows the imposed number concentrations, loosely based on observations from the ABI, which begins at large values of several hundred cm^{-3} near the coast and asymptotes to values ranging between 25 and 200 cm^{-3} in the tropics. Most of this paper will contrast the pristine (25 cm^{-3}) simulation with the polluted (200 cm^{-3}) simulation. Note that the pristine scenario best matches the observations and the polluted scenario should be interpreted as a perturbation from the observed state. Panel *h* shows the expected increases in τ_c with increases in N_c . Panels *d* (polluted) and *i* (pristine) highlight two critical features of the simulations. First, the polluted cloud grows significantly deeper than the pristine cloud and that growth occurs in the overnight and early morning hours. Second, the pristine cloud produces substantially more drizzle than the polluted clouds. Each of these observations is consistent with expectations that increasing N_c should both suppress precipitation and increase the cloud top entrainment efficiency. Next, note that the diurnal evolution of the LWP and f_c (panels *b*, *g*) show general agreement with the observations, while differing in some of the precise details. For example, the LES is not able to produce sufficiently thick and extensive cloud cover over the nighttime hours of days 2-4. We also note that the pristine experiment, which is the most realistic scenario, is well able to simulate the observed cloud top height (CTH), whereas the more polluted experiments show larger growth of the cloud layer (panel *c*), which is not observed in this case but remains a physically plausible outcome under different conditions.

3.1.1 Cloud Radiative Effects

3.2 Cloud Radiative Effects

How does the distinct diurnal variation in cloud properties affect the ERF_{ACI} ? Figure 2 contrasts the pristine and polluted scenarios to understand the relative influence of cloud adjustments relative to the Twomey effect on the CRE. The largest differences in LWP_c occur during the overnight and early morning hours due to the suppression of precipitation (panel a). In contrast, during mid-day, the polluted LWP_c is smaller than the pristine scenario. The f_c adjustment follows a similar diurnal pattern, with the polluted scenario showing a larger f_c overnight into the morning, and a smaller f_c in the afternoon (panel b). Panel c compares the cloud albedo of the pristine and polluted scenarios, including both the Twomey effect and the LWP adjustment. The polluted A_c is generally larger than the pristine, except for a few hours during midday when the reductions in LWP_c more than offset the Twomey effect brightening. What ultimately matters for the energy budget of the system is the CRE shown in panel **d**. Here, a distinct diurnal pattern emerges in the difference between the polluted and pristine scenarios. In the polluted scenario, there is a distinct increase in CRE in the morning, while in the early afternoon, there are modest decreases. Occasionally, a secondary increase in CRE occurs in the evening when the cloud layer is recovering from its afternoon minimum.

It is important to recognize that two factors limit the sensitivity of CRE to N_c at large solar zenith angle, *i.e.*, near sunrise and sunset. First, the incoming solar flux scales as μ_o and second as the cloud albedo approaches unity the Twomey effect tends to zero. As a result, the fairly large cloud adjustment terms in the early morning hours are not very effective at increasing the diurnal average CRE (see Figure 2d/e).

Figure 3 shows a composite diurnal cycle of the ERF_{ACI} averaged over the six-day trajectory. Here panel **a** shows the three individual terms that determine the ERF_{ACI} , calculated from Eqs. 7-9. The Twomey effect (S_N) is always positive with a peak in the late morning. The timing of this peak in S_N results from a combination of the fact that sensitivity is maximum for $A_c = 0.5$ (Platnick and Twomey, 1994) and of the fact that morning hours have larger f_c than afternoon hours, so that the Twomey effect has less leverage in the afternoon than in the morning. The other two cloud adjustment terms (S_{LWP} , S_f) have similar ~~patterns with positive /negative diurnal patterns, each having positive values in the morning /afternoon that mediate the and negative values in the afternoon that over the course of the diurnal cycle partially cancel the~~ Twomey effect. Panel **b** shows the total ERF_{ACI} , which is largely positive in the morning and approximately zero in the afternoon. Note that the sum of the three partial contributions from panel a, approximately calculated using Eqs. 7-9, agrees well with the total adjustment directly calculated from the LES as the difference in $F^\uparrow$ between the purely polluted and pristine cases:

$$S = \frac{dF^\uparrow}{d \ln N_c} = \frac{F^\uparrow(N_{c200}, LWP_{c200}, f_{c200}) - F^\uparrow(N_{c25}, LWP_{c25}, f_{c25})}{\ln N_{c200} - \ln N_{c25}}. \quad (10)$$

Overall, the cloud adjustments ~~are S_{LWP} and S_f average to~~ approximately zero over the diurnal ~~average, acting to enhance cycle, enhancing~~ the Twomey effect in the morning ~~and nearly cancel it but nearly canceling it out~~ in the afternoon. The daylight

240 ~~averages average values~~ of the three terms are $S_N = 0.0068$, $S_{LWP} = -0.0025$, $S_f = 0.0019$ $S_N = 17.0 \text{ W m}^{-2}$, $S_{LWP} = -6.0 \text{ W m}^{-2}$, and $S_f = 5.6 \text{ W m}^{-2}$.

3.2.1 ~~Role of Key Physical Processes/ Key Controls~~

245 Additionally, we calculated these terms for distinct pollution regimes using Eqs. 7–9, that is comparing N_{200} and N_{100} for the more polluted regime, and N_{25} and N_{50} for the more pristine regime, making use of the intermediate simulation results. The results are presented in Table 1. Notably, the S_N term is similar across background microphysical conditions, indicating the relative constancy of the Twomey effect. In contrast, the cloud adjustments are a net increase in albedo in the N_{50-25} experiments and a net decrease in the $N_{200-100}$ experiments. This sign inversion is reminiscent of the 'inverted-V' dependence of LWP on N_c seen in satellite data (Gryspeerd et al., 2022). However, in this specific case the inverted-V in LWP results from a modestly positive S_f term in the pristine conditions, where changes in droplet number concentration more strongly influence autoconversion and a much more strongly negative S_{LWP} term in the polluted conditions, where changes in droplet number concentration more strongly influence cloud top entrainment.

250 The third composite diurnal cycle shown in Fig. 3 (panel *c*) illustrates the evolution of dCRE, which follows a similar pattern to the total susceptibility of shortwave outgoing radiation shown in panel *b*. The strongest effect occurs in the late morning hours, reaching as much as $130\text{--}140 \text{ W m}^{-2}$, with slightly negative values in the early afternoon and approaching zero by the end of the day. Both the evolution of dCRE and the susceptibility of $F^\uparrow$ to aerosols exhibit strong day-to-day variability, with the Twomey effect dominating on day 1, and other adjustments becoming more prominent farther from the continent, where aerosol number concentrations decrease (Fig. 2 *e, f*).

3.3 ~~Role of Key Physical Processes/ Key Controls~~

Why does the diurnal pattern in ERF_{ACI} seen in Figure 3 emerge? To demonstrate the relevant mechanisms, Figure 4 presents a composite diurnal comparison of the cloudy boundary layer structure for the pristine and polluted scenarios. The polluted scenario produces less precipitation than the pristine scenario at all hours of the day (panel *aa*). The polluted cloud entrains more efficiently and grows deeper than the pristine cloud over night (panel *bb*). The polluted cloud is substantially more turbulent than the pristine cloud over night (panel *ee*). Panel *d* shows that while the changes in cloud LWP affect the radiative heating of the cloud layer, the afternoon differences in the shortwave warming are nearly exactly canceled by the differences in the longwave cooling. The resultant difference in radiative heating is primarily due to an overnight increases in longwave cooling of the polluted case. While both clouds are well coupled to the surface fluxes over night, the polluted cloud becomes substantially less coupled than the pristine cloud throughout the sunlit hours (Panel *e* panel *c*). The decoupling index is defined here as the ratio of the ~~cloud base flux to the near-surface flux (van der Dussen et al., 2013)~~ moisture flux at cloud base to that near the surface (van der Dussen et al., 2013), providing a measure of how much of the surface flux reaches the cloud layer.

270 Overall, a picture emerges of a polluted cloud that grows substantially faster over night than the pristine cloud with enhanced LWP due to precipitation suppression and a deeper cloud layer. However, this enhanced growth of the polluted cloud results in a deeper boundary layer that is more easily decoupled from the surface fluxes during the subsequent afternoon hours. These

results explain the consistently positive (early morning) and negative (afternoon) sensitivities in cloud fraction and liquid water path seen in Figures 2 and 3.

275 In our experimental design, modifying N_c influences three model processes directly: radiative transfer, autoconversion, and cloud water sedimentation. We perform a series of additional experiments where we impose the polluted N_c on a particular process rate while all other processes see the pristine N_c to demonstrate the importance of that process on the evolution of the boundary layer and cloud ~~macrophysical-properties-~~microphysical properties. A description of these experiments is provided in Table 2. We show the evolution of four quantities to demonstrate the influence of the various processes. The first two are
280 cloud ~~macrophysical-~~microphysical quantities: LWP and rain water path (RWP). The second two are related to the structure of the boundary layer: inversion height z_{inv} and decoupling index. Figure 5 shows the evolution of these four quantities for the various experiments. Several conclusions can be formed from these results:

- The influence of N_c on radiative transfer has a marginal effect on the evolution of the cloudy boundary layer. This is shown by the similarity between Exp. 3 and the pristine scenario (panels a-d).
- 285 – The autoconversion process has a positive and distinctly diurnal influence on the LWP sensitivity, while the cloud water sedimentation process has a smaller, negative, and relatively constant influence (panel e).
- Both autoconversion and, to a lesser extent, cloud water sedimentation affect the precipitation suppression mechanism (panel f). The latter process indirectly influences rainwater production by removing cloud liquid from the cloud top, thus limiting the efficiency of autoconversion.
- 290 – Both autoconversion and cloud water sedimentation influence the entrainment efficiency and growth of the boundary layer. Autoconversion has a larger effect than cloud water sedimentation, and the two processes interact in a super-linear manner to influence entrainment efficiency (panel g).
- Both autoconversion and cloud water sedimentation contribute to the decoupling of the cloud layer from the surface (panel h), which is consistent with the fact that both processes individually affect the cloud top entrainment rate.

295 A key summary of these conclusions is that the autoconversion and the cloud water sedimentation processes have some similar influences on the development of the cloudy boundary layer. The reason for this is that both processes remove liquid from the cloud top entrainment zone, thereby slowing the rate of precipitation production, decreasing the efficiency of the entrainment, and slowing the decoupling of the boundary layer. This is closely related to the dynamics of the Entrainment Interfacial Layer (EIL; Haman et al., 2007; Kurowski et al., 2009), where the removal of liquid from the cloud top influences
300 the structure of the EIL, leading to changes in boundary layer growth. We also see that these processes interact in a non-linear way, particularly in their influence on the LWP and the entrainment rate. Furthermore, the strong diurnal cycle in the sensitivity of cloud properties is a result primarily of the autoconversion process, whereas the cloud water sedimentation process operates over a longer time scale.

Finally, we comment on the relative role of removal of liquid water from the EIL and sub-cloud evaporative cooling on the
305 evolution of the cloud layer through evaluation of an additional sensitivity study in which the drizzle evaporation process is

turned off under pristine conditions (see supplemental material). In the no-evaporation experiment the surface buoyancy flux weakens due to reduced subcloud-layer cooling and a reduced ocean–atmosphere temperature contrast (Fig. S3). Despite this weaker surface forcing, cloud-layer turbulent kinetic energy tends to be higher at night (Fig. S6), suggesting that it is not directly controlled by surface buoyancy flux but is instead primarily driven by longwave radiative cooling (Wood, 2012). Nonetheless, the entrainment rate is also reduced (Fig. S5) due to lower moisture availability in the cloud layer (Fig. S4). A similar reduction in surface buoyancy flux is observed in the polluted case, although the entrainment rate increases significantly at night because of greater moisture availability compared to the pristine case. Stevens et al. (1998) found that drizzle evaporation stabilizes the subcloud layer and reduces entrainment via decoupling. Uchida et al. (2010) noted that drizzle evaporation below cloud base dampens buoyancy flux, weakens turbulence, and reduces entrainment. However, this dynamical argument is not fully supported by our results as we find that rain evaporation is associated with increased cloud-top entrainment rates. Furthermore, during the day, both cloud-layer turbulence and entrainment decrease more strongly with active evaporation than without evaporation, highlighting a pronounced diurnal modulation in the scenarios analyzed in this study.

~~This paper shows the results of~~

4 Conclusions

This paper analyzes the diurnal susceptibility of shallow subtropical clouds to aerosols using a six-day Lagrangian LES along the ~~stratocumulus-to-cumulus~~ stratocumulus-to-cumulus transition with realistic environmental forcing including a diurnal cycle of solar radiation. Pristine and polluted scenarios are simulated to quantify the ERF_{ACI} and its component terms. The ERF_{ACI} is broken down into the Twomey effect, a LWP_c adjustment, and a f_c adjustment. The daytime average values of the three terms are ~~0.0068, -0.0025, -0.0019~~ approximately 17 W m^{-2} , -6 W m^{-2} , and 6 W m^{-2} , respectively. However, there is a ~~substantial~~ substantial diurnal cycle in the three terms. The Twomey effect is always positive and most efficient in the morning hours because ~~the f_c~~ f_c is larger in the morning than ~~the afternoon~~. ~~More significantly in the afternoon, although it also has a significant positive contribution during the afternoon. More importantly,~~ the LWP_c and f_c adjustments switch signs from positive ~~to negative from morning into in the morning to negative in~~ the afternoon. The ~~resultant~~ resulting diurnal pattern of the ERF_{ACI} is super-Twomey in the morning and near neutral in the afternoon. Results further show evidence of a sign inversion (inverted-V) in the cloud adjustment terms with positive cloud adjustments in pristine conditions and negative cloud adjustments in polluted conditions.

The reason this diurnal pattern in ERF_{ACI} emerges is that the diurnal amplitude of the cloud ~~macrophysical properties~~ extent is increased relative to the pristine case. This occurs because precipitation is suppressed in the polluted cloud relative to the pristine resulting in a thicker more turbulent cloud layer, with enhanced longwave cooling and increased cloud liquid water near the cloud top entrainment zone during the nighttime hours. As a result of the increase cloud top liquid water, the polluted cloud entrains more efficiently and grows substantially faster and deeper overnight. However, this nighttime success of the polluted cloud is not sustainable as it results in a boundary layer that is deeper, drier and more decoupled, which ultimately leads to a stronger mid-day collapse of the cloudy boundary layer the following afternoon.

A key mechanism in the causal chain is the increase in cloud top liquid water with increases in N_c . Through sensitivity
 340 experiments it is shown that both sedimentation of cloud and rain water are effective at reducing the efficiency of the entrainment.
 However, cloud sedimentation and autoconversion interact in a nonlinear manner to result in a combined effect on entrainment
 that is greater than the sum of each term. This could occur due to the non-linearity of the autoconversion process interacting
 with a reduced amount of cloud liquid water at cloud top due to the cloud water sedimentation. Therefore, accurate simulation
 of the entrainment drying mechanism in global models should include both cloud and rain water sedimentation as is the case
 345 in at least one commonly used cloud microphysics parameterization (Morrison and Gettelman, 2008).

The findings of this study are in qualitative agreement with a growing body of literature based on both modeling and
 observations that increasing N_c causes an amplification of the diurnal cycle of cloud properties which subsequently causes
 a morning/afternoon contrast in the sign of the cloud adjustments with adjustments enhancing Twomey brightening in the
 morning and offsetting the brightening in the afternoon. The result in this study is a negligible diurnal average effect of the
 350 adjustments on the ERF_{ACI} . However, this is based on a single suite of simulations and we must be cautious in extrapolating
 these results to more general conditions. In particular, a key mechanism that mediates the diurnal response in these simulations
 is the suppression of precipitation. We have no expectations that increasing in N_c in non-precipitating clouds would have
 the same effect on the diurnal cycle. We could speculate that in that case the cloud adjustments would be robustly negative
 across the diurnal cycle. [Indeed our limited simulations here demonstrate that the cloud adjustments change sign from positive](#)
 355 [to negative as \$N_c\$ is increased.](#) Future research is necessary to extend the Lagrangian approach used here to many more
 trajectories representative of the diversity of atmospheric conditions to fully understand the influence of the diurnal cycle of
 the cloud adjustments on the ERF_{ACI} .

. The System for Atmospheric Modeling code is available upon contacting Dr. Marat Khairoutdinov.

. Trajectory data, model inputs and outputs needed to reproduce the figures are available at: <https://zenodo.org/records/14873449>

360 **Appendix A**

The cloud albedo is calculated using the hybrid model of Meador and Weaver (1980), which includes a dependence on the
 solar zenith angle μ_0

$$\alpha_{cld} = \frac{1}{1 + \gamma_1 \tau_c} \left(\gamma_1 \tau_c + (\beta_o - \gamma_1 \mu_o) \left(1 - \exp \left(\frac{-\tau_c}{\mu_o} \right) \right) \right) \sim \quad (A1)$$

The two γ coefficients of this model are given by

$$365 \quad \gamma_1 = \frac{7 - 3g^2 - \omega_o(4 + 3g) + \omega_o g^2(4\beta_o + 3g)}{4(1 - g^2(1 - \mu_o))} \quad (A2)$$

and

$$\gamma_2 = \frac{1-g^2-\omega_o(4+3g)-\omega_o g^2(4\beta_o+3g-4)}{-4(1-g^2(1-\mu_o))} \quad (A3)$$

where ω_o is the single scatter albedo and g is the asymmetry parameter. The third coefficient is given by

$$\beta_o = \frac{1}{2\omega_o} \int_0^1 P(\mu_o, -\mu') d\mu' \quad (A4)$$

370 which is the ~~the~~ fraction of single scattered radiation out of the solar beam into the backscattering hemisphere. The single scattering phase function (P) is subject to the normalization condition

$$\frac{1}{4\pi} \int_{-1}^1 \int_0^{2\pi} P(\mu, \phi; \mu', \phi') d\phi' d\mu' = \omega_o \quad (A5)$$

The inclusion of the β_o term is a complication as in general it represents an integral that can not be represented analytically. In this work, we parameterize this integral based on numerical integration of the Henyey and Greenstein (1941) phase function

375 for $g = 0.86$ and $\omega_o = 1$ giving the follow approximate formulation

$$\beta_o(g = 0.86, \omega_o = 1) \approx 0.5 \exp -2.7 \mu_o^{0.7} \quad (A6)$$

where the Henyey-Greenstein phase function subject to the proper normalization is given by

$$P_{HG} = \omega_o \frac{1-g^2}{(1+g^2-2g\cos(\Theta))^{\frac{3}{2}}} \quad (A7)$$

. MK and ML designed the experiments and MK carried them out. KS prepared the Lagrangian trajectory and observational data. MK
380 developed the model code and performed the simulations. MK and ML preformed the analysis. MK prepared the manuscript with contributions from all co-authors.

. At least one of the (co-)authors is a member of the editorial board of *Atmospheric Chemistry and Physics*.

. The research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (80NM0018D0004). Texas Advanced Computing Center (TACC) The University of Texas at Austin
385 is acknowledged for providing high-performance computing resources. Work from LLNL is performed under the auspices of the US DOE by LLNL under Contract DE-AC52-07NA27344; IM Number LLNL-ABS-867332. LLNL-JRNL-2002664.

References

- Ackerman, A. S., Toon, O. B., Stevens, D. E., Heymsfield, A. J., Ramanathan, V., and Welton, E. J.: Reduction of tropical cloudiness by soot, *Science*, 288, 1042–1047, <https://doi.org/10.1126/science.288.5468.1042>, 2000.
- 390 Ackerman, A. S., Kirkpatrick, M. P., Stevens, D. E., and Toon, O. B.: The impact of humidity above stratiform clouds on indirect aerosol climate forcing, *Nature*, 432, 1014–1017, <https://doi.org/10.1038/nature03174>, 2004.
- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, 1989.
- Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A.-L., Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle, F., Mauritsen, T.,
395 McCoy, D. T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding Global Aerosol Radiative Forcing of Climate Change, *Reviews of Geophysics*, 58, e2019RG000 660, <https://doi.org/https://doi.org/10.1029/2019RG000660>, e2019RG000660 10.1029/2019RG000660, 2020.
- Berner, A. H., Bretherton, C. S., and Wood, R.: Large-eddy simulation of mesoscale dynamics and entrainment around a pocket of open cells
400 observed in VOCALS-REx RF06, *Atmospheric Chemistry and Physics*, 11, 10 525–10 540, <https://doi.org/10.5194/acp-11-10525-2011>, 2011.
- Boucher, O. and Lohmann, U.: The sulfate-CCN-cloud albedo effect: A sensitivity study with two general circulation models, *Tellus B*, 47, 281–300, <https://doi.org/10.1034/j.1600-0889.47.issue3.1.x>, 1995.
- Boucher, O., Randall, D., Artaxo, P., Bessafi, R., Dufresne, J. L., Forster, P., Kiehl, J., Klein, S. A., Knutti, R., Köhler, M., Lohmann, U.,
405 Ramaswamy, V., Sawyer, V., Shine, K., Simpson, I., Stuber, N., Vogel, B., and Zhang, X.: Clouds and Aerosols, In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 571–657, <https://doi.org/10.1017/cbo9781107415324.016>, 2013.
- Brenguier, J.-L., Pawlowska, H., Schüller, L., Preusker, R., Fischer, J., and Fouquart, Y.: Radiative Properties of Boundary Layer Clouds: Droplet Effective Radius versus Number Concentration, *Journal of the Atmospheric Sciences*, 57, 803 – 821, [https://doi.org/10.1175/1520-0469\(2000\)057<0803:RPOBLC>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<0803:RPOBLC>2.0.CO;2), 2000.
410
- Bretherton, C. S., Blossey, P. N., and Uchida, J.: Cloud droplet sedimentation, entrainment efficiency, and subtropical stratocumulus albedo, *Geophysical Research Letters*, 34, 2007.
- Christensen, M. W. and Stephens, G. L.: Microphysical and macrophysical responses of marine stratocumulus polluted by underlying ships: 2. Impacts of haze on precipitating clouds, *Journal of Geophysical Research: Atmospheres*, 117, 2012.
- 415 Christensen, M. W., Neubauer, D., Poulsen, C. A., Thomas, G. E., McGarragh, G. R., Povey, A. C., Proud, S. R., and Grainger, R. G.: Unveiling aerosol–cloud interactions – Part 1: Cloud contamination in satellite products enhances the aerosol indirect forcing estimate, *Atmospheric Chemistry and Physics*, 17, 13 151–13 164, 2017.
- Coakley, J. A. and Walsh, C. D.: Limits to the Aerosol Indirect Radiative Effect Derived from Observations of Ship Tracks, *Journal of the Atmospheric Sciences*, 59, 668 – 680, 2002.
- 420 Diamond, M. S., Director, H. M., Eastman, R., Possner, A., and Wood, R.: Substantial Cloud Brightening From Shipping in Subtropical Low Clouds, *AGU Advances*, 1, e2019AV000 111, 2020.

- Erfani, E., Blossey, P., Wood, R., Mohrmann, J., Doherty, S. J., Wyant, M., and O, K.-T.: Simulating Aerosol Lifecycle Impacts on the Subtropical Stratocumulus-to-Cumulus Transition Using Large-Eddy Simulations, *Journal of Geophysical Research: Atmospheres*, 127, e2022JD037 258, 2022.
- 425 Fan, J., Wang, Y., Rosenfeld, D., and Liu, X.: Review of Aerosol–Cloud Interactions: Mechanisms, Significance, and Challenges, *Journal of the Atmospheric Sciences*, 73, 4221 – 4252, <https://doi.org/10.1175/JAS-D-16-0037.1>, 2016.
- Feingold, G., Stevens, B., Cotton, W. R., and Frisch, A. S.: The relationship between drop in-cloud residence time and drizzle production in numerically simulated stratocumulus clouds, *Journal of the Atmospheric Sciences*, 56, 1108–1122, [https://doi.org/10.1175/1520-0469\(1999\)056<1108:TRBDIC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<1108:TRBDIC>2.0.CO;2), 1999.
- 430 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419 – 5454, 2017.
- 435 Gryspeerdt, E., Quaas, J., and Bellouin, N.: Constraining the aerosol influence on cloud fraction, *Journal of Geophysical Research: Atmospheres*, 121, 3566–3583, 2016.
- Gryspeerdt, E., Glassmeier, F., Feingold, G., Hoffmann, F., and Murray-Watson, R. J.: Observing short-timescale cloud development to constrain aerosol–cloud interactions, *Atmospheric Chemistry and Physics*, 22, 11 727–11 738, <https://doi.org/10.5194/acp-22-11727-2022>, 2022.
- 440 Haman, K. E., Malinowski, S. P., Kurowski, M. J., Gerber, H., and Brenguier, J.-L.: Small scale mixing processes at the top of a marine stratocumulus—a case study, *Quarterly Journal of the Royal Meteorological Society*, 133, 213–226, 2007.
- Han, Q., Rossow, W. B., Zeng, J., and Welch, R.: Three Different Behaviors of Liquid Water Path of Water Clouds in Aerosol–Cloud Interactions, *Journal of the Atmospheric Sciences*, 59, 726 – 735, 2002.
- Heney, L. G. and Greenstein, J. L.: Diffuse radiation in the galaxy, *Astrophysical Journal*, 93, 70–83, <https://doi.org/10.1086/144246>, 1941.
- 445 Hoffmann, F., Glassmeier, F., Yamaguchi, T., and Feingold, G.: Liquid Water Path Steady States in Stratocumulus: Insights from Process-Level Emulation and Mixed-Layer Theory, *Journal of the Atmospheric Sciences*, 77, 2203 – 2215, 2020.
- Hoffmann, F., Glassmeier, F., Yamaguchi, T., and Feingold, G.: On the Roles of Precipitation and Entrainment in Stratocumulus Transitions between Mesoscale States, *Journal of the Atmospheric Sciences*, 80, 2791–2803, <https://doi.org/10.1175/JAS-D-22-0268.1>, 2023.
- IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK and New York, NY, USA, <https://doi.org/10.1017/9781009157896>, 2021.
- 450 IPCC: Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK and New York, NY, USA, <https://doi.org/10.1017/9781009157926>, 2022.
- 455 Jiang, H., Feingold, G., and Cotton, W. R.: Simulations of aerosol-cloud-dynamical feedbacks resulting from entrainment of aerosol into the marine boundary layer during the Atlantic Stratocumulus Transition Experiment, *Journal of Geophysical Research: Atmospheres*, 107, AAC 17–1–AAC 17–13, <https://doi.org/10.1029/2001JD001502>, 2002.
- Khairoutdinov, M. and Kogan, Y.: A New Cloud Physics Parameterization in a Large-Eddy Simulation Model of Marine Stratocumulus, *Monthly Weather Review*, 128, 229 – 243, 2000.

- 460 Khairoutdinov, M. F. and Randall, D. A.: Cloud Resolving Modeling of the ARM Summer 1997 IOP: Model Formulation, Results, Uncertainties, and Sensitivities, *Journal of the Atmospheric Sciences*, 60, 607 – 625, 2003.
- Kurowski, M. J., P. Malinowski, S., and W. Grabowski, W.: A numerical investigation of entrainment and transport within a stratocumulus-topped boundary layer, *Quarterly Journal of the Royal Meteorological Society*, 135, 77–92, 2009.
- Lebsock, M. D., Stephens, G. L., and Kummerow, C.: Multisensor satellite observations of aerosol effects on warm clouds, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- 465 Lohmann, U. and Feichter, J.: Global indirect aerosol effects: A review, *Atmospheric Chemistry and Physics*, 5, 715–737, <https://doi.org/10.5194/acp-5-715-2005>, 2005.
- Lu, M.-L. and Seinfeld, J. H.: Study of the aerosol indirect effect by large-eddy simulation of marine stratocumulus, *Journal of the Atmospheric Sciences*, 62, 3909–3923, <https://doi.org/10.1175/JAS3551.1>, 2005.
- 470 Manshausen, P., Watson-Parris, D., Christensen, M. W., Stier, P., Neubauer, D., and Grosvenor, D. P.: Invisible ship tracks show large cloud sensitivity to aerosol, *Nature*, 610, 101–106, <https://doi.org/10.1038/s41586-022-05122-0>, 2022.
- Meador, W. E. and Weaver, W. R.: Two-Stream Approximations to Radiative Transfer in Planetary Atmospheres: A Unified Description of Existing Methods and a New Improvement, *Journal of Atmospheric Sciences*, 37, 630 – 643, 1980.
- Moeng, C.-H., Cotton, W. R., Bretherton, C., Chlond, A., Khairoutdinov, M., Krueger, S., Lewellen, W. S., MacVean, M. K., Pasquier, J. R. M., Rand, H. A., Siems, S. T., Stevens, B., and Sykes, R. I.: Simulation of a stratocumulus-topped planetary boundary layer: Intercomparison among different numerical codes, *Bulletin of the American Meteorological Society*, 77, 261–278, [https://doi.org/10.1175/1520-0477\(1996\)077<0261:SOASTP>2.0.CO;2](https://doi.org/10.1175/1520-0477(1996)077<0261:SOASTP>2.0.CO;2), 1996.
- 475 Morrison, H. and Gettelman, A.: A New Two-Moment Bulk Stratiform Cloud Microphysics Scheme in the Community Atmosphere Model, Version 3 (CAM3). Part I: Description and Numerical Tests, *Journal of Climate*, 21, 3642 – 3659, 2008.
- 480 Pincus, R. and Baker, M.: Effect of precipitation on the albedo susceptibility of clouds in the marine boundary layer, *Nature*, 372, 250–252, <https://doi.org/10.1038/372250a0>, 1994.
- Platnick, S. and Twomey, S.: Determining the Susceptibility of Cloud Albedo to Changes in Droplet Concentration with the Advanced Very High Resolution Radiometer, *Journal of Applied Meteorology and Climatology*, 33, 334 – 347, 1994.
- Prabhakaran, P., Hoffmann, F., and Feingold, G.: Effects of intermittent aerosol forcing on the stratocumulus-to-cumulus transition, *Atmospheric Chemistry and Physics*, 24, 1919–1937, 2024.
- 485 Quaas, J., Arola, A., Cairns, B., Christensen, M., Deneke, H., Ekman, A. M. L., Feingold, G., Fridlind, A., Gryspeerdt, E., Hasekamp, O., Li, Z., Lipponen, A., Ma, P.-L., Mülmenstädt, J., Nenes, A., Penner, J. E., Rosenfeld, D., Schrödner, R., Sinclair, K., Sourdeval, O., Stier, P., Tesche, M., van Diedenhoven, B., and Wendisch, M.: Constraining the Twomey effect from satellite observations: issues and perspectives, *Atmospheric Chemistry and Physics*, 20, 15 079–15 099, 2020.
- 490 Sandu, I. and Stevens, B.: On the Factors Modulating the Stratocumulus to Cumulus Transitions, *Journal of the Atmospheric Sciences*, 68, 1865 – 1881, 2011.
- Sandu, I., Brenguier, J.-L., Geoffroy, O., Thouron, O., and Masson, V.: Aerosol Impacts on the Diurnal Cycle of Marine Stratocumulus, *Journal of the Atmospheric Sciences*, 65, 2705 – 2718, 2008.
- Smalley, K. M. and Lebsock, M. D.: Corrections for Geostationary Cloud Liquid Water Path Using Microwave Imagery, *Journal of Atmospheric and Oceanic Technology*, 40, 1049 – 1061, 2023.
- 495 Smalley, K. M., Lebsock, M. D., Eastman, R., Smalley, M., and Witte, M. K.: A Lagrangian analysis of pockets of open cells over the southeastern Pacific, *Atmospheric Chemistry and Physics*, 22, 8197–8219, 2022.

- Smalley, K. M., Lebsock, M. D., and Eastman, R.: Diurnal Patterns in the Observed Cloud Liquid Water Path Response to Droplet Number Perturbations, *Geophysical Research Letters*, 51, e2023GL107323, 2024.
- 500 Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., eds.: *Climate Change 2007: The Physical Science Basis*, Cambridge University Press, Cambridge, UK and New York, NY, USA, contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, 2007.
- Stephens, G. L.: The Parametrization of Radiation for Numerical Weather Prediction and Climate Models, *Monthly Weather Review*, 112, 826–867, [https://doi.org/10.1175/1520-0493\(1984\)112<0826:TPORFN>2.0.CO;2](https://doi.org/10.1175/1520-0493(1984)112<0826:TPORFN>2.0.CO;2), 1984.
- 505 Stevens, B., Cotton, W. R., Feingold, G., and Moeng, C.-H.: Large-Eddy Simulations of Strongly Precipitating, Shallow, Stratocumulus-Topped Boundary Layers, *Journal of the Atmospheric Sciences*, 55, 3616 – 3638, [https://doi.org/10.1175/1520-0469\(1998\)055<3616:LESOSP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1998)055<3616:LESOSP>2.0.CO;2), 1998.
- Stevens, B., Moeng, C.-H., Ackerman, A. S., Bretherton, C. S., Chlond, A., de Roode, S., Edwards, J., Golaz, J.-C., Jiang, H., Khairoutdinov, M., Kirkpatrick, M. P., Lewellen, D. C., Lock, A., Müller, F., Stevens, D. E., Whelan, E., and Zhu, P.: Evaluation
- 510 of Large-Eddy Simulations via Observations of Nocturnal Marine Stratocumulus, *Monthly Weather Review*, 133, 1443–1462, <https://doi.org/10.1175/MWR2930.1>, 2005.
- Toll, V., Christensen, M., Quaas, J., and Bellouin, N.: Weak average liquid-cloud-water response to anthropogenic aerosols, *Nature*, 572, 51–55, <https://doi.org/10.1038/s41586-019-1423-9>, 2019.
- Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of Atmospheric Sciences*, 34, 1149 – 1152, 1977.
- 515 Uchida, J., Bretherton, C. S., and Blossey, P. N.: The sensitivity of stratocumulus-capped mixed layers to cloud droplet concentration: do LES and mixed-layer models agree?, *Atmospheric Chemistry and Physics*, 10, 4097–4109, <https://doi.org/10.5194/acp-10-4097-2010>, 2010.
- van der Dussen, J. J., de Roode, S. R., Ackerman, A. S., Blossey, P. N., Bretherton, C. S., Kurowski, M. J., Lock, A. P., Neggers, R. A. J., Sandu, I., and Siebesma, A. P.: The GASS/EUCLIPSE model intercomparison of the stratocumulus transition as observed during ASTEX: LES results, *Journal of Advances in Modeling Earth Systems*, 5, 483–499, 2013.
- 520 Várnai, T. and Marshak, A.: MODIS observations of enhanced clear sky reflectance near clouds, *Geophysical Research Letters*, 36, 2009.
- Wall, C. J., Norris, J. R., Possner, A., McCoy, D. T., McCoy, I. L., and Lutsko, N. J.: Assessing effective radiative forcing from aerosol–cloud interactions over the global ocean, *Proceedings of the National Academy of Sciences*, 119, e2210481 119, 2022.
- Wall, C. J., Storelvmo, T., and Possner, A.: Global observations of aerosol indirect effects from marine liquid clouds, *Atmospheric Chemistry and Physics*, 23, 13 125–13 141, 2023.
- 525 Walther, A. and Straka, W.: Cloud optical depth from the National Oceanic and Atmospheric Administration (NOAA) comprehensive large array-data stewardship system (CLASS) [Dataset], retrieved from https://www.avl.class.noaa.gov/saa/products/search?sub_id=0&datatype_family=GRABIPRD&submit.x=20&submit.y=10, 2019–2021.
- Wentz, F. J. and Spencer, R. W.: SSM/I Rain Retrievals within a Unified All-Weather Ocean Algorithm, *Journal of the Atmospheric Sciences*, 55, 1613 – 1627, 1998.
- 530 Wood, R.: Stratocumulus clouds, *Monthly Weather Review*, 140, 2373–2423, <https://doi.org/10.1175/MWR-D-11-00121.1>, 2012.
- Yamaguchi, T., Feingold, G., and Kazil, J.: Stratocumulus to Cumulus Transition by Drizzle, *Journal of Advances in Modeling Earth Systems*, 9, 2333–2349, 2017.
- Zhang, J., Chen, Y.-S., Yamaguchi, T., and Feingold, G.: Cloud water adjustments to aerosol perturbations are buffered by solar heating in non-precipitating marine stratocumuli, *Atmospheric Chemistry and Physics*, 24, 10 425–10 440, [https://doi.org/10.5194/acp-24-10425-](https://doi.org/10.5194/acp-24-10425-2024)
- 535 2024, 2024.

Table 1. Composite daytime sensitivity of upward radiative flux to $\ln N_c$ under different aerosol regimes. Note that the full-range S_N is smaller than the pristine and polluted ones, which is due to a stronger stochastic variability between the N_{200} and N_{25} solutions.

	Pristine	Polluted	Full range
$\frac{dF^\uparrow}{d\ln N_c}$	(N_{50} - N_{25})	(N_{200} - N_{100})	(N_{200} - N_{25})
	(W m^{-2})	(W m^{-2})	(W m^{-2})
S_N	21.3	22.8	17.0
S_{LWP}	-0.1	-13.0	-6.0
S_f	11.3	3.2	5.6

Table 2. Summary of sensitivity experiments with varied microphysical processes. N25 refers to droplet number concentration for the pristine case, whereas N200 for the polluted case. Note that all the modifications in the Ex3-Ex5 experiments relate to the baseline pristine case.

Experiment	Radiation	Autoconversion	Sedimentation	Description
Ex1	N200	N200	N200	Polluted case
Ex2	N25	N25	N25	Pristine case
Ex3	N200	N25	N25	Impact of pollution on radiation
Ex4	N25	N200	N25	Impact of pollution on autoconversion
Ex5	N25	N25	N200	Impact of pollution on sedimentation




Figure 1. Overview of the analyzed case: (a) Lagrangian trajectory, and the evolution of (b) liquid water path from LES and observations, (c) cloud top height from LES and observations, (d) curtain plot of cloud water and rain water mixing ratios for the polluted case (N200), (e) observed and prescribed in LES droplet number concentrations, (f) sea surface temperature (SST), (g) observed and simulated cloud fraction, (h) cloud optical thickness from the LESs, (i) curtain plot of cloud water and rain water mixing ratios for the pristine case (N25).




Figure 2. Six diurnal cycles of (a) cloud fraction-liquid water path (LWP_c) and its difference between the polluted and pristine cases, (b) cloud fraction and its difference between the polluted and pristine cases, (c) cloud albedo calculated following Meador-Meador and Weaver (1980) and Weaver (1980)-and its difference between the extreme-polluted and pristine cases, also compared to-Twomey-effect(d) incoming solar shortwave energy flux, and (e) cloud radiative effect along-with, and its difference between the Polluted-polluted and Pristine-pristine cases:- Twomey-effect plotted-in-, and (bf) is calculated as the difference-between-two-values-susceptibility of cloud-albedo calculated for the same cloud-fraction f_c using two-shortwave outgoing radiation to droplet number concentrations-concentration decomposed into the three parts: Twomey effect (i.e-Eq. 7), pristine-and-pollutedLWP adjustment (Eq. 8), and averaged-over-different-cloud fractionsfraction adjustment (Eq. 9).




Figure 3. Composite diurnal cycle of: (a) the sensitivity-susceptibility terms S_N , S_{LWP} , S_f from Eqs. 7-9, calculated offline using the differences between the N200 and N25 simulations, and (b) their sum (magenta) compared against the actual LES model output (black) from Eq. 10. Panel (c) shows the diurnal cycle of $dCRE$.




Figure 4. Composite diurnal eye-cycles for the Pristine-pristine (red) and Polluted-polluted (blue) cases-in-terms-of-scenarios, showing (a) LWP and RWP, (b) cloud-top entrainment rate, (c) PBL decoupling index, (d) cloud-layer shortwave (SW) and longwave (LW) radiative tendencies and their total-differences-difference between the two scenarios (magenta), and (e) cloud-layer turbulent kinetic energy (TKE_c). Text in blue highlights features specific to the Polluted-polluted case.




Figure 5. Results of sensitivity experiments for two **most**-extreme droplet number concentrations, **Polluted-polluted** ($200 \text{ } \mu\text{ecm}^{-3}$) and **Pristine-pristine** ($25 \text{ } \mu\text{ecm}^{-3}$), applied independently to three main model components: radiation (RAD), rain autoconversion (AUT), and cloud water **subsidence-sedimentation** (**SUBSED**), as explained in Tab. 2. The panels show time series of: (a) LWP and (e) its difference between three pairs of key experiments (**Ex1-Ex2**, **Ex1-Ex2**, **Ex4-Ex2**, **Ex4-Ex2**, **Ex5-Ex2**, **Ex5-Ex2**); (b) RWP and (f) its difference; (c) inversion height and (g) its difference; (d) PBL decoupling **index** and (h) its difference, **omitting night time when all (with nighttime values omitted for clarity since the PBLs are coupled in all cases)**. To reduce noise and extract the main signal, the LWP/RWP **and** decoupling index time series are smoothed using a 5-hour **(the former)** and an 8-hour **(the latter)** window, **respectively**.