



## The incubation history of soil samples strongly affects the occlusion of particulate organic matter

Frederick Büks<sup>1</sup>, Sabine Dumke<sup>1</sup>, Julia König<sup>1</sup>

<sup>1</sup>Chair of Soil Science, Dept. of Ecology, Technische Universität Berlin, 10587 Berlin, Germany

Correspondence to: Frederick Büks (frederick.bueks@tu-berlin.de)

**Abstract.** Soil structure is a key proxy for carbon and nutrient storage, stable pore space, erosion stability and rootability. It is often quantified based on the degree of aggregation or the mechanical stability of soil aggregates. This work compares two methods representing basic principles of aggregate measurement. Undisturbed soil samples of loamy sand, clayey silt and silty loam were analyzed by ultrasonication/density fractionation (USD) to quantify different soil organic carbon (SOC) pools and by wet-sieving to measure the amount of water stable aggregates (%WSA). The measurements were carried out on field-fresh soil as well as samples that were air-dried, reset to field capacity (pF 1.8) by capillary action and incubated for 0, 1 and 4 weeks. Our results show, that the strength of POM occlusion sharply decreases after rewetting and is not fully recovered within the following four weeks. In contrast, the amount of water stable aggregates (%WSA) remains largely stable except in clayey silt. In consequence, field-fresh measurements are highly recommended to avoid overestimation of free and weakly bound SOM fractions or erosion stability.



## 1 Introduction

It has been over 90 years since soil structure began to be in the focus of agricultural research (Russell, 1928; Christensen, 1930). First seen under the aspect of soil plowability, the 1960s brought attention on well aggregated soils as a support for root growth and against soil compaction by heavy machinery (Rosenberg, 1964). Today, good soil structure is seen as an eminent proxy of soil quality, as it does not only provide rootability and stable pore space, but but is also related to water holding capacity, drainage of excess rain or flood water, enhanced aeration as well as carbon and nutrient storage within aggregates (Bronick and Lal, 2005). The quantification of soil aggregates was early performed by use of dry and wet sieving (Yoder, 1936; Chepil and Bisal, 1943). From here, the methods branched out into approaches still used to describe the amount and size distribution of soil aggregation. One branch comprised the weight fraction of water-stable aggregates (%WSA) and, based on the same wet-sieving principle, the mean weight diameter (Bryant et al., 1948; Bavel, 1950; Angulo et al., 2024; Meidl et al., 2024). The other branch was aimed to measuring soil structure for its mechanical stability. Wet sieving methods were early applied to quantify the mechanical integrity of soils (aggregate stability) after a certain amount of stress provided by sieve movements (Russel and Feng, 1947). In the second half of the 20th century, ultrasonic dispersion introduced by Edwards and Bremner (1967a) increasingly replaced sieving methods for quantification of aggregate stability, since now the energy input to the soil could be estimated from the power output of the sonotrode (North, 1976). Nowadays, ultrasonication is a common tool to achieve a semi-quantitative view on soil structural stability by comparing the mass of aggregate size fractions or mean weight diameter of water stable aggregates (WSA) after the application of defined quanta of ultrasonication stress (e.g. Lehtinen et al., 2014; Jouquet et al., 2016; Cavael et al., 2020). Beyond that, ultrasonication is combined with density fractionation (USD) of particulate organic matter (POM), which is successively released with increasing energy input and used for the quantification of soil carbon pools (Edwards and Bremner, 1967b; Golchin et al., 1994; Kaiser and Berhe, 2014; Graf-Rosenfellner et al., 2016). However, carbon pool measurements that apply USD extraction are prone to a number of artifacts. A density cut-off at  $1.6 \text{ g cm}^{-3}$  is mandatory, because lower concentrated solutions do not completely separate the respective POM fraction, while higher densities cause co-extraction of the mineral matrix (Cerli et al., 2012). Ultrasound treatments with energy levels  $>50 \text{ J ml}^{-1}$  can cause comminution of POM, sorption to mineral surfaces and lead to a reduced recovery rate, as demonstrated by Büks et al. (2021). This causes a carry-over of POM from fractions with lower to those with higher binding strength and a false estimation of both. Furthermore, adding the dense solution to the soil sample results in a low recovery rate of the free POM fraction (fPOM) due to burying within the matrix (Büks, 2023). This problem is addressed by rinsing the sample into the solution or gentle rotation, which significantly increases the recovery of fPOM and reduces overestimation of occluded POM (oPOM) mass. And be there an air-dried sample, the abrupt addition of water or dense solutions to the soil causes rupture of WSA and release of oPOM by a process called slaking (Emerson, 1967; Bossuyt et al., 2001), which could in turn result in an overestimation of the fPOM fraction. In consequence, comparability of fractionation results is only given under standardized test conditions.

The present work focuses on another parameter, that can potentially cause carbon pool artifacts: The wetting history of the sample. Normally measuring under field-fresh conditions, the extent of a sampling campaign or samples from soil archives can make it necessary to



70 quantify structural characteristics of already air-dried soil. Greacen (1960) showed, that  
aggregate stability is reduced with increasing water content. This is caused by the dissolution  
of various inorganic (and organic) cementation agents within a moist soil matrix (Amézketa,  
1999). On the other hand, a second factor of aggregate stability, that is not only influenced by  
75 the actual water content, but also its near history, is the soil microbiome and its biofilm matrix.  
The composition and activity of soil bacterial and fungal communities adapt to the soil water  
content, and bacterial rather than fungal abundance is significantly reduced under dry  
conditions (Drenovsky et al., 2004; Chowdhury et al., 2011). Since biofilms consist of 90-99 wt  
% water, they reshape during the drying process suggesting altered mechanical capabilities.  
80 The air-drying and long-term storage of soil samples therefore may alter the biofilm matrix  
and fungal hyphae networks, that have been shown to raise the quantity and stability of soil  
aggregates (e.g. Büks and Kaupenjohann, 2016; Bossuyt et al., 2001; Tang et al., 2011).  
This, however, implies a decline of soil structural stability due to air-drying and long-term  
storage, when samples are rewet, compared to field-fresh measurements. Past studies on  
soil structural stability, POM occlusion and the amount of WSA have been conducted with  
85 either field-fresh, air-dried and re-moistened soil samples (e.g. Oztas and Fayetorbay, 2003;  
Annabi et al., 2007; Büks and Kaupenjohann, 2016). The aim of this work is to elucidate  
effects of this different practice on the comparability of the resulting data. We hypothesize that  
air-drying of soil samples causes the weakening of soil structure expressed by the decline of  
parameters such as percentage by weight of WSA and the POM occlusive strength, and that  
90 this effect can be attenuated by long enough re-incubation under conditions of field capacity.



## 2 Material and methods

### 2.1 Sites and sample preparation

Undisturbed soil samples were taken at three different organic farming sites within a homogeneous area of each 1 m<sup>2</sup> (Table 1). The organic litter and top 10 cm were removed, and sampling was carried out in 10-20 cm depth of the mineral topsoil by use of metal rings (Ø=5.6 cm, h=4.0 cm, V=100 ml, n=25 per site). The rings containing the soil core were capped and tightly packed in closed plastic bags to preserve soil humidity for transpiration. Additionally, SOC concentrations of the sites were determined by each five mixed samples (each by three drillings) of the upper 30 cm topsoil along the bed.

**Tab. 1:** Field characteristics of the three sampling sites. For additional information, see Supplements.

| Texture           | SOC<br>(g kg <sup>-1</sup> dry mass) | pH<br>(-) | Region                | Coordinates               |
|-------------------|--------------------------------------|-----------|-----------------------|---------------------------|
| Loamy sand (Sl2)  | 32.2 ± 1.3                           | 5.9 ± 0.1 | Brandenburg/Germany   | 52°46'59.7"N 13°11'55.7"E |
| Clayey silt (Ut3) | 19.7 ± 2.1                           | 7.1 ± 0.1 | Saxony-Anhalt/Germany | 51°10'38.9"N 11°57'27.6"E |
| Silty loam (Lu)   | 18.4 ± 2.2                           | 6.9 ± 0.1 | Thuringia/Germany     | 50°53'14.2"N 11°12'42.7"E |

### 2.2 Setting to field capacity

Back in the laboratory, ten of the cores of each soil were set on a porous plate, saturated via capillary action for 24 hours and then drained by a hydrostatic head of pF 1.8 until constant weight was achieved. Half of these cores were weighed, dried for 24 hours at 105°C, weighed again, and the field capacity was determined from the difference minus the ring weight. The other five were directly measured (field-fresh treatment) for soil organic carbon (SOC) fractions and percentage by weight of water stable aggregates (%WSA). The remaining 3x5 cores were air-dried at 35°C with strong air circulation until constant weight was reached and then set to pF 1.8 as described above. Five of these were directly measured for occlusion of POM carbon and %WSA (W0), the others were incubated at constant water content and 20°C in the dark for 1 and 4 weeks and then treated similarly (Fig. 1).

### 2.3 fPOM, oPOM<sub>50</sub> and residual SOC

For analysis of SOC fractions (Fig. 1), small metal rings (Ø=2.4 cm, h=4.0 cm, V=18.1 ml) were used to subsample approx. 20 g dry soil equivalent from each of the field-fresh and re-incubated soil cores (n=5). The samples were gently removed from the ring by use of a spatula, weighted and rinsed into 200 ml Polyethylene (PE) flasks with 100 ml of sodium polytungstate solution (SPT) following Büks (2023). As the soils differ in their field capacity, the density of the SPT solution was adapted to match 1.6 g cm<sup>-3</sup> after addition to the sample by following Eq. 1

$$\frac{1.6 \text{ g cm}^{-3}}{100 \text{ ml}} = \frac{\rho_{\text{SPT}}}{100 \text{ ml} + m \cdot V} \quad (\text{Eq. 1})$$



with  $m$  the dry mass of the soil sample (g),  $V$  the volume of soil solution per g soil ( $\text{ml g}^{-1}$ ) and  $\rho_{\text{SPT}}$  the required density of added SPT solution. Subsequently, the sample was stored at room temperature for 30 min to allow infiltration of SPT solution into the pore space.

130 The samples were then centrifuged for 26 min at 3,500 g. The floating fPOM was separated by use of a water-jet pump, filtered through a  $0.45 \mu\text{m}$  cellulose acetate filter and cleaned with deionized water until the electrical conductivity of the filtrate dropped below  $50 \mu\text{S cm}^{-1}$ . The samples were flushed with deionized water through a 2 mm sieve to remove coarse material and into aluminum bottles, freeze-dried at  $-20^\circ\text{C}$ , lyophilized, dried for 24 h at  $105^\circ\text{C}$  and stored

135 in an exsiccator with a desiccant battery.

After separation of the fPOM, the PE flasks were refilled to their initial weight with  $1.6 \text{ g cm}^{-1}$  dense SPT solution. Cavitation stress was applied to the samples by use of a sonotrode (Branson® Sonifier 250, sonotrode diameter of 13 mm, frequency of 40 kHz, immersion depth of 15 mm, power output of  $50.02 \pm 1.29 \text{ J s}^{-1}$ ) as described by Büks and Kaupenjohann (2016).  
140 Sonication time corresponding to the applied energy density of  $q=50 \text{ J ml}^{-1}$  was determined based on the energy output of the sonotrode calculated following North (1976). The subsequent extraction and preparation of the respective fraction (oPOM<sub>50</sub>) was conducted similar to the fPOM.

145 The residuum (R), containing the oPOM releasable with  $q>50 \text{ J ml}^{-1}$  and the mineral-associated organic matter (MAOM), was washed approx. 6 times with 100 ml deionized water until the solution dropped below  $50 \mu\text{S cm}^{-1}$  or showed no further decrease of conductivity by washing. Then, these samples were treated similar to the fPOM and oPOM<sub>50</sub> fractions.

All fPOM, oPOM<sub>50</sub> and R samples were weighted, ground and the SOC concentrations were determined by use of an Elementar UNICUBE® CNS analyzer.

150

## 2.4 Water stable aggregates

The %WSA were quantified in each soil and treatment by use of a wet sieving apparatus (Eijkkelkamp) following DIN 19683-16 and the manual instructions (Fig. 1, Eq. 2). Similar to the above preparation, undisturbed subsamples of about 4 g dry weight equivalent were taken  
155 from the five cores of each treatment by use of a smaller metal ring ( $\varnothing=1.6 \text{ cm}$ ,  $h=1.7 \text{ cm}$ ,  $V=3.4 \text{ ml}$ ). The soil was carefully transferred from the ring to a  $250 \mu\text{m}$  sieve as above and the sieve was placed into a beaker with 90 ml of deionized water.

The samples were sieved for 3 min with a frequency of  $34 \text{ min}^{-1}$ . The sievings remained within the first beaker, while the sieve was placed into a new one filled with 70 ml of deionized water.

160 The submerged sample was treated with ultrasound by use of a sonotrode ( $50.02 \pm 1.29 \text{ J s}^{-1}$ ) for 2 min ( $\sim 60 \text{ J ml}^{-1}$ ) to fully disaggregate all macroaggregates. Then, 20 ml of deionized water were added and the samples were sieved again as described above. The matter remaining within the sieve (R) was wet-sieved by use of a 2 mm mesh to remove coarse material. The sievings of the first ( $m_1$ ) and second step ( $m_2$ ) as well as R were dried for 24 h  
165 at  $105^\circ\text{C}$  and set to room temperature in an exsiccator with a desiccant battery.

The mass fraction  $m_1$  represents primary particles and microaggregates with diameters  $<250 \mu\text{m}$ , that were not part of water-stable macroaggregates, whereas  $m_2$  represents all objects  $<250 \mu\text{m}$  that were released through the ultrasonication of water-stable macroaggregates. The fraction R is the sum of objects  $>250 \mu\text{m}$ , that did not pass the mesh  
170 during both steps of separation. The fraction



$$x_m = \frac{m_2}{m_1 + m_2 + R} \quad (\text{Eq. 2})$$


allows for semi-quantitative measurement of water stable macroaggregates. Particles >2 mm  
175 are excluded from the assessment, as samples can differ greatly in their coarse fraction due  
to small stones and litter.

## 2.5 Statistics

180 A pre-search with 1.5 interquartile range method (1.5xIQR) was applied to identify potential  
outliers within the SOC and %WSA data sets and the regular data of each treatment were  
tested for normal distribution by use of the Shapiro-Wilk test ( $\alpha=0.05$ ). Since all WSA and  
SOC data were normally distributed, Grubbs' test for outliers was applied ( $\alpha=0.05$ ) despite of  
the low number of replications ( $n=5$ ). A few outliers were identified across the whole data,  
185 that, however, do not affect the significance of the data (see supplements). Data of %WSA  
and SOC fractions were also positively tested for homogeneity of variance (Levene's test,  
 $p>0.05$ ). For each soil separately, the data were compared by Student's t-test between the  
field-fresh and each of the incubation treatments. In addition, linear regression analyses of the  
%WSA with the SOC fraction data were carried out.



190



**Fig. 1:** Process chart of SOC pool (left) and water-stable macroaggregate measurement (right). Abbreviations refer to free (fPOM) and occluded particulate organic matter (oPOM), non-aggregated particles <250 µm (m<sub>1</sub>), particles <250 µm of water-stable macroaggregates (m<sub>2</sub>) and the residua of both methods (R).






## 3 Results

### 3.1 Shifts of SOC fractions

195 Directly after being rewet (0 weeks), the soils consistently show increased amounts of the fPOM-C and oPOM<sub>50</sub>-C fractions as well as corresponding decreases of the residual fractions, compared with the field-fresh treatments (Fig. 2). This shift is significant ( $p < 0.05$ ) for all samples except the fPOM of clayey silk ( $p = 0.22$ ) and the oPOM<sub>50</sub> of loamy sand ( $p = 0.06$ ). The loamy sand thereby shows the largest increase of fPOM (+3.3 wt%) directly after being rewet, which mainly corresponds to the decrease of the residual fraction. The clayey silt and the silty loam, in contrast, have a rather small release of additional fPOM, and the decrease of the residual fraction corresponds with the oPOM<sub>50</sub> fraction (+5.8 wt% and +14.8 wt%, respectively).



**Fig. 2:** Development of the fPOM-C, oPOM<sub>50</sub>-C and residual C fraction of a loamy sand, a clayey silt and a silty loam after drying and re-incubation for 0, 1 and 4 weeks. The gray bar refers to the respective values of the field-fresh samples (dotted=mean value, outer lines=standard deviation). Large dots are mean values, small dots single measurements. Black dots refer to treatments with values significantly different from the field-fresh samples, gray marks similarity ( $\alpha = 0.05$ ). The logarithmic fits illustrate the trend towards the field-fresh values with full lines within and dotted lines beyond the measured span.





All samples, furthermore, show a consistent trend towards the initial values of the field-fresh samples. The loamy sand reaches non-significant differences to field-fresh values in the fPOM and oPOM<sub>50</sub> fraction within 1 week (t-test  $p \geq 0.05$ ). The clayey silt takes 4 weeks in all SOC fractions, while the silty loam does not restore in any fraction.

### 3.2 Alteration of water stability and correlation with SOC fractions

Of the three soil types, only the clayey silt shows significant ( $p < 0.05$ ) increases of %WSA directly after re-incubation and returns to non-significant differences compared to the field-fresh values within 1 week (Table 1). The peak of aggregation corresponds to the decrease of particle mass, that is not bound into water stable aggregates. The loamy sand have a tendency for increased %WSA directly after being rewet. All soils thereby show higher variation compared to the above SOC measurements (1.4 wt%), with mean standard deviations of 4.2 wt% for the heavy soils and 9.1 wt% for the sand.

The residual fractions of the respective soils have nearly constant mean values across all treatments differing by  $< 1.6$  wt% (see supplements). Deviations within the residuum and a significant increase of the clayey silt residual fraction after 1 week can be better explained by a different composition of the 250–2000  $\mu\text{m}$  mineral and organic matter fraction than by incomplete dispersion of macroaggregates.

**Tab. 2:** Development of the amount of water stable macroaggregates ( $m_2$ ,  $< 250 \mu\text{m}$ ), unbound particles ( $m_1$ ,  $< 250 \mu\text{m}$ ) and residual particles (R, 250–2000  $\mu\text{m}$ ) of loamy sand, clayey silt and silty loam in field-fresh state as well as after drying and re-incubation for 0, 1 and 4 weeks. The \* refers to significant differences compared to the field-fresh samples (Student's t-test,  $p < 0.05$ ).

|             |       | field-fresh     | 0 weeks          | 1 week          | 4 weeks         |
|-------------|-------|-----------------|------------------|-----------------|-----------------|
| Loamy sand  | $m_1$ | $34.0 \pm 12.7$ | $27.4 \pm 5.6$   | $38.4 \pm 8.6$  | $40.5 \pm 10.4$ |
|             | $m_2$ | $51.8 \pm 14.1$ | $57.7 \pm 4.8$   | $48.7 \pm 8.6$  | $46.9 \pm 11.6$ |
|             | R     | $14.2 \pm 1.6$  | $15.0 \pm 3.8$   | $12.9 \pm 1.7$  | $12.6 \pm 3.8$  |
| Clayey silt | $m_1$ | $16.8 \pm 2.5$  | $11.7 \pm 4.3$   | $13.8 \pm 4.7$  | $14.3 \pm 6.3$  |
|             | $m_2$ | $78.9 \pm 2.8$  | $84.8 \pm 4.3^*$ | $82.7 \pm 4.7$  | $81.7 \pm 6.2$  |
|             | R     | $4.3 \pm 0.8$   | $3.6 \pm 0.2$    | $3.5 \pm 0.3^*$ | $4.0 \pm 1.0$   |
| Silty Loam  | $m_1$ | $6.6 \pm 2.9$   | $7.6 \pm 4.0$    | $7.8 \pm 2.5$   | $4.1 \pm 1.1$   |
|             | $m_2$ | $86.3 \pm 2.9$  | $85.3 \pm 4.0$   | $85.6 \pm 3.4$  | $89.3 \pm 1.3$  |
|             | R     | $7.1 \pm 1.2$   | $7.1 \pm 0.7$    | $6.6 \pm 1.0$   | $6.6 \pm 0.7$   |

The regression of %WSA ( $m_2$ ) with oPOM<sub>50</sub>-C plus residual C (the amount of SOC, that should be occluded within water stable aggregates) shows no correlation, neither for loamy sand ( $R^2=0.16$ ) nor for clayey silt ( $R^2=0.10$ ) and silty loam ( $R^2=0.11$ ). Regression of  $m_1$  with the fPOM-C and %WSA with only the oPOM<sub>50</sub>-C or only the residual C lead to similar results ( $R^2 < 0.21$ ,  $R^2 < 0.04$  and  $R^2 < 0.16$ , respectively).



#### 4 Discussion

Our results show, that POM occlusion in sandy, silty and loamy soils is substantially decreased by air-drying and subsequent rewetting. The effect diminishes over time of re-incubation, but did not restore field-fresh values of all SOC fractions within a span of four weeks. Longer periods of incubation were avoided in this experiment, since they increase the risk, that the isolated soil ecosystem within the sampling ring is overgrown by fungal hyphae most probably affecting aggregate characteristics (Tisdall et al., 1997; Fan et al., 2022).

The decreased strength of POM occlusion is indicated by an increased mass of the fPOM-C and weakly bound oPOM<sub>50</sub>-C fraction as well as the corresponding decrease of the residual C fraction, that contains the more strongly bound oPOM and the MAOM. This implies an initial section by section shift from the residual C to the oPOM<sub>50</sub>-C and further to the fPOM-C fraction as a result of air-drying and rewetting. The sandy loam, which has generally lower aggregate stability due to its low clay content, thereby shows a stronger increase of the fPOM-C fraction, compared to the heavier soils. This can be interpreted as partial destruction of soil aggregates causing release of POM. In contrast, clayey silt and silty loam have strong shifts from the residual to the oPOM<sub>50</sub>-C fraction, which accounts for the majority of destabilized POM and indicates a weakening of soil aggregate structure without release of POM. Having decreasing mechanical stability of soil structure in common, all three soils show different pattern of destabilization with mainly destruction (loamy sand), mainly weakening (clayey silt) and both destruction and weakening. In field studies, the observed weakening of soil structure after re-incubation will lead to an overestimation of fPOM and the loosely occluded POM fraction, while the respective residuum is underestimated. This causes overestimation of the labile C pool and underestimation of the binding forces within aggregates. This underpins the need for USD measurements with field-fresh samples.

In contrast to POM occlusion, the percentage by weight of water stable aggregates is less affected by air-drying and gentle rewetting. Only the clayey silt (and in tendency the loamy sand) shows a significant increase of water stable aggregates directly after re-incubation with a subsequent restoration of field-fresh values, while the other treatments have largely constant state. This is in contrast to different works that show a loss of macroaggregates after mainly fast rewetting events (e.g. Denef et al., 2001; Bossuyt et al., 2001; Almajmaie et al., 2017). Haynes and Swift (1990) compared field-fresh and air-dried soil samples after wet-sieving and found the mean weight diameter of macroaggregates in grassland soils to be increased, while arable soils showed decreased values compared to field-fresh samples.

The observed lack of correlation with the SOC measurements seems contradictory at first, as it could be assumed that an unchanged amount of WSA implies unchanged oPOM fractions and, thus, *decreasing* the amount of water-stable macroaggregates causes an increase of fPOM-C at the expense of the oPOM fraction. However, both methods use incongruent parameters, as shown by Almajmaie et al. (2017), who stated poor correlation between different methods of soil structure measurement. The WSA approach represents the ratio of soil particles <250 µm of water-stable macroaggregates to the total sample mass. If measurements are conducted with samples set to pF 1.8, shear forces of wet sieving are the prominent agent of dispersion. In contrast, the USD measurement provides the amount of fPOM, that is fractionated by trickling through SPT solution containing approx. 1.2 M Na<sup>+</sup> as chemical dispersion agent, and oPOM fraction that is released from aggregates under certain levels of cavitation stress in the same Na<sup>+</sup> rich environment. With both methods addressing different binding mechanisms and using different forces of disaggregation, results may, arguably, diverge.



280 As an example, drying could weaken but not fully destroy the structural integrity of existing  
soil aggregates by dehydration of biofilms and fungal hyphae, that work as important  
aggregation agents in soils. This may cause an increased release of weakly bound oPOM  
after the USD treatment, while the amount of non-aggregated material passing the mesh in  
the WSA measurement remains constant as shown for silty loam. At the same time, as a  
285 result of a dehydration treatment, the increase of %WSA may occur through the formation of  
slightly soluble precipitates, that influence soil structural stability directly after rewetting. This  
points out, that also WSA measurements should be carried out with field-fresh samples to  
avoid e.g. overestimation of erosion stability of soils (Barthès and Roose, 2002). If changes of  
structural integrity are associated with altering the size of water stable macroaggregates, they  
can be addressed by measuring the mean weight diameter (MWD) as extension of the %WSA  
290 method (Kemper and Rosenau, 1986).  
Both mechanisms, cementation and partial damage of biofilms, work in opposite directions  
and would occur to different degrees depending on e.g. soil texture or microbial community.  
Also magnitude and duration of the respective effects after rewetting are still not estimated  
and require further studies, which should also include shrinkage and swelling of clay-rich soils  
295 or the span of storage under air-dried conditions.  
In consequence, studies on soil structure should generally use field-fresh samples. If there is  
no option except storing the soil samples, defined re-incubation should be carefully applied for  
≥1 week. Furthermore, Kühnel et al. (2019) showed that similar masses of SOM fractions  
were measured in soil samples both air-dried and frozen for long-term storage, which makes  
300 freezing an additional option of storage, if also microbial analyses are on the schedule. If  
samples are taken during the dry season, rewetting will reflect natural conditions (e.g. after a  
rain event) much better than rinsing dry soil aggregates into SPT solution and thereby risking  
strong slaking and effects of high ionic strength of dissolved Na<sup>+</sup> similar to the solution of  
sodic soils (Rengasamy and Olsson, 1991; Liu et al., 2021).



305 **5 Conclusion**

The present study shows, that measuring structural characteristics of soils with very different soil textures is strongly affected by drying and re-incubation treatments. The strength of POM occlusion decreases after rewetting, even if slaking is avoided, and is not recovered within the following four weeks. In contrast, the amount of water-stable macroaggregates remains stable and only increased significantly in one soil directly after rewetting. This work shows the importance of soil structure measurements with field-fresh samples to avoid overestimation of fee and weakly occluded POM fractions and erosion stability. The structure of measurement campaigns should be adapted to that issue. It further underpins that both investigated methods differ in their measured characteristics and should be used together and not substitute each other.



### **Data availability**

All of the data are published within this paper and in the Supplement.

### **Author Contributions**

320 FB developed the experimental concept, organized and conducted the soil sampling and the laboratory work, analyzed the data and prepared the manuscript. SD planned the extraction process, adapted the USD method, conducted and managed the experiment. JK conducted the laboratory work and data collection.

### **Competing interests**

325 The authors declare that they have no conflict of interest.

### **Acknowledgements**

330 Many thanks go to Sandra Reimann, Carlotta Kollmann and Christine Beusch for their laboratory support. We are also very grateful to the farmers for the opportunity to sample their soils and the German Federal Ministry of Education and Research for funding (BMBF, funding number 01UU2202).



## References

- Almajmaie, A., Hardie, M., Acuna, T. and Birch, C.: Evaluation of methods for determining soil aggregate stability, *Soil Till. Res.*, 167, 39-45, <https://doi.org/10.1016/j.still.2016.11.003>, 2017.
- Amézqueta, E.: Soil Aggregate Stability: A Review, *J. Sustain. Agric.*, 14(2-3), 83–151, [https://doi.org/10.1300/j064v14n02\\_08](https://doi.org/10.1300/j064v14n02_08), 1999.
- Angulo, V., Bleichrodt, R. J., Dijksterhuis, J., Erktan, A., Hefting, M. M., Kraak, B. and Kowalchuk, G. A.: Enhancement of soil aggregation and physical properties through fungal amendments under varying moisture conditions, *Environ. Microbiol.*, 26(5), e16627, <https://doi.org/10.1111/1462-2920.16627>, 2024.
- Annabi, M., Houot, S., Francou, C., Poitrenaud, M. and Bissonnais, Y. L.: Soil Aggregate Stability Improvement with Urban Composts of Different Maturities, *Soil Sci. Soc. Am. J.*, 71(2), 413, <https://doi.org/10.2136/sssaj2006.0161>, 2007.
- Barthès, B. and Roose, E.: Aggregate stability as an indicator of soil susceptibility to runoff and erosion; validation at several levels, *Catena*, 47(2), 133-149, [https://doi.org/10.1016/S0341-8162\(01\)00180-1](https://doi.org/10.1016/S0341-8162(01)00180-1), 2002.
- Bavel, C. V.: Mean weight-diameter of soil aggregates as a statistical index of aggregation, *Proceedings, Soil Science Society of America*, 1949, 1950, Vol. 14, 20-23/1950, 1950.
- Bossuyt, H., Denef, K., Six, J., Frey, S. ., Merckx, R. and Paustian, K.: Influence of microbial populations and residue quality on aggregate stability, *Appl. Soil Ecol.*, 16(3), 195–208, [https://doi.org/10.1016/s0929-1393\(00\)00116-5](https://doi.org/10.1016/s0929-1393(00)00116-5), 2001.
- Bronick, C. J. and Lal, R.: Soil structure and management: a review, *Geoderma*, 124, 3--22, <https://doi.org/10.1016/j.geoderma.2004.03.005>, 2005.
- Bryant, J. C., Bendixen, T. W. and Slater, C. S.: Measurement of the water-stability of soils, *Soil science*, 65(4), 341-346, 1948.
- Büks, F. and Kaupenjohann, M.: Enzymatic biofilm digestion in soil aggregates facilitates the release of particulate organic matter by sonication, *SOIL*, 2, 499–509, <https://doi.org/10.5194/soil-2-499-2016>, 2016.
- Büks, F., Kayser, G., Zieger, A., Lang, F., and Kaupenjohann, M.: Particles under stress: ultrasonication causes size and recovery rate artifacts with soil-derived POM but not with microplastics, *Biogeosciences*, 18, 159–167, <https://doi.org/10.5194/bg-18-159-2021>, 2021.
- Büks, F.: Technical note: The recovery rate of free particulate organic matter from soil samples is strongly affected by the method of density fractionation, *Biogeosciences*, 20, 1529–1535, <https://doi.org/10.5194/bg-20-1529-2023>, 2023.
- Cavael, U., Tost, P., Diehl, K., Büks, F. and Lentzsch, P.: Correlations of Soil Fungi, Soil Structure and Tree Vigour on an Apple Orchard with Replant Soil, *Soil Syst.*, 4(4), 70, <https://doi.org/10.5194/10.3390/soilsystems4040070>, 2020.
- Cerli, C., Celi, L., Kalbitz, K., Guggenberger, G., and Kaiser, K.: Separation of light and heavy organic matter fractions in soil – Testing for proper density cut-off and dispersion level, *Geoderma*, 170, 403–416, <https://doi.org/10.1016/j.geoderma.2011.10.009>, 2012.
- Chepil, W. and Bisal, F.: A Rotary Sieve Method for Determining the Size Distribution of Soil Clods, *Soil Sci.*, 56, 95–100, <https://doi.org/10.1097/00010694-194308000-00002>, 1943.
- Chowdhury, N., Marschner, P., and Burns, R.: Response of microbial activity and community structure to decreasing soil osmotic and matric potential, *Plant Soil*, 344(1-2), 241–254, <https://doi.org/10.1007/s11104-011-0743-9>, 2011.
- Christensen, O.: An Index of Friability of Soils, *Soil Sci.*, 29, 119–136, 1930.
- Denef, K., Six, J., Paustian, K. and Merckx, R.: Importance of macroaggregate dynamics in controlling soil carbon stabilization: short-term effects of physical disturbance induced by dry–wet cycles, *Soil Biol. Biochem.*, 33(15), 2145-2153, [https://doi.org/10.1016/S0038-0717\(01\)00153-5](https://doi.org/10.1016/S0038-0717(01)00153-5), 2001.



- Drenovsky, R. E., Vo, D., Graham, K. J. and Scow, K. M.: Soil water content and organic carbon availability are major determinants of soil microbial community composition, *Microb. Ecol.*, 48(3), 424–430, <https://doi.org/10.1007/s00248-003-1063-2>, 2004.
- 380 Edwards, A. and Bremner, J.: Dispersion of Soil Particles by Sonic Vibration, *J. Soil Sci.*, 18, 47–63, <https://doi.org/10.1111/j.1365-2389.1967.tb01487.x>, 1967a.
- Edwards, A. P. and Bremner, J.: Microaggregates in Soils, *J. Soil Sci.*, 18, 64–73, <https://doi.org/10.1111/j.1365-2389.1967.tb01488.x>, 1967b.
- 385 Emerson, W. W.: A classification of soil aggregates based on their coherence in water, *Soil Res.*, 5(1), 47–57, <https://doi.org/10.1071/SR9670047>, 1967.
- Fan, X., Pan, H., Ping, Y., Jin, G. and Song, F.: The underlying mechanism of soil aggregate stability by fungi and related multiple factor: a review, *Eurasian Soil Sci.*, 55(2), 242–250, <https://doi.org/10.1134/S1064229322020065>, 2022.
- 390 Golchin, A., Oades, J. M., Skjemstad, J. O., and Clarke, P.: Study of free and occluded particulate organic matter in soils by solid state <sup>13</sup>C CP/MAS NMR spectroscopy and scanning electron microscopy, *Soil Res.*, 32, 285–309, <https://doi.org/10.1071/SR9940285>, 1994.
- Graf-Rosenfellner, M., Cierjacks, A., Kleinschmit, B., and Lang, F.: Soil formation and its implications for stabilization of soil organic matter in the riparian zone, *Catena*, 139, 9–18, <https://doi.org/10.1016/j.catena.2015.11.010>, 2016
- 395 Greacen, E.: Water content and soil strength, *J. Soil Sci.*, 11, 313–333, <https://doi.org/10.1111/j.1365-2389.1960.tb01088.x>, 1960.
- Haynes, R.J. and Swift, R.S.: Stability of soil aggregates in relation to organic constituents and soil water content, *J. Soil Sci.*, 41(1), 73–83, <https://doi.org/10.1111/j.1365-2389.1990.tb00046.x>, 1990.
- 400 Jouquet, P., Chintakunta, S., Bottinelli, N., Subramanian, S., and Caner, L.: The influence of fungus-growing termites on soil macro and micro-aggregates stability varies with soil type, *Appl. Soil Ecol.*, 101, 117–123, <https://doi.org/10.1016/j.apsoil.2016.02.001>, 2016.
- Kaiser, M. and Berhe, A. A.: How does sonication affect the mineral and organic constituents of soil aggregates? – A review, *J. Plant. Nutr. Soil Sc.*, 177, 479–495, <https://doi.org/10.1002/jpln.201300339>, 2014.
- 405 Kemper, W. D. and Rosenau, R. C.: Aggregate stability and size distribution. *Methods of soil analysis: Part 1 Physical and mineralogical methods*, 5, 425–442, <https://doi.org/10.2136/sssabookser5.1.2ed.c17>, 1986.
- Kühnel, A., Wiesmeier, M., Spörlein, P., Schilling, B. and Kögel-Knabner, I.: Influence of drying vs. freezing of archived soil samples on soil organic matter fractions, *J. Plant Nutr. Soil Sci.*, 182(5), 772–781, <https://doi.org/10.1002/jpln.201800529>, 2019.
- 410 Lehtinen, T., Lair, G. J., Mentler, A., Gísladóttir, G., Ragnarsdóttir, K. V., and Blum, W. E.: Soil aggregate stability in different soil orders quantified by low dispersive ultrasonic energy levels, *Soil Sci. Soc. Am. J.*, 78(3), 713–723, <https://doi.org/10.2136/sssaj2013.02.0073>, 2014.
- Liu, J., Hu, F., Xu, C., Wang, Z., Ma, R., Zhao, S. and Liu, G.: Comparison of different methods for assessing effects of soil interparticle forces on aggregate stability, *Geoderma*, 385, 114834, <https://doi.org/10.1016/j.geoderma.2020.114834>, 2021.
- 415 Meidl, P., Lehmann, A., Bi, M., Breitenreiter, C., Benkrama, J., Li, E., Riedo, J. and Rillig, M. C.: Combined application of up to ten pesticides decreases key soil processes, *Environ. Sci. Pollut. Res.*, 31(8), 11995–12004, <https://doi.org/10.1007/s11356-024-31836-x>, 2024.
- North, P.: Towards an absolute measurement of soil structural stability using ultrasound, *J. Soil Sci.*, 27, 451–459, <https://doi.org/10.1111/j.1365-2389.1976.tb02014.x>, 1976.
- 420 Oztas, T. and Fayetorbay, F.: Effect of freezing and thawing processes on soil aggregate stability, *CATENA*, 52(1), 1–8, [https://doi.org/10.1016/s0341-8162\(02\)00177-7](https://doi.org/10.1016/s0341-8162(02)00177-7), 2003.
- Rengasamy, P. and Olsson, K.A.: Sodicity and soil structure, *Soil Res.*, 29(6), 935–952, <https://doi.org/10.1071/SR9910935>, 1991.





- 425 Rosenberg, N. J.: Response of plants to the physical effects of soil compaction, *Adv. Agron.*, 16, 181--196,  
[https://doi.org/10.1016/S0065-2113\(08\)60024-3](https://doi.org/10.1016/S0065-2113(08)60024-3), 1964.
- Russell, J.: Report of the subcommittee on soil structure and consistency, *Soil Sci. Soc. Am. J.*, 9, 10--22,  
<https://doi.org/10.2136/sssaj1928.0361599500B920010002x>, 1928.
- 430 Russell, M. B. and Feng, C. L.: Characterization of the stability of soil aggregates, *Soil sci.*, 63(4), 299-304,  
1947.
- Tang, J., Mo, Y., Zhang, J. and Zhang, R.: Influence of biological aggregating agents associated with microbial  
population on soil aggregate stability, *Appl. Soil Ecol.*, 47(3), 153--159,  
<https://doi.org/10.1016/j.apsoil.2011.01.001>, 2011.
- 435 Tisdall, J. M., Smith, S. E. and Rengasamy, P.: Aggregation of soil by fungal hyphae, *Soil Res.*, 35(1), 55-60,  
<https://doi.org/10.1071/S96065>, 1997.
- Yoder, R. E.: A direct method of aggregate analysis of soils and a study of the physical nature of erosion losses,  
*Agron. J.*, 28, 337--351, <https://doi.org/10.2134/agronj1936.00021962002800050001x>, 1936.