



A normalised framework for the Zero Emissions Commitment

Richard G. Williams¹, Philip Goodwin², Paulo Ceppi³, Chris D. Jones^{4,5}, and Andrew MacDougall⁶

¹Department of Earth, Ocean, and Ecological Sciences, School of Environmental Sciences, University of Liverpool, Liverpool, United Kingdom

²School of Ocean and Earth Sciences, University of Southampton, Southampton, UK

³Department of Physics, Imperial College London, London, UK

⁴Hadley Centre, UK Met Office, Exeter, UK

⁵School of Geographical Sciences, University of Bristol, Bristol, UK

⁶St. Francis Xavier University, Nova Scotia, Canada

Correspondence: Richard G. Williams (ric@liverpool.ac.uk)

Abstract. The Zero Emissions Commitment (ZEC) measures the transient climate response after carbon emissions cease, defined by whether there is a continued rise or decrease in global surface temperature. A normalised framework for the ZEC connects the surface temperature response post emissions to carbon, radiative and thermal processes, involving changes in carbon inventories, radiative forcing, planetary heat uptake and climate feedback. The normalised ZEC, defined by the surface temperature change since the pre industrial divided by the temperature change at the time of net zero, is controlled by opposing-
signed contributions: (i) a cooling contribution from a weakening in radiative forcing due to a decrease in atmospheric CO₂ from carbon uptake by the land and ocean versus (ii) surface warming contributions from a decline in the fraction of radiative forcing used for planetary heat uptake augmented by possible amplification by climate feedbacks. From a set of 9 CMIP6 Earth system models following an idealised atmospheric CO₂ experiment, inter-model differences in the post-emission climate response are primarily determined by differences in the ocean heat uptake and the land and ocean uptake of carbon. These inferences as to the controls of the ZEC broadly carry over for diagnostics of a large ensemble, observationally-constrained efficient Earth system model using two different emission scenarios to reach net zero. The large ensembles reveal a partial compensation between the changes in landborne and oceanborne fractions, as well as revealing ensembles with a greater range in amplification of warming by climate feedbacks.

1 Introduction

Climate models reveal a near-linear dependence of the global surface temperature change with cumulative carbon emissions in experiments following idealised CO₂ experiments (Matthews et al., 2009; Allen et al., 2009; Zickfeld et al., 2009; Gillett et al., 2013; Collins et al., 2013). Once carbon emissions cease, climate models suggest either a slight increase or decrease in surface temperature. This delayed climate response to past carbon emissions is important for policy makers as this response affects the maximum amount of carbon that may be emitted before exceeding a warming target (Allen et al., 2022; Matthews and Zickfeld, 2012).



These different phases of the climate response to carbon emissions are represented by two climate metrics. The first climate metric, the Transient Climate Response to Cumulative CO₂ Emissions (TCRE), measures the dependence of surface warming to cumulative carbon emissions (Matthews et al., 2009; Gillett et al., 2013; MacDougall, 2016; Williams et al., 2016; Matthews et al., 2018; Jones and Friedlingstein, 2020; Williams et al., 2020). Individual climate models reveal a nearly constant value for the TCRE over a centennial timescale, although the value of the TCRE varies between individual climate models (Gillett et al., 2013; Williams et al., 2017). The second climate metric, the Zero Emissions Commitment (ZEC), measures the temperature change after the time of net zero and represents the warming that might be in the pipeline from past emissions. Most climate models reveal a slight cooling with a negative ZEC, although some individual models reveal a slight warming with a positive ZEC (MacDougall et al., 2020). There is significant uncertainty as to which processes control the ZEC and different processes dominate according to the timescale of interest, ranging from decades to millennia; see the review by Palazzo Corner et al. (2023).

The climate responses during emissions and post emissions involve a combination of thermal, radiative and carbon processes, including climate and carbon feedback processes. Our aim is to understand the competing effects of these processes in controlling the climate response post emissions in terms of a normalised ZEC, defined by the temperature change relative to the preindustrial divided by the temperature change at the time of net zero. This normalised ZEC is connected to the empirical energy balance at the top of the atmosphere, the dependence of the radiative forcing on atmospheric CO₂ and the global carbon inventory (Section 2).

This framework is applied to diagnostics for a suite of Earth system models following the Zero Emissions Commitment Model-Intercomparison Project (ZECMIP) (Jones et al., 2019; MacDougall et al., 2020), involving a 1% annual rise in atmospheric CO₂ and then emissions cease after a particular cumulative carbon emission is reached (Section 3). This framework reveals the relative importance of thermal, radiative and carbon processes in controlling the climate response post emissions, both for the model-mean response and the inter-model spread. These diagnostics for the climate metrics and their controls are repeated for a large ensemble of observationally-constrained model projections (Goodwin, 2018; Goodwin et al., 2020) following two different choices of emission scenarios, either an annual rise in atmospheric CO₂ or a constant carbon emission (referred respectively to as 1pctCO₂ and flat10 experiments (Sanderson et al., 2024)) until a maximum cumulative carbon emission is reached (Section 4). Finally, the wider implications of the study are discussed and summarised (Section 5).

2 Theory

Theoretical identities are set out for the two key metrics, the TCRE and ZEC, defining the climate response during emissions and post emissions respectively. The TCRE relationship draws upon prior work, but the application to the ZEC has not been set out before.



2.1 Identity for the TCRE

The TCRE measures the dependence of surface warming to cumulative CO₂ emissions and is defined by the change in global-mean, surface air temperature, $\Delta T(t)$ in K, relative to the pre industrial divided by the cumulative carbon emission, $I_{em}(t)$ in EgC, such that

$$\text{TCRE} \equiv \frac{\Delta T(t)}{I_{em}(t)}, \quad (1)$$

where Δ represents the change since the time of the pre industrial. The TCRE is approximately scenario independent and depends only on the cumulative carbon emissions.

The TCRE from (1) may be related to an identity involving the product of two terms, the Transient Climate Response (TCR) affected by climate processes and the airborne fraction affected by the carbon cycle (Matthews et al., 2009; Solomon et al., 2009; Gillett et al., 2013; MacDougall, 2016; Jones and Friedlingstein, 2020), such that

$$\text{TCRE} = \frac{\Delta T(t)}{I_{em}(t)} = \underbrace{\frac{\Delta T(t)}{\Delta I_A(t)}}_{\text{TCR}} \underbrace{\frac{\Delta I_A(t)}{I_{em}(t)}}_{\text{carbon cycle}}, \quad (2)$$

where the TCR is defined by the ratio of the surface temperature change, $\Delta T(t)$, and change in the atmospheric carbon inventory, $\Delta I_A(t)$, and the airborne fraction is defined by the ratio of the change in the atmospheric carbon inventory, $\Delta I_A(t)$, and the cumulative carbon emissions, $I_{em}(t)$.

The TCRE can also be equivalently defined by separating the TCR term in (2) into a product of two terms, involving separate thermal and radiative dependencies, such that

$$\text{TCRE} = \frac{\Delta T(t)}{I_{em}(t)} = \underbrace{\frac{\Delta T(t)}{\Delta F(t)}}_{\text{thermal}} \underbrace{\frac{\Delta F(t)}{\Delta I_A(t)}}_{\text{radiative}} \underbrace{\frac{\Delta I_A(t)}{I_{em}(t)}}_{\text{carbon cycle}}, \quad (3)$$

where the thermal dependence is given by the ratio of the surface temperature change $\Delta T(t)$ and the change in the radiative forcing, $\Delta F(t)$, and the radiative dependence from the ratio of the change in the radiative forcing, $\Delta F(t)$, and the change in the atmospheric carbon inventory, $\Delta I_A(t)$ (Goodwin et al., 2015; Ehlert et al., 2017; Williams et al., 2016, 2017; Katavouta et al., 2018). The benefit of this additional step is to gain insight into the thermal and radiative effects on the TCRE, by drawing upon the energy balance at the top of the atmosphere and the logarithmic dependence of radiative forcing on atmospheric CO₂.

2.2 Identity for the ZEC

The ZEC measures the temperature change relative to the pre industrial, $\Delta T(t)$, minus the temperature change at the time of net zero, t_{ZE} , $\Delta T(t_{ZE})$, and is defined by

$$\Delta T(t) - \Delta T(t_{ZE}). \quad (4)$$



80 This definition of the ZEC measures the absolute value of the temperature change and is likely to be sensitive to the warming level experienced from the emission scenario.

Alternatively, a normalised measure of the ZEC is given by the ratio of the temperature change, $\Delta T(t)$, and the temperature change at the time of net zero, $\Delta T(t_{ZE})$,

$$\frac{\Delta T(t)}{\Delta T(t_{ZE})}. \quad (5)$$

85 This ratio is henceforth referred to as the normalised ZEC and by taking a ratio is less likely to be as sensitive to the warming level. The temperature change, $\Delta T(t)$, may be related to the cumulative carbon emissions, $I_{em}(t)$, by the product of the thermal, radiative and carbon-cycle contributions,

$$\Delta T(t) = \frac{\Delta T(t)}{\Delta F(t)} \frac{\Delta F(t)}{\Delta I_A(t)} \frac{\Delta I_A(t)}{I_{em}(t)} I_{em}(t), \quad (6)$$

so that the normalised ZEC from $\Delta T(t)/\Delta T(t_{ZE})$ may be written as

$$90 \quad \frac{\Delta T(t)}{\Delta T(t_{ZE})} = \underbrace{\left(\frac{\Delta T(t)}{\Delta F(t)} / \frac{\Delta T(t_{ZE})}{\Delta F(t_{ZE})} \right)}_{\text{thermal}} \underbrace{\left(\frac{\Delta F(t)}{\Delta I_A(t)} / \frac{\Delta F(t_{ZE})}{\Delta I_A(t_{ZE})} \right)}_{\text{radiative}} \underbrace{\left(\frac{\Delta I_A(t)}{\Delta I_A(t_{ZE})} \right)}_{\text{carbon cycle}}. \quad (7)$$

The dependence of the emissions is removed after the time of net zero as $I_{em}(t)$ is taken to be fixed for $t > t_{ZE}$. A positive ZEC corresponds to $\Delta T(t)/\Delta T(t_{ZE}) > 0$ and a negative ZEC to $\Delta T(t)/\Delta T(t_{ZE}) < 0$. The normalised temperature changes depend upon the relative size of the normalised thermal, radiative and carbon-cycle terms in (7).

95 Our aim is gain insight into the controls of the ZEC and interpret the continued warming or cooling response in terms of its normalised thermal, radiative and carbon-cycle contributions. Next consider the thermal, radiative and carbon-cycle terms in (7) that determine the normalised ZEC response.

2.2.1 Thermal contribution

100 The thermal contribution may be understood in terms of the energy balance at the top of the atmosphere, where the planetary heat flux into the climate system, ΔN , balances the sum of the radiative forcing into the climate system, ΔF , and the radiative response, ΔR (Gregory et al., 2004; Knutti and Hegerl, 2008; Andrews et al., 2012; Forster et al., 2013),

$$\Delta N(t) = \Delta F(t) + \Delta R(t), \quad (8)$$

where $\Delta F(t)$ and $\Delta R(t)$ are defined as positive when supplying energy into the climate system.

105 The radiative response is parameterised in terms of the product of the climate feedback parameter, $\lambda(t)$, and the change in global mean, surface air temperature, $\Delta T(t)$,

$$\Delta N(t) = \Delta F(t) + \lambda(t)\Delta T(t). \quad (9)$$

The dependence of surface temperature on radiative forcing, $\Delta T(t)/\Delta F(t)$, in (7) is then directly connected from (9) to the product of the inverse of the climate feedback, $\lambda(t)^{-1}$ and the planetary heat uptake divided by the radiative forcing,



$\Delta N(t)/\Delta F(t)$,

$$\frac{\Delta T(t)}{\Delta F(t)} \equiv \frac{1}{\lambda(t)} \frac{\Delta R(t)}{\Delta F(t)} = -\frac{1}{\lambda(t)} \left(1 - \frac{\Delta N(t)}{\Delta F(t)} \right), \quad (10)$$

where $(1 - \Delta N(t)/\Delta F(t))$ represents the fraction of the radiative forcing that escapes back to space, rather than being used for planetary heat uptake.

2.2.2 Radiative contribution

The radiative forcing, $\Delta F(t)$, may be separated into a CO₂ radiative forcing contribution, $\Delta F_{CO_2}(t)$, and a non-CO₂ radiative forcing contribution, $\Delta F_{nonCO_2}(t)$, including the contribution of other greenhouse gases and aerosols,

$$\begin{aligned} \Delta F(t) &= \Delta F_{CO_2}(t) + \Delta F_{nonCO_2}(t) \\ &= \Delta F_{CO_2}(t) (1 + \Delta F_{nonCO_2}(t)/\Delta F_{CO_2}(t)). \end{aligned} \quad (11)$$

The CO₂ radiative forcing contribution, $\Delta F_{CO_2}(t)$, may be related to the change in the logarithm of atmospheric CO₂ relative to the pre industrial,

$$\Delta F_{CO_2}(t) = a \Delta \ln CO_2(t) = a (\ln CO_2(t) - \ln CO_2(t_o)), \quad (12)$$

where a is a radiative forcing coefficient in W m⁻² (that is model dependent) and t_o is the time of the pre-industrial. The change in the logarithm is equivalent to the fractional change, $\delta \ln x = \delta x/x$, so that (12) may be written as

$$\Delta F_{CO_2}(t) = a \frac{\Delta CO_2(t)}{CO_2(t)} \equiv a \frac{\Delta I_A(t)}{I_A(t)}, \quad (13)$$

where $I_A(t)$ is the atmospheric inventory of carbon dioxide (that is dependent on the product of the molar mass of the atmosphere and the mixing ratio of atmospheric CO₂).

The ratio of the change in the radiative forcing from atmospheric CO₂ and atmospheric carbon is then given from (13) by

$$\frac{\Delta F_{CO_2}(t)}{\Delta I_A(t)} = \frac{a}{I_A(t)}, \quad (14)$$

and the normalised radiative contribution from CO₂ to the ZEC in (7) is given by

$$\frac{\Delta F_{CO_2}(t)}{\Delta I_A(t)} / \frac{\Delta F_{CO_2}(t_{ZE})}{\Delta I_A(t_{ZE})} = \frac{I_A(t_{ZE})}{I_A(t)}. \quad (15)$$

and if there are non-CO₂ radiative contributions, then the normalised radiative contribution to the ZEC is then

$$\frac{\Delta F(t)}{\Delta I_A(t)} / \frac{\Delta F(t_{ZE})}{\Delta I_A(t_{ZE})} = \frac{I_A(t_{ZE})}{I_A(t)} \frac{(1 + \Delta F_{nonCO_2}(t)/\Delta F_{CO_2}(t))}{(1 + \Delta F_{nonCO_2}(t_{ZE})/\Delta F_{CO_2}(t_{ZE}))}. \quad (16)$$

2.2.3 Carbon-cycle contribution

The change in atmospheric carbon inventory, $\Delta I_A(t)$, is related to the carbon budget involving the cumulative carbon emission, $I_{em}(t)$, and the changes in the land and ocean carbon inventories, $\Delta I_L(t)$ and $\Delta I_O(t)$,

$$\Delta I_A(t) = I_{em}(t) - \Delta I_L(t) - \Delta I_O(t). \quad (17)$$



This response can be expressed in terms of the airborne fraction, $\Delta I_A(t)/I_{em}(t)$, varying with the landborne and oceanborne fractions, $\Delta I_L(t)/I_{em}(t)$ and $\Delta I_O(t)/I_{em}(t)$ respectively (Jones et al., 2013),

$$\frac{\Delta I_A(t)}{I_{em}(t)} = 1 - \frac{\Delta I_L(t)}{I_{em}(t)} - \frac{\Delta I_O(t)}{I_{em}(t)}. \quad (18)$$

2.2.4 Mechanistic insight from the normalised ZEC identity

140 The ZEC response may be affected by a wide range of thermal, radiative and carbon processes, so that isolating their causal effect and comparing their relative importance is challenging to achieve. The benefit of the normalised ZEC framework is that there is a more direct link provided to the thermal, radiative and carbon processes by utilising the top of the atmosphere energy balance (9) and the radiative dependence (13), such that combining (7) with (10) and (16) leads to

$$\begin{aligned} \frac{\Delta T(t)}{\Delta T(t_{ZE})} = & \underbrace{\left(\frac{\lambda(t_{ZE})}{\lambda(t)} \left(1 - \frac{\Delta N(t)}{\Delta F(t)} \right) / \left(1 - \frac{\Delta N(t_{ZE})}{\Delta F(t_{ZE})} \right) \right)}_{\text{thermal}} \\ & \times \underbrace{\left(\frac{I_A(t_{ZE})}{I_A(t)} \frac{(1 + \Delta F_{nonCO2}(t)/\Delta F_{CO2}(t))}{(1 + \Delta F_{nonCO2}(t_{ZE})/\Delta F_{CO2}(t_{ZE}))} \right)}_{\text{radiative}} \underbrace{\left(\frac{\Delta I_A(t)}{\Delta I_A(t_{ZE})} \right)}_{\text{carbon cycle}} \end{aligned} \quad (19)$$

Hence, whether there is continued warming, $\Delta T(t)/\Delta T(t_{ZE}) > 1$, and a positive ZEC depends on the time evolution of the products of (i) the thermal contribution involving the climate feedback, $\lambda(t)$, and the dependence of the planetary heat uptake on the radiative forcing, $\Delta N(t)/\Delta F(t)$, (ii) the radiative contribution involving the atmospheric carbon inventory, $I_A(t)$, and the ratio of non-CO₂ radiative forcing and CO₂ radiative forcing, $\Delta F_{nonCO2}(t)/\Delta F_{CO2}(t)$, and (iii) the change in atmospheric carbon, $\Delta I_A(t)$, which via the carbon budget (17) is related to the cumulative carbon emissions, $I_{em}(t)$, minus the increase in land and ocean carbon inventories, $\Delta I_L(t) + \Delta I_O(t)$.

155 For idealised experiments with only forcing from atmospheric CO₂, the normalised identity for the ZEC (19) simplifies with the normalised radiative contribution given by the ratio of the atmospheric carbon inventory, $I_A(t_{ZE})/I_A(t)$, so that

$$\begin{aligned} \frac{\Delta T(t)}{\Delta T(t_{ZE})} = & \underbrace{\left(\frac{\lambda(t_{ZE})}{\lambda(t)} \left(1 - \frac{\Delta N(t)}{\Delta F(t)} \right) / \left(1 - \frac{\Delta N(t_{ZE})}{\Delta F(t_{ZE})} \right) \right)}_{\text{thermal}} \\ & \times \underbrace{\left(\frac{I_A(t_{ZE})}{I_A(t)} \right)}_{\text{radiative}} \underbrace{\left(\frac{\Delta I_A(t)}{\Delta I_A(t_{ZE})} \right)}_{\text{carbon cycle}} \end{aligned} \quad (20)$$

This relationship for the normalised ZEC, $\Delta T(t)/\Delta T(t_{ZE})$, can then be used to (i) provide mechanistic insight as to the drivers of the temperature change after net zero and (ii) explain inter-model differences in the response of Earth system models after net zero.




Figure 1. Diagnostics of the climate response for a 1pctCO₂ experiment with an annual 1% increase in atmospheric CO₂ until the cumulative carbon emission reaches 1000 PgC from ZECMIP (Jones et al., 2019): (a) cumulative carbon emission, $I_{em}(t)$ in PgC, versus time in years; (b) change in global-mean surface air temperature relative to the pre industrial, $\Delta T(t)$ in K, versus time; and (c) the change in surface air temperature versus cumulative carbon emissions. The two key climate metrics are defined by these relationships, the TCRE defined by the slope in (c) and the ZEC defined by the temperature change in (b) relative to the time of net zero or by the vertical excursions in (c) after the maximum cumulative carbon emission is reached. The plot includes smoothing of temperature with a 10 year running mean.

165 2.3 Analyses of ZECMIP responses

2.4 Core experiments

The responses of 9 full Earth system models are analysed following the ZECMIP protocols (Jones et al., 2019; MacDougall et al., 2020), involving an annual 1% rise in atmospheric CO₂ until a cumulative carbon emission of 1000 PgC is reached and then there is no further carbon emission (Fig. 1a). A single realisation is analysed for each model.



170 Under the ZECMIP protocol, each individual model experiment branches at the time of net zero, one branch continuing with the 1% rise in atmospheric CO₂ and the other branch continuing with no further emissions. The time of net zero as defined by the branch point varies from 61 to 71 years across the set of Earth system models (MacDougall et al., 2020).

Prior to net zero, the global-mean surface temperature increases nearly linearly with the rise in cumulative carbon emissions (Fig. 1b). There is a nearly constant slope of the temperature change versus cumulative carbon emissions up until the maximum
 175 emission, which defines the climate metric, the TCRE (Fig. 1c).

After net zero, there are a range of temperature responses from a slight cooling to a slight continued warming (Fig. 1b), where the temperature change relative to the temperature at net zero defines the ZEC. The continued temperature change is also evident in the positive and negative excursions in temperature at the maximum carbon emissions in Fig. 1c.

The temperature response after net zero is interpreted in terms of a normalised ZEC involving the continuing temperature
 180 change, $\Delta T(t)$, divided by the temperature change at net zero, $\Delta T(t_{ZE})$. This estimate of the temperature change at net zero is performed using a 20 year averaging window around the time of net zero to reduce the effect of interannual variability. The averaging is performed on the 1% branch experiment that always includes carbon emissions with an approximately linear rise in temperature, rather than combining a forced response up to net zero and an unforced response after net zero; this choice follows MacDougall et al. (2020) to avoid a possible bias in the estimate of the temperature at net zero, $\Delta T(t_{ZE})$. This
 185 averaging approach is applied for all the variables evaluated at net zero in our normalised framework.

The temperature response during and after emissions, and the associated climate metrics TCRE and ZEC involves a variety of competing drivers (Fig. 2) involving changes in carbon inventories, radiative forcing, radiative response and planetary heat uptake.

2.4.1 Changes in carbon inventories

190 The carbon emissions lead to an increase in the atmospheric, ocean and carbon inventories: a temporary increase in the atmospheric carbon inventory (with a model mean and inter-model standard deviation) of 488 ± 33 PgC at years 55-75 and the remainder taken up by the land and ocean inventories, 253 ± 53 PgC and 207 ± 26 PgC respectively (Fig. 2a-c). Post emissions, the cumulative carbon emission of 1006 ± 31 PgC is more equally partitioned between the atmosphere, land and ocean, each holding 341 ± 51 PgC, 354 ± 80 PgC and 311 ± 40 PgC respectively at years 140-160.

195 2.4.2 Changes in radiative response and planetary heat uptake

The changes in atmospheric carbon dioxide drive the changes in radiative forcing, reaching a maximum radiative forcing of 3.1 ± 0.2 W m⁻² at years 55-75 close to net zero (Fig. 2d). Most of the radiative forcing is returned to space with the radiative response reaching -2.0 ± 0.5 W m⁻² and a smaller planetary heat uptake reaching 1.1 ± 0.6 W m⁻² (Fig. 2e,f). Post emissions, the radiative forcing reduces to 2.4 ± 0.3 W m⁻² at years 140-160 with the radiative response only slightly decreasing in
 200 magnitude to -1.9 ± 0.4 W m⁻² and the planetary heat uptake reducing further to 0.5 ± 0.5 W m⁻².

Hence, the temperature response up to and after net zero involves changes in atmospheric carbon due to the land and ocean carbon uptake, and the resulting radiative forcing is either returned to space or used to warm the planet. The goal now is to




Figure 2. Climate response during emissions and post emissions versus time (year) since the pre industrial for the 9 Earth system models: changes in (a) atmospheric carbon inventory, ΔI_A (PgC); (b) land carbon inventory, ΔI_L (PgC); (c) ocean carbon inventory, ΔI_O (PgC); (d) radiative forcing supplying heat to the climate system, F (W m^{-2}); (e) radiative response representing a heat loss to space, $-\Delta R$ (W m^{-2}); and (f) planetary heat uptake, ΔN (W m^{-2}), positive representing a gain in heat. The plot includes smoothing of planetary heat uptake with a 10 year running mean.

draw upon the identity for the normalised ZEC in order to compare the effect of changes in the carbon sinks, radiative response and planetary heat uptake.

205 2.5 Controls of the ZEC and normalised ZEC

The ZEC measures the temperature change after net zero. The timing of net zero varies from years 61 to 71 in the set of models and, in our subsequent analysis, we choose to align their time series so that the timing of net zero coincides. The ZEC, defined



by $\Delta T(t) - \Delta T(t_{ZE})$, reaches -0.03 ± 0.14 K for year 25, and -0.10 ± 0.19 K and -0.10 ± 0.24 K for years 50 and 90 after net zero (Fig. 3a) (MacDougall et al., 2020).

210 Alternatively, the normalised ZEC, given by the ratio of the temperature change relative to the pre industrial, $\Delta T(t)$, and the change for net zero, $\Delta T(t_{ZE})$, varies from 0.97 ± 0.09 for year 25 to 0.93 ± 0.11 and 0.93 ± 0.14 respectively for years 50 and 90 after net zero (Table 1; Fig. 3b).

The model-mean changes in the normalised ZEC are relatively small, accounting for a temperature anomaly decrease of only 7% after net zero. However, the individual model responses are much larger, reaching 20% changes after net zero; as
 215 previously highlighted by MacDougall et al. (2020).

The normalised ZEC response is made up of competing thermal and carbon responses, as revealed in the normalised framework (19): (i) the normalised thermal contribution, $\Delta T(t)/\Delta F(t)$, is large and positive, reaching 1.22 ± 0.11 and 1.33 ± 0.15 after 50 and 90 years respectively (Table 1; Fig. 3c); and (ii) the normalised radiative contribution, $\Delta F(t)/\Delta I_A(t)$, is relatively small, only reaching 1.09 ± 0.02 and 1.11 ± 0.02 after 50 and 90 years respectively (Fig. 3d); and (iii) the normalised carbon
 220 contribution, $\Delta I_A(t)$, is large and negative, reaching 0.70 ± 0.06 and 0.63 ± 0.06 after 50 and 90 years respectively (Table 1; Fig. 3e). Hence, the normalised ZEC is primarily determined by a competition between the normalised thermal and carbon contributions.

For individual models, there are some large variations, with the normalised thermal contribution exceeding a 30% increase for CNRM-ESM2 and UKESM1, and the normalised carbon contribution reaching a 30% decrease for CanESM5, CESM2,
 225 CNRM-ESM2, GFDL-ESM2, and NorESM2 (Fig. 4, red and green lines; Table A1).

The resulting normalised ZEC response involves an interplay of these normalised thermal and carbon contributions. For example, the positive normalised ZEC response for UKESM1 is due to a strong thermal contribution and only a moderate opposing carbon contribution, while the positive normalised ZEC response for CNRM-ESM2 involves a very strong thermal contribution and an opposing strong carbon contribution. Meanwhile the negative normalised ZEC response for NorESM2 is
 230 due to a relatively modest thermal contributions and relatively strong opposing carbon contributions.

The inter-model spread of the normalised ZEC, measured by the coefficient of variation, reaches 0.12 after 50 years and is made up of contributions of 0.09 for the thermal contribution, 0.02 for the radiative contribution and 0.08 for the carbon contribution (Table 1). Hence, the thermal contribution is the most important contributor to the inter-model spread in the ZEC, closely followed by the carbon contribution and the radiative contribution is least important.

235 2.6 Carbon contribution to the ZEC response

The carbon contribution to the normalised ZEC response involves a normalised decrease in the atmospheric carbon inventory, $\Delta I_A(t)$, which is achieved by an increase in both land and ocean carbon inventories.

In order to compare these different carbon sinks, the carbon changes of each inventory are henceforth normalised by the same cumulative carbon emission at net zero, as given by the airborne, landborne and oceanborne fractions. The airborne fraction,
 240 $\Delta I_A(t)/I_{em}(t_{ZE})$, is a maximum at net zero and then declines in time for all models (Fig. 5, black line) due to the increase in the landborne and oceanborne fractions (Fig. 5, green and blue lines). The airborne fraction is 0.52 ± 0.03 at net zero and




Figure 3. Temporal evolution of the ZEC, its normalised value and components after net zero when emissions cease: (a) the ZEC, surface temperature change, $\Delta T(t') - \Delta T(t'_{ZE})$ in K, after net zero is reached (year); (b) the normalised ZEC, from the normalised surface temperature change, $\Delta T(t')/\Delta T(t'_{ZE})$, a value greater than 1 defines a positive ZEC and a value less than 1 defines a negative ZEC; (c) the thermal contribution from the normalised dependence of surface temperature on radiative forcing, $\Delta T(t')/\Delta F(t')$; (d) the radiative contribution from the normalised dependence of radiative forcing on atmospheric carbon, $\Delta F(t')/\Delta I_A(t')$; (e) the carbon contribution from the normalised atmospheric carbon, $\Delta I_A(t')$. The time series for each individual model is aligned so that the timing of net zero coincides. The normalisation is taken from the average value of the variable over a 20 year period centered on net zero based on the linear response of the 1pct continually-forced experiment. The plot includes smoothing of temperature with a 10 year running mean.

decreases to 0.38 ± 0.05 and 0.34 ± 0.05 at years 50 and 90 after net zero (Table 1). The landborne fraction, $\Delta I_L(t)/I_{em}(t_{ZE})$, increases from 0.26 ± 0.04 at net zero reaching 0.34 ± 0.07 and 0.35 ± 0.08 for 50 and 90 years later respectively; and the oceanborne fraction, $\Delta I_O(t)/I_{em}(t_{ZE})$ increases from 0.22 ± 0.03 at net zero reaching 0.28 ± 0.04 and 0.31 ± 0.05 for 50 and




Figure 4. Temporal evolution of the normalised ZEC, the normalised surface temperature change, and its contributions for 9 different Earth system models: the normalised surface temperature change, $\Delta T(t')$, (black line); the thermal contribution from the normalised dependence of surface temperature on radiative forcing, $\Delta T(t')/\Delta F(t')$ (red line); the radiative contribution from the normalised dependence of radiative forcing on atmospheric carbon, $\Delta F(t')/\Delta I_A(t')$ (blue line); and carbon contribution from the normalised atmospheric carbon, $\Delta I_A(t')$ (green line). The plot includes smoothing of temperature, the alignment of the time series and normalisation as in Fig. 3.

245 90 years later. Hence, initially after net zero, the carbon uptake by the terrestrial system dominates over that by the ocean for most models, but they become comparable to each other by 90 years.

The landborne fraction is much larger than the oceanborne fractions for CanESM5, CNRM-ESM2 and GFDL-ESM2, while the landborne and oceanborne fractions are comparable for UKESM1 and the landborne fraction is much smaller than the oceanborne fraction for ACCESS-ESM1.5 (Fig. 5; Table A1).




Figure 5. Temporal evolution of cumulative airborne fraction, $\Delta I_A(t')$ (black), landborne fraction $\Delta I_L(t')$ (green) and oceanborne fraction $\Delta I_O(t')$ (blue) for time (year) relative to net zero for 9 different Earth system models.

250 The inter-model spread is much larger for the landborne fraction than the oceanborne fraction with coefficients of variation of 0.21 and 0.13 respectively after 50 years (Table 1). These inter-model differences in the landborne and oceanborne fractions are partly compensating, since both coefficients of variation are larger than that for the airborne fraction reaching 0.12.

2.7 Thermal response

255 The thermal contribution to the ZEC may be understood in terms of the top of the atmosphere energy balance (8). The radiative forcing, $\Delta F(t)$, peaks close to the time of net zero and then declines for each model (Fig. 6, black line).

The radiative response, $\Delta R(t)$, is negative and so represents the part of the radiative forcing that is returned to space. The radiative response varies between models, most involve a peak in magnitude at the time of net zero and then a slight decline




Figure 6. Temporal evolution of radiative forcing, $\Delta F(t')$ (black), planetary heat uptake, $\Delta N(t')$ (blue), and radiative response, $-\Delta R(t')$ (red) in W m^{-2} for time (year) relative to net zero for 9 different Earth system models. The plot includes smoothing of temperature and planetary heat uptake with a 10 year running mean.

in magnitude, while in some models the radiative response remains relatively constant in time (Fig. 6, red line). The planetary heat uptake, $\Delta N(t)$, represents the mismatch between the radiative forcing and radiative response. The planetary heat uptake is a maximum at the time of net zero and declines in time for all models (Fig. 6, blue line). For these thermal quantities there is significant interannual variability.

The thermal contribution to the normalised ZEC response, the normalised $\Delta T(t)/\Delta F(t)$, may be derived from the top of the atmosphere energy balance (9). This thermal contribution increases after net zero (Fig. 7, black line) and is made up itself by the product of contributions from the fraction of the radiative forcing escaping to space, the normalised $\Delta R(t)/\Delta F(t)$, and the inverse of the climate feedback parameter, the normalised $\lambda(t)^{-1}$ (Fig. 7, blue and red lines respectively).




Figure 7. Temporal evolution of the thermal contribution defined by the normalised temperature dependence on radiative forcing, $\Delta T(t')/\Delta F(t')$ (black), normalised fraction of radiative forcing escaping to space, $\Delta R(t')/\Delta F(t')$ (blue), and normalised reciprocal of climate feedback parameter $\lambda(t')^{-1}$ (red) for time (year) relative to net zero for 9 different Earth system models. The alignment of the time series and normalisation as in Fig. 3. The plot includes smoothing of temperature and planetary heat uptake with a 10 year running mean.

The normalised fraction of the radiative forcing escaping to space, $\Delta R(t)/\Delta F(t)$, increases after net zero and reaches 1.26 ± 0.11 and 1.30 ± 0.15 for 50 and 90 years later respectively (Table 1, Fig. 7, red line). The normalised inverse of the climate feedback parameter, $\lambda(t)^{-1}$, is close to 1, reaching 0.98 ± 0.06 and 1.03 ± 0.05 for 50 and 90 years later respectively. Thus, the dominant contribution to the increase in the thermal contribution to the normalised ZEC response is from an increase in the fraction of the radiative forcing escaping to space or equivalently given by a decrease in the fraction of the radiative forcing used for planetary heat uptake.

For most individual models, the thermal contribution to the normalised ZEC response, the normalised $\Delta T(t)/\Delta F(t)$, is broadly the same as the fraction of radiative forcing escaping to space, $\Delta R(t)/\Delta F(t)$ (Fig. 7, blue line; Table A1). However,



there is an enhancement of the thermal contribution from the increase in the radiative forcing escaping to space by a time-
 275 varying amplification from the climate feedback parameter for UKESM1 and for the latter parts of the temporal record for
 GFDL-ESM2 and NorESM2-LM (Fig. 7, red line).

Intermodel differences in the thermal contribution are dominated by differences in the fraction of radiative forcing returned
 to space, rather than from differences in the inverse of the climate feedback parameter, since their coefficients of variation are
 0.09 and 0.06 respectively after 50 years (Table 1).

280 These diagnostics of the normalised ZEC framework are based on single model ensembles from ZECMIP. The diagnostic
 framework is next applied to a much larger set of model ensembles to span more fully parameter space and reveal how
 representative our diagnostics for ZECMIP are.

3 Analyses of a large ensemble of an efficient Earth system model

3.1 Experiments with the efficient Earth system model

285 The ZEC diagnostics are now repeated for a large ensemble of an efficient Earth system model (WASP) (Goodwin, 2016).
 WASP includes time-varying climate feedbacks, to represent time-varying changes in the pattern effect (Goodwin, 2018;
 Goodwin et al., 2020), and with the addition here that air-sea exchange for CO₂ includes a full carbonate chemistry solver
 for the surface ocean (Follows et al., 2006).

In these experiments, 10 million prior simulations are integrated using historical forcing and following the SSP245 exper-
 290 iment from year 2014. Of these simulations, 1138 posterior solutions are identified that satisfy observable quantities after
 Goodwin (2018).

These 1138 posterior ensemble members are then integrated forward following two different experiments: (i) an annual 1%
 increase in atmospheric CO₂ with emissions ceasing at 1000 PgC (referred to as the 1pctCO₂ case as for ZECMIP) or (ii) a
 constant emission rate of 10 PgC yr⁻¹ for 100 years until there is 1000 PgC emitted (referred to as the flat10 case) (Sanderson
 295 et al., 2024). This comparison is included as flat10 is scenario choice for CMIP7 and has the benefit of a more constant forcing
 regime.

In the 10 million historically forced prior simulations used to determine observational consistency the WASP model sim-
 ulations include an imposed internal variability (Goodwin, 2018). This internal variability is turned off when the posterior
 simulations are then forced with idealised experiments.

300 3.2 ZEC responses for the large ensemble model

The ZEC responses reveal a slight decrease in surface temperature after net zero for the median of the ensembles for both
 the 1pctCO₂ and flat10 experiments (Fig. 8a,b, blue line). For the 1pctCO₂ experiment, the median ZEC and the 5% to 95%
 ensemble range in brackets are -0.10 K (-0.47 K to 0.43 K) after 50 years, increasing in magnitude to -0.09 K (-0.56 K to
 0.82 K) after 100 years (Fig. 8a; Table A2). There is close agreement in these ZEC estimates with the ZECMIP model mean




Figure 8. Simulations for ZEC from a large model ensemble from WASP over time since net zero (year): (a) the 1pctCO₂ experiment, including the median (line), 1-sigma range (dark shading) and 95% range (light shading) together with the maximum and minimum ZECMIP range; (b) the flat10 experiment; and (c) scatterplot comparing the ZEC responses for the 1pctCO₂ and flat10 experiment for 20, 50 and 100 years for each model realisation.

of -0.10 K at 50 years (Table 1) and the ZECMIP range is comparable to the 1 standard deviation range from WASP (Fig. 8, orange line and dark blue shading).

For the flat10 experiment, the median ZEC and the 5% to 95% ensemble range in brackets are slightly smaller: -0.06 K (-0.25 K to 0.41 K) after 50 years, increasing in magnitude to -0.07 K (-0.31 K to 0.73 K) after 100 years and further to -0.19 K (-0.46 K to 1.02 K) after 400 years (Fig. 8b, Table A3). This slightly smaller magnitude response for flat10 is due to there being a stronger radiative forcing for the 1pctCO₂ case as the forcing is more non linear.

There is some slight curvature in the ZEC responses between the 1pctCO₂ and flat10 experiments with a greater range for the 1pctCO₂ ensembles (Fig. 8c).

The changes in carbon inventories, radiative forcing, radiative feedback and planetary heat uptake (Fig. A1; Tables S2 and S3) vary in a broadly similar manner as for the ZECMIP diagnostics over 100 years (Figs. 1 and 2), although include a much greater ensemble spread and extend for much longer to 400 years.

3.3 Normalised contributions to the ZEC

The normalised ZEC slightly decreases after net zero for the ensemble median to 0.92, 0.92 and 0.82 for years 50, 100 and 400 for the 1pctCO₂ experiment, and 0.95, 0.94 and 0.83 for years 50, 100 and 400 after net zero for the flat10 experiment

(Tables A2 and A3). There is a wide inter-ensemble spread with the normalised ZEC varying from 0.71 to 1.50 at year 100 including a tail of ensembles with much higher normalised ZEC (Fig. 9a for the flat10 experiment).




Figure 9. Normalised ZEC and its components following the flat10 experiment from the WASP simulations: (a) the normalised ZEC from the normalised surface temperature change, $\Delta T(t')$, including the median (line), 1-sigma range (dark shading) and 95% range (light shading); (b) the thermal contribution from the normalised dependence of surface temperature on radiative forcing, $\Delta T(t')/\Delta F(t')$; (c) the radiative contribution from the normalised dependence of radiative forcing on atmospheric carbon, $\Delta F(t')/\Delta I_A(t')$; and (d) the carbon contribution from the normalised atmospheric carbon, $\Delta I_A(t')$. In each case, the normalisation is by the value of the variable at the time of net zero.

320

For each ensemble, the normalised ZEC value (Fig. 9a) equates to the product of the ensemble values for each of the ZEC contributions (Fig. 9b-d). The thermal contribution to the normalised ZEC, the normalised $\Delta T/\Delta F$, increases in time for all ensembles to a median of 1.24 (5% and 95% range of 0.97 to 1.75) after 100 years (Fig. 9b). The radiative contribution to the normalised ZEC, the normalised $\Delta F/\Delta I_A$, only slightly increases in time for all ensembles to 1.07 (1.04 to 1.09) after 100 years (Fig. 9c). The carbon contribution to the normalised ZEC given by the normalised change in atmospheric carbon, ΔI_A , decreases for all ensembles to 0.71 (0.67 to 0.86) after 100 years (Fig 9d).

325

The thermal contribution to the normalised ZEC, the normalised $\Delta T(t)/\Delta F(t)$ and its contributions, reveals similar changes as for median ensemble response as in ZECMIP (Fig. 10a, full lines): there is a strengthening in the normalised $\Delta T(t)/\Delta F(t)$ in time, which is primarily due to the strengthening in the normalised fraction of radiative forcing returned to space, $1 - \Delta N(t)/\Delta F(t)$ augmented by a strengthening in the normalised inverse climate feedback, $\lambda(t)^{-1}$.

330

This contribution is made up of the product of two terms, the normalised radiative response divided by the radiative forcing, the normalised $\Delta R(t')/\Delta F(t')$, and the normalised inverse of the climate feedback parameter, the normalised $\lambda(t')^{-1}$. There is a consistent increase in the fraction of the radiative forcing returned to space or equivalently a decrease in the fraction of radiative forcing used for planetary heat uptake with a relatively tight ensemble spread and the median increasing to 1.12 (5%




Figure 10. Temporal evolution of thermal and carbon variables affecting the normalised ZEC for the flat10 experiment from the WASP simulations: (a) thermal contribution to the normalised ZEC from the normalised temperature dependence on radiation, $\Delta T(t')/\Delta F(t')$ (black line for median, grey shading for 95% range), normalised fraction of radiative forcing escaping to space, $\Delta R(t')/\Delta F(t') = (1 - \Delta N(t')/\Delta F(t'))$ (blue line and shading) and normalised inverse climate feedback parameter $\lambda(t')^{-1}$ (orange line and pale shading); and (b) the partitioning of cumulative carbon emissions into the airborne fraction (black line for median and grey shading for 95% range), oceanborne fraction (blue line and shading) and landborne fraction (green line and shading) for time (year) relative to net zero.

335 and 95% spread of 1.00 to 1.19) (Fig. 10a, blue line and shading; Table A3). The normalised inverse of the climate feedback parameter only slightly increases in time for the median to 1.10 after 100 years (Fig. 10a, orange line and pale shading; Table A2), but there is a wide, asymmetrical spread with the 5% to 95% range extending from 0.84 to 1.73, so including ensembles with much larger $\lambda(t)^{-1}$ providing a greater amplification by climate feedbacks compared with the ZECMIP diagnostics.

The carbon contribution to the normalised ZEC, the normalised atmospheric carbon, $\Delta I_A(t)$, may be understood by the changes in airborne fraction, $\Delta I_A(t)/I_{em}(t)$, which progressively decreases in time (Fig. 10b, black line and grey shading; Table A3). The dominant contribution to the changes in airborne fraction alters from being from the landborne fraction close to the time of net zero, $\Delta I_L(t)/I_{em}(t)$, to the oceanborne fraction on timescales greater than 50 years after net zero, $\Delta I_O(t)/I_{em}(t)$ (Fig. 10b, green and blue lines and shading respectively).

345 There is a much larger ensemble spread around the landborne and ocean borne responses than for the airborne fraction, which implies that the changes in land and ocean carbon sinks partly compensate for each other. This partial compensation in carbon sinks is consistent with the coefficient of variation being larger for the landborne and oceanborne fractions than the airborne fraction as diagnosed for ZECMIP (Table 1).

In summary, the ZEC responses and their normalised contributions are broadly similar in the diagnostics of the large ensemble WASP and the smaller set of 9 Earth system models in ZECMIP. The WASP assessment reveals partial compensation 350 between changes in landborne and oceanborne fractions, and a larger spread in the effect of the climate feedback and the possibility of climate amplification of the ZEC.



4 Discussion and Conclusions

The Zero Emissions Commitment (ZEC) measures whether there is an increase or decrease in global mean surface temperature after carbon emissions cease at the time of net zero. This temperature change after net zero represents a transient response to past carbon emissions. The climate response after emissions cease relative to the pre-industrial era may then be viewed in terms of the global temperature rise associated with the amount of cumulative carbon emissions since the pre industrial, as measured by the Transient Climate Response to Cumulative CO₂ Emissions (TCRE), plus the subsequent transient temperature change due to prior carbon emissions, as measured by the Zero Emissions Commitment.

There are a wide range of climate processes that affect the transient climate response after net zero involving radiative forcing and the global cycling of carbon and heat. Gaining insight as to the relative importance of these different carbon and thermal processes in determining the ZEC is challenging due to their complexity and the effect of carbon and climate feedbacks (Palazzo Corner et al., 2023). Insight into the ZEC and the transient climate response after net zero may be obtained by examining two fundamental balances: the top of the atmosphere energy budget (Gregory et al., 2004), representing how the planet warms in response to radiative forcing and climate feedbacks, and how carbon emissions are partitioned between the carbon inventories of the atmosphere, land and ocean (Jones et al., 2013), and the effect of carbon feedbacks.

In order to gain a more mechanistic understanding of the controls of the ZEC, a normalised framework of the ZEC is set out that draws upon the top of the atmosphere energy budget and the change in carbon inventories including the effects of climate and carbon feedbacks. Firstly, a normalised ZEC is defined by the temperature change relative to the pre industrial divided by the temperature change at net zero. Secondly, the normalised ZEC is connected to the product of normalised thermal, radiative and carbon contributions, which are given respectively by the dependence of surface temperature to radiative forcing, the dependence of radiative forcing on atmospheric carbon and the change in atmospheric carbon. Each of the thermal, radiative and carbon contributions may then be interpreted in terms of the underlying mechanisms at work with the thermal contribution connected to the top of the atmosphere energy balance, the radiative balance connected to the logarithmic dependence of radiative forcing on atmospheric CO₂ and the carbon contribution connected to land and ocean carbon sinks.

The normalised ZEC framework may be applied to gain insight for a variety of carbon emission experiments. Here the normalised framework is applied to diagnostics of 9 Earth system models following the ZECMIP protocols with a 1% annual increase in atmospheric CO₂ until a 1000 PgC cumulative carbon emission (Jones et al., 2019; MacDougall et al., 2020) and a large ensemble of an efficient Earth system model (Goodwin et al., 2020), which is also applied to a scenario of a constant carbon emission over 100 years until a 1000 PgC cumulative carbon emission (Sanderson et al., 2024). In both sets of diagnostics, the ZEC response is controlled by a competition between a cooling contribution from a decline in atmospheric CO₂ due the effect of land and ocean carbon sinks versus a warming contribution from a strengthening in the thermal contribution. The thermal contribution representing the dependence of surface temperature on radiative forcing increases in time. This strengthening in thermal contribution is due to a larger fraction of the radiative forcing warming the surface and a smaller fraction being used for planetary heat uptake, which is consistent with a declining efficiency in global ocean heat uptake and



385 ventilation in time. The thermal contribution may also be augmented by the effect of climate feedbacks that can amplify the surface warming, such as by decreases in surface albedo or cloud albedo leading to an increase in solar absorption.

There are caveats and approximations in our ZECMIP analysis. The ZEC is a small signal:noise problem, and diagnosing this signal from Earth system models has inherent uncertainty (Borowiak et al., 2024). The diagnostics focus on a single model realisation and there are errors associated with how representative a single realisation is compared to a set of realisations by the same model. Our estimate of the radiative forcing from atmospheric CO₂ is based on a simple logarithmic closure and there are more accurate closures that may be applied. The estimate of the climate feedback is diagnosed from the radiative response from the energy balance at the top of the atmosphere divided by the changes in global surface temperature; this diagnostic is noisy on an interannual timescale. The Earth system models have inherent limitations in their inclusion and representation of climate and carbon processes, especially including uncertainties in cloud feedbacks, relatively coarse representation of ocean ventilation and a range of different land closures for carbon uptake. Finally, the ZECMIP analysis by design only includes the radiative forcing effects of atmospheric CO₂, while in reality there are additional radiative forcing effects from other greenhouse gases and aerosols.

The ZEC response of an individual climate model may be achieved in different ways, such as through a particularly large thermal or carbon contribution or through these thermal and carbon contributions reinforcing each other. For example, in the ZECMIP suite of models, a positive ZEC for CNRM-ESM2 is due to a particularly strong thermal contribution, while the positive ZEC for UKESM1 is due to a strong thermal contribution being reinforced by a more modest land carbon uptake, and the negative ZEC for NorESM2-LM is due to a large carbon uptake by the land and ocean.

Inter-model differences in the ZEC response may be understood in terms of the differences in the thermal and carbon contributions. Inter-model differences in the post-emission climate response for 9 Earth systems in ZECMIP are primarily determined by differences in the ocean uptake of heat and the land uptake of carbon, and to a lesser extent by the ocean uptake of carbon and climate feedbacks.

The diagnostics of the efficient Earth system model (WASP) reveal ensemble median responses for the ZEC and its contributions that are broadly similar to those of the ZECMIP diagnostics. The larger ensemble spread in WASP more clearly reveals a partial compensation between changes in the landborne and oceanborne fractions, as well as the ocean carbon sink dominating on longer timescales. The larger ensemble spread in WASP also reveals a wider range in the climate feedback parameter and how its temporal variation can lead to an amplification of surface warming and contributing to a positive ZEC. This difference suggests that the limited number of ZECMIP models may not be fully sampling the possible climate feedback responses compatible with historic warming. The efficient Earth system model does though show less variability in the heat uptake response, which may be due to a limitation that its ocean circulation is unchanging with time.

415 In comparison, inter-model differences in the TCRE are primarily controlled by thermal contributions (MacDougall et al., 2017) involving differences in climate feedbacks and ocean heat uptake, and then by carbon contributions involving the land uptake of carbon and a lesser extent the ocean uptake of carbon (Williams et al., 2020).

In summary, the normalised ZEC framework can provide insight as to why different Earth system models respond in different ways after carbon emissions cease. Gaining this process insight as to why the Earth system models have a wide spread in their



420 warming response after net zero is important for policy makers, since the combination of the climate metrics, the ZEC and TCRE, affects estimates of how much carbon may be emitted before exceeding a warming target.

Data availability. The data analysed for ZECMIP are openly available. The data input files are all based upon CMIP data that are available from the Earth System Grid Federation at <https://esgf-node.llnl.gov/projects/esgf-llnl/>. The data sets used for WASP will be achieved at Zenodo.



Table 1. Statistics for the model-mean climate response and the inter-model spread for time relative to net zero (t' , year): (a) Zero Emission Commitment (ZEC), $\Delta T(t) - \Delta T(t_{ZE})$, and the normalised ZEC, $\Delta T(t)/\Delta T(t_{ZE})$, where Δ is the change since the pre-industrial era and Δt_{ZE} is the temperature change at the time of net zero; (b) normalised contributions to the ZEC, $\Delta T/\Delta F$ is the thermal contribution, $\Delta F/\Delta I_A$ is the radiative contribution and ΔI_A is the atmospheric carbon contribution; (b) normalised contributions to the thermal contribution, $\Delta R/\Delta F$ is the fraction of radiative forcing escaping to space and λ^{-1} is the inverse of the climate feedback parameter; (d) Changes in the airborne, landborne and oceanborne fractions, $\Delta I_A/I_{em}$, $\Delta I_L/I_{em}$ and $\Delta I_O/I_{em}$. Model mean $\bar{x}$, inter-model standard deviation σ_x and coefficient of variation $\sigma_x/\bar{x}$ are provided for the 9 CMIP6 models. For rows (b) to (d), the terms with a large normalised spread are underlined.

time after net zero t'		25 y	50 y	90 y
(a) ZEC				
		$\Delta T(t) - \Delta T(t_{ZE})$ (K)		
$\bar{x} \pm \sigma_x$		-0.03±0.14	-0.10±0.19	-0.10± 0.24
normalised ZEC		$\Delta T(t)/\Delta T(t_{ZE})$		
$\bar{x} \pm \sigma_x$		0.97±0.09	0.93±0.11	0.93± 0.14
$\sigma_x/\bar{x}$		0.09	0.12	0.13
(b) normalised contributions to the ZEC				
thermal contribution		normalised $\Delta T(t)/\Delta F(t)$		
$\bar{x} \pm \sigma_x$		1.16±0.09	1.22±0.11	1.33±0.15
$\sigma_x/\bar{x}$		<u>0.08</u>	<u>0.09</u>	<u>0.11</u>
radiative contribution		normalised $\Delta F(t)/\Delta I_A(t)$		
$\bar{x} \pm \sigma_x$		1.06±0.01	1.09±0.02	1.11±0.02
$\sigma_x/\bar{x}$		0.01	0.02	0.02
atmospheric carbon contribution		normalised $\Delta I_A(t)$		
$\bar{x} \pm \sigma_x$		0.79±0.05	0.70±0.06	0.63±0.06
$\sigma_x/\bar{x}$		0.06	0.08	0.09
(c) normalised contributions to $\Delta T(t)/\Delta F(t)$				
fraction of radiative forcing returned to space		normalised $\Delta R(t)/\Delta F(t)$		
$\bar{x} \pm \sigma_x$		1.20±0.08	1.26±0.11	1.30±0.15
$\sigma_x/\bar{x}$		<u>0.07</u>	<u>0.09</u>	<u>0.12</u>
inverse climate feedback		normalised $\lambda(t)^{-1}$		
$\bar{x} \pm \sigma_x$		0.97±0.04	0.98±0.06	1.03 ±0.05
$\sigma_x/\bar{x}$		0.04	0.06	0.05
(d) carbon changes				
time after net zero t'	0 y	25 y	50 y	90 y
airborne fraction		$\Delta I_A(t)/I_{em}(t_{ZE})$		
$\bar{x} \pm \sigma_x$	0.52±0.03	0.42±0.04	0.38±0.05	0.34±0.05
$\sigma_x/\bar{x}$	0.06	0.10	0.12	0.14
landborne fraction		$\Delta I_L(t)/I_{em}(t_{ZE})$		
$\bar{x} \pm \sigma_x$	0.26±0.04	0.32±0.06	0.34±0.07	0.35±0.08
$\sigma_x/\bar{x}$	<u>0.17</u>	<u>0.20</u>	<u>0.21</u>	<u>0.21</u>
oceanborne fraction		$\Delta I_O(t)/I_{em}(t_{ZE})$		
$\bar{x} \pm \sigma_x$	0.22±0.03	0.26±0.03	0.28±0.04	0.31±0.05
$\sigma_x/\bar{x}$	0.12	0.13	0.13	0.14



425 5 Appendix

The Appendices includes:

Table A1 displays the diagnostics for the individual 9 Earth system models making up ZECMIP;

Tables A2 and A3 displays the diagnostics for the large ensemble using WASP for the 1pctCO2 and flat10 experiments;

Figure A1 displays the climate and carbon variables for a large ensemble following the flat10 experiment for up to 400 years

430 after net zero for WASP, including the median response and shading for the 5% to 95% spread in the ensembles.

Table A1. Climate model response post emissions at 50 years (40-59 year average) after net zero: (a) Zero Emission Commitment (ZEC), $\Delta T(t) - \Delta T(t_{ZE})$, and normalised ZEC, $\Delta T(t)/\Delta T(t_{ZE})$, where Δ is the change since the pre-industrial era and Δt_{ZE} is the temperature change at net zero; (b) Normalised changes in the land and ocean carbon inventories, ΔI_L and ΔI_O ; (c) normalised contributions to the ZEC, $\Delta T/\Delta F$ is the thermal contribution, $\Delta F/\Delta I_A$ is the radiative contribution and ΔI_A is the atmospheric carbon contribution; (d) normalised contributions to the thermal contribution, $\Delta R/F$ is the fraction of radiative forcing escaping to space and λ^{-1} is the inverse of the climate feedback parameter. Model mean $\bar{x}$, inter-model standard deviation σ_x and coefficient of variation $\sigma_x/\bar{x}$ are included.

Model	(a) ZEC metrics		(b) normalised carbon changes			
	ZEC	norm. ZEC	airborne fraction	landborne fraction	oceanborne fraction	
	$\Delta T(t) - \Delta T(t_{ZE})$ (K)	$\Delta T(t)/\Delta T(t_{ZE})$	$\Delta I_A(t)/I_{em}(t_{ZE})$	$\Delta I_L(t)/I_{em}(t_{ZE})$	$\Delta I_O(t)/I_{em}(t_{ZE})$	
ACCESS-ESM1.5	0.00	1.0	0.46	0.20	0.34	
CanESM5	-0.12	0.94	0.33	0.43	0.24	
CESM2	-0.29	0.87	0.38	0.35	0.27	
CNRM-ESM2	0.05	1.03	0.39	0.40	0.21	
GFDL-ESM2	-0.27	0.79	0.31	0.40	0.29	
MIROC-ESL2	-0.09	0.93	0.37	0.35	0.28	
MPI-ESM-2-LR	-0.12	0.93	0.36	0.36	0.28	
NorESM2-LM	-0.32	0.77	0.35	0.34	0.31	
UKESM1	0.28	1.11	0.43	0.27	0.30	
$\bar{x} \pm \sigma_x$	-0.10 ± 0.19	0.93 ± 0.11	0.38 ± 0.05	0.34 ± 0.07	0.28 ± 0.04	
$\sigma_x/\bar{x}$		0.12	0.12	0.21	0.13	

Model	(c) norm. contrib. to ZEC		(d) thermal contribution to ZEC		inverse climate feedback norm. $\lambda(t)^{-1}$
	thermal	radiative	carbon	fraction of forcing returned to space	
	norm. $\Delta T(t)/\Delta F(t)$	norm. $\Delta F(t)/\Delta I_A(t)$	norm. $\Delta I_A(t)$	norm. $\Delta R(t)/\Delta F(t)$	
ACCESS-ESM1.5	1.19	1.06	0.79	1.17	1.02
CanESM5	1.34	1.10	0.64	1.33	1.00
CESM2	1.17	1.10	0.67	1.28	0.91
CNRM-ESM2	1.36	1.09	0.69	1.47	0.92
GFDL-ESM2	1.11	1.10	0.65	1.27	0.90
MIROC-ESL2	1.21	1.08	0.72	1.14	1.06
MPI-ESM-2-LR	1.16	1.07	0.75	1.19	0.98
NorESM2-LM	1.09	1.11	0.63	1.17	0.93
UKESM1	1.38	1.07	0.75	1.32	1.05
$\bar{x} \pm \sigma_x$	1.22 ± 0.11	1.09 ± 0.02	0.70 ± 0.06	1.26 ± 0.11	0.98 ± 0.06
$\sigma_x/\bar{x}$	0.09	0.02	0.08	0.09	0.06



Table A2. Climate model response for the large ensemble, WASP for climate and carbon variables following a 1pctCO₂ experiment with a cumulative carbon emission of 1000 PgC. Quantities given are the median, 5th percentile and 95th percentile at years 50, 100 and 400 after net zero.

	t'	50 years	100 years	400 years
(a) temperature changes				
temperature rise	ΔT (K)	1.09 (0.77 to 1.77)	1.10 (0.71 to 2.14)	0.98 (0.60 to 2.54)
ZEC	$\Delta T(t) - \Delta T(t_{ZE})$ (K)	-0.10 (-0.47 to 0.43)	-0.09 (-0.56 to 0.82)	-0.22 (-0.71 to 1.19)
normalised ZEC	$\Delta T(t)/\Delta T(t_{ZE})$	0.92 (0.63 to 1.36)	0.92 (0.57 to 1.66)	0.82 (0.47 to 1.96)
(b) carbon inventory changes				
atmospheric carbon	ΔI_A (PgC)	310 (252 to 396)	288 (235 to 374)	226 (189 to 301)
land carbon	ΔI_L (PgC)	371 (219 to 476)	333 (177 to 435)	252 (80 to 349)
ocean carbon	ΔI_O (PgC)	320 (262 to 399)	380 (316 to 465)	522 (453 to 629)
(c) radiative balance				
radiative forcing	ΔF (Wm^{-2})	2.26 (1.87 to 2.80)	2.14 (1.76 to 2.66)	1.74 (1.46 to 2.23)
radiative response	$-\Delta R$ (Wm^{-2})	1.89 (1.52 to 2.35)	1.84 (1.50 to 2.25)	1.66 (1.40 to 2.05)
planetary heat uptake	ΔN (Wm^{-2})	0.35 (0.15 to 0.73)	0.25 (0.08 to 0.68)	0.07 (0.03 to 0.26)
(d) normalised contributions to ZEC				
thermal contribution	Normalised $\Delta T/\Delta F$	1.28 (0.94 to 1.77)	1.37 (0.91 to 2.17)	1.50 (0.92 to 2.92)
radiative contribution	Normalised $\Delta F/\Delta I_A$	1.09 (1.06 to 1.11)	1.11 (1.07 to 1.13)	1.15 (1.10 to 1.19)
carbon contribution	Normalised ΔI_A	0.66 (0.60 to 0.74)	0.60 (0.55 to 0.74)	0.47 (0.43 to 0.62)
(e) Normalised thermal components				
Radiative response fraction	Normalised $1 - \Delta N/\Delta F$	1.11 (0.97 to 1.28)	1.15 (0.98 to 1.33)	1.26 (1.15 to 1.40)
Inverse climate feedback	Normalised λ^{-1}	1.15 (0.77 to 1.79)	1.18 (0.73 to 2.18)	1.19 (0.72 to 2.45)
(f) Partition of emissions				
Airborne fraction	$\Delta I_A(t)/I_{em}$	0.31 (0.25 to 0.40)	0.29 (0.24 to 0.37)	0.23 (0.19 to 0.30)
Landborne fraction	$\Delta I_L(t)/I_{em}$	0.37 (0.22 to 0.48)	0.33 (0.18 to 0.44)	0.25 (0.08 to 0.35)
Oceanborne fraction	$\Delta I_O(t)/I_{em}$	0.32 (0.26 to 0.40)	0.38 (0.32 to 0.46)	0.52 (0.45 to 0.63)



Table A3. Climate model response for the large ensemble, WASP for climate and carbon variables following a flat10 experiment with a cumulative carbon emission of 1000 PgC. Quantities given are the median, 5th percentile and 95th percentile at years 50, 100 and 400 after net zero.

	t'	50 years	100 years	400 years
(a) temperature changes				
temperature rise	ΔT (K)	1.10 (0.73 to 1.92)	1.09 (0.69 to 2.24)	0.97 (0.59 to 2.53)
ZEC	$\Delta T(t) - \Delta T(t_{ZE})$ (K)	-0.06 (-0.25 to 0.41)	-0.07 (-0.31 to 0.73)	-0.19 (-0.46 to 1.02)
normalised ZEC	$\Delta T(t)/\Delta T(t_{ZE})$	0.95 (0.77 to 1.29)	0.94 (0.71 to 1.50)	0.83 (0.59 to 1.68)
(b) carbon inventory changes				
atmospheric carbon	ΔI_A (PgC)	299 (242 to 383)	280 (228 to 366)	224 (186 to 297)
land carbon	ΔI_L (PgC)	350 (199 to 455)	320 (159 to 421)	249 (78 to 345)
ocean carbon	ΔI_O (PgC)	349 (290 to 430)	399 (336 to 485)	527 (459 to 636)
(c) radiative balance				
radiative forcing	ΔF (Wm^{-2})	2.20 (1.81 to 2.73)	2.09 (1.71 to 2.61)	1.72 (1.44 to 2.21)
radiative response	$-\Delta R$ (Wm^{-2})	1.86 (1.51 to 2.28)	1.82 (1.50 to 2.23)	1.64 (1.39 to 2.03)
planetary heat uptake	ΔN (Wm^{-2})	0.30 (0.12 to 0.70)	0.22 (0.08 to 0.63)	0.07 (0.03 to 0.24)
(d) normalised contributions to ZEC				
thermal contribution	Normalised $\Delta T/\Delta F$	1.17 (0.98 to 1.48)	1.24 (0.97 to 1.75)	1.34 (0.98 to 2.24)
radiative contribution	Normalised $\Delta F/\Delta I_A$	1.06 (1.04 to 1.07)	1.07 (1.04 to 1.09)	1.10 (0.07 to 1.14)
carbon contribution	Normalised ΔI_A	0.77 (0.73 to 0.86)	0.71 (0.67 to 0.86)	0.57 (0.51 to 0.72)
(e) Normalised thermal components				
Radiative response fraction	Normalised $1 - \Delta N/\Delta F$	1.08 (0.98 to 1.15)	1.12 (1.00 to 1.19)	1.20 (1.13 to 1.29)
Inverse climate feedback	Normalised λ^{-1}	1.09 (0.87 to 1.53)	1.10 (0.84 to 1.73)	1.10 (0.83 to 1.83)
(f) Partition of emissions				
Airborne fraction	$\Delta I_A(t)/I_{em}$	0.30 (0.24 to 0.38)	0.28 (0.23 to 0.37)	0.22 (0.19 to 0.30)
Landborne fraction	$\Delta I_L(t)/I_{em}$	0.35 (0.20 to 0.46)	0.32 (0.16 to 0.42)	0.25 (0.08 to 0.35)
Oceanborne fraction	$\Delta I_O(t)/I_{em}$	0.35 (0.29 to 0.43)	0.40 (0.34 to 0.49)	0.53 (0.46 to 0.63)




Figure A1. Temporal evolution of carbon and climate variables relative to the time of net zero (years) for the flat10 experiment from the WASP simulations: (a) cumulative carbon emissions, $\Delta I_{em}(t')$ (PgC); (b) surface temperature change relative to the pre industrial, $\Delta T(t')$ (K); (c, e and g) the change in atmospheric carbon inventory, $\Delta I_A(t')$, land carbon inventory, $\Delta I_L(t')$ and ocean carbon inventory, $\Delta I_O(t')$, (all in PgC); and (d, f and h) the change in radiative forcing, $\Delta F(t')$, radiative response, $-\Delta R(t')$, and planetary heat uptake, $\Delta N(t')$ (all in W m^{-2}).

Author contributions. RW developed the framework and wrote the first draft of the paper. RW diagnosed the ZECMIP models using model data provided by AM. CJ advised on the ZECMIP protocols and PC advised on the climate feedback diagnostics. PG performed the WASP integrations and diagnostics. All authors edited and commented on the manuscript.

Competing interests. There are no competing interests.

435 *Acknowledgements.* The authors acknowledge the World Climate Research Programme's Working Group on Coupled Modelling responsible for CMIP. RGW acknowledge support from the UK Natural Environmental Research Council, NE/N009789/1 and NE/W009501/1. PC was supported by UK Natural Environmental Research Council grants NE/V012045/1, NE/T006250/1 and EP/Y036123/1. CDJ was supported by the Joint UK BEIS/Defra Met Office Hadley Centre Climate Programme (GA01101) and the European Union's Horizon 2020 research and innovation programme under Grant Agreement No 101003536 (ESM2025 - Earth System Models for the Future).



440 References

- Allen, M. R., Frame, D. J., Huntingford, C., Jones, C. D., Lowe, J. A., Meinshausen, M., and Meinshausen, N.: Warming caused by cumulative carbon emissions towards the trillionth tonne, *Nature*, 458, 1163–1166, 2009.
- Allen, M. R., Friedlingstein, P., Girardin, C. A., Jenkins, S., Malhi, Y., Mitchell-Larson, E., Peters, G. P., and Rajamani, L.: Net zero: science, origins, and implications, *Annual Review of Environment and Resources*, 47, 849–887, 2022.
- 445 Andrews, T., Gregory, J. M., Webb, M. J., and Taylor, K. E.: Forcing, feedbacks and climate sensitivity in CMIP5 coupled atmosphere-ocean climate models, *Geophys. Res. Letts*, 39, L09 712, 2012.
- Borowiak, A., King, A. D., Brown, J. R., Jones, C. D., Ziehn, T., Meinshausen, M., and Cassidy, L.: Projected global temperature changes after net zero are small but significant, *Geophysical Research Letters*, 51, e2024GL108 654, 2024.
- Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichet, T., Friedlingstein, P., Gao, X., Gutowski, W. J., Johns, T., Krinner, G.,
 450 et al.: Long-term climate change: projections, commitments and irreversibility, in: *Climate Change 2013-The Physical Science Basis: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 1029–1136, Cambridge University Press, 2013.
- Ehlert, D., Zickfeld, K., Eby, M., and Gillett, N.: The sensitivity of the proportionality between temperature change and cumulative CO₂ emissions to ocean mixing, *Journal of Climate*, 30, 2921–2935, 2017.
- 455 Follows, M. J., Ito, T., and Dutkiewicz, S.: On the solution of the carbonate chemistry system in ocean biogeochemistry models, *Ocean Modelling*, 12, 290–301, <https://doi.org/https://doi.org/10.1016/j.ocemod.2005.05.004>, 2006.
- Forster, P. M., Andrews, T., Good, P., Gregory, J. M., Jackson, L. S., and Zelinka, M.: Evaluating adjusted forcing and model spread for historical and future scenarios in the CMIP5 generation of climate models, *Journal of Geophysical Research: Atmospheres*, 118, 1139–1150, 2013.
- 460 Gillett, N. P., Arora, V. K., Matthews, D., and Allen, M. R.: Constraining the ratio of global warming to cumulative CO₂ emissions using CMIP5 simulations, *J. Climate*, 26, 6844–6858, 2013.
- Goodwin, P.: How historic simulation–observation discrepancy affects future warming projections in a very large model ensemble, *Climate Dynamics*, 47, 2219–2233, 2016.
- Goodwin, P.: On the time evolution of climate sensitivity and future warming, *Earth’s Future*, 6, 1336–1348,
 465 <https://doi.org/10.1029/2018EF000889>, 2018.
- Goodwin, P., Williams, R. G., and Ridgwell, A.: Sensitivity of climate to cumulative carbon emissions due to compensation of ocean heat and carbon uptake, *Nature Geosci.*, 8, 29–34, 2015.
- Goodwin, P., Leduc, M., Partanen, A.-I., Matthews, H. D., and Rogers, A.: A computationally efficient method for probabilistic local warming projections constrained by history matching and pattern scaling, demonstrated by WASP-LGRTC-1.0, *Geoscientific Model Development*,
 470 13, 5389–5399, 2020.
- Gregory, J. M., Ingram, W. J., Palmer, M. A., Jones, G. S., Stott, P. A., Thorpe, R. B., Lowe, J. A., Johns, T. C., and Williams, K. D.: A new method for diagnosing radiative forcing and climate sensitivity, *Geophys Res Lett.*, 31, L03 205, <https://doi.org/10.1029/2003GL018747>, 2004.
- Jones, C., Robertson, E., Arora, V., Friedlingstein, P., Shevliakova, E., Bopp, L., Brovkin, V., Hajima, T., Kato, E., Kawamiya, M., et al.:
 475 Twenty-first-century compatible CO₂ emissions and airborne fraction simulated by CMIP5 Earth system models under four representative concentration pathways, *Journal of Climate*, 26, 4398–4413, 2013.



- Jones, C. D. and Friedlingstein, P.: Quantifying process-level uncertainty contributions to TCRE and Carbon Budgets for meeting Paris Agreement climate targets, *Environmental Research Letters*, <http://iopscience.iop.org/10.1088/1748-9326/ab858a>, 2020.
- Jones, C. D., Frölicher, T. L., Koven, C., MacDougall, A. H., Matthews, H. D., Zickfeld, K., Rogelj, J., Tokarska, K. B., Gillett, N. P., Ilyina, T., et al.: The Zero Emissions Commitment Model Intercomparison Project (ZECMIP) contribution to C4MIP: quantifying committed climate changes following zero carbon emissions, *Geoscientific Model Development*, 12, 4375–4385, 2019.
- Katavouta, A., Williams, R. G., Goodwin, P., and Roussenov, V.: Reconciling atmospheric and oceanic views of the Transient Climate Response to Emissions, *Geophysical Research Letters*, 45, 6205–6214, 2018.
- Knutti, R. and Hegerl, G. C.: The equilibrium sensitivity of the Earth’s temperature to radiation changes, *Nature Geosci.*, 1, 735–743, 2008.
- MacDougall, A. H.: The Transient Response to Cumulative CO₂ Emissions: a review, *Current Climate Change Reports*, 2, 39–47, 2016.
- MacDougall, A. H., Swart, N. C., and Knutti, R.: The uncertainty in the transient climate response to cumulative CO₂ emissions arising from the uncertainty in physical climate parameters, *Journal of Climate*, 30, 813–827, 2017.
- MacDougall, A. H., Frölicher, T. L., Jones, C. D., Rogelj, J., Matthews, H. D., Zickfeld, K., Arora, V. K., Barrett, N. J., Brovkin, V., Burger, F. A., et al.: Is there warming in the pipeline? A multi-model analysis of the Zero Emissions Commitment from CO₂, *Biogeosciences*, 17, 2987–3016, 2020.
- Matthews, H. D. and Zickfeld, K.: Climate response to zeroed emissions of greenhouse gases and aerosols, *Nature Climate Change*, 2, 338–341, 2012.
- Matthews, H. D., Zickfeld, K., Knutti, R., and Allen, M. R.: Focus on cumulative emissions, global carbon budgets and the implications for climate mitigation targets, *Environmental Research Letters*, 13, 010 201, 2018.
- Matthews, H. R., Gillett, N. P., Stott, P. A., and Zickfeld, K.: The proportionality of global warming to cumulative carbon emissions, *Nature*, 459, 829–833, 2009.
- Palazzo Corner, S., Siegert, M., Ceppi, P., Fox-Kemper, B., Frölicher, T. L., Gallego-Sala, A., Haigh, J., Hegerl, G. C., Jones, C. D., Knutti, R., et al.: The Zero Emissions Commitment and climate stabilization, *Frontiers in Science*, 1, 1170 744, 2023.
- Sanderson, B. M., Brovkin, V., Fisher, R., Hohn, D., Ilyina, T., Jones, C., Koenigk, T., Koven, C., Li, H., Lawrence, D., et al.: flat10MIP: An emissions-driven experiment to diagnose the climate response to positive, zero, and negative CO₂ emissions, *EGUsphere*, 2024, 1–39, 2024.
- Solomon, S., Plattner, G.-K., Knutti, R., and Friedlingstein, P.: Irreversible climate change due to carbon dioxide emissions, *Proc. Natl. Acad. Sci. USA*, 106, 1704–1709, 2009.
- Williams, R. G., Goodwin, P., Roussenov, V. M., and Bopp, L.: A framework to understand the Transient Climate Response to Emissions, *Environmental Research Letters*, 11, <https://doi.org/10.1088/1748-9326/11/1/015003>, 2016.
- Williams, R. G., Roussenov, V., Goodwin, P., Resplandy, L., and Bopp, L.: Sensitivity of global warming to carbon emissions: Effects of heat and carbon uptake in a suite of Earth system models, *Journal of Climate*, 30, 9343–9363, 2017.
- Williams, R. G., Ceppi, P., and Katavouta, A.: Controls of the transient climate response to emissions by physical feedbacks, heat uptake and carbon cycling, *Environmental Research Letters*, 15, 0940c1, 2020.
- Zickfeld, K., Eby, M., Matthews, H. D., and Weaver, A. J.: Setting cumulative emissions targets to reduce the risk of dangerous climate change, *PNAS*, 106, 16 129–16 134, 2009.