



Future intensification of compound and cascading drought and heatwave risks in Europe

Samuel Jonson Sutanto¹, Confidence Duku^{1,2}, Merve Gülveren³, Rutger Dankers², and Spyridon Paparrizos¹

¹Earth Systems and Global Change Group, Environmental Sciences Department, Wageningen University and Research, Droevendaalsesteeg 3a, 6708PB, Wageningen, the Netherlands

²Climate Resilience Group, Wageningen Environmental Research, Wageningen, the Netherlands

³Directorate of Climate Change, Ministry of Environment, Urbanization, and Climate Change, Ankara, Türkiye

Correspondence: Samuel Sutanto (samuel.sutanto@wur.nl)

Abstract. The risks of extreme weather events, such as droughts and heatwaves, are expected to rise across Europe due to global warming, leading to more severe and worsening impacts. These impacts become even more pronounced when compound and cascading (CnC) drought and heatwave hazards occur. Yet, most studies on drought and heatwave have focused on single hazard rather than their impacts. This study aims to identify the future characteristics of both single and compound drought and heatwave hazards across Europe. More specifically, we analyzed changes in the total number of events, average duration, total duration, and frequency. Droughts were identified using the Standardized Soil Moisture Index (SMI) and heatwaves were detected using the Variable Threshold Method (VTM). Both hazards were assessed using bias corrected simulations from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP) models from 1953 to 2014 for historical period and from 2039 to 2100 for future climate scenarios under SSP1-2.6 and SSP5-8.5. Furthermore, we employ a machine learning (ML) approach to project the impacts of droughts and heatwaves, using Germany as a case study. The ML models were developed using hazard characteristics as predictors and drought and heatwave impact data as response variables. Results indicate that the number, duration, and frequency of both drought and heatwave events are projected to increase under SSP1-2.6, with even higher increase for SSP5-8.5, not only when analyzed independently but also as CnC hazards. This applies not only in the south but also across multiple other European regions. Drought hotspots were identified in the West Europe, with projections showing an expansion toward the South and East under SSP1-2.6, and across nearly all of Europe under SSP5-8.5 except for the northern regions. Heatwave hotspots were primarily located in eastern and southern Europe, particularly in Russia, Italy, and Portugal. Future scenarios suggest that southern Europe will remain a key hotspot for heatwaves. The occurrence of compound drought and heatwave events was projected to increase sixfold compared to the reference period, while cascading drought and heatwave events might rise by up to 3.5 times under SSP5-8.5. Additionally, results also reveal that drought impacts on economic, non-economic, and ecosystem are projected to double in Germany, while heatwave impacts on human health and mortality may increase ninefold by 2100. Our findings highlight the need to consider CnC hazards and show once more the urgency of climate mitigation in limiting impacts across multiple sectors.



1 Introduction

- 25 The occurrence of extreme weather events, such as droughts and heatwaves, has become more frequent and severe in recent decades due to climate change (Mukherjee and Mishra, 2021; Seneviratne et al., 2021). The impacts of these events are intensified when droughts and heatwaves occur simultaneously (compound events) and/or consecutively (cascading events) (Fischer et al., 2007; Van Lanen et al., 2016; Rakovec et al., 2022). For example, the major European droughts of 2003, 2010, 2018-2020, and 2022 were particularly severe and costly due to their combination with heatwaves and other hazards such as wildfires
- 30 (Robine et al., 2008; Ionita et al., 2017; Blauhut et al., 2022; Biella et al., 2024). These events heavily impacted sectors such as agriculture, energy, human health, and public water supply (Cammalleri et al., 2020; Biella et al., 2024). Recent drought events across Europe have resulted in estimated economic losses of approximately 9 billion EUR, with the highest losses recorded in Spain (1.5 billion EUR/year), followed closely by Italy (1.4 billion EUR/year) and France (1.2 billion EUR/year) (Cammalleri et al., 2020).
- 35 The impacts of drought and heatwave hazards, as single hazards and as compound and cascading hazards (CnC), are expected to increase under global warming due to the increase of temperature and reduced precipitation in many land surfaces (Seneviratne et al., 2021). For example, studies comparing the impacts of the 2003 and 2018 droughts and heatwaves on European ecosystems and forests concluded that the 2018 event was more extreme than that of 2003 (Buras et al., 2020; Schuldt et al., 2020). As a result, the negative impacts of the 2018 hot drought on ecosystems were not only stronger but also affected
- 40 a larger area. Additionally, Cammalleri et al. (2020) projected that agriculture and energy sectors will experience the highest impact compared to others, while Naumann et al. (2021) estimated that economic losses due to drought in Europe could rise to 65 billion EUR by 2100. The severity of these impacts, especially when combined with other hazards, call the urgent need to manage and minimize the impacts associated with future CnC drought and heatwave events as well as to develop effective adaptation strategies.
- 45 Several studies have analyzed the occurrence of droughts and heatwaves in Europe under global warming, but relatively few have discussed the characteristics of CnC drought and heatwave events, such as duration and frequency (Samaniego et al., 2018; Spinoni et al., 2018; Tripathy et al., 2023; Tripathy and Mishra, 2023). In addition, these studies primarily focused on either single or compound events, neglecting cascading events. Importantly, projections of the impacts of droughts and heatwaves in various sectors, beyond just the hazards themselves, are generally lacking. Naumann et al. (2021) study is pioneering in
- 50 analyzing future drought impacts in terms of economic losses by integrating damage functions with hazard and exposure components. Another promising approach for impact prediction is by applying machine learning (ML) algorithms. Previous studies have utilized ML approaches to link reported drought impacts with hazards and to forecast drought impacts (Stagge et al., 2015; Bachmair et al., 2017; Sutanto et al., 2019a). However, the application of ML to predict future drought and heatwave impacts, as opposed to hazards, has so far been limited.



55 In this study, we analyze the characteristics of future agricultural and hydrological droughts represented by soil moisture
 drought and heatwave events including their CnC events across Europe. Furthermore, we also aim to predict the impacts of
 droughts and heatwaves under both low and high Shared Socioeconomic Pathways scenarios (SSP1-2.6 and 5-8.5, respec-
 tively). In our study, we define compound drought and heatwave (CDH) hazards if these events occur simultaneously on the
 same day and same location (Leonard et al., 2014; Liu and Huang, 2015) while cascading drought and heatwave (CaDH) is
 60 defined if these hazards occur successively or cumulatively over time without being interrupted by a zero-hazard day (Sutanto
 et al., 2019b; Vitolo et al., 2019).

The paper is structured as follows: Section 2 describes the methodology used in this study, Section 3 presents the results,
 Section 4 discusses the findings, and finally, Section 5 concludes the study. Section 2 is further divided into six sub sections,
 describing the hydrometeorological data used in the study (Section 2.1), the drought and heatwave impact database (Section
 65 2.2), methods for identifying droughts and heatwaves (Section 2.3), identification of CnC events (Section 2.4), drought and
 heatwave characteristics (Section 2.5), and impact projections using machine learning (Section 2.6). In Section 3, we present
 the characteristics of single hazards in Section 3.1, CnC hazard characteristics in Section 3.2, a regional summary of both
 single and CnC hazards across Europe in Section 3.3, and projections of drought and heatwave impacts in Section 3.4.

2 Methods

70 2.1 Hydrometeorological data

Soil moisture data were obtained from the ERA5-Land and the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP)
 (Warszawski et al., 2013). ERA5-Land monthly soil moisture data were downloaded from the Copernicus Data Store (CDS),
 while in ISIMIP, soil moisture data (ISIMIP3b) were generated using the CWatM hydrological model (Burek et al., 2020),
 driven by five Coupled Model Intercomparison Project Phase 6 (CMIP6) global climate models: GFDL-ESM4, IPSL-CM6A-
 75 LR, MPI-ESM1-2-HR, MRI-ESM2-0, and UKESM1-0-LL. The ISIMIP soil moisture data were then bias corrected using the
 delta method, with ERA5-Land as a benchmark (Hawkins et al., 2013). ISIMIP model data, originally at a spatial resolution of
 0.5° x 0.5°, were downscaled to 0.1° x 0.1° using bilinear interpolation approach to match with ERA5-Land resolution.

ISIMIP hourly near-surface temperature data are available for five CMIP6 global climate models. These hourly temperature
 data were then converted to daily maximum and minimum temperature for heatwave analysis. The ISIMIP temperature data
 80 have already been bias corrected and therefore, we did not perform bias correction analysis (Lange and Büchner, 2021). Same
 as soil moisture data, a bilinear interpolation approach was applied to downscale the temperature data to a 0.1° x 0.1° spatial
 resolution to ensure consistency with ERA5 Land.

Heatwave and drought analyses were performed individually for each model. Changes in dry hazard characteristics as single
 and compound events are determined by calculating the difference between future and historical events (future minus reference
 85 period). For drought, we used historical data from January 1953 to December 2014 (12 months x 62 years) as reference period
 and from January 2039 to December 2100 for future period (62 years). Same period was chosen for heatwave analysis but



we only considered extended summer period from May to October (184 days x 62 years). Detailed data used in the study is presented in the Supplementary Table S1.

2.2 Drought and heatwave impact databases

90 Drought and heatwave impact databases were compiled from the European Drought Impact Inventory (EDII) (Stahl et al., 2016), the international disaster database (EM-DAT) (Jonkman, 2005), and data mined from scientific and grey literature in English. We followed the format of EDII to compile the heatwave and used the same impact categories. Drought impact data was then grouped into three different sectors, which are economic, non-economic, and ecosystem sectors (Biella et al., 2024). The economic sector consists of agriculture and livestock, forestry, aquaculture and fisheries, energy and industry, waterborne
 95 transportation, and tourism and recreation impacts. The non-economic sector includes public water supply, water quality, air quality, health and public safety, and water access conflicts. The ecosystem sector covers impacts on freshwater ecosystems, terrestrial ecosystems, soil systems, and wildfires. This categorization was based on the drought impact divisions established by the Dita network, IAHS group (Biella et al., 2024).

Heatwave impacts reported from data mined include agriculture, air quality, wildfire, and economic in addition to the excess
 100 mortality statistics often reported. The collected impact data other than excess mortality, however, are too limited and can be combined to human impacts. For instance, wildfires triggered by heatwaves are often also reported having an impact of mortality. Thus, we decided to combine this into human impact. Due to limitations in the reported drought and heatwave impact data, this study only focuses on impact analysis for Germany.

2.3 Drought and heatwave indexes

105 Droughts in soil moisture were identified using the Standardized Soil Moisture Index (SMI) following the construction of the Standardized precipitation Index (SPI) (McKee et al., 1993). The SMI provides a measure of soil moisture dryness by quantifying the deviation from the long-term mean, i.e. number of standard deviations. The SMI was calculated by fitting a probabilistic distribution on monthly soil moisture data. To compute the SMI, the monthly historical soil moisture data was transformed into 12 distributions, corresponding to the month of the year. Gamma distribution was employed in this study and
 110 is described by two parameters: α (the shape parameter) and β (the inverse scale parameter). The gamma distribution has quite a flexible shape parameter, which is suitable for a wide range of drought application in EU (Stagge et al., 2015; Sutanto and Van Lanen, 2021). These distribution parameters then were used to calculate observed and future drought events (for further details, refer to Sutanto et al. (2020a)). The use of distribution parameters derived from the historical period to estimate future droughts implies that no adaptation to climate change is assumed.

115 Heatwave events were identified using the daily variable threshold method (VTM) derived from the historical data (Sutanto and Van Lanen, 2021). At each grid point, the threshold of 90th percentile of the daily maximum and minimum temperature for the historical period was calculated and nine days centered moving window approach was applied to consider the temporal variation of temperature. We identified a heatwave event if both maximum and minimum temperatures exceed the thresholds for at least 3 days during the extended summer period, from May to October (Lavaysse et al., 2018; Sutanto et al., 2019b). Two



120 successive heatwave events are considered independent if they are separated by a minimum of four days, otherwise, it is taken as one single event (Mukherjee and Mishra, 2021). Similar to drought, the threshold levels derived from historical data were applied to analyze future heatwaves, meaning that no adaptation to climate change is assumed.

2.4 Compound and cascading (CnC) events

To analyse the CnC events, binary maps consist of the number 1 and 2 for heatwave and drought events, respectively, were
 125 generated if the month is identified as drought or heatwave month, and 0 (no hazard) otherwise. The next step was overlapping the monthly maps of individual hazards by aggregate the grid cell values. Compound event is identified if the summed value is 3 (1+2), meaning that heatwave and drought occurred at the same time and place. A cascading event occurs if two different events as single-single (drought-heatwave or vice versa) or single-compound happen sequentially without being interrupted by a zero-hazard month. For instance, the binary time series of the events like “0-1-2-2-0”, “0-2-2-1-0”, “0-2-3-1-0”, and “0-1-3-0”
 130 indicate cascading events of heatwaves-drought, drought-heatwaves, drought-compound drought heatwaves, and heatwaves-compound drought heatwaves, respectively. Previous studies by Sutanto et al. (2019b) and Vitolo et al. (2019) provide detailed descriptions of the CnC events. Supplementary Figure S1 shows examples of CDH events from May to October 2018 and from June to August 2019, and CaDH events from May to November 2018 and from June to October 2019 in Germany. There were two drought events occurred in February 2018 and April 2019.

135 2.5 Extreme event characteristics

We defined the characteristics of drought and heatwave hazards both as single and CnC as the total number of events, average duration of events, total duration of events, and frequency of events. The total number of events is defined as total hazard events in a particular period, either the reference or future period, summed across the 62 years. The total duration of hazard events is defined as the total duration of hazards during the analyzed period summed across the 62 years, measured in months for
 140 droughts, and days for heatwaves. The average duration of hazard events is calculated by dividing the total number of events with the total duration of events. Lastly, the frequency of events is defined as the average number of hazard events in a year. For CnC analysis, we assigned the heatwave binary month if there was at least one heatwave event in that particular month (see previous sub-section). Supplementary Table S2 provides detailed information of dry hazard characteristics.

2.6 Hazard projections using machine learning

145 Drought index derived from the SMI, and the updated drought impact data for Germany were utilized to develop drought impact prediction models using an Extreme Gradient Boosting (XGBoost) ML approach (Friedman, 2001; Mardian et al., 2023). The ML models to predict drought impacts on economic sector, non-economic sector, and ecosystem sector were trained using soil moisture data, SMI, and drought duration as the predictors and drought impact data as the response variable. The reported drought impact data were categorized into economic sector, non-economic sector and ecosystem sector (see previous
 150 sub-section). However, for heatwave impacts, for which there was a limited data record, the ML model was developed using



reported impact data on human mortality as the response variable and heatwave intensity, duration, and number of events as the predictors. The drought and heatwave impact data were initially pre-processed by ensuring coverage for all months of the year, filling in missing months with zero values to indicate the absence of extreme events. This was achieved by creating a multi-index using the 'year' and 'month' columns and expanding the data to span all possible combinations. The resulting dataset was merged with the original data to ensure a comprehensive timeseries, with missing values filled appropriately.

For both drought and heatwave reported data, the Synthetic Minority Over-sampling Technique (SMOTE) (Chawla et al., 2002; Lemaître et al., 2017) was used to address inherent class imbalances that could potentially lead to biased models. Class imbalance poses a significant challenge to many ML algorithms, particularly those based on classification trees or other forms of supervised learning. When the target variable's distribution is highly skewed, the model tends to be biased towards predicting the majority class, as it can achieve a superficially high accuracy simply by ignoring the minority class. Droughts and heatwaves are essentially rare events occurring infrequently as reported in the datasets. This imbalance can cause the model to overlook or underpredict the occurrence of such extreme events. By applying SMOTE, which effectively mitigates the class imbalance by generating synthetic samples for the minority class, thereby performing an "up-sampling" procedure, we generated synthetic samples for the minority class, thus enabling the model to better capture the underlying patterns of these infrequent but impactful events. This approach enhances the model's ability to predict the likelihood and impacts of droughts and heatwaves, leading to more reliable and robust predictions.

To model the relationships between predictors and responses, separate XGBoost models were trained for each of the three categorized drought impacts, as well as for heat impact. Hyperparameter tuning was performed to optimize model performance, with parameters such as learning rate, gamma, and maximum depth iteratively adjusted to identify the optimal configuration for predictive accuracy. Data splitting into training and testing sets ensured a robust evaluation and validation of the models. The performance of the models in predicting drought impacts for various sectors, as well as heatwave impacts on humans, was evaluated using the Relative Operating Characteristic (ROC) score (Mason, 1982).

3 Result

3.1 European drought and heatwave characteristics

Based on the median ensemble of ISIMIP model simulations (Warszawski et al., 2013) from 2039 to 2100 (62 years), the occurrence of drought events in Europe is projected to increase by at least 40 events under the SSP5-8.5 scenario compared to the reference period of 1953-2014 (84 events, Fig. 1a). Countries such as Spain, France, the Netherlands, Belgium, Italy, the Balkan regions, and Finland will experience a total drought month (duration) exceeding 220 months over 62 years (744 months), nearly twice as long as the reference period (Fig. 1b) (Samaniego et al., 2018). On average, drought duration will increase by up to half a month in these regions (Fig. 1c). Similarly, the frequency of droughts across Europe is expected to increase by 1 event more than in the past (Fig. 1d) (Spinoni et al., 2018). Under the SSP1.2-6 scenario, the increase in drought characteristic across Europe is approximately half of that under SSP5-8.5 and even lower in western Russia for average drought



duration (Supplementary Fig. S2). For instance, drought events are projected to increase by around 20 events in Germany under SSP1-2.6 and around 40 events under SSP5-8.5.




Figure 1. Changes (future-reference periods) in drought characteristics across Europe under SSP5-8.5 based on the median ensemble of the ISIMIP models. a) Changes in the number of drought events, b) changes in total drought duration in month, c) changes in average drought duration in month, and d) changes in frequency.

185 The increase in the number of drought events and frequency (number of events per year) in eastern Europe is clearly visible when comparing the number of drought events during the reference period to the far future under SSP5-8.5 scenario (Fig. 1a and 1d). For the drought hotspot, however, Western Europe was identified as a drought hotspot (90 events), with fewer drought events occurring in eastern Europe (Supplementary Fig. S3a). Drought hotspot regions are projected to spread towards the south and east under SSP1.2-6, experiencing approximately 20-30 more drought events than in the past (Supplementary
 190 Fig. S3b). Under the SSP5.8-5 scenario, almost all of Europe, except for northern Norway and Sweden, is expected to become drought hotspot regions, with at least 40 more events than in the past, representing an increase of over 44% (Supplementary Fig. S3c).

The increase of European temperature due to climate change strongly influences heatwave characteristics (Russo et al., 2015). The number of heatwave events exhibits substantial changes in many regions, especially in southern Europe, where
 195 more than 350 events in 62 years (>5 events per year) are anticipated under SSP5-8.5 (Fig 2a). Sweden and Ireland are



expected to experience 150 more heatwaves than during the reference period. The changes in total and average heatwave durations are particularly pronounced in southern Europe (Amengual et al., 2014; Molina et al., 2020). In this region, more than 4000 days (36%) between 2039 and 2100 (total 11,408 days) are categorized as heatwave days compared to the past period (1953-2014, Fig. 2b). This means that southern Europe will experience heatwaves for around two months each year during the extended summer period from May to October. Future heatwave events will last at least six days longer in Spain, Italy, and the Balkan region (Fig. 2c). On average, two more heatwave events will occur in the future in southern, western, and parts of eastern Europe (Fig. 2d). The changes in characteristics of future heatwave events will be ~50% lower under SSP1-2.6 than SSP 5-8.5 (Supplementary Fig. S4). For instance, Germany may experience 150 more heatwave events than the reference period, with a total heatwave duration of 1000 days under SSP1-2.6.




Figure 2. Changes (future-reference periods) in heatwave characteristics across Europe under SSP5-8.5 based on the median ensemble of the ISIMIP models. a) Changes in the number of heatwave events, b) changes in total heatwave duration in day (total number of day for 62 years is 11,408 days), c) changes in average heatwave duration in day, and d) changes in frequency.

Unlike droughts, which often occurred in western Europe and the UK during the reference period (Supplementary Fig. S3a), the hotspot regions for heatwave events are identified in eastern Europe, mainly in Russia, Italy, and Portugal (Supplementary Fig. S5a). Global warming, however, shifts these hotspot regions into the Mediterranean, extending from Spain, Italy, and the Balkan countries to southeastern Europe. The number of heatwave events in these regions is projected to be approximately

300 and 500 events higher than the reference period under SSP1-2.6 and SSP5-8.5 scenarios, respectively (Supplementary Fig S5b,c). The rate of increase in heatwave events is less pronounced in the UK, Scandinavian countries, and northeastern Europe.

3.2 The characteristics of compound and cascading events

The changes in compound drought and heatwave (CDH) characteristics across Europe in a warming world under SSP5-8.5 are presented in Figure 3. Overall, the number of CDH events increases by 15 and 30 events more than the reference period under SSP1-2.6 and SSP5-8.5, respectively (Supplementary Fig. S6a and Fig. 3a). Almost all of European region, except Sweden and northern UK, will experience similar changes under extreme climate change scenarios. However, some countries, such as France, western Germany, Switzerland, Austria, northeastern Spain, Norway, and Moldova will experience significant changes even under SSP1-2.6 (Supplementary Fig. S6a). The highest changes in CDH duration are found in western, southern, and eastern Europe (Fig. 3b,c), with a pattern similar to changes in total heatwave duration. This indicates that changes in CDH duration are driven by heatwave duration. The frequency of CDH events will increase on average by around 0.4 events per year under SSP5-8.5, with a maximum frequency of 0.76 events per year (Fig. 3d).




Figure 3. Changes (future-reference periods) in compound event characteristics across Europe under SSP5-8.5 based on the median ensemble of the ISIMIP models. a) Changes in the number of compound events, b) changes in total compound duration in month, c) changes in average compound duration in month, and d) changes in compound frequency.



The changes in cascading drought and heatwave (CaDH) characteristics under extreme warming are similar to CDH (Supplementary Fig. S7). However, CaDH characteristics under SSP1-2.6 show higher changes than CDH (see Supplementary Fig. S6 and S8). On average, the occurrence of CaDH events under SSP1-2.6 is 19 events more than the reference period (compared to 15 under SSP5-8.5). Many regions in Europe will experience higher number of CaDH events than CDH under SSP1-2.6, leading to a higher frequency of CaDH events. In terms of duration, the changes are more pronounced for total CaDH than total CDH, with total durations being 77 and 55 months longer than the reference period, respectively (Sutanto et al., 2019b). In contrast, the average duration of CaDH events (0.8 month) is shorter than that of CDH events (1.1 month). Our results indicate that under SSP1-2.6, CaDH events will occur more frequently across Europe but with shorter durations compared to CDH events.

Analyses of individual CnC events from the reference period reveal that northern Portugal, Italy, and eastern Europe used to be hotspot regions for both compound and cascading events (Fig. 4a,d) (see also Niggli et al. (2022)). Under the SSP1-2.6 scenario, some parts of France, the Alps, southern Germany, and Moldova will become new hotspot regions for CnC dry hazards, associated with a higher increase in CnC events in these regions compared to others (Fig. 4b,e). For example, in eastern Europe, the number of CDH events is projected to increase from 10 events in the past to 25 events in the future under SSP1-2.6, more than doubling. In the Alpine region, the number of CDH events is expected to increase sharply from 4 events in the past to 35 events in the future, which is almost ninefold. A significant difference between future CDH and CaDH hotspot regions is observed as the climate becomes more extreme. While some regions in northern Europe may not emerge as hotspots for CDH, most European regions will become hotspot regions for CaDH events (Fig. 4c,f).




Figure 4. Hotspot regions for compound drought and heatwave (CDH) derived from the number of CDH event across Europe. a) Hotspot regions based on historical period (1953-2014), b) hotspot regions based on future period under SSP1-2.6 (2039-2100), and c) hotspot regions based on future period under SSP5-8.5 (2039-2100). d,e,f) Same as a, b, and c but for cascading drought and heatwave (CaDH). The number of CnC events was obtained from the median ensemble of the ISIMIP models.



3.3 Summary of single and CnC dry hazards in Europe

Figure 5 summarizes the characteristics of droughts, heatwaves, and CnC events, averaged from all grid cells in each European region (Supplementary Fig. S9) from past to future scenarios. Western Europe (WE) has historically experienced a high number of drought events (88.9 events), whereas Eastern Europe (EE) has had slightly fewer (77.7 events). However, EE has experienced the longest drought durations, with an average duration 0.2 to 0.3 months longer than that of other regions (Sutanto and Van Lanen, 2020). Future projections under SSP5-8.5 indicate that WE will continue to experience the highest number of drought events, with an average of 130.7 events. Nonetheless, the increase in drought events in WE (41.8 events) is lower compared to EE, where the rate of increase is projected to be 46.2 events. Conversely, Southern Europe (SE) is projected to face the longest drought durations (Sutanto et al., 2019b), with an average total duration extending up to 231.8 months, doubling from historical values. Both WE and SE are expected to see an increase in the frequency of drought events (2.1 events per year).

	Number of events			Total duration			Average duration			Frequency of events		
	Historical	SSP1-2.6	SSP5-8.5	Historical	SSP1-2.6	SSP5-8.5	Historical	SSP1-2.6	SSP5-8.5	Historical	SSP1-2.6	SSP5-8.5
Drought												
NE	83.9	106.3	123.7	118.8	157.6	204.3	1.4	1.5	1.7	1.4	1.7	1.9
WE	88.9	116.9	130.7	118.8	172.0	223.7	1.3	1.5	1.8	1.4	1.9	2.1
EE	77.7	103.1	123.9	120.3	161.6	221.7	1.6	1.6	1.8	1.3	1.7	1.9
SE	84.2	111.9	129.3	115.7	169.3	231.8	1.4	1.5	1.8	1.4	1.8	2.1
Heatwave												
NE	51.9	114.8	248.5	227.6	639.3	1639.5	4.3	4.5	5.6	2.2	2.7	3.0
WE	54.4	221.5	392.8	233.2	1269.6	2985.9	4.3	5.3	7.2	2.2	2.9	3.7
EE	65.9	207.1	385.8	299.4	1141.5	3066.8	4.5	5.4	7.9	2.2	2.9	3.6
SE	65.4	231.4	416.8	303.5	1493.8	4102.5	4.6	5.4	9.0	1.9	2.6	3.5
Compound												
NE	4.9	15.2	30.0	10.4	44.1	128.7	2.1	2.8	4.5	0.1	0.2	0.5
WE	5.0	24.6	35.5	9.9	88.2	190.4	2.0	3.4	5.5	0.1	0.4	0.6
EE	6.6	21.4	34.9	15.0	72.5	188.6	2.3	3.4	5.5	0.1	0.3	0.6
SE	5.1	20.7	33.9	11.3	81.3	215.1	2.2	3.7	6.4	0.1	0.3	0.5
Cascading												
NE	10.5	25.9	38.3	26.3	78.7	164.7	2.5	2.9	4.3	0.2	0.4	0.6
WE	10.7	35.4	40.3	25.5	123.9	212.4	2.4	3.5	5.4	0.2	0.6	0.7
EE	13.2	33.5	40.1	34.3	114.8	218.4	2.6	3.4	5.4	0.2	0.5	0.6
SE	10.7	31.1	38.6	26.9	119.9	242.2	2.5	3.7	6.3	0.2	0.5	0.6

Figure 5. A summary of drought, heatwave, and CnC characteristics in each European region for reference period, far future under SSP1-2.6, and far future under SSP5-8.5. The unit for drought duration is month, for heatwave is day, and for CnC is month. NE stands for northern Europe, WE stands for western Europe, EE stands for eastern Europe, and SE stands for southern Europe. Blue color indicates values under 25th percentile of each hazard characteristic from historical to SSP5-8.5 in NE, WE, EE, and SE; yellow color indicates median values of each hazard characteristic from historical to SSP5-8.5 in NE, WE, EE, and SE; red color indicates values above 75th percentile of each hazard characteristic from historical to SSP5-8.5 in NE, WE, EE, and SE.

Eastern Europe (EE) experienced the highest number of heatwave events (65.9 events) in the historical period, followed closely by SE with 65.4 events, WE with 54.4 events, and Northern Europe (NE) with 51.9 events (Figure 5). Regarding



heatwave duration, SE reported the longest total duration, averaging 303.5 days, with EE closely following at 299.4 days. The average duration of heatwaves was 4.6 days in SE and 4.5 days in EE. Future projections indicate that SE is expected to experience increase in both the number and duration of heatwave events under SSP5-8.5, with occurrences rising sixfold and durations extending thirteenfold compared to the reference period. projected average heatwave durations are anticipated to be 9 days in SE, 7.9 days in EE, 7.2 days in WE, and 5.6 days in NE. However, the highest frequency of heatwaves is predicted to occur in WE, with an average of 3.7 events per year, followed by EE with 3.6 events per year, SE with 3.5 events per year, and NE with 3 events per year. The increase of heatwave events and duration is lesser in NE compared to other regions, especially for SSP1-2.6 (bluish colors).

Figure 5 also indicates that EE faced the highest number of CDH events from 1953 to 2014 (6.6 events), and an even greater number of CaDH events (13.2 events). Interestingly, the occurrences of CnC events under global warming are projected to be slightly higher in WE compared to EE (reddish colors under SSP1-2.6). Overall, the rate of increase in future CDH event will be nearly twice as high as the increase in CaDH events, though the absolute number of CaDH even remains greater. Future CDH events are expected to be 4 to 6 times higher, while CaDH events are projected to rise by 3 and 3.5 times under SSP1-2.6 and SSP5-8.5 scenarios, respectively. Despite the higher number of future CnC events in WE and EE, these regions are not expected to have longer CnC durations. In contrast, SE is projected to experience the longest CnC durations across Europe under extreme warming conditions, with total durations of 215 months for CDH and 242 months for CaDH. The average duration of CnC events in SE is projected to be between 6.3 to 6.4 months. Additionally, the frequency of CnC events is expected to increase around 0.5 events per year in WE and EE for CDH events, and in WE for CaDH events. This analysis highlights that while the number and frequency of future CnC events are higher in WE, SE is projected to experience longer durations of CnC events, consistent with previous study based on re-analysis data (Sutanto et al., 2019b).

3.4 Predicting drought and heatwave impacts in Germany

Predicting the impacts of dry hazards using Machine Learning (ML) approaches required an extensive dataset of documented impact occurrences (Mount et al., 2016). The impact of drought and heatwave events varies depending on their severity and duration, which means that not all events result in significant impacts. What is more, the availability of documented data on drought and heatwave impacts is often limited as in many countries impact data are not collected and reported at all, or only on an ad hoc basis. Germany stands out as the country with a sufficient number of documented drought impacts in European Drought Impact Inventory (EDII) (Stahl et al., 2016; Sutanto et al., 2019a), which guided our selection of Germany as the focus for our impact prediction study.

Using the reported impacts in Germany and the Extreme Gradient Boosting (XGBoost) ML model (Friedman, 2001) trained using historical hazard indices (1953-2014), the models demonstrate robust forecasting performance, with area under the curve (AUC) values of 0.83, 0.76, and 0.81 for economic, non-economic, and ecosystem sectors, respectively (Supplementary Fig. S10a,b,c, see also Method). For heatwaves, the model evaluation shows a perfect score (AUC=1), which may be influenced by the limited amount of reported impact data (Supplementary Fig. S10d). Simulations from drought impact models generally predict a higher number of impacts occurring within 1 month, as illustrated in Supplementary Figure S11a,b,c. However, these



models tend to underestimate the duration of impacts for drought lasting longer than 5 months. For heatwaves, the model failed to predict impacts for the year 1987 and only predicted impacts with a maximum duration of 2 months (Supplementary Figure S11d). Overall, both drought and heatwave impact models provide relatively accurate simulations of past impacts, particularly for impacts occurred after the year 2000 due to more data on impacts.

290 Figure 6 illustrates simulations the number of drought and heatwave impacts under various scenarios for impacts occurring in a particular year. The number of drought impact on the economic sector is projected to increase in the future, rising from a median of 3 months to 4 months under both SSP1-2.6 and SSP 5-8.5, with longer impact up to 6 months (75th percentile) projected for SSP5-8.5 (Fig. 6a). For non-economic sectors, the duration of drought impacts, which lasted between 2 and 4 months (25th and 75th percentiles, respectively) during the reference period, is projected to increase by 1 month under both
 295 SSP1-2.6 and SSP5-8.5 scenarios (Fig. 6b). The impact on the ecosystem sector is anticipated to increase from 3 to 4 months in the future under SSP5-8.5, based on the median ensemble (Fig. 6c). In contrast, the impact of heatwaves on humans, which rarely occurred annually in the past (Supplementary Fig. S11d) is predicted to become twice as long under SSP1-2.6 and four times under SSP5-8.5 (Fig. 6d). The projected increases in drought and heatwave impacts are likely driven by rising temperatures and less precipitation due to global warming, which can exacerbate heatwaves and accelerate soil drying, leading
 300 to increased soil moisture drought (Miralles et al., 2018; Teuling, 2018).

The characteristics of drought and heatwave impacts for both reference and future scenarios are detailed in Supplementary Table S3. The total number of drought impacts on the economic sector is projected to nearly double in the future, increasing from 36 events to 65 and 66 events under SSP1-2.6 and SSP5-8.5, respectively. The increase in the number of impacts corresponds to a longer total impact duration, extending from 128 months to 216 months for SSP1.2-6 and to 248 months for
 305 SSP5-8.5. Concerning average duration, however, SSP1-2.6 indicates a slightly shorter duration (0.2 months), while SSP5-8.5 demonstrates a longer duration by 0.3 months. The frequency (total number of events/total years) of drought impacts on the economic sector is expected to increase from 2.1 to 3.5 and 4 under SSP1-2.6 and SSP5-8.5, respectively. A similar trend is projected for the non-economic sectors, indicating more and longer impacts. The ecosystem sector is projected to be the most affected by droughts, with the number, total duration, and frequency of drought impacts being three times higher than the
 310 reference period. The impact of heatwaves on human health and losses, on the other hand, will sharply increase in the future due to projected higher temperatures (Peng et al., 2011; Amengual et al., 2014). For instance, the number of heatwave impacts is projected to increase tenfold under SSP5-8.5, with the total duration over 62 years being 25 times longer and the average duration three times longer (increasing from 1.1 to 2.9 months). The rise in the number and duration of impacts will also result in a higher annual frequency of events.

315 4 Discussion

Southern Europe is frequently identified as a hotspot for future drought and heatwave events (Forzieri et al., 2014; Prudhomme et al., 2014; Cammalleri et al., 2020). However, analyses on drought characteristics reveal that many regions across Europe, not just the south, will experience an increase in both the number and frequency of drought events in the future, particularly




Figure 6. The predictions of drought and heatwave impacts in Germany under reference period (1953-2014) and future period (2039-2100) for SSP1-2.6 and SSP5-8.5. a) Prediction of drought impact on economic sector, b) Prediction of drought impact on non-economic sector, c) Prediction of drought impact on ecosystem sector, and d) Prediction of heatwave impact on humans. The Y axis shows the occurrence of impact in a year when impacts are predicted from all models. Lower box indicates 25 percentile, middle orange line indicates median, and upper box indicates 75 percentile. The whiskers show the 10 and 90 percentiles. Open circles indicate outliers.

in Eastern European countries (Paparrizos et al., 2018). Our projections suggest that Southern Europe will experience longer
 320 drought periods, reinforcing its status as a drought hotspot (Sutanto et al., 2019b). Our findings on future heatwave charac-
 teristics also confirm that Southern Europe remains a hotspot for heatwaves (Amengual et al., 2014; Molina et al., 2020).
 Interestingly, while Southern Europe was not identified as a heatwave hotspot in the reference period, Eastern Europe and
 Russia were, likely due to the intense 2010 Russian heatwaves (Russo et al., 2015). In our study, we employed a relative metric
 based on 90th percentile of the historical climate, meaning that heatwaves could be identified everywhere in Europe even in the



325 cold climate regions. Changes in CnC characteristics follow similar patterns to drought characteristics, indicating that drought is the primary driver of CnC dry hazards (Sutanto et al., 2019b).

We acknowledge that the use of ML model to predict the occurrence of dry hazard impacts is constrained by the availability of impact data to train the model (Mount et al., 2016). Despite combining impact data from the EDII (Stahl et al., 2016), the EM-DAT database (CRED, 2022), and data mined from international journals and English reports, the compiled data remains
 330 limited, particularly for heatwaves. Additionally, all databases exhibit temporal and geographical biases (Bachmair et al., 2015; Stahl et al., 2016). Previous studies have highlighted missing data in many disaster databases, where either the disaster or its impact was not reported or were omitted due to low impact (Bachmair et al., 2016; Jones et al., 2023). The absence of impact reports does not always mean there were no impacts (Karlsson and Ziebarth, 2018). Consolidating drought impact data into three impact categories enhances the performance of drought impact model, particularly for predicting droughts lasting more
 335 than one month. Since impact models tend to overpredict impacts occurring only for one month duration, predictions of impacts of this duration should be excluded from the result. Moreover, longer drought events generally trigger more significant impacts than shorter ones. We identified only a few reported heatwave impacts, mainly related to human mortality, with very few reports on forest fires and agricultural losses. Overall, we obtained five years of reported heatwave impacts on human mortality, which could be predicted by the XGBoost model, except for the year 1987, resulting in an AUC curve close to 1. The small number
 340 of positive and negative samples allows the model to achieve an ideal measure of separability between these samples (Marcum, 1947).

The XGBoost ML technique offers several advantages over process-based models, such as increased accuracy and improved time and computational efficiency (Rahman et al., 2022; Peach et al., 2023). However, ML models to predict the impacts of CnC events could not be developed due to data constraints regarding both the number of event and their impacts. Droughts
 345 and heatwaves are extreme events, with a recurrence interval of approximately 10% for events with $SMI < -1$ and for the 90th percentile threshold, respectively (McKee et al., 1993; WMO, 2012). Furthermore, the occurrence of droughts and heatwaves, depending on duration and severity, does not always result in impacts (Bachmair et al., 2016), making the occurrence of drought and heatwave impacts even rarer (Seneviratne et al., 2021). Additionally, the occurrence of CnC events analyzed in this study is less than 10%. This scarcity of data is a primary reason why a CnC impact model could not be developed using the ML
 350 technique at the moment. Moreover, there is no existing database that provides information on CnC impacts. While EM-DAT includes information on primary and secondary hazards, it does not cover impacts. Thus, we highly recommend having a comprehensive multi-hazard impact database.

Techniques for predicting drought impacts can be categorized into top-down, bottom-up, and hybrid approaches (Shyrokaya et al., 2023). The top-down approach utilizes a combination of climate models and deterministic impact models, such as
 355 employing crop models to predict yield losses (Ogutu et al., 2018; Sutanto et al., 2024). Sutanto et al. (2024) employed the WOFOST crop model to assess the impact of CnC events on maize yields, demonstrating the potential of this approach for impact prediction. The bottom-up approach relies on statistical models to link observed impacts to climate variables, such as using log regression model to predict drought impacts (Blauhut et al., 2015). The hybrid approach, employed in this study and others (Sutanto et al., 2019a, 2020b), is based on ML models to develop drought impact functions. The ML models can be



utilized to predict various impacts and require less computational power than top-down approach, which is often dedicated to only predict single impact. The hybrid approaches are valuable for developing impact-based prediction models, as predictions of impacts are still missing in many natural hazard early warning systems (EWSs) (Walker et al., 2024). Many EWSs provide information only on the occurrence of hazards and not on the corresponding impacts. Information on impact occurrence is of utmost importance for stakeholders and policymakers, providing actionable information for disaster risk management and risk reduction.

5 Conclusions

This study is pioneering in predicting future drought and heatwave characteristics, both in isolation and as CnC events, including their impacts. Our findings indicate that drought and heatwave characteristics, such as the number, duration, and frequency of events, will increase across Europe under both SSP1-2.6 and SSP5-8.5. The increase in single hazard events will lead to a corresponding increase in CnC events. Historically, drought hotspots have been concentrated in West Europe, while heatwave hotspots have been identified in eastern and southern Europe, mainly in Russia, Italy, and Portugal. However, under future climate scenarios, these hotspots are projected to shift toward southern Europe. Our findings also demonstrate that the whole Europe will experience CnC events in the future, not just the southern regions where much of the focus of hot and dry hazard research has traditionally been concentrated. The occurrence of CDH events is expected to be six times higher than in the reference period, while CaDH events are projected to increase by 3.5 times under SSP5-8.5.

Additionally, this study highlights the potential of using a hybrid approach based on ML for projecting drought and heatwave impacts. We projected that drought impacts on economic, non-economic, and ecosystem sectors in Germany will be double in 2100, while heatwave impacts on human health and mortality will increase ninefold. These findings demonstrate that the hybrid method offers new opportunities not only for impact projection, as demonstrated here, but also for impact forecasting (Sutanto et al., 2019a; Shyrokaya et al., 2023). We emphasize the urgent need for holistic risk assessment frameworks that account for CnC events and subsequently implementing effective adaptation strategies based on co-occurring events. Despite current limitations in hybrid approach for predicting CnC impacts, advancing complementary approaches—including top-down, bottom-up, and hybrid methods—will be essential for building a comprehensive and actionable understanding of future multi hazard risks.

Code availability. All codes used to conduct the analysis presented in this paper will be available online in the 4TU Centre for Research Data with <https://doi.org/xxxx> (Sutanto and Duku, 202x).

Data availability. The ERA5 Land data are accessible through the Copernicus Data Store (CDS), which are freely available ([doi:10.24381/cds.e2161bac](https://doi.org/10.24381/cds.e2161bac)). Soil moisture and projection data are downloaded from the ISIMIP output (ISIMIP3b) for global water sector (<https://data.isimip.org/>). Other



390 data and generated and/or analyzed during this study will be available online in the 4TU Centre for Research Data with <https://doi.org/xxxx>
(Sutanto and Duku, 202x).

Author contributions. All authors conceived and implemented the research. Data analyses, model output analyses, and all figures have been performed by S.J.S. and M.G. The ML models were developed by C.D. S.J.S wrote the initial version of the paper. R.D. and S.P contributed to interpreting the results, discussion, and improving the paper.

Competing interests. The authors declare no competing interests.

395 *Acknowledgements.* The research is supported by the CDHEU project, which is funded by the Wageningen Data Driven Discoveries in Changing Climate (D3-C2). The hydrometeorological outputs came from the Copernicus Data Store and the ISIMIP data repository (see data availability section). All data were processed using High Performance Computing Cluster Anunna hosted by Wageningen University and Research.



References

- 400 Amengual, A., Homar, V., Romero, R., Brooks, H. E., Ramis, C., Gordaliza, M., and Alonso, S.: Projections of heat waves with high impact on human health in Europe, *Global and Planetary Change*, 119, 71–84, <https://doi.org/https://doi.org/10.1016/j.gloplacha.2014.05.006>, 2014.
- Bachmair, S., Kohn, I., and Stahl, K.: Exploring the link between drought indicators and impacts, *Nat. Hazards Earth Syst. Sci.*, 15, 1381–1397, <https://doi.org/https://doi.org/10.5194/nhess-15-1381-2015>, 2015.
- 405 Bachmair, S., Svensson, C., Hannaford, J., and Stahl, K.: A quantitative analysis to objectively appraise drought indicators and model drought impacts, *Hydrol. Earth Syst. Sci.*, 20, 2589–2609, <https://doi.org/https://doi.org/10.5194/hess-20-2589-2016>, 2016.
- Bachmair, S., Svensson, C., Prosdocimi, I., Hannaford, J., and Stahl, K.: Developing drought impact functions for drought risk management, *Nat. Hazards Earth Syst. Sci.*, 17, 1947–1960, <https://doi.org/https://doi.org/10.5194/nhess-17-1947-2017>, 2017.
- Biella, R., Shyrokaya, A., Ionita, M., Vignola, R., Sutanto, S. J., and co authors: The 2022 drought needs to be a turning point for European drought risk management, Preprint egusphere-2024-2069, <https://doi.org/https://doi.org/10.5194/egusphere-2024-2069>, 2024.
- 410 Blauhut, V., Gudmundsson, L., and Stahl, K.: Towards pan-European drought risk maps: quantifying the link between drought indices and reported drought impacts, *Environ. Res. Lett.*, 10, 014 008, <https://doi.org/https://doi.org/10.1088/1748-9326/10/1/014008>, 2015.
- Blauhut, V., Stoelzle, M., Ahopelto, L., Brunner, M. I., Teutschbein, C., Wendt, D. E., and co authors: Lesson from the 2018–2019 European droughts: a collective need for unifying drought risk management, *Nat. Hazards Earth Syst. Sci.*, 22, 2201–2217, <https://doi.org/https://doi.org/10.5194/nhess-22-2201-2022>, 2022.
- 415 Buras, A., Rammig, A., and Zang, C. S.: Quantifying impacts of the 2018 drought on European ecosystems in comparison to 2003, *Biogeosciences*, 17, 1655–1672, <https://doi.org/https://doi.org/10.5194/bg-17-1655-2020>, 2020.
- Burek, P., Satoh, Y., Kahil, T., Tang, T., Greve, P., Smilovic, M., and co authors: Development of the Community Water Model (CWatM v1.04) - a high-resolution hydrological model for global and regional assessment of integrated water resources management, *Geosci. Model Dev.*, 13, 3267–3298, <https://doi.org/https://doi.org/10.5194/gmd-13-3267-2020>, 2020.
- 420 Cammalleri, C., Naumann, G., Mentaschi, L., Bisselink, B., Gelati, E., De Roo, A., and Feyen, L.: Diverging hydrological drought traits over Europe with global warming, *Hydrol. Earth Syst. Sci.*, 24, 5919–5935, <https://doi.org/https://doi.org/10.5194/hess-24-5919-2020>, 2020.
- Chawla, N. V., Bowyer, K. W., Hall, L. O., and Kegelmeyer, W. P.: SMOTE: Synthetic minority over-sampling technique, *Journal of Artificial Intelligence Research*, 16(1), 321–357, <https://doi.org/https://doi.org/10.1613/jair.953>, 2002.
- 425 CRED: The Emergency Events Database (EM-DAT). The last 25 years in research, CredCrunch67, <https://www.cred.be/sites/default/files/CredCrunch67.pdf>, 2022.
- Fischer, E. M., Seneviratne, S. I., Vidale, P. L., Lüthi, D., and Schär, C.: Soil moisture–atmosphere interactions during the 2003 European summer heat wave, *Journal of Climate*, 20(20), 5081–5099, <https://doi.org/https://doi.org/10.1175/JCLI4288.1>, 2007.
- Forzieri, G., Feyen, L., Rojas, R., Flörke, F., Wimmer, F., and Bianchi, A.: Ensemble projections of future streamflow droughts in Europe, *Hydrol. Earth Syst. Sci.*, 18, 85–108, <https://doi.org/https://doi.org/10.5194/hess-18-85-2014>, 2014.
- 430 Friedman, J.: Greedy function approximation: a gradient boosting machine, *Ann. Stat.*, 29, 1189–1232, <https://doi.org/https://www.jstor.org/stable/2699986>, 2001.
- Hawkins, E., Osborne, T. M., Ho, C. K., and Challinor, A. J.: Calibration and bias correction of climate projections for crop modelling: an idealised case study over Europe, *Agricultural and forest meteorology*, 170, 19–31, <https://doi.org/https://doi.org/10.1016/j.agrformet.2012.04.007>, 2013.
- 435



- Ionita, M., Tallaksen, L. M., Kingston, D. G., Stagge, J. H., Laaha, G., Van Lanen, H. A. J., Scholz, P., Chelcea, S. M., and Haslinger, K.: The European 2015 drought from a climatological perspective, *Hydrol. Earth Syst. Sci.*, 21, 1397–1419, <https://doi.org/https://doi.org/10.5194/hess-21-1397-2017>, 2017.
- Jones, R. L., Kharb, A., and Tubeuf, S.: The untold story of missing data in disaster research: a systematic review of the empirical literature utilizing the Emergency Events Database (EM-DAT), *Environ. Res. Lett.*, 18, 103 006, <https://doi.org/https://doi.org/10.1088/1748-9326/acfd42>, 2023.
- Jonkman, S. N.: Global perspectives on loss of human life caused by floods, *Nat. Hazards*, 34, 151–175, <https://doi.org/https://doi.org/10.1007/s11069-004-8891-3>, 2005.
- Karlsson, M. and Ziebarth, N. R.: Population health effects and health-related costs of extreme temperatures: Comprehensive evidence from Germany, *Journals of Environmental Economics and Management*, 91, 93–117, <https://doi.org/https://doi.org/10.1016/j.jeem.2018.06.004>, 2018.
- Lange, S. and Büchner, M.: ISIMIP3b bias-adjusted atmospheric climate input data (v1.1). Potsdam-Institut für Klimafolgenforschung, Germany, <https://doi.org/10.48364/ISIMIP.842396.1>, 2021.
- Lavaysse, C., Cammalleri, C., Dosio, A., Van der Schrier, G., Toreti, A., and Vogt, J.: Towards a monitoring system of temperature extremes in Europe, *Nat. Hazards Earth Syst. Sci.*, 18, 91–104, <https://doi.org/https://doi.org/10.5194/nhess-18-91-2018>, 2018.
- Lemaître, G., Nogueira, F., and Aridas, C. K.: Imbalanced-learn: A Python Toolbox to Tackle the Curse of Imbalanced Datasets in Machine Learning, *Journal of Machine Learning Research*, 18(17), 1–5, <https://doi.org/https://doi.org/10.48550/arXiv.1609.06570>, 2017.
- Leonard, M., Westra, S., Phatak, A., Lambert, M., Van den Hurk, B., McInnes, K., Risbey, J., Schuster, S., Jakob, D., and Stafford-Smith, M.: A compound event framework for understanding extreme impacts, *WIREs Clim. Change*, 5, 113–128, <https://doi.org/https://doi.org/10.1002/wcc.252>, 2014.
- Liu, M. and Huang, M. C.: Compound disasters and compounding processes: Implications for disaster risk management, In *Global Assessment Report on Disaster Risk Reduction United Nations Office for Disaster Risk Reduction (UNDRR)*, <https://www.undrr.org/publication/compound-disasters-and-compounding-processes-implications-disaster-risk-management>, 2015.
- Marcum, J. I.: A Statistical Theory of Target Detection by Pulsed Radar. Santa Monica, CA: RAND Corporation, https://www.rand.org/pubs/research_memoranda/RM754.html, 1947.
- Mardian, J., Champagne, C., Bonsal, B., and Berg, A.: A machine learning framework for predicting and understanding the Canadian drought monitor, *Water Resources Research*, 59, e2022WR033 847, <https://doi.org/https://doi.org/10.1029/2022WR033847>, 2023.
- Mason, I.: A model for assessment of weather forecasts, *Aust. Meteor. Mag.*, 30, 291–303, 1982.
- McKee, T. B., Doesken, N. J., and Kleist, J.: The relationship of drought frequency and duration to time scale, In *Proc. of 8th Conf. on Applied Climatology (Amer. Meteor. Soc., Anaheim, California, pp. 179–184, 1993.*
- Miralles, D. G., Gentile, P., Seneviratne, S. I., and Teuling, A. J.: Land-atmospheric feedbacks during droughts and heatwaves: state of the science and current challenges, *Ann. N.Y. Acad. Sci.*, 1436, 1–17, <https://doi.org/https://doi.org/10.1111/nyas.13912>, 2018.
- Molina, M. O., Sánchez, E., and Gutiérrez, C.: Future heat waves over the Mediterranean from an Euro-CORDEX regional climate model ensemble, *Scientific Reports*, 10, 8801, <https://doi.org/https://doi.org/10.1038/s41598-020-65663-0>, 2020.
- Mount, N. J., Maier, H. R., Toth, E., Elshorbagy, A., Solomatine, D., Chang, F.-J., and Abrahart, R. J.: Data-driven modelling approaches for socio-hydrology: opportunities and challenges within the Panta Rhei Science Plan, *Hydrological Sciences Journal*, 61(7), 1192–1208, <https://doi.org/https://doi.org/10.1080/02626667.2016.1159683>, 2016.



- Mukherjee, S. and Mishra, A. K.: Increase in Compound Drought and Heatwaves in a Warming World, *Geophys. Res. Lett.*, 48, e2020GL090617, <https://doi.org/https://doi.org/10.1029/2020GL090617>, 2021.
- 475 Naumann, G., Cammalleri, C., Mentaschi, L., and Feyen, L.: Increased economic drought impacts in Europe with anthropogenic warming, *Nat. Clim. Chang.*, 11, 485–491, <https://doi.org/https://doi.org/10.1038/s41558-021-01044-3>, 2021.
- Niggli, L., Huggel, C., Muccione, V., Neukom, R., and Salzmann, N.: Towards improved understanding of cascading and interconnected risks from concurrent weather extremes: Analysis of historical heat and drought extreme events, *PLOS Clim.*, 1(8), e0000057, <https://doi.org/https://doi.org/10.1371/journal.pclm.0000057>, 2022.
- 480 Ogotu, G. E. O., Franssen, W. H. P., Supit, I. Omondi, P., and Hutjes, R. W. A.: Probabilistic maize yield prediction over east Africa using dynamic ensemble seasonal climate forecasts, *Agric. For. Meteorol.*, 250–251, 243–261, <https://doi.org/https://doi.org/10.1016/j.agrformet.2017.12.256>, 2018.
- Paparrizos, S., Maris, F., Weiler, M., and Matzarakis, A.: Analysis and mapping of present and future drought conditions over Greek areas with different climate conditions, *Theor. Appl. Climatol.*, 131, 259–270, <https://doi.org/https://doi.org/10.1007/s00704-016-1964-x>, 2018.
- 485 Peach, L., da Silva, G. V., Cartwright, N., and Strauss, D.: A comparison of process-based and data-driven techniques for downscaling offshore wave forecasts to the nearshore, *Ocean Modelling*, 182, 102168, <https://doi.org/https://doi.org/10.1016/j.ocemod.2023.102168>, 2023.
- Peng, R. D., Bobb, J. F., Tebaldi, C., McDaniel, L., Bell, M. L., and Dominici, F.: Toward a quantitative estimate of future heat wave mortality under global climate change, *Environmental Health Perspective*, 119(5), 701–706, <https://doi.org/https://doi.org/10.1289/ehp.1002430>, 2011.
- 490 Prudhomme, C., Giuntoli, I., Robinson, E. L., Clark, D. B., Arnell, N. W., Dankers, R., and co authors: Hydrological droughts in the 21st century, hotspots and uncertainties from a global multimodel ensemble experiment, *PNAS*, 111(9), 3262–3267, <https://doi.org/https://doi.org/10.1073/pnas.1222473110>, 2014.
- Rahman, K. U., Pham, Q. B., Jadoon, K. Z., Shahid, M., Kushwaha, D. P., Duan, Z., Mohammadi, B., Kheder, K. M., and Anh, D. T.: Comparison of machine learning and process-based SWAT model in simulating streamflow in the Upper Indus Basin, *Applied Water Science*, 12, 178, <https://doi.org/https://doi.org/10.1007/s13201-022-01692-6>, 2022.
- 495 Rakovec, O., Samaniego, L., Hari, V., Markonis, Y., Moravec, V., Thober, S., Hanel, M., and Kumar, R.: The 2018–2020 Multi-Year Drought Sets a New Benchmark in Europe, *Earth’s Future*, 10, e2021EF002394, <https://doi.org/https://doi.org/10.1029/2021EF002394>, 2022.
- Robine, J.-M., Cheung, S. L. K., Roy, S. L., Van Oyen, H., Griffiths, C., Michel, J.-P., and Herrmann, F. R.: Death toll exceed 70,000 in Europe during the summer of 2003, *C.R. Biol.*, 331, 171–178, <https://doi.org/https://doi.org/10.1016/j.crv.2007.12.001>, 2008.
- 500 Russo, S., Sillmann, J., and Fischer, E. M.: Top ten European heatwaves since 1950 and their occurrence in the coming decades, *Environ. Res. Lett.*, 10, 124003, <https://doi.org/https://doi.org/10.1088/1748-9326/10/12/124003>, 2015.
- Samaniego, L., Thober, S., Kumar, R., Wanders, N., Rakovec, O., and co authors: Anthropogenic warming exacerbates european soil moisture droughts, *Nat. Clim. Change*, 8, 421–426, <https://doi.org/https://doi.org/10.1038/s41558-018-0138-5>, 2018.
- 505 Schuldt, B., Buras, A., Arend, M., Vitasse, Y., Beierkuhnlein, C., Damm, A., Gharun, M., Grams, T. E. E., Hauck, M., Hajek, P., Hartmann, H., Hiltbrunner, E., Hoch, G., Halloway-Phillips, M., Körner, C., Larysch, E., Lübke, T., Nelson, D. B., Rammig, A., Rigling, A., Rose, L., Ruehr, N. K., Schumann, K., Weiser, F., Werner, C., Wohlgemuth, T., Zang, C. S., and Kahmen, A.: A first assessment of the impact of the extreme 2018 summer drought on Central European forests, *Basic and Applied Ecology*, 45, 86–103, <https://doi.org/https://doi.org/10.1016/j.baae.2020.04.003>, 2020.



- 510 Seneviratne, S. I., Zhang, X., Adnan, M., Badi, W., Dereczynski, C., and co authors: Weather and Climate Extreme Events in a Changing Climate. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)], Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1513–1766, 2021.
- 515 <https://doi.org/https://doi.org/10.1017/9781009157896.013>, 2021.
- Shyrokaya, A., Pappenberger, F., Pechlivanidis, I., Messori, G., Khatami, S., Mazzoleni, M., and Baldassarre, G.: Advances and gaps in the science and practice of impact-based forecasting of droughts, *WIREs Water*, 11, e1698, <https://doi.org/https://doi.org/10.1002/wat2.1698>, 2023.
- Spinoni, J., Vogt, J. V., Naumann, G., Barbosa, P., and Dosio, A.: Will drought events become more frequent and severe in Europe?, *Int. J. Climatol.*, 38, 1718–1736, <https://doi.org/https://doi.org/10.1002/joc.5291>, 2018.
- 520 Stagge, J. H., Kohn, I., Tallaksen, L. M., and Stahl, K.: Modeling drought impact occurrence based on meteorological drought indices in Europe, *J. Hydrol.*, 530, 37–50, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2015.09.039>, 2015.
- Stahl, K., Kohn, I., Blauhut, V., Urquijo, J., De Stefano, L., and co authors: Impacts of European drought events: insights from an international database of text-based reports, *Nat. Hazards Earth Syst. Sci.*, 16, 801–819, <https://doi.org/https://doi.org/10.5194/nhess-16-801-2016>, 2016.
- 525 Sutanto, S. J. and Van Lanen, H. A. J.: Hydrological drought characteristics based on groundwater and runoff across Europe, *Proc. IAHS*, 383, 281–290, <https://doi.org/https://doi.org/10.5194/piahs-383-281-2020>, 2020.
- Sutanto, S. J. and Van Lanen, H. A. J.: Streamflow drought: implication of drought definitions and its application for drought forecasting, *Hydrol. Earth Syst. Sci.*, 25, 3991–4023, <https://doi.org/https://doi.org/10.5194/hess-25-3991-2021>, 2021.
- 530 Sutanto, S. J., van der Weert, M., Wanders, N., Blauhut, V., and Van Lanen, H. A. J.: Moving from drought hazard to impact forecasts, *Nature Communications*, 10, 4945, <https://doi.org/https://doi.org/10.1038/s41467-019-12840-z>, 2019a.
- Sutanto, S. J., Vitolo, C., Di Napoli, C., D’Andrea, M., and Van Lanen, H. A. J.: Heatwaves, droughts, and fires: Exploring compound and cascading dry hazards at the pan-European scale, *Environ. Int.*, 134, 105 276, <https://doi.org/https://doi.org/10.1016/j.envint.2019.105276>, 2019b.
- 535 Sutanto, S. J., Van Lanen, H. A. J., Wetterhall, F., and Llort, X.: Potential of pan-European hydro-meteorological drought forecasts obtained from a Multi-Hazard early warning system, *Bull. Amer. Meteor. Soc.*, 101, 368–393, <https://doi.org/https://doi.org/10.1175/BAMS-D-18-0196.1>, 2020a.
- Sutanto, S. J., van der Weert, M., Blauhut, V., and Van Lanen, H. A. J.: Skill of large-scale seasonal drought impact forecasts, *Nat. Hazards Earth Syst. Sci.*, 20, 1595–1608, <https://doi.org/https://doi.org/10.5194/nhess-20-1595-2020>, 2020b.
- 540 Sutanto, S. J., Zarzoza Mora, S. B., Supit, I., and Wang, M.: Compound and cascading droughts and heatwaves decrease maize yields by nearly half in Sinaloa, Mexico, *npj Nat. Hazards*, 1, 26, <https://doi.org/https://doi.org/10.1038/s44304-024-00026-7>, 2024.
- Teuling, A. J.: A hot future for European droughts, *Nat. Clim. Change*, 8, 364–365, <https://doi.org/https://doi.org/10.1038/s41558-018-0154-5>, 2018.
- Tripathy, K. P. and Mishra, A. K.: How unusual is the 2022 European compound drought and heatwave event?, *Geophysical Research Letters*, 50, e2023GL105 453, <https://doi.org/https://doi.org/10.1029/2023GL105453>, 2023.
- 545 Tripathy, K. P., Mukherjee, S., Mishra, A. K., and Williams, A. P.: Climate change will accelerate the high-end risk of compound drought and heatwave events, *PNAS*, 120(28), e2219825 120, <https://doi.org/https://doi.org/10.1073/pnas.2219825120>, 2023.



- Van Lanen, H. A. J., Laaha, G., Kingston, D. G., Gauster, T., Ionita, M., and co authors: Hydrology needed to manage droughts: The 2015 European case, *Hydrological Processes*, 30(17), 3097–3104, <https://doi.org/https://doi.org/10.1002/hyp.10838>, 2016.
- 550 Vitolo, C., Di Napoli, C., Di Giuseppe, F., Cloke, H. L., and Pappenberger, F.: Mapping combined wildfire and heat-stress hazards to improve evidence-based decision making, *Environ. Int.*, 127, 21–34, <https://doi.org/https://doi.org/10.1016/j.envint.2019.03.008>, 2019.
- Walker, D. W., oliveira, J. L., Cavalcante, L., Kchouk, S., and co authors: It's not all about drought: What "drought impacts" monitoring can reveal, *International Journal of Disaster Risk Reduction*, 103, 104 338, <https://doi.org/https://doi.org/10.1016/j.ijdr.2024.104338>, 2024.
- Warszawski, L., Frieler, K. and Huber, V., Piontek, F., Serdeczny, O., and Schewe, J.: The inter-Sectoral impact Model Intercomparison Project (ISI-MIP): project framework, *PNAS*, 111(9), 3228–3232, <https://doi.org/https://doi.org/10.1073/pnas.1312330110>, 2013.
- 555 WMO: Standardized precipitation index user guide. In *World Meteorological Organization Report* (eds Svoboda, M., Hayes, M. & Wood, E.) WMO, Geneva, Switzerland, <https://library.wmo.int/idurl/4/39629>, 2012.