

Comparing Flood Forecasting and Early Warning Systems in Northwestern Europe

Tim Busker¹, Daniela Rodriguez Castro², Sergiy Vorogushyn³, Jaap Kwadijk^{5,8}, Davide Zoccatelli⁷,
5 ~~Rafaella G. L. Oliveira~~~~Rafaella Loureiro~~¹, Heather J. Murdock⁴, Laurent Pfister⁷, Benjamin Dewals²,
Kymo Slager⁵, Annegret H. Thieken⁴, Jan Verkade⁵, Patrick Willems⁶, and Jeroen C.J.H. Aerts^{1,5}

¹ Institute for Environmental Studies (IVM), VU University Amsterdam, Amsterdam, The Netherlands.

² Research Group of Hydraulics in Environmental and Civil Engineering (HECE), University of Liège, Liège, Belgium.

³ GFZ Helmholtz Centre for Geosciences, Potsdam, Germany.

10 ⁴ Institute of Environmental Science and Geography, University of Potsdam, Potsdam, Germany.

⁵ Deltares, Delft, The Netherlands.

⁶ Hydraulics and Geotechnics Section, KU Leuven, Leuven, Belgium.

⁷ Luxembourg Institute of Science and Technology (LIST), Esch-sur-Alzette, Luxembourg.

⁸ Faculty of Engineering Technology, University of Twente, Enschede, The Netherlands.

15 *Correspondence to:* Tim Busker (tim.busker@vu.nl)

Abstract. This study compares operational Flood Forecasting and Early Warning Systems (FFEWs) in transboundary river basins in Northwestern Europe, covering parts of Luxembourg, Germany, The Netherlands and Belgium. ~~This region was hit by an extreme flood event in 2021 with more than 200 fatalities. This region was hit by an extreme flood event in 2021 with over 200 fatalities. Due to the high death toll and the large number of people who did not receive warnings, the FFEWSs were heavily criticized afterward. Due to the high death toll and the high share of people not receiving warnings, FFEWSs were heavily criticized in the aftermath. This study shows strong improvements in FFEWSs after the flood event in all countries. This study shows strong improvements of the FFEWSs after the flood event in all countries. Expert interviews with experts~~ across the region reveal that warning thresholds ~~have been~~^{are} optimized, some regions included new warning thresholds for extreme events (e.g.g. dark purple), and flood crisis management plans ~~are have been~~^{are} improved. In addition, all countries
25 now use mobile phone-based alerts using cell broadcasting technology. ~~The assessment of the warning systems shows strong differences between and within countries, with different flood warning levels and warning colour codes in use. The assessment of warning systems showed strong differences within and between the countries, with different flood warning levels and warning color codes used. We recommend investigating to what extent international harmonization of flood warnings can improve communication and decision-making. The interviews also reveal that the adoption of operational impact-based forecasts remains a challenge, while these are crucial for translating hydrological forecasts into effective actions. The interviews also revealed that the uptake of operational impact-based forecasts remains challenging, while these are crucial for translating hydrological forecasts to effective actions. For example, the interviewees stress the need for operational flood inundation forecasts, which are currently only provided in Flanders. For example, interviewees highlighted the need for operational flood inundation forecasts. However, Flanders is the only region where such forecasts are provided. One of the~~
30 ~~four main recommendations is to enhance forecasts with impact-based information, including inundation maps delineating the~~
35

~~population and objects at risk. Our study has four concrete recommendations: 1) Investigate the benefit of streamlined warning levels and colour codes between and within different countries; 2) Assess the added value of more extreme warning levels for catastrophic events, such as the purple levels in Luxembourg and some German regions; 3) accelerate the development of operational impact-based forecasting systems, and 4) implement a structural evaluation of warning communication chains. These challenges must be addressed to reduce the gap between early warning and early action during impactful flood events. This can improve the early actions taken by first responders and the affected population.~~

Formatted: Font: Not Italic

40

1 Introduction


The extreme flood event of July 2021 in Northwestern Europe caused more than 200 fatalities and around €40 billion of damage (Lehmkuhl et al., 2022). The extreme flood event of July 2021 in North-Western Europe caused over 200 fatalities and around €40 billion of damage (Lehmkuhl, et al., 2022). The frequency and severity of such flood events may increase in the future because of climate change and socio-economic developments (Tradowsky et al., 2023). To reduce avoidable damage and casualties, more reliable early warning systems are needed to protect Europe from flood impacts (European Environment Agency, 2024). To reduce avoidable damage and casualties, there is a call for more reliable early warning systems to protect Europe from flood impacts (European Environment Agency, 2024). Forecast-based actions are vital to reduce disaster risk, as they can effectively reduce the exposure and vulnerability of communities to floods (Pappenberger et al., 2015). For example, based on forecast information, the government may issue an evacuation order for communities in predefined flood zones or may advise households and businesses to protect properties and infrastructure with emergency flood protection measures (e.g., sandbags and pumps). For example, based on forecast information, the government may issue an evacuation order for communities in predefined flood zones or may advise households and businesses to protect properties and infrastructure with emergency flood protection measures (e.g., sandbags and pumps). The benefits of forecast-based actions are often much higher than their costs (Global Commission on Adaptation, 2019). Pappenberger et al. (2015) even claim a benefit on the order of 400 euros for every euro invested. Pappenberger et al. (2015) even claim a benefit in the order of 400 Euro for every Euro invested. However, the 2021 flood further revealed significant shortcomings in the early warning system chains. While exceptionally extreme precipitation was predicted by weather models (Busker et al., 2025; Mohr et al., 2023), and warnings were issued 1–2 days in advance (DWD, 2021; KNMI, 2021), these signals did not always lead to the necessary actions (Da Costa et al., 2026; Thicken et al., 2023a), showcasing the potential of early warning systems. Extreme precipitation was predicted by weather models, such as the model operated by the German Weather Service (DWD, 2021; Mohr et al., 2023), which issued extreme weather warnings 1–2 days in advance (DWD, 2021; KNMI, 2021). Despite these promising numbers, the impacts of the flood event were devastating and raised questions about the functioning of the operational flood warnings and the effectiveness of early actions taken.

1.1 A Flood Forecasting and Early Warning Systems (FFEWS)

A flood forecasting and early warning system (FFEWS) consists of a chain of components (Fig. 1). Hydro-meteorological data is used as input to weather and flood forecasting models, which jointly predict whether warning thresholds are surpassed with a certain probability (e.g., Alfieri et al., 2019). These thresholds can be, for example, pre-defined rainfall amounts or water levels with a certain return period. Surpassing a threshold may lead to a warning to the public or crisis managers who make decisions on how to respond (Fig. 1). Meteorological and hydrological warning levels are often depicted with in colour, as shown in Fig. 1. Each warning level can be connected to a set of pre-defined actions, outlined in crisis

75 management plans. Responses at the governmental level include, for example, closing a certain parking garage because it might be flooded, or in severe cases issuing an evacuation alert or order. Households may decide to install flood protection measures such as sandbags or mobile protection systems (Cao et al., 2024).

80 Uncertainty is present in each component of the FFEWS. In the forecasting part, uncertainty can be partly-partially displayed by using ensemble weather forecasts (Fig. 1, left). In addition, uncertainty of forecasts increases with lead time (i.e., the time between issuing a forecast and the real event) - usually one or a few days ahead of the event (Jiang et al., 2023). Further



85 uncertainties in the system lie in the communication, decision-making, and response stages, and whether decision-makers can process warning signals into effective actions on the ground (Bischiniotis et al., 2020). A key challenge in developing and implementing a FFEWS is that uncertainty in one component can cascade down to the next component, all the way to the last component (Parker and Priest, 2012). For example, uncertainty in the collection and processing of meteorological data (Fig. 1, Left) may lead to a 'missed' event, where thresholds in the system are not exceeded, while observed water levels reach extreme heights (Cloke and Pappenberger, 2009). By contrast, uncertainties can also lead to so-called false alarms, where predictions suggest that warning thresholds may be exceeded, while in reality they are not. For example, uncertainty in the collection and processing of meteorological data (Fig. 1, Left) may lead to a 'missed' event, where thresholds in the system were not surpassed, while observed water levels reached extreme heights (Cloke and Pappenberger, 2009). By contrast, uncertainties can also lead to so-called false alarms, where the

90

~~predictions suggest that warning thresholds may be exceeded, while effectively they are not.~~

95 **Figure 1: Conceptual representation of a Flood Forecasting and Early Warning System (FFEWS). Rainfall ensemble forecasts (left) are used in hydrological models to predict river water levels (middle). Warnings are issued if a share of the ensembles exceed the specified rainfall or water level threshold, which results in triggering (sometimes pre-determined) early actions (right) (adapted from: Busker, 2024).**

100 ~~Numerical weather prediction models have improved greatly in recent decades (Bauer et al., 2015). For example, the Integrated Forecasting System (IFS) operated by the European Centre for Medium-Range Weather Forecasts (ECMWF) showed strong improvements over the last 25 years (Buizza, 2019). The quality of meteorological those forecasts is challenged by the inherent uncertainties in atmospheric processes and initial conditions, which can lead to errors in spatio-temporal or intensity estimates (e.g., Leutbecher and Palmer, 2008). The first source of error relates to input data for forecasting models, while the second arises from the imperfect physical parametrization and (coarse) resolutions of models (Buizza, 2019). Numerical weather prediction models have improved greatly in recent decades (Bauer et al., 2015). For example, the Integrated Forecasting System (IFS) operated by the European Centre for Medium-Range Weather Forecasts (ECMWF) showed strong improvements over the last 25 years (Buizza, 2019). They started with ensemble forecasting in 1992 (Buizza, 2019), and now produce a 51-member ensemble forecast for both the medium-range (15 days ahead) and seasonal range (7 and 12 months ahead), and a 101-member forecast for the sub-seasonal range (46 days ahead).~~ As with meteorological forecasts, the quality of the hydrological forecasts is affected by various factors, such as meteorological observations or hydrological characteristics (e.g. ground- and soil water volume). Over the last decade, ensemble and post-processing techniques have been increasingly adopted in operational forecasting centers (Speight et al., 2021; Wu et al., 2020), used to estimate forecast the uncertainty in hydrological forecasts. Ensemble forecasts are used to calculate determine the probability of certain events, calculated as the share of ensemble members in case some (or all) ensemble members exceed exceeding a certain pre-determined threshold (e.g. precipitation or discharge) of heavy precipitation intensity. Forecast users can now include this probabilistic information in incorporate the probability of a certain event as additional information in the decision-making processes, which makes them more valuable than deterministic forecasts. Therefore, ensemble forecasts have higher value for forecast users than deterministic forecasts (Richardson, 2003). It is expected that ensemble forecasts will be widely integrated in operational forecasting chains in the near future (Pappenberger et al., 2016). They startThe ECMWF started with with ensemble forecasting in 1992 (Buizza, 2019), and now produce a 51-member ensemble forecast for both the medium-range (15 days ahead) and seasonal range (7 and 12 months ahead), and a 101-member forecast for the sub-seasonal-range (46 days ahead).

1.2 Current challenges in FFEWS

125 During the 2021 floods, a large share of affected people were not warned (Thieken et al., 2023b, Endendijk et al., 2023a). Moreover, For example, following an in-depth investigation of official parliamentary documents, Thieken et al. (2023b) showed that around a third of the flood-related fatalities in the German region North Rhine-Westphalia (NRW) were likely not warned. In addition, from the people that were warned, 85% did not expect severe flooding (Thieken et al. 2023a). In the

Netherlands, 75% of people with flooded homes in the heavily affected Geul catchment were not warned (Endendijk et al., 2023a). In some regions, the 2021 flood extended beyond the defined static flood hazard zones (as defined in spatial planning policies), giving citizens a false sense of security, even if they received warnings. This led to households outside of the flood zone being caught by surprise in the NRW and the Rhineland-Palatinate (RLP) regions in Germany (Rhein and Kreibich, 2025; Thieken et al. 2023b), and the Walloon region in Belgium (Dewals et al., 2021). Since households living outside the official flood zones have lower risk perceptions, their responses are also lower than those living in flood zones (Aerts et al., 2018). Empirical data in the affected areas in The Netherlands (Limburg) show that households who received a flood warning were 23% up to twice as likely to take emergency flood risk reduction measures as those who did not receive a warning (Endendijk et al., 2023a). However, this also means that some warned people did not evacuate or take other precautionary measures. These examples point to the importance of clear and tailored communication of the expected flood, the expected impacts, and of providing recommendations for responses (Thieken et al., 2023a). warned people often did not take action due to an underestimation of the severity of the situation (Thieken et al., 2023a). Therefore, research stresses the importance of providing impact and action information in a forecast (Najafi et al., 2024; Kreibich et al., 2021). Impact-based forecasts are more specific and tangible (Merz et al., 2020), trigger a change in people's risk perception of risks, and consequently their intention to respond (Endendijk et al. 2023b; Red Cross Red Crescent Climate Centre, 2020). Moreover, impact-based warnings provide crisis managers and first responders with more tailored information necessary to act (Apel et al., 2022; Boulton et al., 2022; Red Cross Red Crescent Climate Centre, 2020; Meléndez-Landaverde et al., 2020, Merz et al., 2020, Rhein and Kreibich, 2025, Poolman et al., 2018). However, the practice of impact-based forecasting is still in its infancy (Merz et al., 2020). Only about 30% of the national meteorological and hydrological services in Europe use impact-based models (Kaltenberger et al., 2020; Schroeter et al., 2021).

Furthermore, many countries and regions have not yet homogenized their FFEWS protocols, which can further hamper clear and consistent cross-border communication (Coughlan de Perez et al., 2022). This is caused by, for example, by differences in institutional settings and geographical characteristics in the respective regions. The Council of the European Union states that a better integration of activities and cooperation between countries in cross-border river-basins is crucial to further develop FFEWSs (Council of the European Union, 2024).

Research stresses the importance of providing impact and action information in a forecast (Najafi et al., 2024; Kreibich et al., 2021). Impact-based forecasts are more specific and tangible (Merz et al., 2020), trigger change in people's risk perceptions, and consequently their intention to respond (Endendijk et al. 2023b; Red Cross Red Crescent Climate Centre, 2020). Moreover, impact-based warnings provide crisis managers and first responders with more tailored information necessary to act (Apel et al., 2022; Boulton et al., 2022; Red Cross Red Crescent Climate Centre, 2020; Meléndez-Landaverde et al., 2020, Merz et al., 2020, Rhein and Kreibich, 2025, Poolman et al., 2018). However, the practice of impact-based forecasting is still in its infancy (Merz et al., 2020). Only about 30% of the national meteorological and hydrological services in Europe use impact-based models (Kaltenberger et al., 2020; Schroeter et al., 2021).

160 1.3 Research gap and aim

Given these challenges, it is crucial to compare and assess the current state of operational FFEWSs. Only very few studies assess operational FFEWSs in detail (e.g. Kaltenberger et al., 2020; Schroeter et al., 2021), and to the best of our knowledge none of them has assessed all components of the FFEWSs chain, including communication and crisis management. Moreover, a transboundary analysis (e.g. an international comparison of warning levels) is lacking. Therefore, the main goal of this study

165 is to assess and compare the FFEWSs in transboundary river basins in Northwestern Europe, based on insights from semi-structured interviews with experts across the region and a literature review. We focus on the areas hit by the July 2021 flood, specifically The Netherlands (Limburg), the Rhineland-Palatinate (RLP) and North Rhine-Westphalia (NRW) federal states in Germany, Belgium (Flanders and Wallonia) and Luxembourg. We intend (a) to compare the institutional and operational characteristics of FFEWSs (e.g. differences in warning levels across countries and communication protocols), b) to outline and
170 compare the changes made after the 2021 flood event and (c) to develop recommendations to further improve the systems. This will provide a comprehensive overview of the state of FFEWSs and pathways towards translating forecasts to more effective early action. We intend to gain and share knowledge on FFEWSs in a transboundary context and to develop recommendations based on insights from semi-structured interviews and a literature review. We focus on two questions related to the use of FFEWSs: (a) “what are the differences in the warning levels across countries and how are they defined?”,
175 and (b) “how can the communication of early warning information and the translation into effective action be improved?”. We focus on the areas hit by the July 2021 flood, specifically The Netherlands (Limburg), the Rhineland-Palatinate (RLP) and North Rhine-Westphalia (NRW) federal states in Germany, Belgium (Flanders and Wallonia) and Luxembourg.

2 Case study region and approach

2.1 Case study region

180 *The July 2021 flood event*

The summer flood event of ~~summer~~ 2021 severely hit the area shown in Fig. 2, which is comprised of several countries that share the Meuse and Rhine basins. The event was caused by an atmospheric low-pressure system ~~delivering that delivered~~ a total precipitation amount of about 200 mm in the Ardennes-Eifel region over 48 hours (Lehmkuhl et al., 2022; Mohr et al., 2023). The impacts were the largest in the valleys of smaller tributaries such as the Ahr and Vesdre (Fig. 2, red polygons),
185 with the most severe damage in the Ahr Valley, where 135 people lost their lives (out of in total 190 fatalities in Germany; see Thielen et al., 2023b). In Belgium, parts of the Vesdre valley were devastated, ~~where-while~~ in total 39 people lost their lives (Journée et al., 2023). For The Netherlands, no fatalities were reported, but damage amounted to more than half a billion euros, ~~-particularly in the Geul tributary (ENW, 2021). Also in Luxembourg, river gauges recorded unprecedented water levels exceeding~~ the 100-year return period, which led to widespread flooding that led to widespread inundation.

190

The flood revealed considerable shortcomings in the FFEWSs in the region. Although official evacuation orders or advice were eventually given over most of the region, they were mostly issued too late, just before the peak floodwaters arrived, such as in the Ahr catchment (Thieken et al., 2023a; Rhein and Kreibich, 2025), in the Geul catchment in the south of The Netherlands (Endendijk et al., 2023a), and in the Vesdre catchment in Belgium (Dewals et al., 2021). Moreover, In the Vesdre, Ahr and Geul catchments, a significant share part of the local population in the affected catchments did not receive the warnings (see Introduction). In the Netherlands, 75% of people with flooded homes in the heavily affected Geul catchment were not warned (Endendijk et al., 2023a). Following an in-depth investigation of official parliamentary documents, Thieken et al. (2023b) showed that around a third of the flood-related fatalities in the German region North Rhine-Westphalia (NRW) were likely not warned. In addition, from the people that were warned, 85% did not expect severe flooding (Thieken et al., 2023a). In some regions, the flood extended beyond the defined static flood hazard zones (as defined in spatial planning policies), giving citizens a false sense of security, even if they received warnings. This led to households outside of the flood zone being caught by surprise in the NRW and the RLP regions in Germany (Rhein and Kreibich, 2025; Thieken et al., 2023b), and the Walloon region in Belgium (Dewals et al., 2021). Since households living outside the official flood zones have lower risk perceptions, their responses are also lower than those living in flood zones (Aerts et al., 2018). Empirical data in the affected areas in The Netherlands (Limburg) show that households who received a flood warning were 23% up to twice as likely to take emergency flood risk reduction measures as those who did not receive a warning (Endendijk et al., 2023a). However, this also means that some warned people did not evacuate or take other precautionary measures. These examples show the importance of investing in warning communication. Warning messages should be tailored to users, tailored communication of the expected flood, with information on the expected flood, the expected impacts, and and of with actionable providing recommendations for responses (Thieken et al., 2023a). Although official evacuation orders or advices were eventually given over most of the region, they were mostly issued too late, just before the peak floodwaters arrived, such as in the Ahr catchment (Thieken et al., 2023a, Rhein and Kreibich, 2025), in the Geul catchment in the south of The Netherlands (Endendijk et al., 2023a), and in the Vesdre catchment in Belgium (Dewals et al., 2021). Although official evacuation orders or advices were eventually given over most of the region, they were mostly issued too late, just before the peak floodwaters arrived, such as in the Ahr catchment (Thieken et al., 2023a, Rhein and Kreibich, 2025), in the Geul catchment in the south of The Netherlands (Endendijk et al., 2023a), and in the Vesdre catchment in Belgium (Dewals et al., 2021). In some regions, the 2021 flood extended beyond the defined static flood hazard zones (as defined in spatial planning policies), giving citizens a false sense of security, even if they received warnings. This led to households outside of the flood zone being caught by surprise in the NRW and the Rhineland-Palatinate (RLP) regions in Germany (Rhein and Kreibich, 2025; Thieken et al. 2023b), and the Walloon region in Belgium (Dewals et al., 2021). Since households living outside the official flood zones have lower risk perceptions, their responses are also lower than those living in flood zones (Aerts et al., 2018). Empirical data in the affected areas in The Netherlands (Limburg) show that households who received a flood warning were 23% up to twice as likely to take emergency flood risk reduction measures as those who did not receive a warning (Endendijk et al., 2023a). However, this also means that some warned people did not evacuate or take other precautionary measures.

Formatted: Font: Not Italic, English (United States)



Figure 2: The study area in Northwestern Europe with the transboundary Rhine and Meuse rivers, shared by The Netherlands, Germany, ~~Luxembourg~~ and Belgium. These countries and sub-regions were severely impacted by the extreme flood event in July 2021, especially in smaller transboundary tributaries such as the Geul, Vesdre and Ahr (red polygons).

230 *Governance of Flood Forecasting and Early Warning Systems (FFEWS)*

Strong differences in FFEWS ~~approaches-governance are~~ present ~~aeross-in~~ the study area. In The Netherlands, the national government ~~is-managing-makes forecasts for~~ the large waterways (e.g., Rhine and Meuse), while regional governments (including regional water authorities: “~~W~~waterschappen”) are responsible for the smaller regional water systems. ~~The Netherlands organizes crisis management differently than the other countries: S~~pecial administrative regions are ~~formed~~ ~~formed~~ for the crisis management ~~organizations~~, the so-called “Safety Regions” (Dutch: ~~V~~veiligheidsregioes). ~~In~~ Germany, flood ~~risk-forecasting and~~ management is the responsibility of each of the 16 federal States (e.g., Becker et al., 2015). ~~HereIn~~ ~~this study~~, we ~~particularly-focus~~ ~~particularly~~ on flood forecasting and warning deployed in the riparian states of RLP and NRW in Germany. In Belgium, the two federal states (Flanders and Wallonia) ~~both~~ have the mandate to develop and enforce flood risk management regulations. In Luxembourg, flood forecasting is addressed at the national level by the Water Management ~~Administration (AGE), which is part of the Ministry of the Environment, Climate and Biodiversity.~~

2.2 Approach

~~Most of the information in this study was obtained from key-informant interviews. We collected information on FFEWSs in the riparian regions from literature reviews and interviews with 134 experts in the countries of the case study area. These~~

245 ~~experts consisted of senior employees of~~ Additionally, 13 key informant interviews were held with flood forecasting experts
250 ~~from national hydro-meteorological and crisis-management agencies or with agencies and experts involved in crisis~~
~~management~~ or full professors employed at universities (Appendix A). For every country, we searched for at least one
~~forecasting expert at a governmental forecasting agency~~ expert in a forecasting agency and one ~~person~~ directly involved in
255 ~~crisis management, to ensure diversity in the organizations involved. For the interviews, we developed a semi-structured~~
~~questionnaire (Supplement S1) to inquire about~~ specific questions related to evaluating current FFEWSs. We structured
260 ~~the interview in the five different FFEWS pillars, as outlined by the World Meteorological Organization (WMO, 2023): (1)~~
~~data collection, (2) forecasting, (3) communication, (4) decision making, (5) action/response. The division of the interview~~
~~into five pillars facilitated a thematic analysis (Kiger & Varpio, 2020). Within each pillar, we manually synthesized the~~
~~responses, identifying patterns, commonalities, and differences between the different organizations, based on the audio~~
~~recordings obtained with the participant's consent. All interviewees were comfortable communicating in~~ speaking English,
265 ~~so translation of the recordings was not necessary~~ it was not necessary to translate the recordings. We included an additional
iteration of the synthesized material with most interviewees, to ensure correct interpretation of the interviews. To contextualize
our findings, we integrated insights from the literature, comparing our findings with state-of-the-art scientific research. The
interviewees provided us with a total of 14 country-specific reports in national languages and 16 scientific articles. To
270 ~~supplement this information, Below, we discuss the results for each country. We~~ applied a non-systematic literature review
~~literature search~~ with Google Scholar and Scopus, using (combinations of) these main keywords: #flood, #forecast, #warning,
#communication, #response, #evacuation, #residents, #government, #rainfall, #precipitation, #extreme, #waterlevels, #actions,
#alerts, #perception, #Belgium, #Luxembourg, #Netherlands, and #Germany. From the search results, we selected 40 peer-
275 ~~reviewed scientific papers~~ To describe the current state of flood-early warning systems, we selected 16 scientific peer-reviewed
articles for an in-depth analysis. We also manually analyzed 14 country-specific reports that describe or evaluate flood
forecasting systems in national languages and 5 reports from international organizations (e.g. The Red Cross and the WMO).
This selection was primarily performed by experts from the region, selected by experts from the case study
regions, who were either co-authors or interviewees. Additionally, 13 key informant interviews were held with flood
forecasting experts from national hydro-meteorological agencies or with agencies and experts involved in crisis management
(Appendix A). For every country, we searched for at least one forecasting expert at a governmental forecasting agency and
one person directly involved in crisis management, to ensure diversity in the organizations involved. For the interviews, we
developed a semi-structured questionnaire (Supplement S1) to inquire about specific questions related to evaluating current
FFEWSs. We structured the interview in the five different FFEWS pillars, as outlined by WMO (2023): (1) data collection,
(2) forecasting, (3) communication, (4) decision-making, (5) action/response. The division of the interview into five pillars
facilitated a thematic analysis (Kiger & Varpio, 2020). Within each pillar, we manually synthesized the responses, identifying
275 ~~patterns, commonalities, and differences between the different organizations, based on the audio recordings obtained with the~~
~~participant's consent. All interviewees were comfortable in speaking English, so translation of the recordings was not~~
~~necessary. We included an additional iteration of the synthesized material with most interviewees, to ensure correct~~


~~interpretation of the interviews. To contextualize our findings, we integrated insights from the literature, comparing our findings with state-of-the-art scientific research. Below, we discuss the results for each country.~~

280 3 Key characteristics of the operational FFEWS

In this section, the key characteristics of FFEWS are presented based on recent literature and ~~the~~ expert interviews. First, we present a general overview of the different systems (Section 3.1), after which we discuss the warning levels used (Section 3.2), a detailed overview per country (Sections 3.3-3.6), and finally provide an overview of collaborations between the different regions (Section 3.7).

285 3.1 Overview of the operational FFEWS

Table 1 summarizes the key characteristics of FFEWS ~~over in~~ the region hit by the 2021 flood event, reflecting the state of the systems at the time of this study (5 years after the event). The meteorological agencies use different forecasting systems, and all use Ensemble Prediction Systems (EPS) to represent uncertainty (Table 1). The hydrological forecasting agencies also use different hydrological models for flood discharge forecasting. In the German regions RLP and NRW and in Luxemburg, the LARSIM hydrological model is used. Even if the same hydrological model is used, important parameters in the model setup (e.g. spatial resolution: Haag and Luce, 2007) are often user-dependent and can still differ between the different countries. The main forecast models in The Netherlands are HBV and SOBEK, run on the Delft-FEWS system. In Flanders (Belgium), the NAM and MIKE11 models are used for the larger and navigable rivers on the FEWS-Flanders system, and PDM and InfoWorks ICM for the non-navigable rivers. In Wallonia (Belgium), the Hydromax model is most widely used (Table 1). Beyond differences in model types, large differences are seen in the forecast lead time, ranging from only 24-48 hours (RLP, Flanders and Luxembourg) to 15 days (The Netherlands). Most operational flood forecasting centers use probabilistic forecasts to account for uncertainty, except for some regional water authorities in The Netherlands. Moreover, they use ~~and use~~ post-processing methods to correct forecast outputs, based on historical forecast errors, aimed at increasing the forecast skill. ~~Where the other regions rely on discharge forecasts and thresholds,~~ Flanders is the only region where flood inundation forecasts are run operationally for short-term (48-hour) lead times, whereas all other regions rely on discharge forecasts and thresholds. Flanders is the only region where flood inundation forecasts run operationally for 48h forecasts. The two lowest rows of Table 1 show information related to crisis management. All regions have flood emergency management plans ~~for floods.~~ In Germany and Luxembourg, a cell broadcasting system was installed in response to and after the July 2021 flood, whereas all other regions had such a system in place prior to the 2021 flood. All regions have online portals with flood warning information, which is updated regularly during flood situations. Publicly available flood forecast ~~ing~~ information is available via online portals for all regions, except for NWR and Wallonia, where only gauge-based measurements and respective warnings are published.



315 **Table 1. Characteristics of Flood Forecasting and Early Warning Systems (FFEWS) in different European regions and countries, reflecting the state of the systems at the time of this study, after the 2021 flood event. Characteristics of Flood Forecasting and Early Warning Systems (FFEWS) in different European regions and countries.**

	Country	Germany (RLP)	Germany (NRW)	Luxembourg	The Netherlands	Belgium (Flanders)	Belgium (Wallonia)
Meteo forecasts	Forecast agency	DWD		Meteolux	KNMI	RMI	
	Main ensemble model(s)	ICON-D2-EPS and ICON-EPS		ICON-(D2-)EPS, ECMWF-ENS, AROME-EPS	HarmonEPS and ECMWF-ENS	ECMWF-ENS (AROME-EPS is in development)	
Flood forecasts	Forecasting agency	LFU	LANUK	AGE	WMCN and regional water authorities	HIC and VMM	SPW
	Main hydrological model(s)	LARSIM	LARSIM	LARSIM	HBV and SOBEK	NAM, MIKE11, PDM and InfoWorks ICM	Hydromax
	Discharge (D) or flood inundation (I) forecast (lead time)	D (48 h)	D (10-days)	D (24-48 h)	D (5-days and 15-days)	D (10-days), I (48 h)	D (lead time dependent on catchment size)
	Post-processing technique on discharge forecasts	ProFoUnD analysis (Haag et al., 2013).	Unknown	Yes, bias correction	Ensemble dressing (Verkade et al., 2017)	Non-parametric (Van Steenberghe et al., 2012)	Unknown
	Probabilistic forecasts?	Yes	Yes	Yes	National level (WMCN): Yes Regional water authorities: Not always	Yes	Yes
Online portal(s)	www.hochwasserzentralen.de RLP: https://www.hochwasser.rlp.de NRW: https://hochwasserportal.nrw (only observations; forecasts in preparation)		https://www.inondations.lu/map	https://waterinfo.rws.nl	Flanders: https://www.waterinfo.be Wallonia: https://hydrometrie.wallonie.be (only observations)		
Crisis management	Flood Emergency management plans for floods	Yes	Yes, but not specifically for floods	Yes	Yes	Yes, but not specifically for floods	Yes, but not specifically for floods
	Primary warning communication system (start date of operation)	MoWas (October 2001) and DE-alert (February 2023)		LU-alert (October 2024)	NL-alert (November 2012)	BE-alert (June 2017)	

320 The hydrological forecasting agencies also use different hydrological models for flood discharge forecasting. In the German
regions RLP and NRW and in Luxemburg, the LARSIM hydrological model is used. The main forecast models in The
Netherlands are HBV and SOBEK, run on the Delft FEWS system. In Belgium, different systems are used in Flanders (NAM
and MIKE11 on the FEWS Flanders system) and Wallonia (Hydromax) (Table 1). Most operational flood forecasting centers
use probabilistic forecasts to account for uncertainty (except for some regional water authorities in the Netherlands) and use
325 discharge forecasts and thresholds, Flanders is the only region where flood inundation forecasts run operationally for 48h
forecasts. The two lowest rows of Table 1 show information related to crisis management. All regions have emergency
management plans for floods. In Germany and Luxemburg, a cell broadcasting system was installed in response to and after
the July 2021 flood, whereas all other regions had such a system in place prior to the 2021 flood. All regions have online
portals with flood warning information, which is updated regularly during flood situations. Publicly available flood forecasting
330 information is available via online portals for all regions but NWR and Wallonia, where only gauge-based measurements and
respective warnings are published.

3.2 Warning levels

Meteorological warnings


335 Meteorological warnings are issued based on clearly defined rainfall intensity and duration thresholds (Fig. 3). The thresholds
are chosen based on expected impacts to society and are different across countries. Moreover, as those impacts are highly
dependent on the situation (e.g. day or night, location, large events), the thresholds are mostly used only as guidance ~~in praetiee~~.
The meteorological agencies generally use three rainfall thresholds, except for Luxemburg where four levels are used. In The
Netherlands and Belgium, yellow-orange-red ~~color~~ colours are used as warning levels. Already before 2021, the German
Weather Service (DWD) in Germany ~~deploys-issued~~ an additional purple warning for “extreme weather” events. In response
340 to the 2021 flood, Luxemburg implemented a new warning level for “immediate danger” in November 2024, ~~in purple. with~~
~~a purple color~~. This warning requires “immediate action”, such as with rapidly-developing flash-floods. ~~For the lowest~~
~~threshold level, all countries use values of ~30mm/ hour or ~40mm/ 6 hours, whereas the highest threshold is often related to~~
~~an event of 80 to 100mm/ 24 hours (among other thresholds).~~ ~~For the lowest threshold level, all countries use values of~~
~~~30mm/hr or ~40mm/6hrs, whereas the highest threshold is often related to an event of 80 to 100mm/24hrs (among other~~  
345 ~~thresholds).~~ ~~Only in The Netherlands, the red warning level is not distinguished from orange on the basis of rainfall, but only~~  
~~based on expected impacts.~~ ~~Only in The Netherlands, the red warning level is not distinguished from orange based on rainfall~~  
~~but only based on expected impacts.~~ These impacts are estimated in an impact-team meeting with key stakeholders (see Sect.  
3.53).

### *Hydrological warnings*

350 ~~Figure 3 shows that flood warning levels differ between countries and between regions.~~ ~~Figure 3 shows that flood warning~~  
~~levels differ between countries, as well as between regions.~~ For example, Flanders and Wallonia, or NRW and RLP, have  
different ~~color~~colour codes, which are defined in a different way. The number of hydrological warning levels varies between  
3 in Wallonia ~~and~~to 6 in RLP (Fig. 3). Hydrological thresholds are mostly clearly defined in all regions, based on specific  
355 level, and the warning thresholds can therefore differ among different events. In the other regions, the thresholds are based on  
either return periods (e.g., in RLP), or local flood impacts (e.g., in NRW, Luxembourg, Flanders). Warnings in Flanders slightly  
deviate from the other countries and use different water level thresholds for: (a) Alarm States, based on impact to water  
infrastructures, and (b) Flood Warnings, based on the distance between the water level and ~~the~~the dike crest (Fig. 3 and Sect.  
3.64). Red is the highest warning level in all regions, except for the German federal state RLP. In response to the 2021 flood-,  
360 the flood forecasting center in RLP (Landesamt für Umwelt Rheinland-Pfalz) subdivided the already existing purple category  
for extreme events into purple (> 1/50 year event) and dark purple (>1/100 year event) to create the opportunity to point out  
an “extremely dangerous flood event” (Landesamt für Umwelt Rheinland-Pfalz, 2025).

**WARNING LEVELS AND THRESHOLDS**

| Country                                                                                                  | Pluvial floods (National Meteorological Services)                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | Fluvial floods (Hydrological services)                                                                                                                                                                                                                                                        |                                                                                                                                                                                                                                                                                                                                                            |
|----------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|  <p>The Netherlands</p> | <ul style="list-style-type: none"> <li>⚠️ <math>\geq 30</math> mm in 1 h (thunderstorms) or <math>&gt; 50</math> mm in 24 h</li> <li>⚠️ <math>\geq 50</math> mm in 1 h or <math>\geq 75</math> mm in 24 h (heavy rain)</li> <li>⚠️ On the decision of the Weather Impact Team</li> </ul>                                                                                                                                                                                                                                    | <p><b>Large rivers:</b> based on impact-adjusted return periods</p>  <p>1/25 to 1/50 years RP<br/>1/5 to 1/10 years RP<br/>Yearly RP<br/>sub-yearly RP*</p>                                                  | <p><b>Regional rivers (Local water authorities):</b> Thresholds depending on the water authority</p>                                                                                                                                                                                                                                                       |
|  <p>Belgium</p>         | <ul style="list-style-type: none"> <li>⚠️ 20 to 30 mm in 1 h or 20 to 40 mm in 6 h or 25 to 50 mm in 24 h</li> <li>⚠️ 31 to 50 mm in 1 h or 41 to 60 mm in 6h or 51 to 100 mm in 24 h</li> <li>⚠️ Already flooding problems and heavy rain still forecasted or 50 mm in 1 h or 60 mm in 6h or 100 mm in 24h</li> </ul>                                                                                                                                                                                                      | <p><b>Flanders:</b><br/>Alarm states: water level thresholds, based on impact to water infrastructures<br/>Flood warnings: water level thresholds, based on distance between water level and dike crest</p>  | <p><b>Wallonia:</b><br/>Thresholds not present, warning level selected based on expert-judgement</p>                                                                                                                                                                      |
|  <p>Luxembourg</p>     | <ul style="list-style-type: none"> <li>⚠️ Awareness level yellow / Be aware (low risk / potential danger)<br/>15-30 mm in 6h or 20-40 mm in 12h or 30-50 mm in 24h</li> <li>⚠️ Awareness level orange / Be careful (medium risk / danger)<br/>31-45 mm in 6h or 41-60 mm in 12h or 51-81 mm in 24h</li> <li>⚠️ Awareness level red / Utmost vigilance (high risk / significant danger)<br/>&gt;45 mm in 6h or &gt;60 mm in 12h or &gt;80 mm in 25h</li> <li>⚠️ Imminent danger / Immediate action (flash floods)</li> </ul> |  <p>Based on local flood impact</p>                                                                                                                                                                         |                                                                                                                                                                                                                                                                                                                                                            |
|  <p>Germany</p>       | <ul style="list-style-type: none"> <li>⚠️ 15 to 25 mm in 1 hour or 20 to 35 mm in 6 h or 25 to 40 mm in 12 h or 30 to 50 mm in 24 h or 40 to 60 mm in 48 h or 60 to 90 mm in 72 h</li> <li>⚠️ 25 to 40 mm in 1 hour or 35 to 60 mm in 6 hours or 40-70 mm in 12 h or 50-80 mm in 24 h or 60-90 mm in 48 h or 90-120 mm in 72 h</li> <li>⚠️ &gt; 40 mm in 1 hour or &gt; 60 mm in 6 h or &gt; 70 mm in 12 h or &gt; 80 mm in 24 h or &gt; 90 mm in 48 h or &gt; 120 mm in 72 h</li> </ul>                                    | <p><b>North Rhine Westphalia</b><br/>Based on local flood impact</p>                                                                                                                                       | <p><b>Rhineland-Palatinate</b><br/>Based on return periods</p>  <p><math>\geq 100</math> year RP<br/><math>\geq 50</math> year RP<br/><math>\geq 20</math> year RP<br/><math>\geq 10</math> year RP<br/><math>&gt; 2</math> year RP<br/><math>&lt; 2</math> year RP</p> |

| WARNING LEVELS AND THRESHOLDS |                                                                                                                                                                                                                                            |                                                                                                                                                                                                                                                                                         |                                                                                                                                                |
|-------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------|
| Country                       | Pluvial floods (National Meteorological Services)                                                                                                                                                                                          | Fluvial floods (Hydrological services)                                                                                                                                                                                                                                                  |                                                                                                                                                |
| The Netherlands               |  $\geq 30$ mm in 1 h (thunderstorms)<br>or $\geq 50$ mm in 24 h                                                                                           | <i>Large rivers:</i> based on impact-adjusted return periods<br>                                                                                                                                       | <i>Regional rivers (water boards):</i> thresholds depending on the water board                                                                 |
|                               |  $\geq 50$ mm in 1 h<br>or $\geq 75$ mm in 24 h (heavy rain)                                                                                              |                                                                                                                                                                                                                                                                                         |                                                                                                                                                |
|                               |  On the decision of the Weather Impact Team                                                                                                               |                                                                                                                                                                                                                                                                                         |                                                                                                                                                |
| Belgium                       |  20 to 30 mm in 1 h<br>or 20 to 40 mm in 6 h<br>or 25 to 50 mm in 24 h                                                                                    | <i>Flanders:</i><br>Alarm states: water level thresholds, based on impact to water infrastructures<br>Flood warnings: water level thresholds, based on distance between water level and dike crest<br> | <i>Wallonia:</i><br>Thresholds not present, warning level selected based on expert-judgement                                                   |
|                               |  31 to 50 mm in 1 h<br>or 41 to 60 mm in 6h<br>or 51 to 100 mm in 24 h                                                                                    |                                                                                                                                                                                                                                                                                         |                                                                                                                                                |
|                               |  Already flooding problems and heavy rain still forecasted<br>or 60 mm in 1 h<br>or 60 mm in 6h<br>or 100 mm in 24h                                       |                                                                                                                                                                                                                                                                                         |                                                                                                                                                |
| Luxembourg                    |  Awareness level yellow / Be aware (low risk / potential danger)<br>15-30 mm in 6h<br>or 20-40 mm in 12h<br>or 30-50 mm in 24h                            |                                                                                                                                                                                                       | Based on local flood impact                                                                                                                    |
|                               |  Awareness level orange / Be careful (medium risk / danger)<br>31-45 mm in 6h<br>or 41-60 mm in 12h<br>or 51-81 mm in 24h                                 |                                                                                                                                                                                                                                                                                         |                                                                                                                                                |
|                               |  Awareness level red / Utmost vigilance (high risk / significant danger)<br>>45 mm in 6h<br>or >60 mm in 12h<br>or >80 mm in 25h                        |                                                                                                                                                                                                                                                                                         |                                                                                                                                                |
|                               |  Imminent danger / Immediate action (flash floods)                                                                                                      |                                                                                                                                                                                                                                                                                         |                                                                                                                                                |
| Germany                       |  15 to 25 mm in 1 hour<br>or 20 to 35 mm in 6 h<br>or 25 to 40 mm in 12 h<br>or 30 to 50 mm in 24 h<br>or 40 to 60 mm in 48 h<br>or 60 to 90 mm in 72 h | <i>North Rhine Westphalia</i><br>Based on local flood impact<br>                                                                                                                                     | <i>Rhineland-Palatinate</i><br>Based on return periods<br> |
|                               |  25 to 40 mm in 1 hour<br>or 35 to 60 mm in 6 hours<br>or 40-70 mm in 12 h<br>or 50-80 mm in 24 h<br>or 60-90 mm in 48 h<br>or 90-120 mm in 72 h        |                                                                                                                                                                                                                                                                                         |                                                                                                                                                |
|                               |  > 40 mm in 1 hour<br>or > 60 mm in 6 h<br>or > 70 mm in 12 h<br>or > 80 mm in 24 h<br>or > 90 mm in 48 h<br>or > 120 mm in 72 h                        |                                                                                                                                                                                                                                                                                         |                                                                                                                                                |

**Figure 3: An overview of the operational warning levels for pluvial (left) and fluvial (right) floods, as in place six years after the 2021 floods. Colour-coded warning levels are shown, with the corresponding rainfall thresholds (left) and water level thresholds (right). In case the water level thresholds are based on return periods (RPs), the corresponding RP is shown as well.**

370 ~~overview of the operational warning levels for pluvial (left) and fluvial (right) floods. The colors shown for warning levels correspond to the colors used operationally.~~

In the following, we provide an overview per country with (a) a general description of the FFEWSs and a more in-depth assessment of the current (b) meteorological and (c) hydrological forecasts, and (d) ~~cdiscussion on the characteristics of crisis management and communication, with an emphasis on experiences during the 2021 flood event.~~

### 375 3.3 Germany

#### *General FFEWS setup*

~~At present~~Currently, the legal responsibility for weather forecasting, ~~–flood forecasting, and response in Germany is split between the German Weather Service (DWD), flood forecasting centers in the federal states (Länder), and flood emergency management in the counties (Landkreise), respectively. The DWD is mandated to~~ issue-produce weather forecasts and ~~provides~~ issue severe weather warnings. ~~The F~~federal state flood forecasting centers translate the weather forecast into hydrological forecasts and disseminate warning messages for affected catchments. In April 2024, the Federal Act on the DWD (Gesetz über den Deutschen Wetterdienst) was changed, ~~allowing the DWD so that the DWD is now allowed to also~~ disseminate warnings other than severe weather warnings, which include flood warnings. ~~In 2025, a national natural hazards online portal was released to share this information (www.naturgefahrenportal.de). –A national natural hazards online portal to share this information was released in 2025 (www.naturgefahrenportal.de).~~ Local emergency management authorities interpret the information from forecasting agencies and translate it ~~into~~ impact-specific warnings and are responsible for decision making and triggering specific emergency actions, such as evacuation. Warnings are disseminated to the public via the centralized Modular Warning System (MoWaS), operated by The German Federal Office of Civil Protection and Disaster Assistance (BBK).

#### 390 *Meteorological forecasts*

~~In Germany, DWD uses two ensemble forecast models, the ICON-D2-EPS (for the next 48 hours) and ICON-EPS (for the next 7.5 days).~~In Germany, DWD uses two ensemble forecast models, the ICON-D2-EPS (for the next 48 hours) and ICON-EPS (for the next 7.5 days). More recently, ~~DWD has developed a so-called “seamless” forecasting system to better forecast small-scale convective events and flash floods, called SINFONY. This system integrates ensemble weather forecasts and~~ nowcasting techniques (based on real-time radar measurements) over shorter lead times (Blahak et al., 2024). ~~Forecasts from~~The DWD forecasts are ~~delivered-sent~~ to the environmental offices and flood forecasting agencies of the federal states and to water associations (‘Wasserverbände’). These contain a description of the characteristics of severe weather events and the amount of expected precipitation. The thresholds and warning levels for heavy rainfall as used by the DWD are summarized in Fig. 3.

#### 400 *Hydrological forecasts*

~~The Ensemble flood forecasting models are operational in 10 flood forecasting centers in federal states, ensemble flood forecasting models are operational at 10 flood forecasting centers in the federal states.~~ Next ~~In addition~~ to those, regional flood forecasting centers are in operation, specifically for large navigable rivers such as the Rhine. The locations of gauging stations at which the hydrological discharge and water level forecasts are issued are provided by the German Working Group on water issues of the federal states (LAWA-Commission; LAWA, 2020). ~~The national web portal (Table 1) was established after the severe flood in August 2002 and provides an overview of the gauges with up-to-date flood forecasting and warning information.~~ ~~The national web portal (Table 1) was established after the severe flood in August 2002 and provides an overview of gauges with up-to-date flood forecasting and warning information.~~ Moreover, it serves as a gateway to the specific web services of flood forecasting centers in the federal states. It also ~~disseminates/distributes~~ written interpretations of the forecasts (in ~~the~~ so-called "Lageberichte"). The LAWA (2020) report ~~further-also~~ contains a detailed overview of the frequency of forecast updates for specific gauges in case of flooding, which typically ranges between 2 and 24 hours.

~~In Germany, many regional and federal state forecasting centres use a variety of different hydrological models and warning configurations (Demuth and Rademacher, 2016).~~ ~~In Germany, a variety of different hydrological models and warning configurations are used by many regional and federal state forecasting centers (Demuth and Rademacher, 2016).~~ The environmental office in RLP (Landesamt für Umwelt, LfU) is responsible for flood forecasting. They issue forecasts and warnings with a 48-hour lead time, based on a 20-member ensemble forecast from the DWD ICON-D2-EPS model. LfU uses a post-processing method to correct model outputs, based on ~~the Program to assess~~ the Forecast Uncertainty of Discharge (ProFoUnD) analysis (Haag et al., 2013). The forecasts are updated every three hours, which is the same frequency as the DWD forecasts. In NRW, the responsibility for flood forecasting belongs to the Landesamt für Natur, Umwelt und Klima Nordrhein-Westfalen (LANUK). A new flood forecasting ~~team-center~~ was formed in LANUK, as a response to the 2021 floods. Currently, water level observations at some gauges in NRW are accessible in near-real-time, but forecasts are not publicly available (Table 1). ~~In NRW, LANUK runs operational hydrological ensemble forecasts every 3 hours.~~ ~~In NRW, LANUK runs hydrological ensemble forecasts operationally every 3 hours.~~ These predictions are based on the Large Area Runoff Simulation Model (LARSIM) hydrological model and use weather forecast data from the ICON-D2-EPS (20 ensemble members) and ICON-EPS (40 ensemble members) models. The two regions, NRW and RLP, use different ~~color~~ ~~colour~~ codes and warning levels (Fig. 3). ~~Although the warning thresholds in RLP are based on water level return periods, the thresholds in NRW are based on local impacts on buildings and communities.~~ ~~While the warning thresholds in RLP are based on water level return periods, the thresholds in NRW are based on local impacts to buildings and communities.~~ NRW defines four different warning levels, while RLP uses six. NRW is in the process of adding an additional purple warning level as well, which represents a risk of catastrophic floods. This would better streamline the hydrological warnings of NRW and RLP and the meteorological warnings from DWD, which also include a purple level (Fig. 3). ~~Besides~~ ~~In addition to~~ some prototypes in pilot

areas, near-real-time flood inundation and impact forecasting are not yet implemented in the operational flood forecasting and warning systems in Germany (Merz et al., 2020).

#### 435 ***Crisis management and communication***

Warnings from the federal hydrological flood forecasting centers (e.g. LANUK in NRW) are sent to the county (Landkreis) crisis response centers. Based on laws regarding general civil assistance and disaster management, all counties, municipalities, and independent cities in Germany, which are not affiliated to a county, are required to have a disaster management plan (DKKV, 2024; Ministerium des Innern und für Sport Rheinland-Pfalz, 2020). RLP has developed a specific flood emergency management plan: the Rahmen-, Alarm- und Einsatzplan Hochwasser (Ministerium des Innern und für Sport Rheinland-Pfalz, 2020). The county crisis response centers, ~~not the federal state hydrological forecasting centers~~, are responsible for dissemination of the warnings to the public.

Both DWD and flood forecasting centers can activate MoWaS in case of an expected catastrophic flood and further disseminate information to other channels such as warning apps (Meine Pegel, NINA or KATWARN), radio, and television (IIASA, 2022). Since February 2023, the population can be warned using a new cell broadcasting system (DE-alert), which does not require the installation of an application and is connected to the MoWas system. In terms of online information, residents show a clear preference for maps in warning messages (Lindenlaub et al., 2024a).

### **3.4 Luxembourg**

#### 450 ***General FFEWS setup***

Meteorological forecasts are used by the flood forecasting department (Service de prévision des crues) of the water management authority (Administration de la gestion de l'eau, AGE). The AGE produces forecasts for the main rivers systems (the Alzette, Chiers, Moselle, Sûre and Syre), but also for the Mosel in France, which are publicly available via the web portal (Table 1). In case of a large flood, a crisis team is activated to coordinate emergency management activities. Civil protection measures and the organization of the response during ~~normal~~-floods are carried out by the national fire brigade: the ~~CGDIS~~ (Corps Grand-Ducal D'Incendie et de Secours (CGDIS)). In November 2024, LU-alert was launched, which is a new national warning system based on cell-broadcasting alerts and geo-located SMS services on mobile phones ~~and supports the geo-located SMS services~~ (Le gouvernement du Grand-Duché de Luxembourg, 2025). The SMS service has the additional advantage that it enables one to monitor the number of people in a certain area, which CGDIS can use to monitor how many people have been evacuated. This system was launched in response to the ~~2021~~-flood of 2021. With the launch of this new system, Luxembourg also added a new purple warning level for events with "immediate danger" ~~which that~~ require "immediate action", such as for flash floods. This new purple level can be issued by the meteorological agency (Meteolux) and the crisis management, but not by the hydrological agency AGE (see Sect. 4).

### ***Meteorological forecasts***

465 The national meteorological service Meteolux does not run its own weather model but instead uses different weather models  
| from ECMWF, DWD, and MeteoFrance, like the ICON-(D2-)EPS, ECMWF-ENS, and AROME-EPS. A severe weather risk  
assessment unit (Cellule d'évaluation du risque intempéries, CERI) can be initiated to estimate the societal impacts. The CERI  
includes representatives from the AGE, MeteoLux, the national government (Haut-Commissariat à la protection nationale),  
and CGDIS. As described above, Meteolux adopted new warning levels in November 2024: yellow-orange-red-purple (Fig.  
470 3).

### ***Hydrological forecasts***

Hydrological forecasts and warnings are provided by the flood forecasting department of AGE. They use a wide range of  
meteorological forecasts as input, from DWD, ECMWF, and MeteoFrance. The ensemble forecasts are made with the  
LARSIM model and can be updated every hour. They have a lead time of 24 hours for smaller streams, and 48 hours for the  
475 large rivers (e.g., Moselle, Lower Sauer). Ensemble discharge forecasts are optionally bias-corrected for current observation-  
simulation differences in discharge, soil moisture, precipitation, and temperature. They are generally updated every three hours  
during normal situations and every hour during emergencies. AGE only publishes the deterministic forecasts on the portal,  
while multi-model ensemble forecasts are used internally. Warning levels in Luxembourg have recently been revised, and  
AGE now uses three different levels: yellow ("Cote de vigilance"), orange ("Cote de pré-alerte") and red ("Cote d'alerte").  
480 ~~Those are in line with the warning levels of Meteolux, and the crisis management (see below).~~ However, the hydrological  
service (AGE) is not mandated to issue a purple warning in the new system. This is reserved for the meteorological forecasting  
service or crisis management. Interviewees from Luxembourg stressed that the new warning levels are based on the expected  
impact, derived from flood maps and/or consultations with local authorities and firefighters (Fig. 3).

### ***Crisis management and communication***

485 Luxembourg has a specific emergency management plan for extreme weather and floods (Le gouvernement du Grand-Duché  
de Luxembourg, 2025). During regular floods, the flood response is managed by CGDIS, ~~who which is~~ in constant exchange  
with the flood forecasting service. When the flood risk is deemed high, a Crisis Cell is activated, led by the High Commission  
for National Protection (HCPN) ~~and which comprises~~ comprising representatives from various bodies, such as the Police,  
Army, and Government. During a crisis, the AGE flood forecasting department is in constant exchange with CGDIS and the  
490 High Commission for National Protection (HCPN). The HCPN manages the actions required during a crisis (European  
Commission, 2024). Four crisis management phases are specified: normal, vigilance, pre-alert, and alert (Le gouvernement du  
Grand-Duché de Luxembourg, 2025). Those are in line with the hydro-meteorological warnings described above. In the 'pre-  
alert' phase, local flooding is expected and in the 'alert' phase widespread flooding is expected with a significant impact on  
people and property (Le gouvernement du Grand-Duché de Luxembourg, 2025). These alert phases are also coupled ~~with~~

495 specific water levels. In the '(pre-)alert phase', flood bulletins are circulated ~~which that~~ include warnings and a textual  
interpretation. ~~At~~In the 'alert phase', the prime minister (or his/her delegate) can activate a Crisis Unit Cell (Le gouvernement  
du Grand-Duché de Luxembourg, 2025). ~~The CGDIS pointed out that the procedures have improved greatly since the floods~~  
~~in July 2021. The CGDIS pointed out that the procedures have greatly improved since the July 2021 floods.~~ For example,  
before the 2021 flood event, communication between municipalities and CGDIS often failed. ~~Nowadays~~Today, flood alerts  
500 are sent to ~~the~~-local fire stations along with a request to contact the (deputy) mayor of the municipality.

### 3.5 The Netherlands

#### *General FFEWS setup*

The Royal Netherlands Meteorological Institute (KNMI) is responsible for ~~the~~-severe weather warnings in The Netherlands.  
KNMI and national hydrological forecast information is transmitted to ~~the~~-regional authorities, ~~such as the~~ ~~the~~-regional water  
505 authorities (Dutch: "Waterschappen"), who are responsible for forecasts in ~~the~~-smaller streams and catchments. ~~They advise~~  
~~crisis management organizations (the so-called 'Safety Regions') about risk and implementation of protection measures.~~ ~~They~~  
~~advise the crisis management organizations (the so-called 'Safety Regions') about the risk and the implementation of protection~~  
~~measures.~~

#### *Meteorological forecasts*

510 Among the meteorological models used are the HARMONIE-AROME (Frogner et al., 2019) and ECMWF (Haiden et al.,  
2018) weather models. ~~The HARMONIE model provides hourly forecasts, and the ECMWF model provides 6-hourly~~  
~~forecasts. The HARMONIE model provides hourly, and the ECMWF model 6-hourly forecasts. The version of HARMONIE~~  
~~ensemble model is called HarmonEPS, which produces a subset of ensembles at each run every hour ('a running ensemble').~~  
~~The ensemble model version of HARMONIE is called HarmonEPS, which produces a subset of ensembles at every hourly~~  
515 ~~run ('a running ensemble').~~ This is different from ECWMF-ENS, which runs full ensembles every 6 hours. Meteorological  
warnings are classified into the yellow-orange-red ~~color~~colour code (Fig. 3). The KNMI developed an in-house system to  
calculate maps of exceedance probabilities for an ensemble weather forecast: the Probabilistic Alert System for the  
Concurrence of Adverse Weather Elements (PASCAL) system (Kok, 2022).

520 ~~The KNMI works towards impact-orientated forecasting. The KNMI works towards impact-oriented forecasting.~~ To ~~enhance~~  
~~the~~improve knowledge and experience on those impact-~~orientated~~ oriented forecasts, the KNMI launched an Early Warning  
Centre in 2015. In case an orange or red warning is predicted by the ensemble models (> 60% probability), the KNMI initiates  
an online multi-disciplinary weather-impact team meeting. This team includes key organizations such as ProRail, the National  
Crisis Centre (NCC), the Netherlands Traffic Center (VCNL) and the National Operational Coordination Centre (LOCC). This  
525 team provides advice to the KNMI on the selected warning level, which is in practice always followed. The stakeholders in

the team also fill in an impact questionnaire, which results in an impact score per economic sector. The final warning level is decided by the operational lead at the KNMI. To allow for an early engagement of the impact team, the KNMI can already inform the impact team up to seven days in advance when a warning is not yet generated, but possible (using a 30% probability threshold). In case a rainstorm is predicted only at the last moment, the KNMI issues a warning without first consulting the impact team.

### *Hydrological forecasts*

The national Dutch Water Management Center (WMCN) produces 5-day (validated) and 15-day (automatic) forecasts for the river Rhine (gauge Lobith) and the river Meuse (gauge Sint-Pieter), using the HBV and SOBEK models. Those models are run on the Delft-FEWS platform for the whole (international) Rhine and Meuse basins. Forecasts 5-days ahead are deterministic, while the 15-day forecasts are ensemble products. These forecasts use multiple weather models as input: DWD COSMO-LEPS model, DWD ICON-EU, DWD ICON, the ECMWF-ENS, and ECMWF-HRES (deterministic) forecasts. Meteorological uncertainty is included ~~by~~ using ensemble forecasts, where every ~~ensemble member~~ member of the ensemble is used as input to the hydrological model. ~~For a small portion of the stations~~ ~~For a small portion of stations~~ (approximately 5%), a post-processing technique is used on the discharge forecasts (Verkade et al., 2017) to include the hydrological uncertainty. This is done by correcting (“dressing”) an individual streamflow ensemble member based on historical hydrological model errors. In addition, the WMCN uses forecasts from the European Flood Awareness System (EFAS; Smith et al., 2016), which is part of the Copernicus Emergency Management Service. Regional water authorities (Dutch: “waterschappen”) run forecasts for the smaller Dutch rivers and tributaries. However, not all water authorities run forecasting models operationally and many different models are used.

The Dutch national protocol for high waters and flooding (LDHO) provides an overview of the different warning levels for the Dutch main rivers (WMCN, 2023). Four warning levels are in place (Fig. 3): green, yellow (elevated water levels, water nuisance possible), orange (high waters) and red (extreme high water). These are based on both the water level and the discharge thresholds, and expressed as return periods (WMCN, 2023, Appendix D). Different return periods apply to different rivers, ~~as since~~ the strength of the dikes is different (WMCN, 2023). ~~Therefore, Fig. 3 displays the ranges of return periods.~~ ~~Therefore, Fig. 3 displays ranges of return periods.~~ The exact return periods per river can be found in WMCN (2023, Appendix D). As multiple ensemble forecasts are made (using multiple models), expert judgement is needed to determine which warning level is ~~eventually~~ ultimately selected.

In case of orange or red warnings, an expert group gathers into a national committee: the Landelijke Commissie Overstromingsdreiging (LCO). These experts in meteorology and hydrology translate the water level forecasts into impacts and advise on required interventions. The LCO is mandated to evaluate the ~~color~~ colour code selected by the WMCN and can increase the warning level if deemed necessary (e.g., in case of calamities, such as failures of water infrastructures). As a result

of these decisions, information in a web-portal (Table 1) is updated and disseminated every day, ~~which includes including~~ ~~the~~ hydrological forecasts, warning levels, and a textual interpretation. In case of orange and red warning levels, the online information can be updated multiple times per day. The orange warning level can be issued with a maximum lead time of 48 hours. However, ~~the~~ stakeholders can be informed up to 7 days in advance with a notice that the threshold will potentially be exceeded.

~~At-On~~ the local scale, ~~the~~ regional water authorities (“Waterschappen”) and Safety Regions are responsible for hydrological forecasting and responses, respectively. For example, the water authority in Limburg (Waterschap Limburg) runs operational forecasts to predict discharge and/or water levels, with a lead time of 5 days. Predictions are deterministic, although ~~the Waterschap Limburg stressed in the interview that there is a-a~~ strong need and ~~wish-desire exist~~ to ~~deploy-develop operational~~ probabilistic forecasting approaches. In Limburg, a variety of activities were triggered in response to the 2021 flood. A new large programme on flood safety called “Waterveiligheid en Ruimte Limburg (WRL)” ~~has-been~~ ~~was~~ launched in 2022. In this programme, local authorities and the central government are working together to improve Limburg's flood resilience. As part of WRL, a separate early warning project has been started to improve the warning system of the regional water authority (Waterschap Limburg). This leads to input for developing a new impact-based forecasting system of the Waterschap Limburg and the Safety ~~R~~egions.

#### 575 *Crisis management and communication*

The Safety Regions are regional organizations responsible for ~~the erisis~~-management ~~of crises offer~~ various ~~kinds-types~~ of hazards and the provision of information to the population. The Netherlands is the only country with such administrative units specifically designed for crisis management. Based on the estimation of LCO and the Departementaal Crisis Centrum (DCC) of the Ministry of Infrastructure and Water Management, the Safety Region and other stakeholders can be warned through the National Crisis Management System (LCMS). Actions (e.g., dike inspections) are linked to ~~the~~-warning levels, which are detailed in WMCN (2023) for ~~the~~-large rivers and ~~in the protocols from the~~-regional water authorities ~~protocols~~ for ~~the~~-small rivers. Some regions in The Netherlands have ~~a-a~~ dedicated emergency management plans for floods. Limburg province updated the ~~flood disaster management plan (Rampbestrijdingsplan Hoogwater Limburg; Veiligheidsregio Limburg-Noord and Zuid-Limburg, 2023)~~ after the 2021 floods, by including clear warning thresholds for the smaller rivers. Those are identified based on the expected nuisance to the (urban) surroundings and thus take the strength of the dikes and the exposed assets into account (Veiligheidsregio Limburg-Noord & Zuid-Limburg, 2023). In case of a flood crisis, the ~~D~~epartmental (DCC) or ~~N~~ational ~~C~~risis ~~M~~anagement ~~C~~enter (NCC) coordinates the crisis, and the LCO advises on measures to take (WMCN, 2023).

590 The public is warned through multiple channels, of which the most important are NL-ALERT (Table 1) and the 4278 air raid alarm systems. The KNMI also actively disseminates warnings through the newly developed KNMI app ~~s~~ and through social

media. ~~They have a clear approach to warning that is impact- and action-orientated (see Sect. 4.3). They have a clear impact- and action-oriented warning approach (see Sect. 4.3).~~

### 3.6 Belgium

#### 595 *General FFEWS setup*

~~The M~~eteorological forecast and warnings for Belgium are produced by the Royal Meteorological Institute of Belgium (RMI). River discharge forecasts are ~~separately~~-generated ~~separately~~ for Flanders and Wallonia, using RMI forecasts as input. The National Crisis Center (NCC) is responsible for managing national level crisis situations.

#### *Meteorological forecasts*

600 ~~At~~-In the RMI, the deterministic meteorological forecasts are produced by the ALARO model (Gerard et al., 2009). Currently, ensemble forecasts from the ECMWF-ENS system are used and the RMI develops a new ensemble system (AROME-EPS; Frogner et al., 2019) (Table 1). However, operational meteorologists also use other models from other agencies such as the UK Met Office and DWD.

605 Warnings are generated using ~~the yellow-orange-red color~~ codes ~~on~~-at the provincial level (Fig. 3). ~~Warnings are issued if more than 65% of the ensembles exceed the rainfall threshold for a specific colour over more than 25% of the provincial area (Fig. 3). A warning for a given colour can also be issued if either of the following conditions is met: 1) the threshold for the next higher colour is exceeded by more than 65% of the ensembles, even if this occurs over less than 25% of the area; 2) the threshold for the next higher colour is exceeded by 15–65% of the ensembles over more than 25% of the area. Warnings are issued if more than 65% of the ensembles are exceeding the rainfall thresholds for that specific color (Fig. 3), for > 25% of the provincial area. Warnings for a specific color can also be issued if there is a small chance (< 65% or < 25% of the area) on precipitation amounts belonging to the following color (for example, a yellow warning can be issued if there is a small chance on rainfall belonging to an orange warning).~~ These are guidelines, but the RMI can deviate from those based on the expected impact. The impacts are discussed in online impact team meetings, which can be called 2-3 times a day during severe  
615 weather conditions. These meetings involve the hydrological centers for Wallonia and Flanders (SPW and HIC/VMM, respectively), fire brigades, regional and national crisis centers and provincial governors. Although the RMI takes impacts into account in selecting the warning levels (e.g., a rainstorm approaching a festival), the interviewee stressed that the RMI is not mandated to provide impact- or action-oriented advisories. Four radars provide near real-time observations for nowcasting using the INCA-BE system. This system allows for the generation of so-called “flash warnings”, which are automatically  
620 generated warnings at the local level (i.e., municipal) with <1h lead time (Smet, 2017). These warnings are disseminated via the RMI app.

Formatted: English (United States)

### Hydrological forecasts

In Flanders, the Hydrologische Informatie Centrum (HIC) is responsible for monitoring and forecasting of navigable waters. The non-navigable rivers (category 1) in Flanders are monitored and forecast are issued by the Vlaamse MilieuMaatschappij (VMM). Those observations and forecasts are publicly available on the official Waterinfo web-portal (Table 1). The HIC produces ~~long~~medium-range-term (10 days) and short-term (48 hours) forecasts using the hydrological-hydraulic model NAM-MIKE11, run on the platform FEWS-Flanders. The long-term forecasts are based on deterministic ICON/ECWMF forecasts (2 times daily), while the short-term forecasts are derived from deterministic ALARO model runs (4 times daily). The HIC has six hydrodynamic models for different catchments: 1) IJzer, 2) Leie and its drainage channels, 3) Dender, 4) Zenne-  
630 Zeekanaal, 5) the Scheldt estuary and 6) Demer. These models provide ~~inundation~~-flood forecasts, which are also visualized in the web-portal and disseminated to partners. Apart from the ~~river~~-Demer ~~river~~, HIC runs the FFEWS for Flanders 4 times per day with input from the ALARO model of RMI. The rainfall forecasts used as input for the hydrological model are deterministic. Probabilistic discharge forecasts are made using a hydrological post-processing technique, based on historical lead-time dependent model errors (Van Steenberghe et al., 2012). Two types of warnings can be generated 48 hours in advance  
635 for both the navigable and non-navigable rivers: alarm states and flood states. These warnings -both follow the green-yellow-orange-red ~~color~~colour codes, for both -navigable (provided by HIC) and non-navigable (provided by VMM) rivers. The flood warning states (Fig. 3) are visualized ~~on~~-in the so-called 'Vrijboordkaarten' and show the distance between the water level and the dike crest. The ~~color~~colours reflect the following situations: green: no floods; yellow: pre-warning; orange: warning, non-critical flooding would be possible; red: alarm, critical flooding would be possible. These flood warning states are based on  
640 water-level thresholds determined by the hydrological-hydraulic model forecasting results. The alarm states, on the other hand, are determined based on the impact on water structures (e.g., bridges), but do not reflect overall flood impacts to society. The forecasts and warning levels are displayed in near real-time on the Waterinfo portal (Table 1). In case of high discharges, the warnings are actively ~~disseminated~~-distributed to the ~~water authorities~~-De Vlaamse Waterweg waterways authority, which distributes the warnings to other stakeholders and emergency management. After the 2021 flood, the hydrological warning  
645 thresholds have been changed ~~for the navigable rivers~~ in consultation with De Vlaamse Waterweg, mainly to better streamline thresholds among the different regions ~~of~~in Flanders.

In Wallonia, the Direction de la Gestion Hydrologique (DGH) of the Walloon government public service (Service Public de Wallonie, SPW) is responsible for monitoring, forecasting, and warnings. The DGH operates the Hydromax model using a  
650 network of 150 river ~~gauging~~-gauge stations and 100 rain gauges (the 'Wacondah' network) as input (DGH, 2024). Hydrological forecasts are made using deterministic weather forecasts from different ALARO models. In response to the 2021 flood, ~~also~~-ensemble weather forecasts from ECMWF-ENS are ~~also~~ used. Five different percentiles are selected from the 51-member ECWMF ensemble and used in Hydromax to produce hydrological forecasts, which are further post-processed considering forecast errors -of previous floods (DGH, 2025). ~~The smaller tributaries~~The main non-navigable rivers (e.g., the

655 Vesdre) are monitored by ~~T~~the Direction des Cours d'Eau Non Navigables (DCENN), which operates a measurement network called "Aqualim" (DGH, 2024). Warnings follow a green-yellow-red ~~color~~colour code (Fig. 3) and are issued at the sub- catchment level. Yellow warnings are issued if one (or more) rivers in a basin-sub-catchment are expected to cause localized and non-severe floods, and red warnings if one or more rivers in a basin-sub-catchment are expected to cause major floods with an impact on infrastructure and local residents. The thresholds are not prescribed but are defined by expert judgement ~~depending-based~~ on catchment wetness state, precipitation forecasts, discharge in the upstream rivers and tributaries, and ~~from~~ past experience. Observations and flood warnings are displayed on the web-portal of the Walloon government (Table 1). The portal was updated after the 2021 flood, and now also includes the non-navigable waterways (from the Aqualim network). None of the governmental hydrological and crisis management agencies in Wallonia produce ~~and/or use~~ inundation forecasts operationally. ~~Nonetheless~~However, inundation forecasts are produced for some pilot river sections; but currently these   
660 forecasts are not communicated to the public, ~~since-as~~ they do not consistently cover the whole river network.

#### *Crisis management and communication*

Crisis management in Belgium is organized at three levels: municipal, provincial, and federal state level. At the federal level crisis management is coordinated by the Centre de Coordination des Risques et de la Transmission d'Expertise (CORTEX) for Wallonia and the Crisiscentrum van de Vlaamse overheid (CCVO) for Flanders.

670 In Wallonia, in case of an orange warning, CORTEX calls ~~in~~ a special flood expertise unit (CELEX) by videoconference. The CELEX unit was launched in response to the 2021 flood event. In case of (pre-) alerts, CELEX develops and shares flood reports with stakeholders (but not with the general public). These reports include an outlook onf the consequences and the required actions. In Flanders, each province has an Emergency and Intervention Plan (NIP), which ~~was~~is revised after the 2021   
675 flood and is regularly exercised. In case of a large flood disaster, such as in 2021, the federal level is initiated and the National Crisis Center (NCCN) takes over the coordination. The RMI can directly send meteorological warnings to the NCCN. Furthermore, every municipality in Wallonia ~~needs-to~~should have a general emergency management plan (Plan Général d'Urgence et d'Intervention; PGUI). In May 2025, the Walloon government installed a Regional Crisis Management Plan, which can also be applied during flood crises ~~as well~~ (Government of Wallonia, 2025). Since June 2017, the public can be   
680 warned via the mobile cell broadcasting system: BE-ALERT (Table 1).

#### **3.7 Coordination and data sharing across borders**

All interviewees emphasized that data exchange between countries is already quite well organized. For example, the border between The Netherlands and Flanders is marked by the lower part of the Meuse river ("Grensmaas"). To provide a consistent warning for all households along the Grensmaas, Flanders adopts the warning levels used in the Netherlands. In addition, the   
685 Dutch water management center (WMCN) uses both ~~the~~ European data from the EFAS system and additional hydrological information from neighboring countries. In Germany, the exchange of meteorological and hydrological data and forecasts is

well organized. Data ~~is-are~~ exchanged between different federal states and different neighboring countries (Poland, the Czech Republic, Austria, Switzerland, France, Luxembourg, and The Netherlands) in near real-time (Demuth and Rademacher, 2016; LAWA, 2020). Several international agreements are in place to foster collaboration in transboundary river basins. The  
690 International Commission for the Protection of the Rhine (ICPR) fosters collaboration between the different states in the Rhine basin, and organises annual plenary assemblies, ~~in which developments in forecasting systems are discussed (e.g., on harmonization of forecast uncertainty determination; Demuth and Rademacher, 2016).~~ In 1987, the Internationale Kommissionen zum Schutz der Mosel und der Saar (IKSMS) signed an intergovernmental agreement between France, Germany, and Luxembourg to improve flood detection and forecasting in the Saar and Moselle catchments (IKSMS, 2025).  
695 This strong collaboration results in hourly data sharing from rain gauges, discharge gauging stations, precipitation radar data, and weather forecasts between all flood forecasting centers (IKSMS, 2025). Moreover, joint data assimilation systems (such as the PLATIN data exchange server), models and forecasting systems are developed, technical trainings are organized, and language courses are offered to improve communication (IKSMS, 2025). Other transboundary basins, especially for the smaller rivers, can learn from this strong and long-lasting collaboration agreement.

700 ~~While Although transboundary collaboration was~~ not identified as a major issue in the FFEWS chains, some concerns were raised in the interviews. While sometimes common models are developed (such as mentioned above), too often different meteorological and hydrological models are used in different countries (see Table 1) which can result in inconsistent and conflicting information to stakeholders in transboundary regions. Furthermore, some important data ~~is-are~~ not always shared  
705 (efficiently). For example, HIC in Flanders mentioned the need to improve the exchange of information about upstream reservoirs in Wallonia. Often, information ~~of-about~~ reservoir levels is not shared, especially for energy-producing reservoirs (for energy companies, such as ENGIE). ~~In addition, Waterschap Limburg interviewees stressed the importance of an international exchange of flood warnings, which is currently lacking. Further, interviewees from Waterschap Limburg stressed the importance of an international exchange of flood warnings, which is currently lacking.~~ These challenges highlight the need  
710 to continue efforts to streamline transboundary collaboration.

#### 4. Main improvements and remaining challenges after the 2021 flood

The interviews revealed that the FFEWSs in all regions are under strong and rapid development after the 2021 flood (Table 2). All regions now have flood emergency response plans ~~for floods~~ and use cell-broadcast services ~~for-to alert~~ alerting the population. ~~In NRW, Aa~~ new flood forecasting ~~team-center~~ has been ~~developed-established~~ at LANUK ~~in NRW~~ and 26 new  
715 flood monitoring gauges have been installed (LANUK, 2025 (Table 2)). Moreover, all regions defined clear thresholds for the different hydrological warning levels, except for Wallonia (Fig. 3). In autumn 2023, the flood forecasting center (LfU) in the RLP region added a new warning level for very extreme floods (> 100-year return period) with a dark-purple ~~color~~ colour (Fig. 3 ~~and Table 2, top~~). The flood forecasting center of Flanders (HIC) optimized ~~their-its~~ warning thresholds in close collaboration

with stakeholders, such as the Vlaamse Waterweg waterways authority, and many provinces updated their Emergency and Intervention Plans. In Wallonia, a flood expertise unit (CELEX) has been established ~~since~~ after the 2021 floods, which can provide tailored flood impact advice during crises. Furthermore, ensemble forecasts are now used to produce flood forecasts and the web-portal for floods has been renewed. In Limburg, the flood disaster management plan (“Rampbestrijdingsplan Hoogwater Limburg”) now includes flood scenarios for small rivers (e.g. the Geul river), including clear warning thresholds and the corresponding actions. Moreover, a large programme has been launched in 2022 to improve flood safety: Waterveiligheid en Ruimte Limburg (WRL). In November 2024, Luxembourg implemented a new alerting system (LU-alert) and streamlined the warning system, by adopting the same warning levels and colour codes for meteorological forecasts and crisis management alerts. In November 2024, Luxembourg implemented a new alerting system (LU-alert), harmonizing the warning levels of the meteorological forecasts and the crisis management. This includes a new level for “immediate danger” with a purple ~~color~~ colour. Even after those changes, the interviewees identified several issues that require attention to further improve FFEWSs (Table 2, bottom). We will further outline those challenges below.

**Table 2. Most important changes of the Flood Forecasting and Early Warning Systems (FFEWS) triggered in response to the 2021 flood, and the key challenges remaining in different European regions and countries.**

| Country                                                                                                                                                                                              | Germany (RLP)                                                                                                                                         | Germany (NRW)                                                                                           | Luxembourg                                                                     | The Netherlands                                                                                        | Belgium (Flanders)                                                                                  | Belgium (Wallonia)                                                                                      |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|
| <b>Major changes since the 2021 floods</b>                                                                                                                                                           | DE-alert system, and addition of extra warning level for very extreme floods (dark purple)                                                            | DE-alert system, <u>new flood monitoring gauges</u> and establishment of a new flood forecasting center | LU-alert system, including a new warning level for “immediate danger” (purple) | A new multi-year programme on flood safety in Limburg (WRL) and updated flood disaster management plan | Optimized hydrological warning thresholds (see Fig. 3) and revised Emergency and Intervention Plans | Flood expertise units (CELEX), updated flood portal and move from deterministic to ensemble forecasting |
| <b>Key challenges remaining after 2021</b>                                                                                                                                                           | <b>Streamlined warning levels and thresholds (Sect. 4.1)</b>                                                                                          |                                                                                                         |                                                                                |                                                                                                        |                                                                                                     |                                                                                                         |
|                                                                                                                                                                                                      | Different warning levels, thresholds, and <del>color</del> <u>colour</u> codes are used between and within the different countries and organizations. |                                                                                                         |                                                                                |                                                                                                        |                                                                                                     |                                                                                                         |
|                                                                                                                                                                                                      | <b>Warnings for catastrophic events (Sect. 4.2)</b>                                                                                                   |                                                                                                         |                                                                                |                                                                                                        |                                                                                                     |                                                                                                         |
|                                                                                                                                                                                                      | The highest thresholds are often too conservative (too low) and thus do not reflect major societal flood impacts.                                     |                                                                                                         |                                                                                |                                                                                                        |                                                                                                     |                                                                                                         |
| <b>Impact-based forecasting (Sect. 4.3)</b>                                                                                                                                                          |                                                                                                                                                       |                                                                                                         |                                                                                |                                                                                                        |                                                                                                     |                                                                                                         |
| While impact-based forecasting is recognized as crucial for efficient decision-making, such systems are not yet operationally implemented in most of the regions.                                    |                                                                                                                                                       |                                                                                                         |                                                                                |                                                                                                        |                                                                                                     |                                                                                                         |
| <b>Communication before and during the crisis phase (Sect. 4.4)</b>                                                                                                                                  |                                                                                                                                                       |                                                                                                         |                                                                                |                                                                                                        |                                                                                                     |                                                                                                         |
| Effective communication between different organizations (e.g., between meteorological agencies, flood forecasting centers, crisis management authorities) and to the population remains a challenge. |                                                                                                                                                       |                                                                                                         |                                                                                |                                                                                                        |                                                                                                     |                                                                                                         |

Formatted: Font color: Text 1

#### 735 4.1 Streamlined warning levels and thresholds

The first challenge relates to the transboundary differences in meteorological and hydrological warning levels and their thresholds. In general, the number of warning levels, their color and the thresholds used differ between the different countries (Fig. 3). Moreover, they frequently also differ within the different regions of a country and between the meteorological and hydrological agencies of a country (Fig. 3). For example, the number of warning levels and their color codes are different in the two federal states of Belgium: Flanders and Wallonia (Fig. 3). In Luxembourg, MeteoLux and CGDIS apply a purple warning level for events posing immediate danger (Fig. 3). They are mandated to do so as because both maintain permanent operational offices (Fig. 3). While-Although this level is also used by LU-Alert and the crisis management, the flood forecasting center (AGE) is not mandated to use this level as they-it does not operate 24/7. AlsoIn addition, in Germany, the warning levels are not consistent among federal states. For example, in RLP they are linked to flood return periods and in NRW to flood impacts (NRW, 2024) (Fig. 3) – the latter does not necessarily corresponding to the same return periods as in the neighboring RLP (DKKV, 2024). While-Although RLP now distinguishes six warning levels, other German states use just four (like NRW), including a green level for no warning (Fig. 3). Moreover, the “red” color being the highest warning level in some flood forecasting centers in Germany (e.g., LANUK in NRW) is inconsistent with the highest “purple” weather warnings from DWD. The H interviewees in RLP also state that the number of alerts by DWD have-been-was high in the past during some periods of possible events and that it could be difficult for the recipients to estimate if a warning indeed results in a flood. Furthermore, DWD warnings are generally cleared at the end of a precipitation event, while flood warnings can still be active because the flood did not yet end. According to the interviewees, such-this situation happened occurred in 2021 and confused local stakeholders. Research shows that inconsistent visual and textual warnings from multiple sources are confusing for-to decision-makers and the public, which can lead to ineffective communication and actions (Weyrich et al., 2019). Moreover, it is important to note that the underlying hydrological models are often different between and within (e.g. Wallonia and Flanders) different countries. Those models are deployed on a wide range of different forecast lead times (ranging from 48 hours to 15 days). We recommend evaluating the consequences of the above inconsistencies for improving early warning communication. We recommend evaluating the consequences of the above inconsistencies for improvement of communication of early warnings.

#### 760 4.2 Warnings for catastrophic events

After the 2021 flood, some regions started a discussion to introduce a new (or additional) color code for extreme floods. In the German federal state of the RLP, this discussion has led to the adoption of an additional ‘dark purple’ level in addition to the “purple” level in Autumn 2023, based on a 100-year return period threshold (Landesamt für Umwelt Rheinland-Pfalz, 2025). In the German federal state of RLP, this discussion has led to adopting an additional ‘dark purple’ level on top of the “purple” level in Autumn 2023, based on a 100-year return period threshold (Landesamt für Umwelt Rheinland-Pfalz, 2025).

This adjustment was primarily ~~done-made~~ to improve ~~the~~ communication possibilities for “extremely high danger of flooding”. However, recent research suggests that even the purple DWD warnings for the 2021 event were too abstract and that people did not ~~link-associate~~ the warning with such ~~an~~ extreme flood event (Ommer et al., 2024). The warning levels for Luxembourg also contain a new purple level, which can only be issued by the meteorological service MeteoLux and not by the flood forecasting center AGE (Fig. 3). The highest warning level of AGE is “Cote de vigilance rouge” (Red), which is often linked to small return periods. For example, ~~the~~ gauge Bollendorf in the Sauer has a red warning threshold related to a water level of 4.25 m, which corresponds to less than a 5-year return period event. This suggests that extreme river~~ine~~ floods could be better represented in the new warnings. Our interviewees in Wallonia stated that “...there should be a new discussion about the need for an extra level of warning (e.g.: dark red) for really extreme situations”. For the regional water authority in Limburg (Waterschap Limburg), the highest warning level (red) is described as a “water nuisance” ~~to-for~~ people, rather than a high-impact catastrophic event ([Veiligheidsregio Limburg-Noord & Zuid-Limburg, 2023](#)). However, more extreme thresholds come with more uncertainty. Rare events will typically be ~~foreeas~~-predicted with only a few ensemble members. These low-probability and high-impact forecasts bring considerable decision-making challenges (Speight et al., 2021), as was also observed for the 2021 event. There is a need to better understand such extreme events, to better represent them in ~~the~~-models and forecasts, and ~~to~~ better predict potential impacts. This was raised by respondents in all countries.

Flood hazard maps can be used to estimate which areas are going to be flooded during an expected event (e.g.: Dottori et al., 2017). However, the use of static flood hazard maps showed large deficiencies during the 2021 flood. Vorogushyn et al. (2022) showed that the inundated areas in 2021 exceeded the most extreme official hazard maps in the Ahr Valley and recommend developing more severe extreme scenarios. In Wallonia, responders and households in ‘safe green or unmarked zones’ faced the 2021 flood and did not know what to do. Many of the flooded areas directly along the Vesdre river (a tributary of the Meuse river) were marked as safe green zones or simply left unmarked ~~onin~~ official flood hazard maps. As a result, many residents were not aware of the flood risk and were surprised; their houses were flooded up to the second floors (Rodríguez Castro et al., 2025).

#### 4.3 Impact-based forecasting

Our study confirms that the uptake of impact-based forecasting in operational systems is challenging. ~~The M~~most important reasons are the lack of relevant data (e.g., on flood vulnerability), institutional collaboration across sectors (e.g., between forecasting agencies and emergency management), and lack of computational power to run inundation models, especially for multiple ensemble members. Our interview results show that ~~T~~the move towards impact-based forecasting is crucial to optimize warnings and trigger effective early actions, as mentioned in the interviews. Our interviewees in RLP (Germany) explained that the hydrological thresholds based on return periods (see Fig. 3) do not always reflect the corresponding impacts, and that impact-based thresholds would be more valuable to decision-makers. For example, ~~i~~Also in Wallonia, the same warning level for different events can result in significant differences in impact (DGH, 2024). Moreover, ~~many~~ interviewees

~~stressed that f-interviewees indicated that impact-based forecasting is important in taking effective early actions. First~~  
800 responders require specific and detailed information to execute their operations, such as expected flooded areas, flood depths  
at critical infrastructures (e.g., hospitals or schools) and predictions of expected damages. Therefore, many interviewees  
~~stressed that~~ predictions of potentially flooded areas and affected buildings are a crucial component to move to effective  
impact-based forecasting.

#### The importance of inundation forecasts

The HIC and VMM in Flanders are the only flood forecasting centers included in this study which that produces inundation  
forecasts operationally (Table 1). However, these inundation forecasts are not used in any of the official warning levels (see  
Fig. 3). Flood inundation forecasts can be made using multiple methods, by translating streamflow to event-based flood hazard  
810 maps using a pre-computed catalogue of hydrodynamic simulations (Dottori et al., 2017), or by using operational  
hydrodynamic models (Apel et al., 2022). Recent computational advances in flood modeling, such as the use of machine-  
learning, Graphical Processing Units (GPUs) and cloud computing make operational inundation modelling realistic and  
affordable (Hofmann et al., 2021; Speight et al., 2021; Apel et al., 2022; Agerbeek et al., 2024). Apel et al. (2022) produced  
inundation forecasts (in a hindcast mode) for the 2021 flood event in the Ahr Valley at 10 m spatial resolution in a time span  
815 suitable for operational application (14 min on a GPU) and showed the potential value of these forecasts for operational crisis  
management. Estimates of inundated areas (and other flood variables, e.g., flow velocity) can be overlaid with exposed  
buildings and critical infrastructure, to support rapid mapping of flood impacts (e.g., Apel et al., 2022, Najafi et al., 2024,  
Dottori et al., 2017). Citizens show a clear preference for inundation maps in warning messages they receive, compared to  
other information types (e.g., a warning symbol; Lindenlaub et al., 2024a). Moreover, people tend to assess the flood magnitude  
820 higher when an inundation map is provided (Lindenlaub et al., 2024b). The high value of detailed flood inundation forecasts  
for emergency management is also emphasized in England (Aldridge et al., 2020). However, it is not yet clear whether warning  
messages accompanied by information on the flooded area lead to better protective behaviour of residents (Kuller et al., 2021).

#### Best practices of impact-based forecasting

825 Despite the challenge of operational impact-based forecasting, some best practices in the region should be outlined. Some  
regions, such as NRW in Germany, developed impact-based warning thresholds (Fig. 3). Here, the water level thresholds are  
chosen based on the expected impact to assets and people. To translate hydro-meteorological forecasts into impacts and  
advisories, regions established flood expert teams which interpret and translate the forecasts into impacts and advisories for  
the local authorities. CELEX in Wallonia and ~~the~~ LCO in The Netherlands are examples of such groups, which provide an  
830 interpretation and translation of the meteorological and hydrological forecasts. After the 2021 flood, the regional water  
authority in Limburg (Waterschap Limburg) launched a dedicated multi-year programme to develop an impact-based

Formatted: Font: Not Italic

Formatted: y2iqfc, Font color: Custom Color(32,33,36)

Formatted: y2iqfc, Font: Not Italic, Font color: Custom Color(32,33,36), English (United Kingdom)

Formatted: y2iqfc, Font: Not Italic, Font color: Custom Color(32,33,36)

Formatted: Normal

Formatted: y2iqfc, Font color: Custom Color(32,33,36), English (United Kingdom)

forecasting system. The KNMI is rapidly developing impact-based forecasting techniques through the development of the Early Warning Centre, involving a range of multi-sectoral stakeholders (KNMI, 2025). During the interviews, ~~the~~ KNMI also emphasized that their efforts on impact-based forecasting contribute to stronger partnerships and collaborations, with organizations which are all working towards the same goal of predicting (flood) impacts. ~~(Red Cross Red Crescent Climate Centre, 2020).~~

#### 4.4 Communication before and during the crisis phase

One of the interviewees mentioned: “Forecasts are no longer the problem, the problem is to act”. ~~One of the interviewees mentioned: “Forecasts are not the problem anymore, the problem is to act”.~~ Dasgupta et al. (2024) suggest that a key factor for inaction is ineffective communication of the forecasts, including their uncertainty. Here, we elaborate on this challenge, specifically for communication to crisis managers and the local population.

##### *Communication between forecasters, crisis managers, and first responders*

Our results suggest that there are major challenges exist in communication between hydrological forecasting agencies, crisis management authorities, and ~~the~~ first responders right before and during the crisis phase. The ~~the~~ interviewees stressed that the information needs for emergency management are regularly-often not met. The ~~the~~ interviewees in Wallonia noted that not all municipalities have specific flood emergency management plans, even if they are in areas of high flood risk. In The Netherlands, regional authorities (water authorities Waterschappen and Safety Regions) ~~(organization of first responders)~~ stressed the challenge of interpreting the issued-forecasts issued by national services (KNMI and WMCN). An evaluation of the 2021 floods in Germany already pointed out that there was a lack of actionable information in the forecasts, which is highly needed for first responders and the population (DKKV, 2024; Thieken et al., 2023a). In Wallonia, this resulted in crisis managers not receiving the highly needed information about the expected extents of the flood ~~inundation extents~~. According to our interviewees in the region, ~~this~~ has eroded trust between different stakeholders, which is detrimental ~~for to~~ the functioning of an entire warning system (Seebauer and Babsicky, 2017). ~~Moreover, DKKV (2024) points out that there was a lack of clear and consistent communication between different levels of governments. Our interviewees in Germany pointed out that, although improvements are seen since 2021, the communication between hydrological agencies to emergency management units is still insufficient, resulting from a non-effective communication chain and missing communication protocols. Therefore, the information needs of first responders are still not met.~~

Multiple interviewees point to deficiencies in the communication chain of forecast agencies to crisis managers. Our interviewees in Germany pointed out that, although improvements are seen since 2021, the communication between hydrological agencies to emergency management units is still insufficient, resulting from a non-effective communication chain and missing communication protocols. Therefore, the information needs of first responders are not yet met. In Flanders, the warnings from HIC are sent to the De Vlaamse Waterweg waterways authority, who distribute ~~it~~ them further to other

Formatted: Font: Italic

865 stakeholders. However, ~~our interviewees in Flanders pointed out that~~ flood risk management is not a core business of De Vlaamse Waterweg authority; and therefore, they lack specific expertise on flood risk. Consequently, warnings are sometimes misinterpreted and not well communicated further. Similar findings are reported from experts in Luxembourg, where communication between ~~the~~ crisis management (CGDIS) and meteorological and hydrological agencies (MeteoLux and AGE) is sometimes challenging. ~~The~~ ~~i~~ interviewees from Luxembourg suggested that a central Early Warning Center for Luxembourg will be highly valuable ~~in condensing to condense~~ the different forecasts into actionable warnings. In The Netherlands, 870 evaluations after the 2021 flood show the need for centralized and streamlined information (COT, 2022). Our interviewees in The Netherlands stress that local stakeholders (e.g., municipalities) still suffer from fragmented pieces of information from multiple stakeholders such as the KNMI, ~~water boards~~ local water authorities and the WMCN.

Formatted: Font:

875 Research shows that probabilistic information can lead to better decisions (Verkade & Werner, 2011). Most ~~of the~~ operational forecasting centers in the region use probabilistic forecasting (Table 1). However, probabilistic approaches are often lacking for smaller rivers in The Netherlands. Moreover, ~~the~~ communication of probabilistic forecasts can be challenging (Arnal et al., 2019). ~~Our interviewees in~~ Germany, Luxembourg and the Netherlands ~~it was~~ mentioned that local emergency responders often prefer one value (e.g., water level) to take action upon. However, this value can be misleading ~~due to because of~~ the inherent uncertainty. Therefore, it is crucial to invest in better communication and interpretation of probabilistic forecasts and 880 translate them into action advice. The WMCN in the Netherlands is advancing ~~in~~ the communication of probabilistic forecasts by providing tables with exceedance probabilities for each discharge threshold (WMCN, 2024). Other forecasting centers are recommended to follow this approach. It is recommended for other forecasting centers to follow this approach.

### *Communication to the local population*

885 Germany and Luxembourg installed a cell broadcasting system in response to the 2021 flood. Now, all countries covered in this study have such systems in operation (Table 1). Despite the effectiveness of cell broadcasting, research shows the need ~~for~~ a multi-channel approach due to the risk of failure of one system, ~~for example due to (e.g.,~~ power shortages or hacks) (Mahdavian et al., 2020). This was also emphasized in our interviews, such as with the national fire brigade of Luxembourg (CGDIS). Moreover, the 2021 flood showed that people ~~mentioned~~ reported a wide variety of information sources, including social media, ~~in addition to official sources such as~~ NL-ALERT, radio, and television). Air raid sirens are another means of 890 mass-alerting the population. The countries in our study have a different perspective with regards to the air raid sirens. CGDIS emphasized the importance of the air raid sirens to ensure a multi-channel approach to mass-alerting civilians. However, in Wallonia, the sirens are not used, and The Netherlands plans to stop the system, mainly because of the high maintenance costs which ~~due~~ outweigh the societal benefits (Ministerie van Justitie en Veiligheid, 2024). In Germany, the sirens are being renewed since the 2021 flood. The different approaches of neighbouring countries to mass-alerting the population in case of a 895 flood or other catastrophe are noteworthy ~~remarkable~~.

~~Color~~Colour-coded rainfall warnings for the 2021 floods were sometimes perceived as too abstract or vague to increase people's risk perception and willingness to act (Ommer et al., 2024). People take more (effective) protective actions if warnings are impact-based (Meléndez-Landaverde et al., 2019), and even more if those warnings also include behaviour recommendations (Red Cross Red Crescent Climate Centre, 2020; Golding et al., 2019). Despite the importance of investing in warning communication, multiple organisations during our interview stressed that they lack ~~the~~ capacity or knowledge to develop effective communication protocols to inform the public. The 2021 flood in Limburg showed that, while people aware of an evacuation advice evacuated significantly more often, not everyone follows the advices (Endendijk et al., 2023a). This highlights the importance of tailored and concrete evacuation advice. In Germany, 50 to 75% of flood fatalities happened outside of the officially delineated flood hazard zones in NRW and RLP, respectively (Rhein and Kreibich, 2025; Thieken et al., 2023b). This illustrates the importance of flood risk awareness and risk perception as important drivers of adaptation. Since households not living in flood zones have lower risk perceptions, their response rates are also lower than those living in flood zones and they require tailored communication strategies (Aerts et al., 2018). Multiple interviewees (e.g., KNMI and AGE) stressed the importance of tailoring the communication strategy to different target groups. For example, young people in The Netherlands receive information through social media, such as Instagram, and generally will not download the KNMI phone application. The German national flood portal (<https://hochwasserzentralen.de>, Table 1) recently included action recommendations as addition to the hydrological warnings. The Dutch meteorological office KNMI recently hired a communication expert (specialized in social media) to produce clear infographics on '*what can you expect*' and '*what can you do*' during emergency situations (KNMI, 2024). This is presented in plain language (B1 level). We identify this as a best practice that can be followed by other regions.

## 5. Conclusions and recommendations

### 5.1 Conclusions

920 This study reviews the status of the Flood Forecasting and Early Warning Systems (FFEWs) in the countries in ~~Northwestern~~  
~~western~~ Europe hit by the July 2021 flood: Germany, Luxembourg, Belgium, and The Netherlands. Expert interviews in the  
region reveal that all systems are under ~~strong and~~ rapid development after the 2021 flood event. This includes an optimization  
optimization of warning thresholds (including new thresholds for very extreme hydrological events), improvement of online  
flood forecasting portals, new ensemble-based forecasting techniques, installation of new flood gauges, a new flood forecasting  
teamscenter, new national-scale cell broadcasting systems to send out phone-based alerts in case of immediate danger (such  
925 as LU-Alert) and updated crisis management plans for floods ~~disaster management plans~~ (including for small rivers). Now all  
regions, except Wallonia, have clear pre-defined warning thresholds based on peak discharge or water levels. In most regions,  
those hydrological warning thresholds are determined based on the expected local flood impacts.

There are ~~S~~strong differences in exist in flood warning levels and ~~color~~colour codes aeross-between and within the countries.  
930 Additional research is necessary to determine whether an international harmonization of the warning levels is desirable. As a  
response to the ~~extreme-2021~~ extreme flood event ~~of 2021~~, Luxembourg and some regions of Germany have recently  
introduced an additional purple warning code for the most extreme weather and hydrological events. In countries where red is  
the maximum warning level, some experts suggested also adding a purple level to represent truly catastrophic impacts such as  
the 2021 floods. ~~Despite-Although~~ having an extreme warning level can be helpful ~~for-in~~ indicating the potential disastrous  
935 consequences of an event, it is still under-debated whether more warning levels support ~~a~~ more effective communication to the  
public and responders.

The implementation of operational impact-based forecasting and warning is ~~-a challenge~~challenging, both on a technical and  
institutional level. Although impacts are sometimes considered in the design of the hydrological warning thresholds, the  
940 thresholds are often too conservative (too low) and thus do not reflect major societal flood impacts. Moreover, meteorological,  
and hydrological forecasts often lack specific impact and action information, which is highly needed for first responders at the  
local level. The forecasts of river discharge, return periods, and corresponding warnings are often hard to interpret by local  
decision-makers, emergency services and the local population. To trigger effective early actions, it is strongly suggested ~~to~~  
enrich~~that~~ forecasts are enriched with impact-based information and ~~to~~ provide reliable predictions of inundated areas and risk  
945 hotspots. Flanders is the only region where operational flood inundation (i.e., flood maps) forecasts are provided. Recent  
computational developments in inundation modelling such as cloud computing, GPU-based parallelization and machine-  
learning will significantly reduce run times and pave the way for the operational implementation of inundation forecasts.

## 5.2 Recommendations

Resulting from the four key challenges outlined in this study, our study has four concrete recommendations:

950 1) Streamlined warning levels: investigate whether harmonization of meteorological and hydrological warning levels and colour codes between different countries and within different regions of a country, improves communication and decision-making.

2) Warnings for catastrophic events: Assess the added value of more extreme warning levels, such as the (dark) purple levels in Luxembourg and some German regions.

955 3) Impact-based forecasting: accelerate the development of operational impact-based forecasting systems. This in turn requires an expansion of resources and knowledge on impact-based forecasting and a stronger collaboration across sectors, such as between forecasting agencies and emergency management.

4) Forecast communication: implement a structural evaluation of warning communication chains to ensure that warnings are consistently communicated, correctly interpreted by different organizations and the local population, and that the information is tailored to their needs.

960 ~~1) Streamlined warning levels: investigate whether harmonization of meteorological and hydrological warning levels and color codes between different countries and within different regions of a country, improves communication and decision-making.~~

~~2) Warnings for catastrophic events: Assess the added value of more extreme warning levels, such as the (dark) purple levels in Luxembourg and some German regions.~~

965 ~~3) Impact-based forecasting: accelerate the development of operational impact-based forecasting systems. This in turn requires an expansion of resources and knowledge on impact-based forecasting, and a stronger collaboration across sectors, such as between forecasting agencies and emergency management.~~

~~4) Forecast communication: Implement a structural evaluation of warning communication chains, to ensure that warnings are consistently communicated and correctly interpreted by different organizations, and the local population, and ensure that the information is tailored to their needs.~~

970

These challenges need to be addressed to reduce the gap between early warning and early action during impactful flood events.

Formatted: Font: Italic

975 **Appendix A. Interviewed specialists**

Formatted: Heading 1

**Netherlands**

- Senior operational flood forecasting expert at Watermanagementcentrum Nederland (WMCN)
- Senior operational early warning specialist and program manager at Royal Netherlands Meteorological Institute (KNMI)
- 980 • Senior flood early warning specialist at Waterschap Limburg

**Belgium**

- [Head Professor at University of Gent and lead](#) of the meteorological forecasting department of Royal Meteorological Institute of Belgium (RMI)
- Senior operational flood forecasting expert at Flanders Hydraulics Research - Hydrological Information Centre (HIC)
- 985 • Two senior crisis managers [and one operational flood forecasting expert](#) at [Le](#)-Service public de Wallonie (SPW)

**Germany:**

- Two senior operational flood forecasters at Landesamt für Umwelt Rheinland-Pfalz (LfU)
- Senior operational flood forecaster at Natur, Umwelt & Klima in Nordrhein-Westfalen (LANUK)
- Professor at University of Kaiserslautern-Landau (RPTU)

990 **Luxemburg**

- Senior flood forecasting and data management expert at Administration de la gestion de l'eau (AGE)
- General Director at Corps grand-ducal d'incendie et de secours (CDGIS)

995

#### Author contribution

**TB:** Conceptualization, Investigation, Methodology, Writing, Visualization, **JA:** Conceptualization, Investigation, Methodology, Writing, Funding acquisition, Project administration, **DRC, BD, SV, RL, JK, DZ, HM, AT:** Conceptualization, Investigation, Methodology, Writing, **KS:** Conceptualization, Investigation, Methodology, Funding acquisition, Project administration, **PW:** Investigation, Writing, **LF:** Writing, **JV:** Conceptualization, Investigation, Methodology.

#### Competing interests

The authors declare that they have no conflict of interest.

#### Acknowledgements

This research has been supported by the JCAR ATRACE project (<https://www.jcar-atrace.eu>). The authors from University of Liège have also been partly supported by the Interreg NWE project FlashFloodBreaker (<https://flashfloodbreaker.nweurope.eu/>). We gratefully thank KNMI, (The Netherlands), SPW (Belgium), LfU (Germany), RPTU (Germany), WMCN (The Netherlands), HIC (Flanders, Belgium), AGE (Luxembourg), RMI (Belgium), SPW (Wallonia, Belgium), Waterschap Limburg (The Netherlands), LANUK (Germany) and CGDIS (Luxembourg).

#### Financial support

This research was supported by the JCAR ATRACE project. The JCAR ATRACE project has received funding from the Dutch Ministry of Infrastructure and Water Management.

Formatted: Justified, Line spacing: 1,5 lines

## References

- Aerts, J.C.J.H., Botzen, W.J., Clarke, K., Cutter, S.L., Michel-Kerjan, E., Surminski, S., Mysiak, J., Merz, B., Hall, J., Kunreuther, H.: Including Human Behavior in Flood risk assessment. *Nature Climate Change*, 8, 193-199, 2018.
- 1015 AGE: Hochwasserereignis Juli 2021, Esch-sur-Alzette, Luxembourg, <https://eau.gouvernement.lu/fr/actualites/2021/07-juillet/Hochwasserereignis2021.html> (accessed on 20/02/2025), 2025.
- Agerbeek, B., Knepl e, M., Witsenburg, F., & Jonkman, S. N.: Near real-time flood risk modelling in response to increasing uncertainties in flood predictions: Insights from the Kakhovka Dam breach in Ukraine. *Journal of Coastal and Riverine Flood Risk*, 3, 2024.
- 1020 Aldridge, T., Gunawan, O., Moore, R. J., Cole, S. J., Boyce, G., & Cowling, R.: Developing an impact library for forecasting surface water flood risk. *Journal of Flood Risk Management*, 13(3), e12641, 2020.
- Alfieri, L., Zsoter, E., Harrigan, S., Hirpa, F.A., Lavaysse, C., Prudhomme, C., Salamon P.: Range-dependent thresholds for global flood early warning, *Journal of Hydrology*, 4, 100034, doi.org/10.1016/j.hydroa.2019.100034, 2019.
- Apel, H., Vorogushyn, S., & Merz, B.: Brief communication: Impact forecasting could substantially improve the emergency management of deadly floods: case study July 2021 floods in Germany. *Natural Hazards and Earth System Sciences*, 22(9). <https://doi.org/10.5194/nhess-22-3005-2022>, 2022.
- 1025 Arnal, L., Anspoks, L., Manson, S., Neumann, J., Norton, T., Stephens, E., Wolfenden, L. & Cloke, H. L.: Are we talking just a bit of water out of bank? Or is it Armageddon? Front line perspectives on transitioning to probabilistic fluvial flood forecasts in England. *Geoscience Communication*, 3, 203–232., 2020.
- 1030 Bauer, P., Thorpe, A. & Brunet, G.: The quiet revolution of numerical weather prediction. *Nature* 525, 47–55. <https://doi.org/10.1038/nature14956>, 2015.
- Becker, G., Huitema, D., and Aerts, J. C. J. H.: Prescriptions for adaptive comanagement: the case of flood management in the German Rhine basin. *Ecology and Society* 20 (3): 1. <http://dx.doi.org/10.5751/ES-07562-200301>, 2015.
- 1035 Bischiniotis, K., de Moel, H., van Den Homberg, M., Couasnon, A., Aerts, J., Nobre, G. G., Zsoter, E. & van den Hurk, B.: A framework for comparing permanent and forecast-based flood risk-reduction strategies. *Science of the total environment*, 720, 137572, 2020.
- Blahak, U.: SINFONY – DWD’s New Seamless Forecasting System. [https://www.dwd.de/DE/leistungen/pbfb\\_verlag\\_promet/promet\\_107\\_einzelkapitel\\_pdf/Promet\\_107\\_Kap1\\_pdf.pdf?\\_blob=publicationFile&v=3DWD](https://www.dwd.de/DE/leistungen/pbfb_verlag_promet/promet_107_einzelkapitel_pdf/Promet_107_Kap1_pdf.pdf?_blob=publicationFile&v=3DWD), 2024.
- 1040 Boulton, V. L., Black, E., Saado Abdillahi, H., Bailey, M., Harris, C., Kilavi, M., Kniveton, D., MacLeod, D., Mwangi, E., Otieno, G., Rees, E., Rowhani, P., Taylor, O., & Todd, M. C.: Towards drought impact-based forecasting in a multi-hazard context. *Climate Risk Management*, 35, 100402. <https://doi.org/10.1016/J.CRM.2022.100402>, 2022.

1045 Buizza, R.: Introduction to the special issue on “25 years of ensemble forecasting”. *Quarterly Journal of the Royal Meteorological Society*. <https://doi.org/10.1002/qj.3370>, 2019.

Busker, T., de Moel, H., van den Hurk, B., Aerts, J.C.J.H.: Anticipating Weather and Climate Extremes: The Value of Early Action to Reduce Disaster Impacts. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam]. <https://doi.org/10.5463/thesis.837>, 2024.

1050 Busker, T., van den Hurk, B., de Moel, H., & Aerts, J. C. (2025). The value of precipitation forecasts to anticipate floods. *Bulletin of the American Meteorological Society*, 106(3), E473-E491.

Cao, A., Nakamura, S., Otsuyama, K., Namba, M., Yoshimura K.: Current status and challenges in operating flood early warning systems at the local level in Japan. *International Journal of Disaster Risk Reduction*, 112, 104802, [doi.org/10.1016/j.ijdr.2024.104802](https://doi.org/10.1016/j.ijdr.2024.104802), 2024.

Cloke, H., and Pappenberger, F.: Ensemble flood forecasting: A review, *Journal of Hydrology*, 375, Issues 3–4, pp 613-626, [doi.org/10.1016/j.jhydrol.2009.06.005](https://doi.org/10.1016/j.jhydrol.2009.06.005), 2009.

1055 COT: Rapportage Leerevaluatie: Hoogwater Maas en Roer, [https://www.vrln.nl/sites/vrln/files/2022-10/Leerevaluatie\\_Hoogwater\\_Maas\\_en\\_Roer\\_VR\\_Limburg\\_Noord\\_2021\\_rapportage.pdf](https://www.vrln.nl/sites/vrln/files/2022-10/Leerevaluatie_Hoogwater_Maas_en_Roer_VR_Limburg_Noord_2021_rapportage.pdf), 2022a.

Coughlan de Perez, E., Kristoffer B. Berse, Lianne Angelico C. Depante, Evan Easton-Calabria, Elton Pierre R. Evidente, Theodore Ezike, Dorothy Heinrich, Christopher Jack, Alfredo Mahar Francisco A. Lagmay, Selma Lendelvo, Joalane Marunye, Daniel G. Maxwell, Sonia Binte Murshed, Christopher Garimoi Orach, Meethilde Pinto, Leah B. Poole, Komal Rathod, Shampa, Carolyn Van Sant: Learning from the past in moving to the future: Invest in communication and response to weather early warnings to reduce death and damage, *Climate Risk Management*, 38, 100461, <https://doi.org/10.1016/j.crm.2022.100461>, 2022.

Council of the European Union: Council Conclusions on the comprehensive enhancement of the EU's resilience against flooding. <https://data.consilium.europa.eu/doc/document/ST-16841-2024-INIT/en/pdf>, 2024.

1065 Da Costa, J., Ebert, E., Hoffmann, D., Cloke, H. L., & Neumann, J.: Signals without action: A value chain analysis of Luxembourg's 2021 flood disaster. *Natural Hazards and Earth System Sciences*, 26(1), 343–366. <https://doi.org/10.5194/nhess-26-343-2026>.

Dasgupta, A., Louise Arnal, Rebecca Emerton, Shaun Harrigan, Gwyneth Matthews, Ameer Muhammad, Karen O'Regan, Teresa Pérez-Ciria, Emixi Valdez, Bart van Osnabrugge, Micha Werner, Carlo Buontempo, Hannah Cloke, Florian Pappenberger, Ilias G. Pechlivanidis, Christel Prudhomme, Maria-Helena Ramos, Peter Salamon: Connecting hydrological modelling and forecasting from global to local scales: Perspectives from an international. *J. of Flood Risk Management*, <https://doi.org/10.1111/jfr3.12880>, 2024.

1070 Demuth, N. and Rademacher, S.: Flood Forecasting in Germany – Challenges of a Federal Structure and Transboundary Cooperation, *Flood Forecasting: A Global Perspective*, Elsevier, 125–151, 2016.

Formatted: Dutch (Netherlands)

Formatted: English (United Kingdom)

Formatted: Default Paragraph Font, English (United Kingdom)

- Dewals, Benjamin, Erpicum, Sébastien, Piroton, Michel, Archambeau, Pierre: Extreme floods in Belgium. The July 2021 extreme floods in the Belgian part of the Meuse basin, 2021. <https://henry.baw.de/server/api/core/bitstreams/6b2b809e-becd-465e-bd3b-d24e3a9f467a/content>
- DGH: l'hydrométrie en wallonie. <https://hydrometrie.wallonie.be/home.html>, last access: 21 February 2025, 2024.
- 1080 DGH: Prévission des crues, <https://hydrometrie.wallonie.be/home/en-savoir-plus/prevision-des-crues.html>, last access: 16 October 2025, 2025.
- DKKV: Governance und Kommunikation im Krisenfall des Hochwasserereignisses im Juli 2021, DKKV-Schriftenreihe Nr. 63, Januar 2024, Bonn, 2024.
- Dottori, F., Kalas, M., Salamon, P., Bianchi, A., Alfieri, L., and Feyen, L.: An operational procedure for rapid flood risk assessment in Europe, *Nat. Hazards Earth Syst. Sci.*, 17, 1111–1126, <https://doi.org/10.5194/nhess-17-1111-2017>, 2017.
- 1085 DWD: Bericht zum Ablauf und Umfang der operationellen Warn- und Beratungstätigkeit des Deutschen Wetterdienstes im Vorlauf und während des Unwetters 12–15 Juli 2021 in NRW und Rheinland-Pfalz, ausgelöst durch das Tief “Bernd.” [https://www.dwd.de/DE/presse/hintergrundberichte/berichts\\_warnchronologie.pdf.pdf?\\_\\_blob=publicationFile&v=2](https://www.dwd.de/DE/presse/hintergrundberichte/berichts_warnchronologie.pdf.pdf?__blob=publicationFile&v=2), 2021.
- 1090 European Commission: European Civil Protection and Humanitarian Aid Operations: Luxembourg. [https://civil-protection-humanitarian-aid.ec.europa.eu/what/civil-protection/national-disaster-management-system/luxembourg\\_en](https://civil-protection-humanitarian-aid.ec.europa.eu/what/civil-protection/national-disaster-management-system/luxembourg_en) (accessed on 18/12/2024), 2024.
- European Environment Agency: European Climate Risk Assessment (Publication). Luxembourg: European Environment Agency <https://www.eea.europa.eu/publications/european-climate-risk-assessment>, 2024.
- 1095 Endendijk, T., Botzen, W., de Moel, H., Aerts, J., Duijndam, S., Slager, K., Kok, M.: Experience From the 2021 Floods in the Netherlands: Household Survey Results on Impacts and Responses. *Journal of Coastal and Riverine Flood Risk*, 2. <https://doi.org/10.59490/jcrfr.2023.0009>, 2023a.
- 1100 Endendijk, T., Botzen, W., de Moel, H., Aerts, J., Slager, K., Kok, M.: Flood Vulnerability Models and Household Flood Damage Mitigation Measures: An Econometric Analysis of Survey Data. *Water Resources Research*, doi-org.vu-nl.idm.oclc.org/10.1029/2022WR034192, 2023b.
- ENW: Hoogwater 2021 Feiten en Duiding. [https://www.hkv.nl/wp-content/uploads/2021/09/ENW-Hoogwater-2021-Feiten-en-Duiding\\_BK\\_etal.pdf](https://www.hkv.nl/wp-content/uploads/2021/09/ENW-Hoogwater-2021-Feiten-en-Duiding_BK_etal.pdf), 2021.
- Frogner, I., Andrae, U., Bojarova, J., Callado, A., Escriba, P., Feddersen, H., Hally, A., Kauhane, J., Randriamampianina, R., Singleton, A., Smet, G., van den Veen, S., Vignes, A. : HarmonEPS—The HARMONIE Ensemble Prediction System. *Weather and Forecasting*, 10.1175/WAF-D-19-0030.1, 2019.
- 1105 Gerard, L., Piriou, J. M., Brožková, R., Geleyn, J. F., & Banciu, D.: Cloud and precipitation parameterization in a meso-gamma-scale operational weather prediction model. *Monthly weather review*, 137(11), 3960-3977, 2009.

- 1110 Golding, B., Mittermaier, M., Ross, C., Ebert, B., Panchuk, S., Scolobig, A., & Johnston, D.: A value chain approach to optimising early warning systems. Contributing paper to GAR2019. UNDRR, [www. preventionweb.net/publications/view/65828](http://www.preventionweb.net/publications/view/65828), 2019.
- Government of Wallonia, Plan Régional de Gestion de Crise, <https://cortex.wallonie.be/home/nos-outils/plan-regional-gestion-de-crise.html>, last access: 27 October 2025, 2025.
- Haag, I., Ehret, U., Demuth, N., & Moritz, K.: ProFoUnD: Ein Werkzeug zur statistischen Analyse von Abfluss-und  
1115 Wasserstandsvorhersagen und zur Ermittlung von Vorhersageunsicherheiten, 2013.
- Haiden, T., Janousek, M., Vitart, F., Bouallègue, Z. B., Ferranti, L., Prates, F., & Richardson, D.: Evaluation of ECMWF forecasts, including the 2018 upgrade. Reading, UK: European Centre for Medium Range Weather Forecast, 2018.
- Hofmann, J., & Schüttrumpf, H.: floodGAN: Using Deep Adversarial Learning to Predict Pluvial Flooding in Real Time. *Water*, 13, 2255. <https://doi.org/10.3390/w13162255>, 2021.
- 1120 IIASA: PERC Flood event review ‘Bernd’. [https://pure.iiasa.ac.at/id/eprint/18144/1/Zurich\\_PERC\\_Analysis\\_Bernd\\_ENG%20%282%29.pdf](https://pure.iiasa.ac.at/id/eprint/18144/1/Zurich_PERC_Analysis_Bernd_ENG%20%282%29.pdf), 2022.
- IKSMS: Hochwasservorhersage als transnationale Aufgabe des Technischen Ausschusses, last access: 16 October 2025, 2025.
- Jiang, X., Liping Zhang, Zhongmin Liang, Xiaolei Fu, Jun Wang, Jiaxin Xu, Yuchen Zhang, Qi Zhong: Study of early flood warning based on postprocessed predicted precipitation and Xinanjiang model, *Weather and Climate Extremes*, 42,  
1125 100611, [doi.org/10.1016/j.wace.2023.100611](https://doi.org/10.1016/j.wace.2023.100611), 2023.
- Journée, M., Goudenhoofd, E., Vannitsem, S., & Delobbe, L.: Quantitative rainfall analysis of the 2021 mid-July flood event in Belgium. *Hydrology and Earth System Sciences*, 27(17), 3169-3189, <https://doi.org/10.5194/hess-27-3169-2023>, 2023.
- Kaltenberger, R., Schaffhauser, A., & Staudinger, M.: “What the weather will do” – results of a survey on impact-oriented and  
1130 impact-based warnings in European NMHSs. *Advances in Science and Research*, 17, 29–38. <https://doi.org/10.5194/asr-17-29-2020>, 2020.
- KNMI: Het regent nu harder in Zuid-Limburg door klimaatverandering. <https://www.knmi.nl/over-het-knmi/nieuws/extreme-neerslag-in-zuid-limburg>, last access: 13 January 2024, 2021.
- KNMI: Wat kan ik verwachten en wat kan ik doen bij een weerwaarschuwing? <https://www.knmi.nl/over-het-knmi/nieuws/wat-kan-ik-verwachten-en-wat-kan-ik-doen-bij-een-weerwaarschuwing/>, last access: 4 December 2024,  
1135 2024.
- KNMI: Uitleg over: Early Warning Centre (EWC): <https://www.knmi.nl/kennis-en-datacentrum/uitleg/early-warning-centre>, last access: 21-10-2025, 2025.
- Kok, K.: Beschrijving en toepassingen van het PASCAL-systeem: Een automatisch waarschuwingssysteem voor gevaarlijk  
1140 weer. KNMI: technisch rapport, De Bilt, 2022.

- Kreibich, H., Hudson, P., and Merz, B.: Knowing What to Do Substantially Improves the Effectiveness of Flood Early Warning: Bulletin of the American Meteorological Society. pp E1450–E1463, doi.org/10.1175/BAMS-D-20-0262.1, 2021.
- 1145 Kiger, M. E., & Varpio, L. Thematic analysis of qualitative data: AMEE Guide No. 131. *Medical teacher*, 42(8), 846-854, 2020.
- Kuller, M., Schoenholzer, K., Lienert, J.: Creating effective flood warnings: A framework from a critical review, *Journal of Hydrology*, 602, 126708, doi.org/10.1016/j.jhydrol.2021.126708, 2021.
- Landesamt für Umwelt Rheinland-Pfalz, Hochwasservorhersagezentrale Rheinland-Pfalz, <https://www.hochwasser.rlp.de/erlaeuterungen#warnkarte-legende>, last access: 16 October 2025, 2025.
- 1150 LANUK, [Hochwassermeldepegel](https://www.umwelt.nrw.de/system/files/media/document/file/uebersicht_hochwassermeldepegel.pdf) des LANUK, [https://www.umwelt.nrw.de/system/files/media/document/file/uebersicht\\_hochwassermeldepegel.pdf](https://www.umwelt.nrw.de/system/files/media/document/file/uebersicht_hochwassermeldepegel.pdf), last access: 6 February 2026, 2025.
- LAWA: [Bericht zur Evaluation der Handlungsempfehlungen zur weiteren Verbesserung von Grundlagen und Qualität der Hochwasservorhersage an den deutschen Binnengewässern](#). d, 2020.
- 1155 Lehmkuhl, F., Schüttrumpf, H., Schwarzbauer, J., Brüll, C., Dietze, M., Letmathe, P., Völker, C., & Hollert, H.: Assessment of the 2021 summer flood in Central Europe. *Environmental Sciences Europe*, 34(1), 107. <https://doi.org/10.1186/s12302-022-00685-1>, 2022.
- Leutbecher, M., & Palmer, T. N.: Ensemble forecasting. *Journal of Computational Physics*, 227(7), 3515–3539. <https://doi.org/10.1016/j.jcp.2007.02.014>, 2008.
- 1160 Le gouvernement du Grand-Duché de Luxembourg: Info Crise, <https://infocrise.public.lu/en>, last access: 27 October 2025, 2025.
- Lindenlaub, S., Matthes, K. S., & Thieken, A. H.: Improving the understandability of flood warning messages–Explorative study on design preferences. *International Journal of Disaster Risk Reduction*, 111, 104685, 2024a.
- Lindenlaub, S., A., Heidenreich, H., Murdock, & Thieken, A. H.: Warnsituation in den Einzugsgebieten: Erkenntnisse aus Betroffenenbefragungen. In: (DKKV, Hrsg.): *Governance und Kommunikation im Krisenfall des Hochwasserereignisses im Juli 2021*. DKKV-Schriftenreihe 63, Bonn, S. 38-52, 2024b.
- 1165 Mahdavian, F., Stephen Platt, Marcus Wiens, Miriam Klein, Frank Schultmann: Communication blackouts in power outages: Findings from scenario exercises in Germany and France, *International Journal of Disaster Risk Reduction*, 46, <https://doi.org/10.1016/j.ijdr.2020.101628>, 2020.
- 1170 Meléndez-Landaverde, E. R., Werner, M., & Verkade, J.: Exploring protective decision-making in the context of impact-based flood warnings. *Journal of Flood Risk Management*, 13(1), e12587, 2020.
- Merz, B., Kuhlicke, C., Kunz, M., Pittore, M., Babeyko, A., Bresch, D. N., Domeisen, D. I. V., Feser, F., Koszalka, I., Kreibich, H., Pantillon, F., Parolai, S., Pinto, J. G., Punge, H. J., Rivalta, E., Schröter, K., Strehlow, K., Weisse, R., & Wurpts,

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Formatted: Font:

Formatted: Font: English (United Kingdom)

Formatted: Font:

Formatted: English (United Kingdom)

Formatted: Dutch (Netherlands)

- 1175 A.: Impact Forecasting to Support Emergency Management of Natural Hazards. *Reviews of Geophysics*, 58(4).  
<https://doi.org/10.1029/2020RG000704>, 2020.
- Ministerie van Justitie en Veiligheid: Naar een nieuwe toekomst voor waarschuwing en alarmering.  
<https://open.overheid.nl/documenten/dpc-8d5932adf1c2656e52b2cc870be0c248b7889d69/pdf>, last access: 20  
February 2025, 2024.
- 1180 Ministerium des Innern und für Sport Rheinland-Pfalz: Rahmen- Alarm- und Einsatzplan Hochwasser, [https://bks-portal.rlp.de/system/files/og-group/57/dokumente/RAEP%20Hochwasser%20Stand%2018.08.2020\\_0.pdf](https://bks-portal.rlp.de/system/files/og-group/57/dokumente/RAEP%20Hochwasser%20Stand%2018.08.2020_0.pdf) (In  
German), 2020.
- Mohr, S., U. Ehret, M. Kunz, P. Ludwig, A. Caldas-Alvarez, J.E. Daniell, F. Ehmele, H. Feldmann, M.J. Franca, C. Gattke,  
M. Hundhausen, P. Knippertz, K. Küpfer, B. Mühr, J.G. Pinto, J. Quinting, A.M. Schäfer, M. Scheibel, F. Seidel, C.  
Wisotzky: A multi-disciplinary analysis of the exceptional flood event of July 2021 in Central Europe. Part 1: event  
1185 description and analysis. *Nat. Hazards Earth Syst. Sci.*, 23, 2 (2023), pp. 525-551, 10.5194/nhess-23-525-2023, 2023.
- Najafi, H., Kumar Shrestha, P., Rakovec, O., Apel, H., Vorogushyn, S., Kumar, R., Thober, S., Merz, B., & Samaniego, L.:  
High-resolution impact-based early warning system for riverine flooding. *Nature Communications*, 15, 3726.  
<https://doi.org/10.1038/s41467-024-48065-y>, 2024.
- 1190 Ommer, J., Neumann, J., Kalas, M., Blackburn, S., and Cloke, H. L.: Surprise floods: the role of our imagination in preparing  
for disasters, *Nat. Hazards Earth Syst. Sci.*, 24, 2633–2646, <https://doi.org/10.5194/nhess-24-2633-2024>, 2024.
- Pappenberger, F., Cloke, H. L., Parker, D. J., Wetterhall, F., Richardson, D. S., & Thielen, J.: The monetary benefit of early  
flood warnings in Europe. *Environmental Science and Policy*. <https://doi.org/10.1016/j.envsci.2015.04.016>, 2015.
- 1195 Pappenberger, F., Pagano, T. C., Brown, J. D., Alfieri, L., Lavers, D. A., Berthet, L., Bressand, F., Cloke, H. L., Cranston, M.,  
Danhelka, J., Demargne, J., Demuth, N., de Saint-Aubin, C., Feikema, P. M., Fresch, M. A., Garçon, R., Gelfan, A.,  
He, Y., Hu, Y.-Z., Janet, B., Jurdy, N., Javelle, P., Kuchment, L., Laborda, Y., Langsholt, E., Le Lay, M., Li, Z. J.,  
Mannessiez, F., Marchandise, A., Marty, R., Meißner, D., Manful, D., Organde, D., Pourret, V., Rademacher, S.,  
Ramos, M.-H., Reinbold, D., Tibaldi, S., Silvano, P., Salamon, P., Shin, D., Sorbet, C., Sprokkereef, E., Thiemig, V.,  
Tuteja, N. K., van Andel, S. J., Verkade, J. S., Vehviläinen, B., Vogelbacher, A., Wetterhall, F., Zappa, M., Van der  
1200 Zwan, R. E., and Thielen-del Pozo, J.: Hydrological Ensemble Prediction Systems Around the Globe, in: *Handbook of  
Hydrometeorological Ensemble Forecasting*, Springer, Berlin, Heidelberg, [https://doi.org/10.1007/978-3-642-40457-3\\_47-1](https://doi.org/10.1007/978-3-642-40457-3_47-1), 2016.
- Parker, D. and Priest, S.: The Fallibility of Flood Warning Chains: Can Europe’s Flood Warnings Be Effective?, *Water  
Resources Management: An International Journal*, Published for the European Water Resources Association (EWRA),  
Springer; European Water Resources Association (EWRA), vol. 26(10), pages 2927-2950, August, 2012.
- 1205 Poolman, E., Webster, E., Sebege, E., Rae, K. Phillay, D., van Staden, M.: Project on Developing and Implementing an Impact-  
based Severe Weather Warning System. South African National Disaster Management Centre – South Africa. Available

Field Code Changed

- from: <https://invest.edendm.gov.za/wp-content/uploads/2018/03/Water-Safety-and-Security-Project-onDeveloping-and-Implementingan-Impact-based-Severe-Weather-Warning-System.pdf>, 2018.
- 1210 Red Cross Red Crescent Climate Centre, Met Office, ARRCC, UK AID, Anticipation Hub, & Risk-informed Early Action Partnership: The Future of Forecasts: Impact-based forecasting for early action. <https://www.forecast-based-financing.org/wp-content/uploads/2020/09/Impact-based-forecasting-guide-2020.pdf>, 2020.
- Rhein, B. and Kreibich, H.: Causes of the exceptionally high number of fatalities in the Ahr valley, Germany, during the 2021 flood, *Nat. Hazards Earth Syst. Sci.*, 25, 581–589, <https://doi.org/10.5194/nhess-25-581-2025>, 2025.
- 1215 Richardson, D. S.: Predictability and economic value. Seminar on Predictability of Weather and Climate, 9-13 September 2002, 321–332. <https://www.ecmwf.int/node/11922>, 2003.
- Rodríguez Castro, D., Cools, M., Roucour, S., Archambeau, P., Molinari, D., Scorzini, A. R., Dessers, C., Erpicum, S., Pirotton, M., Teller, J. & Dewals, B.: Can macro-or meso-scale coping capacity variables improve the classification of building flood losses?. *Natural Hazards*, 1-31, 2025.
- 1220 Schroeter, S., Richter, H., Arthur, C., Wilke, D., Dunford, M., Wehner, M., and Ebert, E.: Forecasting the impacts of severe weather, *The Aust. J. Emergency Manage.*, 36, 76–83, <https://doi.org/10.3316/informit.766653620579430>, 2021.
- Scolobig, A., Potter, S., Kox, T., Kaltenberger, R., Weyrich, P., Chasco, J., Golding, B., Hilderbrand, D., Fleischhut, N., Uprety, D., & Rana, B.: Connecting Warning with Decision and Action: A Partnership of Communicators and Users. In *Towards the “Perfect” Weather Warning* (pp. 47–85). Springer International Publishing. [https://doi.org/10.1007/978-3-030-98989-7\\_3](https://doi.org/10.1007/978-3-030-98989-7_3), 2022.
- 1225 Seebauer S and Babicky P.: Trust and the communication of flood risks: comparing the roles of local governments, volunteers in emergency services, and neighbours. *J Flood Risk Manag.* 2018 Sep;11(3):305-316. doi: 10.1111/jfr3.12313. Epub 2017 Jul 31. PMID: 32030096; PMCID: PMC6991925, 2017.
- Smet, G.: RMI-EPS: A prototype convection-permitting EPS for Belgium. ———<https://orfeo.belnet.be/handle/internal/7057>, 2017.
- 1230 Smith, P., Pappenberger, F., Wetterhall, F., Thielen, J., Krzeminski, B., Salamon, P., Muraro, D., Kalas, M., and Baugh, C.: On the operational implementation of the European Flood Awareness System (EFAS), ECMWF Tech. Memorandum 778, ECMWF, Reading, UK., 1–34, 2016.
- Speight, L. J., Cranston, M. D., White, C. J., & Kelly, L.: Operational and emerging capabilities for surface water flood forecasting. *Wiley Interdisciplinary Reviews: Water*, 8(3), e1517, 2021.
- 1235 Thieken, A. H., Bubeck, P., Heidenreich, A., von Keyserlingk, J., Dillenaar, L., & Otto, A.: Performance of the flood warning system in Germany in July 2021 - insights from affected residents. *Natural Hazards and Earth System Sciences*, 23(2), 973–990. <https://doi.org/10.5194/nhess-23-973-2023>, 2023a.
- Thieken, A., Zenker, M.-L. & Bubeck, P.: Flood-related fatalities during the flood of July 2021 in North Rhine-Westphalia, Germany: what can be learnt for future flood risk management? *Journal of Coastal and Riverine Flood Risk.*, 2, p. 5. <https://doi.org/10.59490/jcrfr.2023.0005>, 2023b.
- 1240

- Tradowsky, J. S., Philip, S. Y., Kreienkamp, F., Kew, S. F., Lorenz, P., Arrighi, J., ... & Wanders, N.: Attribution of the heavy rainfall events leading to severe flooding in Western Europe during July 2021. *Climatic Change*, 176(7), 90, 2023.
- Van Steenbergen, N., Ronsyn, J., Willems, P.: A non-parametric data-based approach for probabilistic flood forecasting in support of uncertainty communication. *Environmental Modelling & Software*, 33, 92-105, 2012.
- 1245 Veiligheidsregio Limburg-Noord and Zuid-Limburg: Rampbestrijdingsplan Hoogwater Limburg 2023-2026. [https://www.vrln.nl/sites/vrln/files/2024-04/RBP%20Hoogwater%20Limburg%202023-2026\\_DEF.pdf](https://www.vrln.nl/sites/vrln/files/2024-04/RBP%20Hoogwater%20Limburg%202023-2026_DEF.pdf), 2023.
- Verkade, J. S., & Werner, M. G. F.: Estimating the benefits of single value and probability forecasting for flood warning. *Hydrology and Earth System Sciences*, 15(12), 3751-3765, 2011.
- Verkade, J. S., Brown, J. D., Davids, F., Reggiani, P., & Weerts, A. H.: Estimating predictive hydrological uncertainty by dressing deterministic and ensemble forecasts; a comparison, with application to Meuse and Rhine. *Journal of Hydrology*, 555, 257-277, 2017.
- 1250 Vorogushyn, S., Apel, H., Kemter, M., & Thieken, A. H.: Analyse der Hochwassergefährdung im Ahrtal unter Berücksichtigung historischer Hochwasser. *Hydrologie Und Wasserbewirtschaftung*, 66(5), 244-254. [https://doi.org/10.5675/HyWa\\_2022.5\\_2](https://doi.org/10.5675/HyWa_2022.5_2), 2022.
- 1255 Weyrich, P., Scolobig, A., & Patt, A.: Dealing with inconsistent weather warnings: Effects on warning quality and intended actions. *Meteorological Applications*, 26(4), 569-583, 2019.
- WMCN: Landelijk Draaiboek Hoogwater en Overstromingsdreiging <https://iplo.nl/@250864/landelijk-draaiboek-hoogwater/> (in Dutch), 2023.
- WMO: Assessment Guidelines for End-to-End Flood Forecasting and Early Warning Systems [https://library.wmo.int/viewer/58245?medianame=1286\\_en\\_#page=1&viewer=picture&o=bookmark&n=0&q=](https://library.wmo.int/viewer/58245?medianame=1286_en_#page=1&viewer=picture&o=bookmark&n=0&q=), 2023.
- 1260 [Wu, W., Emerton, R., Duan, Q., Wood, A. W., Wetterhall, F., & Robertson, D. E.: Ensemble flood forecasting: Current status and future opportunities. \*Wiley Interdisciplinary Reviews: Water\*, 7\(3\), e1432, 2020.](#)