

# Gas-particle partitioning of m-xylene and naphthalene oxidation products: temperature and NO<sub>x</sub> influence

Marwa Shahin<sup>1</sup>, Julien Kammer<sup>1</sup>, Brice Temime-Roussel<sup>1</sup>, and Barbara D'Anna<sup>1</sup>

<sup>1</sup>Aix-Marseille Univ., CNRS, LCE, Marseille, France

Correspondence to: Barbara D'Anna ([barbara.danna@univ-amu.fr](mailto:barbara.danna@univ-amu.fr)), Marwa Shahin ([marwa.shahin@etu.univ-amu.fr](mailto:marwa.shahin@etu.univ-amu.fr))

**Abstract.** Volatile organic compounds (VOCs) react with atmospheric oxidants resulting in oxygenated products of lower volatility known as semi and intermediate volatile organic compounds (S/IVOCs) forming secondary organic aerosols (SOA). Those compounds can partition between the gas and particle phases, a critical process influenced by several environmental parameters, yet poorly constrained. This study aims to evaluate the effect of temperature and VOC/NO<sub>x</sub> ratio on SOA formation and partitioning of individual SOA products from m-xylene and naphthalene OH-oxidation. Experiments are carried out in an oxidation flow reactor (OFR) and products are identified and quantified using a proton transfer reaction time-of-flight mass spectrometer (PTR-ToF-MS) coupled to a Chemical Analysis of aeRosol ONline (CHARON) inlet. Results show that lower temperatures significantly enhance SOA formation, while lower VOC/NO<sub>x</sub> ratios reduce it. Gas-phase m-xylene major products are C<sub>3</sub>, C<sub>5</sub> and C<sub>8</sub> whereas particle-product distributions exhibit a progressive increase from C<sub>2</sub> to C<sub>8</sub>. In contrast, naphthalene products partition more readily into the condensed phase, with C<sub>8</sub>-C<sub>10</sub> products dominating. Most of the oxidation products from both precursors exhibit a volatility distribution in the SVOC regime, with fewer in the IVOC regime. The decrease in temperature shifts the effective saturation concentration ( $C_i^*$ ) values towards lower values, though no clear relationship between  $C_i^*$  and oxidation state is observed. A comparison between observed and estimated volatilities using a model based on group contribution method (SIMPOL.1) reveals systematic deviations for both light molecules and heavy compounds, suggesting a need for improved predictive models.

## 1 Introduction

Aromatic hydrocarbons (AHs) are an ubiquitous class of air pollutants and contribute to an important fraction of the total volatile organic compounds (VOCs) in urban environments; the relative contribution may vary and depends on location and season (Calvert et al., 2002; Jiang et al., 2017; Montero-Montoya et al., 2018). Among the different AHs, xylene and naphthalene are important anthropogenic VOCs primarily emitted from petrochemical industries, biomass burning, diesel and gasoline engines, or through solvent evaporation (Fang et al., 2021, 2024; Wu et al., 2020; Xuan et al., 2021). In the atmosphere, AHs react with common oxidants as hydroxyl (OH), nitrate (NO<sub>3</sub>) and chloride (Cl) radicals and ozone (O<sub>3</sub>) leading to the formation of oxygenated reaction products of lower vapor pressures also known as semi-volatile organic compounds (SVOCs) and intermediate-volatile organic compounds (IVOCs) (Seinfeld and Pandis, 2016). These compounds

may partition to the particle phase forming secondary organic aerosols (SOA) that represent approximately 60% of ambient organic aerosol (Huang et al., 2020, 2014) and have an impact on visibility (Li et al., 2014; Liu et al., 2017), climate (Liu and Matsui, 2020; Shrivastava et al., 2017) and human health (Anderson et al., 2012; Berlinger et al., 2024; Singh and Tripathi, 2021; Thangavel et al., 2022). Indeed, particulate air pollution is closely correlated to the progression of numerous respiratory diseases, in addition to cancer, cardiovascular diseases, and neurological damage (Liu et al., 2022a; Singh and Tripathi, 2021; Song et al., 2017; Thangavel et al., 2022). In the EU, each year, around 238,000 premature deaths are attributable to fine particulate matter (PM<sub>2.5</sub>) exposure (European Environment Agency, 2022). Improving our understanding of particle sources and properties is thus pivotal to improve a more sustainable environment.

Over the last decades, many laboratory and modelling studies investigated SOA generated by AHs reaction products (Chen et al., 2018; Forstner et al., 1997; Klodt et al., 2023; Lannuque et al., 2018; Li et al., 2022; Liu et al., 2023, 2022b; Lu et al., 2024; Song et al., 2007; Srivastava et al., 2022; Tian et al., 2023). The OH radical oxidation of monoaromatic compounds operates through two main common pathways (Atkinson et al., 1991; Bloss et al., 2005; Calvert et al., 2002; Forstner et al., 1997; Pan and Wang, 2014). In the case of m-xylene, the first pathway is the OH radical addition to the aromatic ring, predominantly on the ortho position (Fan and Zhang, 2008) to form a hydroxy-methyl-benzyl radical with a subsequent addition of O<sub>2</sub> to form a bicyclic radical, a major ring opening product channel (Zhao et al., 2005). The second reaction pathway is the H-abstraction from the methyl group to form a methyl benzyl radical, with a subsequent addition of O<sub>2</sub> to form a benzyl peroxy radical (Atkinson et al., 1991; Molina et al., 1999). In the presence of NO<sub>x</sub>, the peroxy radical mainly reacts with NO to form an alkoxy radical, and finally an abstraction by O<sub>2</sub> leads to the formation of m-tolualdehyde (Srivastava et al., 2023). In the case of naphthalene, OH radical reaction is initiated exclusively by OH addition, most predominantly at the α-carbon (68 %) adjacent to the fusion of its two aromatic rings (Wang et al., 2007). Then, a hydroxy cyclohexadienyl radical is formed, which further reacts with NO<sub>2</sub> to form nitrogen-containing compounds, or with O<sub>2</sub> to form various products out of which 2-formylcinnamaldehyde is major (Nishino et al., 2009).

The NO<sub>x</sub> level has been shown to have significant yet possibly contrasting effects on SOA formation (Chan et al., 2009; Liu et al., 2024; Ng et al., 2007; Qi et al., 2020; Sarrafzadeh et al., 2016; Song et al., 2005; Zhu et al., 2021). Some studies have reported a decrease in SOA yield under high NO<sub>x</sub> conditions which can be explained by the termination reactions of NO with the peroxy (RO<sub>2</sub> and HO<sub>2</sub>) and OH radicals, limiting the formation of lower volatility compounds (Chan et al., 2009; Ng et al., 2007; Song et al., 2005). Nonetheless, Zhu et al. (2021) reported an increase in SOA mass under elevated levels of NO<sub>x</sub> compared to NO<sub>x</sub> free experiments. Enhanced SOA formation under high NO<sub>x</sub> has also been linked to the formation of organic nitrates and the isomerization of alkoxy radicals into low-volatility products, or related to a change in OH concentration due to the presence of NO<sub>x</sub> (Ng et al., 2007; Schwantes et al., 2019; Shi et al., 2022; Srivastava et al., 2023). Under low NO<sub>x</sub> conditions, RO<sub>2</sub> radicals primarily react with HO<sub>2</sub> to form low-volatility organic hydroperoxides, which contribute to new particle formation and increase SOA mass (Xu et al., 2014; Zhao et al., 2018). These findings illustrate that NO<sub>x</sub> is capable of both inhibiting and promoting SOA formation depending on the NO<sub>x</sub> regime.

63 A key process determining the fate of the oxidation products generated from AH oxidation is their ability to partition between  
64 gas and particle phases, either throughout nucleation forming new particles or by condensing on pre-existing particles.  
65 Partitioning is influenced by many environmental parameters such as temperature, relative humidity, the nature and the  
66 diameter of the pre-existing particles on which they condense and the physio-chemical properties of the S/IVOC condensing  
67 (Kamens et al., 2011; Kim et al., 2007; Lannuque et al., 2018; Loza et al., 2012; Ng et al., 2007; Qi et al., 2010; Sato et al.,  
68 2007; Takekawa et al., 2003; Warren et al., 2009; Xu et al., 2015).

69 It is well recognized that SOA yields increase at lower temperatures, a trend consistently reported for both terpenes and  
70 isoprene (Clark et al., 2016; Deng et al., 2021; Svendby et al., 2008; Virtanen et al., 2010), aromatics (Lannuque et al., 2023;  
71 Svendby et al., 2008), alkanes (n-dodecane in Fan et al., 2025; Li et al., 2020) and amines (Price et al., 2016). This behavior  
72 is attributed primarily to the decrease of the vapor pressures of the compounds, displacing the equilibrium towards the particle  
73 phase. However, the impact of temperature on SOA composition is not fully understood, and seems to depend on other  
74 experimental conditions (precursor, seed acidity, etc.). For biogenic precursors, previous studies reported more oligomer  
75 formation at lower temperatures (Li et al., 2020; Fan et al., 2025), driven by increased SVOC partitioning and condensed-  
76 phase reactions, while others observed the opposite, attributing higher-temperature oligomerization to radical or acid-catalyzed  
77 reactions (Clark et al., 2016; Deng et al., 2021; Price et al., 2016). Additionally, Li et al. (2019) highlighted how lower  
78 temperatures increase SOA viscosity, suppressing evaporation and favoring retention of low-volatility species. Conversely,  
79 Lamkaddam et al., (2017) found a weak temperature sensitivity in SOA formation from n-dodecane, suggesting that extremely  
80 low-volatility products and compensating shifts in product types may offset volatility effects. Regarding aromatic compounds,  
81 only Lannuque et al. (2023) investigated the effect of temperature on SOA chemical composition at molecular level, showing  
82 a general agreement of product distribution between 280 K and 295 K. Thus, more studies on different aromatic precursors  
83 and experimental conditions are needed to complete our understanding of the temperature effect on SOA formation from  
84 aromatic precursors.

85 Although temperature is recognized as a major factor governing gas-particle partitioning, most prior studies have mostly  
86 focused on the bulk SOA yields or selected classes of compounds (Ahn et al., 2021; Bahrami et al., 2024; John et al., 2018;  
87 Rutter and Schauer, 2007; Svendby et al., 2008; Takekawa et al., 2003; Wei et al., 2016; Zhang et al., 2023; Zhou, 2021).  
88 While these approaches have advanced the understanding of SOA formation and representation in models, they often lack  
89 molecular level resolved resolution. Experimental studies that probe the effect of temperature on the partitioning of individual  
90 SOA oxidation products, remain limited and relatively recent (Deng et al., 2021; Fan et al., 2025; Lannuque et al., 2023; Li et  
91 al., 2024). Such measurements are particularly important for refining SOA volatility parameterizations under atmospherically  
92 relevant cold conditions, such as those occurring during nighttime or wintertime episodes.

93 The oxidation flow reactor (OFR, Fig. 1) is a type of continuous-flow reactor that uses substantially elevated oxidant levels to  
94 rapidly simulate atmospheric oxidation chemistry (Kang et al., 2007; Peng and Jimenez, 2020). Lannuque et al. (2023) have  
95 studied the gas-particle partitioning of toluene in a similar OFR system. Experiments carried at 280 K were characterized by

a higher mass loading in the particle phase compared to those at 295 K, as well as a shift in volatility values of oxidation products towards lower volatility values.

Direct measurements of S/IVOCs in real time remains challenging due to their complexity, diversity, and low concentrations. The analytical development of the “CHemical Analysis of aeRosols ONline” (CHARON) inlet coupled with a proton-transfer-reaction-time of flight-mass spectrometry (PTR-ToF-MS) enables alternative online measurements of both gas and particle phases at molecular level (Eichler et al., 2015). It also allows measuring both gas and particle phases with reduced artefacts associated with particle collection and thermal desorption compared to traditional techniques (Peng et al., 2023). Promising applications have been shown in several laboratory and field studies (Lannuque et al., 2023; Müller et al., 2017; Muller et al., 2017; Piel et al., 2021), while its application to gas-particle partitioning investigations is relatively new (Gkatzelis et al., 2018; Lannuque et al., 2023; Peng et al., 2023; Piel et al., 2021).

The aim of the present study is to evaluate gas-particle partitioning of S/IVOCs involved in SOA formation from the photooxidation of AHs in different conditions. For that purpose, the photooxidation of AH compounds was investigated using an online CHARON-PTR-ToF-MS coupled to an oxidation flow reactor. These laboratory experiments aim at 1) identifying the gas and particle phase products of m-xylene and naphthalene at a molecular level; 2) evaluating the partitioning behavior of individual SOA products; and 3) assessing the effect of atmospheric conditions (temperature variation and NO<sub>x</sub>/VOC ratio) on this partitioning. Two compounds, m-xylene and naphthalene, are selected, for the following reasons, i) their ubiquity in urban areas (Fang et al., 2021; Wu et al., 2020; Xuan et al., 2021) ii) their known reactivity with OH radicals that is in the same order of magnitude (Calvert et al., 2015), and iii) their SOA formation potential has been previously demonstrated (Chan et al., 2009; Chen et al., 2016, 2021; Kleindienst et al., 2012; Li et al., 2022; Loza et al., 2012; Lu et al., 2024; Ng et al., 2007; Sato et al., 2022; Song et al., 2005, 2007; Srivastava et al., 2022; Takekawa et al., 2003; Ye et al., 2024; Zhang et al., 2019).

## 2 Methods

### 2.1 The OFR experimental setup

Photooxidation experiments have been conducted in a 19.3 L cylindrical aerosol oxidation flow reactor (OFR, 153 mm internal diameter, 105 cm length) made up of quartz, vertically oriented, and surrounded by 6 UVB lamps (Helios Italquartz) with a continuous emission spectrum in the 280-350 nm range ( $\lambda_{\text{max}} = 310$  nm, Fig. 1). An external air conditioning unit is connected to the tube allowing temperature control in the range 280-295 K. The gas phase stream consists of humid air, the selected VOC precursor (m-xylene or naphthalene), nitrogen dioxide (NO<sub>2</sub>) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). For humidification, two glass bottles containing milliQ water are bubbled at 0.2-1 L min<sup>-1</sup> by pure N<sub>2</sub> in order to maintain relative humidity (RH) around 50 % (from 35 % to 65 %, among all experiments). A constant concentration of m-xylene is generated by flowing a constant flow of 0.15 L min<sup>-1</sup> of N<sub>2</sub> over a permeation tube containing a pure solution ( $\geq 99$  % purity, Sigma Aldrich) placed in an oven and kept at a constant temperature of 308 K. Some solid naphthalene (99% purity, Sigma Aldrich) is kept in an iced water bath

while headspace is flown at  $0.1 \text{ L min}^{-1}$  by pure  $\text{N}_2$  to generate a constant flow of naphthalene in the flow tube.  $\text{NO}_2$  is introduced using a cylinder ( $100 \pm 5 \text{ ppm}$  in  $\text{N}_2$ , Linde) and diluted in  $\text{N}_2$  using different mass flow controllers to the desired mixing ratios prior to entering the tube (varying from 40 to 340 ppbV), depending on the required VOC/ $\text{NO}_x$  conditions. A hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) solution (50 % in  $\text{H}_2\text{O}$ , stabilized, Sigma Aldrich) is used as a hydroxyl radical (OH) precursor and is constantly introduced in the OFR by bubbling pure  $\text{N}_2$  in the solution at a flow of  $0.1\text{-}0.2 \text{ L min}^{-1}$ . For each experiment, the OH radical concentration generated was estimated by fitting the VOC precursor (m-xylene or naphthalene) decay assuming a pseudo first order reaction with OH radicals using temperature-dependent values of the kinetic rate constant as recommended from NIST Kinetics Database. This estimation is also based on the hypothesis that the other reactions with OH do not limit the reaction of the precursor in the first seconds following lamps switching on. The OH radical concentrations range from  $2.2$  to  $5.4 \times 10^7 \text{ molecules cm}^{-3}$  corresponding approximately to 1.3 and 3 days of atmospheric OH-radical exposure, taking into account a diurnal average hydroxyl radical concentration of  $1.5 \times 10^6 \text{ molecules cm}^{-3}$  (Mao et al., 2009). Due to background contaminations, compounds with  $m/z$  61 and below, mainly acetaldehyde, acetic acid, formaldehyde and formic acid, are not considered in further analysis. Monodispersed ammonium sulphate (AS) seeds serve as a pre-existing surface and are generated by nebulizing a  $10^{-2} \text{ M}$  AS solution (99.5 % purity, Acros Organics) using a TSI atomizer (model 3076), dried through a silica diffusion drier and then size selected in an Aerodynamic Aerosol Classifier (AAC, Cambustion) to generate monodisperse aerosols with an average diameter of 200 nm. The overall input flow is  $2.4 \text{ L min}^{-1}$  to ensure a residence time in the tube of 8 minutes. In this configuration, particle losses (or its transmissions through the OFR) were estimated by comparing the concentration of seed particles at the inlet and outlet of the OFR, when generating seed at 200 nm electrical mobility diameters. These losses were daily checked prior to each experiment and were in the range  $10 \pm 5 \%$ . In addition, precursor losses were estimated to be around 5 % for m-xylene and 10-15 % for naphthalene. Losses of gaseous products generated during SOA experiments were not experimentally evaluated. Lannuque et al. (2023) showed, in a toluene SOA experiment, that wall losses introduced a 10-15 % deviation on the SOA yield when considering both precursors and reactions products. For m-xylene, we can reasonably assume lower losses as the residence time is shorter and the flow tube has a larger inner diameter. For naphthalene, it is probable that the wall losses were higher than that of m-xylene.

## 2.2 Instrumentation and data analysis

Figure 1 shows the schematic of the experimental set-up, where several online instruments are used to characterize gas and particle phase chemical compositions as well as particle number and size distribution.


A commercial Proton Transfer Reaction-Time of Flight-Mass Spectrometer (PTR-ToF-MS 6000X2, Ionicon Analytik GmbH, Innsbruck, Austria) coupled to a “CHemical Analysis of aeRosols ONline” (CHARON) inlet is used to follow online the gas and particle phase chemical composition of the organic fraction. The CHARON inlet has been already described in detail elsewhere (Eichler et al., 2015; Leglise et al., 2019; Müller et al., 2017). Briefly, the sampled air travels through three major sections of the CHARON inlet 1) a gas phase denuder of activated charcoal that strips off gaseous organics; 2) an aerodynamic lens system (ADL) that collimates the subsampled flow and subsequently enriches the particle concentration and 3) a

thermodesorption unit (TD) heated at  $T = 150^{\circ}\text{C}$  that vaporizes the particles prior their introduction into the drift tube. The PTR-ToF-MS is used with hydronium ions ( $\text{H}_3\text{O}^+$ ) to ionize organic analytes, and operates at a drift tube pressure of 2.6 mbar, a temperature of  $120^{\circ}\text{C}$ , and a voltage of 230 V. This results in an  $E/N = 68\text{ Td}$  ( $E$  = electric field,  $N$  = Number density of the gas molecules in the drift,  $1\text{ Td} = 10^{-17}\text{ V cm}^2\text{ molecule}^{-1}$ ). It represents a relatively low  $E/N$  compared to classical 120-140 Td reported in most of PTR-ToF-MS studies, as the aim is to minimize the potential fragmentation of parent ions and facilitate the molecular characterization of SOA. The potential higher dependence of the sensitivity to relative humidity variations at such low  $E/N$  can be neglected as all experiments were conducted at fixed relative humidity (Pang, 2015; Tani et al., 2003). Instrument background has been daily performed using pure  $\text{N}_2$ , while sensitivity and particle enrichment factors (EF) are controlled at the end of the experiments. A blank was conducted prior to each SOA experiment, using the same conditions ( $\text{H}_2\text{O}_2$  flow,  $\text{NO}_x$  concentration, temperature, humidity, etc.), in the absence of the VOC precursor (either m-xylene or naphthalene). The products formed during these daily blanks were quantified and subtracted to the signal of the following experiment. EF is determined by CHARON calibration using a vanillic acid solution based on the method recommended by Eichler et al. (2015). Instrument sensitivity is evaluated by calculating the transmission curve using a cylinder containing 14 gas standards (benzene, toluene, ethylbenzene, o-, m-, p-xylene, styrene, 1,2,4-trimethyl-, 1,3,5-trimethylbenzene, chloro-, 1,2-dichloro-, 1,3-dichloro-, 1,4-dichloro-, trichloro benzene, each at  $100 \pm 10\%$ , ppb in  $\text{N}_2$ , RESTEK) covering a mass range up to  $m/z$  181.

IDA (Icon Data Analyzer 2.1.1.4) is used to process data recorded by the PTR-ToF-MS such as mass calibration, peak shape definition, peak identification and integration, rate constant calculation and VOC quantification (Müller et al., 2013). The peaks at  $m/z$  21.022 ( $\text{H}_3^{18}\text{O}^+$ ),  $m/z$  330.847 corresponding to diiodobenzene ( $\text{C}_6\text{H}_5\text{I}_2^+$ ) and its fragment at  $m/z$  203.943 ( $\text{C}_6\text{H}_5\text{I}^+$ ) are used to recalibrate the mass scale (PerMaSCal<sup>®</sup> internal standard, 1,3-diiodobenzene,  $T = 55^{\circ}\text{C}$ ). Molecular formulas composed of C, H, O, N atoms are assigned based on exact mass position, chemical rules (valence of atoms, for example) and isotopic patterns. The molecular formula is used to calculate dipole moments and polarizability as introduced by (Bosque and Sales, 2002) and (Sekimoto et al., 2017), which allows to calculate k-rate constants based on the ion-molecule collision theories (Gioumoussis and Stevenson, 1958; Langevin, 1950; Su and Chesnavich, 1982). Concentrations are then estimated based on the rate constant between a proton and each VOC, the experimental transmission of each compound, and the primary ion intensity.

RStudio (RStudio 2023.06.0 Build 421) is used to perform a non-targeted approach for compound selection based on stability periods before and after oxidation. Time intervals are defined for each experiment corresponding to blank (photooxidation without VOC precursor, and HEPA (High-Efficiency Particulate Air) filter for particle phase), reactant (injection of products before the light has been turned on) and products (stable photooxidation), for both gas and particle phases. A Welch  $t$ -test is then used to statistically identify ions that are more concentrated than those in the blank. Then, among the selected ions, the products are defined as compounds more concentrated during oxidation period, and the invert for reactants, based also on Welch  $t$ -test.

193 A high-resolution Time-of-Flight Aerosol Mass Spectrometer (HR-ToF-AMS, ToFwerk AG, Aerodyne Inc. USA) is employed  
 194 for the quantification of the organic and inorganic aerosol fractions (Canagaratna et al., 2007; DeCarlo et al., 2006), with the  
 195 data recorded by the AMS being analyzed using the software SQUIRREL (ToF-AMS Analysis Toolkit 1.65C). The ionization  
 196 efficiency (IE) with respect to nitrate anions was  $4.58 \times 10^{-8}$ . It was calculated using nebulized 300 nm mobility diameter  
 197 ammonium nitrate particles (BFSP software). The relative IE (RIE) of ammonium was 3.45 based on the mass spectrum of  
 198 ammonium nitrate data from IE calibrations. The RIE of sulphate was determined by comparing the theoretical and the  
 199 measured concentration of a solution of ammonium nitrate and ammonium sulphate and was determined to be 1.75. For the  
 200 organic fraction, the default value of 1.4 was used. The AMS data were corrected by collection efficiency (CE) calculated by  
 201 comparison to the SMPS (Scanning Mobility Particle Sizer, TSI Classifier model 3082, DMA, TSI CPC 3776) volume using  
 202 densities of  $1.7 \text{ g cm}^{-3}$  for ammonium sulphate and  $1.4 \text{ g cm}^{-3}$  for organics. The CE values varied from 0.3 for pure ammonium  
 203 sulphate particles to 0.7 after SOA formation.  
 204 The SMPS is used for measuring particle size distribution and number concentration. Additional instruments account for a  
 205 chemiluminescent  $\text{NO}_x$  analyzer (Envitec, model API200E), a pressure sensor (ATM.ECO, STS) and a temperature-humidity  
 206 probe (HMP9 Vaisala) positioned close the aerosol flow tube outlet, the last two instruments being connected to a data logger  
 207 (FieldLogger, NOVUS).



208 **Figure 1. Schematic of the experimental setup. MFC: mass flow controller.**  
 209  
 210

## 2.3 SOA yield, partitioning, and volatility distribution

The expression to describe the fractional aerosol yield ( $Y$ ) was established by (Odum et al., 1996, 1997) and is described in Eq. (1), where  $\Delta M_0$  is the amount of the total organic aerosol formed (in  $\mu\text{g m}^{-3}$ ), and  $\Delta VOC$  is the amount of VOC precursor reacted (in  $\mu\text{g m}^{-3}$ ) measured as difference between inlet and outlet OFR.

$$Y = \frac{\Delta M_0}{\Delta VOC} \quad (1)$$

The distribution of oxidation products between the gas and particle phases can be explained by the partitioning theory of (Yamasaki et al., 1982):

$$K_{p,i} = \frac{C_{p,i}}{C_{g,i} \times TSP} \quad (2)$$

where  $K_{p,i}$  is the experimental partitioning coefficient (in  $\text{m}^3 \mu\text{g}^{-1}$ ) of a species  $i$ ,  $C_{p,i}$  and  $C_{g,i}$  are concentrations (in  $\mu\text{g m}^{-3}$ ) of the species  $i$  in particle and gas phases, respectively, as measured by the CHARON-PTR-ToF-MS, and TSP is the total suspended particulate matter of the aerosol (in  $\mu\text{g m}^{-3}$ ) as measured by SMPS. Larger  $K_p$  values indicate a preference for a compound to partition in the particle phase. Subsequently, the volatility of the species can be defined by  $\log_{10} C_i^*$ , where  $C_i^*$  is known as the effective saturation concentration (in  $\mu\text{g m}^{-3}$ ) of a species  $i$  which describes the gas-particle partitioning behavior of organic compounds and represents the gas-phase concentration of a compound at which it will partition equally between the gas and particle phases under given atmospheric conditions, and is calculated as the reciprocal of  $K_{p,i}$  (Donahue et al., 2006, 2011):

$$C_i^* = \frac{1}{K_{p,i}} \quad (3)$$

The estimated values have been calculated using the Volcalc model based on molecular properties such as molecular weight, numbers of atoms and functional groups (Meredith et al., 2023; Riemer, 2023). The model is based on SIMPOL.1, a group contribution method (Pankow and Asher, 2008), which implements a structure-activity relationship method to calculate the subcooled pure liquid vapor pressure by summing the contributions of the subcooled liquid vapor pressures of individual chemical functional groups:

$$\log_{10} P_{L,i}^\circ(T) = \sum_k v_{k,i} b_k(T) \quad (4)$$

where  $P_{L,i}^\circ(T)$  is the liquid vapor pressure (atm),  $v_{k,i}$  is the number of groups of type  $k$  in  $i$ , the index  $k$  can take on the entire numbers (1,2,3, etc.), and  $b_k(T)$  is the group contribution term for group  $k$ . No second-order interaction terms are included to account for neighboring functional groups, which means that the model does not consider the potential interactions or effects that adjacent functional groups might have on each other but only sums the contributions of individual functional groups independently.

The saturation concentration ( $C_{i,T}^\circ$ ) of the major identified species  $i$  is calculated at  $T = 280$  and  $295$  K as follows:

$$C_{i,T}^\circ = C_{i,293}^\circ \times \frac{293}{T} \times \exp \exp \left( \left( \frac{-\Delta H}{R} \right) \times \left( \frac{1}{T} - \frac{1}{293} \right) \right) \quad (5)$$



241 where  $C_{i,293}^{\circ}$  is the saturation concentration calculated by Volcalc at  $T = 293$  K,  $\Delta H_i$  is the enthalpy of vaporization of species  
 242  $i$  (computationally predicted values from ChemSpider), and  $R$  is the ideal gas constant =  $8.314 \text{ J mol}^{-1} \text{ K}^{-1}$ .  
 243 The saturation concentration ( $C_i^{\circ}$ ) and the effective saturation concentration ( $C_i^*$ ) are related through the activity coefficient  
 244 ( $\gamma_i$ ) that captures the non-ideal interactions of the compound with the aerosol mixture. Its value generally lies between 0.3  
 245 (readily partitions to particle phase) and 3 (readily partitions to gas phase) for ambient atmospheric aerosol (Donahue et al.,  
 246 2011; Liu et al., 2021). As in previous studies, such as Isaacman-VanWertz et al. (2016) and Nie et al. (2022), in this work we  
 247 assume a  $\gamma$  value of 1.

248 **3 Results and discussion**

249 **3.1 SOA yield formation**


250 Table 1 summarizes experimental conditions, consumed reactants ( $\Delta VOC$ ) and formed organic aerosol mass ( $\Delta M_0$ ) used to  
 251 calculate the SOA yield ( $Y$ ) for each experiment (Eq. (1)). Figure 2 presents SOA yields for (a) m-xylene and (b) naphthalene,  
 252 comparing them with results from selected previous studies (Chan et al., 2009; Chen et al., 2018; Ng et al., 2007; Song et al.,  
 253 2005). Filled markers indicate high  $\text{NO}_x$  conditions ( $\text{VOC}/\text{NO}_x < 8$ ), empty markers refer to low  $\text{NO}_x$  conditions ( $\text{VOC}/\text{NO}_x$   
 254  $> 8$ ) (Dodge, 1977; NARSTO. and Electric Power Research Institute., 2000). The red square and blue triangle markers refer  
 255 to this study at 295 K and 280 K, respectively, while all other experiments are conducted at room temperatures (between 295  
 256 K and 300 K).

257 **Table 1. List of conducted laboratory experiments and associated conditions, such as OFR temperature, RH, VOC/ $\text{NO}_x$  ratio, seeds**  
 258 **mass and SOA yield.**

	T	RH	VOC	NO <sub>x</sub>	VOC/NO <sub>x</sub>	Seeds	$[OH] \times 10^7$	$\Delta VOC$	$\Delta M_0$	$Y$	
	K	%	ppbV	ppb	ppbC ppb <sup>-1</sup>	μg m <sup>-3</sup>	molecules cm <sup>-3</sup>	μg m <sup>-3</sup>	%	μg m <sup>-3</sup>	%
m-xylene	280 ± 1.5	75 ± 5	74 ± 0.65	235	2.5 ± 0.1	51 ± 1.6	3.4 ± 0.5	114 ± 3.3	34	26.4 ± 1.2	23.1 ± 1.2
	280 ± 1.5	50 ± 5	69 ± 0.70	40	13.9 ± 0.5	35 ± 0.5	2.2 ± 0.4	95 ± 3.3	30	26.1 ± 1.9	27.5 ± 2.1
	295 ± 2	60 ± 7	73 ± 0.66	221	2.6 ± 0.1	30 ± 0.8	5.4 ± 0.8	155 ± 3.2	49	12.6 ± 1.4	8.1 ± 1.0
	295 ± 2	55 ± 7	83 ± 0.74	40	16.6 ± 0.6	34 ± 0.7	4.7 ± 0.7	153 ± 3.2	42	21.1 ± 1.4	13.8 ± 1.0
naphthalene	280 ± 1.5	40 ± 3	57 ± 0.35	340	1.7 ± 0.1	46 ± 0.6	2.9 ± 0.2	92 ± 2.3	29	23.3 ± 1.8	25.3 ± 2.0
	280 ± 1.5	35 ± 3	53 ± 0.41	62	9.3 ± 0.4	36 ± 1.0	3.2 ± 0.3	79 ± 2.3	27	33.6 ± 6.9	42.7 ± 8.8
	295 ± 2	40 ± 5	53 ± 0.47	340	1.6 ± 0.1	55 ± 1.5	3.5 ± 0.3	84 ± 2.3	28	12.6 ± 1.7	14.9 ± 2.1

259

260 For m-xylene (Fig. 2a), the SOA yield at 295 K under high NO<sub>x</sub> conditions is approximately 8 % (filled red square), in line  
 261 with values reported by Ng et al. (2007) and Chen et al. (2018). Chen et al. (2018) reported yields with higher VOC/NO<sub>x</sub> ratios,  
 262 slightly lower precursor concentrations (44-59 ppb), and without using seeds. The absence of seed particles reduces the  
 263 available surface area for condensation of oxidized products (Lambe et al., 2015). Ng et al. (2007) conducted experiments with  
 264 AS seeds as in our study, but using nitrous acid (HONO) as OH radical precursor, with varying initial xylene concentrations  
 265 (from 42 to 171 ppb) which may explain SOA yield variability (from 3 to 8 %). The effect of higher initial levels of the VOC  
 266 precursor has been shown to reduce aerosol formation (and amount of reacted precursor) and the formation of oxidation  
 267 products, especially those with m/z > 110 da (Chen et al., 2019). It may be due to the competition between IVOC reactions  
 268 with OH that produces LVOC/ELVOC, and precursor oxidation. In our case, the initial m-xylene concentration was kept  
 269 similar between experiments to isolate the effect of temperature and NO<sub>x</sub> on SOA yield and chemistry. Both studies by Ng  
 270 (2007) and Chen (2018) were conducted in dry conditions. Li et al. (2022) reported SOA yields increasing from 6.3 to 14 %  
 271 when humidity increased from 10 % to 70 %. The order of magnitude of SOA yields in this study is close to what we observed,  
 272 the variability of yields being associated mostly to NO<sub>x</sub> or NH<sub>3</sub> initial concentrations. Under low NO<sub>x</sub>, a SOA yield of 14 %  
 273 (empty red square) is observed, lying on the lower level (12-30 %) of reported values by Song et al. (2007), that used lower  
 274 levels of xylene (39-52 ppb).



275

276 **Figure 2. SOA yields at 295 K and 280 K as function of organic aerosol mass formed for (a) m-xylene and (b) naphthalene in**  
 277 **comparison with previous studies. Filled markers correspond to high NO<sub>x</sub> conditions, open markers to low NO<sub>x</sub>.**

278 The effect of temperature on SOA yield is also depicted in Fig. 2. When decreasing from 295 K to 280 K, SOA yield increases  
 279 from 8 to 23 % under high NO<sub>x</sub> conditions and from 14 to 28 % under low NO<sub>x</sub> conditions. Only a few studies investigated

the role of low temperature on SOA formation from monoaromatic precursors. Regarding xylene, only Takekawa et al. (2003) reported that SOA yield is enhanced by a factor around 2 (from 6 % to 13 %, on average) when temperature decreases from 303 K to 283 K at a VOC/NO<sub>x</sub> around 10 (like low NO<sub>x</sub> conditions in this study). This is in good agreement with our results, where a decrease of 15 K doubles the SOA yield under low NO<sub>x</sub> conditions (Fig. 2). Lannuque et al., (2023) found a similar temperature dependence of toluene-SOA and highlighted that the effect was greater for low concentrations of the precursors.

For naphthalene (Fig. 2b), SOA yield at 295 K with high NO<sub>x</sub> conditions ranges from 15 to 18 %, in agreement with Chen et al. (2016) and Chan et al. (2009) that reported values from 3 % to 47 %. Chen et al. (2016) observed the lowest SOA yield under high NO<sub>x</sub>, attributed to the lowest amount of OH generated and highest NO injection. Even though Chan et al. (2009) used AS seeds, the initial amount of naphthalene was less than half of that used in our experiments and the humidity was below 10 %, highlighting again the important effect of experimental conditions on the prediction of SOA yields. Lower naphthalene SOA yields are observed under high NO<sub>x</sub> conditions (filled vs empty square or triangle, Fig. 2). Under low NO<sub>x</sub> conditions, a yield of 18 % is observed in quite good agreement with Chen et al. (2016), who reported yields varying from 4 % to 29 % as a function of the VOC/NO<sub>x</sub> ratio, with lowest yields for highest NO<sub>x</sub> regimes. A consistent increase in SOA yield is observed when switching from high to low NO<sub>x</sub> regime at 280 K, from 25 to 43 % at 280 K, while at 295 K the variation is limited to a few percent increase. The presence of NO<sub>x</sub> in the system may promote a competition for the termination reaction of RO<sub>2</sub> radicals between peroxy radicals (RO<sub>2</sub> and HO<sub>2</sub>) and NO<sub>x</sub> (NO<sub>2</sub> and NO), leading to less oxidized products of higher volatility (Henze et al., 2008; Kroll and Seinfeld, 2008).


The effect of temperature on naphthalene SOA is reported here for the first time. Reducing the temperature from 295 K to 280 K induces an increase from 14.9 % to 25.3 % for high NO<sub>x</sub> conditions, and from 18.3 % to 42.7 % for low NO<sub>x</sub> conditions. SOA yields from naphthalene are in general higher compared to m-xylene, which is expected considering larger carbon skeleton of naphthalene (Aumont et al., 2013; La et al., 2016), and are in good agreement with previous studies (Chan et al., 2009; Chen et al., 2018). However, the effect of temperature on naphthalene SOA yield is slightly lower compared to the m-xylene SOA system. This may be the result of the larger carbon skeleton of naphthalene, leading to products of lower volatility, that have already partitioned more in the particle phase at 295 K.

## **3.2 Chemical composition of oxidation products in gas and particle phases**

### **3.2.1 m-xylene**

Figure 3 presents the chemical distribution of gas (Figs. 3a,c) and particles (Figs. 3b,d) phase products as a function of the carbon atoms following m-xylene oxidation by OH radicals. The mass fraction (in  $\mu\text{g m}^{-3}$ ) is further sub-classified based on the number of oxygen (Figs. 3a,b) and nitrogen atoms (Figs. 3c,d; according to color scale), and the molecular weight distribution was divided into three groups:  $m/z$  below 100,  $m/z$  101-150, and  $m/z$  above 150. The experiment has been carried out under high NO<sub>x</sub> at 280 K. The total m-xylene carbon balance varies within 26-48 % for the detected gas phase products depending on the oxidation conditions and considering that CO, CO<sub>2</sub> and glyoxal are not measured. The latter compound has

312 strong interferences with the high signal of acetone. The SOA formed corresponds to 2.5 % of the total carbon balance at  
 313 295 K and 7 % at 280 K under high NO<sub>x</sub> conditions.



314  
 315 **Figure 3. m-xylene mass products fraction (y-axis) distribution based on the number of carbon atoms (x-axis) for a high NO<sub>x</sub>**  
 316 **experiment at 280 K, colored by the number of (a,b) oxygen and (c,d) nitrogen atoms. Detected compounds are in the (a,c) gas phase**  
 317 **and (b,d) particle phase. Pie charts correspond to the molecular weight contribution to the overall mass.**

318 The overall product distribution at 295 K is similar to that at 280 K and can be found in the SI (Figs. S1a and S1b). At both  
 319 temperatures, the gas phase is mainly composed of C<sub>3</sub>, C<sub>5</sub> and C<sub>8</sub> products with 1 to 3 O atoms for most of them, while the

particle phase is dominated by C<sub>6</sub>, C<sub>7</sub> and C<sub>8</sub> compounds, containing generally 2 to 5 O atoms. In the gas phase, C<sub>8</sub> compounds are ring retaining compounds related to first-generation products, while C<sub>3</sub> and C<sub>5</sub> products arise from a more advanced chemistry. A larger fraction of C<sub>3</sub> products is present in the gas phase at 295 K when compared to 280 K (Figs. 3 and S1a). This could be due to the higher OH concentration for the experiment at 295 K (Table 1), leading to a higher consumption of C<sub>6</sub> and C<sub>8</sub> oxidation products due to their faster reactivity. This is supported by the time evolution of some compounds such as C<sub>6</sub>H<sub>8</sub>O at *m/z* 97.06, C<sub>6</sub>H<sub>6</sub>O<sub>2</sub> at *m/z* 111.04, C<sub>6</sub>H<sub>8</sub>O<sub>4</sub> at *m/z* 145.05, C<sub>8</sub>H<sub>10</sub>O<sub>5</sub> at *m/z* 187.06 (Fig. S3). This hypothesis is also backed up by previous studies reporting the reaction of oxidation products (Cappa and Wilson, 2012; Isaacman-VanWertz et al., 2018; Jin et al., 2021). While for experiments with lower OH concentration, as at 280 K, the gas phase products (Fig. 3a) are characterized by a higher fraction of C<sub>8</sub> (26%) and C<sub>5</sub> (27%) compounds.

The condensed phase is enriched in compounds with higher molecular weight when compared to the gas phase, along with the presence of four or more oxygen atoms which highly reduces their vapor pressure (Cappa and Wilson, 2012). Particle phase products generally exhibit an increasing trend from C<sub>1</sub> to C<sub>8</sub> (Fig. 3b), similar to a recent study on toluene SOA (Lannuque et al., 2023).

The presence of NO<sub>x</sub> may affect product distribution, directly through the formation of organic nitrates, or indirectly by playing a role in shaping oxidation chemistry depending on experimental conditions. The comparison of the chemical composition of gas phase products in low NO<sub>x</sub> and high NO<sub>x</sub> at 295 K showed a general similar distribution (Fig. S1 and S2). The only noticeable change is the stronger contribution of C<sub>3</sub> and C<sub>5</sub> compounds in the low NO<sub>x</sub> experiment resulting from the slight increase in OH radical production as described above (Table 1), rather than a real NO<sub>x</sub> effect. This is supported by the same product distribution in particle phase, when comparing high and low NO<sub>x</sub>, with only a small increase in organic nitrates at high NO<sub>x</sub>. The very close distribution between high and low NO<sub>x</sub> means that the chemistry was not strongly dependent on the NO<sub>x</sub> regime in our conditions, and that NO<sub>x</sub> did not significantly affect the oxidant levels in the OFR.

Table 2 lists identified ions from the gas and particle phase products for two experiments at high NO<sub>x</sub> at 295 K and 280 K. The columns present the measured *m/z* signals, the attributed chemical formula and a tentative chemical assignment based on results from previous studies on *m*-xylene photooxidation (Atkinson et al., 1991; Forstner et al., 1997; Huang et al., 2008; Jang and Kamens, 2001; Li et al., 2018, 2022; Zhang et al., 2019a, b; Zhao et al., 2005) and on the CHARON assignment procedure recommended by Gkatzelis et al. (2018) where average saturation mass concentration log<sub>10</sub>(C<sub>*i*</sub><sup>\*</sup>) are used to discriminate between parent and fragment ions. The compounds are sorted in decreasing numbers of carbon atoms. For each temperature, two columns are presented: the first indicates the contribution of each compound to total gas phase organic products and the second the contribution to SOA.

**Table 2. Ions and their corresponding formulas of the major *m*-xylene products detected during photooxidation experiments under high NO<sub>x</sub>. Reaction products are given as a fraction of the gas phase products (in % of μg m<sup>-3</sup>) and as a fraction of the SOA products (in % of μg m<sup>-3</sup>).**

Carbon number	Measured <i>m/z</i> and ion sum formula	Tentative assignment	<i>T</i> = 298 K		<i>T</i> = 280 K	
			Gaseous	SOA	Gaseous	SOA

			Products (%)	Products (%)	Products (%)	products (%)
8	121.06 (C <sub>8</sub> H <sub>8</sub> O)H <sup>+</sup>	m-tolualdehyde	13.9	0.6	16.7	0.5
8	155.07 (C <sub>8</sub> H <sub>10</sub> O <sub>3</sub> )H <sup>+</sup>	dimethyl-epoxy-oxo-hexenal/ trihydroxy dimethyl benzene	n.d.	3.8	n.d.	3.3
8	137.06 (C <sub>8</sub> H <sub>8</sub> O <sub>2</sub> )H <sup>+</sup>	toluic acid and possible contribution frag. 155.07	n.d.	8.5	n.d.	5.7
8	171.07 (C <sub>8</sub> H <sub>10</sub> O <sub>4</sub> )H <sup>+</sup>	dihydroxy-dimethyl-cyclohexenedione / dimethyl-hexadienedioic acid	0.3	4.6	0.3	5.6
8	153.06 (C <sub>8</sub> H <sub>8</sub> O <sub>3</sub> )H <sup>+</sup>	hydroxy dimethyl quinone	1.9	2.8	2.4	2.9
8	187.06 (C <sub>8</sub> H <sub>10</sub> O <sub>5</sub> )H <sup>+</sup>	hydroxy-cyclohexene-dicarboxylic acid oxo-cyclohexanedicarboxylic acid	0.1	2.6	0.1	2.5
8	169.05 (C <sub>8</sub> H <sub>8</sub> O <sub>4</sub> )H <sup>+</sup>	dihydroxy-methylbenzoic acid	n.d.	2.0	n.d.	1.6
8	185.05 (C <sub>8</sub> H <sub>8</sub> O <sub>5</sub> )H <sup>+</sup>	trihydroxy-(hydroxymethyl)benzaldehyde	n.d.	1.0	n.d.	0.7
8	168.06 (C <sub>8</sub> H <sub>9</sub> NO <sub>3</sub> )H <sup>+</sup>	<i>dimethyl nitrophenol</i>	1.9	0.4	3.6	0.4
8	152.07 (C <sub>8</sub> H <sub>9</sub> NO <sub>2</sub> )H <sup>+</sup>	<i>nitro-xylene</i>	0.4	0.1	1.8	0.1
7	125.06 (C <sub>7</sub> H <sub>8</sub> O <sub>2</sub> )H <sup>+</sup>	dimethyl-pyranone/ methyl-hexadienedial	n.d.	3.6	1.6	5.0
7	141.05 (C <sub>7</sub> H <sub>8</sub> O <sub>3</sub> )H <sup>+</sup>	methyl-oxo-hexadienoic acid /heptenetrione	0.7	2.8	0.7	4.0
7	139.04 (C <sub>7</sub> H <sub>6</sub> O <sub>3</sub> )H <sup>+</sup>	hydroxy benzoic acid/ hydroxy methyl benzoquinone	n.d.	3.1	n.d.	2.4
7	157.05 (C <sub>7</sub> H <sub>8</sub> O <sub>4</sub> )H <sup>+</sup>	hydroxy-dioxo-heptenal / epoxymethylhexenedial	n.d.	2.0	n.d.	1.8
7	109.06 (C <sub>7</sub> H <sub>8</sub> O)H <sup>+</sup>	cresols/ benzyl alcohol	n.d.	1.4	n.d.	1.5
7	173.06 (C <sub>7</sub> H <sub>8</sub> O <sub>5</sub> )H <sup>+</sup>	hydroxy-dioxo-heptenoic acid	n.d.	1.8	n.d.	1.2
7	138.06 (C <sub>7</sub> H <sub>7</sub> NO <sub>2</sub> )H <sup>+</sup>	<i>nitrotoluene</i>	1.1	0.4	1.2	0.3
7	154.05 (C <sub>7</sub> H <sub>7</sub> NO <sub>3</sub> )H <sup>+</sup>	<i>nitrocresol</i>	0.7	0.3	0.6	0.3
6	127.04 (C <sub>6</sub> H <sub>6</sub> O <sub>3</sub> )H <sup>+</sup>	hydroxymethyl furfural/ hydroxyquinol/dimethylfuranione	1.0	2.4	1.4	3.3
6	113.06 (C <sub>6</sub> H <sub>8</sub> O <sub>2</sub> )H <sup>+</sup>	methyl-oxo-pentalenal / dimethylfuranone	n.d.	1.7	0.2	2.2
6	129.06 (C <sub>6</sub> H <sub>8</sub> O <sub>3</sub> )H <sup>+</sup>	hydroxy-oxo-hexenal methyl-oxo-pentenoic acid	0.4	1.6	0.4	2.7
6	111.04 (C <sub>6</sub> H <sub>6</sub> O <sub>2</sub> )H <sup>+</sup>	methylfuraldehyde / benzenediols possible frag. of 129.06	n.d.	3.6	2.7	6.2
6	143.03 (C <sub>6</sub> H <sub>6</sub> O <sub>4</sub> )H <sup>+</sup>	dioxo-hexenoic acid/ methyl-dioxo- pentenoic acid tetrahydroxybenzene	n.d.	1.4	0.2	1.8
6	159.04 (C <sub>6</sub> H <sub>6</sub> O <sub>5</sub> )H <sup>+</sup>	oxo-hexenedioic acid	n.d.	1.2	n.d.	1.2
6	145.05 (C <sub>6</sub> H <sub>8</sub> O <sub>4</sub> )H <sup>+</sup>	hydroxy-dioxo hexanal	n.d.	1.2	0.1	1.1
6	115.07 (C <sub>6</sub> H <sub>10</sub> O <sub>2</sub> )H <sup>+</sup>	cyclopentylcarboxylic acid	n.d.	0.4	0.4	1.1
6	95.03 (C <sub>6</sub> H <sub>6</sub> O)H <sup>+</sup>	phenol	n.d.	1.4	n.d.	1.2
5	113.02 (C <sub>5</sub> H <sub>4</sub> O <sub>3</sub> )H <sup>+</sup>	methyl-furanione	16.7	1.3	19.3	2.1
5	131.04 (C <sub>5</sub> H <sub>6</sub> O <sub>4</sub> )H <sup>+</sup>	methy-hydroxy-oxo-butandial	2.1	1.4	0.9	0.9
5	117.05 (C <sub>5</sub> H <sub>8</sub> O <sub>3</sub> )H <sup>+</sup>	oxo-pentanoic acid + frag at 99.04	1.2	0.7	0.2	0.4
	99.04 (C <sub>5</sub> H <sub>6</sub> O <sub>2</sub> )H <sup>+</sup>	oxo-pentalenal / methyl-butendial	6.7	3.5	2.8	3.9
5	115.03 (C <sub>5</sub> H <sub>6</sub> O <sub>3</sub> )H <sup>+</sup>	oxo-pentenoic acid	1.4	2.4	0.7	1.8
5	97.03 (C <sub>5</sub> H <sub>4</sub> O <sub>2</sub> )H <sup>+</sup>	furaldehyde	1.1	0.7	2.2	1.7

5	101.06 (C <sub>5</sub> H <sub>8</sub> O <sub>2</sub> )H <sup>+</sup> 83.05 (C <sub>5</sub> H <sub>6</sub> O)H <sup>+</sup>	oxo-pentanal and isomers + frag at 83.05 methyldfuran	n.d 0.6	0.4 1.0	n.d 0.3	0.4 1.3
4	71.05 (C <sub>4</sub> H <sub>6</sub> O)H <sup>+</sup>	dihydrofuran / MACR / MVK	1.1	0.3	0.4	0.3
4	87.04 (C <sub>4</sub> H <sub>6</sub> O <sub>2</sub> )H <sup>+</sup>	butanedial / crotonic acid	n.d.	2.0	n.d.	2.2
4	103.04 (C <sub>4</sub> H <sub>6</sub> O <sub>3</sub> )H <sup>+</sup>	hydroxy-oxo-butanal	0.4	1.5	0.1	1.3
4	85.03 (C <sub>4</sub> H <sub>4</sub> O <sub>2</sub> )H <sup>+</sup>	butenedial	n.d.	1.3	n.d.	1.2
3	73.03 (C <sub>3</sub> H <sub>4</sub> O <sub>2</sub> )H <sup>+</sup>	methylglyoxal	20.9	3.2	9.5	2.7
3	75.04 (C <sub>3</sub> H <sub>6</sub> O <sub>2</sub> )H <sup>+</sup>	propanoic acid	14.3	1.7	13.3	1.6
3	89.02 (C <sub>3</sub> H <sub>4</sub> O <sub>3</sub> )H <sup>+</sup>	pyruvic acid / hydroxy-propanedial	1.5	0.6	n.d.	0.6
2	77.03 (C <sub>2</sub> H <sub>4</sub> O <sub>3</sub> )H <sup>+</sup>	PAN fragment	3.7	0.3	3.6	0.5

\*n.d. = not detected

The C<sub>8</sub> compounds are dominating both gas and particle phases accounting for 27 % of the total products. The most abundant C<sub>8</sub> gas phase product is m-tolualdehyde (C<sub>8</sub>H<sub>8</sub>O detected at *m/z* 121.06), comprising alone 17 % of the reaction products. It is a first generation ring-retaining aromatic compound previously identified in many studies (Atkinson et al., 1991; Forstner et al., 1997; Huang et al., 2008; Srivastava et al., 2023; Zhang et al., 2019b; Zhao et al., 2005). Toluic acid is a second generation ring retaining compound formed by the additional oxidation of m-tolualdehyde by OH radical (Forstner et al., 1997; Srivastava et al., 2022, 2023), and it is a major C<sub>8</sub> particle phase product (C<sub>8</sub>H<sub>8</sub>O<sub>2</sub>, detected at *m/z* 137.06) contributing to 5 % of the SOA mass. Another major SOA component is found at *m/z* 171.07 (C<sub>8</sub>H<sub>10</sub>O<sub>4</sub>) tentatively assigned to dimethyl-hexadienedioic acid, a ring opening product possibly formed by OH-addition to the benzene ring followed by a ring cleavage, or to dihydroxy-dimethyl-cyclohexenedione, a ring retaining compound formed by successive OH radical reactions on the benzene ring. A second ring opening compound is detected at *m/z* 155.07 (C<sub>8</sub>H<sub>10</sub>O<sub>3</sub>) previously identified as dimethyl-epoxy-oxo-hexenal (Zhao et al., 2005). Other C<sub>8</sub> compounds are C<sub>8</sub>H<sub>8</sub>O<sub>3-5</sub> (at *m/z* 153.06, *m/z* 169.05 and *m/z* 185.05) and C<sub>8</sub>H<sub>10</sub>O<sub>5</sub> at *m/z* 187.06 (Table 2).

The contribution of C<sub>7</sub> compounds to gas and particle phases accounts for 6 % and 19 %, respectively. Among the C<sub>7</sub> prominent products in the particle phase we identify C<sub>7</sub>H<sub>8</sub>O<sub>2</sub> at *m/z* 125.06 tentatively assigned to dimethyl-pyranone (Forstner et al., 1997) or methyl-hexadienedial resulting from a ring opening of the phenoxy radical intermediate formed via OH-addition to the ring (Jang and Kamens, 2001). Further oxidation of methyl-hexadienedial will result in methyl-oxo-hexadienoic acid C<sub>7</sub>H<sub>8</sub>O<sub>3</sub> at *m/z* 141.05 (Jang and Kamens, 2001). Another C<sub>7</sub> compound is the C<sub>7</sub>H<sub>6</sub>O<sub>3</sub> at *m/z* 139.04 tentatively assigned to methyl cyclohexene tricarbonyls or hydroxy methyl benzoquinone, both being ring-retaining products. They can be formed upon OH-addition to first- or second-generation products and release of one methyl group (Jang and Kamens, 2001). Other C<sub>7</sub> compounds important in the particle phase include C<sub>7</sub>H<sub>8</sub>O (detected at *m/z* 109.06), C<sub>7</sub>H<sub>8</sub>O<sub>4</sub> (detected at *m/z* 157.05), and C<sub>7</sub>H<sub>8</sub>O<sub>5</sub> (detected at *m/z* 173.04). Some of the C<sub>7</sub>-C<sub>8</sub> products are nitrogen containing-compounds: nitro-m-xylene (C<sub>8</sub>H<sub>9</sub>NO<sub>2</sub>, at *m/z* 152.07), nitrotoluene (C<sub>7</sub>H<sub>7</sub>NO<sub>2</sub> at *m/z* 138.06), dimethyl nitrophenol (C<sub>8</sub>H<sub>9</sub>NO<sub>3</sub> at *m/z* 168.06) and nitrocresol (C<sub>7</sub>H<sub>7</sub>NO<sub>3</sub> at *m/z* 154.05). The latter is the result of NO<sub>2</sub> reaction with the benzyl peroxy radical in the H-abstraction route (Li et al., 2018; Srivastava et al., 2023). Aliphatic nitrogen-containing compounds are easily fragmented in the PTR-MS and thus

are hardly detected. Nevertheless, a known peroxyacetyl nitrate (PAN) fragment  $C_2H_4O_3$  detected at  $m/z$  77.03, also previously assigned as an unspecific fragment from nitro-group containing compounds has been detected (Müller et al., 2012). In total, the sum of all nitrogen-containing products (including PAN fragment) accounted for 9-14 % of the gaseous phase mass loading and 6 % of the particulate phase under high- $NO_x$  conditions, which is close to what was seen in toluene photooxidation experiments by Lannuque et al. (2023). Under low  $NO_x$  conditions, organic nitrates (including PAN fragment) accounted for 4-12 % in the gas phase and 3-4 % in the particle phase out of total products, indicating limited  $NO_x$  influence on the formation of nitrogen-containing compounds (Figs. S1c,d; S2c,d). These results suggest that in our system, the shift from low to high  $NO_x$  did not substantially alter the oxidation pathways or enhance  $NO_x$ -dependent products formation.

The  $C_6$  compounds accounted for only 6 % of the gas phase and for 19% of the organic aerosol fraction. Top products are ring-retaining furan-derived, such as methyl-furaldehyde or benzenediols ( $C_6H_6O_2$ , detected at  $m/z$  111.04) making up 3 % and 6 % in gas and particle phases respectively, and dimethylfurandione ( $C_6H_6O_3$ , detected at  $m/z$  127.04) with 1.4 % in the gas phase, and 3.3 % in the particle phase, alongside with dimethyl-furanone with 2.2 % in the particle phase ( $C_6H_8O_2$ , detected at  $m/z$  113.06). Furanoid products can be formed through a bridged oxide intermediate on a bicyclic ring whereas furandiones are known to originate from conjugated dicarbonyls via reaction with OH-radicals followed by cyclization (Forstner et al., 1997; Jang and Kamens, 2001). Examples of unsaturated dicarbonyls and tricarbonyls in the condensed phase are methyl-oxo-pentenal ( $C_6H_8O_2$ , detected at  $m/z$  113.06), and hydroxy-oxo-hexenal ( $C_6H_8O_3$ , detected at  $m/z$  129.06), also assigned to methyl-oxo-pentenoic acid (Li et al., 2017, 2018; Zhao et al., 2005). Those multi-carbonyls are a result of the decomposition of bicyclic alkoxy radicals (Huang et al., 2008; Zhao et al., 2005). Other  $C_6$  compounds can be found at  $m/z$  143.03 ( $C_6H_6O_4$ ), 97.06 ( $C_6H_8O$ ), 145.05 ( $C_6H_8O_4$ ), and 115.07 ( $C_6H_{10}O_2$ ) (Table 2).

As shown in Fig. 3 and Fig. S1, the  $C_5$  products represent a major contributor of the gas phase with around 30 % of the total organic gaseous products and 13 % of the particle phase. In agreement with previous studies, most of the compounds are furan derivatives (Forstner et al., 1997; Jang and Kamens, 2001; Li et al., 2018). The  $C_5H_4O_3$  (detected at  $m/z$  113.02) assigned to methylfurandione accounts alone for, 17-19 % of the gas phase products, followed by oxo-pentenal and isomers ( $C_5H_6O_2$ , detected at  $m/z$  99.04) with 3-7 % abundance. Other important products include unsaturated or oxo-aldehydes (such as  $C_5H_6O_2$  at  $m/z$  99.04 and  $C_5H_6O_4$  at  $m/z$  131.04), organic acids such oxo-pentenoic acid ( $C_5H_6O_3$  at  $m/z$  115.03) (Table 2). Lannuque et al. (2023) identified dihydroxy-oxopentanoic acid (DHOPA,  $C_5H_8O_5$ ), a known tracer of toluene and xylene SOA, in particle phase. In our study, DHOPA parent ion was below the detection limit which is highly probable considering its very low formation yield (Srivastava et al., 2023) and the low precursor concentration used compared to Lannuque et al. (2023). We therefore assume that DHOPA did not contribute significantly to the fragment ion at  $m/z$  131.04 ( $C_5H_6O_4$ ) expected upon water loss. In general the high contribution of  $C_5$  compounds can be explained by the m-xylene structure ( $C_8$  aromatic) and the presence of two methyl groups on the benzene ring, which implies the formation of methylated furan derivatives such as methyl-furandione, methyl-furanone and furaldehyde, in addition to the high contribution of  $C_3$  compounds such as methylglyoxal, in agreement with previous studies (Birdsall and Elrod, 2011; Fan and Zhang, 2008; Forstner et al., 1997; Jang and Kamens, 2001; Li et al., 2022; Song et al., 2007; Zhang et al., 2019b).




The C<sub>4</sub> compounds are the least abundant in both phases (below 7 %, on average) and account for shorter functionalized aldehydes, furans and acids formed by minor pathways in the aromatic ring cleavage (Table 2). This is in agreement with the above explained ring opening mechanism, that mainly breaks the carbon skeleton of the xylene in C<sub>5</sub> and C<sub>3</sub> compounds, instead of two C<sub>4</sub> moieties (Forstner et al., 1997; Jenkin et al., 2003; Pan and Wang, 2014).

The C<sub>3</sub> compounds account for 23-37 % of the gas phase and 6 % of the particle phase products. Methylglyoxal (C<sub>3</sub>H<sub>4</sub>O<sub>2</sub>, detected at *m/z* 73.03) accounts for 21% of the gas phase products at 295 K while it is close to 10 % at 280 K. It makes up to 3 % of the SOA fraction and is a major second-generation product resulting from the ring cleavage of bicyclic alkoxy radicals (Fan and Zhang, 2008). Propanoic acid (C<sub>3</sub>H<sub>6</sub>O<sub>2</sub>, detected at *m/z* 75.04) accounts for 8 % of the gaseous phase and only 1 % in the particle phase, as it is a quite volatile compound. This reaction product can result from the oxidation of multifunctional ring opening products (Jang and Kamens, 2001).

### 3.2.2 Naphthalene

In analogy with the m-xylene, Fig. 4 presents the chemical distribution of naphthalene gas phase products (Figs. 4a,c) and particle phase products (Figs. 4b,d) for photooxidation experiments under high NO<sub>x</sub> conditions at 280 K, in addition to pie charts showing the molecular weight distribution. Out of the total naphthalene carbon balance, 30-32 % is accounted for the detected gas phase products under the different oxidation conditions, taking into account that CO, CO<sub>2</sub> and glyoxal are not measured. As for the SOA formed, it corresponds to 8 % at 295 K and 14 % at 280 K of the carbon balance under high NO<sub>x</sub> conditions.



**Figure 4. Naphthalene mass products fraction (y-axis) distribution based on the number of carbon atoms (x-axis) for a high  $\text{NO}_x$  experiment at 280 K, colored by the number of (a,b) oxygen and (c,d) nitrogen atoms. Detected compounds are in the (a,c) gas phase and (b,d) particle phase. Pie charts correspond to the molecular weight contribution to the overall mass.**

As the measured carbon distribution is comparable at the two temperature conditions, only the experiment at 280 K is shown in the main text (see Fig. S2 for distributions at 295 K). The gas phase products distribution is dominated by C<sub>10</sub> and C<sub>8</sub> compounds containing mainly 2 or 3 oxygen atoms followed by C<sub>3</sub> and C<sub>4</sub> compounds, such as methylglyoxal, resulting from the further degradation of naphthalene oxidation products. As expected, the particle phase contains more oxygen atoms and is characterized by heavy molecular weight compounds with *m/z* above 150 comprising 75 % of the overall mass, explained by the readily partitioning of major C<sub>10</sub> products containing mainly 2 to 4 oxygen atoms. Table 3 presents the major identified ion fragments, their corresponding chemical formula and a tentative assignment to each compound based on previous studies (Bunce et al., 1997; Chan et al., 2009; Chen et al., 2016; Kautzman et al., 2010; Lee and Lane, 2009; Nishino et al., 2009; Riva, 2013; Sasaki et al., 1997; Tomaz, 2015). Similar to what was observed in the case of *m*-xylene, the overall oxidation product distributions, including nitrogenous species (1-4 % in both high and low NO<sub>x</sub> conditions), remain comparable across NO<sub>x</sub> conditions (Figs. S4 and S5 at 295 K). The strong similarity in product distributions and molecular compositions implies minimal shifts in bulk volatility as well in oxidation conditions.

**Table 3. List of ions and their corresponding formulas of the major naphthalene products detected during photooxidation experiments under high NO<sub>x</sub>. Reaction products are given as a fraction of the gas phase products (in % of μg m<sup>-3</sup>) and as a fraction of the SOA products (in % of μg m<sup>-3</sup>).**

Carbon number	Measured <i>m/z</i> and ion sum formula	Tentative assignment	<i>T</i> = 295 K		<i>T</i> = 280 K	
			Gaseous products (%)	SOA products (%)	Gaseous products (%)	SOA products (%)
10	161.06 (C <sub>10</sub> H <sub>8</sub> O <sub>2</sub> )H <sup>+</sup>	formyl cinnamaldehyde	2.0	8.9	3.5	25.4
10	193.05 (C <sub>10</sub> H <sub>8</sub> O <sub>4</sub> )H <sup>+</sup>	carboxy cinnamic acid	0.8	15.3	1.0	8.1
10	159.04 (C <sub>10</sub> H <sub>6</sub> O <sub>2</sub> )H <sup>+</sup>	naphthoquinone	6.6	5.6	4.9	9.3
10	177.05 (C <sub>10</sub> H <sub>8</sub> O <sub>3</sub> )H <sup>+</sup>	formyl cinnamic acid	0.6	7.7	1.1	7.1
10	175.04 (C <sub>10</sub> H <sub>6</sub> O <sub>3</sub> )H <sup>+</sup>	epoxy-naphthoquinone	1.6	8.4	1.9	4.8
10	145.07 (C <sub>10</sub> H <sub>8</sub> O)H <sup>+</sup>	naphthol	0.4	2.2	0.5	3.6
10	191.04 (C <sub>10</sub> H <sub>6</sub> O <sub>4</sub> )H <sup>+</sup>	dihydroxy naphthoquinone	0.7	3.5	0.8	2.0
10	195.06 (C <sub>10</sub> H <sub>10</sub> O <sub>4</sub> )H <sup>+</sup>	carboxybenzenepropanoic acid benzofurancarboxyaldehyde	0.3	1.5	0.3	1.0
10	174.05 (C <sub>10</sub> H <sub>7</sub> NO <sub>2</sub> )H <sup>+</sup>	<i>nitronaphthalene</i>	1.2	0.3	0.3	0.2
10	190.05 (C <sub>10</sub> H <sub>7</sub> NO <sub>3</sub> )H <sup>+</sup>	<i>nitronaphthol</i>	0.5	2.5	0.1	0.4
9	147.05 (C <sub>9</sub> H <sub>6</sub> O <sub>2</sub> )H <sup>+</sup>	benzopyrone	3.5	8.3	1.9	5.8
9	163.04 (C <sub>9</sub> H <sub>6</sub> O <sub>3</sub> )H <sup>+</sup>	hydroxycoumarin	1.5	6.3	2.2	4.5
9	133.06 (C <sub>9</sub> H <sub>8</sub> O)H <sup>+</sup>	indanone	1.6	1.3	1.3	3.6
9	179.04 (C <sub>9</sub> H <sub>6</sub> O <sub>4</sub> )H <sup>+</sup>	dihydroxy-indanedione	0.5	4.2	0.1	2.8
9	165.06 (C <sub>9</sub> H <sub>8</sub> O <sub>3</sub> )H <sup>+</sup>	hydroxy cinnamic acid	0.9	2.7	0.6	1.8
9	181.05 (C <sub>9</sub> H <sub>8</sub> O <sub>4</sub> )H <sup>+</sup>	dihydroxy cinnamic acid	0.2	1.5	0.2	0.9
8	135.05 (C <sub>8</sub> H <sub>6</sub> O <sub>2</sub> )H <sup>+</sup>	phthaldialdehyde / phthalide	19.3	3.9	5.4	4.3
8	149.03 (C <sub>8</sub> H <sub>4</sub> O <sub>3</sub> )H <sup>+</sup>	phthalic anhydride	13.8	3.3	9.9	2.0

8	151.04 (C <sub>8</sub> H <sub>6</sub> O <sub>3</sub> )H <sup>+</sup>	hydroxy phthaldehyde hydroxy phthalide	0.5	1.2	1.0	3.7
7	123.05 (C <sub>7</sub> H <sub>6</sub> O)H <sup>+</sup>	benzoic acid	1.9	0.2	1.4	0.1
7	107.05 (C <sub>7</sub> H <sub>6</sub> O)H <sup>+</sup>	benzaldehyde	1.0	n.d.	0.9	n.d.
6	111.05 (C <sub>6</sub> H <sub>6</sub> O <sub>2</sub> )H <sup>+</sup>	catechol / benzoquinone	1.2	n.d.	0.9	n.d.
6	115.07 (C <sub>6</sub> H <sub>10</sub> O <sub>2</sub> )H <sup>+</sup>	dimethyloxolan-one / hexanedione	1.3	0.3	1.3	0.6
5	101.06 (C <sub>5</sub> H <sub>8</sub> O <sub>2</sub> )H <sup>+</sup>	methyl furan	0.7	n.d.	1.6	n.d.
4	89.06 (C <sub>4</sub> H <sub>8</sub> O <sub>2</sub> )H <sup>+</sup>	hydroxybutanone / hydroxybutanal	10.1	0.1	11.5	n.d.
4	71.05 (C <sub>4</sub> H <sub>6</sub> O)H <sup>+</sup>	dihydrofuran / MACR / MVK	1.2	n.d.	1.8	n.d.
3	75.04 (C <sub>3</sub> H <sub>6</sub> O <sub>2</sub> )H <sup>+</sup>	propanoic acid	12.3	0.3	19.9	0.2
3	73.03 (C <sub>3</sub> H <sub>4</sub> O <sub>2</sub> )H <sup>+</sup>	methylglyoxal	3.6	0.1	5.4	0.1
2	77.02 (C <sub>2</sub> H <sub>4</sub> O <sub>3</sub> )H <sup>+</sup>	PAN fragment	2.0	0.1	1.3	0.1

\*n.d. = not detected

The C<sub>10</sub> compounds account for around 15 % of the gas phase products at both temperatures and clearly dominate the particulate phase with 59 % and 64 % of the overall aerosol mass at 295 K and 280 K, respectively. The 2-formyl cinnamaldehyde (C<sub>10</sub>H<sub>8</sub>O<sub>2</sub> at *m/z* 161.06) is the most abundant particle phase product accounts for 25 % of the SOA formed at 280 K. It is assumed to be formed via two possible routes i) ring cleavage upon the reaction of the naphthol peroxy radical (from an OH-naphthalene adduct) with NO, and ii) a hydrogen shift from the alcohol group on the naphthol peroxy radical followed by loss of OH and ring opening (Kautzman et al., 2010; Nishino et al., 2009; Qu et al., 2006; Sasaki et al., 1997; Wang et al., 2007). The predominance of one route over the other is depending on the amount of NO<sub>x</sub>. The 2-formyl cinnamaldehyde can further be oxidized, leading to the formation of 2-carboxy cinnamic acid (C<sub>10</sub>H<sub>8</sub>O<sub>4</sub> at *m/z* 193.05), the second most important C<sub>10</sub> compound that accounts for 8-15 % of the SOA mass. The oxidation of 2-formyl cinnamaldehyde also produces 2-formyl cinnamic acid (C<sub>10</sub>H<sub>8</sub>O<sub>3</sub> at *m/z* 177.05) (Bunce et al., 1997), that accounts for 7-8 % of the condensed phase. The addition of O<sub>2</sub> to the OH-naphthalene adduct results in the formation of epoxy-quinone (C<sub>10</sub>H<sub>6</sub>O<sub>3</sub> at *m/z* 175.04), representing 5-8 % of the particle phase. Other important C<sub>10</sub> ring retaining compounds are dihydroxy naphthoquinone (C<sub>10</sub>H<sub>6</sub>O<sub>4</sub> at *m/z* 191.04) and 1,4-naphthoquinone (C<sub>10</sub>H<sub>6</sub>O<sub>2</sub> at *m/z* 159.04). This latter has multiple formation routes, either from reaction of OH radical with naphthalene or with naphthol (C<sub>10</sub>H<sub>8</sub>O at *m/z* 145.07), or from the photodegradation of nitronaphthalene (C<sub>10</sub>H<sub>7</sub>NO<sub>2</sub> at *m/z* 174.05) (Atkinson et al., 1989; Kautzman et al., 2010), potentially explaining the low abundance of the latter (< 1 % in mass) alongside with its reaction product with OH, nitro-naphthol (C<sub>10</sub>H<sub>7</sub>NO<sub>3</sub> at *m/z* 190.05). At both high and low NO<sub>x</sub> conditions, the NO<sub>2</sub><sup>+</sup> signal in the particle phase did not exceed 3 % out of the aerosol mass detected by CHARON-PTR-ToF-MS meaning that the nitro-derivates of naphthalene did not undergo strong fragmentation. In total, the sum of all nitrogen-containing products (including PAN fragment) accounted for 2-3 % of the gaseous phase mass loading and 1-4 % of the particulate phase under high-NO<sub>x</sub> conditions. The observed low yields of nitro derivatives during the photo oxidation of naphthalene is in agreement with previous studies (Kautzman et al., 2010; Lee and Lane, 2009; Sasaki et al., 1997). Nitronaphthol (C<sub>10</sub>H<sub>7</sub>NO<sub>3</sub> at *m/z* 190.05) is a well known tracer of naphthalene SOA in high NO<sub>x</sub> conditions, and has been

observed here as a main nitrogen containing compound. Other previously reported tracers (nitrosalicylic acid, dinitrosalicylic acid or nitrophthalic acid) are not detected by CHARON-PTR-ToF-MS. The NO<sub>x</sub> conditions in our study might explain they are not formed, since the formation of these compounds is NO<sub>x</sub> dependent, and our study used lower NO<sub>x</sub> levels compared to studies using HONO or CH<sub>3</sub>NO as OH radical precursors, for example (Sato et al., 2022). It should also be considered that the UV lights used, which peaked at 310 nm, may induce photolysis of nitronaphthalene and other nitro-derivates, as previously reported in chamber experiments with similar UV lamps (Healy et al., 2012).

The formation of C<sub>9</sub> and C<sub>7</sub> products can be explained by H-abstraction of 2-formyl cinnamaldehyde, leading to a formyl peroxy radical that subsequently reacts with NO to form an alkoxy radical implying the loss of CO<sub>2</sub> group (Kautzman et al., 2010). The C<sub>9</sub> compounds contribute by 6-9 % to the gas phase and are the second highest contributors to the particle phase with 16-20 % of the total organic mass. Identified chemical formulas include C<sub>9</sub>H<sub>6</sub>O<sub>2</sub> (at *m/z* 147.05) which can be either benzopyrone (also known as coumarin) or indene-dione, as well as indanone (C<sub>9</sub>H<sub>8</sub>O at *m/z* 133.06) first detected by Lee and Lane (2009). No formation mechanism has been proposed for these products so far, but their molecular structure clearly indicates a rearrangement following ring opening, to form either a new ring at 6 atoms including one oxygen in the case of coumarin, or at 5 atoms for the indene-dione and indanone. This ring closure is a common feature in the case of dialdehydes oxidation, as explained by Lannuque and Sartelet (2024), which may support the hypothesis that these compounds originated from 2-formyl cinnamaldehyde. Further photo-oxidation of coumarin may lead to hydroxycoumarin (C<sub>9</sub>H<sub>6</sub>O<sub>3</sub> at *m/z* 163.04) accounting for 2.2 % of the gas phase and 4-6 % of the particle phase. A more oxygenated compound, dihydroxy-indanedione (C<sub>9</sub>H<sub>6</sub>O<sub>4</sub> at *m/z* 179.04), was previously identified by Lee et al. (2018). Logically, this compound was mainly present in the condensed phase with 3-4 % of the total SOA mass.

C<sub>8</sub> oxidation products make up a good 18-35 % and 10-11 % of the gas and particle phases, respectively. Phthaldialdehyde and phthalic anhydride largely dominate the C<sub>8</sub> compounds and are of particular importance in the gas phase. Phthaldialdehyde (C<sub>8</sub>H<sub>6</sub>O<sub>2</sub> at *m/z* 135.05), is both a first- and second-generation ring-opening product of the OH radical reaction with naphthalene, with the second-generation pathway expected to originate from the reaction of OH radical with 2-formyl cinnamaldehyde (Sasaki et al., 1997; Wang et al., 2007). Further addition of OH radical to phthaldialdehyde will lead to hydroxy phthaldehyde / hydroxy phthalide (C<sub>8</sub>H<sub>6</sub>O<sub>3</sub> at *m/z* 151.04) that makes up 3.7 % of SOA (Table 3), while the H-abstraction followed by intramolecular cyclization results in phthalic anhydride (C<sub>8</sub>H<sub>4</sub>O<sub>3</sub> at *m/z* 149.03) making up 2-3% of the condensed phase, and 10-14 % in gas phase (Table 3) previously observed by Wang et al. (2007).

C<sub>7</sub> compounds are relatively less abundant accounting for 3 % of the gas phase products and less than 1 % of SOA. They include compounds like benzoic acid (C<sub>7</sub>H<sub>6</sub>O<sub>2</sub> at *m/z* 123.05) and benzaldehyde (C<sub>7</sub>H<sub>6</sub>O at *m/z* 107.05).

Only two C<sub>6</sub> compounds are detected and represent 3-4 % of the total gas products and only 1 % of the SOA yield. The dominant C<sub>6</sub> in the gaseous phase is C<sub>6</sub>H<sub>10</sub>O<sub>2</sub> (at *m/z* 115.07) with 1.3 %, which can be dimethyloxolanone or hexanedione. The former can be formed from the oxidation of phthaldialdehyde, implying the opening of the second aromatic cycle (Kautzman et al., 2010). Another minor C<sub>6</sub> compound is catechol or benzoquinone (C<sub>6</sub>H<sub>10</sub>O<sub>2</sub> at *m/z* 115.07).

C<sub>5</sub> products account for only 3-6 % of the gaseous phase and less than 2 % of SOA. A major detected compound is C<sub>5</sub>H<sub>8</sub>O<sub>2</sub> at *m/z* 101.06 (1.6 %) which can be assigned as methyl furan or 4-oxopentanal. The C<sub>2</sub>-C<sub>4</sub> products are more volatile, all together make less than 2 % of the total SOA mass but present a considerable fraction of the gas phase products with 18 % for C<sub>4</sub>, 26 % for C<sub>3</sub>, and 4 % for C<sub>2</sub>. The major C<sub>4</sub> contributor is C<sub>4</sub>H<sub>8</sub>O<sub>2</sub> (at *m/z* 89.06) representing 12 % of the gas phase products. It is tentatively assigned to hydroxybutanal or butanoic acid. Almost all the C<sub>3</sub> fraction is represented by propanoic acid (C<sub>3</sub>H<sub>6</sub>O<sub>2</sub> at *m/z* 75.04) with 12-20 % contribution to the gas phase products, and methylglyoxal (C<sub>3</sub>H<sub>4</sub>O<sub>2</sub> at *m/z* 73.03) up to 5 %. The C<sub>2</sub> compound is tentatively assigned to the PAN fragment C<sub>2</sub>H<sub>4</sub>O<sub>3</sub> detected at *m/z* 77.02 (1.4 %) (Müller et al., 2012).

### 3.3 Experimentally derived and estimated gas-particle phase partitioning

A two-dimensional space, the 2D-VBS framework (Donahue et al., 2011; Murphy et al., 2012), is used as a means to visualize the compounds distribution as function of the experimentally derived volatility ( $\log_{10}C_i^*$ ) and the O/C ratio (Fig. 5) or the oxidation state of carbon (OSc) (Fig. S4). Such molecular level effect of temperature on SOA for m-xylene and naphthalene is reported here for the first time. Figures 5a and 5b display the distribution of measured volatilities for the major compounds detected in both gas and particle phases for m-xylene and naphthalene, respectively, under high NO<sub>x</sub> conditions. The volatilities of the identified compounds are in the range of SVOCs with  $C_i^*$  values ranging from 0.3 to 300  $\mu\text{g m}^{-3}$  (vapor pressures approximately 10<sup>-8</sup> to 10<sup>-5</sup> torr) to IVOCs with  $C_i^*$  values ranging from 300 to 3 × 10<sup>6</sup>  $\mu\text{g m}^{-3}$  (vapor pressures approximately 10<sup>-5</sup> to 10<sup>-1</sup> torr) (Donahue et al., 2012), indicated by the light green ( $\log_{10}C_i^*$  from 0 to 2.5) and light blue ( $\log_{10}C_i^*$  from 2.5 to 5) background shading, respectively. Data points are indicated for the experiments at 280 K (purple circles) and 295 K (red circles), and the size of the circles is proportional to the mass concentration (in  $\mu\text{g m}^{-3}$ ) of each ion in the particle phase. For m-xylene approximately out of 110 ions detected in the particle phase, half of them are exclusively found in the particle phase. This fraction should populate a low volatility area (as ELVOC), and it is not represented in this 2D-VBS (Fig.5a) but represents more than 30 % of the SOA mass. Among the remaining ions that partition between the two phases, only 18 are considered in Fig.5a, either because they are important in terms of mass or because they are considered parent ions following Gkatzelis et al. (2018) method. About 24 to 58 % of the particle mass populates the SVOCs regime, while 8 to 16 % are in the IVOCs regime, depending on the experimental conditions. In a recent work the SOA components from OH radical oxidation of toluene populated mostly the SVOC range and only 10-17 % was exclusively in the particle phase (Lannuque et al., 2023), probably because of higher initial VOC concentration and lower OH exposure (10-20 h compared to 1.3-3 days in this study) leading to a less advanced oxidation. But overall, the effect of temperature observed is quite similar for the two chemical systems. For naphthalene at 280 K, out of the 110 ions detected in the particle phase, 40 ions are exclusively observed in the particle phase, making up 5 % only of the condensed mass fraction. For the selected 20 compounds that partition between the two phases, approximately 78-95 % of their SOA mass populates the SVOC regime and 3-17 % lies in the IVOC portion. The larger skeleton structure of naphthalene oxidation products, even first generation, can explain the high fraction of SOA in the SVOC regime (predominantly C<sub>8-10</sub> compounds) as confirmed by the O/C range, lower for naphthalene than for m-xylene.

538 The derived saturation concentration ( $C_i^*$ ) values range between 1 to 6919  $\mu\text{g m}^{-3}$ , comparable to other SOA systems (Gkatzelis  
539 et al., 2018; Kostenidou et al., 2024; Lannuque et al., 2023). Previous studies on biogenic VOCs reported a decreasing volatility  
540 with increasing OSc (Gkatzelis et al., 2018; Jimenez et al., 2009; Kroll et al., 2011), while the present works and previous  
541 investigations on anthropogenic VOCs (toluene and gasoline vehicle emissions) did not confirm such a trend (Kostenidou et  
542 al., 2024; Lannuque et al., 2023).

543 Figures 5c and 5d present the difference of  $\log_{10}C_i^*$  between the two experiments conducted at 280 and 295 K,  $\Delta\log_{10}C_i^*$ . Its  
544 values vary from 0.06 to 1.08 as a function of carbon number and oxygen number (tables S2 and S3). For m-xylene oxidation  
545 products (Fig. 5c),  $\Delta\log_{10}C_i^*$  decreases with increasing carbon number. Indeed, the C<sub>5</sub>-C<sub>8</sub> products bearing 3 to 5 oxygen atoms  
546 (darker green markers) lie at the bottom of the plot generally exhibiting  $\Delta\log_{10}C_i^*$  values below 0.4, emphasizing the effect of  
547 oxygen atoms in reducing the volatility of these compounds. The C<sub>3</sub> compounds, tentatively associated to methylglyoxal and  
548 propanoic acid, exhibit also moderate  $\Delta\log_{10}C_i^*$ , probably because they are still volatile even at the lower experimental  
549 temperatures (Table 2). Only few among the identified C<sub>4</sub>-C<sub>5</sub> products exhibit  $\Delta\log_{10}C_i^*$  values above 0.6 and are tentatively  
550 identified as ring opening products, with multiple functional groups increasing their polarity (aldehydes, acids and  
551 furandiones).

552





Figure 5. Panels (a) and (b) present the 2D-VBS framework: the O/C ratio (y-axis) for the detected parent ions of m-xylene and naphthalene photooxidation, respectively, as a function of saturation concentration ( $\log_{10}C_i^*$  in  $\mu\text{g m}^{-3}$ ; x-axis) for high  $\text{NO}_x$  experiments. The size of the circles denotes the mass of each species. Experiments carried out at 280 K are in light violet while the ones carried out at 295 K are in magenta. The light green and the light blue background shadings correspond to the SVOC ( $0 < \log_{10}C_i^* < 2.5$ ) and the IVOC ( $2.5 < \log_{10}C_i^* < 5$ ) regimes, respectively. Panels (c) and (d) present  $\Delta\log_{10}C_i^*$  (the difference of  $\log_{10}C_i^*$  between experiments at 280 K and 295 K; y-axis) as function of carbon number (x-axis), color scaled by oxygen number and sized by the mass (in  $\mu\text{g m}^{-3}$ ) of each compound at 280 K for m-xylene and naphthalene oxidation products, respectively.

Naphthalene oxidation products on the other hand (Fig. 5d) have  $\Delta\log_{10}C_i^*$  over a similar range compared to m-xylene, but exhibit a different behaviour. Most of the relevant reaction products are associated to  $\text{C}_{8-10}\text{H}_{4-10}\text{O}_{1-4}$  products, accounting for 90 % of the SOA mass and have at least one aromatic ring. These compounds belong SVOC with a  $\log_{10}C_i^* < 2$ , and span a broad range of  $\Delta\log_{10}C_i^*$  depending on the number of oxygen atoms (from 1 to 5) and the specific functionalities. Logically, the most oxygenated  $\text{C}_{10}$  are poorly affected by temperature, since they are the least volatile and already mostly in the particle phase at 295 K. Compared to m-xylene, the most important naphthalene SOA-products are less oxygenated and seem to be more temperature-sensitive.



### 3.4 Comparison of experimentally derived and calculated volatilities

The experimentally derived volatilities are here compared to the estimated ones from pure-liquid saturation vapour pressure using Volcalc based on SIMPOL.1 (Meredith et al., 2023; Pankow and Asher, 2008; Riemer, 2023). Volatilities from the two methods are presented in Figs. 6a and 6b for both m-xylene and naphthalene oxidation products at 280 K, respectively. A considerable discrepancy is observed between the experimental and calculated values, the latter ones spanning a larger range of volatilities. For m-xylene SOA, the theoretical approach tends to largely overestimate the volatility of small and oxygenated compounds below  $m/z$  120, similarly to recent investigations using different techniques as SV-TAG coupled to GC-MS, thermal desorption-AMS and FIGAERO-CIMS and CHARON (Stark et al., 2017; Ijaz et al., 2024; Liang et al., 2023). For some light carbonyl compounds present among the m-xylene reaction products, the disagreement can be potentially explained by presence of ammonium ions ( $\text{NH}_4^+$ ) in the seeds that can act as a catalyst for accretion reactions such as aldol condensation (Li et al., 2022, 2011; Nozière et al., 2010; Sareen et al., 2010) as well as acetal and hemiacetal formation (Jang et al., 2002; Li et al., 2022; Loeffler et al., 2006; Shapiro et al., 2009) shifting the equilibrium to the condensed phase for these molecules. Furthermore, Lannuque et al. (2023) have achieved better model representation of experimental SOA mass concentration after including interactions between aldehydes as methylglyoxal and inorganic compounds (such as ammonium). Those reactions are also catalyzed in the presence of water, especially for highly oxidized hydrophilic compounds (Meng et al., 2024). For some other compounds, as functionalized acids and some dialdehydes, disagreements may arise from the fragmentation in the mass spectrometer. Despite a relatively low  $E/N$  value (68 Td) used in this work, fragmentation may still occur, particularly in polyfunctional products (Leglise et al., 2019). We also tested the fragmentation of several compounds in a separate work (Lannuque et al., 2023) where we could observe low or negligible fragmentation for methylglyoxal, furans, furfurals, maleic acids and anhydrides but important fragmentation for small linear aldehydes.



**Figure 6. Theoretical (y-axis) versus experimental  $\log_{10}C_i^*$  values (x-axis) for (a) m-xylene and (b) naphthalene oxidation products at 280 K under high  $\text{NO}_x$  conditions. The black line is the 1:1 fit. The size of the markers is proportional to the oxygen number. The color gradient corresponds to  $m/z$  of the detected compounds.**

For naphthalene SOA products, larger discrepancies are observed for larger compounds ( $m/z \geq 160$ ) associated to  $\text{C}_{8-10}$  compounds bearing multiple functional groups. For such types of compounds, the model predicts very low volatilities (O'Meara et al., 2014) also demonstrating that SIMPOL.1 consistently overestimates temperature-induced volatility changes for highly oxygenated compounds (such as hydroxylated ketones and diacids). These discrepancies are attributed to the additive nature of SIMPOL.1's functional group contribution framework. At 280 K, a carboxylic acid group ( $-\text{COOH}$ ) decreases the saturation vapor pressure by nearly a factor of 6000, a hydroxyl group ( $-\text{OH}$ ) by factor 200, while a ketone group ( $=\text{O}$ ) reduces it by less than a factor of 9. It is also worth noting that multiple structural isomers are not distinguished in our work, and this may also introduce some uncertainties. An example is provided by 1,2-naphthoquinone and 1,4-naphthoquinone owing quite different vapor pressure making the former potentially more interesting in the particle phase (McWhinney et al., 2013). Liang et al. (2023) and Dang et al., (2019) also reported very different gas-particle partitioning behavior and vapor pressures values among isomers. Barley and McFiggans (2010) and Peräkylä et al. (2020) highlighted significant overpredictions for diacids and polyfunctional compounds, attributing these errors to the model's parameterizations and group contributions. This work therefore aligns with previous findings and suggests that vapor pressures of multifunctional compounds of the aerosol are still not well estimated by models.

Figure S7 shows the comparison of estimated versus experimental  $\log_{10}C_i^*$  values at 295 K under high  $\text{NO}_x$  conditions for both m-xylene and naphthalene. These results closely match those at 280 K (Fig. 6), with no significant difference in the distribution.

Figure S8 presents the same analysis under low NO<sub>x</sub> conditions at 295 K, which similarly shows no notable deviation from the trends observed at low NO<sub>x</sub> (Fig. S7). This indicates that neither temperature nor NO<sub>x</sub> regime substantially explain the model-observation discrepancy in our dataset. Temperature-induced volatility changes are quantified in Figs. 5c,d which represents the difference between experimental log<sub>10</sub>*C<sub>i</sub>*<sup>\*</sup> at 280 K and 295 K (i.e. Δlog<sub>10</sub>*C<sub>i</sub>*<sup>\*</sup>) against the C number. The temperature effect on the discrepancy between the experimental and the estimated log<sub>10</sub>*C<sub>i</sub>*<sup>\*</sup> values is presented in Fig. S9. The values of Δlog<sub>10</sub>*C<sub>i</sub>*<sup>\*</sup>, the difference between estimated and experimental log<sub>10</sub>*C<sub>i</sub>*<sup>\*</sup>, for the oxidation products of m-xylene (Fig. S9a) and naphthalene (Fig. S9b) is plotted as a function *m/z*, at 295 K and 280 K. The discrepancies at both temperatures are quite close to each other. Overall, SIMPOL.1's bias in predicting volatility for oxidation products is not strongly affected by temperature. This consistent discrepancy with temperature and NO<sub>x</sub> confirmed that partitioning of organic compounds is more complex than the simple evaporation of compounds isolated, and that surface interactions and bulk chemical reactions are potential important drivers affecting partitioning equilibria.

#### 4 Conclusion

This study presents a detailed experimental investigation of the OH-initiated photooxidation of two anthropogenic aromatic precursors, m-xylene and naphthalene, using an OFR under different NO<sub>x</sub> and temperature conditions. For both precursors, SOA yields are found to strongly increase when temperature and NO<sub>x</sub> decrease, in agreement with previous studies. The CHARON-inlet coupled to a PTR-ToF-MS successfully quantified between 65-80 % of the total organic mass covered by an aerosol mass spectrometer (HR-ToF-AMS). Major products both in the gas and particulate phases are confirmed based PTR-ToF-MS inferred molecular formula and intercomparison with the literature. Major gas phase products of m-xylene SOA are C<sub>3</sub>, C<sub>5</sub> and C<sub>8</sub> compounds whereas the particle phase products are dominated by C<sub>6</sub>-C<sub>8</sub> compounds. This pattern is consistent through different temperatures, indicating similar chemistry in these conditions. For naphthalene experiments, gas phase products are dominated by C<sub>8</sub> and C<sub>10</sub> compounds, while the particle phase composition mainly consists of C<sub>8</sub>, C<sub>9</sub> and C<sub>10</sub> compounds. Nitro-derivatives (nitrogen containing compounds + PAN fragment) measured in both phases did not exceed 7 % in naphthalene experiments whereas they could reach up to 20 % in the m-xylene experiments. The similarity in product distributions under both high and low NO<sub>x</sub> conditions for both m-xylene and naphthalene indicates that the chemical reactions were largely unaffected by the NO<sub>x</sub> levels, implying that NO<sub>x</sub> had minimal influence on oxidant concentrations in the OFR. The volatility properties of the individual compounds are presented in the 2D-VBS framework: 24-58 % of the SOA mass generated by m-xylene populates the SVOC regime and 8-16 % populates the IVOCs regime, while the naphthalene SOA mass is mostly (up to 95 %) found in the SVOC regime. No clear correlation could be observed between the volatility values and the increasing oxidation state (OSc), in agreement with previous studies on anthropogenic monoaromatic precursors. Temperature variation, from 295 to 280 K, induced an expected decrease in volatility (Δlog<sub>10</sub>*C<sub>i</sub>*<sup>\*</sup>), this decrease ranging from 0.06 to 1.08. The magnitude of Δlog<sub>10</sub>*C<sub>i</sub>*<sup>\*</sup> seems to be controlled by multiple parameters, as temperature, carbon number, oxygen number and specific chemical moieties. When experimentally derived volatilities are compared to a group contribution

parameterization model based on pure liquid vapor pressure (SIMPOL.1), large discrepancies are observed. The small discrepancies between estimated and experimental  $\log_{10}C_i^*$  at both temperatures indicate that the temperature has little impact on SIMPOL.1's bias in predicting the volatility of oxidation products. The difference between estimated and experimental volatilities highlight the complexity of gas-particle partitioning, and the limit of parameterizations that should be further validated using larger datasets from various measurement techniques and experimental conditions.

This study advances our understanding of SOA composition and gas-particle partitioning at different and relevant atmospheric conditions and also holds some implications for urban air quality management and climate modelling. The observed temperature-dependent shifts in SOA mass loadings and volatility highlight the importance of accounting for seasonal variations, particularly in urban areas with high anthropogenic VOC emissions. Furthermore, the findings underscore the need to refine current air quality and climate models by incorporating more recent findings and real-world atmospheric conditions, paving the way for more accurate predictions of aerosol impacts on air quality, climate, and public health.

#### **Data availability**

Finalized data are mostly available in supplementary material. More detailed data are available on request.

#### **Supplementary material**

#### **Competing interests**

The authors declare that they have no conflict of interest.

#### **Author contribution**

MS, JK, and BD designed the experimental setup. MS, JK, BTR and BD performed the experiments and data treatment. MS, BD and JK analysed and interpreted the data. MS ran the SIMPOL.1 model. MS drafted the article. All the co-authors revised the article.

#### **Financial Support**

This work was funded by the POLEMICS project of the Agence Nationale de la Recherche (ANR) program (grant ANR-18-CE22-0011), and the MAESTRO-EU6 project (ADEME CORTEA n. 1866C0001), the French government under the France 2030 investment plan, as part of the Initiative d'Excellence d'Aix-Marseille Université – A\*MIDEX ” AMX-21-PEP-016, the French national research agency (ANR-22-CE22-0003-01) and the Environmental Sciences doctoral school (ED 251). BD

acknowledges the MITI program from CNRS for the financial support allowing the design and the construction of the new oxidation flow reactor (OFR) deployed in this study.

## References

- Ahlberg, E., Eriksson, A., Brune, W. H., Roldin, P., and Svenningsson, B.: Effect of salt seed particle surface area, composition and phase on secondary organic aerosol mass yields in oxidation flow reactors, *Atmos. Chem. Phys.*, 19, 2701–2712, <https://doi.org/10.5194/acp-19-2701-2019>, 2019.
- Ahn, J., Rao, G., and Vejerano, E.: Temperature dependence of the gas-particle partitioning of selected VOCs, *Environ. Sci.: Processes Impacts*, 23, 947–955, <https://doi.org/10.1039/D1EM00176K>, 2021.
- Anderson, J. O., Thundiyil, J. G., and Stolbach, A.: Clearing the Air: A Review of the Effects of Particulate Matter Air Pollution on Human Health, *J. Med. Toxicol.*, 8, 166–175, <https://doi.org/10.1007/s13181-011-0203-1>, 2012.
- Aoki, N., Inomata, S., and Tanimoto, H.: Detection of C1–C5 alkyl nitrates by proton transfer reaction time-of-flight mass spectrometry, *International Journal of Mass Spectrometry*, 263, 12–21, <https://doi.org/10.1016/j.ijms.2006.11.018>, 2007.
- Atkinson, R., Aschmann, S. M., and Zielinska, B.: GAS-PHASE ATMOSPHERIC CHEMISTRY OF 1- AND 2-NITRONAPHTHALENE AND 1,4-NAPHTHOQUINONE, 1989.
- Atkinson, R., Aschmann, S. M., and Arey, J.: Formation of ring-retaining products from the OH radical-initated reactions of *o* -, *m* -, and *p* -xylene, *Int J of Chemical Kinetics*, 23, 77–97, <https://doi.org/10.1002/kin.550230108>, 1991.
- Aumont, B., Camredon, M., Mouchel-Vallon, C., La, S., Ouzebidou, F., Valorso, R., Lee-Taylor, J., and Madronich, S.: Modeling the influence of alkane molecular structure on secondary organic aerosol formation, *Faraday Discuss.*, 165, 105, <https://doi.org/10.1039/c3fd00029j>, 2013.
- Bahrami, A., Haghighat, F., and Zhu, J.: Indoor environment gas-particle partitioning models of SVOCs and impact of particle properties on the partitioning: A review., *Building & Environment*, 262, N.PAG, <https://doi.org/10.1016/j.buildenv.2024.111791>, 2024.
- Barley, M. H. and McFiggans, G.: The critical assessment of vapour pressure estimation methods for use in modelling the formation of atmospheric organic aerosol, *Atmos. Chem. Phys.*, 10, 749–767, <https://doi.org/10.5194/acp-10-749-2010>, 2010.
- Baudic, A., Gros, V., Sauvage, S., Locoge, N., Sanchez, O., Sarda-Estève, R., Kalogridis, C., Petit, J.-E., Bonnaire, N., Baisnée, D., Favez, O., Albinet, A., Sciare, J., and Bonsang, B.: Seasonal variability and source apportionment of volatile organic compounds (VOCs) in the Paris megacity (France), *Atmos. Chem. Phys.*, 16, 11961–11989, <https://doi.org/10.5194/acp-16-11961-2016>, 2016.
- Berlinger, B., Fehérvári, P., Kővágó, C., Lányi, K., Mátis, G., Mackei, M., and Könyves, L.: There Is Still a Need for a Comprehensive Investigation of the Health Consequences of Exposure to Urban Air with Special Regard to Particulate Matter (PM) and Cardiovascular Effects, *Atmosphere*, 15, 296, <https://doi.org/10.3390/atmos15030296>, 2024.
- Birdsall, A. W. and Elrod, M. J.: Comprehensive NO-Dependent Study of the Products of the Oxidation of Atmospherically Relevant Aromatic Compounds, *J. Phys. Chem. A*, 115, 5397–5407, <https://doi.org/10.1021/jp2010327>, 2011.

699 Bloss, C., Wagner, V., Jenkin, M. E., Volkamer, R., Bloss, W. J., Lee, J. D., Heard, D. E., Wirtz, K., Martin-Reviejo, M., Rea,  
700 G., Wenger, J. C., and Pilling, M. J.: Development of a detailed chemical mechanism (MCMv3.1) for the atmospheric oxidation  
701 of aromatic hydrocarbons, *Atmos. Chem. Phys.*, 5, 641–664, <https://doi.org/10.5194/acp-5-641-2005>, 2005.

702 Bosque, R. and Sales, J.: Polarizabilities of Solvents from the Chemical Composition, *J. Chem. Inf. Comput. Sci.*, 42, 1154–  
703 1163, <https://doi.org/10.1021/ci025528x>, 2002.

704 Bunce, N. J., Liu, L., Zhu, J., and Lane, D. A.: Reaction of Naphthalene and Its Derivatives with Hydroxyl Radicals in the Gas  
705 Phase, *Environ. Sci. Technol.*, 31, 2252–2259, <https://doi.org/10.1021/es960813g>, 1997.

706 Calvert, J. G., Atkinson, R., Becker, K. H., Kamens, R. M., Wallington, T. H., Seinfeld, J. H., and Yarwood, G.: *The  
707 Mechanisms of Atmospheric Oxidation of Aromatic Hydrocarbons*, Oxford University Press, 2002.

708 Calvert, J. G., Orlando, J. J., Stockwell, W. R., and Wallington, T. J.: *The Mechanisms of Reactions Influencing Atmospheric  
709 Ozone*, Oxford University Press, <https://doi.org/10.1093/oso/9780190233020.001.0001>, 2015.

710 Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B., Drewnick, F., Coe, H.,  
711 Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P.,  
712 and Worsnop, D. R.: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass  
713 spectrometer, *Mass Spectrometry Reviews*, 26, 185–222, <https://doi.org/10.1002/mas.20115>, 2007.

714 Cappa, C. D. and Wilson, K. R.: Multi-generation gas-phase oxidation, equilibrium partitioning, and the formation and  
715 evolution of secondary organic aerosol, *Atmos. Chem. Phys.*, 12, 9505–9528, <https://doi.org/10.5194/acp-12-9505-2012>,  
716 2012.

717 Chan, A. W. H., Kautzman, K. E., Chhabra, P. S., Surratt, J. D., Chan, M. N., Crounse, J. D., Wennberg, P. O., Flagan, R. C.,  
718 and Seinfeld, J. H.: Secondary organic aerosol formation from photooxidation of naphthalene and alkylnaphthalenes:  
719 implications for oxidation of intermediate volatility organic compounds (IVOCs), *Atmos. Chem. Phys.*, 9, 3049–3060,  
720 <https://doi.org/10.5194/acp-9-3049-2009>, 2009.

721 Chen, C.-L., Kacarab, M., Tang, P., and Cocker, D. R.: SOA formation from naphthalene, 1-methylnaphthalene, and 2-  
722 methylnaphthalene photooxidation, *Atmospheric Environment*, 131, 424–433,  
723 <https://doi.org/10.1016/j.atmosenv.2016.02.007>, 2016.

724 Chen, C.-L., Li, L., Tang, P., and Cocker, D. R.: SOA formation from photooxidation of naphthalene and methylnaphthalenes  
725 with m-xylene and surrogate mixtures, *Atmospheric Environment*, 180, 256–264,  
726 <https://doi.org/10.1016/j.atmosenv.2018.02.051>, 2018.

727 Chen, T., Chu, B., Ma, Q., Zhang, P., Liu, J., and He, H.: Effect of relative humidity on SOA formation from aromatic  
728 hydrocarbons: Implications from the evolution of gas- and particle-phase species, *Science of The Total Environment*, 773,  
729 145015, <https://doi.org/10.1016/j.scitotenv.2021.145015>, 2021.

730 Chen, T., Liu, Y., Chu, B., Liu, C., Liu, J., Ge, Y., Ma, Q., Ma, J., and He, H.: Differences of the oxidation process and  
731 secondary organic aerosol formation at low and high precursor concentrations, *Journal of Environmental Sciences*, 79, 256–  
732 263, <https://doi.org/10.1016/j.jes.2018.11.011>, 2019.

733 Clark, C. H., Kacarab, M., Nakao, S., Asa-Awuku, A., Sato, K., and Cocker, D. R.: Temperature Effects on Secondary Organic  
734 Aerosol (SOA) from the Dark Ozonolysis and Photo-Oxidation of Isoprene, *Environ. Sci. Technol.*, 50, 5564–5571,  
735 <https://doi.org/10.1021/acs.est.5b05524>, 2016.

- 737 Dang, C., Bannan, T., Shelley, P., Priestley, M., Worrall, S. D., Waters, J., Coe, H., Percival, C. J., and Topping, D.: The effect  
738 of structure and isomerism on the vapor pressures of organic molecules and its potential atmospheric relevance, *Aerosol*  
739 *Science and Technology*, 53, 1040–1055, <https://doi.org/10.1080/02786826.2019.1628177>, 2019.
- 740 DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T.,  
741 Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-Deployable, High-Resolution, Time-of-Flight Aerosol Mass  
742 Spectrometer, *Analytical Chemistry*, 78, 8281–8289, <https://doi.org/10.1021/ac061249n>, 2006.
- 743 Deng, Y., Inomata, S., Sato, K., Ramasamy, S., Morino, Y., Enami, S., and Tanimoto, H.: Temperature and acidity dependence  
744 of secondary organic aerosol formation from  $\alpha$ -pinene ozonolysis with a compact chamber system, *Atmos. Chem. Phys.*, 21,  
745 5983–6003, <https://doi.org/10.5194/acp-21-5983-2021>, 2021.
- 746 Dodge, M.: Combined use of modeling techniques and smog chamber data to derive ozone-precursor relationships,  
747 *International conference on photochemical oxidant pollution and its control: Proceedings*, 881–889, 1977.
- 748 Donahue, N. M., Robinson, A. L., Stanier, C. O., and Pandis, S. N.: Coupled Partitioning, Dilution, and Chemical Aging of  
749 Semivolatile Organics, *Environ. Sci. Technol.*, 40, 2635–2643, <https://doi.org/10.1021/es052297c>, 2006.
- 750 Donahue, N. M., Epstein, S. A., Pandis, S. N., and Robinson, A. L.: A two-dimensional volatility basis set: 1. organic-aerosol  
751 mixing thermodynamics, *Atmos. Chem. Phys.*, 11, 3303–3318, <https://doi.org/10.5194/acp-11-3303-2011>, 2011.
- 752 Donahue, N. M., Kroll, J. H., Pandis, S. N., and Robinson, A. L.: A two-dimensional volatility basis set – Part 2: Diagnostics  
753 of organic-aerosol evolution, *Atmos. Chem. Phys.*, 12, 615–634, <https://doi.org/10.5194/acp-12-615-2012>, 2012.
- 754 Duncianu, M., David, M., Kartigeyane, S., Cirtog, M., Doussin, J.-F., and Picquet-Varrault, B.: Measurement of alkyl and  
755 multifunctional organic nitrates by proton-transfer-reaction mass spectrometry, *Atmos. Meas. Tech.*, 10, 1445–1463,  
756 <https://doi.org/10.5194/amt-10-1445-2017>, 2017.
- 757 Eichler, P., Müller, M., D’Anna, B., and Wisthaler, A.: A novel inlet system for online chemical analysis of semi-volatile  
758 submicron particulate matter, *Atmos. Meas. Tech.*, 8, 1353–1360, <https://doi.org/10.5194/amt-8-1353-2015>, 2015.
- 759 European Environment Agency: Europe’s air quality status 2021, Publications Office, LU, 2022.
- 760 Fan, J. and Zhang, R.: Density Functional Theory Study on OH-Initiated Atmospheric Oxidation of *m*-Xylene, *J. Phys. Chem.*  
761 *A*, 112, 4314–4323, <https://doi.org/10.1021/jp077648j>, 2008.
- 762 Fan, C., Wang, W., Wang, K., Lei, T., Xiang, W., Hou, C., Li, J., Guo, Y., and Ge, M.: Temperature effects on SOA formation  
763 of n-dodecane reaction initiated by Cl atoms, *Atmospheric Environment*, 346, 121070,  
764 <https://doi.org/10.1016/j.atmosenv.2025.121070>, 2025.
- 765 Fang, H., Luo, S., Huang, X., Fu, X., Xiao, S., Zeng, J., Wang, J., Zhang, Y., and Wang, X.: Ambient naphthalene and  
766 methylnaphthalenes observed at an urban site in the Pearl River Delta region: Sources and contributions to secondary organic  
767 aerosol, *Atmospheric Environment*, 252, 118295, <https://doi.org/10.1016/j.atmosenv.2021.118295>, 2021.
- 768 Fang, H., Wang, W., Xu, H., Huang, Y., Jiang, H., Wu, T., Li, J., Zha, S., Zhang, J., Zhou, R., and Wang, X.: Sources and  
769 secondary transformation potentials of aromatic hydrocarbons observed in a medium-sized city in yangtze river delta region:  
770 Emphasis on intermediate-volatility naphthalene, *Atmospheric Environment*, 318, 120239,  
771 <https://doi.org/10.1016/j.atmosenv.2023.120239>, 2024.

772 Forstner, H. J. L., Flagan, R. C., and Seinfeld, J. H.: Secondary Organic Aerosol from the Photooxidation of Aromatic  
 773 Hydrocarbons: Molecular Composition, *Environ. Sci. Technol.*, 31, 1345–1358, <https://doi.org/10.1021/es9605376>, 1997.

774 Gioumousis, G. and Stevenson, D. P.: Reactions of Gaseous Molecule Ions with Gaseous Molecules. V. Theory, *The Journal*  
 775 *of Chemical Physics*, 29, 294–299, <https://doi.org/10.1063/1.1744477>, 1958.

776 Gkatzelis, G. I., Hohaus, T., Tillmann, R., Gensch, I., Müller, M., Eichler, P., Xu, K.-M., Schlag, P., Schmitt, S. H., Yu, Z.,  
 777 Wegener, R., Kaminski, M., Holzinger, R., Wisthaler, A., and Kiendler-Scharr, A.: Gas-to-particle partitioning of major  
 778 biogenic oxidation products: a study on freshly formed and aged biogenic SOA, *Atmos. Chem. Phys.*, 18, 12969–12989,  
 779 <https://doi.org/10.5194/acp-18-12969-2018>, 2018.

780 Heald, C. L., Jacob, D. J., Park, R. J., Russell, L. M., Huebert, B. J., Seinfeld, J. H., Liao, H., and Weber, R. J.: A large organic  
 781 aerosol source in the free troposphere missing from current models, *Geophysical Research Letters*, 32, 2005GL023831,  
 782 <https://doi.org/10.1029/2005GL023831>, 2005.

783 Healy, R. M., Chen, Y., Kourtchev, I., Kalberer, M., O’Shea, D., and Wenger, J. C.: Rapid Formation of Secondary Organic  
 784 Aerosol from the Photolysis of 1-Nitronaphthalene: Role of Naphthoxy Radical Self-reaction, *Environ. Sci. Technol.*, 46,  
 785 11813–11820, <https://doi.org/10.1021/es302841j>, 2012.

786 Henze, D. K., Seinfeld, J. H., Ng, N. L., and Kroll, J. H.: Global modeling of secondary organic aerosol formation from  
 787 aromatic hydrocarbons: high- vs. low-yield pathways, *Atmos. Chem. Phys.*, 8, 2405–2421, [https://doi.org/10.5194/acp-8-](https://doi.org/10.5194/acp-8-2405-2008)  
 788 2405-2008, 2008.

789 Huang, D. D., Kong, L., Gao, J., Lou, S., Qiao, L., Zhou, M., Ma, Y., Zhu, S., Wang, H., Chen, S., Zeng, L., and Huang, C.:  
 790 Insights into the formation and properties of secondary organic aerosol at a background site in Yangtze River Delta region of  
 791 China: Aqueous-phase processing vs. photochemical oxidation, *Atmospheric Environment*, 239, 117716,  
 792 <https://doi.org/10.1016/j.atmosenv.2020.117716>, 2020.

793 Huang, M., Zhang, W., Hao, L., Wang, Z., Zhao, W., Gu, X., and Fang, L.: Low-Molecular Weight and Oligomeric  
 794 Components in Secondary Organic Aerosol from the Photooxidation of p-Xylene, *J Chinese Chemical Soc.*, 55, 456–463,  
 795 <https://doi.org/10.1002/jccs.200800068>, 2008.

796 Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco,  
 797 F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade,  
 798 G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad, I. E., and Prévôt, A. S. H.: High  
 799 secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218–222,  
 800 <https://doi.org/10.1038/nature13774>, 2014.

801 Ijaz, A., Temime-Roussel, B., Kammer, J., Mao, J., Simpson, W., Law, K. S., and Barbara, D.: In situ measurements of gas-  
 802 particle partitioning of organic compounds in Fairbanks, *Faraday Discuss.*, 10.1039.D4FD00175C,  
 803 <https://doi.org/10.1039/D4FD00175C>, 2024.

804 Isaacman-VanWertz, G., Massoli, P., O’Brien, R., Lim, C., Franklin, J. P., Moss, J. A., Hunter, J. F., Nowak, J. B.,  
 805 Canagaratna, M. R., Misztal, P. K., Arata, C., Roscioli, J. R., Herndon, S. T., Onasch, T. B., Lambe, A. T., Jayne, J. T., Su, L.,  
 806 Knopf, D. A., Goldstein, A. H., Worsnop, D. R., and Kroll, J. H.: Chemical evolution of atmospheric organic carbon over  
 807 multiple generations of oxidation, *Nature Chem*, 10, 462–468, <https://doi.org/10.1038/s41557-018-0002-2>, 2018.

808 Jang, M. and Kamens, R. M.: Characterization of Secondary Aerosol from the Photooxidation of Toluene in the Presence of  
 809 NO<sub>x</sub> and 1-Propene, *Environ. Sci. Technol.*, 35, 3626–3639, <https://doi.org/10.1021/es010676+>, 2001.



810 Jang, M., Czoschke, N. M., Lee, S., and Kamens, R. M.: Heterogeneous Atmospheric Aerosol Production by Acid-Catalyzed  
811 Particle-Phase Reactions, *Science, New Series*, 298, 814–817, <https://doi.org/10.1126/science.1075798>, 2002.

812 Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of the Master Chemical  
813 Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic compounds, Part B, 2003.

814 Jiang, Z., Grosselin, B., Daële, V., Mellouki, A., and Mu, Y.: Seasonal and diurnal variations of BTEX compounds in the  
815 semi-urban environment of Orleans, France, *Science of The Total Environment*, 574, 1659–1664,  
816 <https://doi.org/10.1016/j.scitotenv.2016.08.214>, 2017.

817 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe,  
818 H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson,  
819 K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M.,  
820 Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., E., Dunlea, J., Huffman, J. A., Onasch, T. B., Alfarra, M. R.,  
821 Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D.,  
822 Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel,  
823 J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C.  
824 E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525–1529,  
825 <https://doi.org/10.1126/science.1180353>, 2009.

826 Jin, Z.-H., Yu, D., Liu, Y.-X., Tian, Z.-Y., Richter, S., Braun-Unkhoff, M., Naumann, C., and Yang, J.-Z.: An experimental  
827 investigation of furfural oxidation and the development of a comprehensive combustion model, *Combustion and Flame*, 226,  
828 200–210, <https://doi.org/10.1016/j.combustflame.2020.12.015>, 2021.

829 John, E., Coburn, S., Liu, C., McAughey, J., Mariner, D., McAdam, K. G., Sebestyén, Z., Bakos, I., and Dóbé, S.: Effect of  
830 temperature and humidity on the gas–particle partitioning of nicotine in mainstream cigarette smoke: A diffusion denuder  
831 study, *Journal of Aerosol Science*, 117, 100–117, <https://doi.org/10.1016/j.jaerosci.2017.12.015>, 2018.

832 Kamens, R. M., Zhang, H., Chen, E. H., Zhou, Y., Parikh, H. M., Wilson, R. L., Galloway, K. E., and Rosen, E. P.: Secondary  
833 organic aerosol formation from toluene in an atmospheric hydrocarbon mixture: Water and particle seed effects, *Atmospheric*  
834 *Environment*, 45, 2324–2334, <https://doi.org/10.1016/j.atmosenv.2010.11.007>, 2011.

835 Kang, E., Root, M. J., Toohey, D. W., and Brune, W. H.: Introducing the concept of Potential Aerosol Mass (PAM), *Atmos.*  
836 *Chem. Phys.*, 2007.

837 Kautzman, K. E., Surratt, J. D., Chan, M. N., Chan, A. W. H., Hersey, S. P., Chhabra, P. S., Dalleska, N. F., Wennberg, P. O.,  
838 Flagan, R. C., and Seinfeld, J. H.: Chemical Composition of Gas- and Aerosol-Phase Products from the Photooxidation of  
839 Naphthalene, *J. Phys. Chem. A*, 114, 913–934, <https://doi.org/10.1021/jp908530s>, 2010.

840 Kim, D.-Y., Soda, S., Kendo, A., and Oh, J.-H.: Atmospheric Photochemistry in Low-and High-NO<sub>x</sub> Regimes, *Journal of*  
841 *Environmental Science International*, 16, 1–8, <https://doi.org/10.5322/JES.2007.16.1.001>, 2007.

842 Kleindienst, T. E., Jaoui, M., Lewandowski, M., Offenberg, J. H., and Docherty, K. S.: The formation of SOA and chemical  
843 tracer compounds from the photooxidation of naphthalene and its methyl analogs in the presence and absence of nitrogen  
844 oxides, *Atmos. Chem. Phys.*, 12, 8711–8726, <https://doi.org/10.5194/acp-12-8711-2012>, 2012.

845 Klodt, A. L., Aiona, P. K., MacMillan, A. C., Ji (Julie) Lee, H., Zhang, X., Helgestad, T., Novak, G. A., Lin, P., Laskin, J.,  
846 Laskin, A., Bertram, T. H., Cappa, C. D., and Nizkorodov, S. A.: Effect of relative humidity, NO<sub>x</sub>, and ammonia on the  
847 physical properties of naphthalene secondary organic aerosols, *Environ. Sci.: Atmos.*, 3, 991–1007,  
848 <https://doi.org/10.1039/D3EA00033H>, 2023.

849 Kostenidou, E., Marques, B., Temime-Roussel, B., Liu, Y., Vansevent, B., Sartelet, K., and D'Anna, B.: Secondary organic  
850 aerosol formed by Euro 5 gasoline vehicle emissions: chemical composition and gas-to-particle phase partitioning, *Atmos.*  
851 *Chem. Phys.*, 24, 2705–2729, <https://doi.org/10.5194/acp-24-2705-2024>, 2024.

852 Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in  
853 the atmosphere, *Atmospheric Environment*, 42, 3593–3624, <https://doi.org/10.1016/j.atmosenv.2008.01.003>, 2008.

854 Kroll, J. H., Donahue, N. M., Jimenez, J. L., Kessler, S. H., Canagaratna, M. R., Wilson, K. R., Altieri, K. E., Mazzoleni, L.  
855 R., Wozniak, A. S., Bluhm, H., Mysak, E. R., Smith, J. D., Kolb, C. E., and Worsnop, D. R.: Carbon oxidation state as a metric  
856 for describing the chemistry of atmospheric organic aerosol, *Nature Chem.*, 3, 133–139, <https://doi.org/10.1038/nchem.948>,  
857 2011.

858 La, Y. S., Camredon, M., Ziemann, P. J., Valorso, R., Matsunaga, A., Lannuque, V., Lee-Taylor, J., Hodzic, A., Madronich,  
859 S., and Aumont, B.: Impact of chamber wall loss of gaseous organic compounds on secondary organic aerosol formation:  
860 explicit modeling of SOA formation from alkane and alkene oxidation, *Atmos. Chem. Phys.*, 16, 1417–1431,  
861 <https://doi.org/10.5194/acp-16-1417-2016>, 2016.

862 Lambe, A. T., Chhabra, P. S., Onasch, T. B., Brune, W. H., Hunter, J. F., Kroll, J. H., Cummings, M. J., Brogan, J. F., Parmar,  
863 Y., Worsnop, D. R., Kolb, C. E., and Davidovits, P.: Effect of oxidant concentration, exposure time, and seed particles on  
864 secondary organic aerosol chemical composition and yield, *Atmos. Chem. Phys.*, 15, 3063–3075, <https://doi.org/10.5194/acp-15-3063-2015>, 2015.

866 Lamkaddam, H., Gratien, A., Pangui, E., Cazaunau, M., Picquet-Varrault, B., and Doussin, J.-F.: High-NO<sub>x</sub> Photooxidation  
867 of *n*-Dodecane: Temperature Dependence of SOA Formation, *Environ. Sci. Technol.*, 51, 192–201,  
868 <https://doi.org/10.1021/acs.est.6b03821>, 2017.

869 Langevin, P. (1872–1946): Une formule fondamentale de théorie cinétique, *Ann. Chim. et Physique*, 1905, 5, 245, 269–300,  
870 1950.

871 Languille, B., Gros, V., Petit, J.-E., Honoré, C., Baudic, A., Perrussel, O., Foret, G., Michoud, V., Truong, F., Bonnaire, N.,  
872 Sarda-Estève, R., Delmotte, M., Feron, A., Maisonneuve, F., Gaimoz, C., Formenti, P., Kotthaus, S., Haeffelin, M., and Favez,  
873 O.: Wood burning: A major source of Volatile Organic Compounds during wintertime in the Paris region, *Science of The Total*  
874 *Environment*, 711, 135055, <https://doi.org/10.1016/j.scitotenv.2019.135055>, 2020.

875 Lannuque, V. and Sartelet, K.: Development of a detailed gaseous oxidation scheme of naphthalene for secondary organic  
876 aerosol (SOA) formation and speciation, *Atmospheric Chemistry and Physics*, 24, 8589–8606, [https://doi.org/10.5194/acp-24-](https://doi.org/10.5194/acp-24-8589-2024)  
877 [8589-2024](https://doi.org/10.5194/acp-24-8589-2024), 2024.

878 Lannuque, V., Camredon, M., Couvidat, F., Hodzic, A., Valorso, R., Madronich, S., Bessagnet, B., and Aumont, B.:  
879 Exploration of the influence of environmental conditions on secondary organic aerosol formation and organic species  
880 properties using explicit simulations: development of the VBS-GECKO parameterization, *Atmos. Chem. Phys.*, 18, 13411–  
881 13428, <https://doi.org/10.5194/acp-18-13411-2018>, 2018.

882 Lannuque, V., D'Anna, B., Kostenidou, E., Couvidat, F., Martinez-Valiente, A., Eichler, P., Wisthaler, A., Müller, M.,  
883 Temime-Roussel, B., Valorso, R., and Sartelet, K.: Gas–particle partitioning of toluene oxidation products: an experimental  
884 and modeling study, *Atmos. Chem. Phys.*, 23, 15537–15560, <https://doi.org/10.5194/acp-23-15537-2023>, 2023.

885 Lee, J. Y. and Lane, D. A.: Unique products from the reaction of naphthalene with the hydroxyl radical, *Atmospheric*  
886 *Environment*, 43, 4886–4893, <https://doi.org/10.1016/j.atmosenv.2009.07.018>, 2009.

Leglise, J., Müller, M., Piel, F., Otto, T., and Wisthaler, A.: Bulk Organic Aerosol Analysis by Proton-Transfer-Reaction Mass Spectrometry: An Improved Methodology for the Determination of Total Organic Mass, O:C and H:C Elemental Ratios, and the Average Molecular Formula, *Anal. Chem.*, 91, 12619–12624, <https://doi.org/10.1021/acs.analchem.9b02949>, 2019.

Li, J., Wang, W., Li, K., Zhang, W., Peng, C., Zhou, L., Shi, B., Chen, Y., Liu, M., Li, H., and Ge, M.: Temperature effects on optical properties and chemical composition of secondary organic aerosol derived from *n*-dodecane, *Atmos. Chem. Phys.*, 20, 8123–8137, <https://doi.org/10.5194/acp-20-8123-2020>, 2020.

Li, K., Wang, W., Ge, M., Li, J., and Wang, D.: Optical properties of secondary organic aerosols generated by photooxidation of aromatic hydrocarbons, *Scientific Reports*, 4, 4922, <https://doi.org/10.1038/srep04922>, 2014.

Li, K., Li, J., Liggio, J., Wang, W., Ge, M., Liu, Q., Guo, Y., Tong, S., Li, J., Peng, C., Jing, B., Wang, D., and Fu, P.: Enhanced Light Scattering of Secondary Organic Aerosols by Multiphase Reactions, *Environ. Sci. Technol.*, 51, 1285–1292, <https://doi.org/10.1021/acs.est.6b03229>, 2017.

Li, K., Li, J., Wang, W., Li, J., Peng, C., Wang, D., and Ge, M.: Effects of Gas-Particle Partitioning on Refractive Index and Chemical Composition of *m*-Xylene Secondary Organic Aerosol, *J. Phys. Chem. A*, 122, 3250–3260, <https://doi.org/10.1021/acs.jpca.7b12792>, 2018.

Li, L., Thomsen, D., Wu, C., Priestley, M., Iversen, E. M., Tygesen Skonager, J., Luo, Y., Ehn, M., Roldin, P., Pedersen, H. B., Bilde, M., Glasius, M., and Hallquist, M.: Gas-to-Particle Partitioning of Products from Ozonolysis of  $\Delta^3$ -Carene and the Effect of Temperature and Relative Humidity, *J. Phys. Chem. A*, 128, 918–928, <https://doi.org/10.1021/acs.jpca.3c07316>, 2024.

Li, Y., Zhao, J., Gomez-Hernandez, M., Lavallee, M., Johnson, N. M., and Zhang, R.: Functionality-based formation of secondary organic aerosol from *m*-xylene photooxidation, *Atmos. Chem. Phys.*, 22, 9843–9857, <https://doi.org/10.5194/acp-22-9843-2022>, 2022.

Li, Z., Schwier, A. N., Sareen, N., and McNeill, V. F.: Reactive processing of formaldehyde and acetaldehyde in aqueous aerosol mimics: surface tension depression and secondary organic products, *Atmos. Chem. Phys.*, 11, 11617–11629, <https://doi.org/10.5194/acp-11-11617-2011>, 2011.

Li, Z., Tikkanen, O.-P., Buchholz, A., Hao, L., Kari, E., Yli-Juuti, T., and Virtanen, A.: Effect of Decreased Temperature on the Evaporation of  $\alpha$ -Pinene Secondary Organic Aerosol Particles, *ACS Earth Space Chem.*, 3, 2775–2785, <https://doi.org/10.1021/acsearthspacechem.9b00240>, 2019.

Liang, Y., Wernis, R. A., Kristensen, K., Kreisberg, N. M., Croteau, P. L., Herndon, S. C., Chan, A. W. H., Ng, N. L., and Goldstein, A. H.: Gas–particle partitioning of semivolatile organic compounds when wildfire smoke comes to town, *Atmos. Chem. Phys.*, 23, 12441–12454, <https://doi.org/10.5194/acp-23-12441-2023>, 2023.

Liu, J., Zhu, S., Guo, T., Jia, B., Xu, L., Chen, J., and Cheng, P.: Smog chamber study of secondary organic aerosol formation from gas- and particle-phase naphthalene ozonolysis, *Atmospheric Environment*, 294, 119490, <https://doi.org/10.1016/j.atmosenv.2022.119490>, 2023.

Liu, K., Hua, S., and Song, L.: PM<sub>2.5</sub> Exposure and Asthma Development: The Key Role of Oxidative Stress, *Oxidative Medicine and Cellular Longevity*, 2022, 1–12, <https://doi.org/10.1155/2022/3618806>, 2022a.

Liu, M. and Matsui, H.: Impacts of Climate Change on Particulate Matter (PM), in: *Handbook of Air Quality and Climate Change*, edited by: Akimoto, H. and Tanimoto, H., Springer Nature Singapore, Singapore, 1–18, [https://doi.org/10.1007/978-981-15-2527-8\\_39-1](https://doi.org/10.1007/978-981-15-2527-8_39-1), 2020.

925 Liu, M., Bi, J., and Ma, Z.: Visibility-Based PM<sub>2.5</sub> Concentrations in China: 1957–1964 and 1973–2014, *Environ. Sci.*  
926 *Technol.*, 51, 13161–13169, <https://doi.org/10.1021/acs.est.7b03468>, 2017.

927 Liu, Q., Huang, D. D., Lambe, A. T., Lou, S., Zeng, L., Wu, Y., Huang, C., Tao, S., Cheng, X., Chen, Q., Hoi, K. I., Wang,  
928 H., Mok, K. M., Huang, C., and Li, Y. J.: A Comprehensive Characterization of Empirical Parameterizations for OH Exposure  
929 in the Aerodyne Potential Aerosol Mass Oxidation Flow Reactor (PAM-OFR), <https://doi.org/10.5194/egusphere-2024-2721>,  
930 30 September 2024.

931 Liu, S., Wang, Y., Xu, X., and Wang, G.: Effects of NO<sub>2</sub> and RH on secondary organic aerosol formation and light absorption  
932 from OH oxidation of o-xylene, *Chemosphere*, 308, 136541, <https://doi.org/10.1016/j.chemosphere.2022.136541>, 2022b.

933 Liu, X., Day, D. A., Krechmer, J. E., Ziemann, P. J., and Jimenez, J. L.: Determining Activity Coefficients of SOA from  
934 Isothermal Evaporation in a Laboratory Chamber, *Environ. Sci. Technol. Lett.*, 8, 212–217,  
935 <https://doi.org/10.1021/acs.estlett.0c00888>, 2021.

936 Loeffler, K. W., Koehler, C. A., Paul, N. M., and De Haan, D. O.: Oligomer Formation in Evaporating Aqueous Glyoxal and  
937 Methyl Glyoxal Solutions, *Environ. Sci. Technol.*, 40, 6318–6323, <https://doi.org/10.1021/es060810w>, 2006.

938 Loza, C. L., Chhabra, P. S., Yee, L. D., Craven, J. S., Flagan, R. C., and Seinfeld, J. H.: Chemical aging of  
939 <i>m</i>-xylene secondary organic aerosol: laboratory chamber study, *Atmos. Chem. Phys.*, 12, 151–167,  
940 <https://doi.org/10.5194/acp-12-151-2012>, 2012.

941 Lu, H., Huang, Q., Li, J., Ying, Q., Wang, H., Guo, S., Qin, M., and Hu, J.: Simulation of Regional Secondary Organic Aerosol  
942 Formation From Monocyclic Aromatic Hydrocarbons Using a Near-Explicit Chemical Mechanism Constrained by Chamber  
943 Experiments, *JGR Atmospheres*, 129, e2023JD040690, <https://doi.org/10.1029/2023JD040690>, 2024.

944 Lu, R., Zhou, P., Ma, F., Zhao, Q., Peng, X., Chen, J., and Xie, H.-B.: Multi-generation oxidation mechanism of M-xylene:  
945 Unexpected implications for secondary organic aerosol formation, *Atmospheric Environment*, 327, 120511,  
946 <https://doi.org/10.1016/j.atmosenv.2024.120511>, 2024.

947 Mao, J., Ren, X., Brune, W. H., Olson, J. R., Crawford, J. H., Fried, A., Huey, L. G., Cohen, R. C., Heikes, B., Singh, H. B.,  
948 Blake, D. R., Sachse, G. W., Diskin, G. S., Hall, S. R., and Shetter, R. E.: Airborne measurement of OH reactivity during  
949 INTEX-B, *Atmos. Chem. Phys.*, 9, 163–173, <https://doi.org/10.5194/acp-9-163-2009>, 2009.

950 McWhinney, R. D., Zhou, S., and Abbatt, J. P. D.: Naphthalene SOA: redox activity and naphthoquinone gas–particle  
951 partitioning, *Atmos. Chem. Phys.*, 13, 9731–9744, <https://doi.org/10.5194/acp-13-9731-2013>, 2013.

952 Meng, X., Wu, Z., Chen, J., Qiu, Y., Zong, T., Song, M., Lee, J., and Hu, M.: Particle phase state and aerosol liquid water  
953 greatly impact secondary aerosol formation: insights into phase transition and its role in haze events, *Atmos. Chem. Phys.*, 24,  
954 2399–2414, <https://doi.org/10.5194/acp-24-2399-2024>, 2024.

955 Meredith, L. K., Ledford, S. M., Riemer, K., Geffre, P., Graves, K., Honeker, L. K., LeBauer, D., Tfaily, M. M., and Krechmer,  
956 J.: Automating methods for estimating metabolite volatility, *Front. Microbiol.*, 14, 1267234,  
957 <https://doi.org/10.3389/fmicb.2023.1267234>, 2023.

958 Molina, M. J., Zhang, R., Broekhuizen, K., Lei, W., Navarro, R., and Molina, L. T.: Experimental Study of Intermediates from  
959 OH-Initiated Reactions of Toluene, *J. Am. Chem. Soc.*, 121, 10225–10226, <https://doi.org/10.1021/ja992461u>, 1999.

Montero-Montoya, R., López-Vargas, R., and Arellano-Aguilar, O.: Volatile Organic Compounds in Air: Sources, Distribution, Exposure and Associated Illnesses in Children, *Annals of Global Health*, 84, 225–238, <https://doi.org/10.29024/aogh.910>, 2018.

Müller, M., Graus, M., Wisthaler, A., Hansel, A., Metzger, A., Dommen, J., and Baltensperger, U.: Analysis of high mass resolution PTR-TOF mass spectra from 1,3,5-trimethylbenzene (TMB) environmental chamber experiments, *Atmos. Chem. Phys.*, 12, 829–843, <https://doi.org/10.5194/acp-12-829-2012>, 2012.

Müller, M., Mikoviny, T., Jud, W., D’Anna, B., and Wisthaler, A.: A new software tool for the analysis of high resolution PTR-TOF mass spectra, *Chemometrics and Intelligent Laboratory Systems*, 127, 158–165, <https://doi.org/10.1016/j.chemolab.2013.06.011>, 2013.

Müller, M., Eichler, P., D’Anna, B., Tan, W., and Wisthaler, A.: Direct Sampling and Analysis of Atmospheric Particulate Organic Matter by Proton-Transfer-Reaction Mass Spectrometry, *Anal. Chem.*, 89, 10889–10897, <https://doi.org/10.1021/acs.analchem.7b02582>, 2017.

Muller, M., Eicher, P., D’Anna, B., Tan, W., and Wisthaler, A.: Direct Sampling and Analysis of Atmospheric Particulate Organic Matter by Proton-Transfer-Reaction Mass Spectrometry, *Analytical Chemistry*, 89, 10889–10897, <https://doi.org/10.1021/acs.analchem.7b02582>, 2017.

Murphy, B. N., Donahue, N. M., Fountoukis, C., Dall’Osto, M., O’Dowd, C., Kiendler-Scharr, A., and Pandis, S. N.: Functionalization and fragmentation during ambient organic aerosol aging: application of the 2-D volatility basis set to field studies, *Atmos. Chem. Phys.*, 12, 10797–10816, <https://doi.org/10.5194/acp-12-10797-2012>, 2012.

NARSTO. and Electric Power Research Institute.: An assessment of tropospheric ozone pollution: a North American perspective, EPRI, 2000.

Ng, N. L., Kroll, J. H., Chan, A. W. H., Chhabra, P. S., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from m-xylene, toluene, and benzene, *Atmos. Chem. Phys.*, 7, 3909–3922, <https://doi.org/10.5194/acp-7-3909-2007>, 2007.

Nie, W., Yan, C., Huang, D., Wang, Z., Liu, Y., Qiao, X., Guo, Y., Tian, L., Zheng, P., Xu, Z., Li, Y., Xu, Z., Qi, X., Sun, P., Wang, J., Zheng, F., Li, X., Yin, R., Dallenbach, K., Bianchi, F., Petäjä, T., Zhang, Y., Wang, M., Schervish, M., Wang, S., Qiao, L., Wang, Q., Zhou, M., Wang, H., Yu, C., Yao, D., Guo, H., Ye, P., Lee, S., Li, Y., Liu, Y., Chi, X., Kerminen, V., Ehn, M., Donahue, N., Wang, T., Huang, C., Kulmala, M., Worsnop, D., Jiang, J., and Ding, A.: Secondary organic aerosol formed by condensing anthropogenic vapours over China’s megacities, *NATURE GEOSCIENCE*, 15, 255–+, <https://doi.org/10.1038/s41561-022-00922-5>, 2022.

Nishino, N., Arey, J., and Atkinson, R.: Formation and Reactions of 2-Formylcinnamaldehyde in the OH Radical-Initiated Reaction of Naphthalene, *Environ. Sci. Technol.*, 43, 1349–1353, <https://doi.org/10.1021/es802477s>, 2009.

Nozière, B., Dziedzic, P., and Córdoba, A.: Inorganic ammonium salts and carbonate salts are efficient catalysts for aldol condensation in atmospheric aerosols, *Phys. Chem. Chem. Phys.*, 12, 3864, <https://doi.org/10.1039/b924443c>, 2010.

Odum, J. R., Hoffmann, T., Bowman, F., Collins, D., Flagan, R. C., and Seinfeld, J. H.: Gas/Particle Partitioning and Secondary Organic Aerosol Yields, *Environ. Sci. Technol.*, 30, 2580–2585, <https://doi.org/10.1021/es950943+>, 1996.

Odum, J. R., Jungkamp, T. P. W., Griffin, R. J., Forstner, H. J. L., Flagan, R. C., and Seinfeld, J. H.: Aromatics, Reformulated Gasoline, and Atmospheric Organic Aerosol Formation, *Environ. Sci. Technol.*, 31, 1890–1897, <https://doi.org/10.1021/es960535l>, 1997.

997 O'Meara, S., Booth, A. M., Barley, M. H., Topping, D., and McFiggans, G.: An assessment of vapour pressure estimation  
998 methods, *Phys. Chem. Chem. Phys.*, 16, 19453–19469, <https://doi.org/10.1039/C4CP00857J>, 2014.

999 Pan, S. and Wang, L.: Atmospheric Oxidation Mechanism of *m* -Xylene Initiated by OH Radical, *J. Phys. Chem. A*, 118,  
1000 10778–10787, <https://doi.org/10.1021/jp506815v>, 2014.

1001 Pang, X.: Biogenic volatile organic compound analyses by PTR-TOF-MS: Calibration, humidity effect and reduced electric  
1002 field dependency, *Journal of Environmental Sciences*, 32, 196–206, <https://doi.org/10.1016/j.jes.2015.01.013>, 2015.

1003 Pankow, J. F. and Asher, W. E.: SIMPOL.1: a simple group contribution method for predicting vapor pressures and enthalpies  
1004 of vaporization of multifunctional organic compounds, *Atmos. Chem. Phys.*, 8, 2773–2796, [https://doi.org/10.5194/acp-8-](https://doi.org/10.5194/acp-8-2773-2008)  
1005 2773-2008, 2008.

1006 Peng, Y., Wang, H., Gao, Y., Jing, S., Zhu, S., Huang, D., Hao, P., Lou, S., Cheng, T., Huang, C., and Zhang, X.: Real-time  
1007 measurement of phase partitioning of organic compounds using a proton-transfer-reaction time-of-flight mass spectrometer  
1008 coupled to aCHARON inlet, *ATMOSPHERIC MEASUREMENT TECHNIQUES*, 16, 15–28, [https://doi.org/10.5194/amt-](https://doi.org/10.5194/amt-16-15-2023)  
1009 16-15-2023, 2023.

1010 Peng, Z. and Jimenez, J. L.: Radical chemistry in oxidation flow reactors for atmospheric chemistry research, *Chem. Soc. Rev.*,  
1011 49, 2570–2616, <https://doi.org/10.1039/C9CS00766K>, 2020.

1012 Peräkylä, O., Riva, M., Heikkinen, L., Quéléver, L., Roldin, P., and Ehn, M.: Experimental investigation into the volatilities  
1013 of highly oxygenated organic molecules (HOMs), *Atmos. Chem. Phys.*, 20, 649–669, [https://doi.org/10.5194/acp-20-649-](https://doi.org/10.5194/acp-20-649-2020)  
1014 2020, 2020.

1015 Piel, F., Müller, M., Winkler, K., Skytte af Sättra, J., and Wisthaler, A.: Introducing the extended volatility range proton-  
1016 transfer-reaction mass spectrometer (EVR PTR-MS), *Atmos. Meas. Tech.*, 14, 1355–1363, [https://doi.org/10.5194/amt-14-](https://doi.org/10.5194/amt-14-1355-2021)  
1017 1355-2021, 2021.

1018 Price, D. J., Kacarab, M., Cocker, D. R., Purvis-Roberts, K. L., and Silva, P. J.: Effects of temperature on the formation of  
1019 secondary organic aerosol from amine precursors, *Aerosol Science and Technology*, 50, 1216–1226,  
1020 <https://doi.org/10.1080/02786826.2016.1236182>, 2016.

1021 Qi, L., Nakao, S., Tang, P., and Iii, D. R. C.: Temperature effect on physical and chemical properties of secondary organic  
1022 aerosol from *m*-xylene photooxidation, *Atmos. Chem. Phys.*, <https://doi.org/10.5194/acp-10-3847-2010>, 2010.

1023 Qi, X., Zhu, S., Zhu, C., Hu, J., Lou, S., Xu, L., Dong, J., and Cheng, P.: Smog chamber study of the effects of NO<sub>x</sub> and NH<sub>3</sub>  
1024 on the formation of secondary organic aerosols and optical properties from photo-oxidation of toluene, *Science of The Total*  
1025 *Environment*, 727, 138632, <https://doi.org/10.1016/j.scitotenv.2020.138632>, 2020.

1026 Qu, X., Zhang, Q., and Wang, W.: Mechanism for OH-initiated photooxidation of naphthalene in the presence of O<sub>2</sub> and NO<sub>x</sub>:  
1027 A DFT study, *Chemical Physics Letters*, 429, 77–85, <https://doi.org/10.1016/j.cplett.2006.08.036>, 2006.

1028 Riemer, K.: volcalc: Calculate Volatility of Chemical Compounds, Zenodo, <https://doi.org/10.5281/ZENODO.8015155>, 2023.

1029 Riva, M.: Caractérisation d'une nouvelle voie de formation d'aérosols organiques secondaires (AOS) dans l'atmosphère: Rôle  
1030 des précurseurs polyaromatiques, Université Bordeaux 1, 2013.

1031 Rutter, A. P. and Schauer, J. J.: The effect of temperature on the gas–particle partitioning of reactive mercury in atmospheric  
1032 aerosols, *Atmospheric Environment*, 41, 8647–8657, <https://doi.org/10.1016/j.atmosenv.2007.07.024>, 2007.

- Salvador, C. M., Chou, C. C.-K., Cheung, H.-C., Ho, T.-T., Tsai, C.-Y., Tsao, T.-M., Tsai, M.-J., and Su, T.-C.: Measurements of submicron organonitrate particles: Implications for the impacts of NO<sub>x</sub> pollution in a subtropical forest, *Atmospheric Research*, 245, 105080, <https://doi.org/10.1016/j.atmosres.2020.105080>, 2020.
- Sareen, N., Schwier, A. N., Shapiro, E. L., Mitroo, D., and McNeill, V. F.: Secondary organic material formed by methylglyoxal in aqueous aerosol mimics, *Atmos. Chem. Phys.*, 10, 997–1016, <https://doi.org/10.5194/acp-10-997-2010>, 2010.
- Sarrafzadeh, M., Wildt, J., Pullinen, I., Springer, M., Kleist, E., Tillmann, R., Schmitt, S. H., Wu, C., Mentel, T. F., Zhao, D., Hastie, D. R., and Kiendler-Scharr, A.: Impact of NO<sub>x</sub> and OH on secondary organic aerosol formation from  $\beta$ -pinene photooxidation, *Atmos. Chem. Phys.*, 16, 11237–11248, <https://doi.org/10.5194/acp-16-11237-2016>, 2016.
- Sasaki, J., Aschmann, S. M., Kwok, E. S. C., Atkinson, R., and Arey, J.: Products of the Gas-Phase OH and NO<sub>3</sub> Radical-Initiated Reactions of Naphthalene, *Environ. Sci. Technol.*, 31, 3173–3179, <https://doi.org/10.1021/es9701523>, 1997.
- Sato, K., Hatakeyama, S., and Imamura, T.: Secondary Organic Aerosol Formation during the Photooxidation of Toluene: NO<sub>x</sub> Dependence of Chemical Composition, *J. Phys. Chem. A*, 111, 9796–9808, <https://doi.org/10.1021/jp071419f>, 2007.
- Sato, K., Ikemori, F., Ramasamy, S., Iijima, A., Kumagai, K., Fushimi, A., Fujitani, Y., Chatani, S., Tanabe, K., Takami, A., Tago, H., Saito, Y., Saito, S., Hoshi, J., and Morino, Y.: Formation of secondary organic aerosol tracers from anthropogenic and biogenic volatile organic compounds under varied NO<sub>x</sub> and oxidant conditions, *Atmospheric Environment: X*, 14, 100169, <https://doi.org/10.1016/j.aeaoa.2022.100169>, 2022.
- Schwantes, R. H., Charan, S. M., Bates, K. H., Huang, Y., Nguyen, T. B., Mai, H., Kong, W., Flagan, R. C., and Seinfeld, J. H.: Low-volatility compounds contribute significantly to isoprene secondary organic aerosol (SOA) under high-NO<sub>x</sub> conditions, *Atmos. Chem. Phys.*, 19, 7255–7278, <https://doi.org/10.5194/acp-19-7255-2019>, 2019.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics : from air pollution to climate change*, Third edition., John Wiley & Sons, Inc. Hoboken, New Jersey, Hoboken, New Jersey, 2016.
- Sekimoto, K., Li, S.-M., Yuan, B., Koss, A., Coggon, M., Warneke, C., and De Gouw, J.: Calculation of the sensitivity of proton-transfer-reaction mass spectrometry (PTR-MS) for organic trace gases using molecular properties, *International Journal of Mass Spectrometry*, 421, 71–94, <https://doi.org/10.1016/j.ijms.2017.04.006>, 2017.
- Shapiro, E. L., Szprengiel, J., Sareen, N., Jen, C. N., Giordano, M. R., and McNeill, V. F.: Light-absorbing secondary organic material formed by glyoxal in aqueous aerosol mimics, *Atmos. Chem. Phys.*, 9, 2289–2300, <https://doi.org/10.5194/acp-9-2289-2009>, 2009.
- Shi, D., Liu, J., Wang, Y., Xu, L., Guo, T., Jia, B., and Cheng, P.: Secondary organic aerosol formation from cis-3-hexen-1-ol/NO<sub>x</sub> photo-oxidation: The roles of cis-3-hexen-1-ol concentration, illumination intensity, NO<sub>x</sub> and NH<sub>3</sub>, *Atmospheric Environment*, 278, 119090, <https://doi.org/10.1016/j.atmosenv.2022.119090>, 2022.
- Shrivastava, M., Cappa, C. D., Fan, J., Goldstein, A. H., Guenther, A. B., Jimenez, J. L., Kuang, C., Laskin, A., Martin, S. T., Ng, N. L., Petaja, T., Pierce, J. R., Rasch, P. J., Roldin, P., Seinfeld, J. H., Shilling, J., Smith, J. N., Thornton, J. A., Volkamer, R., Wang, J., Worsnop, D. R., Zaveri, R. A., Zelenyuk, A., and Zhang, Q.: Recent advances in understanding secondary organic aerosol: Implications for global climate forcing, *Reviews of Geophysics*, 55, 509–559, <https://doi.org/10.1002/2016RG000540>, 2017.

1070 Singh, K. and Tripathi, D.: Particulate Matter and Human Health, in: Environmental Health, edited by: Otsuki, T., IntechOpen,  
1071 <https://doi.org/10.5772/intechopen.100550>, 2021.

1072 Song, C., Na, K., and Cocker, D. R.: Impact of the Hydrocarbon to NO<sub>x</sub> Ratio on Secondary Organic Aerosol Formation,  
1073 Environ. Sci. Technol., 39, 3143–3149, <https://doi.org/10.1021/es0493244>, 2005.

1074 Song, C., Na, K., Warren, B., Malloy, Q., and Cocker, D. R.: Secondary Organic Aerosol Formation from *m*-Xylene in the  
1075 Absence of NO<sub>x</sub>, Environ. Sci. Technol., 41, 7409–7416, <https://doi.org/10.1021/es070429r>, 2007.

1076 Song, C., He, J., Wu, L., Jin, T., Chen, X., Li, R., Ren, P., Zhang, L., and Mao, H.: Health burden attributable to ambient  
1077 PM<sub>2.5</sub> in China, Environmental Pollution, 223, 575–586, <https://doi.org/10.1016/j.envpol.2017.01.060>, 2017.

1078 Srivastava, D., Vu, T. V., Tong, S., Shi, Z., and Harrison, R. M.: Formation of secondary organic aerosols from anthropogenic  
1079 precursors in laboratory studies, npj Clim Atmos Sci, 5, 22, <https://doi.org/10.1038/s41612-022-00238-6>, 2022.

1080 Srivastava, D., Li, W., Tong, S., Shi, Z., and Harrison, R. M.: Characterization of products formed from the oxidation of  
1081 toluene and *m*-xylene with varying NO<sub>x</sub> and OH exposure, Chemosphere, 334, 139002,  
1082 <https://doi.org/10.1016/j.chemosphere.2023.139002>, 2023.

1083 Stark, H., Yatavelli, R. L. N., Thompson, S. L., Kang, H., Krechmer, J. E., Kimmel, J. R., Palm, B. B., Hu, W., Hayes, P. L.,  
1084 Day, D. A., Campuzano-Jost, P., Canagaratna, M. R., Jayne, J. T., Worsnop, D. R., and Jimenez, J. L.: Impact of Thermal  
1085 Decomposition on Thermal Desorption Instruments: Advantage of Thermogram Analysis for Quantifying Volatility  
1086 Distributions of Organic Species, Environ. Sci. Technol., 51, 8491–8500, <https://doi.org/10.1021/acs.est.7b00160>, 2017.

1087 Su, T. and Chesnavich, W. J.: Parametrization of the ion–polar molecule collision rate constant by trajectory calculations, The  
1088 Journal of Chemical Physics, 76, 5183–5185, <https://doi.org/10.1063/1.442828>, 1982.

1089 Svendby, T. M., Lazaridis, M., and Tørseth, K.: Temperature dependent secondary organic aerosol formation from terpenes  
1090 and aromatics, J Atmos Chem, 59, 25–46, <https://doi.org/10.1007/s10874-007-9093-7>, 2008.

1091 Takekawa, H., Minoura, H., and Yamazaki, S.: Temperature dependence of secondary organic aerosol formation by photo-  
1092 oxidation of hydrocarbons, Atmospheric Environment, 37, 3413–3424, [https://doi.org/10.1016/S1352-2310\(03\)00359-5](https://doi.org/10.1016/S1352-2310(03)00359-5),  
1093 2003.

1094 Tani, A., Hayward, S., and Hewitt, C. N.: Measurement of monoterpenes and related compounds by proton transfer reaction-  
1095 mass spectrometry (PTR-MS), International Journal of Mass Spectrometry, 223–224, 561–578, [https://doi.org/10.1016/S1387-3806\(02\)00880-1](https://doi.org/10.1016/S1387-3806(02)00880-1), 2003.

1097 Thangavel, P., Park, D., and Lee, Y.-C.: Recent Insights into Particulate Matter (PM<sub>2.5</sub>)-Mediated Toxicity in Humans: An  
1098 Overview, IJERPH, 19, 7511, <https://doi.org/10.3390/ijerph19127511>, 2022.

1099 Tian, L., Huang, D. D., Wang, Q., Zhu, S., Wang, Q., Yan, C., Nie, W., Wang, Z., Qiao, L., Liu, Y., Qiao, X., Guo, Y., Zheng,  
1100 P., Jing, S., Lou, S., Wang, H., Yu, J. Z., Huang, C., and Li, Y. J.: Underestimated Contribution of Heavy Aromatics to  
1101 Secondary Organic Aerosol Revealed by Comparative Assessments Using New and Traditional Methods, ACS Earth Space  
1102 Chem., 7, 110–119, <https://doi.org/10.1021/acsearthspacechem.2c00252>, 2023.

1103 Tomaz, S.: Etude des composés polyaromatiques dans l’atmosphère: caractérisation moléculaire et processus réactionnels en  
1104 lien avec l’aérosol organique, Université de Lille, 2015.



1105 Virtanen, A., Joutsensaari, J., Koop, T., Kannosto, J., Yli-Pirilä, P., Leskinen, J., Mäkelä, J. M., Holopainen, J. K., Pöschl, U.,  
 1106 Kulmala, M., Worsnop, D. R., and Laaksonen, A.: An amorphous solid state of biogenic secondary organic aerosol particles,  
 1107 *Nature*, 467, 824–827, <https://doi.org/10.1038/nature09455>, 2010.

1108 Volkamer, R., Jimenez, J. L., San Martini, F., Dzepina, K., Zhang, Q., Salcedo, D., Molina, L. T., Worsnop, D. R., and Molina,  
 1109 M. J.: Secondary organic aerosol formation from anthropogenic air pollution: Rapid and higher than expected, *Geophysical*  
 1110 *Research Letters*, 33, 2006GL026899, <https://doi.org/10.1029/2006GL026899>, 2006.

1111 Wang, L., Atkinson, R., and Arey, J.: Dicarbonyl Products of the OH Radical-Initiated Reactions of Naphthalene and the C<sub>1</sub>  
 1112 - and C<sub>2</sub> -Alkyl naphthalenes, *Environ. Sci. Technol.*, 41, 2803–2810, <https://doi.org/10.1021/es0628102>, 2007.

1113 Warren, B., Austin, R. L., and Cocker, D. R.: Temperature dependence of secondary organic aerosol, *Atmospheric*  
 1114 *Environment*, 43, 3548–3555, <https://doi.org/10.1016/j.atmosenv.2009.04.011>, 2009.

1115 Wei, W., Mandin, C., Blanchard, O., Mercier, F., Pelletier, M., Le Bot, B., Glorennec, P., and Ramalho, O.: Temperature  
 1116 dependence of the particle/gas partition coefficient: An application to predict indoor gas-phase concentrations of semi-volatile  
 1117 organic compounds, *Science of The Total Environment*, 563–564, 506–512, <https://doi.org/10.1016/j.scitotenv.2016.04.106>,  
 1118 2016.

1119 Wu, K., Duan, M., Zhou, J., Zhou, Z., Tan, Q., Song, D., Lu, C., and Deng, Y.: Sources Profiles of Anthropogenic Volatile  
 1120 Organic Compounds from Typical Solvent Used in Chengdu, China, *J. Environ. Eng.*, 146, 05020006,  
 1121 [https://doi.org/10.1061/\(ASCE\)EE.1943-7870.0001739](https://doi.org/10.1061/(ASCE)EE.1943-7870.0001739), 2020.

1122 Xu, J., Griffin, R. J., Liu, Y., Nakao, S., and Cocker, D. R.: Simulated impact of NO<sub>x</sub> on SOA formation from oxidation of  
 1123 toluene and m-xylene, *Atmospheric Environment*, 101, 217–225, <https://doi.org/10.1016/j.atmosenv.2014.11.008>, 2015.

1124 Xu, L., Kollman, M. S., Song, C., Shilling, J. E., and Ng, N. L.: Effects of NO<sub>x</sub> on the Volatility of Secondary Organic Aerosol  
 1125 from Isoprene Photooxidation, *Environ. Sci. Technol.*, 48, 2253–2262, <https://doi.org/10.1021/es404842g>, 2014.

1126 Xuan, L., Ma, Y., Xing, Y., Meng, Q., Song, J., Chen, T., Wang, H., Wang, P., Zhang, Y., and Gao, P.: Source, temporal  
 1127 variation and health risk of volatile organic compounds (VOCs) from urban traffic in harbin, China, *Environmental Pollution*,  
 1128 270, 116074, <https://doi.org/10.1016/j.envpol.2020.116074>, 2021.

1129 Yamasaki, Hiroyasu., Kuwata, Kazuhiro., and Miyamoto, Hiroko.: Effects of ambient temperature on aspects of airborne  
 1130 polycyclic aromatic hydrocarbons, *Environ. Sci. Technol.*, 16, 189–194, <https://doi.org/10.1021/es00098a003>, 1982.

1131 Ye, F., Li, J., Gao, Y., Wang, H., An, J., Huang, C., Guo, S., Lu, K., Gong, K., Zhang, H., Qin, M., and Hu, J.: The role of  
 1132 naphthalene and its derivatives in the formation of secondary organic aerosol in the Yangtze River Delta region, China, *Atmos.*  
 1133 *Chem. Phys.*, 24, 7467–7479, <https://doi.org/10.5194/acp-24-7467-2024>, 2024.

1134 Zhang, H., Li, H., Zhang, Y., Wang, X., Bi, F., Meng, L., Li, Y., Zhao, L., Zhang, X., Peng, Z., Mu, Y., Mellouki, W., and  
 1135 Chai, F.: Synergistic generation mechanisms of SOA and ozone from the photochemical oxidation of 1,3,5-trimethylbenzene:  
 1136 Influence of precursors ratio, temperature and radiation intensity, *Atmospheric Research*, 293, 106924,  
 1137 <https://doi.org/10.1016/j.atmosres.2023.106924>, 2023.

1138 Zhang, J., Choi, M., Ji, Y., Zhang, R., Zhang, R., and Ying, Q.: Assessing the Uncertainties in Ozone and SOA Predictions  
 1139 due to Different Branching Ratios of the Cresol Pathway in the Toluene-OH Oxidation Mechanism, *ACS Earth Space Chem.*,  
 1140 5, 1958–1970, <https://doi.org/10.1021/acsearthspacechem.1c00092>, 2021.

1141 Zhang, P., Huang, J., Shu, J., and Yang, B.: Comparison of secondary organic aerosol (SOA) formation during o-, m-, and p-  
 1142 xylene photooxidation, *Environmental Pollution*, 245, 20–28, <https://doi.org/10.1016/j.envpol.2018.10.118>, 2019a.

1143 Zhang, Q., Xu, Y., and Jia, L.: Secondary organic aerosol formation from OH-initiated oxidation of  $\alpha$ -pinene: effects of relative humidity on yield and chemical composition, *Atmos. Chem. Phys.*, 19, 15007–15021,  
 1144 <https://doi.org/10.5194/acp-19-15007-2019>, 2019b.

1146 Zhao, D., Schmitt, S. H., Wang, M., Acir, I.-H., Tillmann, R., Tan, Z., Novelli, A., Fuchs, H., Pullinen, I., Wegener, R., Rohrer,  
 1147 F., Wildt, J., Kiendler-Scharr, A., Wahner, A., and Mentel, T. F.: Effects of NO<sub>x</sub> and SO<sub>2</sub> on the secondary organic aerosol  
 1148 formation from photooxidation of  $\alpha$ -pinene and limonene, *Atmos. Chem. Phys.*, 18, 1611–1628, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-18-1611-2018)  
 1149 18-1611-2018, 2018.

1150 Zhao, J., Zhang, R., Misawa, K., and Shibuya, K.: Experimental product study of the OH-initiated oxidation of m-xylene,  
 1151 *Journal of Photochemistry and Photobiology A: Chemistry*, 176, 199–207, <https://doi.org/10.1016/j.jphotochem.2005.07.013>,  
 1152 2005.

1153 Zhou, X.: The gas/particle partitioning behavior of phthalate esters in indoor environment: Effects of temperature and  
 1154 humidity, *Environmental Research*, 2021.

1155 Zhu, M., Huang, M., Xue, B., Cai, S., Hu, C., Zhao, W., Gu, X., and Zhang, W.: Chemical analysis of nitro-aromatic  
 1156 compounds of secondary organic aerosol formed from photooxidation of p-xylene with NO<sub>x</sub>, *J Chinese Chemical Soc*, 68,  
 1157 1697–1708, <https://doi.org/10.1002/jccs.202100105>, 2021.