

# Equatorial Ionospheric Plasma Bubbles During Intense geomagnetic storms of Solar Cycle 25

Nadia Imtiaz<sup>1, 2</sup>, Andres Calabia<sup>3, 4</sup>, Chukwuma Anoruo<sup>5</sup>, Aqsa Zahid<sup>6</sup>, Christine Amory-Mazaudier<sup>7</sup>, and Binod Adhikari<sup>8</sup>

<sup>1</sup>Theoretical Physics Division, PINSTECH, Pakistan

<sup>2</sup>School of Physics and Astronomy, University of Southampton, U.K.

<sup>3</sup>School of Surveying and Land Information Engineering, Henan Polytechnic University, Jiaozuo, China.

<sup>4</sup>Department of Physics and Mathematics, University of Alcalá, 28801-Alcalá de Henares (Madrid), Spain.

<sup>5</sup>Laboratório de Física e Astronomia, Universidade do Vale do Paraíba, Av. Shishima Hifumi, 2911–Urbanova, São José Dos Campos (SP), Brazil

<sup>6</sup>department of Physics, Quaid-i-Azam University, Pakistan

<sup>7</sup>Plasmas (LPP), CNRS, Sorbonne Université, Université Paris Saclay, Observatoire de Paris, Ecole polytechnique, Institut Polytechnique de Paris, Paris, France

<sup>8</sup>department of Physics, Patan M. Campus, Tribhuvan University, Lalitpur, Nepal

**Correspondence:** Nadia Imtiaz (nhussain@ualberta.ca)

**Abstract.** This study examines the low-latitude ionospheric response to four intense geomagnetic storms during Solar Cycle 25 (March, April, November 2023, and May 2024), focusing on Equatorial Ionization Anomaly (EIA) variations and post-sunset plasma irregularities. We use the Weimer (2005) model for Joule Heating (JH), Madrigal TEC maps, and GNSS-derived ROTI to analyze storm-time changes in EIA structure and equatorial plasma bubbles (EPBs). The May 2024 storm exhibited the strongest post-sunset JH, particularly near the June solstice, while March and April storms showed moderate JH and November the lowest. Equinox storms produced nearly symmetric JH patterns, whereas solstice storms revealed interhemispheric asymmetries. JH, together with storm-time electric fields and equatorial meridional winds, influences the location, strength, hemispheric asymmetry, and the generation or suppression of plasma irregularities of the EIA crest. Well-developed double-crest EIAs with steep density gradients favor post-sunset irregularities, while single-crest or merged EIAs are less favourable. Fluctuations in IMF  $B_z$  drive east-west prompt penetration electric fields that dynamically modulate the F-region, altering the plasma fountain effect, EIA structure, and post-sunset irregularity distribution. These results highlight the combined role of storm-driven electrodynamics and neutral winds in low-latitude ionospheric variability, improving understanding of EIA dynamics and plasma irregularity formation during geomagnetic storms.

## 15 1 INTRODUCTION

The study of ionospheric properties, particularly the equatorial ionosphere, is critical because of its influence on radio signal transmission, such as those from Global Navigation Satellite Systems (GNSS). The ionosphere, an ionized layer of the upper atmosphere extending from about 60 to 1000 km. It comprises three main regions—D (50–90 km), E (90–150 km), and F

(150–500 km)—whose boundaries vary with local time, season, and geomagnetic activity. During magnetic storms, thermo-  
spheric and ionospheric parameters such as neutral wind, composition, and electric fields deviate from their normal patterns,  
20 driving upper-atmosphere variability (Balan et al., 2018) and causing GNSS signal disturbances (Zhang et al., 2020). Geo-  
magnetic storms strongly impact low-latitude ionospheric phenomena such as the equatorial plasma fountain (EPF), equatorial  
ionization anomaly (EIA), and plasma irregularities (Kelly, 2009; Kassa et al., 2015). The EPF arises from the upward  $\mathbf{E} \times \mathbf{B}$   
drift that lifts plasma from the magnetic equator and its subsequent field-aligned diffusion. The resulting EIA exhibits enhanced  
25 electron density at about  $15^\circ$  magnetic latitude north and south, with a trough near the magnetic equator (Balan et al., 2018).  
Both the EPF and EIA are highly sensitive to variations in zonal electric fields during geomagnetic storms.

Storm-time electric fields can emerge from either a short-lived prompt penetration electric field (PPEF) or a disturbance dy-  
namo electric field (DDEF). The southward turning of the Interplanetary Magnetic Field  $B_z$  component causes a rapid increase  
30 in high-latitude convection and substorm activity. Convective electric fields at high latitudes can expand much faster than the  
magnetosphere can respond, enabling them to rapidly penetrate the low-latitude ionosphere. The PPEF polarity is east during  
the day and west during the night. When the IMF  $B_z$  component turns northward, the Region 2 currents intensify and create a  
shielding layer in response to the previously enhanced magnetospheric convection. Region 2 currents are field-aligned currents  
flowing at lower latitudes (equatorward of the auroral zone) that connect the inner magnetosphere to the ionosphere and play a  
35 key role in shielding the inner magnetosphere from external electric fields. This shielding layer is a configuration of currents  
and electric fields that normally prevents external magnetospheric electric fields from penetrating to low latitudes. However,  
when the Region 2 currents overcompensate, they produce an overshielding electric field with polarity opposite to the PPEF,  
which rapidly penetrates to low latitudes during the initial phase of enhanced convection when IMF  $B_z$  turns southward. This  
overshielding electric field penetrates into the low-latitude ionosphere (near the equator, roughly  $\pm 30^\circ$  magnetic latitude), and  
40 this overshielding penetration can be as efficient as the electric field penetration that occurs when the IMF  $B_z$  turns southward.  
The DDEF is a manifestation of disturbed thermospheric circulation and equatorial winds. High-latitude heating during storms  
improves meridional neutral winds, resulting in strong equatorial winds. At low latitudes, these disturbance winds generate a  
westward/eastward electric field on the dayside and nightside, contradicting their quiet-time tendencies. Compared to PPEF,  
DDEF takes hours to form and usually lasts several hours (Astafyeva et al., 2018; Blanc and Richmond, 1980; Huang et al.,  
45 2005; Kikuchi et al., 2008). Moreover, the vertical equatorial plasma drifts associated with PPEF and DDEF exhibit seasonal  
and longitudinal dependence (Fejer et al., 1999).

In addition to storm-time electric fields, equatorward thermospheric neutral winds transport plasma between hemispheres,  
modifying the low-latitude ionosphere. High-latitude storm energy input rapidly expands the neutral atmosphere, altering the  
50 global O/ $N_2$  ratio and, in turn, affecting EIA plasma densities (Li et al., 2018; Rishbeth, 2000; Kassa et al., 2015). Thermo-  
spheric circulation also drives EIA hemispheric asymmetry, enhancing the winter-hemisphere crest via trans-equatorial winds  
(Huang et al., 2005; Tulasi Ram et al., 2009). Seasonal and orbital effects, such as increased solar EUV near perihelion, can fur-  
ther raise northern EIA densities by 20% compared to the southern crest (Lin et al., 2005). Storm-time perturbations in electric

fields, neutral winds, and composition strongly influence the evolution of the EIA. PPEFs in the east and disturbed meridional winds can expand the EIA to mid-latitudes and induce significant density changes (Balan et al., 2009; Tsurutani et al., 2004). Although the EIA typically exhibits a double-crest structure, single or merged crests may develop under specific conditions. A stronger fountain effect produces extended, pronounced crests, whereas weaker fountains can suppress them, leading to a single crest, as observed during extremely low solar activity over 110°E (Huang et al., 2013). In contrast, merging of daytime EIA crests during storm recovery can result from neutral equatorward winds, with minor contributions from downward  $\mathbf{E} \times \mathbf{B}$  drift and increased O/N<sub>2</sub> ratios (Balan et al., 2013).

Plasma depletions, or Equatorial Plasma Bubbles (EPBs), are localized reductions in F-region plasma density caused by post-sunset Rayleigh-Taylor (R–T) instabilities, leading to significant radio signal disruptions. Most depletions occur in the EIA region (Cueto et al., 2012). Post-sunset plasma depletions, or EPBs, develop via R-T instabilities, with onset controlled by pre-reversal enhancement (PRE), terminator alignment, and seasonal factors, and are further enhanced during geomagnetic storms (Araujo-Pradere et al., 2006; Liu et al., 2010; Seba and Nigussie, 2016; Amaechi et al., 2018). Morphological and spectral studies show that the irregularities in the E-region are rod-shaped along the field lines, whereas the irregularities in the F-region extend along and across the field (Vaggu et al., 2023). Statistical analysis from Swarm observations reveal that EPB occurrence varies with longitude, season, local time, latitude, solar and geomagnetic activity, and is influenced by vertical  $\mathbf{E} \times \mathbf{B}$  drift, thermospheric winds, and atmospheric gravity waves (Aa et al., 2020; Fuller-Rowell et al., 1997). Overall, the complex interplay of electric fields, neutral winds, thermospheric composition, and geomagnetic activity governs the spatiotemporal variability of the low-latitude ionosphere, particularly the EIA and post-sunset plasma irregularities, highlighting the need for combined modeling and observational studies to understand ionospheric responses during geomagnetic storms.

In low-latitude and equatorial regions, intense geomagnetic storms can generate nighttime ionospheric plasma irregularities, ranging from centimeters to thousands of kilometers (Abdu et al., 2009). The Storm-time uplift of the F-layer enhances vertical drift  $\mathbf{E} \times \mathbf{B}$  and plasma density gradients, favoring R-T instability and post-sunset irregularity formation (González, 2022; Calabria et al., 2024; Imtiaz et al., 2024; Tahir et al., 2024; Kassa et al., 2024). Eastward (westward) storm-time electric fields promote (suppress) irregularities by raising (lowering) the F-region, with PRE providing additional uplift conducive to R–T instability (Aarons, 1991, 1997; Tulasi Ram et al., 2009). Vertical plasma transport, driven by PPEF and DDEF, is a primary mechanism for irregularity formation (Abdu, 2020), while suppression can result from westward PPEFs during northward IMF B<sub>z</sub> or storm-induced DDEFs (Huang, 2019; Kassa et al., 2023; Imtiaz et al., 2024; Kassa et al., 2024). For example, long-term eastward PPEFs can enhance upward ion drift by up to 180 m/s (Huang, 2019), and simulations and observations during the November 3–4, 2023 storm show that eastward/westward fields correspond to upward/downward ion drifts (Wu et al., 2024). Overall, the occurrence and characteristics of ionospheric irregularities depend on solar and geomagnetic activity, latitude, longitude, local time, and season (Abdu, 1997; Dugassa et al., 2019; Araujo-Pradere et al., 2006).

To better understand and model their variability, it is essential to analyze the occurrence characteristics of EPBs under varying geomagnetic conditions. In this study, we examine storm-time ionospheric conditions that influence EIA structures and the formation of post-sunset plasma irregularities or EPBs. Madrigal TEC maps are used to characterize EIA features, such as separated or merged crests, during the main phases of intense geomagnetic storms in Solar Cycle 25. Post-sunset ionospheric irregularities are analyzed using the GNSS-based Rate of Change TEC index (ROTI), while PPEF observations are employed to assess the role of low-latitude electric fields in shaping EIA structures favourable for R–T instability. The paper is structured as follows: Section 2 describes the data and analysis methods, Section 3 presents the results and interpretations, and Section 4 summarizes the findings and conclusions.

## 2 DATASETS and MODELS

We employ 1-minute and hourly data of solar wind parameters such as the IMF  $B_z$ , the solar wind speed ( $V_{sw}$ ), the eastward zonal component  $E_y$  of the interplanetary electric field (IEF), the ring current proxy index SYM-H, the geomagnetic activity index (Kp), and the solar radio flux at 10.7 cm (F10.7). The data is freely available at [https://omniweb.gsfc.nasa.gov/form/omni\\_min.html](https://omniweb.gsfc.nasa.gov/form/omni_min.html) in the NASA OMNIWeb database. The SYM-H index is preferred over Dst due to its 1-minute cadence, allowing better resolution of rapid geomagnetic variations. While both indices represent the symmetric component of the ring current, the higher temporal resolution of SYM-H enables a more accurate depiction of the rapid variations and short-term dynamics of geomagnetic variations. Information concerning the sudden storm commencement (SSC), the end of the main phase (MPE), the minimum values of the SYM-H index, the number of Sun spots (R) and the magnetic quiet days (Q) during different geomagnetic storms is also given in Table 1.

High-latitude ionospheric electrodynamics are analyzed using the CCMC-hosted Weimer-2005 model, driven by ACE solar wind data (Weimer, 2005). This empirical model estimates magnetic field-aligned currents (FACs) and electric potentials as functions of solar wind proxies. FACs and potentials, expressed via Fourier series of longitude at discrete latitudes, allow calculation of Joule Heating (JH) and total Poynting flux. Conductivity variations are implicitly included, enabling derivation of ionospheric currents and JH without a separate conductivity model.

Worldwide TEC maps with  $1^\circ \times 1^\circ$  resolution, updated every 5 minutes, are available from the CEDAR Madrigal database using ground-based GNSS signals (<http://cedar.openmadrigal.org/>) (Rideout and Coster, 2006). Ionospheric irregularities are quantified using the Rate of TEC Index (ROTI), derived from over 9,300 GNSS phase observations at 30-second intervals and averaged into 5-minute values at ionospheric pierce points (Aarons, 1997; Basu et al., 1999; Pi et al., 1997). 2D ROTI maps have a  $0.25^\circ \times 0.25^\circ$  grid, with  $ROTI \geq 0.5$  TECU/min indicating disturbances and  $ROTI \geq 1.0$  TECU/min marking significant irregularities. GNSS-TEC software by Seemala (2011) solves for receiver biases and estimates the TEC from satellites with elevation angles  $\geq 30^\circ$  to minimize multi-path errors. Given TEC data at discrete intervals, the rate of change of TEC (ROT)

**Table 1.** Information concerning the Sudden Storm Commencement (SSC), the MPEs, the minimum values of the SYM-H index, the number of Sun spots (R), and Quiet days (Q) during different geomagnetic storms of the solar cycle 25.

Event	SSC	MPE	SYM-H <sub>min</sub>	R	Q days
March 23-25, 2023	March 23 at 6:32 UT	March 24 at 5:21 UT	-170 nT	150	March 17-19
April 23-25, 2023	April 23 at 8:56 UT	April 24 at 4:03 UT	-233 nT	80	April 20-22
November 4-6, 2023	November 4 at 17:40 UT	November 5 at 16:54 UT	-189 nT	91	November 1-3
May 10-13, 2024	May 10 at 17:15 UT	May 11 at 02:14 UT	-518 nT	200	May 7-9

The information given in this table is obtained from OMNI web database.

120 can be computed as:

$$\text{ROT}(t_i) = \frac{d\text{TEC}}{dt} = \frac{\text{TEC}(t_{i+1}) - \text{TEC}(t_i)}{t_{i+1} - t_i}. \quad (1)$$

In this equation, dTEC is the difference between TEC( $t_{i+1}$ ) and TEC( $t_i$ ) values at times  $t_{i+1}$  and  $t_i$ , respectively. ROTI is then calculated as the standard deviation of ROT over a specified period, usually 5 minutes,

$$\text{ROTI} = \sqrt{\langle \text{ROT}^2 \rangle - \langle \text{ROT} \rangle^2}, \quad (2)$$

125 Table 2 shows the geographic and geomagnetic coordinates of equatorial and low-latitude GNSS receivers in the Asian, African and American sectors, which are used to examine the ROTI temporal patterns.

**Table 2.** Location of GNSS stations used in the analysis. Geographic and geomagnetic coordinates are given in degrees.

Station	Sector	Geographic (Lat, Lon)	Geomagnetic (Lat, Lon)
BAKO	Asia	6.49° S, 106.85° E	16.13° S, 179.93° W
DGAR	Asia	6.49° S, 72.37° E	14.94° S, 143.93° W
IISC	Asia	13.02° N, 77.57° E	4.87° N, 150.96° W
CUSV	Asia	13.74° N, 100.53° E	4.39° N, 173.37° W
HKSL	Asia	22.37° N, 113.93° E	13.19° N, 173.15° W
RABT	Africa	34.1° N, 6.50° W	23.88° N, 69.23° E
YKRO	Africa	6.86° S, 5.24° W	10.92° N, 68.67° E
ABPO	Africa	19.02° S, 47.23° E	23.49° S, 116.71° E
BOGT	America	4.64° N, 74.08° W	14.31° N, 1.46° W
RIOP	America	1.65° S, 78.65° W	7.23° N, 6.46° W
CORD	America	31.53° S, 64.47° W	22.43° S, 7.27° E
SANT	America	33.15° S, 70.67° W	23.77° S, 1.83° E

Table 3 also includes information on the Local Times (LTs) of SSCs and MPEs for the locations of these GNSS receivers. For the categorization of ionospheric irregularities, we used ROTI values as follows: a ROTI value lower than 0.25 TECU/min indicates the absence of irregularities; a weak irregularity is identified when ROTI ranges from 0.25 to 0.5 TECU/min; a

**Table 3.** Information concerning the Local Time (LT) of SSC and the MPEs at GNSS stations used in the analysis.

Station	March 23-25, 2023	April 23-25, 2023	November 4-6, 2023	May 10-13, 2024	UT to LT Conversion
	SSC, MPE (LT)	SSC, MPE (LT)	SSC, MPE (LT)	SSC, MPE (LT)	
BAKO	13:32, 12:21	15:56, 11:03	00:40, 23:54	00:15, 09:14	UT+7
	March 23, March 24	April 23, April 24	November 5	May 11	
DGAR	11:32, 10:21	13:56, 09:03	22:40, 21:54	22:15, 07:14	UT+5
	March 23, March 24	April 23, April 24	November 4, November 5	May 10, May 11	
IISC	11:32, 10:21	13:56, 09:03	22:40, 21:54	22:15, 07:14	UT+5
	March 23, March 24	April 23, April 24	November 4, November 5	May 10, May 11	
CUSV	13:32, 12:21	15:56, 11:03	00:40, 23:54	00:15, 09:14	UT+7
	March 23, March 24	April 23, April 24	November 5	May 11	
HKSL	13:32, 12:21	15:56, 11:03	00:40, 23:54	00:15, 09:14	UT+7
	March 23, March 24	April 23, April 24	November 5	May 11	
RABT	06:32, 05:21	08:56, 04:03	17:40, 16:54	17:15, 02:14	UT+0
	March 23, March 24	April 23, April 24	November 4, November 5	May 10, May 11	
YKRO	06:32, 05:21	08:56, 04:03	17:40, 16:54	17:15, 02:14	UT+0
	March 23, March 24	April 23, April 24	November 4, November 5	May 10, May 11	
ABPO	09:32, 08:21	11:56, 07:03	20:40, 19:54	20:15, 05:14	UT+3
	March 23, March 24	April 23, April 24	November 4, November 5	May 10, May 11	
BOGT	09:32, 00:21	11:56, 23:03	20:40, 11:54	12:15, 21:14	UT-5
	March 23, March 24	April 23, April 24	November 4, November 5	May 10, May 11	
RIOP	01:32, 00:21	03:56, 23:03	12:40, 11:54	12:15, 21:14	UT-5
	March 23, March 24	April 23	November 4, November 5	May 10	
CORD	02:32, 01:21	04:56, 00:03	13:40, 12:54	13:15, 22:14	UT-4
	March 23, March 24	April 23, April 24	November 4, November 5	May 10	
SANT	01:32, 00:21	03:56, 23:03	12:40, 11:54	12:15, 21:14	UT-5
	March 23, March 24	April 23	November 4, November 5	May 10	

130 moderate irregularity is noted when ROTI falls between 0.5 and 1 TECU/min; and a strong irregularity is present when ROTI exceeds 1 TECU/min.

During geomagnetic storms, the zonal electric field can be expressed as  $QP(t) = Q(t) + PPEF(t)$ , where  $Q(t)$  is the quiet-time field and  $PPEF(t)$  is the prompt-penetration electric field. Depending on IMF  $B_z$  polarity and local time, PPEFs can enhance or reverse the eastward daytime field, producing significant storm-time variability. The temporal profile of the equatorial electric field at a given longitude is obtained from the real-time PPEF model (<https://geomag.colorado.edu/online-calculators/real-time-model-ionospheric-electric-fields>), which uses solar wind data and a climatological quiet-time model to compute the

total eastward electric field QP), i.e., the sum of background quiet-time electric field (Q) and the PPEF, as a function of time and longitude.


## 140 3 RESULTS AND DISCUSSION

### 3.1 SOLAR AND MAGNETOSPHERE CONDITIONS

The investigation of the ionospheric response to geomagnetic storms reveals insight into the complex interplay between solar wind and Earth's atmosphere. Here, we investigate the four geomagnetic storms that occurred during the ascending phase of Solar Cycle 25. Figures 1-4 show (a) the  $B_z$  and  $V_{sw}$ , (b) the  $E_y$  and SYM-H, (c) the AE and AU, and (d) Kp and F10.7 during  
145 the storm events considered. In these figures, the SSC and MPE are indicated by vertical red lines, while the yellow rectangles highlight the time interval from 18:00 to 00:00 UT. Following is a brief overview of the geomagnetic conditions during these storms:

**March 23-25, 2023 storm:** As shown in Figure 1a, the IMF  $B_z$  shows two large southward turns throughout the main phase of the storm. The first southward turn occurs on March 23, at approximately 6:32 UT (Universal Time). Between 6:32  
150 and 12:00 UT, the value of IMF  $B_z$  decreased from 2 to -10 nT. The second long-term southward turn of the IMF  $B_z$  is seen on March 23, at 17:22 UT. During this turn, the IMF  $B_z$  decreased from 10 nT to approximately -18 nT. The IMF  $B_z$  remains south for approximately 9 hours before increasing approximately at 5:21 UT on March 24. During the main phase of this storm, the  $V_{sw}$  fluctuated between 400 and 500 km/s. During the recovery phase of this storm,  $V_{sw}$  increased and reached maximum value of 600 km/s on March 25. A northward IMF  $B_z$  generates an electric field westward during the day and eastward at  
155 night. Figure 1 b shows IEF  $E_y$  reached approximately 4 mV/m after the first southward excursion of the IMF  $B_z$ . From 12:00 to 18:00 UT, on March 23, it fluctuated rapidly between +5 and -5 mV/m. During the second southward excursion, it again increases and reached the maximum value of 8.22 mV/m at 17:22 on March 23. After that, it started decreasing to normal conditions. Figure 1b shows main phase of this storm started soon after SYM-H reached a value of 28 nT at the time of SSC. During the extended southward turning of IMF  $B_z$  ( $B_z < 0$ ), enhanced ring current caused by magnetic reconnection dropped  
160 the SYM-H index to -170 nT on March 24, at 5:21 UT (end of the main phase). The SYM-H increases during the recovery phase until it reaches quiet time value. Figure 1 d shows that before SSC, Kp was lower than 4+. On March 24, approximately at 04:00 UT, the Kp value peaked at 8+ and then began to decrease as the storm recovery began approximately at 5:21 UT. The F10.7 index fluctuated between 157.7 and 159.4 sfu indicating moderate solar activity during this storm.


**April 23-25, 2023 storm:** Figure 2(a-d) shows the solar wind and geomagnetic conditions during the period April  
165 22-26, 2023. The interval between the vertical red lines indicates the main phase of the storm, starting at SSC on April 23 at 8:56 UT. At the time of the SSC, the IMF  $B_z$  is at 0 nT which turned southward, and then later reached the value of -20 nT between 18:00 UT to 21:00 UT. During this time,  $V_{sw}$  increased from 400 km/s and reached the maximum value of 751 km/s at 21:39 UT on April 23. After 3 hours, the IMF  $B_z$  turned northward and fluctuated between 0 nT and 20 nT until 01:20 UT  
170 before a second strong southward excursion on April 24. This second long-term southward turn of IMF  $B_z$  is seen on April



**Figure 1.** Temporal variation of: (a) solar wind speed ( $V_{sw}$ ) and  $B_z$  component of IMF, (b)  $E_y$  component of IMF and SYM-H, (c) the auroral electrojet indices AE and AU, and (d) the magnetic index ( $K_p$ ) during March 23-25, 2023 storm.

24 at 1:21 UT. During this turn, a sharp decrease in the IMF  $B_z$  value from 20 nT to approximately -34 nT is observed in Figure 2a. The IMF  $B_z$  remains southward for approximately 3 hours until the end of the main phase at 4:03 UT on April 24. During this period,  $V_{sw}$  remained at approximately 600 km/s until the main phase ended at 4:03. Figure 2b shows a two-step geomagnetic storm, with SYM-H minima values of -179 nT (at 21:59 UT) and -233 nT (at 04:03 UT) on April 23 and 175 24, respectively. During the main phase, the IEF  $E_y$  oscillates rapidly between -21 and 21 mV/m. During the solar wind-magnetosphere interaction process, the  $K_p$  index increased to 8.3, indicating intense geomagnetic storm conditions. Following the strong geomagnetic storm, the  $K_p$  index dropped from 8.3 to 6.7, then surged to 8, before returning to a calm state. During this time, the solar activity level F10.7 remained steady at approximately 135 sfu as shown in Figure 2d. The recovery phase began with the northward turning of the IMF  $B_z$  and returned to 0 nT after 24 h. During recovery phase,  $V_{sw}$  also gradually 180 decreased to normal conditions.


**November 4-7, 2023 storm:** The main phase of this storm began with a southward turning of the IMF  $B_z$ , reaching a lowest value of approximately -24.9 nT at 11:31 UT on November 5. Then at 12:00 UT it turned northward and reached maximum value of approximately 40.1 nT at 13:56 UT on the same day. Afterwards, the IMF  $B_z$  shows a second long-term southward turn reaching -20 nT until the end of the main phase at 16:54 UT. The IMF  $B_z$  increases gradually to attain normal values on November 5. During the main phase, the  $V_{sw}$  is at moderate level at approximately 500 km/s and it increased to 600 185 km/s at 12:00 UT on November 6. Figure 3 (b) shows the IEF  $E_y$  and SYM-H, correspondingly. During the compression phase, the SYM-H sharply increased to 23 nT at 17:09 UT on November 4 and shortly after this the main phase of the storm begins



**Figure 2.** Temporal variation of: (a) solar wind speed ( $V_{sw}$ ) and  $B_z$  component of IMF, (b)  $E_y$  component of IMF and SYM-H, (c) the auroral electrojet indices AE and AU, and (d) the magnetic index ( $K_p$ ) and the solar radio flux at 10.7 cm (F10.7) during April 23-25, 2023 storm.

with decrease in SYM-H. It was also a double step storm with minimum SYM-H value of approximately -100 nT (at 12:00 UT) and -189 nT (at 16:54 UT) observed on November 5. Following the polarity of the IMF  $B_z$ , the IEF  $E_y$  also showed fluctuations between -19.5 and 10.5 mV/m. Before the SSC,  $K_p$  was lower than 3+, indicating quiet conditions, but as the storm begins, it rapidly increased to 5+ due to the initial effect of solar wind shock at 17:40 UT on November 4. From 00:00 to 08:00 UT on November 5, the  $K_p$  decreased to normal levels before rapidly increasing to 7+, predicting an intense geomagnetic storm between 12:00 UT and 19:00 UT on November 5. The solar radio flux remained in the range  $140 < F10.7 < 160$  sfu during this period. The recovery phase started at 16:54 UT with a northward turning of the IMF  $B_z$  on November 5.

**195 May 10-13, 2024 storm:** Figure 4a displays the temporal changes in  $V_{sw}$  and the IMF  $B_z$  during May 10 to 13, 2024. The main phase began approximately at 17:15 UT on May 10, with a sharp southward swing in the IMF  $B_z$ , reaching -30 nT, and an increase in  $V_{sw}$ , reaching approximately 700 km/s. Figure 4b shows variations in SYM-H and IEF  $E_y$  between May 10 and 13. Initially, the SYM-H index increases sharply to 88 nT as the Earth's magnetosphere compresses approximately at 17:15 UT on May 10. Following this initial increase, the main phase begins with a reduction of SYM-H due to increased ring current in the magnetosphere. The main phase lasted until 02:14 UT on May 11, when SYM-H index reached a maximum low of roughly -518 nT. During this phase, the IEF  $E_y$  fluctuates rapidly between -37 and 35 mV/m. The  $K_p$  index indicated substantial geomagnetic activity on May 10 between 12 and 15 UT, reaching a value of 5. shortly after the SSC,  $K_p$  reached to 8 and then began fluctuating between 8 and 9 until the main phase ended at 02:14 UT on May 11. Following storm recovery, the IMF  $B_z$  shows rapid North-South oscillations while the SYM-H increases gradually to reach normal level. During this




**Figure 3.** Temporal variation of parameters shown as: (a) solar wind speed ( $V_{sw}$ ) and  $B_z$  component of IMF, (b)  $E_y$  component of IMF and SYM-H, (c) the auroral electrojet indices AE and AU, and (d) the magnetic index ( $K_p$ ) and the solar radio flux at 10.7 cm (F10.7) during November 4-7, 2023.

205 phase,  $V_{sw}$  increased to 998 km/s and the  $K_p$  index varied between 7 and 9 until late evening on May 11. The F10.7 remained at approximately 220 and 226 sfu indicating strong solar activity during this storm as shown in Figure 4d.

### 3.2 High-Latitude Ionosphere

During geomagnetic storms, enhanced convective electric fields and energetic particle injections increase Pedersen currents and ionospheric conductivity (Buonsanto, 1999; Lukianova, 2020; Dungey, 1961; Axford, 1964). This intensifies Joule heating (JH), particularly in the auroral regions where particle precipitation occurs. The resulting thermal expansion of the neutral atmosphere drives equatorward winds and generates traveling atmospheric and ionospheric disturbances (Richmond and Matsushita, 1975; Fuller-Rowell et al., 1994). The solar wind energy dissipated into the magnetosphere–ionosphere system through JH was estimated for four major geomagnetic storms using the Weimer (2005) model hosted at CCMC. Under quiet conditions, JH ranges from 1 to 10  $\text{mW}, \text{m}^{-2}$ , but it increases substantially during disturbed periods. Table 5 and Fig. 5 present JH values 210 between 18:00 and 00:00 UT during the main phase of each event.

The March 2023 storm ( $B_z = -20$  nT,  $V_{sw} = 600$   $\text{km s}^{-1}$ ) produced symmetric JH peaks of  $\sim 57$   $\text{mW}, \text{m}^{-2}$ . The April 2023 two-step storm (SYM-H = -179, -233 nT) exhibited strong interhemispheric asymmetry with a northern peak near 97  $\text{mW}, \text{m}^{-2}$ . The November 2023 storm showed weaker, nearly symmetric heating, while the May 2024 event ( $B_z = -30$  nT,  $V_{sw} = 700$   $\text{km s}^{-1}$ ) reached 115  $\text{mW}, \text{m}^{-2}$  at 22:00 UT, indicating extreme energy deposition. The relative intensity of post-



**Figure 4.** Temporal variation of: (a) solar wind speed ( $V_{sw}$ ) and  $B_z$  component of IMF, (b)  $E_y$  component of IMF and SYM-H, (c) the auroral electrojet indices AE and AU, and (d) the magnetic index ( $K_p$ ) and the solar radio flux at 10.7 cm (F10.7) during May 10-13, 2024.

220 sunset JH follows: May > March > April > November. Hemispheric asymmetries in JH arise from differences in conductivity, magnetic dipole tilt, and IMF orientation, particularly  $B_y$  (Fuller-Rowell et al., 1997; Smith et al., 2023). Enhanced auroral JH modifies global thermospheric circulation (Blanc and Richmond, 1980), redistributing mass density and influencing equatorial plasma transport, thereby contributing to the modulation of post-sunset ionospheric irregularities (Jin et al., 2022).


### 3.3 Low-Latitude Ionosphere Response

225 During magnetic storms, the low latitude ionosphere exhibits different features due to the complicated interplay of geomagnetic activity, solar radiation, and ionospheric dynamics. Some of these features are discussed here:

#### 3.3.1 Structure of Equatorial Ionization Anomaly (EIA)

JH has a considerable and primarily indirect effect on the post-sunset EIA during the main phase of geomagnetic storms, which is mainly forced by storm-time neutral winds, thermospheric upwelling, and electric field modulation. The strong JH causes an equatorward wind surge, which uses neutral drag to drive plasma along magnetic field lines to higher altitudes. This decreases the downward plasma diffusion and elevates ionospheric heights, reducing chemical loss and supporting positive ionospheric storm effects, which strengthen the EIA intensity. However, the EIA's structure typically undergoes the most significant changes in the evening, at the local sunset. After sunset, the equatorial ionosphere experiences PRE and vertical plasma drifts because of enormous Cowling conductivities and rapid changes in zonal winds. The combination of PPEF and PRE can

230



**Figure 5.** Polar maps of Joule heating from the Weimer 2005 Model computed at CCMC during the main phase of the four geomagnetic storms after sunset, from 18:00 UT to 00:00 UT.

235 significantly increase ambipolar diffusion and upward  $\mathbf{E} \times \mathbf{B}$  drift, resulting in poleward extension of EIA crests. In contrast to poleward expansion, geomagnetically quiet or disturbed conditions can lead EIA crests to merge into a single peak across the geomagnetic equator. The electrodynamics, vertical plasma drift, neutral winds, and an increase in the low-latitude O/N2 ratio all contribute to the structuring of EIA crests (Balan et al., 2018; Luan, 2021).


240 Figure 6 (1-8) show the temporal variation of the ionospheric TEC during the main phase of the March 23-24, 2023 geomagnetic storm after sunset, from 18:00 UT to 00:00 UT. During the March 23-25, 2023 storm, TEC values showed notable inter-hemispheric asymmetries, with higher values in the southern hemisphere during the main phase. Initially, EIA showed well-separated crests, but between 18:00 and 20:00 UT, a significant increase in ionization can be observed in both the crest and trough regions. This behavior differs from the normal EIA structure, in which the trough near the magnetic equator usually  
 245 has lower ionization than the crests. The deviation from the usual fountain effect is most likely caused by the combined effects

of storm-time electric fields and neutral wind dynamics, resulting in unusual plasma redistribution (Luan, 2021). At 22:00 UT, the TEC decreases in the equatorial zone, leading to the formation of the double crests in the EIA. On March 24, from 00:00 UT to 06:00 UT, high TEC values were observed at equatorial and low latitudes, indicating geomagnetic disturbances towards the Asian sector. At 12:00 UT, the double crest in the northern hemisphere became a single crest and the TEC values slightly  
250 decreased. Starting from 14:00 UT, ionospheric TEC values began to increase again, marking the beginning of a second phase of ionospheric irregularities. The increase in TEC values towards the northern hemisphere indicated a period of geomagnetic disturbance until 22:00 UT. The decreasing TEC values are observed during the recovery phase of the geomagnetic storm from 06:00 UT onward (See supporting Figure of madrigal TEC maps for March 23-24, 2023).

Figure 6 (9-14) shows the temporal evolution of ionospheric TEC during the strong geomagnetic storm of April 23-24, 2023.  
255 During the main phase of this storm, a well-defined crest EIA can be observed on April 23, with asymmetric EIA over the pre-midnight sector (America) and expanding poleward at approximately  $\pm 25^\circ$  geomagnetic latitude (Aa et al., 2024). The observed EIA asymmetry in this sector is due to strong southward trans-equatorial wind surges induced by storm-time JH. For example, in the northern hemisphere, equatorial (southward) meridional winds restrict plasma downward movement, resulting in plasma concentration at higher altitudes and strong EIA crests. In contrast, poleward (southward) meridional winds increase  
260 plasma diffusion and prevent plasma accumulation at high altitudes in the southern hemisphere, causing EIA crests to gradually dissipate. After achieving the peak value for the day, the TEC begins to decrease from 22:00 UT forward. The EIA's strength and latitudinal range had significantly decreased, as evidenced by the global ionospheric TEC distribution on April 24. The main phase of this geomagnetic storm terminates at 04:00 UT, with the EIA's double crest over the southern hemisphere merging to a single crest (See the supporting Figure of madrigal TEC maps for April 23-24, 2023). Following the peak TEC  
265 value in the southern crest, a recovery phase begins at 4:03 UT and persisted until 22:00 UT. During the recovery phase, the ionospheric TEC returns to normal state.

Figure 6 (15-21) shows ionospheric TEC maps during show the temporal variation of the ionospheric TEC during the main phase of the November 4-5, 2023 geomagnetic storm after sunset, from 18:00 UT to 00:00 UT. During this period, storm-time ionization was initially confined to the equatorial region, resulting in underdeveloped EIA crests. However, by 20:00 UT, ion-  
270 ization increased in the northern and southern low latitudes, forming a weak EIA with smaller latitudinal extent. Storm-time electrodynamic and variations in thermospheric winds and composition influence the EIA, resulting in decreased TEC at typical crest latitudes. This unusual EIA pattern reflects the complex interaction of disrupted electric fields, neutral winds, and plasma movement during the storm's main phase over the America.

On May 10, 2024, a double crest EIA structure at the equator containing normal TEC values with minor variations observed in  
275 other regions until 18:00 UT (See the supporting Figure of madrigal TEC maps for May 10-11, 2024). The global ionospheric maps in Figure 6 (22-28) illustrate the temporal variation of the ionospheric TEC during the main phase of the May 10-11, 2024 geomagnetic storm after sunset, from 18:00 UT to 00:00 UT. The May 10-11, 2024 storm was particularly intense, with substantial increase in TEC in the EIA crests over American longitudes and significant poleward shifting of the EIA crests up to approximately  $\pm 35^\circ$  geomagnetic latitudes, during the main phase of this storm. In pre-midnight sector, the meridional  
280 wind with large southward (poleward) component in southern hemisphere during the main phase of the storm greatly enhanced



**Figure 6.** Madrigal-based global ionospheric TEC maps during the main phase of the four geomagnetic storms after sunset, from 18:00 UT to 00:00 UT shown as: (1-7) March 23-24, 2023, (8-14) April 23-24, 2023, (15-21) November 4-5, 2023, and (22-28) May 22-28, 2024.

the ambipolar diffusion in the southern hemisphere through neutral drag, resulting in the poleward expansion of the EIA crests between 22:00 to 00:00 UT. Meanwhile, the zonal winds of the storm-time exhibited large surges to the west opposite to the zonal winds of the prevailing quiet-time to the east at the same time. The mid-latitude westward wind surge caused by Coriolis-induced deflection of equatorial wind surges, as well as the ion drag effect of the intense westward subauroral polarization stream that coincided with the primary ionospheric trough. A significant westward wind surge builds up DDEF in the low- and mid-latitude ionosphere, which is typically westward during the day until local dusk. This alters the equatorial zonal electric field, which can minimize upward plasma drifts and reduce the intensity of the EIA. It is very unlikely that the disturbance dynamo was responsible for the excessive poleward expansion observed of the EIA crests during the main phase of the storm (Aa et al., 2024). The recovery phase showed negative with a strong reduction in TEC along with EIA crests suppression

290 and merging into a single equatorial band (See the supporting Figure of meridional TEC maps for May 10-11, 2024). This can be attributed to the effect of composition changes and DDEF that are dominant during the storm recovery phase (Aa et al., 2024). As reported in previous studies, both positive and negative storm phases have an impact on the latitudinal density gradients at the EIA crests and the background ionospheric density, both of which are essential for the growth of equatorial spread F (ESF) through the R-T instability (Sripathi and Singh, 2020; Singh and Sripathi, 2020). These gradients are modified by storm-time electric fields such as PPEF and DDEF, which change the positions and intensities of ESF events. The ESF formation can also be further shaped by low-latitude sporadic E (Es) layers, which can locally increase densities and initiate localized irregularities (Singh and Sripathi, 2020). The observed dependency of plasma irregularities on the storm phase and the underlying ionospheric structures can be explained by these mechanisms. Furthermore, atmospheric waves and tides play a crucial role in the diurnal variability of the PRE, EIA, and the generation of plasma irregularities. Tidal and gravity wave perturbations originating in the lower atmosphere modify neutral winds, which in turn influence the equatorial dynamo and the vertical  $E \times B$  drift (Sripathi and Singh, 2020). This variability alters the magnitude and timing of the PRE, modifies the EIA density gradients, and sustains the R-T instability, thereby controlling the occurrence and intensity of equatorial spread F (ESF) (Singh and Sripathi, 2020). Even under geomagnetically disturbed conditions, these processes highlight the importance of lower atmosphere-ionosphere coupling in regulating plasma irregularities.

305

In each case, the variations in TEC were influenced by the intensity and duration of geomagnetic activity, driven by solar wind parameters and enhanced JH, which affected the distribution and magnitude of ionospheric disturbances at low latitudes. **Also, the F10.7 index also serves as a key indicator of solar activity. Moderate solar flux values ( $100 < F10.7 < 200$ ) are recorded during the equinox (March–April) and winter solstice (November) storms, while enhanced solar activity ( $F10.7 > 200$ ) characterized the May solstice storm (Mother’s Day event). Such elevated F10.7 levels correspond to increased solar EUV irradiance, resulting in a denser and more variable ionosphere due to intensified ionization processes particularly during the mother’s day storm.**

**Previous studies have reported that the seasonal variation of equatorial plasma bubbles (EPBs) is primarily governed by the angle between the magnetic meridian and the solar terminator. The occurrence probability of post-sunset EPBs is generally higher during equinoctial months (March and April) and minimal around the summer solstice (as seen during the May storm). In our study, the occurrence of EPBs during equinox storms can therefore be attributed mainly to seasonal effects. In contrast, the suppression of post-sunset EPBs during the main phase of the November storm can be explained by the fact that the prompt penetration electric field (PPEF) arrived after midnight local time, when post-sunset conditions were no longer favourable for bubble development. The unseasonal generation of post-sunset EPBs during the Mother’s Day storm appears to result from the combined influence of strong solar flux and a significant eastward PPEF near local sunset.**

320

**Table 5.** Joule Heating (JH) in northern and southern hemispheres (NH and SH) and the Equatorial Ionization Anomaly (EIA) over the American longitudes during the main phase of the four geomagnetic storms after sunset, from 18:00 UT to 00:00 UT.

Storm Days	Time (UT)	JH(mW/m <sup>2</sup> )		EIA structure
		NH	SH	
March 23, 2023	18:00 UT	41.3	41.8	merged or single crest (moderate)
–	19:00 UT	40.7	40.7	merged or single crest (strong)
–	20:00 UT	50.9	52.6	merged or single crest (strong)
–	21:00 UT	52.5	53.4	merged or single crest (strong)
–	22:00 UT	50.1	52.0	merged or single crest (strong)
–	23:00 UT	52.2	53.9	double crest (moderate)
March 24, 2023	00:00 UT	56.6	55.9	double crest (moderate)
April 23, 2023	18:00 UT	33.7	22.9	double crest (moderate)
–	19:00 UT	94.7	77.4	double crest (strong)
–	20:00 UT	97.0	85.4	double crest (strong)
–	21:00 UT	73.5	52.2	double crest (strong)
–	22:00 UT	44.7	14.4	double crest (strong)
–	23:00 UT	55.1	24.6	double crest (moderate)
April 24, 2023	00:00 UT	37.0	13.2	double crest (weak)
November 4, 2023	18:00 UT	26.4	24.3	merged or single crest(weak)
–	19:00 UT	30.1	27.7	merged or single crest (weak)
–	20:00 UT	27.9	25.6	merged or single crest (weak)
–	21:00 UT	22.4	22.0	merged or single crest (weak)
–	22:00 UT	25.2	26.0	double crest (weak)
–	23:00 UT	25.0	28.0	double crest (weak)
November 5, 2023	00:00 UT	10.3	19.7	double crest (weak)
May 10, 2024	18:00 UT	63.8	51.8	double crest (moderate)
–	19:00 UT	47.7	30.4	double crest (moderate)
–	20:00 UT	83.1	80.9	double crest (strong)
–	21:00 UT	74.5	80.2	double crest (strong)
–	22:00 UT	114.7	101.4	double crest (strong)
–	23:00 UT	57.1	73.2	double crest (strong)
May 11, 2024	00:00 UT	98.5	116.2	double crest (weak)

### 3.3.2 Ionospheric Plasma Bubbles (EPBs)

Figures 7-10 show LT variations of EPBs in the three longitudes that are America, Africa, and Asia during the four geomagnetic storms. Each storm's unique solar wind conditions, geomagnetic activity, and JH distribution which can influence the extent and location of ionospheric disturbances and plasma irregularities. The storm-time ring current is critical for both the development and suppression of EPBs. For instance, EPBs are generated if the greatest excursion of the ring current (minimum SYM-H index) occurs between midnight and postmidnight. However, the plasma irregularities are suppressed when the maximum excursion of the ring current occurs in the early afternoon (Aarons, 1991). Table 3 shows the LT of maximum excursion of ring current or MPE at each GNSS location. During geomagnetic storms in March, April, and May, the ring current has the greatest excursion after midnight in American sector, indicating the highest likelihood of plasma irregularities in the evenings following the storms' main phases in this sector. For instance, the maximum ring current excursions at the GNSS stations BOGT, RIOP, and SANT (LT = UT-5) occurred at approximately 00:21, 23:03, 11:54, and 21:14 LTs on March 24, April 24, November 5, and May 11, respectively. On the same days, the CORD station (LT = UTC-4) reported the highest excursions at 01:21, 00:03, 12:54, and 22:14 LT. Notably, the highest excursions during the storms of March, April, and May occurred predominantly during the night to midnight which is the most ideal time for the growth of post-sunset EPBs. During November storm, the maximum ring current excursion occurred at approximately 11:54, and 12:54 LTs which is the least favourable time for EPBs to occur. However, the intensity of post-sunset plasma irregularities varies between these storms due to complex interaction of important factors such as storm-time electric fields, meridional winds, background ionospheric conditions, local time and seasons. The largest excursion of the ring current in the November storm occurred before dusk, indicating that post-sunset plasma irregularities were unlikely during the main phase of this storm.

During the the main phase of March 23-25, 2023 storm, strong post-sunset plasma irregularities were observed near magnetic equator (BOGT and RIOP) in the American sector. However, the plasma irregularities reduced over the stations CORD and SANT in the EIA crests region of South America. The unusual EIA structure consisting of high electron density in the crests and moderate in the trough, results in a non-uniform vertical distribution. A steep vertical density gradient at the bottomside F-layer, when combined with equatorward trans-equatorial neutral winds and vertical plasma drifts, can provide favourable conditions for the development of strong plasma irregularities over BOGT and RIOP. During the main phase of April 23, 2023 storm, ionospheric plasma conditions supported post-sunset plasma irregularities to develop near the magnetic equator while inhibiting plasma irregularities over crest regions (CORD and SANT) highlighting the localized impact of weaker crests. The November 4-6, 2023 storm showed highly suppressed post-sunset EPBs across all sectors due to lower ionospheric conductivity and weak electric fields, which reduced vertical plasma drifts and prevented irregularities from growing. The severe May 10-13, 2024 storm had a significant impact on the ionosphere, causing intense post-sunset plasma irregularities, particularly over southern low latitude stations (CORD and SANT) in the American sector. The supper-fountain effect due to huge uplift resulted in intensified EIA crests with steeper plasma density gradients, supporting the formation of strong post-sunset EPBs in the crests region. Overall, post-sunset EPBs occur primarily over the American sector during the main phases of storms in March, April, and May. No EPBs were observed during the main phase of the November storm. Throughout the recovery phase

of these storms, EPBs remained highly suppressed at these locations. The seasonal behavior of EPBs depends on magnetic declination (alignment of the Earth's magnetic field with the solar terminator) and trans-equatorial meridional winds (Seba and Nigussie, 2016; Tsunoda, 1985). Lloyd's season classification includes winter (January, February, November, December),  
360 summer (May to August), and equinox (March, April, September, and October) (Liu et al., 2010). During equinoctial months, the Sun's greater elevation and minimal trans-equatorial thermospheric winds can result in significant ionization and steeper plasma gradients, which can sustain EPBs (Rastogi, 1980; Maruyama, 1988). Intense EPBs can significantly disturb GNSS-based navigation and communication systems by causing amplitude and phase scintillations. These scintillations cause fast fluctuations in signal power and carrier phase, which reduces positioning precision and reliability. During seasons conducive  
365 to the formation of EPBs, notably at the equinoxes, GNSS receivers may encounter large range errors, cycle slipping, or even loss of lock, particularly for single-frequency users. As a result, understanding the seasonal and regional variability of EPBs is crucial for the forecasting scintillation impacts and creating mitigation solutions, including the use of multi-frequency receivers, real-time monitoring, and augmentation systems (Aarons, 1997; Basu et al., 1999).

During geomagnetic storms, equatorial ionospheric plasma density distribution is influenced by electric fields, including DDEF  
370 from storm-time neutral winds and PPEF of magnetospheric origin (Astafyeva et al., 2018). According to (Abdu, 1997; Abdu et al., 2009), the presence of eastward/westward PPEF after sunset during the main phase of a storm can affect the vertical drift of plasma over the equatorial region either strengthening or disturbing the favourable conditions for EPBs. During a storm's recovery phase, the westward DDEF in the sunset terminator may cause the plasma to move downward, and the normal PRE is reversed, reducing EPBs (Abdu, 1997). Figure 11 (a-d) depicts the temporal fluctuations of the PPEF during geomagnetic  
375 storms on March 22-25, 2023, April 22-25, 2023, November 4-7, 2023, and May 10-12, 2024. Several northward and southern turnings of the IMF  $B_z$  occurred during the main phase of these storms. As the IMF  $B_z$  turns southward (northward), the convection electric field and polar cap potentials grow or decrease. The overshielding (under-shielding) electric field caused by the IMF  $B_z$  northward (southward) turning can suppress/trigger EPBs by reversing/enhancing the PRE in the local evening regions. The difference in the pattern of the ionospheric EPBs over the three sectors depends on the local timing of IMF  $B_z$   
380 southward, the disturbed electric field and longitudinal variability of vertical drift (Amaechi et al., 2018; Fejer, 1997; Fejer et al., 1999). At 18:00 UT during the main phase of the March 23 storm, a westward PPEF of magnitude approximately -0.55 mV/m was seen in the Asian sector. At the same time, an eastward PPEF of approximately 0.5 mV/m can be observed in the American and African sectors before dusk.

Following sunset on April 23, multiple variable duration PPEFs are visible in the three sectors. In America, the four PPEFs  
385 are noticeable. The first peak value is 0.33 mV/m from 18:05 to 20:05 UT, the second is -0.78 mV/m from 18:05 to 20:05 UT, the third is 0.61 mV/m from 23:05 to 23:55 UT, and the fourth is 0.45 mV/m from 00:40 to 03:15 UT. In the African sector, two PPEFs of varied duration with maximum values of roughly 0.60 mV/m and -0.65 mV/m are observed before midnight from 21:00 UT to 23:00 UT and 23:05 to 23:55 UT. A strong long-duration westward PPEF with a peak value of approximately -1.36 mV/m is also detected after midnight, between 1:45 and 4:35 UT. In Asia, the first westward PPEF with a peak value of  
390 -0.54 mV/m occurs from 18:05 to 20:05 UT, the second eastward is 0.94 mV/m from 21:10 to 23:00 UT, and the third long duration eastward PPEF with a peak value of 0.66 mV/m occurs from 01:45 to 04:35 UT on April 24. During the main phase

of November 4-7, 2023 storm, the PPEFs of different polarities and duration reach the equatorial ionosphere during the day. In America, the first westward PPEF with maximum value of approximately -0.32 mV/m occurred in the morning hours from 8:30 to 10:25 UT, and the second eastward penetrating field of magnitude approximately 0.44 mV/m is observed at local noon from 12:05 UT to 13:25 UT. In Africa, two notable PPEFs with peak values of -0.53 mV/m and 0.34 mV/m are observed at local noon. In Asia, there is one noticeable westward penetrating field with a peak value of approximately -0.42 mV/m from 12:05 to 13:35 UT. During the mother's day geomagnetic storm on 10-12 May 2024, we observed significant storm-time PPEFs after sunset. For instance, intense eastward-directed PPEFs of different durations between 18:00 and 22:00 UT cause super-fountain effect in the American sector. This results in to notable variations in ionospheric electron density and the positioning of EIA crests as shown in madrigal TEC maps. We can also see eastward penetrating electric fields with maximum values of 2.59 mV/m and 1.80 mV/m in Africa and Asia, respectively, and westward PPEF with a peak value of roughly -2.73 mV/m in America between 22:50 and 23:25 UT. America and Africa experience an eastward PPEF with a peak value of 1.39 mV/m and a westward PPEF with a peak value of -1.95 mV/m, respectively, between 23:30 and 2:00 UT. During the March 23-25, 2023 storm, a westward PPEF of approximately -0.55 mV/m was observed in the Asian sector at 18:00 UT, while an eastward PPEF of approximately 0.5 mV/m was observed in the American and African sectors before dusk. These PPEFs contributed to the merging of EIA crests and increased TEC in the equatorial zone, leading to strong post-sunset EPBs in the American sector. The April 23-25, 2023 storm exhibited multiple PPEFs of varying magnitudes and durations across different sectors, with significant eastward and westward PPEFs in America, Africa, and Asia. These PPEFs influenced the expansion of EIA crests and increased TEC, particularly in the northern hemisphere, and caused noticeable post-sunset EPBs in the American sector. The November 4-6, 2023 storm had PPEFs of different polarities and durations, with westward and eastward PPEFs observed in America, Africa, and Asia. However, lower ionospheric conductivity and weak electric fields suppressed post-sunset EPBs across all sectors. The May 10-13, 2024 storm featured intense eastward-directed PPEFs between 18:00 and 22:00 UT, causing the super fountain effect in the American sector, leading to significant TEC variations and poleward shifting of EIA crests. Strong eastward and westward PPEFs were also observed in Africa and Asia, contributing to enhanced post-sunset EPBs, particularly in the southern low latitude stations.

- The magnitude and spatial distribution of JH varied significantly with storm intensity, local time, duration, and season. A clear solstice–equinox contrast was observed, with the May 2024 storm exhibiting the most intense post-sunset JH and the November 2023 storm showing the weakest. Nearly symmetric JH patterns were observed during equinoctial storms, whereas solstice events exhibited pronounced hemispheric asymmetries.
- Distinct electrodynamic pathways were identified linking JH-driven equatorward winds to the evolution of the equatorial ionization anomaly (EIA). The combined influence of storm-time electric fields and thermospheric winds determined whether single- or double-crest EIAs developed, with asymmetric winds amplifying or suppressing the anomaly in opposite hemispheres.
- The seasonal variation of post sunset EPBs was consistent with previous findings, being primarily controlled by the angle between the magnetic meridian and the solar terminator. The occurrence probability of post-sunset EPBs was highest

during equinoctial storms (March and April) and minimal during the summer solstice (May storm). In our analysis, the EPBs observed during equinox storms reflect seasonal effects, while their suppression during the November 2023 storm is attributed to a delayed PPEF arriving after midnight, when post-sunset conditions were no longer favourable for instability growth. Conversely, the unseasonal generation of post-sunset EPBs during the Mother's Day storm resulted from the combined influence of elevated solar flux and a strong eastward PPEF near local sunset.

430

- The results further highlight the dynamic interplay between overshielding and undershielding PPEFs and thermospheric winds in modulating the PRE that further affects the post-sunset plasma uplift and EPBs onset. Variations in EIA gradients and vertical plasma drifts control EPB intensity and occurrence, exhibiting clear seasonal and hemispheric dependencies.

435

Overall, these findings demonstrate a strong coupling between high-latitude Joule heating and low-latitude ionospheric variability, providing new observational evidence that storm-time thermospheric forcing plays a critical role in shaping the global electrodynamic response. The results strengthen the physical basis for forecasting space weather impacts on GNSS and satellite communication systems. Future work will integrate multi-instrument observations and physics-based modeling to quantify the relative contributions of JH, PPEFs, and neutral dynamics to storm-time ionospheric variability.

#### 440 4 Conclusions

We presented a comprehensive analysis of four geomagnetic storms (March 23–25, April 23–25, and November 4–6, 2023, and May 10–13, 2024) to investigate storm-time coupling between solar wind forcing, Joule heating (JH), thermospheric dynamics, and low-latitude ionospheric responses. This multi-event, multi-season comparison provides one of the first systematic characterizations of how high-latitude energy input through JH modulates equatorial ionospheric structures and post-sunset irregularities.

445

- The magnitude and spatial distribution of JH varied significantly with storm intensity, local time, duration, and season. A clear solstice–equinox contrast was observed, with the May 2024 storm exhibiting the most intense post-sunset JH and the November 2023 storm showing the weakest. Nearly symmetric JH patterns were observed during equinoctial storms, whereas solstice events exhibited pronounced hemispheric asymmetries.

450

- Distinct electrodynamic pathways were identified linking JH-driven equatorward winds to the evolution of the equatorial ionization anomaly (EIA). The combined influence of storm-time electric fields and thermospheric winds determined whether single- or double-crest EIAs developed, with asymmetric winds amplifying or suppressing the anomaly in opposite hemispheres.

455

- The seasonal variation of post sunset EPBs was consistent with previous findings, being primarily controlled by the angle between the magnetic meridian and the solar terminator. The occurrence probability of post-sunset EPBs was highest during equinoctial storms (March and April) and minimal during the summer solstice (May

storm). In our analysis, the EPBs observed during equinox storms reflect seasonal effects, while their suppression during the November 2023 storm is attributed to a delayed PPEF arriving after midnight, when post-sunset conditions were no longer favourable for instability growth. Conversely, the unseasonal generation of post-sunset EPBs during the Mother's Day storm resulted from the combined influence of elevated solar flux and a strong eastward PPEF near local sunset.

460

- The results further highlight the dynamic interplay between overshielding and undershielding PPEFs and thermospheric winds in modulating the PRE that further affects the post-sunset plasma uplift and EPBs onset. Variations in EIA gradients and vertical plasma drifts control EPB intensity and occurrence, exhibiting clear seasonal and hemispheric dependencies.

465

Overall, these findings demonstrate a strong coupling between high-latitude Joule heating and low-latitude ionospheric variability, providing new observational evidence that storm-time thermospheric forcing plays a critical role in shaping the global electrodynamic response. The results strengthen the physical basis for forecasting space weather impacts on GNSS and satellite communication systems. Future work will integrate multi-instrument observations and physics-based modeling to quantify the relative contributions of JH, PPEFs, and neutral dynamics to storm-time ionospheric variability.

470

*Data availability.* Solar wind parameters and magnetic indices are obtained from the GSFC/SPDF OMNIWeb interface ([https://omniweb.gsfc.nasa.gov/form/omni\\_min.html](https://omniweb.gsfc.nasa.gov/form/omni_min.html)). MIT-TEC maps are obtained from the MIT Haystack website (<http://cedar.openmadrigal.org/static/experiments4/2021/gps/>). Global ROTI maps are available at <https://stdb2.isee.nagoya-u.ac.jp/GPS/GPS-TEC/>. ROTI variations videos are obtained from System for Ionosphere Monitoring and Research from GNSS database, at ([https://simurg.iszf.irk.ru/create\\_map](https://simurg.iszf.irk.ru/create_map)). Equatorial electric fields over a specific longitude are obtained from the real-time Prompt Penetration Equatorial Electric Field Model (PPEEFM) available at <https://geomag.colorado.edu/real-time-model-of-the-ionospheric-electric-fields.html>. Weimer model simulations are provided by the Community Coordinated Modeling Center (CCMC) at NASA Goddard Space Flight Center through (<https://ccmc.gsfc.nasa.gov/models/>).

475

480 *Competing interests.* It is declare that the authors have no competing interest.

## References

- Aa, E., Zhu, S., and Liu, S.: Statistical Analysis of Equatorial Plasma Irregularities Retrieved From Swarm 2013–2019 Observations, *Journal of Geophysical Research: Space Physics*, 125, <https://doi.org/10.1029/2020JA028065>, 2020.
- Aa, E., Chen, Y., and Luo, B.: Dynamic Expansion and Merging of the Equatorial Ionization Anomaly During the 10–11 May 2024  
485 Super Geomagnetic Storm, *Remote Sensing*, 16, <https://doi.org/10.3390/rs16224290>, 2024.
- Aarons, J.: The role of the ring current in the generation or inhibition of equatorial F layer irregularities during magnetic storms, *Radio Science*, 26, 1131–1149, 1991.
- Aarons, J.: Global positioning system phase fluctuations at auroral latitudes, *Journal of Geophysical Research: Space Physics*, 102, 17 219–17 231, 1997.
- 490 Abdu, M. A.: Major phenomena of the equatorial ionosphere-thermosphere system under disturbed conditions, *Journal of Atmospheric and Solar-Terrestrial Physics*, 59, 1505–1519, 1997.
- Abdu, M. A.: *The Dynamical Ionosphere*, chap. Equatorial F region irregularities, Elsevier, 2020.
- Abdu, M. A., Batista, I. S., Reinisch, B. W., Souza, J. R. D., Sobral, J. H. A., Pedersen, T. R., Medeiros, A. F., Schuch, N. J.,  
495 Paula, E. R. D., and Groves, K. M.: Conjugate Point Equatorial Experiment (COPEX) campaign in Brazil: Electrodynamics highlights on spread F development conditions and day-to-day variability, *Journal of Geophysical Research: Space Physics*, 114, <https://doi.org/10.1029/2008JA013749>, 2009.
- Amaechi, P. O., Oyeyemi, E. O., and Akala, A. O.: Geomagnetic storm effects on the occurrences of ionospheric irregularities over the African equatorial/low-latitude region, *Advances in Space Research*, 61, 2074–2090, <https://doi.org/10.1016/j.asr.2018.01.035>, 2018.
- 500 Araujo-Pradere, E. A., Fuller-Rowell, T. J., Codrescu, M. V., and Bilitza, D.: Characteristics of the ionospheric variability as a function of season, latitude, local time, and geomagnetic activity, *Radio Sci.*, 40, RS5009, <https://doi.org/10.1029/2004RS003179>, 2006.
- Astafyeva, E., Zakharenkova, I., Hozumi, K., Alken, P., Coisson, P., Hairston, M. R., and Coley, W. R.: Study of the  
505 Equatorial and Low-Latitude Electrodynamic and Ionospheric Disturbances During the 22–23 June 2015 Geomagnetic Storm Using Ground-Based and Space-borne Techniques, *Journal of Geophysical Research: Space Physics*, 123, 2424–2440, <https://doi.org/10.1002/2017JA024981>, 2018.
- Axford, W. I.: Viscous interaction between the solar wind and the earth’s magnetosphere, *Planetary and Space Science*, 12, 45–53, [https://doi.org/10.1016/0032-0633\(64\)90067-4](https://doi.org/10.1016/0032-0633(64)90067-4), 1964.
- Balan, N., Alleyne, H., Otsuka, Y., Lekshmi, D. V., Fejer, B. G., and McCrea, I.: Relative effects of electric field and neutral wind on  
510 positive ionospheric storms, *Earth, Planets and Space*, 61, 2009.
- Balan, N., Otsuka, Y., Nishioka, M., Liu, J. Y., and Bailey, G. J.: Physical mechanisms of the ionospheric storms at equatorial and higher latitudes during the recovery phase of geomagnetic storms, *Journal of Geophysical Research: Space Physics*, 118, <https://doi.org/10.1002/jgra.50275>, 2013.
- Balan, N., Liu, L., and Le, H. J.: A brief review of equatorial ionization anomaly and ionospheric irregularities, *Earth and Planetary  
515 Physics*, 2, 257–275, <https://doi.org/10.26464/epp2018025>, 2018.
- Basu, S., Groves, K. M., Quinn, J. M., and Doherty, P.: A comparison of TEC fluctuations and scintillation at Ascension Island, *Journal of Atmospheric and Solar-Terrestrial Physics*, 61, 1219–1226, 1999.

- Blanc, M. and Richmond, A. D.: The ionospheric disturbance dynamo, *Journal of Geophysical Research*, 85, 1980.
- Buonsanto, M. J.: Ionospheric storms: A review, *Space Science Reviews*, 88, 563–601, 1999.
- 520 Calabia, A., Imtiaz, N., Altadill, D., Yasyukevich, Y., Segarra, A., Prol, F. S., Adhikari, B., del Peral, L., Frias, M. D. R., and Molina, I.: Uncovering the Drivers of Responsive Ionospheric Dynamics to Severe Space Weather Conditions: A Coordinated Multi-Instrumental Approach, *Journal of Geophysical Research: Space Physics*, 129, e2023JA031862, <https://doi.org/10.1029/2023JA031862>, 2024.
- Cueto, M., Magdaleno, S., Cezon, A., and Sardon, E.: Characterization of equatorial ionospheric features on the verge of the next  
525 solar cycle maximum, *Journal of Geophysical Research*, pp. 3055–3065, <https://doi.org/10.1029/2004JA010884>, 2012.
- Dugassa, T., Habarulema, J. B., and Nigussie, M.: Longitudinal variability of occurrence of ionospheric irregularities over the American, African and Indian regions during geomagnetic storms, *Advances in Space Research*, 63, 2609–2622, 2019.
- Dungey, J. W.: Interplanetary Magnetic Field and the Auroral Zones, *Physical Review Letters*, 6, 47–48, <https://doi.org/10.1103/PhysRevLett.6.47>, 1961.
- 530 Fejer, B. G., Scherliess, L., and de Paula, E. R.: Effects of the vertical plasma drift velocity on the generation and evolution of equatorial spread F, *Journal of Geophysical Research*, 104, 19854–19869, 1999.
- Fejer, D. J.: The electrodynamics of the low-latitude ionosphere: Recent results and future challenges, *Journal of Atmospheric and Solar-Terrestrial Physics*, 59, 1465–1482, [https://doi.org/10.1016/S1364-6826\(96\)00149-6](https://doi.org/10.1016/S1364-6826(96)00149-6), 1997.
- Fuller-Rowell, T. J., Codrescu, M. V., Moffett, R. J., and Quegan, S.: Response of the Thermosphere and Ionosphere to Geomagnetic  
535 Storms, *Journal of Geophysical Research*, 99, 3893–3914, 1994.
- Fuller-Rowell, T. J., Codrescu, M. V., Roble, R. G., and Richmond, A. D.: How does the thermosphere and ionosphere react to a geomagnetic storm?, in: *Geophysical Monograph Series*, vol. 98, pp. 203–225, American Geophysical Union, 1997.
- González, G.: Storm-time variability of ionospheric irregularities over South America, *Journal of Atmospheric and Solar-Terrestrial Physics*, 241, 105980, <https://doi.org/10.1016/j.jastp.2022.105980>, 2022.
- 540 Huang, C. S.: Long-Lasting Penetration Electric Fields during Geomagnetic Storms: Observations and Mechanisms, *Journal of Geophysical Research: Space Physics*, 124, 2019.
- Huang, C. S., Foster, J., and Kelley, M.: Long-duration penetration of the interplanetary electric field to the low-latitude ionosphere during the main phase of magnetic storms, *Journal of Geophysical Research*, 110, 2005.
- Huang, L., Huang, J., Wang, J., Jiang, Y., Deng, B., Zhao, K., and Lin, G.: Analysis of the north–south asymmetry of the  
545 equatorial ionization anomaly around 110°E longitude, *Journal of Atmospheric and Solar-Terrestrial Physics*, 102, 354–361, <https://doi.org/10.1016/j.jastp.2013.06.010>, 2013.
- Imtiaz, N., Dugassa, T., Calabia, A., Anorou, C., and Kashcheyev, A.: Westward PPEF plays important role in the suppression of post-midnight plasma irregularities: A case study of the November 2021 geomagnetic storm, *Journal of Geophysical Research: Space Physics*, 129, e2023JA032367, <https://doi.org/10.1029/2023JA032367>, 2024.
- 550 Jin, H., Yan, C., Fang, G. Y., He, H., Xu, X., Zhang, W., Sun, Y., Li, K., Hu, J., and Jiang, J.: Interaction Between Equatorial to Low-Latitude Postmidnight F-Region Irregularities and LSTIDs in China During Geomagnetic Disturbances Based on Ground-Based Instruments, *Journal of Geophysical Research: Space Physics*, 127, 2022.
- Kassa, T., Damtie, B., Bires, A., Yizengaw, E., and Cilliers, P.: Storm-time characteristics of the equatorial ionization anomaly in the East African sector, *Advances in Space Research*, 56, 57–70, <https://doi.org/10.1016/j.asr.2015.04.002>, 2015.

- 555 Kassa, Y., Damtie, B., and Tebabal, A.: Ionospheric and magnetic signatures of extreme space weather events of 17 March and 23 June 2015 over the African sector, *Journal of Atmospheric and Solar-Terrestrial Physics*, 243, 106 003, 2023.
- Kassa, Y., Tebabal, A., and Damtie, B.: Nighttime ionospheric irregularity during intense geomagnetic storm events over the Europe-African longitudinal sector, *Heliyon*, 10, e38 138, 2024.
- Kelly, M. C.: *The Earth's Ionosphere: Plasma Physics and Electrodynamics*, Academic Press, 2nd edn., 2009.
- 560 Kikuchi, T., Hashimoto, K. K., and Nozaki, K.: Penetration of magnetospheric electric fields to the equator during a geomagnetic storm, *Journal of Geophysical Research*, 113, 2008.
- Li, K. F., Lin, L. C., Bui, X. H., and Liang, M. C.: The 11 year solar cycle response of the equatorial ionization anomaly observed by GPS radio occultation, *Journal of Geophysical Research: Space Physics*, 123, <https://doi.org/10.1002/2017JA024634>, 2018.
- Lin, C. H., Richmond, A. D., Heelis, R. A., Bailey, G. J., Lu, G., Liu, J. Y., and Su, S. Y.: Theoretical study of the low- and midlatitude ionospheric electron density enhancement during the October 2003 superstorm: Relative importance of the neutral wind and the electric field, *Journal of Geophysical Research*, 110, <https://doi.org/10.1029/2005JA011304>, 2005.
- 565 Liu, J., Zhao, B., and Liu, L.: Time delay and duration of ionospheric total electron content responses to geomagnetic disturbances, *Annales Geophysicae*, 28, 795–805, <https://doi.org/10.5194/angeo-28-795-2010>, 2010.
- Luan, X.: Equatorial Ionization Anomaly Variations During Geomagnetic Storms, chap. 13, pp. 301–312, *American Geophysical Union (AGU)*, <https://doi.org/10.1002/9781119815617.ch13>, 2021.
- 570 Lukianova, R.: Swarm field-aligned currents during a severe magnetic storm of September 2017, *Annales Geophysicae*, 38, 191–206, <https://doi.org/10.5194/angeo-38-191-2020>, 2020.
- Maruyama, T.: A diagnostic model for equatorial spread F, 1, Model description and application to electric field and neutral wind effects, *Journal of Geophysical Research*, 93, 14 611–14 622, <https://doi.org/10.1029/JA093iA12p14611>, 1988.
- 575 Pi, X., Mannucci, A., Lindqwister, U. J., and Ho, C. M.: Monitoring of global ionospheric irregularities using the worldwide GPS network, *Geophysical Research Letters*, 24, 2283–2286, 1997.
- Rastogi, R. G.: Seasonal variation of equatorial spread F in the American and Indian zones, *Journal of Geophysical Research*, 85, 722–726, <https://doi.org/10.1029/JA085iA02p00722>, 1980.
- Richmond, A. D. and Matsushita, S.: Thermospheric Response to a Magnetic Substorm, *Journal of Geophysical Research*, 80, 2839–2850, 1975.
- 580 Rideout, W. and Coster, A.: Automated GPS processing for global total electron content data, *GPS Solutions*, 10, 219–228, 2006.
- Rishbeth, H.: The equatorial F-layer: progress and puzzles, *Annales Geophysicae*, 18, 730–739, <https://doi.org/10.1007/s00585-000-0730-6>, 2000.
- Seba, E. B. and Nigussie, M.: Investigating the effect of geomagnetic storm and equatorial electrojet on equatorial ionospheric irregularity over East African sector, *Advances in Space Research*, 58, 1708–1719, <https://doi.org/10.1016/j.asr.2016.06.037>, 2016.
- 585 Seemala, G.: GPS-TEC analysis application, Technical report, Institute for Scientific Research, 2011.
- Singh, R. and Sripathi, S.: A statistical study on the local time dependence of equatorial spread F (ESF) irregularities and their relation to low-latitude Es layers under geomagnetic storms, *Journal of Geophysical Research: Space Physics*, 125, e2019JA027 212, <https://doi.org/10.1029/2019JA027212>, 2020.
- 590 Smith, A. R., Ozturk, D. S., Delamere, P., Lu, G., and Kim, H.: Investigating the interhemispheric asymmetry in Joule heating during the 2022 geomagnetic storm, *Journal of Geophysical Research: Space Physics*, 128, e2023JA031 872, <https://doi.org/10.1029/2023JA031872>, 2023.

- 595 Sripathi, S. and Singh, R.: A study on the response of the ionosphere to the three major space weather events of 2015 using meridional chain of ionosondes and GPS receivers over India, *Journal of Sun and Geosphere*, <https://doi.org/10.31401/SunGeo.2018.02.08>, 2020.
- Tahir, A., Wu, F., Shah, M., Amory-Mazaudier, C., Jamjareegulgarn, P., Verhulst, T. G. W., and Ameen, M. A.: Multi-instrument observation of the ionospheric irregularities and disturbances during the 23–24 March 2023 geomagnetic storm, *Remote Sensing*, **16**, 1594, <https://doi.org/10.3390/rs16091594>, 2024.
- 600 Tsunoda, R. T.: High-latitude F-region irregularities: A review and synthesis, *Reviews of Geophysics*, **23**, 577–582, <https://doi.org/10.1029/RG023i005p00577>, 1985.
- Tsurutani, B. T., Mannucci, A., Iijima, B., Abdu, M. A., Sobral, J. H. A., Gonzalez, W. D., Guarnieri, F. L., Tsuda, T., Saito, A., Yumoto, K., Fejer, B. G., Fuller-Rowell, T. J., Kozyra, J. U., Foster, J. C., Coster, A. J., and Vasyliunas, V. M.: Global dayside ionospheric uplift and enhancement associated with interplanetary electric fields, *Annales Geophysicae*, **22**, 567–574, <https://doi.org/10.5194/angeo-22-567-2004>, 2004.
- 605 Tulasi Ram, S., Su, S. Y., and Liu, C. H.: FORMOSAT-3/COSMIC observations of seasonal and longitudinal variations of equatorial ionization anomaly and its interhemispheric asymmetry during the solar minimum period, *Journal of Geophysical Research: Space Physics*, **114**, <https://doi.org/10.1029/2008JA013880>, 2009.
- Vaggu, P. R., Deshpande, K. B., Datta-Barua, S., Bust, G. S., Hampton, D. L., López Rubio, A., and Conroy, J. P.: Morphological and spectral features of ionospheric structures at E- and F-region altitudes over Poker Flat analyzed using modeling and observations, *Sensors*, **23**, –, <https://doi.org/10.3390/sym15101940>, 2023.
- 610 Weimer, D. R.: Improved ionospheric electrodynamic models and application to calculating Joule heating rates, *Journal of Geophysical Research*, **110**, A05 306, <https://doi.org/10.1029/2004JA010884>, 2005.
- Wu, Q., Wang, W., Lin, D., Huang, C., and Zhang, Y.: Penetrating electric field during the Nov 3–4, 2021 geomagnetic storm, *Journal of Atmospheric and Solar-Terrestrial Physics*, **257**, 106 219, <https://doi.org/10.1016/j.jastp.2024.106219>, 2024.
- 615 Zhang, X., Wan, W., and Yue, X.: Storm-time development of the equatorial ionization anomaly in the East Asian sector, *Journal of Geophysical Research: Space Physics*, **125**, e2020JA027 973, 2020.







Figure 7. Local time variation of ionospheric ROTI during March 22-25, 2023 storm.




**Figure 8.** Local time variation of ionospheric ROTI during April 22-25, 2023 storm.



**Figure 9.** Local time variation of ionospheric ROTI during November 4-7, 2023 storm.



**Figure 10.** Local time variation of ionospheric ROTI during May 10-13, 2024 storm.



**Figure 11.** Variations in the ionospheric prompt penetration electric fields (PPEFs) over the three longitudes during the three intense geomagnetic storms of SC 25.