

Please note that the line numbers referenced in this response refer to **the version of the manuscript with tracked changes**, as they may differ from the ones in the clean version.

Response to Referee #1

Major comments

0. First, a common index for subtropical high is omega at 700hPa? I imagined the most common index is the sea surface pressure (SLP).

Yes, SLP is indeed a common index for subtropical highs, and we use it to identify the climatological location of subtropical highs (Section 2.2.1). However, in this study, we use ω_{700} and $d\Phi/dx$ in the Sc area instead of SLP in the subtropical high area, based on our hypothesized free-troposphere and surface pathways. Our updated Section 3.4 adds the relationships between SLP and the two selected factors to show the deviation between the two factors and SLP. Apart from that, we still interested in whether the hypothesized pathways work in the Sc area. Hence, we use ω_{700} and $d\Phi/dx$ in the Sc area to ensure the physical interpretability of the results.

By this comment we realized that the phrase “strength of subtropical highs” might lead readers to associate it with SLP, rather than the subsidence rate or geopotential gradient that we intended to convey through the two pathways. Hence, we’ve revised the manuscript to avoid using “strength/intensity” in this context. An additional explanation is added in L59-65.

1. The title says "Factors Causing Stratocumulus to Deviate from Subtropical High Variability". So I imagined the ultimate target is the relationship between the low cloud fraction and SLP. If so, I wanted to see the relationship between them for the seasonal and interannual time scales first. In addition, I guess this relationship (low cloud fraction and SLP) can be decomposed into the following relationships.

(A)	the low cloud fraction and EIS	x (for seasonal and interannual)
(B)	EIS and θ_{700}	o
(C)	θ_{700} and Q (defined by eq. (5))	o
(D)	Q and ω_{700}	o
(E)	ω_{700} and SLP	x
(F)	EIS and θ_{1000}	o
(G)	θ_{1000} and ocean upwelling	x
(H)	θ_{1000} and latent heat flux	x
(I)	ocean upwelling and gradient of SLP	o
(J)	latent heat flux and gradient of SLP	o
(K)	a gradient of SLP and SLP itself	x

The authors showed the important essential relationships among them (Relationships marked by "o" are (at least partially) shown. Relationships marked by "x" are not shown.). However, could the authors schematically describe in the text which relation is discussed in the present paper (because it is a bit hard to catch what is discussed and what is not discussed in the manuscript.)? The relationships that are not shown could also be helpful for the readers. Could the authors add such figures (at least (A) for seasonal and interannual, and (E).) in supplemental materials? I also want to know the correlation between EIS and ω_{700} (that is not shown in Table 1).

Thanks a lot. We agree that the listed decomposition of relationships is useful. A description of which relation is discussed is added at the end of the introduction. (L69-75)

All the relationships listed above have been analyzed:

(A) is shown in the manuscript only for all monthly means (Fig. 2), but we've added the information for the seasonal and the interannual timescales. Corresponding plots are shown below. We've also added a sentence noting the inclusion of these correlation coefficients in L141-142.

(B)-(D) are already shown in Table 1 and Figures 4-6.


(E) has been added as a replacement for the right column of Figure 9. The revised Figure 9 shows that ω_{700} in the Sc area deviates from that in the subtropical high area, and ω_{700} in the high area covaries with SLP there at different rates. Hence, the simple factor of SLP in subtropical highs cannot represent subsidence in the Sc area.

(F) is analyzed in Table 1.


(G)-(H) are added as the new Figure 8.

(I)-(J) are shown below but are not put in the manuscript as they provide the same information as the current Figure 8—that the hypothesized surface pathway is rejected.

(K) is added as Figure 10, describing that the geopotential gradient in the Sc area doesn't covary with SLP in the subtropical high area.



Plots for (A): scatter plots of low cloud fraction (κ) vs. EIS on the seasonal timescale (left) and on the interannual time scale (right). Correlation coefficients are represented by r and the regression lines (black) are presented for $p - values \leq 0.05$. Equations of regression lines are shown besides correlation coefficients. Each colour represents one region.



Plots for (I)-(J): scatter plots of geopotential gradient and latent heat flux (left) and of geopotential gradient and ocean upwelling velocity (right). The primes indicate deviations from the mean of the respective regions on the corresponding timescales. The top branch is for seasonal and the low branch is for interannual series. Regression lines are presented for $p - \text{values} \leq 0.05$, and thickened when $r^2 \geq 0.25$.

The correlation between EIS and ω_{700} has been added in Table 1.

1'. (Related to item 1.) Could the authors clarify the reason why θ_{1000} are not examined at the beginning of the analysis in the text more explicitly if the authors don't do them? Is this because "Environmental changes associated with variations in the strength of the subtropical high pressure regions correlate better with LTS (and EIS) than they do with the components of the LTS and EIS the variations are thought to influence (L240-241)"?

The reason that we didn't show the dependence of θ_{1000} on those influencing factors is that the dependence of those influencing factors on the geopotential gradient is already rejected (section 3.3). However, as both reviewers are interested in seeing that disconnection explicitly. Therefore, we added a new Fig. 8 showing the relationships between θ_{1000} and geopotential gradient, LHF, and ocean upwelling in section 3.3.

Minor comments

1. L1-2: "Stratocumulus (Sc) covers the eastern flanks of maritime subtropical high pressure systems and exerts an influence on the global energy budget comparable to CO₂."

This sentence seems to be unclear. Do the authors mean the total CO₂ radiative effect

(difference between the current CO₂ concentration level and zero CO₂)? Or the change of (doubling) CO₂ (~4 W/m²)? The influence of Sc is (the global average of) cloud radiative effect of Sc itself (difference between the current Sc existence and clear state. not change in CRE of Sc)?

We meant: comparable to a doubling of CO₂. This sentence has been corrected in L1-2.

2. L4: "gradient"

"difference" is better?

It has been replaced.

3. L39-41: "The first hypothesized mechanism would be that variations in the strength of the high influence the free-tropospheric temperature above cloud top through enhanced adiabatic warming. This mechanism follows the pioneering work of Rodwell and Hoskins (2001), who demonstrate how monsoons can influence the strength of the high to their west"

The first sentence mentions the influence of high-pressure systems on clouds (the cause is the variation in high-pressure systems and the effect is the variation in clouds). The second sentence seems to discuss the influence of the monsoons on the variation of high-pressure systems (the cause is the monsoon variations and the effect is the variations of high-pressure systems.). Are the mechanisms similar?

The connection between the first hypothesized pathway and the monsoon-desert mechanism has been clarified and this part has been rephrased in L42-46.

4. L157-164 (2nd paragraph of 3.1):

I thought the essence of these characteristics was discussed in Klein & Hartmann (1993). So it's appropriate to cite the publication here.

We've added a citation in L193-194 following the discussion of the seasonal timescale.

5. L168: "or aloft, driven by downwelling"

What is "downwelling"? Air (atmospheric) subsidence? Or ocean downwelling? Could you please clarify what downwelling or upwelling you mean when you use these terms?

It means atmospheric subsidence here. We've additionally clarified all the other uses of the terms upwelling/downwelling in the manuscript.

6. L169-171: "For the seasonal cycle, the three regions with the largest variability of the strength of the high (measured by the standard variation of ω_{700} , H in Table 1) are also the regions where θ_{700} varies more than θ_{1000} . However, this correlation does not imply causation, as the interannual series fails to exhibit the same relationship."

Especially mid-latitude (including subtropics), the seasonal cycle of θ_{700} is mainly controlled by the seasonal cycle of solar zenith angle (not by ω_{700}), I thought. I guess people generally imagine that ω_{700} cannot determine θ_{700} (at least) except tropics. Am I wrong?

The Sc regions in our study are not exclusively mid-latitude: several (NP, SP, SA) extend into the tropics, while others reach higher latitudes. Therefore, we believe it worth to explicitly test the correlation between ω_{700} and θ_{700} here.

Our goal in comparing the variability of ω_{700} and θ_{700} across regions on both timescales was to test whether such a relationship might emerge statistically under our hypothesized free-tropospheric pathway. The discussion here on seasonal timescales is not the main focus; instead, it serves structurally as a lead-in to highlight the lack of correlation on interannual timescales. We have expanded the last sentence to clarify that statement in L203-206.

7. L177-179: "The somewhat weaker relationship between EIS and LHF can be expected because while surface winds cool the surface, and hence lower θ_{1000} , increased latent heat fluxes also breakup the cloud decks"

This sentence discusses the relationship between EIS and LHF. So the fact that increased latent heat fluxes break up the cloud decks cannot be the reason for the relationship between EIS and LHF (i.e., cloud cover doesn't affect EIS directly), I guess.

Sorry for the misleading sentence. In the previous version, we implicitly treated EIS as a proxy of Sc cloudiness, which led to the inclusion of that discussion. However, we recognize that using the influence of LHF on cloudiness as support for its impact on EIS is not logically rigorous. Therefore, we have removed this sentence to avoid confusion.

8. L184: "atmospheric downwelling"

Does this mean the air (atmospheric) subsidence?

Yes, it's been changed to subsidence here.

9. Captions of Figure 3 and 6:

Plotted variables look not raw values but anomalies of the variables. In addition, I couldn't find the meaning of the apostrophe attached to the variables.

Yes, the plotted variables are anomalies there. We've added a clarification that "The primes indicate deviations from the mean of the respective regions on the corresponding timescales." in both captions.

10. L230: "cloud-top temperature"

L232: cloud-top

Generally, 700 hPa is not cloud top but above stratocumulus and in the free atmosphere, I guess.

Thanks. Yes, we automatically use the word "cloud" to represent stratocumulus in the manuscript but realized that it may be misleading. Hence, we replaced the first statement with "temperature above Sc" (L271) and the second one with "free troposphere" (L273).

11. L242: "upwelling velocity"

Can you clarify what upwelling this is? Ocean upwelling velocity?

Yes, it's ocean upwelling velocity and has been clarified. (L286)