

Strong relation between atmospheric CO₂ growth rate and terrestrial water storage in tropical forests on interannual timescales

Samantha Petch¹, Liang Feng², Paul Palmer², Robert P. King³, Tristan Quaife¹, and Keith Haines¹

¹National Centre for Earth Observation, Department of Meteorology, University of Reading, Reading, RG6 6BB, UK

²National Centre for Earth Observation, School of GeoSciences, University of Edinburgh, Edinburgh, EH9 3FF, UK

³Hadley Centre, Met Office, Exeter, EX1 3PB, UK

Correspondence: Samantha Petch (s.petch@pgr.reading.ac.uk)

Abstract. The atmospheric CO₂ growth rate (CGR) is characterised by large interannual variability, mainly due to variations in the land carbon uptake, the most uncertain component in the global carbon budget. We explore the relationships between CGR and global terrestrial water storage (TWS) from the GRACE satellites. A strong negative correlation ($r = -0.70$, $p < 0.01$, based on monthly data) between these quantities over 2002-2023 indicates that drier years correspond to a higher CGR, suggesting reduced land uptake. We then show regional TWS-CGR correlations and use a metric to assess their contributions to the global correlation. The tropics can account for the entire global TWS-CGR correlation, with small cancelling contributions from the Northern and Southern Hemisphere extratropics. Tropical America makes the dominant contribution (69%) to the global TWS-CGR correlation, despite occupying $< 12\%$ of the land surface. Aggregating TWS by MODIS land cover type, tropical forests exhibit the strongest CGR correlations and contribute most to the global TWS-CGR correlation (39%), despite semi-arid and cropland/grassland regions both having more interannual TWS variability. Tropical forests exhibit the strongest CGR correlations due to their high productivity and sensitivity to water stress, which strongly influences interannual variations in carbon uptake. An ensemble mean of ~~four~~eight atmospheric CO₂ flux inversion products also indicate a ~~74~~66% tropical contribution to CGR variability, with tropical America/Africa contributing ~~30~~27%/~~27~~28% respectively. Regarding land cover type, semi-arid/tropical forests contribute almost equally (37%/35%) to CGR variability, although tropical forests cover a much smaller surface area (25%/10%). ~~time~~Time series of global and regional TWS and CO₂ flux inversions through 2002-2023 also show changing regional contributions between global CGR events, which are discussed in relation to regional drought and ENSO events. Our study advances previous work by providing a more detailed analysis of regional contributions and doing a temporal breakdown of contributions.

1 Introduction

The atmospheric CO₂ growth rate (CGR) exhibits large amounts of year-to-year variability, which holds high significance in the context of climate change mitigation and projection. This variability is predominantly driven by fluctuations in the land carbon uptake, which has a year-to-year variability of about 1 GtCyr^{-1} , with smaller contributions from oceanic uptake and anthropogenic emissions (Friedlingstein et al., 2023). The land carbon sink results from an imbalance between the uptake of carbon through photosynthesis (GPP), the loss of carbon through ecosystem respiration (ER), and carbon losses through

25 other disturbances (D), such as fire and land use change. The net carbon flux is termed the net biome production (NBP), where $NBP = GPP - ER - D$. This land carbon sink plays a crucial role in offsetting anthropogenic CO₂ emissions, accounting for approximately 30% of these emissions each year (Friedlingstein et al., 2020). However, NBP remains the most uncertain component of the global carbon budget, with modelling studies indicating significant uncertainties in both the magnitude and sign of future projections of the sink (Ahlström et al., 2012). Reducing these uncertainties will require a better understanding of
30 the processes underlying the CO₂ fluxes and how they might change in the future, which will be essential for shaping effective mitigation policies.

It is well documented that the CGR interannual variability (IAV) is closely related to El Niño-Southern ~~Oscillation~~ Oscillations (ENSO) (Gurney et al., 2012). There are noticeable increases in CGR during El Niño events and decreases during La Niña events (Keeling and Revelle, 1985). There is also a widespread consensus that variations in tropical ecosystems exert the
35 most significant influence on global CGR, with numerous studies concentrating their efforts on the Tropics (e.g., Wang et al., 2013, 2014; Liu et al., 2023; Luo and Keenan, 2022). In particular, tropical temperature is found to have a strong positive correlation with CGR (Wang et al., 2013). ~~The Other~~ physical factors governing the land carbon uptake remain elusive. Given that tropical temperatures are typically considered optimal for photosynthesis (Huang et al., 2019), any elevation in these temperatures are anticipated to have a dampening effect on GPP while simultaneously elevating ER. This combined impact may
40 amplify the role of temperature as a significant driver of CGR variability.

While many studies have pointed to the influence of temperature on CGR (e.g., Cox et al., 2013; Wang et al., 2013), recent research is increasingly highlighting the significance of water availability as a primary control. The launch of the Gravity Recovery and Climate Experiment (GRACE) satellites in 2002 has largely helped to support this notion by providing accurate terrestrial water storage (TWS) data based on gravitational measurements. Humphrey et al. (2018) used the GRACE data and
45 demonstrated that there is a strong negative correlation between CGR variability and observed changes in global TWS (monthly $r = -0.65$ and yearly $r = -0.85$), revealing that drier years, especially in the tropics, coincide with higher CGR. The study also highlighted that this water storage relationship is clearer than for precipitation, as previous studies find there is a weaker and lagged response to precipitation. Most vegetation responds to soil moisture (Wang et al., 2016), which is a component of TWS: ~~Precipitation~~, whereas precipitation anomalies only account for water input and do not consider losses from evapotranspiration
50 or runoff. Even when combined, ~~observations of these fluxes have been shown to~~ water flux observations inadequately capture the interannual variations observed by GRACE (Petch et al., 2023a). He et al. (2022) demonstrated that interannual variability in atmospheric VPD was also significantly negatively correlated with net ecosystem productivity and hence to the interannual variability of the CGR, although this analysis relied upon VPD estimates from FluxCOM and TRENDY. In this study, we focus on the influence of TWS on CGR due to the availability of large-scale, observation-based data from the GRACE satellite mission. In contrast, VPD at global or continental scales is typically derived from model-based products or reanalysis data. While TWS is not necessarily a better explanatory variable to temperature or VPD, it offers a complementary perspective and allows us to leverage independent observational datasets.

Separating the individual contributions of various climatic drivers to CGR fluctuations can be challenging due to their often intertwined nature. For example, VPD and soil moisture are coupled, where high VPD usually corresponds to dry soils and

60 low VPD to wet soil. As a result, vegetation responses to water stress are likely to reflect the combined influence of both
variables, making their individual effects difficult to disentangle. Temperature and water availability are also interrelated,
further complicating attribution. Nevertheless, Humphrey et al. (2018) showed that the relationship between CGR and TWS
stands independently of temperature influences, suggesting that TWS captures a distinct and meaningful component of CGR
variability.

65 Wang et al. (2022) then showed that the relative influence of temperature and water on net ecosystem exchange (NEE) IAV
shifts across different regions and seasons. They employed three different approaches: atmospheric inversions, process-based
vegetation models from TRENDY, and a data-driven model (FLUXCOM). Their findings reveal broad agreement that the
tropics are a key driver of global correlations; however, the dominant driver of global NEE IAV varied due to disagreements
70 ing the relative magnitudes of water and temperature contributions in the Northern Hemisphere for determining the dominant
drivers of NEE IAV. Another recent study by Liu et al. (2023) provides evidence that the coupling between interannual CGR
and TWS is becoming increasingly strong. They report an increase of around 35% in CGR sensitivity to tropical water vari-
ations from 1989-2018 compared to 1960-1989, however, existing climate models do not exhibit signs of this rising trend in
tropical water-CGR coupling (Liu et al., 2023). Models are also found to have a tendency to underestimate the strength of the
75 coupling (Humphrey et al., 2018). In general, interactions between TWS variations and the carbon cycle ~~is~~are a key uncertainty
in current climate models that could strongly influence CGR over the coming decades.

Despite the dominant role of terrestrial ecosystems in influencing CGR variability, the heterogeneity of these ecosystems
means pinpointing the specific land regions contributing to ~~the~~-IAV can be challenging. ~~There~~and there are inconsistencies
among previous studies concerning the land cover types contributing to ~~the~~-IAV. Ahlström et al. (2015) used process based
80 models and found that semi-arid ecosystems could explain 39% of the global NBP variability over 1982-2011, thereby exerting
dominant control over carbon sink interannual fluctuations, while the mean sink was primarily influenced by tropical forests.
Zhang et al. (2016) found that semi-arid regions contributed 57% of the detrended interannual variability in global GPP. These
results are supported by Humphrey et al. (2018), who used GRACE to demonstrate that TWS in semi-arid regions contribute
the most to the global storage variability, while also mentioning a possible role of TWS in tropical forests which is also well
85 correlated with the CGR.

Marcolla et al. (2017) examined the IAV in the terrestrial carbon budget using three different datastreams: terrestrial ecosys-
tem level observations from FLUXNET (La Thuile and 2015 releases; Baldocchi et al., 2001), a bottom-up global product
resulting from upscaling site level fluxes (the MPI-MTE; le Maire et al., 2010), and a top-down inversion product (Jena Car-
bonscope; Rödenbeck et al., 2003). They found significant discrepancies among these datastreams regarding the main sources
90 of temporal variability, particularly in the tropics, where there is a lack of atmospheric and ecosystem observations. However,
all products unanimously identified several crucial global features, in particular the high relative IAV in the terrestrial carbon
cycle in water-limited ecosystems. Similarly, Piao et al. (2020) used three different approaches, including land carbon cycle
models, a data-driven model (FLUXCOM; Jung et al., 2019), and atmospheric inversion models (taking the mean of CAMS
and Jena CarbonScope; Chevallier et al., 2010; Rödenbeck et al., 2003), to investigate regional contributions to global terrestrial

95 carbon IAV. They found the share in contributions of tropical semi-arid regions versus tropical ~~non-semiarid~~non-semi-arid regions was similar between approaches. However, they found relatively larger contributions from the extratropics in atmospheric inversions compared to other approaches, possibly due to limited surface CO₂ observations over the tropics. These data are used exclusively in the atmospheric inversions, affecting their ability to discern IAV between the tropics and the extratropics.

Our research first aims to assess the global relationship between TWS and CGR interannual variability, extending Humphrey
100 et al. (2018) analysis in time from 2016 up until 2023 and highlighting the global CGR variability we are seeking to explain. This period notably includes some of the largest values of CGR on record. We then look to regionalise the contributions based on both land cover type and large spatial regions, and look at interannual variability events through the whole period. Additionally, we use eight atmospheric CO₂ inversion products to infer regional contributions of CO₂ fluxes. This approach allows us to make similar assessments without relying on assumptions of the relationships between water and CGR. We assess
105 the agreement between the inversion products and GRACE, as well as the consistency among inversion products. We also look at major temporal events in the CGR record to examine which regions are contributing most at different times and assess the consistency between these approaches. We also use the inversions to estimate regional-scale sensitivity of NBP to TWS variability.

Section 2 discusses the data and analysis methods we use to attribute the regional contributions to global signals. In sec-
110 tion 3, we examine the relationships between GRACE TWS on global and regional scales, and the CGR signal on interannual timescales from 2002-2023. Section 4 looks at ~~four~~eight atmospheric CO₂ inversion models from GBC2023 (Friedlingstein et al., 2023) and their ability to regionalise the terrestrial sources of CO₂ over the same time period. Section 5 discusses these results in relation to meteorological conditions through the period, and section 6 provides a summary and conclusions.

2 Data and Methods

115 2.1 GRACE

Terrestrial water storage data are obtained from the Gravity Recovery and Climate Experiment (GRACE) and its successor mission, GRACE Follow-On (GRACE-FO). The version used here is the Jet Propulsion Laboratory (JPL) Mascon RL06v2 (Wiese et al., 2016). This dataset contains gridded monthly global water storage/height anomalies relative to a time-mean, derived from GRACE and GRACE-FO and processed at JPL using the Mascon approach. The water storage/height anomalies
120 are given in equivalent water thickness units (cm). This version of the data employs a Coastal Resolution Improvement filter that reduces signal leakage errors across coastlines. Data can be obtained from http://grace.jpl.nasa.gov/data/get-data/jpl_global_mascons/. There were a small number of months with missing data which were filled with monthly climatology plus the temporal interpolation of monthly storage anomalies.

2.2 GML surface flasks

125 Measurements of atmospheric CO₂ concentration are taken from the Global Monitoring Laboratory (GML) of the National Ocean and Atmospheric Administration (NOAA) (Lan et al., 2023). This dataset compiles measurements from the Cooperative Global Air Sampling Network, where air samples are collected approximately weekly from a globally distributed network of sites. Data can be found at <https://gml.noaa.gov/ccgg/trends/global.html>.

2.3 Atmospheric CO₂ inversions

130 This study uses ~~four~~eight different atmospheric CO₂ inversion products, summarised in Table 1, selected based on their temporal coverage, all of which span at least from 2002 to ~~2020~~2022. These products are from GCB2023 where the land biosphere fluxes have been adjusted to a common fossil fuel emissions dataset (Friedlingstein et al., 2023). Other similar products were discounted on the grounds they only cover shorter periods. Top-down inverse models provide spatially and temporally resolved estimates of the net CO₂ flux exchanged between the surface and the atmosphere. They are generated as source/sink solu-
135 tions using atmospheric transport models made to fit surface flask based atmospheric CO₂ mole fraction measurements within their uncertainties, at various locations (Ciais et al., 2022). Typically, these inversion products employ a Bayesian statistical approach, where an optimal surface CO₂ flux is determined as the maximum likelihood estimate within a statistical distribution of possible fluxes. The products also use prior surface flux values and their associated uncertainty distributions. We also examined the interannual variability in the prior fluxes, where available, and in all cases found very little relationship with the
140 posterior fluxes, and hence conclude that the prior choices were not directly influencing our results.

~~The products used in this study include CarbonTracker-CT2022, results provided by NOAA-GML, Boulder, Colorado, USA from the website at . The Copernicus Atmosphere Monitoring Service (CAMS) global inversion-optimised CO₂ product version v22r1 generated with the inversion system PYVAR, available from . Long-term global CO₂ fluxes estimated by the NICAM-based Inverse Simulation for Monitoring CO₂ (NISMOM-CO₂) v2023, available from , and the University of Edinburgh (UoE) in situ product (Feng et al., 2016). The differences between products concern the selection of~~
atmospheric inversion products differ in their use of transport models, meteorological inputs, and prior flux assumptions—all of which can significantly influence flux estimates, particularly when comparing tropical and extratropical regions (Peylin et al., 2013; Chevall
. In the tropics, flux estimates are especially sensitive to how transport models represent deep convection and vertical mixing, while extratropical estimates are more influenced by synoptic-scale advection and boundary layer dynamics. Prior flux assumptions,
such as prescribed seasonal cycles or vegetation responses, can also introduce regional biases—especially in the tropics where these assumptions often fail to capture complex climate–ecosystem interactions (Munassar et al., 2022; Gaubert et al., 2019).
Moreover, the relative scarcity of CO₂ data, the choice of prior fluxes, and the transport model used. They also differ in spatial and temporal resolution, assumed correlation structures, and mathematical data assimilation approaches (Friedlingstein et al., 2020)
–observations in tropical regions amplifies the impact of model structure and prior uncertainty, whereas denser observational
155 networks in the extratropics provide stronger constraints on inversion results (Patra et al., 2005; Schuh et al., 2019).

The UoE and ~~CT2022-CT NOAA~~ products both adopt an ensemble Kalman filter (EnKF) approach, whereas NISMUN-CO₂ employ a BFGS-based quasi-Newton method. CAMS on the other hand uses a variational approach and Jena CarboScope employs a Bayesian inversion framework. For the prior terrestrial fluxes UoE and ~~CT2022-CT NOAA~~ use the CASA biogeochemical model introduced by Potter et al. (1993), NISMUN-CO₂ use data from the Vegetation Integrative SIMulator for Trace gases (VISIT) (Ito, 2019), whereas the land priors are climatological in the CAMS product. ~~For fire emission data, versions MIROC uses both CASA and VISIT Chandra et al. (2021), and Jena CarboScope uses the LPJ biosphere model (Sitch et al., 2000). The 8 products share harmonized fossil fuel and fire emissions, all using the same version of the Global Fire Emissions Database (GFED) were used in each product (van der Werf et al., 2017).~~

2.4 MODIS

Land cover types are classified using the Terra and Aqua combined Moderate Resolution Imaging Spectro-radiometer (MODIS) Land Cover Climate Modelling Grid product MCD12Q1 Version 6.1, available from <https://lpdaac.usgs.gov>. We use the International Geosphere-Biosphere Programme (IGBP) legacy classification schemes, and group into six surface classes, shown in ~~Figure ??~~Supplementary Figure 1. These will be used to determine aggregated flux information.

~~(a) Map of spatial regions used in study. Percentage land area coverage: Northern Hemisphere extratropics (54%), Southern Hemisphere extratropics (8%), Tropical America (12%), Tropical Africa (18%), Tropical Eurasia and Australia (8%). (b) Land cover vegetation types developed from MODIS data and used for regional flux analysis. Percentage land area coverage: Grassland and croplands (32%), Semi-arid (25%), Tundra and Arctic shrubland (15%), Sparsely vegetated (12%), Tropical forest (10%), and Extratropical forest (5%).~~

2.5 Data processing and analysis framework

The CGR is derived by taking the differential of monthly atmospheric CO₂ concentration data. We calculate the interannual variability by first removing the mean seasonal cycle from the monthly data, then removing the linear trend. We assume that by removing the linear trend this removes the main fossil fuel driving signal. However, this could also remove any trend in biogenic and ocean fluxes. The CGR time series and CO₂ inversion products are then smoothed with a 12 month moving mean, while the GRACE TWS data is smoothed using a 6 month moving mean, following (Humphrey et al., 2018). Different smoothing windows are used because the CGR data is inherently noisier than the TWS data, therefore requiring slightly stronger smoothing to reveal meaningful interannual variability. To convert the global CGR from ppm to GtC we use the conversion factor 1 ppm by volume of atmosphere CO₂ = 2.13 GtC. ~~All gridded inverse model products are resampled to a 0.5 degree grid in order to fit with the GRACE data.~~ We generate an ensemble mean of inversion products spanning from January 2002 to December 2022, with each product equally weighted. ~~Due to CT2022 only being available until December 2020, the ensemble incorporates only the remaining three products in 2021 and 2022.~~

To assess the importance of different regions to a global total, we use the “contribution index” (f_j), as defined by Ahlström et al. (2015) also adopted by Zhang et al. (2019) and Humphrey et al. (2018). This index quantifies the spatial contributions to

Table 1. Summary of atmospheric CO₂ flux inversion products.

Product	Temporal coverage Transport model	Meteorol. fields	Fossils	Fires Reference
CAMS	1979-2023 LMDZ	ERA5-Interim	GCP-GridFED	GFED —Chevallier (2023), Chevallier et al. (2005)
UoE	2001-2022 —GEOS-Chem	MERRA2	Oda and Maksyutov (2011)	GFEDv3.0 —Feng et al. (2009), Feng et al. (2016)
NISMON-CO ₂	1990-2022 —NICAM-TM	JCDAS, JRA-55	GCP-GridFED	GFEDv4.1s —(Niwa et al., 2022), Niwa et al. (2017)
NOAA Carbon-Tracker CT2022	2000-2020 —TM5	ERA5-Interim	Millier, ODIAC	GFEDv4.1s Jacobson et al. (2023) <u>Jacobson et al. (2023)</u>
<u>CarbonTracker Europe (CTE)</u>	<u>TM5</u>	<u>ERA-Interim</u>	<u>EDGAR4.2</u>	<u>van der Laan-Luijkx et al. (2017)</u>
<u>Jena CarboScope</u>	<u>TM3</u>	<u>NCEP reanalysis</u>	<u>EDGAR</u>	<u>Rödenbeck et al. (2003, 2018)</u>
<u>IAPCAS</u>	<u>GEOS-Chem</u>	<u>GEOS-5</u>	<u>Oda and Maksyutov (2011)</u>	<u>Feng et al. (2016); Yang et al. (2021)</u>
<u>MIROC4-ACTM</u>	<u>MIROC4-ACTM</u>	<u>JRA-55</u>	<u>GridFED</u>	<u>Chandra et al. (2021)</u>

the global monthly time series t , scoring regions based on how consistent their regional flux IAV is with the sign and magnitude of the global IAV. It is given by:

$$f_j = \frac{\sum_t \frac{x_{jt}|X_t|}{X_t}}{\sum_t |X_t|} \quad (1)$$

where x_{jt} represents the detrended regional data j at time t , and $X_t = \sum x_{jt}$ denotes the detrended global value. Regions with higher positive values of f_j contribute more to the global variations. This is used on both the regional GRACE TWS variations, and on regional CO₂ fluxes from the atmospheric inversion products. Regions may represent continental areas or areas with particular land cover classes depicted in [Figure ??-Supplementary Figure 1](#).

To connect the TWS and CGR time series we use the Pearson correlation coefficient (r). Additionally, we look to identify the regional origin of the global TWS (GTWS) correlation with CGR. Let GTWS represent the global TWS signal and TWS _{i} represent the storage signal at grid point i . The relationship between the global correlation $r_{GTWS,CGR}$ and grid point correlations $r_{TWS_i,CGR}$ can be expressed as;

$$r_{GTWS,CGR} \times \sigma_{GTWS} \times A = \sum_i r_{TWS_i,CGR} \times \sigma_{TWS_i} \times a_i, \quad (2)$$

where a_i is the area of grid box i , A is the total land surface area such that $A = \sum_i a_i$, and σ_{GTWS} and σ_{TWS_i} are the temporal standard deviations of the global and grid point storage signals, respectively. This equation holds due to the additive properties of covariance. Hence, the contribution of a particular region (R) to the global GTWS-CGR correlation, which we will denote as g , can be expressed as a percentage as follows;

$$g(R) = \frac{\sum_{i \in R} a_i \times \sigma_{TWS_i} \times r_{TWS_i,CGR}}{A \times \sigma_{GTWS} \times r_{GTWS,CGR}} \times 100. \quad (3)$$

This helps identify regions which may have a smaller contribution to the GTWS variability but which are more strongly correlated to CGR variations. This is similar to the metric used by Wang et al. (2022) in their equation (4).

3 CO₂ Growth Rate (CGR) and Terrestrial Water Storage (TWS)

Extending the results from Figure 1a of Humphrey et al. (2018), our Figure 1 shows the de-trended time series of global terrestrial water storage from GRACE (note the reversed axis direction for TWS), and the CGR derived from surface fluxes, for the period January 2002 to December 2023. [It is important to note that, although atmospheric CO₂ concentrations continue to increase each year, the CGR has been detrended; therefore, negative CGR anomalies reflect periods of below-average growth relative to the long-term mean growth rate.](#) A negative correlation of $r = -0.70$ ($p < 0.01$) is found, implying that reduced terrestrial water availability corresponds to lower net land uptake of carbon, consequently leading to a higher atmospheric CGR. A correlation of $r = -0.69$ is observed for the period from 2002 to 2016 as analysed by Humphrey et al. (2018), and

215 $r = -0.74$ from 2016 to 2023. The strongest CGR peak in 2016 is clearly associated with the strong El Niño that occurred, with associated drying across many land areas (Santoso et al., 2017; Wigneron et al., 2020). We will investigate regional relationships later in the paper, but this figure clearly underscores the persistence of the TWS-CGR relationship beyond 2016 shown by Humphrey et al. (2018).




Figure 1. Comparison of the interannual variability of global terrestrial water storage (GTWS) from GRACE (right inverted axes) and atmospheric CO₂ growth rate (CGR) (left axes) from 2002 to 2023. Time series have been de-seasoned and de-trended. The grey dashed line marks the end of the Humphrey et al. (2018) analysis.

To highlight smaller scale features of this relationship, Figure 2a shows the monthly correlation between GRACE TWS
 220 IAV at each grid point and the global CGR time series. The map has been filtered to only show correlations significant at the level of $p < 0.01$, which corresponds to a minimum correlation coefficient of $r = \pm 0.16$. We also calculated the map using the Spearman's correlation coefficient (results not shown) and the spatial patterns were the same as in Figure 2a. The most prominent negative correlations appear over the eastern Amazon and ~~extending extend~~ across eastern parts of tropical South America. Other areas of negative correlation in the tropics extend through India, Southeast Asia and northern Australia. There is
 225 also a substantial region in northeastern Siberia showing strong negative correlation. However, not all ~~of these areas contribute much areas with high local correlations contribute meaningfully~~ to the total ~~GTWS-CGR correlation simply because their TWS variations may be small.~~ Figure 2b shows GTWS-CGR relationship. This is often due to either a small magnitude of TWS variability or a limited spatial extent, which reduces their overall influence on the global signal. To address this, we evaluate the

contribution of each $1^\circ \times 1^\circ$ grid box to the GTWS-CGR signal from Figure 1, based on equation 1, such that the sum of all grid boxes equals 100%. This enables a visualisation of the relative importance of different regions in governing the GTWS-CGR IAV. Grid boxes with higher positive scores indicate a stronger influence, while those with smaller or negative scores are indicative of regions with lesser or counteracting contributions. From this figure $1^\circ \times 1^\circ$ grid cell to the global GTWS signal using Equation 1, as shown in Figure 2b. This contribution metric effectively downweights regions with minimal influence due to low TWS variance allowing a clearer assessment of the regions that dominate GTWS IAV. In this representation, tropical areas stand out much more strongly, as the metric incorporates the magnitude of TWS variability in addition to correlation strength.

From Figure 2, it can be seen that in certain regions, TWS variability exhibits positive correlations with the global CGR, such as in South Brazil and East China. There could be possible biophysical explanations for this result, for instance, it was suggested that the Brazilian Southeast is a transition region characterized by rainfall anomalies with opposite signals related to ENSO (Coelho et al., 2002). It is also possible that these regions simply do not contribute substantially to the global CGR, and the observed positive correlations may reflect local or regional processes that are largely cancelled out at the global scale. Determining the underlying causes of these correlations would require reliable estimates of regional-scale carbon fluxes, but as will be discussed in Section 4, atmospheric CO_2 inversion products currently do not do a good job at capturing fine scale interannual variability, limiting their utility for such attribution.




Figure 2. (a) Correlation map of GRACE TWS and CGR for the period 2002-2023. Only areas with $p < 0.01$ are shown. (b) Contribution of each $1^\circ \times 1^\circ$ grid point to total global TWS-CGR-TWS signal, expressed as a percentage.

Figure 3 shows correlations between regional TWS and the global CGR in blue on the left axis, and the right axis shows the regional contribution to the GTWS signal (f , based on equation (1) in red, and the regional contributions to the GTWS-CGR correlation (g) following equation (3) in orange. To aid interpretation, the regional domains cover the following proportions global land area: Northern Hemisphere extratropics (54%), Southern Hemisphere extratropics (8%), Tropical America (12%), Tropical Africa (18%), and Tropical Eurasia and Australia (8%). Percentage coverage by vegetation type is as follows:

250 Croplands and Grasslands (32%), Semi-arid (25%), Tundra and Arctic shrubland (15%), Sparsely vegetated (12%), Tropical forest (10%), and Extratropical forest (5%). The continental values in Figure 3a show that the tropics alone can explain all of the GTWS-CGR correlation, with the Northern Hemisphere (NH) and Southern Hemisphere (SH) extratropics providing small cancelling contributions.

255 The tropics play a disproportionately large role in driving the covariance between CGR and regional TWS. This is due to a combination of factors: (i) the tropics exhibit some of the largest magnitudes of TWS IAV (ii) Climate anomalies such as ENSO events often synchronise drought and temperature anomalies across vast tropical regions, creating spatially coherent signals that are amplified at the global scale. In contrast, temperate regions typically experience more localized and less synchronised climate variability, resulting in smaller contributions to global CGR variance.

Notably, tropical America makes the largest contribution to the GTWS-CGR correlation ($g = 69\%$), despite only representing
260 around 12% of the land surface area. Table 2 shows the aggregated correlations and contributions for some additional regions. However, it is important to note when aggregating grid boxes into regional or global averages, compensatory effects can arise, where TWS anomalies in different regions covary negatively (e.g., due to ENSO patterns), potentially cancelling each other out. This can obscure the true influence of individual regions on the global signal, as spatial covariances are not fully accounted for, leading to an oversimplification of their relative contributions. For example, Jung et al. (2017) demonstrates that the effects
265 of soil moisture on NEE operate at relatively small spatial scales (in comparison to temperature), however, this does not seem to prevent the strong impact of TWS on the CGR at the global scale, as seen by Humphrey et al. (2018) and in our results.

Figure 3b captures TWS variability by land cover type across the whole time period. The correlation between TWS and CGR variations is strongest in tropical forests ($r = -0.55$, $p < 0.01$), followed by semi-arid regions ($r = -0.46$, $p < 0.01$), but correlation alone does not consider the actual magnitude of TWS variations. The contribution index to GTWS signal (equation
270 1) and the contribution to the GTWS-CGR correlation (equation 3), also shown in Figure 3b, provide greater insight into the role of each land cover type. Croplands and grasslands, as the dominant land cover type covering 32% of the Earth's land surface area, largely contribute to GTWS signal ($f = 22\%$). However, they have a relatively low correlation with CGR and overall do not contribute much to the GTWS-CGR correlation ($g = 15\%$). Tropical forests on the other hand, emerge as the primary land cover type influencing the GTWS-CGR correlation ($g = 37\%$), despite only covering 10% of the land surface.
275 Semi-arid regions also play an important part, contributing a substantial fraction to both the GTWS signal ($f = 25\%$) and the GTWS-CGR correlation ($g = 26\%$). However, semi-arid regions cover approximately 25% of the land surface area, a much larger portion than tropical forests, highlighting the much stronger role of tropical forests in governing the GTWS-CGR relationship per unit area.

~~(a) Correlation between TWS signal for different regions and global CGR is represented by the blue bar on left axis. Contribution of continental/hemispheric regions to global TWS-CGR correlation is shown in orange, and contribution to global TWS signal is shown in red against the right axis. (b) This shows the contributions from each land cover type defined in Figure ??, using the same colour key.~~

~~Regional correlations between GRACE TWS and global Atmospheric CGR variability (first column), regional contributions to the global TWS signal (second column), and contributions to the global TWS-CGR correlation (third column) from 2002~~




Figure 3. (a) Correlation between TWS signal for different regions and global CGR is represented by the blue bar on left axis. Contribution of continental/hemispheric regions to global TWS-CGR correlation is shown in orange, and contribution to global TWS signal is shown in red against the right axis. (b) This shows the contributions from each land cover type, using the same colour key. The dashed line represents the global TWS-CGR correlation.

285 to 2023. Percentage of total land surface area (fourth column). Region Correlation Contrib. to GTWS Signal Contrib. to GTWS-CGR Correlation Land Area % Global -0.70 -1.0 -1.0 100 Tropics -0.61 -0.74 -0.98 37 Extratropics -0.02 -0.26 -0.02 63 Amazon -0.51 -0.13 -0.28 4 All Africa -0.21 -0.23 -0.15 23 All India -0.33 -0.09 -0.10 2 All Australia -0.11 -0.01 -0.05 5

Table 2. Regional correlations between GRACE TWS and global Atmospheric CGR variability (first column), regional contributions to the global TWS signal (second column), and contributions to the global TWS-CGR correlation (third column) from 2002 to 2023. Percentage of total land surface area (fourth column).

<u>Region</u>	<u>Correlation</u>	<u>Contrib. to GTWS Signal</u>	<u>Contrib. to GTWS-CGR Correlation</u>	<u>Land Area %</u>
<u>Global</u>	<u>-0.70</u>	<u>1.0</u>	<u>1.0</u>	<u>100</u>
<u>Tropics</u>	<u>-0.61</u>	<u>0.74</u>	<u>0.98</u>	<u>37</u>
<u>Extratropics</u>	<u>0.02</u>	<u>0.26</u>	<u>0.02</u>	<u>63</u>
<u>Amazon</u>	<u>-0.51</u>	<u>0.13</u>	<u>0.28</u>	<u>4</u>
<u>All Africa</u>	<u>-0.21</u>	<u>0.23</u>	<u>0.15</u>	<u>23</u>
<u>All India</u>	<u>-0.33</u>	<u>0.09</u>	<u>0.10</u>	<u>2</u>
<u>All Australia</u>	<u>-0.11</u>	<u>-0.01</u>	<u>0.05</u>	<u>5</u>

Figure 4 presents a time series of the regional TWS variations through the whole period 2002-2023. Figure 4a regionalises across the five spatial regions shown in Figure 3a. Each region is accounted for such that the sum of positive and negative contributions equals the GTWS, in black, while the CGR variations are also shown against a separate axis. While the NH extratropics show large TWS variations this often tends to be anti-correlated with TWS elsewhere, and there is little correlation with the CGR. By contrast there is a particularly consistent alignment between the timing of many CGR events and the variations in TWS across tropical America. After 2020 the CGR and TWS signals both become more variable without the clear peaks or troughs seen earlier, although there is still reasonable alignment of the GTWS and CGR signals up to the end of 2022. Overall, this figure demonstrates that tropical regions generally contribute significantly to the GTWS signal and are consistently coherent with CGR variations. Whereas, even substantial variations in the NH extratropics, where there is a lot of anti-correlation with the GTWS, are thus unlikely to be significantly impacting the CGR. This pattern is also true for the SH extratropics, although the magnitude of variations is smaller here than in the NH.

Now we consider regionalising the IAV in GRACE TWS and its correlations with the CGR, according to land cover types, using the MODIS land cover map ~~shown in Figure ??~~([Supplementary Figure 1](#)). Figure 4b shows the contribution to the GTWS IAV for each land cover type plotted against the right (inverted) axis, with the CGR on the left axis. The largest CGR peak due to the 2015/2016 El Niño event very closely corresponds to the water storage variations seen in tropical forests, where there is




Figure 4. GRACE terrestrial water storage (TWS) interannual variability and global CO₂ growth rate (CGR) 2002-2023. (a) aggregated by continental regions (b) by land cover type.

a substantial TWS minimum over this period. The CGR returned to reference levels in 2017, and global TWS recovered shortly afterwards. Note that we might also associate the increase in CGR around 2005 with the tropical forests, although this event is not noticeable in the global storage signal. This increase coincides with the 2005 drought in the Amazon driven, not by El Niño, but by elevated tropical North Atlantic sea surface temperatures, affecting the southern two-thirds of Amazonia (Phillips et al., 2009). Then in 2009, a notable drop in CGR aligns with a peak in tropical forest TWS, an occurrence associated with the exceptional flood season observed in the lower Amazon basin during the first half of 2009, supposedly linked to the 2008–2009 La Niña event (Chen et al., 2010). This is followed by a sharp peak in CGR which coincides with the another drought of 2010 in Amazonia, more severe than the drought in 2005. This event started during El Niño and then was intensified as a consequence of the warming of the tropical North Atlantic (Marengo et al., 2011).

A decrease in CGR can also be observed in 2011, indicative of a strong land carbon sink. This sink anomaly has been corroborated by independent observations from MODIS EVI and GOME-2 SIF showing much enhanced photosynthetic activity over this period (Ma et al., 2016), as well as by GOSAT XCO₂ (Detmers et al., 2015). This feature relates to an increase in TWS in semi-arid regions, primarily resulting from a wet period in Australia during 2010-2011. Large parts of semi-arid Australia are covered by endorheic basins, where TWS variations are highly sensitive to precipitation (Petch et al., 2023b). Our results align with Poulter et al. (2014), who found that nearly 60% of carbon uptake over this period could be attributed to semi-arid Australian ecosystems, which experienced several consecutive seasons of increased precipitation due to prevalent La Niña conditions. However, other features in the CGR variability are less clearly associated with TWS variations partitioned in this way. In 2020, the CGR exhibited an opposite response compared to what we normally observe in relation to the tropical forest TWS anomaly. The global TWS anomaly remained positive, possibly due to conditions in other regions (notably wet in NH extratropics and Tropical Africa from Figure 4a). This could also be an indication of a lack of TWS influence on the CGR and may be caused by external factors, such as reduced human emissions due to the COVID-19 pandemic.

~~GRACE terrestrial water storage (TWS) interannual variability and global CO₂ growth rate (CGR) 2002-2023. (a) aggregated by continental regions (b) by land cover type.~~

A clear difference between the Figures 4a,b TWS signals is that the vegetation cover contributions tend to be more aligned with the global TWS signal without the strong cancellations seen in continental TWS. This is largely explained by croplands and grasslands which often have large compensating TWS anomalies distributed between the NH extratropics and the tropical zones on all continents. This can be demonstrated by the disappearance of the pronounced anomalies depicted in Figure 4a when croplands and grasslands are excluded from the analysis ([see supplementary Figures 2 and 3](#)). This is an example of compensatory effects of water anomalies discussed in Jung et al. (2017), which could lead us to underestimate the [overall CGR](#) role of croplands and grasslands.

4 Regional flux contributions to CGR estimated by atmospheric inverse methods

The previous section looked at regionalising TWS with respect to CGR correlations. In contrast this section will try to regionalise CGR contributions by using atmospheric CO₂ inversion model products. This approach offers the advantage of identifying the regions that [directly](#) contribute most to CGR variability without depending on the coupling between TWS and CGR. Additionally, these products potentially allow us to gain direct insight into the relative magnitude of the CO₂ flux contributions from each region, which ~~previously relied on~~ [were previously only inferred from](#) regional TWS variations. The reliability of this approach will however be limited by the quality of inversion products. Therefore, we employ multiple products to assess their agreement with each other, as well as with the previous GRACE results, to provide further insights ~~into their reliability~~.

Figure 5 compares the global land surface CO₂ flux created from the ensemble of ~~four~~ [eight](#) different inversion products, and the CGR interannual variability derived from surface flasks. The shaded region represents the maximum and minimum across the products around the mean. The products, when integrated globally, show good agreement with each other and with the surface flask based CGR in terms of correlation, although there are substantial variations in the amplitude of the resulting




Figure 5. Interannual variability of observed atmospheric CO₂ growth rate (CGR) derived from GML NOAA surface flasks (red dashed), and global terrestrial carbon flux interannual variability from an ensemble of four-eight atmospheric CO₂ inversion products over 2002-2022. Ensemble mean, black, with product range in grey.

345 terrestrial IAV. These results underscore that the majority of the IAV originates from the land, with minimal contributions coming from variations in ocean uptake and fossil fuel emissions. However, there is reduced agreement between the CGR and the inversion products between 2020 and 2021, e.g. CGR observations and inversion anomalies may be of opposite sign. The low CGR may result from anthropogenic emissions being 8.8% lower in the first half of 2020 compared to the same period in the previous year, as a result of the COVID-19 pandemic impacting human activities (Liu et al., 2020). In this case pure
350 detrending is unlikely to remove the anthropogenic signal in the CGR, which could explain this result.

In order to compare the inversion products at finer spatial scales, Figure 6 shows the contribution of each land grid point to the global land surface flux IAV, from equation (1), for each inversion product over the period January 2002 to December 2022 ~~(or-until-December-2020-for-CT2022). These maps exhibit varying levels of spatial detail due to the original resolutions of the data-products-2022.~~ While there are some similarities on larger scales (see below), the maps also highlight many differences
355 between the products. Such discrepancies will limit the value of these products for fine scale analysis.

The aggregated regional CO₂ contributions for each inversion product, expressed as a percentage of the global CGR, are shown in Figure 7. The ensemble means show agreement with GRACE in attributing most of the terrestrial carbon variability to the tropical regions ($f = 74\%$), based on equation (1). NISMON-CO₂ strongly favours tropical America, while CAMS and UoE show tropical Africa as the dominant contributor, ~~and-CT2022 largely favours~~. IAPCAS and MIROC largely favour the
360 NH extratropics. We will quantify this spread among products later.

There is general consensus among the products regarding contributions from land cover types, Figure 7b, with all products attributing nearly all CGR variability to three vegetation types: semi-arid areas, tropical forests, and grasslands/croplands. The ensemble means highlight semi-arid regions ($f = 38\%$) and tropical forests ($f = 35\%$) as the primary contributors, aligning




Figure 6. Maps of $1^\circ \times 1^\circ$ grid point contributions to global land carbon flux IAV for four-eight atmospheric CO₂ inversion products. **Note,** the NISMON-CO₂ map is depicted over a larger range due to higher contribution of tropical forests.

with the GRACE data, which identified these two regions as the most dominant contributors to the GTWS-CGR correlation in

365 Figure 3b. ~~The-~~

~~However, the NISMOM-CO₂ tropical forests contribution is an outlier. Posterior~~ product stands as an outlier, assigning a particularly large contribution to tropical forests. A contributing factor may also be the use of the NICAM transport model, which has previously been shown to yield distinct flux patterns compared to other models, especially in the tropics (Peylin et al., 2013). For instance, NICAM has produced stronger carbon release signals and broader IAV during major El Niño events such as
370 1997/1998. Additionally, sparse observational constraints in tropical regions may exacerbate differences between inversion systems, allowing model structure and transport errors to exert a strong influence. Nevertheless, this observational limitation is common across systems and is therefore unlikely to be the sole driver of NISMOM's divergence. A more targeted model intercomparison would be necessary to isolate the specific causes of this discrepancy.

~~We also find that posterior~~ estimates for large regions ~~are clearly driven by observations, as their IAVs show no obvious~~
375 ~~similarities to exhibit~~ IAV that is not present in the a-priori biosphere flux estimates, in the products for which ~~these~~ priors were available. In ~~addition some priors have only seasonal variability but no IAV~~ many cases the priors include only seasonal cycles and lack substantial interannual variation, indicating that the IAV in the posterior fluxes is not inherited from the prior. This means that observational constraints and transport model dynamics play a key role in shaping the posterior variability.

Interestingly, there is much less consensus regarding spatial region contributions from Figure 7a. For example, IAPCAS
380 shows minimal contributions from tropical America, while contributions from the northern extratropics vary widely across products. IAPCAS and CTE attribute around 40% of CGR variability to the NH extratropics, whereas CAMS attributes only 12%. This spread reflects a known challenge in attributing carbon variability to the NH extratropics, particularly when relying on surface observation inversions. Previous studies have highlighted that inversions using only in situ CO₂ data may underestimate variability in these regions. Guerlet et al. (2013) for example, suggest that satellite observations show more
385 substantial flux variability in the NH extratropics than is captured by surface-only inversions. ~~However it is possible that varying uncertainties in prior estimates may still be selecting some regions to permit more posterior IAV than others.~~

Figure 8 presents time series illustrating how different regions contribute to the total land CO₂ flux variability, based on the ensemble mean of the inversion products, with the CGR line shown in red. This allows us to see how different regions may serve as dominant drivers of CGR variability at different times throughout the 20-year period, similar to Figure 4. The colour
390 bar inserts on Figure 8 represent the standard deviations of the ~~four eight~~ inversion products around the ensemble mean for the whole 20-year period. ~~These bars are designed to share,~~ sharing the same scale as the main axis used in the figure. The standard deviations for continental regions average 0.32 GtC yr⁻¹, whereas for land cover types ~~it average the average is significantly~~ lower, 0.21 GtC yr⁻¹. We also examine how the contribution from each land cover type varies during different temporal CGR events, in Figure 8b.

395 During the period of elevated CGR levels around 2002 (reduced uptake by the land), similar contributions are observed from semi-arid regions and croplands and grasslands, with some input from tropical forests. This peak is thought to possibly be associated with Northern Hemisphere fires (Jones and Cox, 2005). The decrease in CGR observed in 2004 can primarily be attributed to cropland and grasslands according to the inversion products, although the timing of this event does not exhibit a




Figure 7. Regional contributions to total terrestrial CO₂ flux IAV for four-eight different atmospheric CO₂ inversion products and ensemble mean over 2002-2022. (a) shows continental regions while (b) shows vegetation cover type regions.

strong correlation with the GRACE TWS variations in Figure 4b, suggesting that water availability may not be a key driver. In
400 2005 tropical forests and semi-arid regions emerge as the dominant contributors to the positive CGR anomaly, which agrees
well with the GRACE data showing droughts across these regions, despite there being a positive global TWS anomaly during
this period. The inversion products also align with GRACE in attributing the 2008-2009 decrease in CGR mostly to tropical
forests and the 2011 decrease mostly to semi-arid regions. During the substantial increase in CGR coinciding with the 2015/16
El Niño the inversion products show comparable contributions from semi-arid regions, croplands and grasslands and tropical
405 forest, whereas the GRACE data most strongly highlight the role of tropical forests, illustrated in Figure 4b. Overall the




Figure 8. Contribution to terrestrial IAV from atmospheric CO₂ inversion products 2002-2022. (a) aggregated by continental regions, (b) by land cover type.

inversion products show minimal CGR IAV contributions from sparsely vegetated regions, and little from tundra and Arctic shrublands and extratropical forests, also in agreement with the GRACE GTWS-CGR correlation contributions, Figure 3b.

Contribution to terrestrial IAV from atmospheric CO₂ inversion products 2002-2022. (a) aggregated by continental regions, (b) by land cover type.

410 From Figure 8a it is notable that the NH extratropics exhibits considerably less CO₂ flux variability compared to the TWS variability, which is consistent with the insignificant contribution to the GTWS-CGR correlation. The overall contribution to the global land IAV appears more uniformly distributed across spatial regions compared to land cover types, suggesting that

factors related to land cover type may play a more significant role in determining the terrestrial carbon flux variability than spatial regions *per se*. Examining specific time periods reveals interesting regional patterns. For instance, during 2002-2003, the inversion products indicate a period of low uptake from the land, leading to high CGR, primarily driven by variations in the tropics, particularly tropical Africa, despite GRACE observations showing positive tropical TWS variations at this time, Figure 4. Atmospheric data for the inverse models in these regions were sparse until around 2010, which could be a possible cause of this discrepancy.

The drop in CGR during 2004 appears to coincide with increased land uptake in tropical Eurasia and Australia, and tropical Africa, again with limited evidence of TWS influence in these regions. The large CGR drops in 2009 and 2014 appear to be predominately driven by the NH extratropics, despite GRACE only indicating possible contributions from tropical America with minimal input from the NH. In 2022, tropical America stands out as a significant influence for the decline in CGR in the inversion products, which is also accompanied by a clear increase in TWS in the same region. From Figure 8a it is notable that the NH extratropics exhibit considerably less CO₂ flux variability compared to the TWS variability, which is consistent with the insignificant contribution to the GTWS-CGR correlation. Since the larger continental spreads seen in Figure 4a are not reproduced Figure 8a, this suggests that the inverse models do not generally capture the same anti-correlated behaviour seen in water anomalies. This could be because the CO₂ flux IAV occurs only at larger spatial scales or because the atmospheric inversion models lack the spatial resolution to detect smaller scale anti-correlated fluxes. We now perform a sensitivity analysis to try further bring out regional relations.

We calculate local sensitivities of the terrestrial CO₂ flux to GRACE TWS interannual variations using the inversion products. This provides insight into how changes in TWS translate into carbon flux variations and can help us relate Figure 4 to Figure 8. Sensitivities allow us to assess how water-limited or energy-limited conditions might modulate the influence of TWS anomalies on NBP. For instance, a small TWS anomaly in a water-limited region may have a disproportionately large impact on CGR, while large TWS changes in energy-limited regions may have a smaller effect. Figure 9 shows the sensitivities calculated using the ensemble mean of the inversions for (a) spatial regions and (b) land cover type. The sensitivity is calculated using linear regression. These sensitivities do resemble the contributions seen in Figure 3. Figure 9a shows higher sensitivities in the tropics and smaller (insignificant) opposite sensitivities in the NH extratropics and SH extratropics. Figure 9b indicates strong CO₂ sensitivity to TWS variations in semi-arid regions, as well as high sensitivity in tropical forests, and croplands and grasslands. The high sensitivity in tropical forests indicates that TWS variations in these regions directly influence NBP, rather than merely being correlated with CGR through external factors like ENSO. This reinforces our conclusion that tropical forests play a crucial role in regulating the CGR.

Overall, the total sensitivity is greater when aggregating by land cover type than by spatial regions. This suggests that aggregation by land cover better captures the variation in ecosystem responses to water availability, as different land cover types have distinct physiological and ecological responses to TWS anomalies. Aggregation by spatial regions ~~-, however, can~~ can then obscure these differences, as regions often contain a mix of land cover types with varying sensitivities to water availability. For example, areas of high sensitivity to water availability in parts of tropical America (e.g tropical forests) can be

masked when grouped into larger spatial regions that include ecosystems that are more sensitive to energy and less sensitive to water.

The strong correlation and sensitivity in tropical forests could be attributed to several possible factors. Tropical rainforest trees tend to have relatively shallow rooting systems and are hence more likely to be affected by changes in TWS when prolonged or severe droughts deplete soil moisture (Kleidon, 2004). While some trees can develop deep roots that provide drought resilience, the majority of water uptake in tropical forests occurs from shallower soil layers, making them especially sensitive to reductions in available water. Additionally, tropical forests generally have high water-use-efficiency (WUE), which allows them to maximize carbon uptake under favorable conditions but also makes them vulnerable to rapid declines in productivity when water becomes limiting (Keenan et al., 2013; Saleska et al., 2016). Another reason for this sensitivity could be due to the ongoing drying of parts of the Amazon—especially the southeastern region—which has pushed these forests into a state of heightened vulnerability to drought (Barkhordarian et al., 2019), further amplifying their sensitivity to interannual variability in water storage and the observed coupling between terrestrial CO₂ and water storage variability.




Figure 9. Regional CO₂ flux sensitivity to TWS variations (units: kgC yr⁻¹m⁻³), calculated using the ensemble mean of inversions. (a) Sensitivity by spatial regions, and (b) sensitivity by land cover type.

5 Discussion

The significant negative correlation observed between GTWS and CGR suggests a compelling link between water availability and terrestrial carbon uptake, where drier years coincide with elevated CGR levels, implying a weakening of the land carbon sink. This observation supports the notion that water availability plays a pivotal role as a limiting factor influencing land carbon uptake on interannual timescales. From Figure 4, it is also clear that even when the GTWS signal does not align with the CGR, regional TWS component signals sometimes closely follow the CGR, suggesting that water may still be a key factor influencing CO₂ variations. The evidence for water scarcity restricting land uptake is clear during times of extreme drought. For instance,

the Amazon which typically acts as an important carbon sink, experienced major droughts in the years 2005 and 2010 marked by notable peaks in the CGR. Additionally, the inversion products provide regional evidence supporting the impact of water availability during these events. However, it is important to note that other factors, including light availability and temperature, may also impact the interannual variability in land carbon uptake. This becomes apparent when regional variations in water availability do not consistently align with regional carbon variations from inversion products, suggesting weaker water-carbon coupling. ~~For instance, during~~

During the 2016 El Niño event, the high CGR was largely associated with reduced land uptake in tropical Africa, despite this region showing less dominant TWS variations. In this case, enhanced fire activity, reduced cloud cover (and thus increased radiation), and shifts in vegetation phenology have all been suggested as contributing factors to reduced carbon uptake (Liu et al., 2017; Zhu et al., 2018).

Our sensitivity analysis with the inverse model products indicated that the CO₂ flux was not highly sensitive to water variations in this region, suggesting that the variations in carbon uptake in tropical Africa were more likely driven by other factors beyond water availability during this event. Overall, our results demonstrate that water availability exerts a notable influence on interannual land carbon uptake. However, this impact exhibits spatial variability and is subject to influence from other environmental drivers which are not independent from water availability, adding complexity to the carbon-water relationship.

Given that tropical Africa shows a strong control on the CGR but low correlation with TWS, the value in understanding the relationship between CGR and TWS will primarily be in diagnosing the regional carbon cycle for areas such as the Amazon, for which numerous studies have shown evidence of it being vulnerable to water-related tipping points (Cox et al., 2000, e.g) . The Amazon generates about half of its own rainfall (Salati et al., 1979) and so drought conditions have the potential to be amplified, thus creating a stronger carbon signal than would be observed otherwise. However, even in regions where the TWS-CGR relationship is not strong, knowing this helps deepen our understanding of which processes control the carbon cycle in that region and where modelling efforts can then be focused. An application of this technique will be to observe changes to the strength of the water control of the terrestrial carbon cycle with time, to understand how the processes are changing with shifts in climate.

We explored both land cover types and spatial regions because each offers a unique perspective. Examining different spatial regions allowed us to uncover regional patterns and variations, while considering land cover type can help highlight the role of vegetation in shaping carbon-water dynamics. We found that tropical forests contribute the most to the GTWS-CGR correlation while covering only a small fraction of the Earth's land surface relative to other land cover types. Among tropical forests, the Amazon basin stands out from Figure 2, contributing a disproportionately large amount to the GTWS-CGR correlation ($g = 28\%$) relative to its spatial coverage ($< 5\%$). However, these are not the regions dominating the GTWS signal, nor are they the only regions controlling the land carbon uptake IAV. We found that semi-arid regions and croplands and grasslands contribute more to the GTWS signal, which is ~~partially~~ mainly due to their larger spatial coverage. This is consistent with Humphrey et al. (2018) who also looked at the TRENDY model ~~(Sitch et al., 2015) results~~ results (Sitch et al., 2015), and FluxCOM datasets (Tramontana et al., 2016), as well as GRACE, to assess contributions. Ahlström et al. (2015) also found that semi-arid regions dominate the land carbon sink variability by calculating the contribution index using a biogeochemical dynamic global

vegetation model (DGVM) LPJ-GUESS. They also compared results with an ensemble of TRENDY models and found similar partitioning.

The contribution to the GTWS-CGR correlation metric (equation 3) provides further insights that examining the TWS ~~contribution index variability alone~~ (equation 1) ~~alone does not capture~~ does not. For instance, ~~while the contribution index indicates the strong TWS signal~~ the strong TWS variability signals seen in the NH extratropics ~~and or~~ croplands and grasslands, ~~these regions~~ only have small contributions to the global TWS correlation with CGR. However temporal filtering used in our TWS-CGR analysis may still be underestimating the true impact of water availability on CO₂ fluxes in the extratropics, which will be strongly seasonally sensitive (Keppel-Aleks et al., 2014).

Similarly, focusing solely on the regional TWS-CGR correlations does not provide information about the magnitude of the contributions to the global correlation. Therefore, integrating these three metrics provides a comprehensive analysis, which taken together more strongly indicate the role of tropical forests. This complements the work of Humphrey et al. (2018) who ~~suggested~~ indicated a possible role for tropical forests due to the high correlation with CGR.

~~The correlation and contribution to correlation metrics help identify the regions showing coincident water storage variability with the CGR in an integrated way over the entire study period, they do not provide insights into the CGR anomalies across specific temporal events. However the~~ However all three metrics only consider time-integrated information. The regional time series displays (Figure 4 for TWS, and Figure 8 for regional CO₂ fluxes) allow a more detailed analysis of particular events. Our results underscore the influence of tropical America along with tropical forests ~~over~~ through the 20+ year period considered (Figure 3), however, it is also evident from both GRACE and the inversion products, that the dominant region influencing CGR IAV varies across different temporal events (Figure 4, 8). This is consistent with previous studies that have focused on individual events (e.g., Chen et al., 2010; Ma et al., 2016).

Our results show agreement with previous studies that have employed Earth System models (ESM) to assess CGR IAV. For instance, the dominant role of tropical America is also recognised by ~~?~~ Martín-Gómez et al. (2023), who analysed various ESM simulations from CMIP6 over 1986-2013. They find that tropical regions, particularly tropical South America, tropical southern Africa, Southeast Asia, and parts of Oceania, contribute the most to interannual variations in CGR, with average land variance explained percentages of 22%, 10%, 5%, and 4% respectively. Additionally, Kim et al. (2016) analysed how interannual CGR is associated with ENSO using ESM simulations from CMIP5. During El Niño, strong NBP anomalies appear over most tropical land regions, particularly over Amazonia, Australia, and South Asia, and the Maritime Continent, where tropical rain forests exist. This is consistent with the regions we found to show high TWS-CGR correlations in Figure 2.

The process of aggregating local data into regional or global scales introduces challenges due to the complex spatial ~~interactions at different scales~~ scale interactions. For example, Jung et al. (2017) demonstrated that local-to-global aggregation can result in different conclusions regarding the dominant driver of NEE. This ~~result~~ was attributed to compensatory effects, where ~~soil moisture-controlled NEE anomalies in different regions~~ soil moisture-controlled NEE anomalies showed strong spatial anti-correlation, leading to spatial compensation of positive and negative values. This meant that when aggregated globally, ~~that~~ temperature emerged as the dominant driver of NEE because its variability is ~~is on larger~~ on larger spatial

535 scales, so the effects do not average out. Therefore, when interpreting results such as ~~in~~ Figure 3, it is important to consider the potential role of spatial covariance.

~~The covariance between climatic drivers~~ We note also that the reported correlations between TWS and CGR have not been adjusted for co-varying climate factors such as temperature or VPD. As a result, these relationships may partly reflect indirect effects mediated by such variables. Covariance among climatic drivers—particularly between temperature, VPD, and water availability~~can further complicate the interpretation of results. In particular, Humphrey et al. (2021) notes that disentangling the individual effects of vapour pressure deficit (VPD) —introduces a degree of multicollinearity that complicates attribution of individual effects. For example, Humphrey et al. (2021) highlighted the difficulty of disentangling the respective roles of VPD and temperature on NBP variability~~ is challenging due to their interrelated effects. ~~Their sensitivity analysis revealed that direct nature and showed that~~ soil moisture impacts on NBP are influenced shaped not only by ~~local soil moisture conditions direct~~ water limitation but also by indirect ~~effects mediated through feedbacks involving~~ temperature and VPD ~~feedback mechanisms. This underscores the complexity of attributing regional contributions.~~ However, Humphrey et al. (2018) also found that most of the explanatory power contained in TWS could not be accounted for by temperature alone. Their partial correlation analysis showed that the CGR–TWS relationship remained significant even after controlling for temperature, suggesting that TWS captures a distinct and meaningful component of carbon cycle variability. Thus, while water availability cannot be interpreted in isolation from other variables, it still provides uniquely valuable information about ecosystem–climate interactions. Our findings should therefore be viewed as complementary to other studies that emphasise the roles of temperature or VPD.

~~Atmospheric inversion models are specifically designed to regionalise~~ We therefore turned to direct estimates of carbon fluxes from atmospheric inversion models which, while designed to constrain the atmospheric carbon budget, are often employed to regionalise surface carbon fluxes and investigate temporal variations, based on available atmospheric observations. Our analysis of four-eight such products demonstrate the consistency of their results when considering large scales, such as over continental regions or integrated over land cover types. However, limitations become apparent at smaller scales (Figure 6). As highlighted by Marcolla et al. (2017), fine-scale estimates of inversions are subject to limitations because atmospheric data may only effectively constrain larger-scale patterns comparable to the distances between monitoring stations. Globally, products agree because observationally the CGR is well constrained, set broadly by the difference between the slowly varying anthropogenic fossil fuel input and the well-measured accumulation rate of CO₂ in the atmosphere (Baker et al., 2006). However, the inversion models introduce factors contributing to discrepancies, such as the selection of the atmospheric CO₂ data, prior fluxes, and the choice of transport model. For example, Schuh et al. (2019) demonstrate that the transport models TM5 and GEOS-Chem, used by NOAA ~~CT2022-CT~~ and UoE respectively, lead to systematic space-time differences in ~~modeled~~ modelled distributions of CO₂. Bastos et al. (2020) also found that at regional scales, differences between inversion products contribute the most to uncertainty in regional carbon budgets, whereas differences in DGVMs dominate uncertainty at the global scale. They emphasised that reducing the uncertainty in atmospheric inversions, for example through more observations in the tropics or the use of satellite-observations, is essential to reduce uncertainty in carbon budgets.

Here using the ensemble mean ~~can~~ does offer a more representative picture by incorporating a range of estimates from different models, although different atmospheric inversion products may still be related through the same transport models or

570 meteorological fields, allowing biases to remain ~~in the ensemble means. The four products we have used do all have different transport models but the ensemble could still be improved by adding a greater number of data products that have been carefully compared against observational data.~~ It is important that each individual ensemble member is well evaluated; otherwise, outliers or individual members that do not reproduce ~~the~~ data can cause the ensemble mean to be misleading. Overall, our results emphasise the need to improve consistency among inverse models at finer scales. This would be essential for achieving more
575 reliable evidence of regional contributions to carbon cycle dynamics. In future work it would be beneficial to conduct a more comprehensive comparison of atmospheric CO₂ inversion products involving additional products beyond those considered in our study ~~→~~ e.g. containing satellite CO₂ observations.

6 Conclusion

This study first assesses the relationship between TWS variations at global and regional scales with the interannual CGR over
580 ~~2002–2023~~ 2002–2022. This builds on the work of (Humphrey et al., 2018) by extending the TWS-CGR relationship up until 2023. The correlation between global TWS and CGR was $r = -0.70$ over the entire period, with a correlation of $r = -0.69$ observed for the period from 2002 to 2016, and $r = -0.74$ from 2016 to 2023. We expand on Humphrey et al.'s analysis by separately mapping regional TWS-CGR correlations and their relative contributions to the GTWS-CGR correlation. This allows for better identification of key regions where water limitation is likely to be influencing surface CO₂ fluxes and thus
585 contributing to the GTWS-CGR correlation.

Our analysis reveals that the tropics alone can explain the entire GTWS-CGR correlation. Specifically, tropical America emerges as the largest contributing area, accounting for $g = 69\%$ of the GTWS-CGR correlation, despite only representing $f = 40\%$ of the GTWS variance and covering less than 12% of the Earth's land surface area. Moreover, we observe minor cancelling contributions to the GTWS-CGR correlation from the Northern ($g = +17\%$) and Southern ($g = -15\%$) Hemisphere
590 extratropics, which represent $f = 38\%$ and $f = -12\%$ of GTWS variability signal respectively.

Tropical forests emerge as the primary land cover type influencing the global correlation ($g = 37\%$), despite ~~not being to most being less~~ important for the global TWS signal ($f = 22\%$). Semi-arid regions also play an important role in contributing to both the GTWS-CGR correlation and the GTWS signal ($g = 26\%$, $f = 23\%$). Notably, tropical forests cover only 10% of the
595 land surface area, hence they exert a disproportionately stronger impact relative to their spatial extent compared to semi-arid regions, which encompass 25% of the land surface. Croplands and grasslands contribute largely to GTWS variability ($f = 22\%$) due to large spatial coverage but have a relatively low correlation with the CGR ($r = -0.27$) and consequently ~~do not significantly contribute~~ contribute less strongly to the GTWS-CGR correlations ($g = 15\%$).

We also employed ~~four~~ eight different atmospheric CO₂ transport inversion model products to directly assess the regional
600 contributions to CGR interannual variability. The ensemble mean of the ~~four~~ eight products shows good agreement with the CGR globally, and these products also demonstrate considerable consistency among themselves on large scales, however their agreement diminishes when examined at finer spatial scales. In alignment with GRACE TWS data, the ensemble mean of

the inversion products also attribute the majority of terrestrial carbon interannual variability to the tropics ($f = 74\%$), with smaller contributions from the NH extratropics ($f = 15\%$) as well as a small ($f = 11\%$) positive contribution from the SH extratropics, in contrast to GRACE. The inversion products all show broad agreement with GRACE in the dominant land cover types contributing to the CGR variability. ~~While~~ Although not strongly trusting the regional inverse products we also calculated regional-scale sensitivities of inverse model NBP against TWS, and find consistency with the TWS-CGR correlation contribution results.

Finally for both water storage variability and inverse model derived CO_2 fluxes we look at the interannual variation time series and note that the dominant anomaly regions do ~~shift throughout~~ change over the 20-year period, and we discuss how this ties in with literature on patterns of drought over time. These time series also reveal cases where regional TWS signals often closely follow the CGR even when the global TWS does not, thus providing further evidence of the importance of water availability as a constraint on regional CO_2 fluxes. Overall, our results suggest that terrestrial ecosystem models should prioritize improved representation of water constraints, particularly in tropical forests where water availability plays a critical role in modulating carbon fluxes.

Data availability. The MODIS land cover data is available from <https://lpdaac.usgs.gov>. The CGR data Lan et al. (2016) was retrieved from https://gml.noaa.gov/ccgg/trends/gl_data.html. The GRACE TWS data can be obtained from http://grace.jpl.nasa.gov/data/get-data/jpl_global_mascons/. All the atmospheric CO_2 inversion products are publicly available from <https://meta.icos-cp.eu/objects/FHbD8OTgCb7Tlvs99lUDAp00>

Author contributions. This work was carried out by SP as part of her PhD. All the analysis and figures were generated by her as well as a first draft of all the paper text. KH, TQ and RK acted as PhD supervisors. LF and PP provided data resources. Each contributed edits to the paper text.

Competing interests. The authors declare that they have no conflict of interest

Acknowledgements. This research has been supported by the SCENARIO NERC DTP programme (grant no. NE/S007261/1) and from a CASE award from the UK Met Office for SP. KH and TQ would also like to acknowledge the support of the NERC NCEO International Programme (grant no. NE/X006328/1). LF and PIP received support from the UK Space Agency and the UK National Centre for Earth Observation funded by the National Environment Research Council (grant no. NE/R016518/1).

References

- Ahlström, A., Schurgers, G., Arneth, A., and Smith, B.: Robustness and uncertainty in terrestrial ecosystem carbon response to CMIP5 climate change projections., *Environ. Res. Lett.*, 7, <https://doi.org/10.1088/1748-9326/7/4/044008>, 2012.
- 630 Ahlström, A. et al.: The dominant role of semi-arid ecosystems in the trend and variability of the land CO₂ sink, *Science*, 348, 895–899, <https://doi.org/DOI:10.1126/science.aaa1668>, 2015.
- Baker, D. F., Law, R. M., Gurney, K. R., Rayner, P., Peylin, P., Denning, A. S., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Fung, I. Y., Heimann, M., John, J., Maki, T., Maksyutov, S., Masarie, K., Prather, M., B. Pak, S. T., and Zhu, Z.: TransCom 3 inversion intercomparison: Impact of transport model errors on the interannual variability of regional CO₂ fluxes, 1988–2003, *Global Biogeochem. Cycles*, 20, GB1002, <https://doi.org/doi:10.1029/2004GB002439>, 2006.
- 635 Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee, X., Malhi, Y., Meyers, T., Munger, W., Oechel, W., Paw, K. T., Pilegaard, K., Schmid, H. P., Valentini, R., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: FLUXNET: A New Tool to Study the Temporal and Spatial Variability of Ecosystem–Scale Carbon Dioxide, Water Vapor, and Energy Flux Densities., *B. Am. Meteorol. Soc.*, 82, 2415–2434, [https://doi.org/https://doi.org/10.1175/1520-0477\(2001\)082<2415:FANTTS>2.3.CO;2](https://doi.org/https://doi.org/10.1175/1520-0477(2001)082<2415:FANTTS>2.3.CO;2), 2001.
- 640 Barkhordarian, A., Saatchi, S. S., Behrangi, A., Loikith, P. C., and Mechoso, C. R.: A Recent Systematic Increase in Vapor Pressure Deficit over Tropical South America, *Scientific Reports*, 9, 15 331, <https://doi.org/10.1038/s41598-019-51857-8>, 2019.
- Bastos, A., O’Sullivan, M., Ciais, P., Makowski, D., Sitch, S., and Friedlingstein, P., e. a.: Sources of uncertainty in regional and global terrestrial CO₂ exchange estimates., *Global Biogeochemical Cycles*, 34, <https://doi.org/https://doi.org/10.1029/2019GB006393>, 2020.
- 645 Chandra, N., Patra, P. K., Niwa, Y., Ito, A., Iida, Y., Goto, D., Morimoto, S., Kondo, M., T. M. H. T., and Watanabe, M.: The First Global Carbon Dioxide Flux Map Derived from TanSat Measurements, *Adv. Atmos. Sci.*, 38, 1433–1443, <https://doi.org/https://doi.org/10.1007/s00376-021-1179-7>, 2021.
- Chen, J. L., Wilson, C. R., and Tapley, B. D.: The 2009 exceptional Amazon flood and interannual terrestrial water storage change observed by GRACE., *Water Resour. Res.*, 46, W12 526, <https://doi.org/doi:10.1029/2010WR009383>, 2010.
- Chevallier, F.: Part CO₂ Contribution to documentation of products and services as provided within the scope of this contract - 2023_v1, 2023.
- 650 Chevallier, F., Fisher, M., Peylin, P., Serrar, S., Bousquet, P., Bréon, F.-M., Chédin, A., and Ciais, P.: Inferring CO₂ sources and sinks from satellite observations: Method and application to TOVS data, *J. Geophys. Res.*, 110, <https://doi.org/https://doi.org/10.1029/2005JD006390>, 2005.
- Chevallier, F., Ciais, P., Conway, T. J., Aalto, T., Anderson, B. E., Bousquet, P. ., and Worthy, D.: CO₂ surface fluxes at grid point scale estimated from a global 21 year reanalysis of atmospheric measurements., *Journal of Geophysical Research*, 115, D21 307, <https://doi.org/https://doi.org/10.1029/2010JD013887>, 2010.
- 655 Ciais, P., Bastos, A., Chevallier, F., Lauerwald, R., Poulter, B., Canadell, J. G., Hugelius, G., Jackson, R. B., Jain, A., Jones, M., Kondo, M., Luijkx, I. T., Patra, P. K., Peters, W., Pongratz, J., Petrescu, A. M. R., Piao, S., Qiu, C., Von Randow, C., Regnier, P., Saunio, M., Scholes, R., Shvidenko, A., Tian, H., Yang, H., Wang, X., and Zheng, B.: Definitions and methods to estimate regional land carbon fluxes for the second phase of the REgional Carbon Cycle Assessment and Processes Project (RECCAP-2), *Geosci. Model Dev.*, 15, 1289–1316, <https://doi.org/https://doi.org/10.5194/gmd-15-1289-2022>, 2022.
- Coelho, C., Uvo, C., and T., A.: Exploring the impacts of the tropical Pacific SST on the precipitation patterns over South America during ENSO periods., *Theor Appl Climatol.*, 71, 185–197, <https://doi.org/https://doi.org/10.1007/s00376-021-1179-7>, 2002.

- Cox, P. M., Betts, R. A., Jones, C. D., Spall, S. A., and Totterdell, I. J.: Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model, *Nature*, 408, 184–187, 2000.
- 665Cox, P. M., Pearson, D., Booth, B. B., Friedlingstein, P., Huntingford, C., Jones, C. D., and Luke, C. M.: Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability, *Nature*, 494, 341–344, <https://doi.org/https://doi.org/10.1038/nature11882>, 2013.
- Detmers, R. G., Hasekamp, O., Aben, I., Houweling, S., van Leeuwen, T. T., Butz, A., Landgraf, J., Köhler, P., Guanter, L., and Poulter, B.: Anomalous carbon uptake in Australia as seen by GOSAT, *Geophys. Res. Lett.*, 42, 8177–8184, <https://doi.org/doi:10.1002/2015GL065161>, 2015.
- 670Feng, L., Palmer, P. I., Bösch, H., and Dance, S.: Estimating surface CO₂ fluxes from space-borne CO₂ dry air mole fraction observations using an ensemble Kalman Filter, *Atmos. Chem. Phys.*, 9, 2619–2633, <https://doi.org/https://doi.org/10.5194/acp-9-2619-2009>, 2009.
- Feng, L., Palmer, P. I., Parker, R. J., Deutscher, N. M., Feist, D. G., Kivi, R., Morino, I., and Sussmann, R.: Estimates of European uptake of CO₂ inferred from GOSAT XCO₂ retrievals: sensitivity to measurement bias inside and outside Europe, *Atmos. Chem. Phys.*, 16, 1289–1302, <https://doi.org/https://doi.org/10.5194/acp16-1289-2016>, 2016.
- 675Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Arneeth, A., Arora, V., Bates, N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R. A., Ilyina, T., Jain, A. K., Joetzjer, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozzi, D.,
- 680 Marland, G., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., O’Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., van der Werf, G., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, X., and Zaehle, S.: Global carbon budget 2020, *Earth Syst. Sci. Data*, 12, 3269–3340, <https://doi.org/https://doi.org/10.5194/essd-12-3269-2020>, 2020.
- 685Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le Quéré, C., Luijkx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin, M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, , Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O’Brien, K. M., Olsen, A., Omar, A. M., Ono, T., Paulsen, M., Pierrot, D., Pocock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Séférian, R., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tans, P. P.,
- 695 Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf G. R., van Ooijen, E., Wanninkhof, R., Watanabe, M., Wimart-Rousseau, C., Yang, D., Yang, X., Yuan, W., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2023, *Earth Syst. Sci. Data*, 15, 5301–5369, <https://doi.org/https://doi.org/10.5194/essd-15-5301-2023>, 2023.
- Gaubert, B., Stephens, B. B., Basu, S., Chevallier, F., Deng, F., Kort, E. A., Patra, P. K., Peters, W., Rödenbeck, C., Saeki, T., Schimel, D., Van der Laan-Luijkx, I., Wofsy, S., and Yin, Y.: Global atmospheric CO₂ inverse models converging on neutral tropical land exchange,

- 700 but disagreeing on fossil fuel and atmospheric growth rate, *Biogeosciences*, 116, 117–134, [https://doi.org/https://doi.org/10.5194/bg-16-117-](https://doi.org/https://doi.org/10.5194/bg-16-117-2019)
2019, 2019.
- Guerlet, S., Basu, S., Butz, A., Krol, M., Hahne, P., Houweling, S., et al.: Reduced carbon uptake during the 2010 Northern Hemisphere summer from GOSAT, *Geophysical Research Letters*, 40, 2378–2383, <https://doi.org/10.1002/grl.50402>, 2013.
- Gurney, K. R., Castillo, K., Li, B., and Zhang, X.: A positive carbon feedback to ENSO and volcanic aerosols in the tropical terrestrial biosphere, *Glob. Biogeochem. Cycles*, 26, GB1029, <https://doi.org/https://doi.org/10.1029/2011GB004129>, 2012.
- 705 He, B., Chen, C., Lin, S., Wenping Yuan, H. W. C., Chen, D., Zhang, Y., Guo, L., Zhao, X., Liu, X., Piao, S., Zhong, Z., Wang, R., and Tang, R.: Worldwide impacts of atmospheric vapor pressure deficit on the interannual variability of terrestrial carbon sinks, *National Science Review*, 9, <https://doi.org/https://doi.org/10.1093/nsr/nwab150>, 2022.
- Huang, M., Piao, S., Ciais, P., and et al: Air temperature optima of vegetation productivity across global biomes., *Nat. Ecol. Evol.*, 3, 772–7796, <https://doi.org/https://doi.org/10.1038/s41559-019-0838-x>, 2019.
- 710 Humphrey, V., Zscheischler, J., Ciais, P., Gudmundsson, L., Sitch, S., and Seneviratne, S. I.: Sensitivity of atmospheric CO₂ growth rate to observed changes in terrestrial water storage, *Nature*, 560, 628–631, <https://doi.org/https://doi.org/10.1038/s41586-018-0424-4>, 2018.
- Humphrey, V., Berg, A., Ciais, P., Gentile, P., Jung, M., Reichstein, M., ..., and Frankenberg, C.: Soil moisture-atmosphere feedback dominates land carbon uptake variability., *Nature*, 592, 65–69., <https://doi.org/https://doi.org/10.1038/s41586-021-03325-5>, 2021.
- 715 Ito, A.: Disequilibrium of terrestrial ecosystem CO₂ budget caused by disturbance-induced emissions and non-CO₂ carbon export flows: a global model assessment, *Earth Syst. Dynam.*, 10, 685–709, <https://doi.org/https://doi.org/10.5194/esd-10-685-2019>, 2019.
- Jacobson, A. R., Schuldt, K. N., Tans, P., Andrews, A., Miller, J. B., Oda, T., Mund, J., Weir, B., Ott, L., Aalto, T., Abshire, J. B., Aikin, K., Aoki, S. and Apadula, F., Arnold, S., Baier, B., Bartyzel, J. NS Beyersdorf, A. B. T., and Zimnoch., M.: CarbonTracker CT2022. NOAA Global Monitoring Laboratory., <https://doi.org/https://doi.org/10.25925/Z1GJ-3254>, 2023.
- 720 Jones, C. and Cox, P.: On the significance of atmospheric CO₂ growth rate anomalies in 2002–2003, *Geophys. Res. Lett.*, 32, L14 816, <https://doi.org/doi:10.1029/2005GL023027>, 2005.
- Jung, M., Reichstein, M., Schwalm, C., et al.: Compensatory water effects link yearly global land CO₂ sink changes to temperature., *Nature*, 541, 516–520, <https://doi.org/https://doi.org/10.1038/nature20780>, 2017.
- Jung, M., Schwalm, C., Migliavacca, M., Walther, S., Camps-Valls, G., Koirala, S., Carvalhais, N., et al.: Scaling carbon
725 fluxes from eddy covariance sites to globe: Synthesis and evaluation of the FLUXCOM approach., *Biogeosciences Discussions*, <https://doi.org/https://doi.org/10.5194/bg-2019-368>, 2019.
- Keeling, C. D. and Revelle, R.: Effects of EL Nino/southern oscillation on the atmospheric content of carbon dioxide, *Meteoritics*, 20, 437–450, 1985.
- Keenan, T. F., Hollinger, D. Y., Bohrer, G., Dragoni, D., Munger, J. W., Schmid, H. P., and Richardson, A. D.: Increase in forest water-
730 use efficiency as atmospheric carbon dioxide concentrations rise, *Nature*, 499, 324–327, <https://doi.org/https://doi.org/10.1038/nature12291>, 2013.
- Keppel-Aleks, G., Wolf, A. S., Mu, M., Doney, S. C., Morton, D. C., Kasibhatla, P. S., et al.: Separating the influence of temperature, drought, and fire on interannual variability in atmospheric CO₂, *Global Biogeochemical Cycles*, 28, 1295–1310, <https://doi.org/10.1002/2014GB004890>, 2014.
- 735 Kim, J., Kug, J., Yoon, J., and Jeong, S.: Increased Atmospheric CO₂ Growth Rate during El Niño Driven by Reduced Terrestrial Productivity in the CMIP5 ESMs, *J. Climate*, 29, 8783–8805, <https://doi.org/https://doi.org/10.1175/JCLI-D-14-00672.1>, 2016.

- Kleidon, A.: Global Datasets of Rooting Zone Depth Inferred from Inverse Methods, *J. Climate*, 17, 2714–2722, [https://doi.org/https://doi.org/10.1175/1520-0442\(2004\)017<2714:GDORZD>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0442(2004)017<2714:GDORZD>2.0.CO;2), 2004.
- Lan, X., Tans, P., and Thoning, K.: Trends in globally-averaged CO₂ determined from NOAA Global Monitoring Laboratory measurements, Version 2023-11, <https://doi.org/https://doi.org/10.15138/9N0H-ZH07>, 2016.
- Lan, X., Tans, P., and Thoning, K.: Trends in globally-averaged CO₂ determined from NOAA Global Monitoring Laboratory measurements. Version 2024-03, <https://doi.org/10.15138/9N0H-ZH07>, 2023.
- le Maire, G., Delpierre, N., Jung, M., Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ibrom, A., Kolari, P., Longdoz, B., Moors, E. J., Pilegaard, K., Rambal, S., Richardson, A. D., and Vesala, T.: Detecting the critical periods that underpin interannual fluctuations in the carbon balance of European forests, *J. Geophys. Res.*, 115, G00H03, <https://doi.org/https://doi.org/10.1029/2009JG0012447>, 2010.
- Liu, J., Bowman, K. W., Schimel, D. S., Parazoo, N. C., Jiang, Z., Lee, M., Bloom, A. A., Wunch, D., Frankenberg, C., Sun, Y., O'Dell, C. W., Gurney, K. R., Menemenlis, D., Gierach, M., Crisp, D., and Eldering, A.: Contrasting carbon cycle responses of the tropical continents to the 2015–2016 El Niño, *Science*, 358, eaam5690, <https://doi.org/10.1126/science.aam5690>, 2017.
- Liu, L., Ciais, P., Wu, M., Padrón, R. S., Friedlingstein, P., Schwaab, J., Gudmundsson, L., and Seneviratne, S. I.: Increasingly negative tropical water–interannual CO₂ growth rate coupling, *Nature*, 618, 755–760, <https://doi.org/https://doi.org/10.1038/s41586-023-06056-x>, 2023.
- Liu, Z., Ciais, P., and Deng, Z.: Near-real-time monitoring of global CO₂ emissions reveals the effects of the COVID-19 pandemic., *Nat Commun.*, 11, 5172, <https://doi.org/https://doi.org/10.1038/s41467-020-18922-7>, 2020.
- Luo, X. and Keenan, T.: Tropical extreme droughts drive long-term increase in atmospheric CO₂ growth rate variability, *Nat Commun*, 13, 1193, <https://doi.org/https://doi.org/10.1038/s41467-022-28824-5>, 2022.
- Ma, X., Huete, A., Cleverly, J., et al.: Drought rapidly diminishes the large net CO₂ uptake in 2011 over semi-arid Australia., *Sci Rep*, 6, 37 747, <https://doi.org/https://doi.org/10.1038/srep37747>, 2016.
- Marcolla, B., Rödenbeck, C., and Cescatti, A.: Patterns and controls of inter-annual variability in the terrestrial carbon budget, *Biogeosciences*, 14, 3815–3829, <https://doi.org/https://doi.org/10.5194/bg-14-3815-2017>, 2017.
- Marengo, J. A., Tomasella, J., Alves, L. M., Soares, W. R., and Rodriguez, D. A.: The drought of 2010 in the context of historical droughts in the Amazon region, *Geophys. Res. Lett.*, 38, L12 703, <https://doi.org/doi:10.1029/2011GL047436>., 2011.
- Martín-Gómez, V., Ruprich-Robert, Y., Tourigny, E., et al.: Large spread in interannual variance of atmospheric CO₂ concentration across CMIP6 Earth System Models, *Atmos Sci*, 6, 206, <https://doi.org/https://doi.org/10.1038/s41612-023-00532-x>, 2023.
- Munassar, S., Monteil, G., Scholze, M., Karstens, U., Rödenbeck, C., Koch, F.-T., Totsche, K. U., and Gerbig, C.: Why do inverse models disagree? A case study with two European CO₂ inversions, *Atmos. Chem. Phys.*, 23, 2813–2828, <https://doi.org/https://doi.org/10.5194/acp-23-2813-2023>, 2022.
- Niwa, Y., Tomita, H., Satoh, M., Imasu, R., Sawa, Y., Tsuboi, K., Matsueda, H., Machida, T., Sasakawa, M., Belan, B., and Saigusa, N.: A 4D-Var inversion system based on the icosahedral grid model (NICAM-TM 4D-Var v1.0) – Part 1: Offline forward and adjoint transport models, *Geosci. Model Dev.*, 10, 1157–1174, <https://doi.org/https://doi.org/10.5194/gmd-10-1157-2017>, 2017.
- Niwa, Y., Ishijima, K., Ito, A., and Iida, Y.: Toward a long-term atmospheric CO₂ inversion for elucidating natural carbon fluxes: technical notes of NISMOM-CO₂ v2021.1, *Prog. Earth Planet Sci.*, 9, <https://doi.org/https://doi.org/10.1186/s40645-022-00502-6>, 2022.
- Oda, T. and Maksyutov, S.: A very high-resolution (1 km×1 km) global fossil fuel CO₂ emission inventory derived using a point source database and satellite observations of nighttime lights, *Atmos. Chem. Phys.*, 11, 543–556, <https://doi.org/https://doi.org/10.5194/acp-11-543-2011>, 2011.

Patra, P. K., Maksyutov, S., Ishizawa, M., Nakazawa, T., Takahashi, T., and Ukita, J.: Interannual and decadal changes in the sea-air CO₂ flux from atmospheric CO₂ inverse modeling, *Global Biogeochem. Cycles*, 19, GB4013, <https://doi.org/doi:10.1029/2004GB002257>, 2005.

Petch, S., Dong, B., Quaife, T., King, R. P., and Haines, K.: Water and energy budgets over hydrological basins on short and long timescales, *Hydrol. Earth Syst. Sci.*, 27, 1723–1744, <https://doi.org/https://doi.org/10.5194/hess-27-1723-2023>, 2023a.

Petch, S., Dong, B., Quaife, T., King, R. P., and Haines, K.: Precipitation explains GRACE water storage variability over large endorheic basins in the 21st century, *Front. Environ. Sci.*, 11, <https://doi.org/https://doi.org/10.3389/fenvs.2023.1228998>, 2023b.

Peylin, P., Law, R. M., Gurney, K. R., Chevallier, F., Jacobson, A. R., Maki, T., Niwa, Y., Patra, P. K., Peters, W., Rayner, P. J., Rödenbeck, C., van der Laan-Luijkx, I. T., and Zhang, X.: Global atmospheric carbon budget: Results from an ensemble of atmospheric CO₂ inversions, *Biogeosciences*, 10, 6699–6720, <https://doi.org/https://doi.org/10.5194/bg-10-6699-20137>, 2013.

Phillips, L., Aragao, L., Lewis, S., Fishee, J., Torres-Lezama, A., et al.: Drought Sensitivity of the Amazon Rainforest., *Science*, 323, 1344–1347, <https://doi.org/DOI:10.1126/science.1164033>, 2009.

Piao, S., Wang, X., Wang, K., Li, X., Bastos, A., Canadell, J. G., Ciais, P., Friedlingstein, P., and Sitch, S.: Interannual variation of terrestrial carbon cycle: Issues and perspectives, *Glob Change Biol.*, 26, 300–318, <https://doi.org/https://doi.org/10.1111/gcb.14884>, 2020.

Potter, C. S., Randerson, J. T., Field, C. B., Matson, P. A., Vitousek, P. M., Mooney, H. A., and Klooster, S. A.: Terrestrial ecosystem production - a process model-based on global satellite and surface data., *Global Biogeochemical Cycles*, 7, 811–841, 1993.

Poulter, B., Frank, D., Ciais, P., et al.: Contribution of semi-arid ecosystems to interannual variability of the global carbon cycle., *Nature*, 509, 600–603, <https://doi.org/https://doi.org/10.1038/nature13376>, 2014.

Rödenbeck, C., Houweling, S., Gloor, M., and Heimann, M.: CO₂ flux history 1982–2001 inferred from atmospheric data using a global inversion of atmospheric transport., *Atmos. Chem. Phys.*, 3, 1919–1964, <https://doi.org/https://doi.org/10.5194/acp-3-1919-2003>, 2003.

Rödenbeck, C., Zaehle, S., Keeling, R., and Heimann, M.: History of El Niño impacts on the global carbon cycle 1957–2017: a quantification from atmospheric CO₂ data, *Philos. T. Roy. Soc. B*, 373, 20170 303, <https://doi.org/https://doi.org/10.1098/rstb.2017.0303>, 2018.

Salati, E., Dall'Olio, A., Matsui, E., and Gat, J. R.: Recycling of water in the Amazon basin: an isotopic study, *Water resources research*, 15, 1250–1258, 1979.

Saleska, S. R., Wu, J., Guan, K., Araujo, A. C., Huete, A. R., Nobre, A. D., and Restrepo-Coupe, N.: Dry-season greening of Amazon forests, *Nature*, 531, <https://doi.org/https://doi.org/10.1038/nature16457>, 2016.

Santoso, A., Mcphaden, M. J., and Cai, W.: The defining characteristics of ENSO extremes and the strong 2015/2016 El Niño., *Reviews of Geophysics*, 55, 1079–1129, <https://doi.org/https://doi.org/10.1002/2017RG000560>, 2017.

Schuh, A. E., Jacobson, A. R., Basu, S., Weir, B., Baker, D., Bowman, K., et al.: Quantifying the impact of atmospheric transport uncertainty on CO₂ surface flux estimates., *Global Biogeochemical Cycles*, 33, 484–500, <https://doi.org/https://doi.org/10.1029/2018GB006086>, 2019.

Sitch, S., Prentice, I., Smith, B., Cramer, W., Kaplan, J., Lucht, W., Sykes, M., Thonicke, K., and Venevsky, S.: LPJ – a coupled model of vegetation dynamics and the terrestrial carbon cycle, in "The role of vegetation dynamics in the control of atmospheric CO₂ content", Ph.D. thesis, Lund University, Lund, Sweden, 2000.

Sitch, S. et al.: Recent trends and drivers of regional sources and sinks of carbon dioxide., *Biogeosciences*, 12, 653–679, <https://doi.org/doi:10.5194/bg-12-653-2015>, 2015.

Tramontana, G., Jung, M., Schwalm, C. R., Ichii, K., Camps-Valls, G., Ráduly, B., Reichstein, M., Arain, M. A., Cescatti, A., Kiely, G., Merbold, L., Serrano-Ortiz, P., Sickert, S., Wolf, S., and Papale, D.: Predicting carbon dioxide and energy fluxes across global FLUXNET sites with regression algorithms., *Biogeosciences*, 13, 4291–4313, <https://doi.org/https://doi.org/10.5194/bg-13-4291-2016>, 2016.

- van der Laan-Luijkx, I. T., van der Velde, I. R., van der Veen, E., Tsuruta, A., Stanislawski, K., Babenhauserheide, A., Zhang, H. F., Liu, Y., He, W., Chen, H., Masarie, K. A., Krol, M. C., and Peters, W.: The CarbonTracker Data Assimilation Shell (CTDAS) v1.0: implementation and global carbon balance 2001–2015, *Geosci. Model Dev*, 10, 2785–2800, <https://doi.org/https://doi.org/10.5194/gmd-10-2785-2017>, 2017.
- van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C.,
815 Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016., *Earth Syst. Sci. Data*, 9, 697–720, <https://doi.org/https://doi.org/10.5194/essd-9-697-2017>, 2017.
- Wang, J., Zeng, N., and Wang, M.: Interannual variability of the atmospheric CO₂ growth rate: roles of precipitation and temperature, *Biogeosciences*, 13, 2339–2356, <https://doi.org/doi:10.5194/bg-13-2339-2016>, 2016.
- Wang, K., Bastos, A., Ciais, P., Wang, X., Rödenbeck, C., Gentile, P., Chevallier, F., Humphrey, V., Huntingford, C., O’Sullivan, M., Seneviratne, S. I., Sitch, S., and Piao, S.: Regional and seasonal partitioning of water and temperature controls on global land carbon uptake variability, *Nature Communications*, 13, <https://doi.org/https://doi.org/10.1038/s41467-022-31175-w>, 2022.
- Wang, W., Ciais, P., Nemani, R. R., Canadell, J. G., Piao, S., Sitch, S., White, M. A., Hashimoto, H., Milesi, C., and Myneni, R. B.: Variations in atmospheric CO₂ growth rates coupled with tropical temperature., *Proc. Natl Acad. Sci.*, 110, 13 061–13 066, <https://doi.org/https://doi.org/10.1073/pnas.1219683110>, 2013.
- 825 Wang, X., Piao, S., Ciais, P., Friedlingstein, P., Myneni, R. B., Cox, P., Heimann, M., Miller, J., Peng, S., Wang, T., Yang, H., and Chen, A.: A two-fold increase of carbon cycle sensitivity to tropical temperature variations, *Nature*, 506, 212–215, <https://doi.org/https://doi.org/10.1038/nature12915>, 2014.
- Wiese, D., Landerer, F. W., and Watkins, M. M.: Quantifying and reducing leakage errors in the JPL RL05M GRACE mascon solution, *Water Resources Research*, 52, 7490–7502, <https://doi.org/https://doi.org/10.1002/2016WR019344>, 2016.
- 830 Wigneron, J., Fan, L., Ciais, P., Bastos, A., Brandt, M., Chave, J., Saatchi, S., Baccini, A., and Fensholt, R.: Tropical forests did not recover from the strong 2015–2016 El Niño event, *Sci. Adv.*, 6, eaay4603, <https://doi.org/10.1126/sciadv.aay4603>, 2020.
- Yang, D., Liu, Y., Feng, L., Wang, J., Yao, L., Cai, Z., Zhu, S., Lu, N., and Lyu, D.: The First Global Carbon Dioxide Flux Map Derived from TanSat Measurements, *Adv. Atmos. Sci.*, 38, 1433–1443, <https://doi.org/https://doi.org/10.1007/s00376-021-1179-7>, 2021.
- Zhang, L., Ren, X., Wang, J., He, H., Wang, S., Wang, M., Piao, S., Yan, H., Ju, W., Gu, F., Zhou, L., Niu, Z., Ge, R., Li, Y., Lv, Y., Yan, H., Huang, M., and Yu, G.: Interannual variability of terrestrial net ecosystem productivity over China: regional contributions and climate attribution, *J Environ. Res. Lett.*, 14, 014 003, <https://doi.org/10.1088/1748-9326/aaec95>, 2019.
- 835 Zhang, Y., Xiao, X., Guanter, L., Zhou, S., Ciais, P., Joiner, J., Sitch, S., Wu, X., Nabel, J., Dong, J., Kato, E., Jain, A., Wiltshire, A., and Stocker, B.: Precipitation and carbon-water coupling jointly control the interannual variability of global land gross primary production., *Sci Rep*, 26, 39 748, <https://doi.org/doi:10.1038/srep39748>, 2016.
- 840 Zhu, J., Zhang, M., Zhang, Y., Zeng, X., and Xiao, X.: Response of tropical terrestrial gross primary production to the super El Niño event in 2015., *Journal of Geophysical Research: Biogeosciences*, 123, 23 193–3203, <https://doi.org/https://doi.org/10.1029/2018JG004571>, 2018.