

ROCKE-3D 2.0: An updated general circulation model for simulating the climates of rocky planets

Kostas Tsigaridis^{1,2,3}, Andrew S. Ackerman², Igor Aleinov^{1,2,3}, Mark A. Chandler^{1,2,3}, Thomas L. Clune⁴, Christopher M. Colose^{5,2,3}, Anthony D. Del Genio^{2,6}, Maxwell Kelley², Nancy Y. Kiang^{1,2,3}, Anthony Leboissetier^{5,2,3}, Jan P. Perlwitz^{7,2,3}, Reto A. Ruedy^{5,2}, Gary L. Russell², Linda E. Sohl^{1,2,3}, Michael J. Way^{2,3,8}, Eric T. Wolf^{9,10,3}

¹Center for Climate Systems Research, Columbia University, New York, NY-10025, USA

²NASA Goddard Institute for Space Studies, New York, NY 10025, USA

³Sellers Exoplanet Environments Collaboration, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

10 ⁴NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

⁵Autonomic Integra, Gaithersburg, MD, 20879

⁶Department of Applied Physics and Applied Mathematics, Columbia University, New York, NY 10027, USA

⁷Climate, Aerosol and Pollution Research, LLC, Bronx, NY 10471, USA

⁸Theoretical Astrophysics, Department of Physics and Astronomy, Uppsala University, Uppsala, SE-75120, Sweden

15 ⁹NASA NExSS Virtual Planetary Laboratory, Seattle, WA 98195, USA

¹⁰University of Colorado, Boulder, Laboratory for Atmospheric and Space Physics, USA

Correspondence to: Kostas Tsigaridis (kostas.tsigaridis@columbia.edu)

Abstract. We present the second generation of ROCKE-3D (Resolving Orbital and Climate Keys of Earth and Extraterrestrial Environments with Dynamics), a generalized 3-dimensional General Circulation Model (GCM) for use in Solar System and exoplanetary simulations of rocky planet climates. ROCKE-3D version 2.0 is a descendant of ModelE2.1, the flagship Earth System Model of the NASA Goddard Institute for Space Studies (GISS) used in the most recent Intergovernmental Panel ~~for~~ on Climate Change (IPCC) assessments. ROCKE-3D is a continuous effort to expand the capabilities of GISS ModelE to handle a broader range of planetary conditions, including different atmospheric planet sizes, gravities, pressures, rotation rates, more diverse chemistry schemes and atmospheric compositions, diverse ocean and land distributions and topographies, and potential basic biosphere functions. In this release we present updated physics, and many more supported configurations which can serve as starting points to simulate the atmospheres of rocky terrestrial planets of interest. Two different radiation schemes are supported, the GISS radiation, valid only for atmospheres similar to that of modern Earth, and SOCRATES-, which is more generalized but more computationally expensive. While ROCKE-3D can simulate a very wide range of planetary and atmospheric configurations, we describe here a small subset of them, with the goal of demonstrating the structural capabilities, rather than the scientific breadth, of the model. Three different atmospheric composition options are described (preindustrial Earth, the aerosol and ozone-free atmosphere used in ROCKE-3D 1.0, and an anoxic atmosphere with no aerosols), three ocean configurations (prescribed, Q-flux, and dynamic), and two resolutions: the medium resolution (4x5 degrees in latitude and longitude, previously used in ROCKE-3D 1.0), and the fine resolution, which has double the resolution in the atmosphere and 4 times the horizontal and 3 times the vertical resolution in the ocean. Finally, for the land surface hydrology, we have

20
25
30

35 introduced generalized physics for arbitrary topography in the pooling and evaporation of water and river transport of water between grid cells, and for the vertical stratification of temperature in dynamic lakes. We quantify how the different component choices affect model results, and discuss strengths and limitations of using each component, together with how one can select which component to use. ROCKE-3D is publicly available and tutorial sessions are available for the community, greatly facilitating its use by any interested group.

40 1. Introduction

Three-dimensional General Circulation Models (GCMs) have been in existence for several decades, dating as far back as the 1950s (Phillips, 1956). Their use has mostly been in the climate modeling of Modern Earth, as well as in more generic studies of atmospheric (e.g. Showman et al., 2013) and oceanic circulation (e.g. Bryan, 1969). Initially these models were focused on the atmospheric component and have been called Atmospheric GCMs, or AGCMs (Manabe and Wetherald, 1967). The next major component to be added were the oceans which provided a source of water to the atmosphere, and a variety of computationally efficient ocean models were coupled to the AGCMs and became known as AOGCMs (Manabe and Stouffer, 1993). These computationally efficient oceans included those with fixed sea surface temperatures (SSTs) and later what were termed Q-flux oceans where the heat (Q) fluxes (flux) between oceanic grid cells were fixed at model start (e.g. Russell et al., 1985; Slingo, 1982). Initially the fluxes used came from satellite observations or from numerical ocean models (e.g. Bryan, 1982; Miller et al., 1983). The depths of these oceans had no standard fixed values, but researchers have typically used depths up to ~100m. These computationally efficient types of oceans have a variety of names such as thermodynamic, slab, etc (see Way et al. 2017 for more details). In the past couple of decades the Earth GCM community has moved to fully dynamic oceans that explicitly calculate horizontal and vertical ocean heat transports at resolved spatial scales while also including a range of parameterized processes that operate on smaller unresolved scales~~that include a range of parameterized processes~~, but they are more computationally expensive than the simpler SST or Q-flux oceans. In modern times the exoplanet community has started to use AOGCMs~~a parameterized ocean~~, but they typically use Q-flux type oceans where the fluxes are set to zero (highlighted as Q-flux=0 here when relevant). This is because they have no way to set the ocean heat fluxes ~~because they obviously do not have~~due to the lack of any observations, and have typically been unwilling to use fully dynamic oceanic models (AOGCMs) to estimate the fluxes, due to the increased computational cost and additional free parameters, given the lack of information about oceans in exoplanets. This approach obviously has limitations in accurately modeling the climate with perhaps the most “notorious” being the eyeball world of tidally locked planets around M-dwarf type stars (e.g. Pierrehumbert, 2010). For example, using the ROCKE-3D model Del Genio et al. (2019a) showed in simulations of Proxima Centauri b that large scale circulation patterns (see their Figures 1a,b and 2a,b) and mean surface temperatures (Q-flux=0 had a mean surface T=-37 °C, while the dynamical ocean had T=-21 °C) can differ drastically between a model using a Q-flux=0 and a fully dynamic ocean model. Yet in other scenarios the differences can be minor as seen in the work of Way et al. (2018) in a parameter study of slowly rotating worlds with increasing insolation. They showed that Earth-like planets around solar type stars showed large

mean surface temperature differences when the length of day was similar to that of modern Earth, but at longer length of days and higher insulations the differences were marginal ($<2^{\circ}\text{C}$; See their Fig. 2). ~~As demonstrated in this brief introduction, the~~ ROCKE-3D ~~model~~ is capable of modeling the atmosphere coupled to any of the ocean models described above. ROCKE-3D also has fully coupled cryosphere and land surface models, interactive ocean biogeochemistry and interactive atmospheric chemistry. Not only are these processes vital for a model Earth GCM, they are also important for studying the climates of exoplanets found at the inner and outer edges of the habitable zone (e.g. Kopparapu et al., 2013). Herein we discuss the second release of the ROCKE-3D GCM. Sect. 2 provides a detailed model description; Sect. 3 describes how ROCKE-3D evolved from its parent Earth GCM, ModelE; Sect. 4 presents model configurations than can be utilized by the end user; and Sect. ~~05~~ is the discussion.

2. Model description

ROCKE-3D version 1.0 (Way et al., 2017), ~~named planet_1.0 here,~~ was a descendent of GISS ModelE2, described and thoroughly evaluated by Schmidt et al. (2014). Since then, several bug fixes, updates and upgrades to both ROCKE-3D and ModelE have happened in parallel and are brought together in ROCKE-3D version 2.0, ~~named planet_2.0 here,~~ which is described in this work. There are two development pathways that contributed to ~~planet-ROCKE-3D~~ 2.0: The Earth-centric development in ModelE2.1 is summarized in Sect. 2.1, while the planetary-centric development in ROCKE-3D 2.0 is described in Sect. 2.2. More specifically, several physics changes were introduced in GISS ModelE2.1, which is the version of ModelE used in the Coupled Model Intercomparison Project phase 6 (CMIP6; Eyring et al., 2016), and are thoroughly described elsewhere (Kelley et al., 2020). Additional changes present in ModelE2.1 that are either bug fixes or structural enhancements since the CMIP6 version of the model were also ported to ~~planet-ROCKE-3D~~ 2.0 and are briefly described in ~~the same~~ ~~section~~ Sect. 2.1. These latter enhancements include major changes in the prognostic tracers code present in ROCKE-3D 2.0, which will be described elsewhere, since they are not used here. The planetary-related changes, which include both physics enhancements and diagnostics enrichment relevant to the generalized ROCKE-3D 2.0 model, are described in Sect. 2.2. Here we study the climatology simulated by the model using a fixed atmospheric composition per configuration, in the exact same way that was the case for ROCKE-3D 1.0 (Way et al., 2017).

The hydrostatic ModelE2.1 dynamical core employed in ROCKE-3D utilizes the Arakawa "B"-grid staggering, whose benefits at coarse resolution over the conventional "C"-grid were argued by Hansen et al. (1983); this reference provides other introductory details as well. Momentum and air-mass transport continues to employ the Arakawa centered-differencing approach, with dissipation at small scales applied via a Shapiro filter. A prominent update from the Hansen et al. (1983) version is that potential temperature, humidity, and other scalars are transported using the conservative quadratic upstream scheme (QUS; Prather, 1986) whose implicit numerical diffusion is inherently very scale-selective. The benefits of the QUS for tracers were described in Rind and Lerner (1996). Air mass and momentum fields are advanced using a leapfrog scheme with a timestep respecting the Courant–Friedrichs–Lewy (CFL) condition for the barotropic mode (Lamb wave) whose horizontal

velocity is on the order of the speed of sound. Potential temperature advection is performed on the even leapfrog timesteps, with temporal interpolation as needed for hydrostatic integrations. Since humidity and tracer fields do not directly affect the dynamics in this version, they are advected on a longer, but locally adaptive, timestep, respecting local CFL conditions for wind speeds, which are typically much less than the speed of sound.

The ROCKE-3D atmosphere includes hydrostatic resolved dynamics and parameterized sub-grid vertical transport via turbulent eddies and vertically extensive cumulus-cloud convection; stratiform cloud fraction is diagnosed from relative humidity and convective cloud fraction from vertical mass exchange rates; condensation of water vapor is explicitly calculated in clouds as a source of precipitation and the liquid and ice hydrometeors seen by radiative transfer; optional chemical and aerosol tracer constituents propagate through all these phenomena as they undergo their own processes. The evolutions of these processes are outlined in a series of papers (Bauer et al., 2020; Kelley et al., 2020; Schmidt et al., 2006, 2014) which themselves may refer back to component-specific "original version" descriptions, e.g. Del Genio et al. (1996) for stratiform clouds or Lacis and Oinas (1991) for longwave radiative transfer.

Sea ice in ROCKE-3D 2.0 is treated as two mass layers, each containing two thermal layers: 1) the top mass layer has a fixed thickness of 0.1 m and contains both ice and snow (snow thickness evolves with precipitation), and 2) the bottom mass layer is ice only and can grow infinitely in thickness (with a minimum of 0.1 m). Sea ice thermodynamics are based on the Brine Pockets or "BP" formulation, with both mass and energy budget (Schmidt et al., 2004) and include brine rejection through gravity drainage and flushing of meltwater. Ice sheets are bottomless, but only the upper 3 m are used for thermodynamics computations. Those are represented by two layers of ice: 0.1 m at the top and 2.9 m at the bottom. The ice can accumulate a layer of snow on top of it. The snow in excess of 0.1 m of dense ice equivalent is converted to the ice and pushed down the ice layers. If iceberg calving is enabled (optional) the accumulated ice is distributed as icebergs to the ocean cells according to a prescribed mask (specified in an input file). The albedo of the ice sheet snow is either fixed at 0.8 (default for modern Earth) or is updated according to a snow aging algorithm as described in Hansen et al. (1983).

The vegetation model coupled to ROCKE-3D is the Ent Terrestrial Biosphere Model (Ent TBM). This dynamic global vegetation model (DGVM) provides simulation of biophysics (albedo, photosynthesis, transpiration, plant and soil respiration) given prescribed vegetation boundary conditions of sub-grid cover fractions of up to 13 plant functional types (PFTs), canopy heights, plant densities, and monthly leaf area index (LAI). PFTs are distinguished by growth form (grass, shrub, tree), photosynthetic pathway (C3, C4), phenology (evergreen/deciduous, perennial/annual), and some climatological distinctions (cold vs. arid shrubs, cold vs. warm grasses). The PFTs are based on Earth's vegetation, and can be used to simulate paleovegetation at time periods that had similar PFTs, but these are not generalized in behavior to be suitable for arbitrary exoplanets that have very different climatology, seasonality, orbital period, and atmospheric composition. The model's physics are described in Kim et al. (2015), and global scale performance in coupled carbon cycle simulations is described in Ito et al. (2020).

The public release of the ~~planet~~ ROCKE-3D 2.0 code comes with multiple possible configurations of radiative transfer calculations, atmospheric composition, ocean parameterizations, and model resolution that cross the parameter space with all

possible combinations of them. Their physics, outlined in Sect. 2.3 to 2.6, existed as options in [planet-ROCKE-3D 1.0](#) (Way et al., 2017), but were not provided as standard configurations, and were not examined in parallel as done here. In addition to those template configurations detailed in Sect. 4, we also describe here new and improved physics options that are available in [planet-ROCKE-3D 2.0](#) in Sect. 2.2. These include new land/river/lakes development, the option of including geothermal heat flux, the updates that support thin atmospheres, and calendar and equation of time updates. Finally, more technical changes that are of interest to model users, are discussed in Appendix A. The model code, all output, and the model configurations used here, are available on a zenodo archive (Tsigaridis et al., 2025).

2.1. GISS ModelE2.1 development since ModelE2

As mentioned above, ROCKE-3D Planet 1.0 is based upon GISS ModelE2. The parent Earth GCM upon which ROCKE-3D is based on is continuously being developed and serves as the basis for GISS participation in CMIP (e.g. Bauer et al., 2020; Kelley et al., 2020; Miller et al., 2021; Nazarenko et al., 2022). Successive generations of CMIP GCMs in turn form the basis for assessments of past and future terrestrial climate change as well as current climate variability. The model changes going from the previous GISS ModelE2 to the current ModelE2.1 are described in detail in Kelley et al. (2020). Here we present a summary of the most important ModelE improvements in the E2.1 version and expand on them as needed throughout the manuscript.

Other than trivial structural changes, the GISS radiation scheme was not updated since [planet-ROCKE-3D 1.0](#) in any significant way. Moist convection was improved by increasing entrainment of environmental air into convective updrafts and by increasing the evaporation of falling precipitation into the environment, both of which have the effect of making humidity more sensitive to convection and vice-versa. In ModelE2, glaciation of a supercooled liquid water cloud was determined probabilistically as a function of temperature since the model does allow liquid and ice to co-exist in a gridbox, and once glaciated, remains ice for the lifetime of the cloud. In E2.1, this is replaced by a temperature-dependent autoconversion rate of supercooled water to precipitating ice that increases with decreasing temperature, the effect of which is to increase supercooled liquid cloud occurrence at high latitudes. Another change makes the threshold relative humidity for subgrid cloud formation sensitive to aware of the planetary boundary layer height as simulated by the model, rather than assuming a fixed pressure altitude that was the case in older model versions. Several changes were also made to the model turbulence scheme to increase vertical turbulent transport of water vapor and improve spatial variations in boundary layer depth.

The dynamic ocean that is coupled to the atmosphere for many ROCKE-3D applications was improved via enhancements to the parameterization of mesoscale horizontal diffusivity due to subgrid scale ocean eddies and small-scale vertical diffusivity. The mesoscale diffusivity is now stronger at the ocean surface and decreases exponentially with depth. The baroclinicity scaling of the diffusivity now utilizes a fixed length scale rather than the Rossby radius but retains the Rossby latitudinal dependence, an option that should only be used for Earth simulations. The vertical diffusivity now includes a contribution from tidal dissipation. The physics of interactive sea ice were updated to allow low lead fractions and closures of leads, independent

165 horizontal advection of snow mass, and thermodynamics based on an energy-conserving brine pocket parameterization that allows salt to directly affect specific heat and ice melt rates.

The model has the capability to include irrigation, in which water is withdrawn from lakes and then from an unlimited groundwater pool if lakes are insufficient. This groundwater pool is not dynamic with the full hydrological cycle and does not get recharged from runoff or infiltration ~~Groundwater recharge is not accounted for~~ (Sect. 2.4.1 in Kelley et al., 2020), so there are small increases or decreases in total planet water mass and sea level depending on whether a particular climate simulation draws from or adds to the groundwater supply. For simulations without humans (i.e. modern Earth), irrigation is not included.

2.2. Updates to ROCKE-3D 2.0 since version 1.0

This section describes model development that was done specifically for ROCKE-3D 2.0 and that are not necessarily in ModelE2.1. These includes improvements to land surface hydrology, introduction of a geothermal heat flux, capabilities for thin atmospheres, and improved calendaring options.

2.2.1. Land

In ~~planet-ROCKE-3D~~ 1.0, surface hydrology on land was as described for soils in [Rosenzweig and Abramopoulos \(1997\)](#) and for lakes and rivers in [Russell et al. \(1995\)](#) and [Schmidt et al. \(2014\)](#). Surface soil hydrology physics remain the same in ~~planet-ROCKE-3D~~ 2.0, but that for lakes and rivers has been considerably updated to be generalized for arbitrary topography. The original physics and the new physics ~~are~~ will be described in detail elsewhere ~~in the future~~ (Kiang et al., in preparation). Briefly, as before, each grid cell can have a single dynamic lake of conical shape that can shrink and swell in area and depth, and rivers transport excess lake water between grid cells. The new physics: allow lake size and initial lake water to be initialized separately; introduce lake bathymetry that scales with grid size; allow variable river speed based on lake surface relative altitudes (previously the speed was fixed by topography); allow rivers to transport water in any of 8 gridded directions (sides and corners) from a grid cell based on relative lake surface altitudes (previously only one direction was allowed and was prescribed based on topography only); allow very small, shallow lakes to dry out; and improve prediction of the mixed layer depth and therefore lake temperature. These new physics have been evaluated for modern Earth, and result in improved continental recycling of moisture and precipitation in the tropics, improved seasonality of river flow, and improved surface temperature and seasonality of ice cover for large lakes; in addition, they have been evaluated for sensitivities of idealized flat land planets ~~(Kiang et al., in preparation)~~. A publication about these evaluations is forthcoming. The option to prescribe river directions as in ~~planet-ROCKE-3D~~ 1.0 is retained for cases of known or historical river directions. The ~~planet-ROCKE-3D~~ 2.0 dynamic lakes and rivers make the land surface hydrology suitable for modern Earth, Earth paleo-topography, and other planets, from idealized flat land planets to those with known topographies like Venus or Mars, as well as for climate changes on any planet that would affect the spatial patterns of precipitation. However, the new prognostic river routing can result in

195 very large lakes where there would otherwise be wetlands, effectively removing vegetation because currently wetlands and flooding physics are not represented, so the prognostic rivers should not be used for carbon cycle simulations.

The processes that accumulate water in lakes, combined with the horizontal transport of water across the land surface via runoff and river flow, are important for capturing the mass water balance on land. The geographic distribution of lakes determines the evaporative surface area, the surface cover of liquid water (which can produce glint as a sign of liquid water), the regional formation of clouds, the surface energy balance, and temperature. Previous ROCKE-3D 1.0 experiments could have lakes either without river transport, leading to potential excess buildup of water in some grid cells, or with prescribed river directions in a single direction, which could still result in excess buildup and generally lacked realism in the spread of water across the land surface. Fully prognostic, multi-directional river transport reproduced the retention of water in areas that are in fact wetlands, promoted continental recycling of precipitation, and significantly rectified the warm bias. The generalized lake/river dynamics also enable the simulation of surface hydrology of idealized planets for which only a flat topography is specified, in which case river directions cannot be estimated from topography but must arise from the relative balance of surface water between grid cells.

The new lake and river physics, which were not used in this work but are available in the public release of ~~planet-ROCKE-3D~~ 2.0, can be turned on with these different options: 1) using a new function to scale conical lake shape (cone slope) with grid size, or alternatively an input file that explicitly prescribes every grid cell with its lake cover fraction and sill height; 2) a new variable speed river flow; 3) defining river directions can either be assigned from an input file or use the new prognostic multi-directional river directions; 4) decide whether to allow small lakes to evaporate or not; 5) use the new more variable mixed layer depth. Alternative river direction files may be constructed for a variety of topographies and climates.

2.2.2. Geothermal heat flux

215 Coupled atmosphere-ocean GCM simulations of modern Earth have seldom included geothermal heat flux (GHF) as a background energy source, since the ~~radiative foreingenergy~~ it contributes to Earth's climate system is low ($\sim 0.1 \text{ W m}^{-2}$ on global average; Davies, 2013) compared to anthropogenic greenhouse gases (see e.g. Fig. 7.6 in Intergovernmental Panel on Climate Change (IPCC), 2023). However, the influence of GHF through the ocean floor in particular has long been recognized as a source of warming that disrupts deep ocean stratification and the rate of ocean circulation at depth, whether the GHF is applied uniformly across the ocean floor or in a more realistic spatially-variant pattern (Adcroft et al., 2001; Emile-Geay and Madec, 2009; Hofmann and Maqueda, 2009; Scott et al., 2001). We expect that interior heat flow from any rocky planet interior, whether from primordial radiogenic heat or contemporary tidal heating, would be at least as important for a variety of other rocky worlds. These include Earth-like planets with extensive snow and ice cover, in highly elliptical orbits or otherwise on the outer margins of habitability, as well as icy moons with deep interior oceans (e.g. Běhouňková et al., 2010; Butcher et al., 2017; Colose et al., 2021; Hendrix et al., 2019; Henning and Hurford, 2014).

In ROCKE-3D, GHF is available as an optional boundary condition that can be added in any configuration. Heat fluxes may be prescribed as either a uniform heat flow or a spatially variant field, and they may be prescribed for land only, for ocean only, or for both. If applied to land, fluxes will be added to the lowest soil level and to lakes. If a Q-flux ocean is used, heat will be applied to the mixed layer and sea ice. If a dynamic ocean is used, fluxes will be added to the lowest ocean layer, whether that ocean is open or ice-covered. Heat fluxes are not applied to prescribed sea surface temperature fields, nor will they have an impact on the bases of prescribed land ice sheets.

NetCDF input files for spatially-variant heat fluxes should be prepared at the appropriate model resolution used for a given simulation. We provide sample modern Earth heat flow files at 4x5 degree resolution for both the atmosphere and the ocean, typical for exoplanet runs; and at 2x2.5 and 1x1.25 degrees for when a higher atmosphere/ocean resolution is needed (Fig. 1).




Figure 1. Modern Earth [geothermal](#) heat fluxes at 4x5 (left) and 1x1.25 (right) degree resolutions from a digital 2x2 degree equal area map, developed by Davies (2013) from a collection of more than 38,000 heat flow measurements from around the world.

2.2.3. Thin atmospheres

If atmospheric pressure at the surface of the planet is significantly lower than the one on modern Earth, then special care should be taken to represent certain processes and to ensure the stability of the model in general. Typically, we consider the atmosphere “thin” if the surface pressure falls below the triple point of water (~6 [mbar](#)). Examples of such planets are modern Mars and planets with transient atmospheres induced by impactors or periods of high volcanic activity (Aleinov et al., 2019). Atmospheres below the triple point of water cannot support liquid water, and while most model algorithms (clouds, precipitation) handle this automatically, a special option is provided to exclude any movement of liquid water in the ground. This option has to be enabled (Appendix B) for proper simulation of the hydrological cycle on such planets.

For thin atmospheres, the ground temperature is mainly determined by the competition of absorbed [solar-stellar](#) and outgoing longwave radiation fluxes. As a result, a high temperature gradient is present between the insolated and shaded parts of the surface. This causes the lower atmospheric temperature to decouple from the ground temperature over the shaded regions due to the very stable stratification near the surface, especially for a radiatively neutral atmosphere. In such cases, the conditions

in the lower atmosphere are mainly determined by the sensible heat flux over the insolated regions, which is much lower than other surface fluxes, and special care should be taken to avoid its distortion by numerical artefacts. The algorithms for surface fluxes in ~~planet-ROCKE-3D~~ 2.0 were updated to properly handle such conditions. We also disable the horizontal heat transport in the planetary boundary layer; these fluxes are negligible under thin atmospheric conditions, but they can cause numerical instabilities. At the top of the atmosphere, the atmospheric layers are very thin, and may require very short time step to maintain their stability with respect to radiative heating/cooling. To avoid this problem, we provide an option to treat the upper three layers as isothermal layers by the radiation model (Appendix B). In the current release, the model can handle surface atmospheric pressures down to 10 microbar. ~~This configuration was already used to simulate a~~ A thin CO atmosphere is as a likely candidate for a transient volcanically-induced paleo-lunar atmosphere (Aleinov et al., 2019; Needham and Kring, 2017).

Depending on atmospheric composition, extremely thin atmospheres may require non-Local Thermal Equilibrium (non-LTE) corrections in the radiation algorithm. These have not been implemented yet and may become available in future releases. For exceedingly thin atmospheres (< 10 microbars) non-LTE radiative effects may become important, however, non-LTE physics are not implemented in SOCRATES. For Early Moon simulations we assume CO-dominated atmospheres, in which, fortuitously, non-LTE effects are negligible and can be ignored ~~One should notice though, that in some cases non-LTE corrections may not be needed even for atmospheres thinner than 10 microbar, as is the case for a CO atmosphere~~ (Forget et al., 2017). However, caution should be exercised that for different atmospheric compositions (e.g. CO₂-dominated) non-LTE effects can be important and can have non-negligible impacts on results. ~~A thin CO atmosphere is a likely candidate for a transient volcanically-induced paleo-lunar atmosphere (Aleinov et al., 2019; Needham and Kring, 2017).~~

2.2.4. Calendar and equation of time

The calendar facility in ModelE is designed to generate a custom calendar that is tailored to the various exoplanet orbital parameters. With some important caveats that are elaborated on below, the generated calendar maintains a familiar correspondence to conventional Earth-based calendars, e.g. each orbital year is divided into 12 months of varying duration, but integral-integer numbers of days. An attempt is made to align seasons by forcing the vernal equinox to be at the same fraction of the calendar as is the case for the Earth. Each day is divided evenly into 24 “hours”, which are generally different than the expected 3600 seconds in duration.

To avoid complexities analogous to leap days/years, the rotational period is, by default, quantized to ensure an integral-integer number of days per orbit. More specifically, the rotational period is adjusted in a minimal manner such that the number of rotational periods evenly divides the orbital period. We adjust the rotational period rather than the orbital period, because we generally have superior information about orbital periods of actual observed exoplanets. This quantization can be deactivated, in which case the orbital year and the calendar year are of different durations, which then requires some caution in interpreting some diagnostics, i.e. an “annual mean” quantity might require more than one orbit to properly average out.

The duration of each of the 12 months is determined by a heuristic based upon a reference orbit and calendar for the Earth. More precisely, we determine the longitude relative to the vernal equinox of the start time of each month in the reference 365-day pseudo-Julian calendar used by ModelE for modern Earth simulations. The exoplanet calendar attempts to preserve the same angles for the start times of its months, but makes small adjustments to ensure an integral-integer number of days in each month. In this manner, all other things being equal the exoplanet “February” will generally be a shorter month, simply because it is a short month in a conventional calendar. This aspect is easily dominated by other factors for planets with large eccentricities and/or different longitude of the periapsis as compared to that of the Earth.

For exoplanets with a large separation between rotational and orbital periods, the above calendaring system works quite well, but the implementation requires and allows further adjustments to better support slow rotators. By default, the minimum number of calendar days in a year is 120 (roughly 10 per month). For a slow rotator, this means that calendar days will lose direct correspondence with the solar days. This default can be overridden, which can result in months that have 0 days. Further, for extremely slow rotators such as Venus, the quantization of the rotational period mentioned above is too severe, and the implementation therefore provides an option to deactivate the default quantization. In such cases, the calendar is of relatively little use, and useful diagnostics generally require averages over many orbits to avoid day-night biases within any given orbit. The implementation of calendar in planet-ROCKE-3D 1.0, or more precisely the calculation of the actual hour angle, did not include the effects due to the equation of time (EOT) which arises due to the difference in the length of a solar day at apoapsis versus periapsis and also the obliquity-induced effect between solstices and equinoxes. For the Earth this effect is minor and not usually included when studying modern climate, but for highly elliptical orbits this effect can be significant (Colose et al., 2021). ROCKE-3D 2.0 corrects this oversight, and introduces 3 options for EOT that can be chosen by the user upon model configuration: “off”, “naïve”, and “default”. The “naïve” option implementation neglects the contributions due to obliquity. This option and “off” are provided for backward compatibility, and we recommend that the correct formula provided by “default” is always used.

2.3. Radiation schemes

Two fundamentally different radiation schemes are available in ROCKE-3D: the first is the GISS radiation (denoted as “G” in our naming convention; Sect. 4), which is strongly optimized for present-day and near-term paleoclimate Earth applications (Kelley et al., 2020 and references therein), and the other is the SOCRATES radiation (denoted as “S”; Amundsen et al., 2016, 2017; Edwards, 1996; Edwards and Slingo, 1996), whose implementation in ROCKE-3D has been presented in Way et al. (2017). The GISS radiation is much faster, but has limited applicability, and suffers accuracy losses when the atmosphere and incident stellar energy distribution (SED) under study deviates appreciably from that of preindustrial Earth and present-day Sun combination. This is especially important when it comes to making large changes to the amount of atmospheric absorbers (e.g. H₂O, CO₂, CH₄), large changes in the temperature profile, and when using SEDs from cool stars where the peak in radiation is shifted into the near-infrared. SOCRATES, on the other hand, is much more flexible and can simulate different

atmosphere and SEDs combinations very easily, albeit with preprocessing requirements to create the so-called “spectral file”.

315 A spectral file is a construct native to SOCRATES which contains all aspects of the radiation problem, including the spectral interval and gauss point grids, gas absorption coefficients for major and minor species, Rayleigh scattering coefficients, cloud optical properties, and aerosol optical properties. Different spectral files can be fed in to SOCRATES at runtime, meaning that the radiative transfer problem can be defined for numerous diverse atmospheres without needing to modify the code and recompile SOCRATES. Generally, better accuracy for more exotic star-atmosphere combinations requires increasing the
320 density of spectral intervals and gauss points which results in a slower runtime performance ([also see Sect. 4.5](#)). However, the speed reduction is necessary to obtain highly accurate radiative transfer results for the variety of atmospheres that can be studied with ROCKE-3D. All simulations presented here use the ga7_dsa spectral file (Appendix C).

The motivated user can create their own unique spectral files using a set of shell and python scripts provided publicly on
325 Github (see Code availability), which wrap native SOCRATES routines for creating and manipulating all components of the spectral file. However, as part of ROCKE-3D, we provide numerous pre-computed spectral files, which should satisfy the requirements of most use cases (see Data availability). These are divided into several general classes based on the atmospheric composition which they serve, referencing Earth and Mars time-periods. Each class has numerous spectral files optimized for specific subsets of the atmospheric composition. As the name implies, “Modern Earth” spectral files are only appropriate for
330 modern Earth atmospheric compositions and temperature ranges, while allowing for increases and decreases by several doublings of CO₂ or CH₄. “Archean” spectral files have been constructed to incorporate high amounts of CO₂ and CH₄ in an N₂ dominated anoxic atmosphere while allowing total pressures up to 10 bars. The Paleoproterozoic spectral file includes a combination of low O₂ and moderately elevated CO₂ and CH₄ in an otherwise N₂ dominated atmosphere. We signify “Mars Through Time” to describe spectral files which feature CO₂ dominated anoxic atmospheres, valid for modern Mars but also
335 any generic dense CO₂ atmosphere (Del Genio et al., 2019b; Guzewich et al., 2021; Schmidt et al., 2022). We have also constructed additional spectral files for other worlds, including Titan and a putative early lunar atmosphere (Aleinov et al., 2019). Stellar spectra have been constructed for numerous stars ranging from ultracool M-dwarf stars to late F-dwarfs. An at length description of the technical details of the currently available spectral files and stellar spectra is included with our publicly available materials (ROCKE-3D spectral files). Refer to Appendix C for additional information.

340 2.4. Atmospheres

In [planet-ROCKE-3D](#) 1.0 we only provided template configurations for one type of atmosphere, that of preindustrial Earth but without atmospheric aerosols, O₃, and stratospheric formation of H₂O from CH₄ oxidation. In [planet-ROCKE-3D](#) 2.0, in addition to the [planet-ROCKE-3D](#) 1.0 atmospheric configuration (denoted as “x” in our naming convention; Sect. 4), we added two additional template configurations: one (denoted as “A”) is the exact atmosphere of preindustrial Earth for the year
345 1850, and the other (denoted as “N”) is the same atmosphere but without aerosols, O₃, and stratospheric formation of H₂O

from CH₄ oxidation, and with the O₂ and Ar of the atmosphere replaced by N₂. This makes the N configuration essentially a pure N₂ atmosphere, other than trace components present like CO₂ and N₂O, both of which are at their preindustrial Earth levels. ROCKE-3D can run using a wide range of additional compositions, e.g. pure CO₂ (Mars), pure CO or N₂ (Aleinov et al., 2019), as well as mixtures of non-condensable gases.

2.5. Oceans

The ocean parameterization chosen is key-very important in the simulated climate of any planet with a substantial ocean present. In Earth's modern climate, most of the poleward planetary-heat transport occurs in the atmosphere except in the deep tropics must act to diverge energy from the tropics and converge energy in Polar Regions. Most of this poleward energy transport occurs in the atmosphere (Held, 2001; Klinger and Marotzke, 2000; Trenberth and Caron, 2001), with only a small fraction of the energetic demand taken up by the ocean. Nonetheless, Nnumerous studies have also demonstrated a very strong coupling between the sea ice margin and convergence of ocean heat transport at higher latitudes (e.g. Aylmer et al., 2020; Bitz et al., 2005; Ferreira et al., 2011; Rose et al., 2013; Winton, 2003). An exception is in the tropical belt, where the atmospheric energy transport is a small residual of contributions from the Hadley cell's low-level equatorward branch (where most of the moisture resides) and upper-level poleward flow. Net poleward transport occurs due to the higher potential energy aloft and the rising and sinking of air in the overturning circulation. The equatorial ocean favors a significant role for ocean heat transport into the subtropics (Held, 2001; Klinger and Marotzke, 2000), however, the atmosphere quickly overtakes the ocean heat transport by 15–20° latitude in both hemispheres. Numerous studies have also demonstrated a very strong coupling between the sea ice margin and convergence of ocean heat transport at higher latitudes (Aylmer et al., 2020; Bitz et al., 2005; Ferreira et al., 2011; Rose et al., 2013; e.g. Winton, 2003), which must be small beyond the ice edge due to the insulating effect of ice.

Even further, on tidally locked worldsFor exoplanets, the effect of ocean dynamics can produce results that are qualitatively far different than that assuming Q-flux=0 (Del Genio et al., 2019b; Hu and Yang, 2014). Most notably, on synchronously rotating worlds, the presence of a dynamic ocean can expand the area of deglaciated ocean on planets near temperatures where sea ice is expected to form. Generally, a dynamic ocean forms what is termed a “lobster pattern” in temperature, while a Q-flux=0 ocean forms an “eyeball” world (e.g. Pierrehumbert, 2010; see also Fig. 2a and 2b in Del Genio et al., 2019).

As in planet-ROCKE-3D 1.0, three ocean configurations are available. The first configuration (prescribed sea surface temperature (*ssT*); denoted as “p” in our naming convention; Sect. 4) has no interactive ocean but rather a prescribed sea ice extent and *ssT*. Although no ocean calculations happen in this configuration, the ocean heat transport is implied and is equal to that of preindustrial Earth. The second (Q-flux; denoted as “q”) is a Q-flux ocean (Miller et al., 1983; Russell et al., 1985), in which (contrary to planet-ROCKE-3D 1.0 where we used Earth-like heat fluxes) the heat fluxes throughout the ocean are set to zero. This is a common assumption in exoplanet studies with important consequences to the simulated climate, as shown here. Even further, on tidally locked worlds the effect of ocean dynamics can produce results that are qualitatively far different than that assuming Q-flux=0 (Del Genio et al., 2019b; Hu and Yang, 2014). Most notably, the presence of a dynamic ocean

can expand the area of deglaciated ocean on planets near temperatures where sea ice is expected to form. Generally, a dynamic ocean forms what is termed a “lobster pattern”, while a Q flux=0 ocean forms an “eyeball” world (e.g. Pierrehumbert, 2010; see also Fig. 2a and 2b in Del Genio et al., 2019). In this model setup the atmosphere and sea ice are coupled to a mixed-layer ocean of a specified depth, which allows for time-varying storage and release of heat, and source of evaporation to the atmosphere. The depth of the Q-flux=0 ocean in ~~planet-ROCKE-3D~~ 2.0 is 100 m (the default value was 65 m in ~~planet-ROCKE-3D~~ 1.0), and it is trivial to change to any desired depth. The last configuration (“dynamic ocean”; denoted as “o”) is a fully dynamic ocean, similar to what was used in ~~planet-ROCKE-3D~~ 1.0, with present-day Earth’s bathymetry, but any configuration can be used, e.g. a bathtub ocean with flat bathymetry used in aquaplanet simulations (Sect. 4.4.4). The initial conditions used are based on present-day Earth’s modern ocean configuration (Levitus et al., 1994; Levitus and Boyer, 1994).

2.6. Grid resolutions

The Earth version of ROCKE-3D, ModelE2.1, is routinely run in the fine horizontal atmospheric resolution of 2x2.5 degrees in latitude and longitude, denoted as “F” in our naming convention (Sect. 4), with 40 horizontal layers to 0.1 hPa. This model version is coupled to a dynamic ocean with a 1x1.25 degrees resolution in latitude and longitude and 40 layers to the ocean floor. Virtually all studies done thus far with ROCKE-3D have been performed at medium resolution (denoted as “M”), either using 40 or 20 atmospheric vertical layers and a 4x5 ocean with 13 layers. In ~~planet-ROCKE-3D~~ 2.0 we only kept the 20-layer atmospheric model version as a legacy (working but unsupported) option, and opted to use for all simulations the 40-layer atmospheric version. We also decided to include both the M and F resolutions, in order to be able to resolve with greater accuracy larger-than-Earth planets and fast rotators.

2.7. Template model configurations

In ~~planet-ROCKE-3D~~ 1.0 (Way et al., 2017), we provided template configurations for an Earth-like atmosphere with conditions similar to preindustrial (year 1850), but with zero aerosols, ozone (O₃), and stratospheric water vapor formation from methane (CH₄) oxidation. In ~~planet-ROCKE-3D~~ 2.0 we expanded the available options offered, which now include combinations of two radiation schemes, three different atmospheres, three ocean configurations, and two horizontal resolutions, resulting in a total of 36 supported configurations. Here we will describe the reasons that led us to the choice of such configurations, the decisions made to balance them, their differences, and their limitations. Their naming convention is presented in Fig. 2, which is the one used in the code repository and here, and a detailed discussion of their resulted climates will follow in the next sections.

P2{G,S}{A,x,N}{p,q,o}{F,M}40

425 **Figure 2.** Template configuration naming convention. For the meaning of the letters, see [Table 1](#)~~Table 4~~. As an example, a fine resolution configuration of preindustrial Earth atmosphere with a dynamic ocean and GISS radiation is named P2GApF40.

Table 1~~4~~. ROCKE-3D version 2.0 template configuration naming convention explanation.

Component	Explanation
Model version	P1: Template configuration of ROCKE-3D version 1.0 (Way et al., 2017)
	P2: Template configuration of ROCKE-3D version 2.0 (planet_2.0 ; this work)
Radiation	G: GISS radiation
	S: SOCRATES radiation
Atmosphere ^a	A: Atmosphere of preindustrial Earth (year 1850), as described by GISS ModelE CMIP6 simulations
	x: Same as A, without a) aerosols; b) O3; c) stratospheric H2O formation from CH4 oxidation
	N: Same as x, with O2 and Ar replaced by N2
Ocean	p: prescribed sea surface temperature and sea ice extent and thickness
	q: Q-flux=0 ocean, 100m mixed layer depth
	o: dynamic ocean; see next line for resolution
Resolution	M40: atmosphere: 4°x5° with 40 layers to 0.1 hPa; ocean, if o configuration: 4°x5° with 13 layers
	F40: atmosphere: 2°x2.5° with 40 layers to 0.1 hPa; ocean, if o configuration: 1°x1.25° with 40 layers

^a: In the text these configurations are frequently named Earth's (A), ~~planet-ROCKE-3D~~ 1.0 (x), and anoxic (N) atmospheres.

3. Generalizing ModelE into ROCKE-3D

430 In this work, ~~we use~~ the GCM variable names ~~of the model output are~~ in italics ~~of the model output~~ for many quantities, rather than a more verbose explanation. This helps the model user to exactly understand which model output we've used in our analysis, but also to detail which variables are important to analyze when creating a new world. These variables are explained in Appendix D.

3.1. Radiative balance

435 Bringing the model into radiative balance is a necessary step to create a control simulation in which temperature, as well as shortwave and longwave fluxes, are acceptably close to target values. There are multiple ways one can calibrate the model to achieve this. Our method is to bring the model into radiative balance under preindustrial conditions (described in Sect. 4.1), for which we use a prescribed ocean configuration and make changes to cloud parameters so that the absolute value of the net incoming ~~solar-stellar~~ and outgoing thermal radiation of the whole planet (*net_rad_planet*) is small (within $\pm 0.2 \text{ W m}^{-2}$). ~~For~~
440 ~~Earth-centric science questions, a~~A secondary goal is that the present-day net radiative balance of the planet is on the order of

+1 W m⁻², ~~which-but this~~ is not relevant to this work which focuses on a generalized planetary configuration rather than the current state of the planet and the impact humans have on it. It is important to note that not all configurations can easily (if at all) stay within those limits, mostly because when deviating from the actual preindustrial Earth there is no “real” ocean field to be used, so the atmosphere might ~~have a hard time~~ take much longer, or even fail, to fully adjust. This will be discussed later, on a case-by-case basis, when it occurs.

There are other, secondary criteria that also need to be loosely met, which are known for the present-day atmosphere, but can only be estimated for the preindustrial one. These include global mean surface air temperature (*tsurf*; 14-15 °C at present day, roughly 1 °C cooler at preindustrial); planetary albedo (*plan_alb*; about 29.6% at present day, unknown at preindustrial); ~~solar~~ stellar (shortwave) radiation net flux at the top of the atmosphere (*srnf_toa*; 239±2 W m⁻² at present day, probably the same at preindustrial); and ocean sea ice fraction (*oicefr*; 4% at present day, slightly greater at preindustrial). When balancing the model, we always try to have $|net_rad_planet| \leq 0.2 \text{ W m}^{-2}$, while trying to also maintain *srnf_toa* as close as possible to 239±2 W m⁻², by modifying the radiative balancing factors of clouds (see Sect. 4.1).

We only balanced the model for configurations of interest for the public release of the model. Intermediate configurations use balancing values which might or might not bring the model to radiative balance, depending on how much different model physics affect the modeled climatologies. In Sect. 3.3 below, some model runs will clearly be out of radiative balance, by design.

3.2. Effects of new physics on mean climatology

We checked whether the code merges across the Earth (Sect. 2.1) and planetary (ROCKE-3D 2.0; Sect. 2.2) model versions affected the mean climatology of the model in any significant way. For this, we simulated both the year 2000 and year 1850 climatologies, by performing 30-year simulations with the prescribed ocean (p) model: the first 10 years were used as a spinup, and the latter 20 for the analysis. This is twice as many years as we typically use for balancing the Earth model (both spinup and analysis); we decided to use more years than usual as an additional layer of safety, and as expected it was proven to be unnecessary. The set of simulations performed for the year 2000 showed that all key diagnostics mentioned in Sect. 3.14.1 are both within bounds and extremely close to each other across model versions, ensuring that the merging process was successful. The same applies for the 1850 climatology across model versions, where all simulations showed that they are in radiative balance. From now on, all simulations discussed are done with the merged product, ~~the planet-ROCKE-3D 2.0-version-of ROCKE-3D~~, and with 1850 base climatological conditions.

There are two major changes that as expected introduced major changes in the model physics, and require rebalancing: the change from GISS radiation to SOCRATES (using the ga7_dsa spectral file; Appendix C), and the coarsening of the default Earth resolution from F40 to M40. Note that the M20 resolution used in planet-ROCKE-3D 1.0 is not supported any more. When SOCRATES is used, *net_rad_planet* decreases by 1.3 W m⁻² and *srnf_toa* decreases by 6 W m⁻². When changing


resolution from F40 to M40 *net_rad_planet* increases by 0.8 W m⁻² for GISS and by 0.2 W m⁻² for SOCRATES radiation, while *srnf_toa* decreases by 2.0 and 2.6 W m⁻², respectively.

The change in resolution includes not only the physical change of grid box sizes, but also the ~~disabling-replacement~~ of the gravity wave drag parameterization from that used on the Earth version of the model (Kelley et al., 2020), with the simpler “Rayleigh friction” version used in ~~planet-ROCKE-3D~~ 1.0 (Way et al., 2017). Structurally, the M simulations employ a simplified version of “Rayleigh friction”, which exponentially decays wind components over a user-specified range of layers near the model top. The simplification is to apply a user-specified exponential time constant per layer. This replaces the parameters traditionally associated with momentum loss to a solid surface (e.g. drag coefficient, air density, and quadratic dependence upon wind speed). Furthermore, the Earth-oriented version has a special tuning of its “Rayleigh friction” near the poles; this was especially important for getting the correct stratospheric wind structure in M configurations, because they completely lack the parameterized gravity wave momentum transports that shape stratospheric winds in later model generations (F resolution). ~~no-This~~ geographic adjustment is ~~imposed, unlike the original-dropped in non-~~Earth-oriented ~~versionsimulations~~. An even further improved version of the gravity wave drag that is present to the high model top version of the Earth model (Rind et al., 2020) will become available in the next release of ROCKE-3D, but again only for the F model resolution, not the M, and with many more than 40 layers (102 in Rind et al., 2020). When using the F40 resolution with the simplified gravity wave drag, the *net_rad_planet* change for GISS (SOCRATES) radiation is -0.46 (-0.50) W m⁻², and the corresponding change for *srnf_toa* is -0.24 (-0.29) W m⁻².

Another necessary change between resolutions is the time step used in the Shapiro filter, which is a hyperdiffusion operator that is applied to remove numerically-induced high-frequency noise from the velocity and tracer fields (Shapiro, 1970, 1975): in the fine (F40) resolution this is 225 s, while in the medium resolution (M40) it is 450 s. This change has only a marginal impact in the key diagnostics *net_rad_planet* and *srnf_toa*.

3.3. Creation of a generalized Earth configuration

ModelE2.1 contains a control configuration which is meant to simulate preindustrial Earth (year 1850; E2.1GApF40_1850). This contains some Earth-centric parameterizations, each for a different reason, that are not valid for use in other planetary configurations. In ~~planet-ROCKE-3D~~ 1.0 such configurations also existed, but the changes from the tight Earth configuration to the more general one were not studied systematically. Here we present in detail which options we have either eliminated or generalized, and their individual impact on model results, towards the creation of the several supported template configurations in ~~planet-ROCKE-3D~~ 2.0, described in Sect. 4. We focus on the net planetary radiation (*net_rad_planet*) and the ~~solar-stellar~~ (shortwave) radiation net flux at the top of the atmosphere (*srnf_toa*). Box-and-whisker plots of those diagnostics for the 20 years of the simulation (following 10 years of spinup) are presented in Fig. 3. It is important to mention that many of the differences discussed here would have been reduced by rebalancing the model, but since these are interim simulations that are not meant to be used for production the absolute values of such differences do not matter much.



520

Figure 3. Box-and-whisker plots of net planetary radiation (*net_rad_planet*; left) and solar-stellar (shortwave) radiation net flux at the top of the atmosphere (*srnf_toa*; right) for the intermediate simulations described in Sect. 3.3. The part of the simulation names before the underscore is explained in Sect. 3-4, with the addition that E2.1 and T3 mean the default and the updated E2.1 version of GISS ModelE, respectively. For the part after the underscore, see Table 2Table 2.

525

Table 22. Description of intermediate simulations described in Sect. 3.3 with the GISS radiation. All configurations that start with P2G in the table were also performed using the SOCRATES radiation instead of GISS, in which case we replace G with S in the name, e.g. P2GApM40_1850 becomes P2SApM40_1850, but are omitted below for brevity.

Name	Year	Incremental from	Description
<u>E2.1GApF40_2000</u>	<u>2000</u>	<u>None</u>	<u>Base Earth configuration of ModelE2.1.</u>
<u>T3GApF40_2000</u>	<u>2000</u>	<u>E2.1GApF40_2000</u>	<u>Base Earth configuration of updated E2.1.</u>

Name	Year	Incremental from	Description
<u>P2GApF40_2000</u>	<u>2000</u>	<u>T3GApF40_2000</u>	<u>Starting F40 configuration of planet_</u> planet_ <u>ROCKE-3D 2.0, includes planet_</u> planet_ <u>ROCKE-3D 1.0 and updated E2.1.</u>
<u>P2GApM40_2000</u>	<u>2000</u>	<u>P2GApF40_2000</u>	<u>Starting M40 configuration of planet_</u> planet_ <u>ROCKE-3D 2.0, includes planet_</u> planet_ <u>ROCKE-3D 1.0 and updated E2.1.</u>
E2.1GApF40_1850	1850	None	Base Earth configuration of ModelE2.1.
T3GApF40_1850	1850	E2.1GApF40_1850	Updated Earth configuration of ModelE2.1.
P2GApF40_1850	1850	T3GApF40_1850	Starting point (F40) of planet_ <u>planet_</u> ROCKE-3D 2.0. It includes planet_ <u>planet_</u> ROCKE-3D 1.0 and updated E2.1, for the A atmosphere (<u>Table 1</u> Table 1).
P2GApM40_1850	1850	P2GApF40_1850	Starting point (M40) of planet_ <u>planet_</u> ROCKE-3D 2.0. It includes planet_ <u>planet_</u> ROCKE-3D 1.0 and updated E2.1, for the A atmosphere (<u>Table 1</u> Table 1).
E2.1GApF40_2000	2000	None	Base Earth configuration of ModelE2.1.
T3GApF40_2000	2000	E2.1GApF40_2000	Base Earth configuration of updated E2.1.
P2GApF40_2000	2000	T3GApF40_2000	Starting F40 configuration of planet_2.0, includes planet_1.0 and updated E2.1.
P2GApM40_2000	2000	P2GApF40_2000	Starting M40 configuration of planet_2.0, includes planet_1.0 and updated E2.1.
P2GApF40_nodrag	1850	P2GApF40_1850	Simplified “Rayleigh friction” gravity wave drag.
P2GApM40_DT	1850	P2GApM40_1850	Half time step for Shapiro filter.
P2GApM40_noIRR	1850	P2GApM40_1850	No irrigation.
P2GxpM40_noO3etc	1850	P2GApM40_noIRR	No aerosols, no O ₃ , and no H ₂ O formation from CH ₄ oxidation in the stratosphere.
P2GApM40_notEarth	1850	P2GApM40_1850	No RADN8 ^a , maxctop ^{b,c} , and SW-H ₂ O absorption increase ^d .
P2GApM40_interim P2GApF40_interim	1850	P2GApM40_noIRR P2GxpM40_noO3etc P2GApM40_notEarth	Combination of noIRR, noO3etc, and notEarth, and uniform GHGs distribution, for both M40 and F40 resolutions. These are the starting point for the x and N atmospheres (<u>Table 1</u> Table 1).
P2GxpM40_noRADN4 P2GxpF40_noRADN4	1850	P2GApM40_interim P2GApF40_interim	Disabled second-order optimization of LW radiation specific to Earth conditions.
P2GxpM40_UTCF P2GxpF40_UTCF	1850	P2GApM40_interim P2GApF40_interim	Allows for a more physically-consistent profile of longwave flux divergence near the top of the atmosphere.

- ^a: Earth-centric clouds inhomogeneity correction, but instead use a constant value of 0.12.
- ^b: CPU-saving setting which skips cloud calculations above a fixed pressure level (50hPa for Earth).
- ^c: This is the only change that is relevant for SOCRATES in P2SApM40_notEarth.
- ^d: Shortwave long-path H₂O absorption increase.

530

After having constructed preindustrial Earth model configurations for all combinations between fine and medium resolutions, and GISS and SOCRATES radiations, the next goal was to set up configurations for uninhabited planet simulations. These primarily included configurations that do not include irrigation, aerosols, and O₃, but also other changes related to clouds (P2GApM40_notEarth in Table 2). We quantified the difference they introduce to *smf_toa*, and by assuming linearity in responses (in the absence of a better assumption), we modified our target *smf_toa* value from $239 \pm 2 \text{ W m}^{-2}$ (Sect. [3.1-4.1](#)) for the radiative balancing to a new value, as described below. All simulations have been performed with the M40 resolution, which is anticipated to become the most used one.

535

540

The removal of irrigation does not produce any noticeable change in *net_rad_planet* and *smf_toa*. The same applies to key diagnostics like surface temperature and total cloud fraction, where any regional changes calculated are not statistically significant. On the other hand, and as expected, removing aerosols, O₃, and the formation of H₂O in the stratosphere by CH₄ oxidation produces a large change in *net_rad_planet* for GISS (SOCRATES) radiation of $+1.33 \text{ (-1.60) W m}^{-2}$ and $-3.5 \text{ (-8.7) W m}^{-2}$ for *smf_toa*. The large difference of *smf_toa* between the two radiation schemes leads to a very different response in clouds, which results in a *net_rad_planet* change of a different sign between the two model configurations. Regionally, the removal of aerosols, O₃, and stratospheric H₂O from CH₄ oxidation completely dominates the change in total cloud cover.

545

~~Cloud cover Throughout the tropics cloud cover~~ increases ~~nearly everywhere throughout the tropics~~ by as much as 60% locally, but ~~has no effect~~ changes less than 5% anywhere else (Fig. 4). It also has some statistically significant impact on *tsurf* at polar latitudes and in particular in Antarctica, but the effect is overall smaller than the removal of Earth-centric adjustments (P2GApM40_notEarth in Table 2) that cause a cooling of surface temperatures over land almost everywhere, which largely dominates the net effect of both changes (Fig. 4). Over the ocean the changes of *tsurf* are negligible, primarily because we use

550

the prescribed ocean configuration (p) which does not allow changes to *sst*, a major driver to *tsurf*. It has to be noted that the choice of atmospheric composition described here (no ozone, aerosols, and stratospheric water vapor from methane oxidation; x configuration in Table 1) was made for consistency with the planet-ROCKE-3D 1.0 non-Earth configurations. We decided to keep that legacy configuration in planet-ROCKE-3D 2.0 for continuity, although it contains inconsistencies, in that the atmosphere contains O₂ but not O₃. We also created a set of configurations in planet-ROCKE-3D 2.0 (anoxic Earth) that

555

eliminates that inconsistency, which we recommend to be the default one for future studies. More on this in Sect. 4.




Figure 4. Surface temperature (top row) and total cloud fraction (bottom row) change due to the removal of aerosols, O₃, and stratospheric water vapor from CH₄ oxidation (left column), some Earth-centric corrections (including the cloud-top limit, see text and Table 2; middle column), and the combination of the two (right column). Note that the right column also includes the effect of irrigation removal, which is statistically insignificant in the preindustrial atmosphere used here. Dots show grid boxes where the calculated differences are not statistically significant (p > 0.05).

In another simulation generalized for non-Earth simulations, we removed a clouds heterogeneity parameterization whose spatial distribution is relevant for Earth only (first introduced in ModelE by Schmidt et al., 2006), and used a constant factor instead. This factor scales the cloud optical depth down to account for the fact that variable optical depth within a partially cloudy gridbox causes shortwave extinction to be less than what one would get from a homogeneous cloud. ModelE uses a

map of the inhomogeneity correction appropriate to Earth from the International Satellite Cloud Climatology Project (ISCCP) D1 cloud climatology (Rossow et al., 2002). In order to not impose an Earth-like pattern of inhomogeneity, we used instead a global mean value of 0.12 (see Appendix B for technical instructions). In addition, we removed a shortwave long-path H₂O absorption increase, which was found to be necessary for Earth simulations to correct an underestimate of the direct and diffuse shortwave H₂O band absorption analytical fit used in the GISS radiation when compared against measurements, particularly for high-humidity cases. All of those changes are only relevant when using GISS radiation, not SOCRATES. We also allowed the clouds code to run up to the model top (0.1 hPa), instead of up to 50hPa, a limit inspired by present-day Earth and set for computational efficiency. Other atmospheres, especially those without a stratospheric temperature inversion, might have clouds much higher than where they exist on Earth, so removing this Earth-centric limit is important. This change is relevant for both GISS and SOCRATES simulations. The GISS radiation simulation with those adjustments removed resulted in a change in *net_rad_planet* of -1.27 W m⁻² and *srnf_toa* of -1.9 W m⁻². As expected, these changes require the model to be rebalanced. The clouds change (allow them to reach the model top) in the SOCRATES simulation did not change either of the diagnostics beyond noise, since no clouds are expected to exist in that model configuration above 50 hPa. Although no incremental simulation with was performed with the clouds change allowed to reach the model top alone was performed using the GISS radiation, it is expected that it will not affect the climatology, as was the case for SOCRATES.

After having tested all changes from the standard preindustrial Earth configuration, we combined all pieces together, and generated a candidate configuration for radiative balancing, for both radiation schemes and for both resolutions. These are the interim simulations in Table 2. For the GISS radiation, the total *srnf_toa* change for both resolutions is -5.7 W m⁻², while for SOCRATES it is a bit over -9 W m⁻² (-9.1 for M40 and -9.7 for F40). For the GISS radiation this net change adds up pretty linearly to the individual changes caused by the elimination of O₃ and aerosols (-3.5 W m⁻²) and adjustments in radiation (-1.9 W m⁻²). For SOCRATES, the change from O₃ and aerosols is -8.7 W m⁻² (no Earth-specific adjustments exist to be eliminated), which is also very close to the net effect of M40 (-9.1 W m⁻²). The *net_rad_planet* for the GISS radiation ends up pretty well balanced for M40 (-0.06 W m⁻²), probably by chance, while it is 0.4 W m⁻² for F40. SOCRATES is further away from radiative balance, with -1.8 and -2.3 W m⁻² *net_rad_planet* for M40 and F40, respectively.


As a last configuration test for the GISS radiation only, we removed a longwave correction (noRADN4 in Table 2), which we will only keep enabled in the atmospheres that exactly resemble modern Earth, and also enabled a more physically-consistent profile of longwave flux divergence near the top of the atmosphere (Appendix B). The longwave correction introduces a major radiative imbalance of +8.1 W m⁻² for both resolutions and a +1.0 W m⁻² (M40) and +1.3 W m⁻² (F40) change in *srnf_toa*, while the longwave flux divergence, probably the least impactful modification, only marginally changed results.

3.4. Analysis of additional diagnostics

Although *net_rad_planet* and *srnf_toa* are key diagnostics that affect the simulated climate of any atmosphere, there are many more diagnostics one can examine and understand differences between simulations. A key set of those, but by no means an

exhaustive one, are compared against the default preindustrial configuration of ModelE2.1 in Fig. 5. A key conclusion is the confirmation that virtually all [Earth](#) configurations across model versions ([ModelE2.1 \(E2.1GApF40_1850\)](#), [ModelE2.1 with updates \(T3GApF40_1850\)](#), [ROCKE-3D 2.0 starting point \(P2GApF40_1850\)](#), and [final ROCKE-3D 2.0 generalized configuration \(P2GApF40\)](#); also see Table 2) produce the same climatology. These are marked with a blue arrow in Fig. 5, where there are practically no differences between the simulations.

630



635

Figure 5. Relative differences of key diagnostics of the intermediate simulations listed in [Table 2](#) compared to the original preindustrial Earth simulation in ModelE2.1, marked with an orange arrow. Exactly equivalent simulations with different model versions (see text for details) are marked with blue arrows. Remember that most of the simulations presented here are not in radiative balance. Diagnostics are: *sst*: sea surface temperature; *oicefr*: ocean ice fraction; *lakefr*: lake fraction; *grnd_alb*: ground albedo; *plan_alb*: planetary albedo; *wsurf*: wind speed at surface; *snowdp*: snow depth; *snowfr*: snow fraction; *snowfall*: snow fall; *prec*: precipitation; *cldi*: cloud condensed ice column; *cldw*: cloud condensed water column; *cldtpt*: cloud top temperature; *cldtpp*: cloud top pressure; *clwp*: cloud liquid

water path; *pcldh/pcldm/pcldl/pcldt*: high/middle/low/total cloud cover; *qatm*: atmospheric water vapor column; *qsurf*: surface air specific humidity; *tsurf*: surface air temperature; *srnf_toa*: net ~~solar-stellar~~ radiation at the top of the atmosphere. See Appendix D for more details.

A number of interesting patterns emerge when looking at the other simulations, which will be only briefly discussed here. Those simulations are not in radiative balance (Fig. 3), due to their intermediate nature, as mentioned earlier. First of all, the year 2000 simulations are slightly warmer as expected, which results in less ocean ice (*oicfr*) and reduced snowfall rate and snow amount (both depth and fraction). The warmer temperature also allows the atmosphere to hold more moisture at surface and in the column. Another persistent pattern is for the year 1850 simulations with SOCRATES, which calculate warmer cloud-top temperatures and higher pressures, implying that clouds reach higher altitudes when using the GISS radiation. Further, the simulations without O₃ are cloudier, in particular in the higher altitude clouds region. This is mostly due to the increased presence of ice clouds, a result of colder temperatures from the middle troposphere and upwards. Also note that these simulations do not have a stratospheric ~~ic temperature inversione~~, due to the absence of O₃, as further discussed in Sect. 05. The cloudier atmosphere also results in an increase in the planetary albedo.

4. Working with different model configurations

4.1. Radiative balancing per configuration

For preindustrial conditions, Earth is assumed to be very close to radiative balance, meaning that the absorbed shortwave ~~solar-stellar~~ radiation is approximately equal to the outgoing longwave radiation at the top of the atmosphere, to within a few tenths of a W m⁻², in the range 240 ± 5 W m⁻². For the various configurations of ModelE2.1, small variations of a single parameter (*U00a*) have sufficient differential impacts on SW and LW to serve as the final rebalancing mechanism, after previous tuning steps set *U00b*, *WMUI_multiplier*, and *radiusl_multiplier*. In the search for radiative balance across the multiple configurations of ~~planet-ROCKE-3D~~ 2.0, the four parameters were not required to remain close to one another across configurations, nor to E2.1 values. This freedom was taken to be justified due to the differences in resolution, radiation scheme, etc. being considered as major rather than minor perturbations to the system. Table 3 lists the results. A short explanation of these variables is presented in Appendix D, while more detailed explanations of some of them have been presented elsewhere (Kelley et al., 2020; Schmidt et al., 2006). Note that the appropriate name for *U00a* and *U00b* is U_a and U_b, respectively, and this name has been used in past publications, but we opted to use the variable name found in the model code here, similar to what we do with other quantities discussed throughout the manuscript.

For every model configuration, a new radiative balancing is likely to be required. We call this “balancing”, because we modify some key clouds-related properties in order to achieve radiative balance in the model, based on our best known configuration of a radiatively balanced case, that of preindustrial Earth (year 1850; Sect. 3.1). These are not unique choices, as there is more than one way to achieve radiative balance in the model, all of which are more or less equivalent. For this we use the p ocean

670 (Table 1), in order to let the atmosphere adjust to the best known state of the ocean at preindustrial times. After allowing the model to radiatively equilibrate for about 10 simulation years, we continue the simulation for 20 more years, which we average for evaluation. ~~This is equivalent with a 20 member ensemble mean.~~ When changes were needed, we performed them (not shown; only the final chosen values are presented in Table 3) and repeated the 30-year simulation. When the configuration is in radiative equilibrium (the planet neither gains nor loses energy, averaged over 20 years), we perform a 500-year simulation
675 (a complete overkill, but for consistency with the other simulations) and average the last 100 years for the analysis presented here. This was done for all 12 simulations with the p ocean: (2 radiation schemes) x (3 atmospheres) x (2 resolutions). The balancing factors for each of the p oceans were then used for the similar configurations where the ocean was either q or o (Table 1), without any further adjustments. The final values per configuration are listed in Table 3, and their explanation can be found in Appendix D.

680

Table 33: Radiative balancing parameters of clouds for all p ocean configurations that bring the model to radiative balance. Two additional balancing knobs exist, *WMU_multiplier* and *radiusl_multiplier*, but both equal to 1 in all configurations, so they are omitted from the table for clarity.

	<i>U00a</i>	<i>U00b</i>	<i>WMUI_multiplier</i>	<i>radiusl_multiplier</i>
P2GApM40	0.675	1.1	0.001	1.1
P2GApF40	0.655	1	2	1.1
P2GxpM40	0.6	1.5	0.001	1.16
P2GxpF40	0.64	1.5	0.001	1.16
P2GNpM40	0.6	1.5	0.001	1.16
P2GNpF40	0.635	1.5	0.001	1.16
P2SApM40	0.7	0.6	0.001	1.01
P2SApF40	0.7	0.6	0.001	1.01
P2SxpM40	0.68	0.538	0.001	1.1
P2SxpF40	0.68	0.5	0.01	1.1
P2SNpM40	0.67	0.5	0.001	1.1
P2SNpF40	0.68	0.5	0.001	1.1

685 The use of a prescribed ocean in the balancing is necessary, because otherwise the ocean will try to absorb (or give back) the excess heat (or deficit) when *net_rad_planet* is positive (negative). This means that for a different configuration, one should either use a proper *sst* dataset for the balancing, or in the absence of one should use a Q-flux or dynamic ocean and select the balancing parameters from the atmosphere that is closer to the one under study.

It has to be stressed that we do not have any such information to perform a proper balancing on any non-modern-Earth planet;
690 one has to rely on the values for Earth, and pick the closest set to the new planet configuration. If that planet has an ocean, we

won't know its *sst* in the foreseeable future to balance radiation in the same way, so using a dynamic ocean is strongly recommended. Picking a set of values that structurally make sense from Table 3 (i.e. based on which radiation scheme, atmosphere, and resolution would be used) and then allowing the dynamic ocean to adjust as needed is the only feasible approach.

695 Following balancing, all model configurations have absolute *net_rad_planet* values either nearer zero or very close to the ± 0.2 W m⁻² range (Fig. 6). It is interesting to note that all simulated atmospheres using the GISS radiation, dynamic ocean, and fine resolution (P2G[AxN]oF40) have a much larger variability than any other combination of physics, while their corresponding simulations using the medium resolution (P2G[AxN]M40) have among the least.




Figure 6. Box-and-whisker plot of net planetary radiation (*net_rad_planet*) in W m^{-2} for all ~~p~~-ocean configurations following balancing. ~~using the last 100 years of 200 to 500-year-long simulations were used for the p and q ocean configurations, which were longer (1000 to 2000 years) for the o ocean (more for the dynamic ocean runs).~~ The bars show the 25/75 percentiles, the whiskers the 9/91, the bar is the median, the star is the mean, and outlier ~~years~~ are presented with dots. Also shown the $\pm 0.2 \text{ W m}^{-2}$ values, described in the text.


The balanced P2GApF40 simulation is in all climate-relevant respects the same as ModelE2.1 E2.1GApF40_1850, which is the most heavily tested configuration of GISS ModelE when a prescribed ocean is used (Sect. 3.4). Some regional differences across the model simulations are very minor, which confirms that the model skill and mean climatology ~~does~~ not change with the updated code in planet-ROCKE-3D 2.0 when compared with ModelE2.1.

4.2. Bringing the model to equilibrium

When simulating an atmosphere of any configuration on any given planet, ROCKE-3D needs some time to produce a stable climate for a climatological analysis. The time needed to arrive to equilibrium varies based on the atmosphere, ocean, and land to be simulated, the resolution, and the deviation of initial conditions from the final equilibrium. There are different metrics that define when the model is in equilibrium, which in general might depend on the needs of each individual experiment, since different components of the model equilibrate at different timescales. These timescales can vary from very few years for an atmosphere similar to present-day Earth and a prescribed ocean, centuries for biogeochemical carbon cycling involving land and ocean biology (Jones et al., 2016; Séférian et al., 2020), to several millennia for surface hydrology and a dynamic ocean (van den Hurk et al., 2016; Johns et al., 1997; Sitch et al., 2015; Wood, 1998). For a planet with an abundance of water, even in the absence of an ocean, surface hydrology equilibration times can also be at the several millennia timescales, to allow deep soils to come into full equilibrium. The same applies for dynamic ocean configurations which are several kilometers deep. The simulations listed in Table 2 are with a prescribed ocean and the water cycle does not need to be in complete equilibrium for the radiative equilibration of the atmosphere, so a few years are enough. The metrics used as criteria of an equilibrated climate also vary depending on the research question. Here, we want the net radiation of the planet to be near zero (within $\pm 0.2 \text{ W m}^{-2}$) in a long climatological mean over several years (orbits, for non-Earth planets), and also to have surface air temperature stabilized.

All simulations presented in Table 3 start with a correct set of balancing parameters for the underlying ocean (Sect. 4.1), so they are in near-instant equilibrium (just a few years). For the Q-flux and dynamic oceans, although using the correct balancing parameters, they are not in equilibrium at the beginning of the simulation because of imperfect parameterizations and forcings that can't exactly reproduce the prescribed ocean conditions. The Q-flux ocean takes much longer than the prescribed ocean to reach equilibrium, about a century in planet-ROCKE-3D 2.0 or 20 years in planet-ROCKE-3D 1.0, while for the dynamic ocean it takes a millennium or more: about 1500 using the medium resolution and 1000 for the fine resolution (Fig. 7). Note that even after 2000 years, not everything is in perfect equilibrium, but having some very-slow-equilibrating variables still drifting does not affect the other climatological parameters that have equilibrated already. For a deep dynamic ocean the time

735 for full oceanic equilibrium (e.g. a stable global potential temperature) can take many thousands of years (Fig. 8). Another
example is the total amount of liquid freshwater (*mwI*), which may need over 3000 years to equilibrate, but will likely require
much more for the less Earth-centric simulations in ROCKE-3D (Fig. 7). For that reason, a smart experimental configuration
can save several days or months of simulation time, when restart files are used properly. As an example, if a simulation one
wants to do resembles one of the model runs presented here, they can start from the end of our simulations using the restart
740 files we distribute (see Data availability section), greatly reducing the spin up time required. The same applies for all previously
published ROCKE-3D simulations - if the desired simulation is very similar to a previously published one, it is best to use a
restart file from that simulation.







Figure 7. Annual mean T timeseries of net_rad_planet (first row), $tsurf$ (second row), $gwtr$ (third row), mwl (fourth row), and $snowdp$ (last row) for the prescribed ocean simulations (left), the Q-flux=0 ocean (middle), and the dynamic ocean (right) with SOCRATES and the x atmosphere. The planet-ROCKE-3D 2.0 fine resolution model is shown in blue, the planet-ROCKE-3D 2.0 medium resolution in redpurple, and the planet-ROCKE-3D 1.0 medium resolution in green. Note that the dynamic ocean simulation shown with a blue line in the last column equilibrated much faster than the other two, so only 1000 years are presented.

The differences between planet-ROCKE-3D 1.0 and planet-ROCKE-3D 2.0 in the Q-flux ocean are due to the fact that the heat transfer used in planet-ROCKE-3D 1.0 resembles that of Earth so the initial conditions are much closer to equilibrium compared to Q-flux=0 in planet-ROCKE-3D 2.0. The dynamic ocean in planet-ROCKE-3D 1.0 also equilibrates much faster, in about 51000 years, while the equivalent simulation in planet-ROCKE-3D 2.0 needs 3-times-as-much-time-over 1500 years.

755 The fine resolution model ~~lies between the two~~equilibrates much faster, with about a millennium needed to achieve a stable
~~climate in about 500 years.~~ It is worth reiterating that here we are only referring to atmospheric radiative balance. If one has a
deep dynamic ocean like that of modern Earth and one is interested in science questions related to such oceans then it is vital
to consider whether the ocean itself is in equilibrium. One way in which this can be explored is by looking at the mean global
ocean potential temperature, but it is often useful to look at specific areas in the ocean as well as globally (Fig. 8).

760



765

Figure 8. Time series of ocean potential temperature (*pot_temp*) for a ~~planet-ROCKE-3D~~ 1.0 simulation for Modern Earth. The mean depth at level 11 is 1129 m and at level 13 is 3868 m. Even after 1500 years it is obvious that neither the global mean, nor the specific grid cells/levels appear to have leveled off. It was not until nearly 6000 years that the global ocean potential temperature began to level off. It is important to note that deeper oceans take longer to come into equilibrium.

4.3. Selecting the best template to start building a new world

800 Selecting the most appropriate model configuration for a simulation requires a careful balance of factors that affect both performance and results within the target science questions, but also appropriateness of the configuration for the planet of interest. A number of criteria need to be considered for making the best choice. The planet size can dictate the choice of resolution; for small planets like Mars the medium resolution is adequate, while for super-Earths the fine resolution might be more appropriate. Another factor to consider is whether the chosen resolution properly samples the Rossby radius of deformation, which depends on both the model resolution and the planet spin rate. The generally-assumed notion that a higher resolution should always be preferred might not always be true (depending on the science question), because the computational cost of going from medium to fine is very significant (Sect. 4.5). In the case of exoplanets, for which we may have very few, if any, constraints on their atmosphere, there is often little point in performing fine resolution simulations when the computational time might be better spent sampling over a larger parameter space with the medium resolution instead of gaining little additional scientific insight with a few high-resolution simulations.

810 For atmospheres that closely resemble that of present-day Earth the GISS radiation would suffice, so the simulation can take advantage of faster simulation times, while for very different atmospheres the SOCRATES radiation must be selected. If the planet has an ocean, the choice of a dynamic ocean configuration is always the best, while for a planet that closely resembles modern Earth the use of a prescribed ocean can be justified. For completely unknown planetary configurations the Q-flux ocean is a compromise between performance and oceanic response to climate, but the unknown factor of oceanic heat transport, presented later, can vastly affect (or bias) results. Lastly, the choice of atmospheric composition, both in terms of major and trace gases composition, is the parameter space that most ROCKE-3D users are expected to be routinely modifying. We do provide radiation tables for a wide range of compositions and stellar types (see Data availability section and Appendix C), but for cases outside the presently available tables the users would need to calculate their own. Any choice of planetary parameters (eccentricity, obliquity, etc.) should follow, since they don't affect the choice of the template used to initialize the model.

820 Another thing to consider when starting a new simulation is the spinup time required to bring the model to equilibrium. We provide equilibrated model conditions (restart files) for all simulations presented here (see [Data availability](#) section), or restart files that have been created previously can be used to spin off new simulations. The key thing to ~~consider~~ keep in mind is that the configuration changes between a restart file and a new simulation that the model code allows are limited. It is also important to note that restart files from ~~planet-ROCKE-3D~~ 1.0 are not consistent with those of ~~planet-ROCKE-3D~~ 2.0.

825 ~~planet-ROCKE-3D~~ 2.0.

4.4. Creating a new planet

Although an infinite number of new planet configurations are in theory possible, in practice certain changes can't easily be accommodated. Difficult changes include altering the horizontal resolution or vertical layering, incorporating a dynamic ocean when one was not used in prior simulations (although the reverse, starting a p or q ocean simulation from an o restart file, is

possible), changing the land-ocean mask, topographic relief or ocean bathymetry while a simulation is ongoing, or altering atmospheric composition substantially. Altering the location of even a single land grid cell may require a series of cascading adjustments to many other boundary and initial condition data sets which makes the task very tedious. It is not uncommon when wholesale changes are made to continent and ocean configurations that fine adjustments must be made to deal with sub-grid scale issues related to ocean gateways, vegetation distributions, or continental drainage basins (river flow and lakes). In this section we discuss some of the issues related to generating a “new planet” through the creation of self-consistent boundary and initial condition data sets. We also discuss a variety of pre-existing planetary and ancient Earth model configurations that are available for use with ROCKE-3D. A set of key parameters related with the definition of a new planet are presented in Table 4.

Table 4: Description of parameters that are used to define a planet. Unless explicitly stated (footnote ^c) these parameters are modified via the model configuration “rundeck” file.

<u>Parameter</u>	<u>Description</u>	<u>Units</u>	<u>Earth value</u>
<u>solar_spec (S)^a</u>	<u>Stellar spectrum</u>	<u>N/A</u>	<u>sun</u>
<u>spectral_file_sw</u>	<u>Shortwave spectral file</u>	<u>N/A</u>	<u>ga7^b</u>
<u>spectral_file_lw</u>	<u>Longwave spectral file</u>	<u>N/A</u>	<u>ga7^b</u>
<u>quantizeyearlength</u>	<u>Adjusts orbital period to be an integral multiple of the rotation period</u>	<u>N/A (true or false)</u>	<u>False^c</u>
<u>planet_s0 (S)</u>	<u>Solar constant</u>	<u>W m⁻²</u>	<u>1360.67^c</u>
<u>S0X</u>	<u>Solar constant multiplier relative to Earth</u>	<u>Dimensionless</u>	<u>1^c</u>
<u>siderealorbitalperiod</u>	<u>Length of orbital period</u>	<u>s</u>	<u>31,536,000^c</u>
<u>siderealrotationperiod</u>	<u>Length of rotation period</u>	<u>s</u>	<u>86,400^c</u>
<u>eccentricity</u>	<u>Orbital eccentricity</u>	<u>Dimensionless</u>	<u>0.0167^c</u>
<u>obliquity</u>	<u>Obliquity of axis</u>	<u>Degrees</u>	<u>23.44^c</u>
<u>longitudeAtPeriapsis</u>	<u>Angle of perihelion relative to vernal equinox^d</u>	<u>Degrees</u>	<u>282.9^c</u>
<u>hourAngleOffset</u>	<u>Degrees east from international dateline</u>	<u>Degrees</u>	<u>0^c</u>
<u>geotherm_landx</u> <u>geotherm_ocnx</u>	<u>Land and ocean, respectively, geothermal heat flux multiplier. Acts on GEOTHERM_LAND or GEOTHERM_OCN input files.</u>	<u>Dimensionless</u>	<u>0 if no input file is used, 1 if an input file is provided</u>
<u>CO2X, CH4X, N2OX, O3X, CFC11X, CFC12X, O2X, NO2X,N2CX</u>	<u>Gas multipliers for radiation (multiplies values given in GHG input file)</u>	<u>Dimensionless</u>	<u>1^c</u>
<u>grav^e</u>	<u>Acceleration of gravity</u>	<u>m s⁻²</u>	<u>9.80665</u>

<u>Parameter</u>	<u>Description</u>	<u>Units</u>	<u>Earth value</u>
<u>radius^e</u>	<u>Planet radius</u>	<u>m</u>	<u>6,371,000</u>
<u>mair^e</u>	<u>Molar mass of dry air</u>	<u>g mol⁻¹</u>	<u>28.9655</u>
<u>srat^e</u>	<u>Ratio of specific heats at fixed pressure and volume</u>	<u>Dimensionless</u>	<u>1.401</u>
<u>psf^e</u>	<u>Global mean surface pressure</u>	<u>mbar</u>	<u>984</u>
<u>ptop^e</u>	<u>Pressure over which the model uses constant pressure levels, instead of adaptive to the underlying topography (sigma-coordinate)</u>	<u>mbar</u>	<u>150</u>
<u>pN2, pO2, pAr, pH2^e</u>	<u>Fraction of N₂, O₂, Ar, and H₂, respectively, in dry air.</u>	<u>Dimensionless</u>	<u>0.780840, 0.209476, 0.0093, 0, respectively</u>

^a: Applicable only to the SOCRATES (S) or GISS (G) radiation scheme.

^b: Multiple spectral files are permissible for Earth. sp_sw_ga7/sp_sw_ga7_dsa and sp_lw_ga7/sp_lw_ga7_dsa are used for the Earth runs presented in this paper, but these may be changed to allow for greater flexibility.

^c: These default values do not need to be set explicitly for Earth runs.

^d: This is sometimes called the “longitude of perigee” (see e.g., Appendix B, Berger et al., 1993) that is 180 degrees different from what they call the longitude of perihelion. In the GISS model, modern Earth’s value is 282.9 degrees.

^e: Defined in the source code file shared/PlanetParams_mod.F90, rather than in the rundeck.

4.4.1. Ancient Earths

A variety of boundary condition data sets (Fig. 9) that are based on paleogeographic reconstructions are available for use with ROCKE-3D. These ancient Earth boundary conditions represent various time periods throughout Earth’s history. The specific configurations are not uniformly distributed in geologic time but were chosen to supply testable scenarios for either significant climatic events or because they represent noteworthy continent-ocean distributions that have potentially compelling impacts on the global or regional climate characteristics of a planet. The available boundary conditions (Fig. 9) include: 1) The Sturtian (720 Ma), an equatorial supercontinent that is representative of the Neoproterozoic Earth and has primarily been used with Snowball Earth studies and examination of Earth’s Cryogenian interval; 2) The Late Ordovician (450 Ma), with continents primarily located in the Southern Hemisphere, including a significant-sized land mass covering the South Pole, in contrast to the Northern Hemisphere, which is dominated by open ocean; 3) Early Jurassic (180 Ma), a time period where the Pangean supercontinent, the single supercontinent (and, consequently, a single large ocean as well) was roughly symmetrical about the equator and nearly extended from pole to pole; 4) The Mid-Cretaceous (100 Ma), a period of long-lived extreme warmth, among the warmest intervals of the past billion years, and perhaps the closest Earth has gotten to a super-greenhouse climate since multi-cellular life forms evolved, during which the Pangean supercontinent had broken up, with an early Atlantic Ocean

allowing a flow of water across the equator; 5) The Cretaceous-Paleogene Boundary (66 Ma), which was the time corresponding to the Chicxulub asteroid impact event, thought to have contributed to the extinction of dinosaurs ; 6) The mid-Pliocene (3.2 Ma), during which continents were already approximately in their modern geographic locations, however ice sheets were greatly reduced and consequently sea level was higher, altering global coastlines. This time is thought to be the most recent period in Earth’s history with global temperatures as warm as models predict for Earth’s future climate maximum as impacted by greenhouse gas increases. For much older time periods, where the preserved geologic record is too limited to reconstruct paleogeographic records that could be used to generate realistic boundary condition data sets, we generally use either the Sturtian supercontinent boundary conditions (described above), or an idealized continental configuration (see next section), or simply run the model as a water world (aquaplanet) or land-only world.




Figure 9. Ancient Earth boundary conditions currently available. Top row, left to right: Sturtian (Neoproterozoic Snowball Earth [720 Ma] and general Precambrian equatorial supercontinent configuration); Late Ordovician, 450 Ma; Early Jurassic, 180 Ma (Pangean supercontinent). Bottom row, left to right: Mid-Cretaceous, 100 Ma; Cretaceous-Paleogene boundary, 66 Ma; Mid-Pliocene, 3 Ma.

With the exception of the mid-Pliocene reconstruction, which includes reconstructed bathymetry as outlined by Haywood et al. (2016), other maps of ancient Earth have a uniform ocean depth. This uniform depth is a simplification given that the details of ocean bathymetry are poorly known, if at all, further back in Earth’s history. A reasonable depth for ancient Earth simulations is 3800 meters; for “Earth-like” exoplanet simulations using these land distributions and a dynamic ocean, in the absence of any better constraint, we frequently use a depth of 1000 meters.

4.4.2. Venus

All current versions of ROCKE-3D are unable to model modern day Venus because of its extreme surface temperatures (~750K) and pressures (~92 bar), but a number of studies of a hypothetically ancient habitable Venus have been published (Way et al., 2016; Way and Del Genio, 2020)(Way et al. 2016; Way & Del Genio 2020). In these studies five different sets of

boundary conditions were utilized. Three of them involved the use of modern-day Venusian topography (can be found in Fig. 5 of Way and Del Genio, 2020); a land planet with only 20 cm of water incorporated within the soil (termed Arid-Venus), a planet with 10 m water global equivalent layer (GEL) called 10m-Venus with the water dispersed into the lowest topographic regions as dynamically active lakes (their existence/size determined by a competition between evaporation and precipitation during the model simulation), and a 310 m-Venus with 310 m GEL again dispersed in the lowest topographic regions as oceans and lakes where the oceans are permanent features and the lakes again are dynamic. Two other cases were chosen as they are commonly used in the exoplanet literature: an aquaplanet scenario (with the only land being at the south pole point and set to zero topographic height) and a modern Earth land-sea mask, but only using a 310 m deep ocean to make it comparable to the 310 m-Venus scenario mentioned above. The reader would be right to ask why one uses modern Venusian topography when any ancient topography was likely completely different? First, it is an interesting contrast to modern Earth which has a plethora of land at high latitudes whereas the 310 m-Venus has a lot of land also at low-latitudes, and in contrast to the aquaplanet scenario. Finally, with presently available data there is no way to reconstruct what ancient Venus' topography was like as ~80% of planet has been resurfaced in the past ~300-750 million years (Bougher et al., 1997; Strom et al., 1994). All boundary condition files for the cases described above are located in our on-line repository (see Data availability). In an attempt to compare some of the previously published [planet-ROCKE-3D](#) 1.0 ancient Venus work with [planet-ROCKE-3D](#) 2.0 we ran a number of [planet-ROCKE-3D](#) 2.0 simulations with the exact same setup. Generally for the dryer simulations (arid-Venus and 10m-Venus) the results were very similar (see Sect. 4.4.4 for Proxima Centauri b). For the simulations with oceans this was not always the case. For example, for the 310 m-Venus case at 2.9Ga [planet-ROCKE-3D](#) 2.0 had a mean temperature of 3.4 °C while the published [planet-ROCKE-3D](#) 1.0 results (Way & Del Genio 2020) was 9.6 °C. This difference of 6 °C is likely due to the very different cloud scheme differences between ModelE2.1 ([planet-ROCKE-3D](#) 1.0) and ModelE2.2 ([planet-ROCKE-3D](#) 2.0) as demonstrated in Fig. 10, where the planetary albedo and net cloud radiative ~~forings-effect~~ are quite distinct.




Figure 10. Comparison of [planet-ROCKE-3D 1.0](#) and [planet-ROCKE-3D 2.0](#) simulations of an ancient Venus climate from simulation #25 in Way and Del Genio (2020). The simulation is of a 310 m deep global equivalent layer ocean that includes a number of low-latitude continents (Fig. 5 in Way and Del Genio, 2020). X-axis: number of Venus orbits completed. Y-axis: Net Rad: Net radiative balance at the top of the atmosphere (*net_rad_planet*); T_{surf} : global mean surface temperature (*tsurf*); Albedo=global mean planetary albedo (*plan_alb*), Cloud Rad = net cloud radiative [forcing effect](#) (shortwave+longwave).

4.4.3. Mars and Moon

Mars has been modeled with ROCKE-3D at several epochs in its history, but thus far only ancient Mars simulations have been published. The simulations were aimed at the Noachian (Guzewich et al., 2021) and the Hesperian (Schmidt et al., 2022). In the Noachian case modern Mars topography was used (Guzewich et al., 2021), but a subset of simulations used a topography with an inferred map before the Tharsis region was formed, and its associated true polar wander (Bouley et al., 2016). In Schmidt et al. (2022) only modern Mars topography was considered. Guzewich et al. (2021) utilized varying amounts of GEL

920 water, whereas Schmidt et al. (2022) only considered a northern polar ocean (~10 % of the surface area of the planet) and the Hellas Basin to have water.

As part of its thin atmosphere capability (see Sect. 2.2.3), ROCKE-3D is currently capable of simulating modern Mars with a pure CO₂ atmosphere (~6 mb_{ar} at surface) and fully coupled CO₂, H₂O and dust cycles. For the water cycle it uses a 3-layer snow model developed for the parent Earth model, while the CO₂ condensation is currently handled by a bucket model which
925 keeps track of the condensate amount and its areal cover fraction. The cover fractions of the H₂O and the CO₂ snow are combined together (the greater absorbs the smaller) for the computation of ground albedo. No distinction is currently made between CO₂ and H₂O snow in the albedo algorithm.

Soil dust aerosols have been incorporated into the Mars version of ROCKE-3D as radiatively active tracers by successfully utilizing the soil dust module that was developed for NASA GISS ModelE for Earth's atmospheric conditions, with only minor
930 modifications ~~(Perlwitz et al., in preparation)~~. It is a sectional scheme with five size bins that simulates the emission, atmospheric transport, and deposition of dust. Different to the original scheme in ModelE, dust emission is explicitly dependent on atmospheric surface density (as a function both of pressure and temperature at the surface) to account for the large density variations at the surface of Mars, affecting the buoyancy of dust particles.

Following a hypothesis of [Needham and Kring](#) (2017) that in the past the Moon could have had a thin transient atmosphere
935 due to volcanic outgassing, we used ROCKE-3D to study such an atmosphere (Aleinov et al., 2019). In our experiments, we used modern lunar topography and albedo, and conditions at ~3.5 Ga (peak of lunar volcanic activity) for orbital parameters and insolation. A special algorithm was added to treat Permanently Shadowed Regions (PSRs) as patches of land which receive no shortwave radiation. We investigated a parameter space of surface pressures 1-10 mb_{ar} and compositions which included pure CO and pure CO₂ atmospheres, either completely dry or with 5x10⁻³ kg/kg water. The results have been presented in the
940 past (Aleinov et al., 2019) and the configuration we provide can be used for similar model setups.

4.4.4. Known extrasolar terrestrial planets

The ROCKE-3D model has been used in a number of studies involving extrasolar planets and their theoretical atmospheres. For comparison purposes we examined results from Proxima Centauri b as an aquaplanet with a fully coupled ocean with
945 ~~planet-ROCKE-3D~~ 1.0 and ~~planet-ROCKE-3D~~ 2.0. Here, ~~is~~ it is important to note that there was an update to the sea ice parameterization from ~~planet-ROCKE-3D~~ 1.0 to 2.0 as described in Sect. 2.1 above. Since the Proxima Centauri b control run from Del Genio et al. (2019b) is particularly cold, in order to make the comparison more realistic we did an additional control run with the ~~planet-ROCKE-3D~~ 2.0 sea ice fix in ~~planet-ROCKE-3D~~ 1.0. This can be seen in Fig. 11 below. As in the ancient Venus case discussed in Sect. 4.4.2 the clouds appear to play a key role in making the climate ~8 °C colder in the
950 ~~planet-ROCKE-3D~~ 2.0 case versus in ~~planet-ROCKE-3D~~ 1.0 with the sea ice fix (P1+SF). Without the sea ice fix the difference is nearly 10.5 °C. This difference in the clouds manifests itself markedly in the amount of the ocean that is covered

in ice as seen in Fig. 11. An additional 4% of ice accumulates in `planet-ROCKE-3D` 1.0 with the sea ice fix, and another 13% on top of that with `planet-ROCKE-3D` 2.0.




Figure 11. Same as Fig. 10 where P1=`planet-ROCKE-3D` 1.0 and P2=`planet-ROCKE-3D` 2.0. P1+SF is the `planet-ROCKE-3D` 1.0 version with the sea ice fix mentioned in Sect. 2.1. The bottom row is the fraction of the ocean covered in ice. Since this is an aquaplanet setup, that equals to the planet surface covered in ice.

ROCKE-3D participated in the THAI model intercomparison (Fauchez et al., 2020) that studied the planet Trappist-1e with the same boundary condition files and atmospheric compositions across four models (Fauchez et al., 2022; Sergeev et al., 2022; Turbet et al., 2022). Here we look at differences between `planet-ROCKE-3D` 1.0 (used in the THAI intercomparison) for the dry case (Ben1) versus `planet-ROCKE-3D` 2.0 (Fig. 12). In this setup there is no moisture in the atmosphere or surface, hence there are no clouds. The planetary albedo is almost exactly the same, yet the surface temperatures still differ by $\sim 3^\circ\text{C}$.

This gives us a lower limit for the differences we can expect between the same setup on [planet-ROCKE-3D](#) 1.0 versus 2.0. In general, similar differences were seen in the dry cases in the THAI intercomparison (Turbet et al., 2022), and in related studies (e.g. Sergeev et al., 2024).




Figure 12. Same as Fig. 10 for the planet Trappist-1e with a dry atmospheric setup.

Idealized land-ocean configurations

Some studies focused on early Earth or terrestrial exoplanets will have experiment designs that require some representation of ocean and emergent land area but are not yet at a stage that would benefit from the complexities of the realistic ancient Earth boundary conditions described in Sect. 4.4.1. It may also be the case that the percentage of land area required is notably greater

or less than the land coverage offered by those ancient Earth configurations (which range between 20 to 30% land cover). To fill this need, we have created sets of boundary condition files for 13 idealized continent configurations, in which land is represented at increasingly smaller fractions of the total global area (Fig. 13). Configurations with $\geq 25\%$ land cover are available in three orientations: pole-to-pole meridional, polar cap, and equatorial land belt. The smallest global land cover options, 12.5% and 6.25%, are each available as two- or four-continent distributions to emulate the presence of microcontinents. All 13 configurations have a uniform land elevation of 50 meters, a ground albedo that is calculated by using a 50 % dark and 50 % bright soil mix, and include simple “bathtub” oceans, in which the sea floor slopes steeply downward from the continental edges to a uniform ocean bottom. The ~~with a~~ maximum depth is configurable, and is typically set between 1000 and of nearly 3800 meters for simulations with a dynamic ocean. Together with the two land planet and aquaplanet configurations, we cover a spectrum of land cover possibilities.




Figure 13. Idealized continent configurations.

Aquaplanets

In addition to the Earth-centric configurations described here, we also provide four aquaplanet ones (GISS or SOCRATES radiation scheme, with dynamic or Q-flux=0 oceans). Aquaplanets are commonly used configurations for studies in climate dynamics or for planets with unknown surface boundary conditions, as is the case for exoplanets.

The aquaplanets with a dynamic ocean are substantially warmer than the runs with Earth’s topography (20.9 and 22.2 °C with SOCRATES or GISS radiation, respectively). They also feature a substantially reduced pole-to-equator temperature gradient relative to Earth runs (Fig. 14), and a stronger greenhouse effect owing to the greater abundance of atmospheric water vapor.




Figure 14. Zonal-mean temperature (°C) for aquaplanet simulations with SOCRATES (solid) or GISS (dashed) radiation scheme, and a dynamic ocean (red) or Q-flux=0 ocean (blue) configuration. Two simulations (varying CO₂ amount) are annotated for the SOCRATES Q-flux=0 configuration, which exhibits a large change between 1xCO₂ and 2xCO₂.

The Q-flux=0 aquaplanet runs are colder than their dynamic ocean counterparts, and differ markedly between the GISS and SOCRATES radiation schemes (global mean temperature of 13.3 and −44.5 °C, respectively). This is due to a snowball (ice-albedo) instability that occurs for the SOCRATES but not the GISS run. Although this discrepancy is obviously large, we note that previous studies have shown that the coupling between sea ice and the convergence of ocean heat transport patterns is extraordinarily strong in aquaplanet simulations (e.g. Rose, 2015) and work in climate dynamics using aquaplanets often disable sea ice formation altogether due to the strong confounding influence of ocean heat transport on the stability of the sea ice edge (Rencurrel and Rose, 2018; e.g. Voigt et al., 2016). While sea ice is sometimes disabled in some aquaplanet studies to isolate specific dynamical processes, its inclusion is essential for fully coupled exoplanet simulations. Sea ice plays a critical role in setting the climate state, particularly in transitions between ice-free, waterbelt, and snowball regimes and tipping points.

and strongly influences the planet's energy balance and climate stability (e.g. Brunetti et al., 2019; Hörner and Voigt, 2024).
For the realistic exploration of exoplanet climates, especially near habitability boundaries, sea ice must be actively included.

For the Q-flux=0 aquaplanets, our simulations are very close to an instability threshold, such that simply changing the radiation scheme results in a large bifurcation despite only marginal differences in all other runs reported here. In fact, if the Q-flux=0 aquaplanet simulation using SOCRATES is instead run (from initially warm conditions) with a doubling of atmospheric CO₂, it does not transition into a snowball and is warmer than the GISS counterpart simulation with 1xCO₂. The GISS radiation Q-flux=0 aquaplanet run also collapses into a near-snowball state with 0.5xCO₂, indicating that these simulations are quite sensitive near their current state. Such a CO₂ forcing is a large perturbation in the context of contemporary climate change on Earth, but is small by planetary climate standards. This is consistent with the studies discussed above that aquaplanets are remarkably prone to large snowball instabilities and the Q-flux (or Q-flux=0) configuration, in particular, should be used with caution for climates in which sea ice is expected to form.

Deep ocean worlds

~~In order to explore a wider range of planets and moons with deep oceans (on the order of tens to hundreds of kilometers deep), an upcoming update to ROCKE 3D 2.0 will extend the ocean component from a Hydrostatic (H) model to a Quasi Hydrostatic (QH) model. With an increasing number of planets presumed to have surface oceans or icy moons with subsurface oceans (e.g. Europa, Enceladus) some assumptions made about the H momentum equations for planets with relatively shallow oceans like Earth ($h/R \sim 10^{-4}$, where h is the ocean depth and R is the radius of the planet) need to be revisited to take into account planetary bodies with 1) ocean depths reaching tens of kilometers, and 2) cases where ocean depths are not negligible anymore when compared with the planet or moon radius (Marshall et al., 1997). Hydrostatic Primitive Equations (HPEs) were developed as a less GCM computationally expensive form of the full Non Hydrostatic (NH) momentum equations, by considering Earth's oceans in a hydrostatic balance where the pressure field is balanced by the gravity force. In cases where h/R is not small anymore (e.g. $h/R \sim 10^{-1}$ for Enceladus) the QH setup is considered as being more realistic for deep oceans than the H equations, while not being as computationally expensive as the NH ones. This is achieved by re-establishing some of the Coriolis and metric terms neglected in the momentum equations to get a full treatment of the Coriolis force and relaxing the shallow water approximation in the momentum equations. As an example, in spherical coordinates where 'r' was constant and equal to R, the vertical position of a parcel of water 'r' is now a variable). When this update is finalized, it will be made available as an easy to implement update into ROCKE 3D 2.0.~~

4.5. Model performance

Model Performance was discussed in the planet-ROCKE-3D 1.0 paper (Way et al., 2017), but since its importance should not be underestimated, we further discuss it here. Each model component described here has its own computational requirements.

The exact wall-time for a simulation to complete does change depending on computer load (e.g. input/output contention) and hardware capabilities (e.g. CPU generation, memory availability), and can fluctuate even from one simulation to the other on the same computer. For this, the values presented in Fig. 15 should only be used in a qualitative manner, rather than as exact representations of the whole duration of a simulation on any computer at any given time. Even with these approximate numbers, it is clear that SOCRATES radiation is more computationally expensive than GISS, and the same applies for the fine resolution over the medium one. The prescribed and Q-flux ocean runs are practically indistinguishable when it comes to computation time, while the dynamic ocean is more computationally demanding. This difference amplifies with resolution: while moving from medium to fine resolution roughly increases the simulation time by a factor of 2-3, a direct result of the quadrupling of the atmospheric resolution, when the dynamic ocean is included the computational time increases by about an order of magnitude. This is an additive effect of the 4x atmospheric increase in resolution and the ~40x time-oceanic increase (Table 1). Interestingly, the performance penalty added by the presence of a dynamic ocean in the medium resolution is overwhelmed by the slowdown due to SOCRATES, while for the fine resolution it is more pronounced. Further, the A atmosphere when using SOCRATES has a clear slowdown by about 50%, probably linked with the presence of O₃ and aerosol calculations that significantly add to the radiative transfer performance.




Figure 15. Approximate wall-time required per model configuration. The different atmospheres, oceans, and radiation schemes are listed at the top of the plot, while the two resolutions are separated by color (blue: medium; red: fine).

4.6. Further modifications of the model configuration

1055 There are too many parameters that one can alter to modify the behavior of the model to list here. These involve either sanity-
checking limits, expansion of capabilities, or fixes of known issues, as described in Sect. 4.6.1. In addition, during and after
the end of the simulations presented here, model development continued. That led to some additional model development and
bug fixes (Sect. 4.6.2) that are now available in [planet-ROCKE-3D 2.0](#), but were not available when the simulations presented
here started. All the parameters below can be defined in the model “rundecks”, the configuration files that completely describe
1060 each simulation.

4.6.1. Parameters that can be modified

Beyond what has been presented thus far, there are several more ways one can modify how the model runs. These additional
options can change physics (thus results), can significantly affect model performance, and/or can change what diagnostic
output is produced. A comprehensive list and explanation of all options is far outside the scope of this work, so we only list
1065 here the most important, or most influential, of the parameters that ROCKE-3D users should be aware of. The key ones are
described below, and many more are listed in [Table 5Table 4](#).

The main parameter which defines how the model is run is its physics time step, DT_{src} . It tells the model how often its main
parts (atmosphere, ocean, land, radiation) should exchange fluxes. Once set up at the start of a simulation, it cannot be changed.
Other time stepping parameters can be changed though, and we use this functionality if the model exhibits numerical
1070 instabilities during the course of a simulation. In particular, one can decrease DT or increase N_{lsurf} (typically, by powers of
2) to improve stability. The choice of DT depends on many factors, but in general smaller planets and planets with thinner
atmosphere need smaller DT . For example, for most of our modern Mars runs, we use $DT=50$ s.

A key parameter related with both model results and model performance is NRAD, which stands for “every how many time
steps radiation is called”. Ideally this should equal to 1 (every time step) but radiation is the most computationally expensive
1075 physics part of the model, so traditionally we use the value of 5. The choice of this value is not random: since the model is
based on an Earth model, where the planet has a rotation period of 24 hours, the default 30-minute physics time step means
that there are 48 time steps per day. The value of NRAD=5 is the smallest non-unity value that does not divide 48 exactly,
ensuring a non-uniform (thus, non-biased) sampling over the course of multiple days. In other words, over e.g. a month, every
30 minutes of the day are sampled, thus getting a non-biased average. For planetary applications, care must be taken for the
1080 choice of this variable: the number of time steps per day, which depends on the model time step and the rotation period, should
never be exactly divided by NRAD, if NRAD>1. Very large values of NRAD though will add a sampling bias, so the right
balance between performance and accuracy needs to be maintained. Even beyond planetary rotation periods, for very fast
rotators (e.g. orbital periods of very few days) one might not be able to use daily or “monthly” means if the choice of NRAD
is such that radiation is called very few times over the averaging period.

1085

Table 54: Description of parameters that modify the model's behavior.

Parameter	Definition	Default value	Additional information
<i>cloud_rad_forc</i>	Enables cloud radiative forcing diagnostics	0 (do not simulate)	Results in additional radiation calls, which may have a non-negligible performance penalty.
<i>DT</i>	Timestep for atmospheric dynamics	M resolution: 450 s F resolution: 225 s	The ratio between <i>DTsrc</i> and <i>DT</i> should be an even number (enforced by the model automatically if different).
<i>DTO</i>	Timestep for ocean barotropic dynamics	M resolution: 450 s F resolution: 112.5 s	Recalculated in the model so that the ratio between the model physics timestep <i>DTsrc</i> and <i>DTO</i> is an even number. Advection and mesoscale use longer steps, which for most use cases don't have to be defined explicitly.
<i>DTsrc</i>	Timestep for model <u>physics exchange fluxes across model components</u>	1800 s	Should divide the length of a calendar day exactly. Cannot be changed during the run.
<i>lake_ice_max</i>	Inhibits lake ice to grow indefinitely via snow accumulation	0 (no limit)	Excess ice is placed in GLMELT arrays.
<i>maxclow</i>	Pressure at top of low clouds	680 hPa	Only used for diagnostics, and scales automatically with surface pressure, if other than 1 bar and undefined. It serves as the boundary between <i>pcldl</i> and <i>pcldm</i> in the output.
<i>maxcmid</i>	Pressure at top of middle clouds	440 hPa	Only used for diagnostics, and scales automatically with surface pressure, if other than 1 bar and undefined. It serves as the boundary between <i>pcldm</i> and <i>pcldh</i> .
<i>maxctop</i>	Pressure above which no clouds are allowed	50 hPa for Earth, 0 hPa for other planets	Contrary to <i>maxclow</i> and <i>maxcmid</i> , altering <i>maxctop</i> can affect <u>prognostic</u> model results.

Parameter	Definition	Default value	Additional information
<i>meso_diffusivity_const</i>	Mesoscale ocean diffusivity constant	No default, must be explicitly defined if used.	Frequently 1200 m ² s ⁻¹ is used. It should be considered a tuning factor.
<i>minGroundTemperature</i>	Coldest allowed ground temperature	-150 °C	
<i>NIsurf</i>	Number of substeps of surface flux interactions per physics time step	2	This is specifically relevant for surface turbulent fluxes, such as latent and sensible heat, and momentum drag.
NRAD	Call radiation every NRAD time steps	5	The ratio of (time steps per day) over NRAD should never be an integer, except when NRAD=1.
rtau	Rayleigh friction timescale	No default, not used if undefined.	Enforces linear (“Rayleigh friction”) gravity wave drag, if defined.
use_VMP	Use virtual mixed-phase clouds	Enabled	
wsn_max	Maximum snow depth allowed	2 m water equivalent	Remove the snow in excess of wsn_max and dump its water into GLMELT file. Can only be used if run with an ocean. It should be at least as large as the total snow deposition over a season.

4.6.2. Other updates and future developments

After the completion of the simulations presented in this paper, issues in the calculation of the ocean momentum diffusion terms were discovered in planet-ROCKE-3D 2.0 and were fixed. These include 1) some quantities used in the tridiagonal solver for the ocean velocities that were not defined properly, and 2) the hemispheric symmetry was re-established in the calculation of some intermediary arrays. The impact of these bug fixes was evaluated in the framework of ModelE2.1 at preindustrial (year 1850) conditions and were found to have a minimal impact on the global climatology. They are nonetheless necessary to have full ocean dynamic hemispheric symmetry for aquaplanets runs, and are now enabled by default in ROCKE-3D without user intervention.

Deep ocean worlds

In order to explore a wider range of planets and moons with deep oceans (on the order of tens to hundreds of kilometers deep), an upcoming update to ROCKE-3D 2.0 will extend the ocean component from a Hydrostatic (H) model to a Quasi-Hydrostatic (QH) model. With an increasing number of planets presumed to have surface oceans or icy moons with subsurface oceans (e.g. Europa, Enceladus) some assumptions made about the H momentum equations for planets with relatively shallow oceans like Earth ($h/R \sim 10^{-4}$, where h is the ocean depth and R is the radius of the planet) need to be revisited to take into account planetary bodies with 1) ocean depths reaching tens of kilometers, and 2) cases where ocean depths are not negligible anymore when compared with the planet or moon radius (Marshall et al., 1997). Hydrostatic Primitive Equations (HPEs) were developed as a less GCM-computationally-expensive form of the full Non-Hydrostatic (NH) momentum equations, by considering Earth's oceans in a hydrostatic balance where the pressure field is balanced by the gravity force. In cases where h/R is not small anymore (e.g. $\frac{h}{R} \sim 10^{-1}$ for Enceladus) the QH setup is considered as being more realistic for deep oceans than the H equations, while not being as computationally expensive as the NH ones. This is achieved by re-establishing some of the Coriolis and metric terms neglected in the momentum equations to get a full treatment of the Coriolis force and relaxing the shallow water approximation in the momentum equations. As an example, in spherical coordinates where 'r' was constant and equal to R , the vertical position of a parcel of water 'r' is now a variable. When this update is finalized, it will be made available as an easy-to-implement update into ROCKE-3D 2.0.


5. Discussion

The effect of the different model configurations presented in Table 1 on the modeled climatology are presented below, together with other configurations that are available in ~~planet-ROCKE-3D~~ 2.0 but are not based on Earth. We present the latter ones as means to compare the results of ~~planet-ROCKE-3D~~ 1.0 simulations using ~~planet-ROCKE-3D~~ 2.0. We expect that the most popular configuration will include the SOCRATES radiation (S), using an anoxic atmosphere (N), a dynamic ocean (o), and the medium resolution of the model (M40), so P2SNoM40. When studying differences across configurations below, we'll use that simulation as the point of reference, to the extent possible.

5.1. Quick view

The results across simulations present large differences that show the impact of configuration choices to the overall climate of the atmosphere in question. We will first take a quick look at the global mean results (Fig. 16), before diving deeper into regional changes in subsequent sections. The first thing one notices when comparing Fig. 5 with Fig. 16 is how much more colorful the latter is, implying that the simulated differences across planetary configurations are much more impactful to climate variables than the changes we did towards the generalization of the Earth configuration. In other words, the climatological differences we get when simulating Earth with and without any Earth-centric tunings is much smaller than the

140 differences seen when simulating different planets. This provides an additional assurance that the generalization procedure does not negatively impact model skill.



145 **Figure 16.** Same as in Fig. 5 for the template configurations described in [Table 1](#)Table 1. Note the addition of a legend over the plot, which makes understanding the differences across columns easier. Also note that contrary to Fig. 5, all simulations presented here are in radiative balance.

Starting from comparing the old (Way et al., 2017) against the new (this work) ROCKE-3D versions, the biggest changes are found in the high clouds fraction (*pcldh*), which drops from 58% in [planet-ROCKE-3D 1.0](#) to 29% in [planet-ROCKE-3D 2.0](#), when averaging across all simulations, a value much closer to the value expected for present-day Earth. Together with some modest middle and low level cloudiness, the mean total cloud fraction (*pcldt*) for all simulations decreased from 75% in [planet-ROCKE-3D 1.0](#) to 59% in [planet-ROCKE-3D 2.0](#). The cloud-top pressure in [planet-ROCKE-3D 1.0](#) is lower than [planet-ROCKE-3D 2.0](#), which means that clouds were able to reach higher altitudes in the previous version of ROCKE-3D.


When looking at the [planet-ROCKE-3D 2.0](#) results and comparing them with P2SNoM40, two major patterns appear. One is related with clouds and radiation: when averaging out over simulations that use the GISS radiation, low cloud fraction ([pcldl](#)) is 44%, compared to when averaging over SOCRATES simulations where it is 32%. This is an expected difference that results from the *U00b* cloud balancing values we used (Table 3), which are higher in all GISS simulations: larger *U00b* values reduce the critical relative humidity for cloud formation, making it easier to form clouds in grid cells where the *U00b* scheme is used. Another difference is that with the GISS radiation the tropopause temperature ([ttrop](#)) and pressure ([ptrop](#)) are both consistently lower than the reference simulation (P2SNoM40), while in the simulations using SOCRATES radiation there is no consistent pattern, with both positive and negative changes. [ROCKE-3D uses the definition by the World Meteorological Organization \(WMO\) to calculate the tropopause layer, as the lowest level at which the temperature lapse rate decreases to 2 K km⁻¹ or less.](#) Another major pattern that appears in Fig. 16 is the large changes calculated by the Q-flux=0 ocean simulations. As will be detailed in Sect. 5.4, the absence of heat transport via the ocean makes the ocean much colder, resulting in colder *sst*, increased ocean ice fraction (*oicefr*) and ground albedo (*grnd_alb*), as well as increases in all snow metrics. Surface air temperature (*tsurf*) also drops when using the Q-flux=0 ocean configuration, which results in less water vapor at surface (*qsurf*) and in the column (*qatm*). This is a robust response regardless of the choice of radiation scheme, atmospheric composition, and resolution, although the magnitude of the effect does change across simulations.

5.2. GISS vs. SOCRATES radiation schemes

ModelE uses the GISS radiation scheme by default; with everything else held constant but switching radiation from GISS to SOCRATES (ga7_dsa spectral file), the model calculates a net planetary radiation imbalance of about -1.3 W m⁻² when prescribed *sst* is used for the A atmosphere. An imbalance is expected for such simulations, where physics changes happen in the atmosphere but the ocean is not allowed to respond. The balanced SOCRATES simulation with a dynamic ocean produces small differences in the global mean climatology of surface air temperature (Fig. 17) but has some significant regional differences (Fig. 18). Interestingly, the GISS radiation simulates cooler temperatures at high latitudes compared to SOCRATES when the A atmosphere is used, with the exception over North Atlantic, where a strong heating is calculated. This heating negatively correlates with less cloudiness, so more sunlight penetrates to the surface, and a complete disappearance of sea ice from the region.

~~(Romanou et al., 2023)The x and N atmospheres on the other hand behave differently when the radiation scheme changes: the Southern hemisphere mostly heats up and the Northern midlatitudes cool, while the North Pole heats up when using the N atmosphere (cools when using x). Sea ice changes again anti-correlate with surface air temperature, while clouds respond in a very similar manner between the x and N atmospheres, with a reduction over the tropical belt and an increase everywhere else, and in particular over the upwelling midlatitudes in both hemispheres. This shows the different response of GISS radiation with regard to aerosols and stratospheric O₃, which are the main differences of the A atmosphere with x and N, but also to~~


185 clouds, due to the different cloud balancing values selected (Table 3) which produce more low clouds ($U00b$) and total ($U00a$
and $U00b$) with the GISS radiation choice of values.



1190 **Figure 17.** Same as in Fig. 6 for surface air temperature (t_{surf}).

The behavior observed in the A-atmosphere configuration appears to stem from the absence of an active overturning circulation in the SOCRATES radiation scheme case for the M-resolution, which effectively shuts down the northward transport of heat. In contrast, the simulation using the GISS radiation scheme maintains a robust Atlantic Meridional Overturning Circulation (AMOC) of approximately 23 Sv, accounting for the warmer temperatures over the North Atlantic and the associated reduction in sea ice area relative to the SOCRATES run. While the radiation scheme does not directly control ocean circulation, its influence on surface energy fluxes may indirectly affect AMOC stability. In our model, AMOC strength has been shown to be highly sensitive to freshwater input from ice melt and transport (Romanou et al., 2023), suggesting a possible link. We also note that at the higher F-resolution, both the SOCRATES and GISS simulations exhibit an active AMOC.

The x and N atmospheres on the other hand behave differently when the radiation scheme changes: the Southern hemisphere mostly heats up and the Northern midlatitudes cool, while the North Pole heats up when using the N atmosphere (cools when using x). Sea ice changes again anti-correlate with surface air temperature, while clouds respond in a very similar manner between the x and N atmospheres, with a reduction over the tropical belt and an increase everywhere else, and in particular over the upwelling midlatitudes in both hemispheres. This shows the different response of GISS radiation with regard to aerosols and stratospheric O_3 , which are the main differences of the A atmosphere with x and N, but also to clouds, due to the different cloud balancing values selected (Table 3) which produce more low clouds (*U00b*) and total (*U00a* and *U00b*) with the GISS radiation choice of values. In the x simulations (central column in Fig. 18), the southern hemisphere becomes warmer with the GISS radiation and features increased cloud cover. However, the northern hemisphere becomes colder, also with more clouds. This particular GISS-SOCRATES difference appears to be the superposition of two other GISS-SOCRATES differences: one, the model wants to be cooler at high latitudes with the GISS radiation, excepting the North Atlantic AMOC-related feature, and two, the model wants to be warmer at southern high latitudes when using the GISS radiation (and slightly warmer at northern high latitudes) and when changing the atmospheric composition (and cloud tuning) but holding the ocean circulation fixed. When we combine these two patterns, we get the southern hemisphere warming in Fig. 19 slightly winning out over its cooling in the upper left of Fig. 18, but in the northern hemisphere the Fig. 18 cooling wins over the Fig. 19 warming.







Figure 18. Surface air temperature (top row), total cloud cover (middle row) and ocean ice fraction (bottom row) differences between GISS and SOCRATES radiation simulations for the A (left column), x (middle column), and N atmospheres (right column) with a dynamic ocean. Positive values mean the GISS radiation is warmer, and black dots show areas where the differences are not statistically significant at the 95% confidence.

Focusing on the N atmosphere (Fig. 19), the change in radiation scheme using a dynamic ocean resembles that of the prescribed ocean surprisingly well, with two notable differences: the continental warming is more pronounced and extends to the whole tropical belt, although in a smaller magnitude over tropical oceans, and there is a persistent cooling over the northern mid-latitude oceans. All of those changes are too small, less than 5 degrees everywhere, when compared to the differences between the two Q-flux=0 simulations, in which major temperature differences are calculated between the two radiation schemes. GISS radiation calculates a very wide-spread warming over the polar half of the southern hemisphere, which nearly everywhere exceeds 10 °C, and reaches 18 °C locally. This is consistent with a much less extended sea ice cover in the southern hemisphere when using the GISS radiation, which moves the southern hemisphere ice line about 10 degrees further away from the tropics (Sect. 5.4). The clouds respond in a similar way across ocean configurations when the radiation scheme changes, with the GISS radiation leading to less clouds in the tropics and more in the extratropics. The differences in cloud cover between the

1235


1240

GISS and SOCRATES simulations are generally consistent with the differences in the cloud tuning parameters, at least in the extratropics where there are the most stratiform clouds (the tuning parameters are for stratiform clouds). In particular, the A-to-x changes to those parameters for the GISS radiation are very large compared to typical changes for rebalancing Earth, and go in the direction of making Gx a cloudier world. Interestingly, the SOCRATES radiation did not require such a large change for A-to-x rebalancing, perhaps because its *radiusl_multiplier* was left unchanged, while the A-to-x rebalancing for the G radiation increased *radiusl_multiplier*, which tends to make clouds less opaque and thus requires increases of cloud cover to compensate (decreased *U00a*, increased *U00b*).



1245 **Figure 19.** Surface air temperature (left column), total cloud cover (middle column) and ocean ice fraction (right column) differences between GISS and SOCRATES radiation simulations for the p (top row) and q (bottom row) oceans with the N atmosphere; see Fig. 18 for the o ocean. Positive values mean the GISS radiation is warmer, and black dots show areas where the differences are not statistically significant at the 95% confidence.

1250 The vertical profile of the global mean temperature (Fig. 20) for the different atmospheres used shows the major change in the stratosphere that occur for the x and N atmospheres, a direct result of the absence of stratospheric O₃. The absence of a stratospheric temperature inversion has important implications on several metrics of the model, as presented here and in other sections. Only the dynamic ocean and medium resolution is shown in Fig. 20, but the results discussed here are valid for all ocean configurations for both resolutions. For the preindustrial atmosphere (A), both the GISS and SOCRATES radiations agree very well, while all other atmospheres agree with each other in the troposphere only. Starting from the lower stratosphere, 1255 at around 50 hPa and higher, temperatures start to deviate with altitude, and reach 10 °C at the model top, across radiation configurations. The x and N atmospheres with SOCRATES are virtually identical, which hints that the GISS radiation might consider O₂ and Ar (both absent in the anoxic (N) atmosphere) differently than SOCRATES does. Further, the GISS radiation simulates a very weak inversion that maximizes around 10 hPa, which is not calculated by SOCRATES, with implications for water vapor and high altitude clouds. The planet-ROCKE-3D 1.0 configuration, only available for the x atmosphere and 1260 medium resolution, agrees extremely well with its successor (P2SxoM40) in planet-ROCKE-3D 2.0, except in the region between 100-10 hPa, which shows a much smoother transition in specific humidity, rather than the sharper change of slope with altitude in planet-ROCKE-3D 2.0 at around 50 hPa. ~~This is a direct consequence of the smoother temperature transition that is simulated in the same altitudes.~~






Figure 20. Global mean temperature (*TempL*; left) and specific humidity (*SpHuL*; right) vertical profile using the dynamic ocean configuration for the different atmospheres and radiation schemes. The vertical profile calculated for the SxoM40 configuration from [planet-ROCKE-3D 1.0](#) is also included for comparison.

5.3. Earth's vs. [planet-ROCKE-3D 1.0](#) vs. anoxic atmospheres

The removal of aerosols and O₃ (and H₂O from CH₄ oxidation in the stratosphere to a lesser extent) in the [planet-ROCKE-3D 1.0](#) atmosphere has a very strong impact on the modeled climatology. First and foremost, the vertical temperature profile of both atmospheres without O₃ (x and N) do not create a stratospheric temperature inversion, leading to large changes in the vertical temperature profile (Fig. 20), water vapor amount, and the presence of clouds (Fig. 21). The specific humidity (water vapor concentration) is nearly identical across configurations in the troposphere, but deviates in the stratosphere due to the temperature differences simulated. The A atmosphere maintains a near-constant profile in the stratosphere with concentrations of around 2 ppmm (parts per million by mass; mg H₂O per kg of dry air), while the x and N atmospheres have 2 to 3 orders of magnitude lower concentrations. This is directly explained by the much colder temperatures in [the x and N atmospheres where the stratosphere-at the altitudes where the stratosphere exists used to be](#) in the A atmosphere, which makes it unable to hold a lot of water vapor, despite the absence of the cold trap in the tropopause. In [planet-ROCKE-3D 2.0](#) the transition from a

decreasing concentration with altitude in the lower atmosphere transitions very quickly to a near-constant profile higher up, while in ~~planet-ROCKE-3D~~ 1.0 there was a much more gradual decrease, driven by the temperature profile (Fig. 20).

The vertical distribution of clouds also changes depending on the configuration. In ~~planet-ROCKE-3D~~ 1.0, liquid water clouds are virtually absent above 400 hPa (Fig. 21), while in ~~planet-ROCKE-3D~~ 2.0 there is a persistent liquid clouds presence throughout the column, until the temperatures become too cold and water clouds ~~abruptly~~ give ~~abruptly~~ way to ice. This happens because in ~~planet-ROCKE-3D~~ 2.0 it is possible to form supercooled liquid clouds via a new “virtual” mixed-phase (VMP) clouds parameterization (Kelley et al., 2020), which was not present in ~~planet-ROCKE-3D~~ 1.0, and significantly increases the amount of supercooled water cloud in the Southern Ocean and the Arctic. This parameterization is enabled by default in ~~planet-ROCKE-3D~~ 2.0, but it is optional and can be disabled (Appendix B), although rebalancing of the model would be required in that case. Near the surface, the radiation scheme choice appears to impact low clouds, with the simulations using the GISS radiation calculating more clouds than those with SOCRATES, but this is due to different model balancing choices across radiation configurations (Table 3), not due to radiation scheme itself. Ice clouds are very similar between the two ROCKE-3D model versions (Fig. 21).




Figure 21. Global mean concentration [parts per million by mass] of water clouds (left), ice clouds (middle), and total cloud fraction (right) vertical profiles using the dynamic ocean configuration for the different atmospheres and radiation schemes. The vertical profile calculated for the SxoM40 configuration from ~~planet-ROCKE-3D~~ 1.0 is also included for comparison.

The simulations with the prescribed ocean have the smallest changes in surface air temperature when other model configurations change, in particular over the ocean (see, e.g., [top-left panel of Fig. 19 for the p ocean vs. the bottom-left panel of the same figure for the q ocean vs. the top-left panel of Fig. 18 for the o ocean, for when looking at the changes due to the radiation scheme selection; other panels in those two figures present additional variables](#)). This is because the air-sea heat exchange over the ocean is driven by *sst*, which is constant across configurations, and the atmospheric forcing induced by the different configurations and the different balancing parameters is not strong enough to change surface air temperature a lot, in particular when it comes to global mean temperature. This is, however, not the case for the Q-flux=0 and dynamic ocean configurations, in which the ocean, including its surface temperature, is allowed to respond ([Fig. 18](#)).

When ocean configurations change across simulations, large differences are calculated in the model results. In the Q-flux=0 case, differences are expected to be large by design, due to the zero heat flux assumed, which is clearly different from the heat transport that is implicitly included in the p and o oceans. In the Q-flux=0 case any horizontal heat transport can only occur via the atmosphere. All Q-flux=0 model configurations extend the polar ice caps further away from the poles, in agreement with past studies (e.g., Seager et al., 2002; Winton, 2003), by as much as 16 degrees when compared to the prescribed ocean and 32 degrees when compared to the dynamic ocean, which is more pronounced when the SOCRATES radiation is used (Fig. 22). The more extensive ice caps are a direct result of heat that is mostly deposited near the tropics on Earth ([due to its obliquity](#)) that is not able to move fast enough towards the poles via the ocean, resulting in much colder extratropics and poles, which favor the extended ocean freezing. This effect is expected to be amplified on fast-rotating planetary configurations with obliquities lower than that of Earth (23.44 degrees), including the typical value used in the literature of zero. It has to be remembered that in [planet-ROCKE-3D 1.0](#) the Q-flux ocean was using modern Earth's horizontal heat transport instead of zero, resulting in simulations that resemble more the p and o oceans. Since this has limited applicability in a generalized model where even the slightest change of continental configuration or bathymetry would make such a setup invalid, ~~we~~ we decided to use the more easily manageable and more popular configuration (e.g. Batra and Olson, 2024; Sergeev et al., 2022; Wolf et al., 2022) of zero heat transport, although it is highly likely that it will not be the optimal choice for any planet.

1325




Figure 22. Sea ice cover for the Q-flux=0 ocean configuration when the GISS (left) and SOCRATES (right) radiation is used. Global means are 13.5 % for GISS and 19.2 % for SOCRATES.

1330 The global mean ocean ice fraction (*oicefr*, which includes lake ice) is much higher in all Q-flux=0 ocean simulations than the
 prescribed ocean that uses climatological sea ice cover from the preindustrial period, while the dynamic ocean *oicefr* is similar
 to that of the prescribed ocean (Fig. 23). For the Q-flux=0 case the Earth's atmospheric composition with the GISS radiation
 has much more ice than any other simulation, a direct result of the lower temperatures simulated, which are below freezing
 globally-averaged (Fig. 17).

1335




Figure 23. Same as in Fig. 6 for ocean and lake ice fraction (*oicefr*).

As mentioned earlier, the model-simulated *tsurf* resembles the underlying *sst*, regardless of the ocean parameterization used. For the p ocean the difference between the A and N atmospheres for *sst* is zero (they both use the same input data) and for *tsurf* the differences are very small, in particular between 60° S and 60° N latitudes, and maximizes at 2 °C near the North Pole (blue vs. green lines in Fig. 24). The fact that for the same *sst* there are differences in *tsurf* means that the A atmosphere simulation tries to calculate cooler polar temperatures, which is evident when comparing the two o ocean simulations (red line in Fig. 24 vs. zero, since P2SNoM40 is the control). P2SAoM40 is colder at both poles, by as much as 7 °C at the North Pole, which contributes to the buildup of much more sea ice than the p oceans, resulting in cooler *sst*. It is worth mentioning that for *sst* the difference between all simulations presented in Fig. 24 is practically zero below sea ice at the North Pole, which demonstrates that ice insulates the underlying ocean from the atmosphere above.

Another thing worth mentioning is the presence of water vapor in the atmosphere, which is greater in the Northern hemisphere for the p oceans and lesser in the Southern hemisphere, following *tsurf*. What is also interesting is that the tropical precipitation is further north for the p oceans compared against the dynamic ocean simulations, which means that the Inter-Tropical Convergence Zone (ITCZ; the meteorological equator) shifts northward when the temperatures are warmer there.




Figure 24. Zonal mean difference from P2SNoM40 for sea surface temperature (*sst*; first panel), surface air temperature (*tsurf*; second panel), ocean and lake ice fraction (*oicefr*; third panel), total atmospheric water vapor load (*qatm*; fourth panel), and total atmospheric water vapor load (*qatm*) precipitation (*prec*; last panel).

5.5. Medium vs. fine resolutions

1360 The change in resolution, although technical in nature, introduces some important changes in the results. As a reminder, the resolution change is not the same in the atmosphere and in the ocean: in the atmosphere the $4^\circ \times 5^\circ$ latitude by longitude M resolution doubles to $2^\circ \times 2.5^\circ$ in F with the vertical layering remaining the same, while for the ocean the resolution increases by an order of magnitude, going from $4^\circ \times 5^\circ$ and 13 vertical layers to $1^\circ \times 1.125^\circ$ and 40 layers. The ocean change is particularly important since the F resolution resolves better the coastline and narrow straits, as well as the different bathymetries across the global ocean. Vertical mixing is an important process sustaining the oceans' overturning circulations (Munk and Wunsch, 1998; Wunsch and Ferrari, 2004). For Earth simulations, the circumstances generating this mixing (vertical shear and negative buoyancy stratification) occur less frequently in the model at coarse resolution, and the circulation response to the mixing is also muted. The lack of semi-prescribed tidally induced vertical mixing in these simulations also pushes them toward weaker overturning. To avoid shutdowns of the Atlantic overturning circulation in the M configurations, we increased the condition-
1370 independent "background" vertical diffusion from 0.1 to $0.6 \text{ cm}^2 \text{ s}^{-1}$.

An important result when comparing model resolutions is that all M runs with the p and q oceans are colder (Fig. 17). The difference in surface air temperature is very small in the p ocean configuration, since *tsurf* is still driven by the prescribed *sst* which is the same across simulations, but interpolated on a different grid. The differences are also very small compared to other pairs of simulations discussed in the previous sections. For the q ocean though, the change from M to F resolution results
1375 in a strong global mean surface warming across model configurations, about 2°C for GISS and almost 6°C for SOCRATES, compared to the marginal increase for the p ocean.

The different resolutions also lead to a change in the north-south gradient of *sst* across all dynamic ocean configurations (Fig. 25). The fine resolution simulations result in warmer temperatures across all Northern midlatitudes, with zonal mean values reaching 6°C , while in the Southern hemisphere region they are colder by about the same amount. P2GAoM40 shows
1380 exceptional behavior, with sea surface temperatures in the Northern hemisphere resembling those of other M runs, but in the Southern hemisphere looking more like the F ones. All dynamic ocean simulations agree reasonably well in the tropics. Sea ice follows exactly the same pattern, while surface air temperature behaves similarly with *sst*, but with a little more spread over the North and South Poles (Fig. 25). The air temperature changes also affect water vapor in a similar manner, as was shown earlier in a similar situation (Fig. 24).

1385




Figure 25. Zonal mean difference from P2SNoM40 for sea surface temperature (*sst*; first panel), surface air temperature (*tsurf*; second panel), ocean and lake ice fraction (*oicefr*; third panel), total atmospheric water vapor load (*qatm*; last panel).

5.6. Reproducibility of results

In the development and distribution of the source code of ROCKE-3D, we pay particular attention to model reproducibility, which is a widespread problem across many science domains (e.g. Donoho et al., 2008). For that reason, we only update the code by either adding diagnostics that don't change the model results, or fix key bugs and add functionality using the opt-in approach discussed in Sect. 4.3 and Appendix B. Any major development happens in a separate branch of the code repository, which will eventually be distributed at a future date as a future ROCKE-3D model version (e.g. 2.1 or 3.0). However, exact reproducibility is frequently impossible, for hardware purposes: different computer architectures, different compilers with different optimizations, and different ways to compile the model even on the same computer, introduce numerical noise which, when propagated, does not allow one to exactly reproduce model simulations of any complexity, let alone a 3-dimensional GCM like ROCKE-3D. Even with this numerical noise though is random and does not introduce biases, which means that the climatologies calculated by any computer are virtually identical after long integrations, which the modelers should be doing regardless. Long integrations are needed regardless for a proper analysis of results, so a change of computer is not expected to

affect results qualitatively, and even quantitatively any change will not be statistically significant. This ensures that the model version presented here, including its updates described in Sect. 4.6.2, will generate qualitatively the same, but not identical, results on all computer architectures.

1405 As described in more detail in the Data availability section below, we distribute all results presented here to the public, so that they can be used in any way deemed necessary. This includes, but is not limited to, further analyses not presented here; evaluation of simulations performed by others that their climatologies are indeed the same; and continuation of simulations using small changes in physics or model parameters, in order to avoid long spinup times.

6. Outreach

1410 ROCKE-3D tutorials have been held annually since 2017. Participation has been in-person and remote with extensive online documentation that explains how to install supporting software requirements (ROCKE-3D compilers and libraries, 2022) and the model itself (ROCKE-3D installation, 2022). These are currently live documents hosted on Google Docs that are updated as operating system versions change and the supporting software is updated, often in step with operating system updates. The presentation slides are also hosted on google slides as they are updated year-by-year (ROCKE-3D tutorial slides, 2022). The
1415 tutorials are also recorded and placed on the GISS YouTube Channel (ROCKE-3D tutorial video, 2022). This allows anyone in the world to access all materials at any time of year.

7. Conclusions

The ROCKE-3D simulations presented here have addressed limits to the encoded physics, parameterizations of key processes, and inconsistencies that become more apparent in scenarios beyond modern Earth, possibly impacting future projections and
1420 past simulations. This was achieved by a) systematically studying how Earth-centric parameterizations, or the absence of them, affects model results, and b) exploring the role different parameterizations have on an Earth-like simulation, by studying the role of the radiation scheme chosen (GISS vs. SOCRATES), the impact of changing atmospheric composition (presence/absence of oxygen and aerosols), the role of how ocean is parameterized (prescribed; Q-flux=0; dynamic), and the role of horizontal resolution (4x5 vs. 2x2.5 in the atmosphere and 4x5 vs. 1x1.125 in the ocean in latitude and longitude). We
1425 also presented new components of the model, like the geothermal heat flux, an expansion of the dynamic lakes, the ability to simulate thin atmospheres, an improved and much more capable calendar system, and several other updates since ROCKE-3D v1 (Way et al., 2017).

We discussed the importance of radiative balance and how to achieve it in a new world, explained how to identify how much spinup time is needed when starting from non-equilibrium conditions, and how to create a new planet. We also described
1430 planetary configurations that are available to users, including several ancient Earth options, Venus, Mars, the transient atmosphere of the ancient Moon, other rocky exoplanets like those orbiting the Proxima Centauri b and Trappist systems, and

how to use idealized land-ocean configurations, including aquaplanets. We made extensive analyses about influential parameters that users need to be aware of when using ROCKE-3D, including their role, valid ranges (where applicable), and important caveats when modified. These include ways to continue simulations in the case of a crash. Lastly, we presented some important information about the reproducibility of results.

Several future development tasks are already identified and we are actively pursuing them. We are working on enabling the prognostic calculation of a variable mean molecular weight of the atmosphere, which would allow the inclusion of water vapor and condensate in the mass and heat of the atmosphere. We are also including the option to simulate multiple condensable species, which would allow mixed water/CO₂ ice presence on Mars and CH₄ ice on Titan. We are also working on merging the cloud microphysics calculations from the Earth parent model, that will be used for CMIP7, with ROCKE-3D, which will allow more realistic cloud calculations in other planets. Other minor developments continue to be implemented in the model, in what will eventually become ROCKE-3D v3.0 in the future.

The ROCKE-3D development continues to contribute to making a more robust ModelE for modern Earth climate science, by questioning model assumptions and pushing capabilities through research on the Earth-through-time and exoplanet habitability. The model code and tutorials on how it can be installed and used by new users are publicly available.

Appendix A. Key technical updates since ~~planet-ROCKE-3D~~ 1.0

Other than the science-relevant updates described in the paper, there are many other minor updates, which do affect model users that want to update their code to the newest version. These are listed below. It has to be noted that only the few changes that are ported to ~~planet-ROCKE-3D~~ 2.0 from the updated ModelE 2.1 and affect the model's climate independently of prognostic tracers are listed here. The tracers code which affects the prognostic atmospheric composition has received much larger changes, but these are not the subject of this work. In most cases the tracers updates were also less substantial, although not all.

A structural change that introduced only numerical noise differences from the original implementation is that the handling of greenhouse gases was taken out of the radiation code, and is now present in its own module (GHGMOD). Another difference that was ported in is a unit change of the arrays that holds atmospheric mass (and tracer amounts, if enabled) from kg gridbox⁻¹ to kg m⁻².

A new option in ~~planet-ROCKE-3D~~ 2.0 (PCLD_INDICES_LOCAL) avoids a possible array indexing error in computing the fraction of high level clouds (*pcl_{dh}*) if the model top pressure is greater than 680 mbar, and that it also avoids the possible non-physical artifacts of non-zero *pcl_{dl}* and *pcl_{dm}* being possible in model columns with surface pressure less than 680 and 480 mbar, respectively.

Appendix B. Opt-in configuration changes

The selected options listed below need to be defined in the model rundecks prior to compilation. The model has many more additional ways to change its configuration and behavior which extend far beyond the scope of this manuscript.

Table B1: Opt-in configuration changes that modify the default model behavior in a significant way.

Option and section mentioned	Explanation
RAD_ISO_THERMAL_RQT Sect. 2.2.3	Suppresses the prognostic computation of the temperature of the upper 3 radiative layers (outside the model domain) and sets it to the temperature of the topmost model layer. These layers continue to participate in the radiative transfer, but their temperature is not prognostic. The rationale for this is that for thin atmospheres the extremely thin layers at the top have such a low heat capacity that it is too easy to overshoot when updating their temperature.
GHY_NO_LIQUID Sect. 2.2.3	Suppress the liquid water movement in the soil. All deposited water will be assumed to stay at the surface. This option should be enabled for atmospheres with surface pressure below the triple point of water, since no liquid water can exist in such conditions.
EOT={off,naive,default} Sect. 2.2.4	Choice of equation of time (EOT) to be used. Off: do not use it, only provided for backward compatibility; naïve: does not include the contributions due to obliquity; default: the default and full implementation of EOT.
Clouds inhomogeneity correction Sect. 3.3	Define a RADN8 file (like e.g. in P2GAoF40) for an Earth-centric spatially-varying value. Do not define RADN8 and set instead KCLDEP=1 (tells the model to use a constant global value) and EPSCON=0.12 (tells the model what that value should be; 0.12 used here).
TAPER_UTCF Sect. 3.3	When defined, allows for a more physically-consistent profile of longwave flux divergence near the top of the atmosphere.
use_vmp={0,1} Sect. 5.3	Enables (=1; default and used here) or disables (=0) “virtual” mixed-phase clouds. A model rebalancing is required after changing this value.

Appendix C. Spectral files and stellar spectra

Table A4.C1.: List of spectral files provided with ROCKE-3D for operation with SOCRATES. Unless otherwise indicated in the description, spectral files use HITRAN 2012 and have a maximum pressure and temperature of 1 bar and 400 K, respectively. All spectral files contain H₂O and use the MT-CKD water vapor continuum model (Mlawer et al., 2012). For more details refer to the online documentation (ROCKE-3D spectral files).

Name	LW bins	SW bins	Description
------	---------	---------	-------------

Present-day Earth			
ga7	9	6	default Met Office modern Earth, temperatures +/- 50 K from the mid-latitude summer profile
ga7_dsa	9	6	default Met Office modern Earth, with expanded temperature range
dsa	12	21	increased precision for high temperature, high water vapor, and for M-dwarf spectra
Paleoproterozoic			
etw_pz	12	29	N ₂ O ₂ dominated, up to 5000 ppmv CO ₂ , up to 200 ppmv CH ₄
Archean			
dsa_ar	12	29	N ₂ dominated, anoxic, up to 1% CO ₂ , up to 1% CH ₄
dsa_arcc	17	35	N ₂ dominated, anoxic, 1-10% CO ₂ , up to 1% CH ₄
dsa_arva	17	18	N ₂ dominated, anoxic, up to 1% CO ₂ , up to 10 ppmv CH ₄ , up to 1000 ppmv N ₂ O
dsa_ar10bar	15	43	N ₂ /H ₂ dominated, anoxic, up to 1% CO ₂ , up to 1% CH ₄ , 10 bar maximum pressure
etw_arcc10bar	17	43	N ₂ /H ₂ dominated, anoxic, 1-20% CO ₂ , up to 1% CH ₄ , 10 bar maximum pressure
etw_arcc10bar_highco2	17	43	N ₂ /H ₂ dominated, anoxic, 20-80% CO ₂ , up to 1% CH ₄ , 10 bar maximum pressure
Mars through time			
dsa_mars	17	42	CO ₂ dominated, anoxic, 10 bar maximum pressure
dsa_pm	16	46	CO ₂ /H ₂ dominated, anoxic, up to 1% CH ₄ , Wordsworth et al. (2017) CIAs, 10 bar maximum pressure
dsa_pm_n2o	16	46	CO ₂ /H ₂ dominated, anoxic, up to 1% CH ₄ , up to 1000 ppmv N ₂ O, Wordsworth et al. (2017) CIAs, 10 bar maximum pressure
dsa_pm_hit16	16	46	CO ₂ /H ₂ dominated, anoxic, up to 1% CH ₄ , Turbet et al. (2020) CIAs, 10 bar maximum pressure, HITRAN2016
Other worlds			
dsa_titan	10	14	N ₂ dominated, anoxic, up to 5% CH ₄ , up to 1% H ₂ , 300 K maximum temperature, 2 bar maximum pressure
etw_moon	17	29	CO dominated, up to 30% CO ₂ , up to 10% H ₂

Table A4.C2.: List of stellar spectra provided with ROCKE-3D for operation with SOCRATES. Stellar spectrum are crafted either from exact replication or interpolation of theoretical spectrum from the BT-Settl and BT-NextGen grid of models (e.g. Allard, 2013), with specifications and sources shown in the description column.

Short Name	T _{eff} (K)	Description
lte060-4.5-0.0a+0.0.BT-NextGen	6000	log(g) = 4.5, [Fe/H] = 0, BT-Next Gen .
sun	5776	Solar spectrum, SPARC/SOLARIS (Lean et al., 2005).
sun_0.27gya	5773	Sun scaled by Claire et al. (2012) for 0.27 gya.
sun_0.715gya	5763	Sun scaled by Claire et al. (2012) for 0.75 gya.
sun_1.3gya	5750	Sun scaled by Claire et al. (2012) for 1.3 gya.
sun_2.0gya	5731	Sun scaled by Claire et al. (2012) for 2.0 gya.
sun_2.9gya	5700	Sun scaled by Claire et al. (2012) for 2.9 gya.
sun_3.8gya	5660	Sun scaled by Claire et al. (2012) for 3.8 gya.
lte055-4.5-0.0a+0.0.BT-NextGen	5500	log(g) = 4.5, [Fe/H] = 0, BT-NextGen.
hd22049	5084	HD 22049, VPL Spectral Database (Segura et al., 2003).
lte050-4.5-0.0a+0.0.BT-NextGen	5000	log(g) = 4.5, [Fe/H] = 0, BT-NextGen .
sun_X4875	4875	Sun red giant phase version 1 from PARSEC.
sun_X4805	4803	Sun red giant phase version 2 from PARSEC.
epsindi	4730	Epsilon Indi (Rugheimer et al., 2013).
sun_X4545	4543	Sun red giant phase version 3 from PARSEC.
lte045-4.5-0.0a+0.0.BT-NextGen	4500	log(g) = 4.5,[Fe/H] = 0, BT-NextGen.
hd85512	4305	HD 85512, MUSCLES Database (Loyd et al., 2016).
lte040-4.5-0.0a+0.0.BT-NextGen	4000	log(g) = 4.5, [Fe/H] = 0, BT-NextGen.
hirano	3933	K2-155, log(g)=4.73, [Fe/H] = -0.42, BT-Settl.
K2-240	3810	K2-240, log(g)=4.7, [Fe/H] = -0.1, BT-Settl.
k186	3788	Kepler-186 (Bolmont et al., 2014).
Gl514	3728	Gl 514, log(g) = 4.76, [Fe/H] = -0.14, BT-Settl.
lte037-4.5-0.0a+0.0.BT-NextGen	3700	log(g) = 4.5, [Fe/H] = 0, BT-NextGen.
TOI1266	3600	TOI-1266, log(g) = 4.85, [Fe/H] = -0.5, BT-Settl.
TOI2285	3491	TOI-2285, log(g) = 4.77, [Fe/H] = -0.05, BT-Settl.
toi700	3480	TOI 700, log(g) = 4.81318, [Fe/H] = 0, BT-Settl.
K2-18	3457	log(g) = 4.79044, [Fe/H] = 0.123; BT-Settl.
L98-59	3415	log(g) = 4.86, [Fe/H] = -0.46; BT-Settl.

adleo	3390	Ad Leo,VPL Spectral Database (Segura et al., 2005).
lte033-4.5-0.0a+0.0.BT-NextGen	3300	$\log(g) = 4.5$, $[\text{Fe}/\text{H}] = 0$, , BT-Next Gen.
gj876	3200	GJ 876 (Domagal-Goldman et al., 2014).
k1649	3200	Kepler-1649, $\log(g) = 5$, $[\text{Fe}/\text{H}] = 0$, BT-Settl.
pc	3042	Proxima Centauri (Meadows et al., 2018).
lte030-4.5-0.0a+0.0.BT-NextGen	3000	$\log(g) = 4.5$, $[\text{Fe}/\text{H}] = 0$, BT-NextGen.
LHS_1140	2988	$\log(g) = 5.04$, $[\text{Fe}/\text{H}] = -0.24$, BT-Settl.
teegarden	2904	Teegarden's Star $\log(g) = 5.3288$, $[\text{Fe}/\text{H}] = -0.19$, BT-NextGen.
SPECULOOS-2	2850	SPECULOOS-2, $\log(g) = 5.126$, $[\text{Fe}/\text{H}] = -0.028$, BT-Settl.
trappist-1	2600	TRAPPIST-1, $\log(g) = 5$, $[\text{Fe}/\text{H}] = 0$, BT-Settl.
lte026-4.5-0.0a+0.0.BT-NextGen	2600	$\log(g) = 4.5$, $[\text{Fe}/\text{H}] = 0$, BT-Next Gen.

1475

Appendix D. Explanation of variable names used

Table D1: Explanation and units of variables used in the manuscript.

Variable	Explanation
<i>cldi</i>	Cloud condensed ice column [kg m^{-2}].
<i>cldtp</i>	Cloud top pressure, as seen by radiation [hPa].
<i>cldtpt</i>	Cloud top temperature, as seen by radiation [$^{\circ}\text{C}$].
<i>cldw</i>	Cloud condensed water column [kg m^{-2}].
<i>clwp</i>	Cloud liquid water path [kg m^{-2}].
<i>grnd_alb</i>	Ground albedo.
<i>gwtr</i>	Total Earth water [kg]
<i>icecld</i>	Ice cloud concentration [ppmm]
<i>IWPrad</i>	Cloud ice water path, as seen by radiation [kg m^{-2}].
<i>lakefr</i>	Lake fraction [%].
<i>LWPrad</i>	Cloud liquid water path, as seen by radiation [kg m^{-2}].
<i>mwl</i>	Total amount of water in rivers and lakes [kg]
<i>net_rad_planet</i>	Net radiative balance at the top of the atmosphere [W m^{-2}]. It includes both the incoming solar stellar and outgoing thermal radiation fluxes, with opposing signs. A well equilibrated atmosphere is assumed when $ \textit{net_rad_planet} \leq 0.2 \text{ W m}^{-2}$.
<i>ocean_fkph</i>	Background vertical diffusion coefficient [$\text{cm}^2 \text{ s}^{-1}$]

Variable	Explanation
<i>oicefr</i>	Ocean and lake ice fraction [%].
<i>pcl dh</i>	High cloud cover [%] as seen by radiation. It includes clouds from 440 hPa to the model top.
<i>pcl dl</i>	Low cloud cover [%] as seen by radiation. It includes clouds from the surface to 680 hPa.
<i>pcl dm</i>	Medium cloud cover [%] as seen by radiation. It includes clouds between 680 and 440 hPa.
<i>pcl dt</i>	Total cloud cover [%] as seen by radiation. It includes all clouds.
<i>plan_alb</i>	Planetary albedo.
<i>pot_temp</i>	Ocean potential temperature [°C].
<i>prec</i>	Precipitation [mm day ⁻¹].
<i>qatm</i>	Atmospheric water vapor column [kg m ⁻²].
<i>qsurf</i>	Surface air specific humidity [kg kg ⁻¹].
<i>radiusi_multiplier</i>	Ice cloud particle radius multiplier [dimensionless].
<i>radiusl_multiplier</i>	Liquid cloud particle radius multiplier [dimensionless].
<i>snowdp</i>	Snow depth [mm H ₂ O].
<i>snowfall</i>	Snow fall [mm day ⁻¹].
<i>snowfr</i>	Snow fraction [%].
<i>SpHuL</i>	Specific humidity at model levels [kg kg ⁻¹].
<i>srnf_toa</i>	Solar -Stellar radiation net flux at the top of the atmosphere [W m ⁻²].
<i>sst</i>	Sea surface temperature [°C].
<i>TempL</i>	Air temperature at model levels [K].
<i>trnf_toa</i>	Thermal radiation net flux at the top of the atmosphere [W m ⁻²].
<i>tsurf</i>	Surface air temperature [°C].
<i>U00a</i>	Threshold grid box mean relative humidity for stratiform clouds above the planetary boundary layer top in the absence of moist convection. The threshold increases towards 1.0 as layer thickness decreases, in an attempt to be scale-aware [dimensionless].
<i>U00b</i>	Scaling parameter for stratiform cloud threshold relative humidity within the planetary boundary layer or in the presence of moist convection. Typical values of the resulting threshold are much closer to 1.0 than U00a. The scaling is based on an assumed Gaussian distribution of saturation deficit as suggested by Siebesma et al. (2003) [dimensionless].
<i>WMU_multiplier</i>	Tuning factor applied to the critical liquid cloud mixing ratio above which rapid autoconversion occurs [dimensionless].
<i>WMUI_multiplier</i>	Tuning factor applied to the critical ice cloud mixing ratio above which rapid autoconversion occurs [dimensionless].

Variable	Explanation
<i>wsurf</i>	Wind speed at surface [m s^{-1}].
<i>wtrcld</i>	Water cloud concentration [ppmm]

Code availability

The ROCKE-3D model is publicly available: <https://simplex.giss.nasa.gov/gcm/ROCKE-3D/>. [The model code, all output, and the model configurations used here, are available on a zenodo archive](#) (Tsigaridis et al., 2025). All ROCKE-3D prescribed ocean configurations listed in Table 3, as well as their Q-flux=0 and dynamic ocean equivalent configurations analyzed here, are provided as template “rundecks”, the configuration files of the model, using the same naming convention and modified as described in Sect. 4.3, together with the model code.

Data availability

All NetCDF data used in this and other publications of ROCKE-3D can be downloaded from the NCCS data portal: https://portal.nccs.nasa.gov/GISS_modelE/ROCKE-3D/publication-supplements/. SOCRATES spectral files are also available from the same portal: https://portal.nccs.nasa.gov/GISS_modelE/ROCKE-3D/spectral_files/. A zenodo archive (Tsigaridis et al., 2025) with data from this work includes 1) all rundecks used in this work, for users to be able to reproduce our simulations; 2) all restart files at the end of the equilibration period for all simulations listed in Fig. 6 and throughout the manuscript, for simulation continuations without the need of a spinup; 3) time series of globally and annually averaged quantities in the atmosphere from the start of each simulation, including spinup, a small subset of which have been used to create Fig. 7; 4) spatially-varying climatological annual means of all atmospheric model output which was used for all maps presented in the manuscript. These data are also available in the NCCS data portal (https://portal.nccs.nasa.gov/GISS_modelE/ROCKE-3D/publication-supplements/Tsigaridis2025GMD-planet_2.0/) together with the individual years that constructed the climatologies, which are too voluminous for zenodo to host.

Author contributions

KT and ADDG designed the work. KT, ADDG, and MK performed the model balancing. KT ran all simulations and made most of the plots. Others that made plots include CMC, LS, and MJW. ASA helped with the clouds diagnostics and analysis. IA developed the Moon configuration, and the Mars configuration together with KT and JPP. MAC and LS worked on the alternate continental configurations. LS and RAR worked on the geothermal heat flux. TLC improved the calendar and added the equation of time. CMC and AL worked on the aquaplanet and ocean simulations. MJW worked on Venus and exoplanet simulations. NYK, GLR, RAR, and IA worked on the new surface hydrology physics. ETW worked on SOCRATES radiative

transfer and radiation tables. KT wrote the first version of the manuscript and led subsequent edits with the help of all co-authors.

Competing interests

1505 The authors declare that they have no conflict of interest.

Acknowledgements

This work was supported by NASA's Nexus for Exoplanet System Science (NExSS) and the NASA Interdisciplinary Consortia for Astrobiology Research (ICAR). Further support for this research was provided by NASA Earth and Planetary Science Division Research Programs, through ISFM work package ROCKE-3D at the Goddard Institute for Space Studies. We
1510 acknowledge support from the GSFC Sellers Exoplanet Environments Collaboration (SEEC), which is funded by the NASA Planetary Science Division's Internal Scientist Funding Model. Earth climate modeling at GISS is supported by the NASA Modeling, Analysis and Prediction program (MAP). Thin atmospheres research was supported by NASA Solar System Workings program award 80NSSC21K0163. Resources supporting this work were provided by the NASA High-End Computing (HEC) Program through the NASA Center for Climate Simulation (NCCS) at Goddard Space Flight Center.

1515 **References**

Adcroft, A., Scott, J. R., and Marotzke, J.: Impact of geothermal heating on the global ocean circulation, *Geophysical Research Letters*, 28, 1735–1738, <https://doi.org/10.1029/2000GL012182>, 2001.

Aleinov, I., Way, M. J., Harman, C., Tsigaridis, K., Wolf, E. T., and Gronoff, G.: Modeling a Transient Secondary Paleolunar Atmosphere: 3-D Simulations and Analysis, *Geophysical Research Letters*, 46, 5107–5116,
1520 <https://doi.org/10.1029/2019GL082494>, 2019.

Allard, F.: The BT-Settl Model Atmospheres for Stars, Brown Dwarfs and Planets, *Proceedings of the International Astronomical Union*, 8, 271–272, <https://doi.org/10.1017/S1743921313008545>, 2013.

Amundsen, D. S., Mayne, N. J., Baraffe, I., Manners, J., Tremblin, P., Drummond, B., Smith, C., Acreman, D. M., and Homeier, D.: The UK Met Office global circulation model with a sophisticated radiation scheme applied to the hot Jupiter HD
1525 209458b, *A&A*, 595, A36, <https://doi.org/10.1051/0004-6361/201629183>, 2016.

Amundsen, D. S., Tremblin, P., Manners, J., Baraffe, I., and Mayne, N. J.: Treatment of overlapping gaseous absorption with the correlated-k method in hot Jupiter and brown dwarf atmosphere models, *A&A*, 598, A97, <https://doi.org/10.1051/0004-6361/201629322>, 2017.

Aylmer, J., Ferreira, D., and Feltham, D.: Impacts of Oceanic and Atmospheric Heat Transports on Sea Ice Extent, *Journal of*
1530 *Climate*, 33, 7197–7215, <https://doi.org/10.1175/JCLI-D-19-0761.1>, 2020.

- Batra, K. and Olson, S. L.: Climatic Effects of Ocean Salinity on M Dwarf Exoplanets, *The Astrophysical Journal Letters*, 971, L11, <https://doi.org/10.3847/2041-8213/ad63a5>, 2024.
- Bauer, S. E., Tsigaridis, K., Faluvegi, G., Kelley, M., Lo, K. K., Miller, R. L., Nazarenko, L., Schmidt, G. A., and Wu, J.: Historical (1850-2014) aerosol evolution and role on climate forcing using the GISS ModelE2.1 contribution to CMIP6, *Journal of Advances in Modeling Earth Systems*, e2019MS001978, <https://doi.org/10.1029/2019MS001978>, 2020.
- Běhouňková, M., Tobie, G., Choblet, G., and Čadež, O.: Coupling mantle convection and tidal dissipation: Applications to Enceladus and Earth-like planets, *Journal of Geophysical Research: Planets*, 115, <https://doi.org/10.1029/2009JE003564>, 2010.
- Bitz, C. M., Holland, M. M., Hunke, E. C., and Moritz, R. E.: Maintenance of the Sea-Ice Edge, *Journal of Climate*, 18, 2903–2921, <https://doi.org/10.1175/JCLI3428.1>, 2005.
- Bolmont, E., Raymond, S. N., von Paris, P., Selsis, F., Hersant, F., Quintana, E. V., and Barclay, T.: FORMATION, TIDAL EVOLUTION, AND HABITABILITY OF THE KEPLER-186 SYSTEM, *The Astrophysical Journal*, 793, 3, <https://doi.org/10.1088/0004-637X/793/1/3>, 2014.
- Bougher, S. W., Hunten, D. M., Phillips, R. J., Matthews, M. S., Ruskin, A. S., and Guerrieri, M. L. (Eds.): *Venus II: Geology, Geophysics, Atmosphere, and Solar Wind Environment*, University of Arizona Press, <https://doi.org/10.2307/j.ctv27tct5m>, 1997.
- Bouley, S., Baratoux, D., Matsuyama, I., Forget, F., Séjourné, A., Turbet, M., and Costard, F.: Late Tharsis formation and implications for early Mars, *Nature*, 531, 344–347, <https://doi.org/10.1038/nature17171>, 2016.
- Brunetti, M., Kasparian, J., and Vérard, C.: Co-existing climate attractors in a coupled aquaplanet, *Climate Dynamics*, 53, 6293–6308, <https://doi.org/10.1007/s00382-019-04926-7>, 2019.
- Bryan, K.: A numerical method for the study of the circulation of the world ocean, *Journal of Computational Physics*, 4, 347–376, [https://doi.org/10.1016/0021-9991\(69\)90004-7](https://doi.org/10.1016/0021-9991(69)90004-7), 1969.
- Bryan, K.: Poleward Heat Transport by the Ocean: Observations and Models, *Annual Review of Earth and Planetary Sciences*, 10, 15–38, <https://doi.org/10.1146/annurev.ea.10.050182.000311>, 1982.
- Butcher, F. E. G., Balme, M. R., Gallagher, C., Arnold, N. S., Conway, S. J., Hagermann, A., and Lewis, S. R.: Recent Basal Melting of a Mid-Latitude Glacier on Mars, *Journal of Geophysical Research: Planets*, 122, 2445–2468, <https://doi.org/10.1002/2017JE005434>, 2017.
- Claire, M. W., Sheets, J., Cohen, M., Ribas, I., Meadows, V. S., and Catling, D. C.: THE EVOLUTION OF SOLAR FLUX FROM 0.1 nm TO 160 μ m: QUANTITATIVE ESTIMATES FOR PLANETARY STUDIES, *The Astrophysical Journal*, 757, 95, <https://doi.org/10.1088/0004-637X/757/1/95>, 2012.
- Colose, C. M., Haqq-Misra, J., Wolf, E. T., Del Genio, A. D., Barnes, R., Way, M. J., and Ruedy, R.: Effects of Spin–Orbit Resonances and Tidal Heating on the Inner Edge of the Habitable Zone, *The Astrophysical Journal*, 921, 25, <https://doi.org/10.3847/1538-4357/ac135c>, 2021.
- Davies, J. H.: Global map of solid Earth surface heat flow, *Geochemistry, Geophysics, Geosystems*, 14, 4608–4622, <https://doi.org/10.1002/ggge.20271>, 2013.

- Del Genio, A. D., Yao, M.-S., Kovari, W., and Lo, K. K.-W.: A Prognostic Cloud Water Parameterization for Global Climate Models, *Journal of Climate*, 9, 270–304, [https://doi.org/10.1175/1520-0442\(1996\)009<0270:APCWPF>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<0270:APCWPF>2.0.CO;2), 1996.
- Del Genio, A. D., Way, M. J., Kiang, N. Y., Aleinov, I., Puma, M. J., and Cook, B.: Climates of Warm Earth-like Planets. III. Fractional Habitability from a Water Cycle Perspective, *The Astrophysical Journal*, 887, 197, <https://doi.org/10.3847/1538-4357/ab57fd>, 2019a.
- Del Genio, A. D., Way, M. J., Amundsen, D. S., Aleinov, I., Kelley, M., Kiang, N. Y., and Clune, T. L.: Habitable Climate Scenarios for Proxima Centauri b with a Dynamic Ocean, *Astrobiology*, 19, 99–125, <https://doi.org/10.1089/ast.2017.1760>, 2019b.
- Domagal-Goldman, S. D., Segura, A., Claire, M. W., Robinson, T. D., and Meadows, V. S.: ABIOTIC OZONE AND OXYGEN IN ATMOSPHERES SIMILAR TO PREBIOTIC EARTH, *The Astrophysical Journal*, 792, 90, <https://doi.org/10.1088/0004-637X/792/2/90>, 2014.
- Donoho, D., Maleki, A., Shahram, M., and Stodden, V.: 15 Years of Reproducible Research in Computational Harmonic Analysis, 2008.
- Edwards, J. M.: Efficient Calculation of Infrared Fluxes and Cooling Rates Using the Two-Stream Equations, *Journal of Atmospheric Sciences*, 53, 1921–1932, [https://doi.org/10.1175/1520-0469\(1996\)053<1921:ECOIFA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1996)053<1921:ECOIFA>2.0.CO;2), 1996.
- Edwards, J. M. and Slingo, A.: Studies with a flexible new radiation code. I: Choosing a configuration for a large-scale model, *Quarterly Journal of the Royal Meteorological Society*, 122, 689–719, <https://doi.org/10.1002/qj.49712253107>, 1996.
- Emile-Geay, J. and Madec, G.: Geothermal heating, diapycnal mixing and the abyssal circulation, *Ocean Science*, 5, 203–217, <https://doi.org/10.5194/os-5-203-2009>, 2009.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Faucher, T. J., Turbet, M., Wolf, E. T., Boutle, I., Way, M. J., Del Genio, A. D., Mayne, N. J., Tsigaridis, K., Kopparapu, R. K., Yang, J., Forget, F., Mandell, A., and Domagal Goldman, S. D.: TRAPPIST-1 Habitable Atmosphere Intercomparison (THAI): motivations and protocol version 1.0, *Geoscientific Model Development*, 13, 707–716, <https://doi.org/10.5194/gmd-13-707-2020>, 2020.
- Faucher, T. J., Villanueva, G. L., Sergeev, D. E., Turbet, M., Boutle, I. A., Tsigaridis, K., Way, M. J., Wolf, E. T., Domagal-Goldman, S. D., Forget, F., Haqq-Misra, J., Kopparapu, R. K., Manners, J., and Mayne, N. J.: The TRAPPIST-1 Habitable Atmosphere Intercomparison (THAI). III. Simulated Observables—the Return of the Spectrum, *The Planetary Science Journal*, 3, 213, <https://doi.org/10.3847/PSJ/ac6cf1>, 2022.
- Ferreira, D., Marshall, J., and Rose, B.: Climate Determinism Revisited: Multiple Equilibria in a Complex Climate Model, *Journal of Climate*, 24, 992–1012, <https://doi.org/10.1175/2010JCLI3580.1>, 2011.
- Forget, F., Bertrand, T., Vangichith, M., Leconte, J., Millour, E., and Lellouch, E.: A post-new horizons global climate model of Pluto including the N₂, CH₄ and CO cycles, *Icarus*, 287, 54–71, <https://doi.org/10.1016/j.icarus.2016.11.038>, 2017.
- Guzewich, S. D., Way, M. J., Aleinov, I., Wolf, E. T., Del Genio, A., Wordsworth, R., and Tsigaridis, K.: 3D Simulations of the Early Martian Hydrological Cycle Mediated by a H₂-CO₂ Greenhouse, *Journal of Geophysical Research: Planets*, 126, e2021JE006825, <https://doi.org/10.1029/2021JE006825>, 2021.

- Hansen, J., Russell, G., Rind, D., Stone, P., Lacis, A., Lebedeff, S., Ruedy, R., and Travis, L.: Efficient Three-Dimensional Global Models for Climate Studies: Models I and II, *Monthly Weather Review*, 111, 609–662, 1983.
- 1605 Haywood, A. M., Dowsett, H. J., Dolan, A. M., Rowley, D., Abe-Ouchi, A., Otto-Bliesner, B., Chandler, M. A., Hunter, S. J., Lunt, D. J., Pound, M., and Salzmann, U.: The Pliocene Model Intercomparison Project (PlioMIP) Phase 2: scientific objectives and experimental design, *Climate of the Past*, 12, 663–675, <https://doi.org/10.5194/cp-12-663-2016>, 2016.
- Held, I. M.: The Partitioning of the Poleward Energy Transport between the Tropical Ocean and Atmosphere, *Journal of the Atmospheric Sciences*, 58, 943–948, [https://doi.org/10.1175/1520-0469\(2001\)058<0943:TPOTPE>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<0943:TPOTPE>2.0.CO;2), 2001.
- 1610 Hendrix, A. R., Hurford, T. A., Barge, L. M., Bland, M. T., Bowman, J. S., Brinckerhoff, W., Buratti, B. J., Cable, M. L., Castillo-Rogez, J., Collins, G. C., Diniega, S., German, C. R., Hayes, A. G., Hoehler, T., Hosseini, S., Howett, C. J. A., McEwen, A. S., Neish, C. D., Neveu, M., Nordheim, T. A., Patterson, G. W., Patthoff, D. A., Phillips, C., Rhoden, A., Schmidt, B. E., Singer, K. N., Soderblom, J. M., and Vance, S. D.: The NASA Roadmap to Ocean Worlds, *Astrobiology*, 19, 1–27, <https://doi.org/10.1089/ast.2018.1955>, 2019.
- 1615 Henning, W. G. and Hurford, T.: TIDAL HEATING IN MULTILAYERED TERRESTRIAL EXOPLANETS, *The Astrophysical Journal*, 789, 30, <https://doi.org/10.1088/0004-637X/789/1/30>, 2014.
- Hofmann, M. and Maqueda, M. A.: Geothermal heat flux and its influence on the oceanic abyssal circulation and radiocarbon distribution, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2008GL036078>, 2009.
- Hörner, J. and Voigt, A.: Sea-ice thermodynamics can determine waterbelt scenarios for Snowball Earth, *Earth System Dynamics*, 15, 215–223, <https://doi.org/10.5194/esd-15-215-2024>, 2024.
- 1620 Hu, Y. and Yang, J.: Role of ocean heat transport in climates of tidally locked exoplanets around M dwarf stars, *Proceedings of the National Academy of Sciences*, 111, 629–634, <https://doi.org/10.1073/pnas.1315215111>, 2014.
- van den Hurk, B., Kim, H., Krinner, G., Seneviratne, S. I., Derksen, C., Oki, T., Douville, H., Colin, J., Ducharne, A., Cheruy, F., Viovy, N., Puma, M. J., Wada, Y., Li, W., Jia, B., Alessandri, A., Lawrence, D. M., Weedon, G. P., Ellis, R., Hagemann, S., Mao, J., Flanner, M. G., Zampieri, M., Materia, S., Law, R. M., and Sheffield, J.: LS3MIP (v1.0) contribution to CMIP6: the Land Surface, Snow and Soil moisture Model Intercomparison Project – aims, setup and expected outcome, *Geoscientific Model Development*, 9, 2809–2832, <https://doi.org/10.5194/gmd-9-2809-2016>, 2016.
- 1625 Intergovernmental Panel on Climate Change (IPCC) (Ed.): The Earth’s Energy Budget, Climate Feedbacks and Climate Sensitivity, in: *Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, 923–1054, <https://doi.org/10.1017/9781009157896.009>, 2023.
- Ito, G., Romanou, A., Kiang, N. Y., Faluvegi, G., Aleinov, I., Ruedy, R., Russell, G., Lerner, P., Kelley, M., and Lo, K.: Global Carbon Cycle and Climate Feedbacks in the NASA GISS ModelE2.1, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002030, <https://doi.org/10.1029/2019MS002030>, 2020.
- 1635 Johns, T. C., Carnell, R. E., Crossley, J. F., Gregory, J. M., Mitchell, J. F. B., Senior, C. A., Tett, S. F. B., and Wood, R. A.: The second Hadley Centre coupled ocean-atmosphere GCM: model description, spinup and validation, *Climate Dynamics*, 13, 103–134, <https://doi.org/10.1007/s003820050155>, 1997.
- Jones, C. D., Arora, V., Friedlingstein, P., Bopp, L., Brovkin, V., Dunne, J., Graven, H., Hoffman, F., Ilyina, T., John, J. G., Jung, M., Kawamiya, M., Koven, C., Pongratz, J., Raddatz, T., Randerson, J. T., and Zaehle, S.: C4MIP – The Coupled

- 1640 Climate–Carbon Cycle Model Intercomparison Project: experimental protocol for CMIP6, Geoscientific Model Development, 9, 2853–2880, <https://doi.org/10.5194/gmd-9-2853-2016>, 2016.
- Kelley, M., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Del Genio, A. D., Elsaesser, G. S., Faluvegi, G., Kiang, N. Y., Kim, D., Lacis, A. A., Leboissetier, A., LeGrande, A. N., Lo, K. K., Marshall, J., Matthews, E. E.,
1645 McDermid, S., Mezuman, K., Miller, R. L., Murray, L. T., Oinas, V., Orbe, C., García-Pando, C. P., Perlwitz, J. P., Puma, M. J., Rind, D., Romanou, A., Shindell, D. T., Sun, S., Tausnev, N., Tsigaridis, K., Tselioudis, G., Weng, E., Wu, J., and Yao, M.-S.: GISS-E2.1: Configurations and Climatology, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002025, <https://doi.org/10.1029/2019MS002025>, 2020.
- Kim, Y., Moorcroft, P. R., Aleinov, I., Puma, M. J., and Kiang, N. Y.: Variability of phenology and fluxes of water and carbon with observed and simulated soil moisture in the Ent Terrestrial Biosphere Model (Ent TBM version 1.0.1.0.0), *Geoscientific Model Development*, 8, 3837–3865, <https://doi.org/10.5194/gmd-8-3837-2015>, 2015.
- Klinger, B. A. and Marotzke, J.: Meridional Heat Transport by the Subtropical Cell, *Journal of Physical Oceanography*, 30, 696–705, [https://doi.org/10.1175/1520-0485\(2000\)030<0696:MHTBTS>2.0.CO;2](https://doi.org/10.1175/1520-0485(2000)030<0696:MHTBTS>2.0.CO;2), 2000.
- Kopparapu, R. K., Ramirez, R., Kasting, J. F., Eymet, V., Robinson, T. D., Mahadevan, S., Terrien, R. C., Domagal-Goldman, S., Meadows, V., and Deshpande, R.: HABITABLE ZONES AROUND MAIN-SEQUENCE STARS: NEW ESTIMATES, *The Astrophysical Journal*, 765, 131, <https://doi.org/10.1088/0004-637X/765/2/131>, 2013.
- Lacis, A. A. and Oinas, V.: A Description of the Correlated k Distribution Method for Modeling Nongray Gaseous Absorption, Thermal Emission, and Multiple Scattering in Vertically Inhomogeneous Atmospheres, *J. Geophys. Res.-Atmos.*, 96, 9027–9063, <https://doi.org/10.1029/90jd01945>, 1991.
- 1660 Lean, J., Rottman, G., Harder, J., and Kopp, G.: *SORCE Contributions to New Understanding of Global Change and Solar Variability*, *Solar Physics*, 230, 27–53, <https://doi.org/10.1007/s11207-005-1527-2>, 2005.
- Levitus, S. and Boyer, T. P.: *World ocean atlas 1994. Vol. 4, Temperature*, 1994.
- Levitus, S., Burgett, R., and Boyer, T. P.: *World ocean atlas 1994. Vol. 3, Salinity*, 1994.
- Loyd, R. O. P., France, K., Youngblood, A., Schneider, C., Brown, A., Hu, R., Linsky, J., Froning, C. S., Redfield, S.,
1665 Rugheimer, S., and Tian, F.: THE MUSCLES TREASURY SURVEY. III. X-RAY TO INFRARED SPECTRA OF 11 M AND K STARS HOSTING PLANETS, *The Astrophysical Journal*, 824, 102, <https://doi.org/10.3847/0004-637X/824/2/102>, 2016.
- Manabe, S. and Stouffer, R. J.: Century-scale effects of increased atmospheric CO₂ on the ocean–atmosphere system, *Nature*, 364, 215–218, <https://doi.org/10.1038/364215a0>, 1993.
- 1670 Manabe, S. and Wetherald, R. T.: Thermal Equilibrium of the Atmosphere with a Given Distribution of Relative Humidity, *Journal of the Atmospheric Sciences*, 24, 241–259, [https://doi.org/10.1175/1520-0469\(1967\)024<0241:TEOTAW>2.0.CO;2](https://doi.org/10.1175/1520-0469(1967)024<0241:TEOTAW>2.0.CO;2), 1967.
- Marshall, J., Hill, C., Perelman, L., and Adcroft, A.: Hydrostatic, quasi-hydrostatic, and nonhydrostatic ocean modeling, *Journal of Geophysical Research: Oceans*, 102, 5733–5752, <https://doi.org/10.1029/96JC02776>, 1997.
- 1675 Meadows, V. S., Arney, G. N., Schwieterman, E. W., Lustig-Yaeger, J., Lincowski, A. P., Robinson, T., Domagal-Goldman, S. D., Deitrick, R., Barnes, R. K., Fleming, D. P., Luger, R., Driscoll, P. E., Quinn, T. R., and Crisp, D.: The Habitability of

- Proxima Centauri b: Environmental States and Observational Discriminants, *Astrobiology*, 18, 133–189, <https://doi.org/10.1089/ast.2016.1589>, 2018.
- 1680 Miller, J. R., Russell, G. L., and Tsang, L.-C.: Annual oceanic heat transports computed from an atmospheric model, *Dynamics of Atmospheres and Oceans*, 7, 95–109, [https://doi.org/10.1016/0377-0265\(83\)90012-X](https://doi.org/10.1016/0377-0265(83)90012-X), 1983.
- 1685 Miller, R. L., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Kelley, M., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Del Genio, A. D., Elsaesser, G. S., Faluvegi, G., Kiang, N. Y., Kim, D., Lacis, A. A., Leboissetier, A., LeGrande, A. N., Lo, K. K., Marshall, J., Matthews, E. E., McDermid, S., Mezuman, K., Murray, L. T., Oinas, V., Orbe, C., Pérez García-Pando, C., Perlwitz, J. P., Puma, M. J., Rind, D., Romanou, A., Shindell, D. T., Sun, S., Tausnev, N., Tsigaridis, K., Tselioudis, G., Weng, E., Wu, J., and Yao, M.-S.: CMIP6 Historical Simulations (1850–2014) With GISS-E2.1, *Journal of Advances in Modeling Earth Systems*, 13, e2019MS002034, <https://doi.org/10.1029/2019MS002034>, 2021.
- 1690 Mlawer, E. J., Payne, V. H., Moncet, J.-L., Delamere, J. S., Alvarado, M. J., and Tobin, D. C.: Development and recent evaluation of the MT_CKD model of continuum absorption, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 370, 2520–2556, <https://doi.org/10.1098/rsta.2011.0295>, 2012.
- Munk, W. and Wunsch, C.: Abyssal recipes II: energetics of tidal and wind mixing, *Deep Sea Research Part I: Oceanographic Research Papers*, 45, 1977–2010, [https://doi.org/10.1016/S0967-0637\(98\)00070-3](https://doi.org/10.1016/S0967-0637(98)00070-3), 1998.
- 1695 Nazarenko, L. S., Tausnev, N., Russell, G. L., Rind, D., Miller, R. L., Schmidt, G. A., Bauer, S. E., Kelley, M., Ruedy, R., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Del Genio, A. D., Elsaesser, G. S., Faluvegi, G., Kiang, N. Y., Kim, D., Lacis, A. A., Leboissetier, A., LeGrande, A. N., Lo, K. K., Marshall, J., Matthews, E. E., McDermid, S., Mezuman, K., Murray, L. T., Oinas, V., Orbe, C., García-Pando, C. P., Perlwitz, J. P., Puma, M. J., Romanou, A., Shindell, D. T., Sun, S., Tsigaridis, K., Tselioudis, G., Weng, E., Wu, J., and Yao, M.-S.: Future Climate Change Under SSP Emission Scenarios With GISS-E2.1, *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002871, <https://doi.org/10.1029/2021MS002871>, 2022.
- 1700 Needham, D. H. and Kring, D. A.: Lunar volcanism produced a transient atmosphere around the ancient Moon, *Earth and Planetary Science Letters*, 478, 175–178, <https://doi.org/10.1016/j.epsl.2017.09.002>, 2017.
- Phillips, N. A.: The general circulation of the atmosphere: A numerical experiment, *Quarterly Journal of the Royal Meteorological Society*, 82, 123–164, <https://doi.org/10.1002/qj.49708235202>, 1956.
- 1705 Pierrehumbert, R. T.: A PALETTE OF CLIMATES FOR GLIESE 581g, *The Astrophysical Journal Letters*, 726, L8, <https://doi.org/10.1088/2041-8205/726/1/L8>, 2010.
- Prather, M. J.: Numerical advection by conservation of second-order moments, *Journal of Geophysical Research: Atmospheres*, 91, 6671–6681, <https://doi.org/10.1029/JD091iD06p06671>, 1986.
- Rencurrel, M. C. and Rose, B. E. J.: Exploring the Climatic Response to Wide Variations in Ocean Heat Transport on an Aquaplanet, *Journal of Climate*, 31, 6299–6318, <https://doi.org/10.1175/JCLI-D-17-0856.1>, 2018.
- 1710 Rind, D. and Lerner, J.: Use of on-line tracers as a diagnostic tool in general circulation model development: 1. Horizontal and vertical transport in the troposphere, *Journal of Geophysical Research: Atmospheres*, 101, 12667–12683, <https://doi.org/10.1029/96JD00551>, 1996.
- Rind, D., Orbe, C., Jonas, J., Nazarenko, L., Zhou, T., Kelley, M., Lacis, A., Shindell, D., Faluvegi, G., Romanou, A., Russell, G., Tausnev, N., Bauer, M., and Schmidt, G.: GISS Model E2.2: A Climate Model Optimized for the Middle Atmosphere—

- 1715 Model Structure, Climatology, Variability, and Climate Sensitivity, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032204, <https://doi.org/10.1029/2019JD032204>, 2020.

ROCKE-3D compilers and libraries: <https://tinyurl.com/ROCKE3D-compilerslibraries>, last access: 25 July 2022.

ROCKE-3D installation: <https://tinyurl.com/ROCKE3D-install>, last access: 25 July 2022.

ROCKE-3D spectral files: <https://tinyurl.com/ROCKE3D-spectralfiles>.
- 1720 ROCKE-3D tutorial slides: <https://tinyurl.com/ROCKE3D-tutorialslides>, last access: 25 July 2022.

ROCKE-3D tutorial video: <https://tinyurl.com/ROCKE3D-tutorialvideo>, last access: 25 July 2022.

Romanou, A., Rind, D., Jonas, J., Miller, R., Kelley, M., Russell, G., Orbe, C., Nazarenko, L., Latto, R., and Schmidt, G. A.: Stochastic Bifurcation of the North Atlantic Circulation under a Midrange Future Climate Scenario with the NASA-GISS ModelE, *Journal of Climate*, 36, 6141–6161, <https://doi.org/10.1175/JCLI-D-22-0536.1>, 2023.
- 1725 Rose, B. E. J.: Stable “Waterbelt” climates controlled by tropical ocean heat transport: A nonlinear coupled climate mechanism of relevance to Snowball Earth, *Journal of Geophysical Research: Atmospheres*, 120, 1404–1423, <https://doi.org/10.1002/2014JD022659>, 2015.

Rose, B. E. J., Ferreira, D., and Marshall, J.: The Role of Oceans and Sea Ice in Abrupt Transitions between Multiple Climate States, *Journal of Climate*, 26, 2862–2879, <https://doi.org/10.1175/JCLI-D-12-00175.1>, 2013.
- 1730 Rosenzweig, C. and Abramopoulos, F.: Land-Surface Model Development for the GISS GCM, *Journal of Climate*, 10, 2040–2054, [https://doi.org/10.1175/1520-0442\(1997\)010<2040:LSMDFT>2.0.CO;2](https://doi.org/10.1175/1520-0442(1997)010<2040:LSMDFT>2.0.CO;2), 1997.

Rossow, W. B., Delo, C., and Cairns, B.: Implications of the Observed Mesoscale Variations of Clouds for the Earth’s Radiation Budget, *Journal of Climate*, 15, 557–585, [https://doi.org/10.1175/1520-0442\(2002\)015<0557:IOTOMV>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<0557:IOTOMV>2.0.CO;2), 2002.
- 1735 Rugheimer, S., Kaltenegger, L., Zsom, A., Segura, A., and Sasselov, D.: Spectral Fingerprints of Earth-like Planets Around FGK Stars, *Astrobiology*, 13, 251–269, <https://doi.org/10.1089/ast.2012.0888>, 2013.

Russell, G. L., Miller, J. R., and Tsang, L.-C.: Seasonal oceanic heat transports computed from an atmospheric model, *Dynamics of Atmospheres and Oceans*, 9, 253–271, [https://doi.org/10.1016/0377-0265\(85\)90022-3](https://doi.org/10.1016/0377-0265(85)90022-3), 1985.
- 1740 Russell, G. L., Miller, J. R., and Rind, D.: A coupled atmosphere-ocean model for transient climate change studies, *Atmosphere-Ocean*, 33, 683–730, 1995.

Schmidt, F., Way, M. J., Costard, F., Bouley, S., Séjourné, A., and Aleinov, I.: Circumpolar ocean stability on Mars 3 Gy ago, *Proceedings of the National Academy of Sciences*, 119, e2112930118, <https://doi.org/10.1073/pnas.2112930118>, 2022.

Schmidt, G. A., Bitz, C. M., Mikolajewicz, U., and Tremblay, L.-B.: Ice–ocean boundary conditions for coupled models, *Ocean Modelling*, 7, 59–74, [https://doi.org/10.1016/S1463-5003\(03\)00030-1](https://doi.org/10.1016/S1463-5003(03)00030-1), 2004.
- 1745 Schmidt, G. A., Ruedy, R., Hansen, J. E., Aleinov, I., Bell, N., Bauer, M., Bauer, S., Cairns, B., Canuto, V., Cheng, Y., Del Genio, A., Faluvegi, G., Friend, A. D., Hall, T. M., Hu, Y. Y., Kelley, M., Kiang, N. Y., Koch, D., Lacis, A. A., Lerner, J., Lo, K. K., Miller, R. L., Nazarenko, L., Oinas, V., Perlwitz, J., Rind, D., Romanou, A., Russell, G. L., Sato, M., Shindell, D. T.,

- Stone, P. H., Sun, S., Tausnev, N., Thresher, D., and Yao, M. S.: Present-day atmospheric simulations using GISS ModelE: Comparison to in situ, satellite, and reanalysis data, *Journal of Climate*, 19, 153–192, <https://doi.org/10.1175/jcli3612.1>, 2006.
- 1750 Schmidt, G. A., Kelley, M., Nazarenko, L., Ruedy, R., Russell, G. L., Aleinov, I., Bauer, M., Bauer, S. E., Bhat, M. K., Bleck, R., Canuto, V., Chen, Y.-H., Cheng, Y., Clune, T. L., Del Genio, A., de Fainchtein, R., Faluvegi, G., Hansen, J. E., Healy, R. J., Kiang, N. Y., Koch, D., Lacis, A. A., LeGrande, A. N., Lerner, J., Lo, K. K., Matthews, E. E., Menon, S., Miller, R. L., Oinas, V., Oloso, A. O., Perlwitz, J. P., Puma, M. J., Putman, W. M., Rind, D., Romanou, A., Sato, M., Shindell, D. T., Sun, S., Syed, R. A., Tausnev, N., Tsigaridis, K., Unger, N., Voulgarakis, A., Yao, M.-S., and Zhang, J.: Configuration and
1755 assessment of the GISS ModelE2 contributions to the CMIP5 archive, *Journal of Advances in Modeling Earth Systems*, 6, 141–184, <https://doi.org/10.1002/2013MS000265>, 2014.
- Scott, J. R., Marotzke, J., and Adcroft, A.: Geothermal heating and its influence on the meridional overturning circulation, *Journal of Geophysical Research: Oceans*, 106, 31141–31154, <https://doi.org/10.1029/2000JC000532>, 2001.
- 1760 Séférian, R., Berthet, S., Yool, A., Palmiéri, J., Bopp, L., Tagliabue, A., Kwiatkowski, L., Aumont, O., Christian, J., Dunne, J., Gehlen, M., Ilyina, T., John, J. G., Li, H., Long, M. C., Luo, J. Y., Nakano, H., Romanou, A., Schwinger, J., Stock, C., Santana-Falcón, Y., Takano, Y., Tjiputra, J., Tsujino, H., Watanabe, M., Wu, T., Wu, F., and Yamamoto, A.: Tracking Improvement in Simulated Marine Biogeochemistry Between CMIP5 and CMIP6, *Current Climate Change Reports*, 6, 95–119, <https://doi.org/10.1007/s40641-020-00160-0>, 2020.
- 1765 Segura, A., Krelow, K., Kasting, J. F., Sommerlatt, D., Meadows, V., Crisp, D., Cohen, M., and Mlawer, E.: Ozone Concentrations and Ultraviolet Fluxes on Earth-Like Planets Around Other Stars, *Astrobiology*, 3, 689–708, <https://doi.org/10.1089/153110703322736024>, 2003.
- Segura, A., Kasting, J. F., Meadows, V., Cohen, M., Scalo, J., Crisp, D., Butler, R. A. H., and Tinetti, G.: Biosignatures from Earth-Like Planets Around M Dwarfs, *Astrobiology*, 5, 706–725, <https://doi.org/10.1089/ast.2005.5.706>, 2005.
- 1770 Sergeev, D. E., Fauchez, T. J., Turbet, M., Boutle, I. A., Tsigaridis, K., Way, M. J., Wolf, E. T., Domagal-Goldman, S. D., Forget, F., Haqq-Misra, J., Kopparapu, R. K., Lambert, F. H., Manners, J., and Mayne, N. J.: The TRAPPIST-1 Habitable Atmosphere Intercomparison (THAI). II. Moist Cases—The Two Waterworlds, *The Planetary Science Journal*, 3, 212, <https://doi.org/10.3847/PSJ/ac6cf2>, 2022.
- 1775 Sergeev, D. E., Boutle, I. A., Lambert, F. H., Mayne, N. J., Bendall, T., Kohary, K., Olivier, E., and Shipway, B.: The Impact of the Explicit Representation of Convection on the Climate of a Tidally Locked Planet in Global Stretched-mesh Simulations, *The Astrophysical Journal*, 970, 7, <https://doi.org/10.3847/1538-4357/ad4ecd>, 2024.
- Shapiro, R.: Smoothing, filtering, and boundary effects, *Reviews of Geophysics*, 8, 359–387, <https://doi.org/10.1029/RG008i002p00359>, 1970.
- Shapiro, R.: Linear filtering, *Math. Comp.*, 29, 1094–1097, <https://doi.org/10.1090/S0025-5718-1975-0389356-X>, 1975.
- 1780 Showman, A. P., Wordsworth, R. D., Merlis, T. M., and Kaspi, Y.: Atmospheric Circulation of Terrestrial Exoplanets, in: *Comparative Climatology of Terrestrial Planets*, University of Arizona Press, https://doi.org/10.2458/azu_uapress_9780816530595-ch12, 2013.
- 1785 Siebesma, A. P., Bretherton, C. S., Brown, A., Chlond, A., Cuxart, J., Duynkerke, P. G., Jiang, H., Khairoutdinov, M., Lewellen, D., Moeng, C.-H., Sanchez, E., Stevens, B., and Stevens, D. E.: A Large Eddy Simulation Intercomparison Study of Shallow Cumulus Convection, *Journal of the Atmospheric Sciences*, 60, 1201–1219, [https://doi.org/10.1175/1520-0469\(2003\)60<1201:ALESIS>2.0.CO;2](https://doi.org/10.1175/1520-0469(2003)60<1201:ALESIS>2.0.CO;2), 2003.

- 1790 Sitch, S., Friedlingstein, P., Gruber, N., Jones, S. D., Murray-Tortarolo, G., Ahlström, A., Doney, S. C., Graven, H., Heinze, C., Huntingford, C., Levis, S., Levy, P. E., Lomas, M., Poulter, B., Viovy, N., Zaehle, S., Zeng, N., Arneth, A., Bonan, G., Bopp, L., Canadell, J. G., Chevallier, F., Ciais, P., Ellis, R., Gloor, M., Peylin, P., Piao, S. L., Le Quéré, C., Smith, B., Zhu, Z., and Myneni, R.: Recent trends and drivers of regional sources and sinks of carbon dioxide, *Biogeosciences*, 12, 653–679, <https://doi.org/10.5194/bg-12-653-2015>, 2015.
- Slingo, J. M.: A study of the earth's radiation budget using a general circulation model, *Quarterly Journal of the Royal Meteorological Society*, 108, 379–405, <https://doi.org/10.1002/qj.49710845606>, 1982.
- Strom, R. G., Schaber, G. G., and Dawson, D. D.: The global resurfacing of Venus, *Journal of Geophysical Research: Planets*, 99, 10899–10926, <https://doi.org/10.1029/94JE00388>, 1994.
- 1795 Trenberth, K. E. and Caron, J. M.: Estimates of Meridional Atmosphere and Ocean Heat Transports, *Journal of Climate*, 14, 3433–3443, [https://doi.org/10.1175/1520-0442\(2001\)014<3433:EOMAAO>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<3433:EOMAAO>2.0.CO;2), 2001.
- Tsigaridis, K., Ackerman, A. S., Aleinov, I., Chandler, M. A., Clune, T. L., Colose, C. M., Del Genio, A. D., Maxwell, K., Kiang, N. Y., Leboissetier, A., Perlwitz, J. P., Ruedy, R. A., Russell, G. L., Sohl, L. E., Way, M. J., and Wolf, E. T.: ROCKE-3D v2, <https://doi.org/10.5281/zenodo.14721184>, 2025.
- 1800 Turbet, M., Boulet, C., and Karman, T.: Measurements and semi-empirical calculations of CO₂ + CH₄ and CO₂ + H₂ collision-induced absorption across a wide range of wavelengths and temperatures. Application for the prediction of early Mars surface temperature, *Icarus*, 346, 113762, <https://doi.org/10.1016/j.icarus.2020.113762>, 2020.
- 1805 Turbet, M., Fauchez, T. J., Sergeev, D. E., Boutle, I. A., Tsigaridis, K., Way, M. J., Wolf, E. T., Domagal-Goldman, S. D., Forget, F., Haqq-Misra, J., Kopparapu, R. K., Lambert, F. H., Manners, J., Mayne, N. J., and Sohl, L.: The TRAPPIST-1 Habitable Atmosphere Intercomparison (THAI). I. Dry Cases—The Fellowship of the GCMs, *The Planetary Science Journal*, 3, 211, <https://doi.org/10.3847/PSJ/ac6cf0>, 2022.
- 1810 Voigt, A., Biasutti, M., Scheff, J., Bader, J., Bordoni, S., Codron, F., Dixon, R. D., Jonas, J., Kang, S. M., Klingaman, N. P., Leung, R., Lu, J., Mapes, B., Maroon, E. A., McDermid, S., Park, J., Roehrig, R., Rose, B. E. J., Russell, G. L., Seo, J., Toniazzi, T., Wei, H.-H., Yoshimori, M., and Vargas Zeppetello, L. R.: The tropical rain belts with an annual cycle and a continent model intercomparison project: TRACMIP, *Journal of Advances in Modeling Earth Systems*, 8, 1868–1891, <https://doi.org/10.1002/2016MS000748>, 2016.
- Way, M. J. and Del Genio, A. D.: Venusian Habitable Climate Scenarios: Modeling Venus Through Time and Applications to Slowly Rotating Venus-Like Exoplanets, *Journal of Geophysical Research: Planets*, 125, e2019JE006276, <https://doi.org/10.1029/2019JE006276>, 2020.
- 1815 Way, M. J., Del Genio, A. D., Kiang, N. Y., Sohl, L. E., Grinspoon, D. H., Aleinov, I., Kelley, M., and Clune, T.: Was Venus the first habitable world of our solar system?, *Geophysical Research Letters*, 43, 8376–8383, <https://doi.org/10.1002/2016GL069790>, 2016.
- 1820 Way, M. J., Aleinov, I., Amundsen, D. S., Chandler, M. A., Clune, T. L., Del Genio, A. D., Fujii, Y., Kelley, M., Kiang, N. Y., Sohl, L., and Tsigaridis, K.: Resolving Orbital and Climate Keys of Earth and Extraterrestrial Environments with Dynamics (ROCKE-3D) 1.0: A General Circulation Model for Simulating the Climates of Rocky Planets, *The Astrophysical Journal Supplement Series*, 231, 12, 2017.
- Winton, M.: On the Climatic Impact of Ocean Circulation, *Journal of Climate*, 16, 2875–2889, [https://doi.org/10.1175/1520-0442\(2003\)016<2875:OTCIOO>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<2875:OTCIOO>2.0.CO;2), 2003.

- 1825 Wolf, E. T., Kopparapu, R., Haqq-Misra, J., and Fauchez, T. J.: ExoCAM: A 3D Climate Model for Exoplanet Atmospheres, The Planetary Science Journal, 3, 7, <https://doi.org/10.3847/PSJ/ac3f3d>, 2022.
- Wood, R. A.: Time Step Sensitivity and Accelerated Spinup of an Ocean GCM with a Complex Mixing Scheme, Journal of Atmospheric and Oceanic Technology, 15, 482–495, [https://doi.org/10.1175/1520-0426\(1998\)015<0482:TSSAAS>2.0.CO;2](https://doi.org/10.1175/1520-0426(1998)015<0482:TSSAAS>2.0.CO;2), 1998.
- 1830 Wordsworth, R., Kalugina, Y., Lokshtanov, S., Vigasin, A., Ehlmann, B., Head, J., Sanders, C., and Wang, H.: Transient reducing greenhouse warming on early Mars, Geophysical Research Letters, 44, 665–671, <https://doi.org/10.1002/2016GL071766>, 2017.
- Wunsch, C. and Ferrari, R.: Vertical mixing, energy, and the general circulation of the oceans, Annual Review of Fluid Mechanics, 36, 281–314, <https://doi.org/10.1146/annurev.fluid.36.050802.122121>, 2004.