



Effect of trachyte and basalt powder on the growth and yield of maize (*Zea Mays* L.) in the Sudano-Sahelian zone of Cameroon (Central Africa)

Bienvenu Sidsi^{1,3}, Claudine Vounba¹, Simon Djakba Basga², Aubin Nzeugang Nzeukou³, Merlin Gountie Dedzo⁴, Désiré Tsozué^{3*}

¹ Institute of Agricultural Research for Development (IRAD), P.O. Box 33 Maroua, Cameroon

² Institute of Agricultural Research for Development (IRAD), P.O. Box 2123 Yaoundé, Cameroon

³ Department of Earth Sciences, Faculty of Science, University of Maroua, P.O. Box 814 Maroua, Cameroon

⁴ Department of Life and Earth Sciences, Higher Teachers' Training College, University of Maroua, P.O. Box. 55, Maroua, Cameroon

Correspondence to: Désiré Tsozué (tsozudsir@yahoo.fr)

Abstract. The Sudano-Sahelian zone of Cameroon, characterized by a low annual rainfall, faces challenges in soil fertility preservation due to agricultural intensification and unsustainable practices. This study aims to evaluate the effect of trachyte and basalt powders inputs on soil and maize yield in Guiring experimental farm. Fieldwork involved collecting and describing samples of trachyte, basalt, and soil and setting up the experimental design. In the laboratory, the ground rock samples underwent geochemical analysis, and the soil samples were analysed for their mineralogical and physicochemical properties. The experiment followed a completely randomized block design with three repetitions and six treatments (T0, T1, T2, T3, T4 and T5). The soil consists of kaolinite, smectite, sepiolite, and quartz. Its texture is dominated by sand fraction, with a neutral pH (6.98). The organic matter (1.30 to 3.17%) and total nitrogen contents (0.11 to 0.13%) are relatively low. The concentrations of potassium, magnesium, sodium, and calcium vary from 0.10 to 0.40 cmol_c kg⁻¹, 0.72 to 5.44 cmol_c kg⁻¹, 0.13 to 0.56 cmol_c kg⁻¹, and 2.64 to 6 cmol_c kg⁻¹, respectively. The cation exchange capacity is moderate to high, ranging from 18.70 to 25 cmol_c kg⁻¹, while the available phosphorus content is high, ranging from 12.60 to 30.30 mg kg⁻¹. The studied soils are moderately suitable for maize cultivation. Fertilization trials showed a significant improvement in maize growth and yield, within plots treated with basalt powder yielding higher (2558.64 kg ha⁻¹ and 2931.16 kg ha⁻¹) than those treated with trachyte powder (2362.87 kg ha⁻¹ and 2763.91 kg ha⁻¹) and the control plots (645.83 kg ha⁻¹). Plots treated with NPK fertilizer recorded the highest yield (3164.45 kg ha⁻¹). Although the treatment with conventional fertiliser resulted in a relative higher yield, the advantage of using rock powders lies in their environmental benefits, long-term effectiveness, and more affordable cost.



1 Introduction

In recent decades, population growth and food insecurity have led to intensification of agriculture, soil overexploitation, and use of chemical fertilizers (Pham et al., 2018; Jharia et al., 2021; Khatri et al., 2024). Intensively exploited soils degrade and impoverish rapidly, resulting in a reduction of biological activity, soil acidification and deterioration of soil structure (Kuria et al., 2018; Nanganoa et al., 2019; Rajwar et al., 2021). As a limited resource, it takes between 200 to 1000 years to form a layer of 2.5 cm in thickness, making its exploitation for agricultural purposes fragile due to the increase in the world population and climate change (Moges and Taye, 2016). Soil degradation has become a major global concern due to its harmful effects on biodiversity, agricultural production, and food security (Mekuriaw et al., 2017; Hossain et al., 2020). To maintain soil fertility and productivity, it is necessary to replenish the supply of nutrients (Nkouathio et al., 2008; Garai et al., 2022).

Soil provides physical stability, support for agriculture and contributes to the nutrient cycle, water retention capacity, storage, filtering, buffering, and transformation of compounds (Tsozué et al., 2020a; Toor et al., 2021; Sivaram et al., 2023). Soil health, also known as soil quality, is a key factor in sustainable agriculture, influencing ecosystem quality, such as air and water quality (Are et al., 2018; Tsozué et al., 2020a). Maintaining soil quality is an essential determinant for managing ecosystem services and sustainable soil management (Tsozué et al., 2020a; Wu et al., 2023). Its restoration and assessment require the use of relevant biological, physical, and/or chemical indicators (Tsozué et al., 2020a; Oliveira et al., 2024). It is important to consider current land use and precisely define the type of disturbance that may alter its quality (Peng et al., 2021). In the case of anthropogenic disturbances, land use, including agricultural management systems, strongly influences soil quality, expressed by changes in its physicochemical and biological properties and agricultural yield (Farahani et al., 2019; Tsozué et al., 2020a; Mamabolo et al., 2024).

Intensive agriculture leads to excessive nutrient depletion from the soil, thereby compromising its fertility on a global scale (Padhiary et al., 2024). While nitrogen (N) and phosphorus (P) are often highlighted, potassium (K) also plays a crucial role, and its depletion is becoming an increasing concern (Khan et al., 2023). To address this issue, the use of rock powders has emerged as a sustainable alternative to chemical fertilizers. These natural fertilizers enrich the soil with essential nutrients and enhance its structure, although their performance depends on soil characteristics and climatic conditions (Manning and Theodoro, 2020; Rodrigues et al., 2024b). Numerous studies have explored the use of rock powders for soil fertilization and remineralization. In the Americas, several studies have demonstrated that these mineral powders enhance water retention and soil fertility while contributing to carbon sequestration (Beerling et al., 2018; Theodoro et al., 2021; Lewis et al., 2021; Ramos et al., 2022; Medeiros et al., 2023). In Europe, research has confirmed their positive impact on improving soil fertility and remineralization (Beerling et al., 2018; Vienne et al., 2022). Similarly, in Asia, silicate rocks have shown their ability to increase agricultural productivity (Xiang et al., 2020; Basak et al., 2023; Sniatala et al., 2023). Africa has also embraced this innovation, using rock powders to boost agricultural yields while preserving the environment (Mekuriaw et al., 2017; Muhie, 2022). In Cameroon, in particular, several studies have demonstrated the effectiveness of these powders (Nkouathio et al.,



2008; Mein'da et al., 2022; Tetchou et al., 2022). Although crushed rocks offer agronomic and environmental benefits, some limitations remain, including their slow dissolution rate, high logistical costs, and the need for pedological expertise to maximize their effectiveness. Given the growing concerns over soil degradation and nutrient depletion, it becomes crucial to explore sustainable solutions, particularly in regions like the Sudano-Sahelian zone of Cameroon, where challenging climatic conditions contribute to the deterioration of soil fertility.

The Sudano-Sahelian zone of Cameroon, characterized by a low precipitation and a prolonged dry season, faces challenges in preserving soil fertility. Inadequate agricultural practices and pressure on the soil have led to its degradation, resulting in a decline in fertility and agricultural yield (Prasannakumar et al., 2011; Hishe et al., 2017). According to FAO (2013), global food demand is expected to increase by 60% between 2006 and 2050, making imperative, if not essential, to modify or replace agricultural techniques with more sustainable and environmentally friendly practices to preserve the integrity of various environmental components and ensure long-term agricultural sustainability (Nkouathio et al., 2008; Muhie, 2022). One of these techniques involves effectively regenerating soil fertility using appropriate methods that do not impact the environment, including the use of geological materials (Burbano et al., 2022). As the minerals constituting rocks dissolve, the released chemical elements become available to plants (Silva et al., 2005; Manning and Theodoro, 2020; Basak et al., 2023). To address the challenges of soil fertility decline and rising fertilizer costs in the Sudano-Sahelian zone, exploring alternative solutions such as local rock powders becomes essential for sustainable and resilient agriculture.

The rising cost of fertilizers represents a major constraint for farmers, especially in the Sudano-Sahelian zone of Cameroon, limiting their access to these essential inputs (Sinha et al., 2022). This situation has raised global concerns about food shortages, forcing countries reliant on fertilizer imports to find mechanisms and new technological pathways to reduce their dependence on the international market (Xiang et al., 2020; Sniatala et al., 2023). The use of **local rock powders, or remineralizers**, appears to be a promising alternative (Dalmora et al., 2020; Swoboda et al., 2023). This agroecological approach not only reduces reliance on chemical fertilizers but also sustainably improves soil fertility while being more economical and environmentally friendly (Ramos et al., 2017; Conceição et al., 2020; Swoboda et al., 2022). In light of the challenges posed by increasing fertilizer costs and the need for sustainable agricultural practices, exploring the potential of local geological materials like trachyte and basalt powders to enhance soil fertility and maize production in Cameroon has become a vital research focus.

Maize, one of the most produced and consumed cereals in the world, with about 850 million tons over approximately 162 million hectares, is primarily used for food but also serves for the production of biofuels and the extraction of starch, oil, and other industrial substances. In the northern regions of Cameroon in particular, maize is a staple food in almost every household. In a context marked by soil degradation, the environmental impacts of intensive chemical fertilizers use, and rising input costs, it is essential to assess to what extent the use of local available geological materials such as trachyte and basalt powders can improve soil fertility, promote maize growth and yield, while fostering sustainable and resilient agriculture in the face of climate change. This issue is the focus of studies conducted at the experimental farm of the Agricultural Research Institute for Development (IRAD) in Guiring.



100

2 Materials and methods

2.1 Study area

This study was conducted in the locality of Guiring, situated in the Diamaré Division (Maroua III Subdivision) in the Far-North Region of Cameroon, located between 10°32'21.8" and 10°41'55.7" North latitude and 14°15'33.8" and 14°29'42.0" East longitude (Fig. 1.). The area has a Sudano-Sahelian climate (Suchel, 1987), with a short and irregularly distributed rainy season from June to September, followed by a long dry season lasting from October to May. The average annual rainfall is approximately 792.25 mm, and the average annual temperature reaches 27.7°C (Table 1). According to De Martonne's aridity index (1926), the region experiences four months of rainy season (June to September, with an aridity index > 20) and eight months of dry season (aridity index < 20) (Table 1). The highest temperatures are recorded in April, while the lowest occur in January. The terrain is mainly characterized by a relatively flat topography, marked by several ecological zones, including the Sahelo-Sudanian domain, the elevated Sudanian domain, and the periodically flooded Sahelian domain. The substratum consists of alluvial deposits of Quaternary age (Tament et al., 2015; Gountié et al., 2019). The vegetation consists of scattered trees and shrubs associated with grasses. Among the most representative species are *Acacia seyal*, *Balanites aegyptiaca*, *Faidherbia albida*, and *Ziziphus mauritiana* (Letouzey, 1980).




Fig. 1. (a) Location of Cameroon in Africa, (b) Location of Far-North region of Cameroon, (c) Location of the study area.


Table 1. Monthly averages of precipitation, temperature, and relative humidity in Maroua-Salack from 1980 to 2020.

Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Mean	Total
Precipitation (mm)	0	0	0.87	32.10	60.54	105.77	192.77	232.41	118.20	48.79	0.49	0.31	/	792.25
Temperature (°C)	24.20	27.50	30.90	32.90	31.50	28.90	26.50	25.20	25.90	27.50	27.00	24.40	27.70	/
Relative humidity (%)	19	14	11	20	40	60	73	81	79	56	27	23	42	/
Aridity index	0.00	0.00	0.26	8.97	17.49	31.81	63.38	79.22	30.90	15.61	0.16	0.10	20.65	/

2.2 Experimental design, treatments, and plant material

The experimental design is a completely randomized block design with five (05) treatments (T_1 , T_2 , T_3 , T_4 , T_5) in addition to the control (T_0) as described in table 2 and **three (03)** replications, resulting in twenty four (24) experimental plots (Fig. 2.). It was conducted at the experimental site of IRAD in Guiring. Each experimental unit received one of the recommended treatments, T_0 , T_1 , T_2 , T_3 , T_4 , and T_5 (Table 2). The maize density was 0.80×0.25 m, with 2 to 3 seeds needed per hole. Two weeks after sowing, thinning was performed to leave one plant per cluster. The plant material used is the CMS-9015 variety of maize, developed by IRAD. This variety is specifically adapted to the Sudanese-Sahelian zone of Cameroon. Under normal conditions, it produces grain yields ranging from 4 to 4.5 tons per hectare, with a vegetative cycle of 95 days.

Treatment designation	Description of the treatment
T₀	Control plot without fertilizing inputs
T₁	Treatment with trachyte powder (2000 kg ha ⁻¹ ; 1.96 kg/9.6m ²)
T₂	Treatment with trachyte powder (2000 kg ha ⁻¹ ; 1.96 kg/9.6m ²) + urea (104,17 kg ha ⁻¹ ; 100 g/9.6m ²) at 06 weeks
T₃	Treatment with basalt powder (2000 kg ha ⁻¹ ; 1.96 kg/9.6m ²)
T₄	Treatment with basalt powder (2000 kg ha ⁻¹ ; 1.96 kg/9.6m ²) + urea (104,17 kg ha ⁻¹ ; 100 g/9.6m ²) at 06 weeks
T₅	Treatment with NPK fertilizer (208,33 kg ha ⁻¹ ; 200 g/9.6m ²) + urea (104,17 kg ha ⁻¹ ; 100 g/9.6m ²) at 06 weeks






grinding, thin section preparation, petrographic analyses, and geochemical analyses. In total, 3 samples of trachyte and 3 samples of basalts were subjected to petrographic and geochemical analysis in the laboratory.

The rock powders were obtained from a commercial quarry located in the Meri municipality, Douvagar locality, in the Farth-
135 Nord region of Cameroon. The rock was fragmented and passed through a ball mill, then sieved using a 2 mm mesh to obtain
powder less than 2 mm (Figs. 3 c and d). In total, 100 kg of trachyte powder and 100 kg of basalt powder were obtained.

The petrographic analysis was carried out on thin sections prepared at the Tokyo Institute of Technology, Japan. The
microscopic description was done in the geology laboratory at the University of Maroua using an optic microscope to
examine the rock samples in detail. This phase allowed for the precise identification of the different minerals present in each
140 sample, based on their optical and textural characteristics. The observations focused on mineralogical assemblages, grain
shapes and the relationships between the various minerals.

For the measurements of trace and major elements, all samples were crushed into millimetre-sized fragments using a jaw
crusher, and fragments with saw and hammer marks were carefully separated. After each sample, the jaw crusher was
systematically cleaned with compressed air and deionized water. The major elements and some trace elements, including Sc,
145 Co, V, Ni, and Cr, were analysed by X-ray fluorescence (XRF) using a Rigaku RIX2100 at Osaka City University, Graduate
School of Science. The trace element concentrations in the samples were determined by ICP-MS at the Tokyo Institute of
Technology, Japan, in accordance with the method described by Yokoyama et al. (1999). Approximately 50 mg of rock
powder were digested with HNO₃, HClO₄, HCl, and HF. Before acid digestion, enriched spikes of ¹¹³In and ²⁰³Tl were
added to the sample to serve as internal standards during measurement. The data reported by Makishima and Nakamura
150 (2006) were used to calibrate the trace element concentrations against the basalt and rhyolite standards JB-3 and JR-1,
respectively. The typical analytical reproducibility (2σ) was 4% for Nb and Pb, 5% for Y and Ta, and <3% for other trace
elements.



155 **Fig. 3. Samples and powders of trachyte and basalt:** (a) Trachyte sample from Zamai, (b) Basalt sample from Gawar, (c) Trachyte
 160 powder obtained after grinding, (d) Basalt powder obtained after grinding.

2.4 Soil sampling and laboratory analysis

The fieldwork was carried out in two phases. The first phase consisted of direct observation and a detailed description of the study site and its surroundings. In the second phase, a representative soil profile was excavated and described to facilitate soil classification. The soil description took into account several parameters: colour, porosity, texture, coarse elements,



structure, biological activity, thickness, and the boundaries between horizons. The colour was determined in its dry state using the Munsell Soil Colour Charts (2009). A total of six (06) samples, each weighing approximately 0.5 kg, were collected from the different horizons of the profile. The collected samples were placed in polyethylene bags, carefully labelled, and prepared for laboratory analysis.

165 In the laboratory, the main analyses focused on mineralogical and physicochemical analyses. Particles size distribution was performed using the Robinson pipette method on air-dried, ground, and 2 mm-sieved samples. The pH was measured in a soil-water suspension, then in a normal potassium chloride solution at 1:2.5 ratio using a glass electrode pH meter. Exchangeable bases (Ca^{2+} , Mg^{2+} , K^{+} , and Na^{+}) were extracted from the soil using a pH 7 ammonium acetate solution and measured by atomic absorption spectrophotometry. The sum of exchangeable bases (SBE) was determined by adding the
170 concentrations of Ca^{2+} , Mg^{2+} , K^{+} , and Na^{+} . Cation exchange capacity (CEC) was measured using an ammonium acetate solution at pH 7, following a three-step process: washing the soil with alcohol to remove the ammonium solution saturating the pores, determining NH_4^{+} by Kjeldahl distillation after quantitative desorption with KCl. The CEC of the clay fraction was calculated from the CEC at pH 7, organic carbon content, and clay percentage. Base saturation (V) was calculated using the formula $(\text{SBE}/\text{CEC}) \times 100$. Total nitrogen (N) was determined by the Kjeldahl method. Organic carbon (OC) was measured
175 by wet oxidation according to Walkley and Black (1934) using a sodium dichromate and concentrated H_2SO_4 mixture. Organic matter (OM) content was calculated by multiplying the OC by 1.724, and the C/N ratio was determined. Available phosphorus was measured using the Bray no. 2 method (Bray and Kurtz, 1945).

Mineralogical analysis was carried out using infrared (IR) spectroscopy, a non-destructive analytical method widely used to identify and characterize the functional groups present in samples. This technique provides detailed information about the
180 chemical bonds and molecular structures of minerals. Diffuse reflectance IR spectra were recorded in the range of 4000 to 400 cm^{-1} using a Perkin Elmer 2000 FTIR spectrometer (Perkin Elmer, Waltham, MA, USA), equipped with a DTGS (deuterated triglycine sulphate) detector, optimized for high sensitivity and precise resolution. To avoid contamination from atmospheric moisture and ensure the reliability of the results, measurements were carried out under strictly controlled environmental conditions. This method not only allows for the identification of minerals present in the soil but also helps
185 describe their local chemical environments, providing a deeper understanding of the composition and properties of soils.

2.5 Pedoclimatic Assessment

The climatic and pedological assessment of the study area was based on climatic data (precipitation, temperature, relative humidity, and insolation) collected at the Maroua-Salak station between 1980 and 2020, in accordance with the climatic requirements of crops (Sys, 1985; Sys et al., 1993; Issiné et al 2022). A climatic index (CI) was calculated using the
190 parametric formula:

$$I_C = R_{\min} \times \sqrt{\frac{A}{100} \times \frac{B}{100} \times \dots},$$



where R_{min} is the smallest value, A , B , ... are the other parametric values. Adjustments are made if the CI falls between 25 and 92.5.

The pedological assessment used a table of soil characteristics to determine a land quality index (LI), based on similar parametric values using the formula:

$$I_T = R_{min} \times \sqrt{\frac{A}{100} \times \frac{B}{100} \times \dots},$$

The calculation of the CI and LI allows for an integrated assessment of agroclimatic and soil quality for agricultural production.

2.6 Collection of maize parameters

The growth and yield parameters of maize, including germination rate, plant height, number of leaves per plant, stem diameter, ear length, ear diameter, ear weight, 100-grain weight, and grain yield (kg ha^{-1}), were measured.

The germination rate (GR), assessed 10 days after sowing, is determined using the formula $\text{GR} = \text{number of germinated seeds} \times 100 / \text{total number of seeds sown}$, then multiplying the result by 100 to obtain the germination percentage (Ellis et al., 1986). Plant height was measured using a tape measure, recording the distance from the soil to the top of the stem, with measurements taken every 21 days. These measurements were performed on a representative sample from each plot an average value was considered (Cox and Andrade, 1988). The number of leaves was counted by tallying the leaves deployed on the main stem, with an average estimation made every 3 weeks (Cox and Andrade, 1988). The stem and ear diameter were measured using a calliper respectively before and after harvest (Ngoune Tandzi et al., 2019). The ear length, measured from base to tip, was recorded with a tape measure when the ears were mature. The weight of the ears was obtained by weighing them with a high-precision electronic scale after harvest (Muthaura et al., 2017). The weight of 100 randomly selected dried maize grains was also measured using a precision electronic scale (Soleymani, 2018). The yield per hectare was estimated by harvesting a representative portion of each plot, weighing the harvest, and then extrapolating the results to the hectare scale using a conversion factor (Soleymani, 2018).

2.7 Data Analysis

Statistical analyses focused on the physicochemical properties of soils as well as the growth and yield parameters of maize. The physicochemical properties of soils were subjected to standard statistical analyses, including mean, maximum-minimum values, standard deviation, and coefficient of variation. The Anderson-Darling normality test was used to assess the normality of the distribution. Correlation analysis to evaluate relationships between soil parameters was conducted using Spearman's correlation test. A descriptive analysis of growth and yield parameters of maize was performed to summarize and describe the collected agricultural data using measures of central tendency (mean, median) and dispersion (standard deviation). These analyses were conducted using XLSTAT 2023 version 25.1.141.



3 Results

3.1 Petrographic and geochemical characteristics of rocks

Petrographically, trachyte outcrop in the form of slabs in the Zamai locality and exhibit an aphyric texture, with rare
 225 phenocrysts ($\leq 5\%$). These scarce phenocrysts are mainly composed of sanidine, sometimes associated with clinopyroxene.
 Sanidine, characterized by elongated, euhedral prisms reaching up to 0.6×1.8 mm, displays simple Carlsbad twins. The
 trachytes **are composed of feldspar microliths**, accompanied by a low proportion of clinopyroxene and oxides (Figs. 4 a and
 b). Basalts outcrop as lava flows, exhibiting well-defined columnar joints and a porphyritic texture. Plagioclase, olivine, and
 clinopyroxene constitute the phenocrysts (Figs. 4 c and d). Olivine, present in some samples, is often altered to iddingsite.
 230 Peridotite xenoliths are also observed in some samples, containing approximately 65% olivine and 35% pyroxene.
 Plagioclase microliths, associated with clinopyroxene, olivine, and oxides, are the main components of the basalt matrix.






Fig. 4. Photomicrographs of thin sections of representative samples from the study area. **(a)** and **(b)** Aphyric texture of Minawao trachyte showing mineral assemblages. **(c)** and **(d)** Porphyritic texture of Gawar basalt showing mineral assemblages. Ol: olivine. CPx: clinopyroxene. Pl: plagioclase. Opq: opaque minerals. Sa: sanidine. Amp: amphibole. Mineral abbreviations from Whitney and Evans (2010).

From a geochemical perspective, the concentration of major and trace elements in the studied rocks are summarized in Table 3. Trachyte exhibit variations in major element contents across samples. The SiO₂ content ranges between 65.31% and 66.5%. Meanwhile, the TiO₂ content varies from 0.16% to 0.25%. The Al₂O₃ content remains constant at 16.3%, whereas Fe₂O₃ concentrations fluctuate between 4.432 and 5.6%. The contents of CaO and K₂O show variations ranging from 0.63 to 1.12% and 4.66 to 4.82%, respectively. Regarding basalts, notable variations are also observed in the proportions of major elements. The SiO₂ content ranges from 40.86% to 41.34%, while TiO₂ content is between 4.22% and 4.26%. Similarly, the percentage of Al₂O₃ fluctuates between 13.17% and 13.43%. Fe₂O₃ concentrations vary slightly, ranging from 15.33% to 15.44%, while CaO content is between 10.14% and 10.42%, and K₂O ranges from 1.39% to 1.69%. These subtle yet significant differences highlight the important geochemical variations between trachyte and basalt samples, underscoring the diversity of their mineralogical composition.

Trace element concentrations in the trachyte show significant fluctuation. The chromium (Cr) content remains relatively constant at 2 ppm, while other elements such as nickel (Ni), strontium (Sr), rubidium (Rb), yttrium (Y), barium (Ba), and niobium (Nb) exhibit more pronounced variations, ranging from 7.2 to 9.1 ppm, 7.3 to 20.91 ppm, 136.19 to 148.82 ppm, 60.37 to 97.05 ppm, 16.2 to 48.93 ppm, and 194.15 to 197.25 ppm, respectively. In contrast, basalts show much higher concentrations of Cr (176.1 to 183 ppm), Ni (269.6 to 277.7 ppm), Sr (1648.5 to 1685.5 ppm), and Ba (561.1 to 572.7 ppm). However, the basalts have relatively low levels of Rb (18.9 to 23.5 ppm) and Y (22.9 to 23.2 ppm) compared to the trachyte (Table 3). These variations indicate substantial differences in trace elements between trachytes and basalts, reflecting distinct geochemical signatures.

Table 3. Major (wt.%) and trace (mg kg⁻¹) elements data for the Gawar-Zamay lavas.

2022	Basalte			Trachyte		
	Gaw1	Gaw2	Gaw3	Zam1	Zam2	Zam3
Longitude	13°49'57.4"E	13°49'55.8"E	13°49'58.2"E	13°53'15.9"E	13°53'17.2"E	13°53'15.7"E
Latitude	10°30'39.1"N	10°30'37.3"N	10°30'38.6"N	10°36'25.5"N	10°36'26.8"N	10°36'26.0"N
Altitude (m.a.s.l)	754	751	753	602	604	605
(wt%)						
SiO ₂	41.34	41.28	40.86	66.5	66.5	65.31
TiO ₂	4.26	4.25	4.22	0.16	0.16	0.25
Al ₂ O ₃	13.43	13.39	13.17	16.3	16.3	16.3
Fe ₂ O ₃	15.33	15.34	15.44	4.43	4.43	5.6
MnO	0.18	0.18	0.18	0.13	0.14	0.17
MgO	8.5	8.53	8.66	0.04	0.02	0.04
CaO	10.42	10.5	10.14	0.68	0.68	1.12
Na ₂ O	3.46	3.21	3.42	7.07	7.09	6.1
K ₂ O	1.39	1.69	1.67	4.66	4.82	4.82



P2O5	1.13	1.12	1.12	0.01	0.02	0.02
Total	99.41	99.48	98.87	99.93	99	99
Mg	5.5	55.6	5,8			
(mg kg ⁻¹)						
Sc	24	21.6	23.5	3	b.d.l.	0.9
V	224.1	222.1	223	b.d.l.	b.d.l.	b.d.l.
Cr	178	176.1	183	2	2	2
Co	60.3	59.8	62.1	7.9	7.2	9.1
Ni	271	269.6	271.7	2.4	2.6	1.4
Ga	17	18.1	17.8	39.9	41.7	36.5
Zn	107.7	107.8	104.1	247.6	250.5	208.5
Sr	1668.6	1685.5	1648.5	10.8	7.13	20.91
Rb	18.9	23.5	22.2	148.33	148.82	136.19
Y	23.2	23	22.9	97.05	91.17	60.37
Cs	0.2	0.2	0.2	0.8	0.79	0.3
Ba	572.7	561.1	562.5	48.34	28.93	16.12
La	49.2	48.3	48.1	168.45	160.65	184.76
Ce	106.7	104.9	104.7	420.27	413.43	477.2
Pr	13	12.9	12.8	34.65	33.4	38.1
Nd	56.9	55.9	55.8	125.43	120.77	141.43
Sm	11	10.8	10.8	21.74	21.15	22.02
Eu	3.8	3.7	3.7	1.91	1.84	1.87
Gd	10.7	10.5	10.4	22.93	22.11	22.56
Tb	1.3	1.3	1.3	3	2.89	2.56
Dy	6.4	6.3	6.3	16.48	15.78	12.63
Ho	1.1	1.1	1.1	3.12	2.98	2.28
Er	2.7	2.6	2.6	8.4	8.06	5.98
Tm	0.3	0.3	0.3	1.12	1.08	0.76
Yb	1.9	1.8	1.8	709	6.88	4.95
Lu	0.3	0.3	0.3	1	0.97	0.74
Hf	3.33	3.57	2.94	5.39	33.46	33.35
Ta	3.45	3.48	3.42	1.85	10.75	10.56
Pb	1.8	1.7	1.7	10.51	10.42	7.78
Th	3.8	3.7	3.7	18.61	18.32	12.11
U	1.1	1.1	1.1	4.96	4.91	1.55
Zr	143.8	147.73	125.7	216	1333.53	1314.17
Nb	67.09	67.67	67.24	33.57	197.25	194.15

3.2 Morphological and mineralogical characteristics of the soil

The studied soil, located at latitude 10°33'17" North and longitude 14°22'16" East, and at of 386 m, has a thickness of 270 cm. The relief is flat and the vegetation highly anthropogenized.

260 The studied soil profile consists of six horizons (Fig. 5.) divided into four distinct textural groups: sandy loam, loamy sand, loamy clayey sand, clayey sand and sandy. The colours range from light yellow-brown (10YR 5/4) to dark brown (7.5YR 3/2), passing through light yellow (10YR 5/4) and up to pale yellow (2.5Y 7/4). Each horizon has a fine, granular structure

with significant biological and matrix porosity, promoting good permeability and optimal friability. The transitions between horizons are progressive and regular (Table 4).




Fig. 5. Macroscopic organization of the soil in the study area.



Table 4. Morphological and physical characteristics of the soil profile.

Site	Depth(cm)	Colour (Moist)	Structure	Consistency		Rock fragments	Boundary	Textural class
				Dry	Wet			
Guiring	0-25	10YR 4/3	f&wg	s	s & p	n	g	sandy loam
	25-50	7.5YR 3/2	f&wm	s	s & p	c	g	loamy sand
	50-110	10YR 5/4	m&mma	h	s & p	m	g	loamy clayey sand
	110-160	2.5Y 7/4	c&sabk	l	s & p	d	c	sandy
	160-220	10YR 2/1	m&sma	h	s & p	m	c	loamy clayey sand
	220-270	2.5Y 7/4	c&sabk	l	s & p	d	g	sandy
Soil characteristics								
Structure				Consistency		Rock fragments	Boundary	
Size	Type	Grade		Dry	Wet			
vf = very fine (G5 mm)	g = granular	w = weak (peds barely observable)		l = loose	s = sticky	n = none (0%)		g = gradual
f = fine (5–10 mm)	abk = angular	m = moderate (peds observable)		s = soft	p = plastic	m = many (15%–40%)		c = clear
m = medium (10–20 mm)	blocky	s = strong (peds clearly observable)		h = hard		v = very few (0%–2%)		a = abundant (40%–80%)
c = coarse (20–50 mm)	blocky					c = common (5%–15%)		d = dominant (>80%)
vc = very coarse (>50 mm)	l=lumpy							
	ma=massive							

Mineralogically, the infrared spectra of the soil, illustrated in Fig. 6, reveal the presence of kaolinite, smectites, sepiolite, and quartz. In the high-frequency region, the hydroxyl groups display two bands at $3774 - 3622 \text{ cm}^{-1}$, corresponding to the stretching vibrations of the OH^- groups characteristic of kaolinite according to Famer (1964) and Frost et al. (1998). The low intensity absorption band located at 1621 cm^{-1} is attributed to water molecules, possibly associated with smectites. Sepiolite is primarily distinguished by the band at 531 cm^{-1} . The presence of quartz is manifested by the peaks at 1003, 774, 693, 531, and 459 cm^{-1} , which correspond to Si-O vibrations.




Fig. 6. Infrared spectra of the studied soil.

3.3 Physicochemical characteristics of the soils

The particle size distribution is dominated by sand fraction with proportions varying from 62% to 82%, corresponding to a mean value of $76.17 \pm 9.17\%$ (Table 5). Clay constitutes the second important particle size fraction, with proportions ranging from 13% to 23%, with a mean value of $16.33 \pm 5.16\%$. Silt fraction appears to be the least abundant, with proportions varying from 5% to 15%, with an average of $9.17 \pm 4.92\%$. Sand is strongly negatively correlated with clay ($r = -0.99$, $p < 0.05$). Silt has a moderate positive correlation with clay ($r = 0.53$, $p < 0.05$) and a negative correlation with sand ($r = -0.57$, $p < 0.05$) (Table 6).

The analysed soils exhibit a pH ranging from acidic to slightly basic (6.80 to 7.20), with an average of 6.98. No correlation was observed between pH and other elements, except for clay ($r = 0.91$, $p < 0.05$). The organic matter (OM) and total nitrogen (N) contents are relatively low, with values ranging from 1.30% to 3.17%, and 0.11% to 0.13%, with averages of $2.57 \pm 0.67\%$, and $1.12 \pm 0.01\%$ (Table 5). The C/N ratio, relatively low, varies from 6 to 16, with an average of 12.33 ± 3.61 . Base saturation (V) varies between 23.60% and 42.41%, with an average value of 34.28% (Table 5).

The exchangeable bases (Ca^{2+} , Mg^{2+} , K^{+} , and Na^{+}) exhibit notable variation across the soil profile. Calcium (Ca^{2+}) and magnesium (Mg^{2+}) concentrations are low to moderate, with values ranging from 2.64 to 6.00 $\text{cmol}_c \text{ kg}^{-1}$ in the case of Ca^{2+} and 0.72 to 5.44 $\text{cmol}_c \text{ kg}^{-1}$ for Mg^{2+} , with respective averages of 3.88 ± 1.49 and $2.45 \pm 1.58 \text{ cmol}_c \text{ kg}^{-1}$ (Table 5). Regarding potassium (K^{+}) and sodium (Na^{+}), their contents are very low to low, oscillating between 0.10 and 0.40 $\text{cmol}_c \text{ kg}^{-1}$ for K^{+} and between 0.10 and 0.60 $\text{cmol}_c \text{ kg}^{-1}$ for Na^{+} , with respective averages of 0.18 ± 0.12 and $0.28 \pm 0.17 \text{ cmol}_c \text{ kg}^{-1}$ (Table 5). Mg^{2+} shows a positive correlation with clay ($r = 0.62$, $p < 0.05$) and pH H_2O ($r = 0.66$, $p < 0.05$). K^{+} is strongly



correlated with clay ($r = 0.76$, $p < 0.05$) and moderately with pH H₂O ($r = 0.65$, $p < 0.05$). The sum of exchangeable bases (SEB) ranges from 5.10 to 8.90 cmol_c kg⁻¹, with an average value of 6.88 ± 1.66 cmol_c kg⁻¹ (Table 5). This SEB shows a moderately positive correlation with clay content ($r = 0.79$, $p < 0.05$). The cation exchange capacity (CEC) displays relatively high values, ranging from 18.70 to 25.00 cmol_c kg⁻¹, with an average of 22.05 ± 2.52 cmol_c kg⁻¹ (Table 5). It shows a strong correlation with the clay fraction ($r = 0.82$, $p < 0.05$), as well as with pH H₂O ($r = 0.84$, $p < 0.05$). It is also moderately correlated with potassium (K⁺) ($r = 0.75$, $p < 0.05$) (Table 6). Phosphorus (P) shows concentrations ranging from low to medium, with values between 12.60 and 30.30 mg kg⁻¹, and an average of 19.00 ± 6.95 mg kg⁻¹ (Table 5). Phosphorus shows a strong correlation with clay content ($r = 0.92$, $p < 0.05$) and with CEC ($r = 0.81$, $p < 0.05$) (Table 6). Base saturation (V) presents values ranging from 23.60% to 42.41%, with an average of $31.11 \pm 7.42\%$. Electrical conductivity varies from 0.02 to 1.20 dS/m, with an average of 0.27 ± 0.46 dS/m (Table 5). The latter is strongly correlated with Na, with a coefficient of $r = 0.92$ ($p < 0.05$).

Morphologically, the study soil profile is characterized by an alternation of different soil texture which are sandy loam, loamy sand, loamy clayey sand, clayey sand and sandy, in line with the alluvial nature of the parent rock. This alternation of soil texture is confirmed by the zigzag evolution of the percentage of different particle size distribution fraction noted in table 5. These characteristics correspond to those of Fluvisols. Base saturation ratio in all soil horizons is below 50%, leading to the choice of Dystric as principal qualifier. The light colour of soil and the organic matter content (<3%) lead to classify the studied soils as Ochric Dystric Fluvisols according to the IUSS Working Group WRB (2022).

Table 5. Physicochemical characteristics and summary statistics of the soil profile.

Depth statistics	cm	/	0-25	25-50	50-110	110-160	160-220	220-270	Min	Max	Mean	SD	CV
Sand (%)			82.00	82.00	67.00	88.00	62.00	88.00	62.00	82.00	76.17	9.17	0.12
Silt (%)			15.00	5.00	10.00	8.00	15.00	8.00	5.00	15.00	9.17	4.92	0.54
Clay (%)			13.00	13.00	23.00	5.00	23.00	5.00	5.00	23.00	16.33	5.16	0.32
pH _{H2O}			6.80	7.00	7.20	6.80	7.20	6.90	6.80	7.20	6.98	0.18	0.03
pH _{KCl}			4.40	4.20	6.30	6.60	4.60	5.50	4.20	6.60	5.27	1.02	0.19
OC (%)			0.75	1.84	1.66	1.66	1.36	1.66	0.75	1.84	1.49	0.39	0.26
OM (%)			1.30	3.17	2.86	2.86	2.34	2.86	1.30	3.17	2.57	0.67	0.26
N (%)			0.13	0.13	0.11	0.13	0.12	0.11	0.11	0.13	0.12	0.01	0.09
C/N			6.00	14.00	16.00	12.00	11.00	15.00	6.00	16.00	12.33	3.61	0.29
Ca (cmol _c kg ⁻¹)			2.96	2.64	2.72	5.52	6.00	3.44	2.64	6.00	3.88	1.49	0.38
Mg (cmol _c kg ⁻¹)			2.24	2.48	5.44	0.72	2.00	1.84	0.72	5.44	2.45	1.58	0.65
K (cmol _c kg ⁻¹)			0.15	0.11	0.42	0.13	0.18	0.11	0.10	0.40	0.18	0.12	0.64
Na (cmol _c kg ⁻¹)			0.22	0.22	0.31	0.56	0.13	0.31	0.13	0.56	0.29	0.17	0.61
SBE (cmol _c kg ⁻¹)			5.57	5.45	8.88	7.94	8.32	5.10	5.10	8.90	6.88	1.66	0.24
CEC (cmol _c kg ⁻¹)			20.80	23.17	25.00	18.72	24.38	20.19	18.70	25.00	22.05	2.52	0.11
P (mg kg ⁻¹)			13.10	12.58	24.29	17.52	30.28	16.22	12.60	30.30	19.00	6.95	0.37
V(%)			26.77	23.60	34.52	42.41	34.12	25.26	23.60	42.41	31.11	7.42	23.53



CE (mS cm ⁻¹)	0.07	0.02	0.18	1.18	0.04	0.08	0.02	1.20	0.27	0.46	1.73
---------------------------	------	------	------	------	------	------	------	------	------	------	------

315 **Table 6.** Spearman correlation matrix for linear relationships between soil profile properties.

	Sand	Silt	Clay	pH _{H2O}	pH _{KCl}	OC	OM	N	C/N	Ca	Mg	K	Na	SBE	CEC	P	V	CE
Sand	1																	
Silt	-0.57	1																
Clay	-0.99*	0.53	1															
pH _{H2O}	-0.90*	0.31	0.91	1														
pH _{KCl}	-0.05	-0.40	0.14	-0.05	1													
OC	0.00	-0.81*	0.04	0.29	0.40	1												
OM	0.00	-0.81*	0.04	0.29	0.40	0.99*	1											
N	0.56	-0.21	-0.63	-0.58	-0.32	-0.06	-0.06	1										
C/N	-0.17	-0.66	0.25	0.46	0.46	0.91*	0.91*	-0.46	1									
Ca	-0.37	0.17	0.25	0.07	0.21	-0.01	-0.01	0.22	-0.19	1								
Mg	-0.49	0.21	0.62	0.66	0.13	0.11	0.11	-0.69	0.42	-0.58	1							
K	-0.64	0.28	0.76	0.65	0.44	0.08	0.08	-0.71	0.36	-0.24	0.90*	1						
Na	0.44	-0.62	-0.38	-0.52	0.83	0.35	0.35	0.22	0.22	0.20	-0.29	-0.04	1					
SBE	-0.76	0.28	0.79	0.59	0.55	0.15	0.15	-0.31	0.20	0.50	0.36	0.69	0.20	1				
CEC	-0.79	0.42	0.82*	0.94*	-0.27	0.13	0.13	-0.47	0.30	-0.18	0.75	0.65	-0.67	0.43	1			
P	-0.96*	0.47	0.92*	0.80*	0.21	0.06	0.06	-0.52	0.19	0.58	0.29	0.52	-0.25	0.81*	0.60	1		
V	-0.34	0.02	0.35	0.07	0.73	0.12	0.12	0.04	0.03	0.69	-0.11	0.31	0.63	0.83*	-0.14	0.50	1	
CE	0.27	-0.41	-0.26	-0.45	0.72	0.22	0.22	0.41	0.00	0.50	-0.44	-0.11	0.92	0.37	-0.62	-0.07	0.81*	1

*Significant at $p < 0.05$.

3.4 Suitability of the studied soil for maize cultivation

The study area presents very favourable climatic conditions for the growth and yield of maize. The average temperature of 26.6 °C during the growing cycle presents a slight limitation for the production of maize (S1-1) (Table 7). The precipitation during the growing cycle, approximately 649 mm, has no limitation for the cultivation of maize. The climatic index of 91.41 (S1-0) indicates that there are no major limitations for maize cultivation in this area. This means that climatic conditions are favourable, suggesting the possibility of achieving optimal yields. The low slope (1%), presents a slight limitation (S1-1). The risk of flooding is non-existent (FO), and drainage is of good quality, indicating that there are no constraints related to soil moisture (S1-0) (Table 8). The soil texture, of the sandy-loamy type (SL), classified as S2, has proven to be a moderate limitation with a parametric value of 67.4. The soil depth, greater than 100 cm, is very suitable (S1-0), which is optimal for root development. Presence of coarse fragments displays a slight limitation (S1.1), with a parametric value of 96.3 for maize production (Table 8). The apparent cation exchange capacity (CEC) of the soil, classified as S1-1, indicates a slight limitation, with a parametric value of 90. Base saturation and organic carbon content, both classified as S2, reveal moderate limitations, with respective parametric values of 75.2 and 72.8. The soil pH (S1-1) and the electrical conductivity is suitable (S1-0) for the production of maize although there is a slight limitation related to pH (Table 8). The overall land suitability index is 62.37, which classifies it as S2sf, indicating moderate suitability for cultivation, primarily due to constraints related to the soil's physical properties and fertility.



Table 7. Climatic suitability evaluation for the production of Maize using simple limitation and parametric methods.

Characteristics	Values	Classes	Limitations	Parametric values
Precipitation				
Precipitation during crop cycle (mm)	649.18	S1-0	0	100
Precipitation during 3rd cycle (mm)	232.41	S1-0	0	100
Temperature				
Mean temperature during crop cycle (°C)	26.6	S1-1	1	95
Mean maximum temperature during crop cycle (°C)	31.8	S1-1	1	86.9
Mean minimum temperature during crop cycle (°C)	23	S1-1	1	93
Relative humidity during crop cycle (%)	73.25	S1-0	0	100
n/N during crop cycle	62.6	S1-1	0	100
Calculated climatic index (CR)		S1-1		91.41

335

Table 8. Land suitability evaluation for the maize production using simple limitation and parametric methods.

Characteristics	Values	Classes	Limitations	Parametric values
Topography (t)				
Slope (%)	1	S1-1	1	93.2
Wetness (w)				
Flooding	FO	S1-0	0	100
Drainage	Good	S1-0	0	100
Physical soil characteristics (s)				
Texture/structure	SL	S2	2	67.4
Soil depth (cm)	> 100	S1-0	0	100
Coarse fragments (%)		S1-1	1	96.3
Soil fertility characteristics (f)				
Apparent CEC (cmol _c kg ⁻¹ clay)	22.05	S1-1	1	90
Base saturation (%)	31.3	S2	2	75.2
Organic carbon (%)	0.75	S2	2	72.8
pH-H ₂ O	6.98	S1-1	1	94
CE	0.27	S1-0	0	100
Suitability and calculated land index (Is)		S2sf		62.37

3.4 Evaluation of the effect of trachyte and basalt powder on the growth and yield of maize

3.4.1 Effect of trachyte and basalt powder on maize growth parameters

The maize growth parameters include the germination rate, plant height, number of leaves, stem diameter, as well as ear length and ear diameter.

340



Germination was observed in all plots starting from the 5th day after sowing. After 10 days, the recorded germination rate reached 92.18%, indicating good uniformity and a of plant growth across all experimental plots.

The plant height ranges from 147.32±2.09 cm for treatment T0 to 215.12±2.09 cm for treatment T5, with the latter showing the tallest plants, while T0 (control) shows the shortest height. Treatments T1 (175.15±2.09 cm), T2 (181.47±2.09 cm), T3 (179.57±2.09 cm), and T4 (194.85±2.09 cm) show no significant differences among them, although all are higher than T0. However, treatment T5 (215.12 cm) is significantly taller than all other treatments. A progressive increase in plant height is observed with inputs of trachyte and basalt powders (Table 9).

The evolution of the number of leaves based on treatments shows that treatment T5, with an average of 22 leaves per plant (22±1), yields the best results, showing a significant difference compared to the other treatments. In contrast, treatment T0, with 18 leaves per plant (18±1), recorded the lowest number of leaves. Treatments T1, T2, and T4 produced similar results, each with an average of 20 leaves per plant (20±1). Treatment T3, with an average of 19 leaves per plant (19±1), is slightly lower than T1, T2, and T4, but still significantly higher than T0.

The stem diameter shows a significant increase depending on the treatments (Table 9). T0 recorded the smallest stem diameter (1.54±0.07 cm), serving as the reference without any treatment influence. T1 and T2 had stem diameters of 2.52±0.07 cm and 2.67±0.07 cm, respectively, both larger than T0. Similarly, T3 recorded a stem diameter of 2.56±0.07 cm, and T4 reached 2.72±0.07 cm, both showing wider diameters than T0. T5 exhibited the largest diameter at 3.03±0.07 cm. The plots treated with trachyte and basalt powder clearly stand out from the control plot (T0).

The evolution of ear length shows a significant increase as the treatments are applied (Table 9). Treatment T0, with an ear length of 8.55±0.9 cm, presents the shortest ears and stands out significantly from all other treatments. T1 (13.03±0.9 cm) and T2 (13.98±0.9 cm) display noticeably longer ears than those of T0. For treatments T3 and T4, the ear length increases to 13.45±0.9 cm and 14.07±0.9 cm, respectively. Under the influence of treatment T5, the length reaches its maximum at 15.65±0.9 cm, a value significantly higher than the others (Table 9).

The ear diameter shows a statistically significant increase with the application of the different treatments (Table 9). Treatment T0, with an average diameter of 3.00±0.08 cm, produces the smallest ears. T1 (3.54±0.08 cm) and T2 (3.86±0.08 cm) show a noticeable increase compared to T0. Treatment T3 (3.76±0.08 cm) and T4, with a diameter of 4.20±0.08 cm, also show significant increases, surpassing T0. Under the effect of treatment T5, the diameter reaches its maximum at 4.99±0.08 cm, a value significantly higher than those observed for the other treatments.

3.4.2 Effect of trachyte and basalt powder on maize yield parameters

Several parameters help assess the overall productivity of maize, including the weight of the ears, the weight of 100 maize grains, and the total yield in kilograms per hectare (kg ha⁻¹).

The weight of the ears increases significantly according to the treatments, indicating a positive response over time, resulting from the dissolution of the rock powder (Table 9). Treatment T0, with a weight of 31.74±6.82 g, is significantly lower than



all the others. Under treatment T1, the weight doubles, reaching 75.49 ± 6.82 g, and T2 (93.46 ± 6.82 g) also shows a notable improvement compared to T0. Treatment T3 (82.06 ± 6.82 g) and T4 (97.45 ± 6.82 g), are statistically significant compared to T0. Treatment T5, with a weight of 131.96 ± 6.82 g, shows a substantial and significant increase compared to all other treatments.

The variation in the weight of 100 grains shows that treatment T0, with a weight of 18.85 ± 2.18 g, is significantly lower than all the other treatments. Treatments T1 (24.21 ± 2.18 g) and T2 (27.64 ± 2.18 g) exhibit notable increases compared to T0, Treatments T3 (26.01 ± 2.18 g) and T4 (29.35 ± 2.18) also show significantly higher values than T0. Treatment T5, with an average weight of 32.02 ± 2.18 g for 100 grains, significantly outperforms all other treatments (Table 9).

The yield in kg ha^{-1} based on the treatments is presented in Table 9. Treatment T0, with a yield of $645.83 \pm 27.02 \text{ kg ha}^{-1}$, shows the lowest level among all treatments. In contrast, treatments T1 ($2362.87 \pm 27.02 \text{ kg ha}^{-1}$) and T2 ($2763.91 \pm 27.02 \text{ kg ha}^{-1}$) recorded significantly higher yields, demonstrating notable improvement compared to T0 due to the application of trachyte powder. Similarly, treatment T3 ($2558.64 \pm 27.02 \text{ kg ha}^{-1}$) and treatment T4 ($2931.16 \pm 27.02 \text{ kg ha}^{-1}$) reveal significant increases, highlighting the positive impact of basalt powder on maize yield. Treatment T5, with a maximum yield of $3164.45 \pm 27.02 \text{ kg ha}^{-1}$, statistically outperforms all other treatments. Globally, it is noted that application of basalt powder has more positive effect on maize yields than trachyte powder while adding urea to these treatments increase the maize yield.

Table 9. Statistical analysis of the effect of basalt and trachyte powder on the growth and yield of Maize.

Treatment	T0	T1	T2	T3	T4	T5
Plant height (cm)	$147.32 \pm 12.09a$	$175.15 \pm 12.9b$	$181.47 \pm 12.09b$	$179.57 \pm 12.09b$	$194.85 \pm 12.09c$	$215.12 \pm 12.09d$
Number of leaves	$18 \pm 1a$	$20 \pm 1b$	$20 \pm 1b$	$19 \pm 1c$	$20 \pm 1b$	$22 \pm 1d$
Stem diameter (cm)	$1.54 \pm 0.07a$	$2.52 \pm 0.07b$	$2.67 \pm 0.07c$	$2.56 \pm 0.07b$	$2.72 \pm 0.07d$	$3.03 \pm 0.07e$
Ear length (cm)	$8.55 \pm 0.9a$	$13.03 \pm 0.9b$	$13.98 \pm 0.9c$	$13.45 \pm 0.9bd$	$14.07 \pm 0.9e$	$15.65 \pm 0.99f$
Ear diameter (cm)	$3.00 \pm 0.08a$	$3.54 \pm 0.08b$	$3.86 \pm 0.08c$	$3.76 \pm 0.08cd$	$4.20 \pm 0.08e$	$4.99 \pm 0.08f$
Ear weight (g)	$31.74 \pm 6.82a$	$75.49 \pm 6.82b$	$93.46 \pm 6.82c$	$82.06 \pm 6.82d$	$97.45 \pm 6.82ce$	$131.96 \pm 6.82f$
Weight of 100 grain (g)	$18.85 \pm 2.18a$	$24.21 \pm 2.18b$	$27.64 \pm 2.18c$	$26.01 \pm 2.18cd$	$29.35 \pm 2.18ce$	$32.02 \pm 2.188f$
Yield (kg ha^{-1})	$645.83 \pm 27.02a$	$2362.87 \pm 27.02b$	$2763.91 \pm 27.02c$	$2558.64 \pm 27.02d$	$2931.16 \pm 27.02e$	$3164.45 \pm 27.02f$

T0: Control plot without treatment; T1: Treatment with trachyte powder (1.96 kg of trachyte powder for 9.6m^2); T2: Treatment with trachyte powder (1.96 kg of trachyte powder for 9.6m^2) + urea (100g for 9.6m^2) at 06 weeks; T3: Treatment with basalt powder (1.96 kg of basalt powder for 9.6m^2); T4: Treatment with basalt powder (1.96 kg of trachyte powder for 9.6m^2) + urea (100g for 9.6m^2) at 06 weeks. T5: Treatment with NPK fertilizer (200g for 9.6m^2) + urea (100g for 9.6m^2) at 06 weeks.

Mean value followed by the letters in the same line are not statistically different.



4 Discussion

4.1 Physical and mineralogical characteristics of soils

Particle size analyses revealed that the upper part of the soil profile primarily consists of a sandy-loam horizon, characterized by a high sand content ranging from 67 to 82%. The underlying horizons exhibit a succession of sandy and sandy-clay textures, with similar sand proportions. These observations are consistent with those observed in Fluvisols from various arid regions around the world (Romanens et al., 2019). This succession may reflect successive episodes of alluvial deposits, as well as long periods of drought, indicating pedogenetic evolution under hydrological influence. The predominance of sand fractions warrants particular attention, as it can negatively affect water and nutrient retention capacities while directly impacting plant root development, as also noted by Basga et al. (2018) and Tsozué et al. (2020a). Nevertheless, the limited presence of silt particles may help mitigate these effects by enhancing infiltration and slightly increasing the soil's water and nutrient retention capacity, as suggested by Lu et al. (2020) and Tsozué et al. (2021). The clay fraction, ranging from 13% to 18%, plays a significant role in the soil's structural stability and fertility. According to Kome et al. (2019) and Gautam et al. (2022), an adequate clay proportion can enhance both nutrient availability and water retention in the soil.

From a mineralogical perspective, the studied soils are predominantly composed of quartz, along with clay minerals such as kaolinite, smectites, and sepiolite. Kaolinite, an indicator of advanced weathering processes in tropical environments, results from climatic conditions that promote the leaching of basic cations and the neoformation of secondary minerals. These findings are consistent with the work of Tsozué et al. (2020b) and Lyu and Lu (2024). In contrast, the presence of smectite suggests moderate pedoclimatic conditions with variable humidity, which enhance cation exchange capacity and water retention. However, due to its swelling and shrinking behaviour, smectite could cause drainage problems and surface water stagnation, limiting agricultural use for certain sensitive crops (Hopmans et al., 2021). Sepiolite, on the other hand, is characteristic of semi-arid environments with low leaching rates (Bannari et al., 2021). According to Easwaran et al. (2024), this mineral is notable for its remarkable ability to retain water and nutrients, making it particularly beneficial for soils in arid areas or regions prone to drought. Additionally, the significant presence of quartz in the studied soils reflects the influence of alluvial parent material or exogenous inputs. These factors play a crucial role in shaping the textural and structural characteristics of the soil profiles, as observed by Tsozué et al. (2017) and Kemnitz and Lucke (2019).

4.2 Physicochemical characteristics of the studied soils

From a physicochemical perspective, the pH reveals low acidity, ranging between 6.80 and 7.20. These values are typical of soils under a Sudano-Sahelian climate, consistent with the chemical conditions observed in this region (Tsozué et al., 2020a; Tamto Mamdem et al., 2024). According to Tsozué et al. (2021) and Thiaw et al. (2024), this pH level could influence the availability of nutrients for plants. The slight acidification, particularly in the surface layer, can be attributed to the leaching of bases and the use of chemical fertilizers and pesticides (Pahalvi et al., 2021; Sarkar et al., 2024). The strong correlation



between pH and cation exchange capacity highlights the importance of regulating pH to optimize CEC and, consequently, improve the availability of essential nutrients for crops. These findings are consistent with those of Santos et al. (2023) regarding acidic soils in the amazonian region. According to Poggi et al. (2023), pH variations must be considered when selecting crops based on their tolerance. Organic matter plays a key role in soil fertility; its low proportion could be explained by several factors, including the semi-arid climate characteristic of the study area and the sandy texture of the soils. Comparable results have been reported by Plaza et al. (2018) and De Alencar et al. (2023) in the semi-arid regions of Brazil. The high correlation between OM and the C/N ratio shows that OM plays a crucial role in nutrient cycles and soil structure, which are important for ensuring sustainable agricultural productivity (Yang et al., 2024). Abdelhak (2022) highlighted that the low concentration of OM could result from the overexploitation of agricultural lands and intensive grazing after harvest. Arif et al. (2021) and Choudhury et al. (2024) emphasized in their research that improved management practices, such as adding organic matter, crop rotation, or using cover crops to improve soil fertility and overall health, could be corrective measures. The very low C/N ratio indicates rapid mineralization, likely related to high temperatures and the sandy texture of the soils. Similar results have been reported by Chen et al. (2024) in soils rich in organic carbon. The low content of available phosphorus in the soil could be attributed to its fixation on the clay-humic complex, a phenomenon strongly influenced by the clay content, pH, and calcium levels. Joshi et al. (2021) and Lotse Tedontsah et al. (2022) demonstrated that these factors play a key role in phosphorus availability, which is essential for crop growth and yield. The relatively high CEC is a crucial indicator of soil fertility, illustrating its ability to retain and exchange nutrients, essential for plant growth. According to Obi et al. (2020), this suggests that the soil can sustainably provide nutrients. High concentrations of Ca^{2+} and Mg^{2+} , especially at depth, indicate a good reserve of nutrients for plants (Hechmi et al., 2023). In contrast, the relatively low K^{+} content could limit plant growth, as potassium is important for photosynthesis and water regulation (Johnson et al., 2022). The moderate Na^{+} concentration would be beneficial in preventing salinity issues (Hussain et al., 2021). These exchangeable bases play a vital role in plant health and improving agricultural yields. The strong positive correlation between Mg and K is a favourable sign for soil quality and productivity. Verma et al. (2024), in their work, emphasized the importance of integrated nutrient management to optimize agricultural production. Electrical conductivity is a key indicator of soil salinity. Since the measured values are low, this would not favour optimal absorption of water and nutrients by plant roots (Golia, 2023).

4.3 Kinetics of nutrient element release from trachyte and basalt powders

The release of nutrients from trachyte and basalt powders is a key topic in pedology and geochemistry, particularly in biogeochemical weathering (Podia et al., 2023). These processes enhance the bioavailability of essential plant nutrients (Medeiros et al., 2023). Mineral hydration from water input causes swelling and disintegration, releasing cations like K, Ca, and Mg (Ramos et al., 2022; Wakeel, 2013).



The release of nutrients from trachyte and basalt powders is governed by interdependent physical, chemical, and biological processes (Swoboda et al., 2022). Mineralogical composition plays a fundamental role in nutrient dynamics. Basalt, rich in plagioclase feldspars, pyroxenes, and olivine, provides Ca, Mg, Fe, and trace elements. Its high reactivity stems from mafic minerals, which are more easily altered (Korndörfer et al., 2002; Ghasera and Rashid, 2024). In contrast, trachyte, mainly composed of alkali feldspars, is a major K source but has lower reactivity due to its felsic nature (Meena and Biswas, 2014; Mbissik et al., 2021). Particle size significantly influences mineral dissolution and nutrient release. Studies by Duarte et al. (2013), Burbano et al. (2021), and Rodrigues et al. (2024b) highlight that finer particles increase the specific surface area, accelerating dissolution. Meena and Biswas (2014) demonstrated that rock phosphate and mica release cations more rapidly when finely ground and combined with compost. Wakeel (2013) further emphasized that acidic soils enhance silicate solubilization, an effect amplified by particle fragmentation. Optimizing rock powder size thus improves agronomic efficiency by promoting sustained nutrient release. Soil pH and redox conditions also affect element solubility: acidic pH favours silicate dissolution and K, Mg, and Fe release, while reducing conditions increase Fe and Mn mobility (Wakeel, 2013; Vondráčková et al., 2017). Organic matter and microbial activity play key roles in mineral weathering (Rout et al., 2024). *Bacillus mucilaginosus* produces organic acids that enhance nutrient solubilization (Yang et al., 2016). Root exudates and mycorrhizal associations further improve K and Mg bioavailability, facilitating plant uptake (Ma and Yamaji, 2006; Menna and Biswas, 2014). This microbial activity, coupled with mycorrhizal interactions, enhances mineral breakdown and nutrient uptake (Menna and Biswas, 2014). These symbioses boost root access to nutrients and activate biological processes that promote silicate dissolution (Yadegari et al., 2024). These combined mechanisms make trachyte and basalt powders particularly effective in nutrient-deficient soils, facilitating the release of K, Ca, and Mg. Climatic conditions also influence nutrient release. Basak and Biswas (2009) found that high humidity promotes mineral alteration and microbial activity, while excessive rainfall can cause nutrient leaching (Lima et al., 2010). Nutrient interactions with soil minerals impact accessibility. Clays adsorb cations, limiting their immediate availability, but organic amendments enhance soil cation exchange capacity, reducing leaching (Duarte et al., 2013). An integrated soil management approach, combining organic amendments with optimized environmental conditions, is essential to maximize nutrient efficiency from minerals.

4.4 Suitability of the soils from Guiring to maize production

The climate of the study area offers particularly favourable conditions for maize growth and yield, as evidenced by annual rainfall and the climatic index. These factors highlight the importance of climate in this region, without presenting major constraints for maize cultivation. However, potential variations due to climate change could disrupt rainfall patterns and temperatures, necessitating continuous monitoring to anticipate and mitigate their impacts (Liu et al., 2024). In this context, Rocha et al. (2020) recommend water management strategies such as irrigation during dry periods to ensure stable agricultural productivity. Conversely, some edaphic parameters pose moderate limitations, including the sandy-loam texture and low base saturation, which affect water retention and nutrient availability (Verdoodt et al., 2003; Tsozué et al., 2015;



490 Issiné et al., 2022; Scanlan et al., 2022). To address these limitations, adapted management practices are essential. Adding organic matter, such as compost or crop residues, and using cover crops like legumes can improve soil structure and water retention capacity (Obi et al., 2020). Additionally, the application of lime or dolomite could increase calcium and magnesium saturation, while potassium fertilizers can correct low potassium availability (Hechmi et al., 2023). Improving soil texture is also a critical strategy. Adding clay or biochar, identified by Hussain et al. (2021) as an effective solution, 495 enhances water retention and cation exchange capacity, both of which are fundamental properties for the fertility of sandy soils. The use of organic fertilizers is similarly effective in overcoming fertility limitations while promoting sustainable soil management (Van Leeuwen et al., 2015; Das et al., 2024; Robinson, 2024). Finally, trachyte and basalt powders, used as natural amendments, represent a promising alternative for improving soil fertility, especially in semi-arid regions where soils are often degraded by nutrient depletion and poor structure which can not optimize high nutrients release fertilizers such as 500 NPK and urea. Derived from the weathering or grinding of silicate or carbonate rocks, these powders gradually release essential minerals such as calcium, magnesium, and potassium, thereby increasing exchangeable bases and cation saturation (Swoboda et al., 2022; Basak et al., 2023). Similarly, as demonstrated by Manning and Theodoro (2020) and Ramos et al. (2022), they enhance nutrient retention, improve soil mineral composition, increase cation exchange capacity, and contribute to sustainable agricultural fertility management. As slow-release mineral sources, these powders also help restore nutrient 505 balances, improve water retention, and revitalize soil microbial activity (Anda et al., 2015; Guo et al., 2023). This is particularly relevant in semi-arid zones, where conventional fertilizers are often ineffective due to rapid nutrient leaching and unfavourable climatic conditions (Ramos et al., 2022; Cardozo et al., 2024). By stimulating microbial activity, they enhance long-term soil health and support sustainable agricultural practices (Ramos et al., 2022). This combination of integrated solutions could transform semi-arid soils into productive and resilient resources, adapted to the challenges posed by climate 510 change and edaphic constraints.

4.5 Effect of trachyte and basalt powder on maize growth and yield

Trachyte powder, with relatively high concentrations of K_2O and Na_2O , proves beneficial for soil fertilization, as these elements promote plant growth, particularly potassium, which is essential for photosynthesis and water regulation. Nevertheless, its low levels of Fe and Mg limit its direct nutrient contribution for maize, which primarily requires nitrogen, 515 phosphorus, and potassium for its growth (Reimann et al., 2018; Linhares et al., 2021). In contrast, basalts, characterized by high concentrations of Fe_2O_3 , CaO , TiO_2 , and P_2O_5 , while exhibiting reduced levels of SiO_2 and Al_2O_3 , can serve as a natural nutrient source (Oppon et al., 2023). They are particularly suited to enriching soils deficient in these elements, thereby improving soil fertility and productivity (Kagou Dongmo et al., 2018; Luchese et al., 2021; Conceição et al., 2022). The results show that the application of rock powders and conventional fertilizers led to an increase in essential parameters 520 related to maize growth and yield, including plant height, stem diameter, ear diameter, length, and weight, 100-grain weight, and overall yield, compared to the control. Plots treated solely with trachyte and basalt powders exhibited significantly



higher growth compared to control plots (Ramos et al., 2017, 2022; Reis et al., 2024). These results highlight the potential of these powders as effective amendments for improving maize growth, particularly due to their contents of K_2O , P_2O_5 , CaO , and Fe_2O_3 . Several studies confirm the beneficial effects of applying rock powders for remineralisation and soil amendment in agriculture, including Mein'da et al. (2022) on the use of pyroclastic powders on ferrallitic soils in the Adamaoua region, Ramos et al. (2015, 2017, 2021, 2022), Oliveira et al. (2020), and Luchese et al. (2023) on volcanic rock powders for soil fertilisation in Brazil. The application of these powders positively impacted maize growth parameters, particularly vegetative development and vigour. This improvement could influence photosynthesis, biomass production, and even seed quality, contributing to increased productivity and yield (Oliveira et al., 2020; Viana et al., 2021; Bamberg et al., 2022; Luchese et al., 2023). These findings confirm that the use of trachyte and basalt powders can sustainably improve soil fertility by supplying both macronutrients (N, P, K, Ca, Mg, S) and micronutrients (Fe, Mn, Zn, Cu, B). This contributes to improved crop quality and yields, consistent with the conclusions of Swoboda et al. (2022) and Luchese et al. (2023). In light of the growing impacts of climate change, soil degradation, rising fertilizer costs for farmers, and crop health issues, rock powders represent a promising alternative to conventional fertilizers (Nkouathio et al., 2008; Burbano et al., 2022; Swoboda et al., 2022).

The results also show that overall maize yield improved significantly, increasing from 645 kg ha^{-1} the control plot to 2763 kg ha^{-1} in the plot treated with trachyte powder and 2931 kg ha^{-1} in the plot amended with basalt powder. These findings emphasize the potential of trachyte and basalt powders to boost maize yields (Tamfuh et al., 2020; Ramos et al., 2020; Mbissik et al., 2023). Although the highest yield was obtained with the application of conventional fertilizers (3164 kg ha^{-1}), trachyte and basalt powders could be more advantageous due to their local availability, lower cost, and positive effects on the soil. They release nutrients slowly and sustainably over multiple seasons, unlike conventional fertilizers, which provide nutrients rapidly but for a short duration. These rock powders act as a natural and sustainable source of essential nutrients for plants, gradually releasing minerals into the soil. They help improve soil fertility without causing pollution or harming biodiversity. Moreover, their use has no negative environmental impact, unlike chemical fertilizers, which can alter ecosystems and affect water and soil quality (Nkouathio et al., 2008; Lopes et al., 2014; Oliveira et al., 2020; Luchese et al., 2023).

5 Conclusion

This study aims to evaluate the effect of using trachyte and basalt powders on maize growth and yield in the Guiring area. Geochemically, the Zamai trachytes are rich in silica and potassium, while the basalts are distinguished by their high content of iron, magnesium, and calcium. The soils in Guiring, dominated by a high sand content (62-82%) and low clay (13-23%) and silt (5-15%) contents. Mineralogically, these soils are composed of kaolinite, smectites, sepiolite, and quartz. The pH ranges from 6.80 to 7.20, organic matter (1.30-3.17%), and total nitrogen (0.11-0.13%) are low, with a low C/N ratio. The cation exchange capacity ($18.70\text{-}25.00 \text{ cmol}_c \text{ kg}^{-1}$) is high, while exchangeable bases (Ca^{2+} , Mg^{2+} , K^+ , Na^+) and phosphorus



are low to moderate, with base saturation varying between 23.60 and 42.41%. The study area has very favorable climatic conditions for maize growth. Soil texture, base saturation, and organic carbon content are classified as S2, indicating moderate limitations. Soil depth (S1-0) is highly suitable, and coarse fragments (S1-1) are slightly suitable. Cation exchange capacity and pH also show slight limitations (S1-1). Overall, the land suitability index is 62.37, classifying it as S2sf, indicating moderate suitability for maize cultivation. Growth and yield parameters show significant improvement based on the treatments applied. The control treatment (T0) shows the lowest yield, with 645.83 kg ha⁻¹. Treatments T1 (2362.87 kg ha⁻¹) and T2 (2763.91 kg ha⁻¹) showed notable improvements with trachyte powder application. Treatments T3 (2558.64 kg ha⁻¹) and T4 (2931.16 kg ha⁻¹) highlight the positive effect of basalt powder. Treatment T5 achieves the maximum yield of 3164.45 kg ha⁻¹, surpassing all other treatments highlighting an importance of associating urea to rocks powders. Although chemical fertilizers provide maximum yields in the short term, trachyte and basalt powders offer a sustainable and cost-effective alternative, improving soils while ensuring competitive yields. Their use is particularly beneficial for resilient and environmentally friendly agriculture.

Author contributions

All the authors substantially contributed to this article. The conceptualization of the study was done by Désiré Tsozué and Bienvenu Sidsi, with input from Claudine Vounba, Simon Djakba Basga. The data acquisition, investigation, methodology, and visualization for the paper were performed by Désiré Tsozué, Bienvenu Sidsi, Claudine Vounba Simon, Djakba Basga, with substantial input from Aubin Nzeugang Nzeukou and Merlin Gountie Dedzo. Bienvenu Sidsi and Claudine Vounba wrote the initial draft, and Désiré Tsozué and Simon Djakba Basga were involved in the reviewing, editing, and the validation of the paper. All authors read and agreed to the final version of the manuscript.

Data availability

The research data will be freely available on request from the corresponding author.

Declarations Competing interests

On behalf of all authors, the corresponding author states that there is no conflict of interest.

Acknowledgements

Authors thank the administrative authorities of the Institute of Agricultural Research for Development (IRAD) of Maroua for providing plots for experimentation and support for field campaigns.



6 References

- Abdelhak, M.: Soil improvement in arid and semiarid regions for sustainable development. In Natural resources conservation and advances for sustainability, <https://doi.org/10.1016/B978-0-12-822976-7.00026-02>, 2022.
- Anda, M., Shamsuddin, J., Fauziah, C. I.: Improving chemical properties of a highly weathered soil using finely ground
 585 basalt rocks. *Catena*. 124, 147-161. <https://doi.org/10.1016/j.catena.2014.09.012>, 2015.
- Are, K. S., Oshunsanya, S. O. and Oluwatosin, G. A.: Changes in soil physical health indicators of an eroded land as influenced by integrated use of narrow grass strips and mulch. *Soil and Tillage Research*. 184, 269-280. <https://doi.org/10.1016/j.still.2018.08.0092>, 2018.
- Arif, M., Ikramullah, Jan, T., Riaz, M., Akhtar, K., Ali, S. and Wang, H.: Biochar and leguminous cover crops as an
 590 alternative to summer fallowing for soil organic carbon and nutrient management in the wheat-maize-wheat cropping system under semiarid climate. *Journal of Soils and Sediments*. 21, 1395-1407. <https://doi.org/10.1007/s11368-020-02866-y>, 2021.
- Bannari, A. and Abuelgasim, A.: Potential and limits of vegetation indices compared to evaporite mineral indices for soil salinity discrimination and mapping. *Soil Discussions*. 1-48. <https://doi.org/10.5194/soil-2021-55>, 2021.
- Basak, B. B., Sarkar, B., Maity, A., Chari, M. S., Banerjee, A. and Biswas, D. R.: Low-grade silicate minerals as value-
 595 added natural potash fertilizer in deeply weathered tropical soil. *Geoderma*. 433, 116433. <https://doi.org/10.1016/j.geoderma.2023.116433>, 2023.
- Basga, S. D., Madi, P. O., Balna, J., Abib, C. F., Tsozué, D. and Njiemoun, A.: Sandy soil fertility restoration and crops yields after conversion of long term *Acacia senegal* planted fallows in North Cameroon. <https://doi.org/10.5897/AJAR2018.13283>, 2018.
- Beerling, D. J., Leake, J. R. and Long, S. P.: Farming with crops and rocks to address global climate, food and soil security. *Nature Plants* 4, 138–147. <https://doi.org/10.1038/s41477-018-0108-y>, 2018.
- Burbano, D. F. M., Theodoro, S. H., de Carvalho, A. M. X. and Ramos, C. G.: Crushed volcanic rock as soil remineralizer: a strategy to overcome the global fertilizer crisis. *Natural Resources Research*. 31(5), 2197-2210. <https://doi.org/10.1007/s11053-022-10107-x>, 2022.
- Cardozo, E., Pinto, V., Nadaleti, W., Thue, P., dos Santos, M., Gomes, C. and Vieira, B.: Sustainable agricultural practices: volcanic rock potential for soil remineralization. *Journal of Cleaner Production*. 142876. <https://doi.org/10.1016/j.jclepro.2024.142876>, 2024.



- Chen, Z., Zhang, H., Tu, X., Wang, J., Elrys, A. S., Tang, Q. and Cheng, Y.: C/N ratio of high-organic C materials is a poor predictor of microbial nitrate immobilization potential in a nitrate-rich soil: An 15N incubation study. *Soil and Tillage Research*. 238, 106019. <https://doi.org/10.1016/j.still.2024.106019>, 2024.
- Choudhury, D., Kumar, P., Zhimo, V. Y. and Sahoo, J.: Crop rotation patterns and soil health management. In *Bioremediation of Emerging Contaminants from Soils*. (pp. 565-589). Elsevier. <https://doi.org/10.1016/B978-0-443-13993-2.00025-6>, 2024.
- Conceição, L. T., Silva, G. N., Holsback, H. M. S., de Figueiredo Oliveira, C., Marcante, N. C., de Souza Martins, E. And Santos, E. F.: 2022. Potential of basalt dust to improve soil fertility and crop nutrition. *Journal of Agriculture and Food Research*. 10, 100443. <https://doi.org/10.1016/j.jafr.2022.100443>, 2022.
- Cox, W. J. and Andrade, H. F: Growth, yield, and yield components of maize as influenced by ethephon. *Crop science*. 28(3), 536-542. <https://doi.org/10.2135/cropsci1988.0011183X002800030023x>, 1988.
- Dalmora, A. C., Ramos, C. G., Oliveira, M. L. S., Oliveira, L. F. S., Schneider, I. A. H. and Kautzmann, R. M.: Application of andesite rock as a clean source of fertilizer for eucalyptus crop: Evidence of sustainability. *Journal of Cleaner Production*. 256, 120432. <https://doi.org/10.1016/j.jclepro.2020.120432>, 2020.
- Das, S., Kim, P. J., Nie, M. and Chabbi, A.: Soil organic matter in the anthropocene: Role in climate change mitigation, carbon sequestration, and food security. *Agriculture, Ecosystems and Environment*. 109180. <https://doi.org/10.1016/j.agee.2024.109180>, 2024.
- De Alencar, G. V., Gomes, L. C., Barros, V. M. D. S., Ortiz Escobar, M. E., de Oliveira, T. S. and Mendonça, E. D. S.: Organic farming improves soil carbon pools and aggregation of sandy soils in the Brazilian semi-arid region. *Soil Use and Management*. 40(3), e13097. <https://doi.org/10.1111/sum.13097>, 2024.
- Duarte I. N., Pereira H. S. and Korndörfer G. H.: Lixiviação de potássio proveniente do termopotássio. *Pesqui Agropecu Trop* 43:195–200. <https://doi.org/10.1590/S1983-40632013000200003>, 2013.
- Easwaran, C., Moorthy, G., Christopher, S. R., Mohan, P., Marimuthu, R., Koothan, V. and Nallusamy, S.: Nano hybrid fertilizers: A review on the state of the art in sustainable agriculture. *Science of The Total Environment*. 172533, <https://doi.org/10.1016/j.scitotenv.2024.172533>, 2024.



- Ellis, R. H., Covell, S., Roberts, E. H., & Summerfield, R. J.: The influence of temperature on seed germination rate in grain legumes: II. Intraspecific variation in chickpea (*Cicer arietinum* L.) at constant temperatures. *Journal of Experimental Botany*. 37(10), 1503-1515. <https://doi.org/10.1093/jxb/37.10.1503>, 1986.
- Farahani, E., Mosaddeghi, M. R., Mahboubi, A. A., and Dexter, A. R.: Prediction of soil hard-setting and physical quality using water retention data. *Geoderma*. 338, 343-354. <https://doi.org/10.1016/j.geoderma.2018.12.012>, 2019.
- Garai, S., Mondal, M., Nayak, J., Sarkar, S., Banerjee, H., Brahmachari, K. and Hossain, A.: Input Use Efficiency for Improving Soil Fertility and Productivity. *Input Use Efficiency for Food and Environmental Security*. 305-333. https://doi.org/10.1007/978-981-16-51991_10, 2021.
- Gautam, A., Guzman, J., Kovacs, P. and Kumar, S.: Manure and inorganic fertilization impacts on soil nutrients, aggregate stability, and organic carbon and nitrogen in different aggregate fractions. *Archives of Agronomy and Soil Science*. 68(9), 1261-1273. <https://doi.org/10.1080/03650340.2021.1887480>, 2022.
- Ghasera, K. M. and Rashid, S. A.: Influence of micro-scale factors in weathering and elements mobility: Evidence from a comparative study of granite and basalt weathering profiles across India. *Catena*, 235, 107680. <https://doi.org/10.1016/j.catena.2023.107680>, 2024.
- Golia, E. E.: The impact of heavy metal contamination on soil quality and plant nutrition. Sustainable management of moderate contaminated agricultural and urban soils, using low cost materials and promoting circular economy. *Sustainable Chemistry and Pharmacy*. 33, 101046. <https://doi.org/10.1016/j.scp.2023.101046>, 2023.
- Gountié, D. M., Asobo, N. E., Fozing, E. M., Tchamabé, B. C., Zangmo, T. G., Nguindhama, D., Tchokona, S. D., Kamgang, P., Aka, T. F. and Ohbak, T.: Petrology and geochemistry of lavas from Gawar, Minawao and Zamay volcanoes of the northern segment of the Cameroon volcanic line (Central Africa): Constraints on mantle source and geochemical evolution. *J. Afr. Earth Sci.* 153, 3-41. <https://doi.org/10.1016/j.jafrearsci.2019.02.010>, 2019.
- Günel, H. and Acir, N.: Spatial variability of clay minerals in a semi-arid region of Türkiye. *Geoderma Regional*. e00820. <https://doi.org/10.1016/j.geodrs.2024.e00820>, 2024.
- Gunnarsen, K. C., Jensen, L. S., Rosing, M. T., Dietzen, C.: Greenlandic glacial rock flour improves crop yield in organic agricultural production. *Nutrient Cycling in Agroecosystems*. 126(1), 51-66. <https://doi.org/10.1007/s10705-023-10274-0>, 2023.



- Guo, F., Wang, Y., Zhu, H., Zhang, C., Sun, H., Fang, Z. and Wu, F.: Crop productivity and soil inorganic carbon change mediated by enhanced rock weathering in farmland: A comparative field analysis of multi-agroclimatic regions in central China. *Agricultural Systems*, 210. 103691. <https://doi.org/10.1016/j.agsy.2023.103691>, 2023.
- Hechmi, S., Melki, S., Khelil, M. N., Ghrib, R., Gueddari, M. and Jedidi, N.: Potential risk of soil irrigation with treated wastewater over 40 years: a field experiment under semi-arid conditions in northeastern Tunisia. *Journal of Arid Land*. 15(4), 407-423. <https://doi.org/10.1007/s40333-023-0100-x>, 2023.
- 665 Hishe, S., Lyimo, J.b. and Bewket, W.: Soil and water conservation effects on soil properties in the Middle Silluh Valley, northern Ethiopia. *International Soil and Water Conservation Research*. 5, 231-240. <https://doi.org/10.1016/j.iswcr.2017.06.005>, 2017.
- Hopmans, J. W., Qureshi, A. S., Kisekka, I., Munns, R., Grattan, S. R., Rengasamy, P. and Taleisnik, E.: Critical knowledge gaps and research priorities in global soil salinity. *Advances in agronomy*. 169, 1-191. <https://doi.org/10.1016/bs.agron.2021.03.001>, 2021.
- 670 Hossain, A., Krupnik, T. J., Timsina, J., Mahboob, M. G., Chaki, A. K., Farooq, M. and Hasanuzzaman, M.: Agricultural land degradation: processes and problems undermining future food security. In *Environment, climate, plant and vegetation growth*. 17-61. https://doi.org/10.1007/978-3-030-49732-3_2, 2020.
- Hussain, S., Hussain, S., Ali, B., Ren, X., Chen, X., Li, Q. and Ahmad, N.: Recent progress in understanding salinity tolerance in plants: Story of Na⁺/K⁺ balance and beyond. *Plant Physiology and Biochemistry*. 160, 239-256. <https://doi.org/10.1016/j.plaphy.2021.01.029>, 2021.
- Issine, A., Tsozué, D., Nzeukou, A. N., Yongue, R. F., and Kagonbé, B. P.: Soil Characteristics and Pedoclimatic Evaluation of Rainfed Sorghum (*Sorghum bicolor* (L.) Moench) in the Mayo-Lemé Division, South-Western Chad. *Applied and Environmental Soil Science*, 2022. <https://doi.org/10.1155/2022/7394260>, 2022.
- 680 IUSS Working Group WRB: World Reference Base for Soil Resources. International soil classification system for naming soils and creating legends for soil maps. 4th edition. International Union of Soil Sciences (IUSS), Vienna, Austria, 2022.
- Jhariya, M. K., Banerjee, A., Meena, R. S., Kumar, S., Raj, A. Sustainable intensification for agroecosystem services and management: an overview. *Sustainable Intensification for Agroecosystem Services and Management*. 1-35. https://doi.org/10.1007/978-981-16-3207-5_1, 2021.



- 685 Johnson, R., Vishwakarma, K., Hossen, M. S., Kumar, V., Shackira, A. M., Puthur, J. T. and Hasanuzzaman, M.: Potassium in plants: Growth regulation, signaling, and environmental stress tolerance. *Plant Physiology and Biochemistry*. 172, 56-69. <https://doi.org/10.1016/j.plaphy.2022.01.001>, 2022.
- Joshi, S. R., Morris, J. W., Tfaily, M. M., Young, R. P. and Mc Near, D. H.: Low soil phosphorus availability triggers maize growth stage specific rhizosphere processes leading to mineralization of organic P. *Plant and Soil*. 459, 423-440. <https://doi.org/10.1007/s11104-020-04774-z>, 2021.
- 690 Kagou Dongmo, A., Guedjeo, C. S., Azinwi Tamfuh, P., Wotchoko, P., Chenyi, M. L. and Aziwo B. T.: Geochemical and Geotechnical Characterization of Soils Developed on Volcanic Rocks on the Bamenda Mountain (Cameroon Volcanic Line). *International Journal of Advanced Geosciences*. 6, 184-194. <https://doi.org/10.14419/ijag.v6i2.13505>, 2018.
- Kemnitz, H. and Lucke, B.: Quartz grain surfaces—A potential microarchive for sedimentation processes and parent material identification in soils of Jordan. *Catena*. 176, 209-226. <https://doi.org/10.1016/j.catena.2018.12.038>, 2019.
- 695 Khan, F., Siddique, A. B., Shabala, S., Zhou, M., and Zhao, C.: Phosphorus plays key roles in regulating plants' physiological responses to abiotic stresses. *Plants*, 12(15), 2861. <https://doi.org/10.3390/plants12152861>, 2023.
- Khatri, P., Kumar, P., Shakya, K. S., Kirlas, M. C., and Tiwari, K. K. Understanding the intertwined nature of rising multiple risks in modern agriculture and food system. *Environment, Development and Sustainability*. 26(9), 24107-24150. <https://doi.org/10.1007/s10668-023-03638-7>, 2024.
- 700 Kome, G. K., Enang, R. K., Tabi, F. O. and Yerima, B. P. K.: Influence of clay minerals on some soil fertility attributes: a review. *Open Journal of Soil Science*. 9(9), 155-188. <https://doi.org/10.4236/ojss.2019.99010>, 2019.
- Kuria, A. W., Barrios, E., Pagella, T., Muthuri, C. W., Mukuralinda, A. and Sinclair, F. L.: Connaissance des agriculteurs sur les indicateurs de qualité des sols le long d'un gradient de dégradation des terres au Rwanda. *Geoderma regional*. 16, e00199. <https://doi.org/10.1016/j.geodrs.2018.e00199>, 2019.
- 705 Letouzey R., 1985. Notice explicative de la carte phytogéographique du Cameroun au 1/500000. Domaines sahéliens et soudaniens.
- Lewis, A. L., Sarkar, B., Wade, P., Kemp, S. J., Hodson, M. E., Taylor, L. L and Beerling, D. J.: Effects of mineralogy, chemistry and physical properties of basalts on carbon capture potential and plant-nutrient element release via enhanced weathering. *Applied Geochemistry*, 132, 105023. <https://doi.org/10.1016/j.apgeochem.2021.105023>, 2021.



- Lima C C, Mendonça E. D. S. and Roig, A.: Contribution of humic substances from different composts to the synthesis of humin in a tropical soil. *Rev Bras Ciência do Solo* 34:1041–1048. <https://doi.org/10.1590/S0100-06832010000400004>, 2010.
- 715 Linhares, D., Pimentel, A., Garcia, P. and Rodrigues, A.: Deficiency of essential elements in volcanic soils: potential harmful health effects on grazing cattle. *Environmental Geochemistry and Health*. 1-13. <https://doi.org/10.1007/s10653-021-00874-6>, 2021.
- Liu, J., Fu, Z. and Liu, W.: Impacts of precipitation variations on agricultural water scarcity under historical and future climate change. *Journal of Hydrology*. 617, 128999. <https://doi.org/10.1016/j.jhydrol.2022.128999>, 2023.
- 720 Lopes, O. M. M., Carrilho, E. N. V. M. and Lopes-Assad, M. L. R. C.: Effect of rock powder and vinasse on two types of soils. *Revista Brasileira de Ciência do Solo*. 38, 1547-1557. <https://doi.org/10.1590/S0100-06832014000500020>, 2014.
- Lotse Tedontsah, V. P., Mbog, M. B., Ngon Ngon, G. F., Edzoa, R. C., Tassongwa, B., Bitom, D. and Etame, J.: Physicochemical properties and fertility assessment of soils in fouban (west Cameroon). *Applied and Environmental Soil Science*. 2022(1), 7889430. <https://doi.org/10.1155/2022/7889430>, 2022.
- 725 Lu, J., Zhang, Q., Werner, A. D., Li, Y., Jiang, S. and Tan, Z.: Root-induced changes of soil hydraulic properties—A review. *Journal of Hydrology*. 589, 125203. <https://doi.org/10.1155/2022/7889430>, 2020.
- Luchese, A. V., Pivetta, L. A., Batista, M. A., Steiner, F., Giaretta, A. P. D. S. and Curtis, J. C. D.: Agronomic feasibility of using basalt powder as soil nutrient remineralizer. *African Journal of Agricultural Research*. 17(1), 487-497. <http://www.academicjournals.org/AJAR>, 2021.
- 730 Luchese, A. V., de Castro Leite, I. J. G., da Silva Giaretta, A. P., Alves, M. L., Pivetta, L. A. and Missio, R.: Use of quarry waste basalt rock powder as a soil remineralizer to grow soybean and maize. *Heliyon*, 9(3). <https://doi.org/10.1016/j.heliyon.2023.e14050>, 2023.
- Lyu, H. and Lu, H.: Precipitation is the main control on the global distribution of soil clay minerals. *Earth-Science Reviews*. 104891. <https://doi.org/10.1016/j.earscirev.2024.104891>, 2024.
- 735 Ma J. F. and Yamaji N.: Silicon uptake and accumulation in higher plants. *Trends Plant Sci* 11:392–397. <https://doi.org/10.1016/j.tplants.2006.06.007>, 2006.



- Makishima, A. and Nakamura, E.: Determination of major/minor and trace elements in silicate samples by ICP-QMS and ICP-SFMS applying isotope dilution-internal standardisation (ID-IS) and multi-stage internal standardisation. *Geostand. Geoanal. Res.* 30, 245–271. <https://doi.org/10.1111/j.1751-908X.2006.tb01066.x>, 2006.
- 740 Mamabolo, E., Gaigher, R. and Pryke, J. S.: Conventional agricultural management negatively affects soil fauna abundance, soil physicochemical quality and multifunctionality. *Pedobiologia.* 104, 150961. <https://doi.org/10.1016/j.pedobi.2024.150961>, 2024.
- Manning, D. A. and Theodoro, S. H.: Enabling food security through use of local rocks and minerals. *The Extractive Industries and Society.* 7(2), 480-487. <https://doi.org/10.1016/j.exis.2018.11.002>, 2020.
- 745 Mbissik, A., Elghali, A., Ouabid, M., Raji, O., Bodinier, J. L. and El Messbahi, H.: Alkali-hydrothermal treatment of K-rich igneous rocks for their direct use as potassic fertilizers. *Minerals*, 11(2), 140. <https://doi.org/10.3390/min11020140>, 2021.
- Medeiros, F. D. P., Carvalho, A. M. D., Gindri Ramos, C., Dotto, G. L., Cardoso, I. M. and Theodoro, S. H.: Rock Powder enhances soil nutrition and coffee quality in agroforestry systems. *Sustainability*, 16(1), 354. <https://doi.org/10.3390/su16010354>, 2023.
- 750 Meena M. D. and Biswas D.R.: Changes in biological properties in soil amended with rock phosphate and waste mica enriched compost using biological amendments and chemical fertilizers under wheat-soybean rotation. *J Plant Nutr.* <https://doi.org/10.1080/01904167.2014.920368>, 2014.
- Mein'da, C. D., Maimouna, A., Tatchum, L. T., Nguetnkam, J. P. and Megueni, C.: Effect of Pyroclastic Powder on the Growth and Seeds Nutritional Values of Common Bean (*Phaseolus vulgaris* L.) on Ferralitic Soils in Adamawa (Cameroon, Central Africa). *International Journal of Plant & Soil Science.* 34(21), 211-225. <https://doi.org/10.9734/ijpss/2022/v34i2131257>, 2022.
- 755 <https://doi.org/10.9734/ijpss/2022/v34i2131257>, 2022.
- Mekuriaw, A., Heinimann, A., Zeleke, G., Hurni, H. and Hurni, K.: An automated method for mapping physical soil and water conservation structures on cultivated land using GIS and remote sensing techniques. *Journal of geographical sciences*, 27, 79-94. <https://doi.org/10.1007/s11442-017-1365-9>, 2017.
- 760 Moges, D. M. and Taye, A. A.: Determinants of farmers' perception to invest in soil and water conservation technologies in the North-Western Highlands of Ethiopia. *International Soil and Water Conservation Research.* 5(1), 56-61. <https://doi.org/10.1016/j.iswcr.2017.02.003>, 2017.



- Muhie, S.H.: Novel approaches and practices to sustainable agriculture. *Journal of Agriculture and Food Research*. 10, 100446. <https://doi.org/10.1016/j.jafr.2022.100446>, 2022.
- 765 Muthaura, C., Mucheru-Muna, M., Zingore, S., Kihara, J. and Muthamia, J.: Effect of application of different nutrients on growth and yield parameters of maize (*Zea mays*), case of Kandara Murang'a County. *ARPN Journal of Agricultural and Biological Science*. 12(1), 19-33, 2017.
- Nanganoa, L. T., Okolle, J. N., Missi, V., Tueche, J. R., Levai, L. D. and Njukeng, J. N.: Impact of Different Land-Use Systems on Soil Physicochemical Properties and Macrofauna Abundance in the Humid Tropics of Cameroon. *Applied and Environmental Soil Science*. 2019(1), 5701278. <https://doi.org/10.1155/2019/5701278>, 2019.
- 770 Ngoune Tandzi, L. and Mutengwa, C. S.: Estimation of maize (*Zea mays* L.) yield per harvest area: Appropriate methods. *Agronomy*, 10(1), 29. https://doi.org/10.3390/agronomy_10010029, 2019.
- Nkouathio, D. G., Wandji, P., Bardintzeff, J. M., Tematio, P., Dongmo, A. K. and Tchoua, F.: Utilisation des roches volcaniques pour la remineralisation des sols ferrallitiques des regions tropicales. Cas des pyroclastites basaltiques du graben de Tombel (Ligne volcanique du Cameroun). *Bull. Soc. Vaudoise Sci. Nat.* 91, 1-14, 2008.
- 775 Obi, J. C., Udoh, I. B. and Obi, I. C.: Modelling soil properties from horizon depth functions and terrain attributes: An example with cation exchange capacity. *Eurasian Journal of Soil Science*. 9(1), 10-17. <https://doi.org/10.18393/ejss.623325>, 2020.
- Oliveira, A. K. M. Pina, J. C., Pereira, S. R., Bono, J. A. M., Matias, R., Pires, F. and Assis, T. E.: Effect of basalt rock powder associated with different substrates on the initial development of aroeira seedlings (*Myracrodruon urundeuva*). *Research, Society and Development*. 9, e5591210790, 2020. <https://doi.org/10.33448/rsd-v9i12.10790>, 2020.
- 780 Oliveira, V. H., Díez, S., Dolbeth, M. and Coelho, J. P.: Restoration of degraded estuarine and marine ecosystems: A systematic review of rehabilitation methods in Europe. *Journal of hazardous materials*. 133863. <https://doi.org/10.1016/j.jhazmat.2024.133863>, 2024.
- Oppon, E., Koh, S. L., Eufrazio, R., Nabayiga, H. and Donkor, F.: Towards sustainable food production and climate change mitigation: an attributional life cycle assessment comparing industrial and basalt rock dust fertilisers. *The International Journal of Life Cycle Assessment*. 1-12. <https://doi.org/10.1007/s11367-023-02196-4>, 2023.
- Padhiary, M. and Kumar, R.: Assessing the Environmental Impacts of Agriculture, Industrial Operations, and Mining on Agro-Ecosystems. In: Azrour, M., Mabrouki, J., Alabdulatif, A., Guezzaz, A., Amounas, F. (eds) *Smart Internet of Things*



- for Environment and Healthcare. *Studies in Computational Intelligence*, vol 1165. Springer, Cham.
 790 https://doi.org/10.1007/978-3-031-70102-3_8, 2024.
- Pahalvi, H. N., Rafiya, L., Rashid, S., Nisar, B. and Kamili, A. N.: Chemical fertilizers and their impact on soil health. *Microbiota and Biofertilizers*. 2, 1-20. https://doi.org/10.1007/978-3-030-61010-4_1, 2021.
- Peng, K., Jiang, W., Ling, Z., Hou, P. and Deng, Y.: 2021. Evaluating the potential impacts of land use changes on ecosystem service value under multiple scenarios in support of SDG reporting: A case study of the Wuhan urban
 795 agglomeration. *Journal of Cleaner Production*. 307, 127321. <https://doi.org/10.1016/j.jclepro.2021.127321>, 2021.
- Plaza C., Zaccane C., Sawicka K., Méndez A. M., Tarquis A., Gascó G. and Maestre F. T.: Soil resources and element stocks in drylands to face global issues. *Scientific Reports*. 8(1), 13788. <https://doi.org/10.1038/s41598-018-32229-0>, 2018.
- Podia, M., Yadav, P., Hooda, S., Diwan, P. and Gupta, R. K.: Microbial Weathering of Rocks in Natural Habitat: Genetic Basis and Omics-Based Exploration. *Weathering and Erosion Processes in the Natural Environment*, 265-302.
 800 <https://doi.org/10.1002/9781394157365.ch11>, 2023.
- Poggi, G. M., Corneti, S., Aloisi, I. and Ventura, F.: Environment-oriented selection criteria to overcome controversies in breeding for drought resistance in wheat. *Journal of Plant Physiology*. 280, 153895. <https://doi.org/10.1016/j.jplph.2022.153895>, 2023.
- Prasannakumar, V., Vijith, H., Abinod, S. and Geetha, N.: Estimation of soil erosion risk within a small mountainous sub-
 805 watershed in Kerala, India, using Revised Universal Soil Loss Equation (RUSLE) and geo-information technology. *Geoscience Frontiers*. 3(2), 209-215. <https://doi.org/10.1016/j.gsf.2011.11.003>, 2011.
- Rajwar, J., Joshi, D., Suyal, D. C. and Soni, R.: Factors Affecting Soil Ecosystem and Productivity. In: Soni, R., Suyal, D.C., Bhargava, P., Goel, R. (eds) *Microbiological Activity for Soil and Plant Health Management*. Springer, Singapore. https://doi.org/10.1007/978-981-16-2922-8_18, 2021.
- 810 Ramos, C. G., Querol, X., Oliveira, M. L., Pires, K., Kautzmann, R. M. and Oliveira, L. F.: A preliminary evaluation of volcanic rock powder for application in agriculture as soil a remineralizer. *Science of the Total Environment*. 512, 371-380. <https://doi.org/10.1016/j.scitotenv.2014.12.070>, 2015.
- Ramos, C. G., Dalmora, A. C., Kautzmann, R. M., Hower, J., Dotto, G. L. and Oliveira, L. F. S.: Sustainable release of macronutrients to black oat and maize crops from organically-altered dacite rock powder. *Natural Resources Research*. 30,
 815 1941-1953. <https://doi.org/10.1007/s11053-021-09862-0>, 2021.



- Ramos, C. G., Hower, J. C., Blanco, E., Oliveira, M. L. S. and Theodoro, S. H.: Possibilities of using silicate rock powder: An overview. *Geoscience Frontiers*. 13(1), 101185. <https://doi.org/10.1016/j.gsf.2021.101185>, 2022.
- Ramos, C. G., Querol, X., Dalmora, A. C., de Jesus Pires, K. C., Schneider, I. A. H., Oliveira, L. F. S. and Kautzmann, R. M.: Evaluation of the potential of volcanic rock waste from southern Brazil as a natural soil fertilizer. *Journal of Cleaner Production*. 142, 2700-2706. <https://doi.org/10.1016/j.jclepro.2016.11.006>, 2017.
- Reimann, C., Fabian, K., Flem, B., Andersson, M., Filzmoser, P. and Englmaier, P.: Geosphere-biosphere circulation of chemical elements in soil and plant systems from a 100 km transect from southern central Norway. *Science of the Total Environment*. 639, 129-145. <https://doi.org/10.1016/j.scitotenv.2018.05.070>, 2018.
- Reis, B. R., Vasconcelos, A. L. S., Silva, A. M. M., Andreote, F. D. and Azevedo, A.C.: Changes in soil bacterial community structure in a short-term trial with different silicate rock powders. *Chemical and Biological Technologies in Agriculture*, 11(1), 61. <https://doi.org/10.1186/s40538-024-00586-w>, 2024.
- Robinson, G. M.: Global sustainable agriculture and land management systems. *Geography and Sustainability*. <https://doi.org/10.1016/j.geosus.2024.09.001>, 2024.
- Rocha, J., Carvalho-Santos, C., Diogo, P., Beça, P., Keizer, J. J. and Nunes, J. P.: Impacts of climate change on reservoir water availability, quality and irrigation needs in a water scarce Mediterranean region (southern Portugal). *Science of the Total Environment*. 736, 139477. <https://doi.org/10.1016/j.scitotenv.2020.139477>, 2020.
- Rodrigues, M., Bortolini, P. C., Neto, C. K., de Andrade, E. A., dos Passos, A. I., Pacheco, F. P. and de Melo Teixeira, L.: Unlocking higher yields in *Urochloa brizantha*: the role of basalt powder in enhancing soil nutrient availability. *Discover Soil*. 1(1), 4. <https://doi.org/10.1007/s44378-024-00006-3>, 2024a
- Rodrigues, M., da Silva Junges, L. F., Mozorovicz, C., Ziemmer, G. S., Neto, C. K., de Andrade, E. A. and de Melo Teixeira, L.: Paraná Basin Basalt Powder: A Multinutrient Soil Amendment for Enhancing Soil Chemistry and Microbiology. *Journal of South American Earth Sciences*, 104957. <https://doi.org/10.1016/j.jsames.2024.104957>.
- Romanens, R., Pellacani, F., Mainga, A., Fynn, R., Vittoz, P. and Verrecchia, E. P.: Soil diversity and major soil processes in the Kalahari basin, Botswana. *Geoderma Regional*. 19, e00236. <https://doi.org/10.1016/j.geodrs.2019.e00236>, 2019.
- Rout, S., Abhilash, Meshram, P. and Zhang, P.: A comprehensive review on occurrence and processing of phosphate rock based resources-focus on REEs. *Mineral Processing and Extractive Metallurgy Review*, 45(4), 368-388. <https://doi.org/10.1080/08827508.2022.2161537>, 2024.



- Santos, L. Angélica, R. and Paz, S.: Investigation of mineral commodity residues based on alkalinity, solubility and other physicochemical aspects aiming the management of amazonian acidic soils. *Journal of Environmental Management*. 335, 117558. <https://doi.org/10.1016/j.jenvman.2023.117558>, 2023.
- Sarkar, S., Jaswal, A. and Singh, A.: Sources of inorganic nonmetallic contaminants (synthetic fertilizers, pesticides) in agricultural soil and their impacts on the adjacent ecosystems. In *Bioremediation of Emerging Contaminants from Soils*. 135-161. <https://doi.org/10.1016/B978-0-443-13993-2.00007-4>, 2024.
- Scanlan, C. A., Holmes, K. W. and Bell, R. W.: Sand and gravel subsoils. In *Subsoil Constraints for Crop Production*. 179-198. https://doi.org/10.1007/978-3-031-00317-2_8, 2022.
- Sharma, P., Sharma, P. and Thakur, N.: Sustainable farming practices and soil health: a pathway to achieving SDGs and future prospects. *Discov Sustain*. 5, 250. <https://doi.org/10.1007/s43621-024-00447-4>, 2022.
- Silva, M. B., Hermo, B. S., García-Rodeja, E. and Freire, N.V.: Reutilization of granite powder as an amendment and fertilizer for acid soils. *Chemosphere*. 61(7), 993-1002. <https://doi.org/10.1016/j.chemosphere.2005.03.010>, 2005.
- Sinha, E., Calvin, K. V., Kyle, P. G., Hejazi, M. I., Waldhoff, S. T., Huang, M. and Zhang, X.: Implication of imposing fertilizer limitations on energy, agriculture, and land systems. *Journal of environmental management*. 305, 114391. <https://doi.org/10.1016/j.jenvman.2021.114391>, 2022.
- Sivaram, A. K., Abinandan, S., Chen, C., Venkateswartlu, K. and Megharaj, M.: Microbial inoculant carriers: Soil health improvement and moisture retention in sustainable agriculture. *Advances in Agronomy*. 180, 35-91. <https://doi.org/10.1016/bs.agron.2023.03.001>, 2023.
- Sniatala, B., Kurniawan, T. A., Sobotka, D., Makinia, J. and Othman, M. H. D.: Macro-nutrients recovery from liquid waste as a sustainable resource for production of recovered mineral fertilizer: Uncovering alternative options to sustain global food security cost-effectively. *Science of The Total Environment*. 856, 159283. <https://doi.org/10.1016/j.scitotenv.2022.159283>, 2023.
- Sniatala, B., Kurniawan, T. A., Sobotka, D., Makinia, J. and Othman, M. H. D.: Macro-nutrients recovery from liquid waste as a sustainable resource for production of recovered mineral fertilizer: Uncovering alternative options to sustain global food security cost-effectively. *Science of the Total Environment*, 856, 159283. <https://doi.org/10.1016/j.scitotenv.2022.159283>, 2023.



- Soleymani, A.: Corn (*Zea mays* L.) yield and yield components as affected by light properties in response to plant parameters and N fertilization. *Biocatalysis and agricultural biotechnology*. 15, 173-180. <https://doi.org/10.1016/j.bcab.2018.06.011>, 2018.
- Suchel, J. B.: Les climats du Cameroun (these d'État). Université de Bordeaux III, 1186, 1987.
- Swoboda, P., Döring, T. F. and Hamer, M.: Remineralizing soils? The agricultural usage of silicate rock powders: A review. *Science of The Total Environment*. 807, 150976. <https://doi.org/10.1016/j.scitotenv.2021.150976>, 2022.
- 875 Sys, C.: Land Evaluation, International Training Centre for Postgraduate Soil Scientists, University of Ghent, Ghent, Belgium, 1985.
- Sys, C., Van Ranst, E., Debaveye, J. and Beernaert, F.: Land Evaluation part III. Crop requirements. *Agricultural Publications*. vol. 7, 1993.
- Tamen, J, Nkoumbou, C, Reusser, E. and Tchoua, F.: Petrology and geochemistry of mantle xenoliths from the Kapsiki plateau (Cameroon volcanic line): implications for lithospheric upwelling. *J. Afr. Earth Sci.* 101, 119–134. <https://doi.org/10.1016/j.jafrearsci.2014.09.008>, 2015.
- 880 Tamfuh, P. A., Wotchoko, P., Chotangui, A. H., Magha, A., Nono, D. G. K., Peghetmo, M. N. M. and Bitom, D.: Effect of Amending Acid Oxisols Using Basalt Dust, *Tithonia diversifolia* Powder and NPK 20-10-10 on Garlic (*Allium sativum*) Production in Bafut (Cameroon Volcanic Line). *International Journal of Plant & Soil Science*. 31(4): 1-18, Article no. IJPSS.53715, 2020.
- 885 Tamto Mamdem, E. L., Tsozué, D., Matakou, E., Apiniel Atourakail, M. R., Moudjie Noubissie, N. M., Basga, S. D. and Oyono Bitom, D. L.: Degradation Rate/Vulnerability Potential and Fertility Status of Luvisols in the Mandara Mountains (Far-North Cameroon). *The Scientific World Journal*. 2024 (1), 6565723. <https://doi.org/10.1155/2024/6565723>, 2024.
- Tetchou, A. N. T., Sonmo, J. D., Nguiffo, A., Tchouankoue, J. P., Nguetnkam, J. P. and Theodoro, S. H.: Effect of Micaschist Powder in Sugarcane (*Saccharum officinarum*) Farming on Ferrallitic Soils of Mbandjock (Cameroon, Central Africa). *Asian Journal of Soil Science and Plant Nutrition*, 8(4), 16-33. <https://doi.org/10.9734/ajsspn/2022/v8i4165>, 2022.
- 890 Theodoro, S. H., de Paula Medeiros, F., Ianniruberto, M. and Jacobson, T. K. B.: Soil remineralization and recovery of degraded areas: An experience in the tropical region. *Journal of South American Earth Sciences*, 107, 103014. <https://doi.org/10.1016/j.jsames.2020.103014>, 2021.



- 895 Thiaw, P., Sow, S. A., Faye, C. A. T., Abou Sy, A. and Sy, B. A.: Analyse de la dégradation physico-chimique des sols dans les Niayes de la Commune de Darou Khoudoss (Région de Thiès, Sénégal). *Revue Internationale de la Recherche Scientifique*. 2(3), 1253-1276. <https://doi.org/10.5281/zenodo.11660826>, 2024.
- Toor, G. S., Yang, Y. Y., Das, S., Dorsey, S. and Felton, G.: Soil health in agricultural ecosystems: current status and future perspectives. *Advances in Agronomy*. 168, 157-201. <https://doi.org/10.1016/bs.agron.2021.02.004>, 2021.
- 900 Tsozué D. and Yakouba, O.: Properties, classification, genesis and agricultural suitability of soils in a semiarid pediplain of North Cameroon. *African Journal of Agricultural Research*, 11(36), 3471-3481. <https://doi.org/10.5897/AJAR2016.11375>, 2016.
- Tsozué, D., Nafissa, B., Basga, S. D., and Balna, J.: Soil change in Arenosols under long term cultivation in the sudano-sahelian zone of Cameroon. *Geoderma Regional*. 23, e00338. <https://doi.org/10.1016/j.geodrs.2020.e00338>, 2020a.
- 905 Tsozué, D., Basga, S. D. and Nzeukou, A. N.: Spatial variation of soil weathering processes in the tropical high reliefs of Cameroon (Central Africa). *Eurasian Journal of Soil Science*. 9(2), 92-104. <https://doi.org/10.18393/ejss.659830>, 2020b.
- Tsozué, D., Nghonda, J. P. and Mekem, D. L.: Impact of land management system on crop yields and soil fertility in Cameroon. *Solid Earth*, 6(3), 1087-1101. <https://doi.org/10.5194/se-6-1087-2015>, 2015.
- Tsozué, D., Noubissie, N. M. M., Mamdem, E. L., Basga, S. D., Oyono and D. L. B.: Effects of environmental factors and soil properties on soil organic carbon stock in a natural dry tropical area of Cameroon. *Soil*. 7(2), 677-691. <https://doi.org/10.5194/soil-7-677-2021>, 2021.
- 915 Tsozué, D., Nzeugang, A. N., Mache, J. R., Loweh, S. and Fagel, N.: Mineralogical, physico-chemical and technological characterization of clays from Maroua (Far-North, Cameroon) for use in ceramic bricks production. *Journal of Building Engineering*. 11, 17-24. <https://doi.org/10.1016/j.jobbe.2017.03.008>, 2017.
- Valera, C. A., Pissarra, T. C. T., da Costa, A. M., Fernandes, L. F. S. and Pacheco, F. A. L.: The soil conservation agenda of Brazil: A review of “edge-to-edge” science contributions. *Science of the Total Environment*. 176355. <https://doi.org/10.1016/j.scitotenv.2024.176355>, 2024.
- Van Leeuwen, J. P., Lehtinen, T., Lair, G. J., Bloem, J., Hemerik, L., Ragnarsdóttir, K.V. and De Ruiter, P. C.: An ecosystem approach to assess soil quality in organically and conventionally managed farms in Iceland and Austria. *Soil*. 1(1), 83-101. <https://doi.org/10.5194/soil-1-83-2015>, 2015.
- 920



- Verdoodt, A., Van Ranst, E. and Van Averbeke, W.: Modelling crop production potentials for yield gap analysis under semiarid conditions in Guquka, South Africa. *Soil use and management*. 19(4), 372-380. <https://doi.org/10.1111/j.1475-2743.2003.tb00328.x>, 2003.
- 925 Verma, G., Dhaka, A. K., Singh, B., Kumar, A., Choudhary, A. K., Kumar, A. and Kumar, S.: Productivity, soil health, and carbon management index of soybean-wheat cropping system under double zero-tillage and natural-farming based organic nutrient management in north-Indian plains. *Science of the Total Environment*, 917, 170418. <https://doi.org/10.1016/j.scitotenv.2024.170418>, 2024.
- Verma, P., Singh, D., Pathania, I. P. and Aggarwal, K.: Strategies to improve agriculture sustainability, soil fertility and enhancement of farmers income for the economic development. *Soil fertility management for sustainable development*. 43-930 70. https://doi.org/10.1007/978-981-13-5904-0_4, 2019.
- Vienne, A., Poblador, S., Portillo-Estrada, M., Hartmann, J., Ijehon, S., Wade, P. and Vicca, S.: Enhanced weathering using basalt rock powder: carbon sequestration, co-benefits and risks in a mesocosm study with *Solanum tuberosum*. *Frontiers in Climate*, 4, 869456. <https://doi.org/10.3389/fclim.2022.869456>, 2022.
- Vondráčková, S., Hejzman, M., Tlustoš, P. and Száková, J.: Effect of rock phosphate and superphosphate application on mobility of elements (Cd, Zn, Pb, As, Fe, Mn) in contaminated soils. *Environmental Engineering & Management Journal (EEMJ)*, 16(12). <https://doi.org/10.30638/eemj.2017.299>, 2017.
- 935 Wakeel, A.: Potassium-sodium interactions in soil and plant under saline-sodic conditions. *J Plant Nutr Soil Sci* 176:344–354. https://doi.org/10.1002/jpln.20120_0417, 2013.
- Wang, X., Zhao, J., Li, C., Deng, L., Cui, R., Zhou, T. and Liu, G.: Effects of fertilizer application on the bacterial community and weathering characteristics of typical purple parent rocks. *Frontiers in Microbiology*, 15, 1514646. <https://doi.org/10.3389/fmicb.2024.1514646>, 2025.
- 940 Wu, P., Zhao, G., Huang, H., Wu, Q., Bangura, K., Cai, T. and Gao, Z. Optimizing soil and fertilizer management strategy to facilitate sustainable development of wheat production in a semi-arid area: A 12-year in-situ study on the Loess Plateau. *Field Crops Research*. 302, 109084. <https://doi.org/10.1016/j.fcr.2023.109084>, 2023.
- Xiang, T., Malik, T. H. and Nielsen, K.: The impact of population pressure on global fertiliser use intensity, 1970–2011: An analysis of policy-induced mediation. *Technological Forecasting and Social Change*. 152, 119895. <https://doi.org/10.1016/j.techfore.2019.119895>, 2020.



- Yadegari, A. H. and Etesami, H.: The Multifaceted Role of Silicon and Silicon-Solubilizing Bacteria in Sustainable Agriculture. In: de Mello Prado, R., Etesami, H., Srivastava, A.K. (eds) Silicon Advances for Sustainable Agriculture and Human Health. Sustainable Plant Nutrition in a Changing World. Springer, Cham. https://doi.org/10.1007/978-3-031-69876-7_7, 2024.
- Yang, X., Li, Y., Lu, A., Wang, H., Zhu, Y., Ding, H. and Wang, X.: Effect of *Bacillus mucilaginosus* D4B1 on the structure and soil-conservation-related properties of montmorillonite. *Applied Clay Science*, 119, 141-145. <https://doi.org/10.1016/j.clay.2015.08.033>, 2016.
- 955 Yang, J., Masoudi, A., Li, H., Gu, Y., Wang, C., Wang, M. and Liu, J.: Soil total nitrogen mediated the impact of climatic factors on urban soil organic matter under different land uses. *Journal of Soil Science and Plant Nutrition*. 1-18, 01921-8, <https://doi.org/10.1007/s42729-024-01921-8>, 2024.
- Yokoyama, T., Makishima, A. and Nakamura, E.: Separation of thorium and uranium from silicate rock samples using two commercial extraction chromatographic resins. *Anal. Chem.* 71, 135–141. <https://doi.org/10.1021/ac9805807>, 1999.

960