

Regional modeling of internal-tide dynamics around New Caledonia. Part 2: Tidal incoherence and implications for sea surface height observability

Arne Bendinger^{1,a}, Sophie Cravatte^{1,3}, Lionel Gourdeau¹, Clément Vic², and Florent Lyard¹

¹Université de Toulouse, LEGOS (CNES/CNRS/IRD/UT3), Toulouse, France

²Laboratoire d’Océanographie Physique et Spatiale, Univ. Brest, CNRS, Ifremer, IRD, IUEM, Brest, France

³IRD, Centre IRD de Nouméa, New Caledonia

^anow at: Laboratoire d’Océanographie Physique et Spatiale, Univ. Brest, CNRS, Ifremer, IRD, IUEM, Brest, France

Correspondence: Arne Bendinger (arne.bendinger@univ-brest.fr)

Abstract.

New Caledonia, in the southwestern tropical Pacific, has recently been identified as a hot spot for energetic semidiurnal internal tides. In a companion paper, the life cycle of coherent internal tides, characterized by fixed amplitude and phase, was investigated in the ~~regions~~ region through harmonic analysis of a year-long, hourly time series from numerical simulation output.

- 5 In this study, we investigate temporal variability of the internal tide by decomposing the semidiurnal signals into coherent and incoherent components. ~~We show that the departure from coherence, in both generation and propagation of internal tides, is linked to the presence of mesoscale eddies and attempt to identify the underlying mechanisms. Our results suggest that temporal variability of the~~ Semidiurnal barotropic-to-baroclinic energy conversion is largely governed by the coherent component ($>90\%$) ~~through the astronomically forced fortnightly modulated spring-neap cycle, which induces biweekly conversion~~
- 10 ~~variations, amplified~~ by a factor ~~between of 3 and 7. The incoherent component—~~ to 7 from neap to spring tides through the interaction of M2 and S2 barotropic tidal currents. Incoherent conversion—negligible in the annual mean—can explain on monthly to intraseasonal scales a notable fraction of variability ~~while reducing or enhancing, modifying~~ semidiurnal conversion by up to $\pm 20\%$. ~~We find that it is dominated~~ The latter is largely explained by local effects ~~such as, particularly~~ the work of the barotropic tide on baroclinic bottom pressure amplitude variations, ~~and~~ linked to mesoscale-eddy-induced stratification
- 15 changes. Away from the generation sites, tidal incoherence ~~becomes increasingly important, manifesting as refraction and increases, evident through~~ altered orientation of tidal beams ~~due to and increasing phase variability, caused mainly by~~ interactions with mesoscale currents ~~and varying group and phase speeds. Variations in conversion are not consistently proportional to those in energy flux divergence suggesting that variations in energy dissipation are linked to additional mechanisms that deserve further investigation.~~ The incoherent sea surface height signature, with a ~~root-mean-square~~ root-mean-squares amplitude of 1–2 cm, is widespread across the domain ~~. In tidal beam propagation direction, this incoherent component and~~
- 20 introduces limitations in ~~the spectral observability of mesoscale to submesoscale dynamics below 80 km wavelength. Along disentangling balanced (near-geostrophic) and unbalanced (wave-like) motions in spectral space. Transition scales—the length scale at which unbalanced motions become dominant over balanced motions—are proven meaningful when inferred from al-~~

timetry tracks that ~~are not aligned with the predominant propagation direction~~, the incoherent tide tends to dominate over the
25 ~~coherent tide due to its more isotropic nature. Consequently, transition scales estimating the wavelength at which balanced
motion becomes dominant over unbalanced motion should be interpreted with caution in regions with pronounced~~ align with
the main propagation direction of internal tides. However, ~~when not aligned the method is flawed as it does not take into
account the anisotropy of internal-tide activity and well-defined propagation directions.~~ dynamics.

1 Introduction

30 Internal tides, i.e., internal waves at tidal frequency, are ubiquitous in the global ocean, where major bathymetric obstacles
cause the energy transfer from the barotropic tide to baroclinic waves. These internal waves may either dissipate locally or
propagate towards the open ocean: they are argued to play an essential role in our understanding of open-ocean mixing and
the global oceanic energy budget (Melet et al., 2013; Waterhouse et al., 2014; Melet et al., 2016; de Lavergne et al., 2022).
Well-known internal-tide generation sites for low-vertical modes include Luzon Strait (e.g. Alford et al., 2011; Rainville et al.,
35 2013; Kerry et al., 2016; Wang et al., 2023), the Hawaiian Ridge (e.g. Merrifield and Holloway, 2002; Carter et al., 2008;
Zilberman et al., 2011), the Indonesian/Solomon Seas (e.g. Nagai and Hibiya, 2015; Tchilibou et al., 2020), and the Amazo-
nian Shelf Break (e.g. Tchilibou et al., 2022). From ~~semianalytical~~ semi-analytical theory (Falahat et al., 2014b; de Lavergne
et al., 2019; Vic et al., 2019), satellite altimetry (Ray and Zaron, 2016; Zhao et al., 2016; Zaron, 2019), and global numerical
simulations (Müller et al., 2012; Shriver et al., 2012; Arbic, 2022; Buijsman et al., 2017), the southwestern tropical Pacific is
40 also known to be an internal-tide generation hot spot, but has not received particular attention until very recently (Bendinger
et al., 2023, 2024).

Based on a ~~tailored regional numerical modeling effort and a full-calendar year, hourly time series regional~~ simulation out-
put, Bendinger et al. (2023, hereinafter Part 1) ~~characterized~~ analyzed the internal-tide life cycle around New Caledonia, ~~an~~
45 ~~archipelago at the entrance of the Coral Sea and an internal-tide generation hot spot of the~~ strongly dominated by the semid-
iurnal M2 tide. They ~~found that a total of estimated~~ 15.27 GW ~~is of energy~~ converted from the barotropic to baroclinic tide,
closely associated with the northwest-southeast-extending ridge system, continental slopes, shelf breaks, and seamounts. ~~More
than Over~~ 50 % of ~~the locally generated internal tides were found to dissipate close to the~~ this energy dissipates near gen-
eration sites, ~~suggesting a major contribution to vertical mixing processes. The remaining energy propagates in well-defined~~
50 ~~low-vertical mode while the remainder propagates in~~ tidal beams north and south of ~~New Caledonia towards the open ocean
with depth-integrated baroclinic energy fluxes of the island, with fluxes~~ up to 30 kW m^{-1} . ~~Overall, these tidal energetics are
well-comparable with other prominent~~ These energetics are comparable to other internal-tide generation ~~sites hot spots~~ in the
Pacific such as Luzon Strait ~~or and~~ the Hawaiian Ridge.

55 ~~One limitation of the previous analysis in Part 1 lies in the applied harmonic analysis on which the~~ The above results are
based ~~and which is on harmonic analysis~~ representative of the ~~coherent tide only. The term coherent refers here to tidal~~

stationarity, i.e., ~~phase-coherent tide only, with a~~ fixed amplitude and phase ~~for a given location, in-phase with the astronomical tide forcing, and hence, and~~ well predictable in ~~space and time~~. Further, the incoherent tide, i.e., temporal variability, ~~time and space through the astronomical tide forcing~~. Departures from tidal coherence, i.e. tidal incoherence, was not addressed.

60 ~~Tidal incoherence is defined as any departure from tidal coherence, i.e., Though, global or basin-wide numerical simulation output suggests elevated levels of tidal incoherence, which can explain well above 50 % of the semidiurnal tidal variance (Buijsman et al., 2017; Nelson et al., 2019). The phase-incoherent tide can be understood as~~ temporal variations of amplitude and phase within the tidal frequency band ~~, and therefore characterized by unpredictability and is therefore mostly unpredictable~~ (Nash et al., 2012; Kerry et al., 2013; Buijsman et al., 2017). It can represent an essential fraction of the total variance and an

65 important contribution to the internal-tide's life cycle (Zilberman et al., 2011; Buijsman et al., 2017; Zaron, 2017). As a consequence, internal-tide dissipation ~~—~~ estimated as the residual between local conversion and energy flux divergence and deduced in Part 1 ~~—~~ may have been overestimated as it contained both true baroclinic energy dissipation and energy transferred to the incoherent tide. In addition, some of the conclusions, such as those on sea surface height (SSH), should be revisited while considering the importance of tidal incoherence.

70 Sources of tidal incoherence are numerous and linked to interannual, seasonal, and mesoscale variability of stratification and background currents (Rainville and Pinkel, 2006; Tchilibou et al., 2020; Cai et al., 2024; Kaur et al., 2024). In the ~~near-field~~ near field, i.e., close to internal-tide generation sites, local stratification changes or impinging remotely generated internal tides can induce barotropic-to-baroclinic conversion variations by altering the baroclinic bottom pressure (Kelly and Nash, 2010; Zil-

75 berman et al., 2011; Kerry et al., 2014; Pickering et al., 2015; Kerry et al., 2016). In the ~~far-field~~ far field, i.e., away from the generation site and in tidal energy propagation direction, tidal incoherence is commonly induced by spatiotemporal variations of stratification and background currents such as mid-latitude and equatorial jets as well as the mesoscale eddy field (Park and Watts, 2006; Rainville and Pinkel, 2006; Dunphy and Lamb, 2014; Ponte and Klein, 2015; Kelly and Lermusiaux, 2016; Buijsman et al., 2017; Duda et al., 2018; Guo et al., 2023). The mechanisms ~~responsible for the internal-tide's temporal variability and their governing dynamics are geographically highly variable. As such, the responsible mechanisms governing temporal variability of the internal tide vary geographically, and~~ cannot be generalized ~~, as the importance of seasonality (among others) is not directly comparable across ocean basins. Further, temporal variations may be highly unpredictable, linked to mesoscale eddy variability. Quantifying these dynamics remains a challenge and demands dedicated regional studies. New Caledonia is a particularly challenging region~~ being both as it is a hot spot of ~~internal-tide generation and~~ internal tide generation and a region

85 of strong mesoscale variability, ~~potentially making it~~ subject to eddy-internal tide interactions.

Whether tidal incoherence facilitates tidal energy to dissipate is of current research interest. Generally, a loss of tidal coherence is not directly linked to tidal energy dissipation, but rather with non-linear energy transfer to the incoherent tide. Though, it is often associated with energy being transferred from low to higher vertical modes with the latter that tend to dissipate (Dunphy

90 and Lamb, 2014; Bella et al., 2024). Wang and Legg (2023) ~~explicitly~~ showed that baroclinic eddies are capable of trapping low-mode internal tides and effectively transport energy to higher-vertical modes, resulting in internal-tide dissipation within

eddies. However, a robust estimate of the fraction of energy dissipation associated with such processes does not exist and may also vary ~~from region to region~~ geographically due to the different underlying dynamics. In turn, ~~this energy dissipation associated with the incoherent tide~~ can have important implications for tidal-mixing parameterizations in climate and ocean general circulation models, which do not resolve tidal processes. Current parameterizations rely heavily on semi-analytical theory, which do not take into account temporal variations of the background stratification and currents (Vic et al., 2019; de Lavergne et al., 2019, 2020). Further effort is therefore needed to quantify the internal-tide life cycle and energy dissipation.

Internal tides are of particular interest in the context of the Surface Water Ocean Topography (SWOT) satellite altimetry mission launched in December 2022 and providing high-resolution SSH observations down to 15 km wavelength—an order of magnitude higher resolution than conventional satellite altimetry—for three main reasons (Fu and Ubelmann, 2014; Fu et al., 2024). First, SWOT will contribute to the characterization of the internal-tide SSH signature including those associated with higher vertical modes, which are associated with shorter horizontal scales than low vertical modes (Arbic et al., 2018). Second, combined SSH observations of balanced and unbalanced motions allow us to study non-linear scale interactions down to submesoscales and eventually help understand the energy transfer toward dissipation scales on global scales (Klein et al., 2019; Morrow et al., 2019). Third, accurate knowledge of the internal-tide SSH signature is crucial for the study of mesoscale and submesoscale dynamics. In other words, the extent to which we can exploit SWOT’s potential to study fine-scale oceanic dynamics depends on our ability to disentangle the measured SSH signal associated with balanced and unbalanced motions, i.e., mesoscale to submesoscale dynamics and internal gravity waves. In this regard, the transition scale serves typically as a quantity to estimate the length scale ~~below~~ L_t at which unbalanced motions become dominant over balanced motions (Vergara et al., 2023). Though, this is problematic in regions where balanced and unbalanced motions feature equal SSH variance at similar wavelengths (Callies and Wu, 2019). A correction for the coherent internal tide partially addresses this problem while increasing SSH observability of mesoscale to submesoscale dynamics (Qiu et al., 2018; Carrere et al., 2021; Arbic, 2022). The ~~empirical~~ High-Resolution Empirical Tide (HRET) model derived from ~~the~~ over 20-year long altimeter time-series is currently used to apply such correction (Zaron, 2019). However, this correction does not consider incoherent internal tides that can explain a large fraction of the total SSH variance (Zaron, 2017; Lahaye et al., 2024). This makes the dynamical interpretation of SWOT SSH, i.e., the allocation of different dynamics to SSH, a difficult challenge in regions where tidal incoherence is important.

The objective of ~~this companion paper~~ the presented study is to (1) identify regions of important tidal incoherence around New Caledonia, (2) quantify its relative importance across different time scales, and (3) understand the underlying mechanisms leading to tidal incoherence. What are the processes, both in the near- and far-field, that drive temporal variability in the barotropic-to-baroclinic conversion term and the propagation of energy? ~~What insights can be gained regarding the temporal variability of internal-tide dissipation?~~ Finally, (4) how does the incoherent tide manifest in SSH and what are potential implications for SSH observability of balanced motions in ~~New Caledonia—a region where disentangling SWOT SSH from different~~

The study is organized as follows. In Sect. 2, we give a description of the regional numerical model configuration, previously introduced in Part 1, as well as the underlying methodology used to infer temporal variability of the internal tide. In Sect. 3, we begin with an analysis of the semidiurnal tidal diagnostics, decomposed into their coherent and incoherent components. We then attribute in greater detail how mesoscale eddy variability drives temporal variability in the barotropic-to-baroclinic conversion term (Sect. 4) and energy flux (Sect. 5). In Sect. 6 we address the importance of the incoherent SSH signature around New Caledonia and what implications it can have for SSH observability of mesoscale to submesoscale dynamics. We finish with a summary and perspectives of this work in Sect. 7.

2 Data and methods

2.1 Numerical simulation

We use a regional model configuration that consists of a host grid (TROPICO12, $1/12^\circ$ horizontal resolution and 125 vertical levels) that covers the tropical and subtropical Pacific Ocean basin from 142° E- 70° W and 46° S- 24° N, introduced in Part 1. The oceanic reanalysis GLORYS2V4 prescribes initial conditions for temperature and salinity as well as the forcing with daily currents, temperature, and salinity at the open lateral boundaries. ERA5 produced by the European Centre for Medium-Range Weather Forecasts (ECMWF, Hersbach et al., 2020) provides atmospheric forcing at hourly temporal resolution and a spatial resolution of $1/4^\circ$ to compute surface fluxes using bulk formulae and the model prognostic sea surface temperature. The model is forced by the tidal potential of the five major diurnal (K1, O1) and semidiurnal (M2, S2, N2) tidal constituents. At the open lateral boundaries it is forced by barotropic SSH and barotropic currents of the same five tidal constituents taken from the global tide atlas FES2014 (Finite Element Solution 2014, Lyard et al., 2021).

We build on hourly numerical simulation outputs from a higher-resolution horizontal grid refinement within the host grid in the southwestern tropical Pacific Ocean encompassing New Caledonia (CALED060). This nesting grid features $1/60^\circ$ horizontal resolution or ~ 1.7 km grid spacing initialized by an Adaptive Grid Refinement in Fortran (AGRIF, Debreu et al., 2008). Specifically, we make use of the three-dimensional velocity, temperature, salinity, and pressure fields as well as SSH. Coherent tidal harmonics are taken from Part 1, based on harmonic analysis and vertical mode decomposition. We refer to Sect. 2.2 in Part 1 for more details. Both, full-model variables and tidal harmonics are constrained to a full-calendar year (model year 2014). Constraining the analysis to a full-calendar year relies upon a compromise between high computational expenses and a time series long enough for a robust extraction of the coherent tide.

The model's eligibility of realistically simulating ocean dynamics that range from the large-scale circulation down to high-frequency motion using climatology, satellite altimetry, and in-situ measurements was addressed in Part 1. Regarding the dominant semidiurnal M2 internal tide, confidence in the model's SSH signature is explicitly given by comparison with HRET

revealing reasonable amplitude and large-scale (interference) patterns (see ~~their~~ Fig. 13a-d in Part 1). Furthermore, glider
 160 observations revealed the model's accurate representation of the spatiotemporal variability of the semidiurnal energy flux south
 of New Caledonia, ~~i.e., including~~ the location, magnitude, and vertical structure/extent of the westward internal tide energy
 propagation characterized by narrow tidal beams (Bendinger et al., 2024).

2.2 Tidal analysis and diagnostics

~~The~~ Similarly to Part 1, the internal-tide life cycle, i.e., internal-tide generation, propagation, and dissipation, is typically studied
 165 using the following depth-integrated baroclinic energy equation (Simmons et al., 2004; Carter et al., 2008; Buijsman et al., 2017)
 :

$$\underline{T + \nabla_h \cdot \mathbf{F}_{bc} + A = C - C + D_{bc} = 0}, \quad (1)$$

where $T = \partial/\partial t \left(\int_{-H}^0 (KE_{bc} + APE) dz \right)$ is the tendency term of the total (baroclinic kinetic energy KE_{bc} and available
 170 potential energy APE), $\nabla_h \cdot \mathbf{F}_{bc}$ is the baroclinic energy flux divergence with ∇_h the horizontal gradient operator and $\mathbf{F}_{bc} =$
 (F_x, F_y) the energy flux vector, $A = \int_{-H}^0 \nabla_h \cdot (\mathbf{u}(KE_{bc} + APE + \rho_0 \mathbf{u}_{bt} \cdot \mathbf{u}_{bc})) dz$ the advection term, where $\mathbf{u} = \mathbf{u}_{bt} + \mathbf{u}_{bc}$
 is the horizontal velocity vector decomposed into its barotropic ($_{bt}$) and baroclinic ($_{bc}$) parts, and ρ_0 the reference density. C is
 the barotropic-to-baroclinic conversion term as further defined below, and D_{bc} is the baroclinic energy dissipation. Following
 Simmons et al. (2004), Carter et al. (2008), Buijsman et al. (2014, 2017), and Lahaye et al. (2020), we investigated in Part 1
 the depth-integrated baroclinic energy budget by neglecting the tendency and advection terms, D_{bc} is computed as the residual
 175 of $\nabla_h \cdot \mathbf{F}_{bc}$ and C . Further, we neglect the tendency of the total energy, i.e., \div

$$\underline{\nabla_h \cdot \mathbf{F}_{bc} - C + D_{bc} = 0}.$$

This simplification has been made since tendency and advection terms are $\partial/\partial t \left(\int_{-H}^0 (KE_{bc} + APE) dz \right)$, where KE_{bc}
 is the baroclinic kinetic energy and APE available potential energy. Furthermore, \mathbf{F}_{bc} consists only of contributions from
 hydrostatic pressure work omitting contributions from KE and APE advection, non-hydrostatic pressure work, and diffusion
 180 (Kang and Fringer, 2012; Lahaye et al., 2020). These simplifications were made since the given terms were found negligibly
 small in the above studies when averaged over a set of tidal periods and when compared to $\nabla_h \cdot \mathbf{F}_{bc}$ (Kang and Fringer, 2012)
~~. Nonetheless, and strictly speaking, D_{bc} should here be regarded as a residual of energy flux divergence and conversion.~~

In Part 1, we estimated the coherent M2 internal tide energy budget based on a harmonic analysis referenced to a full-model
 185 calendar year. Here, in Part 2 our major objective is to deduce time variability to Equation 1 for the semidiurnal frequency
 band (D^2 , 10-14 h) through a bandpass-filtering technique following Nash et al. (2012), Pickering et al. (2015), and Buijsman
 et al. (2017). Specifically, time variability is inferred by decomposing the bandpass-filtered contribution in Equation 1 into the
 coherent (coh) and incoherent (inc) parts, ~~i.e., for the horizontal velocity vector \mathbf{u} and the pressure perturbation p :~~

$$\mathbf{u}^{D2} = \mathbf{u}^{\text{coh}} + \mathbf{u}^{\text{inc}}, \quad p^{D2} = p^{\text{coh}} + p^{\text{inc}}. \quad (2)$$

190 \mathbf{u}^{D2} and p^{D2} can be further decomposed into their barotropic ~~and baroclinic~~ (bt) and ~~baroclinic~~ (bc) parts. The former is estimated by the depth-average and the latter by the departure from the barotropic depth-average:

$$\mathbf{u}_{\text{bc}}^{D2} = \mathbf{u}^{D2} - \underbrace{\frac{1}{H} \int_{-H}^{\eta} \mathbf{u}^{D2} dz}_{\mathbf{u}_{\text{bt}}^{D2}}, \quad p_{\text{bc}}^{D2} = p^{D2} - \underbrace{\frac{1}{H} \int_{-H}^{\eta} p^{D2} dz}_{p_{\text{bt}}^{D2}}, \quad (3)$$

where H is the water column depth. The semidiurnal conversion term C^{D2} can then be written as:

$$C^{D2} = \frac{1}{T} \int_T -\nabla_h H \cdot \mathbf{u}_{\text{bt}}^{D2} p_{\text{bc}}^{D2}(-H) dt \quad (4)$$

$$195 = \frac{1}{T} \int_T \left(\underbrace{-\nabla_h H \cdot \mathbf{u}_{\text{bt}}^{\text{coh}} p_{\text{bc}}^{\text{coh}}(-H)}_{C^{\text{coh}}} + \underbrace{-\nabla_h H \cdot \mathbf{u}_{\text{bt}}^{\text{inc}} p_{\text{bc}}^{\text{inc}}(-H)}_{C^{\text{inc}*}} + \underbrace{-\nabla_h H \cdot \mathbf{u}_{\text{bt}}^{\text{coh}} p_{\text{bc}}^{\text{inc}}(-H) - \nabla_h H \cdot \mathbf{u}_{\text{bt}}^{\text{inc}} p_{\text{bc}}^{\text{coh}}(-H)}_{C^{\text{cross}}} \right) dt, \quad (5)$$

decomposed into the coherent (C^{coh}) and incoherent conversion ($C^{\text{inc}} = C^{\text{inc}*} + C^{\text{cross}}$), averaged over a given time period T . ~~Note that we consider the cross terms as incoherent (i.e., $C^{\text{inc}} = C^{\text{inc}*} + C^{\text{cross}}$) since there is no preferred phasing between coherent and incoherent motions (Nash et al., 2012). Further, we define~~ Using least-squares fitting to extract the coherent component, the cross terms $C^{\text{cross}} = C^{\text{cross}1} + C^{\text{cross}2}$, where $C^{\text{cross}1} = -\nabla_h H \cdot \mathbf{u}_{\text{bt}}^{\text{coh}} p_{\text{bc}}^{\text{inc}}(-H)$ and $C^{\text{cross}2} = -\nabla_h H \cdot \mathbf{u}_{\text{bt}}^{\text{inc}} p_{\text{bc}}^{\text{coh}}(-H)$. ~~In this study, we put a particular focus on the importance of each of these terms to better understand temporal variations of the conversion term and the underlying governing processes. C^{inc} is typically considered to be negligible when averaged over long time periods (such as the annual mean) since the incoherent component of the barotropic tide forcing ($\mathbf{u}_{\text{bt}}^{\text{inc}}$) is assumed to be small on the considered time scales. Further, the unpredictable nature and random phase of incoherent processes or the interference of are, by construction, considered incoherent. This classification follows from the orthogonality condition inherent in the least-squares framework: the residual (i.e., the incoherent component) is uncorrelated with the fitted coherent signal and lacks a consistent phase relationship with the tidal forcing. In other words, there is no preferred phasing between coherent and incoherent motions cancel out in the long-term (Nash et al., 2012; Buijsman et al., 2017). As a consequence, bilinear products between coherent and incoherent fields tend to average out or remain negligibly small over the regression interval (Nash et al., 2012; Buijsman et al., 2017; Savage et al., 2020). However, we will as we show in the analysis below that these~~
 205 ~~terms are non-negligible~~, these cross terms can account for a substantial fraction of the variability on shorter time scales and, in fact, do explain an important fraction of the variability. We specifically focus on assessing their relative importance to better understand temporal variations in the conversion term and the physical mechanisms that drive them.

Similarly to C^{D2} , the semidiurnal depth-integrated energy flux \mathbf{F}_{bc}^{D2} can be written as:

$$215 \quad \mathbf{F}_{bc}^{D2} = \frac{1}{T} \int_T \int_{-H}^{\eta} \mathbf{u}_{bc}^{D2} p_{bc}^{D2} dz dt \quad (6)$$

$$= \frac{1}{T} \int_T \left(\underbrace{\int_{-H}^{\eta} \mathbf{u}_{bc}^{coh} p_{bc}^{coh} dz}_{F_{bc}^{coh}} + \underbrace{\int_{-H}^{\eta} (\mathbf{u}_{bc}^{inc} p_{bc}^{inc} + \mathbf{u}_{bc}^{coh} p_{bc}^{inc} + \mathbf{u}_{bc}^{inc} p_{bc}^{coh}) dz}_{F_{bc}^{inc}} \right) dt. \quad (7)$$

The coherent terms \mathbf{u}_{bt}^{coh} , \mathbf{u}_{bc}^{coh} , and p_{bc}^{coh} are taken from the ~~tidal analysis and the underlying~~ harmonic analysis and vertical mode decomposition carried out in Part 1. Here, they are representative of the semidiurnal frequency band ~~, i.e., the sum of~~ including the M2, S2, and N2 tidal frequencies. The barotropic part is ~~estimated by equivalent to~~ mode 0 and the baroclinic part ~~by to~~ the sum of modes 1-9. The incoherent terms are computed as the difference between the above semidiurnal bandpassed and coherent time series, i.e.,

$$\mathbf{u}_{bt}^{inc} = \mathbf{u}_{bt}^{D2} - \mathbf{u}_{bt}^{coh}, \quad \mathbf{u}_{bc}^{inc} = \mathbf{u}_{bc}^{D2} - \mathbf{u}_{bc}^{coh}, \quad p_{bc}^{inc} = p_{bc}^{D2} - p_{bc}^{coh}. \quad (8)$$

~~Dissipation is estimated through the residual between energy flux divergence and conversion as discussed above. As such,~~
 225 ~~semidiurnal energy dissipation is defined by~~ Semidiurnal energy dissipation D_{bc}^{D2} is defined as:

$$D_{bc}^{D2} = \frac{1}{T} \int_T (-\nabla_h \cdot \mathbf{F}_{bc}^{D2} + C^{D2}) dt \quad (9)$$

$$= \frac{1}{T} \int_T \underbrace{(-\nabla_h \cdot \mathbf{F}_{bc}^{coh} + C^{coh})}_{D_{bc}^{coh}} + \frac{1}{T} \int_T \underbrace{(-\nabla_h \cdot \mathbf{F}_{bc}^{inc} + C^{inc})}_{D_{bc}^{inc}} dt, \quad (10)$$

~~assuming that the contributions from the tendency and advection terms are negligibly small. A worth noting~~ A important conclusion concerning the coherent internal-tide analysis in Part 1 was that D_{bc}^{coh} may not be associated entirely with true energy
 230 dissipation. In fact, D_{bc}^{coh} represents both energy dissipation and energy transfer to the incoherent tide. It also includes the energy transfer to higher harmonics (higher frequency waves) which, however, expresses presumably in numerical dissipation due to spatial resolution constraints (Peacock and Tabaei, 2005; Zeng et al., 2021). While D_{bc}^{D2} accounts for actual energy dissipation, D_{bc}^{inc} ~~can be understood as consists of both~~ the fraction by which D_{bc}^{coh} is mistakenly associated with true energy dissipation ~~. In other words, it represents (or~~ the error by which energy dissipation in D_{bc}^{coh} is overestimated) and incoherent
 235 energy dissipation.

2.3 Ray tracing

A ray tracing method following Rainville and Pinkel (2006) is ~~used to interpret the departure from tidal coherence in the far-field, associated with the refraction of the tidal beam due to mesoscale background currents, i.e., mesoscale eddies.~~

Specifically, the theoretical employed, which models the horizontal propagation of internal gravity modes is investigated considering wave modes in the presence of spatially varying topography, climatological stratification, planetary vorticity, and depth-independent currents. Following this approach, we assume that stratification, and depth-averaged currents. Here, the objective is to interpret the tidal beam's departure from tidal coherence is primarily due to varying background currents. The choice of depth-independent currents in the far field, attributed to the refraction by mesoscale eddies. Specifically, we model semidiurnal ray paths for the first baroclinic mode for monthly varying stratification and currents. Note that the choice of depth-averaged currents is simplistic and relies on the general assumption that mesoscale eddies are well represented by a barotropic and a mode-1 baroclinic structure with limited vertical shear (Smith and Vallis, 2001) assumption that vertically sheared background currents associated with mesoscale eddies do not alter the qualitative picture of ray trajectories that are obtained. To distinguish between mesoscale stratification and current effects, we consider three different scenarios: (1) monthly varying stratification and currents, (2) monthly varying stratification and annually averaged currents, and (3) annually averaged stratification and monthly varying stratification. The reference ray path is given by annually averaged stratification and currents, representative of tidal coherence.

Bathymetry is taken from ETOPO2v2 (Smith and Sandwell, 1997). Internal gravity wave speeds are predefined and solved by the Sturm-Liouville problem for stratification from the World Ocean Atlas (Locarnini et al., 2018; Zweng et al., 2019). We model semidiurnal ray paths for modes 1-2, given stratification and bathymetry from CALED060. The semidiurnal rays are initialized at 167.75° E, 18.4° S and 167.65° E, 23.35° S within the two major internal-tide generation hot spots as identified in Part 1, i.e., the North (1) and South (2) domains (see Fig. 1), and for a given propagation angle: northeastward (45°) and southwestward (210°), respectively. In an iterative procedure, the ray tracing considers for each step size (1 km) bathymetry, climatological buoyancy and planetary vorticity effects, and the background currents. Through the dispersion relation from the Helmholtz equation for internal wave modes assuming a local wave expression, the ray's group and phase velocity are obtained, which are then used to update the ray's position and direction (angle of propagation). To mimick the effects of background currents on the ray's path, a no-currents scenario is also given. The ray tracing is applied on the depth-averaged currents as derived from the daily-mean three-dimensional velocity field from the regional model output (CALED060). The eligibility of the simplified ray tracing's methodology to infer tidal beam refraction was demonstrated in Bendinger et al. (2024). Specifically, the assumption of considering only depth-independent currents is validated a posteriori, given the relevance of the qualitative picture of ray trajectories that are obtained. stratification, and currents.

3 Annual mean of semidiurnal, coherent, and incoherent tidal diagnostics

The depth-integrated semidiurnal barotropic-to-baroclinic conversion, energy flux, and dissipation (residual) were computed for a full-model calendar year and in the full-regional in the whole domain, decomposed into the coherent and incoherent components (see Sect. 2.2). The annual mean is shown in Fig. 1. In Part 1, we identified four hot-spot regions of internal-tide




Figure 1. Annual mean, depth-integrated, semidiurnal (a) barotropic-to-baroclinic energy conversion, decomposed into the (b) coherent, and (c) incoherent components. (d-f) and (g-i) same as (a-c) but for the depth-integrated, semidiurnal energy flux and dissipation (residual), respectively. For the incoherent energy flux (f), the 2 kW m^{-1} contour is also shown. The thin black lines represent the 1000, 2000, and 3000 m depth contours. The thick black line is the 100 m depth contour representative of the New Caledonian lagoon. The numbered black boxes represent the hot spots of internal-tide generation (1: North, 2: South, 3: Norfolk Ridge, 4: Loyalty Ridge).

generation: North (1), South (2), Norfolk Ridge (3), and Loyalty Ridge (4) (see Fig. 1). Integrated over the subdomains, the barotropic-to-baroclinic conversion is almost entirely dominated by the coherent component (Fig. 1a-b, Table 1). The incoherent component is negligible (Fig. 1c). However, we will show in Sect. 4 that on shorter time scales the conversion term is

275 subject to temporal variations not linked to the astronomical tide forcing.

As stated in Part 1, the depth-integrated energy flux is characterized in the annual mean by two tidal beams that emerge and diverge from the North (1) and South (2) domains (Fig. 1d). While the coherent component is dominant, the incoherent component does explain an important fraction (Fig. 1e-f). This is particularly the case for the South (2) and Loyalty Ridge (4) domains, where the area-integrated incoherent energy flux divergence accounts for roughly 13 % and 16 % of the semidiurnal energy flux divergence, respectively (Table 1). In the North (1) and Norfolk Ridge (3) domains, it accounts for roughly 4 %. Outside and with increasing distance ~~to from~~ the generation sites, i.e., in the ~~far-field~~far field, F_{bc}^{inc} becomes more important with ratios $F_{bc}^{inc} / F_{bc}^{D2} = 0.5-0.9$ (not shown). However, it is important to note that in these cases F_{bc}^{D2} is considerably reduced ($< 5 \text{ kW m}^{-1}$).

The residual D_{bc}^{D2} is here taken as a proxy for energy dissipation following the discussion in Sect. 2.2. Integrated over the subdomains North (1), South (2), Norfolk Ridge (3), and Loyalty Ridge (4), 38 %, 58 %, 40 %, 29 % of the locally generated energy ~~are is~~ dissipated in the ~~near-field~~near field, respectively (Fig. 1g, Table 1). D_{bc}^{coh} is representative of both coherent energy dissipation and energy being removed from the coherent internal tide through non-linear energy transfers (Fig. 1h). ~~As expected, D_{bc}^{coh} is generally increased compared to D_{bc}^{D2} . The difference between both (D_{bc}^{inc}) can be understood as the fraction by which energy dissipation is overestimated in D_{bc}^{coh} or the energy gained by the consists of incoherent energy dissipation and energy transferred from the coherent tide to the incoherent tide. This amounts to~~ The latter expresses by net negative ratios of D_{bc}^{inc} and D_{bc}^{coh} and accounts for 10 %, 9 %, and 22 % in the South (2), Norfolk Ridge (3), and Loyalty Ridge (4) domains, respectively (Fig. 1i and Table 1). These ratios are equivalent to the fraction by which energy dissipation is overestimated in ~~D_{bc}^{coh} . In the North (1) domain, it D_{bc}^{inc} is slightly positive (3 %) suggesting net energy dissipation linked to the incoherent tide. This is a worth noting conclusion for Part 1, suggesting that the coherent dissipation rates for the largely dominating M2 tidal constituent may be overestimated by $\sim 10-20\%$.~~

4 ~~Semidiurnal~~What drives semidiurnal barotropic-to-baroclinic energy conversion variability?

300 4.1 Coherent vs incoherent contributions

In the annual mean, the semidiurnal barotropic-to-baroclinic conversion is largely dominated by the coherent component. In the meanwhile, it is the coherent tide ~~which that~~ is associated with a substantial amount of variability due to the ~~astronomically forced fortnightly modulated~~ spring-neap cycle, ~~i.e., driven by~~ the interaction of M2 and S2 tidal constituents (Fig. 2). Note that the N2 tidal constituent adds a low-frequency component to the modeled variability with a period of ~ 9 months. Specifically, on time scales of ~ 2 weeks conversion linked to the coherent tide may vary on average by a factor which ranges between 3 and 7 among the subdomains between spring and neap tides. This is in agreement with recent findings over the Reykjanes Ridge (Vic et al., 2021). By doing so, ~~it the coherent conversion~~ explains in the area-integral most of the semidiurnal ~~conversion~~ variability ($\gamma^{coh} = \del{90-99\%}0.90-0.99) within the internal-tide generation hot spots (Table 2). The remaining fraction is explained$

Table 1. Annual mean and standard deviation of the regional semidiurnal barotropic-to-baroclinic conversion C^{D2} , baroclinic energy flux divergence ∇F_{bc}^{D2} , and baroclinic dissipation D_{bc}^{D2} integrated over the North (1), South (2), Norfolk Ridge (3), Loyalty Ridge (4) domains, and decomposed into their coherent and incoherent parts.

| | | North | South | Norfolk Ridge | Loyalty Ridge |
|---------------------------------------|-------------|------------------|------------------|------------------|------------------|
| C^{D2} | semidiurnal | 2.79 ± 1.56 | 4.90 ± 2.58 | 2.45 ± 1.13 | 1.23 ± 0.60 |
| | coherent | 2.81 ± 1.57 | 4.97 ± 2.62 | 2.47 ± 1.13 | 1.19 ± 0.56 |
| | incoherent | -0.03 ± 0.32 | -0.07 ± 0.69 | -0.02 ± 0.42 | 0.04 ± 0.20 |
| $\nabla_h \cdot \mathbf{F}_{bc}^{D2}$ | semidiurnal | 1.72 ± 0.95 | 2.06 ± 1.11 | 1.47 ± 0.72 | 0.89 ± 0.46 |
| | coherent | 1.78 ± 0.97 | 1.80 ± 0.88 | 1.41 ± 0.61 | 0.75 ± 0.35 |
| | incoherent | -0.06 ± 0.19 | 0.26 ± 0.46 | 0.06 ± 0.28 | 0.14 ± 0.26 |
| D_{bc}^{D2} | semidiurnal | 1.06 ± 0.67 | 2.83 ± 1.57 | 0.97 ± 0.48 | 0.35 ± 0.28 |
| | coherent | 1.03 ± 0.62 | 3.17 ± 1.76 | 1.06 ± 0.55 | 0.45 ± 0.21 |
| | incoherent | 0.03 ± 0.31 | -0.33 ± 0.54 | -0.09 ± 0.31 | -0.10 ± 0.21 |

All units are given in [GW].

by the incoherent tide ($\gamma^{\text{inc}} = \text{1-10\%0.01-0.10}$).

310

We explicitly show the spatial contribution to semidiurnal conversion variability by the coherent and incoherent parts for the South (2) domain (Fig. 3). The South (2) domain represents the most prominent internal-tide generation site with conversion rates well above 1 W m^{-2} across the steep slopes such as Pines Ridge (Fig. 3a). Even though $\bar{\gamma}$ it is the coherent tide and, thus, the spring-neap cycle which dominates semidiurnal variability, certain regions stand out by increasing contributions of the incoherent tide (C^{inc}). This is the case close to the lagoon and to the southeast around seamounts, notably Antigonina, Jumeaux Est, Jumeaux Ouest, and Stylaster (Fig. 3c). In these regions, the incoherent tide can explain well above 50 % of the semidiurnal conversion variability. **ThoughYet**, it is worth mentioning that they are generally associated with reduced values of C^{D2} compared to the overwhelmingly strong generation at Pines Ridge. This is crucial knowledge for the design and interpretation of in-situ observations at fixed locations such as moorings (see Sect. 7).

320

To better understand the origin of C^{inc} , we further decompose it into a purely incoherent term ($C^{\text{inc}*}$) and two cross-terms (C^{cross1} , C^{cross2}) following Equation 5. Across all internal-tide generation hot spots, C^{cross1} dominates ($\gamma^{\text{cross1}} > \text{73\%0.73}$; Table 2). This implies that semidiurnal conversion variability (apart from the tidal-forcing induced spring-neap variability) is driven by the work of the barotropic tide on baroclinic bottom pressure variations $p_{bc}^{\text{inc}}(-H)$. C^{cross2} , linked to temporal variations of the barotropic forcing, can account for up to $\gamma^{\text{cross2}} = \text{23\%0.23}$. $C^{\text{inc}*}$ tends to play a negligible role. We note that $C^{\text{inc}*}$ and C^{cross2} can have compensating effects (see Norfolk Ridge (3) in Table 2), but the mechanisms underlying this

325




Figure 2. Time series of semidiurnal (black) barotropic-to-baroclinic conversion C^{D2} and energy flux divergence $\nabla_h \cdot \mathbf{F}_{bc}^{D2}$, decomposed into the coherent (blue) and incoherent (green) components, integrated over (a) North, (b) South, (c) Norfolk Ridge, (d) Loyalty Ridge. The mean factor of change for C^{coh} and $\nabla_h \cdot \mathbf{F}_{bc}^{coh}$ between spring and neap ~~tide~~tides (~ 2 weeks) is also given (in blue in the lower left corner).

Table 2. Domain-averaged explained variability of the coherent ($\gamma^{coh} = \text{cov}(C^{coh}, C^{D2}) / \text{var}(C^{D2})$) and incoherent (γ^{inc}) barotropic-to-baroclinic conversion referenced to the semidiurnal conversion variability (C^{D2}). The domain-averaged explained variability of the purely incoherent term $\gamma^{inc*} = \text{cov}(C^{inc*}, C^{inc}) / \text{var}(C^{inc})$ and the two cross-terms ($\gamma^{cross1}, \gamma^{cross2}$) relative to the incoherent conversion variability (C^{inc}) is also given. We refer to Equation 5 for the decomposition of C^{D2} and C^{inc} .

| | North | South | Norfolk Ridge | Loyalty Ridge |
|-------------------|-------|-------|---------------|---------------|
| γ^{coh} | 0.98 | 0.99 | 0.97 | 0.90 |
| γ^{inc} | 0.02 | 0.01 | 0.03 | 0.10 |
| γ^{inc*} | 0.04 | 0.00 | -0.01 | 0.04 |
| γ^{cross1} | 0.73 | 0.91 | 1.06 | 0.89 |
| γ^{cross2} | 0.23 | 0.09 | -0.05 | 0.07 |

compensation remain unclear.

Similarly to the [analysis](#) above, we show for the South (2) domain the contribution of the different terms ~~-, which- that~~ make up C^{inc} (Fig. 4). The annual mean for $C^{\text{inc}*}$, C^{cross1} , and C^{cross2} are shown in Fig. 4a-c. Note that the colorbar range is an order of magnitude smaller than in Fig. 1a-c, and that the overall contribution in the annual mean remains small. ~~The- While the~~ three terms feature similar amplitudes. ~~Though, the-, their~~ spatial patterns differ ~~from each other. Corresponding to-. Based on~~ the area-integrated explained variability in Table 2 ($\gamma^{\text{cross1}} = 91\% \text{ } 0.91$), and alongside high standard deviations, C^{cross1} dominates C^{inc} (Fig. 4e,h). C^{cross2} plays a minor, but non-negligible role ($\gamma^{\text{cross1}} = 9\% \text{ } 0.09$) of the incoherent variance ~~)~~ with elevated contribution particularly above steep bathymetry (Fig. 4e,h,f,i).

To summarize, incoherent conversion averages approximately to zero in the annual mean. Temporally, though, it features [marked](#) positive and negative ~~anomalies-associated-with-enhanced-and-reduced-semidiurnal-conversion-, respectively-~~ contributions [to the semidiurnal conversion at shorter time scales](#) (see Fig. 2). Decomposing C^{inc} allows for a more detailed view on the origin of conversion variations not linked to spring-neap tide variability. As expected, the work of the barotropic tide on baroclinic bottom pressure variations $p_{\text{bc}}^{\text{inc}}(-H)$, expressed by C^{cross1} is the most important. ~~To our surprise~~ [Interestingly](#), conversion variability induced by temporal variations of the barotropic forcing (C^{cross2}) is non-negligible. Temporal variations of the barotropic tide are generally known to exist through seasonal (Müller et al., 2012; Yan et al., 2020) and climatological (Opel et al., 2024) stratification changes. It may also be possible that these temporal variations represent the energy transfer from the internal tide to the barotropic tide due to pressure work (Zilberman et al., 2009). Yet, to our knowledge it remains to be quantified to what extent they may drive conversion variability. ~~Further effort is needed in this regard.~~ Also, we can not fully exclude uncertainties linked to the applied methodology, i.e., the bandpass filter and depth-average to extract the semidiurnal barotropic tide. In the following, we ~~put the~~ focus on the governing processes ~~-, which- that~~ drive baroclinic bottom pressure variations $p_{\text{bc}}^{\text{inc}}(-H)$.




Figure 3. (a) Annual-mean, depth-integrated, semidiurnal barotropic-to-baroclinic energy conversion as in Fig. 1a, but zoomed into the South (2) domain. Explained variability [by-shown for](#) the (b) coherent ($\gamma^{\text{coh}} = \text{cov}(C^{\text{coh}}, C^{\text{D2}}) / \text{var}(C^{\text{D2}})$) and (c) incoherent ($\gamma^{\text{inc}} = \text{cov}(C^{\text{inc}}, C^{\text{D2}}) / \text{var}(C^{\text{D2}})$) component. Explained variability is only shown for regions, where $|C^{\text{D2}}| > 0.05 \text{ W m}^{-2}$. The depth contour interval is 100 m. The SWOT swaths (solid black lines) during the fast sampling phase (1 d repeat orbit) are also shown.

4.2 Mesoscale-eddy-induced conversion variations

Generally speaking, $C^{\text{cross}1}$ is driven by baroclinic bottom pressure variations, which can be due to local and remote effects through local stratification changes and shoaling of remotely-generated internal tides, respectively. The former express by pressure amplitude variations (dP_A) only, whereas the latter express by both dP_A pressure phase variations (dP_ϕ) (Zilberman et al., 2011). Here, dP_A and dP_ϕ are representative of the amplitude and phase difference between the semidiurnal and coherent tide, determined by complex demodulation of $p_{bc}^{D2}(-H)$ and $p_{bc}^{\text{coh}}(-H)$. We start by focusing on Pines Ridge in the South (2) domain before generalizing our findings (Fig. 5). The monthly time series of conversion variations-anomaly expressed by the ratio of $C^{\text{cross}1}/C^{D2}$ reveals two distinct events around April/May and November 2014, during which conver-




Figure 4. Annual mean and standard deviation of depth-integrated semidiurnal barotropic-to-baroclinic incoherent energy conversion separated into the (a,d) purely incoherent ($C^{\text{inc}*}$) term, (b,e) first ($C^{\text{cross}1}$) and (c,f) second ($C^{\text{cross}2}$) cross-terms. The explained variability of (g) $C^{\text{inc}*}$, (h) $C^{\text{cross}1}$, and (i) $C^{\text{cross}2}$ are referenced to the incoherent conversion C^{inc} .

sion is ~~suppressed~~ decreased by more than 10 % ~~in the domain average~~. Note that this anomaly can be much higher locally.

360 Here, conversion variability through C^{cross1} is largely driven by baroclinic bottom pressure amplitude variations (correlation coefficient $r=0.80$ with a 90 % confidence interval $[0.50, 0.93]$ assuming $N=12$ samples) (Fig. 5a). Moreover, there is no correlation with baroclinic bottom pressure phase variations ($r=-0.02$ $[-0.51, 0.49]$). C^{cross1} does not follow a seasonal cycle. Rather, the modeled variability on monthly to intraseasonal time scales are highly suggestive of mesoscale variability.

365 ~~Following Kunze et al. (2002), stratification and pressure perturbations are linked via $p(z) = \int_z^0 N^2 \eta dz' - (1/H) \int_{-H}^0 \int_z^0 N^2 \eta dz' dz$, where η is the vertical displacement of isopycnal surfaces. Here, The monthly time series of bottom stratification $N^2(-H)$ is (extracted from the bottom most grid cell. The), mesoscale SLA, and mesoscale EKE (similarly computed to Sect. 3.2 in Part 1) suggest that conversion variations through dP_A are linked with mesoscale-eddy-induced stratification changes (Fig. 5a and 5b). Particularly around April/May and November, negative conversion/baroclinic bottom pressure amplitude anomalies in Fig. 5b are associated with negative bottom stratification anomalies and negative mesoscale SLA, alongside. The latter is further associated with elevated mesoscale EKE (Fig. 5b). Mesoscale SLA, surface geostrophic velocity, and surface mesoscale EKE are computed similarly to Sect. 3.2 in Part 1. Briefly, we computed 5-d mean SSH to eliminate high-frequency variability before binning the data onto a 1/4 horizontal grid. Finally, we applied a high-pass filter with the region's characteristic cut-off period of 180 d to account for the mesoscale. The 5-day mean of mesoscale SLA, Mesoscale SLA including the surface geostrophic~~

375 ~~velocity field, exemplarily shown for is exemplarily shown in Fig. 5c for a 5-day mean snapshot on 26 April, reveals revealing a cyclonic eddy approaching Pines Ridge (Fig. 5c).~~

~~Assuming that mesoscale eddies with strong surface signature have a depth-independent structure, i.e., barotropic or weakly baroclinic, we suggest that conversion and mesoscale variability are linked through~~ Pressure perturbations and stratification

380 ~~changes are directly linked assuming a second-order Taylor expansion of pressure around depth z_0 :~~

$$p(z) \sim p(z_0) + \left. \frac{\partial p}{\partial z} \right|_{z_0} (z - z_0) + \left. \frac{\partial^2 p}{\partial z^2} \right|_{z_0} \frac{(z - z_0)^2}{2}, \quad (11)$$

In hydrostatic balance ($\frac{\partial p}{\partial z} = -\rho g$), and expressed via stratification ($\frac{\partial^2 p}{\partial z^2} = \rho_0 N^2$), this can be written as: $p(z) \sim p(z_0) - \rho g(z - z_0) + \rho_0 N^2 \frac{(z - z_0)^2}{2}$. By taking the time derivative and assuming adiabatic motion ($\frac{\partial \rho}{\partial t} = 0$), we obtain the relation:

$$\frac{\partial p}{\partial t} \sim \rho_0 \frac{\partial N^2}{\partial t} \frac{(z - z_0)^2}{2}. \quad (12)$$

385 In practice, this translates to decreasing (increasing) stratification, which corresponds to more widely (closely) spaced isopycnals leading to weaker (stronger) baroclinic pressure anomalies. Here, negative and positive stratification anomalies above the seafloor are likely associated with the upward and downward pumping of isopycnals induced by cyclonic (CE) and anticyclonic (AE) eddy activity, respectively. In turn, the upward and downward pumping favors decreasing (increasing) stratification above

the seafloor and, thus, equivalent variations in baroclinic bottom pressure and conversion (see also Fig. A1). In phase with the local tidal forcing, $p_{bc}^{inc}(-H)$ induced by AE adds constructively to $p_{bc}^{coh}(-H)$, whereas $p_{bc}^{inc}(-H)$ induced by CE is in opposite phase and has a destructive effect. This is supported by positive correlations of pressure amplitude variations with bottom stratification ($r=0.84$ [0.59, 0.94]) and mesoscale SLA ($r=0.57$ [0.10, 0.83]) in Fig. 5b.

Generalizing our results for the whole region is far from being straightforward. We calculate the probability distribution of the correlation coefficients for the monthly time series of C^{cross1} with dP_A , dP_ϕ , $N^2(-H)$, and mesoscale SLA for each subdomain (Fig. 6). The correlation coefficients for the domain-integrated/averaged quantities (filled circles) including their 90 % confidence intervals are also given. Generally, pressure amplitude variations are very pronounced ~~suggestive of local effects~~, suggesting that local effects play an important role (Fig. 6a-d). Correlations with pressure phase variations tend to be less pronounced or more randomly distributed. Exempt therefrom is Norfolk Ridge (3), where pressure phase variations are strongly positively correlated suggesting that remote effects are important (Fig. 6c). Note that pure correlations do not provide




Figure 5. (a) Monthly time series of conversion anomaly (black) integrated over Pines Ridge (red box in (c)), expressed by the ratio C^{cross1}/C^{D2} . Also shown are the domain-averaged baroclinic bottom pressure amplitude (dP_A , red) and phase (dP_ϕ , blue) ~~difference between the semidiurnal and coherent tide differences~~ (including standard deviation). (b) Monthly time series of the domain-averaged bottom stratification ($N^2(-H)$, yellow), mesoscale SLA (gray) and EKE (green). Mesoscale SLA, surface geostrophic velocity, and surface mesoscale EKE are computed similarly to Sect. 3.2 in Part 1. Briefly, we compute 5-d mean SSH to eliminate high-frequency variability before binning the data onto a $1/4^\circ$ horizontal grid. Finally, we applied a high-pass filter with the region's characteristic cut-off period of 180 d to account for the mesoscale. The correlation coefficients of (a) C^{cross1} with dP_A , dP_ϕ and of (b) dP_A with $N^2(-H)$, mesoscale SLA including their 90 % confidence intervals are also given. (c) 5-day mean of mesoscale SLA including the geostrophic velocity field reveals a cyclonic eddy approaching Pines Ridge (red box).

information about the amplitude of C^{cross1} . High correlations indicate that the temporal patterns of variability are consistent, but they do not imply that the magnitude of the conversion is equally significant across all regions.

Assuming that local effects dominate overall, the probability distribution for the correlation of pressure amplitude variations with bottom stratification and mesoscale SLA are shown in (Fig. 6e-h). In Fig. 6f and 6g (representative of the South (2) and Norfolk Ridge (3) domains), the slightly negatively skewed probability distributions are statistically robust with the hypothesis that conversion variations are linked with mesoscale-eddy-induced stratification changes. This becomes more evident, when considering the correlation coefficients based on the area-integrated/averaged time series (see the filled circles above the panels). There are good correlations with bottom stratification (South (2): $r=0.89$ [0.71,0.96]; Norfolk Ridge (3): $r=0.80$ [0.49,0.93]) and mesoscale SLA (South (2): $r=0.45$ [-0.06,0.78]; Norfolk Ridge (3): $r=0.49$ [-0.02, 0.79]). In these regions, mesoscale-eddy-induced stratification changes can induce semidiurnal conversion anomalies by up to 20 % (see relative to the




Figure 6. Histograms of the correlation coefficient between monthly averaged C^{cross1} and baroclinic bottom pressure amplitude (dP_A , red) and phase (dP_ϕ , blue) differences ~~between the semidiurnal and coherent tide, determined by complex demodulation of $p_{bc}^{D2}(-H)$ and $p_{bc}^{\text{coh}}(-H)$~~ for (a) North, (b) South, (c) Norfolk Ridge, (d) Loyalty Ridge. We consider only regions where $|C^{D2}| > 0.2 \text{ W m}^{-2}$ to emphasize areas of strong conversion. (e-h) Same as above but for the correlation between dP_A and bottom stratification ($N^2(-H)$, yellow) and mesoscale SLA (gray). The correlation coefficients for the domain-integrated/averaged quantities (filled circles) including their 90 % confidence intervals are also given above the panels. The associated time series are explicitly shown in Fig. A2. Note that correlation coefficients based on the domain-integrated/averaged time series can largely differ from the probability distributions, as explained in Sect. 4.2.

coherent conversion (see also Fig. A2). These correlations (based on the area-integrated/averaged time series) can largely differ from the regional probability distributions, as the former may give greater weight to regions of enhanced conversion influenced by mesoscale variability. In contrast, the probability distributions represent the correlations at every grid point. We only choose
 415 grid points ~~where~~ where $|C^{D2}| > 0.2 \text{ W m}^{-2}$ to emphasize areas of strong conversion. We note that as the threshold increases, the probability distributions become more negatively skewed, indicating increasingly positive correlations of dP_A with $N^2(-H)$ and mesoscale SLA (not shown).

North (1) and Loyalty Ridge (4) show no clear correlations between pressure amplitude variations with bottom stratification
 420 and mesoscale SLA (Fig. 6e ~~and~~ 6h). Several explanations are possible. First, the assumption of depth-independent or weakly baroclinic structure may not be valid. Particularly, conversion at Loyalty Ridge (4) takes place in deeper waters (1000-3000 m compared to <1000 m in the South (2) domain) and, thus, considerably deeper than the eddies' vertical extent. Mesoscale eddies near North (1) are generally less numerous and much less energetic than south of New Caledonia (Keppler et al., 2018). There are many other factors, which can alter conversion such as seasonal stratification changes. Seasonal stratification
 425 changes have recently been reported to drive conversion variations on global scales (Kaur et al., 2024). In our study region, they seem to play a secondary role and are at best superimposed on mesoscale variability (see also Fig. A2). ~~Having said that, Other~~
local processes include the direct influence of background currents, which induce asymmetries in internal-tide generation being enhanced on the upstream side of bathymetric obstacles (Lamb and Dunphy, 2018; Shakespeare, 2020; Dossmann et al., 2020)
. Nonetheless, remote effects can play an essential role in local conversion variations, too. They include primarily the shoal-
 430 ing of remotely generated internal tides, which undergo phase modulations as they propagate through the open ocean before impinging on bathymetric slopes, which are subject to local internal-tide generation. They are usually out of phase with the local tide forcing. Though, they can theoretically be in phase or in opposite phase as well, making it hard to distinguish local from remote effects. However, remote effects seem to be of relatively small importance around New Caledonia except at minor internal-tide generation sites, which lie in propagation direction of the major tidal beams. Moreover, there are no major remote
 435 sources of internal tides, which potentially shoal on the New Caledonia ridges.

We conclude that mesoscale variability can be an important source of conversion variations by enhancing/reducing semidiurnal energy conversion within the internal-tide generation hot spots around New Caledonia. Generalizing our findings is challenging since the relative importance of the underlying dynamics can strongly vary among the generation sites. Furthermore, ~~various~~
 440 ~~different many~~ processes may be superimposed. Nonetheless, on monthly to intraseasonal time scales we attribute 10-20 % of semidiurnal conversion variations to mesoscale-eddy-induced stratification changes.

5 Tidal-Mesoscale-eddy-induced refraction of tidal beams leads to increasing tidal incoherence in the far-field

Once generated, semidiurnal internal tides propagate in narrow tidal beams towards the open ocean. Within the generation hot spots, the variability of semidiurnal energy flux divergence is closely correlated to the one that of semidiurnal conversion and, thus, follows the spring-neap tide cycle (Fig. 2). In the annual mean, tidal incoherence of energy flux divergence can account for an important fraction, which stands in contrast to the conversion term (see Sect. 3 Fig. 1f, Table 1). We will show in the following that tidal incoherence becomes increasingly important in the far-field, linked to mesoscale eddy variability around New Caledonia. First evidence is given by the 5-day monthly mean of F_{bc}^{inc} and the surface geostrophic velocity field (Fig. ??7a). In the influence area or in propagation direction of the tidal beams, elevated incoherent energy levels $>10 \text{ kW m}^{-1}$ are clearly associated with intensified mesoscale currents. This suggests that the semidiurnal energy flux becomes incoherent as it propagates through the eddy field, especially south of New Caledonia where mesoscale EKE is enhanced (Fig. ??7b). Reduced levels of incoherent energy fluxes north of New Caledonia correspond with weaker mesoscale surface currents.

(a) 5-day mean (31 January 2014, spring tide) of incoherent energy flux (F_{bc}^{inc}) overlaid by the surface geostrophic velocity field representative of the mesoscale eddy field. (b) Zonally averaged mesoscale EKE (including standard deviation) as a function of latitude. (c) Modeled semidiurnal ray paths for mode 1 propagating through the mesoscale eddy field, from their initialization north (163.75° E , 18.4° S) and south (167.65° E , 23.35° S) of New Caledonia based on daily-mean, depth-averaged background currents (model year 2014). The no-currents scenario is also given (highlighted white line).




Figure 7. (a) Monthly-mean semidiurnal incoherent energy flux (F_{bc}^{inc}) overlaid by the surface geostrophic velocity field representative of the mesoscale eddy field. (b) Zonally averaged mesoscale EKE as a function of latitude.

Table 3. Semidiurnal Cumulative refraction anomaly relative to the reference ray 's mean group arrival time (after 500 iterations, equivalent to 500 km distance given a step size of 1 km) for modes 1-2 without (w/o) annually averaged stratification and with (w/) currents) averaged over all semidiurnal rays for the first baroclinic mode initiated from the North (1) and South (2) domains .The average group arrival time delay relative to for the no-currents scenario is also given three scenarios as in Figure 8.

| | |
|---|---|
| North (1) mode-1 mode-2 w/o currents cumulative refraction [days/degrees] | 2.29 stratification + currents |
| w/ currents days North (1) | 2.38-3.00 ± 0.08-2.60 |
| South (2) | mode-1 mode-2 w/o currents days2.68-5.40 w/ currents days2.65-16.51 ± |

5.1 Mesoscale-eddy-induced refraction of tidal beam energy propagation

To better understand the underlying mechanism, we applied In the following, we apply a simplified ray tracing following Rainville and Pinkel (2006) and Bendinger et al. (2024) (see Sect. 2.3). The primary objective is to study to quantify the re-
 fraction of tidal beam energy propagation by considering only effects of depth-independent currents .To do so, we initiated a
 semidiurnal ray for two initialization regions in the North (1) and South (2) domains beams through mesoscale eddies distinguishing
 between stratification and currents (see Sect. 2.3). The theoretical mode-1 ray paths for daily-mean, depth-averaged background
 monthly averaged fields of stratification and/or currents are shown in Fig. ??e. The no-current scenario 8. The reference ray
 path for annually averaged stratification and currents is also shown(w/o currents). The ray tracing yields profoundly different
 results for the tidal beam energy propagation north and south of New Caledonia .(Fig. 8a). Theoretical rays initiated in the
 North (1) domain are confined in well-defined and narrow tidal beams, which closely align with the theoretical propagation
 direction for a semidiurnal ray in the absence of background annually averaged stratification and currents. This stands in con-
 trast to the theoretical rays initiated in the South (2) domain, experiencing notable refraction in propagation direction .For
 both regions, mode 2 is more affected by the background currents (not shown). (Table 3). The extent by which a semidiurnal
 ray is refracted by background stratification and/or currents is quantified by the cumulative refraction, for which we integrate
 absolute anomalies of the ray's propagation angle (computed from the group velocity vector). This anomaly is referenced to
 the propagation angle of the reference ray. This illustrates that rays are up to five times more refracted south of New Caledonia
 as compared to north of New Caledonia (Table 3).

Mesoscale eddy activity is elevated south of New Caledonia, explaining the contrast between the North (1) and South (2) do-
 mains and to what extent a theoretical ray is being refracted .In agreement with Keppler et al. (2018), we find that mesoscale
 eddies south of New Caledonia are more long-lived and more energetic as they feature higher rotational speeds than those north
 of New Caledonia (not shown(see Figure 7b). The physical mechanism lies in the background-currents-induced background-currents/stratifi-
 changes of the wave's group and phase speeds, which in turn cause the tidal beam to refract or change orientation, alongside
 increasing phase variability or phase offset in propagation direction relative to the no-currents scenario. Overall, there is
 increasing dispersion south of New Caledonia representing the stochastic nature of the mesoscale eddy field. Changing group
 speeds express inter alia in the wave's group arrival time at a given location (deviating from the propagation direction of the

reference ray (Fig. 8a). Here, tidal beam refraction is dominated by mesoscale currents accounting for twice as much cumulative refraction along the propagation path compared to mesoscale stratification (Fig. 8b and 8c, Table 3). The mean group arrival time for mode 1 initiated from South (2) is 2.65 ± 0.06 d, corresponding with a delay of -0.86 ± 1.44 d relative to the arrival time in the no-currents scenario. This is in agreement with recent findings (e.g. Guo et al., 2023). Overall, ray refraction by mesoscale currents leads to increasing dispersion in the no-currents scenario. Mode 2 is substantially more delayed than mode 1, roughly by a factor 4 or larger, in good agreement with the findings in Rainville and Pinkel (2006). Corresponding with the mean current direction, the theoretical rays initiated from the South (2) domain reach the far-field earlier than expected in the no-currents scenario. In contrast, the theoretical rays initiated from the North (1) domain reach the far-field later than expected.

It is worth mentioning that mesoscale eddies can be associated with stratification anomalies, which alter ray propagation. Keppler et al. (2018) reported strong anomalies in temperature and salinity within mesoscale eddies down to 1000-m depth south of New Caledonia. It remains unclear to what extent those stratification anomalies impact the propagation of semidiurnal internal tides in our region. Nonetheless, from idealized simulations it is suggested that the predominant contribution to tidal beam refraction lies in background currents, whereas stratification anomalies associated with mesoscale eddies only play secondary role (Guo et al., 2023) far field, which expresses inter alia in increasing phase variability.

Here, we associate the tidal beam's refraction with elevated mesoscale EKE (see Fig. ??b). Mesoscale activity is clearly




Figure 8. Modeled semidiurnal ray paths for the first baroclinic mode initiated at the northern (163.75° E, 18.4° S) and southern (167.65° E, 23.35° S) internal-tide generation sites of New Caledonia. Rays are computed using monthly varying and/or annually averaged fields of stratification and currents over a full calendar year. Refraction is shown for (a) monthly varying stratification and currents, (b) monthly varying stratification with annually averaged currents, and (c) monthly varying currents with annually averaged stratification. The reference ray, based on annually averaged stratification and currents, is also shown (dashed black line). The rays are further shown as a function of phase offset relative to the reference ray.

500 6 Implications of tidal incoherence for SSH observability of balanced and unbalanced motions

Internal tides typically manifest in SSH as variations on the order of a few centimeters. ~~Conventional satellite altimetry has taken advantage of this to derive robust global estimates of the coherent internal tide SSH signature using over 20 years of data.~~ In Part 1, the SSH signature of the coherent M2 internal tide was analyzed, ~~revealing multiple interference patterns as M2 tidal waves emanated from several generation sites, interacting constructively and destructively along their propagation paths.~~
505 ~~The results showed good agreement with empirical estimates from satellite altimetry (HRET; Zaron, 2019), with reasonable amplitudes and spatial patterns for both the dominant mode 1 (up to 6 cm) and mode 2 (up to 2 cm).~~

~~It remains to be investigated how the incoherent internal tides.~~ How the incoherent internal tide manifests in SSH ; particularly regarding the implications of tidal incoherence, i.e., eddy-internal tide interactions, for the observability of balanced and unbalanced motions remained to be investigated. These insights may provide crucial information for the dynamical inter-
510 pretation of SSH measurements from satellite altimeter mission such as SWOT . ~~Disentangling these dynamics represents a major challenge since SSH may contain contributions from both mesoscale to submesoscale motions and internal gravity waves. Furthermore, these dynamics can~~ while helping disentangle contributions from balanced and unbalanced motions. This is especially important in regions where these dynamics have comparable temporal and spatial scales and contribute equally to SSH variance. ~~The New Caledonia region provides an excellent study site, characterized by strong mesoscale variability and~~
515 energetic internal tides, such as New Caledonia.

The transition scale L_t , ~~provides is often used as~~ a quantitative measure to ~~distinguish between balanced and unbalanced motions (Qiu et al., 2018; Vergara et al., 2023).~~ It represents estimate the length scale ~~in spectral space~~ where unbalanced motions begin to dominate over balanced motions. ~~Transition scales are L_t is~~ commonly derived from one-dimensional SSH
520 wavenumber spectra along altimeter tracks or two-dimensional spectra from numerical simulations, revealing strong geographic and seasonal variability. For instance, submesoscale processes are typically more energetic in winter months (Callies et al., 2015; Rocha et al., 2016), while internal tides often feature amplified SSH signatures in summer due to enhanced surface stratification (Lahaye et al., 2020; Kaur et al., 2024). However, ~~the underlying methodologies have inherent limitations; particularly the assumption of isotropy. While isotropy may be a reasonable approximation for mesoscale and submesoscale~~
525 ~~circulation, it does not apply to processes such as internal tides , which propagate in~~ these methods assume isotropy in the horizontal plane for the dynamics of interest. This is reasonable for mesoscale to submesoscale processes but not for internal tides with well-defined directions. As a result, this assumption can lead propagation directions, which potentially leads to biased interpretations ~~because the methodology emphasizes isotropic motions.~~

530 In the following, we investigate the SSH imprint of the semidiurnal incoherent tide. Implications of tidal incoherence for SSH observability of balanced and unbalanced motions are deduced by computing SSH wavenumber spectra both in the direction of tidal energy propagation and along realistic altimetry tracks, we explore the implications of tidal incoherence for the observability of balanced and unbalanced motions around New Caledonia.




Figure 9. Annual root-mean-square (rms) of (a) semidiurnal SSH decomposed into the (b) coherent and (c) incoherent components. For the incoherent SSH, the 1 cm contour is shown (green). Bathymetry contours and the black boxes are given as in Fig. 1. (i) and (ii) in (a) are the transects in tidal energy propagation direction, which are used to compute SSH wavenumber spectra (see Sect. 6.3).

6.1 Semidiurnal SSH decomposed into its coherent and incoherent parts

We extend the analysis in Sect. 2.2 by decomposing semidiurnal SSH into its coherent and incoherent parts (Fig. 9). The overall signature resembles the semidiurnal energy flux in Fig. 1d-f with the predominant beams to the north and south of New Caledonia, clearly visible in SSH with root-mean-squares (rms) $\sigma > 6$ cm and dominated by the coherent tide (Fig. 9a-b). The SSH manifestation of the incoherent tide is characterized by an overall smaller but important rms (1-2 cm; Fig. 9c). The incoherent SSH is less confined to the tidal beams and seems more widespread in the domain, which suggests that the dispersion of internal waves occurs all along their propagation through the domain.

6.2 SSH wavenumber spectra in tidal beam energy propagation direction: an optimal case study

In Part 1, we calculated annually averaged SSH wavenumber spectra along the beam direction for two transects: (i) a northern transect and (ii) a southern transect (see Fig. 9a). The objective was to investigate the underlying dynamical regimes and assess the relative importance of balanced and unbalanced motions. The analysis revealed that the coherent internal tide strongly dominates SSH variance within the mesoscale band (70–250 km). The estimated transition scales for the northern and southern transects were estimated 204 km and 163 km, respectively, suggesting that, limiting SSH observability of balanced motions is primarily limited to large eddy scales. Applying By applying a correction for the coherent internal tide significantly reduced the transition scales (L_t^{corr}) to 92 km and 83 km, respectively, it was shown that this observability can be increased by shifting the transition scale to smaller wavelengths.

550

Here, we revisit this analysis by addressing the incoherent internal-tide to understand its impact on transition scales and potentially for SSH observability of balanced motions. Similarly to Part 1, we investigate SSH wavenumber spectra with regard to different dynamics that are separated in terms of frequency bands: subinertial frequencies ($\omega < f$, $\text{SSH}_{\text{subinertial}}$) for

mesoscale and submesoscale dynamics, as well as superinertial frequencies ($\omega > f$, SSH_{superinertial}) for internal gravity waves while distinguishing between the coherent internal-tide (SSH_{coh}), incoherent internal-tide (SSH_{inc}), and supertidal frequencies ($\omega > 1/10$ h, SSH_{supertidal}). Note that the incoherent internal-tide is determined by applying a bandpass filter in the full semidiurnal-diurnal tidal range (10-28 h). As such, it also comprises contributions from near-inertial, non-tidal internal gravity waves, short-live submesoscale features, but we assume that they are negligibly small.

Seasonally averaged SSH spectra for Southern Hemisphere summer (January-March, JFM) and winter (July-September, JAS) are shown in Fig. 10. By definition, the coherent internal tide is the same in both seasons. The seasonality of the transition scale corrected for the coherent internal tide (L_t^{corr}) is, thus, generally attributed to seasonal variations of subinertial motions, i.e., mesoscale to submesoscale dynamics, and unbalanced wave motions. First, we point out that the incoherent internal-tide (SSH_{inc}) predominantly governs motions at superinertial frequencies if corrected for the coherent internal-tide—at least down to 100 km independent of the season for both transects.

Along the southern transect, seasonal modulations of the SSH spectra become evident for all wavelengths < 300 km (Fig. 10c-d). In summer, it features a more flattened wavenumber slope in the mesoscale to submesoscale range with a characteristic k^{-2} slope corresponding to superinertial motions (internal wave continuum) (Fig. 10c). In winter, it becomes more continuous, characterized by a k^{-4} slope (Fig. 10d). This can be attributed to subinertial motions such as mesoscale to submesoscale processes that undergo seasonal variability. This was explicitly shown for New Caledonia in Sérazin et al. (2020) and Bendinger (2023), in which increasing importance of mixed layer instabilities and frontogenesis was attributed to more available potential energy in the Southern Hemisphere winter months. Seasonal modulations of SSH spectra can also be linked to unbalanced wave motions which are amplified in summer months, particularly for higher-vertical modes due to increasing stratification (Lahaye et al., 2020; Kaur et al., 2024). Superinertial processes dominate subinertial motions in both seasons at scales below 180 km. However, the relative importance of superinertial over subinertial motions is more pronounced in summer months. This can be explained by the seasonality of superinertial and subinertial motions being out of phase, i.e., superinertial motions are enhanced in summer while subinertial motions are considerably reduced and vice versa in winter. The transition scale (L_t) does not feature strong seasonality between summer and winter (Fig. 10c,d). Though, the transition scale is not well defined in winter, where subinertial and superinertial signals have similar variance at wavelengths 90-180 km (Fig. 10d).

~~Seasonally averaged SSH wavenumber spectra, i.e., Southern Hemisphere summer (January-March, JFM) and winter (July-September, JAS), for transects (a)-(b) north and (c)-(d) south of New Caledonia, denoted (i) and (ii), respectively, in Fig. 9a. SSH spectra are presented for the altimetry-like SSH (corrected for the barotropic tide, -, black) with regard to the different dynamics that are separated in terms of frequency bands: subinertial ($\omega < f$, -, white) for mesoscale and submesoscale dynamics, superinertial frequencies ($\omega > f$, -, solid red) for internal gravity waves decomposed into the coherent (-, blue), incoherent (-, yellow) internal tide, and supertidal frequencies ($\omega > 1/10$ h, -, gray). The altimetry-like SSH corrected for both the barotropic and baroclinic tide and filtered for motions at superinertial frequencies (-, dashed red) is also given. The characteristic wavenumber slopes k^{-2}~~

and k^{-4} are represented by the dotted black lines encompassing the mesoscale band (70-250 km), vertical dotted black lines). The transition scale L_t (i.e., where $>$) and the transition scale corrected for (L_t^{corr} , i.e., where $>$) for the annually averaged SSH spectra are specified by the red colored numbers.

Important conclusions are made when correcting for the coherent internal tide. In summer, the transition scale (L_t^{corr}) is only slightly reduced from 175 km to 156 km (Fig. 10c). This is linked to the seasonally enhanced incoherent internal tide which is still contained in the signal featuring equal SSH variance with subinertial signals at 90-180 km wavelength (Fig. 10c). At




Figure 10. Seasonally averaged SSH wavenumber spectra, i.e., Southern Hemisphere summer (January-March, JFM) and winter (July-September, JAS), for transects (a)-(b) north and (c)-(d) south of New Caledonia, denoted (i) and (ii), respectively, in Fig. 9a. SSH spectra are presented for the altimetry-like SSH (corrected for the barotropic tide, SSH, black) with regard to the different dynamics that are separated in terms of frequency bands: subinertial ($\omega < f$, SSH_{subinertial}, white) for mesoscale and submesoscale dynamics, superinertial frequencies ($\omega > f$, SSH_{superinertial}, solid red) for internal gravity waves decomposed into the coherent (SSH_{coh}, blue), incoherent (SSH_{inc}, yellow) internal tide, and supertidal frequencies ($\omega > 1/10 h$, SSH_{supertidal}, gray). The altimetry-like SSH corrected for both the barotropic and baroclinic tide and filtered for motions at superinertial frequencies (SSH_{corr_superinertial}, dashed red) is also given. The characteristic wavenumber slopes k^{-2} and k^{-4} are represented by the dotted black lines encompassing the mesoscale band (70-250 km), vertical dotted black lines). The transition scale L_t (i.e., where $\text{SSH}_{\text{superinertial}} > \text{SSH}_{\text{subinertial}}$) and the transition scale corrected for SSH_{coh} (L_t^{corr} , i.e., where $\text{SSH}_{\text{corr_superinertial}} > \text{SSH}_{\text{subinertial}}$) for the annually averaged SSH spectra are specified by the red colored numbers.

595 scales below 90 km, the incoherent tide even dominates SSH variance over subinertial motions and is equally important to the coherent tide. In winter, the transition scale is largely reduced from 165 km to 78 km (Fig. 10d). This is primarily linked to the fact that subinertial motions are energized in winter while the SSH signature of motions at superinertial frequencies are less energetics ~~as stated above~~. We note a significant contribution of motions at supertidal frequencies for scales smaller than 100 km.

600 The northern domain differs from the southern domain in that SSH variance of subinertial motions is generally reduced and the seasonal cycle less pronounced (shown by the white ~~curves~~ lines in Fig. 10a-b). Motions at superinertial frequencies largely govern over motions at subinertial frequencies throughout the year, dominated by the coherent internal tide. As for the southern transect, the incoherent contribution features seasonality and increases in summer, but remains weaker than the coherent signal. Increasing SSH observability of mesoscale and submesoscale motions by correcting for the coherent internal tide signal

605 proves overall to be more efficient since the incoherent internal tide signal is largely reduced in SSH variance compared to the southern transect (Fig. 10a-b). Specifically, the transition scale is reduced from 211 km to 146 km in summer (Fig. 10a) and from 207 km to 85 km in winter (Fig. 10b). Contributions by motions at supertidal frequencies appear to have larger importance in the northern domain compared to the southern domain. In fact, at scales below 146 km and 85 km for summer and winter respectively, SSH variance is governed by equal contributions from the incoherent internal tide and motions at supertidal

610 frequencies.

Briefly summarized, the dominance of unbalanced motions in the mesoscale to submesoscale band strongly limits SSH observability of geostrophic dynamics around New Caledonia, especially in summer. In other words, SSH observability of mesoscale dynamics is limited to large eddy scales even after a correction for the coherent tide in numerical simulation output. It is to a

615 large part the incoherent internal-tide and non-tidal internal gravity waves at scales < 100 km which may eventually constrain SSH observability of mesoscale and submesoscale dynamics.

6.3 SSH wavenumber spectra in along-track direction: a satellite altimetry perspective

The SSH wavenumber spectra in Fig. 10 and the associated conclusions for transition scales are only valid for a one-dimensional transect in tidal beam propagation direction, in which the internal-tide signature is well captured. Here, we mimic SSH

620 wavenumber spectra from satellite altimetry by interpolating SSH from CALED060 (similarly to Sect. 6.3) onto realistic altimetry tracks, which are not aligned with the tidal beam propagation direction (Fig. 11). Further, the SSH wavenumber spectra are averaged for the region south of New Caledonia (Fig. 11a).

South of New Caledonia, and along the tidal beam propagation direction, the coherent tide was found to clearly dominate

625 over subinertial motions or to be of comparable importance, regardless of the season (see Fig. 10c-d). However, averaged SSH wavenumber spectra along given altimetry tracks reveal a different picture. Spectral peaks associated with tidal motions are less pronounced (Fig. 11b-c). L_t is generally decreased, from 175 km to 115 km in summer months and from 165 km to 41 km in winter months. This reduction is linked to the anisotropic nature of the (coherent) internal tide. Along the altimetry

track direction, only a fraction of the internal tide energy is captured, causing the SSH wavenumber spectra to emphasize the more isotropic balanced flow regime. Moreover, the incoherent internal tide becomes increasingly important, dominating over the coherent tide across all spatial scales. This shift reflects the ~~greater~~ larger isotropy of incoherent SSH as compared to the coherent SSH. Consequently, in along-track spectra, the dominance of balanced motions and the incoherent internal tide renders corrections for the coherent tide ineffective. This is evident when comparing L_t and L_t^{corr} in Fig. 11b and c (from 115 km to 107 km in summer months and 41 km to 40 km in winter months).

635

We conclude that the orientation of altimetry tracks has important implications for the interpretation of transition scales computed ~~from~~ on along-track SSH wavenumber spectra. This effect is particularly pronounced in regions with prominent internal-tide motion and well-defined propagation beams, such as around New Caledonia. In such cases, transition scales may lead to erroneous estimates of the wavelength at which unbalanced motion becomes dominant over balanced motion, as anisotropic motions like internal tides are not effectively captured in along-track direction. Separating balanced from unbalanced motions is critical for the dynamical interpretation of SWOT SSH, particularly for accessing the mesoscale to submesoscale flow regime and derived quantities such as surface geostrophic velocities. ~~Therefore, caution is advised when using transition scales derived from along-track SSH to assess these dynamics.~~

645 ~~Same as in Fig. 10, but for CALED060 SSH interpolated onto satellite ground tracks, which reassemble those of SWOT, and averaged over the region south of New Caledonia as shown by the highlighted tracks in (a), separating between (b) summer and (c) winter months.~~




Figure 11. Same as in Fig. 10, but for CALED060 SSH interpolated onto satellite ground tracks, which reassemble those of SWOT, and averaged over the region south of New Caledonia as shown by the highlighted tracks in (a), separating between (b) summer and (c) winter months.

7 Summary and Perspectives

New Caledonia, an archipelago in the southwestern tropical Pacific, is a ~~low-mode~~ semidiurnal internal-tide generation hot spot as revealed by numerical simulation output from a regional model (Bendinger et al., 2023) and in-situ observations (Bendinger et al., 2024). This region is of particular interest for the SWOT altimeter mission since internal tides coexist with the mesoscale to submesoscale circulation. Being subject to potential eddy-internal tide interactions, New Caledonia represents a challenge for SWOT SSH observations. In this ~~companion paper~~ study, we investigated temporal variability of the semidiurnal internal-tide, not previously considered in Bendinger et al. (2023). Based on hourly numerical simulation output of a full-model calendar year and a bandpass-filtering technique and harmonic analysis, we decomposed the depth-integrated semidiurnal barotropic-to-baroclinic conversion, energy flux, and dissipation (residual) into their coherent and incoherent parts. These findings are summarized below.

7.1 Tidal incoherence in the ~~near-field~~ near field

~~Sketch summarizing potential processes that drive conversion variability, focusing on conversion variations due to baroclinic bottom pressure variations $p_{bc}^{inc}(-H)$. u_{bt}^{coh} is representative of the tidal forcing with the associated baroclinic bottom pressure signal $p_{bc}^{coh}(-H)$. Potential mechanisms are separated by local *1 and remote *2 effects. Local effects include stratification changes induced either by seasonal variability or mesoscale eddies. Note that the schematic illustrating mesoscale eddy-induced stratification changes is valid for the Southern Hemisphere (SH). The blue (red) curve represents the incoherent pressure signal p_{bc}^{inc} induced by CE (AE), which adds destructively (constructively) to p_{bc}^{coh} (gray). Remote effects include the shoaling of remotely generated internal tides (dotted black curve), which are phase-modulated along their propagation path (e.g. tidal beam refraction by mesoscale currents, seasonal stratification changes, etc.). See Sect. 7.1 for a detailed description.~~

In the annual mean, semidiurnal barotropic-to-baroclinic conversion is largely dominated by the coherent tide, which in turn explains a large part of the semidiurnal variability (90-99 %) through the ~~astronomically-forced fortnightly~~ spring-neap ~~tidal~~ cycle, i.e., the interaction of M2 and S2. The incoherent tide is negligibly small in the annual mean, suggesting that incoherent contributions cancel out in the long-term average. Though, locally and on shorter time scales, it can explain a notable fraction of semidiurnal variability. Our objective was to identify the underlying mechanisms responsible for conversion variations not linked to the ~~spring-neap tidal cycle~~ tidal forcing.

Generally speaking, sources of conversion variations are numerous and are difficult to distinguish due to the unpredictable nature of local and remote effects, ~~summarized in Fig. ??~~. To well distinguish between these dynamics, we separated the incoherent conversion C^{inc} into its purely incoherent term (C^{inc*}) and two cross terms (C^{cross1} , C^{cross2}) (see Equation 5). Our analysis suggests that the work done by the barotropic tide on baroclinic bottom pressure variations dominates incoherent conversion (71-90 %), expressed by C^{cross1} . The ~~majority occurrence~~ dominance of baroclinic bottom pressure amplitude variations (over phase variations) imply local effects to govern over remote effects. Locally and on monthly to intraseasonal

680 time scales, mesoscale-eddy-induced stratification changes through upward (CE) and downward (AE) pumping of isopycnal surfaces can induce negative and positive conversion anomalies by more than 20 %, respectively. This is supported by positive correlations of baroclinic bottom pressure amplitude variations with bottom stratification and mesoscale SLA. Seasonal variability seems to play a minor role or is superimposed on the dominant mesoscale variability. The importance of conversion variations through changes in the barotropic tide forcing remains to be investigated, but we showed that they can explain an
685 important fraction ~~by-of~~ up to 23 % in the incoherent conversion term.

7.2 Tidal incoherence in the ~~far-field~~far field

Semidiurnal tidal energy propagating towards the open ocean follows at first order the spring-neap tide cycle, closely coupled to semidiurnal conversion variability. However, it features elevated levels of tidal incoherence, which increases with increasing distance ~~to-from~~ the generation sites. Close to the generation sites, tidal incoherence explains up to 20 % of the
690 semidiurnal variability. In the ~~far-field~~far field, it can account for up to 90 % (in 500-1000 km distance ~~to-from~~ the generation site). Tidal incoherence ~~of the semidiurnal energy flux is here is~~ generally of higher importance south of New Caledonia corresponding with enhanced mesoscale activity, leading to dispersion in the far field alongside increasing phase variability. Further, it is associated with the refraction of the tidal beams in propagation direction. This is in agreement with a simplified ray tracing, which tracks the horizontal propagation of inertia-gravity modes at semidiurnal frequency with varying
695 ~~(depth-independent) background currents. The refraction is linked with background stratification and/or currents. Tidal beam refraction occurs through~~ varying group and phase speeds ~~induced by the background currents, which in turn cause the tidal rays to change orientation. Tidal incoherence is generally of higher importance south of New Caledonia corresponding with enhanced mesoscale activity, leading to increased dispersion and phase variability there. as the semidiurnal rays interact with the mesoscale eddy field. Changing orientation is primarily dominated by background currents accounting for twice as much~~ refraction along the propagation path as stratification.
700 North of New Caledonia, the ~~theoretical semidiurnal~~ rays closely align with the theoretical propagation direction for a semidiurnal ray in ~~the absence of background currents. This, in turn, is in agreement with reduced mesoscale activity. annually averaged stratification and currents.~~

7.3 Incoherent-tide SSH limits SSH observability of mesoscale to submesoscale motions

705 The dynamical interpretation of SSH in regions where balanced and unbalanced motions feature similar SSH variance at comparable wavelengths is challenging. By revisiting Part 1, we computed SSH wavenumber spectra in tidal energy propagation direction and extended the analysis by investigating the relative importance of the incoherent tide in SSH variance. A correction for the coherent SSH is only partly effective to access scales toward smaller wavelengths due to superinertial and subinertial motions being out of phase, i.e., superinertial motions are enhanced in summer while subinertial motions are considerably re-
710 duced and vice versa in winter. Ultimately, it is the incoherent tide, which limits SSH observability of balanced and unbalanced motions to scales above 150 km in summer and above 80 km in winter.

SSH wavenumber spectra computed along altimetry tracks lead to several conclusions. The altimetry tracks are not oriented in tidal energy propagation direction and, therefore, the spectra capture only part of the internal-tide energy. The incoherent tide dominates over the coherent tide across all wavelengths, reflecting its more isotropic nature compared to the coherent tide. Moreover, transition scales derived along altimetry tracks are generally reduced compared to those determined in tidal energy propagation direction since the SSH wavenumber spectra emphasize the more isotropic balanced flow regime. Relying on these transition scales in regions where balanced and unbalanced motions coexist may result in a distorted view of the governing dynamics as anisotropic processes such as internal tides are not properly sampled. Yet, knowing at which wavelengths balanced and unbalanced motions dominate is particularly crucial for SWOT in order to disentangle the different flow regimes in SSH measurements.

7.4 Perspectives of this work

This study provides several routes for future work to ~~increase our understanding of~~ better understand the internal-tide life cycle. ~~An open question that arises~~ One open question arising from our analysis ~~concerns the relationship between conversion variability and the impact on outward energy propagation and local energy dissipation. Specifically, is is how variability in barotropic-to-baroclinic energy conversion influences both the outward propagation of internal-tide energy and local dissipation. According to the baroclinic energy budget (Equation 1), dissipation is defined as the residual between conversion and energy flux divergence directly correlated with conversion, or are their variations. We therefore ask: do variations in conversion directly translate into proportional changes in energy flux divergence—and, by extension, in dissipation—or are they~~ partially




Figure 12. Monthly time series of semidiurnal barotropic-to-baroclinic conversion C^{D2} and energy flux divergence $\nabla_h \cdot \mathbf{F}_{bc}^{D2}$, and D_{bc}^{D2} anomalies relative to the annual mean, integrated over (a) North, (b) South, (c) Norfolk Ridge, (d) Loyalty Ridge.

730 decoupled? ~~Similarly, what can be inferred about dissipation—does higher conversion consistently enhance dissipation, and do negative anomalies reduce it~~ For instance, does increased conversion always imply stronger outward flux and higher dissipation (Falahat et al., 2014a)? Monthly anomalies of C^{D2} , $\nabla_h \cdot \mathbf{F}^{D2}$, and D^{D2} relative to the annual mean are shown in Fig 12. Energy flux divergence anomalies generally follow those of conversion, with positive conversion anomalies typically resulting in increased energy flux divergence, and vice versa. In contrast, dissipation anomalies are less variable. Nevertheless, periods
 735 exist when energy flux divergence anomalies are either more or less pronounced than conversion anomalies, leading to reduced or enhanced dissipation, respectively. The processes governing whether ~~excess or deficit~~ an excess or a deficit of tidal energy conversion is balanced primarily by outward energy propagation or by local dissipation (or other terms in Equation ~~??-1~~) remain to be fully understood.

740 This highlights the need for further investigation into the mechanisms controlling energy partitioning between these pathways. Those findings could have important implications for parameterizations of internal-tide dynamics such as tidal mixing and dissipation for climate and ocean general circulation models, which do not resolve tidal processes. Current parameterizations consider geographically varying tidal mixing (Vic et al., 2019; de Lavergne et al., 2019, 2020). However, temporal variations induced by the spring-neap cycle, mesoscale variability, and seasonal changes are not taken into account.

745 Further insight is expected from an extensive in-situ experiment (SWOTALIS, Cravatte et al., 2024) that was carried out in March-May 2023. It was inter-alia dedicated to the deployment of full-depth oceanographic moorings and located in the hot spots of internal-tide generation and dissipation south of New Caledonia. Successfully recovered in November 2023, these moorings provide a unique dataset to better understand the internal-tide life cycle, while assessing our numerical model output.
 750 Furthermore, this region is located beneath the two swaths of SWOT’s fast-sampling phase (1-day repeat orbit). The moorings and the numerical simulation output will play an essential role in the dynamical interpretation of SWOT SSH by allocating the different dynamics such as balanced and unbalanced motions. Emphasis will be given to the SSH signature of the incoherent internal tide, which represents a major challenge for SWOT SSH observability.

Appendix A

755 The hypothesis of conversion variations driven by mesoscale-eddy-induced stratification changes (see Sect. 4.2) is further supported below. We illustrate how mesoscale eddies, AE and CE, may affect the bottom stratification at the internal-tide generation site. Two examples, shown in Fig. A1, illustrate the presence of either an AE (Fig. A1a) or a CE (Fig. A1b) above Norfolk Ridge (3). These features are represented by positive and negative monthly-averaged sea level anomalies (SLA) for July and August, respectively. The zonal sections of the corresponding isopycnals indicate mesoscale-eddy-induced downward
 760 and upward pumping. The resulting stratification changes are depicted in zonally averaged stratification profiles (gray: annual mean; red: July; blue: August).

In July, the AE causes downward pumping of isopycnals, leading to positive stratification anomalies near the seafloor (Fig. A1a). Conversely, in August, the CE induces upward pumping of isopycnals, which results in negative stratification anomalies near the seafloor (Fig. A1b). It is noteworthy that the stratification anomalies just above the seafloor (at approximately 800 m depth) are significantly smaller than those in the water column above. However, they consistently align with the overall sign of the stratification anomaly.

Monthly-averaged mesoscale SLA for (a) July and (b) August showing qualitatively the impact of mesoscale variability on stratification and conversion through downward and upward pumping induced by AE and CE, respectively. The pumping is illustrated by the isopycnals along a zonal section trough the AE and CE. The zonal mean stratification profile for the given month (red: July; blue: August) as well as the annual mean (gray) is also shown.

The monthly time series of the area-integrated conversion anomaly including domain-averaged baroclinic bottom pressure amplitude and phase variations as well bottom stratification, mesoscale SLA and EKE for each subdomain are shown in Fig. A2. In the South (2) and Norfolk Ridge (3) domains, the above hypothesis is strengthened by the positive correlation coefficients (Fig. A2b and A2c). Even though we primarily link conversion variations to mesoscale variability, the monthly time series partly suggest seasonal variations of conversion. It tends to be enhanced in summer months and reduced in winter months corresponding with seasonally varying stratification. Whatsoever, mesoscale variability clearly enhances, suppresses or even reverses the seasonally-driven anomalies. A clear distinction between seasonally- and mesoscale-driven stratification changes is needed in future work to allocate the exact contribution to conversion variations.




Figure A1. Monthly averaged mesoscale SLA for (a) July and (b) August showing qualitatively the impact of mesoscale variability on stratification and conversion through downward and upward pumping induced by AE and CE, respectively. The pumping is illustrated by the isopycnals along a zonal section trough the AE and CE. The zonal mean stratification profile for the given month (red: July; blue: August) as well as the annual mean (gray) is also shown.

Monthly time series of the area-integrated conversion anomaly (black) induced by C^{cross1} expressed by the ratio of C^{inc} and C^{D2} , domain-averaged baroclinic bottom pressure amplitude (dP_A , red) and phase (dP_ϕ , blue) difference between the semidiurnal and coherent tide (including standard deviation), domain-averaged bottom stratification ($N^2(-H)$, yellow), mesoscale SLA (gray) and EKE (green) for (a) North (1), (b) South (2), (c) Norfolk Ridge (3), Loyalty Ridge (4). The correlation coefficients of C^{cross1} with dP_A , dP_ϕ and of dP_A with $N^2(-H)$, mesoscale SLA including their 90 % confidence intervals are also given.




Figure A2. Monthly time series of the area-integrated conversion anomaly (black) induced by C^{cross1} expressed by the ratio of C^{inc} and C^{D2} , domain-averaged baroclinic bottom pressure amplitude (dP_A , red) and phase (dP_ϕ , blue) difference between the semidiurnal and coherent tide (including standard deviation), domain-averaged bottom stratification ($N^2(-H)$, yellow), mesoscale SLA (gray) and EKE (green) for (a) North (1), (b) South (2), (c) Norfolk Ridge (3), Loyalty Ridge (4). The correlation coefficients of C^{cross1} with dP_A , dP_ϕ and of dP_A with $N^2(-H)$, mesoscale SLA including their 90 % confidence intervals are also given.

Code and data availability. The tidal analysis was performed using the COMODO-SIROCCO tools which are developed and maintained by the SIROCCO national service (CNRS/INSU). SIROCCO is funded by INSU and Observatoire Midi-Pyrénées/Université Paul Sabatier and receives project support from CNES, SHOM, IFREMER and ANR (<https://sirocco.obs-mip.fr/other-tools/prepost-processing/comodo-tools/>, last access: 25 August 2023). The numerical model configuration (CALED060) used in this study is introduced and described in detail in Bendinger et al. (2023). The data to reproduce the figures can be found in Bendinger (2025a) with the associated scripts in Bendinger (2025b). The ray-tracing algorithm is described in full detail in Sect. 3b in Rainville and Pinkel (2006).

Author contributions. AB performed the analysis and drafted the manuscript under the supervision of LG and SC. CV and FL were deeply involved in the discussion and interpretation of scientific results. All co-authors reviewed the manuscript and contributed to the writing and final editing.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work has been supported by the Université Toulouse III - Paul-Sabatier (grant from the Ministère de l'Enseignement supérieur de la Recherche et de l'Innovation, MESRI) carried out within the PhD program of Arne Bendinger at the Faculty of Science and Engineering and the Doctoral School of Geosciences, Astrophysics, Space and Environmental Sciences (SDU2E). Sophie Cravatte and Lionel Gourdeau are funded by the Institut de Recherche pour le Développement (IRD); Clément Vic was funded by the Institut français de recherche pour l'exploitation de la mer (IFREMER); Florent Lyard was funded by the Centre National de la Recherche Scientifique (CNRS); RC was funded by CLS. This study has been partially supported through the grant EUR TESS N°ANR-18-EURE-0018 in the framework of the Programme des Investissements d'Avenir. This work is a contribution to the joint CNES-NASA project *SWOT in the Tropics* and is supported by the French TOSCA (la Terre, l'Océan, les Surfaces Continentales, l'Atmosphère) program and the French national program LEFE (Les Enveloppes Fluides et l'Environnement).

References

- Alford, M. H., MacKinnon, J. A., Nash, J. D., Simmons, H., Pickering, A., Klymak, J. M., Pinkel, R., Sun, O., Rainville, L., Musgrave, R., et al.: Energy flux and dissipation in Luzon Strait: Two tales of two ridges, *Journal of Physical Oceanography*, 41, 2211–2222, <https://doi.org/10.1175/JPO-D-11-073.1>, 2011.
- 810 Arbic, B. K.: Incorporating tides and internal gravity waves within global ocean general circulation models: A review, *Progress in Oceanography*, 206, 102 824, <https://doi.org/https://doi.org/10.1016/j.pocean.2022.102824>, 2022.
- Arbic, B. K., Alford, M. H., Ansong, J. K., Buijsman, M. C., Ciotti, R. B., Farrar, J. T., Hallberg, R. W., Henze, C. E., Hill, C. N., Luecke, C. A., et al.: Primer on global internal tide and internal gravity wave continuum modeling in HYCOM and MITgcm, *New frontiers in operational oceanography*, pp. 307–392, http://purl.flvc.org/fsu/fd/FSU_libsubv1_scholarship_submission_1536242074_55feafcc, 2018.
- 815 Bella, A., Lahaye, N., and Tissot, G.: Internal tide energy transfers induced by mesoscale circulation and topography across the North Atlantic, *Journal of Geophysical Research: Oceans*, 129, e2024JC020 914, <https://doi.org/10.1029/2024JC020914>, 2024.
- Bendinger, A.: Internal tides around New Caledonia : dynamics, eddy-internal tide interactions, and SWOT observability, *Theses, Université Paul Sabatier - Toulouse III*, <https://theses.hal.science/tel-04618408>, 2023.
- Bendinger, A.: Regional modeling of internal-tide dynamics around New Caledonia. Part 2: Tidal incoherence and implications for sea
820 surface height observability [data set] , *Zenodo*, <https://doi.org/10.5281/zenodo.14622456>, 2025a.
- Bendinger, A.: Regional modeling of internal-tide dynamics around New Caledonia. Part 2: Tidal incoherence and implications for sea surface height observability [code] , *Zenodo*, <https://doi.org/10.5281/zenodo.14622386>, 2025b.
- Bendinger, A., Cravatte, S., Gourdeau, L., Brodeau, L., Albert, A., Tchilibou, M., Lyard, F., and Vic, C.: Regional modeling of internal-tide dynamics around New Caledonia – Part 1: Coherent internal-tide characteristics and sea surface height signature, *Ocean Science*, 19,
825 1315–1338, <https://doi.org/10.5194/os-19-1315-2023>, 2023.
- Bendinger, A., Cravatte, S., Gourdeau, L., Rainville, L., Vic, C., Sérazin, G., Durand, F., Marin, F., and Fuda, J.-L.: Internal-tide vertical structure and steric sea surface height signature south of New Caledonia revealed by glider observations, *Ocean Science*, 20, 945–964, <https://doi.org/10.5194/os-20-945-2024>, 2024.
- Buijsman, M. C., Klymak, J. M., Legg, S., Alford, M. H., Farmer, D., MacKinnon, J. A., Nash, J. D., Park, J.-H., Pickering, A., and
830 Simmons, H.: Three-dimensional double-ridge internal tide resonance in Luzon Strait, *Journal of Physical Oceanography*, 44, 850–869, <https://doi.org/10.1175/JPO-D-13-024.1>, 2014.
- Buijsman, M. C., Arbic, B. K., Richman, J. G., Shriver, J. F., Wallcraft, A. J., and Zamudio, L.: Semidiurnal internal tide incoherence in the equatorial p acific, *Journal of Geophysical Research: Oceans*, 122, 5286–5305, <https://doi.org/10.1002/2016JC012590>, 2017.
- Cai, T., Zhao, Z., D’Asaro, E., Wang, J., and Fu, L.-L.: Internal tide variability off Central California: multiple sources, seasonality, and
835 eddying background, *Journal of Geophysical Research: Oceans*, 129, e2024JC020 892, <https://doi.org/10.1029/2024JC020892>, 2024.
- Callies, J. and Wu, W.: Some expectations for submesoscale sea surface height variance spectra, *Journal of Physical Oceanography*, 49, 2271–2289, <https://doi.org/10.1175/JPO-D-18-0272.1>, 2019.
- Callies, J., Ferrari, R., Klymak, J. M., and Gula, J.: Seasonality in submesoscale turbulence, *Nature communications*, 6, 6862, <https://doi.org/10.1038/ncomms7862>, 2015.
- 840 Carrere, L., Arbic, B. K., Dushaw, B., Egbert, G., Erofeeva, S., Lyard, F., Ray, R. D., Ubelmann, C., Zaron, E., Zhao, Z., et al.: Accuracy assessment of global internal-tide models using satellite altimetry, *Ocean Science*, 17, 147–180, <https://doi.org/10.5194/os-17-147-2021>, 2021.

Carter, G. S., Merrifield, M., Becker, J. M., Katsumata, K., Gregg, M., Luther, D., Levine, M., Boyd, T. J., and Firing, Y.: Energetics of M 2 barotropic-to-baroclinic tidal conversion at the Hawaiian Islands, *Journal of Physical Oceanography*, 38, 2205–2223, <https://doi.org/10.1175/2008JPO3860.1>, 2008.

Cravatte, S., Bendinger, A., Carpaneto Bastos, C., Detandt, G., Gourdeau, L., Le Ridant, A., Rodier, M., Varillon, D., and Vic, C.: SWOTALIS-4. N/O Antea, 21 novembre au 29 novembre 2023, Nouméa/Nouméa, Ref. Rapport de mission, <https://archimer.ifremer.fr/doc/00905/101642/>, 2024.

de Lavergne, C., Falahat, S., Madec, G., Roquet, F., Nycander, J., and Vic, C.: Toward global maps of internal tide energy sinks, *Ocean Modelling*, 137, 52–75, <https://doi.org/10.1016/j.ocemod.2019.03.010>, 2019.

de Lavergne, C., Vic, C., Madec, G., Roquet, F., Waterhouse, A. F., Whalen, C., Cuypers, Y., Bouruet-Aubertot, P., Ferron, B., and Hibiya, T.: A parameterization of local and remote tidal mixing, *Journal of Advances in Modeling Earth Systems*, 12, e2020MS002065, <https://doi.org/10.1029/2020MS002065>, 2020.

de Lavergne, C., Groeskamp, S., Zika, J., and Johnson, H. L.: The role of mixing in the large-scale ocean circulation, *Ocean mixing*, pp. 35–63, <https://doi.org/10.1016/B978-0-12-821512-8.00010-4>, 2022.

Debreu, L., Vouland, C., and Blayo, E.: AGRIF: Adaptive grid refinement in Fortran, *Computers & Geosciences*, 34, 8–13, <https://doi.org/10.1016/j.cageo.2007.01.009>, 2008.

Dossmann, Y., Shakespeare, C., Stewart, K., and Hogg, A. M.: Asymmetric internal tide generation in the presence of a steady flow, *Journal of Geophysical Research: Oceans*, 125, e2020JC016503, <https://doi.org/10.1029/2020JC016503>, 2020.

Duda, T. F., Lin, Y.-T., Buijsman, M., and Newhall, A. E.: Internal tidal modal ray refraction and energy ducting in baroclinic Gulf Stream currents, *Journal of Physical Oceanography*, 48, 1969–1993, <https://doi.org/10.1175/JPO-D-18-0031.1>, 2018.

Dunphy, M. and Lamb, K. G.: Focusing and vertical mode scattering of the first mode internal tide by mesoscale eddy interaction, *Journal of Geophysical Research: Oceans*, 119, 523–536, <https://doi.org/10.1002/2013JC009293>, 2014.

Falahat, S., Nycander, J., Roquet, F., Thurnherr, A. M., and Hibiya, T.: Comparison of calculated energy flux of internal tides with microstructure measurements, *Tellus A: Dynamic Meteorology and Oceanography*, 66, 23240, <https://doi.org/10.3402/tellusa.v66.23240>, 2014a.

Falahat, S., Nycander, J., Roquet, F., and Zarroug, M.: Global calculation of tidal energy conversion into vertical normal modes, *Journal of Physical Oceanography*, 44, 3225–3244, <https://doi.org/10.1175/JPO-D-14-0002.1>, 2014b.

Fu, L.-L. and Ubelmann, C.: On the transition from profile altimeter to swath altimeter for observing global ocean surface topography, *Journal of Atmospheric and Oceanic Technology*, 31, 560–568, <https://doi.org/10.1175/JTECH-D-13-00109.1>, 2014.

Fu, L.-L., Pavelsky, T., Cretaux, J.-F., Morrow, R., Farrar, J. T., Vaze, P., Sengenès, P., Vinogradova-Shiffer, N., Sylvestre-Baron, A., Picot, N., et al.: The Surface Water and Ocean Topography Mission: A breakthrough in radar remote sensing of the ocean and land surface water, *Geophysical Research Letters*, 51, e2023GL107652, <https://doi.org/10.1029/2023GL107652>, 2024.

Guo, Z., Wang, S., Cao, A., Xie, J., Song, J., and Guo, X.: Refraction of the M2 internal tides by mesoscale eddies in the South China Sea, *Deep Sea Research Part I: Oceanographic Research Papers*, 192, 103946, <https://doi.org/10.1016/j.dsr.2022.103946>, 2023.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

Kang, D. and Fringer, O.: Energetics of barotropic and baroclinic tides in the Monterey Bay area, *Journal of Physical Oceanography*, 42, 272–290, <https://doi.org/10.1175/JPO-D-11-039.1>, 2012.

- 880 Kaur, H., Buijsman, M. C., Zhao, Z., and Shriver, J. F.: Seasonal variability in the semidiurnal internal tide—a comparison between sea surface height and energetics, *Ocean Science*, 20, 1187–1208, <https://doi.org/10.5194/os-20-1187-2024>, 2024.
- Kelly, S. and Nash, J.: Internal-tide generation and destruction by shoaling internal tides, *Geophysical Research Letters*, 37, <https://doi.org/10.1029/2010GL045598>, 2010.
- Kelly, S. M. and Lermusiaux, P. F.: Internal-tide interactions with the Gulf Stream and Middle Atlantic Bight shelfbreak front, *Journal of Geophysical Research: Oceans*, 121, 6271–6294, <https://doi.org/10.1002/2016JC011639>, 2016.
- 885 Keppler, L., Cravatte, S., Chaigneau, A., Pegliasco, C., Gourdeau, L., and Singh, A.: Observed characteristics and vertical structure of mesoscale eddies in the southwest tropical Pacific, *Journal of Geophysical Research: Oceans*, 123, 2731–2756, <https://doi.org/10.1002/2017JC013712>, 2018.
- Kerry, C. G., Powell, B. S., and Carter, G. S.: Effects of remote generation sites on model estimates of M2 internal tides in the Philippine Sea, *Journal of Physical Oceanography*, 43, 187–204, <https://doi.org/10.1175/JPO-D-12-081.1>, 2013.
- 890 Kerry, C. G., Powell, B. S., and Carter, G. S.: The impact of subtidal circulation on internal tide generation and propagation in the Philippine Sea, *Journal of Physical Oceanography*, 44, 1386–1405, <https://doi.org/10.1175/JPO-D-13-0142.1>, 2014.
- Kerry, C. G., Powell, B. S., and Carter, G. S.: Quantifying the incoherent M2 internal tide in the Philippine Sea, *Journal of Physical Oceanography*, 46, 2483–2491, <https://doi.org/10.1175/JPO-D-16-0023.1>, 2016.
- 895 Klein, P., Lapeyre, G., Siegelman, L., Qiu, B., Fu, L.-L., Torres, H., Su, Z., Menemenlis, D., and Le Gentil, S.: Ocean-scale interactions from space, *Earth and Space Science*, 6, 795–817, <https://doi.org/10.1029/2018EA000492>, 2019.
- Kunze, E., Rosenfeld, L. K., Carter, G. S., and Gregg, M. C.: Internal waves in Monterey submarine canyon, *Journal of physical oceanography*, 32, 1890–1913, [https://doi.org/10.1175/1520-0485\(2002\)032<1890:IWIMSC>2.0.CO;2](https://doi.org/10.1175/1520-0485(2002)032<1890:IWIMSC>2.0.CO;2), 2002.
- Lahaye, N., Gula, J., and Roullet, G.: Internal Tide Cycle and Topographic Scattering Over the North Mid-Atlantic Ridge, *Journal of Geophysical Research: Oceans*, 125, e2020JC016376, <https://doi.org/10.1029/2020JC016376>, 2020.
- 900 Lahaye, N., Ponte, A., Le Sommer, J., and Albert, A.: Internal tide surface signature and incoherence in the North Atlantic, *Geophysical Research Letters*, 51, e2024GL108508, <https://doi.org/10.1029/2024GL108508>, 2024.
- Lamb, K. G. and Dunphy, M.: Internal wave generation by tidal flow over a two-dimensional ridge: Energy flux asymmetries induced by a steady surface trapped current, *Journal of Fluid Mechanics*, 836, 192–221, <https://doi.org/10.1017/jfm.2017.800>, 2018.
- 905 Locarnini, M., Mishonov, A., Baranova, O., Boyer, T., Zweng, M., Garcia, H., Seidov, D., Weathers, K., Paver, C., and Smolyar, I.: World ocean atlas 2018, volume 1: Temperature, NOAA Atlas NESDIS 81, p. 52pp, <https://archimer.ifremer.fr/doc/00651/76338/>, 2018.
- Lyard, F. H., Allain, D. J., Cancet, M., Carrère, L., and Picot, N.: FES2014 global ocean tide atlas: design and performance, <https://doi.org/10.5194/os-17-615-2021>, 2021.
- Melet, A., Hallberg, R., Legg, S., and Polzin, K.: Sensitivity of the ocean state to the vertical distribution of internal-tide-driven mixing, *Journal of Physical Oceanography*, 43, 602–615, <https://doi.org/10.1175/JPO-D-12-055.1>, 2013.
- 910 Melet, A., Legg, S., and Hallberg, R.: Climatic impacts of parameterized local and remote tidal mixing, *Journal of Climate*, 29, 3473–3500, <https://doi.org/10.1175/JCLI-D-15-0153.1>, 2016.
- Merrifield, M. A. and Holloway, P. E.: Model estimates of M2 internal tide energetics at the Hawaiian Ridge, *Journal of Geophysical Research: Oceans*, 107, 5–1, <https://doi.org/10.1029/2001JC000996>, 2002.
- 915 Morrow, R., Fu, L.-L., Arduin, F., Benkiran, M., Chapron, B., Cosme, E., d’Ovidio, F., Farrar, J. T., Gille, S. T., Lapeyre, G., et al.: Global observations of fine-scale ocean surface topography with the Surface Water and Ocean Topography (SWOT) mission, *Frontiers in Marine Science*, 6, 232, <https://doi.org/10.3389/fmars.2019.00232>, 2019.

- Müller, M., Cherniawsky, J., Foreman, M., and von Storch, J.-S.: Global M2 internal tide and its seasonal variability from high resolution ocean circulation and tide modeling, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL053320>, 2012.
- 920 Nagai, T. and Hibiya, T.: Internal tides and associated vertical mixing in the Indonesian archipelago, *Journal of Geophysical Research: Oceans*, 120, 3373–3390, <https://doi.org/10.1002/2014JC010592>, 2015.
- Nash, J. D., Kelly, S. M., Shroyer, E. L., Moum, J. N., and Duda, T. F.: The unpredictable nature of internal tides on continental shelves, *Journal of Physical Oceanography*, 42, 1981–2000, <https://doi.org/10.1175/JPO-D-12-028.1>, 2012.
- Nelson, A. D., Arbic, B. K., Zaron, E. D., Savage, A. C., Richman, J. G., Buijsman, M. C., and Shriver, J. F.: Toward realistic non-stationarity of semidiurnal baroclinic tides in a hydrodynamic model, *Journal of Geophysical Research: Oceans*, 124, 6632–6642, <https://doi.org/10.1029/2018JC014737>, 2019.
- 925 Opel, L., Schindelegger, M., and Ray, R. D.: A likely role for stratification in long-term changes of the global ocean tides, *Communications Earth & Environment*, 5, 261, <https://doi.org/10.1038/s43247-024-01432-5>, 2024.
- Park, J.-H. and Watts, D. R.: Internal tides in the southwestern Japan/East Sea, *Journal of Physical Oceanography*, 36, 22–34, <https://doi.org/10.1175/JPO2846.1>, 2006.
- 930 Peacock, T. and Tabaei, A.: Visualization of nonlinear effects in reflecting internal wave beams, *Physics of Fluids*, 17, <https://doi.org/10.1063/1.1932309>, 2005.
- Pickering, A., Alford, M., Nash, J., Rainville, L., Buijsman, M., Ko, D. S., and Lim, B.: Structure and variability of internal tides in Luzon Strait, *Journal of Physical Oceanography*, 45, 1574–1594, <https://doi.org/10.1175/JPO-D-14-0250.1>, 2015.
- 935 Ponte, A. L. and Klein, P.: Incoherent signature of internal tides on sea level in idealized numerical simulations, *Geophysical Research Letters*, 42, 1520–1526, <https://doi.org/10.1002/2014GL062583>, 2015.
- Qiu, B., Chen, S., Klein, P., Wang, J., Torres, H., Fu, L.-L., and Menemenlis, D.: Seasonality in transition scale from balanced to unbalanced motions in the world ocean, *Journal of Physical Oceanography*, 48, 591–605, <https://doi.org/10.1175/JPO-D-17-0169.1>, 2018.
- Rainville, L. and Pinkel, R.: Propagation of low-mode internal waves through the ocean, *Journal of Physical Oceanography*, 36, 1220–1236, <https://doi.org/10.1175/JPO2889.1>, 2006.
- 940 Rainville, L., Lee, C. M., Rudnick, D. L., and Yang, K.-C.: Propagation of internal tides generated near Luzon Strait: Observations from autonomous gliders, *Journal of Geophysical Research: Oceans*, 118, 4125–4138, <https://doi.org/10.1002/jgrc.20293>, 2013.
- Ray, R. D. and Zaron, E. D.: M2 internal tides and their observed wavenumber spectra from satellite altimetry, *Journal of Physical Oceanography*, 46, 3–22, <https://doi.org/10.1175/JPO-D-15-0065.1>, 2016.
- 945 Rocha, C. B., Gille, S. T., Chereskin, T. K., and Menemenlis, D.: Seasonality of submesoscale dynamics in the Kuroshio Extension, *Geophysical Research Letters*, 43, 11–304, <https://doi.org/10.1002/2016GL071349>, 2016.
- Savage, A. C., Waterhouse, A. F., and Kelly, S. M.: Internal tide nonstationarity and wave–mesoscale interactions in the Tasman Sea, *Journal of Physical Oceanography*, 50, 2931–2951, <https://doi.org/10.1175/JPO-D-19-0283.1>, 2020.
- Sérazin, G., Marin, F., Gourdeau, L., Cravatte, S., Morrow, R., and Dabat, M.-L.: Scale-dependent analysis of in situ observations in the mesoscale to submesoscale range around New Caledonia, *Ocean Science*, 16, 907–925, <https://doi.org/10.5194/os-16-907-2020>, 2020.
- 950 Shakespeare, C. J.: Interdependence of internal tide and lee wave generation at abyssal hills: Global calculations, *Journal of Physical Oceanography*, 50, 655–677, <https://doi.org/10.1175/JPO-D-19-0179.1>, 2020.
- Shriver, J., Arbic, B. K., Richman, J., Ray, R., Metzger, E., Wallcraft, A., and Timko, P.: An evaluation of the barotropic and internal tides in a high-resolution global ocean circulation model, *Journal of Geophysical Research: Oceans*, 117, <https://doi.org/10.1029/2012JC008170>, 2012.
- 955

- Simmons, H. L., Hallberg, R. W., and Arbic, B. K.: Internal wave generation in a global baroclinic tide model, *Deep Sea Research Part II: Topical Studies in Oceanography*, 51, 3043–3068, <https://doi.org/10.1016/j.dsr2.2004.09.015>, 2004.
- Smith, K. S. and Vallis, G. K.: The scales and equilibration of midocean eddies: Freely evolving flow, *Journal of Physical Oceanography*, 31, 554–571, [https://doi.org/10.1175/1520-0485\(2001\)031<0554:TSAEOM>2.0.CO;2](https://doi.org/10.1175/1520-0485(2001)031<0554:TSAEOM>2.0.CO;2), 2001.
- 960 Smith, W. H. and Sandwell, D. T.: Global sea floor topography from satellite altimetry and ship depth soundings, *Science*, 277, 1956–1962, <https://doi.org/10.1126/science.277.5334.1956>, 1997.
- Tchilibou, M., Gourdeau, L., Lyard, F., Morrow, R., Koch Larrouy, A., Allain, D., and Djath, B.: Internal tides in the Solomon Sea in contrasted ENSO conditions, *Ocean Science*, 16, 615–635, <https://doi.org/10.5194/os-16-615-2020>, 2020.
- Tchilibou, M., Koch-Larrouy, A., Barbot, S., Lyard, F., Morel, Y., Jouanno, J., and Morrow, R.: Internal tides off the Amazon shelf during two
965 contrasted seasons: interactions with background circulation and SSH imprints, *Ocean Science*, 18, 1591–1618, <https://doi.org/10.5194/os-18-1591-2022>, 2022.
- Vergara, O., Morrow, R., Pujol, M.-I., Dibarboure, G., and Ubelmann, C.: Global submesoscale diagnosis using along-track satellite altimetry, *Ocean Science*, 19, 363–379, <https://doi.org/10.5194/os-19-363-2023>, 2023.
- Vic, C., Naveira Garabato, A. C., Green, J. M., Waterhouse, A. F., Zhao, Z., Melet, A., de Lavergne, C., Buijsman, M. C., and Stephenson,
970 G. R.: Deep-ocean mixing driven by small-scale internal tides, *Nature communications*, 10, 2099, <https://doi.org/10.1038/s41467-019-10149-5>, 2019.
- Vic, C., Ferron, B., Thierry, V., Mercier, H., and Lherminier, P.: Tidal and near-inertial internal waves over the Reykjanes Ridge, *Journal of Physical Oceanography*, 51, 419–437, <https://doi.org/10.1175/JPO-D-20-0097.1>, 2021.
- Wang, M., Zhu, X.-H., Zheng, H., Chen, J., Liu, Z.-J., Ren, Q., Liu, Y., Nan, F., Yu, F., and Li, Q.: Direct evidence of standing in-
975 ternal tide west of the Luzon Strait observed by a large-scale observation array, *Journal of Physical Oceanography*, 53, 2263–2280, <https://doi.org/10.1175/JPO-D-23-0043.1>, 2023.
- Wang, Y. and Legg, S.: Enhanced Dissipation of Internal Tides in a Mesoscale Baroclinic Eddy, *Journal of Physical Oceanography*, 53, 2293–2316, <https://doi.org/10.1175/JPO-D-23-0045.1>, 2023.
- Waterhouse, A. F., MacKinnon, J. A., Nash, J. D., Alford, M. H., Kunze, E., Simmons, H. L., Polzin, K. L., Laurent, L. C. S., Sun, O. M.,
980 Pinkel, R., et al.: Global patterns of diapycnal mixing from measurements of the turbulent dissipation rate, *Journal of Physical Oceanography*, 44, 1854–1872, <https://doi.org/10.1175/JPO-D-13-0104.1>, 2014.
- Yan, T., Qi, Y., Jing, Z., and Cai, S.: Seasonal and spatial features of barotropic and baroclinic tides in the northwestern South China Sea, *Journal of Geophysical Research: Oceans*, 125, e2018JC014 860, <https://doi.org/10.1029/2018JC014860>, 2020.
- Zaron, E. D.: Mapping the nonstationary internal tide with satellite altimetry, *Journal of Geophysical Research: Oceans*, 122, 539–554,
985 <https://doi.org/10.1002/2016JC012487>, 2017.
- Zaron, E. D.: Baroclinic tidal sea level from exact-repeat mission altimetry, *Journal of Physical Oceanography*, 49, 193–210, <https://doi.org/10.1175/JPO-D-18-0127.1>, 2019.
- Zeng, Z., Brandt, P., Lamb, K., Greatbatch, R., Dengler, M., Claus, M., and Chen, X.: Three-dimensional numerical simulations of internal tides in the Angolan upwelling region, *Journal of Geophysical Research: Oceans*, 126, e2020JC016 460,
990 <https://doi.org/10.1029/2020JC016460>, 2021.
- Zhao, Z., Alford, M. H., Giron, J. B., Rainville, L., and Simmons, H. L.: Global observations of open-ocean mode-1 M 2 internal tides, *Journal of Physical Oceanography*, 46, 1657–1684, <https://doi.org/10.1175/JPO-D-15-0105.1>, 2016.

- Zilberman, N., Becker, J., Merrifield, M., and Carter, G.: Model estimates of M2 internal tide generation over Mid-Atlantic Ridge topography, *Journal of Physical Oceanography*, 39, 2635–2651, <https://doi.org/10.1175/2008JPO4136.1>, 2009.
- 995 Zilberman, N., Merrifield, M., Carter, G., Luther, D., Levine, M., and Boyd, T. J.: Incoherent nature of M 2 internal tides at the Hawaiian Ridge, *Journal of physical oceanography*, 41, 2021–2036, <https://doi.org/10.1175/JPO-D-10-05009.1>, 2011.
- Zweng, M., Seidov, D., Boyer, T., Locarnini, M., Garcia, H., Mishonov, A., Baranova, O., Weathers, K., Paver, C., Smolyar, I., et al.: World ocean atlas 2018, volume 2: Salinity, NOAA Atlas NESDIS 82, p. 50pp, <https://archimer.ifremer.fr/doc/00651/76339/>, 2019.