

Evaluating the Utility of Sentinel-1 in a Data Assimilation System for Estimating Snow Depth in a Mountainous Basin

Bareera N. Mirza¹, Eric E. Small², Mark S. Raleigh¹

¹College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, OR, USA

²Geological Sciences, University of Colorado, Boulder, CO, USA

Correspondence to: Bareera N. Mirza (mirzaba@oregonstate.edu)

Abstract:

Seasonal snow plays a critical role in hydrological and energy systems, yet its high spatial and temporal variability makes accurate characterization challenging. Historically, satellite remote sensing has had limited success in mapping snow depth and snow water equivalent (SWE), particularly in global mountain areas. This study evaluates the temporal and spatial accuracy of recently developed snow depth retrievals from the Sentinel-1 (S1) C-band spaceborne radar and their utility within a data assimilation (DA) system for characterizing mountain snowpack. The DA framework integrates the physics-based Flexible Snow Model (FSM2) with a Particle Batch Smoother (PBS) to produce daily snow depth maps at a 500-meter resolution using S1 snow depth data. The S1 data were evaluated from 2017 to 2021 in and near the East River Basin, Colorado, using daily data at 12 ground-based stations for temporal evaluation and four LiDAR snow depth surveys from the Airborne Snow Observatory (ASO) for spatial evaluation. The analysis revealed significant inconsistencies in temporal and spatial errors of S1 snow depth, with higher spatial errors. Errors increased with time, especially during ablation periods, with an average temporal RMSE of 0.40 m. In contrast, the spatial RMSE exceeded 0.7 m, and S1 had poor spatial agreement with ASO LiDAR ($R^2 < 0.3$). Experiments with DA window sizes showed minimal performance differences for full-season and early-season windows. Joint assimilation of S1 snow depth with MODIS Snow Disappearance Date (SDD) yielded similar temporal errors in snow depth but degraded the performance in space relative to assimilating S1 alone. Assimilation of SDD alone outperformed S1 snow depth assimilation spatially, indicating that S1 has limited utility in a DA system. Future work should address retrieval biases, refine algorithms, and consider other snow datasets in the DA system to improve snow depth and SWE mapping in diverse snow environments globally.

1. Introduction

Seasonal snow is a natural freshwater reservoir for around 20% of the world's population (Barnett et al., 2005) and plays a key role in the global energy budget. Global mapping of snow characteristics like snow water equivalent (SWE) is a critical challenge, especially in steep and complex mountain terrain where SWE has high spatial and temporal variability (Clark et al., 2011; Kinar and Pomeroy, 2015; López-Moreno et al., 2011; Raleigh and Small, 2017). The snow community has been developing ground measurement, modeling, and remote sensing techniques through campaigns (e.g., NASA SnowEx) to advance our capabilities for estimating snow characteristics. Despite these efforts, there is still no globally available snow dataset in mountainous terrain at appropriate resolutions (~100s m or finer). Where point SWE measurements exist (e.g., stations), the high spatiotemporal variability makes it challenging to extrapolate SWE across global mountain environments (Cluzet et al., 2022; Dozier, 2011; Dozier et al., 2016; Elder et al., 1998; Grünwald et al., 2010; Herbert et al., 2024; Molotch and Bales, 2006). Improving our ability to estimate SWE in complex mountain regions requires continued advancements in global snow measurement and modeling techniques.

Optical satellite remote sensing has been used extensively in estimating snow properties. Spaceborne photogrammetry is effective for retrieving high-resolution snow depth (Deschamps-Berger et al., 2022; Marti et al., 2016; McGrath et al., 2019). However, this estimation is unavailable during cloudy periods and in dense forests. Spaceborne laser altimetry, such as ICESat and ICESat-2 has demonstrated some capability in mapping snow depth (Besso et al., 2024; Deschamps-Berger et al., 2023; Hu et al., 2022; Treichler and Kääb, 2017), but errors are high and spatial-temporal sampling is sparse. Optical remote sensing techniques for snow cover area (SCA) mapping, using sensors like Landsat, MODIS, and Sentinel-2, permit monitoring snow cover extent (e.g., Stilling et al., 2023). However, accurate mapping is hindered by cloud cover or dense vegetation (Aalstad et al., 2020). While optical remote sensing cannot measure SWE or snow depth, the information on SCA depletion can guide model-based estimates of SWE (e.g., Margulis et al., 2019; Rittger et al., 2016).

Passive microwave remote sensing retrievals have traditionally been used to retrieve snow depth (Foster et al., 1996; Kelly et al., 2003) and SWE (Chang et al., 1987; Derksen et al., 2005) however,

its coarse spatial resolution (up to ~25km) cannot capture the variability of mountain snowpack, and its accuracy is reduced in wet or deep snow conditions (over 1m; Luojus et al., 2021). In contrast, active microwave remote sensing, such as Sentinel-1's (S1) C-band synthetic aperture radar (SAR) backscatter data, has shown potential in mapping mountain snow depth at a scale of 0.5 km or finer (Lievens et al., 2019, 2022). S1 data have the potential to improve understanding of mountain snow distributions and associated processes like orographic precipitation dynamics and streamflow generation (Brangers et al., 2024; Giroto et al., 2024). Nevertheless, S1 snow depth data has two fundamental challenges: (1) the data are not temporally continuous (i.e., gaps due to satellite repeat or removed pixels flagged as wet snow), and (2) snow density is required to convert snow depth to SWE (snow depth to SWE = density x snow depth). Two possible solutions are: (1) machine learning (Broxton et al., 2024, 2019; Dunmire et al., 2024) or (2) data assimilation (DA) frameworks (Smyth et al., 2022; Alonso-González et al., 2022). Machine learning has been used to produce temporally continuous snow depth data utilizing the S1 snow depth data (Broxton et al., 2024), but may require region-specific training due to geographic variations in snowpack characteristics, making it less suitable for global mapping. In contrast, the assimilation of remotely sensed snow depth can improve snowpack estimation (including estimates of snow density) and provide temporally continuous data (Deschamps-Berger et al., 2022; Giroto et al., 2020; Largeron et al., 2020; Margulis et al., 2015; Smyth et al., 2019, 2020) . However, prior S1 DA studies have typically been limited to early-season assimilation, (Brangers et al., 2024; De Lannoy et al., 2024; Giroto et al., 2024) and little to no work has yet explored the full-season potential of the DA pathway for SWE mapping using S1 snow depth.

While S1 snow depth data have the potential to guide DA systems to produce comprehensive spatiotemporal mountain SWE maps, there have been discrepancies in the reported spatial and temporal errors, which prompt questions about its reliability. Independent evaluations have reported large errors (Broxton et al., 2024; Dunmire et al., 2024; Hoppinen et al., 2024; Sourp et al., 2025) relative to LiDAR snow depth from Airborne Snow Observatory (ASO; Painter et al., 2016) surveys and gridded snow datasets (Broxton et al., 2024). Sourp et al. (2025) found that S1 retrievals exhibit no clear error pattern but consistently underestimate snow depth, particularly before and after the melt period. Broxton et al. (2024) found mean bias from 0.27m to 0.25m when compared against ASO LiDAR flights, depending on the removal of “flagged wet pixels”. Hoppinen et al. (2024) found that these flagged pixels do not fully eliminate wet pixels and remove

some shallow dry snow pixels, resulting in a loss of usable data. Hoppinen et al. (2024) showed little to no spatial correlation between S1 and LiDAR data, contrary to $R \sim 0.52$ reported by Lievens et al. (2019) over the Northern Hemisphere. Evaluation studies (Broxton et al., 2024; Hoppinen et al., 2024) validate S1 data on a handful of dates near/after peak SWE with LiDAR surveys in the western U.S., while Lievens et al. (2022) conducted validation against point-scale time series at 1000s of locations in the European Alps and other global mountain ranges. Previous studies have shown that C-band S1 is ineffective near or after peak SWE due to the high liquid water content in the snowpack, which leads to the attenuation and absorption of microwave energy (Gagliano et al., 2023; Nagler et al., 2016), leading to increased uncertainties in the retrieved snow depth. Thus, the spatial evaluation data (airborne LiDAR) are mostly available later in the snow season when S1 data are less reliable. Gascoin et al. (2024) highlights this limitation and recommends two strategies using S1 snow depth in a DA system: (1) assimilating S1 only during the early season when snow is dry, and (2) implementing a joint assimilation of S1 snow depth along with other remote sensing data (e.g., the snow disappearance date from optical snow-covered area data). To our knowledge, no studies have tested these strategies with S1 snow depth.

This paper aims to understand the spatiotemporal error discrepancies in S1 snow depth and assess how these data can be used in a DA system, both alone and in combination with other remote sensing data. First, we evaluate the S1 snow depth data (500 m resolution) using ground-based data at 12 sites and four airborne LiDAR flights in the well-studied East River Basin (Colorado, USA). Second, we develop a DA system using a physics-based model leveraging the recent development of the open-source toolbox Multiple Snow Data Assimilation System (MuSA) (Alonso-González et al., 2022) and deploy our DA system in a cloud-based environment. Using this system, we test whether assimilation of S1 snow depth across the full season (all observations) versus early season (when it is assumed to be most reliable) improves snow depth estimation in time and space, relative to snow simulations without assimilation. Finally, we test the joint assimilation of S1 snow depth and Moderate Resolution Imaging Spectroradiometer (MODIS) snow disappearance date (SDD) (Crumley et al., 2020) following the recommendation of Gascoin et al. (2024). We address three research questions:

1. How do Sentinel-1 snow depth errors vary across space and time?
2. What is the relative value of using full-season Sentinel-1 data compared to early-season data in a data assimilation framework?
3. Can the joint assimilation of Sentinel-1 snow depth and MODIS snow disappearance date enhance DA accuracy?

2. Study Location and Data:

2.1. Study Location:

The study area is the East River Basin, Colorado (ERB Figure 1a), which has a ~748 km² area and features alpine tundra at higher elevations, montane forest at middle elevations, and prairie snow climates in lower elevation valleys (Sturm and Liston, 2021). The ERB has an average elevation of 3266m with 1,420 m topographic relief. It is characterized by cold winters, with a mean annual temperature of 0°C and mean annual precipitation of 1200 mm, mostly as snow (Daly et al., 1994; Hubbard et al., 2018). This cold, dry climate, along with moderate canopy cover, makes the ERB a representative testbed for assessing S1, which tends to struggle in complex terrain (e.g., wet snow, dense forest; Lievens et al., 2019). Furthermore, it has been well-studied and has extensive validation data through multiple scientific field campaigns, such as NASA SnowEx, ASO, the DOE SAIL (Feldman et al., 2023), and NOAA SPLASH (De Boer et al., 2023). It also includes long-term NRCS SNOTEL sites within the basin (Butte) and nearby (e.g., Schofield Pass). In addition, ground measurements are available through a research site established in late 2018 on Snodgrass Mountain (Bonner et al., 2022), which was included in the 2020 NASA SnowEx campaign. Additionally, the ERB has been mapped by airborne LiDAR flight surveys carried out by ASO (Painter et al., 2016), which provided 50 m snow depth measurements twice in two study years (2018 and 2019). The study period spans water years 2018-2021, coinciding with the availability of S1 data and ASO data.




Figure 1: (a) Elevation map of the East River Basin (Jarvis et al., 2008) with Snodgrass and SNOTEL (USDA,NRCS, 2024) stations within and near the basin. (b) SNOTEL stations within 40km from the boundary of the basin. (c) Location of the East River Basin shown relative to the Western United States.

2.2. Data

2.2.1. Meteorological forcing:

ECMWF ERA5-Land reanalysis data (Copernicus Climate Change Service, 2019; Hersbach et al., 2020) are used as the source of meteorological forcing data as they are available globally at a 0.1-degree (9 km) resolution hourly from 1950 to the present. The data, accessed from Google Cloud Storage, include incoming shortwave and longwave radiation [W m^{-2}], total precipitation (liquid and solid) [$\text{kg m}^{-2} \text{s}^{-1}$], surface atmospheric pressure [Pa], 2 m air temperature [K], 2 m relative humidity [%], and 10 m wind speed [m s^{-1}]. The data were not downscaled to the S1 500 m resolution because the DA approach implicitly accounts for fine-scale variability during assimilation (see below). In this implementation, a model ensemble is first generated at the original 9 km resolution and then regridded by nearest neighbor to 0.5 km resolution before applying the Particle Batch Smoother (PBS), effectively capturing the downscaling process (Bachand et al., 2025; Giroto et al., 2024; Smyth et al., 2019, 2020).

2.2.2. Sentinel-1 Snow Depth:

We analyzed the 500 m S1 snow depth product (Lievens et al., 2022), which is available across mountainous regions of the Northern Hemisphere from 2016 to 2021. This dataset is derived from C-Band (5.4 GHz) SAR backscatter measurements collected by the ESA Sentinel-1A and Sentinel-1B satellites. The snow depth retrieval algorithm utilizes temporal variations in the ratio between cross-polarized and co-polarized backscatter signals, which are sensitive to snow depth in dry snow. Snow depth retrievals have reported mean absolute errors of 0.18m over the Northern Hemisphere (Lievens et al., 2019). However, the S1 performance degrades in challenging conditions (snow that is wet, shallow, patchy, or in forests) with a mean absolute error of about 1.3m at a depth of 3.5m and above (Lievens et al., 2022). Evaluation and development of this dataset is ongoing. The latest 500m dataset includes a quality flag for wet snow conditions, which has potential use for identifying and removing pixels with higher retrieval errors, particularly during the ablation season. Wet snow leads to the absorption and attenuation of radar signals, which makes snow depth mapping unreliable. The frequency of observations varies by region depending on the satellite overpass schedule. However, the dataset offers temporal resolution primarily on a daily to weekly basis, with most intervals being less than one week over the ERB.

2.2.3. Snow Disappearance Date (SDD):

We conduct multiple DA experiments (see Section 3.4), some of which utilize SDD as an input in the assimilation to test whether S1 provides new information. We utilize SDD rather than daily time series of snow-covered area for simplicity, and recognizing that SDD has been found to have high correlation with maximum SWE (Trujillo and Molotch, 2014). We also include SDD in a joint assimilation experiment to inform our model about end-of-season snowpack conditions (i.e., Gascoin et al., 2024), especially since S1 is only available through April 30th and has limitations during the melt season due to signal attenuation (Lievens et al., 2022). We derive SDD from different sources depending on the experiment. For the temporal evaluations, we derived SDD from SNOTEL by finding the first zero snow depth value after peak snow depth. For spatial evaluations, we derived SDD from MODIS snow cover data using annual summary metrics based on the daily Normalized Difference Snow Index (NDSI) from the MOD10A1 product with a threshold of 0.15, leveraging the SnowCloudMetrics algorithm implemented on Google Earth Engine (Crumley et al., 2020). The MODIS SDD is determined by identifying the last five snow-free days (backward approach) preceded by at least five consecutive snow-covered days. This

approach minimizes the impact of transient late-season snow events while ensuring a consistent detection of snow disappearance.

2.2.4. Evaluation Data:

We conduct temporal and spatial evaluations of snow depth, both prior to S1 assimilation and posterior to assimilation. Temporal evaluation utilizes daily snow depth from 11 NRCS SNOTEL stations within 40 km of the basin. Since the S1 change detection algorithm was optimized using SNOTEL data (Lievens et al., 2019), an independent research site at Snodgrass Mountain (Figure 1) was also utilized to assess the reliability of S1 snow depth time series away from SNOTEL sites. The Snodgrass Mountain site provides snow depth data, along with numerous monthly in situ snow pit measurements from February to May. These snow pits, located in both open and forested areas with a spatial extent of 1 km, are ideal for evaluating the 500 m S1 grid. S1 and posterior snow depths are evaluated against observations from these sites for WYs 2018–2021 for SNOTEL sites and 2019–2021 at the Snodgrass site.

The spatial evaluation uses four ASO LiDAR flights, with 50 m snow depth data collected near peak SWE (i.e., late March / early April) and in the mid-melt season (i.e., May or June) in 2018 and 2019 over the ERB. The data were resampled to match the 500 m S1 grid resolution using bilinear interpolation.

3. Methods:

3.1. DA Methodology

DA integrates model estimates with observations to estimate the most representative state of a system (e.g., SWE) with uncertainty. DA can account for observational uncertainty (e.g., S1 snow depth, MODIS SDD), and some DA approaches can provide physically consistent estimates of multiple snow states when implemented with a physically-based snow model. In this study, we develop and deploy a Python-based DA system in the Google Cloud Platform. Within the cloud, the system accesses ERA-5 Land meteorological data and uses it as input forcing into the MuSA toolbox, which communicates with the physically-based Flexible Snow Model version 2 (FSM2) model to generate ensembles of model simulations at a 9-km spatial resolution. The snow model ensembles from MuSA are combined with snow observations (e.g., S1 snow depth and/or SDD)

via the particle batch smoother (PBS). Through assimilation of snow data with a wide ensemble (section 3.2), PBS also serves as a downscaling technique(Bachand et al., 2025) from the 9-km ERA5-Land forcings to the 500 m grid. The flow chart in Figure 2 and the sections below detail the key components.




Figure 2. Flowchart illustrating the input, model, and DA framework, and outputs (top to bottom).

3.2. Snow Model and Ensemble Generation

FSM2 (Essery, 2015; Essery et al., 2024) is a multi-physics model that simulates the mass and energy balances of snow on the ground (including under forest canopies), employing three snow layers by default. The model incorporates conservation equations for liquid water, ice, and internal energy, offering a detailed representation of snowpack processes. FSM2's flexibility permits independent activation or deactivation of parameterizations, facilitating a range of model configurations (Essery, 2015; Essery et al., 2024). We utilize the most complex configuration of FSM2 to simulate internal snowpack processes, which we now briefly summarize. Albedo is calculated based on snow age, decreasing its value with time, and increasing it with fresh snowfalls. Snowpack thermal conductivity is determined by snow density, which is computed based on overburden and thermal metamorphism. Turbulent energy fluxes are computed based on bulk aerodynamic theory, and Monin-Obukhov adjustment for atmospheric stability is activated as

part of the calculation process. Meltwater percolation in the snowpack is computed using gravitational drainage. The selected configuration has demonstrated success in related DA studies (Smyth et al., 2022).

MuSA is an open-source, ensemble-based snow DA tool designed to assimilate multiple observations with FSM2 while considering various sources of forcing and measurement uncertainty (Alonso-González et al., 2022). We use MuSA to generate ensembles of FSM2 at a point scale. We perturbed the 9-km ERA5-Land hourly meteorological forcing data by drawing spatially independent, random perturbation parameters from a logitnormal distribution for precipitation (to avoid negative values) and from a normal distribution for temperature and LW radiation (Table 1). For precipitation, we applied bounds of $[0, 4]$ with $\mu = 0$ and $\sigma = 1$, capturing a wide range of basin-scale variability. For temperature and LW, we applied $\mu = 0$ and $\sigma = 2.5$ and $\mu = 0$ and $\sigma = 20.8$, respectively. Several combinations of bounds, standard deviation, and mean values were tested, and the final configuration was selected to be broader than typical ranges to account for the fact that ERA5-Land was not downscaled prior to assimilation and to better represent spatial variability in the East River Basin, Colorado. We tested and visualized the resulting spread to confirm realistic precipitation, temperature, and LW scenarios, and found that more than 98% of observed S1 snow depth values fell within the ensemble bounds during assimilation. Overall, this perturbation strategy was chosen to ensure the ensemble spans a wide range of meteorological conditions and captures a realistic range of possible snow depths.

Table 1. Generation of perturbed inputs for each particle member.

Variable	Unit	Adjustment	Distribution	Lower Bound	Upper Bound	Std. dev.	Mean
Precipitation	mm/h	Multiplicative	Logit-normal	0	4	1	0
Longwave radiation	W/m ²	Additive	Normal	-	-	20.8	0
Temperature	°C	Additive	Normal	-	-	2.5	0

Once the forcing data are perturbed, MuSA runs FSM2 to generate an ensemble of distinct snow simulations. Similar to Bachand et al. (2025), we used 100 ensemble members (particles) to ensure computational efficiency while adequately capturing the variability in the prior distribution. For

each particle, we record the SDD (i.e., the first snow-free date after peak snow depth), for use in multiple DA experiments (see Section 3.4).

3.3. Particle Batch Smoother – DA Algorithm

Once the ensembles are generated, we assimilate snow observations (e.g., snow depth and/or SDD) using the PBS algorithm. Separately, we tested other DA approaches, namely the Particle filter, both with constant and dynamic errors, the latter of which increases errors in time as revealed in our temporal analysis as SNOTEL sites (Figure 4). Despite having constant observational error, PBS had lower RMSE than both Particle filter implementations (Supplement Figure S1, Table S1), particularly during the ablation seasons, likely due to its smoothing properties. We therefore selected the PBS with temporally constant snow depth error for subsequent analysis. PBS is well-suited as a downscaling tool as well (Bachand et al., 2025), further motivating its selection in this study.

PBS employs a Bayesian approach, representing state variables like snow depth and SDD through a collection of particles, where each particle represents a possible system state. For observations within the assimilation window (e.g., snow season), PBS updates particle weights based on their likelihood of representing the true state, using the likelihood function from Margulis et al. (2015). This process combines the prior probability density function (PDF) with the likelihood to estimate the posterior PDF of snowpack variables, such as snow depth and SWE:

$$\mathbf{p}_{\mathbf{Z}|\mathbf{Y}}(\mathbf{Z}|\mathbf{Y}) = \mathbf{p}_{\mathbf{V}}(\mathbf{Z} - \mathbf{M}_j^-) = \frac{1}{\sqrt{(2\pi)^{N_{\text{obs}}|\mathbf{C}_{\mathbf{V}}|}}} \times \exp\left[-0.5(\mathbf{Z} - \mathbf{M}_j^-)^{\top} \mathbf{C}_{\mathbf{V}}^{-1}(\mathbf{Z} - \mathbf{M}_j^-)\right] \quad \text{Equation (1)}$$

Where $\mathbf{p}_{\mathbf{Z}|\mathbf{Y}}(\mathbf{Z}|\mathbf{Y})$ is the likelihood function, which represents the probability of observing the measurement $\mathbf{p}_{\mathbf{V}}(\mathbf{V})$ is the probability density function (PDF) of the measurement error vector \mathbf{V} , \mathbf{Z} is the observed measurement (e.g., S1 snow depth, MODIS SDD), \mathbf{M}_j is the modeled snow variable (e.g., snow depth or SDD), and $\mathbf{C}_{\mathbf{V}}$ is the error covariance matrix of the measurement error vector \mathbf{V} , which represents the uncertainty in the measurement.

3.4. DA Experiments

To address Research Questions 2 and 3, we conducted four experiments, building on the findings of Research Question 1 (S1 error analysis). The DA experiments are as follows (Figure 3):

1. **Snow Depth (Hs) Full Window (Hs-F):** Assimilation of all Hs, including dry and wet snow pixels, from November 15th to April 30th. S1 has data only until April 30th, and 83% of S1 data points before November 15th have zero or less than 10cm snow depth values. Thus, we call Hs-F a full-window experiment.
2. **Snow Depth (Hs) Early Window (Hs-E):** Assimilation of early-season (November 15th to January 15th) Hs, where the early window has a lower observational error than the full window (based on error analysis, section 4.1).
3. **Joint Assimilation of SDD and Snow Depth in the Full Window (SDD+Hs-F):** Combined assimilation of SDD and Hs over the same window as Hs-F. SDD data were derived from SNOTEL for temporal evaluation and MODIS for spatial evaluation.
4. **Snow Disappearance Date Only (SDD):** Assimilation of SDD data alone, derived from SNOTEL (temporal evaluations) or MODIS (spatial evaluations). This serves as a baseline or control experiment to understand whether S1 snow depth adds new information.

SDD is the baseline experiment, and each of the experiments are evaluated against 12 ground-based stations temporally and 4 ASO LiDAR flights spatially (Section 2.2.2). In experiments Hs-F, Hs-E, and SDD+Hs-F, the observational uncertainty value (Equation 1) for S1 snow depth is the RMSE when evaluated against SNOTEL for all four years. The SDD observational uncertainty is set to five days, consistent with the retrieval algorithm in SnowCloudMetrics (Crumley et al., 2020; Slater et al., 2013). In the joint assimilation (SDD+Hs-F), both datasets are weighted equally to balance their respective influence in DA. The joint likelihood is computed in log space under the assumption of independence between the S1 snow depth and SDD observations. Specifically, we define:

$$\mathbf{p}(\mathbf{Z} | \mathbf{Y}) = \mathbf{p}(\mathbf{Z}_{\text{Hs}} | \mathbf{Y}) + \mathbf{p}(\mathbf{Z}_{\text{SDD}} | \mathbf{Y}) \quad \text{Equation (2)}$$

where $\mathbf{p}(\mathbf{Z}_{\text{Hs}}|\mathbf{Y})$ is the snow depth and $\mathbf{p}(\mathbf{Z}_{\text{SDD}}|\mathbf{Y})$ is the SDD likelihood function. This equal-weight approach for joint assimilation is necessary for two reasons: (1) there is an

imbalance in the number of observations for Hs (dozens of observations) versus SDD (1 value) for each location and year, and (2) the Hs time series exhibits temporal autocorrelation.




Figure 3. Conceptual illustration of four experiments. Black dots indicate S1 observation data, black lines represent posterior estimates, and red stars (with blue dashed line) denote the SDD.

3.5. Evaluation

We evaluated snow depth (i.e., S1 observations and PBS experiments) with standard valuation metrics, including coefficient of determination (R^2), root mean squared error (RMSE), mean bias, mean absolute error (MAE), and relative mean absolute difference (RMAD). These were selected to enable comparisons to other studies (Lievens et al., 2019; Broxton et al., 2024; Hoppinen et al., 2024). To avoid seasonal inflation of R^2 in time series, where snowpack seasonality alone can yield artificially high correlations, we restrict temporal validation to dates where both modeled and observed snow depth are present (i.e., excluding summer/no-snow periods). These dates span the snow season (October 1 to April 30 or until disappearance in the SDD case), ensuring metrics evaluates on aligned timesteps. For spatial validation (e.g., S1 vs. LiDAR), metrics are computed for individual dates to avoid temporal autocorrelation effects. While R^2 is sensitive to underlying

model forcings, we use it alongside RMSE, MAE, and mean bias to provide a comprehensive performance evaluation. We also evaluated MAE and RMAD, which are defined as:

$$MAE = \frac{1}{n} \sum |y_i - \hat{y}_i| \quad \text{Equation (3)}$$

$$RMAD = \frac{\frac{1}{n} \sum |\hat{y}_i - y_i|}{\bar{y}} \quad \text{Equation (4)}$$

where y_i is the observed value (e.g., LiDAR, SNOTEL, field data), \hat{y}_i is the predicted value (e.g., PBS posterior mean), \bar{y} is the mean of all observed/validation data values, and n represents the total number of data points.

We also evaluated the probabilistic performance of the posterior ensemble simulations using the Continuous Ranked Probability Score (CRPS) at 12 point-scale observation sites. CRPS is a proper scoring rule that accounts for both the reliability and sharpness of probabilistic estimates from ensemble simulations, with lower values indicating better performance. We use the discrete form of CRPS as described by (Hersbach, 2000), which is equivalent to the continuous CDF-based definition (Matheson and Winkler, 1976) when applied to finite ensemble samples.:

$$CRPS(y_t, \{\hat{y}_{t,i}\}) = (1/N) \sum_{i=1}^N |\hat{y}_{t,i} - y_t| - (1/(2N^2)) \sum_{i=1}^N \sum_{j=1}^N |\hat{y}_{t,i} - \hat{y}_{t,j}| \quad \text{Equation (5)}$$

Where y_t is the observed snow depth (e.g., from SNOTEL) at time t , $\hat{y}_{t,i}$ is the snow depth predicted by the i -th ensemble member at time t , and N is the total number of ensemble members.

To quantify the added value of assimilation relative to the reference (open-loop) simulation, we computed the CRPS Skill Score (CRPSS) following the symmetric formulation of (Cluzet et al., 2022)

$$CRPSS(E, R) = \begin{cases} 1 - CRPS(E)/CRPS(R), & \text{if } CRPS(E) < CRPS(R); \\ CRPS(R)/CRPS(E) - 1, & \text{otherwise} \end{cases} \quad \text{Equation (6)}$$

This formulation bounds CRPSS between $[-1, 1]$, making it possible to directly compare and average positive (improvement) and negative (degradation) values.

4. Results

4.1 Error Analysis

Errors in the S1 500m snow depth data were evaluated temporally at 11 SNOTEL stations and an independent site (Snodgrass) from 2018–2021 and spatially relative to ASO LiDAR flights in 2018 and 2019. We included wet/flagged pixels in all our analyses/experiments because the removal of flagged pixels resulted in performance degradation (results not shown). The results are shown in Table 2.

Table 2. Average R^2 , RMSE, Mean Bias, MAE, and RMAD values of (a) S1 against 11 SNOTEL sites and the Snodgrass site. (b) S1 against ASO 50m LiDAR snow depth aggregated to 500m.

(a) Temporal evaluation – average across all stations (m)						(b) Spatial evaluation – S1 against ASO LiDAR (m)		
Year	2018	2019	2020	2021	Average	2018	2019	Average
R²	0.74	0.80	0.72	0.72	<u>0.74</u>	0.19	0.27	<u>0.23</u>
RMSE	0.32	0.39	0.39	0.53	<u>0.40</u>	0.74	0.91	<u>0.82</u>
Mean Bias	0.08	-0.07	0.09	0.26	<u>0.09</u>	-0.47	-0.26	<u>-0.36</u>
MAE	0.21	0.26	0.27	0.38	<u>0.28</u>	0.60	0.72	<u>0.66</u>
RMAD	0.61	0.39	0.47	0.68	<u>0.53</u>	0.77	0.43	<u>0.60</u>




Figure 4: (a) Time-varying errors at the 11 SNOTEL sites with the timing of LiDAR flights marked with thick grey lines. The solid lines represent the monthly mean bias, while the shaded region represents the \pm one standard deviation from the from daily bias values, indicating intra-month variability. Each station is represented by a different color. (b) Comparison of ASO LiDAR and Sentinel-1–derived snow depth estimates on temporally proximate dates.

When evaluated temporally against stations, S1 had an average bias of +0.09 m, RMSE of 0.40 m and R^2 of 0.74. In contrast, when evaluated spatially against ASO, S1 had higher bias magnitude (-0.36 m), a higher RMSE (0.82 m), and a lower R^2 (0.23) (Table 2). These results indicated that S1 performed better in capturing temporal variations compared to spatial patterns. S1 errors tended to

increase over time across all stations (Figure 4a), with the lowest errors in the early season and the highest errors ($\sim 0.4\text{m}$) near peak snow depth. The timing of the two LiDAR flights (Fig. 4b) coincided with the time of year when errors tend to be high (thick grey vertical lines, Fig. 4a).

4.2. Temporal Experiments:

Before analyzing the temporal performance of our assimilation windows, we first evaluated the overall skill of the posterior ensembles. We computed the CRPS (Eq. 4) across all sites and water years, which directly measures the distance between ensemble predictions and observations. The mean CRPS was 0.21 m, showing that posterior ensembles were, on average, within ~ 21 cm of observed snow depth and maintained skill across a wide range of snowpack conditions (Figure S2). CRPSS was then used to compare each experiment against the open-loop reference. As an uncertainty-aware metric, positive CRPSS values indicate improvement relative to the reference. Systematics CRPSS averages ranged from 0.08 to 0.22 (Table S2), reflecting modest but consistent gains from assimilation. Higher values for SDD and joint assimilation highlight the added value of snow disappearance information, while lower values for Hs-only experiments underscore the noisier character of Sentinel-1 snow depth.

With this confidence in ensemble performance, we assessed the performance of S1 assimilation with respect to window size, i.e., full window data assimilation (Hs-F) compared to the early window (Hs-E) when errors were lower (Fig. 4a). The results are shown in Figure 5 for two sites: (1) an independent site (Snodgrass), which was not used in the optimization of the S1 dataset, and (2) a representative SNOTEL site. The time series of the two sites shows that the performance of Hs-F and Hs-E is not consistent across the years. Hs-F (purple line) performed marginally better than Hs-E (green line) in WY 2019 and similarly in 2020 and 2021 for the Snodgrass site. Likewise, at Chapman Tunnel, Hs-F performed better than Hs-E in WY 2018, poorly in 2021, and similarly to HS-E in 2019 and 2020.




Figure 5: Temporal analysis results at (top) Snodgrass and (bottom) Chapman Tunnel. Hs-F (purple line) and Hs-E (green line) represent the posterior mean of full and early window experiments, respectively. SNOTEL and Snodgrass station data (blue dashed line) provide snow depth evaluation data. Orange diamonds represent S1 snow depth observations. Shading of respective color indicates one standard deviation of posterior particle spread, and the solid black lines represent the early assimilation window. See Figure S3 for the comparison of these experiments against the model-only run.

The statistical analysis across temporal evaluation sites give insights into the average performance of each DA window experiment. The results show RMSE of 0.29 m for Hs-F and 0.35 m for Hs-E, while R^2 is 0.88 for Hs-F and 0.86 for Hs-E (Table 3). Overall, the differences between Hs-F and Hs-E across all sites and evaluation metrics are minimal, suggesting that the accuracy of S1 snow depth retrievals remains consistent regardless of the temporal window. This implies that the retrieval method performs similarly for both the full and early windows.

Table 3: Error metrics of posterior mean of Hs-F (full window assimilation) and Hs-E (early window assimilation). Metrics are averaged across all temporal evaluation sites and across four years. See Table S2 for all experiments comparison against the model-only run.

Metrics	Experiments	
	Hs-F	Hs-E
R²	0.88	0.86
RMSE (m)	0.29	0.35
Mean Bias (m)	0.07	0.09
MAE	0.20	0.23
RMAD	0.46	0.63

While the temporal analysis shows a marginal difference in Hs-F and Hs-E, spatial analysis reveals a significant disparity in performance, with Hs-F outperforming Hs-E significantly across all LiDAR flights in most metrics (Table 4). For example, Hs-F demonstrates a lower average RMSE (0.66 m vs. 0.91 m), lower mean bias (0.05m vs. -0.61m) along with lower MAE (0.42m vs. 0.75m) and RMAD (0.61 vs. 1.2m) compared to Hs-E. However, Hs-F performs slightly worse in specific survey dates, such as R² on 2019-06-10, where Hs-E performs better than Hs-F (0.08 vs. 0.57m). The temporal and spatial RMSE across all stations and years and LiDAR flights are shown in Figure 6, with Hs-F showing lower RMSE in all cases except in the 2019-06-10 LiDAR flight.

Across all four LiDAR surveys, the spatial patterns in snow depth are not strongly related (i.e., R²<0.60) for either Hs-E or Hs-F (Table 4, Figure 7). The density plots (Figure 7c) compare the posterior mean snow depths from the full-window (Hs-F) and early-window (Hs-E) assimilation experiments with LiDAR measurements. The density distribution aligns with LiDAR only in certain cases (Fig. 7c), reflecting both experiments perform consistently poor relative to the LiDAR snow depths. The overall analysis indicates that S1 has high errors spatially regardless of window, but the full-window assimilation approach (Hs-F) provides relatively lower errors across spatial locations and LiDAR dates (Figure 7). Therefore, Hs-F is utilized as the optimal window for the rest of the data assimilation experiments (see Section 4.3).




Figure 6. Bar chart comparing average RMSE from the Hs-F and Hs-E experiments when evaluated at the 12 snow stations. The average errors are shown for all dates and on each of the four LiDAR flight dates.

Table 4. Error metrics for the posterior mean snow depth from the Hs-F and Hs-E experiments when evaluated against snow depth from four LiDAR surveys. See Table S3 for all experiments comparison against the model-only run.

LiDAR Survey	Exp	R ²	RMSE (m)	Mean Bias (m)	MAE (m)	RMAD (m)
2018-03-31	Hs-F	0.51	0.48	-0.21	0.35	0.45
	Hs-E	0.29	1.24	-1.15	1.15	1.48
2018-05-24	Hs-F	0.33	0.39	0.08	0.21	0.88
	Hs-E	0.37	0.78	-0.59	0.69	2.55
2019-04-07	Hs-F	0.33	0.73	-0.08	0.52	0.31
	Hs-E	0.47	0.85	-0.52	0.66	0.40
2019-06-10	Hs-F	0.08	1.06	0.44	0.60	0.83
	Hs-E	0.57	0.77	-0.21	0.52	0.63
Average	<u>Hs-F</u>	<u>0.31</u>	<u>0.66</u>	<u>0.05</u>	<u>0.42</u>	<u>0.61</u>
	<u>Hs-E</u>	<u>0.42</u>	<u>0.91</u>	<u>-0.61</u>	<u>0.75</u>	<u>1.2</u>




Figure 7: The first column in panels a and b shows snow depth from airborne LiDAR near peak snow accumulation. The second and third columns in panels a and b show the posterior mean snow depth from the Hs-F and Hs-E experiments, respectively. Panel C shows the density plot comparing two experiments (Hs-F and Hs-E) against LiDAR measurements. Since both experiments showed similar correlations with LiDAR, they are combined into a single plot. See Figure S5 for the experiments comparison against the model-only run

4.3. Joint Assimilation Experiments:

Finally, we evaluate the joint assimilation of SDD+Hs-F against SDD alone (baseline) to test the utility of S1 snow depth for adding new information to a DA system. We utilized Hs-F instead of Hs-E because errors tended to be lower with Hs-F (see section 4.2). The temporal results of

experiments 3 and 4 of the most representative and independent site are shown in Figure 8. The results show that the performance of Hs-F (experiment 1), SDD+Hs-F (experiment 3), and SDD alone (experiment 4) varies across the years. Hs-F performed better than SDD-Hs-F and SDD in WY 2019 and 2020 and similarly in 2021 for the Snodgrass site. In the SNOTEL representative site, SDD performed better or similar as Hs-F and SDD+Hs-F in all WYs. The temporal analysis indicates that joint assimilation and SDD alone experiments perform better than Hs-F minimally across all metrics ($\sim 0.01\text{m}$). However, the performance of SDD+Hs-F is approximately the same as the baseline SDD experiment (Table 5), suggesting that joint assimilation of SDD with S1 snow depth does not significantly enhance performance relative to just assimilating SDD alone.




Figure 8: Example results of the joint assimilation experiments at (a) Snodgrass and (b) Nast Lake. Hs-F (purple line) and SDD+Hs-F (green line) represent the posterior mean for each experiment. SDD (black line) is the posterior snow depth based on assimilating only SDD. SNOTEL and Snodgrass in situ (blue dashed line) provide snow depth evaluation data. Orange diamonds represent S1 snow depth observations.

Shading of respective color indicates one standard deviation of posterior particle spread. See Figure S4 for the comparison against the model-only run.

Table 5. Error metrics of the posterior mean of Hs-F (full window assimilation), SDD+HS-F (joint assimilation), and SDD. Metrics are averaged across all temporal evaluation sites and across four years. See Table S2 for all experiments comparison against the model-only run.

Metrics	Experiments	
	SDD+Hs-F	SDD
R²	0.92	0.72
RMSE	0.28	0.28
Mean Bias	0.15	0.14
MAE	0.19	0.19
RMAD	0.42	0.42

Spatial analysis provides more insights on univariate (Hs-F or SDD) and joint assimilation experiments (Hs-F+SDD). Across all metrics, assimilating SDD alone results in higher spatial performance compared to both Hs-F and SDD+Hs-F. The temporal and spatial RMSE across all stations, years, and LiDAR flights are shown in Figure 9. The RMSE is 0.41 m for SDD assimilation averaged across all LiDAR flights and years, which is lower than the RMSE of Hs-F (0.66 m, Table 4) and SDD+Hs-F (0.70 m, Table 6). Additionally, assimilating SDD alone demonstrates a stronger relationship with LiDAR observations ($R^2 = 0.72$) compared to Hs-F ($R^2 = 0.31$) and SDD+Hs-F ($R^2 = 0.46$), with better spatial pattern alignment across both LiDAR dates near peak accumulation (Table 6, Fig. 10). Joint assimilation (SDD+Hs-F) shows lower performance in comparison to Hs-F alone or SDD alone, suggesting that the combination of S1 with SDD does not enhance performance and may potentially degrade it.




Figure 9: Bar chart comparing RMSE for Hs-F, SDD+Hs-F, and SDD experiments of stations (SNOTEL+Snodgrass) averaged across all stations and all four years, and each LiDAR flight.




Figure 10: The first column in panels a and b shows snow depth from airborne LiDAR near peak snow accumulation. The second, third, and fourth columns in panels a and b show the posterior mean snow depth from the Hs-F, SDD+Hs-F, and SDD experiments, respectively. Panel c shows the density plot comparing three experiments (Hs-F, SDD-Hs-F, and SDD) against LiDAR measurements. Since both experiments showed similar correlations (or better in the case of SDD) with LiDAR, they are combined into a single plot. See Figure S5 for the comparison against the model-only run

Table 6. Error metrics of posterior mean snow depth from the Hs-F, SDD+Hs-F, and SDD experiments when evaluated against four LiDAR surveys. See Table S3 for the comparison against the model-only run.

LiDAR Survey	Exp	R ²	RMSE (m)	Mean Bias (m)	MAE (m)	RMAD (m)
2018-03-31	Hs-F	0.51	0.48	-0.21	0.35	0.45
	SDD+Hs-F	0.58	0.65	0.55	0.55	0.71

	SDD	0.74	0.32	-0.25	0.33	0.42
2018-04-24	Hs-F	0.33	0.39	0.08	0.21	0.88
	SDD+Hs-F	0.10	0.44	0.23	0.23	0.99
	SDD	0.70	0.12	-0.01	0.13	0.53
2019-04-07	Hs-F	0.33	0.73	-0.08	0.52	0.31
	SDD+Hs-F	0.69	0.71	0.55	0.62	0.35
	SDD	0.70	0.68	-0.48	0.54	0.32
2019-06-10	Hs-F	0.08	1.06	0.44	0.60	0.83
	SDD+Hs-F	0.49	1.0	0.71	0.71	0.85
	SDD	0.76	0.52	-0.09	0.34	0.40
Average	Hs-F	<u>0.31</u>	<u>0.66</u>	<u>0.05</u>	<u>0.42</u>	<u>0.61</u>
	SDD+Hs-F	<u>0.46</u>	<u>0.70</u>	<u>0.51</u>	<u>0.52</u>	<u>0.72</u>
	SDD	<u>0.72</u>	<u>0.41</u>	<u>0.20</u>	<u>0.33</u>	<u>0.42</u>

5. Discussion

This study evaluates the utility of S1 SAR-derived snow depth data in the mountainous ERB (Colorado) to support spatiotemporal snow depth mapping within a DA framework. Our analysis highlights inconsistencies in temporal versus spatial errors in S1 snow depth data. While S1 demonstrates lower temporal errors (RMSE = 0.40 m, $R^2 = 0.74$), spatial errors were higher (RMSE = 0.82m, $R^2 = 0.23$) relative to LiDAR flights near or after peak accumulation (Table 2). Temporal errors were lower during the early season but increased over time, particularly during the ablation phase when wet snow conditions become more likely (Figure 4). These discrepancies align with prior studies (Broxton et al., 2024; Hoppinen et al., 2024), which report significant errors in S1 snow depth data under wet snow conditions and poor spatial correlations with airborne LiDAR data (RMSE > 0.7 m, $R^2 < 0.3$). The limitation may also stem from the timing of spatial validation data, as most LiDAR flights are conducted near or after peak snow depth when S1 performance is most likely compromised due to wet snow. However, early-season (Hs-E) experiments show high uncertainty, suggesting uncertainty is likely due to noisy observations without a clear error pattern. Unlike prior studies that proposed excluding flagged wet pixels as a mitigation strategy, we found that removing flagged pixels increased errors, potentially due to the omission of shallow dry snow observations (Hoppinen et al., 2024). Lievens et al. (2022) reported significantly lower errors (~0.25 m) in the European Alps, which is contrary to studies (Broxton

et al., 2024; Hoppinen et al., 2024) focused on the Western US, including our study. The lower errors reported by Lievens et al. (2022) could be due to a higher overpass frequency and denser validation datasets and needs further investigation. Therefore, while S1 may be useful in other regions, our findings emphasize its limited reliability in much of the Western US., including ERB, and caution against broad generalizations.

Noisy observations are not inherently problematic for data assimilation, provided their uncertainty is well characterized, our current analysis did not reveal consistent year-to-year error patterns, limiting the development of sophisticated error models (Figure. 4). Thus we informed our experiments based on recommendations by Gascoin et al. (2024) and aimed to determine the optimal assimilation window (full season vs. early season) with lower uncertainty for generating spatiotemporal snow depth maps. . The comparison between the full-window (Hs-F) and early-window (Hs-E) approaches showed minimal improvement in error metrics in the temporal analysis at ground-based stations. Spatial evaluations against LiDAR data resulted in Hs-F performing better than Hs-E. However, Hs-F increases the likelihood of including wet snow conditions and higher errors later in the snow season. The early-window approach benefits from reduced retrieval errors due to its focus on lower snow depth values and assumed higher likelihood of dry snow in December and January. However, this approach is more effective in regions where a significant fraction of the total SWE accumulates early in the season (Lundquist et al., 2023). In contrast, basins like the ERB receive significant snowfall after January, which reduces the early-season window's ability to predict SWE reliably later in the year (e.g., April-onward). This seasonal snow accumulation pattern likely explains why the early-window approach is less effective in our study. This variability underscores the inherent challenges in using early-season conditions to make broader inferences about snowpack dynamics later in the year. Despite these limitations, the results show that errors in S1 snow depth retrievals are relatively independent of the assimilation window in the ERB.

The joint assimilation of S1 snow depth with SDD was conducted to understand what information is added by S1 and to test whether SDD can provide additional constraints on the DA outcome. Previous studies successfully used joint assimilation in a DA system with satellite data such as ICESat-2 (Mazzolini et al., 2024). However, joint assimilation showed limited value with S1 snow

depth data: The errors did not decrease when assimilating S1 depth with SDD. Assimilating SDD alone showed a higher R^2 value and lower errors on average compared to Hs-F and SDD+Hs-F.

There are some caveats and potential limitations in the study. The study did not utilize downscaling before assimilation; however, we did not see extreme bias in reference runs (model only, Figure S3-5), which suggests that limited gains from DA are primarily due to S1 observation uncertainty, rather than a lack of downscaling. Additionally, recent research has shown that assimilation of S1 snow depth can implicitly be used as downscaling (Bachand et al., 2025). The current study focused on a single basin and a limited number of nearby snow pillow stations over a four-year period. A more comprehensive evaluation against LiDAR data has already been conducted with the available LiDAR data in the western U.S. (Broxton et al., 2024; Hoppinen et al., 2024). Similar to previous studies, we were also limited by the evaluation of LiDAR data since it was only available at peak snow depth or in the melt season, when S1 has higher errors. This highlights the importance of expanding the spatial and temporal coverage of evaluation data.

Future research should continue to develop and explore approaches for using Sentinel-1 and other spaceborne remote sensing platforms for mapping SWE and other related snowpack variables. One established capability of S1 backscatter data is for wet snow mapping (Cluzet et al., 2024; Gagliano et al., 2023; Nagler et al., 2016). For snow depth and SWE mapping, the existing Lievens et al. (2019) algorithm may be improved through corrections with machine learning as proposed by Broxton et al. (2024). Alternatively, interferometric SAR (InSAR) techniques (Oveisgharan et al., 2024) have recently been shown to have some potential for mapping SWE with S1 C-band data, although challenges related to the frequency of satellite passes remain a challenge (Deeb et al., 2011) and these techniques are likely more reliable with L-band SAR data. The upcoming NISAR mission will provide L-band SAR data, which may provide new opportunities for accurate snow depth mapping in mountainous regions that may overcome some of the current limitations of C-band SAR. However, other limitations may persist for NISAR (e.g., wet snow).

6. Conclusion

This study underscores the challenges of using C-Band S1 SAR-derived snow depth data within a DA framework. While S1 is currently the only high-resolution (<1 km) remotely sensed snow depth dataset available across the Northern Hemisphere, it demonstrated notable biases and

limitations in snow depth mapping in the East River Basin in the western U.S. These errors were similar across both full-window and early-window assimilation experiments and align with prior studies that reported significant spatial biases and retrieval errors. Importantly, we did not observe a consistent seasonal or interannual pattern in the errors, which limits the development of robust correction or error models. The joint assimilation of S1 snow depth with SDD data showed limited or no improvement, suggesting that assimilating SDD alone yields greater accuracy.

While recognizing that our study focused on a single mountain basin, we conclude that the reliability of the current S1 snow depth retrieval algorithm presents major challenges for a snow DA system. Future research should prioritize algorithm improvements, explore machine learning techniques, and conduct additional testing across a wider range of basins and with spatial data. Enhanced methods for snow depth monitoring would improve understanding of snowpack dynamics in regions where ground-based observations are sparse or unavailable, supporting better water resource management and climate impact assessments.

Author Contributions: Data curation and Analysis: BM. Experiment Design: BM, MR, EE. Writing: The draft was led by BM with key contribution and editing by all co-authors. Funding Acquisition: MR and ES.

Code and data availability: This study used airborne LiDAR snow depth data (ASO, Inc.), MODIS and ERA5 data (GEE), NRCS SNOTEL, and independent validation data from teams at the University of Colorado and Oregon State University. The MuSA code, which integrates FSM, is available at <https://github.com/ealonsogzl/MuSA>, with FSM originally developed by Richard Essery. Computational resources were provided by Oregon State University's College of Engineering.

Competing interests: The authors declare that they have no conflict of interest.

Acknowledgments

This work was supported by the National Aeronautics and Space Administration (NASA) Terrestrial Hydrology Program under Award No. 80NSSC22K0685. The authors thank Esteban Alonso González for making the MuSA code available (<https://github.com/ealonsogzl/MuSA>) and Richard Essery for development of FSM. This work utilized resources from Oregon State University College of Engineering high-performance computing network. The independent temporal validation data was collected by student teams from the University of Colorado (Eric Small Hydrology Group) and Oregon State University (CryoSphere Interactions and Geospatial Hydrology Team). The authors also thank the ASO, Inc. team for providing the airborne LiDAR snow depth and GEE for facilitating the access of MODIS and meteorological forcing data used in the study.

610 **References:**

- 611 Aalstad, K., Westermann, S., and Bertino, L.: Evaluating satellite retrieved fractional snow-
612 covered area at a high-Arctic site using terrestrial photography, *Remote Sensing of*
613 *Environment*, 239, 111618, <https://doi.org/10.1016/j.rse.2019.111618>, 2020.
- 614 Alonso-González, E., Aalstad, K., Baba, M. W., Revuelto, J., López-Moreno, J. I., Fiddes, J.,
615 Essery, R., and Gascoin, S.: The Multiple Snow Data Assimilation System (MuSA v1.0),
616 *Geosci. Model Dev.*, 15, 9127–9155, <https://doi.org/10.5194/gmd-15-9127-2022>, 2022.
- 617 Bachand, C. L., Andrews, L. C., Rouf, T., and Giroto, M.: The Utility of Satellite Snow Depth
618 Observations for Downscaling Hydrologic Variables over the Indus Basin Mountain Ranges,
619 *Journal of Hydrometeorology*, 26, 555–575, <https://doi.org/10.1175/JHM-D-24-0131.1>,
620 2025.
- 621 Barnett, T. P., Adam, J. C., and Lettenmaier, D. P.: Potential impacts of a warming climate on
622 water availability in snow-dominated regions, *Nature*, 438, 303–309,
623 <https://doi.org/10.1038/nature04141>, 2005.
- 624 Besso, H., Shean, D., and Lundquist, J. D.: Mountain snow depth retrievals from
625 customized processing of ICESat-2 satellite laser altimetry, *Remote Sensing of*
626 *Environment*, 300, 113843, <https://doi.org/10.1016/j.rse.2023.113843>, 2024.
- 627 Bonner, H. M., Smyth, E., Raleigh, M. S., and Small, E. E.: A Meteorology and Snow Data Set
628 From Adjacent Forested and Meadow Sites at Crested Butte, CO, USA, *Water Resources*
629 *Research*, 58, <https://doi.org/10.1029/2022WR033006>, 2022.
- 630 Brangers, I., Lievens, H., Getirana, A., and De Lannoy, G. J. M.: Sentinel-1 Snow Depth
631 Assimilation to Improve River Discharge Estimates in the Western European Alps, *Water*
632 *Resources Research*, 60, e2023WR035019, <https://doi.org/10.1029/2023WR035019>, 2024.
- 633 Broxton, P., Ehsani, M. R., and Behrangi, A.: Improving Mountain Snowpack Estimation
634 Using Machine Learning With Sentinel-1, the Airborne Snow Observatory, and University of
635 Arizona Snowpack Data, *Earth and Space Science*, 11, e2023EA002964,
636 <https://doi.org/10.1029/2023EA002964>, 2024.
- 637 Broxton, P. D., Van Leeuwen, W. J. D., and Biederman, J. A.: Improving Snow Water
638 Equivalent Maps With Machine Learning of Snow Survey and Lidar Measurements, *Water*
639 *Resources Research*, 55, 3739–3757, <https://doi.org/10.1029/2018WR024146>, 2019.
- 640 Chang, A. T. C., Foster, J. L., and Hall, D. K.: Nimbus-7 SMMR Derived Global Snow Cover
641 Parameters, *A. Glaciology*, 9, 39–44, <https://doi.org/10.3189/S0260305500200736>, 1987.
- 642 Clark, M. P., Hendrikx, J., Slater, A. G., Kavetski, D., Anderson, B., Cullen, N. J., Kerr, T., Örn
643 Hreinsson, E., and Woods, R. A.: Representing spatial variability of snow water equivalent

644 in hydrologic and land-surface models: A review, *Water Resources Research*, 47,
645 2011WR010745, <https://doi.org/10.1029/2011WR010745>, 2011.

646 Cluzet, B., Lafaysse, M., Deschamps-Berger, C., Vernay, M., and Dumont, M.: Propagating
647 information from snow observations with CrocO ensemble data assimilation system: a 10-
648 years case study over a snow depth observation network, *The Cryosphere*, 16, 1281–1298,
649 <https://doi.org/10.5194/tc-16-1281-2022>, 2022.

650 Cluzet, B., Magnusson, J., Quéno, L., Mazzotti, G., Mott, R., and Jonas, T.: Exploring how
651 Sentinel-1 wet-snow maps can inform fully distributed physically based snowpack models,
652 *The Cryosphere*, 18, 5753–5767, <https://doi.org/10.5194/tc-18-5753-2024>, 2024.

653 Copernicus Climate Change Service: ERA5-Land hourly data from 1950 to present,
654 <https://doi.org/10.24381/CDS.E2161BAC>, 2019.

655 Crumley, R. L., Palomaki, R. T., Nolin, A. W., Sproles, E. A., and Mar, E. J.:
656 SnowCloudMetrics: Snow Information for Everyone, *Remote Sensing*, 12, 3341,
657 <https://doi.org/10.3390/rs12203341>, 2020.

658 Daly, C., Neilson, R. P., and Phillips, D. L.: A Statistical-Topographic Model for Mapping
659 Climatological Precipitation over Mountainous Terrain, *J. Appl. Meteor.*, 33, 140–158,
660 [https://doi.org/10.1175/1520-0450\(1994\)033<0140:ASTMFM>2.0.CO;2](https://doi.org/10.1175/1520-0450(1994)033<0140:ASTMFM>2.0.CO;2), 1994.

661 De Boer, G., White, A., Cifelli, R., Intrieri, J., Rose Abel, M., Mahoney, K., Meyers, T., Lantz,
662 K., Hamilton, J., Currier, W., Sedlar, J., Cox, C., Hulm, E., Riihimaki, L. D., Adler, B., Bianco,
663 L., Morales, A., Wilczak, J., Elston, J., Stachura, M., Jackson, D., Morris, S., Chandrasekar,
664 V., Biswas, S., Schmatz, B., Junyent, F., Reithel, J., Smith, E., Schloesser, K., Kochendorfer,
665 J., Meyers, M., Gallagher, M., Longenecker, J., Olheiser, C., Bytheway, J., Moore, B., Calmer,
666 R., Shupe, M. D., Butterworth, B., Heflin, S., Palladino, R., Feldman, D., Williams, K., Pinto,
667 J., Osborn, J., Costa, D., Hall, E., Herrera, C., Hodges, G., Soldo, L., Stierle, S., and Webb, R.
668 S.: Supporting Advancement in Weather and Water Prediction in the Upper Colorado River
669 Basin: The SPLASH Campaign, *Bulletin of the American Meteorological Society*, 104,
670 E1853–E1874, <https://doi.org/10.1175/BAMS-D-22-0147.1>, 2023.

671 De Lannoy, G. J. M., Bechtold, M., Busschaert, L., Heyvaert, Z., Modanesi, S., Dunmire, D.,
672 Lievens, H., Getirana, A., and Massari, C.: Contributions of Irrigation Modeling, Soil
673 Moisture and Snow Data Assimilation to High-Resolution Water Budget Estimates Over the
674 Po Basin: Progress Towards Digital Replicas, *J Adv Model Earth Syst*, 16,
675 <https://doi.org/10.1029/2024ms004433>, 2024.

676 Deeb, E. J., Forster, R. R., and Kane, D. L.: Monitoring snowpack evolution using
677 interferometric synthetic aperture radar on the North Slope of Alaska, USA, *International*
678 *Journal of Remote Sensing*, 32, 3985–4003, <https://doi.org/10.1080/01431161003801351>,
679 2011.

680 Derksen, C., Walker, A., and Goodison, B.: Evaluation of passive microwave snow water
681 equivalent retrievals across the boreal forest/tundra transition of western Canada, *Remote*
682 *Sensing of Environment*, 96, 315–327, <https://doi.org/10.1016/j.rse.2005.02.014>, 2005.

683 Deschamps-Berger, C., Cluzet, B., Dumont, M., Lafaysse, M., Berthier, E., Fanise, P., and
684 Gascoin, S.: Improving the Spatial Distribution of Snow Cover Simulations by Assimilation
685 of Satellite Stereoscopic Imagery, *Water Resources Research*, 58,
686 <https://doi.org/10.1029/2021WR030271>, 2022.

687 Deschamps-Berger, C., Gascoin, S., Shean, D., Besso, H., Guiot, A., and López-Moreno, J.
688 I.: Evaluation of snow depth retrievals from ICESat-2 using airborne laser-scanning data,
689 *The Cryosphere*, 17, 2779–2792, <https://doi.org/10.5194/tc-17-2779-2023>, 2023.

690 Dozier, J.: Mountain hydrology, snow color, and the fourth paradigm, *EoS Transactions*, 92,
691 373–374, <https://doi.org/10.1029/2011EO430001>, 2011.

692 Dozier, J., Bair, E. H., and Davis, R. E.: Estimating the spatial distribution of snow water
693 equivalent in the world’s mountains, *WIREs Water*, 3, 461–474,
694 <https://doi.org/10.1002/wat2.1140>, 2016.

695 Dunmire, D., Lievens, H., Boeykens, L., and De Lannoy, G. J. M.: A machine learning
696 approach for estimating snow depth across the European Alps from Sentinel-1 imagery,
697 *Remote Sensing of Environment*, 314, 114369, <https://doi.org/10.1016/j.rse.2024.114369>,
698 2024.

699 Elder, K., Rosenthal, W., and Davis, R. E.: Estimating the spatial distribution of snow water
700 equivalence in a montane watershed, *Hydrol. Process.*, 12, 1793–1808,
701 [https://doi.org/10.1002/\(SICI\)1099-1085\(199808/09\)12:10/11<1793::AID-](https://doi.org/10.1002/(SICI)1099-1085(199808/09)12:10/11<1793::AID-HYP695>3.0.CO;2-K)
702 [HYP695>3.0.CO;2-K](https://doi.org/10.1002/(SICI)1099-1085(199808/09)12:10/11<1793::AID-HYP695>3.0.CO;2-K), 1998.

703 Essery, R.: A factorial snowpack model (FSM 1.0), *Geosci. Model Dev.*, 8, 3867–3876,
704 <https://doi.org/10.5194/gmd-8-3867-2015>, 2015.

705 Essery, R., Mazzotti, G., Barr, S., Jonas, T., Quaife, T., and Rutter, N.: A Flexible Snow Model
706 (FSM 2.1.0) including a forest canopy, <https://doi.org/10.5194/egusphere-2024-2546>, 11
707 October 2024.

708 Feldman, D. R., Aiken, A. C., Boos, W. R., Carroll, R. W. H., Chandrasekar, V., Collis, S.,
709 Creamean, J. M., De Boer, G., Deems, J., DeMott, P. J., Fan, J., Flores, A. N., Gochis, D.,
710 Grover, M., Hill, T. C. J., Hodshire, A., Hulm, E., Hume, C. C., Jackson, R., Junyent, F.,
711 Kennedy, A., Kumjian, M., Levin, E. J. T., Lundquist, J. D., O’Brien, J., Raleigh, M. S., Reithel,
712 J., Rhoades, A., Rittger, K., Rudisill, W., Sherman, Z., Siirila-Woodburn, E., Skiles, S. M.,
713 Smith, J. N., Sullivan, R. C., Theisen, A., Tuftedal, M., Varble, A. C., Wiedlea, A., Wielandt,
714 S., Williams, K., and Xu, Z.: The Surface Atmosphere Integrated Field Laboratory (SAIL)

715 Campaign, *Bulletin of the American Meteorological Society*, 104, E2192–E2222,
716 <https://doi.org/10.1175/BAMS-D-22-0049.1>, 2023.

717 Foster, J., Liston, G., Koster, R., Essery, R., Behr, H., Dumenil, L., Verseghy, D., Thompson,
718 S., Pollard, D., and Cohen, J.: Snow Cover and Snow Mass Intercomparisons of General
719 Circulation Models and Remotely Sensed Datasets, *J. Climate*, 9, 409–426,
720 [https://doi.org/10.1175/1520-0442\(1996\)009<0409:SCASMI>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<0409:SCASMI>2.0.CO;2), 1996.

721 Gagliano, E., Shean, D., Henderson, S., and Vanderwilt, S.: Capturing the Onset of
722 Mountain Snowmelt Runoff Using Satellite Synthetic Aperture Radar, *Geophysical*
723 *Research Letters*, 50, e2023GL105303, <https://doi.org/10.1029/2023GL105303>, 2023.

724 Giroto, M., Musselman, K. N., and Essery, R. L. H.: Data Assimilation Improves Estimates
725 of Climate-Sensitive Seasonal Snow, *Curr Clim Change Rep*, 6, 81–94,
726 <https://doi.org/10.1007/s40641-020-00159-7>, 2020.

727 Giroto, M., Formetta, G., Azimi, S., Bachand, C., Cowherd, M., De Lannoy, G., Lievens, H.,
728 Modanesi, S., Raleigh, M. S., Rigon, R., and Massari, C.: Identifying snowfall elevation
729 patterns by assimilating satellite-based snow depth retrievals, *Science of The Total*
730 *Environment*, 906, 167312, <https://doi.org/10.1016/j.scitotenv.2023.167312>, 2024.

731 Grünewald, T., Schirmer, M., Mott, R., and Lehning, M.: Spatial and temporal variability of
732 snow depth and ablation rates in a small mountain catchment, *The Cryosphere*, 4, 215–
733 225, <https://doi.org/10.5194/tc-4-215-2010>, 2010.

734 Herbert, J. N., Raleigh, M. S., and Small, E. E.: Reanalyzing the spatial representativeness of
735 snow depth at automated monitoring stations using airborne lidar data, *The Cryosphere*,
736 18, 3495–3512, <https://doi.org/10.5194/tc-18-3495-2024>, 2024.

737 Hersbach, H.: Decomposition of the Continuous Ranked Probability Score for Ensemble
738 Prediction Systems, *Wea. Forecasting*, 15, 559–570, [https://doi.org/10.1175/1520-0434\(2000\)015<0559:dotcrp>2.0.co;2](https://doi.org/10.1175/1520-0434(2000)015<0559:dotcrp>2.0.co;2), 2000.

740 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas,
741 J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X.,
742 Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P.,
743 Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A.,
744 Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P.,
745 Lopez, P., Lupu, C., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and
746 Thépaut, J.: The ERA5 global reanalysis, *Quart J Royal Meteor Soc*, 146, 1999–2049,
747 <https://doi.org/10.1002/qj.3803>, 2020.

748 Hoppinen, Z., Palomaki, R. T., Brencher, G., Dunmire, D., Gagliano, E., Marziliano, A.,
749 Tarricone, J., and Marshall, H.-P.: Evaluating snow depth retrievals from Sentinel-1 volume

750 scattering over NASA SnowEx sites, *The Cryosphere*, 18, 5407–5430,
 751 <https://doi.org/10.5194/tc-18-5407-2024>, 2024.

752 Hu, Y., Lu, X., Zeng, X., Stamnes, S. A., Neuman, T. A., Kurtz, N. T., Zhai, P., Gao, M., Sun, W.,
 753 Xu, K., Liu, Z., Omar, A. H., Baize, R. R., Rogers, L. J., Mitchell, B. O., Stamnes, K., Huang, Y.,
 754 Chen, N., Weimer, C., Lee, J., and Fair, Z.: Deriving Snow Depth From ICESat-2 Lidar
 755 Multiple Scattering Measurements, *Front. Remote Sens.*, 3, 855159,
 756 <https://doi.org/10.3389/frsen.2022.855159>, 2022.

757 Hubbard, S. S., Williams, K. H., Agarwal, D., Banfield, J., Beller, H., Bouskill, N., Brodie, E.,
 758 Carroll, R., Dafflon, B., Dwivedi, D., Falco, N., Faybishenko, B., Maxwell, R., Nico, P.,
 759 Steefel, C., Steltzer, H., Tokunaga, T., Tran, P. A., Wainwright, H., and Varadharajan, C.: The
 760 East River, Colorado, Watershed: A Mountainous Community Testbed for Improving
 761 Predictive Understanding of Multiscale Hydrological–Biogeochemical Dynamics, *Vadose*
 762 *Zone Journal*, 17, 1–25, <https://doi.org/10.2136/vzj2018.03.0061>, 2018.

763 Jarvis, A., Guevara, E., Reuter, H. I., and Nelson, A. D.: Hole-filled SRTM for the globe :
 764 version 4 : data grid, 2008.

765 Kelly, R. E., Chang, A. T., Tsang, L., and Foster, J. L.: A prototype AMSR-E global snow area
 766 and snow depth algorithm, *IEEE Trans. Geosci. Remote Sensing*, 41, 230–242,
 767 <https://doi.org/10.1109/TGRS.2003.809118>, 2003.

768 Kinar, N. J. and Pomeroy, J. W.: Measurement of the physical properties of the snowpack,
 769 *Reviews of Geophysics*, 53, 481–544, <https://doi.org/10.1002/2015RG000481>, 2015.

770 Largeron, C., Dumont, M., Morin, S., Boone, A., Lafaysse, M., Metref, S., Cosme, E., Jonas,
 771 T., Winstral, A., and Margulis, S. A.: Toward Snow Cover Estimation in Mountainous Areas
 772 Using Modern Data Assimilation Methods: A Review, *Front. Earth Sci.*, 8, 325,
 773 <https://doi.org/10.3389/feart.2020.00325>, 2020.

774 Lievens, H., Demuzere, M., Marshall, H.-P., Reichle, R. H., Brucker, L., Brangers, I., De
 775 Rosnay, P., Dumont, M., Giroto, M., Immerzeel, W. W., Jonas, T., Kim, E. J., Koch, I., Marty,
 776 C., Saloranta, T., Schöber, J., and De Lannoy, G. J. M.: Snow depth variability in the Northern
 777 Hemisphere mountains observed from space, *Nat Commun*, 10, 4629,
 778 <https://doi.org/10.1038/s41467-019-12566-y>, 2019.

779 Lievens, H., Brangers, I., Marshall, H.-P., Jonas, T., Olefs, M., and De Lannoy, G.: Sentinel-1
 780 snow depth retrieval at sub-kilometer resolution over the European Alps, *The Cryosphere*,
 781 16, 159–177, <https://doi.org/10.5194/tc-16-159-2022>, 2022.

782 López-Moreno, J. I., Fassnacht, S. R., Beguería, S., and Latron, J. B. P.: Variability of snow
 783 depth at the plot scale: implications for mean depth estimation and sampling strategies,
 784 *The Cryosphere*, 5, 617–629, <https://doi.org/10.5194/tc-5-617-2011>, 2011.

785 Lundquist, J. D., Kim, R. S., Durand, M., and Prugh, L. R.: Seasonal Peak Snow Predictability
786 Derived From Early-Season Snow in North America, *Geophysical Research Letters*, 50,
787 e2023GL103802, <https://doi.org/10.1029/2023GL103802>, 2023.

788 Luojus, K., Pulliainen, J., Takala, M., Lemmetyinen, J., Mortimer, C., Derksen, C., Mudryk, L.,
789 Moisander, M., Hiltunen, M., Smolander, T., Ikonen, J., Cohen, J., Salminen, M., Norberg, J.,
790 Veijola, K., and Venäläinen, P.: GlobSnow v3.0 Northern Hemisphere snow water
791 equivalent dataset, *Sci Data*, 8, 163, <https://doi.org/10.1038/s41597-021-00939-2>, 2021.

792 Margulis, S. A., Giroto, M., Cortés, G., and Durand, M.: A Particle Batch Smoother
793 Approach to Snow Water Equivalent Estimation, *Journal of Hydrometeorology*, 16, 1752–
794 1772, <https://doi.org/10.1175/JHM-D-14-0177.1>, 2015.

795 Margulis, S. A., Fang, Y., Li, D., Lettenmaier, D. P., and Andreadis, K.: The Utility of
796 Infrequent Snow Depth Images for Deriving Continuous Space-Time Estimates of Seasonal
797 Snow Water Equivalent, *Geophysical Research Letters*, 46, 5331–5340,
798 <https://doi.org/10.1029/2019GL082507>, 2019.

799 Marti, R., Gascoin, S., Berthier, E., De Pinel, M., Houet, T., and Laffly, D.: Mapping snow
800 depth in open alpine terrain from stereo satellite imagery, *The Cryosphere*, 10, 1361–1380,
801 <https://doi.org/10.5194/tc-10-1361-2016>, 2016.

802 Matheson, J. E. and Winkler, R. L.: Scoring Rules for Continuous Probability Distributions,
803 *Management Science*, 22, 1087–1096, <https://doi.org/10.1287/mnsc.22.10.1087>, 1976.

804 Mazzolini, M., Aalstad, K., Alonso-González, E., Westermann, S., and Treichler, D.: Spatio-
805 temporal snow data assimilation with the ICESat-2 laser altimeter,
806 <https://doi.org/10.5194/egusphere-2024-1404>, 24 May 2024.

807 McGrath, D., Webb, R., Shean, D., Bonnell, R., Marshall, H., Painter, T. H., Molotch, N. P.,
808 Elder, K., Hiemstra, C., and Brucker, L.: Spatially Extensive Ground-Penetrating Radar Snow
809 Depth Observations During NASA’s 2017 SnowEx Campaign: Comparison With In Situ,
810 Airborne, and Satellite Observations, *Water Resources Research*, 55, 10026–10036,
811 <https://doi.org/10.1029/2019WR024907>, 2019.

812 Molotch, N. P. and Bales, R. C.: SNOTEL representativeness in the Rio Grande headwaters
813 on the basis of physiographics and remotely sensed snow cover persistence, *Hydrological
814 Processes*, 20, 723–739, <https://doi.org/10.1002/hyp.6128>, 2006.

815 Nagler, T., Rott, H., Ripper, E., Bippus, G., and Hetzenecker, M.: Advancements for
816 Snowmelt Monitoring by Means of Sentinel-1 SAR, *Remote Sensing*, 8, 348,
817 <https://doi.org/10.3390/rs8040348>, 2016.

818 Oveisgharan, S., Zinke, R., Hoppinen, Z., and Marshall, H. P.: Snow water equivalent
819 retrieval over Idaho – Part 1: Using Sentinel-1 repeat-pass interferometry, *The Cryosphere*,
820 2024.

821 Painter, T. H., Berisford, D. F., Boardman, J. W., Bormann, K. J., Deems, J. S., Gehrke, F.,
822 Hedrick, A., Joyce, M., Laidlaw, R., Marks, D., Mattmann, C., McGurk, B., Ramirez, P.,
823 Richardson, M., Skiles, S. M., Seidel, F. C., and Winstral, A.: The Airborne Snow
824 Observatory: Fusion of scanning lidar, imaging spectrometer, and physically-based
825 modeling for mapping snow water equivalent and snow albedo, *Remote Sensing of*
826 *Environment*, 184, 139–152, <https://doi.org/10.1016/j.rse.2016.06.018>, 2016.

827 Raleigh, M. S. and Small, E. E.: Snowpack density modeling is the primary source of
828 uncertainty when mapping basin-wide SWE with lidar, *Geophysical Research Letters*, 44,
829 3700–3709, <https://doi.org/10.1002/2016GL071999>, 2017.

830 Rittger, K., Bair, E. H., Kahl, A., and Dozier, J.: Spatial estimates of snow water equivalent
831 from reconstruction, *Advances in Water Resources*, 94, 345–363,
832 <https://doi.org/10.1016/j.advwatres.2016.05.015>, 2016.

833 Slater, A. G., Barrett, A. P., Clark, M. P., Lundquist, J. D., and Raleigh, M. S.: Uncertainty in
834 seasonal snow reconstruction: Relative impacts of model forcing and image availability,
835 *Advances in Water Resources*, 55, 165–177,
836 <https://doi.org/10.1016/j.advwatres.2012.07.006>, 2013.

837 Smyth, E. J., Raleigh, M. S., and Small, E. E.: Particle Filter Data Assimilation of Monthly
838 Snow Depth Observations Improves Estimation of Snow Density and SWE, *Water*
839 *Resources Research*, 55, 1296–1311, <https://doi.org/10.1029/2018WR023400>, 2019.

840 Smyth, E. J., Raleigh, M. S., and Small, E. E.: Improving SWE Estimation With Data
841 Assimilation: The Influence of Snow Depth Observation Timing and Uncertainty, *Water*
842 *Resources Research*, 56, <https://doi.org/10.1029/2019WR026853>, 2020.

843 Smyth, E. J., Raleigh, M. S., and Small, E. E.: The Challenges of Simulating SWE Beneath
844 Forest Canopies are Reduced by Data Assimilation of Snow Depth, *Water Resources*
845 *Research*, 58, e2021WR030563, <https://doi.org/10.1029/2021WR030563>, 2022.

846 Sourp, L., Gascoin, S., Jarlan, L., Pedinotti, V., Bormann, K. J., and Baba, M. W.: Evaluation
847 of high-resolution snowpack simulations from global datasets and comparison with
848 Sentinel-1 snow depth retrievals in the Sierra Nevada, USA, *Hydrol. Earth Syst. Sci.*, 29,
849 597–611, <https://doi.org/10.5194/hess-29-597-2025>, 2025.

850 Sturm, M. and Liston, G. E.: Revisiting the Global Seasonal Snow Classification: An
851 Updated Dataset for Earth System Applications, *Journal of Hydrometeorology*,
852 <https://doi.org/10.1175/JHM-D-21-0070.1>, 2021.

853 Treichler, D. and Kääb, A.: Snow depth from ICESat laser altimetry — A test study in
854 southern Norway, *Remote Sensing of Environment*, 191, 389–401,
855 <https://doi.org/10.1016/j.rse.2017.01.022>, 2017.

856 Trujillo, E. and Molotch, N. P.: Snowpack regimes of the Western United States, Water
857 Resources Research, 50, 5611–5623, <https://doi.org/10.1002/2013WR014753>, 2014.

858 USDA,NRCS: SNO_{TEL} Data, 2024.

859