

~~Validation~~ Evaluation of the ALARO1-SFX (CY43T2) regional climate model over Belgium across different resolutions

Wout Dewettinck¹, Hans Van de Vyver², Daan Degrauwe², Rafiq Hamdi^{2,3}, Michiel Van Ginderachter², Bert Van Schaeybroeck^{2,4}, Kwinten Van Weverberg^{2,4}, Kobe Vandelanotte^{1,2}, Steven Caluwaerts^{1,2}, and Piet Termonia^{1,2}

¹Department of Physics and Astronomy, Ghent University (UGent), Ghent, Belgium

²Royal Meteorological Institute of Belgium (RMIB), Brussels, Belgium

³Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi 830011, China

⁴Department of Geography, Ghent University (UGent), Ghent, Belgium

Correspondence: Wout Dewettinck (wout.dewettinck@ugent.be)

Abstract. Regional climate modelling is essential for providing reliable information to understand ~~localized~~ localised impacts and guide adaptation strategies in the context of climate change. This study ~~validates~~ evaluates long-term continuous climate simulations over Belgium performed with the ALARO1-SFX model, a novel version of ALARO-1 that incorporates an advanced surface scheme (SURFEX) and improved physiographic datasets.

5 A scale-selective ~~validation~~ evaluation setup is introduced, employing a multi-level dynamical downscaling framework to assess the progressive added value of increasing resolution from the mesoscale to convection-permitting scales. The model's performance is evaluated against a gridded observational dataset and precipitation station measurements, focusing on temperature and precipitation biases, diurnal precipitation cycles, and extreme precipitation statistics.

Results indicate that the higher resolutions (12.5 km and 4 km) ~~, combined with the integration of SURFEX into ALARO1-SFX,~~
10 improve temperature and precipitation biases relative to the 25 km simulation, with the 4 km resolution providing the best representation of hourly extreme precipitation and its diurnal cycle. A dedicated sensitivity experiment isolates the contribution of the SURFEX land-surface scheme from that of resolution, attributing a substantial part of the improvement in near-surface temperature and precipitation to the introduction of SURFEX. However, all simulations exhibit varying degrees of wet and cold biases. The findings underscore the added value of convection-permitting modelling for ~~resolving extreme precipitation events~~
15 ~~and~~ improving diurnal precipitation cycles. Furthermore, the study provides ~~experimental~~ evidence that increasing model resolution ~~adds value for simulating extreme precipitation even when employing a scale-aware deep convection parameterization~~ improves the representation of extreme precipitation.

1 Introduction

As climate change intensifies, the demand for reliable and local climate information has become increasingly critical for under-
20 standing ~~localized~~ localised impacts and guiding effective adaptation strategies (Ranasinghe et al., 2021). To this end, state-of-the-art global circulation models (GCMs) perform simulations within the latest phase of the Coupled Model Intercomparison

Project, CMIP6 (Eyring et al., 2016). These models typically run at a horizontal resolution of 50–250 km (Nguyen et al., 2024; IPCC, 2021; Masson-Delmotte et al., 2021). At such resolutions, GCM simulations accurately represent large-scale dynamics and responses to external forcings. However, data at these resolutions are often too coarse for many applications, including
25 local impact models. Additionally, these models fail to account for certain sub-grid characteristics (e.g., complex topography, land-sea contrasts, or land cover heterogeneities) and mesoscale dynamical processes, such as deep convection (Rummukainen, 2016; Giorgi, 2019; Lucas-Picher et al., 2021). These processes are included in GCMs through ~~parameterization~~
parameterisation schemes. Global model simulations at higher resolutions would alleviate these problems, but these are rendered unfeasible due to the immense computational cost (Fuhrer et al., 2018).

30 Regional climate models (RCMs) address this computational cost by simulating only a limited area instead of the entire globe but at a higher resolution (Giorgi, 2019). Thereby, however, lateral boundary conditions need to be provided to RCMs by other datasets. These boundary conditions can originate from GCM simulations or from (global) reanalysis data, such as the ERA5 dataset (Hersbach et al., 2020). Typically, long-term climate simulations with an RCM employ a resolution of 10–50 km. Compared to GCM output, this finer-scale output adds value particularly in regions featuring diverse topography and
35 varying land-cover properties (Rummukainen, 2016; Doblas-Reyes et al., 2021). Nevertheless, at such scales, deep convective processes are still ~~parameterized~~
parameterised and not explicitly resolved. This is a well-known source of model uncertainty for precipitation, evident in e.g., excessive light precipitation and a too early peaking diurnal convective cycle (Ban et al., 2014; Fossier et al., 2015; Hohenegger et al., 2008; Kendon et al., 2012; Vergara-Temprado et al., 2020). However, for climate models with resolutions higher than ~ 5 km, these processes are in fact (partially) resolved (Prein et al., 2015; Lucas-Picher et al.,
40 2021). Models operating at these scales are referred to as convection-permitting models (CPMs).

CPMs typically turn off their convective ~~parameterization~~
parameterisation schemes (Coppola et al., 2020). However, the boundary between convection-permitting and ~~convection-parameterized~~
convection-parameterised scales is not well defined. In the so-called ‘grey zone of convection’ (~ 4 –10 km), deep convection is only partially resolved, making it unclear whether ~~parameterization~~
parameterisation is required (Vergara-Temprado et al., 2020). In the grey zone, the explicit and
45 parameterised representations compete with one another, which can result in double counting of convective transport or no counting at all (Gross et al., 2018). This can lead to convection being triggered too late by the parameterisation, leading to spurious grid-scale storms, or to convection being triggered too early, systematically shifting convective onset too early in the day (Gross et al., 2018).

The development of scale-aware convective ~~parameterization~~
parameterisation schemes addresses this problem. Rather
50 than simply switching convection on or off, these schemes adapt to the resolution by adjusting their tuning parameters accordingly. Scale awareness can be introduced through several approaches, for example by introducing a prognostic variable for the convective fraction of a grid cell (Gerard et al., 2009), scaling the convective area fraction and mass flux with the grid-box area (Grell and Freitas, 2014; Han et al., 2017), or adjusting scheme parameters explicitly as a function of the grid spacing (Zheng et al., 2016). Therefore, models with these ~~parameterization~~
parameterisation schemes can operate at both
55 coarse and fine resolutions and even in the grey zone.

CPMs have been shown to improve some notable limitations associated with the ~~low-resolution-parameterization~~ parameterisation of deep convection used in lower resolution models. For instance, they improve the diurnal cycle of precipitation, as demonstrated in studies across Europe, including Belgium (De Troch, 2016; Brisson et al., 2016), and more widely across the continent (Berthou et al., 2020). Furthermore, while RCMs generally underestimate precipitation extremes at sub-daily durations (Berg et al., 2019), these extremes are better represented in long-term CPM simulations, as shown by Coppola et al. (2020) and Fosser et al. (2024), and studies over Belgium (De Troch et al., 2013; Tabari et al., 2016; Van De Vyver et al., 2021) and the Alpine-Mediterranean region (Müller et al., 2023, 2024). This improvement in extreme precipitation was also noted in the evaluation of an ensemble of kilometre-scale simulations over Europe and the Mediterranean (Pichelli et al., 2021; Ban et al., 2021; Ha et al., 2024). While precipitation is often the primary focus of added value studies for CPMs, it has also been found that temperature is better represented due to improved representations of topography, clouds, precipitation, and land cover (Leutwyler et al., 2017; Lucas-Picher et al., 2021; Soares et al., 2022).

Despite these advancements, significant model uncertainties remain, particularly in climate projections. These ~~uncertainties~~ model uncertainties arise from the inherent limitations of the models themselves, and differ amongst models because of different choices in their architectures. On short time horizons (~30 years), these model uncertainties are often larger than ~~projection-uncertainties in the short term~~ the uncertainty related to the chosen climate scenario (Lehner et al., 2020). While increasing resolution tends to reduce ~~these model~~ uncertainties, as suggested by Fosser et al. (2024), they remain substantial even in CPMs. This is because higher resolution alone is not a panacea; e.g., microphysics processes remain ~~parameterized~~ parameterised in CPMs, contributing to significant uncertainties (Prein et al., 2015; Lucas-Picher et al., 2021). Therefore, it is crucial to maintain model diversity to reliably estimate these uncertainties (Doblas-Reyes et al., 2021). However, the diversity of RCMs is decreasing, as seen in the current iteration of EURO-CORDEX simulations, due to the high community effort required to develop and maintain such models (Jacob et al., 2020; CORDEX).

In this study, we present ALARO-1, an updated version of the ALARO-0 regional climate model originally introduced by ~~the~~ De Troch et al. (2013), where the surface scheme ~~ISBA is now replaced by the SURFEX (Surface Externalisée) scheme (Hamdi et al., 2014). This is now handled by the externalised SURFEX~~ platform (Masson et al., 2013; Hamdi et al., 2014) rather than by the built-in ISBA ² scheme (Noilhan and Planton, 1989; Noilhan and Mahfouf, 1996). Within SURFEX, natural land surfaces are still treated by ISBA, but SURFEX additionally introduces a tiled representation of the surface with dedicated schemes for urban areas, lakes, and seas. This is necessary to use ALARO-1 for land-use climate studies. The ALARO model was originally developed for numerical weather prediction (NWP) by the ALADIN ~~consortium~~ ³ consortium ⁴, which comprised the national (hydro)meteorological services of 16 European and northern African countries (Termonia et al., 2018a). The ALARO model is one of three canonical model configurations (CMCs) within the ALADIN framework, along with the

¹Surface Externalisée.

²Interaction Soil Biosphere Atmosphere.

³Aire Limitée Adaptation Dynamique Développement International.

⁴The ALADIN consortium merged with the HIRLAM (High Resolution Limited-Area Model) consortium in 2020 to form the ACCORD (A Consortium for Convection-scale modelling Research and Development) consortium.

baseline ALADIN model and the AROME~~model~~⁵ ~~model~~⁶. For a more detailed description, we refer to Termonia et al. (2018a) and to Sect. 2.

The key feature of the ALARO physics package is the 3MT (Modular Multiscale Microphysics and Transport) scheme, which handles convection, turbulence, and microphysics (Gerard and Geleyn, 2005; Gerard, 2007; Gerard et al., 2009). The 3MT scheme has a unique scale-aware character, so that it can determine itself which processes are resolved at a specific resolution. This provides a smooth multiscale transition from ~~fully-parameterized~~ ~~fully-parameterised~~ mesoscale resolutions over the grey zone to fully convection-resolving scales. This multi-scale feature of the ALARO model is unique with respect to other convection-permitting models and makes it the ideal candidate for a multi-resolution model ~~validation~~ ~~evaluation~~ from the mesoscale up to the convection-permitting scales. Climate simulations will therefore be ~~validated~~ ~~evaluated~~ at three different resolutions in this study. ~~Validating~~ Specifically, this evaluation focuses on the climatology of near-surface temperature and precipitation, as well as the representation of precipitation extremes and the diurnal precipitation cycle.

Evaluating models at different resolutions is essential for both practical and scientific reasons. From a practical perspective, the computational demands of high-resolution simulations, particularly at convection-permitting scales, are prohibitive for many research groups. As a result, lower-resolution RCMs remain widely used. Scientifically, examining how model representation of the climate evolves with changing resolution is crucial for improving our understanding and interpretation of simulation results.

~~The ALARO model originally used the ISBA (Interaction Soil-Biosphere-Atmosphere) scheme (Noilhan and Planton, 1989; Noilhan and McFarlane, 1997) for land-surface parameterization, which simulates energy and water exchanges between the surface and the atmosphere. The new ALARO1-SFX configuration relies on SURFEX (Masson et al., 2013) in order to extend ISBA's vegetation-based approach by including multiple surface types: nature, urban areas, lakes, and seas. This step is necessary to use ALARO-1 for land-use climate studies.~~

In this study, we evaluated long-term climate simulations with the ALARO-1 model at three different resolutions: 25 km, 12.5 km, and 4 km. Previous climate simulations with ALARO-0 have been extensively ~~validated~~ ~~evaluated~~, particularly over Belgium (De Troch et al., 2013; Hamdi et al., 2014; Giot et al., 2016; Tabari et al., 2016; Berckmans et al., 2017; Top et al., 2021). These studies employed ALARO cycle 36, following the versioning system aligned with the cycles of the global IFS ~~and ARPEGE~~⁷ ~~and ARPEGE~~⁸ models (Termonia et al., 2018a). In contrast, this study ~~utilized~~ ~~utilises~~ a more recent version, ALARO cycle 43t2. Moreover, from cycle 40 onwards, ALARO underwent substantial updates, leading to the development of ALARO-1. ~~This new version marked a step-wise change compared to the original ALARO-0.~~ It introduced significant advancements, including the ACRANEB2 radiation scheme (Mašek et al., 2016; Geleyn et al., 2017) and the TOUCANS scheme (Đurán et al., 2014; Marquet and Geleyn, 2013) for turbulence and shallow convection. The use of ALARO-1 at higher resolutions in this study marks a novel contribution. The combination of ALARO-1 with SURFEX v8.0 (as used in this study) will be denoted by ALARO1-SFX. More generally, the term ALARO-1 will refer to the current version of the ALARO model,

⁵Application of Research to Operations at Mesoscale.

⁶The name ALARO stands for ALADIN-AROME.

⁷Integrated Forecasting System.

⁸Action de Recherche Petite Echelle Grande Echelle.

irrespective of the surface scheme used. The broader term ALARO is used when referring to the model in general or when the specific cycle is clear from the context.

120 The structure of this paper is as follows. Section 2 provides a discussion of the model and the experimental design. Following this, the observational datasets and verification methods used in this study are described in detail. Section 3 presents the results of the ~~validation~~evaluation, highlighting the model’s performance across different resolutions and comparing it with observational data. In Sect. 4, these results are discussed and placed in a broader context, comparing them with those from previous ALARO-simulations. Finally, Sect. 5 draws the conclusions, ~~summarizing~~summarising the key findings of the study
125 while addressing the limitations and suggesting directions for future research.

2 Data & methods

2.1 Model and experimental design

2.1.1 ALARO1-SFX model

Historically, the ALARO model has its roots in the global ~~Integrated Forecast System (IFS)~~IFS of the ECMWF⁹ and the
130 global ARPEGE¹⁰ model of Météo-France. These global models were adapted to run on a limited area, which resulted in ALADIN. The ALADIN model employs the same physical ~~parameterization~~parameterisation schemes as the global ARPEGE model. ALARO is in turn a further development of ALADIN: it shares the dynamical core of ALADIN, but it contains different physics ~~parameterizations~~parameterisations.

The dynamical core of ALARO can be run both hydrostatically and non-hydrostatically (Bubnová et al., 1995; Termonia
135 et al., 2018a). In both configurations, the dynamical equations are spatially ~~discretized~~discretised using a spectral representation of double-Fourier modes. For the time-step algorithm, a semi-implicit semi-Lagrangian scheme is employed. In the vertical dimension, the model uses a hybrid terrain-following pressure coordinate (Simmons and Burridge, 1981; Laprise, 1992).

The ALARO-1 physics package combines several parameterisations. Radiative transfer is treated by the ACRANEB2 scheme (Mašek et al., 2016; Geleyn et al., 2017), a broadband scheme with a single shortwave and a single longwave spectral
140 interval that retains the full cloud–radiation interaction at every time step through a net-exchanged-rate decomposition. Turbulent transport and shallow convection are parameterised by the prognostic-TKE-based TOUCANS scheme (Đurán et al., 2014; Marquet and Gel
, which is valid across the full range of atmospheric stability and represents the effect of moisture and phase changes through a modified moist Brunt–Väisälä frequency, removing the need for a separate shallow-convection scheme. Deep convection is represented by the scale-aware 3MT scheme (Gerard et al., 2009), a mass-flux formulation with prognostic updraught
145 and downdraught properties that abandons the quasi-equilibrium assumption and mimics cloud-resolving behaviour in the convection-permitting limit. The associated stratiform cloud and microphysical processes are treated within the same 3MT framework, ensuring a consistent transition between resolved and subgrid condensation as resolution increases.

⁹European Centre for Medium-Range Weather Forecasts.

¹⁰~~Action de Recherche Petite Echelle Grande Echelle~~

ALARO-1 can be coupled to the land-surface model SURFEX, which considers four types of land surface: nature, urban areas, lakes and seas. For every grid box, a fraction is defined for each of these four types in a tiling approach. The interaction with the free atmosphere is then calculated by type-specific schemes: ISBA for nature, TEB for urban areas, WATFLUX for lakes and SEAFUX for seas. The separate variables of each tile are finally aggregated with a weighted average to compute grid-box values.


2.1.2 Experimental design

Firstly, a simulation ~~was-is~~ run at a resolution of 25 km (see Fig. 1) with the ALARO-1 model using boundary conditions from the ERA-5 reanalysis dataset (Hersbach et al., 2020). Next, this simulation data ~~was-is~~ dynamically downscaled to two higher resolutions with a multi-level nesting strategy: each higher-resolution run ~~was-is~~ coupled hourly to the previous lower-resolution run at the boundaries. The output of all simulations ~~was-is~~ stored at an hourly frequency. This downscaling ~~resulted~~ results in simulations at 12.5 km and 4 km coupled to, respectively, the 25 km and 12.5 km simulations. Henceforth, we will refer to each simulation as ALARO- x km with x the resolution in ~~kilometers~~kilometres. For the lateral boundary coupling, the Davies relaxation scheme ~~was-is~~ employed (Davies, 1976), with a coupling zone of 8 points (Termonia et al., 2018a). The coupled prognostic variables are temperature, the horizontal wind components, specific humidity, and surface pressure.


All simulations span a period of 32 years from 1 January 1991 to 31 December 2022. The first year ~~was-is~~ discarded for each simulation to ensure the model, specifically the soil, ~~reached-reaches~~ an equilibrium state. The simulations are continuous, with monthly updates to certain (otherwise) constant fields including the sea surface temperature, the surface roughness length, surface albedo, surface emissivity, and the vegetation parameters. This monthly update methodology ~~was-is~~ also applied in Giot et al. (2016) and Top et al. (2021).

The different simulation domains are shown in ~~the top panel of~~ Fig. 1a. The ALARO-25km and ALARO-12km simulations both cover the EURO-CORDEX domain (Jacob et al., 2020). The ALARO-4km simulation covers a smaller domain due to computational constraints. This domain is the operational NWP domain of the Royal Meteorological Institute of Belgium. It is centred on Belgium and covers parts of Western Europe.

The main technical specifications of the simulations are outlined in Table 1. All simulations ~~used the hydrostatic dynamical core and are run with 46 vertical levels. Moreover, they all used the same physics parameterization~~ use the same physics parameterisation framework, including the 3MT scheme for deep convection, which adapts to the model resolution due to its scale-aware nature. Moreover, they all are run with 46 vertical levels and use the hydrostatic dynamical core. Although the 4 km simulation reaches the convection-permitting range, where non-hydrostatic effects can become relevant, we consider the hydrostatic core appropriate at this resolution. For ALARO specifically, Van Genderachter et al. (2020) compared both dynamical cores at 4 km over a domain of intense deep convection and found that the hydrostatic configuration performed better, as the ALARO physics package, including the scale-aware 3MT deep-convection scheme, is tuned to the hydrostatic core. Retaining the hydrostatic core is moreover consistent with the operational use of ALARO at ~ 4 km. Finally, it keeps the dynamical core identical across all three resolutions so that resolution effects are not confounded with a change of core.



(a)



(b)

Figure 1. (a) Map showing the three simulation domains used in this study, with each domain corresponding to an ALARO-1 simulation at a specific resolution: 25 km, 12.5 km, and 4 km, as indicated in the figure legend. (b) Map of Belgium displaying the elevation above sea level, with precipitation measurement stations marked. The marker styles (circles, squares, and triangles) denote the topographic regions each station is assigned to: Low-Belgium (0–50 m), Middle-Belgium (50–200 m), and High-Belgium (>200 m). ©EuroGeographics and ©GTOPO30

Table 1. Technical specifications of simulations with the ALARO-1 model. This table outlines the main differences between ~~each of the three~~ simulations, while the common features are described in the text. ALARO-12km-noSFX is a sensitivity experiment used only to isolate the effect of SURFEX (Sect. 3.3).

Simulation	ALARO-25km	<u>ALARO-12km-noSFX</u>	ALARO-12km	ALARO-4km
Resolution	25 km	12.5 km	<u>12.5 km</u>	4 km
Domain size	251 × 251 points	499 × 499 points	<u>499 × 499 points</u>	421 × 421 points
Time step	450 s	300 s	<u>300 s</u>	180 s
Lateral boundary coupling	ERA-5 reanalysis	ALARO-25km	<u>ALARO-25km</u>	ALARO-12km
Land surface scheme	ISBA	<u>ISBA</u>	SURFEX v8.0	SURFEX v8.0

One significant difference between the simulations concerns the land-surface scheme. SURFEX ~~was~~is not employed for ALARO-25km: instead, the original ISBA scheme ~~was used~~is used. The main reason for omitting SURFEX in the 25 km simulation is that urban effects are minimal at this resolution, making its inclusion less relevant. This setup also aligns more closely with the one used in previous simulations from Giot et al. (2016).

The SURFEX land-surface scheme is configured identically in the ALARO-12km and ALARO-4km simulations. Natural land surfaces are represented by the ISBA scheme in its force-restore formulation, which discretises the soil into three layers: a thin surface layer, a root-zone layer, and a deeper sub-root reservoir. Soil hydraulic and thermal properties are derived from the ECOCLIMAP sand and clay fractions (Masson et al., 2003) through the Clapp and Hornberger (1978) pedotransfer relations. The nature tile is treated as a single aggregated patch, and vegetation is prescribed from the ECOCLIMAP climatology without interactive photosynthesis, so that root fractions follow the standard ISBA force-restore profile. Snow is represented by the single-layer force-restore scheme of Douville et al. (1995). Turbulent fluxes over sea and inland water are computed with the direct Charnock method (Charnock, 1955).

Because ALARO-25km and ALARO-12km differ in both resolution and land-surface scheme, the 25 km to 12 km change conflates two effects. To isolate the impact of coupling ALARO-1 to SURFEX, we have performed a sensitivity experiment with an additional simulation, ALARO-12km-noSFX, identical to ALARO-12km but retaining the original ISBA scheme. Holding resolution fixed at 12.5 km, any difference between ALARO-12km and ALARO-12km-noSFX is attributable to the land-surface scheme alone. This experiment is analysed separately in Sect. 3.3; all other analyses retain the original set of three simulations.

2.2 Observational data

The simulations ~~were validated~~are evaluated using the CLIMATE-GRID (CG) dataset (Journée et al., 2019), a Belgian 5 km gridded observational dataset based on interpolated station data and developed at the Royal Meteorological Institute of

Belgium. The dataset spans the period from 1961 until the present day. It includes daily average temperature and daily total precipitation, which are provided as areal averages over each pixel of the 5 km grid.

205 For sub-daily observations of precipitation, we employed station measurements from five measurement networks: VMM in Flanders, DGH in Wallonia, IBGE in Brussels, and the HYDRO and AWS networks covering all of Belgium. The VMM and DGH networks are managed by the Vlaamse Milieumaatschappij and the Direction de la Gestion Hydrologique of SPW Mobilité et Infrastructures, respectively. The IBGE network, established by the Ministry of the Brussels-Capital Region, has been managed by Brussels Environment since 2007 (Dehem et al., 2010; Journée et al., 2014). The AWS and HYDRO networks
210 are operated by the Royal Meteorological Institute of Belgium (Van de Vyver, 2021). The station locations are shown in the bottom-panel of Fig. 1b, while details about each network are provided in table C1. Rainfall-Table C1. Precipitation data from all networks were are aggregated to an hourly resolution to align with the temporal resolution of the simulation output.

2.3 Validation-Evaluation metrics and data pre-processing

The ALARO-1 simulations were are evaluated by calculating the mean bias for near-surface air temperature (simply called
215 temperature from here on) and accumulated precipitation depth on annual, seasonal, and monthly timescales. For an exact mathematical formulation, we refer to appendix A.

To assess whether the biases are statistically distinguishable from zero, we apply a non-parametric bootstrap at each grid point. For a given field, the 31 years (1992–2022) are resampled with replacement 10^3 times. The climatological bias (simulation minus observations) is then recomputed for each resample. Next, the 95 % confidence interval is taken as the 2.5th–97.5th
220 percentiles of the resulting distribution. A grid point is deemed significant where this interval excludes zero. Each simulation and the CLIMATE-GRID observations are anchored to the same historical years: the simulations are all originally driven by ERA5 reanalysis over the identical 1992–2022 period. Therefore, the resampling is paired: the same resampled set of years is used for the simulated and observed fields within each iteration. The same paired-bootstrap procedure is applied to the domain-averaged monthly biases, where the bias is first averaged over all grid points before the confidence interval is
225 computed. A month is deemed significant where this interval excludes zero.

To ensure a consistent comparison, all data were are regridded to a common latitude-longitude grid with a spacing of 0.07° longitude by 0.045° latitude, which closely aligns with the 5 km resolution of the CG dataset. For precipitation, a conservative remapping was is applied to all simulations to preserve the spatially averaged precipitation depth (Jones, 1999). For temperature, the same conservative procedure was is used to reproject ALARO-4km. For ALARO-12km and ALARO-25km, another
230 regridding method was is employed for temperature. Since low-resolution data were are interpolated to a higher resolution, a conservative regridding method would effectively function as a nearest-neighbour approach in such cases. Therefore, a bilinear interpolation procedure was is used instead, as it was is more suitable for the smooth nature of temperature fields.

2.4 Definition of extreme precipitation

Simulations at convection-permitting scales generally add value through the improved representation of convection and, consequently, of precipitation in general (Prein et al., 2015; Ban et al., 2014; Lucas-Picher et al., 2021; Berthou et al., 2020). There-
235

fore, this ~~validation study also focused~~ evaluation study also focuses on the statistics of extreme precipitation over Belgium. Extreme precipitation events ~~were~~ are here defined as the annual maximum (AM) value for a given duration at a specific location or grid point. An advantage of this definition is its simplicity: there is no need to determine an arbitrary threshold. AM values ~~were~~ are selected for durations of 1, 2, 3, 6, 12, 24, 48, and 72 hours using a rolling-window approach over the hourly
240 precipitation data.

The statistics of these AM values ~~were~~ are calculated by fitting a ~~generalized~~ generalised extreme value (GEV) distribution, as defined in eqs. (B1) - (B2) (Coles, 2001). A GEV distribution is defined by three parameters: the shape ξ , location μ , and scale σ . These parameters ~~were~~ are fitted for each duration separately with the L-moments method as defined by Hosking (1990). Consequently, these parameters ~~were~~ are used to calculate the ~~T -year return level of~~ return level $z_d(T)$, i.e. the d -hourly
245 precipitation intensity $z_d(T)$ expected to be exceeded once every T years, where d is the duration and T the return period. For a mathematical description, we refer to appendix B.

Next, we ~~applied~~ apply the regional frequency analysis framework (RFA), as introduced by Hosking and Wallis (1993). The idea behind this framework is to aggregate ~~rainfall~~ precipitation data from different locations that share the (presumably) identical underlying distribution. A single fitting procedure is then performed for an entire region instead of fitting a separate
250 distribution in each point, allowing for a more robust fit. Note that the ~~rainfall data were~~ precipitation data are aggregated and not averaged over a region. Recent literature, such as the work by Carreau et al. (2017) and Guerreiro et al. (2024), further supports the use of RFA for improving precipitation extreme estimations.

We subdivided Belgium into three distinct regions based on elevation: Low-Belgium (0–50 m), Middle-Belgium (50–200 m) and High-Belgium (> 200 m), as shown in Fig. 1. ~~Elevation was~~ Elevation is used as a discriminatory parameter since
255 it correlates strongly with the annual precipitation totals, which themselves are closely linked to precipitation extremes (Van De Vyver, 2012). This subdivision strikes a balance between reducing uncertainties and maintaining relatively homogeneous climatic conditions within each region, as suggested by Van De Vyver (2012).

Intensity-Duration-Frequency (IDF) statistics jointly characterise the intensity of extreme precipitation as a function of both duration and return period, and are a standard tool for hydrological design and flood-risk assessment (Tabari et al., 2016)
260 The return levels for each region are subsequently presented per return period as a function of the duration to construct ~~Intensity-Duration-Frequency (IDF) curves~~ these IDF statistics. In order to gauge the return-level uncertainty, we derived the 95% confidence interval using 10^4 bootstraps, obtained by resampling yearly data. For each simulation, the observational return levels ~~were~~ are compared with their simulated counterparts by calculating the relative error. The error uncertainty ~~was~~ is acquired with a pairwise calculation of the error between all observational and simulated bootstrap samples, respectively. This
265 calculation ~~resulted~~ results in a distribution of 10^8 samples for the relative error.




Figure 2. Bias in mean annual near-surface air temperature of each simulation (~~25 km, 12.5 km, and 4 km resolutions~~) with respect to the daily gridded observational CLIMATE-GRID dataset over Belgium, for the period 1992–2022. (a) ALARO-25km, (b) ALARO-12km, (c) ALARO-4km. Hatching indicates grid points where the bias is not significantly different from zero at the 95 % confidence level, based on a paired bootstrap resampling of years.

3 Results

3.1 ~~Validation of average~~ Average climatology

Figure 2 shows that ALARO-1 at all three resolutions exhibits a cold temperature bias with respect to the Belgian CLIMATE-GRID observations. This bias is most pronounced in the eastern, more elevated region. Near the coast, all simulations exhibit a small warm bias. The cold bias is statistically significant across almost the entire domain in ALARO-25km, whereas for ALARO-12km and ALARO-4km large parts of the interior are not significantly different from zero (hatched in Fig. 2), consistent with their much smaller domain-mean biases of -0.25 and -0.32 °C. Furthermore, the spatial bias patterns for ALARO-12km and ALARO-4km are very similar. Certain urban areas can be discerned in these simulations presenting themselves as areas with a warm bias.

While the bias difference between ALARO-12km and ALARO-4km is small, a large difference is present between ALARO-25km and ALARO-12km. As outlined in Table 1, apart from the resolution difference, the primary distinction between these two simulations is the inclusion of SURFEX into ALARO-12km. ~~These changes seem to considerably alter the current-day climatology, raising the~~ The dedicated ALARO-12km-noSFX experiment (Sect. 3.3) shows that the inclusion of SURFEX, rather than the resolution increase, is the dominant driver of this difference, raising the annual-mean temperature by nearly 1 °C. Overall, ALARO-12km provides the best representation of mean annual temperature.

Considering the annual cycle of the biases, a more complex picture emerges. The spatially averaged biases per month are plotted in Fig. 3, while spatial bias patterns are shown per season in Fig. D1. In each season, ALARO-12km and ALARO-4km perform similarly, except during summer, when both simulations feature a warm bias which is larger in ALARO-12km, particularly in July and August when the bias exceeds 1 °C. This warm bias partially compensates for the cold bias in other




Figure 3. The annual cycle of the temperature bias (left) and relative precipitation bias (right) for each simulation (25 km, 12.5 km, and 4 km resolutions) with respect to the gridded observational CLIMATE-GRID dataset, averaged over Belgium, for the period 1992–2022. Square markers denote months where the domain-averaged bias is significantly different from zero at the 95 % confidence level (paired bootstrap resampling of years); circular markers denote non-significant months.

seasons on an annual basis. Consequently, ALARO-12km exhibits a smaller annual cold bias than ALARO-4km, despite ALARO-4km performing better in more seasons. Additionally, both ALARO-12km and ALARO-4km strongly underestimate the temperature in spring, which also corresponds to the largest seasonal cold bias for ALARO-25km. The minimal largest negative bias, which is around -1.5°C , is attained in May. Finally, during the The domain-averaged monthly biases of ALARO-25km are significant year-round except in September, whereas for ALARO-12km and ALARO-4km significance is confined to the seasonal extremes: the spring cold bias (March–May/June), the summer warm bias (August), and a November cold bias (Fig. 3a).

During the seasons with warming sea surface temperatures (SST), spring and summer, there is a cold bias near the coast, while the opposite is true for autumn and winter (see Fig. D1). This may be attributed to the monthly replacement of SST, creating a lag effect. In addition, coastal grid points partly consist of sea: the diagnosed 2 m temperature is the tile-weighted average over the land and sea fractions, so the (lagged) sea surface temperature is directly reflected in the near-surface temperature of these mixed cells.

In terms of significance (Fig. D1), the cold bias of ALARO-25km is significant across the whole domain in every season. For ALARO-12km and ALARO-4km the bias is largely insignificant, except in spring, where the cold bias is significant domain-wide. In summer, ALARO-12km exhibits a significant warm bias over much of the domain, whereas in ALARO-4km this warm bias is weaker and largely insignificant, consistent with its cooler summer near-surface temperature.

ALARO-1, at all three resolutions, features a wet bias both annually (Fig. 4) and across all months (Fig. 3) and seasons (Fig. D2). A notable exception is the dry bias in, except for summer (July and August) in ALARO-12km and ALARO-4km,




Figure 4. Relative bias in mean annual precipitation of each simulation (~~25 km, 12.5 km, and 4 km resolutions~~) with respect to the daily gridded observational CLIMATE-GRID dataset over Belgium, for the period 1992–2022. (a) ALARO-25km, (b) ALARO-12km, (c) ALARO-4km. Hatching indicates grid points where the bias is not significantly different from zero at the 95 % confidence level, based on a paired bootstrap resampling of years.

which coincides with a warm bias in these simulations. Similarly to temperature, the bias in annual precipitation is smallest for ALARO-12km. ALARO-4km is slightly wetter, while ALARO-25km features a bias nearly twice as high as ALARO-12km.

305 The wet bias is significant over essentially the whole domain for all three simulations (Fig. 4).

However, once more, ALARO-4km performs best when evaluating the seasons separately. ALARO-12km shows a smaller bias only during springtime. Similarly to temperature, the precipitation bias in spring is substantially larger (in absolute values) compared to the other seasons. Regarding significance (Fig. D2), the wet bias is significant across essentially the whole domain for all three simulations in autumn, winter, and spring. Summer is the exception: ALARO-25km retains a significant wet bias only in the south of the domain, whereas ALARO-12km and ALARO-4km show a significant dry bias in the north. Consistent with its smaller domain-mean dry bias, ALARO-4km shows a significant dry bias over a smaller northern area than ALARO-12km, with the central domain no longer significantly biased.

Figure 5 shows the diurnal cycle of precipitation in Uccle, Belgium. The diurnal cycle is first ~~analyzed~~ analysed across all hours ~~-(Fig. 5a).~~ All simulations overestimate ~~rainfall-precipitation~~ for all hours. Apart from the systematic bias, ALARO-4km
315 neatly reproduces the diurnal ~~rainfall-precipitation~~ variation. The observed afternoon peak around 16 UTC, for instance, is also present in ALARO-4km albeit with a higher intensity. ALARO-12km displays a smaller and slightly delayed peak, while ALARO-25km lacks a clear peak, maintaining roughly uniform precipitation amounts throughout the day. During wet hours (i.e. hours with a precipitation rate > 0.1 mm), on the other hand, there is a dry bias for all models ~~-(Fig. 5b).~~ ALARO-4km again aligns most closely with observations, followed by ALARO-12km and ALARO-25km. The afternoon peak at 16
320 UTC remains visible for the observations but is less pronounced and slightly delayed for ALARO-4km. The delay is even larger for ALARO-12km and ALARO-25km, with progressively smaller peaks as resolution decreases. In summer (Fig. 5c), average precipitation values and peak amplitudes are considerably higher, although the simulated peaks occur slightly later and




Figure 5. Diurnal cycle of precipitation in Uccle, Belgium, for station measurements (observations) and for each simulation (25 km, 12.5 km, and 4 km resolutions), [for the period 1992–2022](#). The dashed lines denote the average values over the considered hours. (a) All hours. (b) Only wet hours ($> 0.1 \text{ mm h}^{-1}$). (c) Only wet hours ($> 0.1 \text{ mm h}^{-1}$), summer. (d) Only wet hours ($> 0.1 \text{ mm h}^{-1}$), winter.

are less intense at lower resolutions. In winter ([Fig. 5d](#)), no clear peak is observed, and the average precipitation intensity is approximately half of the summer values.

Overall, these results indicate that although the simulations reproduce the timing of the diurnal cycle reasonably well, they [tend to overestimate precipitation frequency](#) — raining too often with too little intensity — especially at coarser resolutions. [To quantify this, we decompose the mean hourly precipitation \$\bar{P}\$ into the frequency and intensity contributions from wet hours, defined as hours with a precipitation rate \$> 0.1 \text{ mm h}^{-1}\$ \(Table 2\). All simulations overestimate the wet-hour frequency \$\bar{n}_{\text{wet}}\$ \(11.8 % observed\) while underestimating the wet-hour intensity \$\bar{P}_{\text{wet}}\$ \(0.791 \$\text{mm h}^{-1}\$ observed\), most severely at 25 km \(17.2 % and 0.692 \$\text{mm h}^{-1}\$ \). Increasing resolution reduces the frequency bias and raises the wet-hour intensity, shifting the simulations](#)

Table 2. Decomposition of the mean hourly precipitation in Uccle into wet hour intensity and frequency contributions, for the period 1992–2022. A wet hour is defined as an hour with a precipitation rate > 0.1 mm h⁻¹. $\bar{P}$ is the mean precipitation over all hours; $\bar{P}_{\text{wet}}$ is the mean intensity during wet hours; $\bar{n}_{\text{wet}}$ is the corresponding frequency of occurrence.

<u>Quantity</u>	<u>Observations</u>	<u>ALARO-25km</u>	<u>ALARO-12km</u>	<u>ALARO-4km</u>
$\bar{P}$ [mm h ⁻¹]	0.093	0.124	0.112	0.118
$\bar{P}_{\text{wet}}$ [mm h ⁻¹]	0.791	0.692	0.722	0.768
$\bar{n}_{\text{wet}}$ [%]	11.8	17.2	15.0	14.8
$\bar{P}_{\text{wet}} \times \bar{n}_{\text{wet}}$ [mm h ⁻¹]	0.093	0.119	0.108	0.114




Figure 6. Intensity-Duration-Frequency (IDF) curves-statistics for three topographic regions over Belgium, using all available years for each dataset (observational record lengths are given in Table C1; simulations span 1992–2022): (a) Low-Belgium (<50m), (b) Middle-Belgium (50m–200m), and (c) High-Belgium (>200m). The curves-markers show the return-level-return-level precipitation intensities for different durations as a function of duration on a log-log scale (both axes), calculated using the Regional Frequency Analysis (RFA) framework. The colors represent the dataset -stations- (station observations or simulations), while the markers-represent-marker styles denote the return-levels for different return periods (2, 10, and 50 years).

from frequent-weak towards less-frequent, more-intense precipitation. This confirms that the mean wet bias arises from a too-high precipitation frequency combined with too-weak wet-hour intensities, rather than from excessive intensity.

3.2 Extreme precipitation

The calculated IDF-statistics-IDF statistics are presented in Fig. 6 for each topographic region. The difference in observational
335 return levels between the regions is apparent for both sub-daily durations (< 6h) and durations extending beyond one day (>=

24h). In the former case, the values for Middle- and High-Belgium are similar to one another, but those for Low-Belgium are smaller, while in the latter case, the difference between Middle- and Low-Belgium disappears and the return levels over High-Belgium exceed those over the other two regions.

The observational return levels decrease ~~linearly with rainfall~~ approximately linearly with precipitation duration on a log-
340 log scale (see Fig. 6). ~~For the~~, corresponding to the power-law scaling of extreme precipitation. The simulated return levels
~~, a linear relationship can also be discerned, except for~~ follow the same linear behaviour for durations of 6 h and longer, but
deviate at short durations ($< 6\text{h}$), where ~~the values start to~~ they curve downward, away from the observations. ~~This effect is~~
~~quantitatively assessed by evaluating the linear regression coefficients and~~ To quantify this behaviour, a linear regression of the
log-transformed return levels against the log-transformed duration was performed for each dataset, region, and return period.
345 The resulting slopes and coefficients of determination (R^2 -values, as, the fraction of variance in the return levels explained
by the linear fit) are shown in Fig. D3. ~~The coefficients show that the observational return levels exhibit the steepest slope;~~
~~the fitted lines themselves are not overlaid on Fig. 6. The slope is steepest for the observations,~~ followed by the simulations
in order of decreasing resolution. ~~Furthermore, the fitted linear trend has a higher R^2 -value~~ (ALARO-4km, ALARO-12km,
ALARO-25km). The linear fit is also best for the observations (R^2 closest to 1), again followed by ALARO-4km, ALARO-
350 12km, and ALARO-25km, ~~respectively. Some differences can also be discerned for different return periods: overall, the return~~
~~levels with a longer return period exhibit a steeper linear fit, but have a smaller~~ reflecting the increasing downward curvature at
short durations as resolution decreases. Comparing return periods, longer return periods yield a steeper (more negative) slope
but a lower R^2 -value.

Despite its strong wet bias for average ~~rainfall~~ precipitation, ALARO-25km underestimates the sub-daily ($< 6\text{h}$) return levels
355 for all regions and return periods, as shown in Fig. 7. For the (multi-)daily ($\geq 24\text{h}$) durations, the performance depends on
the return period. The least extreme events i.e. those with the shortest return period, are overestimated, while the events with
the longest return period generally tend to be underestimated, although the uncertainty ~~was~~ is large. Conversely, ALARO-
12km provides smaller return levels than ALARO-25km for sub-daily ($< 6\text{h}$) durations, except for hourly events in Low- and
Middle-Belgium ~~(Fig. 7a,d)~~. Hence, the sub-daily extreme precipitation intensities are also underestimated by this simulation.
360 For (multi-)daily ($\geq 24\text{h}$) durations, this intensity decrease with higher resolution is also present, especially in High-Belgium
~~(Fig. 7g-i)~~. Finally, for ALARO-4km the hourly return levels are substantially closer to the observational values for a 2-
year return period and markedly closer for longer return periods. Generally, the ALARO-4km extremes increase in intensity
compared to the ALARO-12km simulation, leading to improvements for some durations and regions, but not universally.

Figure 8 presents the diurnal and seasonal cycles of the frequency of annual-maximum hourly precipitation intensities over
365 Belgium. In the diurnal cycle, the observations clearly show a primary peak in the afternoon. ALARO-4km captures both the
timing and magnitude of this peak with high accuracy. ALARO-12km also reproduces a peak, but it occurs slightly later (i.e.
1-2 hours) and with a higher frequency than observed. For ALARO-25km, the peak occurs even later (around 22:00 UTC) and
has a lower frequency compared to the observations. ~~Furthermore,~~

3.3 Effect of SURFEX




Figure 7. Relative error of the return-level intensities for each region (rows: Low-, Middle-, High-Belgium) and return period T (columns: 2, 10, 50 years), using all available years for each dataset (observational record lengths are given in Table C1; simulations span 1992–2022). Panels (a)–(i) are ordered row-wise. Each box plot represents the uncertainty as gauged by the bootstrap procedure with $10^4 \times 10^4 \times 10^4 (= 10^8)$ samples. The box spans the interquartile range (25th–75th percentile) around the median, and the whiskers span the width of the 95%-confidence interval around the median (2.5th–97.5th percentile).




Figure 8. Frequency of annual-maximum hourly precipitation intensities over Belgium for hourly station observations (HYDRO network) and for each simulation (25 km, 12.5 km, and 4 km resolutions), [using all available years for each dataset \(observational record lengths are given in Table C1; simulations span 1992–2022\)](#). (left) Diurnal cycle: showing the frequency of annual-maximum hourly precipitation intensities for each hour of the day (UTC). (right) Seasonal cycle: showing the frequency of annual-maximum hourly precipitation intensities for each season (winter, spring, summer, autumn).

Figure 9 shows the annual-mean difference between ALARO-12km and ALARO-12km-noSFX, which isolates the effect of the SURFEX land-surface scheme, as all other components are identical between the two simulations. Both differences are significant across the entire domain (Fig. 9a, b). The temperature difference is roughly constant across the domain, except for some urban areas (e.g., Brussels) where the warming is stronger. On the other hand, the precipitation difference exhibits a clear north-south gradient, with the southern part of Belgium experiencing a stronger reduction in precipitation than the northern part. Moreover, the diurnal cycle of precipitation over Belgium (Fig. 9c) isolates how SURFEX acts on the sub-daily distribution. Both configurations share the same single afternoon peak at 16–17 UTC. Introducing SURFEX leaves the timing and shape of the cycle essentially unchanged except for a reduction in amplitude, damping the afternoon peak more strongly than the overnight minimum. Therefore, ~~the observational data reveal a secondary peak in the early morning, which is absent in all simulations. Although resolution-dependent shift in peak timing seen in Fig. 5 is not a consequence of the land-surface scheme.~~

To identify the mechanism, we examine the surface energy balance (Fig. 10), showing the seasonal diurnal cycle of the latent (a–d) and sensible (e–h) heat fluxes. The simulations separate by land-surface scheme rather than resolution: ALARO-12km-noSFX closely tracks ALARO-25km ~~appears to align with the observed morning peak, this is primarily due to elevated nighttime frequencies in the simulation rather than a distinct early morning peak~~, while the SURFEX simulations (ALARO-12km and ALARO-4km) cluster together. Introducing SURFEX reduces the latent heat flux and increases the sensible heat flux, in both




Figure 9. Effect of introducing SURFEX at fixed 12.5km resolution, from the annual-mean difference between ALARO-12km and ALARO-12km-noSFX over Belgium, 1992–2022. (a) Near-surface air temperature difference and (b) relative precipitation difference, with domain-averaged values reported above each panel. Hatching in (a) and (b) indicates grid points where the difference is not significantly different from zero at the 95 % confidence level, based on a paired bootstrap resampling of years. The difference is significant across the entire domain, so no hatching is visible. (c) Domain-mean diurnal cycle of precipitation (all hours) for the two configurations; dashed lines denote the daily-mean value of each.

cases most strongly in summer, and slightly advances the diurnal peak of each. The net surface radiation is essentially identical across all simulations (Fig. D4), so the available energy at the surface is unchanged. The sum of the latent and sensible heat fluxes is correspondingly near-identical in daily-mean magnitude, but its diurnal cycle is advanced in the SURFEX simulation. This phase shift, together with the shift toward sensible heating, points to a reduced surface thermal inertia, consistent with drier soil responding more rapidly to daytime forcing. This is consistent with the flux differences being driven by the surface state, as the net surface radiation, which sets the total available energy, is essentially unchanged.

As a whole, it is shown that the introduction of SURFEX reduces the latent heat flux and increases the sensible heat flux, especially in summer. This can be partly attributed to the inclusion of urban elements, which contain no moisture and therefore reduce the latent heat flux. However, the effect is also present in non-urban areas, indicating that the difference is not solely due to urbanisation. Although the differing soil discretisations of the two schemes preclude a direct quantitative comparison of soil moisture, the reduced latent heat flux, drier summers, and associated warming are all consistent with reduced soil moisture in the SURFEX configuration. This shift in the partitioning of the surface fluxes is consistent with the observed warming of near-surface air temperature and the reduced precipitation.




Figure 10. Seasonal diurnal cycle of the surface upward latent (a–d) and sensible (e–h) heat fluxes over Belgium for all simulations, 1992–2022. Columns are the four seasons (DJF, MAM, JJA, SON). Dashed lines denote the daily-mean value of each simulation.

4 Discussion

4.1 Validation of average climatology

A clear distinction in climatology can be drawn between ALARO-25km on the one hand, and ALARO-12km and ALARO-4km on the other hand, both for near-surface air temperature and accumulated precipitation. The cold bias from ALARO-25km of -1.14°C over Belgium is significantly improved by about 1°C in the two simulations with the highest resolutions, as shown in Fig. 2. Furthermore, these simulations both exhibit a reduced wet bias in comparison with ALARO-25km. The difference between ALARO-12km and ALARO-4km is mostly small when compared to ALARO-25km. Aside from the difference in resolution, ALARO-12km employs the land-surface model SURFEX, while ALARO-25km still uses the built-in module ISBA. ~~This difference in configuration might explain the~~ The dedicated ALARO-12km-noSFX experiment (Sect. 3.3), which holds resolution fixed, shows that this difference in land-surface scheme, rather than the resolution increase, is responsible for the clear distinction between these two sets of simulations.

In Giot et al. (2016) 30-year simulations over Europe were performed with ALARO-0, a previous version of the ALARO model, and the built-in land surface scheme ISBA. These simulations were performed at resolutions of 0.44° ($\sim 50\text{ km}$) and 0.11° ($\sim 12\text{ km}$) and contained an area-averaged yearly bias in near-surface air temperature of a little less than -1°C over the Middle-European sub-region (which covers Belgium, the Netherlands, most of Germany, and parts of neighbouring countries).

This result is very similar to the yearly bias of ALARO-25km in this work. The seasonal temperature biases of Giot et al. (2016) over Middle-Europe are also in line with the results of ALARO-25km.

Furthermore, in Berckmans et al. (2017), simulations with ALARO-0 for a 10-year period were carried out in three different configurations, one of which consisted of continuous simulations as performed in this work. It should be noted that, contrary to those of Giot et al. (2016), these simulations were performed with the SURFEX model, albeit a previous version, SURFEXv5. Over Middle-Europe, an average yearly temperature bias of -1.6°C was found, while for winter and summer a bias of -1.3°C was presented. This is comparable to the results of Giot et al. (2016) and of ALARO-25km, even though the latter simulations did not use SURFEX.

Lastly, 4 km ALARO-0 simulations without SURFEX were performed in the context of the CORDEX.be I project (Termonia et al., 2018b). In the final report (Termonia et al., 2018c) a cold bias of 1.3°C in the average yearly temperature over Belgium was presented. This value is in line with the results from Giot et al. (2016), Berckmans et al. (2017), and ALARO-25km, but contrasts with the temperature bias of the here-presented ALARO-12km and ALARO-4km, even though these CORDEX.be I simulations were run at 4 km. This ~~suggests~~ is consistent with our ALARO-12km-noSFX experiment (Sect. 3.3), which directly demonstrates that the use of SURFEX, and not the resolution increase, is responsible for the improved temperature bias, through a shift in the partitioning of the surface heat fluxes.

Evaluation against the CG dataset reveals that all ALARO-1 simulations exhibit a consistent annual wet bias over Belgium, with this bias most pronounced in ALARO-25km. Seasonal analyses confirm a wet bias across all seasons, except for a dry bias in summer in both ALARO-12km and ALARO-4km. Differences between ALARO-12km and ALARO-4km remain insubstantial across seasons. ~~These patterns indicate~~ The ALARO-12km-noSFX experiment (Sect. 3.3) confirms that the inclusion of SURFEX ~~exerts a greater,~~ rather than the change in spatial resolution, exerts the dominant influence on average precipitation ~~than changes in spatial resolution.~~

Our results for ~~rainfall~~ precipitation (Fig. 4) are now compared to previous continuous simulations from Giot et al. (2016) (resolutions of 50 and 12.5 km). The annual ~~bias from~~ precipitation bias of ALARO-25km is slightly (33.7%) is markedly larger than the ~~bias~~ annual bias of approximately 20% over Middle-Europe (ME) ~~from Giot et al. (2016), while for reported~~ by Giot et al. (2016) (for both their 50 and 12.5 km simulations), whereas ALARO-12km (15.3%) and ALARO-4km ~~it is~~ (17.0%) are somewhat smaller. On the other hand, the annual precipitation bias over ME in Berckmans et al. (2017) (at 20 km resolution) was 15%, which agrees almost perfectly with the results from ALARO-12km and ALARO-4km. For summer and winter precipitation, the biases of Berckmans et al. (2017) and Giot et al. (2016) are similar to each other and resemble closely those of ALARO-25km. Conversely, the winter, and particularly the summer precipitation totals are much drier in ALARO-12km and ALARO-4km compared to ALARO-25km. The large wet bias in spring, reported in Giot et al. (2016) is also present (and slightly increased) in the simulations in this work. Finally, the large bias in autumn was absent in the simulations of Giot et al. (2016), where the bias was between 0 and 10%.

Previous 4 km simulations showed results similar to ALARO-4km. Compared to the HRES-simulations from CORDEX.be I, the bias in winter precipitation was of comparable magnitude (Termonia et al., 2018c). Biases from De Troch et al. (2013) were slightly smaller for summer precipitation than for ALARO-4km: their 4 km simulation showed a bias of only 3%. Notably,

those simulations employed daily restarts, which could have a significant impact—especially in summer, when soil moisture
450 feedbacks tend to play a more important role. There was a difference of roughly 10 percentage points between their 10 km
and 4 km simulations in summer precipitation bias, which is also present between ALARO-12km and ALARO-4km but with
an opposite sign. Both cases can be interpreted as an improvement in the representation of (convective) precipitation when
increasing the resolution.

In De Troch (2016), the multi-scale behaviour of ALARO-0 was investigated by means of the diurnal cycle of precipitation
455 in summer and winter. An identical distinction between all hours and wet hours is made in this work, as shown in Fig. 5. It was
shown in De Troch (2016) that in summer the timing and amplitude of the diurnal peak were improved for simulations with
ALARO-0 at 10 and 4 km compared to a 40 km simulation, when considering all hours. However, the timing of the afternoon
peak as reproduced by the model was still too early. A similar behaviour is evident in this work, although ALARO-4km
captures the timing better (later) than in De Troch (2016). For winter precipitation, the 4 km simulation from De Troch (2016)
460 exhibited a weak peak in the early afternoon, which is absent in the observations. Our ~~ALARO-4m simulations reproduce~~
ALARO-4km simulation reproduces this constant diurnal pattern better than those in ~~De Troch (2016).~~ De Troch (2016). Since
winter precipitation over Belgium is predominantly stratiform, the spurious afternoon peak in their ALARO-0 simulation points
to convective triggering when little is expected; its near-absence in ALARO-1 suggests less spurious wintertime triggering,
though isolating the responsible change is left for future research.

465 When considering only wet hours (i.e. hours with a precipitation rate > 0.1 mm), the observations show no diurnal variation
(regardless of some noise) in winter ~~-(Fig. 5d).~~ All simulations in this paper reproduce this pattern, with the mean precipitation
increasing towards the observed value as the resolution increases. For wet hours in summer (Fig. 5c), all simulations contain an
afternoon/evening peak, with increasing amplitude for higher resolutions, similar to the study of De Troch (2016). The timing of
the peak does not significantly differ between ALARO-25km and ALARO-12km, and is slightly earlier for ALARO-4km. How-
470 ever, for all simulations, the ~~diurnal peak is summer wet-hour peak occurs~~ later than the ~~observational peak, which is observed~~
peak, in contrast with ~~results from De Troch (2016).~~ the well-captured peak timing reported by De Troch (2016) for ALARO-0.
This contrast reflects the change from ALARO-0 to ALARO-1 and cannot be attributed to the scale-aware 3MT scheme, which
is common to both versions, nor to the land-surface scheme, which the ALARO-12km-noSFX experiment shows does not affect
the peak timing (Sect. 3.3, Fig. 9c). A further candidate is the difference in experimental setup: the ALARO-0 simulations of
475 De Troch (2016) were reinitialised daily, whereas ours run continuously. Such daily restarts can plausibly influence the phase
of the simulated diurnal cycle. However, we cannot isolate the responsible factor from the available output. Overall, for both
all-hour (Fig. 5a) and wet-hour (Fig. 5b) precipitation, the diurnal cycles in ALARO-4km are comparable to those in the work
of De Troch (2016).

In our analysis of the mean diurnal precipitation cycle, notable differences emerge when comparing simulations at varying
480 resolutions with observational data, as shown in Fig. 5. Specifically, while the “all-hour” precipitation cycle ~~tends to exhibit~~
~~an overestimation overestimates precipitation~~ relative to observations, the “wet-hour” cycle ~~shows a general underestimation.~~
~~This implies that the precipitation frequency is overestimated. This discrepancy underestimates it, which Table 2 attributes~~
to an overestimated wet-hour frequency combined with an underestimated wet-hour intensity. This is most pronounced in ~~the~~

ALARO-25km ~~simulation and is minimized in the~~ and minimised in ALARO-4km~~simulation, indicating, showing~~ that the
485 coarser resolution ~~more significantly~~ overestimates precipitation amounts ~~due to through~~ an overly high ~~rainfall frequency~~.
~~These results imply that drizzle contributes substantially to the simulated rainfall, particularly in the lower-resolution model.~~
~~This problem is partially mitigated when using CPMs, but not entirely.~~ precipitation frequency, i.e. a substantial drizzle
contribution.

This drizzle bias is a well-known deficiency of regional climate models (Fosser et al., 2015; Berthou et al., 2020). It arises
490 largely because parameterised convection triggers on weak instability and releases it gradually, producing precipitation that
is too frequent and too light. The improvement with resolution reflects the sharper humidity and updraught fields at higher
resolution, which reduce the widespread simultaneous triggering dominating the coarser runs. The residual bias in ALARO-
4km still exhibits a drizzle bias, even at scales where convection is (partially) permitted by the scale-aware convection
scheme 3MT. This finding is consistent with previous works (Kendon et al., 2012; Fosser et al., 2015; Berthou et al., 2020).
495 ~~Additionally, it has been shown that 3MT still parameterising part of the convective spectrum in the grey zone.~~

Furthermore, both the timing and amplitude of the diurnal precipitation peak are best represented at high resolution. ~~This~~
~~result is,~~ as well established for CPMs, ~~as documented by numerous studies referenced in the review of Lucas-Picher et al. (2021)~~
(Lucas-Picher et al., 2021). Most RCMs ~~tend to~~ exhibit a premature diurnal peak, primarily due to the limited memory of at-
mospheric instabilities. ~~However,~~ this behaviour is notably absent ~~in this study.~~ ~~At lower resolutions, the diurnal peak in~~
500 ~~ALARO here, and at lower resolutions the ALARO peak~~ even occurs later than at higher resolutions. ~~This difference may~~
~~be attributable to the scale-aware character of the 3MT scheme.~~ The ALARO-12km-noSFX experiment (Sect. 3.3, Fig. 9c)
shows that introducing SURFEX reduces the amplitude of the diurnal cycle but leaves the timing of the peak unchanged, so
the resolution-dependent shift in peak timing is not driven by the land-surface scheme. Its origin lies instead in the convection
scheme. Conventional mass-flux schemes that close on CAPE remove instability almost as soon as it is generated by surface
505 heating, so that simulated precipitation peaks around local noon, in phase with the surface fluxes, rather than in the late
afternoon as observed (Guichard et al., 2004; Bechtold et al., 2004). In contrast, 3MT employs a prognostic closure that retains
memory of the updraught mesh fraction and updraught velocity between time steps, with the mesh fraction evolving through
the moisture budget. Instability is therefore released gradually over several time steps rather than instantaneously, shifting the
peak away from noon; none of our configurations peaks near midday. At coarser resolutions this gradual release appears to be
510 too gradual, delaying the convective peak, and the timing sharpens as explicit convection progressively takes over at higher
resolutions. A full attribution of the responsible mechanism would require dedicated sensitivity experiments and is left to future
work.

4.2 Extreme precipitation

IDF ~~curves~~ statistics are presented in Fig. 6 for durations of 1 to 72 hours across three distinct topographic regions in Belgium.
515 The regional differences in return levels, with smaller sub-daily (< 6h) values in Low-Belgium and higher (multi-)daily (≥
24h) values in High-Belgium, are consistent with the findings of Van de Vyver (2013). In each case, the return levels scale
linearly with duration on a log-log scale, for both observations and simulations. This corresponds to a power-law relationship

on a linear scale and reflects the underlying scale invariance and multifractal characteristics of extreme precipitation. These phenomena can be effectively captured using simple scaling techniques, which in turn result in a power-law relationship (Gupta and Waymire, 1990; Veneziano and Furcolo, 2002). However, for short ~~rainfall~~-precipitation durations (1–3 hours), simulated return levels deviate from this linear trend, curving downward, whereas observational return levels do not exhibit this pattern. This discrepancy arises entirely from model error, as precipitation values from both observations and simulations were first aggregated to an hourly resolution before extracting the annual maxima. As shown in Fig. D3b, the smaller R^2 -values indicate a less optimal fit for larger spatial scales. The higher (less negative) linear coefficients at these scales may also reflect this underestimation at short durations. However, previous studies have also shown that annual-maximum precipitation at larger spatial scales has a higher linear coefficient with respect to duration than at smaller scales (Innocenti et al., 2019). Next, these linear coefficients can be compared to those from a simple-scaling model by Van De Vyver (2018). In that work, station-specific values in the range of -0.75 and -0.64 were found for 16 stations over Belgium. The observational coefficients depicted in Fig. D3a correspond to this range, although they are slightly more negative.

At durations of 1 to 3 hours, lower-resolution simulations increasingly underestimate return levels compared to observational values, highlighting the added value of the convection-permitting ALARO-4km simulation. A similar result was found by Tabari et al. (2016), who also constructed IDF ~~curves~~-statistics for ALARO-0 simulations at different resolutions. In that study, the 4 km and 10 km ALARO-0 simulations accurately captured sub-daily intensities, with a slight underestimation, while ALARO-40km significantly underestimated sub-daily precipitation amounts. For longer durations (>24 hours h), ALARO-25km tends to overestimate the return levels, particularly for short return periods ~~-(Fig. 7a-c)~~. De Troch (2016) reported a similar finding, as all ALARO-0 simulations (4 km, 10 km, and 40 km) overestimated the 2-year return level for daily precipitation. Additionally, while De Troch (2016) observed increasing overestimation for longer return periods in the 4 km ALARO-0 simulation, this effect is notably absent - and even reversed - in ALARO-4km.

~~Because of the~~ The scale-aware design of the 3MT scheme ~~, the statistics of extreme precipitation should, in principle, be independent of model resolution. It~~ provides a smooth transition between parameterised and resolved convection, which would lead one to expect a gradual change of the extreme-precipitation statistics with resolution, as found in previous studies (De Troch, 2016; Tabari et al., 2016), ~~return levels exhibited a gradual change with resolution, which is broadly consistent with the expected resolution independence.~~ In contrast, the return levels presented in this study display a more abrupt, step-wise change when transitioning from 12 km to 4 km resolution. This behaviour is more similar to that observed in other models without a scale-aware convective scheme, where the step-wise shift is typically attributed to the deactivation of the deep convective ~~parameterization~~-parameterisation at high resolutions. For 3MT, however, no such deactivation occurs, and the apparent abruptness may be partly an artefact of comparison: the 25 km to 12 km change is small, so the 12 km to 4 km change stands out as step-wise by contrast. We propose two contributing effects. First, the difference between ALARO-25km and ALARO-12km conflates resolution with the land-surface scheme: the drying effect of SURFEX (Sect. 3.3) reduces precipitation and may partly offset the resolution-driven increase in extreme intensity at 12 km, suppressing the apparent 25 km to 12 km change. Second, the step from 12 km to 4 km crosses the grey zone of convection (~ 4 –10 km) (Prein et al., 2015; Vergara-Temprado et al., 2020), where 3MT transitions from a parameterised towards a resolved representation of deep convection. This transition is expected

to affect the extreme-precipitation statistics most strongly. Isolating the contributions of these two effects is left to future work.

555 The diurnal cycle of hourly AM precipitation frequencies is well captured by ALARO-4km (see Fig. 8a). Both the frequency peak amplitude and timing are in agreement between ALARO-4km and the observational data. The peak occurs later in ALARO-12km and is further delayed in ALARO-25km, while the peak amplitude increases in ALARO-12km but decreases in ALARO-25km. This peak delay is also present in 0.11° EURO-CORDEX simulations (Meredith et al., 2021). It should be noted that the relatively large diurnal variation in extreme ~~rainfall~~ precipitation frequency of ALARO-25km is quite remarkable, as most models exhibit almost no diurnal variation at similar resolutions (Meredith et al., 2021; Thomassen et al., 2023). The diurnal cycle of extremes from previous ALARO-0 simulations at 12 km and 4 km was shown in Van De Vyver et al. (2021). However, they did not find a large difference in the diurnal cycle between both resolutions, in contrast to the simulations of this work. As with the peak-timing differences discussed above, this stronger resolution sensitivity of ALARO-1 relative to ALARO-0 most plausibly reflects the broader physics update between the two versions rather than the scale-aware convection scheme common to both, and is left to future work.

565 Similarly to the diurnal cycle, the seasonal cycle of hourly extreme-precipitation frequency is well captured by our three simulations, as shown in Fig. 8b. The frequency peak in summer due to the convective nature of hourly extreme precipitation is apparent. This result corresponds to previous simulations from Van De Vyver et al. (2021), who also did not find a difference in the seasonal cycle between 12 km and 4 km ALARO-0 simulations. In contrast, Innocenti et al. (2019) reported the slight outperformance of a 12 km RCM simulation compared to a 4 km convection-permitting WRF simulation in capturing the seasonal cycle of hourly AM occurrences over North America.

5 Conclusions

This study evaluated the performance of the ALARO-1 model, including a scale-aware deep convection ~~parameterization~~ parameterisation, over Belgium using a multi-level dynamical downscaling framework at three horizontal resolutions: 25 km, 12.5 km, and 4 km. The ALARO-1 model was coupled online to the land-surface model SURFEX v8.0 for ALARO-12km and ALARO-4km. These simulations, spanning a 32-year period (~~1991–2022~~ 1991–2022), were compared against both a gridded observational dataset (CLIMATE-GRID) and hourly precipitation values from station measurements.

580 The analysis of average climatology demonstrated that across all resolutions the ALARO-1 model exhibits an annual cold bias between -1.14 °C and -0.25 °C and an annual wet bias between 15.3 % and 33.7 % over Belgium. Comparing resolutions, ALARO-12km and ALARO-4km outperformed ALARO-25km, which exhibited larger biases in both temperature and precipitation. While ALARO-12km and ALARO-4km provided similar results in terms of average climatology, ALARO-4km captured the diurnal cycle of precipitation better. A dedicated sensitivity experiment (ALARO-12km-noSFX) showed that this improvement relative to ALARO-25km is not attributable to the increase in resolution alone: the coupling to SURFEX accounts for a substantial part of the reduction in the temperature and precipitation biases.

585 A key focus of this study was the representation of extreme precipitation, evaluated through Intensity-Duration-Frequency
~~curves~~ statistics derived from annual maximum precipitation. While all simulations followed the expected log-log linear be-
 haviour between return-level precipitation intensity and duration, lower-resolution models underestimated return levels at sub-
 daily timescales. ALARO-4km significantly improved upon this, producing more realistic hourly return levels and better cap-
 turing the diurnal timing of hourly extreme precipitation. These findings provide clear ~~experimental~~ evidence that increasing
 590 model resolution enhances the simulation of extreme precipitation.

Despite these improvements, this study has some limitations. The research domain was limited to Belgium due to compu-
 tational constraints. Another limitation is the absence of simulations at even higher resolutions, which could further improve
 model accuracy but are currently also constrained by computational demands.

Future research should address these challenges by conducting simulations at resolutions higher than 4 km and applying
 595 the ALARO-1 model to regional downscaling of global climate models (GCMs) and climate projections. Additionally, a more
 detailed investigation of the urban effect on temperature and precipitation, as represented by the TEB model in SURFEX, is
 warranted. Expanding the application of the ALARO-1 model to other regions would further validate its performance. Finally,
 given the high computational cost of convection-permitting simulations, future work should explore alternative, more targeted
 downscaling approaches to reduce computational demands while maintaining or improving model accuracy, specifically for
 600 extreme precipitation.

Appendix A: Definition of bias

The temperature and precipitation bias are calculated in the following way. Let X_{kni} be the value of variable X at time k , with
 $k = 1, \dots, K$, in grid point n , with $n = 1, \dots, N$, and in year i , with ~~$i = 1, \dots, I$~~ $i = 1, \dots, I$. We define the total value of X over a
 period t as:

$$605 \quad X_{ni}^t = \sum_{k \in t} X_{kni}$$

The average value over a period t is defined as:

$$\bar{X}_{ni}^t = \frac{1}{K_t} \sum_{k \in t} X_{kni}$$

with K_t the number of values in period t : $K_t = \sum_{k \in t} 1$

In this work, we consider three types of periods t :

- (1) months, denoted as $m \in M$ with $M = \{\text{Jan, Feb, Mar, Apr, May, Jun, Jul, Aug, Sep, Oct, Nov, Dec}\}$,
- 610 (2) seasons, denoted as $s \in S$ with $S = \{\text{DJF, MAM, JJA, SON}\}$,
- (3) years, denoted as y .

The yearly total value of X is related to the seasonal values as:

$$X_{ni}^y = \sum_s X_{ni}^s$$

and the yearly averaged value:

$$\bar{X}_{ni}^y = \frac{1}{4} \sum_s \bar{X}_{ni}^s$$

615 Let $\bar{T}_{ni}^t$ now be the temperature averaged over the period t , with $t \in (M \cup S \cup \{y\})$. The temperature bias is consequently calculated by taking the difference between the model temperature $T_{ni}^{t,m}$ and the observational temperature $T_{ni}^{t,o}$ with $t \in (M \cup S \cup \{y\})$ and averaging it over all years i (with I the number of years). The bias in grid point n is hence:

$$\bar{T}_n^{x,b} = \frac{1}{I} \sum_i (\bar{T}_{ni}^{x,m} - \bar{T}_{ni}^{x,o})$$

The spatially averaged bias is then acquired by averaging over all grid points (with N the number of grid points):

620
$$\bar{T}^{t,b} = \frac{1}{N} \sum_n \bar{T}_n^{t,b}$$

The seasonal and annual values can be straightforwardly related to each other:

$$\bar{T}^{y,b} = \frac{1}{4} \sum_s \bar{T}^{s,b}$$

For precipitation, we consider the relative bias. Let P_{ni}^t now be the total precipitation over the period t with $t \in (M \cup S \cup \{y\})$. The relative bias over this period is then calculated as:

625
$$P_n^{t,r} = \frac{\sum_i (P_{ni}^{t,m} - P_{ni}^{t,o})}{\sum_i P_{ni}^{t,o}}$$

The spatially averaged value then becomes:

$$P^{t,r} = \frac{1}{N} \sum_n P_n^{t,r}$$

For the relative bias, there is no straightforward relation between annual and seasonal values:

$$P^{y,r} \neq \sum_s P^{s,r}$$

630 **Appendix B: Definition of GEV distribution**

~~The~~ For $\xi \neq 0$, the cumulative GEV distribution function (~~for $\xi \neq 0$~~) $F(x)$ is given by:

$$F(x) = \exp \left\{ - \left[1 + \xi \left(\frac{x - \mu}{\sigma} \right) \right]^{-1/\xi} \right\} \quad (\text{B1})$$

The distribution with $\xi = 0$ is then interpreted as the limit of Eq. (B1) as $\xi \rightarrow 0$:

$$F(x) = \exp \left\{ - \exp \left[- \left(\frac{x - \mu}{\sigma} \right) \right] \right\} \quad (\text{B2})$$

635 GEV distribution functions are constructed for each duration separately. We denote the duration-specific GEV parameters as ξ_d , μ_d , and σ_d . Subsequently, these distribution functions are used to calculate return levels $z_d(T)$ for the precipitation intensity. A return level $z_d(T)$ (expressed in mm h⁻¹) is specified for a certain return period T and duration d . It is defined as the precipitation intensity which has a probability of $1/T$ of being exceeded in any given year: so on average once every T years. The formula for these return levels can be derived from equations (B1) and (B2):

$$640 \quad z_d(T) = \mu_d - \frac{\sigma_d}{\xi_d} \left\{ 1 - \left[-\ln \left(1 - \frac{1}{T} \right) \right]^{-\xi_d} \right\} \quad (\text{B3})$$

Appendix C: Station data specifications

Table C1. Specifications of observational networks used in this study.

Network name	Number of years (start year - end year)	Number of locations
DGH	20 (2004 – 2023)	95
VMM	11 (2013 – 2023)	36
AWS	24 (2000 – 2023)	13
HYDRO	38 (1967 – 2004)	18
IBGE	19 (1992 – 2010)	14

Appendix D: Additional figures

D1 Validation of average Average climatology




Figure D1. Bias in mean seasonal near-surface air temperature of each simulation (25 km, 12.5 km, and 4 km resolutions) with respect to the daily gridded observational CLIMATE-GRID dataset over Belgium, for the period 1992–2022. Columns: ALARO-25km (a, d, g, j), ALARO-12km (b, e, h, k), ALARO-4km (c, f, i, l). Rows: winter (DJF; a–c), spring (MAM; d–f), summer (JJA; g–i), autumn (SON; j–l). Hatching indicates grid points where the bias is not significantly different from zero at the 95 % confidence level, based on a paired bootstrap resampling of years.




Figure D2. Bias in mean seasonal precipitation of each simulation (~~25 km, 12.5 km, and 4 km resolutions~~) with respect to the daily gridded observational CLIMATE-GRID dataset over Belgium, for the period 1992–2022. Columns: ALARO-25km (a, d, g, j), ALARO-12km (b, e, h, k), ALARO-4km (c, f, i, l). Rows: winter (DJF; a–c), spring (MAM; d–f), summer (JJA; g–i), autumn (SON; j–l). Hatching indicates grid points where the bias is not significantly different from zero at the 95 % confidence level, based on a paired bootstrap resampling of years.




Figure D3. Log-log linear regression analysis of the return levels as a function of duration, performed for each dataset (observations and ALARO-simulations), topographic region, and return period separately, using all available years for each dataset (observational record lengths are given in Table C1; simulations span 1992–2022). (lefta) The linear regression coefficients. (rightb) The R^2 -values. The dashed line denotes a value of 1, which corresponds to a perfect fit.




Figure D4. Seasonal diurnal cycle over Belgium of the sum of the surface upward latent and sensible heat fluxes (a–d) and the net surface radiative flux (e–h), for all simulations, 1992–2022. Columns are the four seasons (DJF, MAM, JJA, SON). Dashed lines denote the daily-mean value of each simulation.

Code availability. The Python notebooks used for the analysis will be made publicly available upon acceptance of the manuscript. The ALARO1-SFX code, along with all its related intellectual property rights, is owned by the members of the ACCORD consortium. Each member of this consortium can license the shared code to academic institutions of their home country for noncommercial research. Access to the codes of the ACCORD system can be obtained by contacting one of the member institutes mentioned in Termonia et al. (2018a) or by submitting a request in the Contact link below the page of the ACCORD website (<https://www.umr-cnrm.fr/accord/>) and the access will be subject to signing a standardised ACCORD license agreement.

Data availability. The VMM precipitation observations over Flanders are available on <https://waterinfo.vlaanderen.be/>. The DHG precipitation observations may be requested from the Service public de Wallonie - Mobilité et Infrastructures, Direction de la Gestion hydrologique, PEREX, Rue Del’Grête, 22, 5020 NAMUR (Daussoulx). The IBGE data may be requested at the Bruxelles Environnement (<https://environnement.brussels>). The automatic weather station (AWS) precipitation observations from the Royal Meteorological Institute of Belgium are available on <https://opendata.meteo.be/download>. Annual maxima extracted from the HYDRO-network from the Royal Meteorological Institute of Belgium are available on <https://doi.org/10.5281/zenodo.4741177>. Further information about the CLIMATE-GRID dataset can be found on https://opendata.meteo.be/geonetwork/srv/eng/catalog.search#/metadata/RMI_DATASET_GRIDDEDOBS,

although the data are currently not yet publicly available. The simulation data used in this study will be made publicly available in a Zenodo repository upon acceptance of the manuscript.

Author contributions. SC and PT conceptualised the study. SC, PT, and WD acquired funding. WD conducted the investigation and performed the formal analysis. WD, KV and MVG were responsible for software implementation. HVdV, DD, RH, BVS, KVV, SC, and PT supervised the research. WD created the visualisations and prepared the original draft. All authors contributed to the review and editing of the manuscript and approved the final version.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The authors acknowledge the Flemish Supercomputer Center (VSC) for providing the computational resources and services used to perform the ALARO1-SFX regional climate simulations. The authors also thank the Vlaamse Milieumaatschappij (VMM), the Direction de la Gestion Hydrologique (DGH) of SPW Mobilité et Infrastructures, and Brussels Environment for providing precipitation observations over Flanders, Wallonia, and Brussels, respectively. The authors thank the two anonymous referees and the topic editor for their careful reading and constructive comments, which have substantially improved the manuscript. Finally, we acknowledge the use of AI-assisted tools, which were used exclusively for language refinement and text rewriting. All scientific content was developed and validated by all authors.

Financial support. This research has been supported by the Research Foundation – Flanders (FWO) (grant no. 1157523N). KV is supported by the Belgian Science Policy (BELSPO) under Contract B2/223/P1/CORDEXbeII. KVV is funded through Belspo grant FED-tWIN2021-prf024 - ARCHWAY. SC is supported by the BELSPO project FED-tWIN 2020-018_AURA.

References

- Ban, N., Schmidli, J., and Schär, C.: Evaluation of the convection-resolving regional climate modeling approach in decade-long simulations, *Journal of Geophysical Research: Atmospheres*, 119, 7889–7907, <https://doi.org/10.1002/2014JD021478>, 2014.
- Ban, N., Caillaud, C., Coppola, E., Pichelli, E., Sobolowski, S., Adinolfi, M., Ahrens, B., Alias, A., Anders, I., Bastin, S., Belušić, D.,
680 Berthou, S., Brisson, E., Cardoso, R., Chan, S., Christensen, O., Fernández, J., Fita, L., Frisius, T., and Zander, M.: The first multi-model ensemble of regional climate simulations at kilometer-scale resolution, part I: evaluation of precipitation, *Climate Dynamics*, 57, <https://doi.org/10.1007/s00382-021-05708-w>, 2021.
- Bechtold, P., Chaboureaud, J.-P., Beljaars, A., Betts, A. K., Köhler, M., Miller, M., and Redelsperger, J.-L.: The Simulation of the Diurnal Cycle of Convective Precipitation over Land in a Global Model, *Quarterly Journal of the Royal Meteorological Society*, 130, 3119–3137,
685 <https://doi.org/10.1256/qj.03.103>, 2004.
- Berckmans, J., Giot, O., De Troch, R., Hamdi, R., Ceulemans, R., and Termonia, P.: Reinitialised versus continuous regional climate simulations using ALARO-0 coupled to the land surface model SURFEXv5, *Geoscientific Model Development*, 10, 223–238, <https://doi.org/10.5194/gmd-10-223-2017>, publisher: Copernicus GmbH, 2017.
- Berg, P., Christensen, O. B., Klehmet, K., Lenderink, G., Olsson, J., Teichmann, C., and Yang, W.: Summertime precipitation extremes in a EURO-CORDEX 0.11° ensemble at an hourly resolution, *Natural Hazards and Earth System Sciences*, 19, 957–971, <https://doi.org/10.5194/nhess-19-957-2019>, publisher: Copernicus GmbH, 2019.
- Berthou, S., Kendon, E. J., Chan, S. C., Ban, N., Leutwyler, D., Schär, C., and Fosser, G.: Pan-European climate at convection-permitting scale: a model intercomparison study, *Clim Dyn*, 55, 35–59, <https://doi.org/10.1007/s00382-018-4114-6>, 2020.
- Brisson, E., Van Weverberg, K., Demuzere, M., Devis, A., Saeed, S., Stengel, M., and van Lipzig, N. P. M.: How well can a convection-permitting climate model reproduce decadal statistics of precipitation, temperature and cloud characteristics?, *Clim Dyn*, 47, 3043–3061,
695 <https://doi.org/10.1007/s00382-016-3012-z>, 2016.
- Bubnová, R., Hello, G., Bénard, P., and Geleyn, J.-F.: Integration of the Fully Elastic Equations Cast in the Hydrostatic Pressure Terrain-Following Coordinate in the Framework of the ARPEGE/Aladin NWP System, [https://doi.org/10.1175/1520-0493\(1995\)123<0515:IOTFEE>2.0.CO;2](https://doi.org/10.1175/1520-0493(1995)123<0515:IOTFEE>2.0.CO;2), section: Monthly Weather Review, 1995.
- 700 Carreau, J., Naveau, P., and Neppel, L.: Partitioning into hazard subregions for regional peaks-over-threshold modeling of heavy precipitation, *Water Resources Research*, 53, 4407–4426, <https://doi.org/10.1002/2017WR020758>, 2017.
- Charnock, H.: Wind stress on a water surface, *Quarterly Journal of the Royal Meteorological Society*, 81, 639–640, <https://doi.org/10.1002/qj.49708135027>, 1955.
- Clapp, R. B. and Hornberger, G. M.: Empirical Equations for Some Soil Hydraulic Properties, *Water Resources Research*, 14, 601–604,
705 <https://doi.org/10.1029/WR014i004p00601>, 1978.
- Coles, S.: An Introduction to Statistical Modeling of Extreme Values, Springer Series in Statistics, Springer, London, ISBN 978-1-84996-874-4 978-1-4471-3675-0, <https://doi.org/10.1007/978-1-4471-3675-0>, 2001.
- Coppola, E., Sobolowski, S., Pichelli, E., Raffaele, F., Ahrens, B., Anders, I., Ban, N., Bastin, S., Belda, M., Belusic, D., Caldas-Alvarez, A., Cardoso, R. M., Davolio, S., Dobler, A., Fernandez, J., Fita, L., Fumiere, Q., Giorgi, F., Goergen, K., Güttler, I., Halenka, T., Heinzeller, D., Hodnebrog, o., Jacob, D., Kartsios, S., Katragkou, E., Kendon, E., Khodayar, S., Kunstmann, H., Knist, S., Lavín-Gullón, A., Lind,
710 P., Lorenz, T., Maraun, D., Marelle, L., van Meijgaard, E., Milovac, J., Myhre, G., Panitz, H.-J., Piazza, M., Raffa, M., Raub, T., Rockel, B., Schär, C., Sieck, K., Soares, P. M. M., Somot, S., Srncic, L., Stocchi, P., Tölle, M. H., Truhetz, H., Vautard, R., de Vries, H., and

- Warrach-Sagi, K.: A first-of-its-kind multi-model convection permitting ensemble for investigating convective phenomena over Europe and the Mediterranean, *Clim Dyn*, 55, 3–34, <https://doi.org/10.1007/s00382-018-4521-8>, 2020.
- 715 CORDEX, W.: CMIP6 Downscaling Plans, https://wcrp-cordex.github.io/simulation-status/CMIP6_downscaling_plans.html, accessed: 2025-02-18.
- Davies, H. C.: A lateral boundary formulation for multi-level prediction models, *Quarterly Journal of the Royal Meteorological Society*, 102, 405–418, <https://doi.org/10.1002/qj.49710243210>, 1976.
- De Troch, R.: The application of the ALARO-0 model for regional climate modeling in Belgium : extreme precipitation and unfavorable
720 conditions for the dispersion of air pollutants under present and future climate conditions, dissertation, Ghent University, <http://hdl.handle.net/1854/LU-7247081>, 2016.
- De Troch, R., Hamdi, R., Van De Vyver, H., Geleyn, J.-F., and Termonia, P.: Multiscale Performance of the ALARO-0 Model for Simulating Extreme Summer Precipitation Climatology in Belgium, *Journal of Climate*, 26, 8895–8915, <https://doi.org/10.1175/JCLI-D-12-00844.1>, 2013.
- 725 Dehem, D., Tricot, C., Wylleman, P., and Hamdi, R.: Validation des données du réseau pluviométrique géré par l’IBGE, répartition des précipitations, analyse qualitative des sites de mesure et projet de micro-climatologie en Région bruxelloise, Tech. rep., Institut royal météorologique de Belgique, Bruxelles, Belgique, 2010.
- Doblas-Reyes, F., Sörensson, A., Almazroui, M., Dosio, A., Gutowski, W., Haarsma, R., Hamdi, R., Hewitson, B., Kwon, W.-T., Lamptey, B., Maraun, D., Stephenson, T., Takayabu, I., Terray, L., Turner, A., and Zuo, Z.: Linking Global to Regional Climate Change, p. 1363–1512,
730 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.012>, 2021.
- Douville, H., Royer, J. F., and Mahfouf, J. F.: A New Snow Parameterization for the Météo-France Climate Model, *Climate Dynamics*, 12, 21–35, <https://doi.org/10.1007/BF00208760>, 1995.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958,
735 <https://doi.org/10.5194/gmd-9-1937-2016>, publisher: Copernicus GmbH, 2016.
- Fosser, G., Khodayar, S., and Berg, P.: Benefit of convection permitting climate model simulations in the representation of convective precipitation, *Clim Dyn*, 44, 45–60, <https://doi.org/10.1007/s00382-014-2242-1>, 2015.
- Fosser, G., Gaetani, M., Kendon, E. J., Adinolfi, M., Ban, N., Belušić, D., Caillaud, C., Careto, J. A. M., Coppola, E., Demory, M.-E., de Vries, H., Dobler, A., Feldmann, H., Goergen, K., Lenderink, G., Pichelli, E., Schär, C., Soares, P. M. M., Somot, S., and
740 Tölle, M. H.: Convection-permitting climate models offer more certain extreme rainfall projections, *npj Clim Atmos Sci*, 7, 1–10, <https://doi.org/10.1038/s41612-024-00600-w>, publisher: Nature Publishing Group, 2024.
- Fuhrer, O., Chadha, T., Hoefler, T., Kwasniewski, G., Lapillonne, X., Leutwyler, D., Lüthi, D., Osuna, C., Schär, C., Schulthess, T. C., and Vogt, H.: Near-global climate simulation at 1 km resolution: establishing a performance baseline on 4888 GPUs with COSMO 5.0, *Geoscientific Model Development*, 11, 1665–1681, <https://doi.org/10.5194/gmd-11-1665-2018>, publisher: Copernicus
745 GmbH, 2018.
- Geleyn, J., Mašek, J., Brožková, R., Kuma, P., Degrauwe, D., Hello, G., and Pristov, N.: Single interval longwave radiation scheme based on the net exchanged rate decomposition with bracketing, *Quart J Royal Meteor Soc*, 143, 1313–1335, <https://doi.org/10.1002/qj.3006>, 2017.
- Gerard, L.: An integrated package for subgrid convection, clouds and precipitation compatible with meso-gamma scales, *Quarterly Journal*
750 *of the Royal Meteorological Society*, 133, 711–730, <https://doi.org/10.1002/qj.58>, 2007.

- Gerard, L. and Geleyn, J.-F.: Evolution of a subgrid deep convection parametrization in a limited-area model with increasing resolution, *Quarterly Journal of the Royal Meteorological Society*, 131, 2293–2312, <https://doi.org/10.1256/qj.04.72>, 2005.
- Gerard, L., Piriou, J.-M., Brožková, R., Geleyn, J.-F., and Banciu, D.: Cloud and Precipitation Parameterization in a Meso-Gamma-Scale Operational Weather Prediction Model, <https://doi.org/10.1175/2009MWR2750.1>, section: Monthly Weather Review, 2009.
- 755 Giorgi, F.: Thirty Years of Regional Climate Modeling: Where Are We and Where Are We Going next?, *Journal of Geophysical Research: Atmospheres*, 124, 5696–5723, <https://doi.org/10.1029/2018JD030094>, 2019.
- Giot, O., Termonia, P., Degrauwe, D., De Troch, R., Caluwaerts, S., Smet, G., Berckmans, J., Deckmyn, A., De Cruz, L., De Meutter, P., Duerinckx, A., Gerard, L., Hamdi, R., Van Den Bergh, J., Van Genderachter, M., and Van Schaeybroeck, B.: Validation of the ALARO-0 model within the EURO-CORDEX framework, *Geosci. Model Dev.*, 9, 1143–1152, <https://doi.org/10.5194/gmd-9-1143-2016>, 2016.
- 760 Grell, G. A. and Freitas, S. R.: A Scale and Aerosol Aware Stochastic Convective Parameterization for Weather and Air Quality Modeling, *Atmospheric Chemistry and Physics*, 14, 5233–5250, <https://doi.org/10.5194/acp-14-5233-2014>, 2014.
- Gross, M., Wan, H., Rasch, P. J., Caldwell, P. M., Williamson, D. L., Klocke, D., Jablonowski, C., Thatcher, D. R., Wood, N., Cullen, M., Beare, B., Willett, M., Lemarié, F., Blayo, E., Malardel, S., Termonia, P., Gassmann, A., Lauritzen, P. H., Johansen, H., Zarzycki, C. M., Sakaguchi, K., and Leung, R.: Physics–Dynamics Coupling in Weather, Climate, and Earth System Models: Challenges and Recent
765 Progress, *Monthly Weather Review*, 146, 3505–3544, <https://doi.org/10.1175/MWR-D-17-0345.1>, 2018.
- Guerreiro, S. B., Blenkinsop, S., Lewis, E., Pritchard, D., Green, A., and Fowler, H. J.: Unravelling the Complex Interplay Between Daily and Sub-Daily Rainfall Extremes in Different Climates, *Weather and Climate Extremes*, p. 100735, <https://doi.org/10.1016/j.wace.2024.100735>, 2024.
- Guichard, F., Petch, J. C., Redelsperger, J.-L., Bechtold, P., Chaboureaud, J.-P., Cheinet, S., Grabowski, W., Grenier, H., Jones, C. G., Köhler, M., Piriou, J.-M., Tailleux, R., and Tomasini, M.: Modelling the Diurnal Cycle of Deep Precipitating Convection over Land with Cloud-Resolving Models and Single-Column Models, *Quarterly Journal of the Royal Meteorological Society*, 130, 3139–3172, <https://doi.org/10.1256/qj.03.145>, 2004.
- 770 Gupta, V. K. and Waymire, E.: Multiscaling properties of spatial rainfall and river flow distributions, *Journal of Geophysical Research: Atmospheres*, 95, 1999–2009, <https://doi.org/10.1029/JD095iD03p01999>, 1990.
- Ha, M. T., Bastin, S., Drobinski, P., Fita, L., Polcher, J., Bock, O., Chiriaco, M., Belušić, D., Caillaud, C., Dobler, A., Fernandez, J., Goergen, K., Hodnebrog, o., Kartsios, S., Katragkou, E., Lavin-Gullon, A., Lorenz, T., Milovac, J., Panitz, H.-J., Sobolowski, S., Truhetz, H., Warrach-Sagi, K., and Wulfmeyer, V.: Precipitation frequency in Med-CORDEX and EURO-CORDEX ensembles from 0.44° to convection-permitting resolution: impact of model resolution and convection representation, *Clim Dyn*, 62, 4515–4540, <https://doi.org/10.1007/s00382-022-06594-6>, 2024.
- 780 Hamdi, R., Degrauwe, D., Duerinckx, A., Cedilnik, J., Costa, V., Dalkilic, T., Essaouini, K., Jerczynski, M., Kocaman, F., Kullmann, L., Mahfouf, J.-F., Meier, F., Sassi, M., Schneider, S., Váňa, F., and Termonia, P.: Evaluating the performance of SURFEXv5 as a new land surface scheme for the ALADINcy36 and ALARO-0 models, *Geoscientific Model Development*, 7, 23–39, <https://doi.org/10.5194/gmd-7-23-2014>, publisher: Copernicus GmbH, 2014.
- Han, J., Wang, W., Kwon, Y. C., Hong, S.-Y., Tallapragada, V., and Yang, F.: Updates in the NCEP GFS Cumulus Convection Schemes with
785 Scale and Aerosol Awareness, *Weather and Forecasting*, 32, 2005–2017, <https://doi.org/10.1175/WAF-D-17-0046.1>, 2017.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J.,

- Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Vil-
790 laume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049,
<https://doi.org/10.1002/qj.3803>, 2020.
- Hohenegger, C., Brockhaus, P., and Schär, C.: Towards climate simulations at cloud-resolving scales, *metz*, 17, 383–394,
<https://doi.org/10.1127/0941-2948/2008/0303>, 2008.
- Hosking, J. R. M.: L-Moments: Analysis and Estimation of Distributions Using Linear Combinations of Order Statistics, *Journal of the Royal*
795 *Statistical Society. Series B (Methodological)*, 52, 105–124, <https://www.jstor.org/stable/2345653>, publisher: [Royal Statistical Society,
Oxford University Press], 1990.
- Hosking, J. R. M. and Wallis, J. R.: Some statistics useful in regional frequency analysis, *Water Resources Research*, 29, 271–281,
<https://doi.org/10.1029/92WR01980>, 1993.
- Innocenti, S., Mailhot, A., Frigon, A., Cannon, A. J., and Leduc, M.: Observed and Simulated Precipitation over Northeastern North America:
800 How Do Daily and Subdaily Extremes Scale in Space and Time?, *Journal of Climate*, 32, 8563–8582, <https://doi.org/10.1175/JCLI-D-19-0021.1>, 2019.
- IPCC: Annex II: Models [Gutiérrez, J M., A.-M. Tréguier (eds.)], p. 2087–2138, Cambridge University Press, Cambridge, United Kingdom
and New York, NY, USA, <https://doi.org/10.1017/9781009157896.016>, 2021.
- Jacob, D., Teichmann, C., Sobolowski, S., Katragkou, E., Anders, I., Belda, M., Benestad, R., Boberg, F., Buonomo, E., Cardoso, R. M.,
805 Casanueva, A., Christensen, O. B., Christensen, J. H., Coppola, E., De Cruz, L., Davin, E. L., Dobler, A., Domínguez, M., Fealy, R.,
Fernandez, J., Gaertner, M. A., García-Díez, M., Giorgi, F., Gobiet, A., Goergen, K., Gómez-Navarro, J. J., Alemán, J. J. G., Gutiérrez,
C., Gutiérrez, J. M., Güttler, I., Haensler, A., Halenka, T., Jerez, S., Jiménez-Guerrero, P., Jones, R. G., Keuler, K., Kjellström, E., Knist,
S., Kotlarski, S., Maraun, D., van Meijgaard, E., Mercogliano, P., Montávez, J. P., Navarra, A., Nikulin, G., de Noblet-Ducoudré, N.,
Panitz, H.-J., Pfeifer, S., Piazza, M., Pichelli, E., Pietikäinen, J.-P., Prein, A. F., Preuschmann, S., Rechid, D., Rockel, B., Romera, R.,
810 Sánchez, E., Sieck, K., Soares, P. M. M., Somot, S., Srncic, L., Sørland, S. L., Termonia, P., Truhetz, H., Vautard, R., Warrach-Sagi, K., and
Wulfmeyer, V.: Regional climate downscaling over Europe: perspectives from the EURO-CORDEX community, *Reg Environ Change*,
20, 51, <https://doi.org/10.1007/s10113-020-01606-9>, 2020.
- Jones, P. W.: First- and Second-Order Conservative Remapping Schemes for Grids in Spherical Coordinates, *Monthly Weather Review*,
127, 2204–2210, [https://doi.org/10.1175/1520-0493\(1999\)127<2204:FASOCR>2.0.CO;2](https://doi.org/10.1175/1520-0493(1999)127<2204:FASOCR>2.0.CO;2), publisher: American Meteorological Society
815 Section: *Monthly Weather Review*, 1999.
- Journée, M., Ingels, R., and Bertrand, C.: Overview and validation of observational gridded data products for Belgium, in: *EMS Annual*
Meeting Abstracts, vol. 16, pp. EMS2019–99, Copernicus, 2019.
- Journée, M., Tricot, C., Verhumst, K., Hamdi, R., and Dehem, D.: Validation des données du réseau pluviométrique géré par l’IBGE,
répartition des précipitations et projet d’étude “changement climatique et ressources en eau” en Région bruxelloise, Tech. rep., Institut
820 royal météorologique de Belgique, Bruxelles, Belgique, 2014.
- Kendon, E. J., Roberts, N. M., Senior, C. A., and Roberts, M. J.: Realism of Rainfall in a Very High-Resolution Regional Climate Model,
<https://doi.org/10.1175/JCLI-D-11-00562.1>, section: *Journal of Climate*, 2012.
- Laprise, R.: The Euler Equations of Motion with Hydrostatic Pressure as an Independent Variable, https://journals.ametsoc.org/view/journals/mwre/120/1/1520-0493_1992_120_0197_teeomw_2_0_co_2.xml, section: *Monthly Weather Review*, 1992.

- 825 Lehner, F., Deser, C., Maher, N., Marotzke, J., Fischer, E. M., Brunner, L., Knutti, R., and Hawkins, E.: Partitioning climate projection uncertainty with multiple large ensembles and CMIP5/6, *Earth System Dynamics*, 11, 491–508, <https://doi.org/10.5194/esd-11-491-2020>, publisher: Copernicus GmbH, 2020.
- Leutwyler, D., Lüthi, D., Ban, N., Fuhrer, O., and Schär, C.: Evaluation of the convection-resolving climate modeling approach on continental scales, *Journal of Geophysical Research: Atmospheres*, 122, 5237–5258, <https://doi.org/10.1002/2016JD026013>, 2017.
- 830 Lucas-Picher, P., Argüeso, D., Brisson, E., Tramblay, Y., Berg, P., Lemonsu, A., Kotlarski, S., and Caillaud, C.: Convection-permitting modeling with regional climate models: Latest developments and next steps, *WIREs Climate Change*, 12, e731, <https://doi.org/10.1002/wcc.731>, 2021.
- Marquet, P. and Geleyn, J.: On a general definition of the squared Brunt–Väisälä frequency associated with the specific moist entropy potential temperature, *Quart J Royal Meteorol Soc*, 139, 85–100, <https://doi.org/10.1002/qj.1957>, 2013.
- 835 Masson, V., Champeaux, J.-L., Chauvin, F., Meriguet, C., and Lacaze, R.: A Global Database of Land Surface Parameters at 1-Km Resolution in Meteorological and Climate Models, *Journal of Climate*, 16, 1261–1282, [https://doi.org/10.1175/1520-0442\(2003\)16<1261:AGDOLS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)16<1261:AGDOLS>2.0.CO;2), 2003.
- Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouyssel, F., Brousseau, P., Brun, E., Calvet, J.-C., Carrer, D., Decharme, B., Delire, C., Donier, S., Essaouini, K., Gibelin, A.-L., Giordani, H., Habets, F., Jidane, M.,
- 840 Kerdraon, G., Kourzeneva, E., Lafaysse, M., Lafont, S., Lebeaupin Brossier, C., Lemonsu, A., Mahfouf, J.-F., Marguinaud, P., Mokhtari, M., Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., Tanguy, G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of earth surface variables and fluxes, *Geoscientific Model Development*, 6, 929–960, <https://doi.org/10.5194/gmd-6-929-2013>, publisher: Copernicus GmbH, 2013.
- Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.,
- 845 Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., eds.: Framing, Context, and Methods, p. 147–286, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.003>, 2021.
- Mašek, J., Geleyn, J., Brožková, R., Giot, O., Achom, H. O., and Kuma, P.: Single interval shortwave radiation scheme with parameterized optical saturation and spectral overlaps, *Quart J Royal Meteorol Soc*, 142, 304–326, <https://doi.org/10.1002/qj.2653>, 2016.
- 850 Meredith, E. P., Ulbrich, U., Rust, H. W., and Truhetz, H.: Present and future diurnal hourly precipitation in 0.11° EURO-CORDEX models and at convection-permitting resolution, *Environ. Res. Commun.*, 3, 055 002, <https://doi.org/10.1088/2515-7620/abf15e>, publisher: IOP Publishing, 2021.
- Müller, S. K., Caillaud, C., Chan, S., de Vries, H., Bastin, S., Berthou, S., Brisson, E., Demory, M.-E., Feldmann, H., Goergen, K., Kartsios, S., Lind, P., Keuler, K., Pichelli, E., Raffa, M., Tölle, M. H., and Warrach-Sagi, K.: Evaluation of Alpine-Mediterranean precipitation events in convection-permitting regional climate models using a set of tracking algorithms, *Clim Dyn*, 61, 939–957, <https://doi.org/10.1007/s00382-022-06555-z>, 2023.
- 855 Müller, S. K., Pichelli, E., Coppola, E., Berthou, S., Brienens, S., Caillaud, C., Demory, M.-E., Dobler, A., Feldmann, H., Mercogliano, P., Tölle, M., and de Vries, H.: The climate change response of alpine-mediterranean heavy precipitation events, *Clim Dyn*, 62, 165–186, <https://doi.org/10.1007/s00382-023-06901-9>, 2024.
- 860 Nguyen, P. L., Alexander, L. V., Thatcher, M. J., Truong, S. C. H., Isphording, R. N., and McGregor, J. L.: Selecting CMIP6 global climate models (GCMs) for Coordinated Regional Climate Downscaling Experiment (CORDEX) dynamical downscaling over Southeast Asia

- using a standardised benchmarking framework, *Geoscientific Model Development*, 17, 7285–7315, <https://doi.org/10.5194/gmd-17-7285-2024>, publisher: Copernicus GmbH, 2024.
- Noilhan, J. and Mahfouf, J. F.: The ISBA land surface parameterisation scheme, *Global and Planetary Change*, 13, 145–159, [https://doi.org/10.1016/0921-8181\(95\)00043-7](https://doi.org/10.1016/0921-8181(95)00043-7), 1996.
- Noilhan, J. and Planton, S.: A Simple Parameterization of Land Surface Processes for Meteorological Models, [https://doi.org/10.1175/1520-0493\(1989\)117<0536:ASPOLS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<0536:ASPOLS>2.0.CO;2), section: Monthly Weather Review, 1989.
- Pichelli, E., Coppola, E., Sobolowski, S., Ban, N., Giorgi, F., Stocchi, P., Alias, A., Belušić, D., Berthou, S., Caillaud, C., Cardoso, R. M., Chan, S., Christensen, O. B., Dobler, A., de Vries, H., Goergen, K., Kendon, E. J., Keuler, K., Lenderink, G., Lorenz, T., Mishra, A. N., Panitz, H.-J., Schär, C., Soares, P. M. M., Truhetz, H., and Vergara-Temprado, J.: The first multi-model ensemble of regional climate simulations at kilometer-scale resolution part 2: historical and future simulations of precipitation, *Clim Dyn*, 56, 3581–3602, <https://doi.org/10.1007/s00382-021-05657-4>, 2021.
- Prein, A. F., Langhans, W., Fosser, G., Ferrone, A., Ban, N., Goergen, K., Keller, M., Tölle, M., Gutjahr, O., Feser, F., Brisson, E., Kollet, S., Schmidli, J., van Lipzig, N. P. M., and Leung, R.: A review on regional convection-permitting climate modeling: Demonstrations, prospects, and challenges, *Reviews of Geophysics*, 53, 323–361, <https://doi.org/10.1002/2014RG000475>, 2015.
- Ranasinghe, R., Ruane, A., Vautard, R., Arnell, N., Coppola, E., Cruz, F., Dessai, S., Islam, A., Rahimi, M., Ruiz Carrascal, D., Sillmann, J., Sylla, M., Tebaldi, C., Wang, W., and Zaaboul, R.: *Climate Change Information for Regional Impact and for Risk Assessment*, p. 1767–1926, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.014>, 2021.
- Rummukainen, M.: Added value in regional climate modeling, *WIREs Climate Change*, 7, 145–159, <https://doi.org/10.1002/wcc.378>, 2016.
- Simmons, A. J. and Burridge, D. M.: An Energy and Angular-Momentum Conserving Vertical Finite-Difference Scheme and Hybrid Vertical Coordinates, [https://doi.org/10.1175/1520-0493\(1981\)109<0758:AEAAMC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1981)109<0758:AEAAMC>2.0.CO;2), section: Monthly Weather Review, 1981.
- Soares, P., Careto, J., Cardoso, R., Goergen, K., Katragkou, E., Sobolowski, S., Coppola, E., Ban, N., Belušić, D., Berthou, S., Caillaud, C., Dobler, A., Hodnebrog, o., Kartsios, S., Lenderink, G., Lorenz, T., Milovac, J., Feldmann, H., Pichelli, E., and Bastin, S.: The added value of km-scale simulations to describe temperature over complex orography: the CORDEX FPS-Convection multi-model ensemble runs over the Alps, *Climate Dynamics*, 62, <https://doi.org/10.1007/s00382-022-06593-7>, 2022.
- Tabari, H., De Troch, R., Giot, O., Hamdi, R., Termonia, P., Saeed, S., Brisson, E., Van Lipzig, N., and Willems, P.: Local impact analysis of climate change on precipitation extremes: are high-resolution climate models needed for realistic simulations?, *Hydrology and Earth System Sciences*, 20, 3843–3857, <https://doi.org/10.5194/hess-20-3843-2016>, publisher: Copernicus GmbH, 2016.
- Termonia, P., Fischer, C., Bazile, E., Bouyssel, F., Brožková, R., Bénard, P., Bochenek, B., Degrauwe, D., Derková, M., El Khatib, R., Hamdi, R., Mašek, J., Pottier, P., Pristov, N., Seity, Y., Smolíková, P., Španiel, O., Tudor, M., Wang, Y., Wittmann, C., and Joly, A.: The ALADIN System and its canonical model configurations AROME CY41T1 and ALARO CY40T1, *Geoscientific Model Development*, 11, 257–281, <https://doi.org/10.5194/gmd-11-257-2018>, publisher: Copernicus GmbH, 2018a.
- Termonia, P., Van Schaeybroeck, B., De Cruz, L., De Troch, R., Caluwaerts, S., Giot, O., Hamdi, R., Vannitsem, S., Duchêne, F., Willems, P., Tabari, H., Van Uytven, E., Hosseinzadehtalaei, P., Van Lipzig, N., Wouters, H., Vanden Broucke, S., van Ypersele, J.-P., Marbaix, P., Villanueva-Birriel, C., Fettweis, X., Wyard, C., Scholzen, C., Doutreloup, S., De Ridder, K., Gobin, A., Lauwaet, D., Stavrakou, T., Bauwens, M., Müller, J.-F., Luyten, P., Ponsar, S., Van den Eynde, D., and Pottiaux, E.: The CORDEX.be initiative as a foundation for climate services in Belgium, *Climate Services*, 11, 49–61, <https://doi.org/10.1016/j.cliser.2018.05.001>, 2018b.

- Termonia, P., Van Schaeybroeck, B., De Cruz, L., De Troch, R., Hamdi, R., Vannitsem, S., Duchêne, F., Willems, P., Tabari, H., Van Uytven, E., Hosseinzadehtalaei, P., Van Lipzig, N., Wouters, H., Vanden Broucke, S., van Ypersele, J.-P., Marbaix, P., Villanueva-Birriel, C., Fettweis, X., Wyard, C., Scholzen, C., Doutreloup, S., De Ridder, K., Gobin, A., Lauwaet, D., Stavrakou, T., Bauwens, M., Müller, J.-F., Luyten, P., Ponsar, S., and Pottiaux, E.: Combining Regional Downscaling Expertise in Belgium: CORDEX and Beyond, Tech. Rep. CORDEX.be Final Report, Belgian Science Policy Office (BELSPO), Brussels, Belgium, https://www.belspo.be/belspo/brain-be/projects/FinalReports/CORDEXbe_FinRep_AD.pdf, 2018c.
- Thomassen, E. D., Arnbjerg-Nielsen, K., Sørup, H. J. D., Langen, P. L., Olsson, J., Pedersen, R. A., and Christensen, O. B.: Spatial and temporal characteristics of extreme rainfall: Added benefits with sub-kilometre-resolution climate model simulations?, *Quarterly Journal of the Royal Meteorological Society*, 149, 1913–1931, <https://doi.org/10.1002/qj.4488>, 2023.
- Top, S., Kotova, L., De Cruz, L., Aniskevich, S., Bobylev, L., De Troch, R., Gnatiuk, N., Gobin, A., Hamdi, R., Kriegsmann, A., Remedio, A. R., Sakalli, A., Van De Vyver, H., Van Schaeybroeck, B., Zandersons, V., De Maeyer, P., Termonia, P., and Caluwaerts, S.: Evaluation of regional climate models ALARO-0 and REMO2015 at 0.22° resolution over the CORDEX Central Asia domain, *Geoscientific Model Development*, 14, 1267–1293, <https://doi.org/10.5194/gmd-14-1267-2021>, publisher: Copernicus GmbH, 2021.
- Van De Vyver, H.: Spatial regression models for extreme precipitation in Belgium, *Water Resources Research*, 48, 2011WR011707, <https://doi.org/10.1029/2011WR011707>, 2012.
- Van de Vyver, H.: Practical Return Level Mapping for Extreme Precipitation in Belgium, Scientific and Technical Publication 062, Royal Meteorological Institute of Belgium, <https://orfeo.belnet.be/handle/internal/8489>, 2013.
- Van De Vyver, H.: A multiscaling-based intensity–duration–frequency model for extreme precipitation, *Hydrological Processes*, 32, 1635–1647, <https://doi.org/10.1002/hyp.11516>, 2018.
- Van de Vyver, H.: Observed Annual Maximum Sub-Daily Precipitation from the Belgian Hydro-Meteorological Network (1967–2004) and Uccle (1898–2007), <https://doi.org/10.5281/ZENODO.4741178>, 2021.
- Van De Vyver, H., Van Schaeybroeck, B., De Troch, R., De Cruz, L., Hamdi, R., Villanueva-Birriel, C., Marbaix, P., Van Ypersele, J.-P., Wouters, H., Vanden Broucke, S., Van Lipzig, N. P., Doutreloup, S., Wyard, C., Scholzen, C., Fettweis, X., Caluwaerts, S., and Termonia, P.: Evaluation framework for sub-daily rainfall extremes simulated by regional climate models, *Journal of Applied Meteorology and Climatology*, <https://doi.org/10.1175/JAMC-D-21-0004.1>, 2021.
- Van Ginderachter, M., Degrauwe, D., Vannitsem, S., and Termonia, P.: Simulating Model Uncertainty of Subgrid-Scale Processes by Sampling Model Errors at Convective Scales, *Nonlinear Processes in Geophysics*, 27, 187–207, <https://doi.org/10.5194/npg-27-187-2020>, 2020.
- Veneziano, D. and Furcolo, P.: Multifractality of rainfall and scaling of intensity-duration-frequency curves, *Water Resources Research*, 38, 42–1–42–12, <https://doi.org/10.1029/2001WR000372>, 2002.
- Vergara-Temprado, J., Ban, N., Panosetti, D., Schlemmer, L., and Schär, C.: Climate Models Permit Convection at Much Coarser Resolutions Than Previously Considered, <https://doi.org/10.1175/JCLI-D-19-0286.1>, section: Journal of Climate, 2020.
- Zheng, Y., Alapaty, K., Herwehe, J. A., Del Genio, A. D., and Niyogi, D.: Improving High-Resolution Weather Forecasts Using the Weather Research and Forecasting (WRF) Model with an Updated Kain–Fritsch Scheme, *Monthly Weather Review*, 144, 833–860, <https://doi.org/10.1175/MWR-D-15-0005.1>, 2016.
- Đurán, I. B., Geleyn, J.-F., and Váňa, F.: A Compact Model for the Stability Dependency of TKE Production–Destruction–Conversion Terms Valid for the Whole Range of Richardson Numbers, *Journal of the Atmospheric Sciences*, 71, 3004–3026, <https://doi.org/10.1175/JAS-D-13-0203.1>, 2014.