

Dynamic and Thermal Analysis of Sandstorm Processes Based on Vertical Observation Data

Yifei Wang^{1,2,3,4,5}, Wen Huo^{1,3,4,5}, Yongqiang Liu^{1,2,3,4,5}, Mayibaier Maihamuti^{1,2,3,4,5},
Fan Yang^{1,3,4,5}, Chenglong Zhou^{1,3,4,5}, Xinghua Yang^{1,3,4,5,6}, Ali Mamtimin^{1,3,4,5}

¹Institute of Desert Meteorology, China Meteorological Administration, Urumqi 830002, China

²College of Geography and Remote Sensing Sciences, Xinjiang University, Urumqi 830046, China

³National Observation and Research Station of Desert Meteorology, Taklimakan Desert of Xinjiang, Urumqi 830002, China

⁴Taklimakan Desert Meteorology Field Experiment Station, China Meteorological Administration, Urumqi 830002, China

⁵Xinjiang Key Laboratory of Desert Meteorology and Sandstorm, Urumqi 830002, China

⁶School of Geographical Sciences, Shanxi Normal University, Taiyuan 030032, China

Correspondence to: Wen Huo (huowenpet@idm.cn)

Abstract. The Taklamakan Desert (TD) is a major dust-source region in East Asia, yet the dynamic and thermal processes associated with dust-storm development remain insufficiently understood from vertically resolved observations. This study combined dual-gradient tower observations from Tazhong (TZ) and Xiaotang (XT), ERA5 reanalysis, and backward trajectories calculated with the Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model to examine eight dust-storm events in spring and summer 2024. Three main pre-storm airflow pathways were identified: east-inflow transport through the northeastern Tarim Basin, mountain-crossing transport across the Tianshan–Pamir region, and south-to-north transport across the Tibetan Plateau and Kunlun Mountains. ERA5 anomaly diagnostics show that spring events maintained stronger mean sea-level pressure-gradient and 850 hPa geopotential-height-gradient anomalies throughout the pre-onset period, while 10 m wind-speed anomalies increased rapidly after approximately –12 h. In contrast, summer events showed stronger thermal signals before approximately –18 h, including enhanced surface–air temperature contrast, positive net shortwave radiation anomalies, and boundary-layer growth. Tower observations further indicate stronger summer warming and closer-to-onset wind-speed and wind-shear enhancement. Differences between TZ and XT suggest that local terrain modulates the near-surface response to regional dynamic and thermal forcing. These findings provide observational evidence for seasonal differences in TD dust-storm processes.

Keywords: Dust storm; Taklimakan Desert; HYSPLIT; ERA5; dynamic and thermal anomalies; vertical observation

1 Introduction

Dust storms are among the major natural hazards in arid and semi-arid regions of East Asia, where they are often accompanied by strong winds, air pollution, and reduced visibility, affecting transportation, agriculture, ecosystems, and public health (Achakulwisut et al., 2018; Bao and Fang, 2007; Manisalidis et al., 2020; Shao and Dong, 2006; Tong et al., 2023; Wen et al., 2024). Mineral dust can also influence atmospheric radiation, cloud processes, snow and ice albedo, and regional climate through long-range transport (Kok et al., 2023; Prospero, 1999).

The Taklimakan Desert (TD) is located in the arid Tarim Basin and is surrounded by the Tianshan, Pamir, and Kunlun Mountains. Because of its extensive mobile sand surfaces, scarce precipitation, and unique basin topography, the TD is one of the major dust-source regions in East Asia (Ge et al., 2014; Shao and Dong, 2006; Wang et al., 2015; Xu et al., 2020; Yang et al., 2016; Zhang et al., 2025).

The formation and development of dust storms are jointly controlled by atmospheric forcing and surface conditions. Among these controls, dynamic and thermal processes are particularly important because they determine whether surface particles can be lifted and whether the lifted dust can be transported vertically and horizontally (Bi et al., 2022; Su et al., 2024; Zhang et al., 2024). Dynamic forcing, represented by pressure-gradient-driven winds, cold fronts, cyclones, and low-level jets, provides the direct momentum for dust emission (Shao and Dong, 2006; Xu et al., 2020). Thermal forcing, mainly associated with strong surface heating, boundary-layer growth, turbulent mixing, and land–atmosphere temperature contrasts, can enhance vertical exchange and promote dust uplift, especially over inland deserts with intense surface heating (Wang et al., 2025; Xiong et al., 2023). Surface factors such as particle-size distribution, soil moisture, vegetation cover, and aerodynamic roughness further regulate the threshold and intensity of dust emission (Iversen and White, 1982; Marticorena and Bergametti, 1995; Raupach, 1992; Shao and Dong, 2006). Therefore, dust-storm processes cannot be explained by a single factor alone, but should be examined through the coupled effects of dynamic forcing, thermal instability, transport pathways, and terrain-modified local surface conditions.

Previous studies have shown that dynamic forcing is a key driver of many spring dust storms in East

Asia. Mongolian cyclones, cold fronts, cold high-pressure systems, and strong pressure gradients can generate strong near-surface winds and rapidly activate dust sources over Mongolia, northern China, and the Tarim Basin (Gao et al., 2024; Liu et al., 2004; Wang et al., 2009; Xu et al., 2020). S. Chen et al. (2023a) showed that severe dust events in northern China during March and April 2023 were closely related to cold fronts and Mongolian cyclones, with Mongolia and the Taklimakan Desert both contributing substantially to dust concentrations. Li et al. (2023) further demonstrated that dust radiative feedback can influence the intensity and movement of Mongolian cyclones and cold fronts, indicating that dust storms are not only a response to weather systems but may also interact with them. More recently, Mu et al. (2025) quantitatively assessed the role of atmospheric depressions and Mongolian cyclones in spring dust activity over East Asia and found that Mongolian cyclones are a key driver of high-impact dust events. These studies highlight the importance of synoptic-scale dynamic forcing in East Asian dust-storm development. However, many of them focus on northern China and the Gobi source region, whereas the pre-storm dynamic evolution over the Taklimakan Desert, especially its connection with local tower-based observations, remains less well documented.

Compared with the relatively well-recognized role of dynamic forcing, the influence of thermal forcing is more complex and is particularly important for inland desert regions such as the TD. Strong solar heating over mobile sand surfaces can increase the surface–air temperature difference, deepen the atmospheric boundary layer, enhance turbulent mixing, and create favorable conditions for vertical dust uplift (Chen et al., 2023b; Xiong et al., 2023; Yuan et al., 2019; Zhang et al., 2025). Yuan et al. (2019) investigated the influence of dynamic and thermal forcing on the meridional transport of Taklimakan dust in spring and summer and emphasized that seasonal differences in dust transport are closely linked to changes in forcing mechanisms. Y. Chen et al. (2023b) used WRF-Chem simulations to investigate the radiative effects of dust aerosols on weather conditions and dust transport over the TD. Their results suggested that dust–radiation interactions can modify temperature, wind fields, and vertical dust transport, indicating the potential importance of thermal feedbacks in dust-storm evolution. X. Zhang et al. (2025) further indicated that spring Taklimakan dust transported toward the Tibetan Plateau is controlled by circulation patterns, dynamic lifting, thermal anomalies, and terrain effects. These findings suggest that distinguishing between dynamic and thermal controls is useful for understanding seasonal differences in dust-storm development. Nevertheless, how these seasonal differences are reflected in near-surface wind, temperature, and vertical meteorological profiles still requires further observational evidence,

particularly from direct comparisons between spring and summer events.

Transport pathways provide an important way to connect dynamic and thermal forcing with dust-storm evolution. They reflect not only where air masses and dust originate, but also how synoptic circulation, basin-scale airflow, and thermal lifting organize dust movement before and during dust storms. Kai and Huiwang (2007) identified major East Asian dust transport pathways toward eastern China, Korea, Japan, the western Pacific, and Central Asia. T. Bao et al. (2023) examined East Asian dust sources and transport ranges from 1980 to 2020 using ground observations and remote sensing data, and showed that southern Mongolia and the Taklimakan Desert are major source areas. For the TD, Aili et al. (2021) found that air masses during the dusty season at the northeastern desert margin mainly originated from the eastern and southern parts of the desert, indicating the importance of intra-basin transport. Aili et al. (2024) further used HYSPLIT trajectories to establish an early-warning framework for dusty weather in peri-desert regions of the TD. These studies demonstrate the value of trajectory analysis, but pathway classification alone cannot fully explain why different dust storms develop under different dynamic or thermal backgrounds.

Topography further modifies dust-storm occurrence, transport, and near-surface wind and temperature structures. The TD is enclosed by high mountains and connected to surrounding regions through limited basin openings, so airflow pathways are strongly constrained by terrain. The Tarim Basin and surrounding mountain systems can channel near-surface winds, favor dust accumulation within the basin, and promote vertical lifting under suitable dynamic and thermal conditions (Sun and Liu, 2006; Xing et al., 2024; Yuan et al., 2019; Zhang et al., 2025). Thus, topography is closely linked to both transport pathways and dynamic–thermal forcing rather than being only a static background condition. At the local scale, dune relief and underlying-surface differences can alter wind profiles, near-surface thermal structure, and sediment availability. (Aili et al., 2023).

2 Study Area, Data, and Methodology

2.1 Study Area

The Taklamakan Desert (TD) is located within the Tarim Basin in the southwestern part of the Xinjiang Uyghur Autonomous Region in China. It is the second-largest shifting sand desert in the world. Surrounded by mountain ranges such as the Tianshan and Kunlun Mountains, the region experiences a

highly arid climate. Key characteristics of the desert include its remoteness from the ocean, sparse vegetation, diverse dune types with high mobility, extensive and thick shifting sand areas, and fine sand particles. This region is subject to frequent dust storms year-round and is one of the significant sources of dust storms in East Asia(Sun and Liu, 2006).




Figure 1. Location and topography of the Taklimakan Desert, including the Tazhong (TZ) and Xiaotang (XT) observation towers and site photographs.

2.2 Observational Data

The observational data were obtained from two stations: Xiaotang (XT), located on the northern edge of the TD, representing flat terrain, and TZ, situated in the desert interior, representing undulating terrain. The observation tower height is 80 m at TZ and 100 m at XT. Both stations are equipped with meteorological gradient observation systems and dust collection systems. The observation levels at TZ are at 1 m, 2 m, 5 m, 8 m, 16 m, 24 m, 32 m, 47 m, 63 m, and 80 m. while those at XT are 1 m, 2 m, 5 m, 10 m, 24 m, 32 m, 47 m, 63 m, 80 m, and 100 m.

The meteorological gradient observation system records meteorological variables at each layer, including hourly and minute-based measurements of temperature, relative humidity, wind speed, wind direction, and atmospheric pressure. The dust collection system gathers dust samples during dust storms using BSNE dust collectors installed at each layer. In the present analysis, the meteorological gradient observations were mainly used to examine wind-speed, wind-shear, and temperature-profile responses during dust-storm development.

Nine dust storm events were observed between 31 March and 18 June 2024, and the corresponding

dust samples were collected from 3 April to 10 July 2024 (as detailed in Table 1). However, during the second observation, data from TZ were missing, preventing a complete paired record. Therefore, the remaining eight complete dust storm events observations became the primary samples for this study. To reduce the influence of seasonal variation and diurnal cycles on the station-based meteorological profiles, a background-removal procedure was applied to the observational data from both TZ and XT. For each dust-storm event, observation height, and variable, the background value was calculated separately for the two stations using records from 2023 and 2025. Specifically, for a given observation time in 2024, all available records at the same hour of the day within a ± 15 -day calendar window in 2023 and 2025 were averaged to represent the corresponding background state. The anomaly was then obtained by subtracting this station-specific background value from the observed value during the 2024 dust-storm process. This approach reduces the effects of normal seasonal progression and daily temperature or wind-speed cycles, allowing the analysis to focus on meteorological anomalies directly associated with dust-storm development.

Table 1: Summary of observed dust-storm events and sample collection information

Event ID	Station	Start time of dust storm	Duration (h)	Sample collection time
Dust1	TZ	2024-03-31 12:00	22	2024-04-01 09:00
	XT	2024-03-31 12:40	17.3	2024-04-01 06:00
Excluded event	XT	2024-04-05 01:00	5	2024-04-05 06:00
Dust2	XT	2024-04-12 04:00	7.5	2024-04-12 14:00
	TZ	2024-04-14 04:00	11	2024-04-14 15:00
Dust3	XT	2024-04-17 11:00	23	2024-04-18 10:00
	TZ	2024-04-18 10:00	36	2024-04-19 22:00
Dust4	TZ	2024-04-26 21:00	16.5	2024-04-27 13:30
	XT	2024-04-26 02:00	31	2024-04-27 09:00
Dust5	TZ	2024-05-12 07:00	24	2024-05-12 14:30
	XT	2024-05-12 07:30	7	2024-05-12 14:30
Dust6	TZ	2024-05-20 11:00	14	2024-05-20 14:30
	XT	2024-05-20 11:00	3.5	2024-05-20 14:30

Dust7	TZ	2024-06-04 04:00	12	2024-06-04 16:00
	XT	2024-06-04 04:00	3	2024-06-04 07:00
Dust8	TZ	2024-06-18 07:00	2	2024-06-18 09:00
	XT	2024-06-18 07:20	3.7	2024-06-18 11:00

2.3 ERA5 Reanalysis Data

To further investigate the meteorological conditions associated with dust storm occurrence, this study used the ERA5 reanalysis dataset. ERA5, provided by the European Centre for Medium-Range Weather Forecasts (ECMWF), is the fifth generation of global atmospheric reanalysis data. It has a spatial resolution of $0.25^\circ \times 0.25^\circ$ and a temporal resolution of 1 hour.

In this study, ERA5 data were used in two ways. First, the original ERA5 fields were used to describe the synoptic-scale meteorological background before and during each dust-storm event, including mean sea-level pressure, 10 m wind vectors, 2 m air temperature, and 850 hPa geopotential height. These fields were mainly displayed over the regional domain of $70\text{--}100^\circ \text{E}$ and $30\text{--}50^\circ \text{N}$ to identify the pressure configuration, airflow pathway, and thermal background around the Taklamakan Desert.

Second, selected ERA5-derived diagnostic indicators were used for anomaly analysis after background removal. The dynamic indicators included mean sea-level pressure-gradient magnitude, 850 hPa geopotential-height-gradient magnitude, and 10 m wind speed. These indicators were averaged over the regional domain of $70\text{--}100^\circ \text{E}$ and $30\text{--}50^\circ \text{N}$ to represent the large-scale dynamic background around the TD and its surrounding mountains. The thermal indicators included 2 m air temperature, skin temperature, surface–air temperature difference, surface net shortwave radiation, and boundary-layer height. These indicators were averaged over the Taklamakan Desert core region. The desert core was defined using a Taklamakan Desert polygon mask within $72\text{--}92^\circ \text{E}$ and $35\text{--}43^\circ \text{N}$. Grid cells outside the desert mask were excluded from the thermal averaging.

For each dust-storm event and each lead time relative to storm onset, the corresponding 2024 ERA5 value was extracted. The background value was calculated from ERA5 records during 2015–2025, excluding 2024, using the same hour of the day within a ± 3 -day calendar window. The anomaly was then obtained by subtracting this background value from the corresponding 2024 event value. Area-mean values were calculated using latitude-weighted averaging with $\cos(\text{latitude})$ as the weight. This treatment

reduced the influence of seasonal progression, diurnal variation, and unequal grid-cell area, thereby highlighting the dynamic and thermal departures related to dust-storm development.

2.4 HYSPLIT Model

The study used the Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model to analyze the transport pathways of each dust storm event. This model uses grid data from the Global Data Assimilation System (GDAS) dataset to compute backward trajectories (Stein et al., 2015). The GDAS1 data have a horizontal resolution of $1^\circ \times 1^\circ$ and a temporal interval of 3 h. Model vertical velocity was used for the vertical motion calculation.

For each event, backward trajectories were initialized at the dust-storm onset time. The trajectory endpoint was set at 39.43° N, 85.97° E, corresponding to the observation region used in this study. Three starting heights, 100, 500, and 1500 m above ground level (AGL), were selected to represent airflow transport at different lower-tropospheric levels. Each trajectory was traced backward for 48 h. The trajectory ending times were 12:00 UTC on 31 March, 04:00 UTC on 12 April, 11:00 UTC on 17 April, 21:00 UTC on 26 April, 07:00 UTC on 12 May, 11:00 UTC on 20 May, 04:00 UTC on 4 June, and 07:00 UTC on 18 June 2024 for Dust1–Dust8, respectively.

The method assumes that the trajectory of a particle moving within the wind field is calculated by integrating its position changes over time and space, with its vector velocity determined by linear interpolation in both space and time. The first-guess position of the air mass is calculated using the following equation:

$$P_1(t + \Delta t) = P(t) + V(P, t)\Delta t \quad (1)$$

The corrected position is then calculated as:

$$P(t + \Delta t) = P(t) + 0.5[V(P, t) + V(P_1, t + \Delta t)]\Delta t \quad (2)$$


In this Eq. (1) and (2), P represents the initial position, P_1 is the first guessed position, Δt is the time interval, and V is the velocity of the air mass or particle. The simulation was conducted online using the HYSPLIT platform provided by NOAA.

The resulting trajectories were used to characterize the main pre-storm airflow histories reaching the observation region and to support the interpretation of airflow transport pathways together with ERA5 10 m wind fields.

206 3 Results and Analysis

207 3.1 Dust Storm Trajectory Analysis

208



209

210 **Figure 2. Backward trajectories of the eight dust-storm events.**




Figure 3. ERA5 10 m wind-vector fields during the eight dust-storm events.

As shown in Fig 2, the HYSPLIT backward trajectories indicate that the analyzed dust-storm events were associated with three main types of pre-storm airflow transport pathways. Dust1, Dust3, Dust4, Dust5, and Dust6 mainly showed east-inflow transport, with air masses entering the TD through the northeastern opening of the Tarim Basin. Dust2 was characterized by a south-to-north pathway related to airflow crossing the Tibetan Plateau and Kunlun Mountains. Dust7 and Dust8 were associated with northwestern transport, with air masses passing over southern Russia and eastern Uzbekistan before crossing the Tianshan Mountains toward the TD.

The ERA5 10 m wind fields provide regional support for the trajectory results (Fig 3). For most east-inflow events, the near-surface wind field shows organized airflow entering the basin through the eastern or northeastern opening. For Dust7 and Dust8, part of the airflow shows a north-to-south component, consistent with mountain-crossing transport across the Tianshan Mountains. Dust2 differs from the other events, with southerly airflow from the Tibetan Plateau toward southern Xinjiang. These results suggest that the TD dust-storm events were influenced by multiple airflow pathways, among which east-inflow transport was the most frequent in the analyzed cases.

3.2 Dust Storm Dynamic and Thermal Meteorological Background

Figure 4 shows the regional mean sea-level pressure, 850 hPa geopotential height, and 10 m wind vectors before the analyzed dust-storm events. In the spring cases, especially Dust1, Dust3, and Dust4, clearer pressure differences appear between northern Xinjiang, surrounding high-pressure regions, and the TD. These pressure and circulation patterns are consistent with the east-inflow and mountain-crossing pathways identified from the trajectory analysis. Dust2 shows a different circulation background, with

more evident influence from the southern and southwestern margins of the basin, corresponding to its south-to-north transport pathway.




Figure 4. Mean sea-level pressure, 10 m wind vectors, and 850 hPa geopotential height before the analyzed dust-storm events.




Figure 5. 2 m air temperature, 10 m wind vectors, and 850 hPa geopotential height during the analyzed dust-storm events.

In the summer cases, especially Dust7 and Dust8, the pressure-gradient signal is weaker and the dominant wind direction is less clear from the pressure field alone. This indicates that dynamic forcing was less spatially organized than in the spring cases. At the same time, Fig 5 shows that 2 m air temperature over the TD was higher and more spatially continuous in the May–June events. Stronger surface heating may have enhanced the surface–air temperature difference, promoted boundary-layer development, and provided favorable thermal conditions for dust mobilization.

These background fields indicate a seasonal contrast in the pre-storm environment. The March–April events were more closely associated with stronger pressure-gradient and lower-tropospheric circulation signals, whereas the May–June events showed a more evident thermal background. Therefore,

the analyzed events were grouped into spring cases and summer cases for the following dynamic and thermal anomaly analysis.

3.3 Analysis of Dynamic and Thermodynamic Factors

Figure 6 and 7 show the sea-level pressure differences at 24, 12, 6, and 3 h before dust-storm onset in spring and summer, respectively. The pressure difference was calculated as the sea-level pressure at onset minus that at each preceding time. Positive values indicate pressure increase toward onset, whereas negative values indicate pressure decrease.




Figure 6. Sea-level pressure differences before spring dust-storm onset.




Figure 7. Sea-level pressure differences before summer dust-storm onset.

In spring, the pressure-difference fields show stronger and more continuous spatial contrasts than in summer. For Dust1, Dust3, and Dust4, evident pressure changes appear around northern Xinjiang, the northern TD, or the eastern basin margin, indicating that the pressure-gradient system had already adjusted before onset. This agrees with the large-scale dynamic background described above, in which stronger pressure gradients and 850-hPa circulation differences favored airflows entering the TD through the eastern basin gap or crossing the Tianshan Mountains. Dust2 differs from the other spring cases, with more evident pressure changes along the southern and southwestern margins of the TD, corresponding to its south-to-north transport pathway across the plateau and mountain regions.

In summer, the pressure-difference fields are generally weaker and less spatially organized. Dust5 and Dust6 still show moderate pressure changes near the eastern or northeastern TD, consistent with their partial east-inflow characteristics. However, Dust7 and Dust8 show scattered and weak pressure-difference signals, suggesting that pressure-gradient forcing alone cannot explain their occurrence. This is consistent with the previous trajectory and background-field analysis, which indicated that summer dust storms were more likely affected by local thermal adjustment and boundary-layer development.

Figures 8 and 9 show the changes in surface–air temperature difference before dust-storm onset. The surface–air temperature difference was defined as the difference between skin temperature and 2 m air temperature, and its change was calculated as the onset value minus the value at 24, 12, 6, and 3 h before onset. Thus, positive values indicate that the surface–air thermal contrast strengthened toward onset, while negative values indicate that the contrast weakened.




Figure 8. Changes in surface–air temperature difference before spring dust-storm onset.




Figure 9. Changes in surface–air temperature difference before summer dust-storm onset.

In spring, the changes in surface–air temperature difference are relatively irregular. Dust1 shows a localized increase in the thermal contrast at -12 h, but the signal weakens later. Dust2 presents a clearer increase near the southern TD, which may be related to its plateau-crossing pathway and local thermal instability. Dust3 and Dust4 show mixed positive and negative areas, indicating that the near-surface thermal structure was not consistently strengthened before all spring events. This suggests that thermal contrast may have contributed to individual spring cases, but it was not the common dominant pre-onset signal.

In summer, the increase in surface–air temperature difference is more evident. Dust5 and Dust6 show positive changes over the TD or its surrounding regions during -12 to -6 h, while Dust7 and Dust8 show broader positive areas closer to onset. These results indicate that, compared with spring, the summer cases experienced a stronger enhancement of surface–air thermal contrast before dust-storm occurrence. This pattern is consistent with the large-scale thermal background discussed above, in which stronger surface heating in May and June favored unstable stratification, turbulent mixing, and boundary-layer development.

Figure 10 further compares the composite evolution of dynamic and thermal anomalies from -24 to +3 h relative to dust-storm onset. The dynamic indicators are consistent with the spatial pressure-difference maps. The mean sea-level pressure-gradient anomaly and 850-hPa height-gradient anomaly remain higher in spring throughout the pre-onset period, indicating a persistent dynamic background. The 10-m wind-speed anomaly increases more sharply near onset, especially after about -12 h, suggesting that the earlier pressure-gradient adjustment was gradually transferred to near-surface wind enhancement.




Figure 10. Composite evolution of dynamic and thermal anomalies from -24 to +3 h relative to dust-storm onset.

The thermal curves support the spatial characteristics shown in the surface-air temperature-difference maps. In summer, the net shortwave-radiation anomaly is strongly positive during the early pre-onset stage, and the surface-air temperature-difference anomaly is also positive before about -18 h. Meanwhile, the boundary-layer-height anomaly reaches approximately 200–300 m, indicating that stronger radiation and surface heating promoted boundary-layer growth before summer dust storms. After about -12 h, the summer thermal anomalies weaken, while the near-surface wind anomaly begins

to increase. This suggests that summer thermal forcing mainly provided early preconditioning, and the final onset still required near-surface wind enhancement.

Overall, the spring dust storms were more closely linked to persistent dynamic forcing, including pressure-gradient adjustment, 850-hPa circulation differences, and subsequent wind enhancement. These features correspond well to the east-inflow and mountain-crossing pathways. In contrast, the summer cases showed weaker pressure signals but stronger early thermal preconditioning, expressed as enhanced surface–air temperature contrast, positive radiation anomalies, and boundary-layer growth. Therefore, the seasonal difference reflects a shift from dynamic-dominated spring processes to thermally preconditioned summer processes, rather than a complete separation between dynamic and thermal mechanisms.

3.4 Tower-observed wind and temperature anomaly profiles

The tower observations further support the regional dynamic and thermal characteristics identified from the ERA5-based analysis. Fig 11 shows the composite evolution of 10 m wind-speed anomaly, 2–10 m wind-shear anomaly, 0.5–10 m air-temperature-difference anomaly, and 2 m air-temperature anomaly from –24 to +3 h relative to dust-storm onset. The 2 m air-temperature anomaly is generally positive before onset, indicating that near-surface warming had already developed during the pre-onset stage. However, this warming signal weakens as onset approaches, whereas the 10 m wind-speed and wind-shear anomalies increase rapidly near onset. This pattern suggests a transition from early thermal preconditioning to enhanced near-surface momentum and wind shear immediately before dust-storm occurrence.




Figure 11. Composite evolution of tower-observed wind-speed, wind-shear, air-temperature-difference, and 2 m air-temperature anomalies from -24 to +3 h relative to dust-storm onset.

Figure 12 shows the vertical profiles of wind-speed anomalies at TZ and XT. Wind-speed anomalies generally increase before and around dust-storm onset, but their vertical structures differ between the two stations. At XT, especially in summer, positive anomalies become more evident in the lower and middle layers, suggesting stronger near-surface momentum exchange over relatively flat terrain. At TZ, the profiles are more irregular, which may reflect the influence of undulating dune topography and local flow disturbance.




Figure 12. Vertical profiles of wind-speed anomalies at TZ and XT during spring and summer dust-storm events.

Figure 13 presents the vertical profiles of air-temperature anomalies. In summer, both TZ and XT show stronger positive temperature anomalies than in spring, particularly at XT, where the warming signal is more vertically continuous. This is consistent with the stronger surface heating and boundary-layer development indicated by the ERA5 thermal diagnostics. In spring, the temperature profiles are more irregular, suggesting that thermal forcing was not a common pre-onset signal for all spring events.




Figure 13. Vertical profiles of air-temperature anomalies at TZ and XT during spring and summer dust-storm events.

Overall, the tower observations indicate that summer events were more clearly thermally preconditioned in the early stage, while wind-speed and wind-shear enhancement became more evident closer to onset. This station-scale pattern is consistent with the ERA5-based interpretation and shows how regional dynamic and thermal anomalies were expressed in the near-surface layer.

4 Discussion

The HYSPLIT pathway analysis, ERA5 diagnostics, and tower observations provide complementary evidence for interpreting dust-storm development over the Taklimakan Desert. HYSPLIT identifies the pre-storm airflow pathways reaching the observation sites, ERA5 explains the regional pressure, circulation, and thermal backgrounds behind these pathways, and the tower observations show how these regional conditions are expressed in near-surface wind, temperature, wind shear, and vertical profiles.

The pathway types are closely related to the ERA5 circulation patterns. East-inflow events are generally accompanied by clearer pressure differences between northern Xinjiang, surrounding high-pressure regions, and the Tarim Basin, together with organized 10 m winds entering the desert through the northeastern basin opening. Mountain-crossing events correspond more closely to lower-tropospheric circulation differences around the Tianshan–Pamir region, while the south-to-north pathway is related to pressure and wind configurations along the southern and southwestern margins of the desert. These results indicate that the transport pathways were not isolated trajectory features, but were linked to regional dynamic backgrounds and basin topographic constraints.

ERA5 further reveals a seasonal shift in the pre-onset conditions. Spring events show stronger and more persistent mean sea-level pressure-gradient and 850 hPa height-gradient anomalies, consistent with the more distinct dynamic pathways. Summer events show weaker pressure-gradient organization but stronger early thermal anomalies, including enhanced net shortwave radiation, surface–air temperature difference, and boundary-layer height. This suggests that spring dust storms were more closely related to regional dynamic forcing, whereas summer events were more strongly influenced by early thermal preconditioning.

The tower observations provide station-scale support for this interpretation. Summer events show more continuous vertical warming, while wind-speed and wind-shear anomalies strengthen closer to onset, indicating that thermal preconditioning alone is insufficient and that near-surface momentum enhancement remains necessary. The differences between TZ and XT further suggest that local terrain can modify the near-surface response to regional forcing. XT shows more regular vertical structures, whereas TZ exhibits more variable profiles, likely related to dune-induced flow disturbance and heterogeneous roughness.

Several uncertainties remain. The analysis is based on eight events from one field season, and the identified pathway and seasonal characteristics need to be tested using longer-term samples. In addition, the lack of direct turbulence, vertical velocity, and lidar observations limits the interpretation of vertical transport processes. Future studies should combine multi-year observations, turbulence measurements, lidar data, and high-resolution simulations to better distinguish the roles of synoptic forcing, thermal instability, and local terrain heterogeneity.

5 Conclusion

This study examined eight complete dust-storm events over the Taklimakan Desert using HYSPLIT trajectories, ERA5 reanalysis, and dual-gradient tower observations. The main conclusions are as follows.

The analyzed events exhibited three main types of pre-storm airflow transport pathways: east-inflow transport through the northeastern opening of the Tarim Basin, mountain-crossing transport across the Tianshan–Pamir region, and south-to-north transport related to airflows crossing the Tibetan Plateau and Kunlun Mountains.

ERA5 analysis shows that these pathways were closely linked to regional dynamic and thermal backgrounds. Spring events were associated with stronger pressure-gradient and 850 hPa height-gradient anomalies, corresponding to more distinct dynamically driven pathways. Summer events showed weaker pressure-gradient signals but stronger early thermal anomalies, indicating more evident thermal preconditioning before onset.

The tower observations were consistent with the ERA5-based interpretation. Summer events showed stronger positive temperature anomalies and more continuous vertical warming, while wind-speed and wind-shear anomalies strengthened closer to onset. This indicates a staged process from early background adjustment to near-surface wind enhancement.

The two stations showed different vertical responses. XT displayed more regular wind and temperature anomaly structures, whereas TZ showed more variable profiles, suggesting that local terrain heterogeneity may modulate the near-surface response to regional dynamic and thermal forcing.

Overall, Taklimakan dust-storm development is shaped by the interaction among regional airflow pathways, dynamic–thermal background conditions, near-surface wind response, and local terrain effects.

Acknowledgements

We gratefully acknowledge the National Earth System Science Data Center (<http://www.geodata.cn>) and the Institute of Desert Meteorology (China Meteorological Administration, Urumqi) for providing essential datasets. We also extend our thanks to the College of Geography and Remote Sensing Science, Xinjiang University (Urumqi), for their technical support.

Author contributions

Y.W. : Conceptualization , Writing – original draft , Formal Analysis.

WH.:Methodology, Supervision, Writing - review. & editing.

YL.:Validation.

M.M : Investigation, Data Curation.
F.Y: Experimental Design, Supervision.
C.Z..X.Y.A.M.: Investigation , Data Curation.

Financial support

This research was funded by the Xinjiang Uygur Autonomous Region Natural Science Foundation for Outstanding Young Scientists (Grant No. 2025D01E64), the Tianshan Talent Project of Xinjiang (Grant No. 2023TSYCCX0075), the Key Project of the Innovation and Development Special Program, CMA(CXFZ2026J007), the China Meteorological Administration's Young Innovative Team Project (Project No. CMA2024QN13), and the Supported by S&T Development Fund of CAMS (2021KJ034) .

Data availability

The meteorological observational data used in this study are maintained and owned by the Institute of Desert Meteorology, China Meteorological Administration, located in Urumqi, Xinjiang, China, and are subject to institutional regulations and access restrictions. ERA5 reanalysis datasets were obtained from the European Centre for Medium-Range Weather Forecasts (ECMWF) through the Copernicus Climate Data Store (<https://cds.climate.copernicus.eu>). Researchers or interested parties who wish to obtain access to the observational datasets may submit a formal request to the Institute of Desert Meteorology, outlining the purpose and scope of their intended use. Approval of such requests will be at the discretion of the institute, in accordance with its data-sharing policies.

Competing interests

The authors declare no competing interests.

References

Achakulwisut, P., Mickley, L. J., and Anenberg, S. C.: Drought-sensitivity of fine dust in the US Southwest: Implications for air quality and public health under future climate change, *Environ. Res. Lett.*, 13, 054025, <https://doi.org/10.1088/1748-9326/aabf20>, 2018.

Aili, A., Xu, H., Kasim, T., and Abulikemu, A.: Origin and Transport Pathway of Dust Storm and Its Contribution to Particulate Air Pollution in Northeast Edge of

Taklimakan Desert, China, Atmosphere, 12, 113,
<https://doi.org/10.3390/atmos12010113>, 2021.

Aili, A., Xu, H., Xu, Q., and Liu, K.: Aeolian dust movement and deposition under
local atmospheric circulation in a desert-oasis transition zone of the northeastern
Taklimakan desert, Ecological Indicators, 157, 111289,
<https://doi.org/10.1016/j.ecolind.2023.111289>, 2023.

Aili, A., Waheed, A., Zhao, X., and Xu, H.: Establishing an Early Warning System
for Dust Storms in Peri-Desert Regions, Environments, 11, 61,
<https://doi.org/10.3390/environments11040061>, 2024.

Bao, C. and Fang, C.: Water resources constraint force on urbanization in water
deficient regions: A case study of the Hexi Corridor, arid area of NW China, Ecological
Economics, 62, 508–517, <https://doi.org/10.1016/j.ecolecon.2006.07.013>, 2007.

Bao, T., Xi, G., Hao, Y., Chang, I.-S., Wu, J., Xue, Z., Jin, E., Zhang, W., and Bao,
Y.: The Transport Path and Vertical Structure of Dust Storms in East Asia and the
Impacts on Cities in Northern China, Remote Sensing,
<https://doi.org/10.3390/rs15123183>, 2023.

Bi, J., Li, Z., Zuo, D., Yang, F., Li, B., Ma, J., Huang, Z., and He, Q.: Dust Aerosol
Vertical Profiles in the Hinterland of Taklimakan Desert During Summer 2019, Front.
Environ. Sci., 10, 851915, <https://doi.org/10.3389/fenvs.2022.851915>, 2022.

Chen, S., Zhao, D., Huang, J., He, J., Chen, Y., Chen, J., Bi, H., Lou, G., Du, S.,
Zhang, Y., and Yang, F.: Mongolia Contributed More than 42% of the Dust
Concentrations in Northern China in March and April 2023, Adv. Atmos. Sci., 40,
1549–1557, <https://doi.org/10.1007/s00376-023-3062-1>, 2023a.

Chen, Y., An, J., Qu, Y., Xie, F., and Ma, S.: Dust radiation effect on the weather
and dust transport over the Taklimakan Desert, China, Atmospheric Research, 284,
106600, <https://doi.org/10.1016/j.atmosres.2022.106600>, 2023b.

Gao, X., Wang, D., Gui, H., and et al: Dominant circulation pattern and moving path of the Mongolian Cyclone for the severe sand and dust storm in China, Atmospheric Research, <https://doi.org/10.1016/j.atmosres.2024.107272>, 2024.

Ge, J. M., Huang, J. P., Xu, C. P., Qi, Y. L., and Liu, H. Y.: Characteristics of Taklimakan dust emission and distribution: A satellite and reanalysis field perspective: Taklimakan dust Characteristics, J. Geophys. Res. Atmos., 119, 11,772-11,783, <https://doi.org/10.1002/2014JD022280>, 2014.

Gillette, D. A. and Passi, R.: Modeling dust emission caused by wind erosion, J. Geophys. Res., 93, 14233–14242, <https://doi.org/10.1029/JD093iD11p14233>, 1988.

Huo, W., Song, M., Wu, Y., Zhi, X., Yang, F., Ma, M., Zhou, C., Yang, X., Mamtimin, A., and He, Q.: Relationships between Near-Surface Horizontal Dust Fluxes and Dust Depositions at the Centre and Edge of the Taklamakan Desert, Land, 11, 959, <https://doi.org/10.3390/land11070959>, 2022.

Iversen, J. D. and White, B. R.: Saltation threshold on Earth, Mars and Venus, Sedimentology, 29, 111–119, <https://doi.org/10.1111/j.1365-3091.1982.tb01713.x>, 1982.

Kai, Z. and Huiwang, G.: The characteristics of Asian-dust storms during 2000–2002: From the source to the sea, Atmospheric Environment, 41, 9136–9145, <https://doi.org/10.1016/j.atmosenv.2007.08.007>, 2007.

Kok, J. F., Storelvmo, T., Karydis, V. A., Adebiyi, A. A., Mahowald, N. M., Evan, A. T., He, C., and Leung, D. M.: Mineral Dust Aerosol Impacts on Global Climate and Climate Change, Nat. Rev. Earth Environ., 4, 71–86, <https://doi.org/10.1038/s43017-022-00379-5>, 2023.

Li, J., Han, Y., Liu, W., Cao, L., and Lu, Z.: Contributions of gust under stationery weather conditions to the dust aerosol budget in northwestern China, Atmospheric Research, 293, 106914, <https://doi.org/10.1016/j.atmosres.2023.106914>, 2023.

Liu, X., Yin, Z., Zhang, X., and Yang, X.: Analyses of the spring dust storm frequency of northern China in relation to antecedent and concurrent wind, precipitation, vegetation, and soil moisture conditions, *J. Geophys. Res.*, 109, 2004JD004615, <https://doi.org/10.1029/2004JD004615>, 2004.

Manisalidis, I., Stavropoulou, E., Stavropoulos, A., and Bezirtzoglou, E.: Environmental and Health Impacts of Air Pollution: A Review, *Front. Public Health*, 8, 14, <https://doi.org/10.3389/fpubh.2020.00014>, 2020.

Marticorena, B. and Bergametti, G.: Modeling the atmospheric dust cycle: 1. Design of a soil-derived dust emission scheme, *J. Geophys. Res.*, 100, 16415–16430, <https://doi.org/10.1029/95JD00690>, 1995.

Mu, F. and Fiedler, S.: How much do atmospheric depressions and Mongolian cyclones contribute to spring dust activities in East Asia?, *npj Clim Atmos Sci*, 8, 51, <https://doi.org/10.1038/s41612-025-00929-w>, 2025.

Prospero, J. M.: Long-range transport of mineral dust in the global atmosphere: Impact of African dust on the environment of the southeastern United States, *Proc. Natl. Acad. Sci. U.S.A.*, 96, 3396–3403, <https://doi.org/10.1073/pnas.96.7.3396>, 1999.

Raupach, M. R.: Drag and drag partition on rough surfaces, *Boundary-Layer Meteorol*, 60, 375–395, <https://doi.org/10.1007/BF00155203>, 1992.

Shao, Y. and Dong, C. H.: A review on East Asian dust storm climate, modelling and monitoring, *Global and Planetary Change*, 52, 1–22, <https://doi.org/10.1016/j.gloplacha.2006.02.011>, 2006.

Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.

Su, L., Lu, C., Yuan, J., Wang, X., He, Q., and Xia, H.: Measurement report: The promotion of low-level jet and thermal-effect on development of deep convective boundary layer at the southern edge of the Taklimakan Desert, <https://doi.org/10.5194/egusphere-2024-1010>, 15 May 2024.

Sun, J. and Liu, T.: The Age of the Taklimakan Desert, *Science*, 312, 1621–1621, <https://doi.org/10.1126/science.1124616>, 2006.

Tong, D., Feng, I., Gill, T. E., Schepanski, K., and Wang, J.: How Many People Were Killed by Windblown Dust Events in the United States?, *Bulletin of the American Meteorological Society*, 104, E1067–E1084, <https://doi.org/10.1175/BAMS-D-22-0186.1>, 2023.

Wang, B., Ji, H., Zhang, Z., Liang, J., Zhang, L., Li, M., Qiu, R., Luo, H., An, W., Tian, P., and Amonov, M. O.: Surface-Dependent Meteorological Responses to a Taklimakan Dust Event During Summer near the Northern Slope of the Tibetan Plateau, *Remote Sensing*, 17, 1561, <https://doi.org/10.3390/rs17091561>, 2025.

Wang, H., Jia, X., Li, K., and Li, Y.: Horizontal wind erosion flux and potential dust emission in arid and semiarid regions of China: A major source area for East Asia dust storms, *CATENA*, 133, 373–384, <https://doi.org/10.1016/j.catena.2015.06.011>, 2015.

Wang, X., Zhai, P., and Wang, C.: Variations in extratropical cyclone activity in northern East Asia, *Adv. Atmos. Sci.*, 26, 471–479, <https://doi.org/10.1007/s00376-009-0471-8>, 2009.

Wang, Y., Gao, J., Mamtimin, A., Sayit, H., Zhou, C., Li, R., Dawut, M., Yang, F., Huo, W., Wen, C., Song, M., and Aihaiti, A.: Evolution law of atmospheric boundary layer in Gurbantünggüt Desert based on reanalysis dataset and in situ observation data, *Heliyon*, 9, e14147, <https://doi.org/10.1016/j.heliyon.2023.e14147>, 2023.

Wen, H.-J., Wang, S.-L., Wu, C.-D., and Liang, M.-C.: Association of Asian dust storms and PM_{2.5} with clinical visits for respiratory diseases in children, *Atmospheric Environment*, 333, 120631, <https://doi.org/10.1016/j.atmosenv.2024.120631>, 2024.

Xing, Y., Chen, Y., Yan, S., Cao, X., Zhou, Y., Zhang, X., Shi, T., Niu, X., Wu, D., Cui, J., Zhou, Y., Wang, X., and Pu, W.: Dust storms from the Taklamakan Desert significantly darken snow surface on surrounding mountains, *Atmos. Chem. Phys.*, 24, 5199–5219, <https://doi.org/10.5194/acp-24-5199-2024>, 2024.

Xiong, Z., Xu, X., Yang, Y., and Luo, T.: Diurnal vertical distribution and transport of dust aerosol over and around Tibetan Plateau from lidar on International Space Station, *Atmospheric Research*, 294, 106939, <https://doi.org/10.1016/j.atmosres.2023.106939>, 2023.

Xu, C., Guan, Q., Lin, J., Luo, H., Yang, L., Tan, Z., Wang, Q., Wang, N., and Tian, J.: Spatiotemporal variations and driving factors of dust storm events in northern China based on high-temporal-resolution analysis of meteorological data (1960–2007), *Environmental Pollution*, 260, 114084, <https://doi.org/10.1016/j.envpol.2020.114084>, 2020.

Yang, X., Shen, S., Yang, F., He, Q., Ali, M., Huo, W., and Liu, X.: Spatial and temporal variations of blowing dust events in the Taklimakan Desert, *Theor Appl Climatol*, 125, 669–677, <https://doi.org/10.1007/s00704-015-1537-4>, 2016.

Yuan, T., Chen, S., Huang, J., Wu, D., Lu, H., Zhang, G., Ma, X., Chen, Z., Luo, Y., and Ma, X.: Influence of Dynamic and Thermal Forcing on the Meridional Transport of Taklimakan Desert Dust in Spring and Summer, *J. Climate*, 32, 749–767, <https://doi.org/10.1175/JCLI-D-18-0361.1>, 2019.

Zeng, X. and Dickinson, R. E.: Effect of Surface Sublayer on Surface Skin Temperature and Fluxes, *J. Climate*, 11, 537–550, [https://doi.org/10.1175/1520-0442\(1998\)011%3C0537:EOSSOS%3E2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011%3C0537:EOSSOS%3E2.0.CO;2), 1998.

581 Zhang, H., Cisse, M., Dauphin, Y. N., and Lopez-Paz, D.: mixup: Beyond
582 Empirical Risk Minimization, <https://doi.org/10.48550/ARXIV.1710.09412>, 2017.

583 Zhang, L., Zhang, H., Cai, X., Song, Y., Mamtimin, A., and He, Q.: Physical
584 Mechanisms of Deep Convective Boundary Layer Leading to Dust Emission in the
585 Taklimakan Desert, *GEOPHYSICAL RESEARCH LETTERS*, 51, e2024GL108521,
586 <https://doi.org/10.1029/2024gl108521>, 2024.

587 Zhang, X., Wang, T., Wang, S., Jiao, Y., Tang, J., Li, J., Yang, F., Amonov, M.
588 O., and Abdullaev, S. F.: Conductive circulation patterns and transport mechanisms for
589 spring dust from Taklimakan Desert to the Tibetan Plateau, *Environment International*,
590 197, 109356, <https://doi.org/10.1016/j.envint.2025.109356>, 2025.

591