

1   **The NASA-GISS ModelE2.1-CC2 ESM: development and simulations**

2   Anastasia Romanou<sup>1,2</sup>, Paul Lerner<sup>1,2</sup>, Nancy Y. Kiang<sup>1,4</sup>, Igor Aleinov<sup>4</sup>, Maxwell Kelley<sup>1</sup>, Roland L. Miller<sup>1</sup>, Gary Russell<sup>1</sup>, Reto  
3   Ruedy<sup>3</sup>, Gavin A. Schmidt<sup>1</sup>, Maria Z. Hakuba<sup>5</sup>, Ou Wang<sup>5</sup>

4   <sup>1</sup>*NASA-Goddard Institute for Space Studies, New York City, NY 10025, USA*

5   <sup>2</sup>*Dept. of Applied Phys. and Applied Math., Columbia University, New York, NY 10025, USA*

6   <sup>3</sup>*Autonomic Integra LLC, New York, NY 10025, USA*

7   <sup>4</sup>*Climate Systems Research, Columbia University, New York, NY 10025, USA*

8   <sup>5</sup>*Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA*

9

10   Correspondence to: Anastasia Romanou ([anastasia.romanou@nasa.gov](mailto:anastasia.romanou@nasa.gov))

11   **Abstract.** This paper describes the NASA Goddard Institute for Space Studies (GISS) ModelE2.1-Carbon  
12   Cycle version 2 Earth System Model (NASA GISS E2.1-CC2 ESM), assesses its skill against observations  
13   and reports on its performance under future climate scenarios. The first version of GISS ESM (GISS-E2.1-  
14   G-CC) was introduced in CMIP6, and therefore ModelE2.1-CC2 is a post CMIP6 release that includes the  
15   same physical climate model E2.1-G, updated ocean and land carbon cycle and longer equilibrium  
16   simulations. While the focus here is on the land and ocean carbon components and their interactions  
17   with atmosphere and ice, we also describe in detail the physical coupled model for consistency and ease  
18   of reference. We detail parameterizations, tuning, and conservation diagnostics that are relevant to the  
19   global (land & ocean) carbon cycle. We also describe a full suite of simulations performed with this  
20   model, including pre-industrial control, historical, and future climate scenarios. The model is an  
21   improvement to prior releases, has been better tuned and exhibits smaller drifts. However some  
22   persistent biases remain, particularly with regards to low ocean and land primary production. Results  
23   shown here are compared to previous versions of the model, as well as the state-of-the art in Earth  
24   system modeling via community intercomparisons. ADD MORE TEXT JUSTIFICATION


Commented [AR1]: removed part on future simulations

25   **1 Introduction**

26   Earth system models (ESMs) integrate the interactions of atmosphere, ocean, land and ice with the  
27   biosphere to estimate the state of regional and global climate under a wide variety of conditions  
28   (Heavens et al., 2013) and focus on comprehensiveness of Earth system interactions, feedbacks and  
29   interdependencies. ESMs include all the processes present in climate models but they also simulate  
30   the interaction between the physical climate, the biosphere and the chemical constituents of the

31 atmosphere and the ocean (Fig. 1). They can also include the impact of human decision-making, e.g.  
32 changes in land use, in emissions mitigation etc. They are thus particularly useful for estimating  
33 remaining carbon budgets before anthropogenic warming exceeds certain thresholds, studying the  
34 feasibility, scalability and impacts of climate intervention practices and geoengineering activities, such  
35 as carbon dioxide removal (CDR) through soil carbon sequestration, biomass carbon removal and  
36 storage, enhanced mineralization, ocean alkalinity enhancement, afforestation/reforestation etc and  
37 studies of the Earth's climate evolution after curbing emissions.

38 The NASA Goddard Institute for Space Studies (NASA-GISS) ModelE version 2.1 with Carbon Cycle  
39 version 2 (ModelE2.1-CC2) is an Earth system model with a fully interactive land and ocean carbon  
40 cycle which is an update of the GISS submission to the 6th generation Climate Model Intercomparison  
41 Project (CMIP6) which was identified as GISS-E2-1-CC in the CMIP6 repository. We will use the  
42 shorthand GISS-ESM CMIP6+ alternatively with GISS-E2-1-CC2 to denote the difference from the  
43 previous model.



44  
45 **Figure 1. Schematic diagram of the Earth as a complex interrelated system. ESMs strive to include as many of the**  
46 **components and their interactions as possible. Image credit: Kevin W. Miller (Global Science and Technology, Inc.)**  
47 <https://svs.gsfc.nasa.gov/30988>

48 In the following sections we describe as comprehensively as possible the physical and biogeochemical  
49 components of the model, the relevant parts of the radiation, atmospheric chemistry and clouds. We  
50 then present results from the spinups and historical simulations, and assess them against  
51 observations. Lastly we lay out the model response to future emission and mitigation pathways. Lastly

Commented [AR2]: removed

Commented [AR2R2]: removed



we offer a summary of the model, comparison with other models across different experiments in the post-CMIP6 assessments and lay out our future modeling plans.

## 2 Model Description

The underlying physical model, ModelE2.1, has been described in Kelley et al. (2020) and includes updates to earlier versions of the NASA GISS model (Schmidt et al. 2006; Hansen et al. 2007; Schmidt et al. 2014). For ease of access and comprehensiveness of model presentation, we will describe key components of the model here as well, although complete information and evaluations of the physical model should be sought in these publications.

### 2.1 Model configuration

The atmospheric model has horizontal resolution of 2 by 2.5 degrees in latitude and longitude respectively and 40 layers in the vertical from the surface to 0.1 hPa in the lower mesosphere; it is coupled to an ocean with 1 by 1.25 horizontal resolution in latitude and longitude respectively, and has also 40 layers in the vertical.

Atmospheric dynamics uses Arakawa's B-grid scheme where eastward and northward velocity components are located at primary cell corners; vertical discretization follows terrain-following (sigma) layers in the troposphere which transition to constant-pressure layers in the stratosphere. The transition is smooth and centered at 100 hPa with a half-width of approximately 30 hPa. The vertical discretization in the ocean is 40 vertical layers in mass with higher resolution of 5-20m near the surface and within the thermocline and about 200 m at depth.

Prognostic variables in the atmosphere are dry air mass, potential temperature, water vapor mixing ratio, and horizontal velocity components. Virtual potential temperature is used for all density/buoyancy-related calculations. Ocean prognostic variables include seawater mass, potential enthalpy that accounts for variations in specific heat capacity, salt and horizontal velocity components on both the C-grid (perpendicular to primary cell edges) and D-grid (parallel to primary cell edges).

Subsurface reservoirs are split into four types: open water (including lakes and oceans), ice-covered water (including lake ice and sea ice), ground (including bare soil and vegetated regions), and land ice/glaciers. Within each type there are further subdivisions (fraction of plant functional types, fractional snow cover, melt pond fraction over sea ice, etc.), which are communicated to the atmospheric model

83 as weighted mean quantities (for example the albedo in the radiation computations). The non-ocean  
84 reservoirs include prognostic variables for water mass and heat content, and the sea ice also includes  
85 salt content.

86  
87 The model uses a 30-min time step for physics calculations (i.e. surface interactions, condensation etc).  
88 The radiation code is called every five physics time steps (every 2.5 h), however, the zenith angle and  
89 sun light intensity are updated every physics time step. The model uses a 225 s time step for the  
90 atmospheric dynamics (transports) and 112.5 s time step for ocean gravity wave computations.

91  
92 Simplifications in atmospheric thermodynamics ignore the sensible heat of condensate. Although latent  
93 heat of water vapor is treated correctly, its mass is not added to atmospheric dry mass. Consequently,  
94 precipitation and evaporation occur at 0°C, specific potential enthalpy equals potential temperature  
95 times dry specific heat capacity, and humidity affects atmospheric density only in the pressure gradient  
96 force computation.

## 97 **2.2 The ocean component**

98 The ocean model is a dynamic ocean model E2.1-G (Russell et al. 1995; Schmidt et al. 2006; 2014;  
99 Kelley et al. 2020), non-Boussinesq (i.e. includes density variations in the horizontal as well as vertical  
00 momentum equations and in the continuity equation, following Schmidt et al., 2006). It conserves  
01 potential enthalpy, freshwater mass, and salt mass, and uses natural boundary conditions (for heat  
02 and freshwater). It has a free surface and thus it is not volume conserving. The number of vertical  
03 layers varies from 2 to 40. After the dynamics, the mass ratio of layer n and its layer above is fixed in  
04 time and space, but this does vary with n. The thickness of layer 1 varies and is approximately 10 m.  
05 The vertical coordinate is mass per unit area, including the mass of air, ice and water above any level.

06  
07 The ocean model solves the dynamical equations via an invariant form. The Coriolis term is computed  
08 via the C and D-grid velocities and has latitude dependence. The quadratic upstream advection scheme  
09 (Prather, 1986) uses mean, three-dimensional gradients, and second order moments to advect scalar  
10 tracers, thus effectively increasing the resolution, in both horizontal and vertical direction. Bottom  
11 friction is applied at the lowest layer of each ocean column. The ocean model employs the K-profile  
12 parameterization (KPP) for vertical mixing (Large et al., 1994) and the Gent-McWilliams  
13 parameterization for eddy induced tracer transports (Griffies 1998a; 1998b). The mesoscale diffusivity  
14 allows for surface-intensified eddies, by employing an exponential decay of diffusivity with depth that  
15 depends locally on the horizontal gradient of potential density (Kelley et al., 2020) which permits eddies

16 to restratify the Southern Ocean over a large depth range, consistent with the observed density  
17 structure there, while not overacting in other regions of the World Ocean (such as the North Atlantic).  
18 The Rossby radius is replaced by a geographically constant nominal length scale and the diffusivity is a  
19 function of the Brunt-Vaisala frequency and the slope of isopycnal surfaces.

20  
21 Mesoscale transport is expressed in local quasi-isopycnal layers, avoiding difficulties associated with  
22 skew-flux representation 1) in regions with weak vertical stratification, e.g. the surface mixed layer and  
23 at the ocean surface, where the GM parameterization assumes that eddies transport tracers along  
24 isopycnal surfaces. However, near the surface, these surfaces can be highly tilted or intersect the ocean  
25 surface, which violates the assumptions of the parameterization. Applying the GM scheme directly in  
26 these regions can lead to spurious mixing and incorrect representation of processes like deep convection  
27 and mixed-layer re-stratification, 2) near lateral boundaries and the ocean bottom, where the eddy  
28 fluxes must be parallel to these boundaries, but the GM scheme's isopycnal-following nature doesn't  
29 naturally enforce this, and 3) in areas of strong density fronts (e.g., the Gulf Stream or the Antarctic  
30 Circumpolar Current) where the "small-slope approximation" which assumes that the slope of isopycnal  
31 surfaces is small and is valid in the deep ocean, breaks down in areas where the isopycnals are steeply  
32 tilted. Ventilation of marginal seas through their connecting straits has been increased via two  
33 mechanisms in E2.1-G, reducing salinity biases there. While there is no explicit overflow  
34 parameterization in the ocean model, for straits deep enough that density contrasts can drive strong  
35 opposing flows at the surface and at depth, the finer upper-ocean layering in E2.1-G resolves this  
36 structure, in conjunction with a slight tuning of strait depths. Second, horizontal diffusivity was increased  
37 in straits that are shallow or have weaker density contrasts. The first mechanism impacted the Red and  
38 Black seas, and the second the Baltic and Hudson (Kelley et al., 2020). The first is the sole ventilation  
39 mechanism for straits narrower than the nominal resolution, which are parameterized using the Russell  
40 et al. (1995) one-dimensional channel scheme that lacks horizontal mixing.


41  
42 In addition to KPP, the vertical diffusivity also includes a contribution from tidal dissipation. The ocean  
43 model Atlantic meridional overturning circulation (AMOC) is found to be sensitive to tidally driven  
44 mixing, which occurs in the shallow waters bordering the North Atlantic following dissipation  
45 distributions generated by Jayne (2009).

46  
47 The Equation of State employed in the ocean model computes specific volume ( $\text{m}^3/\text{kg}$ ) as a function of  
48 specific potential enthalpy ( $\text{J/kg}$ ), salinity (psu), and pressure (Pa). Using constant uniform gravity,  
49 pressure at any level is gravity times mass per unit area above the level. The one-to-one relationship

50 between temperature and specific potential enthalpy is computed using the adiabatic lapse rate of sea  
51 water as function of temperature, salinity and pressure (IES-1980) and specific enthalpy of sea water  
52 as function of temperature and salinity at atmospheric pressure. Using these formulas, the model's  
53 Equation of State can be computed from the specific volume function of temperature, salinity and  
54 pressure (IES-1980). Complex computations for Equation of State, freezing temperature, specific heat  
55 capacity, and others are evaluated offline and are stored in tables; the model interpolates the table  
56 values as functions of specific potential enthalpy, salinity and pressure (Chen and Millero, 1976). Table  
57 resolutions are 4000 (J/kg) of specific potential enthalpy, 1 (psu) of salinity, and  $2 \times 10^6$  (Pa) of pressure.  
58 Under adiabatic flow, potential enthalpy is a conserved quantity, therefore the ocean model conserves  
59 energy (heat) as well as mass and salt.

#### 61 **Ocean Carbon Cycle**

62 The ocean carbon cycle model is based on the NASA Ocean Biogeochemistry Model (NOBM), originally  
63 developed at the NASA-Global Modeling and Assimilation Office (Gregg and Casey 2007) and later  
64 coupled coupled to the GISS climate model under prescribed atmospheric concentrations (Romanou et  
65 al., 2013; 2014) and under prescribed emissions (Ito et al, 2020). The model includes four phytoplankton  
66 species (diatoms, chlorophytes, cyanobacteria, and coccolithophores), four nutrient species (nitrate,  
67 silicate, ammonia, and iron), three detrital pools (nitrate/carbon, silicate, and iron) and one heterotroph  
68 species (Fig. 2). Carbon cycling is represented through dissolved organic (DOC) and inorganic carbon  
69 (DIC), where carbonate chemistry is solved using MOCSY (Orr and Epitalon, 2015). Ocean carbon  
70 interacts with atmospheric CO<sub>2</sub> through gas exchange parameterization, following the CMIP6 protocol  
71 (Orr et al., 2017). The current version of NOBM (NOBMg) has been expanded to include oxygen and  
72 alkalinity (Lerner et al., 2021; 2024) and an updated treatment of iron scavenging (see below). The  
73 model equations and parameters used are laid out in Appendix A and in Tables A1-A11 therein. A  
74 schematic diagram of the NOBMg and how it is linked to other components of ModelE is shown in Fig.  
75 3. A short description of the model equations is provided below.




Commented [AR3]: removed

Figure 2. Schematic diagram of the ocean carbon cycle model (NOBMg). credit: Jay Friedlander, NASA-GSFC.

All tracers in NOBMg are changed via physical transport and mixing which are following parameterizations of these processes in the physical ocean model.

Phytoplankton species are also subject to sinking, growth via nutrient uptake, respiration and exudation of DOC, senescence and zooplankton grazing. Nutrient distributions are changed via uptake by phytoplankton species for growth, remineralization of particulate organic carbon or nitrogen, and detrital breakdown. Cyanobacteria can also grow via nitrogen fixation, where growth is limited only by light, iron, and temperature. Nitrogen assimilated into organic matter is obtained from  $N_2$ . Cyanobacterial growth via nitrogen fixation follows the same Michaelis-Menten formulation as cyanobacterial growth that is nitrate limited, with two exceptions. First, the iron half-saturation constant for nitrogen fixation is 33% higher than that for nitrate-limited growth, as nitrogen fixation generally relies upon an increased availability of iron. Second, nitrogen fixation is inhibited by the presence of nitrate and ammonium, following laboratory culture experiments using *Trichodesmium* (Holl and Montoya, 2005). Ammonium is produced via the herbivore grazing and heterotrophic loss. Silicate is taken up only by diatoms. Iron is produced by zooplankton, atmospheric deposition of dust, and is lost to scavenging. NOBMg also includes an updated marine iron cycle, with regionally varying prescribed ligands, and a significant increase in the rate of ligand-bound iron scavenging, to be consistent with observed rates for trace metals with similar scavenging characteristics as iron (see section 3.1.1 on Tunings). At each time step, the prognostic iron pool is split into iron that is complexed by organic ligands

98 and a “free” iron pool that represents the sum of all inorganic iron species. Iron complexation chemistry  
 99 is computed at each time step using vertical and regionally varying prescribed ligand concentrations (see  
 100 section 3.1.2 Forcings) and a conditional stability constant ( $K_{FeL} = 10^{11}$  L/mol. Then, free and ligand-  
 101 complexed iron are computed by solving the system of equations for total iron, total ligand  
 102 concentration, and the conditional stability constant (Aumont et al., 2015).



Commented [AR4]: replaced with one single new figure that describes the ocean carbon cycle

103  
 104 **Figure 3. Schematic diagram of the NOBM in GISS ModelE2.1-CC2**

105  
 106 Herbivores grow via grazing of phytoplankton and are lost to heterotrophic consumption and excretion  
 107 to DOC.

108  
 109 Carbon and nitrate particulates (detritus) are linked via the constant C:N ratio. Therefore only one  
 110 equation is used to describe both. Their change is due to advection and diffusion, sinking,  
 111 remineralization and detrital breakdown. Sinking of particulates is expressed via an exponential function  
 112 of depth (see equation A15-A17). Silicate detritus includes an additional source term due to senescence  
 113 of phytoplankton and grazing from herbivores. The iron detritus includes zooplankton production and  
 114 scavenging production. DOC is produced during phytoplankton growth, herbivore excretion and detrital  
 115 breakdown while the only DOC sink is bacterial degradation. DIC sources are phytoplankton and  
 116 herbivore respiration, bacterial degradation of DOC, remineralization of carbon detritus and air sea  
 117 exchange while the only sink is the growth of phytoplankton.

118  
 119 The model includes prognostic alkalinity which is advected and mixed like all other tracers and follows  
 120 the Ocean Carbon Model Intercomparison Project parameterizations (OCMIP-2, Najjar et al., 2007), i.e.

21 it is produced via conversion of nitrate to organic N and carbonate dissolution, while it is lost to  
22 remineralization of detrital organic carbon/nitrogen and DOC and carbonate precipitation (see eq. A43-  
23 A47).

24  
25 Oxygen interacts with the rest of the ocean biogeochemical tracers via remineralization of organic  
26 carbon and by determining when the model switches to slower rates of remineralization that are  
27 consistent with denitrification (eq. A48-A56; Lerner et al. 2024).

## 28 **2.4 The Land component**

### 29 **Vegetation**

30 The Ent Terrestrial Biosphere Model (Ent TBM) (<https://www.giss.nasa.gov/projects/ent/>) is a demographic  
31 dynamic global vegetation model (DGVM) (Kiang, et al. 2006; Kim et al., 2015; Kiang et al., in prep.)  
32 with subgrid vegetation patches distinguished by plant functional type (PFT), currently with single-  
33 cohort structure for patch communities (Fig. 4). The 12 plant functional types are: evergreen  
34 broadleaf trees, evergreen needleleaf trees, cold deciduous broadleaf trees, drought deciduous  
35 broadleaf trees, deciduous needleleaf trees, cold-adapted shrubs, arid-adapted shrubs, C3 perennial  
36 grass, C3 annual grass, C3 arctic grass, C4 grass, and herb crops (with physics of C3 perennial grass).  
37 Soil biogeochemistry is a version of the Carnegie-Stanford-Ames' (CASA') model (Potter et al. 1996;  
38 Randerson et al. 2009; Doney et al. 2006), modified for Ent for an improved response of soil respiration  
39 to soil moisture, as described in Kim et al. (2015). Land ecosystem carbon pools that are calculated  
40 are: foliage, hardwood stem, sapwood stem, fine roots, coarse roots, non-structural carbohydrate  
41 (which we also call labile carbon), and 9 soil carbon pools.

42  
43 Leaf biophysics is a coupled conductance/photosynthesis model that simulates vegetation stomatal  
44 control of both transpiration and uptake of CO<sub>2</sub>. The coupled leaf photosynthesis/ stomatal  
45 conductance model consists of the photosynthesis model of Farquhar and von Caemmerer (1982) and  
46 the Ball-Berry model of stomatal conductance (Ball and Berry, 1987), with leaf boundary layer  
47 conductances following the approach by Collatz et al. (1991) but with the boundary layer conductance  
48 derived from the canopy surface layer conductance of the land surface model; for the cubic equations  
49 of coupled photosynthesis/conductance, a numerical solver for net assimilation of CO<sub>2</sub> and leaf  
50 internal CO<sub>2</sub> concentration was developed. C4 photosynthesis follows the scheme of Collatz and Berry  
51 (1992). Leaf-to-canopy scaling is achieved through the canopy radiative transfer scheme of Spitters et  
52 al. (1986) (introduced to ModelE in CMIP5 in Schmidt et al., 2006) combined with a numerical scheme  
53 for vertical layering of the canopy by converging the total canopy fluxes. Vegetation canopy

.54 absorbance and transmittance of photosynthetically active radiation thus depend on vertical layering  
.55 of leaf area index, solar zenith angle, and the fractions of diffuse vs. direct radiation, providing vertical  
.56 profiles of sunlit and shaded leaf area fractions.

.57  
.58 Physiological algorithms for water stress, cold-hardening in evergreen needleleaf trees, and  
.59 temperature sensitivity of photosynthetic capacity are detailed in Kim et al. (2015). To account for  
.60 variation in parameters of the leaf biophysics by climate and seasonality of vegetation, temperature  
.61 dependence of leaf photosynthetic capacity,  $V_{cmax}$ , is modeled with a Q10 function (Hegarty, 1973),  
.62 with a reference temperature of 25 C and parameters from Collatz et al. (1991). In addition,  
.63 photosynthetic capacity is subject to seasonal cold-hardening in evergreen needleleaf trees, as a  
.64 function of local air temperature trend, and is modeled based on the work of Repo et al. (1990);  
.65 Hanninen and Kramer (2007); and Makela, et al. (2004; 2006). Autotrophic maintenance respiration is  
.66 subject to temperature acclimation, in which the reference optimal temperature in the equation of  
.67 Levis et al. (2004) is allowed to vary between 10-30 C, tracking a 10-day running average of surface air  
.68 temperature; 10-day averages less than 10 C or greater than 30 C are considered outside the range of  
.69 acclimation.

.70  
.71 Water stress is modeled after the approach of Rodriguez-Iturbe et al. (2001), in which stress is a factor  
.72 between 0 (full stress) and 1 (unstressed), as a function of relative extractable water (volume water -  
.73 minus hygroscopic water)/(pore volume minus hygroscopic water). The factor modifies the leaf  
.74 stomatal conductance, and different plant functional types have different critical soil moisture values  
.75 for the onset of stress and wilting points. Because transpiration depends on the exposed leaf area  
.76 index, and wet leaf fractions will not transpire, a scheme for canopy interception of precipitation was  
.77 added following the algorithm presented in Koster and Suarez (1996) which is similar to the modified  
.78 Shuttleworth scheme of Wang et al. (2007). The scheme primarily involves modification of the wet  
.79 fraction of the canopy by computing the drip rate taking into account the fraction of precipitation that  
.80 falls onto a previously wetted portion of the canopy with the grid cell. This scheme also distinguishes  
.81 between stratiform and convective precipitation, because the stratiform type covers a greater portion  
.82 of the grid cell than the convective type (e.g., Koster and Suarez, 1996; Wang et al., 2007).

.83  
.84 For surface albedo, Ent's 12 different vegetation types have different spectral and snow masking  
.85 depth properties. In E2.1 Ent PFTs are mapped to the 8 biome types of Matthews (1984) for seasonal  
.86 albedo for radiative purposes.

.87



.88 In Schmidt et al. (2006) and henceforth, all surface conductance of water vapor for vegetated land was  
.89 assigned to the vegetation, which meant that there was no soil evaporation under canopies. The  
.90 rationale for this approach (Abramopoulos et al., 1988) was that vegetated land was simply a surface  
.91 with a particular albedo and conductance. Numerous studies have demonstrated the importance of  
.92 soil evaporation as a component of evapotranspiration, especially for semiarid ecosystems (e.g.,  
.93 Lawrence et al., 2007). While evaporation from bare soil was included in previous versions of the  
.94 model, in E2.1 we have added a simple representation of soil evaporation based on the work of Zeng  
.95 et al. (2005) and Lawrence et al. (2007), by including the evaporative flux from the vegetated soil into  
.96 the total evapotranspiration scaled as a function of the current leaf area index. Total  
.97 evapotranspiration never exceeds the potential evaporation.


#### .98 .99 **Irrigation and Groundwater**

.00 In E2.1, irrigation is implemented as a prognostic field (Cook et al., 2011; 2014; Krakauer et al., 2016;  
.01 Puma and Cook, 2010; Shukla et al., 2014) where water demand for irrigation (Wada et al., 2014) is  
.02 estimated using prescribed irrigation areal extent (Siebert et al., 2015). The water is drawn first from  
.03 rivers and lakes locally, and if that is insufficient, it is drawn from an external groundwater source  
.04 (which is tracked diagnostically). Groundwater is assumed to have the same temperature as the soil  
.05 and default tracer concentrations. Groundwater recharge is not accounted for, and so there is a small  
.06 increase in total water mass (and eventually, sea level) associated with the net global groundwater  
.07 draw in these simulations. These effects have a complex impact on freshwater delivery to the oceans  
.08 (and hence sea level). Irrigation from local surface water sources leads to increased soil moisture and  
.09 reduced river outflow, but this is dominated by net additions of groundwater which add freshwater to  
.10 the climate system, about 0.2 mm/yr of global sea-level-equivalent in 2010 (Miller et al., 2021). As  
.11 discussed in Miller et al. (2021) this freshwater imbalance is driven by an underestimate of surface  
.12 water in lakes and rivers (which means that more groundwater is needed) and also the lack of  
.13 counteracting increase in reservoirs.

#### .14 .15 **The Land Carbon Coupling**

.16 The land carbon dynamics in the fully coupled carbon cycle physics configuration is the same as  
.17 described in Ito et al. (2020) for the ModelE submission to CMIP6 C4MIP. The leaf biophysics is  
.18 sensitive to the surface atmospheric CO<sub>2</sub> concentration. The model tracks CO<sub>2</sub> exchanges between  
.19 the land and atmosphere in diagnostics for GPP, autotrophic respiration, soil respiration, and carbon  
.20 fluxes due to land cover change. Land carbon change associated with the annual land cover change is  
.21 distributed as CO<sub>2</sub> fluxes to the atmosphere over the following year to avoid having to add a pulse of  
.22 CO<sub>2</sub>.

23  
24



25  
26  
27

Figure 4. Schematic diagram of the Ent in GISS ModelE2.1-CC2

28 **2.6 The sea ice and land ice models**

29 Sea ice and lake ice are computed on the atmospheric grid and the information is mapped on to the  
30 ocean grid. Temperature and salinity profiles in the ice are computed prognostically and sea ice  
31 distribution trends and seasonal cycles are realistic (Kelley et al. 2020).  
32  
33 Salinity in the ice affects basal ice formation/melt and heat diffusion (Schmidt et al., 2006) through the  
34 use of brine-pocket (BP) thermodynamics (Bitz and Lipscomb, 1999; Schmidt et al., 2014) and updated  
35 in Kelley et al. (2020). Sea ice dynamics use the viscous-plastic formulation of Zhang and Rothrock  
36 (2000). Lead fraction is treated differently in the Arctic and Antarctic wintertime thick-ice growth to  
37 produce more realistic heat fluxes and sea ice growth rates. Processes relevant to the salt budget (e.g.,  
38 gravity drainage and flushing of meltwater) are consistently treated with the BP physics.

39

40 The mass and energy associated with net snow accumulation over each hemisphere's ice sheets is  
41 used to adjust the implicit iceberg calving fluxes into the adjacent oceans over a 10 year relaxation  
42 time, and this diagnostic is preserved as glacial melt. This ensures that the model can reach a long-  
43 term equilibrium under changed climate forcings, but does not imply realism in the response of ice  
44 sheets to climate, since the ice sheets are assumed to remain in mass balance.  
45

## 46 **2.7 Radiation, Clouds & Atmospheric Composition**

47 The total solar irradiance at the top of the model's atmosphere is based on satellite calibrations (Kopp  
48 and Lean, 2011) with a base value of  $1,361 \text{ W/m}^2$ . Spectral irradiance values are taken from Coddington  
49 et al. (2016). The radiation model includes explicit multiple scattering calculations for solar radiation  
50 (shortwave, SW) and explicit integrations over both the SW and thermal (longwave, LW) spectral regions  
51 (Schmidt et al., 2006; Schmidt et al., 2014; Kelley et al., 2020). SW radiation is absorbed by  $\text{H}_2\text{O}$ ,  $\text{CO}_2$ ,  
52  $\text{O}_3$ ,  $\text{O}_2$ , and  $\text{NO}_2$  using the k-distribution approach (Lacis and Oinas, 1991) that utilizes 15 noncontiguous  
53 spectral intervals to model overlapping cloud-aerosol and gaseous absorption and is scattered by clouds  
54 and aerosols using size-dependent Mie scattering, ray tracing and T-matrix theory (Mishchenko et al.,  
55 1996).

56  
57 Longwave radiation is absorbed by  $\text{H}_2\text{O}$ ,  $\text{CO}_2$ , and  $\text{O}_3$  using the correlated k distribution with 33 intervals  
58 (Lacis and Oinas 1991; Oinas et al. 2001), designed to match line-by-line computed fluxes and cooling  
59 rates throughout the atmosphere to within about 1%. Weaker bands of  $\text{H}_2\text{O}$ ,  $\text{CO}_2$ , and  $\text{O}_3$ , as well as  
60 absorption by  $\text{CH}_4$ ,  $\text{N}_2\text{O}$ , CFC-11, and CFC-12 are included in an approximate fashion as overlapping  
61 absorbers, but with coefficients tuned to reproduce line-by-line radiative forcing over a broad range of  
62 absorber amounts.

63  
64 Variations in incoming solar forcing in the 20th Century simulation follow Matthes et al. (2017) and  
65 include updated spectral solar irradiance. Orbital parameters that determine the seasonal and  
66 latitudinal variation in insolation are calculated each year (updated on 1st January), based on Berger  
67 (1978).  
68

69 The radiation model also includes the effects of 3D cloud heterogeneity via the 3D cloud  
70 parameterization (Cairns et al., 2000) in order to get more realistic albedos from realistic water paths  
71 and particle sizes (Schmidt et al., 2006). The procedure rescales optical depth, asymmetry parameter,  
72 and single scattering albedo according to the relative variance of the cloud particle density distribution.

73  
74 The partitioning between convective precipitation that descends and has the potential to evaporate in  
75 the environment rather than in the downdraft is increased from 0% to 50%, thus increasing the  
76 sensitivity of humidity to convection. The downdraft buoyancy is based on virtual temperature including  
77 condensate loading. Glaciation in the mixed-phase temperature range has been updated in the  
78 stratiform cloud parameterization.

79  
80 Stratiform cloud formation is done more realistically because there is no restriction in coverage, no  
81 dependence on vertical motion, the altitude is taken to be relative to local planetary boundary layer  
82 (PBL) height rather than a fixed 850 hPa, better demarcating cloud-topped boundary layers from the  
83 free troposphere where the threshold relative humidity is used for the TOA radiation balancing process  
84 (described in section 3.1.1) to maintain top-of-the-atmosphere radiative balance.

85  
86 The advection of humidity (and other tracers) is done only once every physics time step (30 min) with  
87 iterative time stepping to avoid any Courant–Friedrichs–Levy violations. The smaller time steps are used  
88 primarily in the stratosphere and upper troposphere where zonal winds are strong or as a result of  
89 extremely high flow deformation.

90  
91 The atmospheric chemistry configuration used in GISS ModelE2.1-CC2 is the basic non-interactive  
92 composition (NINT f2 used in CMIP6) in which the background fields of ozone and aerosol  
93 concentrations (dust, sea-salt, sulfate, nitrate, ammonium and carbonaceous aerosols e.g. black and  
94 organic carbon) are derived from simulations of the interactive one-moment aerosols (OMA) version of  
95 the model, run as an atmosphere only model with prescribed SST (Kelley et al., 2020). All anthropogenic  
96 and biomass burning emissions of short-lived species were updated to CMIP6 specifications (Hoesly et  
97 al., 2018; van Marle et al., 2017) and were prescribed annually or monthly.

98  
99 Specifically, with regards to dust, the six dust bins (0.1–0.2, 0.2–0.5, 0.5–1, 1–2, 2–4, and 4–8  $\mu\text{m}$  particle  
00 diameter), are used for coatings on dust particles, the dust particle densities, and the weights that are  
01 used to partition the total clay which is advected as a bulk species in the model. The weights reflect the  
02 size distribution of dust. The effect of wind biases on dust transport over the dust source regions is  
03 alleviated in the model by employing an additional term (“gustiness”) to express the high frequency  
04 variability in the surface winds that lift dust into the atmosphere. This variability results from stirring of  
05 the atmospheric boundary layer by dry convective thermals or frictional eddies, along with downdrafts  
06 spreading outward from the base of convective events. The magnitude of the associated wind gusts are

parameterized as described by Cakmur et al., (2004). The longwave optical depth for dust was increased by 30% to account for the longwave scattering effect. The lensing effect of sulfate and nitrate coatings on BC was parameterized by increasing the shortwave optical depth for BC by 50%. For more information see Pérez García-Pando et al., (2016).

Currently in NOBMg within GISS ModelE2.1-CC2, dust iron deposition is obtained from the historical annual cycle climatology extracted from the GISS E2.1-G online dust simulations which include eight externally mixed minerals (illite, kaolinite, smectite, carbonates, quartz, feldspar, iron oxides, and gypsum) plus internal mixtures between seven minerals and iron oxides (Perlwitz et al., 2015a; Perlwitz et al., 2015b). The masses of free and structural iron and their fractions of total iron have been evaluated using measurements from Izaña Observatory (Pérez García-Pando et al., 2016). For more on the dust deposition, see section 3.1.2 on Forcings. In future versions of the model, dust deposition will be interactive with the global carbon cycle.

A second order scheme is used for the transport (Schmidt et al., 2006) of all the tracer fields in the atmosphere which are advected using the highly nondiffusive quadratic upstream scheme (QUS; Prather 1986), which keeps track of nine subgrid-scale moments as well as the mean within each grid box. This increases the effective resolution of the tracer fields and allows the GISS model to produce reasonable climate fields with relatively coarse nominal resolution.

## **2.8 Surface albedo**

The radiation model generates spectrally dependent direct/diffuse flux ratios for use in biosphere feedback interactions. The surface albedo utilizes six spectral intervals, in nanometers: nm: VIS (300-770), NIR1 (770-860), NIR2 (860-1250), NIR3 (1250-1500), NIR4 (1500-2200), NIR5 (2200-4000), and is solar zenith angle dependent for ocean, snow, and ice surfaces.

The spectral and solar zenith angle dependence of ocean albedo is based on calculations of Fresnel reflection from wave surface distributions as a function of wind velocity (Cox and Munk 1956) taking also into account the effects of foam and hydrosols on ocean albedo (Gordon and Jacobs 1977). The spectral and solar zenith angle dependence of snow and sea ice is modeled following Warren and Wiscombe (1980). Snow “ages” following Loth and Graf (1998) and has a different albedo for wet or dry snow. Ocean ice albedo is spectrally dependent and is a function of ice thickness and parameterized melt pond extent (Schmidt et al., 2006 and references therein).

40  
41 Light profiles from the atmospheric radiation module are propagated underwater into the ocean and  
42 spectrally decomposed to 33 wavebands that are used to compute growth of the phytoplankton groups,  
43 sinking profiles, as well as changes in the vertical distribution of ocean temperatures due to biologically  
44 mediated absorption and scattering in the water column (Gregg and Conkright, 2002). NOBMg is  
45 coupled prognostically to the atmospheric radiation module which is part of ModelE2.1. Direct and  
46 diffuse sunlight in 6 wavebands (5 in the near IR and 1 in the visible part of the spectrum) that are  
47 computed by the radiation module are decomposed into the 33 wavelengths (250–3700 nm) needed  
48 for the absorption and scattering by the different phytoplankton species. The decomposition factors are  
49 based on the fractions of direct (diffuse) sunlight at the equator for each wavelength relative to the total  
50 radiation as derived from a monthly climatology provided by the Ocean–Atmosphere Spectral Irradiance  
51 Model (OASIM) and forced by MODIS data for aerosols and clouds (Gregg and Casey, 2009).

52  
53 The spectral albedo of vegetation is prescribed to be seasonally dependent on PFT and Lambertian (i.e.  
54 assuming flat diffusing reflective surface). Bare soil is grey, with values from Matthews (1984). For  
55 vegetation, seasonal spectral albedos are specified by mapping the Ent PFTs to the 8 biome types of  
56 Matthews: tundra (Ent cold-adapted shrubs), C3 grass (Ent C3 perennial, annual and arctic grasses),  
57 shrub (Ent arid-adapted shrubs), deciduous forest (Ent all deciduous trees), evergreen needleleaf forest  
58 (Ent evergreen needleleaf trees), tropical rainforest (Ent evergreen broadleaf trees), and crops (Ent  
59 crops). The total land albedo is computed as a weighted average over the corresponding land cover  
60 fractions and linearly interpolated in time between the prescribed seasonal values. In the presence of  
61 snow, the snow albedo is included into the weighted average in accordance with its masking fraction.  
62

## 63 **2.9 Interface exchanges**

### 64 **2.9.1 Surface fluxes**

65 Separate atmospheric surface fluxes are computed for each subsurface reservoir: open water, sea or  
66 lake ice, land ice, snow free and snow covered ground. ModelE calculates atmospheric turbulence over  
67 the entire vertical column. The surface fluxes are calculated using a submodule that is embedded  
68 between the surface and the midpoint of the first resolved model layer. In the atmospheric planetary  
69 boundary layer (PBL), a formulation for the temperature, moisture, and scalar fluxes is employed and  
70 consists of a local (diffusive) term and a countergradient term derived from large-eddy simulation (LES)  
71 data (Holtslag and Moeng 1991) as well as a nonlocal vertical transport scheme for virtual potential  
72 temperature, specific humidity, and other scalars (Holtslag and Boville 1993).

.73  
.74 A level-2.5 turbulence model (Cheng et al. 2002; Yao and Cheng, 2012) is applied to this region (using  
.75 eight sublayers) independently over each surface type where the mixing profile depends on the height  
.76 of the PBL, which is closely related to the large eddy size and thus characterizes the nonlocality, and the  
.77 buoyancy and shear effects at the surface. The turbulence length scale formulation is obtained from the  
.78 large eddy simulation data by Nakanishi (2001).  
.79  
.80 Above the PBL a second-order closure (SOC) model developed by Cheng et al. (2002) is used, which is  
.81 based on Mellor and Yamada (1982). The Reynolds stress and heat flux equations have been solved with  
.82 a more advanced parameterization of the pressure–velocity and pressure–temperature correlations. In  
.83 addition, a few turbulent time scales are determined by a new two-point turbulence closure model  
.84 (Canuto and Dubovikov 1996a,b). The more realistic “Richardson number criterion” rather than the “TKE  
.85 criterion” to calculate the PBL height, following Troen and Mahrt (1986) and Holtslag and Boville (1993).  
.86  
.87 The lower boundary conditions for the wind and the potential temperature equations are determined  
.88 assuming continuity of the respective turbulent fluxes at the surface. To calculate the fluxes through the  
.89 surface layer, drag and transfer coefficients are set using similarity theory following Hartke and Rind  
.90 (1997). The roughness length for momentum ( $z_{0m}$ ) over land is specified as in Hansen et al. (1983). The  
.91 roughness lengths for temperature ( $z_{0h}$ ) and moisture ( $z_{0q}$ ) over land and land ice are taken from  
.92 Brutsaert (1982) to be proportional to  $z_{0m}$ . Ocean and ocean ice are treated as rough bluff surface, with  
.93  $z_{0m}$  combining the smooth surface value (Brutsaert 1982) with the Charnock relation for the  
.94 aerodynamic roughness length; as for  $z_{0h}$  and  $z_{0q}$  Eqs. (5.26)–(5.27) of Brutsaert (1982) with  
.95 background values  $1.4 \times 10^{-5}$  and  $1.3 \times 10^{-4}$  m, respectively, are used.  
.96  
.97 We reduce the saturation specific humidity by 2% over the oceans (to account for sea salt aerosol  
.98 effects; Gill 1982).  
.99  
.00 Over land, the surface evaporative flux is determined by the vegetation, but to improve estimates of  
.01 surface humidity by the submodule described above, we calculate the maximum available  
.02 evapotranspiration and do not allow the atmosphere to draw more water than is available. This requires  
.03 a change in the humidity surface boundary condition from a variable to a fixed flux in such situations.  
.04  
.05 Continental runoff occurs at the mouths of rivers as well as small tributaries. Prognostic river runoff is  
.06 computed within ModelE2.1-CC as part of the simulated global hydrological cycle, thus increases

07 (decreases) due to changes in precipitation and evaporation over land, and delivers accordingly  
08 nutrients and carbon components to the coastal ocean (for more details see section 3.1.2 on Forcings).

## 09 10 **2.10 Coupling: Ocean, lake, ice, and land surface coupling**

11 Coupling between the different components in ModelE is synchronous and at the frequency of the  
12 physics time step (30 min). It is always implemented via fluxes of the fundamentally conserved  
13 quantities (mass, energy, and tracer). Given the surface conditions, the atmospheric model calculates  
14 the precipitation, radiative, and other surface fluxes (surface wind stress, evaporation, and sensible and  
15 latent heat) over each surface type. Radiation and precipitation fluxes are first applied directly to the  
16 land surface (soils, vegetated ground, and glaciers) and the land surface model computes latent and  
17 sensible heat fluxes, as well as land carbon fluxes. Using the atmosphere–ice wind stress, the sea ice  
18 dynamics calculates the horizontal ice velocities and the resulting ice–ocean stress. The ocean has finer  
19 horizontal resolution than the atmosphere and thus area conserving schemes of horizontal interpolation  
20 of fluxes from the atmospheric and the ocean grid are designed to conserve mass, energy, and tracer  
21 fluxes.

22  
23 River flow occurs on the atmospheric grid and is deposited into the ocean by the coupler (Schmidt et al.  
24 2014). Ice-ocean fluxes involving melt, heat conduction, and momentum drag are computed on the ice  
25 grid, while ice formation in the ocean occurs on the ocean grid and is then interpolated back to the  
26 atmosphere-ice grid ensuring conservation of fluxes.

## 27 28 **2.11 Conservation**

29 More than 100 vertically integrated diagnostics for every process in the atmosphere and the ocean are  
30 accumulated on each horizontal surface cell. They include fresh and salt water mass, atmospheric  
31 kinetic and static energy, ocean energy, and water and energy for each continental reservoir, as well as  
32 all ocean tracer fields. Offline programs utilize these monthly accumulated diagnostics, as well as water  
33 mass, tracer and energy amounts for each reservoir at beginning and end of each month, and compute  
34 the change in reservoirs and compare to flux changes. For both global and regional domains, errors are  
35 zero within the restrictions of machine accuracy (real\*4 for fluxes and real\*8 for reservoirs).



### 36    **2.11.1 Physical model conservation**

37    In the atmosphere, the advection scheme is mass conserving for humidity and tracers, and potential  
38    enthalpy conserving for heat. All processes including dynamics, cloud schemes, gravity wave drag, and  
39    turbulence, conserve energy and air, water, and tracer mass to machine accuracy. All dissipation of  
40    atmospheric kinetic energy through various mixing processes is converted to heat globally. In the long-  
41    term mean, net top-of-atmosphere (TOA) radiative flux approaches zero in the pre-industrial control  
42    simulation.

### 43    **2.11.2 Land carbon conservation**

44    In atmosphere-coupled experiments, the Ent TBM is currently run in biophysics-only mode without  
45    prognostic leaf area index (LAI) or stem growth. Vegetation structure is prescribed from satellite-  
46    derived vegetation cover, type, height, and maximum LAI. Maximum LAI is used to compute  
47    vegetation demography and cohort densities based on Ent's allometric relations by PFT. For each PFT,  
48    we use observed monthly LAI from Moderate Resolution Imaging Spectroradiometer (MODIS) which is  
49    interpolated at a daily time step. For land cover/land use change (LCLUC), Ent prescribes only annual  
50    historical change in cropland cover, rescaling the natural vegetation cover fractions in a grid cell (Miller  
51    et al., 2021; Ito et al. 2020); land use change such as deforestation is included only implicitly as part of  
52    the total cultivated land cover change.

53  
54    Currently, we run Ent in biophysics-only mode where vegetation structure is prescribed and all carbon  
55    pools are prescribed except the soil pools and the labile (non-structural carbohydrate) vegetation  
56    carbon pool which are prognostic. The labile carbon pool is dynamic with uptake of carbon through  
57    photosynthesis, withdrawal for respiration and leaf growth, and retranslocation with senescence. Any  
58    carbon excess, that is due to the fact that LAI is prescribed and no stem growth is allowed, is  
59    redirected to the litter pool in the soil, and that ensures carbon conservation. As mentioned earlier,  
60    land carbon change associated with the annual land cover change is distributed as CO<sub>2</sub> fluxes to the  
61    atmosphere over the course of the following year. Because the prescribed vegetation structure may  
62    not be in equilibrium with the simulated climate due to climate model biases or scenarios of climate  
63    change, respiration is restricted when labile carbon balances approach zero or slightly negative,  
64    allowing recovery of the labile pool during the growing season. This is also done to ensure carbon  
65    conservation.

### 66 2.11.3 Ocean carbon conservation

67 NOBMg conserves carbon, iron, ammonium and silicate but does not conserve nitrate, due to the fact  
68 that the nitrogen cycle is not closed. For nitrogen fixation, we assume an unlimited reservoir  $N_2$  is  
69 available for diazotrophs. Moreover, while there is an oxygen-dependent switch from faster (aerobic)  
70 to slower (anaerobic) remineralization rates, denitrification (the conversion of nitrate to  $N_2$ ) is not  
71 represented in NOBMg. Alkalinity is also not conserved since we do not include an explicit calcium  
72 carbonate tracer. Hence, rather than exchanges between dissolved and solid-phase carbonate ions  
73 being present, dissolved carbonate ions are either removed in near-surface waters by implicit calcium  
74 carbonate production (assumed to be proportional to primary productivity), or released by implicit  
75 carbonate dissolution in deep waters (assumed to follow a fixed exponential profile).

76

## 77 3 Model setup

### 78 3.1 Model coupled-carbon components

79 The general procedure for setting up and running the GISS ModelE2.1-CC2 consists of the following steps  
80 (Fig. 5):

- 81 1) running the physical climate model coupled to the one-moment aerosols (OMA) composition  
82 submodel in atmosphere with prescribed SST in order to obtain distributions of aerosols for the  
83 non-interactive aerosol (NINT) runs
- 84 2) running the fully coupled climate model with aerosols prescribed from (1) to equilibrium, i.e. till  
85 the top of the atmosphere radiative forcing is near 0 (approximately 3000 model years)
- 86 3) running the fully coupled climate model after the completion of step (2) + the land carbon to  
87 equilibrium i.e. the air-land flux of  $CO_2$  is near 0 (within  $\pm 0.1$  PgC/yr from 0, approximately 50  
88 years for radiative equilibrium only), this involves a preliminary soil carbon spinup as described  
89 in section 3.1.3.
- 90 4) running the fully coupled model after the completion of step (2) + the ocean carbon cycle to  
91 equilibrium, i.e. the air-sea flux is near 0 (within  $\pm 0.1$  PgC/yr from 0, about 500 years).

5) running the fully coupled climate model (3)+(4) (including soil carbon) till carbon equilibrium (within  $\pm 0.1$  PgC/yr from 0, spinup run)

Tuning is done separately in each of steps 2-4 by controlling parameters within each submodel in order to ensure that CO<sub>2</sub> flux to the atmosphere is near zero. The tuning procedure for the ESM model is described below.




Figure 5. Steps in coupling the different ModelE components into ModelE-ESM.

### 3.1.1 Model tuning

#### Tuning of the atmosphere-ocean coupled model

The value of the critical relative humidity for clouds is used for the TOA radiation balancing process described to maintain top-of-the-atmosphere radiative balance (Schmidt et al., 2014).

#### Tuning of the land carbon

Commented [AR5]: figure dropped

08 The Ent TBM photosynthetic capacity, slope of the Ball-Berry stomatal conductance equation, and  
09 respiration parameters were tuned offline on Fluxnet sites to predict observed net ecosystem exchange  
10 (NEE) and attain equilibrium behavior of seasonal labile carbon pools, as documented in Kim et al.  
11 (2015). The Fluxnet sites were: Tapajo National Forest, Brazil (evergreen broadleaf trees); Hyttiala,  
12 Finland (evergreen needleleaf trees); Morgan Monroe State Forest, Indiana, and Harvard Forest,  
13 Massachusetts, USA (cold deciduous broadleaf trees); Tonzi Ranch, California, USA (drought deciduous  
14 broadleaf trees); Vaira Ranch, Lone, California, USA (C3 annual grassland). Other vegetation parameters  
15 tuned to summaries from the literature were: relative soil moisture at water stress onset and wilting  
16 point; temperature at which frost hardening commences and begins; linear response of soil respiration  
17 to relative soil moisture and unstressed level. All allometric relations were also derived from literature  
18 sources. For applicability at the global scale, specific leaf area (SLA) by PFT was derived from the TRY  
19 database (<https://www.try-db.org/TryWeb/Database.php>) of leaf traits (Kattge et al. 2011).

21 The SLA was further tuned for each PFT in the coupled model spinup using values within the tuning  
22 parameter range in the TRY database of leaf traits to ensure that the plant labile carbon pool (non-  
23 structural carbohydrate) remains stable at the given size of mature trees. This stable behavior of the  
24 labile carbon pool depends on prescribed plant size boundary conditions being at equilibrium with the  
25 simulated climatology. Carbon starvation, therefore, may occur locally in ESM simulations, in which  
26 case the plant respiration is artificially reduced or stopped to prevent negative carbon balances; such  
27 plants would otherwise experience mortality. Excess labile carbon can be allocated to prognostic stem  
28 growth, or when the model is run in biophysics-only mode with fixed canopy structure it is simply  
29 senesced as extra litterfall to the soil.

#### 31 **Tuning of the ocean carbon cycle model**

33 Tuning is performed in order to ensure that the preindustrial air-sea CO<sub>2</sub> is at equilibrium and  
34 secondarily to produce realistic patterns and magnitudes of ocean primary production. Firstly,  
35 phytoplankton maximum growth rates for chlorophytes and coccolithophores were reduced to be  
36 within the range reported in previous studies. For chlorophytes, the growth rate ( $\mu_{chlo}$ ) was reduced  
37 from 2.52 to 1.89 d<sup>-1</sup>. (Law et al., 2011; Grant et al., 2013; Liefer et al., 2018) For coccolithophores, the  
38 growth rate ( $\mu_{cocc}$ ) was reduced from 2.27 to 1.70 d<sup>-1</sup> (Buitenhuis et al., 2008; Krumhold et al., 2017)  
39 (Table A5). From this change, several tuning experiments were conducted where the maximum carbon  
40 remineralization rate ( $\alpha_c$ ) was perturbed between 0.06 and 0.6 d<sup>-1</sup>. A maximum remineralization rate of  
41 0.3 d<sup>-1</sup> resulted in the air-sea CO<sub>2</sub> flux being at equilibrium, as well as net primary productivity increasing

from 15 Pg C/yr to 23 Pg C/yr (Lerner et al., 2024). To tune primary production, we first increased the stoichiometric ratio of iron to chlorophyll ( $b_{Fe}$ ) from  $2 \times 10^{-5}$  to  $8 \times 10^{-5}$   $\mu\text{g Fe}/\mu\text{g chl}$ , the latter value falls within the upper fourth quartile of the compilation of stoichiometries for different phytoplankton groups (Finkel et al., 2010). We also increased the silica half-saturation constant ( $k_{Si}$ ) for diatom growth from 0.2 to 2.85  $\mu\text{M}$ , the revised value being a compromise between the much lower values found for low-latitude species (0.02-1.4  $\mu\text{M}$ ), and the much higher values found for species in the Southern Ocean (4.2-88.7  $\mu\text{M}$ ; Nelson and Trüger, 1992). Finally, we tuned the remineralization rate of detrital iron and the scavenging rates of ligand-bound in order to increase NPP while achieving an equilibrated air-sea  $\text{CO}_2$  flux. For detrital iron, we explored a range of maximum detrital iron remineralization rates from 0.01 to 0.4  $\text{d}^{-1}$ , while for the ligand-bound iron scavenging rate ( $k_{Fe}$ ), the range explored was  $2.75 \times 10^{-4}$  to  $2.75 \times 10^{-2} \text{ d}^{-1}$ . The best combination of parameters that both increased NPP while allowing the air-sea  $\text{CO}_2$  flux to achieve equilibrium was  $k_{Fe} = 2.75 \times 10^{-3} \text{ d}^{-1}$  and  $\alpha_{Fe} = 0.25 \text{ d}^{-1}$ , for which NPP increased from 23 to 30 Pg C/yr. Note that  $k_{Fe}$  found by this tuning is two orders of magnitude greater than that in the CMIP6 incarnation of the model (Romanou et al., 2013; 2014). However, such an increase is consistent with the observed rates of trace metals with similar scavenging behavior to that of iron (Quigley et al., 2002; Parekh et al., 2004; Lerner et al., 2017). In addition to tuning the iron cycle, we tuned the iron half-saturation constant of  $\text{N}_2$ -fixation so that the globally-integrated rate would fall between 68 and 134 Tg N/yr. We explored a range between 0.12-12  $\mu\text{M}$ , and found that the lower limit provided the best globally integrated value (83 Tg N/yr), while still allowing the air-sea flux to be at equilibrium.

### 3.1.2 Forcing fields for carbon cycle


We use greenhouse gas concentrations from transient historical forcing prescribed for CMIP6 (Meinshausen et al., 2016). In the ESM simulations presented here, radiation experiences atmospheric emission-based  $\text{CO}_2$  abundance changes, as described below, while aerosol and ozone constituents are prescribed based on output of an ensemble of GISS-E2.1 AMIP simulations, which used full atmospheric chemistry and aerosol schemes (OMA; Kelley et al., 2020). The GISS CCv1 model version is labeled GISS-E2.1-G-CC in the CMIP6 repository.

#### Ocean carbon cycle forcing fields

Forcing (external) fields for the ocean carbon cycle include dust iron deposition, ligands, and riverine nutrients and carbon cycle components delivered to the coastal ocean. These are described below separately.

**Dust Iron Deposition**

Atmospheric dust falls to the surface through wet deposition, gravitational settling and turbulent removal. Wet deposition is the result of collisions with precipitation either within or below the cloud. As in many models (Adebiyi and Kok 2020; Meng et al., 2022), dust falls out too fast in ModelE, so that its concentration, especially of larger particles, is unrealistically low downwind of sources (Castellanos et al. 2024), as revealed by ocean deposition measurements over the tropical Atlantic (van der Does et al., 2016; 2020).



**Figure 6. Seasonal averages of dust iron flux deposited at the surface of the ocean (nm, Fe/m<sup>2</sup>/s).**

The ocean iron cycle is forced with the historical annual cycle dust climatology extracted from the consistent (ModelE2.1) online dust simulations using the GISS dust model (Miller et al., 2006), which has been expanded to include eight externally mixed minerals (illite, kaolinite, smectite, carbonates, quartz,

Commented [AR6]: added in supplement


feldspar, iron oxides, and gypsum) plus internal mixtures between minerals and iron oxides (Perlwitz et al., 2015a, 2015b). The masses of free and structural iron and their fractions of total iron have been evaluated using measurements for location at Izaña Observatory (Perez García-Pando et al., 2016). From the PI simulation with the prognostic dust, we extract 50 years of daily dust fluxes and compute a monthly climatology that is used to force the model. In the ModelE dust module, there are 6 size categories, Clay, Silt1, Silt2, Silt3 and Silt4 and Silt5. The last two describe the largest particles which make small contributions for ocean deposition, since they fall out near their source. For each category, we compute dry and wet deposition fluxes, gravitational settling and ice-ocean flux (in units of  $10^{-13}$  kg/m<sup>2</sup>/s). Silt1 particles contribute the largest proportion to the global average dust flux, the annual emission or the total deposition. This flux affects the timing of dust delivery to the ocean, since dust falling on ice may not be released to the ocean until the ice melts a few months later. Furthermore, we assume that the iron content varies within each dust size class, with clay containing 3.5% iron and silt containing 1.2% iron (Fung et al., 2000). Lastly, a constant iron solubility of 2% is also assumed (Romanou et al., 2013).

Figure 6 shows that the peak in dust deposition happens in the spring and summer and mainly in the tropical/subtropical regions of Africa, Saudi Arabia and south-eastern Asia.

#### **Iron Ligands**

The fraction of total iron that is complexed with organic ligands (see Appendix, eq. A27-A29) is computed using spatially varying but temporally constant prescribed ligand concentrations. Ligands are prescribed from a compilation of ligand measurements obtained at various depths from open ocean waters between 1992-2014. (Caprara et al., 2016). Due to the spatial sparseness of the dataset, we employ a simplistic method of translating the data onto the model grid. First, we split the model's ocean domain into thirteen distinct oceanographic regions, including the Antarctic, the Southern Indian ocean, the Southern Pacific Ocean, the Southern Atlantic Ocean, the Equatorial Indian ocean, the Equatorial Pacific ocean, the Equatorial Atlantic ocean, the Northern Indian ocean, the Northern Central Pacific ocean, the Northern Central Atlantic ocean, the Northern Pacific ocean, the Northern Atlantic ocean, and the Mediterranean Sea. Then we combine all ligand samples present within each region into a single composite vertical profile. We then bin-average the composite profile using 100 m bins above 1000 m and using 500 m bins below 1000 m. The bin-averaged profile is then smoothed using LOWESSsmoothing, where the weighted local linear regressions consider 40% of the data, and three iterations of regressions are performed where successive iterations apply residual-based reweightings. Finally, the smoothed profile is linearly interpolated on the vertical coordinates of the model. Since a

single composite profile is applied to an entire region, the resulting spatial distribution of ligands resembles “patches” (Fig. 7). In future model development we will remove the patchiness.



Commented [AR7]: added in supplement

**Figure 7. Spatial distribution of total ligand concentration (nM) used to compute iron speciation in the ocean carbon cycle model. Spatial distributions are shown in the ocean at the surface (upper left), 150 m (upper right), 1000 m (bottom left), and 3000 m (bottom right).**

#### Riverine inputs of biogeochemical constituents

As described earlier, continental runoff is prognostic in ModelE2.1-CC and delivers chemical constituents to the coastal zone. The concentrations of particulate organic carbon (POC), dissolved organic carbon (DOC), dissolved inorganic carbon (DIC), nitrate, silicate, and iron at all the major and many minor river estuaries are obtained from an annual climatology (da Cunha et al., 2007), and they modulate the biogeochemical characteristics of the freshwater outflow into the ocean at these sites. Therefore, as continental runoff increases (decreases) due to changes in precipitation and evaporation over land, the delivery of these ocean biogeochemical tracers will increase (decrease). Particulate carbon burial into the sediment, although prescribed for all areas shallower than 150 m, takes place in the estuaries where detritus delivered with the riverine flow tends to accumulate. Riverine sources of silica, iron, DIC and



'44 DOC concentrations are larger at northern high latitudes (Fig. 8), whereas nitrate and POC show larger  
'45 values in subtropical regions.

'46  
'47 **Land carbon cycle forcing fields**

'48 The prescription of crop cover change here was updated to the Land Use Harmonization Version 2  
'49 Global Carbon Budget 2020 dataset (GCB-LUH2 2020) (Friedlingstein et al. 2020), which provides land  
'50 cover changes through 2020. All crop types in the data set were merged into one crop type and  
'51 treated as a C3 crop. As in Ito et al. (2020), the natural vegetation boundary conditions and crop  
'52 monthly LAI are provided from the Ent Global Vegetation Structure Dataset (Ent GVSD) v1.0 (Kiang et  
'53 al., Ent GVSD v.10 Technical Report, in prep.). The Ent GVSD v1.0 is derived from satellite data sources  
'54 and includes land cover types and monthly varying LAI from the Moderate Resolution Imaging  
'55 Spectroradiometer (MODIS) (Gao et al., 2008; Myneni et al., 2002; Tian et al., 2002a, 2002b; Yang et  
'56 al., 2006) for the year 2004, and tree heights from Simard et al. (2011), who utilized 2005 data from  
'57 the Geoscience Laser Altimeter System (GLAS) aboard the ICESat (Ice, Cloud, and land Elevation  
'58 Satellite). Specific leaf area (carbon mass per leaf area) data from the TRY database of leaf traits  
'59 (Kattge et al., 2011) was classified for the Ent TBM 12 plant functional types (PFTs). With historical  
'60 crop cover change, the natural vegetation fractions are rescaled. The Ent GVSD also uses nearest-  
'61 neighbor extensions of LAI for grid cells where the land cover change results in cover types that were  
'62 not there in the 2004 MODIS land cover. These boundary conditions allow for geographic variation in  
'63 subgrid fraction, height, and LAI for each PFT, such that biomes can be different mixes of PFTs with  
'64 different demography and canopy structure.

'65 The use of the natural vegetation boundary conditions and an older historical crop cover data  
'66 set with the spinups in the ModelE C4MIP contribution (Ito et al. 2020) resulted in annual carbon  
'67 fluxes and stocks well in the middle of CMIP5 predictions, as well as in the middle of emerging CMIP6  
'68 comparisons. However, the prescribed leaf area index and stem biomass do not allow for prediction  
'69 of phenomena like boreal greening with climate warming, and the land model does not yet have  
'70 deforestation or fire as major components of the land carbon budget. The CO2 fertilization effect is  
'71 manifested by enhanced photosynthetic uptake, boosts to litterfall to the soil, and increased water use  
'72 efficiency.

'73  
'74

'75  
'76  
'77  
'78  
'79  
'80  
'81




Figure 8. a) Sites of continental runoff release in ModelE. b) Zonally averaged concentrations of nitrate, silica, dissolved iron, POC, DIC and DOC.

Commented [AR8]: added in the supplement

'82  
'83  
'84  
'85  
'86  
'87  
'88




Figure 9. Historical global total natural vegetation and crop cover area 1850-2020 prescribed for the simulations in this paper. The natural vegetation cover is derived from MODIS for the year 2004, which is rescaled given historical crop cover. Crop cover is the sum of the LUH2 cropland and pasture cover to capture historical land cover changes. Individual simulations will vary in cover fractions due to dynamic lakes having complementary subgrid cover to bare soil and vegetation.

Commented [AR9]: added in the supplement

### 3.1.3 Initial conditions

#### Land carbon initialization and spinup

At initialization, plant labile carbon has default values in relation to plant individual maximum foliage biomass by PFT, twice this amount for woody plants and four times this amount for grasses, sufficient to grow this foliage in the next growing season. Soil carbon may start from zero or be initialized from an input file for the upper 30 cm of soil derived from the International Soil Reference and Information Centre - World Soil Information (ISRIC-WISE) data set (Batjes 1996). All coupled carbon cycle simulations require that these plant labile and soil carbon pools be spun up to equilibrate with climate.

The plant labile carbon pool typically spins up in 30 years. To spin up soil carbon, the same method is used as described in Ito et al. (2020): the equilibrated preindustrial control simulation is run for 9 years and the simulated meteorology (soil moisture and temperature) and litterfall is saved in memory at hourly time steps and then iterated over the soil 750 times for a total of 6750 simulated years. The whole cycle typically takes less than 24 world clock hours on a modern cluster using 88 CPU cores. This is typically repeated 2-3 times to achieve a balance an order of magnitude smaller than the 0.1 GtC/year in the protocol for C4MIP.

#### Ocean carbon initial conditions and spinup

For the ocean carbon cycle, macronutrients (nitrate and silicate), DIC, Alkalinity, and O<sub>2</sub> were initialized from the Global Ocean Data Analysis Project, version 2 climatology (GLODAPv2; Lauvset et al., 2016), which maps fully quality controlled and internally consistent ocean biogeochemical data collected between 1972-2013 onto a 1°x1° lat/lon grid and 33 vertical layers. Phytoplankton were initialized using equations derived from the synoptic relationships found between satellite-derived chlorophyll-a and diagnostic pigments (Hirata et al., 2011). These equations are:

$$\text{Diatoms} = 1.3272 + \exp(-3.9828x + 0.4003) \quad (\text{Eq. 1})$$

$$\text{Chlorophytes} = (0.2490/y) \exp(-1.2621(x - 0.5523)^2) \quad (\text{Eq. 2})$$

$$\text{Cyanobacteria} = (0.0067/(0.6154 * y)) \exp(-19.5190(x + 0.9643)^2 / 0.3787) + 0.1027x^2 - 0.1189x + 0.0626 \quad (\text{Eq. 3})$$

$$\text{Coccolithophores} = 1 - (0.9117 + \exp(-2.7330x + 0.4003))^{-1} - (0.1529 + \exp(1.0306x - 1.5576))^{-1} + 1.8597x - 2.9954 - (0.2490/y) \exp(-1.2621(x - 0.5523)^2) \quad (\text{Eq. 4})$$

where  $x$  is the base-10 logarithm of the chlorophyll-a concentration and  $y$  is chlorophyll-a concentration. Chlorophyll-a used to compute initial phytoplankton functional groups is obtained from averaging the SeaWiFS seasonal climatologies between 1998-2010 ([https://oceandata.sci.gsfc.nasa.gov/SeaWiFS/Mapped/Seasonal\\_Climatology/9km/](https://oceandata.sci.gsfc.nasa.gov/SeaWiFS/Mapped/Seasonal_Climatology/9km/)). Equations (1-4) above were used to initialize phytoplankton in the upper 75 m of the water column; below this depth, phytoplankton concentrations are initialized at 0 mg Chl m<sup>-3</sup>. Ammonium, zooplankton, and detrital material (carbon, silica, and iron) are initialized with spatially invariant values 0.5 uM, 0.05 mg chl m<sup>-3</sup>, and 0 ug L<sup>-1</sup>, respectively. Finally, dissolved iron is initialized by splitting the domain into the same 13 geographic regions used to derive the prescribed ligand concentrations (see Forcings). Within each region, an iron:nitrate ratio is specified that is laterally and vertically invariant. The iron:nitrate ratios are obtained from a compilation of *in-situ* measurements (Fung et al., 2000), and are shown in Table 1. After iron:nitrate ratios are specified, iron concentrations are initialised by multiplying, in each grid cell, this ratio by the initial nitrate concentration from GLODAPv2.

**Table 1. Iron:Nitrate ratios used to estimate initial conditions for iron concentrations. Ratios taken from Fung et al, 2000.**

| Region                     | Antarctic | South Indian Ocean | South Pacific Ocean | South Atlantic Ocean | Equat. Indian Ocean | Equat. Pacific Ocean | Equat. Atlantic Ocean | North Indian Ocean | North-Cent. Pacific Ocean | North-Cent. Atlantic Ocean | North Pacific Ocean | North Atlantic Ocean | Mediterranean Sea |
|----------------------------|-----------|--------------------|---------------------|----------------------|---------------------|----------------------|-----------------------|--------------------|---------------------------|----------------------------|---------------------|----------------------|-------------------|
| Ratio (x10 <sup>-3</sup> ) | 3.5       | 20                 | 4.5                 | 20                   | 22.5                | 4                    | 20                    | 25                 | 6                         | 20                         | 6                   | 10                   | 20                |

Commented [AR10]: added in the supplement

After the tuning, the ocean carbon model spin ups were carried out for approximately 300 years off an equilibrated simulation of the pre-industrial ocean-atmosphere-ice system which was itself spun up for 7000 years.

### 3.2 Experimental design

For model simulations, we conducted a preindustrial control with 1850 conditions, a transient historical run over 1850-2021, future scenarios for SSP2-4.5, SSP5-8.5, and SSP5-3.4 for 2022-2100 and a compound 1% increase in CO<sub>2</sub>. For the historical simulations and the SSPs we used CO<sub>2</sub> emissions (not atmospheric concentrations) from input4mips files (Hoesly et al, 2018; Gidden et al., 2019) for the

period 1850-2014 and constant thereafter, while for non-CO2 GHGs we follow CMIP6 specifications for prescribed concentrations from 1850-2014 (Meinshausen et al., 2017) and use concentrations from the NOAA Global Monitoring Laboratory from 2015 to 2021 (Lan et al., 2022b). For the eA0 (the compound 1% run) we back-calculated implied emissions of CO2 from a concentration driven compound 1 percent run and we used those to force the eA0 scenario. All other GHGs and aerosols are prescribed following the CMIP6 specifications.

Table 2. ModelE2.1-CC experiments described in this paper

| experiment                                | short name | description                                                                                                                                                         |
|-------------------------------------------|------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Preindustrial                             | ePI        | Preindustrial spin-up at quasi-1850 conditions                                                                                                                      |
| Historical                                | eHI        | Historical run 1850-2021 (forcings post 2014 constant)                                                                                                              |
| SSP2-4.5                                  | e245       | 2022-2100, “middle of the road” scenario                                                                                                                            |
| SSP5-8.5                                  | e585       | 2022-2100, “very high emissions”                                                                                                                                    |
| SSP5-3.4                                  | e534       | 2021-2100, “overshoot” scenario, it follows SSP5-8.5 till 2040 and then describes rapid mitigation to zero emissions by 2070 and net negative emissions thereafter. |
| 1% atmospheric CO2 concentration increase | eA0        | 1850-1979, “equivalent 1% CO2 emissions”, emissions derived from a run with prescribed 1%/year CO2 increase in atmospheric concentrations.                          |

Commented [AR11]: new table for PI and Historical simulations

3.3 Auxiliary datasets used for model assessment

Table 3. Observational and reanalysis data used for model assessments


| Data set name                             | fields used        | spatial coverage & resolution | temporal coverage & resolution          | Reference                                                                                             |
|-------------------------------------------|--------------------|-------------------------------|-----------------------------------------|-------------------------------------------------------------------------------------------------------|
| RFROM OHCA maps (Argo+EN.4.2.2+satellite) | ocean heat content | 0.25°x0.25°                   | 1993-2024 anomalies from 1993-2022 mean | <a href="https://www.pmel.noaa.gov/rfrom/">https://www.pmel.noaa.gov/rfrom/</a><br>Lyman and Johnson, |

|                                      |                                                           |               |                                                                           |                                                                                                                             |
|--------------------------------------|-----------------------------------------------------------|---------------|---------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------|
|                                      |                                                           |               |                                                                           | 2023                                                                                                                        |
| ECCO-V4r4                            | derived ocean heat content and uptake                     | 1°x1°         | 1992-2017                                                                 | Forget et al. 2015a, Forget and Ponte 2015; Storto et al., 2019; Fukumori et al., 2021                                      |
| Spawn et al 2020                     | land biomass                                              | 300 m         | year 2010                                                                 | Spawn et al., 2020                                                                                                          |
| Carbon Tracker (which includes OCO2) | land/ocean carbon fluxes, CO2 concentrations              | 1°x1°         | monthly resolution (yearly for uncertainties in carbon fluxes), 2000-2022 | Jacobson et al., 2023                                                                                                       |
| Global Carbon Budget 2024            | land and ocean carbon flux components, CO2 concentrations | global totals | annual                                                                    | Friedlingstein et al. 2025<br><a href="https://globalcarbonbudget.org/gcb-2024">https://globalcarbonbudget.org/gcb-2024</a> |
| MCD43C3                              | land albedo                                               | 0.05°x0.05°   | 16-day, monthly, 2002-2026                                                | Schaaf and Wang, 2021                                                                                                       |
| GLODAP                               | Nitrate, silicate, DIC, Alkalinity                        | 1°x1°         | annual climatology, climatological period from 1972-2013                  | Lauvset et al., 2016                                                                                                        |
| GISTEMP                              | SAT                                                       | 1°x1°         | monthly resolution, 1880-present                                          | GISTEMP Team, 2025<br>Lennsen et al., 2024                                                                                  |
| MERRA2                               | SLP, surface wind speed, surface air temperature          | 0.5°x0.625°   | monthly resolution, 1980-present                                          | Gelar et al., 2017                                                                                                          |
| SOM-FNN                              | ocean carbon flux                                         | 1°x1°         | monthly resolution, 1982-2023                                             | Landschützer et al., 2016                                                                                                   |
| FluxSat v2                           | land GPP                                                  | 0.05°x0.05°   | daily resolution, 2000-03-01 to 2020-08-01                                | Joiner & Yoshida (2021)                                                                                                     |
| World Ocean Atlas 2023               | SST, SSS                                                  | 1°x1°         | decadal resolution, 1991-present                                          | Locarini et al., 2024<br>Reagan et al., 2024                                                                                |
| CAFE                                 | NPP                                                       | 0.25°x0.25°   | 8-day resolution, 1998-2023                                               | Silsbe et al, 2016<br>Ryan-Keogh et al., 2023                                                                               |
| ERA5                                 | surface wind speed, sea level pressure                    | 0.25°x0.25°   | 1980-present                                                              | Hersbach et al., 2023                                                                                                       |

|        |                         |           |           |                                                                               |
|--------|-------------------------|-----------|-----------|-------------------------------------------------------------------------------|
| GPCPv6 | precipitation over land | 0.5°x0.5° | 1951-2010 | Schneider et al., 2011<br><a href="http://gpcc.dwd.de">http://gpcc.dwd.de</a> |
|--------|-------------------------|-----------|-----------|-------------------------------------------------------------------------------|

59 4 Results

60 4.1 Preindustrial control simulation



61  
62 Figure 10. Temporal evolution of preindustrial control fields: a) net radiation at top of the atmosphere (TOA), b) surface air temperature,  
63 c) land carbon flux, d) gross primary production (GPP), e) autotrophic respiration, f) soil respiration, g) net primary production on land  
64 (NPP = GPP - autotrophic respiration) h) net ecosystem production (NEP = NPP - soil respiration), i) excess carbon flux due to vegetation  
65 fractions change, j) net heat uptake by ocean, k) ocean air-sea carbon flux, l) net ocean primary production (NPP ocean). Annual mean  
66 fields are shown in gray. 20yr averages are shown in black. The horizontal axis is in years from the PI control run. Units are shown on  
67 the vertical axes. All fluxes are positive down into the surface (land or ocean).

Commented [AR12]: figure redone to include the previous version of the model (CMIP6). The text now also reflects that change



Surface exchanges and properties are in equilibrium for at least the last 500 years of the pre-industrial control simulation (Fig. 10).

Figure 10 shows the temporal evolution of different fields under 1850 conditions and Table 4 lists their long-term mean, standard deviation and trend. Radiative balance at the top of the atmosphere (TOA) is  $-0.04 \pm 0.34 \text{ W/m}^2$  (Fig. 10a & Table 4). Surface air temperature (Fig. 10b) has a small drift of about  $0.02^\circ\text{C}/100 \text{ years}$ . Net land carbon flux is  $0 \pm 0.65 \text{ PgC}$  (Fig. 10c), with a long term trend of  $-0.05 \text{ PgC}/100 \text{ years}$  (Table 4) while gross primary production (GPP), autotrophic and soil respiration (Fig. 10d,e,f respectively) have small trends, and so do net ecosystem production and net primary production over land (Fig. 10g and h). The largest variability arises from GPP while the largest trend from soil respiration (Table 4). The excess carbon flux due to change in vegetation cover (Fig. 10i) results from ModelE's dynamic lakes varying in cover, which is complementary to the continental land cover by vegetation and soil. Net heat flux (Fig. 10j) is out of the ocean by  $0.07 \text{ W/m}^2$  with a large interannual variability of the order of  $0.49 \text{ W/m}^2$  (Table 4) and a small positive trend of  $2.6 \times 10^{-2} \text{ W/m}^2/100 \text{ years}$ . Net primary productivity in the ocean is about  $30 \text{ PgC/year}$  which is low compared to  $50\text{-}60 \text{ PgC/year}$  from observations (Westberry et al., 2023).

Table 4. Mean, standard deviation and trend over the last 100 years of the PI control run

|                          | long term mean                                 | standard deviation                            | trend (last 100 years)                               |
|--------------------------|------------------------------------------------|-----------------------------------------------|------------------------------------------------------|
| Net Rad TOA              | $-0.04 \text{ W/m}^2$                          | $0.34 \text{ W/m}^2$                          | $-4.42 \times 10^{-6} \text{ W/m}^2 / \text{yr}$     |
| Surface air Temp         | $13.61^\circ\text{C}$                          | $0.14^\circ\text{C}$                          | $2 \times 10^{-4}^\circ\text{C} / \text{yr}$         |
| Ocean Carbon Flux        | $0.00 \text{ Pg Carbon/year}$                  | $0.12 \text{ Pg Carbon/year}$                 | $4 \times 10^{-4} \text{ PgCarbon} / \text{yr}$      |
| Land Carbon Flux         | $0.00 \text{ Pg Carbon/year}$                  | $0.65 \text{ Pg Carbon/year}$                 | $-5.0 \times 10^{-4} \text{ PgCarbon} / \text{yr}$   |
| GPP                      | $101.32 \text{ Pg Carbon/year}$                | $0.98 \text{ Pg Carbon/year}$                 | $-8.1 \times 10^{-4} \text{ PgCarbon} / \text{yr}$   |
| auto resp                | $67.32 \text{ Pg Carbon/year}$                 | $0.50 \text{ Pg Carbon/year}$                 | $-8.2 \times 10^{-4} \text{ PgCarbon} / \text{yr}$   |
| soil resp                | $34.00 \text{ Pg Carbon/year}$                 | $0.39 \text{ Pg Carbon/year}$                 | $5.2 \times 10^{-4} \text{ PgCarbon} / \text{yr}$    |
| ecvf                     | $-1.12 \times 10^{-12} \text{ Pg Carbon/year}$ | $3.46 \times 10^{-13} \text{ Pg Carbon/year}$ | $-3.82 \times 10^{-15} \text{ PgCarbon} / \text{yr}$ |
| NEP                      | $-1 \times 10^{-3} \text{ Pg Carbon/year}$     | $0.65 \text{ Pg Carbon/year}$                 | $-6 \times 10^{-4} \text{ PgCarbon} / \text{yr}$     |
| NPP land                 | $34.00 \text{ Pg Carbon/year}$                 | $0.62 \text{ Pg Carbon/year}$                 | $1.1 \times 10^{-5} \text{ PgCarbon} / \text{yr}$    |
| Net Heat Uptake at Ocean | $-0.07 \text{ W/m}^2$                          | $0.49 \text{ W/m}^2$                          | $2.6 \times 10^{-4} \text{ W/m}^2 / \text{yr}$       |

|           |                      |                     |                       |
|-----------|----------------------|---------------------|-----------------------|
| Surface   |                      |                     |                       |
| NPP ocean | 30.22 Pg Carbon/year | 0.19 Pg Carbon/year | -9.0e-4 PgCarbon / yr |




Figure 11. Distributions of time average fields and standard deviation. Averaging is over the last 100 years from the PI simulation. Maps of mean a) atmospheric CO<sub>2</sub> concentration, b) surface air temperature, c) air-land carbon flux, d) air-sea carbon flux, and e) ocean heat uptake.


Commented [AR13]: Figure moved to supplement. Some discussion on PI patterns remains in the text.

Spatial distributions of the preindustrial annual mean atmospheric CO<sub>2</sub> concentration peak at the equator (Fig. 11a) with values around 284.5 ppm and lowest values at high latitudes about 283.7 ppm.

'91 The largest interannual variability (Fig. 11b) appears over tropical land regions and in the equatorial  
'92 Atlantic and is associated with the large interannual variability due to ENSO (Kelley et al., 2020).  
'93 Surface air temperature (Fig. 11c and d) shows a strong zonal variation with warmer temperatures  
'94 over low latitudes and significantly colder at high latitudes, North America and Asia. The largest  
'95 variability appears over the eastern tropical Pacific and in the Arctic Ocean rim associated with natural  
'96 modes of variability which are stronger than observed in our simulations (Kelley et al., 2020). The air-  
'97 to-land flux is largest in the continental North America and the northern Eurasian continent (Fig. 11e).  
'98 The Amazon is a weak source due to Model's warm and dry bias in this region, and the high latitudes  
'99 are net sinks due slow soil carbon storage in these cold regions, while the largest variability is present  
'00 in Europe and North America, tropical Africa and the Amazon regions (Fig. 11f). The ocean carbon flux  
'01 has a strong sink in the North Atlantic and the Southern Ocean (Fig. 11g), particularly the Labrador Sea  
'02 and the Greenland-Iceland-Norway (GIN) Seas. World ocean subtropical gyres are regions of carbon  
'03 uptake whereas the tropical belt is outgassing CO<sub>2</sub> to the atmosphere with peak values in the eastern  
'04 Equatorial Pacific and secondary peaks in the Benguela and Canaries current upwelling regions. The  
'05 largest variability in the ocean sink is exhibited in the North Atlantic subpolar gyre (mostly in the GIN  
'06 seas and the Gulf Stream separation region) as well as in the tropical upwelling regions (Fig. 11h). The  
'07 net ocean heat flux peaks in the tropical belt (Fig. 11i) where the ocean takes up heat from the  
'08 atmosphere and is lowest in the North Atlantic, specifically along the Gulf Stream, the Labrador and  
'09 Irmingers Seas and the GIN seas, where the ocean releases heat to the atmosphere. The largest ocean  
'10 heat flux variability emerges in the eastern tropical Pacific Ocean, and the North Atlantic subpolar  
'11 regions (Fig. 11j).

'12

13 4.2 Historical simulations



14  
15 **Figure 12. Annual Time series of a) atmospheric CO2 concentration from the model (eHI) and observations from**  
16 **input4mips (Meinshausen et al., 2017) from 1850 to 2015, and from the NOAA Global Monitoring Laboratory from 2015**  
17 **to 2021 (Lan et al., 2022a,b), b) surface air temperature (annual mean timeseries) and superimposed estimates from**  
18 **ERA5 and MERRA2 (see Table 3), c) change in surface air temperature from the average over the period 1951-1980 and**  
19 **GISTEMP, d) air-to-land carbon flux from the model (eHI) compared to observations from the Carbon Tracker (shaded is**  
20 **the uncertainty of the annual mean based on the averaging of inversions that consider different prior estimates of**  
21 **fluxes and their uncertainties; Jacobson et al., 2023), e) components of land carbon fluxes to the atmosphere (positive is**  
22 **flux upward) due to land use change, comparing eHI net emissions to the mean estimates of the Global Carbon Budget**  
23 **2024 (GCB2024, Friedlingstein et al. 2025). The legend tabulates the GCB2024 means and estimated minimum and**  
24 **maximum bounds around the mean over the period 1959-2021 where the eHI simulation and GCB2024 overlap in time,**

Commented [AR14]: redid figure to include the previous version of the model that clearly shows the improvements

f) air-sea carbon flux from the model compared to observations from SOM-FFN and Carbon Tracker (2000-2022) (shaded is the uncertainty around the annual mean, stemming from a combination of uncertainties including (i) the gridding of SOCAT observations, (ii) the RMS error between the SOM-FNN estimates and the gridded observations, and (iii) the uncertainty stemming from the chosen air-sea flux parametrization; Landschützer et al., 2016), g) cumulative heat uptake in the model and comparison to estimates from Argo, both timeseries are anomalies from the 1993-2022 mean, h) ocean heat flux and comparison to estimate from ECCO-V4r4, years 1992-2017. It should be noted here that model forcings post-2014 remain constant at 2014 levels except for atmospheric non-CO2 greenhouse gases which follow observations from NOAA-GML (Lan et al., 2022b), a fact to be considered in model-to-observation comparisons post 2014 shown in this figure.

Atmospheric CO2 concentrations agree well with observations (Fig. 12a) with small discrepancies of up to 12.5 ppm that peak between 1925 and 1950 (Fig. 12b). The small underestimation of atmospheric CO2 stems from insufficient land or maybe even the ocean carbon uptake. These are both underestimated in comparison to CarbonTracker (Fig. e,f) over 2000-2022, but may be overestimated over the longer historical period (more discussion on land and ocean carbon sinks below) while the ocean sink is overestimated compared to SOM-FNN. Spatially positive biases up to 1.5 ppm are found over the northern hemisphere land and ocean especially the high latitudes and similar negative biases are found in the Southern Hemisphere, especially over the tropical forest regions (Fig. 13a1,a2).

Surface air temperature agrees well with reanalysis products in terms of trend (Fig. 12b). The model overestimates the cooling associated with Pinatubo (1991-1992) and generally shows larger interannual variability than the observations after 1950 possibly related to the strong ENSO variability (Kelley et al., 2020). Surface temperature anomalies from preindustrial track the observations very well (Fig. 12c) particularly after the 1950s. Between 1875 and 1950 the model is warmer than the baseline than the observations, and this is despite the fact that there is somewhat less atmospheric CO2 in the model than was observed. Spatially, the biases are more pronounced in the Arctic (by about 4°C), the GINSeas and the Weddell Sea (by about 2°C), while the Eastern Pacific is warmer in the model by about 1-2 degrees C, the subpolar North Atlantic is colder by about 1-2 degrees, the Southern Ocean in general is warmer by about 1 degree C, and the Indian Ocean is colder by by 1 degree C (Fig. 13b1,b2).

Precipitation (Fig. 13c1,c2) in the historical eHI simulation compared to land surface precipitation observations by the Global Precipitation Climatology Centre (GPCC), National Oceanic and Atmospheric Administration (NOAA), Full Data Reanalysis Version 6.0, over 2001-2010, shows that the model has an extensive dry bias over the tropics, a persistent model bias also documented in Kelley et al. (2020) for the ModelE-2.1 contribution to CMIP6. This bias is particularly pronounced in the Amazon Basin, which is a persistent issue in GCMs in general (Respati et al. 2024). The ModelE dry

59 bias is also strong in India and Indonesia. The model has a wet bias in Western China and the West  
60 Coast of South America along the Andes.

61 Average sea level pressure (Fig. SUPP1a,b) is higher than observations by more than 10 mb over  
62 Greenland and Antarctica and the mountainous ranges in south-east Asia and South America. It is  
63 systematically underestimated over the ocean by about 5-10 mb. Surface winds are weaker than in  
64 observations (Fig. SUPP1c,d) by more than 5m/s over Greenland and Antarctica, 3-5 m/s over the mid-  
65 and southwest US and Patagonia, Europe and Northern Africa, and by 2-4 m/s over ocean subpolar  
66 gyres, an issue with all previous modelE versions, and which led to the inclusion of the “gustiness”  
67 term (see Section 2.7).

68 Global mean air-to-land carbon (net biome productivity, NBP) flux (Figure 12d) hovers at a net flux of  
69 zero in the 20-year moving average over 1850-1942, after which the trend becomes a net sink. Over  
70 2000-2022, it is underestimated compared to Carbon Tracker as it lies below the mean and outside the  
71 envelope of interannual variability in this dataset. Interannual variability is also much higher amplitude  
72 in the model, due sensitivity of the land carbon to ModelE’s higher amplitude interannual surface  
73 temperature variability from simulated El Niños. A more detailed breakdown of eHI time series of the  
74 land fluxes (GPP, autotrophic respiration, soil respiration, emissions due to land use change, NPP, NBP,  
75 NEP) and soil carbon stocks are provided in Figure SUPP2, which shows both the annual totals and  
76 their 20-year running means to show trends. These show all natural fluxes accelerating after ~1960.  
77 Regionally, the underestimation (which exceeds 20 mole CO<sub>2</sub>/m<sup>2</sup>/yr) stems from weaker Amazon  
78 and African rainforest uptake as well as weaker boreal forest uptake (Fig. 13d1,d2). The cumulative  
79 land carbon sink 1850-2023 estimated by the Global Carbon Budget 2024 is 220±60 PgC (Friedlingstein  
80 et al. 2025), whereas in the historical simulation the model reaches a cumulative sink of only 56.6 PgC  
81 by 2021 (Figure SUPP2b). CarbonTracker and other flux inversion models estimate generally lower  
82 NBP than DGVMs, which Bastos et al. (2020) attribute to disagreement on NBP interannual variability  
83 responses to El Niño and uncertainties in land use emissions, particularly in South America and  
84 Southeast Asia. Meanwhile, based on remote sensing observations of vegetation change, Randerson  
85 et al. (2025) find that forest growth is far less than predicted by the ensemble of DGVMs in the  
86 GCB2023; the remote sensing data yield an estimated net land carbon sink over 2000-2019 of 0.8±0.7  
87 PgC/year, whereas the GCB2024 DGVMs estimate a high 1.32±0.2 PgC/year due to unrealistic  
88 predictions of forest greening or densification. Our model estimates 1.12±1.04 PgC/year over this  
89 period. The GCB2024 DGVM estimates of earlier preindustrial through 20th century deforestation  
90 emissions also rely on highly uncertain predictions of forest biomass and land use change.  
91 Observational constraints of vegetation biomass remain a challenge: inventory and satellite surface

observations of global total vegetation biomass estimates still range over a wide 380-536 PgC (Erb et al. 2018), while more recent radar and lidar satellite data products of forest heights are still being processed to produce new global biomass estimates, which we detail more later.

In Figure 12e, the emissions from land cover and land use change (LCLUC) are plotted together with the estimates of the Global Carbon Budget 2024 (GCB 2024) (Friendlingstein et al. 2025), which span 1960-2024 only. These show that our land cover change (LCC) emissions strongly underestimate the mean estimate of the GCB2024 by more than 1 PgC/year, and is outside their low (less negative) estimate; the model underestimate is very likely because our model does not include deforestation /forest regrowth, wood harvest, or peat emissions, but only land cover conversion from natural vegetation to croplands. Therefore, our model's LCC emissions from the preindustrial period to 1960 likely also underestimate total LCLUC emissions by a similar magnitude, leading to our underestimate of atmospheric CO<sub>2</sub> (Fig. 12a).

Based on our model physics, we would expect the simulated underprediction of atmospheric CO<sub>2</sub> to be explained by the ModelE land model's lack of full representation of certain components of land use change (LUC). Fig. 12e shows the eHI net carbon emissions due to LUC compared to the breakdown of components of LUC change emissions estimated by the Global Carbon Budget 2024 (GCB2024, Friedlingstein et al. 2025). Although the model represents deforestation that converts forest to croplands or other cover types, it does not represent wood harvest or deforestation that results in degraded forest; also, the model is not a fire-enabled model, so fire emissions associated with deforestation and peatlands are not represented. Therefore, the eHI emissions miss on average 1.0±0.3 PgC/year over 1959-2021 compared to GCB2024 mean net LUC emissions. Since deforestation emissions were higher over the earlier historical period and the model misses these emissions, we would expect the underpredicted net emissions to the atmosphere to result in the model's slightly lower atmospheric CO<sub>2</sub> concentrations, which catch up to the observed as deforestation declined in recent decades. Given the much lower NBP estimated by Randerson et al. (2025), the high bias of GCB2024, and uncertainties in flux inversion models, despite the eHI simulation's comparatively low cumulative land carbon sink, it may be that ModelE's land carbon sink is overpredicted in the context



of the model's simulated flux components (Fig. SUPP2), explaining our low predicted atmospheric CO<sub>2</sub>. These discrepancies highlight the continued large uncertainties in land carbon stocks and fluxes.

Model SST (averaged over 1991-2020) is colder than World Ocean observations by about 2-4°C in the subArctic regions (Fig. 13f1,f2). The tropical eastern Pacific Ocean and Tropical Atlantic are warmer than observations by about 2-4°C, whereas the Southern Ocean is warmer than observations by about 2-4°C. The North Atlantic Current region and the Arctic are colder by about 2-5°C. The model historical climatology 1991-2020 of the sea surface salinity (SSS, Fig. SUPP1) is fresher than observations (Fig. SUPP1) by about 4 psu in the Eurasian side of the Arctic Ocean, and about 2 psu in the tropical regions of the central Pacific Ocean, the Atlantic Ocean and the Indian Ocean. The model overestimates salinity by about 2 psu in the tropical eastern Pacific Ocean and in the Arabian Sea and the Bay of Bengal in the Indian Ocean. The SSS biases largely reflect biases in evaporation and precipitation patterns over the global ocean (Kelley et al., 2020; Miller et al., 2021) which are in turn related to biases in surface pressure and surface winds (Fig. SUPP1).

The ocean carbon flux (Fig. 12f) lies between the observation-based estimates SOM-FFN and Carbon Tracker and within both datasets' interannual variability. Spatially, ocean carbon uptake negative biases (up to 5 mole CO<sub>2</sub>/m<sup>2</sup>/yr) are associated with strong wind-driven circulation and fronts (e.g. over the Gulf Stream, the Equatorial trade winds and the Southern Ocean subtropical front (Fig. 13g1,g2) whereas positive biases (up to 6 mole CO<sub>2</sub>/m<sup>2</sup>/yr) are linked to convection sites in the North Atlantic (Labrador Sea, Irminger Sea and GIN Seas) and the Antarctic Circumpolar Current (ACC) region. Subtropical gyres and the tropical Atlantic and Indian Oceans also show positive biases. Tropical ocean outgassing is also stronger in the GISS model, despite the weaker than observed winds and warmer SSTs there, because the pCO<sub>2</sub> disequilibrium is larger than observed, due to a combination of low productivity (as we describe below) and also because the vertical sinking of organic carbon below the thermocline is too small, which is discussed in more detail in Lerner et al. (2024).


Net primary productivity is underestimated everywhere in the world's oceans compared to CAFE observations by about 100-200 mgC/m<sup>2</sup>/day (Fig. 13h1,h2), except in the subtropical gyre in the North Atlantic and in the Benguela and Canaries current regions. The generally low bias in NPP results from at least two issues. First, there was an error in our ocean radiative transfer routines that compute the direct and diffuse radiation that enters and leaves each ocean layer. For each layer, we incorrectly used the layer depth (distance from the ocean surface to the layer), rather than the layer thickness (distance between upper and lower layer boundaries), when computing the amount by which direct radiation should be reduced from the top to the bottom of each layer. This resulted in too little

shortwave radiation reaching subsurface waters, reducing NPP. Second, phytoplankton in our model currently do not acclimate in ambient light levels, so that chl:C ratios in the sun-lit surface are the same as those deeper in the water column. Acclimation is expected to increase the growth of phytoplankton and NPP under low-light conditions (Geider et al., 1998), so the absence of this process in our model likely also contributes to lower productivity levels. We also should note that our comparison only encompasses a single NPP product, that derived from the CAFE algorithm applied to the Ocean Colour Climate Change Initiative merged remote sensing data product (Ryan-Keogh et al., 2023). NPP products are known to vary widely depending on the algorithm applied, particularly in the Southern Ocean but also to a significant extent in the Northern high latitudes. For example, using the same dataset, the Eppley-VGPM algorithm yields roughly the productivity of the CAFE algorithm in the Southern Ocean (Ryan-Keogh et al., 2023). Therefore, despite improvements that need to be incorporated, how well our model performs in some ocean environments, particularly the Southern Ocean, remains unclear. Finally, we note that a few regions, in particular the Benguela and Mauritanian upwelling regions, display positive NPP biases. The cause of these biases remain a subject of future investigation, although a potential contributing factor is the fact that our


167 model lacks water column (Lam and Kyupers, 2012) and benthic denitrification (Dale et al., 2014), which  
168 could result in too much upwelling of nitrate in these regions .

169


a1) atmCO2, CTracker 2000-2014




a2) eHI - CTracker 2000-2014




b1) Tsurf, GISTEMP 2000-2014




b2) eHI - GISTEMP 2000-2014




c1) Precip, GPCPv6 2001-2010




c2) eHI-Precip, 2001-2010




d1) CO2 fflux, CTracker 2000-2014




d2) eHI - CTracker 2000-2014




e1) CO2 GPP, FluxSat 2000-2009




e2) eHI - FluxSat 2000-2009



f1) SST, WOCE2023 1991-2020



f2) eHI - WOCE2023 1991-2020






Figure 13. Observational and observation-based estimates and model spatial biases averaged over the period of the observations. a1) Atmospheric CO<sub>2</sub> concentration anomalies from the global mean from Carbon Tracker averaged over 2000-2014. a2) Atmospheric CO<sub>2</sub> concentration anomalies, model biases from Carbon Tracker, b1) GISTEMP average 2000-2014, b2) Surface air temperature model biases from GISTEMP v4 (anomalies from 1951-1980) averaged over 2000-2014, c1) Precipitation average 2001-2010 from GPCPv4, c2) historical precipitation biases over land from GPCPv4, d1) Land carbon flux taken from Carbon Tracker between 2000-2014, d2) land carbon flux model biases in the model compared to Carbon Tracker, e1) land GPP taken from FluxSat between 2000-2009, e2) GPP model biases in the model compared to FluxSat, f1) Sea surface temperature (SST) averaged from the period of 1991-2020 from World Ocean Atlas 2023 (WOA2023), f2) historical SST biases in the model compared to WOA2023, g1) ocean carbon flux observations from SOM-FFN averaged over the period 2000-2014, g2) Ocean carbon flux biases in the model compared to SOM-FFN, h1) Ocean net primary productivity (NPP) averaged from 2000-2014 from CAFE (Silbe et al., 2016), h2) ocean NPP biases in

the model compared to CAFE, i1) ocean heat content from ECCO-V4r4, averaged over the period of 1993-2017, i2) Ocean heat uptake bias in the model compared to ECCO-V4r4.

Ocean heat uptake anomaly from a late-20th century baseline is well in agreement with estimates from Argo floats (cumulative; Fig. 12g) and with ECCO-V4r4 (non-cumulative anomaly from preindustrial; Fig. 12h) as well as variability in the latter dataset. Spatially, the patterns of average differences between the GISS model and ECCO-V4r4 for the period 1993-2017 are within  $\pm 50$  W/m<sup>2</sup>, with positive differences (GISS minus ECCO-V4r4) dominating the low latitudes and negative differences the high latitudes and the eastern Pacific Ocean (Fig. 13i1,i2). More specifically, ocean heat uptake averaged over 1993-2017 is underestimated in the model by more than 40 W/m<sup>2</sup> in the North Atlantic subpolar gyre and the ACC region (Fig. 13i1,i2) due to the lower wind speeds there, whereas it is overestimated over other regions with strong currents (e.g. the eastern Pacific and eastern Atlantic equatorial current regions, the Brazilian and Benguela current regions, the Agulhas retroflection region, the eastern part of the western Australian current region, and the Kuroshio region) because the sea surface temperatures are much higher (Fig. 13f1,f2).

**Seasonality**

The seasonal cycle of the air-to-sea flux of CO<sub>2</sub> and the partial pressure of CO<sub>2</sub> in the sea water have been extensively evaluated against several observational data sets (Lerner et al., 2021).

For the land carbon, seasonal GPP fluxes are shown in Fig. SUPP3, compared to estimates by FluxSat v2 (Joiner and Yoshida 2021). ModelE-CC estimates annual GPP as 118.6 $\pm$ 1.5 PgC/yr over 2000-2009, and 124.7 $\pm$ 2.0 PgC/yr over 2014-2021, which is currently at the lower end of both multi-model and observation-based estimates ranging 110-170 PgC/yr (Anav et al. 2015; Cheng et al. 2017; Xu and Chen, 2024). The latest Global Carbon Budget 2024 cites a mean annual GPP of 130 PgC over 2014-2023 (Friedlingstein et al. 2023).

The land seasonal regional GPP patterns qualitatively agree with FluxSat estimates (Fig SUPP3). However, there are stark regional differences. ModelE-CC simulates much higher GPP in the Amazon and Congo Basins in all seasons compared to FluxSat. Since ModelE has a warm/dry bias in these regions, we would expect GPP to be low. The FluxSat data are expected to be a lower bound estimate when based on flux tower data and not utilizing solar-induced fluorescence (SIF) data. Therefore, the higher ModelE GPP may be appropriate. In arid and semi-arid regions outside the Amazon and Congo in South America and Africa, ModelE predicted GPP is much lower than FluxSat; these mostly mixed shrubland and grassland areas are strongly controlled by soil moisture availability, and ModelE has a dry bias in these areas, leading to lower GPP. The shrubland PFT is also one for which the Ent TBM lacked a Fluxnet


19 site to evaluate simulated fluxes, so parameterizations of leaf photosynthetic capacity and respiration  
20 may be improved. In the northern high latitudes, the season MAM exhibits delayed spring greenup and  
21 the summer has low productivity due to ModelE's extended cold bias, while mid-latitude croplands in  
22 the Eastern U.S. and Europe are predicted to have much higher GPP than FluxSat estimates. The  
23 Northern Hemisphere differences between ModelE and FluxSat decline in the fall, SON, when deciduous  
24 trees senesce.

25 **4.3 Future Climate responses**

26 Future climate simulations were also conducted in emissions driven mode, in which atmospheric, ocean,  
27 and land carbon inventories respond to prescribed anthropogenic emissions of CO<sub>2</sub>, instead of  
28 atmospheric concentrations and the physical climate responds to changes in the marine and terrestrial  
29 carbon cycle. For the future scenarios eSSP2-4.5, eSSP5-3.4 and eSSP5-8.5, following the CMIP6  
30 protocol, CO<sub>2</sub> emissions are obtained from harmonized emissions trajectories produced by integrated  
31 assessment models (Gidden et al., 2019). For the eA0 experiment, emissions were obtained as the  
32 residual between the growth of the atmospheric fraction of CO<sub>2</sub> and the integral of the land and ocean  
33 carbon sinks and they were applied to the model in the emissions driven runs, where the atmospheric  
34 concentration of CO<sub>2</sub> is now calculated. The cumulative emissions applied in these simulations are  
35 shown in Fig. 14a. The comparison of the calculated atmospheric CO<sub>2</sub> concentration in each scenario  
36 and the prescribed timelines from CMIP6 ScenarioMIP are shown in Fig. 14b. The emissions driven  
37 simulations SSP5-3.4 and in SSP5-8.5 estimate lower concentrations of CO<sub>2</sub> in the atmosphere by about  
38 10 ppm at year 2100 than their concentration driven counterparts.  
39

**Commented [AR15]:** This section and all sections of future changes are moved to a subsequent paper on the Future reponses.





**Figure 14. Model globally averaged responses under different warming scenarios (described in Table 2). The implied emissions for the historical were taken from the Community Emissions Data System (CEDS; Hoesly et al., 2018), whereas the emissions for the SSP scenarios were from harmonized emissions trajectories produced by integrated assessment models (Gidden et al., 2019). (a) Cumulative emissions, (b) atmospheric concentration of CO<sub>2</sub> in concentration driven runs (dashed lines) as well as in the emissions driven runs, (c) surface air temperature, (c) precipitation, (e) direct visible radiation over land and ocean, (f) diffuse visible radiation over land and ocean, (g) GPP, (h) land NPP, i) land NEP. In panels b-h, variables are smoothed using a 20-year moving average filter.**

All panels in Fig. 14 reveal that the compound 1% scenario (eA0 purple lines) exhibits a far greater response than the historical simulation because the imposed emissions are much larger, as has been discussed in Sanderson et al. (2025). The atmospheric CO<sub>2</sub> concentrations as computed in the emissions driven simulations are very close to the ones given in CEDS (Fig. 14b purple line).

Furthermore, the model prediction of atmospheric CO<sub>2</sub> concentrations at 2014 in the historical simulation differs from the CEDS estimate by -1.9 ppm, and at 2100 by +4.4 ppm, -17.5 ppm and -40.9 ppm in the SSP2-4.5, SSP5.34, and SSP5-8.5 respectively. The eA0 shows surface warming of about 1.4°C, at the time of doubling of CO<sub>2</sub> in the atmosphere and 3.4°C near the time of quadrupling of CO<sub>2</sub>.


59 At 2100 eSSP2-4.5 has warmed by 2.5°C, SSP5-3.4 by 1.7°C (after an overshoot of 2.5°C), and SSP5-8.5  
60 by 4.2°C (Fig. 14c).

61  
62 Global mean precipitation (Fig. 14d) follows the trajectory of surface air temperature with increases of  
63 about 0.2 mm/day in the eA0 scenario, 0.08 mm/day in SSP2-4.5, 0.07 mm/day (after an overshoot of  
64 0.08 mm/day) in SSP5-3.4 and 0.93 mm/day in SSP5-8.5.

65  
66 Direct and diffuse radiation ( $E_d$  and  $E_s$  respectively), globally averaged (Fig. 14e and f) are mirror-images  
67 of each other in the historical simulation, in the sense that when  $E_d$  increases,  $E_s$  decreases. Historical  
68 direct radiation shows a peak at 50.1 W/m<sup>2</sup> in the 1940-1950 period, and a trough at 47.5 W/m<sup>2</sup> in  
69 1990-2000 period. Also direct radiation has a downward trend over the historical period, whereas  
70 diffuse is more or less constant. These variations are associated with analogous variations in SO<sub>2</sub> forcing  
71 fields. eA0 exhibits linear decreases in  $E_d$  and  $E_s$ , but SSP2-4.5 and SSP5-3.4 show increases in  $E_d$  and  
72 decreases in  $E_s$ , linked to the differing cloud patterns in these runs.

73  
74 GPP and NPP for all runs follow the trends in surface air temperature, which is fairly linear, while  
75 atmospheric CO<sub>2</sub> has an upward curve. e534's slowdown in GPP growth after 2040 similarly tracks its  
76 temperature. The land NEP trends are mostly controlled by NPP fluxes on land (Fig. 18b,c). In eA0,  
77 autotrophic respiration increases fairly linearly due to the model's fixed biomass and acclimation to  
78 temperature, while GPP has non-linear trends, such that the difference results in NEP increasing rapidly  
79 in the latter 19th century, slowing down in the early 20th c., and then declining starting around 1940.

80



**Figure 15.** Ocean responses under different warming scenarios: a) sea surface temperature (SST), b) sea surface salinity (SSS), c) cumulative ocean heat flux, note the different axes: left axis for historical, SSP2-4.5, 5-3.4 and 5-8.5, right axis for eA0, d) cumulative air-to-sea ocean carbon flux, e) Atlantic Meridional Overturning Circulation (AMOC) at 26°N (maximum below 500 m), and f) net freshwater flux into the ocean. For all panels except c and d, variables are smoothed using a 20-year moving average filter. Panels c and d are cumulative changes. Units are shown on the vertical axes. Colorscheme is shown on panel (a).

Global mean sea surface temperature (Fig. 15a) follows the increase of surface air temperature (Fig. 14c). Relative to 1850, warming at 2100 is 4.7°C, 1.6°C for SSP2-4.5, 1.1°C for SSP5-3.4 (after an overshoot of 1.5°C), and 2.7°C for SSP5-8.5.

Surface ocean salinity declines in all future scenarios (Fig. 15b): by about 0.6 psu in eA0, 0.13 psu in SSP2-4.5, 0.12 in SSP5-3.4, and 0.2 in SSP5-8.5 consistent with increases in precipitation (Fig. 15c). Ocean heat uptake (Fig. 15c) increases faster than surface air temperature in all scenarios (Fig. 14c) and SST (Fig. 15a). By 2100, the ocean has taken up 2500 zJ in SSP2-4.5, 2300 zJ in SSP5-3.4, and 3100 zJ in

97 SSP5-8.5. In eA0, it has taken up about 3000 zJ by year 129. Ocean carbon uptake increases in all  
98 simulations by 2100 (Fig. 15d): 400 in SSP4-4.5, 350 PgC in SSP5-3.4, and 480 PgC in SSP5-8.5. In eA0,  
99 ocean carbon uptake increases to 470 PgC by year 129.

00  
01 The Atlantic meridional overturning circulation (AMOC) strength at 26N weakens substantially in all the  
02 scenarios (Fig. 15e). In the historical simulation, the decline starts after 1980 and continues till 2014.  
03 SSP5-3.4 declines till 2040 are comparable to SSP5-8.5, some apparent discrepancies shown in Fig. 15e  
04 are due to the 20-year smoothing used in the plot. SSP2-4.5 shows the slowest decline after 2014. In  
05 contrast to the historical, the eA0 shows an immediate decline comparable to the SSPs. SSP2-4.5 and  
06 SSP5-3.4 show a plateau in the decline of AMOC after 2080 while SSP5-8.5 weakens most amongst the  
07 three SSP scenarios. Surface warming in the subpolar North Atlantic as well as meltwater from sea ice  
08 variations under future warming are large in SSP2-2.5 lead to a weak AMOC (Nazarenko et al., 2022;  
09 Romanou et al, 2023) which is more pronounced than the majority but not all the CMIP6 models.

10  
11 Freshwater fluxes into the ocean are showing a near-linear increase in eA0 (Fig. 15f) until about 2050  
12 and a slower increase thereafter. In SSP2-4.5 the plateau occurs around 2070 while in SSP5-3.4 around  
13 2040 (the time of the overshoot). SSP5-8.5 shows a peak in freshening around 2080 and a decline  
14 thereafter. Surface freshening does not entirely follow precipitation changes or surface salinity trends  
15 but also reflects changes in evaporation and freshwater inputs from sea-ice melt and river runoff.

16  
17 Ocean carbon cycle evolution under future scenarios is shown in Fig. 16. Primary production (NPP, Fig.  
18 16a) declines near-steadily in the historical simulation by about 0.004 PgC/yr, or about 0.7 PgC over the  
19 period 1850 to 2014. NPP continues to decline in SSP2-4.5 until 2050, partly due to iron declines and  
20 partly due to light availability declines, and then stabilizes around 29.4 PgC. In contrast, NPP in SSP5-8.5  
21 continues to decline till 2100, with some intermediate plateaus around 2040-2060 and 2080-2100. SSP5-  
22 3.4 plateaus after 2015 till 2060 and then recovers till 2100. Scenario eA0 shows the slowest decline in  
23 NPP, and over the first 50 years (1850-1900) there is a plateau in NPP in that run. Then, for the period  
24 1900-1979, NPP declines almost as fast as in the historical simulation. It is evident that NPP in NOBMg  
25 (version 12) is showing decadal shifts and abrupt changes which point to instabilities that merit further  
26 consideration.


27  
28 Further at depth of about 100m, export production (Fig. 16b) exhibits very interesting behavior as well.  
29 In eA0, it declines continuously until 1979, whereas NPP does not begin declining until 1900. . In the  
30 other scenarios, EP follows NPP but does not undergo the shifts and abrupt changes that NPP does, e.g.

after 2050 in SSP2-4.5 and SSP5-8.5. Surface pH (Fig. 16c) follows the SST trajectories (Fig. 15a) and shows declines in all scenarios, with a reversal only in the overshoot scenario.

Upper ocean nitrate (Fig. 16d) and oxygen concentrations (Fig. 16e) follow surface ocean temperatures with continuous declines except late in the overshoot simulations. These trajectories partly explain the behavior of NPP but notably do not exhibit a plateau in the first 50 years of the AO run as shown for NPP in Fig. 16a.

The decline in surface iron (Fig. 16f) during the earlier part of the historical simulation (1850-1900) is a spurious result stemming from the fact that our pre-industrial iron concentrations were too high due to differences in atmospheric deposition fluxes in the preindustrial and the historical simulations. We run the model with preindustrial iron from CMIP5 GISS simulations which were substantially lower than the historical simulations in CMIP6. In future, we will ensure that PI and historical simulations will be using consistent iron deposition rates and we expect this spurious effect will disappear. All SSP scenarios as well as eAO show increases in dissolved iron concentrations. These trends are not associated with dust deposition trends since dust aerosols are prescribed and constant in time in these simulations. Instead, they could partially be explained by the reduction in biological iron uptake due to reduced NPP (Fig. 16a) and the shoaling of mixed layer depths and that prevents mixing of iron from below.

49



50

Figure 16 Ocean carbon cycle related responses in different scenarios. (a) Net primary production in the ocean (NPP), (b) export production at 100m depth, (c) surface ocean pH, (d) nitrate concentration averaged between 0-100 m, (e) dissolved iron concentration averaged between 0-100 m, and (f) dissolved oxygen concentration


53


averaged between 100-600 m. Different color lines represent the PI control run (ePI, gray), the historical (eHI, blue), and scenarios eSSP2-4.5, eSSP5-8.5, and eA0 (green, yellow, red and purple respectively). For all panels except c,d, variables are smoothed using a 20-year moving average filter.

Table 6 shows responses to climate change at the end of scenarios SSP2-4.5 and SSP5-8.5 (2080-2090 mean) as departures from the early historical period (1870-1899 mean) and compares them to the CMIP6 multi-model mean and spread as computed by Kwaitkowski et al. (2020). The GISS model estimates a lower increase in sea surface temperatures in both scenarios and in fact falls slightly outside the multi-model mean. However, it also predicts a greater decline in ocean NPP compared to the multi-model mean in SSP2-4.5 while for SSP5-8.5 it is slightly underestimated but in both cases the GISS response falls within the spread of all the CMIP6 models. This behavior is also reflected in changes in global mean upper ocean nitrate concentration. With regards to changes in O<sub>2</sub> concentration, the model underestimates the global mean decline but is within the multi-model spread. Changes in ocean pH are very close to the multi-model mean in both scenarios.

**Table 6: Global mean changes in multiple impact drivers for two scenarios in GISS-E2.1-CC2 and from the CMIP6 ensemble (Kwaitkowski et al., 2020). The error ranges show inter-ensemble standard deviation. Global mean changes are computed as the difference between the driver at the end of the scenario (2080-2099 mean) and the beginning of the historical simulation (1870-1899 mean). For NPP, the difference relative to the historical value is shown. For O<sub>2</sub> and NO<sub>3</sub>, changes are computed using averages in the depth interval between 100-600 m (0-100 m).**

|                                         | SSP2-4.5/E2.1-CC2 | SSP2-4.5/CMIP6 | SSP5-8.5/E2.1-CC2 | SSP5-8.5/CMIP6 |
|-----------------------------------------|-------------------|----------------|-------------------|----------------|
| ΔSST (°C)                               | 1.59              | 2.10 ± 0.43    | 2.59              | 3.47 ± 0.78    |
| ΔNPP (%)                                | -2.23             | -1.13 ± 5.81   | -2.87             | -2.99 ± 9.11   |
| ΔO <sub>2</sub> (mmol/m3)               | -7.34             | -8.14 ± 4.08   | -11.75            | -13.27 ± 5.28  |
| ΔNO <sub>3</sub> <sup>-</sup> (mmol/m3) | -0.73             | -0.65 ± 0.32   | -0.99             | -1.06 ± 0.45   |
| ΔpH                                     | -0.260            | 0.260 ± 0.003  | -0.442            | 0.440 ± 0.005  |





**Figure 17. Response to future forcing: changes in SSP2-4.5 (left column), SSP3-3.4 (middle column), SSP5-8.5 (right column). Shown are differences of averaged fields over the last 20 years (2080-2100) from late historical (2015-2020). a1,2,3) surface air temperature (°C), b1,2,3) precipitation (mm/day), c1,2,3) soil liquid water change (kg/m²), d1,2,3) sea surface temperature (°C), e1,2,3) ocean heat uptake (W/m²), f1,2,3) air-to-land carbon flux (gC/m²/day), g1,2,3) air-to-sea carbon flux (gC/m²/day).**

Spatial patterns of changes in the distributions at the end of the 21st century compared to the late historical period (2015-2020) in the three scenarios (SSP2-4.5, SSP3-3.4 and SSP5-8.5) are shown on Fig. 17. In all scenarios, surface air temperatures increase (Fig. 17a1,a2,a3), except for a region of




.85 cooling in the subpolar North Atlantic likely associated with AMOC reduction. The tropics and mid-  
.86 latitudes see reduced precipitation (Fig. 17b1,b2,b3) and thus reduced soil moisture (Fig. 17c1,c2,c3);  
.87 however, the high latitudes see strongly increased soil moisture without a significant increase in  
.88 precipitation, because longer thawed periods allow for greater availability of liquid soil water.  
.89 Compared to SSP2-4.5, sea surface temperatures in SSP5-3.4 (Fig. 17d1,d2,d3) are characterized by  
.90 significant increases in the upwelling in the tropical Pacific and Atlantic as well as in the subtropical  
.91 convergence zone in the Southern Ocean, leading to less warming in these regions. In contrast, SSP5-  
.92 8.5 shows significant warming particularly in the upwelling regions and around the globe and even  
.93 reduced cooling in the subpolar North Atlantic compared to SSP2-4.5.

.94 With regard to ocean heat flux (Fig. 17e1,e2,e3), the largest differences from the late historical  
.95 simulation are found at high latitudes where the ocean takes up more heat in SSP5-8.5 than in the  
.96 other two SSPs. The other regions show little change.

.97 Land carbon (Fig. 17f1,f2,f3) is lost in most of the tropical and subtropical regions in SSP2-4.5 and  
.98 SSP5-8.5 due to increased surface temperatures enhancing both autotrophic and soil respiration as  
.99 well as greater water stress reducing GPP. In contrast, land carbon is gained in the overshoot scenario  
00 SSP5-3.4 in these regions due to less warming and less drying than in the other two scenarios, and  
01 also the eastern US and European forests take up more carbon. Remarkably, all scenarios show  
02 hotspots of increased land carbon uptake along the Yellow Sea coastal region of China, southern  
03 peninsulas of Australia, and the western tip of South America between Ecuador and Peru. These areas  
04 experience milder warming without declines in surface moisture.

05 The tropical and south Indian Ocean source of CO<sub>2</sub> is stronger in SSP5-3.4OVER- than in the other  
06 SSPs (Fig. 17g1,g2,g3) since it is more negative compared to the late historical period, while the sinks  
07 in the North Atlantic and in the Southern Ocean are also weaker. Thus the lower ocean uptake of CO<sub>2</sub>  
08 at the end of SSP5-3.4 seen in Fig. 15d is due to the strengthening of the sources and weakening of the  
09 sinks. In SSP5-8.5 the tropical Atlantic and subtropical Pacific sinks are nearly unchanged but the

uptake is increased everywhere in the rest of the global ocean explaining the increases in global mean uptake of CO<sub>2</sub> (Fig. 15d).



**Figure 18. Land carbon fluxes (sign is positive for flux from air to land): a) cumulative air-to-land land carbon flux, b) Soil carbon uptake, c) autotrophic respiration (negative is flux up to the atmosphere), d) soil respiration, e) land use and land cover change associated carbon flux, and f) NBP. For all panels except a, variables are smoothed using a 20-year moving average filter.**

Figure 18 dives deeper in the land carbon fluxes. The cumulative air-to-land carbon flux (Fig. 18a) is sensitive to: temperature (leaf photosynthetic capacity at, autotrophic and soil respiration); the CO<sub>2</sub>

20 fertilization effect; soil moisture (plant water stress, soil respiration); and crop cover change. Hence the  
21 rise in uptake seen in Fig. 18a over the historical period and future scenarios. The CO<sub>2</sub> fertilization effect  
22 may not be as pronounced as in other models, because we have kept the vegetation canopy structure  
23 and seasonal LAI fixed. In Fig. 18b, soil carbon overall increases in all scenarios, whereas other models  
24 may tend to show greater losses with warming; in ModelE-CC, the soil carbon incorporates the CO<sub>2</sub>  
25 fertilization effect, since we run the vegetation in biophysics-only mode without prognostic stem  
26 growth. With our static natural vegetation structure, the model does not capture the greening that may  
27 be occurring at high latitudes (but still subject to uncertainty due to complex ecosystem dynamics  
28 (Myers-Smith et al. 2020), while at the same time some observational estimates indicate that soil carbon  
29 may be a greater sink than biomass increases in the tropics (Xu et al. 2021), due to genetic and  
30 community structure limitations on tree growth, such that our soil carbon trajectories may be realistic.  
31 Autotrophic respiration (Fig. 18c) and soil respiration (Fig. 18d) increase with warming in all scenarios,  
32 with SSP5-3.5 responding with a decline when temperatures cool. Autotrophic respiration in the model  
33 does not, however, increase unlimited with higher temperature, but respiration acclimates to higher  
34 temperatures as is observed in nature. Soil respiration in the model is also responsive to increased soil  
35 carbon substrate from the CO<sub>2</sub> fertilization effect.

36  
37 Net biome productivity (NBP) in Fig. 18f is the net of GPP, autotrophic and soil respiration, and emissions  
38 due to LULCC. The eA0 1% CO<sub>2</sub> scenario, because of the combination of the CO<sub>2</sub> fertilization effect,  
39 extreme warming, higher precipitation and longer unfrozen season in the high latitudes, as well as  
40 absence of LULCC, exhibits dramatically higher GPP than the eHI scenario, which is reflected in increase  
41 in soil carbon. However, this enhanced GPP begins to saturate by 2050; this may be due to a combination  
42 of a limit on temperature enhancement of leaf photosynthetic capacity as well as drying of continental  
43 interiors.

#### 44 45 **4.4 Heat and Carbon storage on land and in the ocean**


46 Long term storage of heat and carbon in the ocean and on land is important for the control they exert  
47 on future climate change.

#### 48 49 **Ocean storage**

50 Figure. 19 shows a comparison of heat storage trends at different levels in the ocean in the GISS model  
51 and the ECCO-V4r4 ocean state estimate. Generally, the GISS ocean model stores more heat in the


Commented [AR16]: From this section we only keep the historical comparisons with observations.

52 subtropical gyres in the North Atlantic as well as the Atlantic sector of the Antarctic convergence zone  
53 at all levels. ECCO-V4r4 stores more heat in the subpolar North Atlantic and in the deep Southern Ocean  
54 around Antarctica. The deep western boundary current which is clearly outlined in the GISS ocean model  
55 at depths below 600m is not as clearly identified in ECCO-V4r4. The GISS model also stores more heat  
56 in the upper North Pacific Ocean compared to ECCO-V4r4, however the latter stores more heat in the  
57 tropical Pacific. Differences in the storage regions and depths will influence the response of the ocean  
58 model to future forcing changes, as heat stored closer to the surface and at higher latitudes will more  
59 easily be released to the atmosphere compared to heat stored at lower latitudes.



61 **Figure 19. Ocean heat content trend ( $\text{J/m}^2/\text{s}$ ) for the period 1993-2017 in ECCO-V4r4 and GISS2.1-CC2, vertically**  
 62 **integrated: a, b) from surface to 600m, c,d) 600-2000m, e,f) 2000m-bottom.**

63  
 64




65  
 66 **Figure 20. Ocean heat content trend ( $\text{J/m}^2/\text{s}$ ) in GISS2.1-CC2 from three different SSP scenarios, vertically integrated: a,**  
 67 **b,c) from surface to 600m, d,e,f) 600-2000m, g,h,i) 2000m-bottom. Left column is for SSP2-4.5, middle column for SSP5-**  
 68 **3.4, and the right column for SSP5-8.5.**

69

70 In future climates (Fig. 20), heat content in the GISS model continues to grow in the Atlantic and parts  
 71 of the Southern Ocean and at depths down to 2000m. The bottom layer is shown to differ little among  
 72 the different scenarios, except in the North Atlantic region. Clearly, the largest storage over the  
 73 centennial time scale of the 21st Century is taking place in mid-depths 600-2000m faster while little is  
 74 penetrating to greater depths.

75



**Figure 21. Integrated DIC (kg/m<sup>2</sup>) from GLODAP and GISS eHI for the period 1993-2014 at different levels: a) surface, and integrated over b) 0-600m, c) 600-2000m, d) 2000m - bottom.**

Compared to GLODAP, the model overestimates carbon stored in the upper ocean (0-600 m) of the tropical/subtropical regions, especially in the Atlantic (Fig. 21c,d). The model also underestimates the storage of carbon in intermediate waters (600-2000 m), particularly in the North Pacific (Fig. 21e,f). These findings are consistent with those of Lerner et al., 2024 that the flux of carbon beneath 100 m is too high and the transfer efficiency (ratio of carbon export at depth to carbon export at 100 m) is too

85 small, although that study was limited to the equatorial Pacific. High export fluxes near the surface may  
 86 be due to overgrazing by zooplankton, as zooplankton loss terms (via linear and quadratic mortality: eq.  
 87 AA32) produce the bulk of the detrital organic carbon in the model. On the other hand, the lack of  
 88 remineralized DIC in deeper waters below 600 m may due to the absence of processes that could  
 89 contribute to higher detrital sinking speeds, such as a lack of ballasting by inorganic material (carbonates  
 90 and lithogenics; Armstrong et al., 2021), and our use of a single-particle class model rather than multiple  
 91 size classes that would encompass a wide range of sinking speeds as found in the real ocean (Burd et al.,  
 92 2015).




Figure 22. Trends in DIC (kg/m<sup>2</sup>/year) from GISS SSP2-4.5 (first column), SSP5-3.4 (second column), SSP5-8.5 (right column) and at different levels. Top row: surface, second row: 0-600m, third row: 600-2000m, bottom row: 2000m - bottom.

In SSP2-4.5 and 5-3.4OVER the largest increase in the amount of carbon stored is in intermediate waters between 600 and 2000 m in the Atlantic (Fig. 22g,h). Upper ocean (0-600 m) carbon storage has also grown but to a lesser extent (Fig 22d,e). These features are consistent with the model having a reduced but still vigorous (>12 Sv) AMOC that transport carbon from the region of strongest uptake in the high latitude North Atlantic (Fig. 17g1-g2) to waters beneath the wintertime mixed layer depth (>1000 m). Carbon storage also increases mostly between 600-2000 m in the Southern Ocean north of the Polar Front in these scenarios, though to a lesser extent than in the Atlantic (Figs. 22g,h), suggesting this carbon is at least partially transported to this region of the Southern Ocean by the AMOC, where it is then transported zonally by the ACC. In contrast to the other scenarios, SSP5-8.5 shows (i) that carbon storage increases almost equally between the upper 600 m and 600-2000 m in the Atlantic, and (ii) a similar increase in carbon storage is seen in most other ocean regions in the upper 600 m (Figs 22f,i). The fact that the increased carbon storage is no longer concentrated in the intermediate waters of the Atlantic suggests that the role of AMOC (which has weakened substantially; Fig. 15e) in transporting carbon beneath 600 m has drastically declined, as ocean carbon is mostly increasing in the upper ocean (containing mostly surface and mode waters). It may also be that not enough time has passed for the more recently emitted carbon to be transported to deeper water, as in this scenario cumulative emissions are continuously increasing whereas they are plateauing in SSP2-4.5 and declining in SSP5-3.4OVER (Fig 14a). The fact that the two bottom-most layers have little change in carbon storage across the SSP shows that the GISS model stores carbon in the upper 2000m layer in decadal timescales (Fig 22j-l).

#### Land carbon storage Plant labile carbon and soil carbon distribution

With canopy structure and seasonal LAI otherwise prescribed in Ent, there are two prognostic carbon pools: labile carbon (also known as non-structural carbohydrate, NSC) and soil carbon. The labile carbon pool is the net of carbon uptake from photosynthesis plus retranslocation from senescing foliage minus autotrophic respiration. In the Ent TBM, autotrophic respiration for most PFTs is tuned to Fluxnet sites to achieve carbon balances with well-quantified site canopy structure (Kim et al. 2015). The global prescribed canopy structure in the simulations in this study may not be in climatological equilibrium with the simulated climate for several reasons. The LAI is from observations from the specific years 2004 for LAI and 2005 for canopy heights. These satellite observations have errors or simplifications,




where the spikes in MODIS LAI are largely smoothed but the interpolated monthly LAI updates in ModelE will miss the shorter time scale of green-up and senescence. The forest canopy heights derived from ICESat/GLAS used in these runs are now known to be ~6 m too tall, given the recent more accurate observations of the Global Ecosystem Dynamic Investigation (GEDI) mission (Dubayah et al. 2020); this results in excess stem biomass from tall trees in the prescribed canopy structure, resulting in 533 PgC in aboveground biomass (AGB) in the eHI experiment (Fig. 23a). This is nearly twice that estimated from harmonized inventory and satellite cover data by Spawn et al. (2020; Fig 23b,e)), which is 287 PgC AGB and total 409 PgC including belowground biomass for the year 2010, or from the GEDI lidar data (Fig. 23c,f; Dubayah et al., 2022), which is 283 PgC AGB and limited to south of 53 N. The GEDI data product predicts lower tropical biomass than Spawn et al. (2020) and higher in the low- to mid-latitudes. Liu et al. (2025) estimate global AGB to be 378 PgC from satellite radar. Fig. 23e,f difference maps show that the tall bias of Ent TBM forests in the Amazon and Congo Basins and in the boreal zone results in high AGB compared to the Spawn et al. (2020) and GEDI estimates. Grassland areas in eHI (Great Plains, United States; Sahel, Africa) have higher AGB than the observational estimates due to a bug in labile carbon storage for grasses in Ent. Besides only recently available satellite data on vertical canopy structure at appropriate spatial scales for vegetation heterogeneity, plant allometry parameterization remains a large issue for all DGVMs for estimating biomass (Grant et al. 2025; Liu et al., 2024; Jucker et al. 2021; Massoud et al. 2019), resulting in large uncertainties in autotrophic respiration that is a function of the plant biomass. Finally, the GCM simulated climate has regional biases; in ModelE2, the tropics have strong warm/dry bias, and in the high latitudes, a strong cold bias causes prolonged frozen soil, hence prolonged simulated winter-spring water stress and low carbon uptake. In the interest of maintaining canopy structure as close to observations as possible rather than predicting wrong vegetation cover and mortality with climate biases, we opt for managing imbalances in the labile carbon pool, instead.


The map in Fig. 24a of plant labile carbon identifies regions where there is insufficient carbon uptake in both boreal evergreen needleleaf and tropical evergreen broadleaf forests. The carbon deficit in forests in these regions is due both to simulated climate biases and overestimated forest stem biomass. The boreal zone is 8-10 C too cold in ModelE, with frozen soils persisting from the winter into what should be the greenup period, causing water stress and reduced GPP; the Amazon Basin has a warm/dry bias causing water stress where there should not be and also reducing GPP. Excess stem biomass from the overestimation of height from the ICESat/GLAS estimate results in high mass-based autotrophic respiration. Thus, the underpredicted GPP and overestimated autotrophic respiration in these forest types results in plant carbon deficits and labile carbon near zero. The label carbon map also highlights

an error in parameterization of grasses that results in too high labile carbon by a few kg/m<sup>2</sup>, particularly in the U.S. interior as was noted also for Fig. 23e,f.

In Fig. 24b, Soil carbon distributions reflect high storage in the high latitudes where permafrost tends to be (ModelE does not explicitly simulate permafrost). The total soil carbon averaged over years 2002-2020 of eHIs 1693 PgC, which is virtually the same as the 1700 PgC from inventory estimates compiled by Canadell et al. 2021 and cited by the GCB2024.



**Figure 23. Land carbon stocks.** Aboveground plant biomass (AGB) from a) the historical simulation (eHI), b) harmonized inventory and satellite-based estimate by Spawn et al. 2020 (year 2010), c) satellite observations from GEDI (3/25/2019-5/22/2019), d) difference between Spawn and GEDI, e) the historical simulation eHI (year 2010) - Spawn dataset, f) the historical simulation eHI - Spawn dataset (year 2010). Globally integrated AGB is 533 PgC for eHI, whereas it is 287 PgC for SPAWN and 283 PgC for GEDI.



**Figure 24.** Plant labile and soil carbon in historical (eHI, averaged years 2000-2020) and change under future climate scenarios, expressed as the difference of average fields in 2080-2100 with respect to 2000-2020 mean. a) Labile carbon, b) soil carbon, c,d) labile and soil carbon change from historical in SSP2-4.5, e,f) labile and soil carbon change from historical

in SSP5-3.4, and g,h) labile and soil carbon change from historical in SSP5-8.5. Units are in carbon density over land area (kg C/m<sup>2</sup>).

Table 8: Plant labile and soil carbon (PgC), giving the eHI 2002-2010 averages, and the change in the future scenarios 2200-2300 climatological average +/- the standard deviation from the historical in labile carbon pools (Fig. 24a,b,c) and in soil carbon pools (Fig. 24d,e,f).

|              | eHI 2002-2010<br>mean(PgC) | SSP2-4.5<br>Δ (PgC) | SSP5-3.4<br>Δ (PgC) | SSP5-8.5<br>Δ (PgC) |
|--------------|----------------------------|---------------------|---------------------|---------------------|
| Labile (PgC) | 211 ± 1.2                  | 16.6 ± 0.5          | 20 ± 0.3            | 22.4 ± 0.9          |
| Soil (PgC)   | 1693 ± 6                   | 135 ± 6             | 106 ± 2             | 211 ± 21            |

Figure 24 shows departures in plant labile (Fig. 24c,e,g) and soil carbon (Fig. 24d,f,h) of the future scenarios (averaged over years 2200-2300 of the runs) from the historical eHI simulation (averaged over years 2002-2010). Both carbon pools overall increase globally the SSP scenarios, but there are pronounced regional declines and enhancements. Plant labile carbon for all scenarios declines in the areas of mixed broadleaf deciduous forest and croplands in the Eastern U.S. and Europe. Relative to the historical eHI, these regions undergo losses in soil moisture (Fig. 17, row c) in addition to warming (Fig. 17, row a) by as much as 2.7 C in SSP2-4.5 and SSP5-3.4 and as high as 4.4 C in SSP5-8.5. Such labile carbon declines imply increased stress for vegetation and bode poorly for tree mortality and crop failure. Plant labile carbon declines also in all scenarios in the steppe/savannah belt between the Sahara and mid-continental Africa, but with pockets of greening. These ecosystems are controlled by soil moisture and seasonal drought, so small changes in soil moisture can induce greening or senescence; the more extensive drying in our simulations could tip these systems toward desertification. Increased plant labile carbon in India, Central Africa, Southern Brazil, and parts of the boreal forest regions of Canada and Eurasia. Except for India, all of these areas experience increased soil moisture in the future scenarios. In India, with its monsoon climate and low water stress, the increased plant labile carbon results from the CO<sub>2</sub> fertilization effect.

Soil carbon in the future scenarios exhibits clear patterns of enhancements corresponding to areas of stimulated GPP and increased litterfall, and major declines in the high latitudes due to thawing of frozen soils and increased soil respiration. In the Eastern U.S. and Europe, however, where there are declines in plant labile carbon, because our model does not simulate mortality and prescribes LAI, the increased soil carbon is due to the model artificially maintaining the vegetation despite low labile carbon for

growth (for these plants the model reduces or even shuts down autotrophic respiration to maintain carbon mass conservation).




Figure 25. a) Combined emissions driven historical and SSP5-8.5 yearly global mean trajectories of cumulative land and ocean carbon uptake and atmospheric carbon storage. The gray dashed line corresponds to the year 2014. b) Components of the carbon budget in terms of CO<sub>2</sub> flux from the emission driven historical and SSP5-8.5 experiments. Positive flux is defined as CO<sub>2</sub> emitted from fossil fuel/industry and land use change, and negative flux is defined as partitioning of CO<sub>2</sub> into the atmosphere, land, and ocean reservoirs. The land component excludes land use change. Left and right of the gray dashed line correspond to the historical and future periods, respectively.

All storage reservoirs grow (Fig. 25) but more is stored in the atmosphere by the end of the 21st century than in the land and the ocean.

#### 4.5 Köppen-Geiger Climate Classifications

In Fig. 26, we show the Köppen-Geiger classifications of observed 2001-2009 (ERA5) climate and the simulated climates, following the classification algorithms of Kottek et al. (2006), which produce 31 climate classes, listed in Appendix Table B. For the tropics, we implemented a slight modification. Where Kottek et al. (2006) defined a 6-month summer, we used 4-month seasons for tropical climates As and Aw (northern hemisphere, summer = June-September, winter = November-February). This modification identifies the rain shadow areas in East Africa, Sri Lanka, and maybe a little too much of northeastern Brazil.

The observed climatology was classified at 0.5°x0.5°, because the coarser ModelE grid cannot capture a less common climate type, Dsd. We note also that because the Köppen-Geiger scheme uses only precipitation and temperature data, it cannot capture wetland areas like those in the southern end of the Amazon Basin, which are fed by stream- and riverflow; these areas classified as Aw (pink,

Commented [AR17]: Figure and the text associated with it will go to a subsequent paper.

36 equatorial savannah) are in reality Af or Am (dark red, equatorial rainforest, or bright red, equatorial  
37 monsoon). Compared to observations, the PI and historical runs show that ModelE2 has a warm/dry  
38 bias in the Amazon and Congo basins, producing the climates more suited to savannah than tropical  
39 forest. ModelE2 also has a cold bias in the northern high latitudes; where observed climate shows Dfb  
40 and Dfc (snow fully humid), ModelE2 simulates EF and ET (polar frost and polar tundra). ModelE  
41 simulates the Western U.S. as moister (Cs, warm temperature with dry summer) than observed (Bw  
42 and Bw, arid), and does not produce the northeast Asia Dwc climate (subarctic/boreal taiga/boreal  
43 forest). ModelE simulates Central and Western Australia as BSh (arid steppe hot), more conducive to  
44 shrubs and grasses than observed BWh (arid desert hot).

45 The simulated climate change between the PI and the end of the historical period shows a slight  
46 migration of high latitude climate type Dfc (bright purple, boreal forest) northward, with no changes in  
47 the mid-latitudes and tropics. This northward migration continues with the SSP future scenarios, along  
48 with encroachment of lower elevation classes to higher elevation around the Tibetan Plateau, and  
49 moistening of southern Chile. In SSP-585, climate Dfb (mixed conifer/deciduous forest) expands  
50 northward, replacing the southern edge of Dfc, and Australia dries compared to the historical and  
51 other scenarios.

52




Figure 26. Köppen-Geiger climate classifications of observed 2001-2009 climatology (ERA5) and the 5 runs. Colorscheme is from Kottek et al. (2006). Observed climate from ERA5 at 0.5°x0.5° resolution.

Commented [AR18]: Figure now includes only historical comparisons. The rest of the figure will go to a subsequent paper

#### 4.6 TCRE & Carbon Cycle Feedbacks

Commented [AR19]: Section removed. Will go to a subsequent paper.

To quantify the response of surface temperature to carbon emissions, we estimate the slope of surface warming to cumulative emissions at different warming and emissions levels. From the eA0 simulation

we obtain the transient climate response to emissions (TCRE) to be 1.6 °C/1000 GtC after 70 years and 1.4 °C/1000 GtC after 140 years. Therefore TCRE is declining in eA0, due to the weakening of the AMOC (Romanou et al, in prep). Compared to Ito et al. (2020) TCRE has increased in GISS2.1-G-CC2 (it used to be 1.1 C/1000GtC). Other CMIP6 models generally show slightly higher TCRE compared to here at 1.6-2 C/1000 GtC (Sanderson et al., 2025) whereas AR5 estimated 1.65 C/1000 GtC pointing to smaller remaining carbon budgets for any given temperature target. IPCC AR6 suggested a range for TCRE between 1C-2.3C per 1000 GtC with the best estimate of about 1.65.

Similarly, the sensitivity parameter associated with CO<sub>2</sub> (i.e. the fraction of TCR/change in concentration of CO<sub>2</sub> in the atmosphere) for E2.1-G-CC2 is found to be 0.006, as estimated from the slope of the change in surface temperature over the change in atmospheric CO<sub>2</sub> concentration for doubling of CO<sub>2</sub> from the eA0 simulation. This value is in accordance with other CMIP6 models with ECS on the lower range of values whereas “hot models” give sensitivity up to 0.11C (Intergovernmental Panel on Climate Change, 2023).

To quantify the response of the carbon cycle storage on land and ocean to the changes in atmospheric CO<sub>2</sub> concentrations and climate change we follow the integrated flux method (Arora et al., 2020; Arora et al, 2013; Friedlingstein et al., 2006). Our results are compared to other CMIP6 and CMIP5 models reported in Arora et al. (2020) and to Ito et al. (2020) since the physical model and forcings are the same with the exception of different land use change, balanced land flux and an updated ocean biogeochemistry. All changes to the carbon fluxes and the atmospheric concentration of CO<sub>2</sub> are computed with respect to the initial year. The change in total carbon land and ocean storage is computed based on the cumulative flux at a given year of the eA0 simulation. Surface air temperature change is computed with respect to the mean of the initial 10 years and the mean of the 10 years prior to the desired year and the timeseries are smoothed using a 10-year running mean.

For the computation of the feedback parameters  $\beta$  and  $\gamma$  we followed C4MIP recommendation (Jones et al., 2016) performed two additional simulations, a biogeochemically coupled simulation in which radiation responds to preindustrial CO<sub>2</sub> concentrations in the atmosphere (BGC run) and a radiatively coupled simulation in which the carbon cycle responds to preindustrial carbon concentrations in the atmosphere (RAD run). Then


$$\beta = \frac{\Delta C_{BGC}}{\Delta C_{BGC}^A} \quad (1)$$



$$\gamma = \frac{\Delta C_{RAD}}{\Delta T_{RAD}} \quad (2)$$

$$\gamma = \frac{\Delta C_{COU} - \Delta C_{BGC}}{\Delta T_{COU}} \quad (3)$$

where,  $\Delta T_{\{BGC,RAD,COU\}}$  is the change in carbon land or ocean content (PgC) in the BGC, RAD or the fully coupled simulation (COU),  $\Delta C_{\{BGC,RAD\}}^A$  is the change in the atmospheric carbon concentration (ppm) in the BGC run, and  $\Delta T_{\{RAD,COU\}}$  is the surface air temperature change (°C) in the RAD and COU experiments. The changes are computed from the preindustrial. The results are shown in Fig. 27.



**Figure 27. Carbon cycle feedbacks: a) Carbon-concentration feedback,  $\beta$ , b) carbon-climate feedback,  $\gamma$ , as computed from equation (2) dashed lines and (3) continuous lines, and c) fractional contribution of each feedback and the airborne fraction to the total carbon in the system.**

The carbon-concentration parameter,  $\beta$ , is positive for both land and ocean and indicates a negative feedback as the increased uptake in the land and ocean tends to reduce the atmospheric concentration. As Fig. 27a shows,  $\beta$  increases with time and peaks after 80 years from the preindustrial values for the ocean and 90 years for land and then declines as the land and the ocean uptake slows down and the atmospheric fraction increases.  $\beta$ 's peak because after 80 years the atmospheric fraction increases

20 faster than the ocean and land storage terms due to significant slowdown of the land uptake after  
21 year 60 (see Fig. 14i and 18f for the eA0 scenario).

22  
23 The carbon-climate parameter,  $\gamma$ , is negative for both the land and the ocean so the feedback of the  
24 land and the ocean carbon uptake to surface warming is a positive feedback, as decreasing uptake  
25 warms the surface and leads to less uptake. We find that  $\gamma$  continues to decline throughout the  
26 simulation and is substantially larger for land than for the ocean (Fig. 27b).  $\gamma$ 's are decreasing because  
27 warming leads to less uptake by the ocean and by the terrestrial carbon.

28  
29 Compared to Ito et al. (2020) we find that  $\beta_{\text{land}}$  and the max  $\beta_{\text{ocean}}$  are lower by about 0.1 1GtC/ppm  
30 at 2xCO<sub>2</sub> (70 years into the simulation eA0). It is most likely attributed to the better spinup of the ocean  
31 carbon cycle than in Ito et al., (2020) since we reduced the drift in the air-sea flux of CO<sub>2</sub> which led to  
32 less uptake in the early years of the historical simulation. On the other hand,  $\gamma_{\text{land}}$  has a steeper decline  
33 than in Ito et al. (2020) owing to the better spin up of the soil carbon, while  $\gamma_{\text{ocean}}$  includes both the  
34 better spin up and the improved tunings of the ocean carbon cycle e.g. the the remineralization  
35 dependence on temperature is lower in these runs than in Ito et al. (2020) leading to smaller transfer of  
36 organic to inorganic carbon per degree warming and hence less outgassing smaller and smaller  $\gamma_{\text{ocean}}$ .  
37 The fact is that  $\gamma_{\text{ocean}}$  has changed substantially with much lower values that are in agreement with  
38 other CMIP models (Arora et al., 2020).

39  
40 CMIP5/6 models estimated that at 4xCO<sub>2</sub> (140 years into the eA0 simulation)  $\beta$  for land is about  $0.97 \pm$   
41  $0.40$  PgC/ppm and for ocean  $0.79 \pm 0.07$  PgC/ppm. For  $\gamma$ , the CMIP6 models' estimate is  $-45.1 \pm 50.6$   
42 PgC/C for land and  $-17.2 \pm 5$  PgC/C for ocean (Arora et al., 2020).

43  
44 The GISS CMIP6 value for  $\beta$  was  $0.85$  GtC/ppm for the ocean and  $0.68$  GtC/ppm for land (Ito et al.,  
45 2020) while the current value is  $0.8$  and  $0.7$  GtC/ppm respectively. Both values fall within the uncertainty  
46 range estimated from CMIP6 models discussed above. The ocean carbon-climate parameter,  $\gamma$ , for the  
47 GISS CMIP6 version was found to be  $-17.7$  GtC/K whereas here it is  $-15$  GtC/K whereas  $\gamma_L$  in the GISS  
48 CMIP6 release was  $-54.8$  GtC/K whereas here it is  $-65$  GtC/K and within the uncertainty of the CMIP6  
49 models (Arora et al., 2020).

50 The relative contributions of the two carbon cycle feedbacks to the atmospheric carbon content are  
51 shown in Fig. 27c. The different components are determined as in Ito et al. (2020) and we find here as

well that the contribution of the  $\beta$  from land and ocean is much larger than the contribution of the  $\gamma$ 's, with the land components always larger than the ocean components.

## 5 Synopsis and Conclusions

In this paper we described the development, skill and characteristics of the NASA GISS Earth system model coupled to emissions of CO<sub>2</sub>. With respect to the CMIP6 model with emissions (Ito et al, 2020), GISS2.1-CC2 has the same physical climate formulation with non-interactive aerosols which are prescribed from fully interactive runs. Additionally, GISS2.1-CC2 includes interactive oxygen and improved iron distributions. The model also has been run to equilibrium more consistently than in CMIP6. The land carbon physics are the same as in CMIP6, but the improved LUHv2 historical crop cover dataset has been used to replace older estimates used in CMIP6.

The model has improved pre-industrial carbon fields and lower carbon drift than the previous version of the model (Ito et al., 2020). It also includes a better crops and land use forcing product (GCB-LUH2 2020) and various improvements in the ocean carbon cycle (iron ligands, oxygen coupled to ocean biogeochemistry, improved nitrogen fixation). However, ocean primary production remains as low as in the CMIP6 version of the model. While some of the model biases (Latto and Romanou 2018; Marshall et al., 2017; Romanou et al., 2017) are certainly attributed to biases in the underlying ocean model (e.g. low wind speed, Arctic circulation and hydrography) many of the biases in the carbon cycle stem from biases in different parameterizations (e.g. remineralization and nitrogen fixation), missing denitrification sinks for nitrate, sea ice effects on primary production, and a discrepancy between the iron distributions in the preindustrial control climate and the historical simulations which leads to a strong decline in productivity early in the 20th century simulation (Lerner et al, 2021; Lerner et al, 2022).

The land carbon model overall predicts global land GPP, NPP, and soil carbon well within the mean of estimates from multi-model intercomparisons (Friedlingstein et al. 2025) of fluxes, and nearly exactly matches inventory compilations for soil carbon (Canadell et al. 2021). Predicted GPP and NPP may be at the lower end compared to other current DGVMs, but the model's global GPP of 127 Pg/yr at 2021 is within the 146.1±21.3 uncertainties estimated by Cheng et al. 2017 (the primary source for IPCC Sixth Assessment Report Working Group, Canadell et al. 2021, and the GCB2024). The model grossly overpredicts vegetation biomass due to prescription of forest heights from early satellite lidar estimates from ICESat/GLAS (Simard et al, 2011), which we plan to replace with new more accurate estimates from GEDI and ICESat-2. There is a bug in the storage of plant labile carbon for grasses, causing excess

**Commented [AR20]:** Text adjusted to include only PI and historical simulations analysis and evaluation against observations. Also, comparison with the previous version of the model.

accumulation in the plants instead of senescing as litterfall. Meanwhile, the model's configuration in biophysics-only mode with only prescribed vegetation structure does not allow for vegetation change through greening, mortality, or competition; however, the prognostic carbon pools of plant labile carbon and soil carbon provide useful diagnostics for vegetation responses to climate changes and CO<sub>2</sub> fertilization, and the assumption of mature forest canopy structure may be closer to reality than the overpredicted forest growth currently from DGVMs in the GCB2024. The model does not represent several LUC processes, notably deforestation, wood harvest, vegetation and peat fires, resulting in missing about 1 PgC/year in LUC-related emissions compared to GCB2024 models over 2009-2019, and that deficit is likely larger over the preindustrial through 20th century. The cumulative land sink since 1850 is still very poorly constrained, so it is better not to reference other estimates currently, which differ wildly, but rather to scrutinize self-consistency within the model. We conclude that the model's lack of these LUC processes causes a deficit in simulated cumulative fluxes to the atmosphere, relative to the model's other flux components, and may explain for ModelE's slightly underpredicted atmospheric CO<sub>2</sub> over most of the historical period.

Overall the ModelE2.1-CC2 predicts the atmospheric concentrations of CO<sub>2</sub> consistent with prescribed emissions given by Community Emissions Data System (CEDS) and much the mean inter-model difference of 7 ppm found in Hajima et al. (2025).

The current version (NASA ModelE2.1-CC2) has been included in several model intercomparison studies where its performance can be directly assessed against other state-of-the-art Earth system models. In coordinated experiments with a data-assimilated model of the historical ocean, the Ocean Circulation Inverse Model (OCIM; Holzer et al, 2021; Devries 2014) historical and future emissions of CO<sub>2</sub> were estimated from trawling disturbances to the ocean floor (Atwood et al., 2024). From the GISS model both concentration driven as well as emissions driven simulations were used and the results showed that the GISS model (ModelE2.1-CC2) estimates of the fraction of CO<sub>2</sub> that escapes to the atmosphere agree very well with OCIM and OCIM for both concentration and emissions driven runs, with the latter providing significant uncertainty associated with internal climate variability of CO<sub>2</sub> which is not included in concentration driven simulations.

In a 13-model intercomparison study and an ambitious effort to estimate properly the remaining carbon budgets (Silvy et al, 2024) till 1.5 and 2 degrees warming levels, was shown to be consistently within the model spread, except perhaps the land carbon sink, which is however very poorly constrained by observations.

'18  
'19 Similarly, in Sanderson et al., 2025 which investigated 10 comprehensive climate models' and 3  
'20 intermediate complexity models' response to positive, zero and negative emissions of CO<sub>2</sub>, the GISS  
'21 ModelE2.1-CC2 response was within the uncertainty range of the ensemble mean and only exhibited  
'22 the strongest response in the later phase of the carbon dioxide removal experiments due to the  
'23 weakened AMOC.

'24  
'25 Future development in the ocean biogeochemistry will focus on adding diel migration as an  
'26 improvement to the biological pump parameterization in the model, coupling ocean biogeochemistry  
'27 with sea ice, the addition of denitrification and closing the nitrogen cycle, adding particulate inorganic  
'28 carbon to improve alkalinity and the CaCO<sub>3</sub> pump, improvements in the underwater light distributions  
'29 and photoadaptation of phytoplankton, and coupling to physical ocean via albedo and changes in  
'30 stratification.

'31  
'32 Future developments for the land model include releasing a geometric optical radiative transfer  
'33 (GORT) canopy model for global production runs. This canopy radiative transfer model, the Analytical  
'34 Clumped Two-Stream (ACTS) model (Ni-Meister et al. 2010; Yang et al., 2010) leverages the cohort  
'35 structure of the Ent TBM analytically to solve for foliage clumping in heterogeneous canopies in two-  
'36 stream scheme. This approach simulates vertical foliage profiles (VFP) as observed by satellite lidar,  
'37 which will enable ingestion of these profiles from GEDI and ICESat-2. The combination of foliage  
'38 clumping prediction and VFPs then allows prediction of the effects of zenith angle and gaps on canopy  
'39 spectral albedo, vertical light profiles for light competition, and transmittance of radiation to the  
'40 ground for predicting soil temperature and snowmelt. We have also completed an analysis (Grant et  
'41 al. 2025) of the Tallo dataset of tree allometry (Jucker et al., 2022), in which we identified significant  
'42 distinctions between plant functional types; these new allometric parameterizations will be  
'43 introduced into the Ent TBM along with more recent satellite lidar observations of forest canopy  
'44 heights. Combining these canopy physics and updated canopy structure and allometry will allow  
'45 reduction in uncertainty in vegetation biomass; it will provide a fusion between vegetation  
'46 demography for simulating plant competition for light with emerging satellite lidar constraints on  
'47 vertical canopy structure. This framework will allow addressing persistent issues in vegetation-  
'48 atmosphere coupling, including the snow-vegetation albedo feedback and heterogeneity of open tree-  
'49 grass systems. In addition, fire-enabled dynamics coupling both the land carbon cycle and chemistry  
'50 are under development, with fuel loads from Ent exchanged with an updated version of ModelE's fire  
'51 chemical emissions model, pyrE (Mezuman et al. 2020), adding CO<sub>2</sub> as an emission. These new fire

'52 dynamics include natural fires and peat fires, and mortality of vegetation will be enabled in tiers of  
'53 ecological dynamics, from simple stand thinning to full community dynamics. The patch dynamics  
'54 implemented for fires will later enable implementation of deforestation.

'55  
'56 With regards to the global carbon-climate interactivity future versions will include interactive dust  
'57 utilizing modelE's state-of-the-art atmospheric composition model (MATRIX, Bauer et al. 2008) and  
'58 emissions of CO<sub>2</sub> from deforestation from wildfires using GISS pyrE (Mezuman et al. 2020) as well as  
'59 land use change, peats and wetlands. These future model development efforts will aim to develop  
'60 more comprehensive simulations of the Earth System subject to assessments against pioneering NASA  
'61 observational products such as from PACE and SWOT, as well as the hyperspectral Earth Surface  
'62 Mineral Dust Source Investigation (EMIT) (Ochoa et al. 2025).

'65 **Acknowledgments**

'66 We thank James Lui for assistance with processing land carbon and Fluxsat data and John Mekus for  
'67 assistance in the analysis and the model simulations. AR, NYK, PL were supported by the NASA  
'68 Modeling, Analysis, and Prediction (MAP) program. Resources supporting this work were provided by  
'69 the NASA High-End Computing (HEC) Program through the NASA Center for Climate Simulation (NCCS)  
'70 at Goddard Space Flight Center.

'71 **References**

'72 Abramopoulos, F., C. Rosenzweig and B. J. Choudhury: "Improved ground hydrology calculations for global climate models  
'73 (GCMs): soil water movement and evapotranspiration." *Journal of Climate* 1: 921-941, 1988.  
'74  
'75 Adebisi, A. A., & Kok, J. F.: Climate models miss most of the coarse dust in the atmosphere. *Science Advances*, 6(15).  
'76 eaaz9507. [https:// doi.org/10.1126/sciadv.aaz9507](https://doi.org/10.1126/sciadv.aaz9507), 2020.  
'77  
'78 Armstrong, R. A., Lee, C., Hedges, J. I., Honjo, S., and Wakeham, S. G. A new, mechanistic model for organic carbon fluxes in  
'79 the ocean based on the quantitative association of POC with ballast minerals. *Deep Sea Research Part II: Topical Studies in*  
'80 *Oceanography*, 49 (1-3), 219-236, [https://doi.org/10.1016/S0967-0645\(01\)00101-1](https://doi.org/10.1016/S0967-0645(01)00101-1), 2001  
'81  
'82 Atwood TB, Romanou, A., DeVries, T., Lerner, P., Mayorga JS, Bradley D, Cabral RB, Schmidt GA and Sala E: Atmospheric  
'83 CO<sub>2</sub> emissions and ocean acidification from bottom-trawling. *Front. Mar. Sci.* 10:1125137. doi:  
'84 10.3389/fmars.2023.1125137, 2024.  
'85

'86 Ball, J. T. and J. A. Berry. A model predicting stomatal conductance and its contribution to photosynthesis under different  
'87 environmental conditions. *Progress in Photosynthesis Research*. I. Biggins. Nijhoff, Dordrecht, Netherlands. IV: 110-112,  
'88 1987.

'89

'90 Bastos, A., Sullivan, M., Ciais, P., Makowski, D., Sitch, S., Friedlingstein, P., Chevalier, F., Rodenbeck, C., Pongratz, J., Lujckx, I.,  
'91 Patra, P., Peylin, P., Canadell, J., Lauerwald, R., Li, W., Smith, N., Peters, W., Goll, D., Jain, A., Kato, E., Lienert, S., Lombardozzi,  
'92 D., Haverd, V., Nabel, J., Tian, H., Walker, A., and Zaehle, S.: Aggregated regional estimates of net atmosphere-land CO<sub>2</sub>  
'93 fluxes from the five atmospheric inversions and 16 Dynamic Global Vegetation Models, supplemental data to Bastos et al,  
'94 2019 (<https://doi.org/10.1029/2019GB006393>), Integrated Carbon Observation System, [http://dx.doi.org/10.18160/1SVH-](http://dx.doi.org/10.18160/1SVH-3DNB)  
'95 3DNB, 2020.

'96

'97 Batjes, N. H.. "Total carbon and nitrogen in the soils of the world." *European Journal of Soil Science* 47(2): 151-163, 1996.

'98

'99 Berger, A. L.. Long-Term Variations of Daily Insolation and Quaternary Climatic Changes. *J. Atmos. Scie.*, 35, 12, 2362–2367.  
'00 [https://doi.org/10.1175/1520-0469\(1978\)035<2362:LTVODI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2362:LTVODI>2.0.CO;2), 1978

'01

'02 Bauer, S.E., D. Wright, D. Koch, E.R. Lewis, R. McGraw, L.-S. Chang, S.E. Schwartz, and R. Ruedy: MATRIX (Multiconfiguration  
'03 Aerosol TRacker of mIXing state): An aerosol microphysical module for global atmospheric models. *Atmos. Chem. Phys.*, 8,  
'04 6603-6035, doi:10.5194/acp-8-6003-2008, 2008.

'05

'06 Bitz, C. M., and Lipscomb, W. H.. An energy-conserving thermodynamic model of sea ice. *J. Geophys. Res. -Oceans*, 104, C7,  
'07 15669-15677. doi: 10.1029/1999JC900100, 1999.

'08

'09 Brutsaert, W.: *Evaporation into the Atmosphere: Theory, History, and Applications*. Springer, Dordrecht, 299.  
'10 <http://dx.doi.org/10.1007/978-94-017-1497-6>, 1982.

'11

'12 Buitenhuis, E. T., Pangerc, T., Franklin, D. J., Le Quéré, C., & Malin, G. (2008). Growth rates of six coccolithophorid strains as  
'13 a function of temperature. *Limnology and Oceanography*, 53(3), 1181-1185.

'14

'15 Jackson, G. A., and Burd, A. B. Simulating aggregate dynamics in ocean biogeochemical models. *Progress in Oceanography*,  
'16 133, 55-65, <https://doi.org/10.1016/j.pocean.2014.08.014>, 2015.

'17

'18 Cairns, B., Lacis, A. A., and Carlson, B. E.. Absorption within Inhomogeneous Clouds and Its Parameterization in General  
'19 Circulation Models. *J. Atmos. Scie*, 57, 5, 700–714. [https://doi.org/10.1175/1520-0469\(2000\)057<0700:AWICAI>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<0700:AWICAI>2.0.CO;2),  
'20 2000.

'21

'22 Cakmur, R.V., Miller, R.L., and Torres, O.: Incorporating the effect of small scale circulations upon dust emission in an AGCM.  
'23 *J. Geophys. Res.*, 109, D07201, doi:10.1029/2003JD004067, 2004.

'24

'25 Canadell, J. G., P.M.S. Monteiro, M.H. Costa, L. Cotrim da Cunha, P.M. Cox, A.V. Eliseev, S. Henson, M. Ishii, S. Jaccard, C.  
'26 Koven, A. Lohila, P.K. Patra, S. Piao, J. Rogelj, S. Syampungani, S. Zaehle, and K. Zickfeld,: Global Carbon and Other  
'27 Biogeochemical Cycles and Feedbacks, in: *Climate Change 2021 – The Physical Science Basis: Working Group I Contribution*  
'28 *to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Masson-Delmotte, V., P. Zhai,  
'29 A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R.  
'30 Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (Cambridge University Press, Cambridge, 673–816, DOI:  
'31 10.1017/9781009157896.007, 2021.

'32

Canuto, V.M., and M.S. Dubovikov: A dynamical model for turbulence. I. General formalism. *Phys. Fluids*, 8, 571-585, doi:10.1063/1.868842, 1996a.

Canuto, V.M., and M.S. Dubovikov: A dynamical model for turbulence. II. Shear-driven flows. *Phys. Fluids*, 8, 587-598, doi:10.1063/1.868843, 1996b.

Caprara, S., Buck, K. N., Gerringa, L. J., Rijkenberg, M. J., & Monticelli, D.: A compilation of iron speciation data for open oceanic waters. *Frontiers in Marine Science*, 3, 221, <https://doi.org/10.3389/fmars.2016.00221>, 2016.

Castellanos, P., P. Colarco, W.R. Espinosa, S.D. Guzewich, R.C. Levy, R.L. Miller, M. Chin, R.A. Kahn, O. Kempainen, H. Moosmüller, E.P. Nowotnick, A. Rocha-Lima, M.D. Smith, J.E. Yorks, and H. Yu: Mineral dust optical properties for remote sensing and global modeling: A review. *Remote Sens. Environ.*, 303, 113982, doi:10.1016/j.rse.2023.113982, 2024.

Chen, C.-T., and Millero, F. J., 1976. The specific volume of seawater at high pressures. *Deep-Sea Research* 23:595–612.

Cheng, Y., V.M. Canuto, and A.M. Howard: An improved model for the turbulent PBL. *J. Atmos. Sci.*, 59, 1550-1565, doi:10.1175/1520-0469(2002)059<1550:AIMFTT>[2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<1550:AIMFTT>2.0.CO;2), 2002.

Cheng, L., Zhang, L., Wang, Y.-P., Canadell, J. G., Chiew, F. H. S., Beringer, J., Li, L., Miralles, D. G., Piao, S., and Zhang, Y.: Recent increases in terrestrial carbon uptake at little cost to the water cycle, *Nature Communications*, 8, 110, 10.1038/s41467-017-00114-5, 2017.

Coddington, O., Lean, J.L, Pilewskie, P., Snow, M., and Lindholm, D.: A Solar Irradiance Climate Data Record. *BAMS*, 97,7, 1265–1282. <https://doi.org/10.1175/BAMS-D-14-00265.1>, 2016.

Collatz, G. J., J. T. Ball, C. Grivet and J. A. Berry. "Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer." *Agricultural and Forest Meteorology* 54: 107-136, 1991.

Collatz, M.-R.C. and Berry, J.A.: Coupled Photosynthesis-Stomatal Conductance Model for Leaves of C4 Plants. *Australian Journal of Plant Physiology* 19(5) 519 - 538, <https://doi.org/10.1071/PP9920519>, 1992.

Cook, B.I., Puma, M.J., and Krakauer, N.Y.: Irrigation induced surface cooling in the context of modern and increased greenhouse gas forcing. *Clim. Dyn.*, 37, 1587-1600, doi:10.1007/s00382-010-0932-x, 2011.

Cook, B.I., J.E. Smerdon, R. Seager, and S. Coats: Global warming and 21st century drying. *Clim. Dyn.*, 43, no. 9-10, 2607-2627, doi:10.1007/s00382-014-2075-y, 2014.

Cox, C., and Munk, W.: Slopes of the sea surface deduced from photographs of sun glitter, *Bull. Scripps Inst. Oceanogr.*, 6, 401–488, 1956.

da Cunha, L. C., Buitenhuis, E. T., Le Quéré, C., Giraud, X., Ludwig, W.: Potential impact of changes in river nutrient supply on global ocean biogeochemistry. *Global Biogeochemical Cycles*, 21, 4. doi: 10.1029/2006GB002718 2007.

Dale, A. W., Sommer, S., Ryabenko, E., Noffke, A., Bohlen, L., Wallmann, K., Stolpovsky, K., Greinert J., and Pfannkuche, O. Benthic nitrogen fluxes and fractionation of nitrate in the Mauritanian oxygen minimum zone (Eastern Tropical North Atlantic). *Geochimica et Cosmochimica Acta*, 134, 234-256. <https://doi.org/10.1016/j.gca.2014.02.026>, 2014



DeVries, T.: The oceanic anthropogenic CO<sub>2</sub> sink: Storage, air-sea fluxes, and transports over the industrial era. *Global Biogeochem. Cycles* 28, 631–647. doi: 10.1002/2013GB004739, 2014.

Doney, S. C., K. Lindsay, I. Fung and J. John. "Natural variability in a stable, 1000-yr global coupled climate-carbon cycle simulation." *Journal of Climate* 19: 3033-3052, 2006.

Dubayah, R. O., Armston, J., Healey, S. P., Yang, Z., Patterson, P. L., Saarela, S., Stahl, G., Duncanson, L., and Kellner, J. R.: GEDI L4B Gridded Aboveground Biomass Density, Version 2, ORNL DAAC, Oak Ridge, Tennessee, USA. [dataset], 2022.

Dubayah, R. O., J. B. Blair, S. Goetz, L. Fatoyinbo, M. Hansen, S. Healey, M. Hofton, G. Hurtt, J. Kellner, S. Luthcke, J. Armston, H. Tang, L. Duncanson, S. Hancock, P. Jantz, S. Marselis, P. L. Patterson, W. Qi and C. Silva (2020). "The Global Ecosystem Dynamics Investigation: High-resolution laser ranging of the Earth's forests and topography." *Science of Remote Sensing* 1: 100002.

ECCO Consortium, Fukumori, I., Wang, O., Fenty, I., Forget, G., Heimbach, P., & Ponte, R. M. (DATE ACCESSED). ECCO Central Estimate (Version 4 Release 4). Retrieved from URL ACCESSED.\*  
 \*DATE ACCESSED and URL ACCESSED to be specified by user.

ECCO Consortium, Fukumori, I., Wang, O., Fenty, I., Forget, G., Heimbach, P., & Ponte, R. M. (2021, February 10). Synopsis of the ECCO Central Production Global Ocean and Sea-Ice State Estimate (Version 4 Release 4). <https://doi.org/10.5281/zenodo.4533349>

Farquhar, G. D. and S. von Caemmerer. Modelling of Photosynthetic Response to Environmental Conditions. *Physiological Plant Ecology II: Water Relations and Carbon Assimilation*. O. L. Lange, P. S. Nobel, C. B. Osmond and H. Ziegler. Berlin, Heidelberg, Springer Berlin Heidelberg: 549-587, 1982.

Finkel, Z. V., Beardall, J., Flynn, K. J., Quigg, A., Rees, T. A. V., Raven, J. A.: Phytoplankton in a changing world: cell size and elemental stoichiometry. *Journal of plankton research*, 32(1), 119-137, doi:10.1093/plankt/fbp098, 2010.

Forget, G., J.-M. Campin, P. Heimbach, C. N. Hill, R. M. Ponte, and C. Wunsch, 2015: ECCO version 4: An integrated framework for non-linear inverse modeling and global ocean state estimation. *Geoscientific Model Development*, 8. <https://www.geosci-model-dev.net/8/3071/2015/>

Friedlingstein, P., and co-authors: Global Carbon Budget 2020. *Earth System Science Data (ESSD)*, 12, 4, 3269–3340. <https://doi.org/10.5194/essd-12-3269-2020>, 2020.

Friedlingstein, P., M. O'Sullivan, M. W. Jones, R. M. Andrew, J. Hauck, P. Landschützer, C. Le Quééré, H. Li, I. T. Lujikx, A. Olsen, G. P. Peters, W. Peters, J. Pongratz, C. Schwingshackl, S. Sitch, J. G. Canadell, P. Ciais, R. B. Jackson, S. R. Alin, A. Arneeth, V. Arora, N. R. Bates, M. Becker, N. Bellouin, C. F. Berghoff, H. C. Bittig, L. Bopp, P. Cadule, K. Campbell, M. A. Chamberlain, N. Chandra, F. Chevallier, L. P. Chini, T. Colligan, J. Decayeux, L. M. Djeutchouang, X. Dou, C. Duran Rojas, K. Enyo, W. Evans, A. R. Fay, R. A. Feely, D. J. Ford, A. Foster, T. Gasser, M. Gehlen, T. Gkritzalis, G. Grassi, L. Gregor, N. Gruber, Ö. Gürses, I. Harris, M. Hefner, J. Heinke, G. C. Hurtt, Y. Iida, T. Ilyina, A. R. Jacobson, A. K. Jain, T. Jarníková, A. Jersild, F. Jiang, Z. Jin, E. Kato, R. F. Keeling, K. Klein Goldewijk, J. Knauer, J. I. Korsbakken, X. Lan, S. K. Lauvset, N. Lefèvre, Z. Liu, J. Liu, L. Ma, S. Maksyutov, G. Marland, N. Mayot, P. C. McGuire, N. Metzl, N. M. Monacchi, E. J. Morgan, S. I. Nakaoka, C. Neill, Y. Niwa, T. Nützel, L. Olivier, T. Ono, P. I. Palmer, D. Pierrot, Z. Qin, L. Resplandy, A. Roobaert, T. M. Rosan, C. Rödenbeck, J. Schwinger, T. L. Smallman, S. M. Smith, R. Sospedra-Alfonso, T. Steinhoff, Q. Sun, A. J. Sutton, R. Sférian, S. Takao, H. Tatebe, H. Tian, B.

27 Tilbrook, O. Torres, E. Tourigny, H. Tsujino, F. Tubiello, G. van der Werf, R. Wanninkhof, X. Wang, D. Yang, X. Yang, Z. Yu, W.  
 28 Yuan, X. Yue, S. Zaehle, N. Zeng and J. Zeng (2025). "Global Carbon Budget 2024." *Earth Syst. Sci. Data* 17(3): 965-1039.  
 29  
 30 Fung, I.Y., S.K. Meyn, I. Tegen, S.C. Doney, J.G. John, and J.K.B. Bishop: Iron supply and demand in the upper ocean. *Glob.*  
 31 *Biogeochem. Cycles*, **14**, 281-296, doi:10.1029/1999GB900059, 2000. Gao, F., J. T. Morissette, R. E. Wolfe, G. Ederer, J. Pedelty,  
 32 E. Masuoka, R. Myneni, B. Tan and J. Nightingale: "An algorithm to produce temporally and spatially continuous MODIS-LAI  
 33 time series." *IEEE Geoscience and Remote Sensing Letters* 5(1): 60-64, 2008.  
 34  
 35 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C., Darmenov, A., Bosilovich, M. G., Reichle,  
 36 R., Wargan, K., Coy, L., Cullathar, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G-Kong, Koster,  
 37 R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S.D., Sienkiewicz,  
 38 M., and Zhao, B. The modern-era retrospective analysis for research and applications, version 2 (MERRA-2). *Journal of*  
 39 *climate*, 30(14), 5419-5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017  
 40  
 41 Gidden, M. J., Riahi, K., Smith, S. J., Fujimori, S., Luderer, G., Kriegler, E., van Vuuren, D. P., van den Berg, M., Feng, L., Klein,  
 42 D., Calvin, K., Doelman, J. C., Frank, S., Fricko, O., Harmsen, M., Hasegawa, T., Havlik, P., Hilaire, J., Hoesly, R., Horing, J., Popp,  
 43 A., Stehfest, E., and Takahashi, K.: Global emissions pathways under different socioeconomic scenarios for use in CMIP6: a  
 44 dataset of harmonized emissions trajectories through the end of the century, *Geoscientific Model Development*, 12, 1443–  
 45 1475, <https://doi.org/10.5194/gmd-12-1443-2019>, 2019.  
 46  
 47 Gill, A. E.: *Atmosphere-Ocean Dynamics*. Academic Press, 662 pp., 1982.  
 48  
 49 GISTEMP Team, 2025: GISS Surface Temperature Analysis (GISTEMP), version 4. NASA Goddard Institute for Space Studies.  
 50 Dataset accessed 2025-09-28 at <https://data.giss.nasa.gov/gistemp/>.  
 51  
 52 Gordon, H. R., and Jacobs, M. M. Albedo of the ocean–atmosphere system: influence of sea foam, *Applied Optics*, 16(8),  
 53 2257-2260, <https://doi.org/10.1364/AO.16.002257>, 1977.  
 54  
 55 Grant, I., Ni-Meister, W., and Kiang, N. Y.: Multiscale analysis of global variation in tree allometric relationships: parameter  
 56 sets for global vegetation models, *Environmental Research: Ecology*, 2025.  
 57  
 58 Grant, S. R., Bienfang, P. K., & Laws, E. A. (2013). Steady-state bioassay approach applied to phosphorus-limited continuous  
 59 cultures: A growth study of the marine chlorophyte *Dunaliella salina*. *Limnology and oceanography*, 58(1), 314-324.  
 60  
 61 Gregg, W.W. and Casey, N. W.: Modeling coccolithophores in the global oceans. *Deep Sea Research Part II: Topical Studies*  
 62 *in Oceanography*, 54, Issues 5–7, 447-477. doi: 10.1016/j.dsr2.2006.12.007, 2007.  
 63  
 64 Gregg, W. W., and Conkright, M. E.: Decadal changes in global ocean chlorophyll. *Geophys. Res. Lett.*, 29, 15, 201-204. doi:  
 65 10.1029/2002GL014689, 2002.  
 66  
 67 Griffies, S. M.: The Gent–McWilliams Skew Flux. *J. Phys. Oceanog.*, 28, 5, 831–841  
 68 doi: 10.1175/1520-0485(1998)028<0831:TGMSF>[2.0.CO](https://doi.org/10.1175/1520-0485(1998)028<0831:TGMSF>2.0.CO;2);2, 1998a.  
 69  
 70 Griffies, S. M., Gnanadesikan, A., Pacanowski, R. C., Larichev, V. D., Dukowicz, J. K., , and Smith, R. D.: Isoneutral Diffusion in  
 71 a z-Coordinate Ocean Model. *J. Phys. Oceanog.*, 28, 5, doi: 10.1175/1520-0485(1998)028<0805:IDIAZC>[2.0.CO](https://doi.org/10.1175/1520-0485(1998)028<0805:IDIAZC>2.0.CO;2);2, 1998b.  
 72  
 73 Hajima, T., Kawamiya, M., Ito, A., Tachiiri, K., Jones, C. D., Arora, V., Brovkin, V., Seferian, R., Liddicoat, S., Friedlingstein, P.,

74 Shevliakova, E.: Consistency of global carbon budget between concentration- and emission-driven historical experiments  
 75 simulated by CMIP6 Earth system models and suggestions for improved simulation of CO2 concentration. *Biogeosciences*,  
 76 22, 5, 1447--1473. <https://bg.copernicus.org/articles/22/1447/2025/>, 2025.  
 77  
 78 Hanninen, H. and K. Kramer: "A framework for modelling the annual cycle of trees in boreal and temperate regions." *Silva*  
 79 *Fennica* 41(1): 167-205, 2007.  
 80  
 81 Hansen, J., Sato, M., Ruedy, R., Kharecha, P., Lacis, A., Miller, R., Nazarenko, L., Lo, K., Schmidt, G. A., Russell, G., Aleinov, I.,  
 82 Bauer, S., Baum, E., Cairns, B., Canuto, V., Chandler, M., Cheng, Y., Cohen, A., del Genio, A., ... Zhang, S.: Climate simulations  
 83 for 1880-2003 with GISS ModelE. *Climate Dynamics*, 29(7–8), 661–696. <https://doi.org/10.1007/s00382-007-0255-8>, 2007.  
 84  
 85 Hansen, J., Russell, G., Rind, D., Stone, P., Lacis, A., Lebedeff, S., Ruedy, R., and Travis, L.: Efficient three-dimensional global  
 86 models for climate studies: Models I and II. *Mon. Weather Rev.*, **111**, 609-662, doi:10.1175/1520-  
 87 0493(1983)111<0609:ETDGMF>[2.0.CO](https://doi.org/10.1175/1520-0493(1983)111<0609:ETDGMF>2.0.CO;2);2, 1983.  
 88  
 89 Hartke, G. J. and Rind, D., . Improved surface and boundary layer models for the Goddard Institute for Space Studies general  
 90 circulation model. *J. Geophys. Res. -Atmospheres*, 102, D14, 16407-16422. doi: 10.1029/97JD00698, 1997.  
 91  
 92 Heavens, N. G., Ward, D. S. & Mahowald, N. M.: Studying and Projecting Climate Change with Earth System Models. *Nature*  
 93 *Education Knowledge* 4(5):4, 2013.  
 94  
 95 Hegarty, T. W.: "Temperature Coefficient (Q10), Seed Germination and Other Biological Processes." *Nature* 243(5405): 305-  
 96 306, 1973.  
 97  
 98 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I.,  
 99 Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J-N: ERA5 monthly averaged data on single levels from 1940 to present.  
 100 Copernicus Climate Change Service (C3S) Climate Data Store (CDS), doi: [10.24381/cds.f17050d7](https://doi.org/10.24381/cds.f17050d7), 2023 (Accessed on 28-09-  
 101 2025).  
 102  
 103 Hirata, T., Hardman-Mountford, N. J., Brewin, R. J. W., Aiken, J., Barlow, R., Suzuki, K., Isada, T.: Synoptic relationships  
 104 between surface Chlorophyll-a and diagnostic pigments specific to phytoplankton functional types. *Biogeosciences*, 8(2), 311-  
 105 327., <https://doi.org/10.5194/bg-8-311-2011>, 2011.  
 106  
 107 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt,  
 108 R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang,  
 109 Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data  
 110 System (CEDS). *GMD*, 11, 369–408. doi: 10.5194/gmd-11-369-2018, 2018.  
 111  
 112 Holl, C. M., and Montoya, J. P.: Interactions between nitrate uptake and nitrogen fixation in continuous cultures of the marine  
 113 diazotroph *Trichodesmium* (cyanobacteria). *Journal of Phycology*, 41(6), 1178-1183, <https://doi.org/10.1111/j.1529-8817.2005.00146.x>, 2007.  
 114  
 115  
 116 Holtslag, A. A. M., and Moeng, C.-H.: Eddy Diffusivity and Countergradient Transport in the Convective Atmospheric Boundary  
 117 Layer. *J. Atm. Scie*, 48,14, 1690–1698. doi: 10.1175/1520-0469(1991)048<1690:EDACT>[2.0.CO](https://doi.org/10.1175/1520-0469(1991)048<1690:EDACT>2.0.CO;2);2, 1991.  
 118  
 119 Holtslag, A. A. M. and Boville, B. A.: Local Versus Nonlocal Boundary-Layer Diffusion in a Global Climate Model. *J. Cli.*, 6, 10,  
 120 1825-1842. doi:10.1175/1520-0442(1993)006<1825:LVNBLD>[2.0.CO](https://doi.org/10.1175/1520-0442(1993)006<1825:LVNBLD>2.0.CO;2);2, 1993.

121 Holzer, M., DeVries, T., and de Lavergne, C.: Diffusion controls the ventilation of a Pacific Shadow Zone above abyssal  
 122 overturning. *Nat. Commun.* 12, 1–13. doi: 10.1038/s41467-021-24648-x, 2021.  
 123  
 124 IES-80. Millero, F. J., Poisson, A., Tung, C. C., Bradshaw, A. L., Schleicher, K.: Background papers and supporting data on the  
 125 International Equation of State of Seawater, 1980. Conference: Joint Panel on Oceanographic Tables and Standards, 11th,  
 126 Sidney, Canada, 1980. UNESCO technical papers in marine science, 38 [67], 192 p., 1981.  
 127  
 128 Intergovernmental Panel on Climate Change: Technical Summary, in: *Climate Change 2021 – The Physical Science Basis:*  
 129 *Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge  
 130 University Press, 35–144, 2023.  
 131  
 132 Ito, G., Romanou, A., Kiang, N. Y., Faluvegi, G., Aleinov, I., Ruedy, R., Russell, G., Lerner, P., Kelley, M., & Lo, K., 2020. Global  
 133 Carbon Cycle and Climate Feedbacks in the NASA GISS ModelE2.1. *Journal of Advances in Modeling Earth Systems*, 12(10),  
 134 1–44. <https://doi.org/10.1029/2019MS002030>, 2020.  
 135  
 136 Jacobson, A. R., Schuld, K. N., Tans, P., Andrews, A., Miller, J. B., Oda, T., et al.: CarbonTracker CT2022. NOAA Global  
 137 Monitoring Laboratory. <https://doi.org/10.25925/Z1GJ-3254>, 2023.  
 138  
 139 Jayne, S. R.: The Impact of Abyssal Mixing Parameterizations in an Ocean General Circulation Model. *J. Phys. Oceanog.*, 39,  
 140 1756–1775, 10.1175/2009JPO4085.1, 2009.  
 141  
 142 Joiner, J., and Y. Yoshida, Y., 2021: Global MODIS and FLUXNET-derived Daily Gross Primary Production, V2. ORNL DAAC, Oak  
 143 Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAAC/1835>, 2021. Date Accessed: 2025-08-28  
 144  
 145 Joiner, J., and Yoshida Y.: Satellite-based reflectances capture a large fraction of variability in global gross primary production  
 146 (GPP) at weekly time scales. *Agricultural and Forest Meteorology* 291:108092.  
 147 <https://doi.org/10.1016/j.agrformet.2020.108092>. 2020.  
 148  
 149 Joiner, J., Yoshida, Y., Zhang, Y., Duveiller, G., Jung, M., Lyapustin, A., Wang, Y. and Tucker, C.J.: Estimation of Terrestrial  
 150 Global Gross Primary Production (GPP) with Satellite Data-Driven Models and Eddy Covariance Flux Data. *Remote Sensing*  
 151 10(9):1346. <https://doi.org/10.3390/rs10091346>, 2018  
 152  
 153 Jucker, T., Fischer, F. J., Chave, J., Coomes, D. A., Caspersen, J., Ali, A., Loubota Panzou, G. J., Feldpausch, T. R., Falster, D.,  
 154 Usoltsev, V. A., Adu-Bredu, S., Alves, L. F., Aminpour, M., Angoboy, I. B., Anten, N. P. R., Antin, C., Askari, Y., Muñoz, R.,  
 155 Ayyappan, N., Balvanera, P., Banin, L., Barbier, N., Battles, J. J., Beeckman, H., Bocko, Y. E., Bond-Lamberty, B., Bongers, F.,  
 156 Bowers, S., Brade, T., van Breugel, M., Chantrain, A., Chaudhary, R., Dai, J., Dalponte, M., Dimobe, K., Domec, J.-C., Doucet,  
 157 J.-L., Duursma, R. A., Enriquez, M., van Ewijk, K. Y., Farfán-Rios, W., Fayolle, A., Forni, E., Forrester, D. I., Gilani, H., Godlee, J.  
 158 L., Gourlet-Fleury, S., Haeni, M., Hall, J. S., He, J.-K., Hemp, A., Hernández-Stefanoni, J. L., Higgins, S. I., Holdaway, R. J.,  
 159 Hussain, K., Hutley, L. B., Ichie, T., Iida, Y., Jiang, H.-s., Joshi, P. R., Kaboli, H., Larsary, M. K., Kenzo, T., Kloeppel, B. D., Kohyama,  
 160 T., Kunwar, S., Kuyah, S., Kvasnica, J., Lin, S., Lines, E. R., Liu, H., Lorimer, C., Loumeto, J.-J., Malhi, Y., Marshall, P. L., Mattsson,  
 161 E., Matula, R., Meave, J. A., Mensah, S., Mi, X., Momo, S., Moncrieff, G. R., Mora, F., Nissanka, S. P., O'Hara, K. L., Pearce, S.,  
 162 Pelissier, R., Peri, P. L., Ploton, P., Poorter, L., Pour, M. J., Pourbabaie, H., Dupuy-Rada, J. M., Ribeiro, S. C., Ryan, C., Sanaei,  
 163 A., Sanger, J., Schlund, M., Sellan, G., Shenkin, A., Sonké, B., Sterck, F. J., Svátek, M., Takagi, K., Trugman, A. T., Ullah, F.,  
 164 Vadeboncoeur, M. A., Valipour, A., Vanderwel, M. C., Vovides, A. G., Wang, W., Wang, L.-Q., Wirth, C., Woods, M., Xiang, W.,  
 165 Ximenes, F. d. A., Xu, Y., Yamada, T., and Zavala, M. A.: Tallo: A global tree allometry and crown architecture database, *Glob.*  
 166 *Change Biol.*, 28, 5254–5268, <https://doi.org/10.1111/gcb.16302>, 2022.

Kattge, J., Diaz, S., Lavorel, S., Prentice, C., Leadley, P., Bonisch, G., Garnier, E., Westoby, M., Reich, P. B., Wright, I. J., Cornelissen, J. H. C., Violle, C., Harrison, S. P., van Bodegom, P. M., Reichstein, M., Enquist, B. J., Soudzilovskaia, N. A., Ackerly, D. D., Anand, M., Atkin, O., Bahn, M., Baker, T. R., Baldocchi, D., Bekker, R., Blanco, C. C., Blonder, B., Bond, W. J., Bradstock, R., Bunker, D. E., Casanoves, F., Cavender-Bares, J., Chambers, J. Q., Chapin, F. S., Chave, J., Coomes, D., Cornwell, W. K., Craine, J. M., Dobrin, B. H., Duarte, L., Durka, W., Elser, J., Esser, G., Estiarte, M., Fagan, W. F., Fang, J., Fernandez-Mendez, F., Fidelis, A., Finegan, B., Flores, O., Ford, H., Frank, D., Freschet, G. T., Fyllas, N. M., Gallagher, R. V., Green, W. A., Gutierrez, A. G., Hickler, T., Higgins, S. I., Hodgson, J. G., Jalili, A., Jansen, S., Joly, C. A., Kerkhoffs, A. J., Kirkup, D., Kitajima, K., Kleyer, M., Klotz, S., Knops, J. M. H., Kramer, K., Kuhn, I., Kurokawa, H., Laughlin, D., Lee, T. D., Leishman, M., Lens, F., Lenz, T., Lewis, S. L., Lloyd, J., Llusia, J., Louault, F., Ma, S., Mahecha, M. D., Manning, P., Massad, T., Medlyn, B. E., Messier, J., Moles, A. T., Muller, S. C., Nadrowski, K., Naeem, S., Niinemets, U., Nollert, S., Nuske, A., Ogaya, R., Oleksyn, J., Onipchenko, V. G., Onoda, Y., Ordóñez, J., Overbeck, G., Ozinga, W. A., Patino, S., Paula, S., Pausas, J. G., Penuelas, J., Phillips, O. L., Pillar, V., Poorter, H., Poorter, L., Poschlod, P., Prinzing, A., Proulx, R., Rammig, A., Reinsch, S., Reu, B., Sack, L., Salgado-Negre, B., Sardans, J., Shiodera, S., Shipley, B., Siefert, A., Sosinski, E., Soussana, J. F., Swaine, E., Swenson, N., Thompson, K., Thornton, P., Waldram, M., Weiher, E., White, M., White, S., Wright, S. J., Yguel, B., Zaehle, S., Zanne, A. E., and Wirth, C.: TRY - a global database of plant traits, *Glob. Change Biol.*, 17, 2905–2935, 10.1111/j.1365-2486.2011.02451.x, 2011.

Kattge, J. and co-authors: TRY – a global database of plant traits. *Global Change Biology*, 17, 9, pp 2905-2935 <https://doi.org/10.1111/j.1365-2486.2011.02451.x>, 2011

Kelley, M., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., del Genio, A. D., Elsaesser, G. S., Faluvegi, G., ... Yao, M. S.: GISS-E2.1: Configurations and Climatology. *Journal of Advances in Modeling Earth Systems*, 12(8). <https://doi.org/10.1029/2019MS002025>, 2020.

Kiang, N. Y., I. Aleinov, W. Ni-Meister, Y. Kim, M. J. Puma and R. D. Koster. The Ent Terrestrial Biosphere Model: Technical Description. NASA Technical Report. NASA, Goddard Space Flight Center. In prep.

Kiang, N. Y., R. D. Koster, P. R. Moorcroft, W. Ni-Meister and D. Rind. Ent: A Dynamic Global Terrestrial Ecosystem Model for Coupling with GCMs. American Geophysical Union Fall Meeting. San Francisco, CA, 2006.

Kim, Y., P. R. Moorcroft, I. Aleinov, M. Puma and N. Y. Kiang. "Variability of phenology and fluxes of water and carbon with observed and simulated soil moisture in the Ent Terrestrial Biosphere Model (Ent TBM version 1.0.1.0.0)." *Geoscientific Model Development* 8: 3837-3865, 2015.

Koster, R. D. and M. J. Suarez. Energy and water balance calculations in the Mosaic LSM, NASA, 1996.

Kopp, G., and Lean, J. L.: A new, lower value of total solar irradiance: Evidence and climate significance. *Geophys. Res. Lett.*, 38, 1. doi: 10.1029/2010GL045777, 2011.

Krakauer, N.Y., M.J. Puma, B.I. Cook, P. Gentile, and Nazarenko, L.: Ocean-atmosphere interactions modulate irrigation's climate impacts. *Earth Syst. Dyn.*, 7, 863-876. doi:10.5194/esd-7-863-2016, 2016.

Krumhardt, K. M., Lovenduski, N. S., Iglesias-Rodriguez, M. D., & Kleypas, J. A. (2017). Coccolithophore growth and calcification in a changing ocean. *Progress in oceanography*, 159, 276-295.

Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., Christian, J. R., Dunne, J., et al.: Twenty-first century ocean warming, acidification, deoxygenation, and upper-ocean nutrient and primary production decline from CMIP6 model

projections. *Biogeosciences*, 17(13), 3439–3470., <https://doi.org/10.5194/bg-17-3439-2020>, 2020.

Lacis, A. A., and Oinas, V.: A description of the correlated k distribution method for modeling nongray gaseous absorption, thermal emission, and multiple scattering in vertically inhomogeneous atmospheres. *J. Geophys. Res. -Atmospheres*, 96, D5, 9027–9063. <https://doi.org/10.1029/90JD01945>, 1991.

Lam, P., and Kuypers, M. M. (2011). Microbial nitrogen cycling processes in oxygen minimum zones: *Annual review of marine science*, 3(1), 317–345, <https://doi.org/10.1146/annurev-marine-120709-142814>, 2011

Landschützer, P., Gruber, N. and Bakker, D. C. E.: Decadal variations and trends of the globalocean carbon sink, *Global Biogeochem. Cycles*, 30, 1396–1417,doi:10.1002/2015GB005359, 2016.

Lan, X., Tans, P. and K.W. Thoning: Trends in globally-averaged CO2 determined from NOAA Global Monitoring Laboratory measurements. <https://doi.org/10.15138/9NOH-ZH07>

Lan, X., K.W. Thoning, and E.J. Dlugokencky: Trends in globally-averaged CH4, N2O, and SF6 determined from NOAA Global Monitoring Laboratory measurements. <https://doi.org/10.15138/P8XG-AA10>

Large, W. G., McWilliams, J. C., and Doney, S. C.: Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parameterization. *Rev. Geophys.* 32, 4, 363–403. doi: 10.1029/94RG01872, 1994.

Latto, R. and Romanou, A.: The Ocean Carbon States Database: a proof-of-concept application of cluster analysis in the ocean carbon cycle, *Earth Syst. Sci. Data*, 10, 609–626, <https://doi.org/10.5194/essd-10-609-2018>, 2018.

Lauvset, S. K., Key, R. M., Olsen, A., Van Heuven, S., Velo, A., Lin, X., et al., Schirnick, C.: A new global interior ocean mapped climatology: The 1× 1 GLODAP version 2. *Earth System Science Data*, 8(2), 325–340., <https://doi.org/10.5194/essd-8-325-2016>, 2016.

Laws, E. A., Pei, S., Bienfang, P., & Grant, S. (2011). Phosphate-limited growth and uptake kinetics of the marine prasinophyte *Tetraselmis suecica* (Kyllin) Butcher. *Aquaculture*, 322, 117–121.

Lawrence, D., Thornton, P. E., Oleson, K. W., Bonan, G. B.: The partitioning of evapotranspiration into transpiration, soil evaporation, and canopy evaporation in a GCM: Impacts on land-atmosphere interaction. *Journal of Hydrometeorology*, 8, 4, 862–880. doi:10.1175/JHM596.1, 2007.

Lenssen, N., G.A. Schmidt, M. Hendrickson, P. Jacobs, M. Menne, and R. Ruedy: A GISTEMPv4 observational uncertainty ensemble. *J. Geophys. Res. Atmos.* 129 (17), 1–18, doi:10.1029/2023JD040179, 2024.

Lerner, P., Marchal, O., Lam, P. J., Buesseler, K., & Charette, M.: Kinetics of thorium and particle cycling along the US GEOTRACES North Atlantic Transect. *Deep Sea Research Part I: Oceanographic Research Papers*, 125, 106–128, <https://doi.org/10.1016/j.dsr.2017.05.003>, 2017.

Lerner, P., A. Romanou, D. Nicholson, M. Kelley, R. Ruedy, and G. Russell, G.: The sensitivity of the Equatorial Pacific ODZ to particulate organic matter remineralization in a climate model under pre-industrial conditions. *Ocean Model.*, 188, 102303, doi:10.1016/j.ocemod.2023.102303, 2024.

Lerner, P., A. Romanou, M. Kelley, J. Romanski, R. Ruedy, and Russell, G. : Drivers of air-sea CO2 flux seasonality and its long-term changes in the NASA-GISS model CMIP6 submission. *J. Adv. Model. Earth Syst.*, 13, no. 2, e2019MS002028,

doi:10.1029/2019MS002028, 2021.

Levis, S., G. B. Bonan, M. Vertenstein and K. W. Oleson: The Community Land Model's Dynamic Global Vegetation Model (CLM-DGVM): Technical Description and User's Guide. Boulder, Colorado, Terrestrial Sciences Section Climate and Global Dynamics Division, National Center for Atmospheric Research, 2004.

Liefer, J. D., Garg, A., Campbell, D. A., Irwin, A. J., & Finkel, Z. V. (2018). Nitrogen starvation induces distinct photosynthetic responses and recovery dynamics in diatoms and prasinophytes. *PLoS One*, 13(4), e0195705.

Liu, G., Ciais, P., Tao, S., Yang, H., and Bastos, A.: A High-Resolution, Long-Term Global Radar-Based Above-Ground Biomass Dataset from 1993 to 2020, *Earth Syst. Sci. Data Discuss.*, 2025, 1–26, 10.5194/essd-2025-330, 2025.

Locarnini, R.A., A.V. Mishonov, O.K. Baranova, J.R. Reagan, T.P. Boyer, D. Seidov, Z. Wang, H.E. Garcia, C. Bouchard, S.L. Cross, C.R. Paver, and D. Dukhovskoy: World Ocean Atlas 2023, Volume 1: Temperature. A. Mishonov, Technical Editor. NOAA Atlas NESDIS 89, 52 pp. <https://doi.org/10.25923/54bh-1613>, 2024

Loth and Graf: Modeling the snow cover in climate studies: 1. Long-term integrations under different climatic conditions using a multilayered snow-cover model. *J. Geophys. Res. - Atmospheres*, 103, D10, 11313-11327. doi: 10.1029/97JD01411, 1998.

Lyman, J. M. and G. C. Johnson: Global High-Resolution Random Forest Regression Maps of Ocean Heat Content Anomalies Using In Situ and Satellite Data. *J. Atmos. Oceanic Tech.*, DOI: 10.1175/JTECH-D-22-0058.1, 2023.

Makela, A., P. Hari, F. Berninger, H. Hanninen and E. Nikinmaa: "Acclimation of photosynthetic capacity in Scots pine to the annual cycle of temperature." *Tree Physiology* 24: 369-376, 2004.

Makela, A., P., J. Kolari, E. Karimaki, M. Nikinmaa, M. Peramaki and P. Hari: "Modelling five years of weather-driven variation of GPP in a boreal forest." *Agricultural and Forest Meteorology* 139(3-4): 382-398, 2006.

Massoud, E. C., Xu, C. G., Fisher, R. A., Knox, R. G., Walker, A. P., Serbin, S. P., Christoffersen, B. O., Holm, J. A., Kueppers, L. M., Ricciuto, D. M., Wei, L., Johnson, D. J., Chambers, J. Q., Koven, C. D., McDowell, N. G., and Vrugt, J. A.: Identification of key parameters controlling demographically structured vegetation dynamics in a land surface model: CLM4.5(FATES), *GEOSCIENTIFIC MODEL DEVELOPMENT*, 12, 4133–4164, 10.5194/gmd-12-4133-2019, 2019.

Matthes, K., Funke, B., Andersson, M.E., Barnard, L., Beer, J., Charbonneau, P., Clilverd, M. A., Dudok de Wit, T., Haberleiter, M., Hendry, A., Jackman, C. H., Kretzschmar, M., Kruschke, T., Kunze, M., Langematz, U., Marsh, D. R. M., Maycock, A. C., Misios, S., Rodger, C. J., Scaife, A. A., Seppälä, A., Shangquan, M., Sinnhuber, M., Tourpali, K., Usoskin, I., van de Kamp, M., Verronen, P. T., and Versick, S.: Solar forcing for CMIP6 (v3.2). *Geosci. Model Dev.*, 10, 2247–2302. <https://doi.org/10.5194/gmd-10-2247-2017>, 2017.

Matthews E.: Prescription of land-surface boundary conditions in GISS GCM II: A simple method based on fine-resolution data bases. In: NASA Technical Memo, NASA Goddard Space Flight Center, Institute for Space Studies, New York, NY, pp 21 pp., 1984.

Myers-Smith, I. H., J. T. Kerby, G. K. Phoenix, J. W. Bjerke, H. E. Epstein, J. J. Assmann, C. John, L. Andreu-Hayles, S. Angers-Blondin, P. S. A. Beck, L. T. Berner, U. S. Bhatt, A. D. Björkman, D. Blok, A. Bryn, C. T. Christiansen, J. H. C. Cornelissen, A. M. Cunliffe, S. C. Elmendorf, B. C. Forbes, S. J. Goetz, R. D. Hollister, R. de Jong, M. M. Loran, M. Macias-Fauria, K. Maseyk, S.

Normand, J. Olofsson, T. C. Parker, F.-J. W. Parmentier, E. Post, G. Schaepman-Strub, F. Stordal, P. F. Sullivan, H. J. D. Thomas, H. Tømmervik, R. Treharne, C. E. Tweedie, D. A. Walker, M. Wilmking and S. Wipf (2020). "Complexity revealed in the greening of the Arctic." *Nature Climate Change* 10(2): 106-117.

Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D., Fraser, P., Montzka, S. A., Rayner, P., Trudinger, C., Krummel, P., Beyerle, U., Canadell, J. G., Daniel, J. S., Enting, I., Law, R. M., O'Doherty, S., Prinn, R. G., Reimann, S., Rubino, M., Vollmer, M. K., and Weiss, R., (2016), Historical greenhouse gas concentrations, doi:10.5194/gmd-2016-169, 2016.

Mellor, G. L., and Yamada, T.: Development of a turbulence closure model for geophysical fluid problems. *Rev. Geophys.*, 20, 4, 851-875. <https://doi.org/10.1029/RG020i004p00851>, 1982.

Meng, J., Huang, Y., Leung, D. M., Li, L., Adebisi, A. A., Ryder, C. L., et al.: Improved parameterization for the size distribution of emitted dust aerosols reduces model underestimation of super coarse dust. *Geophysical Research Letters*, 49, e2021GL097287. <https://doi.org/10.1029/2021GL097287>, 2022.

Mezuman, K., Tsigaridis, K., Faluvegi, G., and Bauer, S. E.: The interactive global fire module pyrE (v1.0), *Geosci. Model Dev.*, 13, 3091–3118, 10.5194/gmd-13-3091-2020, 2020.

Miller, R.L., Cakmur, R. V., Perlwitz, J.P., Geogdzhayev, I.V., Ginoux, P., Kohfeld, K.E., Koch, D., Prigent, C., Ruedy, R., Schmidt, G.A., and Tegen, I.: *Mineral dust aerosols in the NASA Goddard Institute for Space Sciences ModelE atmospheric general circulation model*. *J. Geophys. Res.*, **111**, D06208, doi:10.1029/2005JD005796, 2006.

Miller, R. L., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Kelley, M., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., del Genio, A. D., Elsaesser, G. S., ... Yao, M. S.: CMIP6 Historical Simulations (1850–2014) With GISS-E2.1. *Journal of Advances in Modeling Earth Systems*, 13(1), 1–59. <https://doi.org/10.1029/2019MS002034>, 2021.

Mishchenko, M.I., Travis, L. D., and Mackowski, D.W.: T-matrix computations of light scattering by nonspherical particles: A review. *J. Quant. Spectrosc. Radiat. Transfer*, 55, 535-575, doi:10.1016/0022-4073(96)00002-7, 1996.

Myneni, R. B., S. Hoffman, Y. Knyazikhin, J. L. Privette, J. Glassy, Y. Tian, Y. Wang, X. Song, Y. Zhang, G. R. Smith, A. Lotsch, M. Friedl, J. T. Morisette, P. Potava, R. R. Nemani and S. W. Running: "Global products of vegetation leaf area and fraction absorbed PAR from year one of MODIS data." *Remote Sensing of Environment* 83(1-2): 214-231, 2002.

Najjar, R.G., Jin, X., Louanchi, F., Aumont, O., Caldeira, K., Doney, S.C., Dutay, J.-C., Follows, M., Gruber, N., Joos, F., Lindsay, K., Maier-Reimer, E., Matear, R. J., Matsumoto, K., Monfray, P., Mouchet, A., Orr, J. C., Plattner, G.-K., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Weirig, M.-F., Yamanaka, Y., and Yool, A.: Impact of circulation on export production, dissolved organic matter, and dissolved oxygen in the ocean: Results from Phase II of the Ocean Carbon-cycle Model Intercomparison Project (OCMIP-2). *Global Biogeo. Cycles*, 21, 3. doi: 10.1029/2006GB002857, 2007.

Nakanishi, M.: Improvement of the Mellor–Yamada turbulence closure model based on large eddy simulation data. *Bound. Layer Meteor.*, 99, 349–378. <https://doi.org/10.1023/A:1018915827400>, 2001.

Nazarenko, L.S., N. Tausnev, G.L. Russell, D. Rind, R.L. Miller, G.A. Schmidt, S.E. Bauer, M. Kelley, R. Ruedy, A.S. Ackerman, I. Aleinov, M. Bauer, R. Bleck, V. Canuto, G. Cesana, Y. Cheng, T.L. Clune, B.I. Cook, C.A. Cruz, A.D. Del Genio, G.S. Elsaesser, G. Faluvegi, N.Y. Kiang, D. Kim, A.A. Lacis, A. Leboissetier, A.N. LeGrande, K.K. Lo, J. Marshall, E.E. Matthews, S. McDerimid, K.



Mezuman, L.T. Murray, V. Oinas, C. Orbe, C.P. García-Pando, J.P. Perlwitz, M.J. Puma, A. Romanou, D.T. Shindell, S. Sun, K. Tsigaridis, G. Tselioudis, E. Weng, J. Wu, and M.-S. Yao: [Future climate change under SSP emission scenarios with GISS-E2.1](#). *J. Adv. Model. Earth Syst.*, **14**, no. 7, e2021MS002871, doi:10.1029/2021MS002871, 2022.

Nelson, D. M., and Tréguer, P.: Role of silicon as a limiting nutrient to Antarctic diatoms: evidence from kinetic studies in the Ross Sea ice-edge zone. *Marine Ecology Progress Series*, **80**, 255-264, 1992.

Ni-Meister, W., Yang, W. Z., and Kiang, N. Y.: A clumped-foliage canopy radiative transfer model for a global dynamic terrestrial ecosystem model. I: Theory, *Agricultural and Forest Meteorology*, **150**, 881–894, 10.1016/j.agrformet.2010.02.009, 2010.

Ochoa, F., Brodrick, P. G., Okin, G. S., Ben-Dor, E., Meyer, T., Thompson, D. R., and Green, R. O.: Soil and vegetation cover estimation for global imaging spectroscopy using spectral mixture analysis, *Remote Sens. Environ.*, **324**, 114746, <https://doi.org/10.1016/j.rse.2025.114746>, 2025.

Oinas, V., Lacis, A.A., Rind, D., Shindell, D.T., and Hansen, J.E.: Radiative cooling by stratospheric water vapor: Big differences in GCM results. *Geophys. Res. Lett.*, **28**, 2791-2794, doi:10.1029/2001GL013137, 2001.

Orr, J. C. and Epitalon, J.-M.: Improved routines to model the ocean carbonate system: mocsy 2.0. *Geoscientific Model Development*, **8**, 3, 485–499, doi: 10.5194/gmd-8-485-2015, 2015.

Orr, J. C. and Najjar, R. G. and Aumont, O. and Bopp, L. and Bullister, J. L. and Danabasoglu, G. and Doney, S. C. and Dunne, J. P. and Dutay, J.-C. and Graven, H. and Griffies, S. M. and John, J. G. and Joos, F. and Levin, I. and Lindsay, K. and Matear, R. J. and McKinley, G. A. and Mouchet, A. and Oschlies, A. and Romanou, A. and Schlitzer, R. and Tagliabue, A. and Tanhua, T. and Yool, A.: Biogeochemical protocols and diagnostics for the CMIP6 Ocean Model Intercomparison Project (OMIP). *Geoscientific Model Development*, **10**, 6, 2169–2199. doi: 10.5194/gmd-10-2169-2017, 2017.

Parekh, P., Follows, M. J., & Boyle, E.: Modeling the global ocean iron cycle. *Global biogeochemical cycles*, **18**(1), 1-5, <https://doi.org/10.1029/2003GB002061>, 2004.

Prather, M., J.: Numerical advection by conservation of second-order moments. *Journal of Geophys. Res. - Atmospheres*, **91**, D6, 6671-6681, doi: 10.1029/JD091iD06p06671, 1986.

Pérez García-Pando, C., Miller, R.L., Perlwitz, J.P., Rodríguez, S. and Prospero, J.M.: [Predicting the mineral composition of dust aerosols: Insights from elemental composition measured at the Izaña Observatory](#). *Geophys. Res. Lett.*, **43**, no. 19, 10520-10529, doi:10.1002/2016GL069873, 2016.

Perlwitz, J.P., C. Pérez García-Pando, and R.L. Miller, 2015: [Predicting the mineral composition of dust aerosols — Part 1: Representing key processes](#). *Atmos. Chem. Phys.*, **15**, 11593-11627, doi:10.5194/acp-15-11593-2015, 2015.

Perlwitz, J.P., C. Pérez García-Pando, and Miller, R.L.: [Predicting the mineral composition of dust aerosols — Part 2: Model evaluation and identification of key processes with observations](#). *Atmos. Chem. Phys.*, **15**, 11629-11652, doi:10.5194/acp-15-11629-2015, 2015.

Potter, C. S., P. A. Matson and P. M. Vitousek. "Process modeling of controls on nitrogen trace gas emissions from soils worldwide." *Journal of Geophysical Research* 101(D1): 1361-1377, 1996.

Puma, M.J., and Cook, B.I.: Effects of irrigation on global climate during the 20th century. *J. Geophys. Res.*, 115, D16120, doi:10.1029/2010JD014122, 2010.

Quigley, M. S., Santschi, P. H., Hung, C. C., Guo, L., & Honeyman, B.D.: Importance of acid polysaccharides for 234Th complexation to marine organic matter. *Limnology and Oceanography*, 47(2), 367-377. <https://doi.org/10.4319/lo.2002.47.2.0367>, 2002.

Randerson, J. T., F. M. Hoffman, P. E. Thornton, N. M. Mahowald, K. Lindsay, Y. H. Lee, C. D. Nevison, S. C. Doney, G. Bonan, R. Stockli, C. Covey, S. W. Running and I. Y. Fung. "Systematic assessment of terrestrial biogeochemistry in coupled climate-carbon models." *Global Change Biology* 15(10): 2462-2484, 2009.

Randerson, J. T., Li, Y., Fu, W., Primeau, F., Kim, J. E., Mu, M., et al. (2025). The weak land carbon sink hypothesis. *Science Advances*, 11(37), eadr5489. <https://doi.org/10.1126/sciadv.adr5489>

J.R. Reagan, D. Seidov, Z. Wang, D. Dukhovskoy, T.P. Boyer, R.A. Locarnini, O.K. Baranova, A.V. Mishonov, H.E. Garcia, C. Bouchard, S.L. Cross, and C.R. Paver: *World Ocean Atlas 2023*, Volume 2: Salinity. A. Mishonov, Technical Editor, NOAA Atlas NESDIS 90, 52pp. <https://doi.org/10.25923/70qt-9574>, 2024.

Repo, T., A. Makela and H. Hanninen, Eds.: *Modelling frost resistance of trees. Silva Carelica, Modelling to understand forest function*, 1990.

Respati, M. R., Dommenget, D., Segura, H., and Stassen, C.: Diagnosing drivers of tropical precipitation biases in coupled climate model simulations, *Climate Dynamics*, 62, 8691–8709, 10.1007/s00382-024-07355-3, 2024.

Rodriguez-Iturbe, I., A. Porporato, F. Laio and L. Ridolfi: "Plants in water-controlled ecosystems: active role in hydrologic processes and response to water stress - I. Scope and general outline." *Advances in Water Resources* 24(7): 695-705, 2001.

Romanou, A., Gregg, W. W., Romanski, J., Kelley, M., Bleck, R., Healy, R., Nazarenko, L., Russell, G., Schmidt, G. A., Sun, S., & Tausnev, N.: Natural air-sea flux of CO<sub>2</sub> in simulations of the NASA-GISS climate model: Sensitivity to the physical ocean model formulation. *Ocean Modelling*, 66, 26–44. <https://doi.org/10.1016/j.ocemod.2013.01.008>, 2013.

Romanou, A., Romanski, J., & Gregg, W. W.: Natural ocean carbon cycle sensitivity to parameterizations of the recycling in a climate model. *Biogeosciences*, 11(4), 1137–1154. <https://doi.org/10.5194/bg-11-1137-2014>, 2014.

Romanou, A., D. Rind, J.A. Jonas, R.L. Miller, M. Kelley, G. Russell, C. Orbe, L. Nazarenko, R. Latta, and G.A. Schmidt: Stochastic bifurcation of the North Atlantic Circulation under a mid-range future climate scenario with the NASA-GISS ModelE. *J. Climate*, 36, no. 18, 6141-6161, doi:10.1175/JCLI-D-22-0536.1, 2023.

Russell, G. L., Miller, J. R., & Rind, D.: A Coupled Atmosphere-Ocean Model for Transient Climate Change Studies. In *ATMOSPHERE-OCEAN*, Vol. 33, Issue 4, 1995.

Ryan-Keogh, T. J., Thomalla, S. J., Chang, N., and Moalusi, T.: A new global oceanic multi-model net primary productivity data product. *Earth System Science Data*, 15(11), <https://doi.org/10.5194/essd-15-4829-2023> 4829-4848, 2023

Sanderson, B.M., V. Brovkin, R.A. Fisher, D. Hohn, T. Ilyina, C.D. Jones, T. Koenig, C. Koven, H. Li, D.M. Lawrence, P. Lawrence, S. Liddicoat, A.H. Macdougall, N. Mengis, Z. Nicholls, E. O'Rourke, A. Romanou, M. Sandstad, J. Schwinger, R. Séférian, L.T. Sentman, I.R. Simpson, C. Smith, N.J. Steinert, A.L.S. Swann, J. Tjiputra, and T. Ziehn: flat10MIP: An emissions-driven

experiment to diagnose the climate response to positive, zero, and negative CO<sub>2</sub> emissions. *Geosci. Model Dev.*, 18, no. 17, 5699–5724, doi:10.5194/gmd-18-5699-2025, 2025.

Schaaf, C., and Wang, Z.: MODIS/Terra+Aqua BRDF/Albedo Daily L3 Global 0.05Deg CMG V061 [Data set]. NASA Land Processes Distributed Active Archive Center. <https://doi.org/10.5067/MODIS/MCD43C3.061> Date Accessed: 2025-09-28

Schmidt, G. A., Kelley, M., Nazarenko, L., Ruedy, R., Russell, G. L., Aleinov, I., Bauer, M., Bauer, S. E., Bhat, M. K., Bleck, R., Canuto, V., Chen, Y. H., Cheng, Y., Clune, T. L., del Genio, A., de Fainchtein, R., Faluvegi, G., Hansen, J. E., Healy, R. J., ... Zhang, J.: Configuration and assessment of the GISS ModelE2 contributions to the CMIP5 archive. *Journal of Advances in Modeling Earth Systems*, 6(1), 141–184. <https://doi.org/10.1002/2013MS000265>, 2014.

Schmidt, G. A., Ruedy, R., Hansen, J. E., Aleinov, I., Bell, N., Bauer, M., Bauer, S., Cairns, B., Canuto, V., Cheng, Y., Genio, A. del, Faluvegi, G., Friend, A. D., Hall, T. M., Hu, Y., Kelley, M., Kiang, N. Y., Koch, D., Lacis, A. A., ... Russell, G. L., 2006. Present-Day Atmospheric Simulations Using GISS ModelE: Comparison to In Situ, Satellite, and Reanalysis Data. *Source: Journal of Climate*, 19(2), 153–192. <https://doi.org/10.2307/26253812>, 2006.

Schneider, U., Becker, A., Finger, P., Meyer-Christoffer, A., Rudolf, B., and Ziese, M.: GPCC Full Data Reanalysis Version 6.0 at 0.5°: Monthly Land-Surface Precipitation from Rain-Gauges built on GTS-based and Historic Data. , Global Precipitation Climatology Centre, National Oceanic and Atmospheric Administration [dataset], DOI: 10.5676/DWD\_GPCC/FD\_M\_V7\_050, 2011.

Shukla, S.P., Puma, M.J., and Cook, B.I.: The response of the South Asian Summer Monsoon circulation to intensified irrigation in global climate model simulations. *Clim. Dyn.*, 42, no. 1-2, 21–36, doi:10.1007/s00382-013-1786-9, 2014.

Siebert, S. and Kumm, M. and Porkka, M. and D'oll, P. and Ramankutty, N. and Scanlon, B. R.: A global data set of the extent of irrigated land from 1900 to 2005. *Hydrology and Earth System Sciences*, 19, 3, 1521–1545, doi: 10.5194/hess-19-1521-2015, 2015.

Silsbe, G. M., M. J. Behrenfeld, K. H. Halsey, A. J. Milligan, and Westberry, T.K.: The CAFE model: A net production model for global ocean phytoplankton, *Global Biogeochem. Cycles*, 30, 1756–1777, doi:10.1002/2016GB005521, 2016.

Silvy, Y., T.L. Frölicher, J. Terhaar, F. Joos, F.A. Burger, F. Lacroix, M. Allen, R. Bernadello, L. Bopp, V. Brovkin, J.R. Buzan, P. Cadule, M. Dix, J. Dunne, P. Friedlingstein, G. Georgievski, T. Hajima, S. Jenkins, M. Kawamiya, N.Y. Kiang, V. Lapin, D. Lee, P. Lerner, N. Mengis, E.A. Monteiro, D. Paynter, G.P. Peters, A. Romanou, J. Schwinger, S. Sparrow, E. Stofferahn, J. Tjiputra, E. Tourigny, and T. Ziehn: AERA-MIP: Emission pathways, remaining budgets and carbon cycle dynamics compatible with 1.5°C and 2°C global warming stabilization. *Earth Syst. Dyn.*, 15, no. 6, 1591–1608, doi:10.5194/esd-15-1591-2024, 2024.

Simard, M., N. Pinto, J. B. Fisher and A. Baccini: "Mapping forest canopy height globally with spaceborne lidar." *Journal of Geophysical Research-Biogeosciences* 116: G04021, 2011.

Spawn, S. A., C. C. Sullivan, T. J. Lark and H. K. Gibbs (2020). "Harmonized global maps of above and belowground biomass carbon density in the year 2010." *Scientific Data* 7(1): 112.

Spitters, C.J.T., and H.A.J.M. Toussaint, Goudriaan, J.: Separating the diffuse and direct component of global radiation and its implications for modeling canopy photosynthesis Part I. Components of incoming radiation, *Agricultural and Forest*

Meteorology, 38, Issues 1–3, pp 217-229, ISSN 0168-1923, [https://doi.org/10.1016/0168-1923\(86\)90060-2](https://doi.org/10.1016/0168-1923(86)90060-2), 1986.

Tian, Y. H., C. E. Woodcock, Y. J. Wang, J. L. Privette, N. V. Shabanov, L. M. Zhou, Y. Zhang, W. Buermann, J. R. Dong, B. Veikkanen, T. Hame, K. Andersson, M. Ozdogan, Y. Knyazikhin and R. B. Myneni: "Multiscale analysis and validation of the MODIS LAI product - I. Uncertainty assessment." Remote Sensing of Environment 83(3): 414-431, 2002a.

Tian, Y. H., C. E. Woodcock, Y. J. Wang, J. L. Privette, N. V. Shabanov, L. M. Zhou, Y. Zhang, W. Buermann, J. R. Dong, B. Veikkanen, T. Hame, K. Andersson, M. Ozdogan, Y. Knyazikhin and R. B. Myneni: "Multiscale Analysis and Validation of the MODIS LAI Product. II. Sampling Strategy. Remote Sens. Environ., 83:431-441." Remote Sensing of Environment 83(3): 431-441, 2002b

Troen, I.B., Mahrt, L. A simple model of the atmospheric boundary layer; sensitivity to surface evaporation. Boundary-Layer Meteorol 37, 129–148. <https://doi.org/10.1007/BF00122760>, 1986.

Van Marle, M.J.E., R.D. Field, G.R. van der Werf, I.A. Estrada de Wagt, R.A. Houghton, L.V. Rizzo, P. Artaxo, and Tsigaridis, K.: Fire and deforestation dynamics in Amazonia (1973-2014). Glob. Biogeochem. Cycles, 31, no. 1, 24-38, <https://doi.org/10.1002/2016GB005445>, 2017.

van der Does, M., Knippertz, P., Zschenderlein, P., Harrison, R.G., Stuut, J.-B.W.: Particle size traces modern Saharan dust transport and deposition across the equatorial North Atlantic. Atmos. Chem. Phys. 16, 13697–13710. <https://doi.org/10.5194/acp-16-13697-2016>, 2016.

van der Does, M., Brummer, G.-J. A., van Crimpen, F. C. J., Korte, L. F., Mahowald, N. M., Merkel, U., et al.: Tropical rains controlling deposition of Saharan dust across the North Atlantic Ocean. Geophysical Research Letters, 47, e2019GL086867. <https://doi.org/10.1029/2019GL086867>, 2020.

Wada, Y. and Wisser, D. and Bierkens, M. F. P.: Global modeling of withdrawal, allocation and consumptive use of surface water and groundwater resources. Earth System Dynamics, 5, 1, 15-40. doi: 10.5194/esd-5-15-2014, 2014.

Wang, Y.-P., D. D. Baldocchi, R. Leuning, E. M. Falge and T. Vesala: "Estimating parameters in a land-surface model by applying nonlinear inversion to eddy covariance flux measurements from eight FLUXNET sites." Global Change Biology 13: 652–670, 2007.

Warren, S. G. and Wiscombe, W. J.: A Model for the Spectral Albedo of Snow. II: Snow Containing Atmospheric Aerosols. 37, 12, 2734–2745. doi: 10.1175/1520-0469(1980)037<2734:AMFTSA>[2.0.CO](https://doi.org/10.1175/1520-0469(1980)037<2734:AMFTSA>2.0.CO;2);2, 1980.

Westberry, T.K, Silsbe, G. M., Behrenfeld, M. J.: Gross and net primary production in the global ocean: An ocean color remote sensing perspective, Earth-Science Reviews, 237, 104322, ISSN 0012-8252, doi: 10.1016/j.earscirev.2023.104322, 2023.

Xu, X. and Chen, D.: Estimating global annual gross primary production based on satellite-derived phenology and maximal carbon uptake capacity, Environmental Research, 252, 119063, <https://doi.org/10.1016/j.envres.2024.119063>, 2024.

Xu, L., S. S. Saatchi, Y. Yang, Y. Yu, J. Pongratz, A. A. Bloom, K. Bowman, J. Worden, J. Liu, Y. Yin, G. Domke, R. E. McRoberts, C. Woodall, G.-J. Nabuurs, S. de-Miguel, M. Keller, N. Harris, S. Maxwell and D. Schimel (2021). "Changes in global terrestrial live biomass over the 21st century." Science Advances 7(27): eabe9829.

Yang, W. Z., Ni-Meister, W., Kiang, N. Y., Moorcroft, P. R., Strahler, A. H., and Oliphant, A.: A clumped-foliage canopy radiative

transfer model for a Global Dynamic Terrestrial Ecosystem Model II: Comparison to measurements, *Agricultural and Forest Meteorology*, 150, 895–907, 10.1016/j.agrformet.2010.02.008, 2010.

Yang, W. Z., B. Tan, D. Huang, M. Rautiainen, N. V. Shabanov, Y. Wang, J. L. Privette, K. F. Huemmrich, R. Fensholt, I. Sandholt, M. Weiss, D. E. Ahl, S. T. Gower, R. R. Nemani, Y. Knyazikhin and R. B. Myneni: "MODIS leaf area index products: From validation to algorithm improvement." *IEEE Transactions on Geoscience and Remote Sensing* 44(7): 1885-1898, 2006.

Yao, M.-S., and Cheng, Y.: Cloud Simulations in Response to Turbulence Parameterizations in the GISS Model E GCM. *J. Cli*, 25, 14, 4963–4974. <https://doi.org/10.1175/JCLI-D-11-00399.1>, 2012.

Zeng, X., Dickinson, R. E., Barlage, M., Dai, Y., Wang, G.: Treatment of undercanopy turbulence in land models. *Journal of Climate*, 18, 23, 5086-5094. doi:10.1175/JCLI3595.1, 2005.

Zhang, J., and Rothrock, D.: Modeling Arctic sea ice with an efficient plastic solution. *J. Geophys. Res. -Oceans*, 105, C2, 3325-3338. <https://doi.org/10.1029/1999JC900320>, 2000.

## Appendix

### A. Carbon Cycle Model Formulation

#### NOBMg

#### Appendix A: Model Equations

Phytoplankton species (diatoms, chlorophytes, cyanobacteria, coccolithophores):

$$\frac{\partial P_i}{\partial t} = \nabla(K\nabla P_i) - \nabla \cdot \mathbf{u}P_i - \nabla w_{s,i}P_i + [\mu_i - (\delta + \Omega)]P_i - \gamma H - \kappa P_i \quad (\text{A1})$$

The terms on the right-hand side represent mixing, advection, sinking, growth via nutrient uptake, exudation, respiration of DOC, zooplankton grazing, and senescence. The subscript *i* indicates one of the phytoplankton functional groups. Parameterization of the sinking rates ( $w_{s,i}$ ), specific growth rates ( $\mu_i$ ), and the grazing rate ( $\gamma$ ) are described below. The constants  $\delta$ ,  $\Omega$ , and  $\kappa$  are the exudation, respiration, grazing, and senescence rates, respectively, as defined in Gregg and Casey, 2007.

The total specific growth rate ( $d^{-1}$ ) is given by:

$$\mu_i = \mu_{m,i} \min[\omega(E_T)_i, \omega(N_T)_i, \omega(Si)_i, \omega(Fe)_i] RG_i + \mu_{i,cyan} \min[\omega(E_T)_i, \omega(Fe)_{i,Nfx}] RG_i N_{m,i} \quad (\text{A2})$$

where  $\mu_{m,i}$  is the maximum growth rate at 20C (see Table 1). Maximum growth rates were adjusted as described in the section "Tuning."  $\omega(E_T)$  is the growth rate fraction that is function only of the total irradiance ( $\mu\text{mol quanta}/\text{m}^2/\text{s}$ ) and is given below

$$\omega(E_T)_i = \frac{E_T}{E_T + k_{E,i}} \quad (A3)$$

where  $E_T$  is the total irradiance, and  $k_E$  is the irradiance at which  $\mu = 0.5\mu_m$  and  $k_{E,i} = 0.5I_{k,i}$ , where  $I_k$  is the light saturation parameter.

$$I_{k,i} = \begin{cases} I_{k,i,hi}, & E_T > 250 \\ I_{k,i} = I_{k,i,low}, & E_T \leq 25 \\ \left(1 - \frac{E_T - (0.5(250-25)+25)}{250 - (0.5(250-25)+25)}\right) I_{k,i,med} + \left(\frac{E_T - (0.5(250-25)+25)}{250 - (0.5(250-25)+25)}\right) I_{k,i,hi}, & E_T \geq 0.5(250-25) + 25 \\ I_{k,i} = \left(1 - \frac{E_T - 25}{(0.5(250-25)+25) - 25}\right) I_{k,i,low} + \left(\frac{E_T - 25}{(0.5(250-25)+25) - 25}\right) I_{k,i,med}, & E_T < 0.5(250-25) + 25 \end{cases} \quad (A4)$$

Here,  $I_{k,i,hi,med,low}$  are light saturation constants for low, medium, and high irradiance categories and are set following Gregg and Casey, 2007 (see Table 1).

The nutrient dependent growth rates are:

$$\omega(NH_4)_i = \frac{A}{A + k_{N,i}} \quad (A5)$$

$$\omega(NO_3)_i = \frac{N}{N + k_{N,i}} \quad (A6)$$

$$f(NO_3)_i = \min[(NO_3)_i, 1 - \omega(NH_4)_i] \quad (A7)$$

$$\omega(N_T)_i = \omega(NH_4)_i + f(NO_3)_i \quad (A8)$$

$$\omega(Fe)_i = \frac{I}{I + k_{Fe,i}} \quad (A9)$$

$$\omega(Si)_i = \frac{S}{S + k_{Si,i}} \quad (A10)$$

Where  $k_{N,Fe,Si,i}$  are half-saturation constants for nitrogen (ammonium and nitrate), iron, and silica limitation, respectively. Note that while  $k_N$  remains set to the value used by Gregg and Casey, 2007,  $k_{Fe,Si}$  have been adjusted as described in the section "Tuning".

The temperature dependence of the growth rate is from Bissinger (2008):

$$R = e^{-b(T-20)} \quad (A11)$$

Where the constant  $b$ , originally set to 0.0631 to fit phytoplankton data from the Liverpool Plankton Database, has been reduced to 0.031 to improve the regional distribution of net primary productivity (in particular, increase NPP in the subtropics). This also yields a Q10 of 1.36, which is closer to the Q10 obtained from a more recent compilation of phytoplankton community growth rates (1.47; Sherman et al., 2015) than the Q10 value found by Bissinger (2008)

Specifically for cyanobacteria there is an additional temperature dependence which reduces their growth rate in cold waters (colder than 15C, Agawin et al., 1998, 2000).

$$G_i = \begin{cases} 0.0294T + 0.558, & i = \text{cyanobacteria} \\ 1, & i = \text{other PFTs} \end{cases} \quad (\text{A12})$$

Our treatment of nitrogen fixation, which amends the specific growth rate for cyanobacteria (second term in eq. 2), has been revised from that of Gregg and Casey, 2007, with two major differences. In Gregg and Casey 2007, the limitation of growth supported by nitrogen fixation included the same iron limitation (i.e., same half-saturation constant for iron-limitation of cyanobacteria) as non-nitrogen fixation supported growth. This has now been, i.e.:

$$\omega(Fe)_{i,Nfix} = \frac{I}{I + k_{Fe,Nfix}} \quad (\text{A13})$$

Where the half-saturation for iron limitation for nitrogen fixation ( $k_{Fe,Nfix}$ ) was tuned to obtain a global nitrogen fixation rate that is within observational uncertainty, as described in the section “Tuning”.

Additionally, we have replaced the cyanobacterial biomass inhibition of nitrogen fixation with inhibition by the presence of nitrate and ammonium, following findings from laboratory studies using *Trichodesmium* cultures Holl and Montoya, 2005. The inhibition factor ( $N_{in,i}$ ) is set to 0 for all non-cyanobacteria remaining phytoplankton groups as these groups do not fix nitrogen and hence the term representing growth due to nitrogen fixation (second term in eq. 2) is set to zero.

$$N_{in,i} = \begin{cases} \frac{r_{kni}}{N+A+r_{kni}}, & i = \text{cyanobacteria} \\ 0, & i = \text{other PFTs} \end{cases} \quad (\text{A14})$$

Where  $r_{kni}$  is the inhibition constant for nitrogen and ammonium from Holl and Montoya, 2005.

Phytoplankton sinking is treated as additional vertical advection term (following Gregg and Casey 2007) but with additional exponential dependence on diatom concentration for the diatoms and limiting term for coccolithophores which depends on the coccolithophore growth rate. For the other species is a function of viscosity (++++)

$$w_{s,i} = \begin{cases} (0.451 + 0.0178T)a_{diat}e^{P_i b_{diat}}, & i = \text{diatoms} \\ w_{sch} (0.451 + 0.0178T), & i = \text{coccolithophores} \\ (0.451 + 0.0178T)w_{s,i}, & i = \text{other PFTs} \end{cases} \quad (\text{A15})$$

$$w_{sch} = \min(\max(0.752g_{cmax} + 0.225, 0.3), 1.4) \quad (\text{A16})$$

$$g_{cmax} = \max(\mu_{occc}, g_{cmax}) \quad (\text{A17})$$

Note that  $g_{cmax}$ , which is initialized to the value of the coccolithophore growth rate, is updated each time step to either match the coccolithophore growth rate (if the growth rate is larger), or retain it's current value (if the growth rate larger).

Zooplankton grazing increases with temperature and the total biomass of phytoplankton.

$$\gamma = \gamma_m R_H (1 - e^{-\Lambda \sum_i P_i}) \quad (\text{A18})$$

$$R_H = 0.06e^{0.1T} + 0.70 \quad (\text{A19})$$

Where  $\Lambda$  and  $\gamma_m$  are the Ivlev constant and maximum grazing rate at 20°C, respectively (Gregg and Casey, 2007):.

**Nutrients** (nitrate, ammonium, silicate, iron):

The governing equation for nitrate is:

$$\frac{\partial N}{\partial t} = \nabla(K\nabla N) - \nabla \cdot \mathbf{u}N - b_N [\sum_i f_{NO3} \mu_i P_i] + R\alpha_N \frac{D_C}{C:N} + \lambda_D \frac{D_C}{C:N} \quad (\text{A20})$$

Where the terms represent, from left to right, mixing, advection, uptake of nitrate via phytoplankton growth, detrital organic carbon/nitrogen remineralization, and detrital organic carbon/nitrogen breakdown to DOC. Here,  $b_N$  is the nitrogen to chlorophyll ratio, C:N is the stoichiometric ratio of carbon to nitrogen in phytoplankton, and  $\lambda_D$  is the rate of detrital breakdown to DOC. Each of these parameters are set to the values in Gregg and Casey, 2007. Note that  $R$  follows equation (11) except that we retain  $b=0.0631$  (the original value from Bissinger *et al.*, 2008).  $\alpha_N$ , the remineralization rate of organic carbon/nitrogen, follows:

$$\alpha_N = r_{aer} H(1 - e^{-O_2/k_{O2}}) \quad (\text{A21})$$

Where  $r_{aer}$  is the maximum rate of remineralization at 20°C,  $H$  is the step function that equals 1 if  $O_2 > 2$  umol/kg, and 0 otherwise, and  $k_{O2}$  is the e-folding length scale for the oxygen-dependence of remineralization. The latter dependence was included following a study by Keil *et al.*, 2016 that found the remineralization rate to decline exponentially with decreasing oxygen concentration (see also Lerner *et al.*, 2024 for a discussion on how the value of  $k_{O2}$  was assigned).

$f_{NO3}$ , the fraction of nitrogen assimilated into phytoplankton biomass that is obtained from nitrate, follows:

$$f_{NO3} = N/(A + N) \quad (\text{A22})$$

The governing equation for ammonium is:

$$\frac{\partial A}{\partial t} = \nabla(K\nabla A) - \nabla \cdot \mathbf{u}A - b_N [\sum_i f_{NH4} \mu_i P_i] + b_N \epsilon [\gamma H + \eta_2 H^2] \quad (\text{A23})$$

Where the terms represent, from left to right, mixing, advection, uptake of ammonium via phytoplankton growth, and zooplankton excretion. Here,  $\epsilon$  is the zooplankton excretion parameter, and  $f_{NH4}$ , the fraction of nitrogen assimilated into phytoplankton biomass that is obtained from ammonium, follows:

$$f_{NO3} = A/(A + N) \quad (\text{A24})$$

The governing equation for dissolved silica is:

$$\frac{\partial S}{\partial t} = \nabla(K\nabla S) - \nabla \cdot \mathbf{u}S - b_S \mu_1 P_1 + R\alpha_S D_S \quad (\text{A25})$$



Where the terms represent, from left to right, mixing, advection, uptake of silica via phytoplankton growth, dissolution of detrital silica. Here,  $b_s$  is the silica to chlorophyll ratio, and  $\alpha_s$  is the detrital silica dissolution rate  $b_s$  is set to the value in Gregg and Casey, 2007, while  $\alpha_s$  was adjusted as described in the “Tuning” section.

For dissolved iron, the governing equation is:

$$\frac{\partial I}{\partial t} = \nabla(K\nabla I) - \nabla \cdot \mathbf{u}I - b_I[\sum_i \mu_i P_i] + b_I\varepsilon[\gamma H + \eta_2 H^2] + R\alpha_I D_I + \frac{A_{Fe}}{L} - \theta_1 I + \theta_2 I' \quad (A26)$$

Where the terms represent, from left to right, mixing, advection, uptake of iron via phytoplankton growth, zooplankton excretion, remineralization of detrital iron, atmospheric deposition, and the scavenging of ligand-bound and free iron. Here,  $b_I$  is the iron to chlorophyll ratio, which is equal to the value used by Gregg and Casey, 2007.  $\alpha_I$ ,  $\theta_1$ , and  $\theta_2$  are the remineralization rate, ligand-bound and free iron scavenging rates, respectively, whose values are adjusted as described in the “Tuning” section.  $A_{Fe}$  is the atmospheric deposition of iron bound to dust, which is prescribed as detailed in the “Forcings” section, while  $L$  is the thickness of the surface layer.  $I'$  is solved for at each timestep from the system of equations that describe the complexation of iron with a single ligand class:

$$[I] = [IL] + [I'] \quad (A27)$$

$$[L_{tot}] = [IL] + [L'] \quad (A28)$$

$$K_{FeL} = [IL]/([I'][L']) \quad (A29)$$

$L_{tot}$  is from the prescribed ligand pool described in “Forcings.” The stability constant  $K_{FeL}$  is set to a constant value of  $10^{11}$  nmol/L, taken from the average value from the compilation of Caprara et al., 2016.

#### Herbivores:

The governing equations for herbivores is

$$\frac{\partial H}{\partial t} = \nabla(K\nabla H) - \nabla \cdot \mathbf{u}H + (1 - \varepsilon)\gamma H - \eta_1 H - \eta_2 H^2 - \zeta H - \Theta H \quad (A30)$$

Where the terms represent, from left to right, mixing, advection, grazing, zooplankton linear and quadratic mortality, zooplankton excretion of DOC, and zooplankton respiration. Here,  $\eta_1$  and  $\eta_2$  are the linear and quadratic mortality rates, and  $\Theta$  is the herbivore respiration rate. These constants are the same as those used by Gregg and Casey, 2007.  $\zeta$  is the excretion rate of DOC by herbivores, and follows:

$$\zeta = r_H \frac{H}{H_0 + H} R \quad (A31)$$

Where  $r_H$  is the excretion rate of herbivores at 20°C, and  $H_0$  is the half-saturation constant of herbivore excretion, both of which are set equal to the values from Gregg and Casey, 2007.  $R$  is the temperature-dependence of herbivore excretion, which is equal to  $R$  for detrital nitrogen/carbon remineralization (eq. 20).

Particulate organic components (carbon/nitrate detritus, silica detritus, iron detritus):

The governing equation for detrital carbon/nitrogen is:

$$\frac{\partial D_C}{\partial t} = \nabla(K\nabla D_C) - \nabla \cdot \mathbf{u} D_C - \nabla \cdot \mathbf{w}_{d,C} D_C - R\alpha_C D_C + \Phi[\kappa \sum_i P_i + \eta_1 H] + \Phi(1 - \varepsilon)\eta_2 H^2 - \lambda_D D_C \quad (\text{A32})$$

Where the terms represent, from left to right, mixing, advection, sinking, remineralization of detrital iron, phytoplankton senescence, mortality of zooplankton, and breakdown of detrital carbon to DOC.  $\Phi$  is the carbon to chlorophyll ratio, which is a constant value equal to the median (50 g C g chl<sup>-1</sup>) of the three values used to represent three photoadaptation states in Gregg and Casey, 2007.  $\alpha_C$  is the remineralization rate of organic matter, defined as:

$$\alpha_C = \begin{cases} r_{aer}(1 - e^{-O_2/k_{O_2}}), & O_2 \geq 2\mu\text{mol/kg} \\ r_{den}, & O_2 < 2\mu\text{mol/kg} \end{cases} \quad (33)$$

Where  $r_{aer}$  is as defined in equation 21, and  $r_{den}$  is rate of remineralization at 20°C under anaerobic conditions, where oxygen is no longer used as the terminal electron acceptor.

For detrital silica, the governing equation is:

$$\frac{\partial D_S}{\partial t} = \nabla(K\nabla D_S) - \nabla \cdot \mathbf{u} D_S - \nabla \cdot \mathbf{w}_{d,S} D_S - R\alpha_S D_S + b_S[\kappa P_1 + \gamma H] \quad (\text{A34})$$

Where the terms represent, from left to right, mixing, advection, sinking, dissolution of detrital silica, phytoplankton senescence and herbivore grazing.

$$\frac{\partial D_I}{\partial t} = \nabla(K\nabla D_I) - \nabla \cdot \mathbf{u} D_I - \nabla \cdot \mathbf{w}_{d,I} D_I - R\alpha_I D_I + b_I[\kappa \sum_i P_i + \eta_1 H] + b_I(1 - \varepsilon)\eta_2 H^2 + \theta_1 I + \theta_2 I' \quad (\text{A35})$$

Where the terms represent, from left to right, mixing, advection, sinking, remineralization of detrital iron, phytoplankton senescence, linear and quadratic herbivore mortality, and the scavenging of ligand-bound and free-iron.

For equations (32-34), the detrital sinking rates follow an exponential dependence, similar to that for diatoms:

$$\mathbf{w}_{d,C,S,I} = (0.451 + 0.0178T)\mathbf{a}_{C,I,S}e^{P_I b_{C,I,S}} \quad (\text{A36})$$

Where the coefficients  $\mathbf{a}_{C,I,S}$  and  $\mathbf{b}_{C,I,S}$  have been previously described in (Ito et al., 2020).

Carbon components (dissolved organic carbon, dissolved inorganic carbon, alkalinity):

The governing equation for dissolved organic carbon (DOC) is

$$\frac{\partial \text{DOC}}{\partial t} = \nabla(K\nabla \text{DOC}) - \nabla \cdot \mathbf{u} \text{DOC} + \Phi\delta \sum_i \mu_i P_i + \Phi\zeta H + \lambda_D D_C - \varphi \text{DOC} \quad (\text{A37})$$

Where the terms represent, from left to right, mixing, advection, phytoplankton exudation, zooplankton excretion, breakdown of detrital carbon, and DOC remineralization. The DOC remineralization rate follows:

$$\varphi = H \frac{N}{N+k_1} \frac{DOC}{DOC+k_2} DOCR \quad (A38)$$

Where  $H$  is the same step function used in eq. 21.  $\lambda_{DOC}$  is the maximum remineralization rate at 20°C, and  $k_1$  and  $k_2$  are half-saturation constants for limitation of DOC remineralization by nitrogen and DOC, respectively. The constants are the same as those used by Gregg and Casey, 2007. The temperature-dependent factor  $R$  follows the same parameterization as that used for the remineralization of detrital carbon/nitrogen (eq. 20).

The governing equation for dissolved inorganic carbon (DIC) is:

$$\frac{\partial DIC}{\partial t} = \nabla(K \nabla DIC) - \nabla \cdot \mathbf{u} DIC - \Phi \sum_i \mu_i P_i + \Phi \Omega \sum_i \mu_i P_i + \Phi \Theta H + \varphi DOC + R \alpha_C \frac{D_C}{C:N} + \mathcal{F}_{AO\ CO_2} \quad (A39)$$

Where the terms represent, from left to right, mixing, advection, uptake of DIC by phytoplankton, phytoplankton respiration, zooplankton respiration, DOC remineralization, detrital carbon remineralization, and the air-sea exchange of  $CO_2$ . The last term follows the OMIP-BGC protocol for CMIP6 (Orr et al., 2017);

$$\mathcal{F}_{AO\ CO_2} = k_{w,CO_2} ff (xCO_2 - CO_{2,s}) fsice / L \quad (A40)$$

Where  $k_{w,CO_2}$  is the piston velocity of air-sea  $CO_2$  gas exchange,  $ff$  is the solubility of  $CO_2$ ,  $xCO_2$  is the dry-air fraction of atmospheric  $CO_2$  at the surface of the ocean,  $CO_{2,s}$  is the partial pressure of  $CO_{2,s}$  at the surface of the ocean, and  $fsice$  is the fraction of the surface ocean grid cell covered by sea ice. The piston velocity is computed following

$$k_{w,CO_2} = a_{wan} (sc_{CO_2} / 660)^{-0.5} w_s^2 \quad (A41)$$

Where  $a_{wan}$  is the Wanninkhof coefficient (Wanninkhof 2014, Orr et al., 2017),  $w_s$  is the surface wind speed computed by the model, and  $sc_{CO_2}$  is the Schmidt number that follows:

$$sc_{CO_2} = 2116.8 - 135.6 * T + 5.2122T^2 - 0.10939T^3 + 0.00094743T^4 \quad (A42)$$

While the solubility of  $CO_2$  is computed following

$$ff = e^{-A + \frac{B}{tk_{100}} + C \log(tk_{100}) - D tk_{100}^2 + S(E - F tk_{100} + G tk_{100}^2)}$$

Where  $tk_{100} = 100/T$ , and the constant A-G are defined in Orr et al., 2016 (see also Table 1).

#### Alkalinity:

The governing equation for alkalinity is:

$$\frac{\partial Alk}{\partial t} = \nabla(K \nabla Alk) - \nabla \cdot \mathbf{u} Alk + b_N [\sum_i \mu_i P_i] - R \alpha_C \frac{D_C}{C:N} - \lambda_D \frac{D_C}{C:N} + 2 J_{Ca} \quad (A43)$$

Where the terms represent, from left to right, mixing, advection, the increase in alkalinity due to the conversion of nitrate (a proton donor) to organic N, the decrease in alkalinity due to the conversion of organic N to nitrate during remineralization of detrital organic carbon/nitrogen and DOC, and carbonate dissolution and precipitation. Carbonate dissolution and precipitation ( $J_{Ca}$ ) follow the OCMIP-2 protocol:

$$J_{Ca} = \begin{cases} -rr C:P * (1 - \sigma_{Ca}) PP, & z \leq z_c \\ -\frac{dF_{Ca}}{dz}, & z > z_c \end{cases} \quad (A44)$$

Where  $rr$  is the rain ratio of  $\text{CaCO}_3$  to organic phosphorous (Yamanaka and Tajika 1996),  $C:P$  is the organic carbon to organic phosphorous ratio,  $z_c$  represents the compensation depth which we set to a constant value, and  $\sigma_{Ca}$  is the fraction of calcium carbonate that dissolved within the compensation depth (Table 1). The change in the downward flux of calcium carbonate with depth uses a fixed exponential profile:

$$F_{Ca} = rr C:P F_c e^{-\frac{(z-z_c)}{d}} \quad (A45)$$

Where  $d$  is a fixed exponential length scale for carbonate dissolution, and  $F_c$  is the primary productivity integrated from the surface to the compensation depth that is associated with the net production of calcium carbonate within the same depth interval:

$$F_c = (1 - \sigma_{Ca}) \int_0^{z_c} PP \, dz \quad (A46)$$

$PP$  is computed simply as the difference between the specific growth rate and the sum of respiration and exudation, integrated over all phytoplankton functional types:

$$PP = \sum_i \mu_i - (\delta + \Omega) \quad (A47)$$

Oxygen:

The governing equation for dissolved oxygen is:

$$\frac{\partial O_2}{\partial t} = \nabla(K \nabla O_2) - \nabla \cdot \mathbf{u} O_2 + b_o \sum_i \mu_i P_i - b_o \Omega \sum_i \mu_i P_i - b_o \Theta H - \phi \frac{DOC}{C:O} - Ra_c \frac{D_C}{C:O} + \mathcal{F}_{AO \, O_2} \quad (A48)$$

Where the terms represent, from left to right, mixing, advection, production of  $O_2$  by phytoplankton, phytoplankton respiration, zooplankton respiration, DOC remineralization, detrital carbon remineralization, and the air-sea exchange of  $O_2$ .  $b_o$  and  $C:O$  are, respectively, the Redfield ratio of  $O_2$  produced or consumed by autotrophic and heterotrophic processes to chlorophyll and carbon in organic matter, respectively. As for  $CO_2$ , The air-sea exchange of  $O_2$  follows the protocol described in Orr et al., 2017:

$$\mathcal{F}_{AO \, O_2} = k_{w,O_2} (O_{2,sat} - O_{2,s}) f_{sice} / L \quad (A49)$$

Where  $k_{w,O_2}$  is the piston velocity for  $O_2$  exchange between the atmosphere and ocean,  $O_{2,sat}$  is the  $O_2$  concentration at saturation, and  $O_{2,s}$  is the surface concentration of  $O_2$ . The piston velocity is computed following:

$$k_{w,O_2} = a_{wan} (sc_{O_2} / 660)^{-0.5} w_s^2 \quad (A50)$$

Where  $sc_{O_2}$  is the Schmidt number of  $O_2$ , which is computed following:

$$sc_{O_2} = 1920.4 - 136.25T + 4.7353T^2 - 0.092307T^3 + 0.000755T^4$$

The  $O_2$  saturation concentration is also computed from the protocol of Orr et al., 2017:

$$O_{2,sat} = K_{sol}(SLP - pH_2O)xO_2 \quad (A51)$$

Where  $SLP$  is sea level pressure from the model,  $xO_2$  is the  $O_2$  dry-air air fraction, and the solubility coefficient  $K_{sol}$  is computed following:

$$K_{sol} = O_{2,sat0}/(xO_2(1 - pH_2O)) \quad (A52)$$

Where the partial pressure of  $H_2O$  at the surface of the ocean is computed from model temperature and salinity:

$$pH_2O = e^{24.4543 - 67.4509tk_{100} - 4.8489 \log\left(\frac{1}{tk_{100}}\right) - 0.000544S} \quad (A53)$$

$$ta = \log((298.15 - T)/(Ts + 273.15)) \quad (A54)$$

$$tk_{100} = 100/(T + 273.15) \quad (A55)$$

And the saturation concentration of  $O_2$  at standard sea level pressure is computed following the equation of Garcia and Gordon, 1992:

$$O_{2,sat0} = (1e^{-3}) * e^{A+Bta+Cta^2+Dta^3+Eta^4+Fta^5-S(G-Hta+Ita^2-Jta^3)-KS^2} \quad (56)$$

**State variables:**

Table A1

| Variable  | Description                | partitioning                                           |
|-----------|----------------------------|--------------------------------------------------------|
| $P_i$     | Phytoplankton species      | Diatoms, chlorophytes, cyanobacteria, coccolithophores |
| $H$       | Herbivores                 |                                                        |
| $N$       | nitrate                    |                                                        |
| $A$       | ammonium                   |                                                        |
| $S$       | silicate                   |                                                        |
| $I$       | iron                       |                                                        |
| $D_{C/N}$ | Detritus C/N               |                                                        |
| $D_S$     | Detritus Silica            |                                                        |
| $D_I$     | Detritus Iron              |                                                        |
| DOC       | Dissolved Organic Carbon   |                                                        |
| DIC       | Dissolved Inorganic Carbon |                                                        |
| Alk       | Alkalinity                 |                                                        |

'96

'97

'98

'99

'00

'01

'05

'06

'13

'14

'15

'16

|           |                                    |  |
|-----------|------------------------------------|--|
| $O_2$     | Oxygen                             |  |
| $O_{2,s}$ | Surface ocean oxygen concentration |  |

Atmospheric variables needed by NOBMg

Table A2

| Variable   | Description                        |
|------------|------------------------------------|
| $X_{CO_2}$ | Mole fraction of oxygen in dry air |
| $w_s$      | Surface wind speed                 |

Sea-ice variables needed by NOBMg

Table A3

| Variable   | Description      |
|------------|------------------|
| $f_{sice}$ | Sea ice fraction |

Physical parameters

Table A4

| parameter | description                         | partitioning | value                     |
|-----------|-------------------------------------|--------------|---------------------------|
| $K$       | Horizontal and vertical diffusivity |              |                           |
| $u$       | Ocean velocity                      |              |                           |
| $E_T$     | Total Irradiance                    |              |                           |
| $a_{wan}$ | gas exchange coefficient            |              | 0.251 cm hr <sup>-1</sup> |

Phytoplankton parameters

Table A5

| parameter | description                           | partitioning                                                 | value |
|-----------|---------------------------------------|--------------------------------------------------------------|-------|
| $w_s$     | Sinking rate of phytoplankton at 31°C | Diatoms<br>Chlorophytes<br>Cyanobacteria<br>coccolithophores |       |

|             |                                                                                                                  |                                                                                 |                                                                                              |                                                      |
|-------------|------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|------------------------------------------------------|
| $a_{diot}$  | diatom sinking rate linear factor                                                                                |                                                                                 | 0.05 d <sup>-1</sup>                                                                         | 2717                                                 |
| $b_{diot}$  | diatom sinking rate exponential factor                                                                           |                                                                                 | 5.0                                                                                          | 2719                                                 |
| $\mu_{m,i}$ | Specific growth rate max at 20°C                                                                                 | Diatoms<br>Chlorophytes<br>Cyanobacteria<br>coccolithophores                    | 3.00 d <sup>-1</sup><br>1.89 d <sup>-1</sup><br>0.72 d <sup>-1</sup><br>1.70 d <sup>-1</sup> | 2721<br>2722<br>2723<br>2724                         |
| $b$         | Temperature dependence for growth, grazing, DOC degradation, and remineralization,                               | growth<br>grazing<br>DOC degradation<br>remineralization                        | 0.03 d <sup>-1</sup><br>0.06 d <sup>-1</sup><br>0.06 d <sup>-1</sup><br>0.06 d <sup>-1</sup> | 2725<br>2726<br>2727<br>2728                         |
| $\delta$    | Phytoplankton DOC exudation rate                                                                                 |                                                                                 | 0.05 d <sup>-1</sup>                                                                         | 2729                                                 |
| $\Omega$    | Phytoplankton respiration rate                                                                                   |                                                                                 | 0.05 d <sup>-1</sup>                                                                         | 2730                                                 |
| $\kappa$    | Senescence rate                                                                                                  |                                                                                 | 0.05 d <sup>-1</sup>                                                                         | 2731                                                 |
| $r_{kni}$   | Factor determining nitrogen fixation inhibition by nitrate abundance                                             |                                                                                 | 7.0                                                                                          | 2732                                                 |
| $r_{ik}$    | Half-saturation constants for light as a function of light level ( $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$ ) | Low medium high<br>Diatoms<br>Chlorophytes<br>Cyanobacteria<br>coccolithophores | 90.0 93.0<br>96.9 87.0<br>65.1 66.0<br>56.1 71.2                                             | 2733<br>2734<br>2735<br>2736<br>2737<br>2738<br>2739 |

Zooplankton parameters

Table A6

| parameter | description                                | partitioning | value                   |
|-----------|--------------------------------------------|--------------|-------------------------|
| $\mu_m$   | Grazing rate maximum at 20°C               |              | 1.2 d <sup>-1</sup>     |
| $\Lambda$ | Ilev constant                              |              | 1 d <sup>-1</sup>       |
| $r_H$     | Zooplankton excretion rate of DOC at 20 °C |              | 0.05 d <sup>-1</sup>    |
| $H_0$     | Half-saturation constant for               |              | 0.24 mg m <sup>-3</sup> |

'50  
'51  
'52  
  
'53  
'54  
'55  
'56  
'57  
'58  
'59  
'60  
'61  
'62  
'63

|               |                                    |                  |                                            |      |
|---------------|------------------------------------|------------------|--------------------------------------------|------|
|               | zooplankton DOC excretion at 20 °C |                  |                                            | 2744 |
| $\theta$      | Herbivore respiration rate         |                  | 0.05 d <sup>-1</sup>                       | 2745 |
| $\eta_{1,2}$  | heterotrophic mortality rates      | Linear quadratic | 0.1 d <sup>-1</sup><br>0.5 d <sup>-1</sup> | 2746 |
| $\varepsilon$ | Zooplankton excretion rate         |                  | 0.05 d <sup>-1</sup>                       | 2747 |
|               |                                    |                  |                                            | 2748 |
|               |                                    |                  |                                            | 2749 |

**Nutrient parameters**

Table A7

| parameter     | description                                            | partitioning                                                                                           | value                                                              |
|---------------|--------------------------------------------------------|--------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------|
| $b_{N,S,I}$   | Nutrient: Chlorophyll ratio for nitrogen, silica, iron |                                                                                                        | (C:N) <sup>-1</sup> ( $\Phi$ )                                     |
| $C:N$         | Carbon : nitrogen ratio                                |                                                                                                        | 106:16 (mol C/mol N)                                               |
| $C:S$         | Carbon : silica ratio                                  |                                                                                                        | 106:16(mol Si/mol N)                                               |
| $C:Fe$        | Carbon : iron ratio                                    |                                                                                                        | 50,000 (mol Fe /mol N)                                             |
| $C:P$         | Carbon : phosphorous ratio                             |                                                                                                        | 117:1(mol P/mol N)                                                 |
| $k_{N,Fe,Si}$ | Nutrient half-saturation constants                     | diatoms N, Fe, Si<br>chlorophytes N, Fe, Si<br>cyanobacteria. N, Fe, Si<br>coccolithophores. N, Fe, Si | 1.0,0.12,2.85 μM<br>0.67,0,0.09 μM<br>2,0,0.08 μM<br>0.5,0,0.08 μM |
| $k_{Fe,Nfix}$ | Half-saturation constant for iron in nitrogen fixation |                                                                                                        | 0.12 μM                                                            |



Detrital Parameters

Table A8

| parameter      | description                                                | partitioning                      | value                                                                          |
|----------------|------------------------------------------------------------|-----------------------------------|--------------------------------------------------------------------------------|
| $\alpha_{S,I}$ | Remineralization rate at 20°C for silica, iron             |                                   | 0.002 d <sup>-1</sup>                                                          |
| $a_{C,S,I}$    | Detrital sinking rate linear factor                        | Carbon/Nitrogen<br>Silica<br>Iron | 2 d <sup>-1</sup><br>3 d <sup>-1</sup><br>1 d <sup>-1</sup>                    |
| $b_{C,S,I}$    | Detrital sinking rate exponential factor                   | Carbon/Nitrogen<br>Silica<br>Iron | 3 d <sup>-1</sup><br>6 d <sup>-1</sup><br>1 d <sup>-1</sup>                    |
| $r_{aer}$      | Aerobic remineralization rate at 20°C for Carbon/nitrate   |                                   | 0.3 d <sup>-1</sup>                                                            |
| $r_{den}$      | anaerobic remineralization rate at 20°C for Carbon/nitrate |                                   | 0.12 d <sup>-1</sup>                                                           |
| $\theta_{1,2}$ | Dissolved iron scavenging rate                             | Total<br>Free                     | 2.74x10 <sup>-3</sup> d <sup>-1</sup><br>2.74x10 <sup>-2</sup> d <sup>-1</sup> |
| $\lambda_D$    | Detrital breakdown rate at 20°C                            |                                   | 0.05 d <sup>-1</sup>                                                           |

Carbon Parameters

| parameter       | description                                                        | partitioning               | value                                                                 |
|-----------------|--------------------------------------------------------------------|----------------------------|-----------------------------------------------------------------------|
| $\Phi$          | Carbon: chlorophyll ratio                                          |                            | 50 mg C (mg chl) <sup>-1</sup>                                        |
| $\lambda_{DOC}$ | DOC remineralization rate                                          |                            | 0.05 d <sup>-1</sup>                                                  |
| $K_{1,2}$       | Half-saturation constant for DOC remineralization                  | N constant                 | 3.0 μM<br>15.0 μM                                                     |
| $A,B,C,D,E,F,G$ | coefficients for CO <sub>2</sub> solubility from Orr et al., 2016. | A<br>B<br>C<br>D<br>E<br>F | -160.733<br>215.4152<br>-1.47759<br>0.029941<br>0.027455<br>0.0053407 |

Alkalinity parameters

Table A9

| parameter            | description                                    | partitioning                                                                                             | value                                         |
|----------------------|------------------------------------------------|----------------------------------------------------------------------------------------------------------|-----------------------------------------------|
| $rr$                 | Rain ratio                                     |                                                                                                          | $0.07 \text{ mol CaCO}_3 (\text{mol P})^{-1}$ |
| $\sigma_{\text{Ca}}$ | DOC remineralization rate                      | fraction of net downward flux of organic matter across the compensation depth that is in dissolved form. | $0.67 \text{ d}^{-1}$                         |
| $d$                  | Length scale for calcium carbonate dissolution |                                                                                                          | 3500 m                                        |
| $z_c$                | Compensation depth                             |                                                                                                          | 75 m                                          |

Oxygen parameters

Table A10

| parameter                         | description                                              | partitioning                                        | value                                                                                                                                     |
|-----------------------------------|----------------------------------------------------------|-----------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| $k_{\text{O}_2}$                  | e-folding length scale for aerobic remineralization rate |                                                     | $40 \text{ } \mu\text{mol O}_2 \text{ kg}^{-1}$                                                                                           |
| $x_{\text{O}_2}$                  | Mole fraction of oxygen in dry air                       |                                                     | 0.20946                                                                                                                                   |
| $A, B, C, D, E, F, G, H, I, J, K$ | coefficients for oxygen saturation at reference pressure | A<br>B<br>C<br>D<br>E<br>F<br>G<br>H<br>I<br>J<br>K | 5.80818<br>3.20684<br>4.11890<br>4.93845<br>1.01567<br>1.41575<br>0.00701211<br>0.00725958<br>0.00793334<br>0.000554491<br>0.000000132412 |

Table A11

| parameter    | description             | partitioning  | value    | parameter    |
|--------------|-------------------------|---------------|----------|--------------|
| $A_{Fe}$     | Dust iron deposition    | Forcing field | Variable | 2824         |
| $L$          | Layer thickness         |               | Variable | 2825         |
| $\Delta z_s$ | Surface layer thickness |               | variable | 2826<br>2827 |

2828

Ent Terrestrial Biosphere Model

The complete formulation of the land carbon dynamics is given in Kim et al., 2015. In this study, prognostic growth is not turned on, but the vegetation model is run in “biophysics-only” mode, as described in Section 3.

B. Köppen-Geiger classes

Table B1. Köppen-Geiger biome/climate classes, with color scheme from <http://koeppen-geiger.vu-wien.ac.at/>


|    | Class | Color             | Description                                         |
|----|-------|-------------------|-----------------------------------------------------|
| 1  | Af    | dark red          | Equatorial rainforest                               |
| 2  | Am    | bright red        | Equatorial monsoon                                  |
| 3  | As    | dark pink         | Equatorial savannah with dry summer                 |
| 4  | Aw    | pale pink         | Equatorial savannah with dry winter                 |
| 5  | BWk   | pale yellow       | Arid desert cold                                    |
| 6  | BWh   | medium yellow     | Arid desert hot                                     |
| 7  | BSk   | cool brown        | Arid steppe cold steppe                             |
| 8  | BSh   | warm yellow-brown | Arid steppe hot                                     |
| 9  | Csa   | bright green      | Warm temperate with dry hot summer                  |
| 10 | Csb   | yellow green      | Warm temperate with dry warm summer                 |
| 11 | Csc   | yellower green    | Warm temperate with dry cool summer and cold winter |
| 12 | Cwa   | warm brown        | Warm temperate with dry winter and hot summer       |
| 13 | Cwb   | medium cool brown | Warm temperate with dry winter and warm summer      |
| 14 | Cwc   | dark brown        | Warm temperate with dry cold winter and cool summer |
| 15 | Cfa   | dark green        | Warm temperate fully humid with hot summer          |

|    |     |                    |                                                             |
|----|-----|--------------------|-------------------------------------------------------------|
| 16 | Cfb | medium green       | Warm temperate fully humid with warm summer                 |
| 17 | Cfc | medium vivid green | Warm temperate fully humid with cool summer and cold winter |
| 18 | Dsa | magenta            | Snow with dry hot summer                                    |
| 19 | Dsb | pale magenta       | Snow with dry warm summer                                   |
| 20 | Dsc | paler magenta      | Snow with dry cool summer and cold winter                   |
| 21 | Dwa | pale grey purple   | Snow with dry winter and hot summer                         |
| 22 | Dwb | medium grey purple | Snow with dry winter and warm summer                        |
| 23 | Dwc | grey purple        | Snow with dry cold winter and cool summer                   |
| 24 | Dwd | deep grey purple   | Snow with dry winter extremely continental                  |
| 25 | Dfa | black purple       | Snow fully humid with hot summer                            |
| 26 | Dfb | red purple         | Snow fully humid with warm summer                           |
| 27 | Dfc | bright purple      | Snow fully humid with cool summer and cold winter           |
| 28 | Dfd | red fuchsia        | Snow fully humid extremely continental                      |
| 29 | EF  | grey blue          | Polar frost                                                 |
| 30 | ET  | turquoise          | Polar tundra                                                |
|    | UA  | light gray 1       | Unknown equatorial cool                                     |
|    | UAu | light gray 2       | Unknown equatorial dry                                      |
|    | UB  | light gray 3       | Unknown arid                                                |
|    | UE  | light gray 4       | Unknown polar                                               |
|    | Ufu | light gray 5       | Unknown warm temp. with snow                                |
|    | Uuu | light gray 6       | Unknown other                                               |


37

38

39



41  
42 Figure SUPP1: SLP, wsurf and SSS from eHI and observations  
43  
44



**Figure SUPP2: Land fluxes and soil carbon from eHI, with annual (thin line), 20-year moving average (thick line), and zero line (dashed line). a) air-to-land flux, b) cumulative land uptake since 1850, c) soil carbon stocks, d) land gross primary productivity (GPP), e) land autotrophic respiration, f) land soil respiration, g) emissions due to land use change (LUC), h) land net primary productivity (NPP), i) land net ecosystem productivity (NEP), and j) land net biome productivity (NBP).**




Figure SUPP3: Land carbon seasonality from eHI and FluxSat.




Figure SUPP4: Averaged climatologies of the last 20 years in SSPs.