

# Deployment and Evaluation of a Low-Cost Sensor System for Atmospheric CO<sub>2</sub> Monitoring on a Sea-Air Interface Buoy

Jialu Liu<sup>1,2</sup>, Pengfei Han<sup>3\*</sup>, Huiling Ouyang<sup>1,4</sup>, Ning Zeng<sup>5,6</sup>, Zhenfeng Wang<sup>1</sup>, Jian Wang<sup>1</sup>, Weiwei Fu<sup>1,2</sup>, Honggang Lv<sup>7</sup>, Wenhao Lin<sup>8</sup>, Zheng Xia<sup>1,9,10</sup>, Bo Yao<sup>1,2, 11\*</sup>

<sup>1</sup>Department of Atmospheric and Oceanic Sciences and Institute of Atmospheric Sciences, Fudan University, Shanghai, 200438, China

<sup>2</sup>Shanghai Frontiers Science Center of Atmosphere-Ocean Interaction, Key Laboratory of Polar Atmosphere-Ocean-Ice System for Weather and Climate of Ministry of Education, and Shanghai Key Laboratory of Ocean-Land-Atmosphere Boundary Dynamics, Fudan University, Shanghai, 200438, China

<sup>3</sup>State Key Laboratory of Atmospheric Environment and Extreme Meteorology, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, 100029, China

<sup>4</sup>Observation and Research Station of Huaniaoshan East China Sea Ocean-Atmosphere Integrated Ecosystem, Ministry of Natural Resources, Shanghai, 200136, China

<sup>5</sup>Department of Atmospheric and Oceanic Science, University of Maryland, College Park, Maryland, 20742, USA

<sup>6</sup>Earth System Science Interdisciplinary Center, University of Maryland, College Park, Maryland, 20742, USA

<sup>7</sup>Key Laboratory of Marine Hazards Forecasting, National Marine Environmental Forecasting Center, Ministry of Natural Resources (MNR), Beijing, 100081, China; National Marine Environmental Forecasting Center, Beijing, 100081, China

<sup>8</sup>Zhejiang Environmental Monitoring Engineering, Co.Ltd., 310013, China

<sup>9</sup>Zhejiang Ecological and Environmental Monitoring Center, Hangzhou, 310012, China

<sup>10</sup>Zhejiang Key Laboratory of Ecological and Environmental Monitoring, Forewarning and Quality Control, Hangzhou, 310012, China

<sup>11</sup>National Observation and Research Station for Wetland Ecosystems of the Yangtze Estuary, Shanghai, 201112, China

*Correspondence to:* Bo Yao (yaobo@fudan.edu.cn); Pengfei Han (pphan@mail.iap.ac.cn)

**Abstract.** Direct in-situ observation of marine CO<sub>2</sub> concentrations is crucial for estimating air-sea CO<sub>2</sub> fluxes, yet such observations remain scarce. Drawn on experiences from urban CO<sub>2</sub> monitoring and buoy-based measurements, this study deployed a sea-air interface buoy platform in the northern South China Sea, near Maoming, Guangdong Province, China. This platform was equipped with three low-cost SenseAir K30 sensors to enable continuous atmospheric CO<sub>2</sub> measurement. This paper presents the first detailed account of the methodology, encompassing hardware design, environmental corrections, land-based validation tests, offshore deployment procedures, and initial observational results. These findings thus provide valuable insights for advancing marine CO<sub>2</sub> observations practices. To mitigate the impacts of temperature, humidity, and pressure on sensor readings — while simultaneously compensating for zero-drift — an environmental correction method was implemented. This approach significantly improved data accuracy: after 1-minute temporal averaging of raw data and a subsequent 1-hour moving average, the root mean square error was reduced from 8.03 ppm to 3.64 ppm in land tests and from 24.26 ppm to 1.59 ppm in marine observations. ~~This approach significantly improved data accuracy: in land tests, the root-mean-square errors was reduced from 8.03 ppm to 3.64 ppm; in marine observations, the root-mean-square errors decreased from 24.26 ppm to 1.59 ppm.~~ Importantly, this level of precision meets the requirements for resolving sea surface CO<sub>2</sub> dynamics (~420-480 ppm). Observed concentrations were consistent with HYSPLIT-simulated long-range atmospheric transport, revealing the stable and homogeneous nature of the marine atmospheric boundary layer, with diurnal variations of approximately 3 ppm, and capturing localized or short-term fluctuations due to terrestrial carbon sources. These results demonstrate the effectiveness of the method, offering a low-cost, high-density solution for marine atmospheric CO<sub>2</sub> monitoring and providing key inputs for inversely estimating ocean carbon sink.

Keywords: marine atmospheric CO<sub>2</sub>; low-cost sensors; environmental correction; buoy measurements

带格式的: 左侧: 1.27 厘米, 右侧: 1.27 厘米, 顶端: 1.27 厘米, 底端: 1.27 厘米

## 1 Introduction

20 Since the Industrial Revolution, less than half of the carbon emitted into the atmosphere by human activities remains in the atmosphere (Friedlingstein et al., 2020; Costa et al., 2023), highlighting the pivotal role of terrestrial and ocean sinks in regulating atmospheric CO<sub>2</sub> concentrations. From 2013 to 2022, the ocean absorbed and stored CO<sub>2</sub> approximately 26% of total anthropogenic CO<sub>2</sub> emissions (Friedlingstein et al., 2023). The ocean stores a vast amount of CO<sub>2</sub>, with inorganic carbon reservoir approximately 50 times greater than those in the atmosphere (Sabine et al., 2004). Therefore, studying oceanic CO<sub>2</sub> sources and sinks is crucial for  
25 developing mitigation strategies and mitigate climate change. The most widely and extensively applied, and long-established method for ocean carbon sink investigation involves measuring the partial pressure difference of CO<sub>2</sub> ( $\Delta p\text{CO}_2$ ) across the air-sea interface (Song et al., 2023). Continuous observations used to calculate the air-sea CO<sub>2</sub> flux, providing the most direct characterization of the ocean carbon cycle system (Wanninkhof et al., 2019; Song et al., 2023).

Owing to the past several decades of continuous observations, a large amount of sea surface  $p\text{CO}_2$  data has been accumulated,  
30 yet it remains insufficient relative to the vast ocean area. The Surface Ocean CO<sub>2</sub> Atlas (SOCAT 2023) revealed that the ocean area covered by monthly CO<sub>2</sub> measurements has decreased by nearly half since 2017, reflecting the decline in global open-ocean CO<sub>2</sub> observation capacity (Bakker et al., 2024). Although recently studies have increasingly employed artificial intelligence and big data technologies to investigate the dynamics of ocean carbon sinks (Landschützer et al., 2013; Xu et al., 2019; Yu et al., 2023), the fundamental limitation of in-situ field observation remains unresolved. Due to the limited spatial and temporal coverage of  $\Delta p\text{CO}_2$   
35 measurements, as well as uncertainties in wind forcing and transport velocity parameterization, the uncertainty in global and regional fluxes estimated from  $\Delta p\text{CO}_2$  measurements can reach up to  $\pm 50\%$  (Wanninkhof et al., 2013; Rhein et al., 2013). In addition to  $\Delta p\text{CO}_2$  data, air-sea CO<sub>2</sub> fluxes can also be estimated using a top-down inversion method that integrates atmospheric CO<sub>2</sub> concentrations with atmospheric transport models (Jacobson et al., 2007; Wanninkhof et al., 2019). Spatial and temporal variations in atmospheric CO<sub>2</sub> concentrations reflect the pattern of sources and sinks across large spatial scales. Consequently, top-down  
40 atmospheric inversion methods are suitable for assessing global and regional CO<sub>2</sub> fluxes and are currently widely adopted to estimate CO<sub>2</sub> emissions from fossil fuels and carbon sinks in terrestrial ecosystems (Piao et al., 2022; Han et al., 2024). However, due to the sparse sampling of CO<sub>2</sub> concentration over the open ocean, significant uncertainties persist in those flux estimations, limiting its applicability (Rödenbeck et al., 2006; Wanninkhof et al., 2019). In summary, the scarcity of marine atmospheric CO<sub>2</sub> concentration observations is the primary obstacle to accurately quantify the oceanic carbon sink.

45 For marine atmosphere, buoy observations excel at meeting the requirements of expanding field observation coverage and significantly increasing data volume compared to research vessel observations constrained by voyage frequency, range, and cost, or satellite observations limited by operational cycles and atmospheric conditions (e.g., clouds, aerosols). Currently, both the ARGO

Global Ocean Observing System and the Southern Ocean Carbon and Climate Observation and Modeling (SOCCOM) project have proposed utilizing buoys for seawater CO<sub>2</sub> observations (Sarmiento et al., 2023), but none of them have conducted atmospheric CO<sub>2</sub> concentrations over the ocean. Urban CO<sub>2</sub> monitoring efforts provide valuable experience for selecting CO<sub>2</sub> observation instruments suitable for deployment on buoys. In recently years, high-density monitoring networks based on low-cost CO<sub>2</sub> sensors have been established in numerous cities worldwide (Karion et al., 2020), as supplements to the land-based observations from the sites within the World Meteorological Organization's (WMO) Global Atmosphere Watch Programme (GAW). For instance, Shusterman et al. (2016) established a CO<sub>2</sub> observation network (BEACO<sub>2</sub>N) which consists of 34 sensors in and around Oakland, California. After applying environmental parameter and drift corrections, the network achieved an accuracy of  $\pm 1.2$ – $2.0$  ppm for 1-minute average dry-air concentrations between, effectively capturing CO<sub>2</sub> variations across multiple temporal scales in urban areas and abnormal short-term CO<sub>2</sub> emission events. Delaria et al. (2021) further corrected the temperature-dependent zero bias of the BEACO<sub>2</sub>N sensor, reducing the error to 1.6 ppm or less. Han et al. (2024) established a 134-station SenseAir K30 sensor observation network and developed a CO<sub>2</sub> calibration system. Data accuracy was enhanced through averaging raw observations, environmental corrections, and calibration with standard gases. After applying long-term drift correction, the sensors (SENSE - IAP) maintained a root mean square error (RMSE) of  $2.4 \pm 0.2$  ppm after 30 months of operation (Cai et al., 2025). Compared to high-precision instruments, low-cost sensors exhibited relatively lower accuracy (several ppm versus  $\sim 0.1$  ppm) but at drastically reduced cost (under 15,000 dollars versus over 150,000 dollars). This cost-performance balance enables the construction of dense observation networks to reveal significant spatial variations of CO<sub>2</sub> induced by emission sources, vegetation carbon sinks, and meteorological conditions (Shusterman et al., 2016; Bakker et al., 2024; Han et al., 2024; Cai et al., 2025).

Considering the advantages of low-cost sensors in land-based CO<sub>2</sub> monitoring and inversion, and the relative maturity of buoy-based observation technologies, we designed a sea-air interface buoy platform equipped with SenseAir K30 sensors (SenseAir AB, Delsbo, Sweden) in the coastal waters off Maoming, Guangdong, China, and evaluated its performance in monitoring marine atmospheric CO<sub>2</sub> concentration. Satisfying measurement accuracy was obtained after instrumental calibration, data processing, and correction strategies, validating the feasibility of marine CO<sub>2</sub> monitoring with low-cost sensors. Here we presented the preliminary results of the buoy-based low-cost sensor system for atmospheric CO<sub>2</sub> monitoring at the Sea-Air interface. Section 2 introduces the instruments and data correction methods; Section 3 presents the results of the land-based experiments for calibration and correction; ~~and~~ Section 4 demonstrates a short-term marine observation case study; and Section 5 presents and discusses the results of post-deployment laboratory validation of the sensors.

## 2 Data and Methods

### 2.1 Instruments and observation site

This study employed a low-cost sensor system integrating three CO<sub>2</sub> Sensor Modules (referred to as CM1, CM2, and CM3) (Figure 1c), capable of simultaneously measuring atmospheric CO<sub>2</sub> concentration and meteorological parameters including temperature, pressure, and humidity. Each CM integrates a CO<sub>2</sub> sensor, environmental parameter sensors, and a Micro-controller Unit (MCU) processor onto a single circuit board housed within a waterproof cube enclosure. Three CM modules are individually mounted in cylindrical housings bolted to a cube enclosure, with silicone seals at the connection points. Both sides and the bottom of the individual housings are wrapped with a membrane that is both breathable and waterproof, ensuring the CMs can operate normally in a marine environment. The CM feature an open-type design that allows ambient air to flow directly through the sensing chamber, without a sealed sampling volume typical of high-precision analysers. The sensors are paired with a data acquisition instrument, and data is collected by a micro-processor named BeagleBone Green Wireless (BBGW) and transmitted back to the server via 4G communication from the base station.

The CO<sub>2</sub> sensor is the K30 sensor module from SenseAir of Sweden, operating on a non-dispersive infrared principle (NDIR). Compared to other low-cost sensors (such as the COZIR Environmental Sensor (CO<sub>2</sub>Meter, Inc., Orlando, FL, USA) and TelaireT6615 (Amphenol Advanced Sensors, USA), etc.), it demonstrates higher raw accuracy, with  $\pm 30 \text{ ppm} \pm 3\%$  (Martin et al., 2017), with a measurement range of 1-10,000 ppm and a resolution of 0.01 ppm, featuring easy configuration and maintenance-free operation. To account for the influence of external environmental variations on the K30 sensor's response, the CM is equipped with a BME680 sensor (Bosch Sensortec GmbH, Reutlingen, Germany) that simultaneously monitors temperature ( $T$ , °C), relative humidity ( $RH$ , %), and atmospheric pressure ( $P$ , hPa) of the internal air mass. The measurement accuracies are  $\pm 0.1 \text{ }^\circ\text{C}$ ,  $\pm 3\%$ , and  $\pm 0.6 \text{ hPa}$ , with corresponding resolutions of 0.01 °C, 0.01%, and 0.01 hPa, respectively. These measurements enable real-time correction of the CO<sub>2</sub> response values for environmental parameters, thereby enhancing the overall accuracy of the observations. The CMs adopt a standard RS485 output mode and are powered by the buoy's 12 V DC battery, operating continuously with a 2s sampling interval. In the marine environment, pitching, strong winds, wave impacts, and rainy conditions are common. Combined with the high humidity and salinity of surface air, these factors often cause condensation and salt deposition on instrument surfaces. To mitigate these effects, the sensors and data logger were connected and securely mounted inside a Stevenson screen (Figure 1c), which was installed near the buoy's center of gravity within the supporting frame (Figure 1b), at an approximate height of 3 m above the sea surface, to provide a relatively stable observation environment.





Figure 1: Deployment and performance evaluation of buoy-mounted sensors for marine atmospheric CO<sub>2</sub> observation. (a) Location of the offshore observation site. Basemap source: Esri World Topographic Map (Esri, <https://server.arcgisonline.com>). (b) On-site scene of the offshore buoy during observation. (c) Schematic of the CO<sub>2</sub> sensor modules (CMs), data logger, and Stevenson screen.

The sea-air coupled monitoring buoy system is composed of a buoy platform and a land-based station (Figure 2). The buoy consists of a buoy body, mooring system, sensors, data acquisition system, power system, safety system, and communication system. Real-time data transmission between the buoy and shore station is achieved via BeiDou (Figure 1b). The buoy body has a diameter of 3 meters, a depth of 0.9 meters, and a total height exceeding 5 meters. The power system comprises high-capacity, compact, rechargeable batteries, and solar panels. The batteries are housed within the instrument compartment, while solar panels are mounted around the buoy tower. These panels charge the batteries, supplying a single operating voltage to the buoy system. The system can sustain normal power supply to the buoy observation system for 15 consecutive days of overcast or rainy weather, ensuring continuous and reliable operation even under severe sea conditions. The buoy data communication system employs dual-mode BeiDou and Iridium satellite communication, with redundant data transmission to ensure an effective data reception rate of better than 95%. The shore station reception and processing system features reception, post-processing, and report generation capabilities, enabling modification and configuration of parameters such as buoy sampling frequency and transmission cycle. It also provides low-voltage, water ingress, and displacement alarm functions for buoys.

设置了格式: 默认段落字体



120 **Figure 2: Schematic diagram of a deep-sea air-sea coupled monitoring buoy system**

To calibrate the CMs, it was first placed in the laboratory and a meteorological observation field within Jiangwan Campus of Fudan University, for side-by-side observations with a cavity ring-down spectrometer (Picarro G2301, Picarro Inc., Santa Clara, USA). (Figure 3). The CMs equipped with Stevenson screen was placed outside the station building, while the sampling gas tube of Picarro was extended into the Stevenson screen through a duct connecting the station building to the outside environment, ensuring simultaneous observations with CMs. Following land-based field observations, the instrument was deployed for field observations at sea. The offshore field observation point is located in the coastal area of Dianbai District, Maoming City, Guangdong Province (21.44 °N, 111.39 °E) (Figure 1a), belonging to northern shelf coastal section of the South China Sea nearshore areas feature port zones, shallow bays, and small islands (such as Dazhuzhou), while offshore lies the broad continental shelf and slope transition zone of the northern South China Sea. Integrated multi-source observations indicate that most of the South China Sea is a weak to moderate CO<sub>2</sub> source with seasonal variations (Zhai et al., 2013; Li et al., 2020; Chen et al., 2024; Zhang et al., 2024). The annual average flux in the northern continental slope region is approximately 0.46 mol m<sup>-2</sup> yr<sup>-1</sup>, with higher values in the central and southern areas (about 1.37 mol m<sup>-2</sup> yr<sup>-1</sup>) (Zhai et al., 2013). During summer, the coastal upwelling brings up subsurface water rich in dissolved inorganic carbon and low in temperature to the surface layer, typically tending to increase atmospheric CO<sub>2</sub>

125  
130

emissions. However, spatiotemporal variations in wind events, biological consumption, and estuarine runoff can cause significant short-term or inter seasonal reversals (Xu et al., 2013; Li et al., 2021). In summary, the northern South China Sea, where the observation point in this study are located, is generally characterized as a minor CO<sub>2</sub> source but exhibits strong spatiotemporal variability (Zhang et al., 2024).




Figure 3: Land-based field observation experiments. (a) Configuration of CMs, Stevenson screen, and Picarro sampling channel. (b) Meteorological observation station, experimental cabin, and field deployment of CMs. (c) Field setup of the Picarro G2301.

## 2.2 CMs data correction method

The original signals had a sampling interval of 2 s and a background noise level of  $\pm 30$  ppm. The raw data from the CMs were filtered and resampled. After quality control, outliers deviating by more than  $4\sigma$  from the mean were removed — this detection was implemented on the native 2 s resolution data within consecutive 1-minute windows (30 raw data points per window) — and temporal averaging was applied to reduce the noise level. The  $4\sigma$  threshold is applied to achieve a compromise between eliminating extreme outliers and retaining the inherent variability of the dataset (Cai et al., 2025). Allan variance, which quantifies the time-averaged stability of continuous measurements, was used to determine the optimal averaging interval that minimizes noise while preserving the integrity of the data signals (Martin et al., 2017). Langridge (2008) indicated that the optimal averaging time for the Allan variance of the K30 sensor to reach its minimum is approximately 3 min, after which extending the averaging time step no longer significantly reduces the noise. Cai et al. (2025) evaluated the noise characteristics of the SenseAir K30 by continuously introducing standard gases. The results showed that at a measurement interval of 2 s, the noise level was 4 ppm, and 1-2 min showed 0.4-0.7 ppm noise level, and from 2 min to 1 h, the noise level decreased to approximately 0.2 ppm. Although the 3-minute interval yields a marginally lower Allan variance, a 1-minute averaging time was adopted in this study because the Allan variance is only slightly higher than at 3 min while allowing resolution of shorter-timescale atmospheric variability (Martin et al., 2019). This choice ensures sufficiently low noise (0.4-0.7 ppm) to resolve

marine CO<sub>2</sub> dynamics while preserving higher-frequency variability associated with rapid coastal atmospheric and air-sea CO<sub>2</sub> exchanges that would be smoothed over with 3-minute averaging, and supports detailed process analysis with the flexibility to aggregate to coarser scales as needed. Therefore, in this study, we selected a 1-minute time step to average the raw data before proceeding with subsequent corrections. This approach significantly reduces the noise level in the data while enabling the resolution of atmospheric changes on a relatively short timescale.

Using CMI as an example, results from observations conducted by CMs and Picarro in laboratory and terrestrial field are presented in Figure 4. After quality control and resampling of raw data, the standard deviation (SD) for the three CMs in laboratory tests improved from 13.05, 17.32, 18.28 ppm to 4.68, 5.26, and 5.48 ppm, respectively. For field tests, the values improved from 15.14, 21.93, 17.06 ppm to 9.33, 14.83, and 8.83 ppm, respectively (Figure 4; Figures S1-S2). Data accuracy improved after minute averaging (Figure 4 a-d), but differences still exist compared to the reference instrument Picarro ( $\Delta\text{CO}_2$ ). A comparison between the laboratory and field results (Figure 4 ae, d) shows that the CMs performed better under laboratory conditions. In the stable laboratory environment,  $\Delta\text{CO}_2$  exhibited no pronounced diurnal variation (Figure 4 be, e), fluctuating steadily around a constant value (system bias). By contrast, in the field, where atmospheric conditions naturally vary (Figure 4f), the environmental parameters showed clear diurnal cycles, and  $\Delta\text{CO}_2$  also displayed diurnal oscillations. In addition, during the field deployment,  $\Delta\text{CO}_2$  exhibited a gradual downward “temporal trend” over the observation period (Figure 4e), which is closely related to changing environmental conditions. Apart from the diurnal cycle, ambient temperature exhibited a marked upward trend over several consecutive days (Figure 4f). As will be discussed in Section 3, the  $\Delta\text{CO}_2$  of CMI is negatively correlated with temperature, indicating that rising temperatures lead to a decrease in  $\Delta\text{CO}_2$ . Therefore, the observed downward trend in  $\Delta\text{CO}_2$  is physically consistent with the gradual rise in temperature, reflecting a temperature-driven response rather than inherent sensor instability. This indicates that the CMs are highly sensitive to environmental conditions, and their measurement accuracy is affected by both atmospheric variability and the CO<sub>2</sub> concentration baseline. Therefore, further calibration with respect to environmental parameters is required. The most used environmental correction method for NDIR sensors is multiple linear regression. This involves establishing empirical regression models using environmental parameters such as temperature, air pressure, and water vapor under laboratory or side-by-side observations with standard instrument (Martin et al., 2017; Han et al., 2024; Cai et al., 2025). In recent years, numerous studies have attempted to model environmental nonlinear effects using machine learning methods such as random forests, gradient boosting, or neural networks, achieving lower root mean square error (RMSE) than linear regression (Biagi et al., 2024; Dubey et al., 2024). However, these approaches still exhibit limitations in interpretability and transferability.

设置了格式: 非突出显示

设置了格式: 非突出显示

设置了格式: 非突出显示

设置了格式: 非突出显示

设置了格式: 非突出显示




Figure 4: Time series of CM1 data during laboratory (a-c) and land-based field (d-f) observations. (a, d) CM1-measured CO<sub>2</sub> concentration at second-level resolution (grey dots) and minute-level resolution (blue dots), alongside Picarro-measured CO<sub>2</sub> concentration (black line). (b, e) Time series of CO<sub>2</sub> concentration difference ( $\Delta\text{CO}_2\text{-CM1-Picarro}$ ) at second-level (grey dots) and minute-level (blue dots) resolution. (c, f) Time series of ambient temperature (T, red line) and relative humidity (RH, blue line).

The 1-min  $\Delta\text{CO}_2$  exhibits linear relationships with  $T$ ,  $P$ , and  $RH$ , and this environment-related dependence differs among individual CMs (Figure 5). Based on these characteristics, this study adopts a multiple linear regression approach for environmental calibration as Cai et al., (2025), where  $\Delta\text{CO}_2$  and the environmental parameters satisfy the following relationship:

$$\Delta C = Y_{CM} - y_0 \quad (1)$$

$$\Delta C = a_0 + f(a_t, T) + f(a_p, P) + f(a_{RH}, RH) + f(a_c, CO_2) \quad (2)$$

Here,  $Y_{CM}$  represents the CMs measurements, and  $y_0$  is the true atmospheric CO<sub>2</sub> concentration. The coefficients  $a_t$ ,  $a_p$ ,  $a_{RH}$ , and  $a_c$  are the correction parameters associated with  $T$ ,  $P$ ,  $RH$ , and the CO<sub>2</sub> concentration, respectively; including the CO<sub>2</sub> term in the calibration equation serves to eliminate the zero-point bias of the CMs. For each CM, the corresponding values of  $a_t$ ,  $a_p$ ,  $a_{RH}$ , and  $a_c$  are unique.  $a_0$  is the baseline concentration correction parameter, calibrated using measurements from reference instruments at the observation site or from nearby atmospheric background stations. All these parameters can be determined through multiple linear regression using the `Linear Regression` function in Python. It should be specifically noted that temporal drift of the sensor was evaluated (Figure S3). Previous studies have incorporated linear time terms into calibration models to account for sensor

格式化表格

设置了格式: 字体: (默认) Times New Roman

aging and drift (Arzoumanian et al., 2019). In the present study, however, the apparent temporal trend in raw error is associated with the gradual increase in temperature during the measurement period. After correction, no consistent linear relationship was observed between sensor error and time. Therefore, a linear time term ( $t$ ) was not included in the regression model. For long-term monitoring, periodic re-calibration every 3-6 months using stable atmospheric CO<sub>2</sub> observations from background reference sites (e.g., MLO) during quiescent atmospheric conditions will be required to address non-linear temporal drift.

The corrected data of CO<sub>2</sub> after environmental correction is:

$$y_{corrected} = Y_{cm} - a_0 - f(a_t, T) - f(a_p, P) - f(a_{RH}, RH) - f(a_c, CO_2) \quad (3)$$

When standard instrument (e.g., Picarro) co-located observations are available, these measurements shall be considered the true values for atmospheric CO<sub>2</sub> concentrations. The specific correction results will be described in Section 3. For marine observations, the baseline correction of CMs is performed using data from the Mauna Loa atmospheric background station in Hawaii, USA (Thoning et al., 2025), corresponding to periods where CMs observations are stable and close to background values. The specific method and results will be introduced in Section 4.

### 3 Environmental correction results for land-based field observations

Temperature variations affect the sensor's light source intensity, detector response, and absorption cross-section, leading to systematic drifts in output (Yasuda et al., 2012). Pressure influences gas density and infrared absorption line broadening, making corrections based on the equation of state or sensor sensitivity particularly important in regions with strong pressure fluctuations (Chen et al., 2010; Curcoll et al., 2022). Water vapor exerts the most complex effects on NDIR sensors: it not only dilutes CO<sub>2</sub> mole fractions in moist air relative to dry air but also causes spectral line broadening within the CO<sub>2</sub> absorption band, introducing biases (Chen et al., 2010; Dubey et al., 2024). Accordingly, the multivariate linear calibration of CMs focuses on three key environmental factors -  $T$ ,  $P$ , and  $RH$ . Figure 5 shows the variations of  $\Delta CO_2$  between the three CMs and the collocated Picarro measurements as a function of environmental parameters ( $T$ ,  $P$ , and  $RH$ ) during the land-based field observations, where the orange and green colors correspond to the data before and after environmental correction, respectively. Figure 6 presents the linear relationships between the CMs and the Picarro, along with the histograms of  $\Delta CO_2$ .

Prior to correction, the  $\Delta CO_2$  of all three CMs exhibited linear relationships with environmental parameters (Figure 5): negative correlation with  $T$ , and positive correlation with  $P$  and  $RH$ . Among these, CM2 demonstrated the most pronounced correlations. The correlation coefficients ( $r$ ) between  $\Delta CO_2$  and  $T$ ,  $P$ , and  $RH$  were -0.79, 0.71, and 0.43, respectively, while the  $r$  values for CM1 and CM3 were -0.49, 0.52, 0.22, and -0.39, 0.44, 0.16, respectively, all significant at the 0.01 level ( $p < 0.01$ ). After correction, the systematic drift of  $\Delta CO_2$  across the three CMs due to environmental parameters was successfully eliminated.

The corresponding  $r$  values were all 0, indicating no significant correlations, with  $\Delta CO_2$  fluctuating around the zero line. Beyond

格式化表格

environmental factors, the multiple linear regression also incorporated the true CO<sub>2</sub> concentration in the atmosphere, represented by Picarro co-located measurements in field observations. The values of the first three CMs before correction all exhibit zero bias relative to the true values (the fit between CO<sub>2</sub> observations and true values includes intercepts) (Figure 6 a-c), with respective *r* values of 0.98, 0.93, and 0.98, all significant at the 0.01 level ( $p < 0.01$ ). After correction, the correlations all improved to 0.99 ( $p < 0.01$ ), and data points largely converged on either side of the 1:1 line. The bias between CMs and Picarro shifted from -3.7 ppm, +43.6 ppm, and -16 ppm to +0.1 ppm, +1.6 ppm, and +0.6 ppm, respectively, shifting from significantly biased to essentially unbiased (Figure 6 d-f). The results above suggest that all three CMs are influenced by environmental variables, but to markedly different extents. Whether for CM2, which inherently exhibits substantial systematic errors, or CM1, which shows minimal data offset prior to correction, our environmental correction method significantly enhances observational accuracy, improves data quality, and demonstrates good universality.




Figure 5: Variations of  $\Delta\text{CO}_2$  ( $\Delta\text{CO}_2=\text{CM-Picarro}$ ) for CM1 (a-c), CM2 (d-f), and CM3 (g-i) during land-based field observations, as functions of temperature (a, d, g), atmospheric pressure (b, e, h), and relative humidity (c, f, i). Orange dots represent data before environmental correction, while green dots represent data after environmental correction.




Figure 6: Direct comparison of hourly moving averages of CO<sub>2</sub> concentrations among CM1, CM2, CM3 and Picarro during land-based field observations (a-c), and histograms of ΔCO<sub>2</sub> distributions (d-f) (ΔCO<sub>2</sub>=CM-Picarro, where CM corresponds to CM1, CM2, and CM3, respectively). Orange dots and bars represent data before environmental correction, while green dots and bars represent data after environmental correction.

Environmental correction effectively reduced the offset values between CMs and Picarro (Figure 7). Before correction, CMs captured CO<sub>2</sub> concentration trends like Picarro but exhibited significant deviations. For the best performing CM1, this deviation was particularly noticeable at low CO<sub>2</sub> concentrations, while CM2 and CM3 showed overall high and low biases, respectively. The ΔCO<sub>2</sub> of CMs exhibited a certain "downward" drift trend over the one-week observation period, CM1 and CM3 moved from zero toward negative values, while CM2 shifted from a relatively high positive value around 50 ppm toward zero. The calibrated results showed high consistency with Picarro, with the RMSE decreasing from 11.04 ppm, 30.51 ppm, and 18.70 ppm to 4.03 ppm, 3.96 ppm, and 3.88 ppm, respectively. The correction effectively eliminated the linear drift trend of ΔCO<sub>2</sub> over time, stabilizing it to fluctuate around the zero line. During the land-based field observations in Shanghai in early spring, the atmospheric temperature and humidity exhibited pronounced diurnal variations, fluctuating between 5-30 °C and 10-60%, respectively. The average RMSE of corrected CMs was 3.64 ppm, which is sufficient to capture terrestrial CO<sub>2</sub> variations (400-600 ppm), even during periods of significant CO<sub>2</sub> fluctuations with pronounced peaks and troughs, such as cases on March 22nd-23rd and March 25th-26th, where the environmental correction method performed well.




Figure 7: Hourly moving average time series of CM1 (a-c), CM2 (d-f), and CM3 (g-i) during land-based field observations: CO<sub>2</sub> concentration of CMs before environmental correction (orange line), after environmental correction (green line), and CO<sub>2</sub> concentration from Picarro (black dashed line) (a, d, g);  $\Delta\text{CO}_2$  ( $\Delta\text{CO}_2 = \text{CM} - \text{Picarro}$ ) of CMs before and after calibration (b, e, h); Ambient temperature (T, red line) and relative humidity (RH, blue line) (c, f, i).

#### 4 Marine observation results

Based on the instrument deployment and marine environment described in Section 2, and the environmental correction method validated through land-based field observations in Section 3, the CMs-equipped air-sea interface buoy observation platform commenced operation in May 2025 in the northern South China Sea off the coast of Dianbai District, Guangdong Province.

Following signal debugging and regular equipment maintenance, observational data were obtained over three months from May 28th to August 28th. The hourly moving-averaged time series of CO<sub>2</sub> concentration,  $T$ , and  $RH$  from the three CMs are shown in Figure 8. During environmental calibration, measurements are first corrected using the environmental coefficients  $a_t$ ,  $a_p$ ,  $a_{RH}$ , and  $a_c$  obtained from land-based calibration; baseline correction is then applied by updating  $a_0$  with reference CO<sub>2</sub> concentrations from the marine atmospheric boundary layer. Over the open ocean, strong horizontal atmospheric mixing results in small zonal variations in marine boundary layer CO<sub>2</sub> concentrations, indicating high zonal uniformity at similar latitudes (Bakwin et al., 2004; Palter et al., 2023). Given the strong real-time nature of this study's observations and the limited availability of co-located and near-surface observation resources, CO<sub>2</sub> observation data from the Mauna Loa atmospheric background station in Hawaii, USA (MLO, 19.54°N, 155.58°W) - located at a latitude similar to the observation site - served as the reference value. The CO<sub>2</sub> datasets were obtained from

the NOAA Global Monitoring Laboratory (GML) (<https://gml.noaa.gov/ccgg/trends/data.html>); Thoning et al., 2025). The relatively stable period (June 17th-24th) of CMs concentrations during the observation was regarded as the atmospheric background state. Both values were substituted into a multiple linear regression calculation to obtain  $a_0$ . After environmental correction, the RMSE of the CMs significantly decreased from 9.27 ppm, 52.39 ppm, and 11.24 ppm to 1.57 ppm, 1.86 ppm, and 1.52 ppm, respectively (Figure 8a; Figures S43-S65). The lower RMSE during the marine test can be partly explained by the substantially lower ambient CO<sub>2</sub> variability over the ocean, as reflected by the smaller standard deviation of the MLO reference data (1.81 ppm) compared to that of the Picarro in-situ measurements at the land site (29.29 ppm).

During the three-month marine observations, the atmosphere at the observation site exhibited high temperatures and humidity, with temperatures ranging from 25 to 37.5 °C and humidity levels between 50% and 100% (Figure 8b). Both showed pronounced diurnal and weekly variations. The mean CMs values before and after correction were 459.48 ppm and 436.94 ppm, with medians of 456.53 ppm and 433.75 ppm, respectively (Figure 8c). The means consistently exceeded the medians, and the ranges surpassed 100 ppm in both cases, which indicates many signal peaks in CO<sub>2</sub> during the observation, and the observation site is susceptible to terrestrial anthropogenic CO<sub>2</sub> emissions. After correction, the overall concentration was approximately 22 ppm lower than the original values, consistent with the CO<sub>2</sub> range at the atmospheric background station MLO (mean 429.32 ppm). The correction eliminated systematic overestimation, bringing the results closer to background concentrations.

The SD of the raw data is 9.79 ppm, showing a slight difference from the corrected value of 9.62 ppm. The first quartile (Q1) changed from 453.08 ppm to 430.10 ppm, and the third quartile (Q3) changed from 463.29 ppm to 441.32 ppm, with an interquartile range (IQR) of 10.21 ppm, slightly below the corrected value of 11.22 ppm. These indicate that the correction process reduced the overall CO<sub>2</sub> concentrations but did not significantly decrease data variability. In fact, the distribution of the middle 50% of data points became wider. Whether considering the extreme fluctuation range (extreme value difference) influenced by

设置了格式: 默认段落字体

anthropogenic land effects, the typical fluctuation range IQR after removing most extreme signals, or the average fluctuation amplitude SD of overall concentrations, the accuracy of CMs corrected data is sufficiently to capture the corresponding signals.




Figure 8: Offshore buoy observation results of CMs. (a) Hourly moving average time series of CO<sub>2</sub> concentrations from CMs before correction (orange line) and after correction (green line), together with daily mean CO<sub>2</sub> series from Mauna Loa Observatory (MLO, red line). The light red and light blue shaded backgrounds correspond to CO<sub>2</sub> fluctuation periods and stable periods, respectively. (b) Time series of ambient temperature (T, red line) and relative humidity (RH, blue line). (c) Histograms and boxplots showing the distributions of CO<sub>2</sub> concentrations before (orange bars) and after correction (green bars).

During the three-month observation period, CO<sub>2</sub> at the monitoring site exhibited short-term fluctuation peaks as well as periods of stable concentrations, both of which can be analyzed and interpreted from the perspective of atmospheric transport. Using the NOAA Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPPLIT) model, 36-hour backward trajectories were calculated for the observation point to identify the primary transport pathways influencing the air mass sources at the location (Cohen et al., 2015). The trajectory origin is set at the observation point, with a time resolution of 6 hours. Meteorological driving data were from the Global Data Assimilation System (GDAS1, 1° × 1°) reanalysis data provided by NOAA (Rolph et al., 2017). After obtaining a series of backward trajectories, the HYSPPLIT clustering module was employed to classify the trajectories. This approach mitigates the impact of uncertainty inherent in individual trajectories and extracts key transport pathway characteristics, providing a basis for subsequent analysis of the relationship between air mass transport and observational results (Cohen et al., 2015). The entire period of marine observations was divided into four segments for clustering. The periods from May 28th to June 16th (Figure 9a) and June 25th to August 1st (Figure 9c) constituted CO<sub>2</sub> fluctuation phases (the background is light red in Figure 8a), with many concentration peaks occurring during these periods. The periods from June 17th to 24th (Figure 9b) and August 2nd to 13rd (Figure 9d) constituted

CO<sub>2</sub> stable phases (the background is light blue in Figure 8a). During these phases, CO<sub>2</sub> concentrations fluctuated minimally and approached the background levels of the marine boundary layer. It is particularly noteworthy that during the two-week period from August 14th to August 28th (the background is white in Figure 8a), CO<sub>2</sub> exhibited alternating patterns of relatively dense peaks and sustained background concentration levels, with each state lasting no more than five days. The limited number of trajectories obtained from segmented analysis makes it inconvenient for cluster analysis.

The trajectory clustering results (Figure 9) indicate that the atmospheric transport pathways corresponding to concentration fluctuation periods are relatively complex, significantly influenced by air masses transported from land. Trajectories from May 28th to June 16th were classified into 17 categories, with 40% originating from land and 60% from the ocean. The most typical inland air mass, represented by Trajectory 4, was transported from northern Guangdong all the way to western Guangdong. Trajectory 12, although originating from the sea, reached the observation point via the western coast of Guangdong within the first 6 hours of the observation period. From June 25th to August 1st, trajectories ending over land accounted for 54%, while those ending over the ocean accounted for 46%. Trajectory 2, corresponding to short-range inland transport, accounted for 13%. The clustering results indicate that local urban emissions from land areas significantly contributed to CO<sub>2</sub> concentrations at the observation point, effectively explaining the observed large fluctuations and multiple peaks during the corresponding period. Correspondingly, air masses at observation point during concentration stable periods were predominantly transported from clean marine atmospheres. The trajectories from June 17th to 24th were clustered into 11 groups, all originating from the South China Sea. Consequently, the observed CO<sub>2</sub> concentrations during this period remained near background levels, which can be considered the CO<sub>2</sub> concentration level in clean air without anthropogenic pollution. This also demonstrates that using CO<sub>2</sub> observations from this period combined with MLO atmospheric background concentrations for baseline correction of CMs is reasonable. Trajectories from August 2nd to 13rd were categorized into 4 types, with 92% originating from marine sources and 8% from land sources. Consequently, although CO<sub>2</sub> concentrations at observation point during this period were slightly elevated above background values, they remained generally stable.




Figure 9: Clustered results of 36-hour backward trajectories from Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPPLIT) model analyses at the offshore observation site during four time periods of buoy observations. Each trajectory is labeled with its cluster number and proportion, and markers on the trajectories indicate 12-hour intervals. (a) 28th May-16th June, (b) 17th-24th June, (c) 25th June-1st August, (d) 2nd-13rd August. (a) and (c) correspond to CO<sub>2</sub> concentration fluctuation periods, while (b) and (d) correspond to CO<sub>2</sub> concentration stable periods.

By comparing the corrected CMs data during the concentration stable periods (including 19 complete days) with the diurnal variations in CO<sub>2</sub> concentration during the summer of 2024 (June to August) at the MLO station (Figure 10), we can further understand how CMs captures the diurnal variation of oceanic boundary layer CO<sub>2</sub>. Hourly CO<sub>2</sub> mole fractions at MLO were obtained from NOAA GML (<https://gml.noaa.gov/data/dataset.php?item=mlo-co2-observatory-hourly>; Thoning et al., 2025). Both CMs and MLO exhibit daily variations in background CO<sub>2</sub> concentrations characterized by lower daytime values and higher nighttime values. This pattern likely stems from the primary influence of air-sea CO<sub>2</sub> fluxes and atmospheric convective transport on oceanic CO<sub>2</sub> concentrations (Lv et al., 2015). During the day, solar radiation heats the Earth's surface, which enhances photosynthesis in marine ecosystems and promotes the uptake of atmospheric CO<sub>2</sub>. As atmospheric temperatures rise, enhanced turbulent activity thickens the atmospheric boundary mixed layer, diluting CO<sub>2</sub> through vertical atmospheric mixing and reducing its concentration. At night, photosynthesis ceases in regional marine ecosystems, leaving only respiration. The sea surface cools, and weakened turbulent mixing restricts vertical air exchange. CO<sub>2</sub> struggles to diffuse into the upper layers, accumulating in the near-surface layer and increasing in

设置了格式: 默认段落字体  
设置了格式: 默认段落字体

concentration. The daily amplitude of CO<sub>2</sub> at MLO station during summer is approximately 2 ppm, while CMS is 3 ppm. This indicates that the environmentally calibrated CMS is sufficiently sensitive to capture the daily variation signal of CO<sub>2</sub> in clean background air unaffected by terrestrial anthropogenic emissions, demonstrating its potential for observing atmospheric CO<sub>2</sub> concentrations in open ocean environments. The daily amplitude of CO<sub>2</sub> at MLO Station during summer is approximately 2 ppm, while CMS is 3 ppm. This indicates that the environmentally calibrated CMS is sufficiently sensitive to capture the daily variation signal of CO<sub>2</sub> in clean background air unaffected by terrestrial anthropogenic emissions, demonstrating its potential for observing atmospheric CO<sub>2</sub> concentrations in open ocean environments.

Observations of the CMS show that the daily maximum and minimum values of CO<sub>2</sub> exhibit a 3-hour lag compared to the MLO station, which may be attributed to differences in the surrounding environments of the two sites. The MLO station is situated on high altitude land, exhibiting a typical mountainous diurnal variation in CO<sub>2</sub> concentration. Mountainous terrain induces strong upslope and downslope airflows, leading to earlier atmospheric mixing. During the day, upslope flow and mixing intensify (NOAA GML, 2024a), resulting in the lowest concentrations around 4 p.m. when vertical mixing is strongest. At night, the mountain air becomes isolated from the free atmosphere, with downslope flows carrying high CO<sub>2</sub> air (NOAA GML, 2024a), reaching peak concentrations around 6 a.m. The observation site in this study is over an ocean surface, which possesses high thermal capacity, exhibits small diurnal temperature variations, and experiences delayed turbulence enhancement and boundary layer development (Nemoto et al., 2009). Consequently, the nocturnal accumulation of CO<sub>2</sub> persists until 8 a.m., several hours after sunrise. Daytime sea breezes and mixing intensify later, with the lowest CO<sub>2</sub> values occurring around 7 p.m. The average CO<sub>2</sub> concentrations at MLO and CMS were 424.85 ppm and 431.38 ppm, respectively, differing by approximately 6.5 ppm. This discrepancy may be due to the fact that the comparison uses 2024 data. According to the NOAA, the annual growth rates of CO<sub>2</sub> concentrations for MLO 2023 and 2024 were 3.36 ppm/yr and 3.33 ppm/yr (NOAA GML, 2024b). Furthermore, even after filtering out short term terrestrial sources, the East Asian monsoon transport can still cause regional increases in atmospheric CO<sub>2</sub>. This monsoon-driven transport represents mesoscale or large-scale processes, not local pollution peaks, and thus systematic differences between CO<sub>2</sub> concentrations and MLO can still be observed during stable periods (Fang et al., 2014; Lin et al., 2018).




Figure 10: Diurnal variations of CO<sub>2</sub> concentrations from CMs (environmentally corrected, green line) during stable concentration periods and from Mauna Loa Observatory (MLO, Jun-Aug 2024, red line), along with corresponding temperature (T, red dash line) and relative humidity (RH, blue dash line) variations.

## 5 Post-deployment laboratory validation

To evaluate long-term sensor stability, post-deployment laboratory calibrations were performed against the Picarro reference in March 2026, approximately 9 months after the initial June 2025 calibration. The results are summarized in Table 1.

Table 1: Post-deployment laboratory validation results

Data Type	Raw		Corrected		Re-Corrected	
	Bias	RMSE	Bias	RMSE	Bias	RMSE
CM1	5.90	6.54	-3.89	4.81	0.09	2.82
CM2	44.73	45.04	-14.62	15.14	0.25	3.92
CM3	1.90	2.45	-9.33	9.46	0.16	1.58
mean	17.51	17.79	-9.28	9.67	0.17	2.70

After marine deployment, noticeable sensor temporal drift is evident in the Corrected data (mean bias: -9.28 ppm, mean RMSE: 9.67 ppm). Performance varied among sensors; CM3 showed reduced accuracy under the original correction compared to its raw data, likely due to individual sensor variability over time. Crucially, sensor sensitivities to temperature, humidity, and pressure remained consistent throughout the deployment. Re-correction by updating the baseline parameter  $a_0$  effectively compensates for this temporal drift, reducing the mean bias to 0.17 ppm and mean RMSE to 2.70 ppm. These results confirm that the environmental

设置了格式: 字体: 10 磅

带格式的: 标题 1

带格式的: 缩进: 首行缩进: 2 字符

设置了格式: 字体: (中文) Times New Roman, 9 磅, 字体颜色: 自动设置

带格式的: 题注

设置了格式: 字体: 10 磅

格式化表格

设置了格式: 字体: 10 磅

设置了格式: 字体: 10 磅

设置了格式: 字体: 10 磅

设置了格式: 字体: 10 磅

设置了格式: 字体: 10 磅

设置了格式: 字体: 10 磅

带格式的: 正文, 缩进: 首行缩进: 2 字符

correction approach remains reliable for short-term deployments, while periodic re-calibration is essential for maintaining measurement accuracy in long-term marine observations.

设置了格式: 默认段落字体

## 5.6 Conclusions

We successfully established a sea-air interface buoy platform along the coast of Maoming, Guangdong Province, employing low-cost sensors to observe marine CO<sub>2</sub> concentrations in the nearshore region of the northern South China Sea. Environmental correction methods effectively eliminated the impact of environmental factors such as temperature, pressure, and humidity fluctuations on CO<sub>2</sub> measurements while correcting for zero bias. With land-based Picarro G2301 co-located observations for comparison, CO<sub>2</sub> accuracy improved from 8.03 ppm to 3.64 ppm (corresponds to the 1-hour moving average of minute-by-minute data). Over ocean, baseline correction using MLO atmospheric background station data improved accuracy from 24.26 ppm to 1.59 ppm, meeting the precision requirements for capturing marine CO<sub>2</sub> concentration signals (e.g., 420-480 ppm in this research). Systematic errors were eliminated, ensuring observed overall concentration levels align with MLO background stations, thereby meeting CO<sub>2</sub> concentration levels in the marine boundary layer. The temporal variations in CO<sub>2</sub> observed by the CMs, including both fluctuating and steady-state phases, can be explained by the long-range atmospheric transport simulated by HYSPLIT. Moreover, the CMs successfully captured the diurnal variations of background CO<sub>2</sub> in the marine atmospheric boundary layer, with an amplitude of approximately 3 ppm.

In summary, this study proposes an environmental correction method for calibrating low-cost sensors, demonstrating its reliability and scientific validity. It has successfully facilitated the application of low-cost sensors aboard buoys for observing marine atmospheric CO<sub>2</sub> concentrations, providing valuable experience for field deployment of marine atmospheric CO<sub>2</sub> monitoring. To our knowledge, this marks the first application of this method, holding significant importance for acquiring CO<sub>2</sub> concentration data in under-observed marine regions. This approach significantly reduces the cost of observing CO<sub>2</sub> in the ocean, opening new possibilities for achieving the goal of substantially increasing observation data. At the same time, high-precision instruments demand stringent environmental conditions. When deployed on buoy platforms in the harsh field observation environment of the ocean, with its powerful winds and waves, maintenance becomes extremely difficult. Low-cost sensors, however, overcome these technical challenges to a large extent.

This study represents the first trial in deploying low-cost sensors aboard buoys to monitor marine CO<sub>2</sub> concentrations. The buoy platform equipped with CMs has withstood several typhoon events, demonstrating excellent watertightness, mechanical robustness, and stability under wave conditions. During deployment, the sensors maintained nearly 100% operational uptime. Post-recovery inspection revealed only minor corrosion on external metallic components, while internal sensor modules remained intact and free of corrosion, further confirming their robustness and good mechanical strength in the harsh marine environment.

platform equipped with CMs has withstood several typhoon events, demonstrating excellent watertightness, mechanical robustness, and stability under wave conditions, which are essential for reliable long-term marine monitoring. The successful detection of daily variations in the CO<sub>2</sub> stable periods further demonstrates the method's potential for deployment in open ocean observations. To achieve the goal of significantly increasing the number of marine observations, low-cost sensors must be deployed on small drifting buoys in the following studies. The current short-term deployment does not allow for full assessment of long-term sensor drift, which will require correction in extended observations. Long-term operation of NDIR sensors is expected to introduce non-linear temporal drift associated with light source aging and detector degradation. Since such drift exhibits no consistent linear relationship with time, a linear correction coefficient was not adopted in this study. Instead, periodic re-calibration every 3-6 months using stable background CO<sub>2</sub> observations under quiescent atmospheric conditions is recommended for future long-term deployments. While the current short-term deployment confirms satisfactory real-world performance, a comprehensive assessment of long-term operational durability and lifespan requires extended multi-month to multi-year deployments, which will be conducted in future work with periodic recalibration and intercomparison. Future work will also aim at long-term buoy deployments to capture seasonal variability and further validate sensor performance under varying environmental conditions. Additionally, the data in this paper covers a relatively short time scale and has not yet accounted for long-term sensor drift. As the duration of observations continues to increase, it will be necessary to correct for this long-term drift. Therefore, this method requires continued refinement and enhancement. In the future, if large-scale deployment of buoys for observation can be realized to obtain extensive regional oceanic CO<sub>2</sub> observational data, these data could be utilized for "top-down" atmospheric inversions. This would provide new perspectives and methodologies for estimating air-sea CO<sub>2</sub> fluxes, representing a groundbreaking endeavor. It holds significant importance for accurately estimating oceanic carbon sinks and quantifying the dynamics of the carbon cycle.

设置了格式: 非突出显示

设置了格式: 非突出显示

### Competing interests

The authors declare that they have no conflicts of interest.

### Funding

This research was supported by the National Key R&D Program of China (No. 2023YFC3705500).

### Author contributions

BY and PFH designed the study. JLL collected and analyzed the datasets. PFH, BY and JLL discussed the sensor results. JLL, BY and PFH led the writing of the paper with contributions from all the coauthors. All authors contributed to the descriptions and discussions of the manuscript.

### Acknowledgments

450 We thank Mr. Zhimin Zhang and Ms. Qixiang Cai, for their help in the instrument development, calibration and deployments. We gratefully acknowledge NOAA for the Mauna Loa Observatory CO<sub>2</sub> data.

## References

- 455 [Arzoumanian, E., Vogel, F. R., Bastos, A., Gaynullin, B., Laurent, O., Ramonet, M., and Ciais, P.: Characterization of a commercial lower-cost medium-precision non-dispersive infrared sensor for atmospheric CO<sub>2</sub> monitoring in urban areas, \*Atmos. Meas. Tech.\*, \*\*12\*\*, 2665–2677, <https://doi.org/10.5194/amt-12-2665-2019>, 2019.](#)
- Bakker, D. C. E., Alin, S. R., Bates, N. R., Becker, M., Gkritzalis, T., Jones, S. D., Kozyr, A., Lauvset, S. K., Metzl, N., Nakaoka, S.-i., O'Brien, K., Olsen, A., Pierrot, D., Steinhoff, T., Sutton, A., Takao, S., Tilbrook, B., Wada, C., and Wanninkhof, R.: SOCAT version 2024: Ocean CO<sub>2</sub> observing effort down to levels of a decade ago, 2024.
- 460 Bakwin, P. S., Davis, K. J., Yi, C., Wofsy, S. C., Munger, J. W., Haszpra, L., and Barcza, Z.: Regional carbon dioxide fluxes from mixing ratio data, *56*, 301–311, 10.3402/tellusb.v56i4.16446, 2004.
- Biagi, R., Ferrari, M., Venturi, S., Sacco, M., Montegrossi, G., and Tassi, F.: Development and machine learning-based calibration of low-cost multiparametric stations for the measurement of CO(2) and CH(4) in air, *Heliyon*, **10**, e29772, 10.1016/j.heliyon.2024.e29772, 2024.
- 465 Cai, Q., Zeng, N., Yang, X., Xu, C., Wang, Z., and Han, P.: A 30-month field evaluation of low-cost CO<sub>2</sub> sensors using a reference instrument, *Atmospheric Measurement Techniques*, **18**, 4871–4884, 10.5194/amt-18-4871-2025, 2025.
- Chen, H., Winderlich, J., Gerbig, C., Hofer, A., Rella, C. W., Crosson, E. R., Van Pelt, A. D., Steinbach, J., Kolle, O., Beck, V., Daube, B. C., Gottlieb, E. W., Chow, V. Y., Santoni, G. W., and Wofsy, S. C.: High-accuracy continuous airborne measurements of greenhouse gases CO<sub>2</sub> and CH<sub>4</sub> using the cavity ring-down spectroscopy (CRDS) technique, *Atmos. Meas. Tech.*, **3**, 375–386, 10.5194/amt-3-375-2010, 2010.
- 470 Chen, Y., Zhao, H., and Gao, H.: Spatiotemporal Evolution of Air–Sea CO<sub>2</sub> Flux in the South China Sea and Its Response to Environmental Factors, *Remote Sensing*, **16**, 10.3390/rs16244724, 2024.
- Cohen, M. D., Stunder, B. J. B., Rolph, G. D., Draxler, R. R., Stein, A. F., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, **96**, 2059–2077, 10.1175/bams-d-14-00110.1, 2015.
- 475 Costa, M. H., Cunha, L. C. d., Cox, P. M., Eliseev, A. V., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P. K., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S., and Zickfeld, K.: Global Carbon and Other Biogeochemical Cycles and Feedbacks, in: *Climate Change 2021 – The Physical Science Basis*, 673–816, 10.1017/9781009157896.007, 2023.
- Curcoll, R., Morguñ, J. A., Kamnang, A., Cañas, L., Vargas, A., and Grossi, C.: Metrology for low-cost CO<sub>2</sub> sensors applications: the case of a steady-state through-flow (SS-TF) chamber for CO<sub>2</sub> fluxes observations, *Atmos. Meas. Tech.*, **15**, 2807–2818, 10.5194/amt-15-2807-2022, 2022.
- 480 Delaria, E. R., Kim, J., Fitzmaurice, H. L., Newman, C., Wooldridge, P. J., Worthington, K., and Cohen, R. C.: The Berkeley Environmental Air-quality and CO<sub>2</sub> Network: field calibrations of sensor temperature dependence and assessment of network scale CO<sub>2</sub> accuracy, *Atmospheric Measurement Techniques*, **14**, 5487–5500, 10.5194/amt-14-5487-2021, 2021.
- Dubey, R., Telles, A., Nikkel, J., Cao, C., Gewirtzman, J., Raymond, P. A., and Lee, X.: Low-Cost CO(2) NDIR Sensors: Performance Evaluation and Calibration Using Machine Learning Techniques, *Sensors (Basel)*, **24**, 10.3390/s24175675, 2024.
- 485 Fang, S. X., Zhou, L. X., Tans, P. P., Ciais, P., Steinbacher, M., Xu, L., and Luan, T.: In situ measurement of atmospheric CO<sub>2</sub> at the four WMO/GAW stations in China, *Atmospheric Chemistry and Physics*, **14**, 2541–2554, 10.5194/acp-14-2541-2014, 2014.
- Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Armeth, A., Arora, V., Bates, N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R.

A., Ilyina, T., Jain, A. K., Joetzier, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozi, D., Marland, G., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., O'Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., van der Werf, G., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, X., and Zaehle, S.: Global Carbon Budget 2020, *Earth System Science Data*, 12, 3269–3340, 10.5194/essd-12-3269-2020, 2020.

Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le Quééré, C., Luijckx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin, M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O'Brien, K. M., Olsen, A., Omar, A. M., Ono, T., Paulsen, M., Pierrot, D., Pockock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Séférian, R., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tans, P. P., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., van Ooijen, E., Wanninkhof, R., Watanabe, M., Wimart-Rousseau, C., Yang, D., Yang, X., Yuan, W., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2023, *Earth System Science Data*, 15, 5301–5369, 10.5194/essd-15-5301-2023, 2023.

Han, P., Yao, B., Cai, Q., Chen, H., Sun, W., Liang, M., Zhang, X., Zhao, M., Martin, C., Liu, Z., Ye, H., Wang, P., Li, Y., and Zeng, N.: Support Carbon Neutral Goal with a High-Resolution Carbon Monitoring System in Beijing, *Bulletin of the American Meteorological Society*, 105, E2461–E2481, 10.1175/bams-d-23-0025.1, 2024.

Jacobson, A. R., Mikaloff Fletcher, S. E., Gruber, N., Sarmiento, J. L., and Gloor, M.: A joint atmosphere-ocean inversion for surface fluxes of carbon dioxide: 2. Regional results, *Global Biogeochemical Cycles*, 21, 10.1029/2006gb002703, 2007.

Karion, A., Callahan, W., Stock, M., Prinzivalli, S., Verhulst, K. R., Kim, J., Salameh, P. K., Lopez-Coto, I., and Whetstone, J.: Greenhouse gas observations from the Northeast Corridor tower network, *Earth System Science Data*, 12, 699–717, 10.5194/essd-12-699-2020, 2020.

Landschützer, P., Gruber, N., Bakker, D. C. E., Schuster, U., Nakaoka, S., Payne, M. R., Sasse, T. P., and Zeng, J.: A neural network-based estimate of the seasonal to inter-annual variability of the Atlantic Ocean carbon sink, *Biogeosciences*, 10, 7793–7815, 10.5194/bg-10-7793-2013, 2013.

Li, D., Ni, X., Wang, K., Zeng, D., Wang, B., Jin, H., Li, H., Zhou, F., Huang, D., and Chen, J.: Biological CO<sub>2</sub> Uptake and Upwelling Regulate the Air-Sea CO<sub>2</sub> Flux in the Changjiang Plume Under South Winds in Summer, *Frontiers in Marine Science*, 8, 10.3389/fmars.2021.709783, 2021.

Li, Q., Guo, X., Zhai, W., Xu, Y., and Dai, M.: Partial pressure of CO<sub>2</sub> and air-sea CO<sub>2</sub> fluxes in the South China Sea: Synthesis of an 18-year dataset, *Progress in Oceanography*, 182, 10.1016/j.pocean.2020.102272, 2020.

Lin, X., Ciais, P., Bousquet, P., Ramonet, M., Yin, Y., Balkanski, Y., Cozic, A., Delmotte, M., Evangelio, N., Indira, N. K., Locatelli, R., Peng, S., Piao, S., Saunio, M., Swathi, P. S., Wang, R., Yver-Kwok, C., Tiwari, Y. K., and Zhou, L.: Simulating CH<sub>4</sub> and CO<sub>2</sub> over South and East Asia using the zoomed chemistry transport model LMDz-INCA, *Atmospheric Chemistry and Physics*, 18, 9475–9497, 10.5194/acp-18-9475-2018, 2018.

Lv, H., Wang, H., Jiang, Y., Chen, H., Qiao, R., and Wang, Z.: Study on the concentration variation of CO<sub>2</sub> in the background area of Xisha, *Haiyang Xuebao*, 37, 21–30, 10.3969/j.issn.0253-4193.2015.06.003, 2015.

Martin, C. R., Zeng, N., Karion, A., Dickerson, R. R., Ren, X., Turpie, B. N., and Weber, K. J.: Evaluation and environmental correction of ambient CO<sub>2</sub> measurements from a low-cost NDIR sensor, *Atmospheric Measurement Techniques*, 10, 2383–2395, 10.5194/amt-10-2383-2017, 2017.

Nemoto, K., Midorikawa, T., Wada, A., Ogawa, K., Takatani, S., Kimoto, H., Ishii, M., and Inoue, H. Y.: Continuous observations of atmospheric and oceanic CO<sub>2</sub> using a moored buoy in the East China Sea: Variations during the passage of typhoons, *Deep Sea Research Part II: Topical Studies in Oceanography*, 56, 542–553, 10.1016/j.dsr2.2008.12.015, 2009.

540 NOAA GML (2024a): About CO<sub>2</sub> measurements, National Oceanic and Atmospheric Administration, Global Monitoring Laboratory (GML), available at: [https://gml.noaa.gov/ccgg/about/co2\\_measurements.html](https://gml.noaa.gov/ccgg/about/co2_measurements.html), last access: 12 November 2025.

NOAA GML (2024b): Trends in atmospheric carbon dioxide – Mauna Loa, National Oceanic and Atmospheric Administration, Global Monitoring Laboratory (GML), available at: <https://gml.noaa.gov/ccgg/trends/gr.html>, last access: 12 November 2025.

Palter, J. B., Nickford, S., and Mu, L.: Ocean Carbon Dioxide Uptake in the Tailpipe of Industrialized Continents, *Geophysical Research Letters*, 50, 10.1029/2023gl1104822, 2023.

545 Piao, S., He, Y., Wang, X., and Chen, F.: Estimation of China’s terrestrial ecosystem carbon sink: Methods, progress and prospects, *Science China Earth Sciences*, 65, 641–651, 10.1007/s11430-021-9892-6, 2022.

Rhein, M., Rintoul, S. R., Aoki, S., Campos, E., Chambers, D., Feely, R. A., Gulev, S., Johnson, G. C., Josey, S. A., Kostianoy, A., Mauritzen, C., Roemmich, D., Talley, L. D., and Wang, F.: Observations: Ocean, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 255–316, 10.1017/CBO9781107415324.010, 2013.

550 Rödenbeck, C., Conway, T. J., and Langenfelds, R. L.: The effect of systematic measurement errors on atmospheric CO<sub>2</sub> inversions: a quantitative assessment, *Atmos. Chem. Phys.*, 6, 149–161, 10.5194/acp-6-149-2006, 2006.

Rolph, G., Stein, A., and Stunder, B.: Real-time Environmental Applications and Display sYstem: READY, *Environmental Modelling & Software*, 95, 210–228, 10.1016/j.envsoft.2017.06.025, 2017.

Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T.-H., Kozyr, A., Ono, T., and Rios, A. F.: The Oceanic Sink for Anthropogenic CO<sub>2</sub>, *Science*, 305, 367–371, doi:10.1126/science.1097403, 2004.

560 Sarmiento, J. L., Johnson, K. S., Arteaga, L. A., Bushinsky, S. M., Cullen, H. M., Gray, A. R., Hotinski, R. M., Maurer, T. L., Mazloff, M. R., Riser, S. C., Russell, J. L., Schofield, O. M., and Talley, L. D.: The Southern Ocean carbon and climate observations and modeling (SOCCOM) project: A review, *Progress in Oceanography*, 219, 10.1016/j.pocean.2023.103130, 2023.

Shusterman, A. A., Teige, V. E., Turner, A. J., Newman, C., Kim, J., and Cohen, R. C.: The Berkeley Atmospheric CO<sub>2</sub> Observation Network: initial evaluation, *Atmospheric Chemistry and Physics*, 16, 13449–13463, 10.5194/acp-16-13449-2016, 2016.

565 Song, J., Qu, B., Li, X., Yuan, H., and Duan, L.: Estimate of Ocean Carbon Sink: Current Status and Reflections, *Advances in Marine Science*, 41, 577–592, 10.12362/j.issn.1671-6647.20230210001, 2023.

Thoning, K. W., Crowell, A. M., and Mund, J. W.: Atmospheric Carbon Dioxide Dry Air Mole Fractions from continuous measurements at Mauna Loa, Hawaii, Barrow, Alaska, American Samoa, and South Pole, 1973–present [dataset], 10.5194/amt-10-2383-2017, 2025.

570 Wanninkhof, R., Park, G.-H., Takahashi, T., Feely, R. A., Bullister, J. L., and Doney, S. C.: Changes in deep-water CO<sub>2</sub> concentrations over the last several decades determined from discrete pCO<sub>2</sub> measurements, *Deep Sea Research Part I: Oceanographic Research Papers*, 74, 48–63, 10.1016/j.dsr.2012.12.005, 2013.

Wanninkhof, R., Pickers, P. A., Omar, A. M., Sutton, A., Murata, A., Olsen, A., Stephens, B. B., Tilbrook, B., Munro, D., Pierrot, D., Rehder, G., Santana-Casiano, J. M., Müller, J. D., Trinanés, J., Tedesco, K., O’Brien, K., Currie, K., Barbero, L., Telszewski, M., Hoppema, M., Ishii, M., González-Dávila, M., Bates, N. R., Metzl, N., Suntharalingam, P., Feely, R. A., Nakaoka, S.-i., Lauvset, S. K., Takahashi, T., Steinhoff, T., and Schuster, U.: A Surface Ocean CO<sub>2</sub> Reference Network, SOCONET and Associated Marine Boundary Layer CO<sub>2</sub> Measurements, *Frontiers in Marine Science*, 6, 10.3389/fmars.2019.00400, 2019.

575 Xu, J., Cai, S., Xuan, L., Qiu, Y., and Zhu, D.: Study on coastal upwelling in eastern Hainan Island and western Guangdong in summer, 2006, *Acta Oceanologica Sinica*(in Chinese), 35, 11–18, 2013.

580

Xu, S., Park, K., Wang, Y., Chen, L., Qi, D., and Li, B.: Variations in the summer oceanic pCO<sub>2</sub> and carbon sink in Prydz Bay using the self-organizing map analysis approach, *Biogeosciences*, 16, 797–810, 10.5194/bg-16-797-2019, 2019.

Yasuda, T., Yonemura, S., and Tani, A.: Comparison of the characteristics of small commercial NDIR CO<sub>2</sub> sensor models and development of a portable CO<sub>2</sub> measurement device, *Sensors (Basel)*, 12, 3641–3655, 10.3390/s120303641, 2012.

585 Yu, S., Song, Z., Bai, Y., Guo, X., He, X., Zhai, W., Zhao, H., and Dai, M.: Satellite-estimated air-sea CO<sub>2</sub> fluxes in the Bohai Sea, Yellow Sea, and East China Sea: Patterns and variations during 2003–2019, *Science of The Total Environment*, 904, 10.1016/j.scitotenv.2023.166804, 2023.

Zhai, W. D., Dai, M. H., Chen, B. S., Guo, X. H., Li, Q., Shang, S. L., Zhang, C. Y., Cai, W. J., and Wang, D. X.: Seasonal variations of sea–air CO<sub>2</sub> fluxes in the largest tropical marginal sea (South China Sea) based on multiple-year underway measurements, *Biogeosciences*, 10, 7775–7791, 10.5194/bg-10-7775-2013, 2013.

590 Zhang, G., Han, B., Yang, Q., Zhu, X., Wang, X., He, H., Li, H., Wang, X., Xie, W., and Chen, D.: Seamounts Enhance the Local Emission of CO<sub>2</sub> in the Northern South China Sea, *Geophysical Research Letters*, 51, 10.1029/2023gl107264, 2024.